

Consistent Approximations in Composite Optimization

Johannes O. Royset

Operations Research Department

Naval Postgraduate School

joroyset@nps.edu

Abstract. Approximations of optimization problems arise in computational procedures and sensitivity analysis. The resulting effect on solutions can be significant, with even small approximations of components of a problem translating into large errors in the solutions. We specify conditions under which approximations are well behaved in the sense of minimizers, stationary points, and level-sets and this leads to a framework of consistent approximations. The framework is developed for a broad class of composite problems, which are neither convex nor smooth. We demonstrate the framework using examples from stochastic optimization, neural-network based machine learning, distributionally robust optimization, penalty and augmented Lagrangian methods, interior-point methods, homotopy methods, smoothing methods, extended nonlinear programming, difference-of-convex programming, and multi-objective optimization. An enhanced proximal method illustrates the algorithmic possibilities. A quantitative analysis supplements the development by furnishing rates of convergence.

Keywords: set-convergence, epi-convergence, graphical convergence, approximation theory.

Date: January 17, 2022

1 Introduction

A fundamental approach to optimization is to replace an actual problem by an approximating one, which is then solved using an existing algorithm. In sensitivity analysis, the actual problem of interest is also replaced by approximating ones for the purpose of identifying the effect of perturbations. These situations raise the questions: If some portion of an optimization problem is changed and the resulting problem is solved, would the obtained solution be a reasonable approximation of a solution of the actual problem? How large would the solution error be relative to the magnitude of the initial change? These questions are further complicated by the need for considering both optimal and stationary points in the nonconvex setting. Thus, we aspire to construct approximations that have minimizers *and* stationary points near the corresponding points for the actual problem. We may even seek approximating objective functions and feasible sets that are near those of the actual problem in a broad sense.

In this paper, we provide a comprehensive framework for constructing and analyzing approximations of optimization problems with a composite structure. We provide sufficient conditions for *consistent approximations*, which guarantee that the approximating problems eventually become accurate relative to the actual problem in the sense of minimizers, stationary points, and level-sets. Examples from stochastic optimization, neural-network based machine learning, distributionally robust optimization, penalty and augmented Lagrangian methods, interior-point methods, homotopy methods, smoothing

methods, extended nonlinear programming, difference-of-convex programming, and multi-objective optimization demonstrate the framework. The algorithmic possibilities are illustrated by a proximal composite method for solving an array of nonconvex composite problems via a sequence of convex problems. A quantitative analysis supplements the development by furnishing rates of convergence.

We consider the broad class of composite optimization problems in the form

$$\underset{x \in \mathbb{R}^n}{\text{minimize}} \quad \varphi(x) = \iota_X(x) + h(F(x)), \quad (1.1)$$

where the *objective function* $\varphi : \mathbb{R}^n \rightarrow \overline{\mathbb{R}} = \mathbb{R} \cup \{-\infty, \infty\}$ is defined in terms of a nonempty closed set $X \subset \mathbb{R}^n$, often representing “basic” constraints such as bounds on the variables, and a locally Lipschitz continuous mapping $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ that models m quantities of interest with f_1, \dots, f_m being the *component functions*, i.e., $F(x) = (f_1(x), \dots, f_m(x))$. For any set C , $\iota_C(x) = 0$ if $x \in C$ and $\iota_C(x) = \infty$ otherwise. The quantities of interest are combined by a convex function $h : \mathbb{R}^m \rightarrow \overline{\mathbb{R}}$, which might assign an infinite penalty to certain values of $F(x)$. For example, $h = \iota_{\{0\}^m}$ sets $h(F(x)) = \infty$ whenever $F(x)$ deviates from the zero vector and thus encodes the equality constraint $F(x) = 0$. The actual problem (1.1) captures classical nonlinear programming as well as many other problem formulations. In general, φ is nonconvex. The convexity assumption imposed on h is not as restrictive as it might appear and helps us to leverage convexity properties when present.

The study of (1.1) and similar problems goes at least back to [43, 44]. Recent efforts [33, 11, 20, 22, 17, 18, 21, 13] suggest that the problem class might be among the most important ones in the nonconvex setting. Its structural properties are theoretically and computationally attractive, especially to address large-scale problems in machine learning and elsewhere. Nevertheless, the actual problem may not be solvable directly and often needs approximations. The mapping F may be approximated by a smooth or even an affine mapping. It might be specified by integrals, suprema, or other expressions that necessitate imprecise, numerical evaluation. The function h could be nonsmooth and extended real-valued, and then approximated by real-valued functions, piecewise affine or smooth, acting as penalties. The set X might be approximated by a polyhedron or other simplifications. The approximations can be computational motivated or introduced as part of sensitivity, stability, or error analysis.

Our approach is traced back to 1902 when P. Painlevé defined set-convergence. It is now well-known that set-convergence of epigraphs (i.e., epi-convergence) of approximating functions and set-convergence of graphs (i.e., graphical convergence) of approximating set-valued mappings furnish the desired convergence guarantees about the solutions of the corresponding minimization problems and generalized equations; see [48, Chap. 4, 5, and 7]. If the set-valued mappings represent optimality conditions for the approximating problems, then graphical convergence translates into convergence of stationary points. We define approximating problems and their optimality conditions as being *consistent approximations* of the actual problem (1.1), paired with an optimality condition, when they exhibit such epi-convergence and graphical convergence.

In the convex setting, epi-convergence of approximating functions φ^ν to the actual objective function φ suffices for the subgradient mappings $\partial\varphi^\nu$ to graphically converge to $\partial\varphi$ by Attouch’s theorem; see for example [48, Thm. 12.35]. Thus, epi-convergence of the approximating functions ensures that the approximations are consistent. In the nonconvex case, one explicitly needs to impose conditions related

to the optimality conditions to guarantee convergence of stationary points and this is recognized by the added requirement about graphical convergence of set-valued mappings in the definition of consistent approximations. If the approximating functions φ^ν are smooth, their epi-convergence to φ does provide some relation between $\nabla\varphi^\nu$ and $\partial\varphi$ but in a less-than-ideal “outer” sense; cf. [10, Lem. 3.4] and [48, Cor. 8.47]. A far-reaching extension of Attouch’s theorem for nonconvex functions is provided by [42], which implies in the context of (1.1) that epi-convergence of certain approximating functions h^ν to h combined with a smooth F and a constraint qualification suffice for graphical convergence in a local sense of the resulting subgradient mappings [11].

Approximations of (1.1) may stem from the smoothing of h and/or F . The specific function $\gamma \mapsto \max\{0, \gamma\}$ can be approximated by smooth functions via convolution [15]. The approximations exhibit both epi-convergence as well as certain convergence of gradients to the subgradients of the max-function [14]; this reference includes many examples of such smoothing. Further developments in this direction is furnished by [10, 11], where in the context of (1.1), h is approximated by a smooth function constructed using inf-convolution; see also [12] for details about smoothing of finite max-functions.

In contrast to these efforts, we consider broad sets of approximations; X , h , and F in (1.1) may all be approximated and not necessarily smoothly. In sensitivity and error analysis, nonsmooth approximations are especially relevant as computational concerns tend to be secondary. We also deviate from the focus on the subgradient mapping of φ and that of its approximations, and instead express optimality conditions using (generalized) *multipliers*, which provide additional flexibility for absorbing inaccuracies. Thus, consistent approximations emerge as widely available under mild assumptions.

There is an extensive literature on general approximations and related stability analysis; see the monographs [40, 48, 7, 35, 16] as well as efforts based on metric regularity and calmness [29, 38], tilt-stability [23, 34, 19], full-stability [36], convergence of abstract iteration schemes [30], and the truncated Hausdorff distance [1, 2, 3, 50, 51]. The latter approach is closely related to the present paper, with direct relevance to our rate of convergence analysis.

Our choice of the term “consistent approximations” is motivated by E. Polak’s concept in [40, Chap. 3-4]. There epi-convergence of approximating objective functions combined with an outer epi-limit of approximating optimality functions (after a sign change) are defined as consistent; see [49, 52, 25, 39, 54] for applications in stochastic optimization, semi-infinite programming, nonsmooth optimization, and optimal control. Since optimality functions are essentially gap-functions of generalized equations representing optimality conditions, Polak’s concept is, roughly, equivalent to weak consistency as we define it below. A related concept based on epigraphical nesting of directional derivatives is defined in [27]. We deviate from these earlier developments by viewing optimality conditions as generalized equations defined by set-valued mappings and thus bypass the need for defining optimality functions. This brings into play well-developed calculus rules and computational procedures for generalized equations. The flexibility of this approach is demonstrated by a set of sufficient conditions for consistency (Theorem 2.4). An introductory treatment of consistent approximations under the assumption of a smooth mapping F is, in parallel, provided by the forthcoming textbook [53]. While we limit the discussion to finite-dimensional problems, many of the concepts can be extended; see [39, 32] for such efforts.

We continue in Section 2 by stating the approximating problems, defining consistency, and speci-

fying sufficient conditions. Section 3 furnishes 12 examples. Section 4 describes a specific algorithm. Section 5 applies the framework to machine learning problems. Section 6 discusses rates of convergence.

Terminology. A ball under norm $\|\cdot\|$ is denoted by $\mathbb{B}(\bar{x}, \rho) = \{x \in \mathbb{R}^n \mid \|x - \bar{x}\| \leq \rho\}$. A function $f : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$ has a *domain* $\text{dom } f = \{x \in \mathbb{R}^n \mid f(x) < \infty\}$ and an *epigraph* $\text{epi } f = \{(x, \alpha) \in \mathbb{R}^{n+1} \mid f(x) \leq \alpha\}$. The function is *lower semicontinuous* (lsc) if $\text{epi } f$ is closed and is *convex* if $\text{epi } f$ is convex. It is *proper* if $\text{epi } f$ is nonempty and $f(x) > -\infty$ for all $x \in \mathbb{R}^n$. It is *continuous relative to* $C \subset \mathbb{R}^n$ if $x^\nu \in C \rightarrow x \in C$ implies that $f(x^\nu) \rightarrow f(x)$. It is *locally Lipschitz continuous* (lLc) at \bar{x} when there are $\delta \in (0, \infty)$ and $\kappa \in [0, \infty)$ such that $|f(x) - f(x')| \leq \kappa \|x - x'\|_2$ whenever $x, x' \in \mathbb{B}(\bar{x}, \delta)$. If f is lLc at every $\bar{x} \in \mathbb{R}^n$, then f is lLc. A mapping $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is lLc (at \bar{x}) if its component functions f_1, \dots, f_m are lLc (at \bar{x}). Functions and mappings are *smooth* (at \bar{x}) if they are continuously differentiable (at \bar{x}). Moreover, $\inf f = \inf\{f(x) \mid x \in \mathbb{R}^n\}$ and ε -argmin $f = \{x \in \text{dom } f \mid f(x) \leq \inf f + \varepsilon\}$, which is simply written as $\text{argmin } f$ if $\varepsilon = 0$. A *lower level-set* $\{f \leq \alpha\} = \{x \in \mathbb{R}^n \mid f(x) \leq \alpha\}$. The *convex hull* of a set C is denoted by $\text{con } C$. The j th component of a vector $x \in \mathbb{R}^n$ is typically indicated by x_j . We let $\text{bdry } C$ be the *boundary* of a set $C \subset \mathbb{R}^n$, i.e., $\text{bdry } C = C \setminus \text{int } C$, where $\text{int } C$ is the interior of C . We adopt the usual rules for extended arithmetic, including $\infty - \infty = \infty$.

Sequences of points, sets, and so forth are usually indexed by superscript $\nu \in \mathbb{N} = \{1, 2, \dots\}$. The collection of subsequences of \mathbb{N} is denoted by $\mathcal{N}_\infty^\#$, with convergence of $\{x^\nu, \nu \in \mathbb{N}\}$ to x along a subsequence $N \in \mathcal{N}_\infty^\#$ being denoted by $x^\nu \xrightarrow{N} x$. The symbols \exists and \forall mean “there exist” and “for all,” respectively. The *inner limit* of a sequence of sets $\{C^\nu \subset \mathbb{R}^n, \nu \in \mathbb{N}\}$ is $\text{LimInn } C^\nu = \{x \in \mathbb{R}^n \mid \exists x^\nu \in C^\nu \rightarrow x\}$. The *outer limit* is $\text{LimOut } C^\nu = \{x \in \mathbb{R}^n \mid \exists N \in \mathcal{N}_\infty^\# \text{ and } x^\nu \in C^\nu \xrightarrow{N} x\}$. Thus, C^ν *set-converges* to $C \subset \mathbb{R}^n$, denoted by $C^\nu \xrightarrow{s} C$, if $\text{LimOut } C^\nu \subset C \subset \text{LimInn } C^\nu$. The functions $\{f^\nu : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}, \nu \in \mathbb{N}\}$ *epi-converge* to $f : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$, denoted by $f^\nu \xrightarrow{e} f$, if $\text{epi } f^\nu \xrightarrow{s} \text{epi } f$. This takes place if and only if

$$\forall x^\nu \rightarrow x, \liminf f^\nu(x^\nu) \geq f(x) \quad (1.2)$$

$$\forall x, \exists x^\nu \rightarrow x \text{ with } \limsup f^\nu(x^\nu) \leq f(x). \quad (1.3)$$

A set-valued mapping $T : \mathbb{R}^n \rightrightarrows \mathbb{R}^m$ has subsets of \mathbb{R}^m as its “values” and a *graph* written as $\text{gph } T = \{(x, y) \in \mathbb{R}^{n+m} \mid y \in T(x)\}$. A sequence of set-valued mappings $\{T^\nu : \mathbb{R}^n \rightrightarrows \mathbb{R}^m, \nu \in \mathbb{N}\}$ *converges graphically* to $T : \mathbb{R}^n \rightrightarrows \mathbb{R}^m$, denoted by $T^\nu \xrightarrow{g} T$, if $\text{gph } T^\nu \xrightarrow{s} \text{gph } T$. The set of solutions to the generalized equation $v \in T(x)$ is written as $T^{-1}(v) = \{x \mid v \in T(x)\}$.

We denote the normal cone to $C \subset \mathbb{R}^n$ at $x \in C$ by $N_C(x)$ and let $N_C(x) = \emptyset$ for $x \notin C$. Likewise, the set of subgradients of $f : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$ at a point x with $f(x) \in \mathbb{R}$ is $\partial f(x)$; $\partial f(x) = \emptyset$ when $f(x) \notin \mathbb{R}$. These quantities are understood in the general (Mordukhovich) sense; see [48, Chap. 6 and 8].

2 Consistent Approximations

Parallel to the actual problem (1.1), we defined the *approximating problems*

$$\left\{ \begin{array}{l} \text{minimize } \varphi^\nu(x) = \iota_{X^\nu}(x) + h^\nu(F^\nu(x)), \\ x \in \mathbb{R}^n \end{array} \quad \nu \in \mathbb{N} \right\} \quad (2.1)$$

where $X^\nu \subset \mathbb{R}^n$, $h^\nu : \mathbb{R}^m \rightarrow \overline{\mathbb{R}}$, and $F^\nu : \mathbb{R}^n \rightarrow \mathbb{R}^m$ are approximations of the corresponding components of the actual problem. We seek to confirm that small discrepancies between the respective components of the approximating and actual problems indeed translate into small errors in the solutions of the approximating problems relative to those of the actual problem. It is clear that pointwise convergence $\varphi^\nu(x) \rightarrow \varphi(x)$ for all $x \in \mathbb{R}^n$ would not suffice; see [48, Figure 7-1]. Uniform convergence of φ^ν to φ on \mathbb{R}^n ensures that cluster points of $\{x^\nu \in \operatorname{argmin} \varphi^\nu, \nu \in \mathbb{N}\}$ indeed are minimizers of φ , but such a requirement is too stringent, especially when X^ν is different than X or $\operatorname{dom} h^\nu$ is different than $\operatorname{dom} h$. We also would like to address the convergence of stationary points for the approximating problems to those of the actual problem and then neither pointwise nor uniform convergence of φ^ν to φ is sufficient.

We make the following *universal* assumptions for the remainder of the paper:

$$X, X^\nu \text{ are nonempty and closed; } \quad h, h^\nu \text{ are proper, lsc, and convex; } \quad F, F^\nu \text{ are lLc.}$$

This limits the scope to a well-structured class of problems for which convenient calculus rules can be brought in, while still addressing a vast array of applications. In particular, [48, Thm. 10.46, Exer. 10.52] lead to optimality conditions for the actual and approximating problems.

If both h and F were smooth, then $0 \in \nabla F(x)^\top \nabla h(F(x)) + N_X(x)$ is a necessary condition for a local minimizer of φ by [48, Thm. 6.12]. Here, $\nabla F(x)$ is the Jacobian matrix of F at x with rows corresponding to the gradients of f_1, \dots, f_m at x . This condition is equivalently written as

$$0 = F(x) - z, \quad 0 = \nabla h(z) - y, \quad 0 \in \sum_{i=1}^m y_i \nabla f_i(x) + N_X(x),$$

where $y = (y_1, \dots, y_m) \in \mathbb{R}^m$ and $z \in \mathbb{R}^n$ are *multiplier vectors*. This condition naturally extends to nonsmooth h and F by essentially “replacing” gradients by subgradients. Thus, we express optimality conditions for the actual and approximating problems concisely as the generalized equations

$$0 \in S(x, y, z) \quad \text{and} \quad 0 \in S^\nu(x, y, z)$$

using the set-valued mappings $S, S^\nu : \mathbb{R}^{n+2m} \rightrightarrows \mathbb{R}^{2m+n}$ given by

$$\begin{aligned} S(x, y, z) &= \{F(x) - z\} \times \{\partial h(z) - y\} \times \left(\sum_{i=1}^m y_i \operatorname{con} \partial f_i(x) + N_X(x) \right) \\ S^\nu(x, y, z) &= \{F^\nu(x) - z\} \times \{\partial h^\nu(z) - y\} \times \left(\sum_{i=1}^m y_i \operatorname{con} \partial f_i^\nu(x) + N_{X^\nu}(x) \right). \end{aligned}$$

If x satisfies $0 \in S(x, y, z)$ for some $y \in \mathbb{R}^m$ and $z \in \mathbb{R}^n$, then x is a *stationary point* for the actual problem, with similar terminology being adopted for the approximating problems. The set of subgradients $\partial f_i(x)$ is convex when f_i is epi-regular¹ at x , which, for example, is the case if f_i is smooth at x or if f_i is convex [48, Thm. 8.30]. Under such circumstances, “con” in the expression for S is superfluous. Since f_i is lLc, $\operatorname{con} \partial f_i(x)$ coincides with the set of Clarke subgradients [48, Thm. 8.49] of f_i at x . Aligned with the approach in [48], we shy away from this terminology. We view the convexification in the optimality condition as stemming from a slight relaxation of otherwise valid optimality conditions and resulting in an additional “buffer” to absorb approximations.

¹A function f is epi-regular at x if its epigraph is Clarke regular [48, Def. 6.4] at $(x, f(x))$.

The optimality conditions generalize many familiar ones. For example, $0 \in S(x, y, z)$ if and only if the KKT condition hold for the actual problem with $X = \mathbb{R}^n$, $h(z) = z_1 + \sum_{i=2}^s \iota_{\{0\}}(z_i) + \sum_{i=s+1}^m \iota_{(-\infty, 0]}(z_i)$, and F being smooth. We see this by deriving $N_X(x) = \{0\}$, $\partial h(z) = \{1\} \times N_{\{0\}^{s-1}}(z_2, \dots, z_s) + N_{(-\infty, 0]^{m-s}}(z_{s+1}, \dots, z_m)$, and

$$\sum_{i=1}^m y_i \operatorname{con} \partial f_i(x) \} + N_X(x) = \sum_{i=1}^m y_i \nabla f_i(x)$$

so that $y_1 = 1$, y_2, \dots, y_s are unrestricted in sign, and, for $i = s + 1, \dots, m$, $y_i = 0$ if $f_i(x) < 0$ and $y_i \geq 0$ if $f_i(x) = 0$. Since even the KKT condition requires a constraint qualification (such as the Mangasarian-Fromovitz), it is not surprising that $0 \in S(x, y, z)$ likewise requires a qualification for it to be a necessary optimality condition. This is formalized in the next proposition.

2.1 Proposition (optimality condition). *Suppose that the following qualification holds at x^* :*

$$y \in N_{\operatorname{dom} h}(F(x^*)) \quad \text{and} \quad 0 \in \sum_{i=1}^m y_i \operatorname{con} \partial f_i(x^*) + N_X(x^*) \quad \implies \quad y = 0. \quad (2.2)$$

If x^ is a local minimizer of (1.1), then $0 \in S(x^*, y^*, z^*)$ for some $y^* \in \mathbb{R}^m$ and $z^* \in \mathbb{R}^n$.*

Proof. Guided by [48, Exer. 10.52], this follows straightforwardly when recalling that $l_F(x, y) = \langle F(x), y \rangle$ implies

$$\partial_x l_F(x, y) \subset \sum_{i=1}^m \partial(y_i f_i)(x) \subset \sum_{i=1}^m \operatorname{con} \{ \partial(y_i f_i)(x) \} = \sum_{i=1}^m y_i \operatorname{con} \partial f_i(x)$$

via [48, Cor. 10.9, Thm. 9.61]. □

The proposition likewise confirms that $0 \in S^\nu(x, y, z)$ is a necessary optimality condition for an approximating problem under a qualification parallel to (2.2). We are now in a position to state what it means for the approximating problems, paired with their optimality conditions, to be well-justified surrogates of the actual problem and its optimality condition.

2.2 Definition (consistent approximations). *The pairs $\{(\varphi^\nu, S^\nu), \nu \in \mathbb{N}\}$ are consistent approximations of (φ, S) when*

$$\varphi^\nu \xrightarrow{e} \varphi \quad \text{and} \quad S^\nu \xrightarrow{g} S.$$

If the graphical convergence is relaxed to merely $\operatorname{LimOut}(\operatorname{gph} S^\nu) \subset \operatorname{gph} S$, then $\{(\varphi^\nu, S^\nu), \nu \in \mathbb{N}\}$ are weakly consistent approximations of (φ, S) .

Under consistency, even in its weak form, we are on solid ground because the approximating problems indeed produce approximating solutions of the actual problem as formalized next using the following notation. For $T : \mathbb{R}^n \rightrightarrows \mathbb{R}^m$, the set of δ -solutions to the generalized equation $0 \in T(x)$ is defined as

$$T^{-1}(\mathbb{B}(0, \delta)) = \bigcup_{v \in \mathbb{B}(0, \delta)} T^{-1}(v) = \bigcup_{v \in \mathbb{B}(0, \delta)} \{x \in \mathbb{R}^n \mid v \in T(x)\}.$$

Thus, a δ -solution $\hat{x} \in T^{-1}(\mathbb{B}(0, \delta))$ “almost” satisfies $0 \in T(x)$ in the sense that $v \in T(\hat{x})$ for some v with $\|v\| \leq \delta$. We observe that such near-solutions may depend on the choice of norm $\|\cdot\|$.

2.3 Proposition (consequences of consistency). *Suppose that $\{(\varphi^\nu, S^\nu), \nu \in \mathbb{N}\}$ are weakly consistent approximations of (φ, S) , the tolerances ε^ν and δ^ν vanish, and $\alpha < \beta$. Then, the following hold:*

- (a) $\text{LimOut}(\varepsilon^\nu\text{-argmin } \varphi^\nu) \subset \text{argmin } \varphi$, provided that $\text{dom } \varphi \neq \emptyset$.
- (b) $\text{LimOut}\{\varphi^\nu \leq \alpha\} \subset \{\varphi \leq \alpha\} \subset \text{LimInn}\{\varphi^\nu \leq \beta\}$.
- (c) $\text{LimOut}(S^\nu)^{-1}(\mathbb{B}(0, \delta^\nu)) \subset S^{-1}(0)$.

Moreover, if the approximations are consistent (and not merely weakly consistent), then there are $\gamma^\nu \rightarrow 0$ such that $\text{LimInn}(S^\nu)^{-1}(\mathbb{B}(0, \gamma^\nu)) \supset S^{-1}(0)$.

Proof. This is a compilation of well-known facts; see [48, Thm. 7.31] for (a); [48, Prop. 7.7] for (b); and [48, Thm. 5.37] for (c) and the final statement. \square

Consistency indeed guarantees a comprehensive sense of approximation tailored to the need of minimization problems. From (a) in the proposition, we see that every cluster point of a sequence of near-minimizers of the approximating problems is a minimizer of the actual problem provided that the tolerances vanish. That is, if $x^\nu \in \varepsilon^\nu\text{-argmin } \varphi^\nu \xrightarrow{N} \bar{x}$ along a subsequence $N \in \mathcal{N}_\infty^\#$, then $\bar{x} \in \text{argmin } \varphi$. This a valuable property, but remains somewhat conceptual in nonconvex settings because there might be no practical algorithm for computing near-minimizers of the approximating problems. Item (b) shows that the lower level-sets of φ^ν eventually become essentially indistinguishable from those of φ . For example, if $\varphi^\nu(x^\nu) \leq \alpha$ and $x^\nu \rightarrow \bar{x}$, then $\varphi(\bar{x}) \leq \alpha$ by the first inclusion in (b). The second inclusion in (b) establishes that any \bar{x} with $\varphi(\bar{x}) \leq \alpha$ can be approached by points x^ν with $\varphi^\nu(x^\nu) \leq \beta$. Thus, φ^ν accurately “predicts” the values of φ as $\nu \rightarrow \infty$.

Item (c) confirms that any cluster point $(\bar{x}, \bar{y}, \bar{z})$ of a sequence $\{(x^\nu, y^\nu, z^\nu), \nu \in \mathbb{N}\}$, with $0 \in S^\nu(x^\nu, y^\nu, z^\nu)$, satisfies $0 \in S(\bar{x}, \bar{y}, \bar{z})$. This holds even if the points (x^ν, y^ν, z^ν) are computed with tolerances δ^ν as long as they vanish. The approximating problems and their optimality conditions are often constructed such that computing nearly stationary points are indeed possible in finite time; see Section 3 for examples. Combining the facts from (b) and (c), we summarize: If $N \in \mathcal{N}_\infty^\#$ and $(x^\nu, y^\nu, z^\nu) \in (S^\nu)^{-1}(\mathbb{B}(0, \delta^\nu)) \xrightarrow{N} (\bar{x}, \bar{y}, \bar{z})$, then

$$(\bar{x}, \bar{y}, \bar{z}) \in S^{-1}(0) \quad \text{and} \quad \varphi(\bar{x}) \leq \liminf_{\nu \in N} \varphi^\nu(x^\nu).$$

Thus, \bar{x} is stationary for the actual problem, with an objective function value at least as good as predicted by the approximations.

These consequences occur under *weak* consistency. Passing to consistency, we achieve enhanced robustness in the sense that every stationary point of the actual problem can be approached by nearly stationary points of the approximating problems. In fact, the proposition guarantees that for some tolerances $\gamma^\nu \rightarrow 0$, one has $(S^\nu)^{-1}(\mathbb{B}(0, \gamma^\nu)) \xrightarrow{s} S^{-1}(0)$.

The proposition justifies the following broad algorithmic framework for solving the actual problem.

Consistent Approximation Algorithm.

Data. $\delta^\nu \geq 0$, with $\delta^\nu \rightarrow 0$.

Step 0. Set $\nu = 1$.

Step 1. Minimize $h^\nu \circ F^\nu$ over X^ν until one obtains x^ν , with corresponding (y^ν, z^ν) , satisfying

$$(x^\nu, y^\nu, z^\nu) \in (S^\nu)^{-1}(\mathbb{B}(0, \delta^\nu)).$$

Step 2. Replace ν by $\nu + 1$ and go to Step 1.

A multitude of implementation details remain unsettled. What approximations are most suitable for the actual problem at hand? What subroutine should be used in Step 1 and with what δ^ν ? We list many examples of approximations in Section 3 and describe a concrete algorithm in Section 4.

Verification of consistency is supported by sufficient conditions for epi-convergence of φ^ν to φ and graphical convergence of S^ν to S . There are several known results. For example, if $X = X^\nu = \mathbb{R}^n$, $F = F^\nu$ are smooth with ∇F having rank m , and $h^\nu \xrightarrow{e} h$, then $\varphi^\nu \xrightarrow{e} \varphi$ by [10, Thm. 3.2]; see also [48, Chap. 5 and 7]. The following conditions appear especially versatile by allowing for approximations of X , h , and F under relatively mild assumptions.

2.4 Theorem (sufficient conditions for consistent approximations). *Suppose that $h^\nu \xrightarrow{e} h$, $X^\nu \xrightarrow{s} X$ with these sets being convex, and, for $i = 1, \dots, m$, one has the property:*

$$\left. \begin{array}{l} x^\nu \in X^\nu \rightarrow x \in X \\ v^\nu \in \partial f_i^\nu(x^\nu) \end{array} \right\} \implies \left\{ \begin{array}{l} f_i^\nu(x^\nu) \rightarrow f_i(x) \\ \{v^\nu, \nu \in \mathbb{N}\} \text{ is bounded with all its cluster points in } \text{con } \partial f_i(x). \end{array} \right. \quad (2.3)$$

Then, each one of the following conditions is sufficient for $\{(\varphi^\nu, S^\nu), \nu \in \mathbb{N}\}$ to be weakly consistent approximations of (φ, S) :

- (a) (real-valuedness): h is real-valued.
- (b) (pointwise convergence): $h^\nu(z) \rightarrow h(z)$ for all $z \in \text{bdry}(\text{dom } h)$ and, in addition, $F^\nu(x) = F(x)$ for all $x \in X$ and $X \subset X^\nu$ for each ν .
- (c) (monotonicity): For each ν : $h^\nu(z) \leq h^\nu(\bar{z})$ when $z \leq \bar{z}$, $h^\nu(z) \leq h(z)$ for all z , $X \subset X^\nu$, and $F^\nu(x) \leq F(x)$ for all $x \in X$.
- (d) (interior points): h is continuous relative to $\text{dom } h$, $X \subset X^\nu$ for each ν , and for all $x \in X$ with $F(x) \in \text{bdry}(\text{dom } h)$, there is $x^\nu \in X \rightarrow x$ such that $F(x^\nu) \in \text{int}(\text{dom } h)$.
- (e) (separability): For $\sum_{k=1}^r m_k = m$, $\text{llc } F_k^\nu, F_k : \mathbb{R}^n \rightarrow \mathbb{R}^{m_k}$, and proper, lsc, and convex $h_k^\nu, h_k : \mathbb{R}^{m_k} \rightarrow \overline{\mathbb{R}}$, $k = 1, \dots, r$, one can express $F^\nu(x) = (F_1^\nu(x), \dots, F_r^\nu(x))$, $F(x) = (F_1(x), \dots, F_r(x))$, $h^\nu(z) = \sum_{k=1}^r h_k^\nu(z_k)$, and $h(z) = \sum_{k=1}^r h(z_k)$ with h_k being either real-valued or h_k satisfying $h_k^\nu(z_k) \rightarrow h_k(z_k)$ for all $z_k \in \text{bdry}(\text{dom } h_k)$ and $F_k^\nu(x) = F_k(x)$ for all $x \in X$. Moreover, $X \subset X^\nu$.

We also have the refinements:

If $X^\nu = X$, then each of (a)-(e) remains sufficient without the convexity assumption on X .

If $f_i^\nu = f_i$, then (2.3) holds automatically.

If $X^\nu = X$ and $f_i^\nu = f_i$, then each of (a)-(e) is sufficient for consistent approximations.

If f_i^ν, f_i are smooth, then (2.3) is equivalent to $f_i^\nu(x^\nu) \rightarrow f_i(x)$ and $\nabla f_i^\nu(x^\nu) \rightarrow \nabla f_i(x)$ as $x^\nu \in X^\nu \rightarrow x$.

If f_i^ν, f_i are smooth, then each of (a)-(e) is sufficient for consistent approximations.

Proof. We start by establishing $\varphi^\nu \xrightarrow{s} \varphi$. To prove the liminf-condition (1.2) for φ^ν and φ , let $x^\nu \rightarrow x$. Since neither $\iota_X(x)$ nor $h(F(x))$ is equal to $-\infty$,

$$\liminf \left(\iota_{X^\nu}(x^\nu) + h^\nu(F^\nu(x^\nu)) \right) \geq \liminf \iota_{X^\nu}(x^\nu) + \liminf h^\nu(F^\nu(x^\nu)).$$

Certainly, $X^\nu \xrightarrow{s} X$ implies that $\iota_{X^\nu} \xrightarrow{s} \iota_X$ and then also $\liminf \iota_{X^\nu}(x^\nu) \geq \iota_X(x)$. If $x^\nu \notin X^\nu$, then $\iota_{X^\nu}(x^\nu) = \infty$. Thus, we assume without loss of generality that $x^\nu \in X^\nu$. This implies that $x \in X$ because $X^\nu \xrightarrow{s} X$. By assumption, $F^\nu(x^\nu) \rightarrow F(x)$ and $h^\nu \xrightarrow{s} h$. These facts ensure that $\liminf h^\nu(F^\nu(x^\nu)) \geq h(F(x))$. When combined with the earlier inequalities, this relation establishes the liminf-condition (1.2) for φ^ν and φ , which thus holds without any of the additional conditions (a)-(e).

Next, we establish the limsup-condition (1.3) for φ^ν and φ . Let $x \in \mathbb{R}^n$ be arbitrary. If $x \notin X$ or $F(x) \notin \text{dom } h$, then $\varphi(x) = \infty$ and the limsup-condition holds trivially. If $x \in X$ and $F(x) \in \text{dom } h$, then we argue as follows:

First, suppose that (a) holds. Since $X^\nu \xrightarrow{s} X$, there exists $x^\nu \in X^\nu \rightarrow x$. Moreover, h^ν converges uniformly to h on compact sets because h is convex and real-valued; see [48, Thm. 7.17]. Consequently, $h^\nu(F^\nu(x^\nu)) \rightarrow h(F(x))$ and one has

$$\limsup \left(\iota_{X^\nu}(x^\nu) + h^\nu(F^\nu(x^\nu)) \right) \leq \limsup \iota_{X^\nu}(x^\nu) + \limsup h^\nu(F^\nu(x^\nu)) = h(F(x)).$$

Second, suppose that (c) holds. Since $X \subset X^\nu$, we construct $\{x^\nu = x, \nu \in \mathbb{N}\}$ and simply need to prove that $\limsup h^\nu(F^\nu(x)) \leq h(F(x))$. This holds because $F^\nu(x) \leq F(x)$ and then $h^\nu(F^\nu(x)) \leq h^\nu(F(x)) \leq h(F(x))$ for all $\nu \in \mathbb{N}$.

Third, suppose that (d) holds. We consider two cases. Suppose that $F(x) \in \text{int}(\text{dom } h)$. Then, construct $\{x^\nu = x, \nu \in \mathbb{N}\}$ and note that $h^\nu(F^\nu(x)) \rightarrow h(F(x))$ because h^ν converges uniformly to h on bounded subsets of the interior of $\text{dom } h$; see [48, Thm. 7.17]. Next, suppose that $F(x) \in \text{bdry}(\text{dom } h)$. Then, by condition (d), there are $\bar{x}^k \in X \rightarrow x$ with $F(\bar{x}^k) \in \text{int}(\text{dom } h)$. Consequently, there exist $\delta_k \in (0, \infty) \rightarrow 0$ such that $\mathbb{B}(F(\bar{x}^k), \delta_k) \subset \text{int}(\text{dom } h)$ for all k . Let $\varepsilon_k \in (0, \infty) \rightarrow 0$. Fix k . Since $\bar{x}^k \in X \cap X^\nu$, $F^\nu(\bar{x}^k) \rightarrow F(\bar{x}^k)$ as $\nu \rightarrow \infty$. Let $\nu_k \in \mathbb{N}$, with $\nu_k \geq k$, be such that $F^\nu(\bar{x}^k) \in \mathbb{B}(F(\bar{x}^k), \delta_k)$ for all $\nu \geq \nu_k$. Since h^ν converges uniformly to h on $\mathbb{B}(F(\bar{x}^k), \delta_k)$ by virtue of that set being in the interior of $\text{dom } h$ (again cf. [48, Thm. 7.17]), there is $\bar{\nu}_k \geq \nu_k$ such that $h^\nu(F^\nu(\bar{x}^k)) \leq h(F(\bar{x}^k)) + \varepsilon_k$ for all $\nu \geq \bar{\nu}_k$. We repeat these arguments for all $k \in \mathbb{N}$. Now, we construct $x^\nu = \bar{x}^1$ for $\nu = 1, \dots, \bar{\nu}_2$ and $x^\nu = \bar{x}^k$ for $\bar{\nu}_k < \nu \leq \bar{\nu}_{k+1}$. Then, $x^\nu \rightarrow x$ and $h^\nu(F^\nu(x^\nu)) \leq h(F(x^\nu)) + \varepsilon_k$ for all $\bar{\nu}_k < \nu \leq \bar{\nu}_{k+1}$. As $\nu \rightarrow \infty$, k tends to ∞ as well in this expression. Thus, its right-hand side converges to $h(F(x))$; recall that h is continuous on its domain. We have shown that $\limsup h^\nu(F^\nu(x^\nu)) \leq h(F(x))$.

Fourth, suppose that (e) holds. Since $X \subset X^\nu$, one has $x \in X^\nu$ for all ν and we construct $\{x^\nu = x, \nu \in \mathbb{N}\}$. Certainly, $F^\nu(x) \rightarrow F(x)$ and

$$\limsup \left(\iota_{X^\nu}(x^\nu) + \sum_{k=1}^r h_k^\nu(F_k^\nu(x^\nu)) \right) \leq \limsup \iota_{X^\nu}(x) + \sum_{k=1}^r \limsup h_k^\nu(F_k^\nu(x)).$$

Suppose that h_k is real-valued. Then, h_k^ν converges uniformly to h_k on compact sets because h_k is convex; see [48, Thm. 7.17]. This implies that $h_k^\nu(F_k^\nu(x)) \rightarrow h_k(F_k(x))$. Alternatively, suppose that $h_k^\nu(z_k) \rightarrow h_k(z_k)$ for all $z_k \in \text{bdry}(\text{dom } h_k)$ and $F_k^\nu(x') = F_k(x')$ for all $x' \in X$. Since $F(x) \in \text{dom } h$, we have $F_k(x) \in \text{dom } h_k$. If $F_k(x) \in \text{bdry}(\text{dom } h_k)$, then, by assumption, $h_k^\nu(F_k(x)) \rightarrow h_k(F_k(x))$. If $F_k(x) \in \text{int}(\text{dom } h_k)$, then $h_k^\nu(F_k(x)) \rightarrow h_k(F_k(x))$ as well because these functions are convex. In summary,

$$\sum_{k=1}^r \limsup h_k^\nu(F_k^\nu(x)) \rightarrow \sum_{k=1}^r h_k(F_k(x))$$

and the needed limsup-condition holds.

Fifth, suppose that (b) holds. This is a special case of (e) with $r = 1$.

We have confirmed that each one of (a)-(e) suffices for $\varphi^\nu \xrightarrow{e} \varphi$ and, in fact, this holds even without the convexity of X^ν and X .

We next turn to the relation between $\text{gph } S^\nu$ and $\text{gph } S$ and start by showing $\text{LimOut}(\text{gph } S^\nu) \subset \text{gph } S$. Let $(\bar{x}, \bar{y}, \bar{z}, \bar{u}, \bar{v}, \bar{w}) \in \text{LimOut}(\text{gph } S^\nu)$. Then, there are $N \in \mathcal{N}_\infty^\#$, $x^\nu \xrightarrow{N} \bar{x}$, $y^\nu \xrightarrow{N} \bar{y}$, $z^\nu \xrightarrow{N} \bar{z}$, $u^\nu \xrightarrow{N} \bar{u}$, $v^\nu \xrightarrow{N} \bar{v}$, and $w^\nu \xrightarrow{N} \bar{w}$ with $(x^\nu, y^\nu, z^\nu, u^\nu, v^\nu, w^\nu) \in \text{gph } S^\nu$. Consequently,

$$u^\nu = F^\nu(x^\nu) - z^\nu, \quad v^\nu \in \partial h^\nu(z^\nu) - y^\nu, \quad w^\nu \in \sum_{i=1}^m y_i^\nu \text{con } \partial f_i^\nu(x^\nu) + N_{X^\nu}(x^\nu).$$

The existence of w^ν implies that $x^\nu \in X^\nu$ because otherwise $N_{X^\nu}(x^\nu)$ would have been an empty set. Since $X^\nu \xrightarrow{s} X$, we also have $\bar{x} \in X$. Thus, $F^\nu(x^\nu) \xrightarrow{N} F(\bar{x})$ and we conclude that $\bar{u} = F(\bar{x}) - \bar{z}$. Attouch's theorem [48, Thm. 12.35] states that $h^\nu \xrightarrow{e} h$ implies $\partial h^\nu \xrightarrow{a} \partial h$. Consequently, $\bar{v} \in \partial h(\bar{z}) - \bar{y}$.

The inclusion for w^ν implies that there exist

$$a_i^\nu \in \text{con } \partial f_i^\nu(x^\nu), \quad i = 1, \dots, m, \quad \text{such that} \quad w^\nu - \sum_{i=1}^m y_i^\nu a_i^\nu \in N_{X^\nu}(x^\nu). \quad (2.4)$$

By Caratheodory's theorem, there are $\lambda_{ij}^\nu \geq 0$ and $b_{ij}^\nu \in \partial f_i^\nu(x^\nu)$, $j = 0, 1, \dots, n$, such that $\sum_{j=0}^n \lambda_{ij}^\nu = 1$ and $a_i^\nu = \sum_{j=0}^n \lambda_{ij}^\nu b_{ij}^\nu$. The sequence $\{(\lambda_{i0}^\nu, \dots, \lambda_{in}^\nu), \nu \in N\}$ is contained in a compact set and thus has a convergent subsequence. By assumption (2.3), $\{b_{ij}^\nu, \nu \in N\}$ is bounded with all its cluster points in $\text{con } \partial f_i(\bar{x})$. Consequently, there exist $\bar{\lambda}_i = (\bar{\lambda}_{i0}, \dots, \bar{\lambda}_{in}) \in \mathbb{R}^{1+n}$, $\bar{b}_{i0}, \dots, \bar{b}_{in}$, and a subsequence of N , which we also denote by N , such that

$$(\lambda_{i0}^\nu, \dots, \lambda_{in}^\nu) \xrightarrow{N} \bar{\lambda}_i \geq 0, \quad \text{with} \quad \sum_{j=0}^n \bar{\lambda}_{ij} = 1, \quad \text{and} \quad b_{ij}^\nu \xrightarrow{N} \bar{b}_{ij} \in \text{con } \partial f_i(\bar{x}).$$

This means that

$$a_i^\nu \xrightarrow{N} \bar{a}_i = \sum_{j=0}^n \bar{\lambda}_{ij} \bar{b}_{ij} \in \text{con } \partial f_i(\bar{x}).$$

Via Attouch's theorem [48, Thm. 12.35], $\iota_{X^\nu} \xrightarrow{e} \iota_X$ implies $N_{X^\nu} \xrightarrow{a} N_X$ because X^ν and X are convex. Using this fact as well as the observations that $y_i^\nu \xrightarrow{N} \bar{y}_i$ and $w^\nu - \sum_{i=1}^m y_i^\nu a_i^\nu \xrightarrow{N} \bar{w} - \sum_{i=1}^m \bar{y}_i \bar{a}_i$, we obtain from (2.4) that $\bar{w} - \sum_{i=1}^m \bar{y}_i \bar{a}_i \in N_X(\bar{x})$. Thus, because $\bar{a}_i \in \text{con } \partial f_i(\bar{x})$, one has

$$\bar{w} \in \sum_{i=1}^m \bar{y}_i \text{con } \partial f_i(\bar{x}) + N_X(\bar{x})$$

and we conclude that $(\bar{u}, \bar{v}, \bar{w}) \in S(\bar{x}, \bar{y}, \bar{z})$. We have established that $\text{LimOut}(\text{gph } S^\nu) \subset \text{gph } S$ and this holds without leveraging any of the conditions (a)-(e).

Under the modified assumption that $X^\nu = X$, but not necessarily convex, we still have $N_{X^\nu} = N_X \xrightarrow{g} N_X$ because N_X is outer semicontinuous; see [48, Prop. 6.6]. Thus, the above argument carries over.

When $f_i^\nu = f_i$, then (2.3) holds automatically because f_i is lLc and ∂f_i is outer semicontinuous; see [48, Prop. 8.7].

Under the assumption that $X^\nu = X$ and $f_i^\nu = f_i$, we show that $\text{LimInn}(\text{gph } S^\nu) \supset \text{gph } S$. Let $(\bar{x}, \bar{y}, \bar{z}, \bar{u}, \bar{v}, \bar{w}) \in \text{gph } S$. This means that

$$\bar{u} = F(\bar{x}) - \bar{z}, \quad \bar{v} \in \partial h(\bar{z}) - \bar{y}, \quad \bar{w} \in \sum_{i=1}^m \bar{y}_i \text{con } \partial f_i(\bar{x}) + N_X(\bar{x}).$$

Set $x^\nu = \bar{x}$, $y^\nu = \bar{y}$, and $w^\nu = \bar{w}$. Since $h^\nu \xrightarrow{e} h$, it follows by Attouch's theorem [48, Thm. 12.35] that $\partial h^\nu \xrightarrow{g} \partial h$ and there are $v^\nu \rightarrow \bar{v}$ and $z^\nu \rightarrow \bar{z}$ such that $v^\nu + \bar{y} \in \partial h^\nu(z^\nu)$. Let $u^\nu = F(\bar{x}) - z^\nu$. This means that $(x^\nu, \bar{y}, z^\nu, u^\nu, v^\nu, w^\nu) \in \text{gph } S^\nu \rightarrow (\bar{x}, \bar{y}, \bar{z}, \bar{u}, \bar{v}, \bar{w})$. Consequently, $(\bar{x}, \bar{y}, \bar{z}, \bar{u}, \bar{v}, \bar{w}) \in \text{LimInn}(\text{gph } S^\nu)$ and $\text{LimInn}(\text{gph } S^\nu) \supset \text{gph } S$ holds. In view of the earlier results, $S^\nu \xrightarrow{g} S$ and we conclude that the approximations are consistent.

For the case with smooth f_i^ν, f_i , the claim about (2.3) is trivial and it only remains to show $\text{LimInn}(\text{gph } S^\nu) \supset \text{gph } S$. Now, we obtain the simplification

$$\sum_{i=1}^m y_i \text{con } \partial f_i(x) = \left\{ \sum_{i=1}^m y_i \nabla f_i(x) \right\} = \{ \nabla F(x)^\top y \},$$

with a similar expression for the approximating functions. Let $(\bar{x}, \bar{y}, \bar{z}, \bar{u}, \bar{v}, \bar{w}) \in \text{gph } S$. This means that $\bar{u} = F(\bar{x}) - \bar{z}$, $\bar{v} \in \partial h(\bar{z}) - \bar{y}$, and $\bar{w} \in \nabla F(\bar{x})^\top \bar{y} + N_X(\bar{x})$. In particular, $\bar{x} \in X$. Set $y^\nu = \bar{y}$. Since $h^\nu \xrightarrow{e} h$, it follows by Attouch's theorem [48, Thm. 12.35] that $\partial h^\nu \xrightarrow{g} \partial h$ and there are $v^\nu \rightarrow \bar{v}$ and $z^\nu \rightarrow \bar{z}$ such that $v^\nu + \bar{y} \in \partial h^\nu(z^\nu)$.

Since $N_{X^\nu} \xrightarrow{g} N_X$, as argued above, there's $(x^\nu, t^\nu) \in \text{gph } N_{X^\nu}$ with $x^\nu \in X^\nu \rightarrow \bar{x}$ and $t^\nu \rightarrow \bar{w} - \nabla F(\bar{x})^\top \bar{y}$. Construct $w^\nu = t^\nu + \nabla F^\nu(x^\nu)^\top \bar{y}$. We then have $w^\nu \rightarrow \bar{w}$ and $w^\nu - \nabla F^\nu(x^\nu)^\top \bar{y} \in N_{X^\nu}(x^\nu)$. Also, construct $u^\nu = F^\nu(x^\nu) - z^\nu$, which converges to \bar{u} . In summary, we've constructed $(x^\nu, \bar{y}, z^\nu, u^\nu, v^\nu, w^\nu) \in \text{gph } S^\nu \rightarrow (\bar{x}, \bar{y}, \bar{z}, \bar{u}, \bar{v}, \bar{w})$. This means that $(\bar{x}, \bar{y}, \bar{z}, \bar{u}, \bar{v}, \bar{w}) \in \text{LimInn}(\text{gph } S^\nu)$ and, thus, $\text{LimInn}(\text{gph } S^\nu) \supset \text{gph } S$ holds. \square

While we believe weak consistency is the natural goal in most applications, in some cases one might be willing to sacrifice some assurances with the benefit of relaxed assumptions.

2.5 Corollary (approximations without full epi-convergence). *Suppose that $h^\nu \xrightarrow{e} h$, $X^\nu \xrightarrow{s} X$ with these sets being convex, and (2.3) holds for $i = 1, \dots, m$. Then, for $\alpha \in \overline{\mathbb{R}}$ and vanishing δ^ν , one has*

$$\text{LimOut}\{\varphi^\nu \leq \alpha\} \subset \{\varphi \leq \alpha\} \quad \text{and} \quad \text{LimOut}(S^\nu)^{-1}(\mathbb{B}(0, \delta^\nu)) \subset S^{-1}(0).$$

If $X^\nu = X$, then the assertions hold without the convexity assumption on X .

Proof. Following the proof of Theorem 2.4, we see that the present assumption suffices for $\liminf \varphi^\nu(x^\nu) \geq \varphi(x)$ to hold whenever $x^\nu \rightarrow x$. By [48, Prop. 7.7], the assertion about level-sets follows. Again following the proof, we deduce that $\text{LimOut}(\text{gph } S^\nu) \subset \text{gph } S$ holds without the conditions (a)-(e) in the theorem and we invoke [48, Thm. 5.37] to reach the conclusion. \square

The assumptions of the corollary permit the approximating φ^ν to be arbitrarily “high” relative to φ , i.e., $\text{LimOut}(\text{epi } \varphi^\nu)$ could be a strict subset of $\text{epi } \varphi$. This is a main reason why we avoid calling the approximations consistent in this case. Still, the mild assumptions make the corollary widely applicable.

3 Examples

An array of examples illustrate the breadth of the framework, but numerous possibilities are omitted including those involving polyhedral approximations of X as in [6]. We recall that X, X^ν and f_i, f_i^ν are nonconvex unless specified otherwise.

3.1 Example (goal optimization). For parameters $\alpha_i \in [0, \infty)$ and $\tau_i \in \mathbb{R}$, we consider the problem

$$\text{minimize}_{x \in X} \sum_{i=1}^m \alpha_i \max\{0, f_i(x) - \tau_i\},$$

which aims to lower each $f_i(x)$ down to the goal of τ_i , with α_i being the per-unit penalty for failing to do so. The problem is of the form (1.1) with $h(z) = \sum_{i=1}^m \alpha_i \max\{0, z_i - \tau_i\}$. A possible approximation is to set $X^\nu = X$ and $F^\nu = F$, but replace h by

$$h^\nu(z) = \sum_{i=1}^m \alpha_i \psi^\nu(z_i - \tau_i), \quad \text{where } \psi^\nu(\gamma) = \frac{1}{\theta^\nu} \ln(1 + \exp(\theta^\nu \gamma)), \quad \theta^\nu \in (0, \infty).$$

Since h^ν is convex and differentiable any number of times, it offers computational benefits over h . The resulting approximating problems and their optimality conditions are consistent approximations provided that $\theta^\nu \rightarrow \infty$. We refer to [14, 12] for more general smoothing schemes.

Detail. By Theorem 2.4, condition (a), and the refinements, we only need to show $h^\nu \xrightarrow{e} h$. Since $\sup_{\gamma \in \mathbb{R}} |\psi^\nu(\gamma) - \max\{0, \gamma\}| \leq (\ln 2)/\theta^\nu$ as seen from [41], (1.2) and (1.3) hold.

For an illustration of the optimality conditions, suppose that f_1, \dots, f_m are smooth. Then, $0 \in S^\nu(x, y, z)$ simplifies to $F(x) = z$, $y = \nabla h^\nu(z)$, $-\nabla F(x)^\top y \in N_X(x)$. In fact, this can be written as $-\nabla F(x)^\top \nabla h^\nu(F(x)) \in N_X(x)$, with

$$\nabla h^\nu(z) = \begin{bmatrix} \alpha_1 \nabla \psi^\nu(z_1 - \tau_1) \\ \vdots \\ \alpha_m \nabla \psi^\nu(z_m - \tau_m) \end{bmatrix} \quad \text{and} \quad \nabla \psi^\nu(\gamma) = \frac{\exp(\theta^\nu \gamma)}{1 + \exp(\theta^\nu \gamma)}.$$

Similarly, the optimality condition $0 \in S(x, y, z)$ specializes to $F(x) = z$, $y \in \partial h(z)$, $-\nabla F(x)^\top y \in N_X(x)$. Since $\partial h(z) = C_1 \times \dots \times C_m$, where $C_i = \{0\}$ if $z_i < \tau_i$, $C_i = [0, \alpha_i]$ if $z_i = \tau_i$, and $C_i = \{\alpha_i\}$ otherwise, the optimality condition simplifies further to $-\nabla F(x)^\top y \in N_X(x)$ with $y_i = 0$ if $f_i(x) < \tau_i$, $y_i \in [0, \alpha_i]$ if $f_i(x) = \tau_i$, and $y_i = \alpha_i$ otherwise. These conditions are necessary for optimality in the

actual and approximating problems by Proposition 2.1; the qualification (2.2) holds in each case because $\text{dom } h = \text{dom } h^\nu = \mathbb{R}^m$. \square

3.2 Example (stochastic optimization). For probabilities $p = (p_1, \dots, p_m) \in P$, where $P \subset \mathbb{R}^m$ is the set of nonnegative vectors with components summing to one, consider the problem

$$\text{minimize}_{x \in X} \sum_{i=1}^m p_i f_i(x),$$

which appears in stochastic optimization and machine learning. The problem is of the form (1.1) with $h(z) = \sum_{i=1}^m p_i z_i$. In practice, the probabilities might not be fully known and this leads to approximating problems with p replaced by $p^\nu \in P$, i.e., $h^\nu(z) = \sum_{i=1}^m p_i^\nu z_i$ in (2.1). The change may also be part of a sensitivity analysis such as when developing influence functions [31, 4]. The approximations are consistent provided that $p^\nu \rightarrow p$.

Detail. Since $h^\nu(z^\nu) \rightarrow h(z)$ whenever $z^\nu \rightarrow z$, we obtain $h^\nu \xrightarrow{e} h$; see (1.2) and (1.3). Thus, the consistency follows via condition (a) of Theorem 2.4. The optimality condition $0 \in S(x, y, z)$ is necessary by Proposition 2.1 because the qualification (2.2) holds. In fact, the optimality condition simplifies to $0 \in \sum_{i=1}^m p_i \text{con } \partial f_i(x) + N_X(x)$ because $y = \nabla h(z) = p$. \square

3.3 Example (distributionally robust optimization). Let $P \subset \mathbb{R}^m$ be as in the previous example and let A and A^ν be nonempty closed subsets of P . We consider the problem

$$\text{minimize}_{x \in X} \max_{p \in A} \sum_{i=1}^m p_i f_i(x)$$

and its approximation obtained by replacing A by A^ν . The set A might consist of a single, true probability vector and be approximated by a set A^ν “centered” on a current best estimate of the true probability vector with some “radius” reflecting the uncertainty about this estimate. The resulting problems fit the forms of (1.1) and (2.1) with $h(z) = \max_{p \in A} \langle p, z \rangle$ and $h^\nu(z) = \max_{p \in A^\nu} \langle p, z \rangle$. If $A^\nu \xrightarrow{s} A$, then the approximations are consistent.

Detail. Let $\hat{A}(z) = \text{argmax}_{p \in A} \langle p, z \rangle$ and $\hat{A}^\nu(z) = \text{argmax}_{p \in A^\nu} \langle p, z \rangle$. The functions h and h^ν are convex. They are also real-valued, and thus continuous, because A and A^ν are compact. To establish $h^\nu \xrightarrow{e} h$, we argue using (1.2) and (1.3). Suppose that $z^\nu \rightarrow z$. Let $\bar{p} \in \hat{A}(z)$. Since $A^\nu \xrightarrow{s} A$, there exists $p^\nu \in A^\nu \rightarrow \bar{p}$. This implies that

$$\liminf (\max_{p \in A^\nu} \langle p, z^\nu \rangle) \geq \liminf \langle p^\nu, z^\nu \rangle = \langle \bar{p}, z \rangle = \max_{p \in A} \langle p, z \rangle.$$

Thus, (1.2) holds for h and h^ν . For fixed z , let $p^\nu \in \hat{A}^\nu(z)$. Then, for any $\{\bar{p}^\nu \in A, \nu \in \mathbb{N}\}$, one has

$$\max_{p \in A^\nu} \langle p, z \rangle = \langle p^\nu, z \rangle = \langle p^\nu - \bar{p}^\nu, z \rangle + \langle \bar{p}^\nu, z \rangle \leq \|p^\nu - \bar{p}^\nu\|_2 \|z\|_2 + \max_{p \in A} \langle p, z \rangle.$$

Since there are $\bar{p}^\nu \in A$ such that $\|p^\nu - \bar{p}^\nu\|_2 \rightarrow 0$ because $A^\nu \xrightarrow{s} A$, $\limsup h^\nu(z) \leq h(z)$. Consequently, $h^\nu \xrightarrow{e} h$ and we again invoke condition (a) of Theorem 2.4 and the refinements. \square

3.4 Example (augmented Lagrangian methods). With $h(z) = z_1 + \sum_{i=2}^m \iota_{\{0\}}(z_i)$, the problem

$$\underset{x \in X}{\text{minimize}} f_1(x) \quad \text{subject to} \quad f_2(x) = 0, \dots, f_m(x) = 0$$

fits the mold (1.1). An approximation stemming from augmented Lagrangian methods utilizes

$$h^\nu(z) = z_1 + \sum_{i=2}^m (y_i^\nu z_i + \frac{1}{2} \theta^\nu z_i^2),$$

where $y^\nu = (y_2^\nu, \dots, y_m^\nu) \in \mathbb{R}^{m-1}$ and $\theta^\nu \in [0, \infty)$. If $\{y^\nu, \nu \in \mathbb{N}\}$ is bounded and $\theta^\nu \rightarrow \infty$, then the resulting approximations are consistent.

Detail. The function h is not real-valued and we turn to condition (b) of Theorem 2.4. For $z \in \text{dom } h = \mathbb{R} \times \{0\}^{m-1}$, one has $h^\nu(z) = z_1 = h(z)$ and the condition holds. This also confirms the limsup-condition (1.3) for h^ν and h . For the liminf-condition (1.2), suppose that $z^\nu \rightarrow z$. Then,

$$\liminf h^\nu(z^\nu) \geq z_1 + \sum_{i=2}^m \liminf (y_i^\nu z_i^\nu) + \frac{1}{2} \sum_{i=2}^m \liminf (\theta^\nu (z_i^\nu)^2).$$

If any z_2, \dots, z_m is nonzero, then the right-hand side equals ∞ , which coincides with $h(z)$. If $z_2 = \dots = z_m = 0$, then the right-hand side is no smaller than $z_1 = h(z)$. Thus, $h^\nu \xrightarrow{e} h$. \square

3.5 Example (min-functions). For smooth $g_{ik} : \mathbb{R}^n \rightarrow \mathbb{R}$, $k = 1, \dots, s_i$, $i = 1, \dots, m$, let

$$f_i(x) = \min_{k=1, \dots, s_i} g_{ik}(x).$$

We adopt smooth approximations of f_i by defining

$$f_i^\nu(x) = -\frac{1}{\theta^\nu} \ln \left(\sum_{k=1}^{s_i} \exp(-\theta^\nu g_{ik}(x)) \right),$$

where $\theta^\nu \in (0, \infty)$; see [37, 14, 5, 12] for related smoothing techniques. If $\theta^\nu \rightarrow \infty$, then (2.3) holds. Moreover, if in addition $X^\nu \supset X$, $X^\nu \xrightarrow{s} X$, with these sets being convex, $h^\nu \xrightarrow{e} h$, $h^\nu(z) \leq h^\nu(\bar{z})$ when $z \leq \bar{z}$, and $h^\nu(z) \leq h(z)$ for all z , then the approximations are weakly consistent.

Detail. We leverage condition (c) of Theorem 2.4 and observe that $F^\nu(x) \leq F(x)$ because (see [41])

$$0 \leq f_i(x) - f_i^\nu(x) \leq \frac{\ln s_i}{\theta^\nu} \quad \forall x \in \mathbb{R}^n. \quad (3.1)$$

Moreover, we need to confirm (2.3). Let $x^\nu \rightarrow \bar{x}$. The previous inequalities show that $f_i^\nu(x^\nu) \rightarrow f_i(\bar{x})$ because $\theta^\nu \rightarrow \infty$. It is apparent that f_i^ν is smooth and, in fact,

$$\nabla f_i^\nu(x) = \sum_{k=1}^{s_i} \mu_{ik}^\nu(x) \nabla g_{ik}(x), \quad \text{with} \quad \mu_{ik}^\nu(x) = \frac{\exp(\theta^\nu (f_i(x) - g_{ik}(x)))}{\sum_{j=1}^{s_i} \exp(\theta^\nu (f_i(x) - g_{ij}(x)))}.$$

Since $\mu_{ik}^\nu(x) \in (0, 1)$ regardless of x , we have that $\{\nabla f_i^\nu(x^\nu), \nu \in \mathbb{N}\}$ is bounded. For some $N \in \mathcal{N}_\infty^\#$, suppose that $\nabla f_i^\nu(x^\nu) \xrightarrow{N} \bar{v}$. We would like to show $\bar{v} \in \text{con } \partial f_i(\bar{x})$. We observe that

$$\text{con } \partial f_i(x) = -\text{con } \partial(-f_i)(x) = -\text{con} \{ \nabla(-g_{ik})(x) \mid k \in \mathbb{A}_i(x) \} = \text{con} \{ \nabla g_{ik}(x) \mid k \in \mathbb{A}_i(x) \},$$

where $\mathbb{A}_i(x) = \operatorname{argmin}_{k=1, \dots, s_i} g_{ik}(x)$. If $k \notin \mathbb{A}_i(\bar{x})$, then the continuity of f_i and g_{ik} implies that

$$\exp(\theta^\nu(f_i(x^\nu) - g_{ik}(x^\nu))) \xrightarrow{N} 0.$$

The denominator in the defining expression for $\mu_{ik}^\nu(x^\nu)$ is greater than one because $f_i(x^\nu) - g_{ij}(x^\nu) = 0$ for $j \in \mathbb{A}_i(x^\nu)$. Thus, $\mu_{ik}^\nu(x^\nu) \xrightarrow{N} 0$.

For any k , $\mu_{ik}^\nu(x^\nu) \in (0, 1)$. Consequently, after passing to another subsequence which we also denote by N , there are $\mu_{ik}^\infty \in [0, 1]$, $k = 1, \dots, s_i$, such that $\mu_{ik}^\nu(x^\nu) \xrightarrow{N} \mu_{ik}^\infty$. Since $\sum_{k=1}^{s_i} \mu_{ik}^\nu(x^\nu) = 1$ for all ν , we must also have $\sum_{k=1}^{s_i} \mu_{ik}^\infty = 1$, with $\mu_{ik}^\infty = 0$ if $k \notin \mathbb{A}_i(\bar{x})$ as already seen. We conclude that

$$\nabla f_i^\nu(x^\nu) = \sum_{k=1}^{s_i} \mu_{ik}^\nu(x^\nu) \nabla g_{ik}(x^\nu) \xrightarrow{N} \bar{v} = \sum_{k \in \mathbb{A}_i(\bar{x})} \mu_{ik}^\infty \nabla g_{ik}(\bar{x}) \in \operatorname{con} \partial f_i(\bar{x})$$

and (2.3) holds. An instance of h and h^ν satisfying condition (c) of Theorem 2.4 follows next. \square

3.6 Example (penalty methods). With $h(z) = z_1 + \iota_{(-\infty, 0]^{m-1}}(z_2, \dots, z_m)$, we consider the problem

$$\underset{x \in X}{\text{minimize}} f_1(x) \text{ subject to } f_2(x) \leq 0, \dots, f_m(x) \leq 0,$$

which is of the form (1.1). An approximation stemming from penalty methods utilizes

$$h^\nu(z) = z_1 + \theta^\nu \sum_{i=2}^m (\max\{0, z_i\})^2,$$

where $\theta^\nu \in [0, \infty)$. If $X^\nu = X$, $F^\nu = F$, and $\theta^\nu \rightarrow \infty$, then the resulting approximations are consistent.

Detail. We turn to condition (c) of Theorem 2.4. Certainly, $h^\nu(z) \leq h^\nu(\bar{z})$ when $z \leq \bar{z}$ and the other requirements hold as well. To confirm $h^\nu \xrightarrow{e} h$, we leverage the characterization (1.2) and (1.3). Let $z^\nu \rightarrow z$. We note that $\liminf h^\nu(z^\nu) \geq h(z)$ when $z = (z_1, \dots, z_m)$ has $z_i \leq 0$ for $i = 2, \dots, m$. If any of these z_i are positive, then $z_i^\nu > 0$ for sufficiently large ν . Thus, $h^\nu(z^\nu) \rightarrow \infty = h(z)$ and (1.2) holds for h^ν and h . For any z , $\limsup h^\nu(z) \leq h(z)$ and (1.3) holds as well. Consequently, $h^\nu \xrightarrow{e} h$.

Condition (c) of Theorem 2.4 also permits outer approximations of X and lower bounding approximations of F (see Example 3.5) while retaining weak consistency. \square

3.7 Example (interior-point methods). For the actual problem in Example 3.6, we consider the approximation

$$h^\nu(z) = \begin{cases} z_1 - \frac{1}{\theta^\nu} \sum_{i=2}^m \ln(-z_i) & \text{if } z_2 < 0, \dots, z_m < 0 \\ \infty & \text{otherwise,} \end{cases}$$

where $\theta^\nu \in (0, \infty)$. This logarithmic penalty approach is the basis for (primal) interior-point methods. If $\theta^\nu \rightarrow \infty$ and for all $x \in X$, with $f_i(x) \leq 0$ for all $i = 2, \dots, m$, there exists $x^\nu \in X \rightarrow x$ such that $f_i(x^\nu) < 0$ for all $i = 2, \dots, m$, then the approximations are consistent.

Detail. We leverage condition (d) of Theorem 2.4. Trivially, h is continuous relative to its domain. We establish $h^\nu \xrightarrow{e} h$ via (1.2) and (1.3). Let $z^\nu \rightarrow z$. If $z_i > 0$ for some $i = 2, \dots, m$, then $z_i^\nu \geq 0$ for sufficiently large ν and $\liminf h^\nu(z^\nu) \geq h(z)$ because both sides are ∞ . If $z_i \leq 0$ for $i = 2, \dots, m$, then $h(z) = z_1$ and the \liminf -condition holds again. The condition $\limsup h^\nu(z^\nu) \leq h(z)$ holds because for any $z_i = 0$, one can select $z_i^\nu = -\exp(-\sqrt{\theta^\nu})$. Then, $z_i^\nu \rightarrow z_i$ and $-(\ln(-z_i^\nu))/\theta^\nu = \sqrt{\theta^\nu}/\theta^\nu \rightarrow 0$. \square

3.8 Example (expectation functions). *Expectation functions arise in stochastic optimization and machine learning and then the component functions f_1, \dots, f_m may take the form*

$$f_i(x) = \mathbb{E}[g_i(\boldsymbol{\xi}, x)],$$

where $g_i : \Xi \times \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$ is defined in terms of a probability space $(\Xi, \mathcal{B}, \mathbb{P})$. An independent and identically distributed sample $\boldsymbol{\xi}_1, \dots, \boldsymbol{\xi}_\nu$ according to \mathbb{P} defines a sample average approximation

$$f_i^\nu(x) = \frac{1}{\nu} \sum_{j=1}^{\nu} g_i(\boldsymbol{\xi}_j, x).$$

If both g_i and $-g_i$ are random lower semicontinuous and locally inf-integrable (see [53, Sec. 8.G] for definitions), then, with probability one, $f_i^\nu(x^\nu) \rightarrow f_i(x)$ whenever $x^\nu \rightarrow x$ by [53, Thm. 8.56].

Detail. If there is a set $B \in \mathcal{B}$ such that $\mathbb{P}(B) = 1$ and $g_i(\xi, \cdot)$ is smooth for all $\xi \in B$, then one can attempt to repeat the above arguments with f_i replaced by the partial derivatives of f_i and conclude that $\nabla f_i^\nu(x^\nu) \rightarrow \nabla f_i(x)$ whenever $x^\nu \rightarrow x$. This would allow us to satisfy the requirement (2.3). \square

3.9 Example (oracle functions). *Suppose that h is real-valued, but not available in an explicit form. If for each z we can compute $h(z)$ and a subgradient $v \in \partial h(z)$, then the approximation*

$$h^\nu(z) = \max_{k=1, \dots, \nu} h(z^k) + \langle v^k, z - z^k \rangle, \quad \text{with } z^k \in \mathbb{R}^m, \quad v^k \in \partial h(z^k),$$

remains available. Now, $h^\nu \xrightarrow{e} h$ if $\{z^k, k \in \mathbb{N}\}$ is a countable dense subset of \mathbb{R}^m .

Detail. By [48, Thm. 7.17], $h^\nu \xrightarrow{e} h$ whenever the functions converge pointwise on a countable dense subset of \mathbb{R}^m . Let $\bar{z} \in \{z^k, k \in \mathbb{N}\}$. Obviously, $h^\nu(\bar{z}) \leq h(\bar{z})$ because h is the pointwise supremum of its affine supports; see [48, Thm. 8.13]. Moreover, there is $\bar{\nu}$ such that $\bar{z} \in \{z^1, \dots, z^{\bar{\nu}}\}$. Thus, $h^\nu(\bar{z}) \geq h(\bar{z})$ for all $\nu \geq \bar{\nu}$. By [48, Thm. 7.17], this also means that $h^\nu(z) \rightarrow h(z)$ for all $z \in \mathbb{R}^m$. \square

3.10 Example (homotopy method). *For proper, lsc, and convex $\hat{h} : \mathbb{R}^{m-1} \rightarrow \overline{\mathbb{R}}$ and lLc $\hat{F} : \mathbb{R}^n \rightarrow \mathbb{R}^{m-1}$, consider the problem*

$$\underset{x \in X}{\text{minimize}} \hat{h}(\hat{F}(x)),$$

which is of the form (1.1) with $h(z) = \hat{h}(z_1, \dots, z_{m-1})$ for $z = (z_1, \dots, z_m)$ and $F(x) = (\hat{F}(x), f_m(x))$ for some lLc $f_m : \mathbb{R}^n \rightarrow \mathbb{R}$. A homotopy method for the problem solves the approximations

$$\underset{x \in X}{\text{minimize}} (1 - \lambda^\nu) \hat{h}(\hat{F}(x)) + \lambda^\nu f_m(x)$$

as $\lambda^\nu \rightarrow 0$. If f_m is chosen wisely, then the approximating problems might be simpler to solve than the actual one while benefitting from warm starts. The approximations are of the form (2.1) with

$$h^\nu(z) = (1 - \lambda^\nu)\hat{h}(z_1, \dots, z_{m-1}) + \lambda^\nu z_m.$$

If $\lambda^\nu \rightarrow 0$, then the approximations are consistent.

Detail. To establish $h^\nu \xrightarrow{e} h$ via (1.2) and (1.3), we note that for $z^\nu \rightarrow z$, one has

$$\liminf(1 - \lambda^\nu)\hat{h}(z_1^\nu, \dots, z_{m-1}^\nu) + \lambda^\nu z_m^\nu \geq \liminf \hat{h}(z_1^\nu, \dots, z_{m-1}^\nu) + \liminf \lambda^\nu z_m^\nu \geq \hat{h}(z_1, \dots, z_{m-1}) = h(z).$$

Moreover, for $z \in \text{dom } h$, $(1 - \lambda^\nu)\hat{h}(z_1, \dots, z_{m-1}) + \lambda^\nu z_m \rightarrow \hat{h}(z_1, \dots, z_{m-1})$. We therefore have both epi-convergence and pointwise convergence (on $\text{dom } h$) and condition (b) of Theorem 2.4, with refinements, establishes consistency. \square

3.11 Example (monitoring functions). In extended nonlinear programming [47], one utilizes

$$h(z) = \sup_{y \in Y} \left\{ \langle z, y \rangle - \frac{1}{2} \langle y, By \rangle \right\} \quad \text{and} \quad h^\nu(z) = \sup_{y \in Y^\nu} \left\{ \langle z, y \rangle - \frac{1}{2} \langle y, B^\nu y \rangle \right\},$$

where $Y, Y^\nu \subset \mathbb{R}^m$ are nonempty polyhedral sets and B, B^ν are symmetric positive semidefinite $m \times m$ -matrices. If $B^\nu \rightarrow B$ and $Y^\nu \xrightarrow{s} Y$, then $h^\nu \xrightarrow{e} h$. If in addition B is positive definite or Y is bounded, then h is real-valued and $h^\nu(z) \rightarrow h(z)$ for $z \in \mathbb{R}^m$. Thus, main steps toward (weak) consistency via Theorem 2.4 are immediately accomplished.

Detail. The epi-convergence $h^\nu \xrightarrow{e} h$ is established via the corresponding conjugate functions $(h^\nu)^*$ and h^* . By Wijsman's Theorem [48, Thm. 11.34], it suffices to show that $(h^\nu)^* \xrightarrow{e} h^*$. Since $h^*(y) = \frac{1}{2} \langle y, By \rangle + \iota_Y(y)$, $(h^\nu)^*(y) = \frac{1}{2} \langle y, B^\nu y \rangle + \iota_{Y^\nu}(y)$, and $\iota_{Y^\nu} \xrightarrow{e} \iota_Y$, we conclude that $(h^\nu)^* \xrightarrow{e} h^*$.

It follows by [48, Thm. 7.17] that the pointwise convergence $h^\nu(z) \rightarrow h(z)$ holds when h is real-valued. This is the case when B is positive definite or when Y is bounded. \square

3.12 Example (difference-of-convex functions). For a proper, lsc, and convex $f : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$ and convex $g : \mathbb{R}^n \rightarrow \mathbb{R}$, consider the problem

$$\underset{x \in X}{\text{minimize}} \quad f(x) - g(x),$$

which involves an objective function of the difference-of-convex kind. As the only approximation, suppose that g is replaced by smooth functions $g^\nu : \mathbb{R}^n \rightarrow \mathbb{R}$ that also satisfy the property:

$$x^\nu \in X \rightarrow x \implies g^\nu(x^\nu) \rightarrow g(x) \text{ and } \{\nabla g^\nu(x^\nu), \nu \in \mathbb{N}\} \text{ is bounded with all cluster points in } \partial g(x).$$

This produces weakly consistent approximations.

Detail. For $z = (z_1, z_2)$, with $z_1 \in \mathbb{R}^n$ and $z_2 \in \mathbb{R}$, we set $h(z) = h_1(z_1) + h_2(z_2)$, where $h_1(z_1) = f(z_1)$ and $h_2(z_2) = -z_2$. Moreover, let $F(x) = (x, g(x))$ and $F^\nu(x) = (x, g^\nu(x))$ so that $m = n + 1$, $m_1 = n$, $m_2 = 1$, and $r = 2$ in the notation of condition (e) of Theorem 2.4. Since g is real-valued and convex, it is lLc and then F is also lLc. With $h^\nu = h$, we trivially obtain $h^\nu \xrightarrow{e} h$. The requirement (2.3) translates into the assumed property. We then invoke condition (e) of Theorem 2.4 to conclude weak consistency. For instance $g(x) = \max_{k=1, \dots, s} g_k(x)$, with $g_k : \mathbb{R}^n \rightarrow \mathbb{R}$ being smooth, could be approximated parallel to Example 3.5. In this case, $0 \in S(x, y, z)$ reduces to $0 \in \partial f(x) - \partial g(x) + N_X(x)$. \square

4 Enhanced Proximal Composite Algorithm

As an example of an implementable version of the Consistent Approximation Algorithm, we consider the setting where X^ν is convex, h^ν is real-valued, and F^ν is twice continuously differentiable. Then, the approximating problems (2.1) are solvable by proximal composite methods, which can be traced back to [24, 45, 8]; see also [46, 55, 9] for trust-region versions and [21] for details about rates of convergence under the assumption that h^ν and ∇F^ν are Lipschitz continuous. Let $\text{prj}_{X^\nu}(x)$ be the projection of x on X^ν .

Enhanced Proximal Composite Algorithm (EPCA).

Data. $x^0 \in \mathbb{R}^n$, $\tau \in (1, \infty)$, $\sigma \in (0, 1)$, $\bar{\lambda} \in (0, \infty)$, $\lambda^0 \in (0, \bar{\lambda}]$, $\{\delta^\nu > 0, \nu \in \mathbb{N}\} \rightarrow 0$.

Step 0. Set $\nu = 1$.

Step 1. Set $k = 0$ and $\bar{x}^0 = \text{prj}_{X^\nu}(x^{\nu-1})$.

Step 2. Compute $x^* \in \arg\min_{x \in X^\nu} h^\nu(F^\nu(\bar{x}^k) + \nabla F^\nu(\bar{x}^k)(x - \bar{x}^k)) + \frac{1}{2\lambda^k} \|x - \bar{x}^k\|_2^2$.

If $x^* = \bar{x}^k$, then go to Step 4.

Step 3. If $h^\nu(F^\nu(\bar{x}^k)) - h^\nu(F^\nu(x^*)) \geq \sigma \left(h^\nu(F^\nu(\bar{x}^k)) - h^\nu(F^\nu(\bar{x}^k) + \nabla F^\nu(\bar{x}^k)(x^* - \bar{x}^k)) \right)$,

then set $\lambda^{k+1} = \min\{\tau\lambda^k, \bar{\lambda}\}$ and go to Step 5.

Else, replace λ^k by λ^k/τ and go to Step 2.

Step 4. Set $x^\nu = x^*$, $z^\nu = F^\nu(x^\nu)$, and y^ν such that $y^\nu \in \partial h^\nu(z^\nu)$ and $-\nabla F^\nu(x^\nu)^\top y^\nu \in N_{X^\nu}(x^\nu)$.

Replace ν by $\nu + 1$ and go to Step 1.

Step 5. Set $\bar{x}^{k+1} = x^*$, $\bar{z}^{k+1} = F^\nu(\bar{x}^k) + \nabla F^\nu(\bar{x}^k)(\bar{x}^{k+1} - \bar{x}^k)$, and \bar{y}^{k+1} such that

$$\bar{y}^{k+1} \in \partial h^\nu(\bar{z}^{k+1}) \quad \text{and} \quad -\nabla F^\nu(\bar{x}^k)^\top \bar{y}^{k+1} - \frac{1}{\lambda^k} (\bar{x}^{k+1} - \bar{x}^k) \in N_{X^\nu}(\bar{x}^{k+1}).$$

Set $\bar{u}^{k+1} = F^\nu(\bar{x}^{k+1}) - \bar{z}^{k+1}$ and $\bar{w}^{k+1} = (\nabla F^\nu(\bar{x}^{k+1}) - \nabla F^\nu(\bar{x}^k))^\top \bar{y}^{k+1} - \frac{1}{\lambda^k} (\bar{x}^{k+1} - \bar{x}^k)$.

If $\max\{\|\bar{u}^{k+1}\|_2, \|\bar{w}^{k+1}\|_2\} \leq \delta^\nu$, then set $x^\nu = \bar{x}^{k+1}$, $y^\nu = \bar{y}^{k+1}$, $z^\nu = \bar{z}^{k+1}$, replace ν by $\nu + 1$,

and go to Step 1.

Else, replace k by $k + 1$, and go to Step 2.

4.1 Theorem (enhanced proximal composite algorithm). *Suppose that $X^\nu \xrightarrow{s} X$ with these sets being convex, $h^\nu \xrightarrow{e} h$ with h^ν real-valued, and (2.3) holds with f_i^ν twice continuously differentiable for $i = 1, \dots, m$. If $\{(x^\nu, y^\nu, z^\nu), \nu \in \mathbb{N}\}$ is generated by EPCA and $(x^\nu, y^\nu, z^\nu) \xrightarrow{N} (\hat{x}, \hat{y}, \hat{z})$ for some $N \in \mathcal{N}_\infty^\#$, then*

$$\hat{x} \in X, \quad h(F(\hat{x})) \leq \liminf_{\nu \in N} h^\nu(F^\nu(x^\nu)), \quad 0 \in S(\hat{x}, \hat{y}, \hat{z}).$$

Proof. The two first claims follow by Corollary 2.5. The construction in Steps 4 and 5 ensures that $(x^\nu, y^\nu, z^\nu) \in (S^\nu)^{-1}(\mathbb{B}(0, \delta^\nu))$ for all ν when the norm on \mathbb{R}^{2m+n} is $(u, v, w) \mapsto \max\{\|u\|_2, \|v\|_2, \|w\|_2\}$; see the discussion below. Thus, Corollary 2.5 also confirms that $0 \in S(\bar{x}, \bar{y}, \bar{z})$. \square

EPCA solves the sequence of approximating problems using a composite proximal method of the kind proposed in [33] for each problem. The outer loop indexed by ν represents refinement of the approximating problems. The inner loop indexed by k corresponds to the iterative solution of each approximating problem by means of solving the convex subproblems in Step 2 obtained by linearization of F^ν . (For inexact solution of such subproblems, we refer to [21].) The proximal parameter λ^k is controlled adaptively using the test in Step 3.

A concern is whether EPCA terminates after a finite k in Step 4 or 5 and thus always produces the next (x^ν, y^ν, z^ν) . This turns out to be the case as long as

$$\{\iota_{X^\nu} + h^\nu \circ F^\nu \leq h^\nu(F^\nu(\bar{x}^0))\} \text{ is bounded,} \quad (4.1)$$

where $\bar{x}^0 = \text{prj}_{X^\nu}(x^{\nu-1})$. We see this as follows:

An optimality condition for the subproblem in Step 2 states that x^* satisfies

$$-\nabla F^\nu(\bar{x}^k)^\top y - \frac{1}{\lambda^k}(x^* - \bar{x}^k) \in N_{X^\nu}(x^*) \text{ for some } y \in \partial h^\nu(F^\nu(\bar{x}^k) + \nabla F^\nu(\bar{x}^k)(x^* - \bar{x}^k)). \quad (4.2)$$

Thus, when $x^* = \bar{x}^k$, one has $y \in \partial h^\nu(F^\nu(\bar{x}^k))$ and $-\nabla F^\nu(\bar{x}^k)^\top y \in N_{X^\nu}(x^*)$. This means that there exists y^ν in Step 4. The best way to compute such y^ν depends on the nature of h^ν and X^ν , but it can be achieved by convex optimization since it at most involve finding points in two convex sets. Then, $(x^\nu, y^\nu, z^\nu) \in (S^\nu)^{-1}(0)$. In Step 5, there exists likewise \bar{y}^{k+1} by (4.2), again computable by convex optimization. The construction in Step 5 ensures that $(\bar{u}^{k+1}, 0, \bar{w}^{k+1}) \in S^\nu(\bar{x}^{k+1}, \bar{y}^{k+1}, \bar{z}^{k+1})$.

For the sake of contradiction, suppose that the algorithm iterates indefinitely without generating the next (x^ν, y^ν, z^ν) . Let $\{\bar{x}^k, k \in \mathbb{N}\}$ be the resulting sequence, which has a cluster point in view of (4.1), say \bar{x} . We deduce from [33], especially Theorem 5.4 and its proof, that $\bar{x}^{k+1} - \bar{x}^k \rightarrow 0$ as $k \rightarrow \infty$ and $(\bar{x}^{k+1} - \bar{x}^k)/\lambda^k \rightarrow 0$ as $k \rightarrow \infty$ along the subsequence corresponding to \bar{x} . Then, for k along the subsequence corresponding to \bar{x} , $\bar{z}^{k+1} \rightarrow F^\nu(\bar{x})$, which means that the associated sets of subgradients $\partial h^\nu(\bar{z}^{k+1})$ are contained in a compact set and then the same holds for \bar{y}^{k+1} . This fact as well as the recognition that $F^\nu(\bar{x}^{k+1}) \rightarrow F^\nu(\bar{x})$ and $\nabla F^\nu(\bar{x}^{k+1}) - \nabla F^\nu(\bar{x}^k) \rightarrow 0$ imply that $\bar{u}^{k+1} \rightarrow 0$ and $\bar{w}^{k+1} \rightarrow 0$ as $k \rightarrow \infty$ along the subsequence. Consequently, there's k^* such that $\max\{\|\bar{u}^{k^*+1}\|_2, \|\bar{w}^{k^*+1}\|_2\} \leq \delta^\nu$. We have shown that EPCA terminates for some finite k in either Step 4 or 5.

5 Inverse Problems in Machine Learning

An inverse problem in machine learning is that of determining an input to a collection of neural networks such that their outputs best match a given quantity [26, 28]. Specifically, we are given s neural networks represented by the mappings $F_i : \mathbb{R}^{n_0} \rightarrow \mathbb{R}^{n_q}$, $i = 1, \dots, s$. Each network associates an input vector $x \in \mathbb{R}^{n_0}$ with an output vector $F_i(x) \in \mathbb{R}^{n_q}$. The goal is to determine an input vector x such that the output vectors $F_1(x), \dots, F_s(x)$ are optimized in the sense of a convex function $\hat{h} : \mathbb{R}^{sn_q} \rightarrow \mathbb{R}$, i.e.,

$$\underset{x \in \hat{X}}{\text{minimize}} \quad \hat{h}(F_1(x), \dots, F_s(x)), \quad (5.1)$$

where $\hat{X} \subset \mathbb{R}^{n_0}$ is a nonempty closed set that could impose restrictions on the choice of input vector. For some target output vector $t \in \mathbb{R}^{n_q}$ and nonnegative weights p_1, \dots, p_s , we may simply have

$$\hat{h}(F_1(x), \dots, F_s(x)) = \sum_{i=1}^s p_i \|t - F_i(x)\|_2^2.$$

Suppose that the neural networks are of the feed-forward kind producing their output from passing through q affine mappings composed with an activation function. For each layer $k = 1, \dots, q$, there is a given $n_k \times n_{k-1}$ -matrix $A_{i,k}$, a given vector $b_{i,k} \in \mathbb{R}^{n_k}$, and $F_{i,k} : \mathbb{R}^{n_{k-1}} \rightarrow \mathbb{R}^{n_k}$ with

$$F_{i,k}(x) = G_{i,k}(A_{i,k}x + b_{i,k}) \quad \text{for } x \in \mathbb{R}^{n_{k-1}},$$

where $G_{i,k} : \mathbb{R}^{n_k} \rightarrow \mathbb{R}^{n_k}$ has $G_{i,k}(y) = (g_{i,1,k}(y_1), \dots, g_{i,n_k,k}(y_{n_k}))$ for $y = (y_1, \dots, y_{n_k}) \in \mathbb{R}^{n_k}$ and $g_{i,j,k} : \mathbb{R} \rightarrow \mathbb{R}$ are given epi-regular lLc functions. For example, the ReLU function $g_{i,j,k}(\gamma) = \max\{0, \gamma\}$ is a common such activation function; it is epi-regular and lLc. Then, $F_i = F_{i,q} \circ \dots \circ F_{i,1}$.

It is convenient to express (5.1) using additional variables as follows: We think of $x_{1,i,k} \in \mathbb{R}^{n_k}$ as the output of the k th layer for neural network i . Let $r = \sum_{k=1}^q n_k$, $X = \hat{X} \times \mathbb{R}^{sr}$, and $x = (x_0, x_1)$, with $x_0 \in \mathbb{R}^{n_0}$ and $x_1 = (x_{1,1,1}, \dots, x_{1,s,q}) \in \mathbb{R}^{sr}$. Set

$$F(x) = (x_{1,1,q}, \dots, x_{1,s,q}, H_1(x), \dots, H_s(x)) \in \mathbb{R}^{sn_q + sr} \quad \text{and} \quad h(z) = \hat{h}(z_0) + \iota_{\{0\}^{sr}}(z_1, \dots, z_s),$$

where, for $z = (z_0, z_1, \dots, z_s)$, with $z_0 \in \mathbb{R}^{sn_q}$ and $z_i \in \mathbb{R}^r$, $i = 1, \dots, s$, and

$$H_i(x) = \begin{bmatrix} G_{i,1}(A_{i,1}x_0 + b_{i,1}) - x_{1,i,1} \\ G_{i,2}(A_{i,2}x_{1,i,1} + b_{i,2}) - x_{1,i,2} \\ \vdots \\ G_{i,q}(A_{i,q}x_{1,i,q-1} + b_{i,q}) - x_{1,i,q} \end{bmatrix}.$$

Thus, the inverse problem (5.1) is equivalently expressed in the form (1.1) with $n = n_0 + sr$ and $m = sn_q + sr$; h is proper, lsc, and convex and F is lLc. For theoretical and computational reasons, several approximations may arise. Suppose that $A_{i,j,k}^\nu$, $b_{i,k}^\nu$, convex $\hat{h}^\nu : \mathbb{R}^{sn_q} \rightarrow \mathbb{R}$, and epi-regular lLc $g_{i,j,k}^\nu : \mathbb{R} \rightarrow \mathbb{R}$ approximate $A_{i,j,k}$, $b_{i,k}$, \hat{h} , and $g_{i,j,k}$, respectively. The approximating quantities define h^ν and F^ν via H_i^ν and $G_{i,k}^\nu$ in the manner laid out for h , F , H_i , and $G_{i,k}$. For simplicity, \hat{X} is not approximated so that $X^\nu = X$. It turns out that weak consistency follows naturally.

5.1 Proposition (weak consistency in inverse machine learning). *In the notation of this section, suppose that for each (i, j, k) , the following property holds:*

$$\left. \begin{array}{l} \gamma^\nu \in \mathbb{R} \rightarrow \gamma \\ \alpha^\nu \in \partial g_{i,j,k}^\nu(\gamma^\nu) \end{array} \right\} \implies \left\{ \begin{array}{l} g_{i,j,k}^\nu(\gamma^\nu) \rightarrow g_{i,j,k}(\gamma) \\ \{\alpha^\nu, \nu \in \mathbb{N}\} \text{ is bounded with all its cluster points in } \partial g_{i,j,k}(\gamma). \end{array} \right. \quad (5.2)$$

If $A_{i,j,k}^\nu \rightarrow A_{i,j,k}$, $b_{i,k}^\nu \rightarrow b_{i,k}$, and $\hat{h}^\nu \xrightarrow{e} \hat{h}$, then the resulting pairs $\{(\varphi^\nu, S^\nu), \nu \in \mathbb{N}\}$ are weakly consistent approximations of (φ, S) .

Proof. We establish $\varphi^\nu \xrightarrow{e} \varphi$ using (1.2) and (1.3). The liminf-condition holds because $h^\nu \xrightarrow{e} h$ and $H_i^\nu(x^\nu) \rightarrow H_i(x)$ whenever $x^\nu \rightarrow x$. For the limsup-condition, it suffices to consider $\bar{x} = (\bar{x}_0, \bar{x}_1)$ such that $\bar{x}_0 \in \hat{X}$ and $H_i(\bar{x}) = 0$ for $i = 1, \dots, s$. Construct $x^\nu = (x_0^\nu, x_1^\nu)$ with $x_0^\nu = \bar{x}_0$ and $x_1^\nu = (x_{1,1,1}^\nu, \dots, x_{1,s,q}^\nu)$, where for each $i = 1, \dots, s$,

$$x_{1,i,1}^\nu = G_{i,1}^\nu(A_{i,1}^\nu \bar{x}_0 + b_{i,1}^\nu), \quad x_{1,i,2}^\nu = G_{i,2}^\nu(A_{i,2}^\nu x_{1,i,1}^\nu + b_{i,2}^\nu), \quad \dots, \quad x_{1,i,q}^\nu = G_{i,q}^\nu(A_{i,q}^\nu x_{1,i,q-1}^\nu + b_{i,q}^\nu).$$

Thus, we have $H_i^\nu(x^\nu) = 0$ and $x^\nu \rightarrow \bar{x}$. Since \hat{h} is real-valued and convex, $\hat{h}^\nu(x_{1,1,q}^\nu, \dots, x_{1,s,q}^\nu) \rightarrow \hat{h}(\bar{x}_{1,1,q}, \dots, \bar{x}_{1,s,q})$ by [48, Thm. 7.17]. This implies that $\varphi^\nu(x^\nu) \rightarrow \varphi(\bar{x})$.

Next, we turn to the optimality conditions. Suppose that $(\bar{x}, \bar{y}, \bar{z}, \bar{u}, \bar{v}, \bar{w}) \in \text{LimOut}(\text{gph } S^\nu)$. Then, there are $N \in \mathcal{N}_\infty^\#$ and $(x^\nu, y^\nu, z^\nu, u^\nu, v^\nu, w^\nu) \xrightarrow{N} (\bar{x}, \bar{y}, \bar{z}, \bar{u}, \bar{v}, \bar{w})$ with $(u^\nu, v^\nu, w^\nu) \in S^\nu(x^\nu, y^\nu, z^\nu)$.

Let $A_{i,j,k}$ be the j th row of $A_{i,k}$ and $b_{i,j,k}$ be the j th component of $b_{i,k}$, with parallel definitions of $A_{i,j,k}^\nu$ and $b_{i,j,k}^\nu$. The j th component of $x_{1,i,k} \in \mathbb{R}^{n_k}$ is denoted by $x_{1,i,j,k}$. Moreover, the affine function $a_{i,j,k} : \mathbb{R}^{n_0+sr} \rightarrow \mathbb{R}$ has

$$a_{i,j,k}(x) = \langle A_{i,j,k}, x_{1,i,k-1} \rangle + b_{i,j,k}$$

with the convention that $x_{1,i,0} = x_0$. The function $f_{i,j,k} : \mathbb{R}^{n_0+sr} \rightarrow \mathbb{R}$ has

$$f_{i,j,k}(x) = g_{i,j,k}(a_{i,j,k}(x)) - x_{1,i,j,k}.$$

The approximations $a_{i,j,k}^\nu$ and $f_{i,j,k}^\nu$ are defined similarly.

Since $u^\nu = F^\nu(x^\nu) - z^\nu$, we conclude that $\bar{u} = F(\bar{x}) - \bar{z}$; recall that $g_{i,j,k}^\nu(\gamma^\nu) \rightarrow g_{i,j,k}(\gamma)$ when $\gamma^\nu \rightarrow \gamma$. The fact that $v^\nu - y^\nu \in \partial h^\nu(z^\nu)$ implies that $v_0^\nu - y_0^\nu \in \partial \hat{h}^\nu(z_0^\nu)$ and $z_i^\nu = 0$, $v_i^\nu - y_i^\nu \in \mathbb{R}^r$, $i = 1, \dots, s$, where $v^\nu = (v_0^\nu, v_1^\nu, \dots, v_s^\nu)$, with $v_0^\nu \in \mathbb{R}^{s n_q}$ and $v_i^\nu \in \mathbb{R}^r$ for $i = 1, \dots, s$; the vectors \bar{v} , \bar{y} , \bar{z} , \bar{y}^ν , and \bar{z}^ν are partitioned similarly. By Attouch's Theorem [48, Thm. 12.35], we conclude that $\bar{v}_0 - \bar{y}_0 \in \partial \hat{h}(\bar{z}_0)$. Thus, $\bar{v} - \bar{y} \in \partial h(\bar{z})$.

It remains to confirm the last portion of $(\bar{u}, \bar{v}, \bar{w}) \in S(\bar{x}, \bar{y}, \bar{z})$ involving subgradients of the component functions of F . Since $g_{i,j,k}$ is lLc and epi-regular, the component function $f_{i,j,k}$ has

$$\partial f_{i,j,k}(x) = c_{i,j,k}^\top \partial g_{i,j,k}(a_{i,j,k}(x)) - e_{i,j,k},$$

where $c_{i,j,k} = \nabla a_{i,j,k}(x)$ and $e_{i,j,k} \in \mathbb{R}^{n_0+sr}$ is a vector with a single 1 placed appropriately and with zero elsewhere. Likewise,

$$\partial f_{i,j,k}^\nu(x) = (c_{i,j,k}^\nu)^\top \partial g_{i,j,k}^\nu(a_{i,j,k}^\nu(x)) - e_{i,j,k},$$

where $c_{i,j,k}^\nu = \nabla a_{i,j,k}^\nu(x)$. The assumption (5.2) implies that any sequence $\{d_{i,j,k}^\nu \in \partial f_{i,j,k}^\nu(x^\nu), \nu \in N\}$ is bounded and thus has a cluster point, with any such point in $\partial f_{i,j,k}(\bar{x})$. Since $N_{\hat{X}}$ is outer semicontinuous, we deduce that the last requirement holds and, thus, $(\bar{u}, \bar{v}, \bar{w}) \in S(\bar{x}, \bar{y}, \bar{z})$. \square

A typical activation function is the ReLU with $g_{i,j,k}(\gamma) = \max\{0, \gamma\}$. A smooth approximation could be $g_{i,j,k}^\nu(\gamma) = \frac{1}{\theta^\nu} \ln(1 + \exp(\theta^\nu \gamma))$. The assumption (5.2) holds in this case; see Example 3.1.

6 Rates and Error Estimates

Consistency furnishes guarantees about the limiting behavior of approximations, but it also can be beneficial to quantify the rate of convergence. In this section, we refine results from [51] and estimate the discrepancy between near-solutions of the optimality condition $0 \in S^\nu(x, y, z)$ and those of $0 \in S(x, y, z)$. Chapter 8 of [16] addresses similar issues using different techniques, especially for problems with affine structure. For error estimates of minimizers, minima, and level-sets, we refer to [50, 51].

The *point-to-set distance* between $\bar{x} \in \mathbb{R}^n$ and $C \subset \mathbb{R}^n$ under norm $\|\cdot\|$ is denoted by

$$\text{dist}(\bar{x}, C) = \inf_{x \in C} \|x - \bar{x}\| \quad \text{when } C \neq \emptyset \quad \text{and} \quad \text{dist}(\bar{x}, \emptyset) = \infty.$$

The *excess* of $C \subset \mathbb{R}^n$ over $D \subset \mathbb{R}^n$ is defined as

$$\text{exs}(C; D) = \begin{cases} \sup_{x \in C} \text{dist}(x, D) & \text{if } C \neq \emptyset, D \neq \emptyset \\ \infty & \text{if } C \neq \emptyset, D = \emptyset \\ 0 & \text{otherwise.} \end{cases}$$

We concentrate on a truncated version given by

$$\text{exs}_\rho(C; D) = \text{exs}(C \cap \mathbb{B}(0, \rho); D) \quad \text{for } \rho \in [0, \infty).$$

All these concepts rely on the choice of norm. The Euclidean norm remains the default, but in the context of $S, S^\nu : \mathbb{R}^{n+2m} \rightrightarrows \mathbb{R}^{2m+n}$ we adopt the norms given by

$$\|(x, y, z)\|_{\text{in}} = \max\{\|x\|_2, \|y\|_2, \|z\|_2\} \quad \text{and} \quad \|(u, v, w)\|_{\text{out}} = \max\{\|u\|_2, \|v\|_2, \|w\|_2\} \quad (6.1)$$

for the argument and value spaces, respectively, where $x \in \mathbb{R}^n$, $y \in \mathbb{R}^m$, $z \in \mathbb{R}^m$, $u \in \mathbb{R}^m$, $v \in \mathbb{R}^m$, and $w \in \mathbb{R}^n$. The graphs of S, S^ν are subsets of $\mathbb{R}^{n+2m} \times \mathbb{R}^{2m+n}$, for which we adopt the norm given by

$$\max\{\|(x, y, z)\|_{\text{in}}, \|(u, v, w)\|_{\text{out}}\}. \quad (6.2)$$

Thus, $S^{-1}(\mathbb{B}(0, \varepsilon))$ is the set of near-solutions of $0 \in S(x, y, z)$ with the tolerance now being specified by $\|\cdot\|_{\text{out}}$, i.e., $(\bar{x}, \bar{y}, \bar{z}) \in S^{-1}(\mathbb{B}(0, \varepsilon))$ if and only if $(\bar{u}, \bar{v}, \bar{w}) \in S(\bar{x}, \bar{y}, \bar{z})$ and $\|(\bar{u}, \bar{v}, \bar{w})\|_{\text{out}} \leq \varepsilon$.

6.1 Proposition (solution error in optimality conditions). *Suppose that $0 \leq \delta^\nu \leq \rho < \infty$ and $\varepsilon \geq \delta^\nu + \text{exs}_\rho(\text{gph } S^\nu; \text{gph } S)$. Then, under the norms (6.1) and (6.2), one has*

$$\text{exs}_\rho\left((S^\nu)^{-1}(\mathbb{B}(0, \delta^\nu)); S^{-1}(\mathbb{B}(0, \varepsilon))\right) \leq \text{exs}_\rho(\text{gph } S^\nu; \text{gph } S).$$

Proof. A slight modification of the proof of Theorem 5.1 in [51] yields this fact. \square

The proposition ensures that if $(\hat{x}, \hat{y}, \hat{z}) \in \mathbb{B}(0, \rho)$ nearly satisfies the optimality condition $0 \in S^\nu(x, y, z)$ with tolerance $\delta^\nu \in [0, \infty)$, i.e., $\text{dist}(0, S^\nu(\hat{x}, \hat{y}, \hat{z})) \leq \delta^\nu$, then $(\hat{x}, \hat{y}, \hat{z})$ is no further way than $\text{exs}_\rho(\text{gph } S^\nu; \text{gph } S)$ from a point that nearly satisfies the optimality condition $0 \in S(x, y, z)$ with tolerance ε . Thus, the rate of convergence of near-solutions of the approximating problems is governed by the rate of decay of $\text{exs}_\rho(\text{gph } S^\nu; \text{gph } S)$ as $\nu \rightarrow \infty$.

6.2 Theorem (estimate of excess). *Suppose that $\rho \in [0, \infty)$ and $X = X^\nu$. Let*

$$\begin{aligned} \eta_0^\nu &= \sup \left\{ \|F^\nu(x) - F(x)\|_2 \mid x \in X \cap \mathbb{B}(0, \rho) \right\} \\ \eta^\nu &= \sup \left\{ \text{exs}(\partial f_i^\nu(x); \text{con } \partial f_i(x)) \mid x \in X \cap \mathbb{B}(0, \rho), i = 1, \dots, m \right\}. \end{aligned}$$

If the norm on $\mathbb{R}^{n+2m} \times \mathbb{R}^{2m+n}$ is (6.2) and the norm on $\mathbb{R}^m \times \mathbb{R}^m$ is $\max\{\|z\|_2, \|v\|_2\}$, then one has

$$\text{exs}_\rho(\text{gph } S^\nu; \text{gph } S) \leq \max \left\{ \sqrt{m}\rho\eta^\nu, \eta_0^\nu + \text{exs}_{2\rho}(\text{gph } \partial h^\nu; \text{gph } \partial h) \right\}.$$

Proof. Suppose that $(\bar{x}, \bar{y}, \bar{z}, \bar{u}, \bar{v}, \bar{w}) \in \text{gph } S^\nu \cap \mathbb{B}(0, \rho)$, where the ball is specified by (6.2). Then,

$$\bar{u} = F^\nu(\bar{x}) - \bar{z}, \quad \bar{v} + \bar{y} \in \partial h^\nu(\bar{z}), \quad \bar{w} \in \sum_{i=1}^m \bar{y}_i \text{con } \partial f_i^\nu(\bar{x}) + N_X(\bar{x}).$$

We construct a point $(\bar{x}, \bar{y}, z, u, v, w) \in \text{gph } S$ that is near $(\bar{x}, \bar{y}, \bar{z}, \bar{u}, \bar{v}, \bar{w})$ in the norm (6.2).

Since $(\bar{x}, \bar{y}, \bar{z}, \bar{u}, \bar{v}, \bar{w}) \in \mathbb{B}(0, \rho)$, one has $\max\{\|\bar{z}\|_2, \|\bar{v} + \bar{y}\|_2\} \leq 2\rho$. Moreover, $\text{gph } \partial h$ is nonempty, which follows largely from the definition of subgradients; cf. [48, Corollary 8.10]. These facts ensure that there are $z, v \in \mathbb{R}^m$ such that $(z, v + \bar{y}) \in \text{gph } \partial h$ and

$$\max \left\{ \|z - \bar{z}\|_2, \|(v + \bar{y}) - (\bar{v} + \bar{y})\|_2 \right\} \leq \text{exs}_{2\rho}(\text{gph } \partial h^\nu; \text{gph } \partial h).$$

There are $\bar{a}_i \in \text{con } \partial f_i^\nu(\bar{x})$ such that $\bar{w} - \sum_{i=1}^m \bar{y}_i \bar{a}_i \in N_X(\bar{x})$. By Caratheodory's theorem, there exist $\lambda_i^k \geq 0$ and $a_i^k \in \partial f_i^\nu(\bar{x})$, $k = 0, 1, \dots, n$, such that $\sum_{k=0}^n \lambda_i^k = 1$ and $\bar{a}_i = \sum_{k=0}^n \lambda_i^k a_i^k$. By assumption, there exists $b_i^k \in \text{con } \partial f_i(\bar{x})$ such that $\|a_i^k - b_i^k\|_2 \leq \eta^\nu$ and this holds for each i and k . Let $\bar{b}_i = \sum_{k=0}^n \lambda_i^k b_i^k$, which then also is in $\text{con } \partial f_i(\bar{x})$. We construct

$$w = \bar{w} + \sum_{i=1}^m \bar{y}_i (\bar{b}_i - \bar{a}_i) \quad \text{and} \quad u = F(\bar{x}) - z.$$

It now follows that $(\bar{x}, \bar{y}, z, u, v, w) \in \text{gph } S$ and it remains to compute the distance between this point and $(\bar{x}, \bar{y}, \bar{z}, \bar{u}, \bar{v}, \bar{w})$. We already have $\max\{\|z - \bar{z}\|_2, \|v - \bar{v}\|_2\} \leq \text{exs}_{2\rho}(\text{gph } \partial h^\nu; \text{gph } \partial h)$. Moreover,

$$\|u - \bar{u}\|_2 \leq \|F^\nu(\bar{x}) - F(\bar{x})\|_2 + \|z - \bar{z}\|_2 \leq \eta_0^\nu + \text{exs}_{2\rho}(\text{gph } \partial h^\nu; \text{gph } \partial h).$$

Finally, one has $\|w - \bar{w}\|_2 \leq \sum_{i=1}^m |\bar{y}_i| \|\bar{b}_i - \bar{a}_i\|_2 \leq \sqrt{m}\rho\eta^\nu$. \square

The theorem allows us to bound the solution error associated with a particular approximation in terms of the error in the individual components h^ν and F^ν . (We refer to [51] for error estimates

involving $X^\nu \neq X$ in the simplified setting with smooth F .) Thus, the focus turns to estimating the component errors. For that purpose, the following fact, which is a direct application of Theorem 5.2 in [1], is useful. It leverages the *truncated Hausdorff distance* between the sets C and D :

$$\hat{d}_\rho(C, D) = \max \{ \text{exs}_\rho(C; D); \text{exs}_\rho(D; C) \}.$$

6.3 Proposition (approximation of subgradients). *Suppose that the norm on \mathbb{R}^{m+1} is $\max\{\|z\|_2, |\alpha|\}$, the norm on $\mathbb{R}^m \times \mathbb{R}^m$ is $\max\{\|z\|_2, \|v\|_2\}$, and $\rho > \max\{\text{dist}(0, \text{epi } h^\nu), \text{dist}(0, \text{epi } h)\}$. Then, there are $\alpha, \bar{\rho} \in [0, \infty)$, dependent on ρ , such that*

$$\text{exs}_\rho(\text{gph } \partial h^\nu; \text{gph } \partial h) \leq \hat{d}_\rho(\text{gph } \partial h^\nu, \text{gph } \partial h) \leq \alpha \sqrt{\hat{d}_{\bar{\rho}}(\text{epi } h^\nu, \text{epi } h)} \leq \alpha \sqrt{\sup_{z \in \mathbb{B}(0, \bar{\rho})} |h(z) - h^\nu(z)|}.$$

The truncated Hausdorff distance between epigraphs can be computed using results from [51], with the sup-expression given above being among the crudest possibilities.

6.4 Example (goal optimization; cont.). *In Example 3.1, smoothing of h^ν causes a solution error*

$$\text{exs}_\rho \left((S^\nu)^{-1}(\mathbb{B}(0, \delta^\nu)); S^{-1}(\mathbb{B}(0, \varepsilon)) \right) \leq \beta / \sqrt{\theta^\nu}$$

for a constant β when $0 \leq \delta^\nu \leq \rho < \infty$, $\varepsilon \geq \delta^\nu + \beta / \sqrt{\theta^\nu}$, and $\rho > \frac{1}{2} \max\{\sqrt{\sum_{i=1}^m \min\{0, \tau_i\}^2}, \bar{\alpha} / \theta^\nu\}$, where $\bar{\alpha} = (\ln 2)(\sum_{i=1}^m \alpha_i)$.

Detail. Since $\sup_{z \in \mathbb{R}^m} |h^\nu(z) - h(z)| \leq \bar{\alpha} / \theta^\nu$ by Example 3.1, we obtain via Proposition 6.3 that

$$\text{exs}_{2\rho}(\text{gph } \partial h^\nu; \text{gph } \partial h) \leq \alpha \sqrt{\bar{\alpha} / \theta^\nu}$$

for some $\alpha \in [0, \infty)$ when ρ is sufficiently large as indicated. The claim then follows by Theorem 6.2 and Proposition 6.1. \square

6.5 Example (distributionally robust optimization; cont.). *For Example 3.3, there is a constant $\beta \in [0, \infty)$ such that*

$$\text{exs}_\rho \left((S^\nu)^{-1}(\mathbb{B}(0, \delta^\nu)); S^{-1}(\mathbb{B}(0, \varepsilon)) \right) \leq \beta \sqrt{\alpha^\nu}$$

as long as $0 \leq \delta^\nu \leq \rho < \infty$ and $\varepsilon \geq \delta^\nu + \beta \sqrt{\alpha^\nu}$, where $\alpha^\nu = \max\{\text{exs}(A, A^\nu), \text{exs}(A^\nu, A)\}$.

Detail. Since $\text{dist}(0, \text{epi } h^\nu) = \text{dist}(0, \text{epi } h) = 0$, it follows by Proposition 6.3 that

$$\hat{d}_{2\rho}(\text{gph } \partial h^\nu, \text{gph } \partial h) \leq \alpha \sqrt{\sup_{z \in \mathbb{B}(0, \bar{\rho})} |h(z) - h^\nu(z)|} \tag{6.3}$$

for some $\alpha, \bar{\rho} \in [0, \infty)$. Fix $z \in \mathbb{B}(0, \bar{\rho})$ and let $p^\nu \in \hat{A}^\nu(z)$; see Example 3.3 for notation. Then, there exists $\bar{p} \in A$ such that $\|p^\nu - \bar{p}\|_2 \leq \alpha^\nu$. Thus, $h^\nu(z) - h(z) \leq \langle p^\nu - \bar{p}, z \rangle \leq \|p^\nu - \bar{p}\|_2 \|z\|_2 \leq \bar{\rho} \alpha^\nu$. We repeat this argument with the roles of h and h^ν reversed and conclude that $|h(z) - h^\nu(z)| \leq \bar{\rho} \alpha^\nu$. In combination with (6.3), Theorem 6.2, and Proposition 6.1, the claim follows. \square

6.6 Example (augmented Lagrangian methods; cont.). For Example 3.4, we obtain that

$$\text{exs}_\rho \left((S^\nu)^{-1}(\mathbb{B}(0, \delta^\nu)); S^{-1}(\mathbb{B}(0, \varepsilon)) \right) \leq \beta^\nu / \theta^\nu, \quad \text{where } \beta^\nu = (2\rho + \|y^\nu\|_\infty) \sqrt{m-1},$$

provided that $0 \leq \delta^\nu \leq \rho < \infty$ and $\varepsilon \geq \delta^\nu + \beta^\nu / \theta^\nu$.

Detail. In this case, $\text{gph } \partial h = (\mathbb{R} \times \{0\}^{m-1}) \times (\{1\} \times \mathbb{R}^{m-1})$ and

$$\text{gph } \partial h^\nu = \{(z, v) \in \mathbb{R}^{2m} \mid v = (1, y_2^\nu + \theta^\nu z_2, \dots, y_m^\nu + \theta^\nu z_m)\}.$$

In view Proposition 6.1 and Theorem 6.2, it suffices to bound $\text{exs}_{2\rho}(\text{gph } \partial h^\nu; \text{gph } \partial h)$. Let $(z^\nu, v^\nu) \in \text{gph } \partial h^\nu \cap \mathbb{B}(0, 2\rho)$. We construct $(\bar{z}, \bar{v}) \in \text{gph } \partial h$ by setting $\bar{v}_1 = 1$, $\bar{z}_1 = z_1^\nu$, $\bar{z}_i = 0$, $\bar{v}_i = y_i^\nu + \theta^\nu z_i^\nu$ for $i = 2, \dots, m$. Since $\|v^\nu\|_2 \leq 2\rho$, one has $|v_i^\nu| \leq 2\rho$ and $|y_i^\nu + \theta^\nu z_i^\nu| \leq 2\rho$ for $i = 2, \dots, m$. Moreover, $\|v^\nu - \bar{v}\|_2^2 = 0$ and

$$\|z^\nu - \bar{z}\|_2^2 = \sum_{i=2}^m (z_i^\nu - \bar{z}_i)^2 = \sum_{i=2}^m (z_i^\nu)^2 \leq \sum_{i=2}^m \left(\frac{2\rho + |y_i^\nu|}{\theta^\nu} \right)^2 \leq (\beta^\nu / \theta^\nu)^2.$$

Thus, the distance (in the appropriate norm) between (z^ν, v^ν) and (\bar{z}, \bar{v}) is at most β^ν / θ^ν . \square

6.7 Example (exact penalty methods). We next approach the problem in Example 3.4 using an exact penalty method. Thus, $h(z) = z_1 + \sum_{i=2}^m \iota_{\{0\}}(z_i)$ as before, but for $\theta^\nu \in [0, \infty)$ we set

$$h^\nu(z) = z_1 + \sum_{i=2}^m \theta^\nu |z_i|.$$

If $\theta^\nu \rightarrow \infty$, then the resulting approximations are consistent; the details are omitted. Interestingly,

$$(S^\nu)^{-1}(\mathbb{B}(0, \delta^\nu)) \cap \mathbb{B}(0, \rho) \subset S^{-1}(\mathbb{B}(0, \varepsilon))$$

provided that $0 \leq \delta^\nu \leq \rho < \infty$, $\varepsilon \geq \delta^\nu$, and $\theta^\nu \geq 2\rho$. This means that the approximating problem has a certain ‘‘exactness’’ property: for sufficiently large θ^ν , satisfaction of the optimality condition for the approximation problem locally in $\mathbb{B}(0, \rho)$ implies satisfaction of the condition for the actual problem.

Detail. Example 6.6 furnishes $\text{gph } \partial h$. Moreover, $\partial h^\nu(z) = \{1\} \times C_2 \times \dots \times C_m$, where $C_i = \{\theta^\nu\}$ if $z_i > 0$, $C_i = [-\theta^\nu, \theta^\nu]$ if $z_i = 0$, and $C_i = \{-\theta^\nu\}$ otherwise.

Suppose that $\theta^\nu \geq 2\rho$. Let $(z^\nu, v^\nu) \in \text{gph } \partial h^\nu \cap \mathbb{B}(0, 2\rho)$. Then, $v_1^\nu = 1$ and $v_i^\nu \in C_i$ for $i = 2, \dots, m$. Since $\|v^\nu\|_2 \leq 2\rho$, one has $|v_i^\nu| < 2\rho \leq \theta^\nu$ for $i = 2, \dots, m$. This means that $z_i^\nu = 0$ for such i in view of the definition of C_i . We conclude that $(z^\nu, v^\nu) \in \text{gph } \partial h$ and $\text{exs}_{2\rho}(\text{gph } \partial h^\nu; \text{gph } \partial h) = 0$. This in turn implies that $\text{exs}_\rho(\text{gph } S^\nu; \text{gph } S) = 0$ by Theorem 6.2. The claim follows from Proposition 6.1. \square

6.8 Example (homotopy method; cont.). In Example 3.10, we obtain for $\lambda^\nu \in (0, 1)$ that

$$\text{exs}_\rho \left((S^\nu)^{-1}(\mathbb{B}(0, \delta^\nu)); S^{-1}(\mathbb{B}(0, \varepsilon)) \right) \leq \beta^\nu \lambda^\nu, \quad \text{where } \beta^\nu = \sqrt{1 + \frac{4\rho^2 - (\lambda^\nu)^2}{(1 - \lambda^\nu)^2}},$$

provided that $0 \leq \delta^\nu \leq \rho < \infty$, $\varepsilon \geq \delta^\nu + \beta^\nu \lambda^\nu$, and $\rho \geq \lambda^\nu / 2$.

Detail. In this case, $\partial h(z) = \partial \hat{h}(z_1, \dots, z_{m-1}) \times \{0\}$ and $\partial h^\nu(z) = (1 - \lambda^\nu) \partial \hat{h}(z_1, \dots, z_{m-1}) \times \{\lambda^\nu\}$. To bound $\text{exs}_{2\rho}(\text{gph } \partial h^\nu; \text{gph } \partial h)$, let $(z^\nu, v^\nu) \in \text{gph } \partial h^\nu \cap \mathbb{B}(0, 2\rho)$. Then $v^\nu = ((1 - \lambda^\nu)\hat{y}^\nu, \lambda^\nu)$ with $\hat{y}^\nu \in \partial \hat{h}(z_1^\nu, \dots, z_{m-1}^\nu)$. We construct $(\bar{z}, \bar{v}) \in \text{gph } \partial h$ by setting $\bar{v} = (\hat{y}^\nu, 0)$ and $\bar{z} = z^\nu$. Then, one has $\|z^\nu - \bar{z}\|_2 = 0$ and $\|v^\nu - \bar{v}\|_2^2 = (1 + \|\hat{y}^\nu\|_2^2)(\lambda^\nu)^2$. Since $\|v^\nu\|_2^2 = (1 - \lambda^\nu)^2 \|\hat{y}^\nu\|_2^2 + (\lambda^\nu)^2 \leq 4\rho^2$, we obtain that $1 + \|\hat{y}^\nu\|_2^2 \leq (\beta^\nu)^2$. From this we conclude that $\text{exs}_{2\rho}(\text{gph } \partial h^\nu; \text{gph } \partial h) \leq \beta^\nu \lambda^\nu$. The result then follows by Theorem 6.2 and Proposition 6.1. \square

Acknowledgement. The author is thankful to K. Balasubramanian (UC Davis) for introducing him to the inverse problem in Section 5. The research is supported in part by ONR (N0001421WX01496) and AFOSR (18RT0599).

References

- [1] H. Attouch and R. J-B Wets. Quantitative stability of variational systems: I. The epigraphical distance. *Transactions of the American Mathematical Society*, 328(2):695–729, 1991.
- [2] H. Attouch and R. J-B Wets. Quantitative stability of variational systems: II. A framework for nonlinear conditioning. *SIAM J. Optimization*, 3:359–381, 1993.
- [3] H. Attouch and R. J-B Wets. Quantitative stability of variational systems: III. ε -approximate solutions. *Mathematical Programming*, 61:197–214, 1993.
- [4] S. Basu, P. Pope, and S. Feizi. Influence functions in deep learning are fragile. *arXiv:2006.14651*, 2020.
- [5] A. Beck and M. Teboulle. Smoothing and first order methods: A unified framework. *SIAM J. Optimization*, 22(2):557–580, 2012.
- [6] A. Ben-Tal and A. Nemirovski. On polyhedral approximations of the second-order cone. *Mathematics of Operations Research*, 26(2):193–205, 2001.
- [7] J.F. Bonnans and A. Shapiro. *Perturbation Analysis of Optimization Problems*. Springer, 2000.
- [8] J.V. Burke. Descent methods for composite nondifferentiable optimization problems. *Mathematical Programming*, 33(3):260–279, 1985.
- [9] J.V. Burke and M.C. Ferris. A Gauss-Newton method for convex composite optimization. *Mathematical Programming*, 71:179–194, 1995.
- [10] J.V. Burke and T. Hoheisel. Epi-convergent smoothing with applications to convex composite functions. *SIAM J. Optimization*, 23(3):1457–1479, 2013.
- [11] J.V. Burke and T. Hoheisel. Epi-convergence properties of smoothing by infimal convolution. *Set-Valued and Variational Analysis*, 25:1–23, 2017.

- [12] J.V. Burke, T. Hoheisel, and C. Kanzow. Gradient consistency for integral-convolution smoothing functions. *Set-Valued and Variational Analysis*, 21(2):359–376, 2013.
- [13] J.V. Burke, T. Hoheisel, and Q.V. Nguyen. A study of convex convex-composite functions via infimal convolution with applications. *Mathematics of Operations Research*, 46(4):1324–1348, 2021.
- [14] X. Chen. Smoothing methods for nonsmooth, nonconvex minimization. *Mathematical Programming*, 134:71–99, 2012.
- [15] X. Chen and O. L. Mangasarian. A class of smoothing functions for nonlinear and mixed complementarity problems. *Mathematical Programming*, 71:51–70, 1995.
- [16] Y. Cui and J.-S. Pang. *Modern Nonconvex Nondifferentiable Optimization*. SIAM, 2021.
- [17] Y. Cui, J.-S. Pang, and B. Sen. Composite difference-max programs for modern statistical estimation problems. *SIAM J. Optimization*, 28(4):3344–3374, 2018.
- [18] D. Davis and D. Drusvyatskiy. Stochastic model-based minimization of weakly convex functions. *SIAM J. Optimization*, 29(1):207–239, 2019.
- [19] D. Drusvyatskiy and A.S. Lewis. Tilt stability, uniform quadratic growth, and strong metric regularity of the subdifferential. *SIAM J. Optimization*, 23:256–267, 2013.
- [20] D. Drusvyatskiy and A.S. Lewis. Error bounds, quadratic growth, and linear convergence of proximal methods. *Mathematics of Operations Research*, 43(3):919–948, 2018.
- [21] D. Drusvyatskiy and C. Paquette. Efficiency of minimizing compositions of convex functions and smooth maps. *Mathematical Programming*, 178:503–558, 2019.
- [22] J.C. Duchi and F. Ruan. Stochastic methods for composite and weakly convex optimization problems. *SIAM J. Optimization*, 28(4):3229–3259, 2018.
- [23] A.C. Eberhard and R. Wenczel. A study of tilt-stable optimality and sufficient conditions. *Nonlinear Analysis*, 75:1260–1281, 2012.
- [24] R. Fletcher. Nondifferential and variational techniques in optimization. In D.C. Sorensen and R. J-B Wets, editors, *Nondifferential and variational techniques in optimization*, Mathematical Programming Study 17, page 67–76. Academic Press, 1982.
- [25] J.C. Foraker, J.O. Royset, and I. Kaminer. Search-trajectory optimization: Part 1, formulation and theory. *J. Optimization Theory and Applications*, 169(2):530–549, 2016.
- [26] P. Hand and V. Voroninski. Global guarantees for enforcing deep generative priors by empirical risk. In *31st Annual Conference On Learning Theory*, 2018.
- [27] J.L. Hige and S. Sen. Epigraphical nesting: A unifying theory for the convergence of algorithms. *J. Optimization Theory and Applications*, 84(2):339–360, 1995.

- [28] W. Huang, P. Hand, R. Heckel, and V. Voroninski. A provably convergent scheme for compressive sensing under random generative priors. *J. Fourier Analysis and Applications*, 27(2):1–34, 2021.
- [29] A.D. Ioffe and J.V. Outrata. On metric and calmness qualification conditions in subdifferential calculus. *Set-Valued and Variational Analysis*, 16(2-3):199–227, 2008.
- [30] D. Klatte, A. Kruger, and B. Kummer. From convergence principles to stability and optimality conditions. *J. Convex Analysis*, 19(4):1043–1072, 2012.
- [31] P.W. Koh and P. Liang. Understanding black-box predictions via influence functions. *arXiv:1703.04730*, 2017.
- [32] D.P. Kouri and T.M. Surowiec. Epi-regularization of risk measures. *Mathematics of Operations Research*, 45(2):774–795, 2020.
- [33] A.S. Lewis and S.J. Wright. A proximal method for composite minimization. *Mathematical Programming*, 158:501–546, 2016.
- [34] A.S. Lewis and S. Zhang. Partial smoothness, tilt stability, and generalized Hessians. *SIAM J. Optimization*, 23:74–94, 2013.
- [35] B.S. Mordukhovich. *Variational Analysis and Generalized Differentiation I: Basic Theory*. Grundlehren der mathematischen Wissenschaften. Springer, 2. edition, 2013.
- [36] B.S. Mordukhovich, R.T. Rockafellar, and M.E. Sarabi. Characterizations of full stability in constrained optimization. *SIAM J. Optimization*, 23:1810–1849, 2013.
- [37] Y. Nesterov. Smooth minimization of non-smooth functions. *Mathematical Programming*, 103:127–152, 2005.
- [38] J.P. Penot. Error bounds, calmness and their applications in nonsmooth analysis. *Contemporary Mathematics*, 514:225–247, 2010.
- [39] C. Phelps, J.O. Royset, and Q. Gong. Optimal control of uncertain systems using sample average approximations. *SIAM J. Control and Optimization*, 54(1):1–29, 2016.
- [40] E. Polak. *Optimization. Algorithms and Consistent Approximations*, volume 124 of *Applied Mathematical Sciences*. Springer, 1997.
- [41] E. Polak, J.O. Royset, and R.S. Womersley. Algorithms with adaptive smoothing for finite minimax problems. *J. Optimization Theory and Applications*, 119(3):459–484, 2003.
- [42] R.A. Poliquin. An extension of Attouch’s theorem and its application to second-order epi-differentiation of convexly composite functions. *Transactions of the American Mathematical Society*, 332(2):861–874, 1992.

- [43] M.J.D. Powell. Algorithms for nonlinear constraints that use Lagrangian functions. *Mathematical Programming*, 14(2):224–248, 1978.
- [44] M.J.D. Powell. A fast algorithm for nonlinearly constrained optimization calculations. In G.A. Watson, editor, *Numerical Analysis*, Lecture Notes in Mathematics, vol. 630, page 144–157. Springer, 1978.
- [45] M.J.D. Powell. General algorithms for discrete nonlinear approximation calculations. In *Approximation Theory, IV*, page 187–218. Academic Press, 1983.
- [46] M.J.D. Powell. On the global convergence of trust region algorithms for unconstrained minimization. *Mathematical Programming*, 29(3):299–303, 1984.
- [47] R.T. Rockafellar. Extended nonlinear programming. In G. Di Pillo and R. Giannessi, editors, *Nonlinear Optimization and Related Topics*, pages 381–399. Kluwer, 1999.
- [48] R.T. Rockafellar and R. J-B Wets. *Variational Analysis*. Springer, 3rd printing-2009 edition, 1998.
- [49] J.O. Royset. Optimality functions in stochastic programming. *Mathematical Programming*, 135(1):293–321, 2012.
- [50] J.O. Royset. Approximations and solution estimates in optimization. *Mathematical Programming*, 170(2):479–506, 2018.
- [51] J.O. Royset. Stability and error analysis for optimization and generalized equations. *SIAM J. Optimization*, 30(1):752–780, 2020.
- [52] J.O. Royset and E.Y. Pee. Rate of convergence analysis of discretization and smoothing algorithms for semi-infinite minimax problems. *J. Optimization Theory and Applications*, 155(3):855–882, 2012.
- [53] J.O. Royset and R. J-B Wets. *An Optimization Primer*. Springer, forthcoming.
- [54] J.O. Royset and R. J-B Wets. Optimality functions and lopsided convergence. *J. Optimization Theory and Applications*, 169(3):965–983, 2016.
- [55] Y. Yuan. On the superlinear convergence of a trust region algorithm for nonsmooth optimization. *Mathematical Programming*, 31(3):269–285, 1985.