


# Covering Many (or Few) Edges with $k$ Vertices in Sparse Graphs\*

Tomohiro Koana ✉ 


Technische Universität Berlin, Algorithmics and Computational Complexity, Germany

Christian Komusiewicz ✉ 

Philipps-Universität Marburg, Fachbereich Mathematik und Informatik, Marburg, Germany

André Nichterlein ✉ 

Technische Universität Berlin, Algorithmics and Computational Complexity, Germany

Frank Sommer ✉ 

Philipps-Universität Marburg, Fachbereich Mathematik und Informatik, Marburg, Germany

---

## Abstract

---

We study the following two fixed-cardinality optimization problems (a maximization and a minimization variant). For a fixed  $\alpha$  between zero and one we are given a graph and two numbers  $k \in \mathbb{N}$  and  $t \in \mathbb{Q}$ . The task is to find a vertex subset  $S$  of exactly  $k$  vertices that has value at least (resp. at most for minimization)  $t$ . Here, the value of a vertex set computes as  $\alpha$  times the number of edges with exactly one endpoint in  $S$  plus  $1 - \alpha$  times the number of edges with both endpoints in  $S$ . These two problems generalize many prominent graph problems, such as DENSEST  $k$ -SUBGRAPH, SPARSEST  $k$ -SUBGRAPH, PARTIAL VERTEX COVER, and MAX  $(k, n - k)$ -CUT.

In this work, we complete the picture of their parameterized complexity on several types of sparse graphs that are described by structural parameters. In particular, we provide kernelization algorithms and kernel lower bounds for these problems. A somewhat surprising consequence of our kernelizations is that PARTIAL VERTEX COVER and MAX  $(k, n - k)$ -CUT not only behave in the same way but that the kernels for both problems can be obtained by the same algorithms.

**2012 ACM Subject Classification** Theory of computation  $\rightarrow$  Graph algorithms analysis; Theory of computation  $\rightarrow$  Parameterized complexity and exact algorithms


**Keywords and phrases** Parameterized Complexity, Kernelization, Partial Vertex Cover, Densest  $k$ -Subgraph, Max  $(k, n - k)$ -Cut, Degeneracy

**Funding** *Tomohiro Koana*: Supported by the Deutsche Forschungsgemeinschaft (DFG), project DiPa, NI 369/21.

*Frank Sommer*: Supported by the Deutsche Forschungsgemeinschaft (DFG), project EAGR, KO 3669/6-1.

---

\* An extended abstract appears in the Proceedings of the 39th International Symposium on Theoretical Aspects of Computer Science (STACS '22). This version contains all missing proofs.



■ **Figure 1** Problem definition cheat sheet.

## 1 Introduction

Fixed-cardinality optimization problems are a well-studied class of graph problems where one seeks for a given graph  $G$ , a vertex set  $S$  of size  $k$  such that  $S$  optimizes some objective function  $\text{val}_G(S)$  [7, 9, 8, 29]. Prominent examples of these problems are DENSEST  $k$ -SUBGRAPH [6, 17, 29], SPARSEST  $k$ -SUBGRAPH [20, 21, 36], PARTIAL VERTEX COVER [1, 19, 25], and MAX  $(k, n - k)$ -CUT [8, 34, 35].

A common thread in these examples is that all these problems are formulated in terms of the number of edges that have one or two endpoints in  $S$ : In the decision version of DENSEST  $k$ -SUBGRAPH we require that there are at least  $t$  edges with both endpoints in  $S$ ; CLIQUE is the special case where  $t = \binom{k}{2}$ . Conversely, in SPARSEST  $k$ -SUBGRAPH we require that at most  $t$  edges have both endpoints in  $S$  and INDEPENDENT SET is the special case with  $t = 0$ . In PARTIAL VERTEX COVER we require that at least  $t$  edges have at least one endpoint in  $S$ . Finally, in MAX  $(k, n - k)$ -CUT we require that at least  $t$  edges have exactly one endpoint in  $S$ .

We study the following general problem first defined by Bonnet et al. [4] that contains all of the above problems as special case.<sup>1</sup>

MAX  $\alpha$ -FIXED CARDINALITY GRAPH PARTITIONING (MAX  $\alpha$ -FCGP)

**Input:** A graph  $G$ ,  $k \in \mathbb{N}$ , and  $t \in \mathbb{Q}$ .

**Question:** Is there a set  $S$  of exactly  $k$  vertices such that

$$\text{val}(S) := (1 - \alpha) \cdot m(S) + \alpha \cdot m(S, V(G) \setminus S) \geq t ?$$

Here,  $\alpha \in [0, 1]$ , and  $m(S)$  denotes the number of edges with two endpoints in  $S$  and  $m(S, V(G) \setminus S)$  denotes the number of edges with exactly one endpoint in  $S$ . Naturally, one may also consider the minimization problem, denoted as MIN  $\alpha$ -FIXED CARDINALITY GRAPH PARTITIONING (MIN  $\alpha$ -FCGP), where we are looking for a set  $S$  such that  $\text{val}(S) \leq t$ .

The value of  $\alpha$  describes how strongly edges with exactly one endpoint in  $S$  influence the value of  $S$  relative to edges with two endpoints in  $S$ . For  $\alpha = 1/3$ , edges with two endpoints in  $S$  count twice as much as edges with one endpoint in  $S$  and, thus, every vertex contributes exactly its degree to the value of  $S$ . Hence, in this case, we simply want to find a vertex set with a largest or smallest degree sum.

More importantly, MAX  $\alpha$ -FCGP and MIN  $\alpha$ -FCGP contain all of the above-mentioned problems as special cases (see Figure 1). For example, PARTIAL VERTEX COVER (MAXPVC) is MAX  $\alpha$ -FCGP with  $\alpha = 1/2$  as all edges with at least one endpoint in  $S$  count the same. MAX  $(k, n - k)$ -CUT is MAX  $\alpha$ -FCGP with  $\alpha = 1$  since edges with both endpoints in  $S$  are

<sup>1</sup> On the face of it, the definition of Bonnet et al. [4] seems to be more general as it has separate weight parameters for the internal and outgoing edges. It can be reduced to our formulation by adapting the value of  $t$  and thus our results also hold for this formulation.




Figure 2 Overview over our results. Each box displays the parameterized results (see also bottom right) with respect to  $k$  and the corresponding parameter  $p$  for all variants (maximization, minimization, and all  $\alpha \in [0, 1]$ , see bottom left). Note that the split of the boxes is not proportional to the corresponding values of  $\alpha$ . See Section 2 (paragraph “Graph parameter definitions.”) for the definitions of the parameters. A line from a box for parameter  $p$  to a box above for parameter  $p'$  implies that  $p \in \mathcal{O}(p')$  on all graphs. Thus, hardness results hold also for all parameters below and tractability results for all parameters above.

ignored. SPARSEST  $k$ -SUBGRAPH is MIN  $\alpha$ -FCGP with  $\alpha = 0$  as only the edges with both endpoints in  $S$  count. Consequently, there exist values of  $\alpha$  such that MAX  $\alpha$ -FCGP and MIN  $\alpha$ -FCGP are NP-hard and W[1]-hard on general graphs with respect to the natural parameter  $k$  [8, 10, 14, 19]. This hardness makes it interesting to study these problems on input graphs with special structure and Bonnet et al. [4] and Shachnai and Zehavi [35] studied this problem on bounded-degree graphs.

We continue this line of research and give a complete picture of the parameterized complexity of MIN  $\alpha$ -FCGP and MAX  $\alpha$ -FCGP on several types of sparse graphs that are described by structural parameters. In particular, we provide kernelization algorithms and kernel lower bounds for these problems, see Figure 2 for an overview.

**Known results.** MAXPVC can be solved in  $\mathcal{O}^*((\Delta + 1)^k)$  time where  $\Delta$  is the maximum degree of the input graph [33]. For the degeneracy  $d$ , Amini et al. [1] gave an  $\mathcal{O}^*((dk)^k)$ -time algorithm which was recently improved to an algorithm with running time  $\mathcal{O}^*(2^{\mathcal{O}(dk)})$  [31]. Bonnet et al. [4] showed that in  $\mathcal{O}^*(\Delta^k)$  time one can solve MAX  $\alpha$ -FCGP for all  $\alpha > 1/3$  and MIN  $\alpha$ -FCGP for all  $\alpha < 1/3$ . Bonnet et al. [4] call these two problem cases *degrading*. This name reflects the fact that in MAX  $\alpha$ -FCGP with  $\alpha > 1/3$ , adding a vertex  $v$  to a set  $S$  increases the value at least as much as adding  $v$  to a superset of  $S$ .<sup>2</sup> This is because here one edge with both endpoints in  $S$  is less valuable than two edges each with one endpoint in  $S$ . In MIN  $\alpha$ -FCGP this effect is reversed since we aim to minimize val. The other problem cases are called *non-degrading*. For non-degrading problems, Bonnet et al. [4] achieved a running time of  $\mathcal{O}^*((\Delta k)^{\mathcal{O}(k)})$  and asked whether they can also be solved in  $\mathcal{O}^*(\Delta^{\mathcal{O}(k)})$  time. This question was answered positively by Shachnai and Zehavi [35], who showed that MAX

<sup>2</sup> Note that this matches the definition of submodularity.

$\alpha$ -FCGP and MIN  $\alpha$ -FCGP can be solved in  $\mathcal{O}^*(4^{k+o(k)}\Delta^k)$  time.

Kernelization has been studied only for special cases. MAX  $(k, n - k)$ -CUT admits a polynomial problem kernel when parameterized by  $t$  [34]. This also gives a polynomial kernel for  $k + \Delta$  since instances with  $t > \Delta k$  are trivial no-instances. It is also known that SPARSEST  $k$ -SUBGRAPH admits a kernel with  $\gamma k^2$  vertices [26]. Here,  $\gamma$  is a parameter bounded by  $\max(c, d + 1)$  [18]. In contrast, DENSEST- $k$  SUBGRAPH is unlikely to admit a polynomial problem kernel when parameterized by  $\Delta + k$  since CLIQUE is a special case.

More broadly, for graph problems that are W[1]-hard for the standard parameter solution  $k$  size, the study of kernelization on sparse input graphs has received much attention in recent years [11, 12, 28, 32].

Independent of our work, a polynomial *compression* for MAXPVC (the special case of MAX  $\alpha$ -FCGP with  $\alpha = 1/2$ ) of size  $(dk)^{\mathcal{O}(d)}$  was recently discovered by Panolan and Yaghoubzade [31].

**Our results.** We provide a complete picture of the parameterized complexity of MAX  $\alpha$ -FCGP and MIN  $\alpha$ -FCGP for all  $\alpha$  with respect to the combination of  $k$  and five parameters describing the graph structure: the maximum degree  $\Delta$  of  $G$ , the  $h$ -index of  $G$ , the degeneracy of  $G$ , the  $c$ -closure of  $G$ , and the vertex cover number  $\text{vc}$  of  $G$ . With the exception of the  $c$ -closure, all parameters are sparseness measures. The  $c$ -closure, first described by Fox et al. [18], measures how strongly a graph adheres to the triadic closure principle. Informally, the closure of a graph is small whenever all vertices with many common neighbors are also neighbors of each other. For a formal definition of all parameters refer to Section 2.

Our results are summarized by Figure 2. On a very general level, our main finding suggests that the degrading problems are much more amenable to FPT algorithms and kernelizations than their non-degrading counterparts. No such difference is observed when considering the running time of FPT algorithms for the parameter  $k + \Delta$  but it becomes striking in the context of kernelization and when using secondary parameters that are smaller than  $\Delta$ . Given the importance of the distinction between the degrading and non-degrading cases, we distinguish these subcases of MAX  $\alpha$ -FCGP and MIN  $\alpha$ -FCGP by name (DEGRADING MAX  $\alpha$ -FCGP, NON-DEGRADING MAX  $\alpha$ -FCGP, DEGRADING MIN  $\alpha$ -FCGP, NON-DEGRADING MIN  $\alpha$ -FCGP).

On a technical level, by introducing an annotated version of the problem that keeps track of removed vertices, we separate and unify arguments that deal with vertices identified as (not) being part of a solution. In particular, we show that by introducing vertex weights (called *counter*) we can deal with vertices whose contribution is substantially below or above the average contribution that is necessary to reach the threshold  $t$ . More precisely, if the contribution of a vertex  $v$  is much above  $t/k$ , then we can add  $v$  to the solution and if it is much below  $t/k$ , then we can remove  $v$ . As a consequence, we can show that the weights can be bounded in the maximum degree of the annotated instance. This gives the kernels for the parameter  $k + \Delta$ .

The main step in the more sophisticated kernelizations for the degeneracy  $d$  and the  $c$ -closure is now to decrease the maximum degree of the instance as this allows us to use the kernel for  $k + \Delta$ . To decrease the maximum degree, for these parameters, we make use of Ramsey bounds. More precisely, the Ramsey bounds help to find a large independent set  $I$  such that all vertices outside of  $I$  have only a bounded number of neighbors in  $I$ . This then allows to prove by pigeonholing the following for the vertex  $v$  of  $I$  with the currently worst contribution to the objective function: No matter what the optimal solution selects outside of  $I$ , there is always some vertex of  $I \setminus \{v\}$  that gives at least as good a contribution to the

final solution as  $v$ . For the parameter  $c$ , we also need an additional pigeonhole argument excluding large cliques in order to apply the Ramsey bound. For the parameter  $d$ , we establish a new constructive Ramsey bound for  $K_{i,j}$ -free graphs that may be of independent interest.

We remark that when we describe the kernel size for  $\alpha > 0$  (for instance, Proposition 4.2), the factor  $1/\alpha$  is hidden in the  $\mathcal{O}$  notation. We would like to emphasize, however, that the exponents in the kernel size such as  $\mathcal{O}(c)$  and  $\mathcal{O}(d)$  do not depend on  $1/\alpha$ . On the other hand, the lower bounds such as Theorem 4.11 hold indeed for all  $\alpha$  in the range corresponding to the case.

We believe that this general approach could be useful for other parameterizations that are not considered in this work. A somewhat surprising consequence of our kernelizations is that PARTIAL VERTEX COVER and MAX  $(k, n - k)$ -CUT not only behave in the same way but that the kernels for both problems can also be obtained by the same algorithms.

Several proofs (marked with  $(\star)$ ) are deferred to the appendix.

## 2 Preliminaries

For  $q \in \mathbb{N}$ , we write  $[q]$  to denote the set  $\{1, 2, \dots, q\}$ . For a graph  $G$ , we denote its *vertex set* by  $V(G)$  and its *edge set* by  $E(G)$ . Let  $X, Y \subseteq V(G)$  be vertex subsets. We use  $G[X]$  to denote the *subgraph induced* by  $X$ . We let  $G - X$  denote the graph obtained by *removing* the vertices in  $X$ . We denote by  $N_G(X) := \{y \in V(G) \setminus X \mid xy \in E(G), x \in X\}$  the *open neighborhood* and by  $N_G[X] := N_G(X) \cup X$  the *closed neighborhood* of  $X$ . By  $E_G(X, Y) := \{xy \in E(G) \mid x \in X, y \in Y\}$  we denote the set of edges *between*  $X$  and  $Y$ . As a shorthand, we set  $E_G(X) := E_G(X, X)$ . Furthermore, we denote by  $m_G(X, Y) := |E_G(X, Y)|$  and  $m_G(X) := |E_G(X)|$  the *sizes* of these edge sets. For all these notations, when  $X$  is a singleton  $\{x\}$  we may write  $x$  instead of  $\{x\}$ . Let  $v \in V(G)$ . We denote the *degree* of  $v$  by  $\deg_G(v)$ . We drop the subscript  $\cdot_G$  when it is clear from context.

**Graph parameter definitions.** For more information on parameterized complexity, we refer to the standard monographs [10, 14]. We denote the size of a smallest vertex cover (a set of vertices that covers all edges) of a graph  $G$  by  $\text{vc}_G$ . The maximum and minimum degree of  $G$  are  $\Delta_G := \max_{v \in V(G)} \deg_G(v)$  and  $\delta_G := \min_{v \in V(G)} \deg_G(v)$ , respectively. The degeneracy of  $G$  is  $d_G := \max_{S \subseteq V(G)} \delta_{G[S]}$ . The  *$h_G$ -index* of a graph  $G$  is the largest integer  $h$  such that  $G$  has at least  $h$  vertices of degree at least  $h$  [16]. We say that  $G$  is  $c$ -closed for  $c := \max(\{0\} \cup \{|N_G(u) \cap N_G(v)| \mid uv \notin E(G)\}) + 1$  [18].

**Ramsey numbers.** Ramsey's theorem states that for every  $p, q \in \mathbb{N}$ , there exists an integer  $R(p, q)$  such that any graph on at least  $R(p, q)$  vertices contains either a clique of size  $p$  or an independent set of size  $q$ . The numbers  $R(p, q)$  are referred to as *Ramsey numbers*. Although the precise values of Ramsey numbers are not known, some upper bounds have been proven. For instance, it holds that  $R(p, q) \leq \binom{p+q-2}{p-1}$  (see e.g. [24]). The proof for this upper bound is constructive. More precisely, given a graph  $G$  on at least  $\binom{p+q-2}{p-1}$  vertices, we can find in time  $n^{\mathcal{O}(1)}$  either a clique of size  $p$  or an independent set of size  $q$ .

## 3 A Data Reduction Framework via Annotation

In this section, we introduce an annotated variant which allows easier handling for kernelization by giving more options for encoding information in the instances. Moreover, to avoid

repeating certain basic arguments, we provide general data reduction rules and statements used in the subsequent sections. Finally, we describe how to reduce from the annotated to the non-annotated problem variants in polynomial time.

In the annotated problem variant, we have additionally as input a (possibly empty) partial solution  $T \subseteq V(G)$  and **counter**:  $V \rightarrow \mathbb{N}$  which encodes for each vertex, the number of neighbors in the original graph that are guaranteed to be not in a solution. For a set  $S \subseteq V(G)$ , we set

- $\text{counter}(S) := \sum_{v \in S} \text{counter}(v)$  and
- $\text{val}_G(S) := \alpha(m(S, V(G) \setminus S) + \text{counter}(S)) + (1 - \alpha)m(S)$ .

For  $v \in S$  we set  $\text{deg}^{+c}(v) := \text{deg}(v) + \text{counter}(v)$ .

ANNOTATED MAX  $\alpha$ -FCGP

**Input:** A graph  $G$ ,  $T \subseteq V(G)$ , **counter**:  $V(G) \rightarrow \mathbb{N}$ ,  $k \in \mathbb{N}$ , and  $t \in \mathbb{Q}$ .

**Question:** Is there a vertex set  $S$ ,  $T \subseteq S \subseteq V(G)$ , of size  $k$  with  $\text{val}_G(S) \geq t$  (MAX) or  $\text{val}_G(S) \leq t$  (MIN), respectively?

For a partial solution  $T \subseteq V(G)$ , we define the *contribution* of a vertex  $v$ . Note that our definition slightly differs from that of Bonnet et al. [4]:

$$\begin{aligned} \text{cont}(v, T) &:= \alpha \cdot (|N(v) \setminus T| + \text{counter}(v)) + (1 - 2\alpha)|N(v) \cap T| \\ &= \alpha \text{deg}^{+c}(v) + (1 - 3\alpha)|N(v) \cap T| \end{aligned}$$

This definition is chosen so that the value  $\text{val}(S)$  of a vertex set  $S$  computes as follows.

► **Lemma 3.1** (\*). *Let  $G$  be a graph and  $S := \{v_1, \dots, v_n\} \subseteq V(G)$  a vertex set. Then, it holds that  $\text{val}(S) = \sum_{i \in [|S|]} \text{cont}(v_i, \{v_1, \dots, v_{i-1}\})$ .*

**Main reduction rules.** Annotations are helpful for data reductions in the following way: If we identify a vertex  $v$  that is (or is not) in a solution, then, we can simplify the instance as follows using the annotations.

► **Reduction Rule 3.2** (Inclusion Rule). If there is a solution  $S$  with  $v \in S \setminus T$ , then add  $v$  to  $T$ . If there is a vertex  $v \in T$  with  $\text{counter}(v) > 0$ , then reduce  $t$  by  $\alpha \cdot \text{counter}(v)$  and set  $\text{counter}(v) := 0$ .

► **Reduction Rule 3.3** (Exclusion Rule). If there is a solution  $S$  with  $v \notin S$ , then for each  $u \in N(v)$  increase  $\text{counter}(u)$  by one and remove  $v$  from  $G$ .

The reduction rules themselves are simple. The difficulty lies in identifying vertices that are included in or excluded from some solution. In the respective arguments, we use the subsequently discussed notion of better vertices.

**Better vertices.** The following notion captures a situation that frequently appears in our arguments and allows for simple exchange arguments (see following lemma).

► **Definition 3.4.** *A vertex  $v \in V(G)$  is better than  $u \in V(G)$  with respect to a vertex set  $T \subseteq V(G)$  if  $\text{cont}(v, T) \geq \text{cont}(u, T)$  for the maximization variant (if  $\text{cont}(v, T) \leq \text{cont}(u, T)$  for the minimization variant).*

*A vertex  $v \in V(G)$  is strictly better than  $u \in V(G)$  if for all  $T \subseteq V(G)$  of size at most  $k$  we have  $\text{cont}(v, T) \geq \text{cont}(u, T)$  for the maximization variant ( $\text{cont}(v, T) \leq \text{cont}(u, T)$  for the minimization variant).*

When we simply say that  $v$  is better than  $u$ , we mean that  $v$  is better than  $u$  with respect to the empty set. The following lemma immediately follows from Lemma 3.1.

► **Lemma 3.5.** *Let  $S$  be a solution of an instance of  $\alpha$ -FCGP. Suppose that there are two vertices  $v \in S$  and  $v' \notin S$  such that  $v'$  is better than  $v$  with respect to  $S \setminus \{v\}$  or  $v'$  is strictly better than  $v$ . Then,  $S' := (S \setminus \{v\}) \cup \{v'\}$  is also a solution.*

**Proof.** We give a proof for the maximization variant; the minimization variant follows analogously. By Lemma 3.1, we have  $\text{val}(S') = \text{val}(S \setminus \{v\}) + \text{cont}(v', S \setminus \{v\}) \geq \text{val}(S \setminus \{v\}) + \text{cont}(v, S \setminus \{v\}) = \text{val}(S)$ . ◀

Observe that the contribution of any vertex  $v$  differs from  $\alpha \deg^{+c}(v)$  by at most  $|(1-3\alpha)k|$ . This observation allows us to identify some strictly better vertices in the following. This is helpful when we wish to apply the second part of Lemma 3.5 on strictly better vertices.

► **Lemma 3.6** ( $\star$ ). *Let  $u, v \in V(G)$ . Vertex  $v$  is strictly better than  $u$  if*

- (Maximization:)  $\alpha \deg^{+c}(u) \leq \alpha \deg^{+c}(v) - |(1-3\alpha)k|$ .
- (Minimization:)  $\alpha \deg^{+c}(u) \geq \alpha \deg^{+c}(v) + |(1-3\alpha)k|$ .

**Reduction to the non-annotated problem.** The following two lemmas (for maximization and minimization variant respectively) remove annotations and generate an equivalent instance of  $\alpha$ -FCGP. We remark that the resulting instance size depends on  $\Gamma := \max_{v \in V(G)} \text{counter}(v) + 1$ . We obtain an upper bound on  $\Gamma$  in terms of  $k + \Delta$  in the next section.

► **Lemma 3.7.** *Given an instance  $\mathcal{I} := (G, T, \text{counter}, k, t)$  of ANNOTATED MAX  $\alpha$ -FCGP with  $\alpha \in (0, 1]$ , we can compute an equivalent instance  $\mathcal{I}'$  of MAX  $\alpha$ -FCGP of size  $\mathcal{O}((\Delta + \Gamma) \cdot |V(G)| + k \cdot |T|)$  in polynomial time.*

**Proof.** We may assume that  $G$  has at least  $k$  vertices (otherwise the lemma holds for a trivial No-instance  $\mathcal{I}'$ ). We construct an equivalent instance  $\mathcal{I}' := (G', k, t')$  of MAX  $\alpha$ -FCGP. The graph  $G'$  is obtained from  $G$  as follows:

1. Add  $\text{counter}(v) + \lfloor 1/\alpha \rfloor$  degree-one neighbors to every vertex  $v \in V(G)$ .
2. Additionally, add  $\ell := \Delta + \Gamma + \lfloor 1/\alpha - 3 \rfloor \cdot k + \lfloor 1/\alpha \rfloor$  degree-one neighbors to every vertex  $v \in T$ .

By  $L_v$  we denote the set of degree-one vertices added to vertex  $v \in V(G)$  and by  $L := \bigcup_{v \in V(G)} L_v$  we denote the set of all newly added leaf vertices. To conclude the construction of  $\mathcal{I}'$ , we set  $t' := t + \alpha(\ell \cdot |T| + \lfloor 1/\alpha \rfloor \cdot k)$ . Since  $G$  has at most  $\Delta \cdot |V(G)|$  edges and we add at most  $\mathcal{O}((\Gamma + 1) \cdot |V(G)| + (\Delta + k) \cdot |T|)$  edges, we see that  $G'$  has at most  $\mathcal{O}((\Delta + \Gamma) \cdot |V(G)| + k \cdot |T|)$  edges.

Next, we prove the equivalence between  $\mathcal{I}$  and  $\mathcal{I}'$ . For a solution  $S$  of  $\mathcal{I}$ , its value in  $G'$  is increased by  $\alpha \cdot \lfloor 1/\alpha \rfloor$  for every vertex in  $S$  and additionally, by  $\alpha \cdot \ell$  for every vertex in  $T$ , amounting to  $t + \alpha(\ell \cdot |T| + \lfloor 1/\alpha \rfloor \cdot k)$ .

Conversely, consider a solution  $S'$  of  $\mathcal{I}'$ . First, we show that there is a solution containing all vertices of  $T$  and no leaf vertex of  $L$  using Lemma 3.5. Suppose that for some vertex  $v \in V(G)$ , one of its degree-one neighbors  $v' \in L_v$  is in  $S'$  but not  $v$  itself. We then have  $\text{cont}(v', S' \setminus \{v\}) = \alpha$  and  $\text{cont}(v, S' \setminus \{v\}) \geq \alpha$ , implying that  $(S' \setminus \{v'\}) \cup \{v\}$  is also a solution by Lemma 3.5. Thus, in the following we can assume that  $S' \cap L_v = \emptyset$  for every vertex  $v \in V(G) \setminus S'$ . If there is a vertex  $v' \in S' \cap L_v$  for some  $v \in V(G)$ , then by the assumption that  $|V(G)| \geq k$ , the pigeonhole principle gives us a vertex  $w \in V(G) \setminus S'$  with  $S' \cap L_w = \emptyset$ . Since  $|L_w| \geq \text{counter}(w) + \lfloor 1/\alpha \rfloor \geq \lfloor 1/\alpha \rfloor$ , we have  $\text{cont}(w, S' \setminus \{v'\}) \geq \alpha \cdot \lfloor 1/\alpha \rfloor \geq \alpha(1/\alpha -$



$1) = 1 - \alpha$ . We thus have  $\text{cont}(v', S' \setminus \{v'\}) = 1 - \alpha \leq \text{cont}(w, S' \setminus \{v'\})$ . Hence,  $(S' \setminus \{v'\}) \cup \{w\}$  is a solution, again by Lemma 3.5. Thus, in the following, we can assume that  $S' \cap L = \emptyset$ .

So we may assume that  $S'$  consists only of vertices in  $G$ . Suppose that some vertex  $v \in T$  is not in  $S'$ . For any vertex  $v' \in S' \setminus T$ , we have  $\deg(v') \leq \deg_G(v) + \text{counter}(v) + \lfloor 1/\alpha \rfloor \leq \Delta + \Gamma + \lfloor 1/\alpha \rfloor$ . So we have  $\deg(v') \geq \Delta + \Gamma + \lfloor 1/\alpha \rfloor \cdot k + \lfloor 1/\alpha \rfloor \geq \deg(v) + \lfloor 1/\alpha - 3 \rfloor \cdot k$ . Applying Lemma 3.6 with  $\text{counter}(v) = \text{counter}(v') = 0$ , we obtain that  $v$  is strictly better than  $v'$ . Now, it follows from Lemma 3.5 that  $\mathcal{I}'$  has a solution  $S'$  such that  $T \subseteq S' \subseteq V(G')$ . Hence,  $S'$  is also a solution for  $\mathcal{I}$ .  $\blacktriangleleft$

► **Lemma 3.8** ( $\star$ ). *Given an instance  $\mathcal{I} := (G, T, \text{counter}, k, t)$  of ANNOTATED MIN  $\alpha$ -FCGP for  $\alpha \in (0, 1]$ , we can compute an equivalent instance  $\mathcal{I}'$  of MIN  $\alpha$ -FCGP of size  $\mathcal{O}((\Delta + \Gamma)^3 \cdot |V(G)|)$  in polynomial time.*

## 4 Parameterization By Maximum Degree

### 4.1 Polynomial Kernels in Degrading Cases

Recall that in the degrading cases we have  $\alpha \in (1/3, 1]$  for maximization and  $\alpha \in [0, 1/3)$  for minimization. Furthermore, recall that for two vertices  $u$  and  $v$ ,  $v$  is said to be better than  $u$  if  $\text{cont}(v, T) \geq \text{cont}(u, T)$  (vice versa for the minimization variant).

For the annotated version we define  $\Delta_{\overline{T}} := \max_{v \in V(G) \setminus T} \deg(v)$ . Clearly,  $\Delta_{\overline{T}} \leq \Delta$ .

► **Reduction Rule 4.1** ( $\star$ ). Let  $\mathcal{I}$  be an instance of ANNOTATED DEGRADING  $\alpha$ -FCGP. If there are more than  $\Delta_{\overline{T}}k + 1$  vertices that are better than  $v$  with respect to  $T$ , then apply the Exclusion Rule (Reduction Rule 3.3) to  $v$ .

Next, we show that the exhaustive application of Reduction Rule 4.1 yields a polynomial kernel for DEGRADING  $\alpha$ -FCGP.

► **Proposition 4.2.** *DEGRADING  $\alpha$ -FCGP has a kernel of size*

- $\mathcal{O}(\Delta^2 k)$  for maximization and  $\alpha \in (1/3, 1]$ , and
- $\mathcal{O}(\Delta^4 k)$  for minimization and  $\alpha \in (0, 1/3)$ .

**Proof.** Given an instance of DEGRADING  $\alpha$ -FCGP, we transform it into an equivalent instance of ANNOTATED DEGRADING  $\alpha$ -FCGP and apply Reduction Rule 4.1 exhaustively. Observe that  $|V(G)| \leq \Delta_{\overline{T}}k + 1 \leq \Delta k + 1$ . Moreover, we have  $T = \emptyset$  and  $\Gamma \leq \Delta$  since each neighbor of a vertex can increase its counter by at most one. By Lemma 3.7 (maximization) resp. Lemma 3.8 (minimization), we obtain an equivalent instance of  $\alpha$ -FCGP of size  $\mathcal{O}(\Delta^2 k)$  (maximization) resp.  $\mathcal{O}(\Delta^4 k)$  (minimization).  $\blacktriangleleft$

Note that Proposition 4.2 does not cover the case  $\alpha = 0$  for minimization. Note that this problem is referred to as SPARSEST  $k$ -SUBGRAPH. We remark that this is complemented by Proposition 6.4, in which we provide a kernel of size  $\mathcal{O}(d^2 k)$ . Since  $d \leq \Delta$ , this implies also a kernel of size  $\mathcal{O}(\Delta^2 k)$ .

Proposition 4.2 basically shows that given an instance of DEGRADING  $\alpha$ -FCGP, we can find in polynomial time an equivalent instance of DEGRADING  $\alpha$ -FCGP of size  $\mathcal{O}(\Delta + k)^{\mathcal{O}(1)}$ . In the following, we will show that an equivalent instance of DEGRADING  $\alpha$ -FCGP that has size  $(\Delta + k)^{\mathcal{O}(1)}$  can be constructed even if an instance of ANNOTATED DEGRADING  $\alpha$ -FCGP is given (see Proposition 4.10). Proposition 4.10 plays an important role in kernelizations in subsequent sections. Essentially, the task of kernelization boils down to bounding the maximum degree  $\Delta$  by Proposition 4.10.



As shown in the proof of Proposition 4.2, the instance size becomes polynomial in  $k + \Delta$  by exhaustively applying Reduction Rule 4.1. Recall that in Section 3, we presented a polynomial-time procedure to remove annotations with an additional polynomial factor in  $\Delta + \Gamma$  on the instance size, where  $\Gamma$  denotes the maximum counter. To prove Proposition 4.10, it remains to bound  $\Gamma$  for ANNOTATED DEGRADING  $\alpha$ -FCGP.

**Bounding the largest counter  $\Gamma$ .** Throughout the section, let  $k' := k - |T|$  and  $t' := t - \text{val}(T)$ . First, we identify vertices which are contained in a solution, if one exists.

► **Definition 4.3.** Let  $\mathcal{I}$  be a Yes-instance of ANNOTATED DEGRADING  $\alpha$ -FCGP. A vertex  $v \in V(G) \setminus T$  is called *satisfactory* if

- (Maximization:)  $\text{cont}(v, T) \geq t'/k' + (3\alpha - 1)(k - 1)$  and  $\alpha \in (1/3, 1]$ .
- (Minimization:)  $\text{cont}(v, T) \leq t'/k' - (1 - 3\alpha)(k - 1)$  and  $\alpha \in (0, 1/3)$ .

► **Reduction Rule 4.4** (\*). Let  $\mathcal{I}$  be an instance of ANNOTATED DEGRADING  $\alpha$ -FCGP with  $\alpha > 0$  and let  $v \in V(G) \setminus T$  be a satisfactory vertex. Apply the Inclusion Rule (Reduction Rule 3.2) on vertex  $v$ .

We henceforth assume that Reduction Rule 4.4 is exhaustively applied on every satisfactory vertex. Next, we identify vertices which are not contained in any solution.

► **Definition 4.5.** Let  $\mathcal{I}$  be a Yes-instance of ANNOTATED DEGRADING  $\alpha$ -FCGP. A vertex  $v \in V(G) \setminus T$  is called *needless* if

- (Maximization:)  $\text{cont}(v, T) \leq t'/k' - (3\alpha - 1)(k - 1)^2$  for maximization and  $\alpha \in (1/3, 1]$ .
- (Minimization:)  $\text{cont}(v, T) \geq t'/k' + (1 - 3\alpha)(k - 1)^2$  for minimization and  $\alpha \in (0, 1/3)$ .

► **Reduction Rule 4.6** (\*). Let  $\mathcal{I}$  be an instance of ANNOTATED DEGRADING  $\alpha$ -FCGP with  $\alpha > 0$  and let  $v \in V(G) \setminus T$  be a needless vertex. Apply the Exclusion Rule (Reduction Rule 3.3) on vertex  $v$ .

We henceforth assume that Reduction Rule 4.6 is applied on every needless vertex. The following reduction rule decreases the counter of each vertex in  $V(G) \setminus T$ . After this rule is exhaustively applied, we may assume that  $\text{counter}(v) = 0$  for at least one vertex  $v \in V(G) \setminus T$ . Recall that we already have  $\text{counter}(v) = 0$  for every vertex in  $T$ .

► **Reduction Rule 4.7.** If  $\text{counter}(v) > 0$  for every vertex  $v \in V(G) \setminus T$ , then decrease  $\text{counter}(v)$  by 1 for every vertex  $v \in V(G) \setminus T$  and decrease  $t$  by  $\alpha k'$ .

Next, we show that after the exhaustive application of Reduction Rule 4.7 the counter of each vertex is bounded polynomially in terms of  $\Delta$  and  $k$ .

► **Lemma 4.8.** Let  $\mathcal{I}$  be a Yes-instance of ANNOTATED DEGRADING  $\alpha$ -FCGP with  $\alpha > 0$ . We have  $\text{counter}(v) \in \mathcal{O}(\Delta + k^2)$  for every vertex  $v \in V(G) \setminus T$ .

**Proof.** First and foremost, observe that there exists at least one vertex  $u \in V(G) \setminus T$  with  $\text{counter}(u) = 0$ , since otherwise Reduction Rule 4.7 is still applicable. Since Reduction Rule 3.3 is applied to each needless vertex, we conclude that every vertex in  $V(G) \setminus T$  has contribution at least  $t'/k' - (3\alpha - 1)(k - 1)^2$  for maximization. Furthermore, since Reduction Rule 3.2 is applied to each satisfactory vertex, we conclude that every vertex  $v \in V(G) \setminus T$  has contribution at least  $t'/k' - (1 - 3\alpha)(k - 1)$  for minimization. In particular, we have

$$\begin{aligned} \text{cont}(u, T) &\geq t'/k' - (3\alpha - 1)(k - 1)^2 \text{ for maximization, and} \\ \text{cont}(u, T) &\geq t'/k' - (1 - 3\alpha)(k - 1) \text{ for minimization.} \end{aligned}$$

Since also  $\text{cont}(u, T) = \alpha \cdot \deg(u) + (1 - 3\alpha)|N(u) \cap T|$  we obtain that

$$t'/k' \leq \alpha \cdot \deg(u) + (1 - 3\alpha)|N(u) \cap T| + (3\alpha - 1)(k - 1)^2 \text{ for maximization, and} \quad (1)$$

$$t'/k' \geq \alpha \cdot \deg(u) + (1 - 3\alpha)[(k - 1) + |N(u) \cap T|] \text{ for minimization.} \quad (2)$$

Moreover, since Reduction Rule 3.2 is applied to each satisfactory vertex, we conclude that every vertex  $v \in V(G) \setminus T$  has contribution at most  $t'/k' + (3\alpha - 1)(k - 1)$  for maximization. Furthermore, since Reduction Rule 3.3 is applied to each needless vertex, we conclude that every vertex in  $V(G) \setminus T$  has contribution at most  $t'/k' + (1 - 3\alpha)(k - 1)^2$  for minimization. This implies that in particular

$$\alpha \cdot \text{counter}(v) \leq t'/k' + (3\alpha - 1)(k - 1) \text{ for maximization, and.} \quad (3)$$

$$\alpha \cdot \text{counter}(v) \leq t'/k' + (1 - 3\alpha)(k - 1)^2 \text{ for minimization.} \quad (4)$$

For maximization and  $\alpha \in (1/3, 1]$  it then follows from Equations (1) and (3) that

$$\text{counter}(v) \leq \deg(u) + \frac{3\alpha - 1}{\alpha}[k(k - 1) - |N(v) \cap T|] \in \mathcal{O}(\Delta + k^2).$$

For minimization and  $\alpha \in (0, 1/3)$  it then follows from Equations (2) and (4) that

$$\text{counter}(v) \leq \deg(u) + \frac{1 - 3\alpha}{\alpha}[k(k - 1) + |N(v) \cap T|] \in \mathcal{O}(\Delta + k^2) \text{ for minimization.}$$

This concludes the proof.  $\blacktriangleleft$

**Putting everything together.** We use the following proposition in our kernels for DEGRADING  $\alpha$ -FCGP. Therefore, we first transform the instance into an equivalent instance of ANNOTATED DEGRADING  $\alpha$ -FCGP. Second, we apply our reduction rules. Third, we reduce back to the unannotated version.

- **Proposition 4.9** ( $\star$ ). *Let  $\alpha > 0$ . Given an instance  $(G, T, \text{counter}, k, t)$  of*
  - ANNOTATED DEGRADING MAX  $\alpha$ -FCGP, *we can compute in polynomial time an equivalent DEGRADING MAX  $\alpha$ -FCGP instance of size  $\mathcal{O}(|V(G)|^2 + |V(G)|k^2) \subseteq \mathcal{O}(|V(G)|^3)$ , and*
  - ANNOTATED DEGRADING MIN  $\alpha$ -FCGP, *we can compute in polynomial time an equivalent DEGRADING MIN  $\alpha$ -FCGP instance of size  $\mathcal{O}(|V(G)| \cdot (|V(G)| + k^2)^3) \subseteq \mathcal{O}(|V(G)|^7)$ .*
- **Proposition 4.10.** *Given an instance  $(G, T, \text{counter}, k, t)$  of ANNOTATED DEGRADING  $\alpha$ -FCGP with  $\alpha > 0$ , we can compute in polynomial time an equivalent DEGRADING  $\alpha$ -FCGP instance of size  $\mathcal{O}((\Delta + k)^{\mathcal{O}(1)})$ .*

**Proof.** Follows from Lemmas 3.7–3.8, and Proposition 4.2.  $\blacktriangleleft$

## 4.2 No Polynomial Kernels in Non-Degrading Cases

Note that if  $\alpha = 0$ , then MAX  $\alpha$ -FCGP corresponds to DENSEST  $k$ -SUBGRAPH. It is already known that DENSEST  $k$ -SUBGRAPH does not admit a polynomial kernel for  $\Delta + k$  [29]. We strengthen and generalize this result: First, we observe that DENSEST  $k$ -SUBGRAPH does not admit a polynomial kernel for  $k$ , even when  $\Delta = 3$ . Second, we extend this negative result to NON-DEGRADING MAX  $\alpha$ -FCGP and NON-DEGRADING MIN  $\alpha$ -FCGP when  $\Delta$  is a constant.

- **Theorem 4.11** ( $\star$ ). *Unless  $\text{coNP} \subseteq \text{NP/poly}$ ,*

1. NON-DEGRADING MAX  $\alpha$ -FCGP on subcubic graphs does not admit a polynomial kernel for  $k$ , and
2. NON-DEGRADING MIN  $\alpha$ -FCGP on graphs with constant maximum degree does not admit a polynomial kernel for  $k$ .

## 5 Parameterization by $c$ -closure

**Hardness for the non-degrading case.** We start with showing that the Non-Degrading case is intractable.

► **Theorem 5.1** ( $\star$ ). NON-DEGRADING MAX  $\alpha$ -FCGP remains  $W[1]$ -hard with respect to the solution size  $k$  even on 2-closed and 2-degenerate graphs.

► **Theorem 5.2** ( $\star$ ). NON-DEGRADING MIN  $\alpha$ -FCGP remains  $W[1]$ -hard with respect to the solution size  $k$  even on 2-closed graphs.

**A  $k^{\mathcal{O}(c)}$ -size kernel for the degrading case.** In contrast to the non-degrading case, we develop a kernel of size  $k^{\mathcal{O}(c)}$  for the degrading case. To this end, we apply a series of reduction rules to obtain an upper bound of  $k^{\mathcal{O}(c)}$  on the maximum degree. Then, the kernel of size  $k^{\mathcal{O}(c)}$  follows from Proposition 4.10. In order to upper-bound the maximum degree, we rely on a polynomial Ramsey bound for  $c$ -closed graphs [27].

► **Lemma 5.3** ([27, Lemma 3.1]). Any  $c$ -closed graph  $G$  on at least  $R_c(a, b) := (c-1) \cdot \binom{b-1}{2} + (a-1)(b-1) + 1$  vertices contains a clique of size  $a$  or an independent set of size  $b$ . Moreover, a clique of size  $a$  or an independent set of size  $b$  can be found in polynomial time.

Using a similar approach as Reduction Rule 4.1 (but exploiting the  $c$ -closure instead of the maximum degree) yields the following.

► **Reduction Rule 5.4.** Let  $\mathcal{I}$  be an instance of ANNOTATED DEGRADING  $\alpha$ -FCGP. Let  $v \in V(G)$  be some vertex and let  $X_v \subseteq N(v)$  be the set of vertices better than  $v$ . If  $|X_v| > (c-1)k$ , then apply the Exclusion Rule (Reduction Rule 3.3) to  $v$ .

► **Lemma 5.5.** Reduction Rule 5.4 is correct.

**Proof.** We provide a proof for the maximization version; the minimization version follows analogously. Let  $S$  be a solution. Assume that  $v \in S$  (we are done otherwise). We show that there is a vertex  $v' \neq v$  such that  $S' := (S \setminus \{v\}) \cup \{v'\}$  constitutes a solution. By Lemma 3.5, it suffices to show that  $\text{cont}(v', S \setminus \{v\}) \geq \text{cont}(v, S \setminus \{v\})$ . Let  $S'_v := S \setminus N[v]$ . Each vertex in  $S'_v$  is, by definition, nonadjacent to  $v$ , and hence it shares at most  $c-1$  neighbors in common with  $v$ . This implies  $|X_v \setminus N(S'_v)| \geq |X_v| - (c-1) \cdot |S'_v| > 0$  as  $X_v \subseteq N(v)$ . Thus, there exists a vertex  $v' \in X_v \setminus N(S'_v)$ , that is,  $N(v') \cap S'_v = \emptyset$ . Then, we have  $N(v') \cap (S \setminus \{v\}) \subseteq S \cap N(v)$  and thus  $|N(v') \cap (S \setminus \{v\})| \leq |N(v) \cap (S \setminus \{v\})|$ . Moreover, we have  $\alpha \deg^{+c}(v') \geq \alpha \deg^{+c}(v)$  (recall that  $v'$  is better than  $v$ ). Since  $\alpha \in (1/3, 1]$ , it follows that

$$\begin{aligned} \text{cont}(v', S \setminus \{v\}) &= \alpha \deg^{+c}(v') + (1 - 3\alpha) |N(v') \cap (S \setminus \{v\})| \\ &\geq \alpha \deg^{+c}(v) + (1 - 3\alpha) |N(v) \cap (S \setminus \{v\})| = \text{cont}(v, S \setminus \{v\}). \end{aligned}$$

This concludes the proof. ◀

Note that if there is a clique of size  $(c-1)k+1$ , then Reduction Rule 5.4 applies to one of the vertices with the smallest contribution. Thus, applying Reduction Rule 5.4 exhaustively removes all cliques of size  $(c-1)k+1$ .

► **Lemma 5.6.** *Let  $v \in V(G)$  be a vertex such that  $\deg(v) \geq R_c((c-1)k+1, (k+1)k^{c-2})$ . Then, we can find in polynomial time a set  $X$  of  $i \in [c-1]$  vertices and an independent set  $I$  with the following properties:*

- (i) *The set  $X$  contains  $v$ .*
- (ii) *The set  $I \subseteq \bigcap_{x \in X} N(x)$  is an independent set of size at least  $(k+1)k^{c-i}$ .*
- (iii) *For every vertex  $u \in V(G) \setminus X$ , it holds that  $|N(u) \cap I| \leq (k+1)k^{c-i-1}$ .*

**Proof.** Since there is no clique of size  $(c-1)k+1$ , there is an independent set  $I_v$  of size  $(k+1)k^{c-2}$  in  $N(v)$  by Lemma 5.3 (which can be found in polynomial time). Let  $X$  be an inclusion-wise maximal set of  $i$  vertices including  $v$  such that  $|\bigcap_{x \in X} N(x) \cap I_v| > (k+1)k^{c-i}$ . One of such sets can be found by the following polynomial-time algorithm: We start with  $X := \{v\}$  and  $i := 1$ . We will maintain the invariant that  $|X| = i$ . If there exists a vertex  $v' \in V(G) \setminus X$  with  $|N(v') \cap \bigcap_{x \in X} N(x) \cap I_v| > (k+1)k^{c-i-1}$ , then we add  $v'$  to  $X$  and increase  $i$  by 1. We keep doing so until there remains no such vertex  $v'$ .

We show that this algorithm terminates for  $i = |X| \leq c-1$ . Assume to the contrary that the algorithm continues for  $i = c-1$ . We then have that  $|N(v') \cap \bigcap_{x \in X} N(x) \cap I_v| > (k+1)k^{c-i-1} = k+1 \geq 2$  for some vertex  $v' \in V(G) \setminus X$ . Since  $I_v$  is an independent set, the set  $N(v') \cap \bigcap_{x \in X} N(x) \cap I_v$  contains two nonadjacent vertices. Note, however, that these two vertices have at least  $|X \cup \{v'\}| = c$  common neighbors, contradicting the  $c$ -closure of  $G$ .

Finally, we show that a set  $X$  found by this algorithm and  $I := \bigcap_{x \in X} N(x) \cap I_v$  satisfy the three properties of the lemma. We have  $|\bigcap_{x \in X} N(x) \cap I_v| = |N(v') \cap \bigcap_{x \in X \setminus \{v\}} N(x) \cap I_v| > (k+1)k^{c-(i-1)-1} = (k+1)k^{c-i}$ , where  $v'$  is the last vertex added to  $X$ . Moreover, since  $X$  is inclusion-wise maximal, we have  $|N(u) \cap I| = |N(u) \cap \bigcap_{x \in X} N(x) \cap I_v| \leq (k+1)k^{c-i-1}$  for every vertex  $u \in V(G) \setminus X$ . ◀

► **Reduction Rule 5.7.** Let  $\mathcal{I}$  be an instance of ANNOTATED DEGRADING  $\alpha$ -FCGP. Let  $X, I$  be as specified in Lemma 5.6 and let  $v \in I$  be a vertex such that every other vertex in  $I$  is better than  $v$ . If  $k \geq 2$ , then apply the Exclusion Rule (Reduction Rule 3.3) to  $v$ .

► **Lemma 5.8.** *Reduction Rule 5.7 is correct.*

**Proof.** Again, we show the proof for the maximization variant; the minimization variant follows analogously. For the sake of contradiction, assume that every solution  $S$  contains  $v$ . By Lemma 5.6, every vertex  $u \in V(G) \setminus X$  has at most  $(k+1)k^{c-i}$  neighbors in  $I$ . Moreover, since  $I$  is an independent set, we have  $|I \cap N[v']| = 1$  for every vertex  $v' \in I$  (including  $v$ ). For  $S' := S \setminus X$ , we have

$$\begin{aligned} |I \setminus N[S']| &\geq |I| - |I \cap N[v]| - |I \cap N[S' \setminus \{v\}]| \\ &\geq (k+1)k^{c-i} - (k-1)(k+1)k^{c-i-1} - 1 = k^{c-i} + k^{c-i-1} - 1 > 0. \end{aligned}$$

Let  $v'$  be an arbitrary vertex in  $I \setminus N[S']$ . We show that  $\text{cont}(v', S \setminus \{v\}) \geq \text{cont}(v, S \setminus \{v\})$ . By Lemma 3.5, this would imply that  $(S \setminus \{v\}) \cup \{v'\}$  is a solution not containing  $v$ . Since  $v$  and  $v'$  are both adjacent to all vertices of  $X$  and  $\alpha \in (1/3, 1]$ , we have  $|N(v) \cap (S \setminus \{v\})| > |X \cap (S \setminus \{v\})|$ . We thus have

$$\begin{aligned} \text{cont}(v', S \setminus \{v\}) &= \alpha \deg^{+c}(v') + (1-3\alpha)|X \cap (S \setminus \{v\})| \\ &\geq \alpha \deg^{+c}(v) + (1-3\alpha)|X \cap (S \setminus \{v\})| \\ &\geq \alpha \deg^{+c}(v) + (1-3\alpha)|N(v) \cap (S \setminus \{v\})| = \text{cont}(v, S \setminus \{v\}). \end{aligned}$$

Here, the first inequality follows from the fact that  $v'$  is better than  $v$ . ◀

By applying these reduction rules, we can ensure that  $\Delta \leq R_c((c-1)k+1, (k+1)k^{c-2}) \in k^{\mathcal{O}(c)}$ . Proposition 4.10 leads to the following:

► **Proposition 5.9.** *DEGRADING  $\alpha$ -FCGP has a kernel of size  $k^{\mathcal{O}(c)}$ .*

**Matching lower bounds.** Next, we show that the kernels provided in Theorem 5.9 cannot be improved under standard assumptions.

► **Proposition 5.10** ( $\star$ ). *DEGRADING MAX  $\alpha$ -FCGP has no kernel of size  $\mathcal{O}(k^{c-3-\epsilon})$  unless  $\text{coNP} \subseteq \text{NP/poly}$ .*

► **Proposition 5.11** ( $\star$ ). *For each  $\alpha \in (0, 1/3)$ , MIN  $\alpha$ -FCGP does not admit a kernel of size  $\mathcal{O}(k^{c-3-\epsilon})$  unless  $\text{coNP} \subseteq \text{NP/poly}$ .*

Note that MIN  $\alpha$ -FCGP for  $\alpha = 0$  is equivalent to SPAREST  $k$ -SUBGRAPH which admits a kernel of size  $\mathcal{O}(c^2k^3)$  [26].

Now, Propositions 5.9–5.11 imply the following.

► **Theorem 5.12.** *DEGRADING  $\alpha$ -FCGP admits a kernel of size  $k^{\mathcal{O}(c)}$ . For  $\alpha > 0$ , DEGRADING  $\alpha$ -FCGP does not admit a kernel of size  $k^{\mathcal{O}(c)}$  unless  $\text{coNP} \subseteq \text{NP/poly}$ .*

## 6 Parameterization by Degeneracy

**Minimization variant.** We start with the minimization variant which turns out to be easier than the maximization variant. This is most likely because of the following bound on  $t$ .

► **Lemma 6.1** ( $\star$ ). *In non-trivial instances  $(G, k, t)$  of MIN  $\alpha$ -FCGP we have  $t \leq dk$ .*

Shachnai and Zehavi [35] showed that MIN  $\alpha$ -FCGP with  $\alpha \in (0, 1]$  admits an FPT-algorithm with respect to  $k + t$ . Hence, we obtain the following.

► **Corollary 6.2.** *MIN  $\alpha$ -FCGP for  $\alpha > 0$  is FPT parameterized by  $d + k$ .*

Naturally, we may now ask whether this FPT result can be strengthened to a polynomial kernel. As shown by Theorem 4.11, the non-degrading case of MIN  $\alpha$ -FCGP does not admit a polynomial kernel even on graphs with constant maximum degree which implies constant degeneracy. In contrast, the degrading has a kernel whose size is polynomial in  $d + k$ .

► **Theorem 6.3** ( $\star$ ). *DEGRADING MIN  $\alpha$ -FCGP admits a kernel of size  $(d + k)^{\mathcal{O}(1)}$ .*

► **Proposition 6.4** ( $\star$ ). *SPARSEST  $k$ -SUBGRAPH admits a kernel with  $\mathcal{O}(dk)$  vertices and of size  $\mathcal{O}(d^2k)$ .*

**Maximization variant.** Recall, that MAXPVC is the special case of MAX  $\alpha$ -FCGP with  $\alpha = 1/2$ . Amini et al. [1] showed that MAXPVC can be solved in  $\mathcal{O}^*((dk)^k)$  time. Adapting this algorithm leads to an FPT-algorithm for  $\alpha$ -FCGP with respect to  $d + k$  for  $\alpha \neq 0$ :

► **Proposition 6.5** ( $\star$ ). *DEGRADING  $\alpha$ -FCGP can be solved in  $\mathcal{O}^*((dk)^k)$  time for  $\alpha \neq 0$ .*

The rest of this section is devoted to the proof of the next theorem.

► **Theorem 6.6.** *DEGRADING MAX  $\alpha$ -FCGP admits a kernel of size  $k^{\mathcal{O}(d)}$  but, unless  $\text{coNP} \subseteq \text{NP/poly}$ , no kernel of size  $\mathcal{O}(k^{d-2-\epsilon})$ .*

In particular, this implies that MAXPVC admits a kernel of size  $k^{\mathcal{O}(d)}$ . We remark that a compression of size  $(dk)^{\mathcal{O}(d)}$  was obtained independently by Panolan and Yaghoubzade [31].

**A kernel for biclique-free graphs in the degrading case.** We next develop a kernel of size  $k^{\mathcal{O}(d)}$ . In fact, our algorithm works for *biclique-free graphs*—graphs that do not have a biclique  $K_{a,b}$  as a subgraph for  $a \leq b \in \mathbb{N}$ . Note that a  $d$ -degenerate graph has no  $K_{d+1,d+1}$  as a subgraph, since otherwise every vertex in  $K_{d+1,d+1}$  has at least degree  $d+1$ .

Note that a clique of size  $a+b$  contains  $K_{a,b}$  as a subgraph. So given a graph  $G$  with no occurrence of  $K_{a,b}$  on at least  $\binom{a+b+k-2}{k-1} \in k^{\mathcal{O}(a+b)}$  vertices, one can find an independent set of size  $k$  in polynomial time (see Section 2). We show that this upper bound on the number of vertices can be improved: the sum  $a+b$  in the exponent can be replaced by  $\min\{a,b\}$ .

► **Lemma 6.7.** *For  $a \leq b \in \mathbb{N}$ , let  $G$  be a graph that contains no  $K_{a,b}$  as a subgraph. If  $G$  has at least  $R(k)$  vertices, then we can find in polynomial time an independent set of size  $k$ , where  $R(k) \in (a+b)^{\mathcal{O}(a)} \cdot k^a$*

**Proof.** We first show that if  $G$  has at least  $k + b \binom{k}{a} + \sum_{\ell \in [a-1]} R(a+b, \ell+1) \binom{k}{\ell}$  vertices, then it contains an independent set of size  $k$ . We give an algorithm to find an independent set of size  $k$  in polynomial time later. Let  $I$  be a maximum independent set in  $G$ . We assume for contradiction that  $|I| < k$ . We prove that there are at most  $t \binom{k}{a}$  vertices that have at least  $a$  neighbors in  $I$  and that there are at most  $\sum_{\ell \in [a-1]} R(a+b, \ell+1)$  vertices that have at most  $a-1$  neighbors in  $I$ .

For each subset  $X \subseteq I$  of size exactly  $a$ , note that there are at most  $b$  vertices  $v$  such that  $N(v) \supseteq X$ , since otherwise there is a  $K_{a,b}$  in  $G$ . It follows that the number of vertices with at least  $a$  neighbors in  $I$  is at most  $t \binom{|I|}{a} \leq b \binom{k}{a}$ . Consider a set  $X \subseteq I$  of size  $\ell \in [a-1]$ . Let  $V_X := \{v \in V(G) \setminus I \mid N(v) \cap I = X\}$ . Then, there is no independent set  $I'$  of size  $\ell+1$  in  $V_X$ , since otherwise  $(I \setminus X) \cup I'$  is an independent set of size at least  $|I|+1$ , contradicting the fact that  $I$  is an independent set of maximum size. Moreover, there is no clique of size  $a+b$  in  $V_X$ . Thus,  $|V_X| < R(a+b, \ell+1)$ . The number of vertices with at most  $a-1$  neighbors in  $I$  is then at most  $\sum_{X \subseteq I, |X|=\ell \in [a-1]} R(a+b, \ell+1) \binom{|I|}{\ell} \leq \sum_{\ell \in [a-1]} R(a+b, \ell+1) \binom{k}{\ell}$ .

We turn the argument above into a polynomial-time algorithm as follows. Suppose that we have an independent set  $I'$  of size smaller than  $k$ . As discussed above, there are at most  $b \cdot \binom{k}{a}$  vertices that have at least  $s$  neighbors in  $I'$ . Hence, there is a vertex set  $X \subseteq I'$  of size  $\ell$  such that  $|V_X| > R(a+b, \ell+1)$ . Note that  $X$  can be found in polynomial time, for instance, by counting the number of vertices  $v'$  such that  $N(v') \cap I' = N(v) \cap I'$  for each vertex  $v \in V(G)$ . We can then find an independent set  $I''$  of size  $\ell+1$  in  $X$  (this can be done in polynomial time as discussed in Section 2). This way, we end up with an independent set  $(I' \setminus X) \cup I''$  of size at least  $|I'|+1$ . Note that this procedure of finding an independent set of greater size is repeated at most  $k$  times, and thus the overall running time is polynomial. ◀

We remark that for fixed  $a \leq b \in \mathbb{N}$ , Lemma 6.7 gives us an  $\mathcal{O}(n^{1-1/a})$ -approximation algorithm for INDEPENDENT SET that runs in  $n^\ell$  time with a constant  $\ell$  not depending on  $a$  or  $b$ . An  $\mathcal{O}(n^{1-1/a})$ -approximation algorithm is known on graphs where  $K_{a,b}$  is excluded as an *induced subgraph* [5, 15]. However, these algorithms have running time  $n^{\Omega(a)}$ .

We then apply Lemma 6.7 to obtain a lemma analogous to Lemma 5.6.

► **Lemma 6.8.** *Let  $v \in V(G)$  be a vertex such that  $\deg(v) \geq R(bk^{a-1})$ . Then, we can find in polynomial time a set  $X$  of  $i \in [a-1]$  vertices and an independent set  $I$  with the following properties:*

- (i) *The set  $X$  contains  $v$ .*
- (ii) *The set  $I \subseteq \bigcap_{x \in X} N(x)$  is an independent set of size at least  $bk^{a-i} + 1$ .*
- (iii) *For every vertex  $u \in V(G) \setminus X$ , it holds that  $|N(u) \cap I| \leq bk^{a-i-1}$ .*



**Proof.** By Lemma 6.7, there is an independent set  $I_v$  of size  $bk^{b-1}$  in  $N(v)$  (which can be found in polynomial time). Let  $X$  be an inclusion-wise maximal set of  $i$  vertices including  $v$  with  $|\bigcap_{x \in X} N(x) \cap I_v| > bk^{a-i}$ . Such a set can be found by the following polynomial-time algorithm: We start with  $X = \{v\}$  and  $i = 1$ . We will maintain the invariant that  $|X| = i$ . If there exists a vertex  $v' \in V(G) \setminus X$  with  $|N(v') \cap \bigcap_{x \in X} N(x) \cap I_v| > bk^{a-i-1}$ , then we add  $v'$  to  $X$  and increase  $i$  by 1. We keep doing so until there remains no such vertex  $v'$ .

We show that this algorithm terminates for  $i = |X| \leq a - 1$ . Assume to the contrary that the algorithm continues for  $i = a - 1$ . We then have that  $|N(v') \cap \bigcap_{x \in X} N(x) \cap I_v| > bk^{a-i-1}$  for some vertex  $v' \in V(G) \setminus X$ . It follows that the set  $X \cup \{v'\}$  (which is of size  $a$ ) has more than  $b$  common neighbors, contradicting the fact that  $G$  has no  $K_{a,b}$  as a subgraph.

Finally, we show that a set  $X$  found by this algorithm and  $I := \bigcap_{x \in X} N(x) \cap I_v$  satisfy the three properties of the lemma. We have  $|I| = |\bigcap_{x \in X} N(x) \cap I_v| = |N(v') \cap \bigcap_{x \in X \setminus \{v'\}} N(x) \cap I_v| > bk^{a-(i-1)-1} = bk^{a-i}$ , where  $v'$  is the last vertex added to  $X$ . Moreover, since  $X$  is inclusion-wise maximal, we have  $|N(u) \cap I| = |N(u) \cap \bigcap_{x \in X} N(x) \cap I_v| \leq bk^{a-i-1}$  for every vertex  $u \in V(G) \setminus X$ .  $\blacktriangleleft$

► **Reduction Rule 6.9.** Let  $\mathcal{I}$  be an instance of ANNOTATED DEGRADING  $\alpha$ -FCGP. Let  $X, I$  be as specified in Lemma 6.8 and let  $v \in I$  be a vertex such that every other vertex in  $I$  is better than  $v$ . Then, apply the Exclusion Rule (Reduction Rule 3.3) to  $v$ .

► **Lemma 6.10.** *Reduction Rule 6.9 is correct.*

**Proof.** We show the proof for the maximization variant; the minimization variant follows analogously. For the sake of contradiction, assume that every solution  $S$  contains  $v$ . By Lemma 6.8, every vertex  $u \in V(G) \setminus X$  has at most  $bk^{a-i}$  neighbors in  $I$ . Moreover, since  $I$  is an independent set, we have  $|I \cap N[v']| = 1$  for every vertex  $v' \in I$  (including  $v$ ). For  $S' := S \setminus X$ , we have

$$\begin{aligned} |I \setminus N[S']| &\geq |I| - |I \cap N[v]| - |I \cap N[S' \setminus \{v\}]| \\ &\geq (bk^{a-i} + 1) - (k-1)bk^{a-i-1} - 1 = bk^{a-i-1} > 0. \end{aligned}$$

Let  $v'$  be an arbitrary vertex in  $I \setminus N[S']$ . We show that  $\text{cont}(v', S \setminus \{v\}) \geq \text{cont}(v, S \setminus \{v\})$ . By Lemma 3.5, this would imply that  $(S \setminus \{v\}) \cup \{v'\}$  is a solution not containing  $v$ . Since  $v$  and  $v'$  are both adjacent to all vertices of  $X$  and  $\alpha \in (1/3, 1]$ , we have

$$\begin{aligned} \text{cont}(v', S \setminus \{v\}) &= \alpha \deg^{+c}(v') + (1 - 3\alpha)|X \cap (S \setminus \{v\})| \\ &\geq \alpha \deg^{+c}(v) + (1 - 3\alpha)|X \cap (S \setminus \{v\})| \\ &\geq \alpha \deg^{+c}(v) + (1 - 3\alpha)|N(v) \cap (S \setminus \{v\})| = \text{cont}(v, S \setminus \{v\}). \end{aligned}$$

Here, the first inequality follows from the fact that  $v'$  is better than  $v$ .  $\blacktriangleleft$

By applying Reduction Rule 6.9 exhaustively, we end up with an instance with maximum degree  $\Delta \leq R(bk^{a-1})$ . The following proposition then follows from Proposition 4.10 using the bound in Lemma 6.7:

► **Proposition 6.11.** *For any  $a \leq b \in \mathbb{N}$ , DEGRADING  $\alpha$ -FCGP on graphs that do not contain  $K_{a,b}$  as a subgraph has a kernel of size  $(R(bk^{a-1}) + k)^{\mathcal{O}(1)} \in b^{\mathcal{O}(a)} k^{\mathcal{O}(a^2)}$ .*

Note that a  $d$ -degenerate graph contains no  $K_{d+1, d+1}$  as a subgraph. We obtain the following result using the folklore fact that any  $d$ -degenerate graph on at least  $(d+1)k$  vertices has an independent set of size  $k$ .



► **Lemma 6.12** (★). *DEGRADING  $\alpha$ -FCGP admits a kernel of size  $k^{\mathcal{O}(d)}$ .*

Now, we show that significant improvement in Lemma 6.12 is unlikely. This, together with Lemma 6.12, implies Theorem 6.6.

► **Proposition 6.13** (★). *DEGRADING MAX  $\alpha$ -FCGP admits no kernel of size  $\mathcal{O}(k^{d-2-\epsilon})$  unless  $\text{coNP} \subseteq \text{NP/poly}$ .*

## 7 Parameterization by $\text{vc}$ and $h$ -index

To complete the picture of the parameterized complexity landscape, we consider two parameters that are larger than the degeneracy of  $G$ : the  $h$ -index of  $G$  and the vertex cover number of  $G$ . We start with the maximization variant and the  $h$ -index. As NON-DEGRADING MAX  $\alpha$ -FCGP does not admit a polynomial kernel with respect to  $k$  even if  $\Delta$  is constant (see Thm. 4.11), the same holds for the  $h$ -index. The degrading case admits a polynomial kernel.

► **Proposition 7.1** (★). *DEGRADING MAX  $\alpha$ -FCGP admits a kernel of size  $\mathcal{O}(h^2k^2 + k^4)$ .*

We complement this with showing fixed-parameter tractability for  $k + h$ .

► **Proposition 7.2** (★). *NON-DEGRADING MAX  $\alpha$ -FCGP is fixed-parameter tractable with respect to  $k + h$ .*

For the larger parameter vertex cover number  $\text{vc}$ , we achieve a kernel for all  $\alpha > 0$ .

► **Proposition 7.3** (★). *If  $\alpha \neq 0$ , then MAX  $\alpha$ -FCGP admits a kernel of size  $\mathcal{O}(\text{vc}^4 + \text{vc} \cdot k^3)$ .*

For  $\alpha = 0$ , MAX  $\alpha$ -FCGP corresponds to DENSEST  $k$ -SUBGRAPH and CLIQUE is one of its special cases ( $t = \binom{k}{2}$ ). Since CLIQUE does not admit a polynomial kernel with respect to  $\text{vc}$  [3] (and any clique is of size at most  $\text{vc} + 1$ ), DENSEST  $k$ -SUBGRAPH does not admit a polynomial kernel. However, DENSEST  $k$ -SUBGRAPH can be solved by a straightforward algorithm in  $\mathcal{O}^*(2^{\text{vc}})$  time. Thus, DENSEST  $k$ -SUBGRAPH admits a kernel of size  $\mathcal{O}(2^{\text{vc}})$ .

**Minimization variant.** Note that DEGRADING MIN  $\alpha$ -FCGP has a polynomial kernel with respect to  $d + k$  (see Theorem 6.3) and, thus, also with respect to  $h + k$  and  $\text{vc} + k$ . As NON-DEGRADING MIN  $\alpha$ -FCGP does not admit a polynomial kernel with respect to  $k$  even if  $\Delta$  is constant (see Thm. 4.11), the same holds for the  $h$ -index. It remains to consider NON-DEGRADING MIN  $\alpha$ -FCGP parameterized by  $\text{vc} + k$ .

► **Proposition 7.4** (★). *MIN  $\alpha$ -FCGP admits a kernel of size  $\mathcal{O}(\text{vc}^8 + \text{vc} \cdot k^7)$  for  $\alpha > 0$  and of size  $\mathcal{O}(\text{vc}^2 + \text{vc} \cdot k)$  for  $\alpha = 0$ .*

## 8 Conclusion

We provided a systematic parameterized complexity analysis for  $\alpha$ -FCGP (see Figure 2). Although we settled the existence of polynomial kernels with respect to various parameters combined with the solution size  $k$ , several open questions remain. First, our polynomial kernels are not optimized and thus the polynomials are of high degree. Looking for smaller kernels is thus an obvious first task. Second, can our positive results for  $c$ -closure and degeneracy be extended to the smaller parameter weak closure [18]? Furthermore, while we looked at parameters that are small in sparse graphs, can similar results be achieved for dense graphs as considered e. g. by Lochet et al. [30]? Finally, we believe that an experimental verification of our data reduction rules would demonstrate their practical usefulness.

---

References

---

- 1 Omid Amini, Fedor V. Fomin, and Saket Saurabh. Implicit Branching and Parameterized Partial Cover Problems. *Journal of Computer and System Sciences*, 77(6):1159–1171, 2011.
- 2 Sanjeev Arora and Boaz Barak. *Computational Complexity - A Modern Approach*. Cambridge University Press, 2009.
- 3 Hans L. Bodlaender, Bart M. P. Jansen, and Stefan Kratsch. Kernelization lower bounds by cross-composition. *SIAM Journal on Discrete Mathematics*, 28(1):277–305, 2014.
- 4 Edouard Bonnet, Bruno Escoffier, Vangelis Th. Paschos, and Emeric Tourniaire. Multi-parameter Analysis for Local Graph Partitioning Problems: Using Greediness for Parameterization. *Algorithmica*, 71(3):566–580, 2015.
- 5 Édouard Bonnet, Stéphan Thomassé, Xuan Thang Tran, and Rémi Watrigant. An Algorithmic Weakening of the Erdős-Hajnal Conjecture. In *Proceedings of the 28th Annual European Symposium on Algorithms (ESA '20)*, volume 173 of *LIPICs*, pages 23:1–23:18. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020.
- 6 Nicolas Bourgeois, Aristotelis Giannakos, Giorgio Lucarelli, Ioannis Milis, and Vangelis Th. Paschos. Exact and superpolynomial approximation algorithms for the DENSEST  $k$ -SUBGRAPH PROBLEM. *European Journal of Operational Research*, 262(3):894–903, 2017.
- 7 Maurizio Bruglieri, Matthias Ehrgott, Horst W. Hamacher, and Francesco Maffioli. An annotated bibliography of combinatorial optimization problems with fixed cardinality constraints. *Discrete Applied Mathematics*, 154(9):1344–1357, 2006.
- 8 Leizhen Cai. Parameterized Complexity of Cardinality Constrained Optimization Problems. *The Computer Journal*, 51(1):102–121, 2008.
- 9 Leizhen Cai, Siu Man Chan, and Siu On Chan. Random Separation: A New Method for Solving Fixed-Cardinality Optimization Problems. In *Proceedings of the Second International Workshop on Parameterized and Exact Computation (IWPEC '06)*, volume 4169 of *LNCS*, pages 239–250. Springer, 2006.
- 10 Marek Cygan, Fedor V. Fomin, Lukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michal Pilipczuk, and Saket Saurabh. *Parameterized Algorithms*. Springer, 2015.
- 11 Marek Cygan, Fabrizio Grandoni, and Danny Hermelin. Tight Kernel Bounds for Problems on Graphs with Small Degeneracy. *ACM Transactions on Algorithms*, 13(3):43:1–43:22, 2017.
- 12 Marek Cygan, Marcin Pilipczuk, Michal Pilipczuk, and Jakub Onufry Wojtaszczyk. Kernelization hardness of connectivity problems in  $d$ -degenerate graphs. *Discrete Applied Mathematics*, 160(15):2131–2141, 2012.
- 13 Holger Dell and Dániel Marx. Kernelization of packing problems. In *Proceedings of the 23rd Annual ACM-SIAM Symposium on Discrete Algorithms (SODA '12)*, pages 68–81. SIAM, 2012.
- 14 Rodney G. Downey and Michael R. Fellows. *Fundamentals of Parameterized Complexity*. Texts in Computer Science. Springer, 2013.
- 15 Pavel Dvořák, Andreas Emil Feldmann, Ashutosh Rai, and Pawel Rzazewski. Parameterized Inapproximability of Independent Set in  $H$ -Free Graphs. In *Proceedings of the 46th International Workshop on Graph-Theoretic Concepts in Computer Science, (WG '20)*, volume 12301 of *Lecture Notes in Computer Science*, pages 40–53. Springer, 2020.
- 16 David Eppstein and Emma S. Spiro. The  $h$ -Index of a Graph and its Application to Dynamic Subgraph Statistics. *Journal of Graph Algorithms and Applications*, 16(2):543–567, 2012.
- 17 Uriel Feige and Michael Seltser. On the densest  $k$ -subgraph problem. Technical report, Weizmann Institute of Science. Department of Applied Mathematics and Computer Science, 1997.
- 18 Jacob Fox, Tim Roughgarden, C. Seshadhri, Fan Wei, and Nicole Wein. Finding Cliques in Social Networks: A New Distribution-Free Model. *SIAM Journal on Computing*, 49(2):448–464, 2020.
- 19 Jiong Guo, Rolf Niedermeier, and Sebastian Wernicke. Parameterized Complexity of Vertex Cover Variants. *Theory of Computing Systems*, 41(3):501–520, 2007.

- 20 Satoshi Hara, Takayuki Katsuki, Hiroki Yanagisawa, Takafumi Ono, Ryo Okamoto, and Shigeki Takeuchi. Consistent and Efficient Nonparametric Different-Feature Selection. In *Proceedings of the 20th International Conference on Artificial Intelligence and Statistics (AISTATS '17)*, volume 54 of *Proceedings of Machine Learning Research*, pages 130–138. PMLR, 2017.
- 21 Satoshi Hara, Tetsuro Morimura, Toshihiro Takahashi, Hiroki Yanagisawa, and Taiji Suzuki. A Consistent Method for Graph Based Anomaly Localization. In *Proceedings of the Eighteenth International Conference on Artificial Intelligence and Statistics (AISTATS '15)*, volume 38 of *Proceedings of Machine Learning Research*, pages 333–341. PMLR, 2015.
- 22 Danny Hermelin and Xi Wu. Weak compositions and their applications to polynomial lower bounds for kernelization. In *Proceedings of the 23rd Annual ACM-SIAM Symposium on Discrete Algorithms (SODA '12)*, pages 104–113. SIAM, 2012.
- 23 Shlomo Hoory, Nathan Linial, and Avi Wigderson. Expander graphs and their applications. *Bulletin of the American Mathematical Society*, 43(04):439–562, 2006.
- 24 Stasys Jukna. *Extremal Combinatorics - With Applications in Computer Science*. Texts in Theoretical Computer Science. An EATCS Series. Springer, 2011.
- 25 Joachim Kneis, Alexander Langer, and Peter Rossmanith. Improved Upper Bounds for Partial Vertex Cover. In *Proceedings of the 34th International Workshop on Graph-Theoretic Concepts in Computer Science (WG '08)*, volume 5344 of *LNCS*, pages 240–251, 2008.
- 26 Tomohiro Koana, Christian Komusiewicz, and Frank Sommer. Computing Dense and Sparse Subgraphs of Weakly Closed Graphs. In *Proceedings of the 31st International Symposium on Algorithms and Computation, (ISAAC '20)*, volume 181 of *LIPICs*, pages 20:1–20:17. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020.
- 27 Tomohiro Koana, Christian Komusiewicz, and Frank Sommer. Exploiting  $c$ -Closure in Kernelization Algorithms for Graph Problems. In *Proceedings of the 28th Annual European Symposium on Algorithms (ESA '20)*, volume 173 of *LIPICs*, pages 65:1–65:17. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020.
- 28 Tomohiro Koana, Christian Komusiewicz, and Frank Sommer. Essentially Tight Kernels For (Weakly) Closed Graphs. In *Proceedings of the 32nd International Symposium on Algorithms and Computation, (ISAAC '21)*, volume 212 of *LIPICs*, pages 35:1–35:15. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021.
- 29 Christian Komusiewicz and Manuel Sorge. An Algorithmic Framework for Fixed-Cardinality Optimization in Sparse Graphs Applied to Dense Subgraph Problems. *Discrete Applied Mathematics*, 193:145–161, 2015.
- 30 William Lochet, Daniel Lokshtanov, Saket Saurabh, and Meirav Zehavi. Exploiting Dense Structures in Parameterized Complexity. In *Proceedings of the 38th International Symposium on Theoretical Aspects of Computer Science (STACS '21)*, volume 187 of *LIPICs*, pages 50:1–50:17. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021.
- 31 Fahad Panolan and Hannane Yaghoubizade. Partial Vertex Cover on Graphs of Bounded Degeneracy. *CoRR*, abs/2201.03876, 2022.
- 32 Geevarghese Philip, Venkatesh Raman, and Somnath Sikdar. Polynomial Kernels for Dominating Set in Graphs of Bounded Degeneracy and Beyond. *ACM Transactions on Algorithms*, 9(1):11:1–11:23, 2012.
- 33 Venkatesh Raman and Saket Saurabh. Short Cycles Make W-hard Problems Hard: FPT Algorithms for W-hard Problems in Graphs with no Short Cycles. *Algorithmica*, 52(2):203–225, 2008.
- 34 Saket Saurabh and Meirav Zehavi.  $(k, n - k)$ -MAX-CUT: An  $\mathcal{O}^*(2^p)$ -time algorithm and a polynomial kernel. *Algorithmica*, 80(12):3844–3860, 2018.
- 35 Hadas Shachnai and Meirav Zehavi. Parameterized Algorithms for Graph Partitioning Problems. *Theory of Computing Systems*, 61(3):721–738, 2017.
- 36 Rémi Watrigant, Marin Bougeret, and Rodolphe Giroudeau. Approximating the Sparsest  $k$ -Subgraph in Chordal Graphs. *Theory of Computing Systems*, 58(1):111–132, 2016.

## A Appendix – Preliminaries for proofs in the appendix

We call  $v$  *isolated* if  $\deg_G(v) = 0$  and *non-isolated* otherwise. We also say that  $v$  is a *leaf vertex* if  $\deg_G(v) = 1$  and a *non-leaf vertex* if  $\deg_G(v) \geq 2$ .

**Weak compositions.** Our kernel lower bounds are based on weak  $q$ -compositions. For more information on parameterized complexity, we refer to the standard monographs [10, 14].

► **Definition A.1** ([13, 22]). *Let  $q \geq 1$  be an integer, let  $L_1 \subseteq \{0, 1\}^*$  be a classic (non-parameterized) problem, and let  $L_2 \subseteq \{0, 1\}^* \times \mathbb{N}$  be a parameterized problem. A weak  $q$ -composition from  $L_1$  to  $L_2$  is a polynomial time algorithm that on input  $x_1, \dots, x_{t^q} \in \{0, 1\}^n$  outputs an instance  $(y, k') \in \{0, 1\}^* \times \mathbb{N}$  such that:*

- $(y, k') \in L_2 \Leftrightarrow x_i \in L_1$  for some  $i \in [t^q]$ , and
- $k' \leq t \cdot n^{\mathcal{O}(1)}$ .

► **Lemma A.2** ([10, 13, 22]). *Let  $q \geq 1$  be an integer, let  $L_1 \subseteq \{0, 1\}^*$  be a classic NP-hard problem, and let  $L_2 \subseteq \{0, 1\}^* \times \mathbb{N}$  be a parameterized problem. The existence of a weak  $q$ -composition from  $L_1$  to  $L_2$  implies that  $L_2$  has no compression of size  $\mathcal{O}(k^{q-\epsilon})$  for any  $\epsilon > 0$ , unless  $\text{coNP} \subseteq \text{NP/poly}$ .*

## B Appendix – Proofs

### B.1 Proof of Lemma 3.1

**Proof.** Let  $S_i = \{v_1, \dots, v_{i-1}\}$  and  $\overline{S}_i = \{v_i, \dots, v_{|S|}\}$  for each  $i \in [|S|]$ . Observe that

$$\begin{aligned} m(S, V(G) \setminus S) &= \sum_{i \in [|S|]} |N(v_i) \setminus S| = \sum_{i \in [|S|]} |N(v_i) \setminus S_{i-1}| - |N(v_i) \cap \overline{S}_i| \text{ and} \\ m(S) &= \sum_{i \in [|S|]} |N(v_i) \cap S_{i-1}| = \sum_{i \in [|S|]} |N(v_i) \cap \overline{S}_i|. \end{aligned}$$

We thus have

$$\begin{aligned} \text{val}(S) &= \alpha(m(S, V(G) \setminus S) + m(S) + \text{counter}(S)) + (1 - 2\alpha) \cdot m(S) \\ &= \sum_{i \in [|S|]} \left( \alpha[ (|N(v_i) \setminus S_{i-1}| - |N(v_i) \cap \overline{S}_i|) + |N(v_i) \cap \overline{S}_i| + \text{counter}(v)] \right. \\ &\quad \left. + (1 - 2\alpha)|N(v_i) \cap S_{i-1}| \right) \\ &= \sum_{i \in [|S|]} \text{cont}(v_i, \overline{S}_i), \end{aligned}$$

which proves the lemma. ◀

### B.2 Proof of Lemma 3.6

**Proof.** We give a proof for the maximization variant; the minimization variant follows analogously. By the definition of strictly better vertices, it suffices to show that  $\text{cont}(v, T) -$

$\text{cont}(u, T) \geq 0$  for each  $T \subseteq V(G)$  of size at most  $k$ :

$$\begin{aligned} & \text{cont}(v, T) - \text{cont}(u, T) \\ &= \alpha \deg^{+c}(v) + (1 - 3\alpha)|N(v) \cap T| - \alpha \deg^{+c}(u) - (1 - 3\alpha)|N(u) \cap T| \\ &= \alpha(\deg^{+c}(v) - \deg^{+c}(u)) + (1 - 3\alpha)(|N(v) \cap T| - |N(u) \cap T|) \\ &\geq |(1 - 3\alpha)k| + (1 - 3\alpha)(|N(v) \cap T| - |N(u) \cap T|) \geq |(1 - 3\alpha)k| + (1 - 3\alpha)k \geq 0. \end{aligned}$$

This completes the proof.  $\blacktriangleleft$

### B.3 Proof of Lemma 3.8

**Proof.** We may assume that  $G$  has at least  $k$  vertices (otherwise the lemma holds for a trivial No-instance  $\mathcal{I}$ ). We construct an equivalent instance  $\mathcal{I}' := (G', k, t')$  of MIN  $\alpha$ -FCGP.

Let  $\ell$  be the smallest integer greater than  $(1/\alpha - 1)\Delta + \Gamma$ . Let  $G'$  be the graph obtained from  $G$  as follows: For every vertex  $v \in V(G) \setminus T$ , introduce  $\ell + \text{counter}(v)$  cliques on  $2\ell + 1$  vertices. Let  $\mathcal{C}_v$  be the set of these cliques. Next, for each clique in  $\mathcal{C}_v$ , add an edge between  $v$  and one of its vertices. Furthermore, let  $C_v$  be the set of vertices in these cliques and let  $C := \bigcup_{v \in V(G) \setminus T} C_v$ . To conclude the construction of  $\mathcal{I}'$ , we set  $t' := t + \alpha\ell(k - |T|)$ . Observe that for each vertex we add  $\mathcal{O}(\ell)$  cliques and each of these cliques consists of  $\mathcal{O}(\ell)$  vertices. Hence, we add at most  $\mathcal{O}(\ell^3) = \mathcal{O}((\Delta + \Gamma)^3)$  edges for every vertex in  $V(G) \setminus T$  and thus the claimed size follows.

Next, we show that  $\mathcal{I}$  and  $\mathcal{I}'$  are equivalent. For a solution  $S$  of  $\mathcal{I}$  its value in  $G'$  is increased by  $\alpha\ell$  for every vertex  $v \in V(G) \setminus T$ , amounting to  $t + \alpha\ell(k - |T|)$ .

Conversely, suppose that  $\mathcal{I}'$  has a solution  $S'$ . We show that there is a solution that contains all vertices of  $T$  but no vertex of  $C$ . For an arbitrary chosen vertex  $v' \in S'$ , the following holds:

1.  $\text{cont}(v, S' \setminus \{v'\}) \leq (1 - \alpha)\Delta + \Gamma$  for any vertex  $v \in T$ .
2.  $\alpha\ell \leq \text{cont}(v, S' \setminus \{v'\}) \leq (1 - \alpha)\Delta + \alpha(\Gamma + \ell)$  for any vertex  $v \in V(G) \setminus T$ .
3.  $\text{cont}(v, S' \setminus \{v'\}) \geq 2\alpha\ell$  for any vertex  $v \in C$ .

From  $\ell \geq (1/\alpha - 1)\Delta + \Gamma$ , we conclude that  $(1 - \alpha)\Delta + \Gamma \leq \alpha\ell$  and that  $(1 - \alpha)\Delta + \alpha(\Gamma + \ell) \leq 2\alpha\ell$ . The first inequality implies that each vertex in  $T$  has a lower contribution than each vertex in  $V(G) \setminus T$  and the second inequality implies that each vertex in  $V(G) \setminus T$  has a lower contribution than each vertex in  $C$ . Thus, by Lemma 3.5, we obtain that  $\mathcal{I}'$  admits a solution  $S'$  with  $T \subseteq S' \subseteq V(G')$ . Hence,  $S'$  is also a solution for  $\mathcal{I}$ .  $\blacktriangleleft$

### B.4 Proof of Reduction Rule 4.1

► **Lemma B.1.** *Reduction Rule 4.1 is correct.*

**Proof.** Let  $\mathcal{I}'$  be the reduced instance. Clearly, if  $\mathcal{I}'$  has a solution  $S'$ , then  $S'$  is also a solution for  $\mathcal{I}$ . Conversely, suppose that  $\mathcal{I}$  has a solution  $S$ . If  $v \notin S$ , then  $S$  is also a solution for  $\mathcal{I}'$ . In the following, we assume that  $v \in S$ .

By the pigeonhole principle, there exists a vertex  $v'$  better than  $v$  such that  $v \notin N[S \setminus T]$ . We claim that  $S' := (S \setminus \{v\}) \cup \{v'\}$  is a solution for  $\mathcal{I}'$ . First, we consider maximization. By Lemma 3.5, it suffices to show that  $\text{cont}(v', S \setminus \{v\}) \geq \text{cont}(v, S \setminus \{v\})$ . Since  $v' \notin N[S \setminus T]$ , we have  $\text{cont}(v', S \setminus \{v\}) = \text{cont}(v', T) \geq \text{cont}(v, T) \geq \text{cont}(v, S \setminus \{v\})$ . Here, the last inequality follows from the fact that the contribution is degrading. Moreover, for minimization, the last two  $\geq$  would be  $\leq$ .  $\blacktriangleleft$

## B.5 Proof of Reduction Rule 4.4

► **Lemma B.2.** *Reduction Rule 4.4 is correct.*

**Proof.** Let  $\mathcal{I}'$  be the reduced instance. Clearly, if  $\mathcal{I}'$  has a solution  $S'$ , then  $S'$  is also a solution for  $\mathcal{I}$ . Conversely, suppose that  $\mathcal{I}$  has a solution  $S$ . The lemma clearly holds for  $v \in S$ . So we will assume that  $v \notin S$ . We start with an auxiliary claim:

▷ **Claim B.3.** There is an ordering  $(v_1, \dots, v_{k'})$  of the vertices of  $S \setminus T$  with

- $\ell_1 \geq \ell_2 \geq \dots \geq \ell_{k'}$  for maximization and  $\alpha \in (1/3, 1]$ , and
- $\ell_1 \leq \ell_2 \leq \dots \leq \ell_{k'}$  for minimization and  $\alpha \in (0, 1/3)$ ,

where  $S_i := T \cup \{v_1, \dots, v_{i-1}\}$  and  $\ell_i := \text{cont}(v_i, S_i)$  for every  $i \in [k']$ .

**Proof.** Consider an ordering of  $S \setminus T$ , where the  $i$ -th vertex  $v_i$  is chosen in such a way that  $\text{cont}(v_i, S_i)$  is maximized (minimized). Then, since the contribution is degrading, we have  $\text{cont}(v_i, S_i) \geq \text{cont}(v_{i+1}, S_i) \geq \text{cont}(v_{i+1}, S_{i+1})$  for maximization, and  $\text{cont}(v_i, S_i) \leq \text{cont}(v_{i+1}, S_i) \leq \text{cont}(v_{i+1}, S_{i+1})$  for minimization, respectively, for every  $i \in [k' - 1]$ . ◀

Note that  $S_{k'} = S \setminus \{v_{k'}\}$ . Now, consider the vertex set  $S' := S_{k'} \cup \{v\}$ . We show that  $\text{val}(S') \geq t$  for maximization and that  $\text{val}(S') \leq t$  for minimization. By Lemma 3.1, we have  $\text{val}(S') = \text{val}(S) - \ell_{k'} + \text{cont}(v, S_{k'})$ . The definition of contribution yields that

$$\begin{aligned} \text{cont}(v, S_{k'}) &= \alpha \deg^{+c}(v) + (1 - 3\alpha)|N(v) \cap S_{k'}| \\ &= \text{cont}(v, T) + (1 - 3\alpha)|N(v) \cap (S_{k'} \setminus T)|. \end{aligned}$$

Since  $|S_{k'} \setminus T| = k' - 1$  and  $\text{cont}(v, T) \geq t'/k' + (3\alpha - 1)(k - 1)$  for maximization and  $\text{cont}(v, T) \leq t'/k' - (1 - 3\alpha)(k - 1)$  for minimization, we have

$$\begin{aligned} \text{cont}(v, S_{k'}) &\geq [t'/k' + (3\alpha - 1)(k - 1)] + (1 - 3\alpha)(k' - 1) \geq t'/k' \\ &\quad \text{for maximization and } \alpha \in (1/3, 1], \text{ and} \\ \text{cont}(v, S_{k'}) &\leq [t'/k' - (1 - 3\alpha)(k - 1)] + (1 - 3\alpha)(k' - 1) \leq t'/k' \\ &\quad \text{for minimization and } \alpha \in (0, 1/3). \end{aligned}$$

For maximization, if  $\ell_{k'} \leq t'/k'$ , then we have  $\ell_{k'} \leq \text{cont}(v, S_{k'})$  and thus  $\text{val}(S') \geq \text{val}(S) \geq t$ . Analogously, for minimization, if  $\ell_{k'} \geq t'/k'$ , then we have  $\ell_{k'} \geq \text{cont}(v, S_{k'})$  and thus  $\text{val}(S') \leq \text{val}(S) \leq t$ . Thus, in the following, we assume that  $\ell_{k'} > t'/k'$  for maximization and that  $\ell_{k'} < t'/k'$  for minimization. We obtain that

$$\begin{aligned} \text{val}(S_{k'}) &= \text{val}(T) + \sum_{i=1}^{k'-1} \ell_i. \text{ Thus,} \\ \text{val}(S_{k'}) &\geq \text{val}(T) + (k' - 1)\ell_{k'} \geq \text{val}(T) + \frac{(k' - 1)t'}{k'} \text{ for maximization, and} \\ \text{val}(S_{k'}) &\leq \text{val}(T) + (k' - 1)\ell_{k'} \leq \text{val}(T) + \frac{(k' - 1)t'}{k'} \text{ for minimization.} \end{aligned}$$

Hence, for maximization we have  $\text{val}(S') = \text{val}(S_{k'}) + \text{cont}(v, S_{k'}) \geq [\text{val}(T) + (k' - 1)t'/k'] + t'/k' = t$  and for minimization we have  $\text{val}(S') = \text{val}(S_{k'}) + \text{cont}(v, S_{k'}) \leq [\text{val}(T) + (k' - 1)t'/k'] + t'/k' = t$ . ◀

## B.6 Proof of Reduction Rule 4.6

► **Lemma B.4.** *Reduction Rule 4.6 is correct.*

**Proof.** Let  $\mathcal{I}'$  be the reduced instance. Clearly, if  $\mathcal{I}'$  has a solution  $S'$ , then  $S'$  is also a solution for  $\mathcal{I}$ . Conversely, suppose that  $\mathcal{I}$  has a solution  $S$ . The lemma clearly holds for  $v \notin S$ . So we will assume that  $v \in S$ . Since Reduction Rule 3.2 is exhaustively applied to each satisfactory vertex  $v'$ , we obtain that  $\text{cont}(v', T) < t'/k' + (3\alpha - 1)(k - 1)$  for maximization and  $\text{cont}(v', T) > t'/k' - (1 - 3\alpha)(k - 1)$  for minimization, for each vertex  $v' \in V(G) \setminus T$ . Together with Lemma 3.1 we thus obtain for maximization that

$$\begin{aligned} \text{val}(S) &\leq \text{val}(T) + \text{cont}(v, T) + (k' - 1)[t'/k' + (3\alpha - 1)(k - 1)] \\ &< \text{val}(T) + [t'/k' - (3\alpha - 1)(k - 1)^2] + (k' - 1)[t'/k' + (3\alpha - 1)(k - 1)] \\ &= \text{val}(T) + t' - (3\alpha - 1)(k - 1)(k - k') \leq t \text{ since } \alpha \in (1/3, 1]. \end{aligned}$$

Similar, for minimization, we obtain that

$$\begin{aligned} \text{val}(S) &\geq \text{val}(T) + \text{cont}(v, T) + (k' - 1)[t'/k' - (1 - 3\alpha)(k - 1)] \\ &> \text{val}(T) + [t'/k' + (1 - 3\alpha)(k - 1)^2] + (k' - 1)[t'/k' - (1 - 3\alpha)(k - 1)] \\ &= \text{val}(T) + t' + (1 - 3\alpha)(k - 1)(k - k') \geq t \text{ since } \alpha \in (0, 1/3). \end{aligned}$$

For maximization, this is a contradiction to  $\text{val}(S) \geq t$ , and for minimization this is a contradiction to  $\text{val}(S) \leq t$ . Thus,  $S$  cannot contain the needless vertex  $v$ . ◀

## B.7 Proof of Prop. 4.9

**Proof.** Using Reduction Rules 3.2, 3.3, and 4.7 exhaustively yields, by Lemma 4.8, an instance where  $\text{counter}(v) \in \mathcal{O}(|V(G)| + k^2)$ . For maximization, by Lemma 3.7 we get an equivalent instance of DEGRADING MAX  $\alpha$ -FCGP of size  $\mathcal{O}(|V(G)|^2 + |V(G)|k^2) \subseteq \mathcal{O}(|V(G)|^3)$ . For minimization, by Lemma 3.8 we get an equivalent instance of DEGRADING MIN  $\alpha$ -FCGP of size  $\mathcal{O}(|V(G)| \cdot (|V(G)| + k^2)^3) \subseteq \mathcal{O}(|V(G)|^7)$ . ◀

## B.8 Proof of Thm. 4.11

**Proof.** We present a polynomial parameter transformation from CLIQUE parameterized by the size of the largest connected component (this parameterization does not admit a polynomial kernel which can be seen by an or-composition of disjoint CLIQUE instances with the same clique size) to NON-DEGRADING MAX  $\alpha$ -FCGP (and NON-DEGRADING MIN  $\alpha$ -FCGP) parameterized by the solution size  $k$ . The reduction is based on a reduction of Feige and Seltser [17] that shows NP-hardness of DENSEST  $k$ -SUBGRAPH on subcubic graphs with a subsequent reduction to NON-DEGRADING MAX  $\alpha$ -FCGP and NON-DEGRADING MIN  $\alpha$ -FCGP.

The reduction from CLIQUE to DENSEST  $k$ -SUBGRAPH works as follows. Let  $(G, \ell)$  be an instance of CLIQUE. Without loss of generality, we may assume that every connected component of  $G$  has exactly  $\hat{n}$  vertices and more than  $\binom{\ell}{2}$  edges. For each connected component  $H$  of  $G$  perform the following construction. Let  $\{v_1, \dots, v_{\hat{n}}\}$  denote the vertex set of  $H$ . For each vertex  $v_i$  of  $H$  create a cycle  $C_i := (v_i^1, \dots, v_i^{\hat{n}})$  of length  $\hat{n}$ . This cycle is called the *vertex cycle of  $v_i$* . Then, add a path on  $\hat{n}^2 + 1$  edges between  $v_r^q$  and  $v_q^r$  if  $v_r$  and  $v_q$  are adjacent in  $H$ . In the following, this path is called *connector path of  $v_r^q$  and  $v_q^r$* . To each newly added vertex on this path, attach a degree-one vertex. Furthermore, attach a



degree-one vertex to each vertex on a vertex cycle which has no connector path. We call these newly added degree-one vertices *pendent*. Observe that every nonpendent vertex has degree three. Let  $G'$  denote the constructed graph.

▷ **Claim B.5.** The graph  $G$  contains a clique of size  $\ell$  if and only if  $G'$  contains a  $k' := \ell\hat{n} + \binom{\ell}{2}\hat{n}^2$ -vertex subgraph with at least  $\delta' := \ell\hat{n} + \binom{\ell}{2}(\hat{n}^2 + 1)$  edges.

*Proof.* Suppose that  $G$  contains a clique  $K$  of size  $\ell$ . Then, consider the induced subgraph that contains the  $\ell$  vertex cycles of the clique vertices and all connector paths between them. This subgraph has the claimed number of vertices and edges since the number of cycles is  $\ell$  and since each pair of cycles is connected via a path because  $K$  is a clique.

Conversely, assume that  $G'$  contains a  $k'$ -vertex subgraph  $G'[S]$  with at least  $\delta'$  edges. First, observe that we may assume that  $S$  contains no pendent vertices: By construction and the fact that each connected component of  $G'$  has more than  $\binom{\ell}{2}$  edges, each nonpendent vertex of  $G'$  is connected to more than  $k'$  nonpendent vertices. Hence, every pendent vertex of  $S$  can be replaced by a nonpendent neighbor  $u \notin S$  of some nonpendent vertex  $v \in S$  without decreasing the number of edges in  $G'[S]$ .

After this observation, the correctness proof can be carried out in the same manner as in the proof of Feige and Seltser [17]. We sketch the details for sake of completeness. First, observe that every degree-one vertex  $u$  of  $G'[S]$  may reach at most one degree-3 vertex  $v$  of  $G'[S]$  via some path that contains only vertices that have degree 2 in  $G'[S]$ . We say that the degree-one vertex  $u$  is *associated* with the degree-3 vertex  $v$ . We call a degree-3 vertex *good* if it is not associated with any degree-one vertex. A simple calculation shows that  $G'[S]$  must contain at least  $2\binom{\ell}{2}$  good degree-3 vertices to achieve the claimed number of edges (refer to Feige and Seltser [17] for further details). If a degree-3 vertex is good, then  $G'[S]$  contains its connector path completely. By the choice of  $k'$  and the path length,  $G'[S]$  may contain at most  $\binom{\ell}{2}$  connector paths completely and thus both endpoints of a connector path that is contained completely in  $G'[S]$  are good. Now, let  $\mathcal{P}$  denote the set of connector paths that are completely contained in  $G'[S]$  and let  $\mathcal{C}$  denote the collection of vertex cycles that contain a good vertex or, equivalently, that contain an endpoint of a connector path in  $\mathcal{P}$ .

We show that we may assume that every vertex cycle of  $\mathcal{C}$  is completely contained in  $G'[S]$ . Assume, towards a contradiction, that  $G'[S]$  does not contain all vertex cycles of  $\mathcal{C}$  completely and that  $G'[S]$  is a  $k'$ -vertex and  $\delta'$ -edge subgraph of  $G$  with the largest possible number of vertices of  $\mathcal{C}$ . Choose some vertex cycle  $C_v$  of  $\mathcal{C}$  that is not contained completely in  $G'[S]$ . By definition of  $\mathcal{C}$ , this vertex cycle contains a good vertex  $u$ . Since  $C_v$  is not contained completely in  $G'[S]$ , some vertex  $y$  of  $C_v$  has degree one in  $G'[S]$  and is connected to  $u$  via a path of vertices of  $C_v$ . Since  $u$  is good, there must be some vertex  $w$  on this path which has degree 3 in  $G'[S]$  but which is not good. Thus,  $G'[S]$  contains some but not all vertices of the connector path starting at  $w$ . Let  $x$  be the degree-one vertex on this connector path that is contained in  $G'[S]$  and reachable from  $w$  via the connector path vertices. Removing  $x$  from  $G'[S]$  and adding the missing cycle neighbor of  $y$  gives a graph  $\tilde{G}$  with the same number of edges and vertices but with more cycle vertices, a contradiction to the choice of  $G'[S]$ .

Consequently, we may assume that  $G'[S]$  contains all cycles of  $\mathcal{C}$  completely. By the choice of  $k'$  we have that  $|\mathcal{C}| = \ell$ . Thus,  $G'[S]$  contains  $\ell$  vertex cycles with  $\binom{\ell}{2}$  connector paths between them. The vertex set of  $G$  corresponding to the vertex cycles is thus a clique of size  $\ell$ . ◁

The proof of Claim B.5 also implies the correctness of the following observation

▷ **Claim B.6.** If  $G'$  contains an  $\ell\hat{n} + \binom{\ell}{2}\hat{n}^2$ -vertex subgraph with at least  $\ell\hat{n} + \binom{\ell}{2}(\hat{n}^2 + 1)$  edges, then it contains one such subgraph that has no pendant vertices. In other words, such a size  $\ell\hat{n} + \binom{\ell}{2}\hat{n}^2$ -vertex subgraph contains only vertices of degree 3.

**No polynomial kernel for NON-DEGRADING MAX  $\alpha$ -FCGP.** Observe that the size parameter  $k'$  of the DENSEST- $k$ -SUBGRAPH instance depends only on  $\hat{n}$  and that the maximum degree of  $G'$  is 3. Moreover, Claim B.5 shows the correctness of the reduction. We thus have shown the theorem statement for DENSEST- $k$ -SUBGRAPH, the special case of NON-DEGRADING MAX  $\alpha$ -FCGP with  $\alpha = 0$ . We now provide a polynomial-time transformation from the DENSEST- $k$ -SUBGRAPH instance  $(G', k', \delta')$  to an instance  $(G', k', t)$  of NON-DEGRADING MAX  $\alpha$ -FCGP with arbitrary  $\alpha \in [0, 1/3)$  by setting  $t := (1 - \alpha)\delta' + \alpha(3k' - 2\delta')$ . Since  $G'$  and  $k'$  are not changed by this transformation, it only remains to show its correctness.

Assume that  $(G', k', \delta')$  is a yes-instance, and let  $S$  be a size- $k'$  vertex set such that  $G'[S]$  has at least  $\delta'$  edges. By Claim B.6, we may assume that  $S$  contains only vertices that have degree 3 in  $G'$ . Thus, in  $G'$ , we have  $\text{val}(S) = (1 - \alpha)m(S) + \alpha m(S, V(G') \setminus S) = (1 - \alpha)m(S) + \alpha(3k' - 2m(S)) \geq (1 - \alpha)\delta' + \alpha(3k' - 2\delta')$  since  $m(S) \geq \delta'$  and  $(1 - \alpha) > 2\alpha$ .

Conversely, let  $S$  be a size- $k'$  set with  $\text{val}(S) \geq t = (1 - \alpha)\delta' + \alpha(3k' - 2\delta')$  in  $G'$ . First, observe that we may assume that  $S$  does not contain pendent vertices: Each such vertex has degree one in  $G'$  and thus degree at most one in  $G'[S]$ . Such a vertex exists since each connected component of  $G$  has at least  $\binom{\ell}{2}$  edges and hence each connected component of  $G'$  has at least  $k'$  vertices. Consequently, it can be replaced by some nonpendent neighbor of a vertex in  $S$ . Consequently,  $\text{val}(S) = (1 - \alpha)m(S) + \alpha(3k' - 2m(S))$ . As in the proof of the forward direction, it follows that  $m(S) \geq \delta'$ .

**No polynomial kernel for NON-DEGRADING MIN  $\alpha$ -FCGP.** Finally, we provide a polynomial-parameter transformation from the DENSEST- $k$ -SUBGRAPH instance  $(G', k', \delta')$  to an instance  $(G'', k', t)$  of NON-DEGRADING MIN  $\alpha$ -FCGP. Next, we distinguish whether  $\alpha \in (1/3, 1)$  or  $\alpha = 1$ . In both cases, we add a gadget to each pendent vertex of  $G'$  such that each size- $k'$  set  $S$  with minimal value  $\text{val}(S)$  in  $G''$  contains only vertices that have degree at most 3 in  $G''$ , which will be exactly the vertices of  $G'$ . Afterwards, the proof of the equivalence of  $(G, \ell)$  and  $(G'', k', t)$  is the same as for NON-DEGRADING MAX  $\alpha$ -FCGP. Hence, for both cases it remains to show that any solution with minimal value  $\text{val}(S)$  does only contain nonpendent vertices of  $G'$ .

First, we show the statement for  $\alpha \in (1/3, 1)$ . Construct  $G''$  from  $G'$  by adding for each degree-one vertex  $u$  in  $G'$  a set  $K_u$  on  $\lceil 15/(1 - \alpha) \rceil$  vertices and making  $K_u \cup \{u\}$  a clique. Set  $t := (1 - \alpha)\delta' + \alpha(3k' - 2\delta')$ . Assume towards a contradiction that some set  $S$  of minimal value contains a vertex  $u$  with  $\deg(u) > 3$  in  $G''$ . By construction  $u$  is part of some clique  $K_y$  where  $y$  is some pendent vertex of  $G'$ . Let  $Y := K_y \cap S$  and let  $r := |Y|$ .

Next, we show that for  $S' := (S \setminus Y) \cup Z$  for any vertex set  $Z \subseteq V(G') \setminus S$  consisting of non-pendent vertices we have  $\text{val}(S') < \text{val}(S)$ , contradicting the minimality of  $\text{val}(S)$ . Observe that such a vertex set  $Z$  exists since each connected component of  $G'$  has at least  $\binom{\ell}{2}$  edges and thus each connected component of  $G''$  has at least  $k''$  nonpendent vertices. Since  $|K_y| = \lceil 15/(1 - \alpha) \rceil$ , we obtain that  $\text{val}(S \setminus Y) \leq \text{val}(S) - (1 - \alpha)\binom{r}{2} - \alpha r \lceil 15/(1 - \alpha) \rceil$ . Furthermore, observe that adding the vertices in  $Z$  increases the objective value by at most  $3r$  since each nonpendent vertex has degree 3 in  $G''$ . Also, note that removing  $Y$  from  $S$  may result in a new out-going edge of  $S$  which is incident with the neighbor of  $y$  in  $V(G')$ . Hence, we obtain that

$$\text{val}((S \setminus Y) \cup Z) \leq \text{val}(S) - (1 - \alpha) \binom{r}{2} - \alpha r \lceil 15/(1 - \alpha) \rceil + 3r + \alpha.$$

Note that if  $r \leq \lceil 15/(1 - \alpha) \rceil - 4$ , then  $-\alpha r \lceil 15/(1 - \alpha) \rceil + 3r + \alpha < 0$  and thus  $\text{val}((S \setminus Y) \cup Z) < \text{val}(S)$ , a contradiction to the minimality of  $S$ . Otherwise, if  $r \geq \lceil 15/(1 - \alpha) \rceil - 3$ , then  $(1 - \alpha) \binom{r}{2} > (1 - \alpha) \lceil 11/(1 - \alpha) \rceil \lceil 10/(1 - \alpha) \rceil / 2 > 55/(1 - \alpha) > 3 \cdot \lceil 15/(1 - \alpha) \rceil + \alpha$  and thus  $\text{val}((S \setminus Y) \cup Z) < \text{val}(S)$ , a contradiction to the minimality of  $S$ . Hence,  $S$  contains only vertices of  $G'$ .

Second, we show the statement for  $\alpha = 1$ . Construct  $G''$  from  $G'$  by adding for each degree-one vertex  $u$  in  $G'$  a graph  $H$  to  $G''$  and for an arbitrary vertex  $h \in V(H)$  add the edge  $uh$  to  $G''$ . The graph  $H$  is an  $(|V(G')|, d, \rho)$ -edge expander, that is,  $H$  has  $|V(G')|$  vertices, is  $d$ -regular for some constant  $d$  and every  $T \subseteq V(H)$  of size at most  $|V(G')|/2$  fulfills  $m(T, V(H) \setminus T) \geq \rho d |T|$ . We choose the  $(|V(G')|, d, \rho)$ -edge expander  $H$  in such a way that  $\rho d \geq 4$ . Note that such a graph  $H$  exists and can be constructed in polynomial-time [2, 23]. Finally, we  $t := (1 - \alpha)\delta' + \alpha(3k' - 2\delta')$ . Assume towards a contradiction that some set  $S$  of minimal value contains a vertex  $u$  with  $\deg(u) > 3$  in  $G''$ . By construction  $u$  is part of some copy of  $H$  which was attached to a pendent (degree-one) vertex  $y$  of  $G'$ . Let  $Y := H \cap S$  and let  $r := |Y|$ .

Since  $H$  is an  $(|V(G')|, d, \rho)$ -edge expander and since  $k' < |V(G'')|/2$  we obtain that  $m(Y, V(G'') \setminus S) \geq 4r$ . Since each vertex in  $G'$  has degree at most 3, we thus obtain for any set  $Z \subseteq V(G')$  of size  $r$  that  $\text{val}((S \setminus Y) \cup Z) \leq \text{val}(S \setminus Y) - 4r + 3r < \text{val}(S)$ , a contradiction to the minimality of  $S$ . Hence,  $S$  contains only vertices of  $G'$ . ◀

## B.9 Proof of Thm. 5.1

**Proof.** We reduce from the W[1]-hard problem DENSEST- $k$ -SUBGRAPH in 2-closed and 2-degenerate graphs [33]. In DENSEST- $k$ -SUBGRAPH one asks for a vertex subset  $S \subseteq V(G)$  of exactly  $k$  vertices such that  $G[S]$  has at least  $t$  edges. Recall, that DENSEST  $k$ -SUBGRAPH is the special case of MAX  $\alpha$ -FCGP with  $\alpha = 0$ . Hence, it remains to show the theorem for  $\alpha \in (0, 1/3)$ .

Let  $(G, k, t)$  be an instance of DENSEST  $k$ -SUBGRAPH. By  $n(G) := |V(G)|$  we denote the number of vertices of  $G$ . We construct an equivalent instance  $(G', k', t')$  of MAX  $\alpha$ -FCGP as follows: The graph  $G'$  consists of a copy of  $G$ . By  $Z$  we denote the set of all vertices which are a copy of a vertex in  $G$ . Furthermore, we add exactly  $n(G) - \deg_G(v)$  many leaf-vertices to each vertex  $v \in Z$ . We denote these vertices by  $I_v$ . By  $I := \bigcup_{v \in V(G)} I_v$  we denote the set of all these leaf-vertices. Finally, we set  $k' := k$  and  $t' := \alpha(kn(G) - 2t) + (1 - \alpha)t$ .

By construction, each vertex in  $I$  has degree 1. Since  $G$  is 2-closed and 2-degenerate, we conclude that also  $G'$  is 2-closed and 2-degenerate.

Let  $S \subseteq V(G)$  be such that  $|S| = k$  and that  $m(G[S]) \geq t$ . Since each vertex  $v \in V(G')$  which is a copy of a vertex in  $G$  has degree exactly  $n(G)$ ,  $N(x) \cap N(y) \subseteq Z$  for each two vertices  $x, y \in Z$ , and  $m(G[S]) \geq t$  we conclude that exactly  $x \geq t$  edges have both endpoints in  $S$  and that exactly  $kn(G) - 2x \leq kn(G) - 2t$  edges have exactly one endpoint in  $S$ . Thus,  $\text{val}(S) \geq \alpha(kn(G) - 2t) + (1 - \alpha)t = t'$  since  $\alpha \in (0, 1/3)$ .

Conversely, suppose that  $S' \subseteq V(G')$  is set of exactly  $k$  vertices with  $\text{val}(S') \geq t'$ . Observe that by construction each vertex in  $I$  has degree 1. Hence,  $\text{cont}(v) \leq \max(\alpha, 1 - \alpha) < 1$  for each vertex  $v \in I$ . Furthermore, observe that for each vertex  $z \in Z$  we have  $\text{cont}(z) \geq \alpha n(G) > 1$  since  $\alpha$  is a constant. Thus,  $\text{cont}(z) > \text{cont}(v)$  for each vertex  $z \in Z$  and each  $v \in I$ . Hence, we can assume that  $S' \subseteq Z$ . Let  $x$  be the number of edges with both endpoints in  $S'$ . Since each vertex in  $S'$  has degree exactly  $n(G)$ , we conclude that exactly  $kn(G) - 2x$  edges have

exactly one endpoint in  $S'$ . Hence,  $\text{val}(S') = \alpha(kn(G) - 2x) + (1 - \alpha)x$ . Since  $\text{val}(S') \geq t'$  and  $\alpha \in (0, 1/3)$ , we conclude that  $x \geq t$ . Hence,  $G[S']$  contains at least  $t$  edges.  $\blacktriangleleft$

### B.10 Proof of Thm. 5.2

**Proof.** We reduce from the W[1]-hard problem DENSEST- $k$ -SUBGRAPH in 2-closed graphs [33]. In DENSEST- $k$ -SUBGRAPH one asks for a vertex subset  $S \subseteq V(G)$  of exactly  $k$  vertices such that  $G[S]$  has at least  $t$  edges.

Let  $(G, k, t)$  be an instance of DENSEST  $k$ -SUBGRAPH. We construct an equivalent instance  $(G', k', t')$  of MIN  $\alpha$ -FCGP as follows: The graph  $G'$  consists of a copy of  $G$ . By  $Z$  we denote the set of all vertices which are a copy of a vertex in  $G$ . Furthermore, for each vertex  $v \in V(G)$  we add a set  $W_v$  of exactly  $n(G) - \deg_G(v)$  many vertices to  $G'$  such that each vertex of  $W_v$  is connected with vertex  $v$ . By  $W := \bigcup_{v \in V(G)} W_v$  we denote the set of all these vertices. Next, for each vertex  $w \in W$  we add a clique  $C_w$  of size  $4t'$  to  $G'$  such that each vertex of  $C_w$  is connected with  $w$ . By  $C := \bigcup_{w \in W} C_w$  we denote the set of all these vertices. Finally, we set  $k' := k$  and  $t' := \alpha(kn(G) - 2t) + (1 - \alpha)t$ .

Observe that  $G'[W \cup C]$  is a cluster graph. Furthermore, observe that each vertex in  $W \cup C$  has at most one neighbor in  $Z$ . Since  $G[Z]$  is 2-closed, we conclude that  $G'$  is 2-closed.

Let  $S \subseteq V(G)$  be such that  $|S| = k$  and that  $m(G[S]) \geq t$ . Since each vertex  $v \in V(G')$  which is a copy of a vertex in  $G$  has degree exactly  $n(G)$ ,  $N(x) \cap N(y) \subseteq Z$  for each two vertices  $x, y \in Z$ , and  $m(G[S]) \geq t$  we conclude that exactly  $x \geq t$  edges have both endpoints in  $S$  and that exactly  $kn(G) - 2x \leq kn(G) - 2t$  edges have exactly one endpoint in  $S$ . Thus,  $\text{val}(S) \leq \alpha(kn(G) - 2t) + (1 - \alpha)t = t'$  since  $\alpha \in (1/3, 1]$ .

Conversely, suppose that  $S' \subseteq V(G')$  is set of exactly  $k$  vertices with  $\text{val}(S') \leq t'$ . Observe that by construction each vertex in  $W \cup C$  has degree at least  $2t'$ . If  $S'$  contains a vertex  $v$  of  $W \cup C$ , then  $v$  has at least  $3t'$  neighbors which are not in  $S'$  since  $t' > k$ . Thus,  $\text{val}(S') > \alpha 3t' > t'$ , a contradiction to the assumption  $\text{val}(S') \leq t'$ . Hence,  $S' \cap (W \cup C) = \emptyset$ . Let  $x$  be the number of edges with both endpoints in  $S'$ . Since each vertex in  $S'$  has degree exactly  $n(G)$ , we conclude that exactly  $kn(G) - 2x$  edges have exactly one endpoint in  $S'$ . Hence,  $\text{val}(S') = \alpha(kn(G) - 2x) + (1 - \alpha)x$ . Since  $\text{val}(S') \leq t$  and  $\alpha \in (1/3, 1]$ , we conclude that  $x \geq t$ . Hence,  $G[S']$  contains at least  $t$  edges.  $\blacktriangleleft$

### B.11 Proof of Prop. 5.10

We first prove the following technical lemma.

**► Lemma B.7.** *Let  $(G, k, t)$  be an instance of DEGRADING  $\alpha$ -FCGP with a solution  $S$  fulfilling  $T \subseteq S \subseteq T \cup P$  where  $P$  is a set of vertices of the same degree. Let  $P^* := S \cap P$ . Then:*

1. *For maximization,  $\text{val}(S)$  is maximal if  $m(P^*)$  and  $m(T, P^*)$  are minimal.*
2. *For minimization,  $\text{val}(S)$  is minimal if  $m(P^*)$  and  $m(T, P^*)$  are minimal.*

**Proof.** Let  $z := \deg(p)$  for each  $p \in P$ , let  $P^* := \{p_1, \dots, p_\ell\}$ , and let  $P_i := \{p_j \mid j < i\}$ . By Lemma 3.1 we obtain that

$$\begin{aligned} \text{val}(S) &= \text{val}(T) + \sum_{i=1}^{\ell} \text{cont}(p_i, T \cup P_i) \\ &= \text{val}(T) + \alpha z \ell + (1 - 3\alpha) \sum_{i=1}^{\ell} |N(p_i) \cap T| + |N(p_i) \cap P_i| \\ &= \text{val}(T) + \alpha z \ell + (1 - 3\alpha)(m(P^*, T) + m(P^*)) \end{aligned}$$

Note that  $\text{val}(T) + \alpha z \ell$  is a constant. Thus, for maximization in the degrading case we obtain that  $\text{val}(S)$  is maximized if  $m(P^*, T)$  and  $m(P^*)$  are minimized since  $(1 - 3\alpha) < 0$ . Furthermore, for minimization in the degrading case we obtain that  $\text{val}(S)$  is minimized if  $m(P^*, T)$  and  $m(P^*)$  are minimized since  $(1 - 3\alpha) > 0$ .  $\blacktriangleleft$

**Proof of Proposition 5.10.** We provide a weak  $q$ -composition from INDEPENDENT SET on 2-closed graphs to DEGRADING MAX  $\alpha$ -FCGP in  $(q + 3)$ -closed graphs. Here, we assume that  $k > 3q + \lceil \frac{\alpha}{3\alpha - 1} \rceil$ .

INDEPENDENT SET

**Input:** A graph  $G = (V, E)$  and an integer  $k$ .

**Question:** Is there an independent set of size exactly  $k$ ?

Let  $[t]^q$  be the set of  $q$ -dimensional vectors whose entries are in  $[t]$ . For a vector  $x \in [t]^q$  we denote by  $x_i$  the  $i$ -th entry of  $x$ . Assume that  $q \geq 2$  is a constant and that we are given exactly  $[t]^q$  instances  $\mathcal{I}_x := (G_x, k)$  of INDEPENDENT SET on 2-closed graphs for  $x \in [t]^q$ . Let  $V_x := V(G_x)$  for each  $x \in [t]^q$  and let  $D := \bigcup_{x \in [t]^q} V_x$ . We construct an equivalent instance  $(H, k', t')$  of DEGRADING MAX  $\alpha$ -FCGP as follows.

**Construction:** First for each  $x \in [t]^q$  we add the graph  $G_x$  to  $H$ . We then add a clique  $C$  consisting of  $tq$  vertices to  $H$ . The vertices of  $C$  are denoted by  $w_j^i$  with  $i \in [q]$  and  $j \in [t]$ . Furthermore, for each  $x \in [t]^q$  and  $v \in V_x$ , we add the edge  $vw_{x_i}^i$  for each  $i \in [q]$ . Fix an integer  $\ell \geq t^q \cdot n + (\alpha k + k + 1)/\alpha$ . We add leaf vertices so that  $\deg(w_j^i) = \ell + \lceil (3\alpha - 1)tq/\alpha \rceil$  for each vertex  $w_j^i$  and  $\deg(v) = \ell$  for each vertex  $v \in D$ . We denote the union of all these added leaf vertices by  $L$ . Finally, we set  $k' := k + (t - 1)q$  and

$$t' := (1 - \alpha) \frac{(t - 1)q((t - 1)q - 1)}{2} + \alpha \left[ k\ell + (t - 1)q \left( \ell + \lceil \frac{(3\alpha - 1)tq}{\alpha} \rceil \right) - (t - 1)q((t - 1)q - 1) \right].$$

**Closure:** We show that  $H$  is  $(q + 3)$ -closed. Since leaf vertices have degree one and  $C$  is a clique, we only have to consider nonadjacent vertex pairs where one vertex  $u$  is in  $D$  and the other vertex  $v$  is in  $C \cup D$ . Without loss of generality we assume that  $u \in V_x$  for some  $x \in [t]^q$ . Recall that  $N(u) \subseteq V_x \cup C \cup L$  and  $u$  has exactly  $q$  neighbors in  $C$ . First, consider the case  $v \in C$ . By construction we obtain from  $uv \notin E$  that  $u'v \notin E$  for each  $u' \in V_x$ . Thus,  $|N(u) \cap N(v)| = q$ . Second consider the case  $v \in D$ . If  $v \notin V_x$ , then  $|N(u) \cap N(v)| \leq q - 1$ . Otherwise, if  $v \in V_x$ , we obtain  $|N(u) \cap N(v)| \leq q + 2$  from the fact that  $\mathcal{I}_x$  is 2-closed.

**Correctness:** In the following, we prove that there exists an independent set of size exactly  $k$  for some instance  $\mathcal{I}_x$  with  $x \in [t]^q$  if and only if there exists a vertex set  $S$  of size exactly  $k'$  in  $H$  such that  $\text{val}(S) \geq t'$ .

Suppose that instance  $\mathcal{I}_x$  has an independent set  $I$  of size exactly  $k$  for some  $x \in [t]^q$ . By  $C^* := C \setminus \{w_{x_i}^i \mid i \in [q]\}$  we denote the non-neighbors of  $V_x$  in  $C$ . Note that  $|C^*| = (t - 1)q$ . We show that  $S := I \cup C^*$  is a solution of  $(H, k', t')$ . Clearly,  $|S| = k + (t - 1)q = k'$ . Since no vertex of  $I$  is connected with any vertex in  $C^*$ ,  $I$  is an independent set, and since  $C^*$  is a clique of size  $(t - 1)q$ , we conclude that  $m_H(S) = (t - 1)q((t - 1)q - 1)/2$ . Furthermore, since each vertex in  $I$  has degree exactly  $\ell$ , and since each vertex in  $C^*$  has degree exactly  $\ell + \lceil (3\alpha - 1)tq/\alpha \rceil$ , we conclude that  $m_H(S, V(H) \setminus S) = k\ell + (t - 1)q(\ell +$

$\lceil (3\alpha - 1)tq/\alpha \rceil - (t-1)q((t-1)q-1)$ . Thus,  $\text{val}(S) = t'$  and hence  $(H, k', t')$  is a Yes-instance of DEGRADING MAX  $\alpha$ -FCGP.

Conversely, suppose that  $(H, k', t')$  has a solution  $S \subseteq V(H)$  of size exactly  $k'$  with  $\text{val}(S) \geq t'$ . First, we show that we can assume that  $S \cap L = \emptyset$ . Assume that there exists a vertex  $v \in S \cap L$  and let  $w \in (C \cup D) \setminus S$ . We show that for  $S' := S \setminus \{v\} \cup \{w\}$  we have  $\text{val}(S') > \text{val}(S)$ . Observe that  $\text{val}(S \setminus \{v\}) \geq \text{val}(S) - 1$ . Furthermore, note that adding  $w$  may result into at most  $k$  new inner edges and hence the value decreases by at most  $k$ . Simultaneously, since  $\deg(w) \geq \ell$ , at least  $\ell - k$  new outer edges emerge such that the value increases by at least  $\alpha(\ell - k)$ . Since  $\ell > (\alpha k + k + 1)/\alpha$ , we obtain  $\text{val}(S') > \text{val}(S)$ .

Thus, in the following we can assume that  $S \cap L = \emptyset$ . Let  $C^* := C \cap S$ ,  $|C^*| = z$ , and  $D^* := S \setminus C^* \subseteq D$ . In the following, we show that  $z = (t-1)q$  and that there exists an  $x \in [t]^q$  such that  $N(V_x) \cap C^* = \emptyset$ . Therefore, we consider the cases that  $z < (t-1)q$  and that  $z > (t-1)q$ . In both cases we verify that  $\text{val}(S) < t$  for each solution with exactly  $z$  vertices in  $C$ .

**Case 1:**  $z \leq (t-1)q - 1$ . In other words,  $z = (t-1)q - p$  for some  $p \in [(t-1)q]$ . Recall that  $\deg(v) = \ell$  for each vertex  $v \in D$ . Thus, by Lemma B.7,  $\text{val}(S)$  is maximized if both  $m_H(D^*)$  and  $m_H(C^*, D^*)$  are minimized. Since  $z = |C^*| < (t-1)q$ , it is possible that no vertex of  $D^*$  is adjacent to any vertex in  $C^*$ . Thus,  $\text{val}(S)$  is maximized if  $D^*$  is an independent set and if  $E_H(C^*, D^*) = \emptyset$ . Hence,

$$\begin{aligned} \text{val}(S) = f(p) &\leq (1 - \alpha) \frac{((t-1)q - p)((t-1)q - p - 1)}{2} \\ &\quad + \alpha \left[ (k+p)\ell + ((t-1)q - p) \left( \ell + \left\lceil \frac{(3\alpha - 1)tq}{\alpha} \right\rceil \right) \right] \\ &\quad - \alpha((t-1)q - p)((t-1)q - p - 1). \end{aligned}$$

Now, we obtain that

$$f'(p) = (1 - 3\alpha)p + \frac{3\alpha - 1}{2}(2(t-1)q - 1) - \alpha \left\lceil \frac{(3\alpha - 1)tq}{\alpha} \right\rceil.$$

Since  $(1 - 3\alpha) < 0$  for  $\alpha \in (1/3, 1]$ , we obtain for  $p \leq (t-1)q$  that

$$\begin{aligned} f'(p) &\leq \frac{3\alpha - 1}{2}(2(t-1)q - 1) - \alpha \left\lceil \frac{(3\alpha - 1)tq}{\alpha} \right\rceil \\ &\leq \frac{3\alpha - 1}{2}(2(t-1)q - 1) - (3\alpha - 1)tq = (1 - 3\alpha)(q + 1/2) < 0. \end{aligned}$$

Since  $\alpha \in (1/3, 1]$  we conclude that  $f(p)$  is a concave quadratic function. And since  $f'(p) < 0$  for each  $p \leq (t-1)q$ , we thus conclude that  $f(0) > f(p)$  for each  $p \in [(t-1)q]$ , a contradiction to the assumption that  $f(p) = \text{val}(S) \geq t'$ .

**Case 2:**  $z \geq (t-1)q + 1$ . Let  $z = (t-1)q + p$  for some  $p \in [q]$ . By the pigeonhole principle there exist at least  $p$  indices  $i \in [q]$  such that  $w_j^i \in S$  for each  $j \in [t]$ . Recall that by construction, each vertex  $v \in D$  has exactly one neighbor in the set  $\{w_j^i, j \in [t]\}$ . Since  $|D^*| = k - p$  we conclude that  $m_H(C^*, D^*) \geq (k-p)p$ . Recall that  $\deg(v) = \ell$  for each vertex  $v \in D$ . Thus, by Lemma B.7,  $\text{val}(S)$  is maximized if  $D^*$  is an independent set and



if  $m_H(C^*, D^*) = (k - p)p$ . Thus,

$$\begin{aligned} \text{val}(S) = f(p) &\leq (1 - \alpha) \left[ (k - p)p + \frac{((t - 1)q + p)((t - 1)q + p - 1)}{2} \right] + \alpha(k - p)(\ell - p) \\ &\quad + \alpha p \left[ \ell + \left\lceil \frac{(3\alpha - 1)tq}{\alpha} \right\rceil - (t - 1)q - k + 1 \right] \\ &\quad + \alpha(t - 1)q \left[ \ell + \left\lceil \frac{(3\alpha - 1)tq}{\alpha} \right\rceil - (t - 1)q - p + 1 \right]. \end{aligned}$$

Now, we obtain that

$$f'(p) = (3\alpha - 1)p + \left(1 - 3\alpha\right) \left(k + (t - 1)q - \frac{1}{2}\right) + \alpha \left\lceil \frac{(3\alpha - 1)tq}{\alpha} \right\rceil.$$

Since  $\alpha \in (1/3, 1]$  and since  $p \in [q]$ , we obtain that

$$\begin{aligned} f'(p) &\leq (3\alpha - 1)q + (1 - 3\alpha)(k + (t - 1)q - 1/2) + \alpha \left( \frac{(3\alpha - 1)tq}{\alpha} + 1 \right) \\ &\leq (3\alpha - 1) \left( 2q + 1/2 + \frac{\alpha}{3\alpha - 1} - k \right). \end{aligned}$$

Since  $k > 3q + \lceil \frac{\alpha}{3\alpha - 1} \rceil$  and since  $\alpha \in (1/3, 1]$ , we obtain that  $f'(p) < 0$  for  $p \leq q$ . Furthermore, since  $\alpha \in (1/3, 1]$  we see that  $f(p)$  is a convex quadratic function. Thus, we conclude that  $f(0) > f(p)$  for each  $p \in [q]$ , a contradiction to the assumption that  $f(p) = \text{val}(S) \geq t'$ .

Hence,  $|C^*| = (t - 1)q$  and thus  $|D^*| = k$ . According to Lemma B.7,  $\text{val}(S)$  is maximal if both  $m_H(D^*)$  and  $m_H(C^*, D^*)$  are minimal. Observe that if  $D^*$  is an independent set and if  $E_H(C^*, D^*) = \emptyset$ , then  $\text{val}(S) = t'$ . Otherwise, if  $E_H(C^*, D^*) \neq \emptyset$  or if  $D^*$  is no independent set, then  $\text{val}(S) < t'$ . Thus,  $E_H(C^*, D^*) = \emptyset$  and  $D^*$  is an independent set. Note that, if there exists two vertices  $u, v \in D^*$  such that  $u \in V_x$  and  $v \in V_y$  with  $x \neq y$  for  $x, y \in [t]^q$ , then  $(N(u) \cup N(v)) \cap C \geq q + 1$ . Since  $|C^*| = (t - 1)q$  this implies that  $E_H(C^*, D^*) \neq \emptyset$ , a contradiction. Hence,  $D^* \subseteq V_x$  for some  $x \in [t]^q$  and by Lemma B.7 the set  $D^*$  is an independent set. Thus, the instance  $x$  contains an independent set of size at least  $k$ .

Hence, we have a weak- $q$ -composition from INDEPENDENT SET to DEGRADING MAX  $\alpha$ -FCGP in  $(q + 3)$ -closed graphs. Now, the theorem follows by Lemma A.2.  $\blacktriangleleft$

## B.12 Proof of Prop. 5.11

**Proof.** The proof is similar to the proof of Theorem 5.10. But for completeness we add this proof.

We provide a weak  $q$ -composition from INDEPENDENT SET on 2-closed graphs to MIN  $\alpha$ -FCGP in  $(q + 3)$ -closed graphs. Here, we assume that  $k > 3q + \lceil \frac{\alpha}{1 - 3\alpha} \rceil$ .

INDEPENDENT SET

**Input:** A graph  $G = (V, E)$  and an integer  $k$ .

**Question:** Is there an independent set of size exactly  $k$ ?

Let  $[t]^q$  be the set of  $q$ -dimensional vectors whose entries are in  $[t]$ . For a vector  $x \in [t]^q$  we denote by  $x_i$  the  $i$ -th entry of  $x$ . Assume that  $q \geq 2$  is a constant and that we are given exactly  $[t]^q$  instances  $\mathcal{I}_x := (G_x, k)$  of INDEPENDENT SET on 2-closed graphs for  $x \in [t]^q$ . Let  $V_x := V(G_x)$  for each  $x \in [t]^q$  and let  $D := \bigcup_{x \in [t]^q} V_x$ . We construct an equivalent instance  $(H, k', t')$  of MIN  $\alpha$ -FCGP as follows.



**Construction:** First for each  $x \in [t]^q$  we add the graph  $G_x$  to  $H$ . We then add a clique  $C$  consisting of  $tq$  vertices to  $H$ . The vertices of  $C$  are denoted by  $w_j^i$  with  $i \in [q]$  and  $j \in [t]$ . Furthermore, for each  $x \in [t]^q$  and  $v \in V_x$ , we add the edge  $vw_{x_i}^i$  for each  $i \in [q]$ . Fix an integer  $\ell \geq t^q \cdot n + \lceil (1 - 3\alpha)tq/\alpha \rceil$ . We add leaf vertices so that  $\deg(w_j^i) = \ell - \lceil (1 - 3\alpha)tq/\alpha \rceil$  for each vertex  $w_j^i$  and  $\deg(v) = \ell$  for each vertex  $v \in D$ . Next, for each added leaf vertex  $v$  in the last step, we add a clique  $C_v$  (including  $v$ ) of size  $\lceil 3t'/\alpha \rceil$  to  $H$ . By  $L$  we denote the set of newly added vertices with the cliques  $C_v$  for the new leaf vertices. Finally, we set  $k' := k + (t - 1)q$  and

$$t' := (1 - \alpha) \frac{(t - 1)q((t - 1)q - 1)}{2} + \alpha \left[ k\ell + (t - 1)q \left( \ell - \left\lceil \frac{(1 - 3\alpha)tq}{\alpha} \right\rceil \right) - (t - 1)q((t - 1)q - 1) \right].$$

**Closure:** We show that  $H$  is  $(q + 3)$ -closed. Note that  $H[L]$  is a cluster graph and thus has closure one. Furthermore, observe that each vertex in  $L$  has at most one neighbor in  $V(H) \setminus L$ . Also, since  $C$  is a clique, we only have to consider nonadjacent vertex pairs where one vertex  $u$  is in  $D$  and the other vertex  $v$  is in  $C \cup D$ . Without loss of generality we assume that  $u \in V_x$  for some  $x \in [t]^q$ . Recall that  $N(u) \subseteq V_x \cup C \cup L$  and  $u$  has exactly  $q$  neighbors in  $C$ . First, consider the case  $v \in C$ . By construction we obtain from  $uv \notin E$  that  $u'v \notin E$  for each  $u' \in V_x$ . Thus,  $|N(u) \cap N(v)| = q$ . Second consider the case  $v \in D$ . If  $v \notin V_x$ , then  $|N(u) \cap N(v)| \leq q - 1$ . Otherwise, if  $v \in V_x$ , we obtain  $|N(u) \cap N(v)| \leq q + 2$  from the fact that  $\mathcal{I}_x$  is 2-closed.

**Correctness:** In the following, we prove that there exists an independent set of size exactly  $k$  for some instance  $\mathcal{I}_x$  with  $x \in [t]^q$  if and only if there exists a vertex set  $S$  of size exactly  $k'$  in  $H$  such that  $\text{val}(S) \leq t'$ .

Suppose that instance  $\mathcal{I}_x$  has an independent set  $I$  of size exactly  $k$  for some  $x \in [t]^q$ . By  $C^* := C \setminus \{w_{x_i}^i \mid i \in [q]\}$  we denote the non-neighbors of  $V_x$  in  $C$ . Note that  $|C^*| = (t - 1)q$ . We show that  $S := I \cup C^*$  is a solution of  $(H, k', t')$ . Clearly,  $|S| = k + (t - 1)q = k'$ . Since no vertex of  $I$  is connected with any vertex in  $C^*$ ,  $I$  is an independent set, and since  $C^*$  is a clique of size  $(t - 1)q$ , we conclude that  $m_H(S) = (t - 1)q((t - 1)q - 1)/2$ . Furthermore, since each vertex in  $I$  has degree exactly  $\ell$ , and since each vertex in  $C^*$  has degree exactly  $\ell - \lceil (1 - 3\alpha)tq/\alpha \rceil$ , we conclude that  $m_H(S, V(H) \setminus S) = k\ell + (t - 1)q(\ell - \lceil (1 - 3\alpha)tq/\alpha \rceil) - (t - 1)q((t - 1)q - 1)$ . Thus,  $\text{val}(S) = t'$  and hence  $(H, k', t')$  is a Yes-instance of  $\text{MIN } \alpha\text{-FCGP}$ .

Conversely, suppose that  $(H, k', t')$  has a solution  $S \subseteq V(H)$  of size exactly  $k'$  with  $\text{val}(S) \leq t'$ . First, we show that we can assume that  $S \cap L = \emptyset$ . Assume that there exists a vertex  $v \in S \cap L$ . Since  $\deg(v) = \lceil 3t'/\alpha \rceil$  we conclude that  $m_H(\{v\}, S \setminus \{v\}) \geq 2t'\alpha$  and since  $\text{val}(S) \geq \alpha \cdot m_H(\{v\}, S \setminus \{v\})$  we obtain that  $\text{val}(S) > t'$ , a contradiction.

Thus, in the following we can assume that  $S \cap L = \emptyset$ . Let  $C^* := C \cap S$ ,  $|C^*| = z$ , and  $D^* := S \setminus C^* \subseteq D$ . In the following, we show that  $z = (t - 1)q$  and that there exists an  $x \in [t]^q$  such that  $N(V_x) \cap C^* = \emptyset$ . Therefore, we consider the cases that  $z < (t - 1)q$  and that  $z > (t - 1)q$ . In both cases we verify that  $\text{val}(S) > t'$  for each solution with exactly  $z$  vertices in  $C$ .

**Case 1:**  $z \leq (t - 1)q - 1$ . In other words,  $z = (t - 1)q - p$  for some  $p \in [(t - 1)q]$ . Recall that  $\deg(v) = \ell$  for each vertex  $v \in D$ . Thus, by Lemma B.7,  $\text{val}(S)$  is minimized if both  $m_H(D^*)$  and  $m_H(C^*, D^*)$  are minimized. Since  $z = |C^*| < (t - 1)q$ , it is possible

that no vertex of  $D^*$  is adjacent to any vertex in  $C^*$ . Thus,  $\text{val}(S)$  is minimized if  $D^*$  is an independent set and if  $E_H(C^*, D^*) = \emptyset$ . Hence,

$$\begin{aligned} \text{val}(S) = f_1(p) &\geq (1 - \alpha) \frac{((t-1)q - p)((t-1)q - p - 1)}{2} \\ &\quad + \alpha \left[ (k+p)\ell + ((t-1)q - p) \left( \ell - \left\lceil \frac{(1-3\alpha)tq}{\alpha} \right\rceil \right) \right] \\ &\quad - \alpha((t-1)q - p)((t-1)q - p - 1). \end{aligned}$$

Now, we obtain that

$$f'(p) = (1 - 3\alpha)p + \frac{3\alpha - 1}{2} (2(t-1)q - 1) + \alpha \left\lceil \frac{(1-3\alpha)tq}{\alpha} \right\rceil.$$

Since  $\alpha \in (0, 1/3)$  and since  $p \leq (t-1)q$ , we obtain that

$$f'(p) \geq \frac{3\alpha - 1}{2} (2(t-1)q - 1) + \alpha \frac{(1-3\alpha)tq}{\alpha} = (1 - 3\alpha)(q + 1/2) > 0.$$

Since  $\alpha \in [0, 1/3]$  we conclude that  $f(p)$  is a convex quadratic function. And since  $f'(p) > 0$  for each  $p \leq q$ , we thus conclude that  $f(0) < f(p)$  for each  $p \in [q]$ , a contradiction to the assumption that  $f(p) = \text{val}(S) \leq t'$ .

**Case 2:**  $z \geq (t-1)q + 1$ . Let  $z = (t-1)q + p$  for some  $p \in [q]$ . By the pigeonhole principle there exist at least  $p$  indices  $i \in [q]$  such that  $w_j^i \in S$  for each  $j \in [t]$ . Recall that by construction, each vertex  $v \in D$  has exactly one neighbor in the set  $\{w_j^i, j \in [t]\}$ . Since  $|D^*| = k - p$  we conclude that  $m_H(C^*, D^*) \geq (k - p)p$ . Recall that  $\deg(v) = \ell$  for each vertex  $v \in D$ . Thus, by Lemma B.7,  $\text{val}(S)$  is minimal if  $D^*$  is an independent set and if  $m_H(C^*, D^*) = (k - p)p$ . Thus,

$$\begin{aligned} \text{val}(S) = f(p) &\geq (1 - \alpha) \left[ (k - p)p + \frac{((t-1)q + p)((t-1)q + p - 1)}{2} \right] + \alpha(k - p)(\ell - p) \\ &\quad + \alpha p \left[ \ell - \left\lceil \frac{(1-3\alpha)tq}{\alpha} \right\rceil - (t-1)q - k + 1 \right] \\ &\quad + \alpha(t-1)q \left[ \ell - \left\lceil \frac{(1-3\alpha)tq}{\alpha} \right\rceil - (t-1)q - p + 1 \right]. \end{aligned}$$

Now, we obtain that

$$f'(p) = (3\alpha - 1)p + (1 - 3\alpha) \left( k + (t-1)q - \frac{1}{2} \right) - \alpha \left\lceil \frac{(1-3\alpha)tq}{\alpha} \right\rceil.$$

From  $p \leq q$  and  $3\alpha - 1 < 0$  for  $\alpha \in (0, 1/3)$ , we obtain that

$$\begin{aligned} f'(p) &\geq (3\alpha - 1)q + (1 - 3\alpha) \left( k + (t-1)q - 1/2 \right) - \alpha \left( \frac{(1-3\alpha)tq}{\alpha} + 1 \right) \\ &= (1 - 3\alpha) \left( k - 2q - 1/2 - \frac{\alpha}{1 - 3\alpha} \right). \end{aligned}$$

Since  $k > 3q + \left\lceil \frac{\alpha}{1-3\alpha} \right\rceil$  and since  $\alpha \in (0, 1/3)$ , we obtain that  $f'(p) > 0$  for  $p \leq q$ . Furthermore, since  $\alpha \in (0, 1/3)$  we see that  $f(p)$  is a concave quadratic function. Thus, we

conclude that  $f(0) < f(p)$  for each  $p \in [q]$ , a contradiction to the assumption that  $f(p) = \text{val}(S) \leq t'$ .

Hence,  $|C^*| = (t-1)q$  and thus  $|D^*| = k$ . According to Lemma B.7,  $\text{val}(S)$  is minimal if both  $m_H(D^*)$  and  $m_H(C^*, D^*)$  are minimal. Observe that if  $D^*$  is an independent set and if  $E_H(C^*, D^*) = \emptyset$ , then  $\text{val}(S) = t'$ . Otherwise, if  $E_H(C^*, D^*) \neq \emptyset$  or if  $D^*$  is no independent set, then  $\text{val}(S) > t'$ . Thus,  $E_H(C^*, D^*) = \emptyset$  and  $D^*$  is an independent set. Note that, if there exists two vertices  $u, v \in D^*$  such that  $u \in V_x$  and  $v \in V_y$  with  $x \neq y$  for  $x, y \in [t]^q$ , then  $(N(u) \cup N(v)) \cap C \geq q+1$ . Since  $|C^*| = (t-1)q$  this implies that  $E_H(C^*, D^*) \neq \emptyset$ , a contradiction. Hence,  $D^* \subseteq V_x$  for some  $x \in [t]^q$  and by Lemma B.7 the set  $D^*$  is an independent set. Thus, the instance  $x$  contains an independent set of size at least  $k$ .

Hence, we have a weak- $q$ -composition from INDEPENDENT SET to MIN  $\alpha$ -FCGP in  $(q+3)$ -closed graphs. Now, the theorem follows by Lemma A.2.  $\blacktriangleleft$

### B.13 Proof of Lemma 6.1

**Proof.** If  $|V(G)| < k$ , then we have a trivial no-instance and the statement follows immediately. Hence, in the following we assume that  $|V(G)| \geq k$ . Next, we show that there exists a solution  $S$  such that  $|F| \leq dk$  where  $F$  is the set of edges with at least one endpoint in  $S$ . Then, it follows that  $\text{val}(S) \leq t$  in non-trivial instances of MIN  $\alpha$ -FCGP and the statement is proven. Let  $\sigma$  be a degeneracy ordering of  $G$  and let  $S := \{v_1, \dots, v_k\}$  consist of the first  $k$  vertices of  $\sigma$ . Each edge in  $F$  is of the form  $xy$  where  $x < y$  with respect to  $\sigma$ . Since  $S$  consists of the first  $k$  vertices of  $\sigma$  and  $G$  is  $d$ -degenerate, we conclude that for  $x = v_i$  with  $i \in [k]$  there are at most  $d$  edges in  $F$  of the form  $xy$ . Since  $|S| = k$ ,  $F$  contains at most  $dk$  edges.  $\blacktriangleleft$

### B.14 Proof of Thm. 6.3

First, we consider the case  $\alpha = 0$ . Recall that SPARSEST  $k$ -SUBGRAPH is the special case of  $\alpha$ -FCGP for minimization and  $\alpha = 0$ .

### B.15 Proof of Prop. 6.4

**Proof.** Let  $(G, k, t)$  be an instance of SPARSEST  $k$ -SUBGRAPH. Assume that  $|V(G)| \geq dk$ . Now, since  $G$  is  $d$ -degenerate, we obtain an independent set  $I$  of size at least  $k$  in  $G$ . Observe that  $\text{val}(S) = 0$  and thus  $(G, k, t)$  is a trivial yes-instance. Hence,  $|V(G)| < dk$ . Since the number of edges in a  $d$ -degenerate graph is bounded by  $d \cdot |V(G)|$  the statement follows.  $\blacktriangleleft$

Now, we consider  $\alpha \in (0, 1/3)$ .

► **Lemma B.8.** MIN  $\alpha$ -FCGP for  $\alpha \in (0, 1/3)$  admits a kernel of size  $\mathcal{O}(d^4 k^5)$ .

**Proof.** Let  $I$  be an instance of MIN  $\alpha$ -FCGP for a fixed  $\alpha \in (0, 1/3)$ . First, we transform  $I$  into an equivalent instance  $I' := (G, \emptyset, \text{counter}, k, t)$  of ANNOTATED MIN  $\alpha$ -FCGP with parameter  $\alpha$ . Note that the upper bound for  $t$  from Lemma 6.2 still holds for  $I'$ . Observe that if  $G$  contains a vertex  $v$  such that  $\alpha \deg^{+c}(v) \geq t + k$ , then we have for each solution  $S$  containing  $v$  that  $\text{val}(S) > t$ . Hence, if there is a solution  $S$  for  $I'$ , then  $v \notin I$ . Thus, it is safe to apply Reduction Rule 3.3 to vertex  $v$ . In the following, we assume that Reduction Rule 3.3 is applied exhaustively.

Note that now we have  $\Delta(G) < (t+k)/\alpha$  and also  $\Gamma(G) < (t+k)/\alpha$ . Next, we can apply Reduction Rule 4.1 exhaustively. Similar to the proof of Proposition 4.2, after the exhaustive application of Reduction Rule 4.1 we have  $|V(G)| \leq \Delta k$ . Afterwards, with the reduction

described in Lemma 3.8, we can construct an equivalent instance  $(G', k, t')$  of MIN  $\alpha$ -FCGP with parameter  $\alpha$  of size  $\mathcal{O}((\Delta(G) + \Gamma(G))^3 |V(G)|) = \mathcal{O}(d^4 k^5)$ . ◀

Now, Proposition 6.4 and Lemma B.8 lead to Theorem 6.3.

### B.16 Proof of Prop. 6.5

**Proof.** First, we show the result for maximization and  $\alpha \in (1/3, 1]$ . Let  $L$  denote the vertices of  $G$  of degree at least  $t/(\alpha k)$ . Clearly, if  $|L| \geq dk$ , then there exists a subset  $I \subseteq L$  such that  $I$  induces an independent set in  $G$ . Since  $\deg(v) \geq t/(\alpha k)$  for each vertex  $v \in I$  we obtain that  $\text{val}(I) \geq t$ . Otherwise,  $|L| < dk$ . Since  $\alpha \in (1/3, 1]$  implies Degrading, we conclude that  $\text{cont}(v) > \text{cont}(u)$  for each vertex  $v \in L$  and each vertex  $u \in V(G) \setminus L$ . Since  $k$  vertices from  $V(G) \setminus L$  are not sufficient to reach value  $t$ , we conclude that each solution contains at least one vertex of  $L$ . Now, this implies an  $\mathcal{O}^*((dk)^k)$  time algorithm.

Second, we show the result for minimization and  $\alpha \in (0, 1/3)$ . Let  $L$  denote the vertices of  $G$  of degree at most  $t/(\alpha k)$ . Clearly, if  $|L| \geq dk$ , then there exists a subset  $I \subseteq L$  such that  $I$  induces an independent set in  $G$ . Since  $\deg(v) \leq t/(\alpha k)$  for each vertex  $v \in I$  we obtain that  $\text{val}(I) \leq t$ . Otherwise,  $|L| < dk$ . Since  $\alpha \in (0, 1/3)$  implies Degrading, we conclude that  $\text{cont}(v) < \text{cont}(u)$  for each vertex  $v \in L$  and each vertex  $u \in V(G) \setminus L$ . Since  $k$  vertices from  $V(G) \setminus L$  are lead to a value larger than  $t$ , we conclude that each solution contains at least one vertex of  $L$ . Now, this implies an  $\mathcal{O}^*((dk)^k)$  time algorithm. ◀

### B.17 Proof of Lemma 6.12

**Proof.** Since a  $d$ -degenerate graph has no  $K_{d,d}$  as a subgraph, there is a kernel of size  $(R(dk^{d-1} + k)^{O(1)})$  by Proposition 6.11. Recall that for  $\ell \in \mathbb{N}$ ,  $R(\ell)$  denotes an integer such that any graph on  $R(\ell)$  vertices has an independent set of size  $\ell$ . Since  $R(\ell) \leq (d+1)\ell$  for any  $d$ -degenerate graphs, there is a kernel of size  $(R(dk^{d-1} + k)^{O(1)}) \in k^{\mathcal{O}(d)}$ . ◀

### B.18 Proof of Prop. 6.13

**Proof.** We provide a weak  $q$ -composition from INDEPENDENT SET on 2-degenerate graphs to DEGRADING MAX  $\alpha$ -FCGP in  $(q+2)$ -degenerate graphs. Here, we assume that  $k > |c(\alpha)|/(3\alpha - 1)$  for some later specified constant  $c(\alpha) > 0$ .

INDEPENDENT SET

**Input:** A 2-degenerate graph  $G = (V, E)$  and an integer  $k$ .

**Question:** Is there an independent set of size exactly  $k$ ?

Let  $[t]^q$  be the set of  $q$ -dimensional vectors whose entries are in  $[t]$ . For a vector  $x \in [t]^q$  we denote by  $x_i$  the  $i$ th entry of  $x$ . Next, assume that  $q \geq 2$  is a constant and that we are given exactly  $t^q$  instances  $\mathcal{I}_x := (G_x, k)$  of INDEPENDENT SET on 2-degenerate graphs for  $x \in [t]^q$ . We construct an equivalent instance  $(H, k', t')$  of DEGRADING MAX  $\alpha$ -FCGP as follows.

**Construction:** We add an independent set  $J$  consisting of  $tq$  vertices to  $H$ . The vertices of  $J$  are denoted by  $w_j^i$  with  $i \in [q]$  and  $j \in [t]$ . Next, for each  $x \in [t]^q$  we add the graph  $G_x$  to  $H$ . By  $D$  we denote the union of these vertices. Furthermore, for each vertex  $v \in V_x$  for each  $x \in [t]^q$  we add the edge  $vw_{x_i}^i$  for each  $i \in [q]$ . Now, we fix an integer  $\ell > t^q \cdot n > k$ . For each vertex  $w_j^i$  in the independent set  $D$  we add leaf vertices such that  $\deg(w_j^i) = \ell + 1$ . Next,

for each vertex  $v \in D$  we add leaf vertices such that  $\deg(v) = \ell$ . By  $L$  we denote the union of all these added leaf vertices. Finally, we set  $k' := k + (t-1)q$  and  $t' := \alpha[k\ell + (t-1)q(\ell+1)]$ .

**Degeneracy:** Next, we show that  $d(H) = q + 2$ . Let  $S$  be any subgraph of  $H$ . We have to show that  $S$  contains a vertex with degree at most  $d(H) = q + 2$ . Clearly, if  $S$  contains a leaf vertex of  $L$ , then  $S$  contains a vertex of degree 1. Thus, in the following we can assume that  $S \cap L = \emptyset$ . If  $S \subseteq J$  then  $S$  is an independent set. Hence, let  $S^D := S \cap D \neq \emptyset$  and let  $S_x^D = S^D \cap V_x \neq \emptyset$  for some  $x \in [t]^q$ . Since by assumption  $\mathcal{I}_x$  is 2-degenerate, there exists a vertex  $v \in S_x^D$  such that  $|N(v) \cap S_x^D| \leq 2$ . By construction  $v$  has only neighbors in  $V_x$  and exactly  $q$  neighbors in  $J$ . Thus,  $v$  has degree at most  $q + 2$ .

**Correctness:** In the following, we prove that there exists an independent set of size exactly  $k$  for some instance  $\mathcal{I}_x$  with  $x \in [t]^q$  if and only if there exists a vertex set  $S$  of size exactly  $k'$  in  $H$  such that  $\text{val}(S) \geq t'$ .

Suppose that instance  $\mathcal{I}_x$  has an independent set  $I$  of size exactly  $k$  for some  $x \in [t]^q$ . By  $J^* := J \setminus \bigcup_{i \in [q]} \{w_{x_i}^i\}$  we denote the non-neighbors of  $V_x$  in  $J$ . Note that  $|J^*| = (t-1)q$ . We show that  $S := I \cup J^*$  is a solution of  $(H, k', t)$ . Clearly,  $|S| = k + (t-1)q = k'$ . Since no vertex of  $I$  is connected with any vertex in  $J^*$ ,  $I$  is an independent set, and since  $J^*$  is an independent set, we conclude that  $I \cup J^*$  is an independent set. Thus,  $E_H(S) = \emptyset$ . Furthermore, since each vertex in  $I$  has degree exactly  $\ell$ , and since each vertex in  $J^*$  has degree exactly  $\ell+1$ , we conclude that  $m_H(S, V(H) \setminus S) = k\ell + (t-1)q(\ell+1)$ . Thus,  $\text{val}(S) = t'$  and hence  $(H, k', t')$  is a Yes-instance of DEGRADING MAX  $\alpha$ -FCGP.

Conversely, suppose that  $(H, k', t')$  has a solution  $S \subseteq V(H)$  of size exactly  $k'$  with  $\text{val}(S) \geq t'$ . First, we show that  $S$  cannot contain any leaf-vertex in  $L$ . Assume towards a contradiction that  $S \cap L \neq \emptyset$ . Let  $v \in S \cap L$  and let  $S' := S \setminus \{v\}$ . According to Lemma B.7,  $\text{val}(S)$  is maximal if  $S'$  is an independent set and  $E(v, S') = \emptyset$ . Since the maximum degree in  $H$  is  $\ell + 1$  and since  $\deg(v) = 1$ , we obtain that  $\text{val}(S) \leq \alpha[1 + (k' - 1)(\ell + 1)]$ . Hence,

$$\begin{aligned} t - \text{val}(S) &\geq \alpha k\ell + \alpha(t-1)q(\ell+1) - [\alpha + \alpha k\ell + \alpha k + \alpha(t-1)q(\ell+1) - \alpha(\ell+1)] \\ &= \alpha(\ell - k) > 0 \end{aligned}$$

since  $\ell > k$ . Hence, in the following we can assume that  $S \cap L = \emptyset$ . Let  $J^* := J \cap S$ ,  $|J^*| = z$ , and  $D^* := S \setminus J^* \subseteq D$ . In the following, we show that  $z = (t-1)q$  and that there exists an  $x \in [t]^q$  such that  $N(V_x) \cap J^* = \emptyset$ . Therefore, we consider the cases that  $z < (t-1)q$  and that  $z > (t-1)q$ . In both cases we verify that  $\text{val}(S) < t'$  for each solution with exactly  $z$  vertices in  $J$ .

**Case 1:**  $z \leq (t-1)q - 1$ . Let  $p := (t-1)q - z$ . Note that  $p \in [(t-1)q]$ . Recall that  $\deg(v) = \ell$  for each vertex  $v \in D$ . Thus, by Lemma B.7,  $\text{val}(S)$  is maximized if  $m_H(D^*)$  and  $m_H(C^*, D^*)$  are minimized. Since  $|J^*| < (t-1)q$ , it is possible that no vertex of  $D^*$  is adjacent to any vertex in  $J^*$ . Thus,  $\text{val}(S)$  is maximal if  $S$  is an independent set with exactly  $(t-1)q - p$  vertices in  $J$ . Hence,  $\text{val}(S) \leq \alpha[(k+p)\ell + ((t-1)q - p)(\ell+1)]$ . Now, we obtain that

$$\begin{aligned} t - \text{val}(S) &\geq \alpha k\ell + \alpha(t-1)q(\ell+1) - [\alpha k\ell + \alpha p\ell + \alpha(t-1)q(\ell+1) - \alpha p(\ell+1)] \\ &= \alpha p > 0. \end{aligned}$$

Thus,  $t' - \text{val}(S) > 0$ , a contradiction.

**Case 2:**  $z \geq (t-1)q + 1$ . Let  $p := z - (t-1)q$ . In other words  $p \in [q]$ . By the pigeonhole principle there exist at least  $p$  indices  $i \in [q]$  such that  $w_j^i \in S$  for each  $j \in [t]$ . Recall that by construction, each vertex  $v \in D$  has exactly one neighbor in the set  $\{w_j^i, j \in [t]\}$ . Since  $|D^*| = k - p$  we conclude that  $m_H(C^*, D^*) \geq (k-p)p$ . Recall that  $\deg(v) = \ell$  for each vertex  $v \in D$ . Thus, by Lemma B.7,  $\text{val}(S)$  is maximized if  $m_H(D^*)$  and  $m_H(C^*, D^*)$  are minimized. Hence,  $\text{val}(S)$  is maximal if  $m_H(C^*, D^*) = (k-p)p$  and  $D^*$  is an independent set. Since  $H[J]$  is an independent set we obtain that

$$\text{val}(S) \leq (1-\alpha)(k-p)p + \alpha[(k-p)(\ell-p) + (t-1)q(\ell+1) + p(\ell+1-k+p)].$$

Now, we obtain that

$$\begin{aligned} t - \text{val}(S) &\geq \alpha k \ell + \alpha(t-1)q(\ell+1) - kp \\ &\quad + p^2 + 3\alpha kp - 3\alpha p^2 - \alpha k \ell - \alpha(t-1)q(\ell+1) - \alpha p \\ &= -kp + p^2 + 3\alpha kp - 3\alpha p^2 - \alpha p \\ &= (3\alpha - 1)kp + (1 - 3\alpha)p^2 - \alpha p. \end{aligned}$$

From  $p \geq 0$  and  $3\alpha - 1 > 0$  for  $\alpha \in (1/3, 1]$ , we obtain that

$$t - \text{val}(S) \geq (3\alpha - 1)k + (1 - 3\alpha)p^2 - \alpha p = (3\alpha - 1)k + c'(\alpha, p)$$

where  $c'(\alpha, p) < 0$  is a constant only depending on the parameters  $\alpha$ , and  $p$ . We set  $c(\alpha) := \max_{p \in [q]} c'(\alpha, p)$  which is a constant smaller than 0 only depending on  $\alpha$ . In other words,  $t - \text{val}(S) \geq (3\alpha - 1)k + c(\alpha) > 0$  since by assumption  $k > |c(\alpha)|/(3\alpha - 1)$ . A contradiction to the fact that  $\text{val}(S) \geq t'$ .

Hence,  $|J^*| = (t-1)q$  and thus  $|D^*| = k$ . According to Lemma B.7,  $\text{val}(S)$  is maximal if  $m_H(D^*)$  and  $m_H(J^*, D^*)$  are minimal. Observe that if  $E_H(J^*, D^*) = \emptyset$  and  $D^*$  is an independent set, then  $\text{val}(S) = t'$ . Otherwise, if  $E_H(J^*, D^*) \neq \emptyset$  or if  $D^*$  is no independent set, then  $\text{val}(S) < t'$ . Thus,  $E_H(J^*, D^*) = \emptyset$  and  $D^*$  is an independent set. Note that, if there exists two vertices  $u, v \in D^*$  such that  $u \in V_x$  and  $v \in V_y$  with  $x \neq y$  for  $x, y \in [t]^q$ , then  $(N(u) \cup N(v)) \cap J \geq q + 1$ . Since  $|J^*| = (t-1)q$  this implies that  $E_H(J^*, D^*) \neq \emptyset$ , a contradiction. Hence,  $D^* \subseteq V_x$  for some  $x \in [t]^q$ . Furthermore,  $J^* = J \setminus N(D^*)$ . Thus, the instance  $x$  contains an independent set of size at least  $k$ .

Hence, we have a weak- $q$ -composition from INDEPENDENT SET to DEGRADING MAX  $\alpha$ -FCGP in  $(d+2)$ -closed graphs. Now, the theorem follows by Lemma A.2.  $\blacktriangleleft$

## B.19 Proof of Prop. 7.1

The result is based on two data reduction rules. The first rule discards (according to the Exclusion Rule (Reduction Rule 3.3)) vertices with “low” contribution assuming there are “sufficient” vertices of high contribution. The second rule adds vertices with a “very high” contribution to a solution (according to the Inclusion Rule (Reduction Rule 3.2)) assuming there are only “few” vertices of “high” contribution. Subsequently, we specify concrete values for “sufficient”, “high” etc.

**► Definition B.9.** *Let  $\mathcal{I}$  be an instance of ANNOTATED  $\alpha$ -FCGP and let  $x \in \mathbb{N}$ . Then  $V_x := \{v \in V(G) \mid \deg^{+c}(v) \geq x\}$ .*

We start with some lemmas used in the proof of Lemma B.12.

► **Lemma B.10.** *Let  $\mathcal{I}$  be an yes-instance of ANNOTATED MAX  $\alpha$ -FCGP, let  $x \in \mathbb{N}$  with  $|V_x| \geq k$ , and let  $v \in V(G)$  be a vertex with  $\alpha \cdot \deg^{+c}(v) < \alpha x - |(1 - 3\alpha)k|$ . Then there is a solution  $S$  with  $v \notin S$ .*

**Proof.** Assume towards a contradiction that  $v$  is contained in each solution  $S$ . Observe that  $v \notin V_x$ . Since  $|S| = k$  it follows that there is a vertex  $u \in V_x \setminus S$ . We claim that  $S' := (S \setminus \{v\}) \cup \{u\}$  is also a solution. This follows from Lemmas 3.5 and 3.6. ◀

► **Lemma B.11.** *Let  $\mathcal{I}$  be an yes-instance of ANNOTATED MAX  $\alpha$ -FCGP, let  $x \in \mathbb{N}$  with  $|V_x| \leq k$ , and let  $v \in V(G)$  be a vertex with  $\alpha \deg^{+c}(v) \geq \alpha x + |(1 - 3\alpha)k|$ . Then there is a solution  $S$  with  $v \in S$ .*

**Proof.** Assume towards a contradiction that  $v$  is not contained in any solution  $S$ . Since  $|S| = k$ ,  $v \in V_x \setminus S$ , and  $|V_x| \leq k$  it follows that there is a vertex  $u \in S \setminus V_x$ . We claim that  $S' := (S \setminus \{u\}) \cup \{v\}$  is also a solution. This follows from Lemmas 3.5 and 3.6. ◀

► **Lemma B.12.** *Let  $\mathcal{I} := (G, k, t)$  be an instance of MAX  $\alpha$ -FCGP.*

1. *If  $\alpha > 0$  and there are at least  $k$  vertices with degree at least  $h + 1 + |(1 - 3\alpha)k/\alpha|$ , then there is an equivalent instance of size  $\mathcal{O}(h^3)$  computable in polynomial time.*
2. *If  $\alpha > 1/3$  and there are less than  $k$  vertices with degree at least  $h + 1 + |(1 - 3\alpha)k/\alpha|$ , then there is an equivalent instance of size  $\mathcal{O}(h^2k^2 + k^4)$  computable in polynomial time.*

**Proof of Lemma B.12.** First, we transform  $\mathcal{I}$  into an equivalent instance  $(G, \emptyset, \text{counter}, k, t)$  of ANNOTATED MAX  $\alpha$ -FCGP where  $\text{counter}(v) = 0$  for all  $v \in V(G)$ . Then  $V_{h+1}$  is the set of vertices of degree more than  $h$ , that is,  $|V_{h+1}| \leq h$ . We set  $Y := V(G) \setminus V_{h+1}$  to be the vertices with degree at most  $h$ . Moreover, we set  $x := h + 1 + |(1 - 3\alpha)k/\alpha| > h$  (recall  $\alpha > 0$ ). Thus,  $V_x \subseteq V_{h+1}$ .

**Part 1**  $|V_x| \geq k$  and  $\alpha > 0$ : For each vertex  $v \in Y$  we have  $\deg(v) = \deg^{+c}(v) \leq h < x - |(1 - 3\alpha)k/\alpha|$ . Hence, by Lemma B.10, there is a solution not containing  $v$ . We thus can apply Reduction Rule 3.3 on  $v$ . As this application does not change  $\deg^{+c}(u)$  for any  $u \in V(G) \setminus \{v\}$ , we can apply Reduction Rule 3.3 on all vertices in  $Y$  and obtain a graph with  $h$  vertices (the vertices in  $V_{h+1}$ ). Using Proposition 4.9, we get an instance for MAX  $\alpha$ -FCGP of size  $\mathcal{O}(h^3)$ .

**Part 2**  $|V_x| < k$  and  $\alpha > 1/3$ : Consider the set  $V_{x+|(1-3\alpha)k/\alpha|}$  of vertices with degree at least  $h + 2|(1 - 3\alpha)k/\alpha|$ . By Lemma B.11 for each  $v \in V_{x+|(1-3\alpha)k/\alpha|}$  there is a solution containing  $v$ . Thus, we can apply Reduction Rule 3.2 on all vertices in  $V_{x+|(1-3\alpha)k/\alpha|}$ . Afterwards, we have  $\Delta_{\overline{T}} \leq x + |(1 - 3\alpha)k/\alpha|$ . The exhaustive application of Reduction Rule 4.1 results in a graph with at most  $xk + |(1 - 3\alpha)k/\alpha|k = hk + |2/\alpha - 6|k^2 = \mathcal{O}(hk + k^2)$  vertices. Using Proposition 4.9, we get an instance for MAX  $\alpha$ -FCGP of size  $\mathcal{O}((hk + k^2)^2 + (hk + k^2)k^2) = \mathcal{O}(h^2k^2 + k^4)$ . ◀

Lemma B.12 implies the kernel for  $\alpha > 1/3$ . As NON-DEGRADING MAX  $\alpha$ -FCGP does not admit a polynomial kernel with respect to  $k$  even if  $\Delta$  is constant (see Thm. 4.11), it follows that the condition  $\alpha > 1/3$  in part 2 of Lemma B.12 cannot be improved to  $\alpha > 0$ .



## B.20 Proof of Prop. 7.2

**Proof.** We first transform  $\mathcal{I}$  into an equivalent instance  $(G, \emptyset, \text{counter}, k, t)$  of ANNOTATED MAX  $\alpha$ -FCGP where  $\text{counter}(v) = 0$  for all  $v \in V(G)$ . To make the description of the algorithm easier, we slightly change the definition of  $\text{val}$  by setting

$$\text{val}_G(V') := \alpha(m(V', V(G) \setminus V')) + \text{counter}(V') + (1 - \alpha)m(V'),$$

that is, we do not multiply the counter by  $\alpha$  but instead add correct multipliers when updating the counter.

Now, let  $V_{>h}$  be the set of vertices of degree at least  $h + 1$ . For each subset  $T$  of size at most  $k$  of  $V_{>h}$ , branch into the case that  $T = S \cap V_{>h}$ . In each case, we may now apply the Exclusion Rule (Rule 3.3) to remove the vertices of  $V_{>h} \setminus T$ . Now the contribution of  $T$  depends on which vertices of  $V(G) \setminus V_{>h}$  are contained in  $S$ . This can be incorporated into the counters of  $V(G) \setminus V_{>h}$  as follows. Pick any vertex  $u \in T$ . Then, decrease  $t$  by  $\text{counter}(u) + (1 - \alpha)|N(u) \setminus T| + \alpha|N(u) \cap T|$ . Now, for each vertex  $v \in N(u) \setminus T = N(u) \setminus V_{>h}$ , add  $(1 - \alpha) - \alpha = 1 - 2\alpha$  to  $\text{counter}(v)$ . The correctness of this step can be seen as follows: if  $v$  is not contained in  $S$ , then the contribution of  $uv$  is  $\alpha$  and this contribution is already recorded in the decrease of  $t$ . However, if we also add  $v$  to  $u$ , then the contribution of  $uv$  is  $1 - \alpha$ , so this gives a value of  $1 - \alpha$  for the internal edge  $uv$  minus  $\alpha$  for the fact that  $uv$  is no longer an outgoing edge. Finally, remove  $u$  from  $G$ .

After this removal of vertices in  $T$  has been applied exhaustively, the remaining graph has only vertices of degree at most  $h$ . Our aim is to find in this graph a vertex set  $S'$  of size  $k - |T|$  that maximizes  $\text{val}(S')$ . Now this problem can be solved in  $f(h, k) \cdot n^{\mathcal{O}(1)}$  time since  $\text{val}$  fulfills a property of fixed-cardinality optimization problems called *component linear* by Komusiewicz and Sorge [29]: First,  $\text{val}(S \cup T) \leq \text{val}(S) + \text{val}(T)$  because an internal edge counts twice as much as an outgoing edge. Second,  $\text{val}(S \cup T) \geq \text{val}(S) + \text{val}(T)$  if there are no edges between  $S$  and  $T$  in  $G$ .

Altogether, the running time is  $h^k \cdot n^{\mathcal{O}(1)} \cdot f(h, k)$  since we create  $h^k$  many cases in the branching on  $V_{>h}$ .  $\blacktriangleleft$

## B.21 Proof of Prop. 7.3

**Proof.** We follow the proof of Lemma B.12. As  $\text{vc} \geq h$ , we only need to work on Part 2, that is  $\alpha \in (0, 1/3)$ . Let  $(G, k, t)$  be an instance of MAX  $\alpha$ -FCGP. First, we transform  $(G, k, t)$  into an equivalent instance  $(G, \emptyset, \text{counter}, k, t)$  of ANNOTATED MAX  $\alpha$ -FCGP where  $\text{counter}(v) = 0$  for all  $v \in V(G)$ . Then, let  $X$  be a vertex cover of size  $\text{vc}$ . We set  $I := V(G) \setminus X$  to be the independent set. Note that each vertex in  $I$  has degree less at most  $\text{vc}$ . Moreover, we set  $x := \text{vc} + \lfloor (1 - 3\alpha)k/\alpha \rfloor > \text{vc}$  since  $\alpha > 0$ . Thus,  $V_x \subseteq X$ .

**Case 1**  $|V_x| \geq k$ : This case follows from Lemma B.12, Part 1.

**Case 2**  $|V_x| < k$ : Consider the set  $V_{x + \lfloor (1 - 3\alpha)k/\alpha \rfloor}$  of vertices with degree at least  $\text{vc} + 2\lfloor (1 - 3\alpha)k/\alpha \rfloor$ . By Lemma B.11 for each  $v \in V_{x + \lfloor (1 - 3\alpha)k/\alpha \rfloor}$  there is a solution containing  $v$ . Thus, we can apply Reduction Rule 3.2 on all vertices in  $V_{x + \lfloor (1 - 3\alpha)k/\alpha \rfloor}$ . Hence, we have  $\Delta_{\overline{T}} \leq x + \lfloor (1 - 3\alpha)k/\alpha \rfloor$ . Recall that  $T$  is the set of vertices fixed in the solution. Thus, there are at most  $\text{vc} \cdot \Delta_{\overline{T}} \in \mathcal{O}(\text{vc}^2 + \text{vc} \cdot k)$  vertices in  $I$  that have at least one neighbor in  $X \setminus T$ . Denote these vertices by  $I_{\overline{T}}$ . The remaining vertices in  $I \setminus I_{\overline{T}}$  have neighbors only in  $T$ . Hence, their contribution is fixed and we can get rid of all but the  $k$  vertices in  $I \setminus I_{\overline{T}}$  with highest contribution using Reduction Rule 3.3. Thus, we are left with the vertices in  $X$ , in  $I_{\overline{T}}$ , and

the  $k$  vertices with highest contribution of  $I \setminus I_{\overline{T}}$ . These are  $\mathcal{O}(\text{vc}^2 + \text{vc} \cdot k/\alpha)$  many vertices. Using Proposition 4.9, we get an instance for MAX  $\alpha$ -FCGP of size  $\mathcal{O}(\text{vc}^4 + \text{vc}^2 \cdot k^2 + \text{vc} \cdot k^3) = \mathcal{O}(\text{vc}^4 + \text{vc} \cdot k^3)$ .  $\blacktriangleleft$

## B.22 Proof of Prop. 7.4

**Proof.** Let  $X$  be a vertex cover of size  $\text{vc}$  and let  $I := V(G) \setminus X$  be the independent set. Without loss of generality we can assume that  $|I| \geq k$  since otherwise we already have a trivial  $\text{vc} + k$ -vertex kernel. If  $\alpha = 0$ , we have a trivial yes-instance as  $\text{val}(I') = 0$  for all  $I' \subseteq I$  of size  $k$ . Thus, in the following, we assume that  $\alpha > 0$ . Let  $(G, k, t)$  be an instance of MIN  $\alpha$ -FCGP. We transform  $(G, k, t)$  into an equivalent instance  $(G, \emptyset, \text{counter}, k, t)$  of ANNOTATED MIN  $\alpha$ -FCGP where  $\text{counter}(v) = 0$  for all  $v \in V(G)$ .

Let  $x := \text{vc} + \lfloor (1 - 3\alpha)k/\alpha \rfloor$ . Observe that there is a solution that does not contain any vertex in  $V_x$ : To this end, observe that  $\deg(v) \leq \text{vc}$  for each  $v \in I$ . Hence, by Lemma 3.6, each vertex in  $I$  is strictly better than each vertex in  $V_x$ . Since  $|I| \geq k$ , it follows from Lemma 3.5 that there is a solution not containing any vertex from  $V_x$ . Hence, we can apply Reduction Rule 3.3 on each vertex in  $V_x$ . As a result, the remaining vertices in  $X$  form still a vertex cover and have degree less than  $\text{vc} + \lfloor (1 - 3\alpha)k/\alpha \rfloor$ . Thus, less than  $\text{vc}(\text{vc} + \lfloor (1 - 3\alpha)k/\alpha \rfloor)$  vertices in  $I$  have neighbors in  $X$ ; the remaining vertices are isolated vertices. As the contribution of each isolated vertex  $v$  in any solution is exactly  $\alpha \cdot \text{counter}(v)$ , we can simply sort the isolated vertices by their contribution and remove all but the  $k$  vertices with the lowest contribution. Thus, we end up with at most  $\text{vc} + \text{vc}(\text{vc} + \lfloor (1 - 3\alpha)k/\alpha \rfloor) + k = \mathcal{O}(\text{vc}^2 + \text{vc} \cdot k)$  vertices. Using Proposition 4.9, we get an instance for MAX  $\alpha$ -FCGP of size  $\mathcal{O}((\text{vc}^2 + \text{vc} \cdot k) \cdot ((\text{vc}^2 + \text{vc} \cdot k) + k^2)^3) = \mathcal{O}(\text{vc}^8 + \text{vc} \cdot k^7)$ .  $\blacktriangleleft$