

# Noether currents and generators of local gauge transformations in the covariant canonical formalism

Satoshi Nakajima<sup>\*†</sup>

## Abstract

We investigate generators of local transformations in the covariant canonical formalism (CCF) for a gauge field and the second order formalism of gravity. The CCF treats space and time on an equal footing regarding the differential forms as the basic variables. The conjugate forms  $\pi_A$  are defined as derivatives of the Lagrangian  $d$ -form  $L(\psi^A, d\psi^A)$  with respect to  $d\psi^A$ , namely  $\pi_A := \partial L / \partial d\psi^A$ , where  $\psi^A$  are  $p$ -form dynamical fields. The form-canonical equations are derived from the form-Legendre transformation of the Lagrangian form  $H := d\psi^A \wedge \pi_A - L$ . We show that the Noether current form is the generator of an infinitesimal transformation  $\psi^A \rightarrow \psi^A + \delta\psi^A$  if the transformation of the Lagrangian form is given by  $\delta L = dl$  and  $\delta\psi^A$  and  $l$  are described by only  $\psi^A$  and the parameters. As an instance, we study the local gauge transformation for the gauge field and the local Lorentz transformation for the second order formalism of gravity.

## 1 Introduction

In the traditional analytical mechanics of fields, the canonical formalism gives special weight to time. The covariant canonical formalism (CCF) [1–11] is a covariant extension of the traditional canonical formalism. The form-Legendre transformation and the form-canonical equations are derived from a Lagrangian  $d$ -form with  $p$ -form dynamical fields  $\psi^A$ . The conjugate forms are defined as derivatives of the Lagrangian form with respect to  $d\psi^A$ . One can obtain the form-canonical equations of gauge theories or those of the second order formalism of gravity without fixing a gauge nor introducing the Dirac bracket or any other artificial tricks. Although the second order formalism of gravity (of which the dynamical variable is only the frame form (vielbein)) is a non-constrained system in the CCF, the first order formalism (of which the dynamical variables are both the frame form and the connection form) is a constrained system even in the CCF. In Refs. [1–4], the CCFs of the first order formalism of gravity and supergravity have been studied. Only for  $d = 4$ , the CCF of the second order formalism of gravity without Dirac field [6] and with Dirac field [7] have been studied. In Refs. [6, 7], the method to derive the form-canonical equations used special characteristics of  $d = 4$ . Without using those, we derive the form-canonical equations in this paper.

---

<sup>\*</sup>Department of Physics Engineering, Faculty of Engineering, Mie University, Tsu, Mie 514-8507, Japan

<sup>†</sup>subarusatosi@gmail.com

Poisson brackets of the CCF are proposed in Refs. [1, 10] and in Ref. [9] independently. These are equivalent. Although the form-canonical equations of the CCF are equivalent to modified De Donder-Weyl equations [8], the Poisson bracket of the CCF is not equivalent to it of the De Donder-Weyl theory [12, 13]. Reference [10] introduced the generator of the CCF and studied it of the local Lorentz transformation of gravity in the first order formalism. The generators of the local Lorentz transformation and the supersymmetry for supergravity have been studied [11] in the first order formalism. Relations between the Noether current and the generator of the CCF were not clear.

The structure of the paper is as follows. First, we review the covariant canonical formalism (§2). Next, in §3.1, we consider the Noether current form  $N$  for an infinitesimal local transformation  $\psi^A \rightarrow \psi^A + \delta\psi^A$ ,  $L = dl$  where  $\psi^A$  are differential forms of the dynamical fields and  $L$  is the Lagrangian form. If  $\delta\psi^A$  and  $l$  are described by only  $\psi^A$  and the parameters, the Noether current  $N$  described by  $\psi^A$  and  $\pi_A := \partial L / \partial d\psi^A$  is the generator of the transformation of the CCF (§3.2). As an instance, we study the local gauge transformation for the gauge field coupled with a matter field (§3.4) and the local Lorentz transformation for the second order formalism of gravity coupled with Dirac fields (§4). Our demonstrations are for non-constrained systems, in contrast to previous studies for constraint systems. In Appendix A, we study the canonical equations of the second order formalism of gravity in the arbitrary dimension  $d \geq 3$ . In Appendix B, several formulas are listed.

## 2 Covariant canonical formalism

In this section, we review the covariant canonical formalism.

Let us consider  $d$  dimension space-time. Suppose a  $p$ -form  $\beta$  is described by forms  $\{\alpha^i\}_{i=1}^k$ . If there exists the form  $\omega_i$  such that  $\beta$  behaves under variations  $\delta\alpha^i$  as  $\delta\beta = \delta\alpha^i \wedge \omega_i$ , we call  $\omega_i$  the *derivative* of  $\beta$  by  $\alpha^i$  and denote

$$\frac{\partial\beta}{\partial\alpha^i} := \omega_i. \quad (2.1)$$

In this case,  $\beta$  is *differentiable* by  $\alpha^i$ .

The Lagrangian  $d$ -form  $L$  is given by  $L = \mathcal{L}\eta$  where  $\mathcal{L}$  is the Lagrangian density and  $\eta = *1$  is the volume form<sup>1)</sup> and described by  $\psi$  and  $d\psi$ ,  $L = L(\psi, d\psi)$ , where  $\psi$  is a set the forms of the dynamical fields. For simplicity, we treat  $\psi$  as single  $p$ -form in this section. The Euler-Lagrange equation is given by

$$\frac{\partial L}{\partial\psi} - (-1)^p d \frac{\partial L}{\partial d\psi} = 0. \quad (2.2)$$

---

<sup>1)</sup>The Hodge operator  $*$  maps an arbitrary  $p$ -form  $\omega = \omega_{\mu_1 \dots \mu_p} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_p}$  ( $p = 0, 1, \dots, d$ ) to a  $(d-p)$ -form as

$$*\omega = \frac{1}{(d-p)!} E^{\mu_1 \dots \mu_p}_{\nu_1 \dots \nu_{d-p}} \omega_{\mu_1 \dots \mu_p} dx^{\nu_1} \wedge \dots \wedge dx^{\nu_{d-p}}.$$

Here,  $E_{\mu_1 \dots \mu_d}$  is the complete anti-symmetric tensor such that  $E_{01 \dots d-1} = \sqrt{-g}$  ( $g$  is the determinant of the metric  $g_{\mu\nu}$ ). And  $**\omega = -(-1)^{p(d-p)}\omega$  holds.

The above Euler-Lagrange equation has been used since the 1970's [14–16].

The *conjugate form*  $\pi$  is defined by

$$\pi := \frac{\partial L}{\partial d\psi}. \quad (2.3)$$

This is a  $q$ -form where  $q := d - p - 1$ . The *Hamilton  $d$ -form* is defined by

$$H(\psi, \pi) := d\psi \wedge \pi - L \quad (2.4)$$

and described by  $\psi$  and  $\pi$ . The variation of  $H$  is given by

$$\delta H = (-1)^{(p+1)q} \delta\pi \wedge d\psi - \delta\psi \wedge \frac{\partial L}{\partial\psi}. \quad (2.5)$$

Then, we obtain

$$\frac{\partial H}{\partial\psi} = -\frac{\partial L}{\partial\psi}, \quad \frac{\partial H}{\partial\pi} = (-1)^{(p+1)q} d\psi. \quad (2.6)$$

By substituting the Euler-Lagrange equation (2.2), we obtain the *canonical equations*

$$d\psi = (-1)^{(p+1)q} \frac{\partial H}{\partial\pi}, \quad d\pi = -(-1)^p \frac{\partial H}{\partial\psi}. \quad (2.7)$$

The *Poisson bracket* proposed in Ref. [9] is given by

$$\{A, B\} = (-1)^{p(a+d+1)} \frac{\partial A}{\partial\psi} \wedge \frac{\partial B}{\partial\pi} - (-1)^{(d+p-1)(a+1)} \frac{\partial A}{\partial\pi} \wedge \frac{\partial B}{\partial\psi}. \quad (2.8)$$

Here,  $A$  and  $B$  are differentiable by  $\psi$  and  $\pi$ , and  $A$  is an  $a$ -form. The Poisson bracket proposed in Ref. [10], denoted by  $\{A, B\}_F$ , is given by  $\{A, B\}_F = -\{B, A\}$ . If  $A$ ,  $B$  and  $C$  are  $a$ -form,  $b$ -form and  $c$ -form respectively and differentiable by  $\psi$  and  $\pi$ ,

$$\{B, A\} = -(-1)^{(a+d+1)(b+d+1)} \{A, B\}, \quad (2.9)$$

$$\{A, B \wedge C\} = \{A, B\} \wedge C + (-1)^{(a+d+1)b} B \wedge \{A, C\}, \quad (2.10)$$

and

$$\begin{aligned} (-1)^{(a+d+1)(c+d+1)} \{A, \{B, C\}\} + (-1)^{(b+d+1)(a+d+1)} \{B, \{C, A\}\} \\ + (-1)^{(c+d+1)(b+d+1)} \{C, \{A, B\}\} = 0 \end{aligned} \quad (2.11)$$

hold. The canonical equations can be written as

$$d\psi = -\{H, \psi\}, \quad d\pi = -\{H, \pi\}. \quad (2.12)$$

The fundamental brackets are

$$\{\psi, \pi\} = (-1)^{pd}, \quad \{\pi, \psi\} = -1, \quad \{\psi, \psi\} = 0 = \{\pi, \pi\}. \quad (2.13)$$

If a form  $F$  is differentiable by  $\psi$  and  $\pi$ , and  $F$  does not depend positively on space-time points,

$$\begin{aligned} dF &= d\psi \wedge \frac{\partial F}{\partial\psi} + d\pi \wedge \frac{\partial F}{\partial\pi} \\ &= (-1)^{(p+1)q} \frac{\partial H}{\partial\pi} \wedge \frac{\partial F}{\partial\psi} - (-1)^p \frac{\partial H}{\partial\psi} \wedge \frac{\partial F}{\partial\pi} \\ &= -\{H, F\} \end{aligned} \quad (2.14)$$

holds.

### 3 Generator of local gauge transformation

Let us consider that an infinitesimal transformation of dynamical fields  $\psi^A$  and its conjugate forms  $\pi_A$ :

$$\psi^A \rightarrow \psi^A + \delta\psi^A, \quad \pi_A \rightarrow \pi_A + \delta\pi_A. \quad (3.1)$$

Here,  $A$  is the label of the fields. If there exists  $(d-1)$ -form  $G$  such that

$$\delta\psi^A = \{\psi^A, G\}, \quad \delta\pi_A = \{\pi_A, G\}, \quad (3.2)$$

we call  $G$  the *generator* of the transformation [10]. If a form  $F$  is differentiable by  $\psi^A$  and  $\pi_A$ , the transformation of  $F$  is given by

$$\delta F = \{F, G\}. \quad (3.3)$$

In §3.1, we review the Noether current. In §3.2, we show that the Noether current is the generator under a few assumptions. In §3.3, we study Noether currents given by  $\varepsilon^r N_r + d\varepsilon^r \wedge F_r$ . In §3.4, we study the local gauge transformation for gauge field coupled with a matter field.

#### 3.1 Noether current

We explain the Noether current. For an infinitesimal transformation of  $p$ -form dynamical fields  $\psi^A \rightarrow \psi^A + \delta\psi^A$ , an identical equation

$$\delta L \equiv \delta\psi^A \wedge [L]_A + d\left(\delta\psi^A \wedge \frac{\partial L}{\partial d\psi^A}\right) \quad (3.4)$$

holds. Here,  $[L]_A := \partial L / \partial \psi^A - (-1)^p d(\partial L / \partial d\psi^A)$  and  $\equiv$  denotes identical equation which holds without using the Euler-Lagrange equations. If  $\delta L = dl$  holds, under the Euler-Lagrange equations  $[L]_A = 0$ , the Noether current

$$N := \delta\psi^A \wedge \frac{\partial L}{\partial d\psi^A} - l \quad (3.5)$$

is conserved

$$dN = 0. \quad (3.6)$$

#### 3.2 The Noether current is the generator

In the following of this paper, we suppose that  $\delta\psi^A$  and  $l$  are described by only  $\psi^A$  and the parameters: These do not include  $d\psi^A$  or  $\pi_A$ .

In this subsection, first, we calculate the transformation formula of  $\pi_A$ . Next, we show that the Noether current is the generator of the transformation.

The variation of  $\psi'^A$  is given by

$$\begin{aligned}\Delta\psi'^A &= \psi'^A + \Delta\psi^B \wedge \frac{\partial\delta\psi^A}{\partial\psi^B} \\ &= (\delta_B^A + \xi_B^A)\Delta\psi^B.\end{aligned}\tag{3.7}$$

Here, we supposed that nonzero components of

$$\xi_B^A := \frac{\partial\delta\psi^A}{\partial\psi^B}\tag{3.8}$$

are 0-forms. Using (3.7), we have

$$\Delta\psi^A = (\delta_B^A - \xi_B^A)\Delta\psi'^B,\tag{3.9}$$

$$\Delta d\psi^A = (\delta_B^A - \xi_B^A)\Delta d\psi'^B - (-1)^p\Delta\psi'^B \wedge d\xi_B^A.\tag{3.10}$$

Here, we used  $\Delta d\psi^A = d\Delta\psi^A$ . Substituting the above two equations into

$$\Delta L = \Delta\psi^A \wedge \frac{\partial L}{\partial\psi^A} + \Delta d\psi^A \wedge \frac{\partial L}{\partial d\psi^A},\tag{3.11}$$

we have

$$\frac{\partial L}{\partial d\psi'^A} = (\delta_A^B - \xi_B^A)\pi_B.\tag{3.12}$$

The derivative of  $L' = L + dl$  by  $d\psi'^A$  is given by

$$\begin{aligned}\pi'_A &= \frac{\partial L'}{\partial d\psi'^A} \\ &= \frac{\partial L}{\partial d\psi'^A} + \frac{\partial l}{\partial\psi'^A} \\ &= (\delta_A^B - \xi_B^A)\pi_B + \frac{\partial l}{\partial\psi^A}.\end{aligned}\tag{3.13}$$

We show that the Noether current  $N$  is the generator of the transformation:

$$\begin{aligned}\{\psi^A, N\} &= (-1)^{p(d+1)} \frac{\partial N}{\partial\pi_A} \\ &= \delta\psi^A,\end{aligned}\tag{3.14}$$

$$\begin{aligned}\{\pi_A, N\} &= -(-1)^{(d+p+1)(d-p)} \frac{\partial N}{\partial\psi^A} \\ &= -\xi_B^A\pi_B + \frac{\partial l}{\partial\psi^A} \\ &= \delta\pi_A.\end{aligned}\tag{3.15}$$

Here, we used (3.13).

### 3.3 $\varepsilon^r N_r + d\varepsilon^r \wedge F_r$ type Noether current

In the following of this paper, we consider the (local) Noether currents given by

$$N = \varepsilon^r N_r + d\varepsilon^r \wedge F_r, \quad (3.16)$$

where  $\varepsilon^r$  are the infinitesimal parameters (0-forms). Here,  $N_r$  are the (global) Noether currents for global transformation. Under the Euler-Lagrange equations  $[L]_A = 0$ ,  $N_r$  are conserved

$$dN_r = 0. \quad (3.17)$$

Using (3.6) and (3.17), we have

$$d\varepsilon^r \wedge (N_r - dF_r) = 0. \quad (3.18)$$

This leads to

$$N_r = -\{F_r, H\}. \quad (3.19)$$

Here, we used (2.14). According to Ref. [10], (3.19) holds without using the Euler-Lagrange equations.

For the infinitesimal local gauge transformation for gauge field and the infinitesimal local Lorentz transformation for the second order formalism of gravity, the Noether currents are given by (3.16). In the next subsection, we study a gauge field and calculate  $N_r$  and  $F_r$  and confirm (3.19). In §4, we study the Noether current and the generator of the infinitesimal Lorentz transformation for the second order formalism of gravity.

### 3.4 Local gauge transformation

We consider a gauge field  $A^r$  coupled with  $p$ -form matter field  $\psi^A$ . For instance, the matter field is the scalar field. To simplify, we suppose that the matter field is not the Dirac field because the conjugate form of the Dirac field is not independent. The total Lagrangian form is given by

$$L = L_0(\psi^A, (D\psi)^A) + L_1, \quad (3.20)$$

where  $(D\psi)^A$  is the covariant derivative. Here,  $L_0$  and  $L_1$  are the Lagrangian forms for the matter field and the gauge field respectively. The infinitesimal gauge transformation is given by

$$\delta\psi^A = \varepsilon^r (\mathbf{G}_r)^A_B \psi^B, \quad \delta A^r = \varepsilon^s f^r_{st} A^t - d\varepsilon^r, \quad \delta L_0 = 0, \quad \delta L_1 = 0. \quad (3.21)$$

Here,  $\varepsilon^r$  are infinitesimal parameters,  $\mathbf{G}_r$  are representations of the generators of a linear Lie group  $\mathcal{G}$ . The matrices  $\mathbf{G}_r$  satisfy

$$[\mathbf{G}_r, \mathbf{G}_s] = f^t_{rs} \mathbf{G}_t, \quad (3.22)$$

where  $[A, B] := AB - BA$  and  $f^t_{rs}$  are the structure constants of  $\mathcal{G}$ . The covariant derivative is given by  $(D\psi)^A := d\psi^A + A^r (\mathbf{G}_r)^A_B \wedge \psi^B$ .  $L_1$  is given by

$$L_1 = -\frac{1}{2k} \mathcal{F}^r \wedge * \mathcal{F}_r, \quad (3.23)$$

where  $k$  is a positive constant,  $\mathcal{F}^r := dA^r + \frac{1}{2}f^r_{bc}A^b \wedge A^c$  is the curvature of the gauge field, and  $\mathcal{F}_r := \kappa_{rs}\mathcal{F}^s$  where  $\kappa_{rs} := -f^a_{rb}f^b_{sa}(= \kappa_{sr})$  is the Killing form. The global Noether currents  $N_r$  are given by

$$N_r = N_r^{(0)} + N_r^{(1)}, \quad (3.24)$$

$$N_r^{(0)} = (\mathbf{G}_r)^A_B \psi^B \wedge \pi_A, \quad (3.25)$$

$$N_r^{(1)} = f^s_{rt} A^t \wedge \pi_s \quad (3.26)$$

Here,  $\pi_A$  and  $\pi_r$  are conjugate forms of  $\psi^A$  and  $\pi_r$  respectively. The global Noether currents satisfy

$$\{G_r, G_s\} = f^t_{rs} G_t \quad (G_r = N_r, N_r^{(0)}, N_r^{(1)}). \quad (3.27)$$

$F_r$  of (3.16) are given by

$$F_r = -\pi_r. \quad (3.28)$$

We can confirm that the local Noether current  $N$  is the generator of the transformation:

$$\delta\psi^A = \{\psi^A, N\} = \varepsilon^r (\mathbf{G}_r)^A_B \psi^B, \quad (3.29)$$

$$\delta A^r = \{A^r, N\} = \varepsilon^s f^r_{st} A^t - d\varepsilon^r, \quad (3.30)$$

$$\delta\pi_A = \{\pi_A, N\} = -\varepsilon^r (\mathbf{G}_r)^B_A \pi_B, \quad (3.31)$$

$$\delta\pi_r = \{\pi_r, N\} = -\varepsilon^s f^t_{sr} \pi_t. \quad (3.32)$$

We confirm (3.19). The conjugate form of  $A^a$  is given by

$$\pi_a = -\frac{1}{k} * \mathcal{F}_a. \quad (3.33)$$

The Hamilton form is given by

$$H = H_1 - L_0(\psi^A, (D\psi)^A), \quad (3.34)$$

$$H_1 := -\frac{1}{2} f^a_{bc} A^b \wedge A^c \wedge \pi_a + \frac{k}{2} \pi_a \wedge * \pi^a, \quad (3.35)$$

with  $\pi^a := (\kappa^{-1})^{ab} \pi_b = -\frac{1}{k} * \mathcal{F}^a$ . The derivatives of  $H$  are given by

$$\frac{\partial H}{\partial A^a} = -f^c_{ab} A^b \wedge \pi_c - J_a, \quad (3.36)$$

$$\frac{\partial H}{\partial \pi_a} = k * \pi^a - \frac{1}{2} f^a_{bc} A^b \wedge A^c. \quad (3.37)$$

Here,

$$J_r := \frac{\partial L_0(\psi^A, (D\psi)^A)}{\partial A^r} = (\mathbf{G}_r)^A_B \psi^B \wedge \pi_A = N_r^{(0)}. \quad (3.38)$$

Then, we have

$$\begin{aligned}
-\{F_r, H\} &= -\frac{\partial H}{\partial A^r} \\
&= f^c{}_{rb} A^b \wedge \pi_c + J_r \\
&= N_r.
\end{aligned} \tag{3.39}$$

Here, we used (3.38).

We remark that the canonical equations  $dA^a = \partial H/\partial \pi_a$  and  $d\pi_a = \partial H/\partial A^a$  become

$$dA^a = k * \pi^a - \frac{1}{2} f^a{}_{bc} A^b \wedge A^c, \tag{3.40}$$

$$d\pi_a = -f^c{}_{ab} A^b \wedge \pi_c - J_a. \tag{3.41}$$

The above two equations can be rewritten as

$$\mathcal{F}^a = k * \pi^a, \quad D\pi_a := d\pi_a + f^c{}_{ab} A^b \wedge \pi_c = -J_a. \tag{3.42}$$

The former is equivalent to the definition of  $\pi_a$ . The latter is equivalent to the Yang-Mills-Utiyama equation. The covariant canonical formalism does not need the gauge fixing.

## 4 Second order formalism of gravity

In this section, we study the Noether current and the generator of the infinitesimal local Lorentz transformation for the second order formalism of gravity coupled with Dirac fields in the arbitrary dimension ( $d \geq 3$ ). In §4.1, we introduce the notations. In §4.2, we review the Lagrange formalism of the second order formalism of gravity. In §4.3, we study the local Noether current. Our study is for a non-constrained system (the second order formalism), in contrast to previous studies [10,11] for constraint systems (the first order formalism).

### 4.1 Notations

We explain the notations used in this paper. Let  $g$  be the metric of which has signature  $(-+\cdots+)$ , and let  $\{\theta^a\}_{a=0}^{d-1}$  denote an orthonormal frame (vielbein). We have  $g = \mathring{g}_{ab} \theta^a \otimes \theta^b$  with  $\mathring{g}_{ab} := \text{diag}(-1, 1, \cdots, 1)$ . All indices are lowered and raised with  $\mathring{g}_{ab}$  or its inverse  $\mathring{g}^{ab}$ . The first structure equation is

$$d\theta^a + \omega^a{}_b \wedge \theta^b = \Theta^a, \tag{4.1}$$

where  $\omega^a{}_b$  is the connection form and  $\Theta^a$  is the torsion 2-form. In this paper, we suppose  $\omega_{ba} = -\omega_{ab}$ . Let  $A^a{}_b$  be the Levi-Civita connection. Then,  $\omega^a{}_b$  is given by

$$\omega^a{}_b = A^a{}_b + K^a{}_b. \tag{4.2}$$

For the infinitesimal local Lorentz transformation

$$\delta\theta^a = \varepsilon^a{}_b \theta^b, \tag{4.3}$$



$\omega^{ab}$ ,  $A^{ab}$ , and  $K^{ab}$  behave as

$$\delta\omega^{ab} = \varepsilon^a_c \omega^{cb} + \varepsilon^b_c \omega^{ac} - d\varepsilon^{ab}, \quad (4.4)$$

$$\delta A^{ab} = \varepsilon^a_c A^{cb} + \varepsilon^b_c A^{ac} - d\varepsilon^{ab}, \quad (4.5)$$

$$\delta K^{ab} = \varepsilon^a_c K^{cb} + \varepsilon^b_c K^{ac}. \quad (4.6)$$

Here,  $\varepsilon^{ab}$  are infinitesimal parameters which satisfy  $\varepsilon^{ab} = -\varepsilon^{ba}$ .

We put

$$\eta^a = *\theta^a, \quad \eta^{ab} = *(\theta^a \wedge \theta^b), \quad \eta^{abc} = *(\theta^a \wedge \theta^b \wedge \theta^c), \quad \eta^{abcd} = *(\theta^a \wedge \theta^b \wedge \theta^c \wedge \theta^d). \quad (4.7)$$

In Appendix B, several identities about  $\theta^b \wedge \eta_{a_1 \dots a_r}$  ( $r = 1, 2, 3, 4$ ),  $\delta\eta_{a_1 \dots a_r}$  ( $r = 0, 1, 2, 3$ ) and  $d\eta_{a_1 \dots a_r}$  ( $r = 1, 2, 3$ ) are listed.

The curvature 2-form  $\Omega^a_b$  is given by

$$\Omega^a_b = d\omega^a_b + \omega^a_c \wedge \omega^c_b. \quad (4.8)$$

Expanding the curvature form as

$$\Omega^a_b = \frac{1}{2} R^a_{bcd} \theta^c \wedge \theta^d, \quad (4.9)$$

we define

$$R_{ab} := R^c_{acb}, \quad R := R^a_a. \quad (4.10)$$

Because of (B.2),  $R\eta$  can be written as

$$R\eta = \Omega^{ab} \wedge \eta_{ab}. \quad (4.11)$$

## 4.2 Lagrange formalism

The Lagrangian form of the gravity in the second order formalism is given by

$$L(\theta, d\theta) = L_G(\theta, d\theta) + L_{\text{mat}}(\theta, d\theta). \quad (4.12)$$

Here,  $L_G$  is the Lagrangian form for the pure gravity given by

$$L_G(\theta, d\theta) = \frac{1}{2\kappa} W', \quad W' := R\eta - d(\omega^{ab} \wedge \eta_{ab}), \quad (4.13)$$

and  $L_{\text{mat}}(\theta, d\theta) := L_{\text{mat}}(\theta, \omega(\theta, d\theta))$  is the Lagrangian form of ‘‘matters’’ which are scalar fields, Dirac fields and gauge fields. Here,  $\kappa$  is the Einstein constant. Only the Dirac fields couple to  $\omega^{ab}$ .

We derive the Euler-Lagrange equation of the gravity. The variation of  $L$  is given by <sup>2)</sup>

$$\begin{aligned} \delta L(\theta, d\theta) = & \delta\theta^c \wedge \left( \frac{1}{2\kappa} [\Omega^{ab} \wedge \eta_{abc} - d(\omega^{ab} \wedge \eta_{abc})] + T_c \right) + \delta d\theta^c \wedge \frac{1}{2\kappa} \omega^{ab} \wedge \eta_{abc} \\ & + \delta\omega^{ab}(\theta, d\theta) \wedge \left( \frac{1}{2\kappa} [d\eta_{ab} - \omega^c{}_a \wedge \eta_{cb} - \omega^c{}_b \wedge \eta_{ac}] + \frac{\partial L_{\text{mat}}}{\partial \omega^{ab}} \right), \end{aligned} \quad (4.15)$$

where

$$T_a := \frac{\partial L_{\text{mat}}(\theta, \omega)}{\partial \theta^a} \quad (4.16)$$

is the energy-momentum form. We suppose that

$$\frac{1}{2\kappa} [d\eta_{ab} - \omega^c{}_a \wedge \eta_{cb} - \omega^c{}_b \wedge \eta_{ac}] + \frac{\partial L_{\text{mat}}}{\partial \omega^{ab}} \equiv 0, \quad (4.17)$$

which is the same as the Euler-Lagrange equation for  $\omega^{ab}$  of the first order formalism <sup>3)</sup>. Under this supposition, (4.15) leads to

$$\frac{\partial L}{\partial \theta^c} = \frac{1}{2\kappa} [\Omega^{ab} \wedge \eta_{abc} - d(\omega^{ab} \wedge \eta_{abc})] + T_c, \quad \frac{\partial L}{\partial d\theta^c} = \frac{1}{2\kappa} \omega^{ab} \wedge \eta_{abc}. \quad (4.18)$$

The Euler-Lagrange equation  $\partial L / \partial \theta^c + d(\partial L / \partial d\theta^c) = 0$  becomes the Einstein equation

$$-\frac{1}{2\kappa} \Omega^{ab} \wedge \eta_{abc} = T_c. \quad (4.19)$$

We remark that (4.17) can be rewritten as

$$\frac{1}{2\kappa} \Theta^c \wedge \eta_{abc} \equiv -\frac{\partial L_{\text{mat}}}{\partial \omega^{ab}} \quad (4.20)$$

using (B.4). If we expand  $T_c$  as  $T_c = T_c{}^b \eta_b$ , (4.19) leads to

$$R^a{}_b - \frac{1}{2} R \delta_b^a = \kappa T_b{}^a. \quad (4.21)$$

---

<sup>2)</sup>  $\delta W' = \delta(\omega^a{}_c \wedge \omega^{cb} \wedge \eta_{ab} + \omega^{ab} \wedge d\eta_{ab})$  and

$$\omega^{ab} \wedge \delta d\eta_{ab} = \delta d\theta^c \wedge \omega^{ab} \wedge \eta_{abc} + \delta\theta^c \wedge \omega^{ab} \wedge d\eta_{abc} \quad (4.14)$$

hold. Here, we used the following formulas:

$$d\eta_{ab} = d\theta^c \wedge \eta_{abc}, \quad \delta\eta_{abc} = \delta\theta^d \wedge \eta_{abcd}, \quad d\eta_{abc} = d\theta^d \wedge \eta_{abcd}.$$

<sup>3)</sup> In the second order formalism, (4.17) is not an equation of motion.  $\omega^{ab}$  is determined by (4.17). The method to obtain (4.18) in this subsection is also called the 1.5 order formalism.

### 4.3 Generator of the local Lorentz transformation

The local Noether current  $N$  for the transformation (4.3) is given by

$$N = \frac{1}{2}\varepsilon^{ab}N_{ab} + \frac{1}{2}d\varepsilon^{ab} \wedge F_{ab}, \quad (4.22)$$

with

$$N_{ab} = N_{ab}^{(0)} + N_{ab}^{(1)}, \quad (4.23)$$

$$N_{ab}^{(0)} = 2\frac{\partial L_{\text{mat}}}{\partial \omega^{ab}}, \quad (4.24)$$

$$N_{ab}^{(1)} = 2\theta_{[b} \wedge \pi_{a]} = \theta_b \wedge \pi_a - \theta_a \wedge \pi_b. \quad (4.25)$$

Here, we used (3.38) for  $N_{ab}^{(0)}$ .  $[\ ]$  denotes the antisymmetrization. The global Noether currents  $N_{ab}^{(1)}$  satisfy

$$\{N_{ab}^{(1)}, N_{cd}^{(1)}\} = \mathring{g}_{bc}N_{ad}^{(1)} - \mathring{g}_{ac}N_{bd}^{(1)} + \mathring{g}_{ad}N_{bc}^{(1)} - \mathring{g}_{bd}N_{ac}^{(1)}. \quad (4.26)$$

In the next paragraph, we will show that

$$F_{ab} = -\frac{1}{\kappa}\eta_{ab}. \quad (4.27)$$

We can confirm that  $N$  is the generator of the local Lorentz transformation:

$$\delta\theta^a = \{\theta^a, N\} = \varepsilon^a_b\theta^b, \quad (4.28)$$

$$\delta\pi_a = \{\pi_a, N\} = -\varepsilon^b_a\pi_b - d\varepsilon^{bc} \wedge \frac{1}{2\kappa}\eta_{abc}. \quad (4.29)$$

We derive (4.27).  $W'$  of (4.13) can be rewritten as

$$W' = W^* + \mathcal{K}, \quad (4.30)$$

$$W^* := A^a_c \wedge A^{cb} \wedge \eta_{ba}, \quad (4.31)$$

$$\mathcal{K} := K^a_c \wedge K^{cb} \wedge \eta_{ab}. \quad (4.32)$$

Then, by introducing  $L_G^* := W^*/(2\kappa)$ , we have

$$\begin{aligned} \delta L &= \delta L_G^* \\ &= \delta\theta^a \wedge \frac{\partial L_G^*}{\partial \theta^a} + d(\delta\theta^a) \wedge \frac{\partial L_G^*}{\partial d\theta^a} \\ &= \varepsilon^a_b \left( \theta^b \wedge \frac{\partial L_G^*}{\partial \theta^a} + d\theta^b \wedge p_a \right) + d\varepsilon^{ab} \wedge \theta_{[b} \wedge p_{a]}, \end{aligned} \quad (4.33)$$

where

$$p_a := \frac{\partial L_G^*}{\partial d\theta^a} = \frac{1}{2\kappa}A^{bc} \wedge \eta_{abc}. \quad (4.34)$$

Because of the global Lorentz invariance, the first term of (4.33) vanishes. Using (B.4), we have

$$\theta_{[b} \wedge p_{a]} = \frac{1}{\kappa} A_{[b}^c \wedge \eta_{a]c} = \frac{1}{2} dF_{ab}. \quad (4.35)$$

Then, we have

$$\delta L = \frac{1}{2} d\varepsilon^{ab} \wedge dF_{ab} = dl, \quad l := -\frac{1}{2} d\varepsilon^{ab} \wedge F_{ab}. \quad (4.36)$$

The above equation and (3.5) lead to (4.27).

We confirm (3.19). The conjugate form of  $\theta^a$  is given by

$$\pi_a = \frac{1}{2\kappa} \omega^{bc} \wedge \eta_{abc}. \quad (4.37)$$

The Hamilton form is given by

$$H(\theta, \pi) = d\theta^a \wedge \pi_a - L = H_G(\theta, \pi) - L_{\text{mat}}(\theta, \pi). \quad (4.38)$$

Here,  $H_G$  is the Hamilton form for gravity. Then, we have

$$\begin{aligned} -\{F_{ab}, H\} &= \frac{1}{\kappa} \{\eta_{ab}, H_G\} - \frac{1}{\kappa} \{\eta_{ab}, L_{\text{mat}}\} \\ &= -\frac{1}{\kappa} \frac{\partial \eta_{ab}}{\partial \theta^c} \wedge \frac{\partial H_G}{\partial \pi_c} + \frac{1}{\kappa} \frac{\partial \eta_{ab}}{\partial \theta^c} \wedge \frac{\partial L_{\text{mat}}}{\partial \pi_c} \\ &= -\frac{1}{\kappa} \eta_{abc} \wedge (-\omega^c_d \wedge \theta^d) - \frac{1}{\kappa} \eta_{abc} \wedge \Theta^c \\ &= \frac{1}{\kappa} \omega^c_d \wedge \theta^d \wedge \eta_{abc} + 2 \frac{\partial L_{\text{mat}}}{\partial \omega^{ab}}. \end{aligned} \quad (4.39)$$

We used (A.12), (B.3) and

$$-\frac{\partial L_{\text{mat}}}{\partial \pi_a} = \Theta^a \quad (4.40)$$

in the third line, and (4.20) in the fourth line. (4.40) is derived from (4.20). Because

$$\kappa N_{ab}^{(1)} = -\omega^{cd} \wedge \theta_{[b} \wedge \eta_{a]cd} = -2\omega^c_{[b} \wedge \eta_{a]c} = \omega^c_d \wedge \theta^d \wedge \eta_{abc} \quad (4.41)$$

holds using (B.1), we have

$$N_{ab} = -\{F_{ab}, H\}. \quad (4.42)$$

## 5 Summary

In the covariant canonical formalism (CCF), we showed that the Noether current is the generator of the infinitesimal local transformation of  $p$ -form dynamical fields  $\psi^A \rightarrow \psi^A + \delta\psi^A$  if the transformation of the Lagrangian form is given by  $\delta L = dl$  and  $\delta\psi^A$  and  $l$  are described by only  $\psi^A$  and the parameters. As an instance, we studied the local gauge transformation for the gauge field and the local Lorentz transformation for the second order formalism of gravity. Our demonstrations are for non-constrained systems in the CCF, in contrast to previous studies [10, 11] for constraint systems.

## A Covariant canonical formalism of gravity

We study the canonical equations of the second order formalism of gravity in the arbitrary dimension  $d \geq 3$ .

In (4.13),  $W'$  can be rewritten as

$$\begin{aligned} W' &= \omega^a_c \wedge \omega^{cb} \wedge \eta_{ab} + \omega^{ab} \wedge d\eta_{ab} \\ &= W + \Theta^a \wedge \omega^{bc} \wedge \eta_{abc} \end{aligned} \quad (\text{A.1})$$

with

$$W := \omega^a_c \wedge \omega^{cb} \wedge \eta_{ba}. \quad (\text{A.2})$$

In the second line of (A.1), we used (B.4). Using (4.1) and (B.1),  $W$  can be rewritten as

$$W = d\theta^a \wedge \frac{1}{2}\omega^{bc} \wedge \eta_{abc} - \Theta^a \wedge \frac{1}{2}\omega^{bc} \wedge \eta_{abc}. \quad (\text{A.3})$$

The conjugate form of  $\theta^a$  is given by (4.37). The Hamilton form is given by (4.38) with

$$H_G(\theta, \pi) = \frac{W}{2\kappa}. \quad (\text{A.4})$$

Here, we used (A.1) and (A.3).

The canonical equations are given by

$$\begin{aligned} d\theta^a &= \frac{\partial H_G}{\partial \pi_a} - \frac{\partial L_{\text{mat}}}{\partial \pi_a} \\ &= \frac{\partial H_G}{\partial \pi_a} + \Theta^a, \end{aligned} \quad (\text{A.5})$$

$$d\pi_a = \frac{\partial H_G}{\partial \theta^a} - \frac{\partial L_{\text{mat}}(\theta, \pi)}{\partial \theta^a}. \quad (\text{A.6})$$

Here, we used (4.40).

To calculate  $\partial H_G/\partial \pi_a$ ,  $\partial H_G/\partial \theta^a$ , we represent  $W$  by  $\theta^a$  and  $\pi_a$ . Using (B.2),  $W$  can be rewritten as

$$W = (\omega_{abc}\omega^{bca} + \omega_a\omega^a)\eta. \quad (\text{A.7})$$

Here, we expand  $\omega_{ab}$  as  $\omega_{ab} = \omega_{abc}\theta^c$  and put  $\omega_a := \omega^b_{ab}$ . We can represent  $\omega_{abc}$  by  $\theta^a$  and  $\pi_a$  as

$$\omega_{abc} = \kappa \left[ v_{c,ab} + \frac{1}{d-2} (\mathring{g}_{ac}v_b - \mathring{g}_{bc}v_a) \right], \quad (\text{A.8})$$

$$v_{c,ab} := - * V_{c,ab}, \quad V_{c,ab} := \pi_c \wedge \theta_a \wedge \theta_b \quad (\text{A.9})$$

with  $v_a := v^b_{ab}$ .

Next, we derive the canonical equations. For an arbitrary  $d$ -form  $\xi$ ,

$$\begin{aligned} \delta v_{c,ab}\xi &= -\delta v_{c,ab}\xi * \eta = -\delta v_{c,ab}\eta * \xi = (-\delta[v_{c,ab}\eta] + v_{c,ab}\delta\eta) * \xi \\ &= (-\delta V_{c,ab} + v_{c,ab}\delta\eta) * \xi \end{aligned} \quad (\text{A.10})$$

holds [6]. Then, we have

$$\delta v_{c,ab}\eta = \delta V_{c,ab} - v_{c,ab}\delta\eta. \quad (\text{A.11})$$

Using this, we have

$$\frac{\partial H_G}{\partial \pi_c} = -\omega^c_a \wedge \theta^a, \quad (\text{A.12})$$

$$\frac{\partial H_G}{\partial \theta^d} = \frac{1}{2\kappa}(\omega^c_d \wedge \omega^{ab} \wedge \eta_{abc} + \omega^a_c \wedge \omega^{bc} \wedge \eta_{bad}). \quad (\text{A.13})$$

Substituting (A.12) into (A.5), we have

$$d\theta^a = -\omega^a_b \wedge \theta^b + \Theta^a, \quad (\text{A.14})$$

which is equivalent to the first structure equation (4.1). By the way, using (4.20), we can show that

$$-\frac{\partial L_{\text{mat}}(\theta, \pi)}{\partial \theta^c} = -T_c - \frac{1}{2\kappa}\omega^{ab} \wedge \Theta^d \wedge \eta_{abcd}. \quad (\text{A.15})$$

Substituting (A.13) and (A.15) into (A.6), we have

$$d\pi_c = \frac{1}{2\kappa}(\omega^d_b \wedge \omega^{ab} \wedge \eta_{adc} + \omega^d_c \wedge \omega^{ab} \wedge \eta_{abd} - \omega^{ab} \wedge \Theta^d \wedge \eta_{abcd}) - T_c. \quad (\text{A.16})$$

## B Formulas

Several useful formulas are listed. For  $\theta^b \wedge \eta_{a_1 \dots a_r}$  ( $r = 1, 2, 3, 4$ ),

$$\theta^b \wedge \eta_{a_1 \dots a_r} = (-1)^{r-1} r \delta_{[a_1}^b \eta_{a_2 \dots a_r]} \quad (\text{B.1})$$

hold [16]. Using this, we have

$$\theta^a \wedge \theta^b \wedge \eta_{cd} = (\delta_c^a \delta_d^b - \delta_d^a \delta_c^b) \eta. \quad (\text{B.2})$$

For  $\delta \eta_{a_1 \dots a_r}$  ( $r = 0, 1, 2, 3$ ),

$$\delta \eta_{a_1 \dots a_r} = \delta \theta^b \wedge \eta_{a_1 \dots a_r b} \quad (\text{B.3})$$

hold. For  $d\eta_{a_1 \dots a_r}$  ( $r = 1, 2, 3$ ),

$$\begin{aligned} d\eta_{a_1 \dots a_r} &= (r+1) A_{[a_1}^b \wedge \eta_{b a_2 \dots a_r]} \\ &= (r+1) \omega_{[a_1}^b \wedge \eta_{b a_2 \dots a_r]} + \Theta^b \wedge \eta_{a_1 \dots a_r b} \end{aligned} \quad (\text{B.4})$$

hold.  $\eta_{a_1 \dots a_r}$  ( $r = 1, 2, 3, 4$ ) can be written as [16]

$$\eta_{a_1 \dots a_r} = e_{a_r} \rfloor \eta_{a_1 \dots a_{r-1}}. \quad (\text{B.5})$$

Here,  $\rfloor$  is the interior product and  $\{e_a\}$  is the dual basis of  $\{\theta^a\}$  ( $e_a \rfloor \theta^b = \delta_a^b$ ).

## References

- [1] A. D’Adda, J. E. Nelson and T. Regge, “Covariant canonical formalism for the group manifold”, *Annals of Physics* **165**, 384 (1985).
- [2] J. E. Nelson and T. Regge, “Covariant Canonical Formalism for Gravity”, *Annals of Physics* **166**, 234 (1986).
- [3] A. Lerda, J. E. Nelson and T. Regge, “Covariant Canonical Formalism for Supergravity”, *Phys. Lett.* **161B**, 294 (1985).
- [4] A. Lerda, J. E. Nelson and T. Regge, “The Group Manifold Hamiltonian for Supergravity”, *Phys. Lett.* **161B**, 297 (1985).
- [5] J. M. Nester, “General pseudotensors and quasilocal quantities”, *Classical and Quantum Gravity* **21**, S261 (2004).
- [6] Y. Kaminaga, “Covariant Analytic Mechanics with Differential Forms and Its Application to Gravity”, *Electron. J. Theor. Phys.* **9**, 199 (2012).
- [7] S. Nakajima, “Application of covariant analytic mechanics with differential forms to gravity with Dirac field”, *Electron. J. Theor. Phys.* **13**, 95 (2016).
- [8] S. Nakajima, “Reconsideration of De Donder-Weyl theory by covariant analytic mechanics”, arXiv:1602.04849v2.
- [9] Y. Kaminaga, “Poisson Bracket and Symplectic Structure of Covariant Canonical Formalism of Fields”, *Electron. J. Theor. Phys.* **14**, 55 (2018).
- [10] L. Castellani and A. D’Adda, “Covariant Hamiltonian for gravity coupled to  $p$ -forms”, *Phys. Rev. D* **101**, 025015 (2020).
- [11] L. Castellani, “Covariant hamiltonian for supergravity in  $d = 3$  and  $d = 4$ ”, *J. High Energy Phys.* **2020**, 69 (2020).
- [12] I. V. Kanatchikov, “Canonical structure of classical field theory in the polymomentum phase space”, *Rep. Math. Phys.* **41**, 49 (1998).
- [13] F. Gieres, “Covariant canonical formulations of classical field theories”, arXiv:2109.07330v2
- [14] A. Trautman, “On the Einstein-Cartan equations”, *Bull. Acad. Pol. Sci* **20**, 185 (1972).
- [15] W. Thirring, “A Course in Mathematical Physics 2”, Springer (second edition, 1978).
- [16] F. W. Hehl, J. D. McCrea, E. W. Mielke and Y. Ne’eman, “Metric-affine gauge theory of gravity: field equations, Noether identities, world spinors, and breaking of dilation invariance”, *Phys. Rep.* **258**, 1 (1995).