

EVERY COMMUTATIVE JB*-TRIPLE SATISFIES THE COMPLEX MAZUR–ULAM PROPERTY

DAVID CABEZAS, MARÍA CUETO-AVELLANEDA, DAISUKE HIROTA, TAKESHI MIURA,
AND ANTONIO M. PERALTA

ABSTRACT. We prove that every commutative JB*-triple satisfies the complex Mazur–Ulam property. Thanks to the representation theory, we can identify commutative JB*-triples as spaces of complex-valued continuous functions on a principal \mathbb{T} -bundle L in the form

$$C_0^{\mathbb{T}}(L) := \{a \in C_0(L) : a(\lambda t) = \lambda a(t) \text{ for every } (\lambda, t) \in \mathbb{T} \times L\}.$$

We prove that every surjective isometry from the unit sphere of $C_0^{\mathbb{T}}(L)$ onto the unit sphere of any complex Banach space admits an extension to a surjective real linear isometry between the spaces.

1. INTRODUCTION

New recent advances continue improving our understanding of Tingley’s problem by enlarging the list of positive solutions, and the range of spaces satisfying the Mazur–Ulam property. As introduced in [7], a Banach space X satisfies the Mazur–Ulam property if every surjective isometry from its unit sphere onto the unit sphere of any other Banach space admits an extension to a surjective real linear isometry between the spaces. The new advances are struggling to provide new tools and techniques to attack this intriguing open question on the possibility of extending a surjective isometry between the unit spheres of two Banach spaces to a surjective real linear isometry between the spaces. A remarkable outstanding advance has been obtained by T. Banach in [2], who proved that every 2-dimensional Banach space X satisfies the *Mazur–Ulam property*, that is, every surjective isometry from the unit sphere of X onto the unit sphere of any other Banach space extends to a real linear isometry between the involved Banach spaces. This is in fact the culminating point of deep technical advances (see [1, 6, 3]).

The abundance of unitary elements in unital C^* -algebras, real von Neumann algebras and JBW*-algebras is a key property to prove that these spaces together with all JBW*-triples satisfy the Mazur–Ulam property (cf. [16, 4, 13]). A prototypical example of non-unital C^* -algebra is given by the C^* -algebra $K(H)$, of all compact operators on an infinite dimensional complex Hilbert space H , or more generally, by a compact C^* -algebra (i.e. a c_0 -sum of $K(H)$ -spaces). Compact C^* -algebras and weakly compact JBW*-triples are in the list of complex Banach spaces satisfying the Mazur–Ulam property (see [18]).

Another research line is exploring Tingley’s problem in the case of certain function algebras and spaces. The first positive solution to Tingley’s problem for a Banach space consisting of analytic functions, apart from Hilbert spaces, was obtained by O. Hatori, S. Oi and R. Shindo Togashi in [12], where it is proved that each surjective isometry between the unit spheres of two uniform

2010 *Mathematics Subject Classification.* 46J10, 46B04, 46B20, 46J15, 47B49, 17C65.

Key words and phrases. extension of isometries, Tingley’s problem, abelian JB*-triple.

algebras (i.e. closed subalgebras of $C(K)$ containing the constants and separating the points of K) can be always extended to a surjective real-linear isometry between the uniform algebras. O. Hatori has gone further by showing that every uniform algebra satisfies the complex Mazur–Ulam property [11, Theorem 4.5]. The “*complex Mazur–Ulam property*” has been coined by O. Hatori to denote those complex Banach spaces for which every surjective isometry from its unit sphere onto the unit sphere of a complex Banach spaces admits an extension to a real linear mapping between the spaces.

The non-unital analogue of uniform algebras is materialized in the notion of uniformly closed function algebra on a locally compact Hausdorff space L . We recently showed that each surjective isometry between the unit spheres of two uniformly closed function algebras on locally compact Hausdorff spaces admits an extension to a surjective real linear isometry between these algebras (see [8]). In the just quoted reference we also proved that Tingley’s problem admits a positive solution for any surjective isometry between the unit spheres of two commutative JB^* -triples, which are not, in general, subalgebras of the algebra $C_0(L)$ of all complex-valued continuous functions on L vanishing at infinity (see Section 3 for the detailed representation of commutative JB^* -triples). In this note we shall employ a recent tool developed by O. Hatori in [11] to infer that a stronger conclusion holds, namely, every commutative JB^* -triple satisfies the complex Mazur–Ulam property. Among the consequences we derive that every commutative C^* -algebra enjoys the complex Mazur–Ulam property. Commutative unital C^* -algebras satisfy the stronger Mazur–Ulam property by a result due to M. Mori and N. Ozawa [16]. Some other previous achievements in this line were obtained by R. Liu for $C(K, \mathbb{R})$ [15, Corollary 6], and by O. Hatori for closed subspaces of $C_0(L, \mathbb{R})$ separating the points of L and satisfying a topological-geometric hypothesis called condition (r) (see [11, Theorem 5.4] in the preprint version).

2. PRELIMINARIES

We shall briefly recall some basic terminology to understand the sufficient condition, established by O. Hatori in [11, Proposition 4.4], to guarantee that a Banach space satisfies the complex Mazur–Ulam property.

Let X be a real or complex Banach space, and let X^* , $S(X)$ and \mathcal{B}_X denote the dual space, the unit sphere and the closed unit ball of X , respectively. It is known, thanks to Hahn–Banach theorem or Eidelheit’s separation theorem, that maximal convex subsets of $S(X)$ and maximal proper norm closed faces of \mathcal{B}_X define the same subsets (cf. [21, Lemma 3.3] or [22, Lemma 3.2]). The set of all maximal convex subsets of $S(X)$, equivalently, all maximal proper norm closed faces of \mathcal{B}_X , will be denoted by \mathfrak{F}_X . For each $F \in \mathfrak{F}_X$ there exists an extreme point φ of the closed unit ball \mathcal{B}_{X^*} such that $F = \varphi^{-1}\{1\} \cap S(X)$ (cf. [21, Lemma 3.3]). The set of all extreme points φ of \mathcal{B}_{X^*} for which $\varphi^{-1}\{1\} \cap S(X)$ is a maximal convex subset of $S(X)$ will be denoted by \mathcal{Q}_X . On the latter set we consider the equivalence relation defined by

$$\varphi \sim \phi \Leftrightarrow \exists \gamma \in \mathbb{T} = S(\mathbb{K}) \text{ with } \varphi^{-1}\{1\} \cap S(X) = (\gamma\phi)^{-1}\{1\} \cap S(X),$$

where $\mathbb{K} = \mathbb{R}$ if X is a real Banach space and $\mathbb{K} = \mathbb{C}$ if X is a complex Banach space. A *set of representatives* for the quotient set \mathcal{Q}_X / \sim (or for \mathfrak{F}_X) will consist in a subset \mathbf{P}_X of \mathcal{Q}_X which is formed by precisely one, and only one, element in each equivalent class of \mathcal{Q}_X / \sim . According to the just commented notation and results, for each $F \in \mathfrak{F}_X$ there exists a unique $\varphi \in \mathbf{P}_X$ and $\gamma \in \mathbb{T}$ such that $F = F_{\varphi, \gamma} := (\gamma\varphi)^{-1}\{1\} \cap S(X)$ (cf. [11, Lemma 2.5]). That is, the elements in \mathfrak{F}_X are bijectively labelled by the set $\mathbf{P}_X \times \mathbb{T}$, and we can define a bijection $\mathcal{I}_X : \mathfrak{F}_X \rightarrow \mathbf{P}_X \times \mathbb{T}$ labelling the set \mathfrak{F}_X .

For example, by the classical description of the extreme points of the closed unit ball of the dual of a $C(K)$ space as those functionals of the form $\lambda\delta_t(f) = \lambda f(t)$ ($f \in C(K)$) with $t \in K$, $\lambda \in \mathbb{T}$, the set $\mathbf{P}_{C(K)} = \{\delta_t : t \in K\}$ is a set of representatives for $\mathfrak{F}_{C(K)}$. It is shown in [11, Example 2.4] that for a uniform algebra A over a compact Hausdorff space K , the set $\{\delta_t : t \in \text{Ch}(A)\}$ is a set of representatives for A , where $\text{Ch}(A)$ denotes the Choquet boundary of A .

Let A, B be non-empty closed subsets of a metric space (E, d) . The usual Hausdorff distance between A and B is defined by

$$d_H(A, B) = \max\{\sup_{a \in A} d(a, B), \sup_{b \in B} d(b, A)\}.$$

We shall employ this Hausdorff distance to measure distances between elements in \mathfrak{F}_X .

According to [11], a Banach space X satisfies the *condition of the Hausdorff distance* if the elements in \mathfrak{F}_X satisfy the following rules:

$$(2.1) \quad d_H(F_{\varphi, \lambda}, F_{\varphi', \lambda'}) = \begin{cases} |\lambda - \gamma\lambda'|, & \text{if } \varphi^{-1}\{1\} \cap S(X) = (\gamma\varphi')^{-1}\{1\} \cap S(X), \\ 2, & \text{if } \varphi \not\sim \varphi'. \end{cases}$$

for $\varphi, \varphi' \in \mathcal{Q}_X$ and $\lambda, \lambda' \in \mathbb{T}$. Let $\mathbf{P}_X \subset \mathcal{Q}_X$ be a set of representatives for \mathfrak{F}_X . Having in mind the properties of the mapping \mathcal{I}_X^{-1} for \mathbf{P}_X , the condition of the Hausdorff distance in (2.1) can be rewritten in the form

$$(2.2) \quad d_H(F_{\varphi, \lambda}, F_{\varphi', \lambda'}) = \begin{cases} |\lambda - \lambda'|, & \text{if } \varphi = \varphi', \\ 2, & \text{if } \varphi \neq \varphi', \end{cases}$$

for $\varphi, \varphi' \in \mathbf{P}_X$ and $\lambda, \lambda' \in \mathbb{T}$. Under the light of [11, Lemma 3.1] to conclude that a complex Banach space X together with a set of representatives \mathbf{P}_X satisfies (2.2) it suffices to prove that

$$(2.3) \quad F_{\varphi, \lambda} \cap F_{\varphi', \lambda'} \neq \emptyset \text{ for any } \varphi \neq \varphi' \text{ in } \mathbf{P}_X, \lambda, \lambda' \text{ in } \mathbb{T}.$$

Let us go back to the set \mathcal{Q}_X determining the set \mathfrak{F}_X of all maximal proper norm closed faces of \mathcal{B}_X . For $\varphi \in \mathcal{Q}_X$ and $\alpha \in \mathbb{D} = \mathcal{B}_{\mathbb{K}}$ ($\mathbb{K} = \mathbb{R}$ or \mathbb{C}), we set

$$M_{\varphi, \alpha} = \left\{ x \in S(X) : d\left(x, F_{\varphi, \frac{\alpha}{|\alpha|}}\right) \leq 1 - |\alpha|, d\left(x, F_{\varphi, -\frac{\alpha}{|\alpha|}}\right) \leq 1 + |\alpha| \right\},$$

where $\frac{\alpha}{|\alpha|} = 1$ if $\alpha = 0$. Note that in this definition the inequalities “ \leq ” can be replaced with equalities. It is known that for each φ in a set of representatives \mathbf{P}_X , the inclusion

$$M_{\varphi, \alpha} \subseteq \{x \in S(X) : \varphi(x) = \alpha\}$$

holds for all $\alpha \in \mathbb{D}$ (cf. [11, Lemma 4.3]). The equality of these two sets is the second condition required in the following result, which can be applied to guarantee that a Banach space enjoys the Mazur–Ulam property [11].

Proposition 2.1. [11, Proposition 4.4] *Let X be a complex Banach space. Suppose that \mathbf{P}_X is a set of representatives for \mathfrak{F}_X . Assume that X satisfies the following two hypotheses:*

- (i) X satisfies the condition of the Hausdorff distance,
- (ii) For each φ in \mathbf{P}_X and $\alpha \in \mathbb{D}$ we have

$$M_{\varphi, \alpha} = \{x \in S(X) : \varphi(x) = \alpha\}.$$

Then X satisfies the complex Mazur–Ulam property.

3. MAZUR-ULAM PROPERTY FOR COMMUTATIVE JB^* -TRIPLES

The aim of this section is to show that every abelian JB^* -triple satisfies the Mazur–Ulam property. We shall avoid the axiomatic definition of these objects and we shall simply recall their representation as function spaces. It follows from the Gelfand theory for JB^* -triples (see [14, Corollary 1.11]) that each abelian JB^* -triple can be identified with the norm closed subspace of $C_0(L)$ defined by

$$C_0^{\mathbb{T}}(L) := \{a \in C_0(L) : a(\lambda t) = \lambda a(t) \text{ for every } (\lambda, t) \in \mathbb{T} \times L\},$$

where L is a principal \mathbb{T} -bundle L , that is, a subset of a Hausdorff locally convex complex space such that $0 \notin L$, $L \cup \{0\}$ is compact, and $\mathbb{T}L = L$ (see also [8]).

We can state next the main result of the paper.

Theorem 3.1. *Let L be a principal \mathbb{T} -bundle. Then, $C_0^{\mathbb{T}}(L)$ satisfies the complex Mazur–Ulam property, that is, for each complex Banach space X , every surjective isometry $\Delta : S(C_0^{\mathbb{T}}(L)) \rightarrow S(X)$ admits an extension to a surjective real linear isometry $T : C_0^{\mathbb{T}}(L) \rightarrow X$.*

The proof will be obtained after a series of technical results witnessing that every function space of the form $C_0^{\mathbb{T}}(L)$ satisfies the hypotheses of Proposition 2.1. Before dealing with the technical details, we shall present an interesting corollary.

Although for each locally compact space \tilde{L} , the Banach space $C_0(\tilde{L})$ is isometrically isomorphic to a $C_0^{\mathbb{T}}(L)$ space (cf. [17, Proposition 10]), there exist principal \mathbb{T} -bundles L for which the space $C_0^{\mathbb{T}}(L)$ is not isometrically isomorphic to a $C_0(L)$ space (cf. [14, Corollary 1.13 and subsequent comments]). So, there exist abelian JB^* -triples which are not isometrically isomorphic to commutative C^* -algebras. The next corollary is a weaker consequence of our previous theorem.

Corollary 3.2. *Every abelian C^* -algebra (that is, every $C_0(\tilde{L})$ space) satisfies the complex Mazur–Ulam property.*

Compared with previous results, we observe that as a consequence of the result proved by O. Hatori for uniform algebras in [11, Theorem 4.5] every unital abelian C^* -algebra satisfies the complex Mazur–Ulam property. Actually, all unital C^* -algebras enjoy the Mazur–Ulam property [16]. In the case of real-valued continuous functions, R. Liu proved that for each compact Hausdorff space K , $C(K, \mathbb{R})$ satisfies the Mazur–Ulam property (see [15, Corollary 6]). Previous attempts were conducted by G. Ding [9] and X.N. Fang and J.H. Wang [10].

Let \tilde{L} be a locally compact Hausdorff space. A closed subspace E of $C_0(\tilde{L}, \mathbb{K})$, separates the points of \tilde{L} if for any $t_1 \neq t_2$ in \tilde{L} there exists a function $a \in E$ such that $a(t_1) \neq a(t_2)$. Following [11], we shall say that E satisfies the condition (r) if for any t in the Choquet boundary of E , each neighborhood V of t , and $\varepsilon > 0$ there exists $u \in E$ such that $0 \leq u \leq 1 = u(t)$ on \tilde{L} and $0 \leq u \leq \varepsilon$ on $\tilde{L} \setminus V$. The proof of Corollary 5.4 in the preprint version of [11] affirms that each closed subspace E of $C_0(\tilde{L}, \mathbb{R})$ separating the points of \tilde{L} and satisfying a stronger assumption than condition (r) has the Mazur–Ulam property. After some private communications with O. Hatori we actually learned that property (r) is enough to conclude that any such closed subspace E satisfies the Mazur–Ulam property. Actually the desired conclusion can be derived from [5, Theorem 2.4] by just observing that condition (r) implies that the isometric identification of E in $C_0(\overline{\text{Ch}(E)})$ is C -rich, and hence a lush space. Corollary 3.9 in [20] implies that E has the Mazur–Ulam property.

We focus now on the main goal of this section. Henceforth, let L be a principal \mathbb{T} -bundle and $L_0 \subset L$ a maximal non-overlapping set, that is, L_0 is maximal satisfying that for each $t \in L_0$ we have $L_0 \cap \mathbb{T}t = \{t\}$ (its existence is guaranteed by Zorn’s lemma).

Assume that a Banach space Y satisfies the following property: for every extreme point $\varphi \in \partial_e(\mathcal{B}_{Y^*})$, the set $\{\varphi\}$ is a weak*-semi-exposed face of \mathcal{B}_{Y^*} . It is clear that each extreme point $\varphi \in \partial_e(\mathcal{B}_{Y^*})$ is determined by the set $\{\varphi\}_r = \varphi^{-1}(1) \cap S(Y)$. Hence, the equivalence relation \sim defined in Section 2 (cf. [11, Definition 2.1]) can be characterized in the following terms: for $\varphi, \psi \in \partial_e(\mathcal{B}_{Y^*})$, we have

$$\varphi \sim \psi \iff \varphi = \gamma\psi \text{ for some } \gamma \in \mathbb{T}.$$

Since $Y = C_0^{\mathbb{T}}(L)$ satisfies the mentioned property, the set $\{\delta_t : t \in L_0\}$ is a set of representatives for the relation \sim . We know that each maximal proper face F of the closed unit ball of $C_0^{\mathbb{T}}(L)$ is of the form

$$F = F_{\delta_{t_0}, \lambda} = F_{t_0, \lambda} := \{a \in S(C_0^{\mathbb{T}}(L)) : \delta_{t_0}(a) = a(t_0) = \lambda\}$$

for some $(t_0, \lambda) \in L_0 \times \mathbb{T}$ (cf. [8, Lemma 3.5]).

The next remark, which has been borrowed from [8, Remark 3.4] states a kind of Urysohn's lemma for the space $C_0^{\mathbb{T}}(L)$.

Remark 3.3. [8, Remark 3.4] Suppose L is a principal \mathbb{T} -bundle. Let W be a \mathbb{T} -invariant open neighbourhood of t_0 in L which is contained in a compact \mathbb{T} -invariant subset. Then, there exists a function $h \in S(C_0^{\mathbb{T}}(L))$ satisfying $h(t_0) = 1$ and $h(t) = 0$ for all $t \in X \setminus W$.

We recall next, for later purposes, the following version of [19, Theorem 2.7] for \mathbb{T} -symmetric compact subsets of principal \mathbb{T} -bundles.

Remark 3.4. Let L be a principal \mathbb{T} -bundle. Let K be a \mathbb{T} -symmetric compact subset of L contained in an open \mathbb{T} -symmetric set $U \subset L$. Then, there exists an open \mathbb{T} -symmetric set V with compact \mathbb{T} -symmetric closure such that

$$K \subset V \subset \overline{V} \subset U.$$

The proof follows straightforwardly from [19, Theorem 2.7] by just observing that for each open (respectively, closed or compact) subset $\mathcal{O} \subset L$, the set $\mathbb{T}\mathcal{O}$ is open (respectively, closed or compact).

We can now begin with the technical details for our arguments.

Lemma 3.5. *If $t_1 \neq t_2$ in L_0 , then there exist open \mathbb{T} -symmetric subsets $V_1, V_2 \subset L$ satisfying:*

- $\mathbb{T}t_j \subset V_j$ for $j = 1, 2$;
- $\overline{V_j}$ is compact for $j = 1, 2$;
- $V_1 \cap V_2 = \overline{V_1} \cap \overline{V_2} = \emptyset$.

Proof. Since L_0 is non-overlapping, we know that $\mathbb{T}t_1$ and $\mathbb{T}t_2$ are disjoint compact subsets of L . Hence

$$\mathbb{T}t_1 \subset L \setminus \mathbb{T}t_2,$$

where $L \setminus \mathbb{T}t_2$ is \mathbb{T} -symmetric and open. By Remark 3.4, there exists a \mathbb{T} -symmetric open set $V_1 \subset L$ with \mathbb{T} -symmetric compact closure satisfying

$$\mathbb{T}t_1 \subset V_1 \subset \overline{V_1} \subset L \setminus \mathbb{T}t_2.$$

Now, having in mind that $\overline{V_1}$ is \mathbb{T} -symmetric and compact with $\overline{V_1} \cap \mathbb{T}t_2 = \emptyset$, we deduce that $L \setminus \overline{V_1}$ is an open \mathbb{T} -symmetric set containing $\mathbb{T}t_2$. We can find, again via Remark 3.4, an open \mathbb{T} -symmetric subset V_2 with \mathbb{T} -symmetric compact closure satisfying

$$\mathbb{T}t_2 \subset V_2 \subset \overline{V_2} \subset L \setminus \overline{V_1}.$$

□

Corollary 3.6. *Let $t_1 \neq t_2$ in L_0 and $\lambda_1, \lambda_2 \in \mathbb{T}$. Then, there exist a function $a \in S(C_0^\mathbb{T}(L))$ such that $a(t_j) = \lambda_j$ for $j = 1, 2$.*

Proof. Lemma 3.5 assures the existence of disjoint open \mathbb{T} -symmetric neighbourhoods with compact closure W_1 and W_2 of t_1 and t_2 , respectively. By Remark 3.3, there exist functions $a_1, a_2 \in S(C_0^\mathbb{T}(L))$ such that $a_j(t_j) = 1$ and $a_j|_{L \setminus W_j} \equiv 0$ for $j = 1, 2$.

Since W_1 and W_2 are disjoint, we have $0 \in \{a_1(t), a_2(t)\}$ for each $t \in L$. Hence, $a := \lambda_1 a_1 + \lambda_2 a_2$ lies in $S(C_0^\mathbb{T}(L))$ and satisfies the required conclusion. \square

Remark 3.7. The previous corollary shows that $F_{t_1, \lambda} \cap F_{t_2, \lambda'} \neq \emptyset$ for any $t_1 \neq t_2$ in L_0 and $\lambda, \lambda' \in \mathbb{T}$. Therefore, the space $C_0^\mathbb{T}(L)$ satisfies (2.3) and hence the condition of the Hausdorff distance (cf. (2.1) and (2.3) or [11, Lemma 3.1]).

We shall next show that $C_0^\mathbb{T}(L)$ satisfies the second hypothesis in Proposition 2.1.

Along this note the set $\mathcal{B}_\mathbb{C} \setminus \{0\}$ will be regarded as a principal \mathbb{T} -bundle, and we shall write $C_0^\mathbb{T}(\mathcal{B}_\mathbb{C})$ for the abelian JB*-triple associated with this principal \mathbb{T} -bundle, that is,

$$C_0^\mathbb{T}(\mathcal{B}_\mathbb{C}) = \{f \in C_0(\mathcal{B}_\mathbb{C} \setminus \{0\}) : f(0) = 0, f(\lambda t) = \lambda f(t), \forall t \in \mathcal{B}_\mathbb{C}, \lambda \in \mathbb{T}\}.$$

Lemma 3.8. *For each non-zero $a \in \mathcal{B}_{C_0^\mathbb{T}(L)}$ and each $0 < \varepsilon < \|a\| \leq 1$, there exists a function $u_\varepsilon \in S(C_0^\mathbb{T}(L))$ satisfying*

$$u_\varepsilon(s) = \frac{a(s)}{|a(s)|}$$

for every $s \in L$ with $|a(s)| \geq \varepsilon$.

Proof. Let us observe that each function $f \in C_0^\mathbb{T}(\mathcal{B}_\mathbb{C})$ is uniquely determined by its values on $[0, 1]$. Consider the function $h_\varepsilon \in C_0^\mathbb{T}(\mathcal{B}_\mathbb{C})$ defined on $[0, 1]$ by

$$h_\varepsilon(t) = \begin{cases} 0, & t \leq \varepsilon/2, \\ \text{affine}, & \varepsilon/2 < t < \varepsilon, \\ 1, & t \geq \varepsilon. \end{cases}$$

Since $\|a\| > \varepsilon$, the function $(h_\varepsilon)_t(a) = h \circ a$ lies in $S(C_0^\mathbb{T}(L))$.

We also know that for each $s \in L$ such that $|a(s)| \geq \varepsilon$ we have

$$u_\varepsilon(s) = h_\varepsilon(a(s)) = h_\varepsilon\left(\frac{a(s)}{|a(s)|}|a(s)|\right) = \frac{a(s)}{|a(s)|} h_\varepsilon(|a(s)|) = \frac{a(s)}{|a(s)|} \cdot 1,$$

which concludes the proof. \square

The next technical result is the key tool for our purposes.

Lemma 3.9. *Let $t_0 \in L$. Suppose that $a \in S(C_0^\mathbb{T}(L))$ satisfies $0 < |a(t_0)| < 1$. Then, for each $\varepsilon > 0$ with $\varepsilon < \min\{|a(t_0)|, 1 - |a(t_0)|\}$, there exist $a_\varepsilon, b_\varepsilon \in S(C_0^\mathbb{T}(L))$ such that $\|a - a_\varepsilon\| \leq \frac{\varepsilon}{2}$, and for each $0 < r < 1$ the function*

$$c_{r, \varepsilon} := r a_\varepsilon - (1 + r|a(t_0)|) b_\varepsilon$$

lies in $F_{t_0, -\frac{a(t_0)}{|a(t_0)|}}$ with $\|a_\varepsilon - c_{r, \varepsilon}\| \leq 2 - r + r|a(t_0)|$.

Proof. To simplify the notation we shall write $\alpha = a(t_0)$. Consider the function $g_\varepsilon \in C_0^\mathbb{T}(\mathcal{B}_\mathbb{C})$ defined on $[0, 1]$ by

$$g_\varepsilon(t) = \begin{cases} t, & 0 \leq t \leq |\alpha| - \varepsilon, \\ \text{affine}, & |\alpha| - \varepsilon < t < |\alpha| - \varepsilon/2, \\ |\alpha|, & |\alpha| - \varepsilon/2 \leq t \leq |\alpha| + \varepsilon/2, \\ \text{affine}, & |\alpha| + \varepsilon/2 < t < |\alpha| + \varepsilon, \\ t, & |\alpha| + \varepsilon \leq t \leq 1. \end{cases}$$

By setting $a_\varepsilon = (g_\varepsilon)_t(a) = g_\varepsilon \circ a \in C_0^\mathbb{T}(L)$, we have $\|a_\varepsilon\| = 1$ and $a_\varepsilon(t_0) = \alpha$.

Let $\iota : \mathcal{B}_\mathbb{C} \hookrightarrow \mathbb{C}$ denote the inclusion mapping –note that $\iota \in C_0^\mathbb{T}(\mathcal{B}_\mathbb{C})$. Clearly, $\|g_\varepsilon - \iota\| = \varepsilon/2$, thus

$$\|a - a_\varepsilon\| = \|(g_\varepsilon)_t(a) - \iota_t(a)\| \leq \varepsilon/2.$$

Let us find, by the continuity of a , an open neighborhood $\mathcal{O}_\varepsilon \subset L$ of t_0 satisfying $|a(s) - a(t_0)| < \varepsilon/2$, for all $s \in \mathcal{O}_\varepsilon$. The set $W_\varepsilon := \mathbb{T}\mathcal{O}_\varepsilon$ is open and \mathbb{T} -symmetric. For each $s \in W_\varepsilon$, we have $s = \lambda_s t_s$ for some $\lambda_s \in \mathbb{T}$ and $t_s \in \mathcal{O}_\varepsilon$, hence

$$\| |a(s)| - |a(t_0)| \| = \| |a(\lambda_s t_s)| - |a(t_0)| \| = \| |a(t_s)| - |a(t_0)| \| \leq |a(t_s) - a(t_0)| < \varepsilon/2,$$

which leads to

$$(3.1) \quad |\alpha| - \varepsilon/2 = |a(t_0)| - \varepsilon/2 < |a(s)| < |a(t_0)| + \varepsilon/2 = |\alpha| + \varepsilon/2$$

for all $s \in W_\varepsilon$. Consequently,

$$(3.2) \quad a_\varepsilon(s) = g_\varepsilon(a(s)) = \frac{a(s)}{|a(s)|} g_\varepsilon(|a(s)|) = \frac{a(s)}{|a(s)|} |\alpha|$$

for all $s \in W_\varepsilon$.

Since W_ε is open, \mathbb{T} -symmetric and contains t_0 , Remark 3.4 assures the existence of an open and \mathbb{T} -symmetric set V_ε with compact \mathbb{T} -symmetric closure satisfying

$$(3.3) \quad \mathbb{T}t_0 \subset V_\varepsilon \subset \overline{V_\varepsilon} \subset W_\varepsilon.$$

By Lemma 3.8, there exists $u_\varepsilon \in S(C_0^\mathbb{T}(L))$ satisfying

$$(3.4) \quad u_\varepsilon(s) = \frac{a_\varepsilon(s)}{|a_\varepsilon(s)|}, \quad \text{for all } s \in L \text{ with } |a_\varepsilon(s)| \geq |\alpha| - \varepsilon.$$

Since $|a_\varepsilon(t_0)| = |\alpha| > |\alpha| - \varepsilon$, we have $u_\varepsilon(t_0) = \frac{\alpha}{|\alpha|}$.

Fix, via Remark 3.3, a function $f_\varepsilon \in S(C_0^\mathbb{T}(L))$ such that $f_\varepsilon(t_0) = 1$ and $f_\varepsilon|_{L \setminus V_\varepsilon} \equiv 0$.

For each $r \in]0, 1[$, let us define the function

$$c_{r,\varepsilon} := r a_\varepsilon - (1 + r|\alpha|) |f_\varepsilon| u_\varepsilon \in C_0^\mathbb{T}(L).$$

Clearly, we have

$$(3.5) \quad c_{r,\varepsilon}(t_0) = r a_\varepsilon(t_0) - (1 + r|\alpha|) |f_\varepsilon(t_0)| u_\varepsilon(t_0) = r\alpha - (1 + r|\alpha|) \frac{\alpha}{|\alpha|} = -\frac{\alpha}{|\alpha|}.$$

Given $s \in L \setminus V_\varepsilon$, we have $f_\varepsilon(s) = 0$. Hence, $c_{r,\varepsilon}(s) = r a_\varepsilon(s)$, so $|c_{r,\varepsilon}(s)| \leq r$. If $s \in V_\varepsilon \subset W_\varepsilon$, by (3.2) we have

$$a_\varepsilon(s) = \frac{a(s)}{|a(s)|} |\alpha|.$$

In particular, $|a_\varepsilon(s)| = |\alpha| > |\alpha| - \varepsilon$, and (3.4) gives

$$u_\varepsilon(s) = \frac{a_\varepsilon(s)}{|a_\varepsilon(s)|} = \frac{|\alpha| \frac{a(s)}{|a(s)|}}{|\alpha|} = \frac{a(s)}{|a(s)|}.$$

Therefore,

$$\begin{aligned} c_{r,\varepsilon}(s) &= r a_\varepsilon(s) - (1 + r|\alpha|) |f_\varepsilon(s)| u_\varepsilon(s) = r|\alpha| \frac{a(s)}{|a(s)|} - (1 + r|\alpha|) \frac{a(s)}{|a(s)|} |f_\varepsilon(s)| \\ &= \frac{a(s)}{|a(s)|} (r|\alpha| - (1 + r|\alpha|) |f_\varepsilon(s)|). \end{aligned}$$

Since $|f_\varepsilon(s)| \in [0, 1]$, we have

$$\begin{aligned} -1 - r|\alpha| &\leq -(1 + r|\alpha|) |f_\varepsilon(s)| \leq 0, \\ -1 &\leq r|\alpha| - (1 + r|\alpha|) |f_\varepsilon(s)| \leq r|\alpha| \leq 1, \end{aligned}$$

which implies that $|r|\alpha| - (1 + r|\alpha|) |f_\varepsilon(s)| \leq 1$ for all $s \in V_\varepsilon$. It follows that $|c_{r,\varepsilon}(s)| \leq 1$, for all $s \in L$.

We have shown that $\|c_{r,\varepsilon}\| \leq 1$, and (3.5) implies that $c_{r,\varepsilon} \in F_{t_0, -\frac{\alpha}{|\alpha|}} \subset S(C_0^\mathbb{T}(L))$.

Finally, we compute the distance between a_ε and $c_{r,\varepsilon}$ to get

$$\begin{aligned} \|a_\varepsilon - c_{r,\varepsilon}\| &= \|(1 - r)a_\varepsilon + (1 + r|\alpha|) |f_\varepsilon(s)| u_\varepsilon(s)\| \\ &\leq 1 - r + 1 + r|\alpha| = 2 - r + r|\alpha|. \end{aligned}$$

□

We are now in a position to prove that every abelian JB*-triple satisfies the second hypothesis of Proposition 2.1 for L_0 as the set of representatives.

Proposition 3.10. *For every $t_0 \in L_0$ and $\alpha \in \overline{\mathbb{D}}$ the equality*

$$M_{t_0, \alpha} = M_{\delta_{t_0}, \alpha} = \{a \in S(C_0^\mathbb{T}(L)) : a(t_0) = \alpha\}$$

holds.

Proof. We only need to show the inclusion \supseteq , because the other implication always holds. Take any $a \in S(C_0^\mathbb{T}(L))$ such that $a(t_0) = \alpha$. We shall discuss first the case where $|\alpha| = 1$, in such case we have $a \in F_{t_0, \frac{\alpha}{|\alpha|}} = F_{t_0, \alpha}$ and $-a \in F_{t_0, -\frac{\alpha}{|\alpha|}} = F_{t_0, -\alpha}$. Thus, $d(a, F_{t_0, \frac{\alpha}{|\alpha|}}) \leq d(a, a) = 0 = 1 - |\alpha|$ and $d(a, F_{t_0, -\frac{\alpha}{|\alpha|}}) \leq d(a, -a) = 2 = 1 + |\alpha|$.

We assume next that $\alpha = 0$. For each $\varepsilon > 0$ we can find an open neighbourhood U_ε of t_0 and an element $a_\varepsilon \in S(C_0^\mathbb{T}(L))$ such that $\|a - a_\varepsilon\| < \varepsilon$ and $a_\varepsilon|_{U_\varepsilon} \equiv 0$. Keeping in mind Remark 3.4, we may assume that U_ε is \mathbb{T} -symmetric and has compact \mathbb{T} -symmetric closure. Then, by Remark 3.3, there exists a function $b_\varepsilon \in F_{t_0, 1}$ (and thus $-b_\varepsilon \in F_{t_0, -1}$) with $b_\varepsilon|_{L \setminus U_\varepsilon} \equiv 0$. Putting all together, we have

$$d(a, F_{t_0, \pm 1}) \leq \|a \mp b_\varepsilon\| \leq \|a - a_\varepsilon\| + \|a_\varepsilon \mp b_\varepsilon\| < \varepsilon + \max\{\|b_\varepsilon\|, \|a_\varepsilon\|\} \leq 1 + \varepsilon.$$

Therefore, $d(a, F_{t_0, \pm 1}) \leq 1 \mp |\alpha|$.

We finally assume that $0 < |\alpha| = |a(t_0)| < 1$. For each $\varepsilon > 0$, Lemma [8, Lemma 3.15], gives $b_\varepsilon \in F_{t_0,1}$ and $a_\varepsilon \in S(C_0^\mathbb{T}(L))$ such that

$$c_{r,\varepsilon} = ra_\varepsilon + (1 - r|a(t_0)|) \frac{a(t_0)}{|a(t_0)|} b_\varepsilon \in F_{t_0, \frac{\alpha}{|\alpha|}} \quad (\forall 0 < r < 1),$$

$a_\varepsilon(t_0) = a(t_0)$, and $\|a - a_\varepsilon\| < \varepsilon$.

On the one hand, we have

$$\begin{aligned} \|a - c_{r,\varepsilon}\| &= \|a - a_\varepsilon\| + \|a_\varepsilon - c_{r,\varepsilon}\| < \varepsilon + \left\| (1 - r)a_\varepsilon - (1 - r|\alpha|) \frac{\alpha}{|\alpha|} b_\varepsilon \right\| \\ &\leq \varepsilon + (1 - r) + 1 - r|\alpha| = 2 - r - r|\alpha| + \varepsilon \end{aligned}$$

for each $\varepsilon > 0$ and $0 < r < 1$. Letting $\varepsilon \rightarrow 0$ and $r \rightarrow 1$ we get $d(a, F_{t_0, \frac{\alpha}{|\alpha|}}) \leq 1 - |\alpha|$.

Similarly, by Lemma 3.9, there exist $a'_\varepsilon, b'_\varepsilon \in S(C_0^\mathbb{T}(L))$ such that $\|a - a'_\varepsilon\| \leq \varepsilon/2$, and for each $0 < r < 1$, the function

$$c'_{r,\varepsilon} := ra'_\varepsilon - (1 + r|\alpha|) b'_\varepsilon$$

lies in $F_{t_0, -\frac{\alpha}{|\alpha|}}$ with $\|a_\varepsilon - c'_{r,\varepsilon}\| \leq 2 - r + r|\alpha|$. Thus,

$$\|a - c'_{r,\varepsilon}\| = \|a - a'_\varepsilon\| + \|a'_\varepsilon - c'_{r,\varepsilon}\| \leq \varepsilon/2 + 2 - r + r|\alpha|$$

for all $\varepsilon > 0$ and $0 < r < 1$. By taking limits in $\varepsilon \rightarrow 0$ and $r \rightarrow 1$ we arrive at $d(a, F_{t_0, -\frac{\alpha}{|\alpha|}}) \leq 1 + |\alpha|$. Therefore, $a \in M_{t_0, \alpha}$. \square

Proof of Theorem 3.1. Remark 3.7 and Proposition 3.10 guarantee that $C_0^\mathbb{T}(L)$ satisfies the hypotheses in Proposition 2.1 [11, Proposition 4.4] for the set of representatives given by L_0 . The just quoted proposition gives the desired conclusion. \square

Acknowledgements We would like to thank Professor O. Hatori from Niigata University for sharing with us some of his recent achievements, besides some fruitful discussions and useful comments.

Second author partially supported by EPSRC (UK) project ‘‘Jordan Algebras, Finsler Geometry and Dynamics’’ ref. no. EP / R044228 / 1 and by the Spanish Ministry of Science, Innovation and Universities (MICINN) and European Regional Development Fund project no. PGC2018-093332-B-I00, Junta de Andalucıa grants FQM375 and A-FQM-242-UGR18. Fourth author partially supported by JSPS KAKENHI (Japan) Grant Number JP 20K03650. Fifth author partially supported by MCIN / AEI / 10. 13039 / 501100011033 / FEDER ‘‘Una manera de hacer Europa’’ project no. PGC2018-093332-B-I00, Junta de Andalucıa grants FQM375, A-FQM-242-UGR18 and PY20_ 00255, and by the IMAG–Marıa de Maeztu grant CEX2020-001105-M / AEI / 10.13039 / 501100011033.

REFERENCES

- [1] T. Banach, Any isometry between the spheres of absolutely smooth 2-dimensional Banach spaces is linear, *J. Math. Anal. Appl.* **500** (2021), no. 1, Paper No. 125104.
- [2] T. Banach, Every 2-dimensional Banach space has the Mazur-Ulam property, *Linear Algebra Appl.* **632** (2022), 268-280.
- [3] T. Banach, J. Cabello Sanchez, Every non-smooth 2-dimensional Banach space has the Mazur-Ulam property, *Linear Algebra Appl.* **625** (2021), 1-19.
- [4] J. Becerra Guerrero, M. Cueto-Avellaneda, F.J. Fernandez-Polo, A.M. Peralta, On the extension of isometries between the unit spheres of a JBW*-triple and a Banach space, *J. Inst. Math. Jussieu* **20**, no. 1 (2021), 277–303.

- [5] K Boyko, V Kadets, M Martińn, D Werner, Numerical index of Banach spaces and duality, *Math. Proc. Cambridge Phil. Soc.* **142** (1) (2007), 93-102
- [6] J. Cabello-Sánchez, A reflection on Tingley's problem and some applications, *J. Math. Anal. Appl.* **476** (2) (2019), 319–336.
- [7] L. Cheng, Y. Dong, On a generalized Mazur-Ulam question: extension of isometries between unit spheres of Banach spaces, *J. Math. Anal. Appl.* **377** (2011), 464-470.
- [8] Cueto-Avellaneda, María and Hirota, Daisuke and Miura, Takeshi and Peralta, Antonio M, Exploring new solutions to Tingley's problem for function algebras, *arXiv preprint arXiv:2110.11120* (2021).
- [9] G. Ding, On the extension of isometries between unit spheres of E and $C(\Omega)$, *Acta Math. Sin. (Engl. Ser.)* **19** (2003) 793-800.
- [10] X.N. Fang, J.H. Wang, Extension of isometries between the unit spheres of normed space E and $C(\Omega)$, *Acta Math. Sinica (Engl. Ser.)*, **22** (2006), 1819–1824.
- [11] O. Hatori, The Mazur-Ulam property for uniform algebras, to appear in *Studia Math.* arXiv:2107.01515
- [12] O. Hatori, S. Oi, R. Shindo Togashi, Tingley's problems on uniform algebras, *J. Math. Anal. Appl.* **503** (2021), no. 2, Paper No. 125346, 14 pp.
- [13] O.F.K. Kalenda, A.M. Peralta, Extension of isometries from the unit sphere of a rank-2 Cartan factor, *Anal. Math. Phys.* **11**, 15 (2021).
- [14] W. Kaup, A Riemann Mapping Theorem for bounded symmetric domains in complex Banach spaces, *Math. Z.* **183** (1983), 503–529.
- [15] R. Liu, On extension of isometries between unit spheres of $\mathcal{L}^\infty(\Gamma)$ -type space and a Banach space E , *J. Math. Anal. Appl.* **333** (2007), 959-970.
- [16] M. Mori, N. Ozawa, Mankiewicz's theorem and the Mazur-Ulam property for C^* -algebras, *Studia Math.* **250**, no. 3 (2020), 265–281.
- [17] G.H. Olsen, On the classification of complex Lindenstrauss spaces, *Math. Scand.* **35** (1974), 237-258.
- [18] A.M. Peralta, On the extension of surjective isometries whose domain is the unit sphere of a space of compact operators, to appear in *FILOMAT.* arXiv:2005.11987v1
- [19] W. Rudin, Real and complex analysis. Third edition. McGraw-Hill Book Co., New York, 1987.
- [20] D. Tan, X. Huang, R. Liu, Generalized-lush spaces and the Mazur-Ulam property, *Studia Math.* **219** (2013), 139–153.
- [21] R. Tanaka, The solution of Tingley's problem for the operator norm unit sphere of complex $n \times n$ matrices, *Linear Algebra Appl.* **494** (2016), 274-285.
- [22] R. Tanaka, Tingley's problem on finite von Neumann algebras, *J. Math. Anal. Appl.*, **451** (2017), 319-326.

(D. Cabezas) DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, FACULTAD DE CIENCIAS, UNIVERSIDAD DE GRANADA, 18071 GRANADA, SPAIN.

Email address: dcabezas@ugr.es

(M. Cueto-Avellaneda) SCHOOL OF MATHEMATICS, STATISTICS AND ACTUARIAL SCIENCE, UNIVERSITY OF KENT, CANTERBURY, KENT CT2 7NX, UK

Email address: emecueto@gmail.com

(D. Hirota) GRADUATE SCHOOL OF SCIENCE AND TECHNOLOGY, NIIGATA UNIVERSITY, NIIGATA 950-2181, JAPAN

Email address: hirota@m.sc.niigata-u.ac.jp

(T. Miura) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, NIIGATA UNIVERSITY, NIIGATA 950-2181, JAPAN

Email address: miura@math.sc.niigata-u.ac.jp

(A.M. Peralta) INSTITUTO DE MATEMÁTICAS DE LA UNIVERSIDAD DE GRANADA (IMAG), DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, FACULTAD DE CIENCIAS, UNIVERSIDAD DE GRANADA, 18071 GRANADA, SPAIN.

Email address: aperalta@ugr.es