

COARSE EMBEDDINGS OF SYMMETRIC SPACES AND EUCLIDEAN BUILDINGS

OUSSAMA BENSAID

ABSTRACT. Introduced by Gromov in the 80's, coarse embeddings are a generalization of quasi-isometric embeddings when the control functions are not necessarily affine. In this paper, we will be particularly interested in coarse embeddings between symmetric spaces and Euclidean buildings. The quasi-isometric case is very well understood thanks to the rigidity results for symmetric spaces and buildings of higher rank by Anderson–Schroeder, Kleiner, Kleiner–Leeb and Eskin–Farb in the 90's. In particular, it is well known that the rank of these spaces is monotonous under quasi-isometric embeddings. This is no longer the case for coarse embeddings as shown by horospherical embeddings. However, we show that in the absence of a Euclidean factor in the domain, the rank is monotonous under coarse embeddings. This answers a question by David Fisher and Kevin Whyte. We can also relax the condition on the domain by allowing it to contain a Euclidean factor of dimension 1, answering a question by Gromov.

CONTENTS

1. Introduction	2
1.1. Coarse embeddings in geometric group theory	2
1.2. Main results	3
1.3. Idea of the proof	6
1.4. Organization of the paper	7
1.5. Acknowledgements	8
2. Background	8
2.1. Coarse embeddings	8
2.2. Coarse geometry of symmetric spaces and Euclidean buildings	9
2.3. Metric currents	12
2.4. Filling functions	13
2.5. Connect the dots argument	15
3. The domain X is a product of spaces with exponential growth	16
3.1. Preliminary lemmas	16
3.2. Proof of Theorem 1.9	18
4. The domain X is a model space with no Euclidean factor	28
4.1. Preliminary result on Weyl sectors	28
4.2. Parallelepipeds and Parallelograms	30
4.3. k -sequences	37

4.4. Proof of Theorem 1.10	38
5. When the domain X has a one-dimensional Euclidean factor	49
5.1. Coarse embeddings of Euclidean spaces into lower rank	49
6. Further questions	54
References	55

1. INTRODUCTION

The purpose of the present paper is to provide obstructions to the existence of coarse embeddings between symmetric spaces and Euclidean buildings. In order to put our main results in perspective, we start this introduction with a brief survey on the notion of coarse embedding and its applications in geometric group theory.

1.1. Coarse embeddings in geometric group theory. We recall that a map $f : (X, d_X) \rightarrow (Y, d_Y)$ is a coarse embedding if there exist functions $\rho_{\pm} : [0, \infty) \rightarrow [0, \infty)$ such that $\rho_{-}(r) \rightarrow \infty$ as $r \rightarrow \infty$ and for all $x, y \in X$

$$\rho_{-}(d_X(x, y)) \leq d_Y(f(x), f(y)) \leq \rho_{+}(d_X(x, y)).$$

Coarse embeddings have been introduced under the name of “placements” or “placings” in [Gro88]. They also appear in the literature as uniform embeddings in [Gro93],[Sha04],[FW18], effectively proper Lipschitz maps in [BW92] or uniformly proper embeddings [MSW03].

Coarse embeddings from groups or infinite graphs to Hilbert spaces (or more general Banach spaces) have received a lot of attention due to their connection with the Baum–Connes and the Novikov conjecture (see [Yu00], [STY02], [KY06]). However, they have originally been much less studied than quasi-isometric embeddings among finitely generated groups, probably due to their much greater flexibility. Note that a subgroup inclusion between finitely generated groups is always a coarse embedding, while it is a quasi-isometric embedding only when the subgroup in question is undistorted. Very few geometric invariants are known to be monotonous under coarse embedding: the only examples (which takes infinitely many different values) being the volume growth, the asymptotic dimension [Gro93], the separation profile [BST12] and the Poincaré profiles [HMT20].

The volume growth allows to say for example that a metric space with exponential growth cannot be coarsely embedded into a space with polynomial growth, but it does not distinguish between spaces of exponential growth. More recently the Følner function has been shown to be monotonous under coarse embedding between amenable groups [DKLMT20]. This provides a much more refined invariant to distinguish among solvable groups.

The asymptotic dimension was introduced by Gromov in [Gro88] and [Gro93] and is a large-scale analogue of Lebesgue covering dimension. For symmetric spaces of nonpositive curvature and Euclidean buildings, it simply coincides with the dimension. For instance it allows to rule out coarse embeddings from \mathbb{H}^n to \mathbb{H}^p for $p < n$, but it does not prevent for

instance the existence of a coarse embedding of the complex hyperbolic plane to \mathbb{H}^4 .

The separation profile is a powerful monotone coarse invariant introduced by Benjamini, Schramm and Timár [BST12]. The separation profile of an infinite, bounded degree graph at $n \in \mathbb{N}$ is the supremum over all subgraphs of size $\leq n$, of the number of vertices needed to be removed from the subgraph, in order to cut it to connected pieces of size at most $n/2$. The separation profile of a metric space with bounded geometry can be defined as the separation profile of any graph that is quasi-isometric to it. For rank 1 symmetric spaces, it was shown to detect the conformal dimension of the boundary [HMT20], providing for instance an obstruction to the existence of a coarse embedding from the complex hyperbolic plane to \mathbb{H}^3 . Later Hume–Mackay–Tessera [HMT20] introduced the L^p -Poincaré profiles, generalizing the separation profile. These invariants are very efficient in distinguishing between spaces of the form $X \times \mathbb{R}^n$ where X is a hyperbolic space. However, for higher rank symmetric spaces all L^p -Poincaré profiles are $\simeq n/\log n$: this provides an obstruction to coarse embeddings from higher rank to $X \times \mathbb{R}^n$, where X is rank one, but does not provide any obstruction among higher rank symmetric spaces.

Hume–Sisto gave in [HS17] an obstruction for a group to admit a coarse embedding into a real hyperbolic space: admitting exponentially many fat bigons. It is shown for example that direct products of two infinite groups one of which has exponential growth, solvable groups that are not virtually nilpotent, and uniform higher-rank lattices do not coarsely embed into a hyperbolic space.

Using a combination of results among the ones mentioned above and a result by Le Coz and Gournay [CG19], Tessera recently proved the following statement [Tes20]: an amenable group admits a coarse embedding into a hyperbolic group if and only if it is virtually nilpotent. We conclude this survey by mentioning an interesting application of the previous result in pseudo-riemannian geometry. The story starts with an early observation of Gromov [Gro88, Section 4.1] according to which the isometry group of a compact $(n+1)$ -dimensional Lorentzian manifold admits a coarse embedding into the real hyperbolic space \mathbb{H}^n . Charles Frances [Fra21] exploited this point of view and the above result to prove a Tits alternative for discrete subgroups of the isometry group of compact Lorentz manifolds. Namely, a discrete, finitely generated, subgroup of the isometry group of a compact Lorentz manifold of dimension $n \geq 2$ either contains a free subgroup in two generators, in which case it is virtually isomorphic to a discrete subgroup of $\mathrm{PO}(1, d)$, or is virtually nilpotent of growth degree $\leq n - 1$.

We deduce from this short survey that the only known obstruction for coarse embeddings among higher rank symmetric spaces is given by their dimension. Our goal is to address this problem by showing that under certain conditions, the rank is also a coarse monotonous invariant.

1.2. Main results.

Let S be a product of symmetric spaces of non-compact type and B a product of thick Euclidean buildings with cocompact affine Weyl group, with bounded geometry and no Euclidean factor. The spaces that we are going to consider are of the form $X = \mathbb{R}^n \times S \times B$.

We will call such space a *model space*. Below are a few known examples of embeddings between such model spaces:

- Every regular tree T_d quasi-isometrically embeds into the hyperbolic plane \mathbb{H}^2 .
- There are quasi-isometric embeddings $\mathbb{H}^3 \rightarrow \mathbb{H}^2 \times \mathbb{H}^2$ and $\mathbb{H}^5 \rightarrow \mathbb{H}^3 \times \mathbb{H}^3$ (see [BF98] for a more general statement).
- There exist coarse embeddings $\mathbb{H}^2 \rightarrow \mathbb{H}^3$ with arbitrarily small lower control [BH21].
- \mathbb{H}^n quasi-isometrically embeds into a product of n binary trees $T_3 \times \cdots \times T_3$ [BDS07].
- There exist quasi-isometric embeddings of the product of n copies of the hyperbolic plane $\mathbb{H}^2 \times \cdots \times \mathbb{H}^2$ into the symmetric spaces $\mathrm{SL}_{n+1}(\mathbb{R})/\mathrm{SO}_{n+1}(\mathbb{R})$ and $\mathrm{Sp}_{2n}(\mathbb{R})/\mathrm{U}_n(\mathbb{R})$ [FW18].

We define the *rank* of such a model space as the maximal dimension of an isometrically embedded copy of a Euclidean space. We are interested in the following natural question.

Question 1.1. Is the rank monotonous under coarse embeddings? In other words, if X and Y are two model spaces and there is a coarse embedding $f : X \rightarrow Y$, does it imply that $\mathrm{rank}(X) \leq \mathrm{rank}(Y)$?

The answer is positive and well-known if one replaces the word coarse by quasi-isometric. It was shown by Anderson–Schroeder [AS86] that if X is a symmetric space of non-compact type and $f : \mathbb{R}^n \rightarrow X$ is a quasi-isometric embedding, then X contains an n -flat, i.e. there exists an isometric embedding of \mathbb{R}^n into X . Later, Kleiner [Kle99] generalized this to all locally compact cocompact Hadamard spaces, which includes all our model spaces. This answers the question for quasi-isometric embeddings. Indeed, if $f : X \rightarrow Y$ is a quasi-isometric embedding between two model spaces and $\mathrm{rank}(X) = p$ then \mathbb{R}^p quasi-isometrically embeds into Y , hence $\mathrm{rank}(Y) \geq p$. Moreover, it was shown by Kleiner–Leeb [KL97] and by Eskin–Farb [EF97] that maximal quasi-flats in higher rank symmetric spaces of non-compact type satisfy a generalization of Morse Lemma: if Y is a symmetric space of non-compact type of rank $n \geq 2$, and $f : \mathbb{R}^n \rightarrow Y$ is a quasi-isometric embedding then there exist $\delta > 0$ and $k \in \mathbb{N}$ (that depends on the quasi-isometry constants) such that $f(\mathbb{R}^n)$ lies in the δ -neighborhood of a union of k flats in Y . Note that this is a general phenomenon that occurs in all asymptotic hierarchically hyperbolic spaces [BHS21].

The monotonicity of the rank is no longer satisfied in the setting of coarse embeddings in general. Counter-examples are given by horospherical embeddings. Let \mathbb{H}^2 denote the upper half-plane model for the hyperbolic plane. We have

$$d_{\mathbb{H}^2}((x, 1), (x + r, 1)) = \arg \cosh(1 + r^2/2) \underset{\ell \rightarrow \infty}{\sim} 2 \ln r.$$

So the map $f : \mathbb{R} \rightarrow \mathbb{H}^2$, $x \mapsto (x, 1)$, which is a parametrization of a horocycle, is a coarse embedding that is exponentially distorted. This carries over to the hyperbolic space of any dimension $n \geq 1$, and provides coarse embeddings $f : \mathbb{R}^n \rightarrow \mathbb{H}^{n+1}$ whose images are horospheres. These embeddings do not respect the monotonicity of the rank since $\mathrm{rank}(\mathbb{R}^n) = n$ and $\mathrm{rank}(\mathbb{H}^{n+1}) = 1$. Therefore, as long as a model space X has a Euclidean factor \mathbb{R}^p with $p \geq 2$, it can be embedded into a model space with lower rank. Indeed, if $X = \mathbb{R}^p \times S \times B$ and $\mathrm{rank}(X) = p + r$, then it coarsely embeds into $\mathbb{H}^{p+1} \times S \times B$, whose rank is $1 + r$. This naturally leads us to ask the following questions.

Question 1.2. When X has no Euclidean factor, can we still have a coarse embedding from X to a model space Y such that $\text{rank}(X) > \text{rank}(Y)$? What about when X has a Euclidean factor of dimension 1? i.e. if $X = \mathbb{R} \times S \times B$, can we have a coarse embedding from X to a model space Y such that $\text{rank}(X) > \text{rank}(Y)$?

We can show that, in the absence of a Euclidean factor in the domain X , the rank is monotonous under coarse embeddings, answering a question of Fisher and Whyte [FW18].

In all the following results, the Euclidean buildings in the model space in the target are not necessarily thick, nor have cocompact affine Weyl group. It is only required for the Euclidean buildings in the domain.

Theorem 1.3 (See Theorem 1.9). *Let $X = X_1 \times \cdots \times X_k$ be a product of geodesic metric spaces of exponential growth, and let $Y = \mathbb{R}^n \times S \times B$ be a model space.*

If $k > \text{rank}(Y)$, then there is no coarse embedding from X to Y .

This result has been originally announced by Gromov in [Gro93], who outlined a strategy of proof. Our proof relies mainly on his sketch. Moreover, we can extend this strategy to prove a similar result when X is a general model space. Namely, we show

Theorem 1.4 (see Theorem 1.10). *Let $X = S \times B$ and $Y = \mathbb{R}^n \times S' \times B'$ be model spaces. If $\text{rank}(X) > \text{rank}(Y)$, then there is no coarse embedding from X to Y .*

This implies for example that there is no coarse embedding from $T_3 \times T_3 \times T_3$ into $\mathbb{H}^p \times \mathbb{H}^q$ for any integers $p, q \geq 1$, nor into the symmetric space $\text{SL}_3(\mathbb{R})/\text{SO}_3(\mathbb{R})$.

When X has a Euclidean factor of dimension 1, we can show that the monotonicity of the rank still holds, answering a question by Gromov [Gro93].

Theorem 1.5 (see Theorem 1.11). *Let X be either $X_1 \times \cdots \times X_k$ as in Theorem 1.3, or a model space $S \times B$ of rank k as in Theorem 1.4, and let $Y = \mathbb{R}^n \times S' \times B'$ be of rank $\leq k$.*

Then there is no coarse embedding from $X \times \mathbb{R}$ to Y .

We also give a result about coarse embeddings of Euclidean spaces into symmetric spaces of lower rank, which uses the generalization of Morse lemma for higher-rank symmetric spaces:

Theorem 1.6. *Let $p > k \geq 1$ be integers. Let Y be a symmetric space of non-compact type such that $\text{rank}(Y) = k$, and let $f : \mathbb{R}^p \rightarrow Y$ be a coarse embedding. Then no Euclidean subspace $E \simeq \mathbb{R}^k \subset \mathbb{R}^p$ is sent quasi-isometrically by f .*

In the rank one case, we can extend the result to Gromov-hyperbolic geodesic metric spaces with bounded geometry thanks to a result by Bonk–Schramm [BS11].

Corollary 1.7. *Let Y be a Gromov-hyperbolic geodesic metric space with bounded geometry, and let $p \geq 2$ be an integer. A coarse embedding $f : \mathbb{R}^p \rightarrow Y$ is uniformly compressing, i.e. it admits a sublinear upper control function ρ^+ .*

Now, one can naturally ask: If X does not coarsely embed into Y , does adding a Euclidean factor in the target makes the embedding possible?

We can answer this question when $\text{rank}(X) > \text{rank}(Y) + 1$. In fact, when $\text{rank}(X) > \text{rank}(Y) + 1$, even adding a nilpotent factor in the target does not make this embedding possible:

Corollary 1.8. *Let X be either $X_1 \times \cdots \times X_k$ as in Theorem 1.3, or a model space $S \times B$ of rank k as in Theorem 1.4, and let $Y = \mathbb{R}^n \times S' \times B'$ be a model space of rank $< k - 1$. Then for any nilpotent connected Lie group P , there is no coarse embedding from X to $Y \times P$.*

Proof. By Assouad's embedding theorem [Ass82], P coarsely embeds into some \mathbb{Z}^d , which in turn coarsely embeds into \mathbb{H}^{d+1} . Therefore, we get a coarse embedding from X to $Y \times \mathbb{H}^{d+1}$, but $\text{rank}(Y \times \mathbb{H}^{d+1}) = \text{rank}(Y) + 1 < k$. Such coarse embedding is not possible by Theorems 1.3 and 1.4. \square

The main tools that we are going to use are filling functions using Lipschitz chains. They measure the difficulty of filling cycles of a given size. The key point is that filling functions detect the rank. In other words, they behave differently whether the dimension of the cycle is smaller or greater than the rank of the space. In fact, if a cycle Σ in a model space X has dimension $k < \text{rank}(X)$ and $\text{Vol}_k^X(\Sigma) = l$, then it can be filled similarly as in a Euclidean space, i.e. it has a filling with $(k+1)$ -volume $\lesssim l^{\frac{k+1}{k}}$. On the other hand, if $k > \text{rank}(X)$, then it can be filled linearly like in a hyperbolic space, i.e. it admits a filling with $(k+1)$ -volume $\lesssim l$. This result is due to Leuzinger [Leu14]. Our proof will also rely on the co-area formula which requires some slicing techniques. However, Lipschitz chains do not behave well under slicing, that is why we will consider metric currents in complete metric spaces that were introduced by Ambrosio–Kirchheim [AK⁺00]. They can be seen as a generalization of Lipschitz chains since any Lipschitz chain induces naturally a current, which behaves well under slicing.

We can restate our results in terms of filling functions, and in a more general framework, allowing the target space Y to be any complete uniformly contractible metric space (see Section 2.5 for a definition) with at most exponential growth:

Theorem 1.9. *Let $X = X_1 \times \cdots \times X_k$ be a product of geodesic metric spaces with exponential growth, and let Y be a uniformly contractible complete metric space with at most exponential growth. If $\text{FV}_k^Y(l) \sim l$, then there is no coarse embedding from X to Y .*

Theorem 1.10. *Let $X = S \times B$ be a model space and let Y be a uniformly contractible complete metric space with at most exponential growth. If $\text{FV}_k^Y(l) \sim l$, then there is no coarse embedding from X to Y .*

Theorem 1.11. *Let X be either $X_1 \times \cdots \times X_k$ as in Theorem 1.3, or a model space $S \times B$ of rank k , and let Y be a uniformly contractible complete metric space with at most exponential growth. If $\text{FV}_{k+1}^Y(l) \sim l$, then there is no coarse embedding from $X \times \mathbb{R}$ to Y .*

Theorems 1.3, 1.4 and 1.5 immediately follow from these by Leuzinger's theorem, since all our model spaces are uniformly contractible, complete, and with at most exponential growth.

1.3. Idea of the proof. Let us first look at Theorem 1.9, which implies Theorem 1.3 by Leuzinger's Theorem. The case $k = 1$ is a simple volume obstruction observation: a space of exponential growth cannot be coarsely embedded in a space with at most exponential growth with sublinear upper control ρ_+ . Let us give a sketch of the proof when $k = 2$, which already contains most of the conceptual difficulties of the general case. When $X = X_1 \times X_2$ is a product of two geodesic metric spaces of exponential growth and the 2-dimensional filling function of Y is linear, suppose that a coarse embedding $f : X \rightarrow Y$ exists. By our observation, no copy of X_1 inside of X is sent sublinearly, which means that there exist

“undistorted directions”, more precisely there exist pairs of points $(a_n)_n, (b_n)_n$ in X_1 such that $d_X(a_n, b_n) \rightarrow \infty$ and which are mapped in Y quasi-isometrically. Now consider the rectangles R_n with one geodesic segment from a_n to b_n in X_1 as one side, and a geodesic segment in a copy of X_2 of length at most $d_X(a_n, b_n)^{1/2}$ as the orthogonal side. By the linear filling in Y , $f(\partial R_n)$ can be filled in Y by a chain of volume $\leq Cd_X(a_n, b_n)$. Since (a_n) and (b_n) are sent quasi-isometrically, the height of $f(\partial R_n)$ is comparable to that of ∂R_n , however the filling of $f(\partial R_n)$ in Y is much smaller than that of ∂R_n in X . This implies, by the co-area formula for currents, that the width of $f(\partial R_n)$ is highly compressed. From this observation, we construct a sequence of sets in X whose “coarse”-volume, that will be defined in the background material, is not coarsely preserved by f , which is not possible for coarse embeddings. For $k \geq 3$, the proof is done in a similar way, by induction on k .

To prove Theorem 1.10, which implies Theorem 1.4, we will adapt the previous proof. Since the domain is non longer a product space, we will consider parallelograms instead of rectangles, and adapt the size of its sides to make the proof work. More importantly, instead of looking for “undistorted directions”, we need to find “maximally singular undistorted directions”. Maximally singular directions are directions following which a space of rank k contains a subset that factorizes as a product $\mathbb{R} \times X'$, where X' has rank $k - 1$. To find these directions, we will decompose the cycle along the walls of a suitable Weyl sector, and use a pigeonhole-like reasoning.

Theorem 1.5 is a consequence of the first two theorems, combined with a result about coarse embeddings of Euclidean spaces into symmetric spaces of lower rank. This result allows us to derive an upper bound for fillings of images of k -parallelograms, starting from the linear filling of dimension $k + 1$ in Y .

The proof of Theorem 1.6 is done by contradiction. If there exists such quasi-isometric embedding, then by the quasi-flats Theorem of Kleiner–Leeb [KL97] and Eskin–Farb [EF97], such a quasi-flat is a bounded distance from a finite union of maximal flats. This implies that there exist arbitrarily big balls that are sent uniformly close to a maximal flat. By considering parallelograms inside these balls and the filling of the projection of their images in this flat, we get a contradiction. Namely, we get a sequence of images of parallelograms that are sent quasi-isometrically in a Euclidean space, and that are filled linearly, which gives a contradiction by applying the co-area formula. Finally, a result of Bonk–Schramm [BS11] says that any Gromov hyperbolic geodesic metric space with bounded geometry quasi-isometrically embeds into a hyperbolic space \mathbb{H}^n . By composing by this quasi-isometry and applying the previous theorem we get the Corollary 1.7.

1.4. Organization of the paper. We will start in section 2 by giving some background material that will be used throughout the paper. More precisely, section 2.1 is dedicated to coarse embeddings. In section 2.2 we collect definitions and some large-scale properties of the model spaces that we consider, like parallel sets and cross sections. In 2.3 we give a brief overview of the theory of metric currents introduced by Ambrosio–Kirchheim, and state the Slicing Theorem that will play a crucial role in the proofs. We define the filling functions in subsection 2.4 and state Leuzinger’s theorem that shows that the filling functions can detect the rank of the model spaces. Finally, connect the dots argument in 2.5 will allow us to always assume that our coarse embeddings are Lipschitz, which is very convenient when

working with Lipschitz chains or currents. Before proving Theorem 1.9 in section 3.2, which implies Theorem 1.3, we will give some useful lemmas about coarse embeddings and volume preservation in section 3.1. In section 4, we give a preliminary result about Weyl sectors in either symmetric spaces of non-compact type or Euclidean buildings, and we prove a decomposition result for parallelograms that will be used in the proof of Theorem 1.10 right after. The goal of section 5 is to give a proof of Theorem 1.11, thus implying Theorem 1.5. It starts first by giving a result about coarse embeddings of Euclidean spaces into symmetric spaces of lower rank, which also implies Theorem 1.6 and Corollary 1.7. Finally, we collect some further questions in section 6.

1.5. Acknowledgements. I am extremely grateful to my advisor Romain Tessera for introducing me to this subject, for all the discussions that we had and all his support. I wish also to thank Charles Frances for explaining to me the motivations and applications of coarse embeddings in Lorentzian geometry. I have benefited from many useful discussions with David Hume, John Mackay, Pierre Pansu, and Linus Kramer. I would also like to thank Amandine Escalier and Mingkun Liu for many discussions and for helping me with the drawings.

2. BACKGROUND

2.1. Coarse embeddings. A map $f : (X, d_X) \rightarrow (Y, d_Y)$ is a coarse embedding if there exist functions $\rho_{\pm} : [0, \infty) \rightarrow [0, \infty)$ such that $\rho_{-}(r) \rightarrow \infty$ as $r \rightarrow \infty$ and for all $x, y \in X$

$$\rho_{-}(d_X(x, y)) \leq d_Y(f(x), f(y)) \leq \rho_{+}(d_X(x, y)).$$

Equivalently, for every pair $(x_n)_{n \geq 0}, (x'_n)_{n \geq 0}$ of sequences of points in X ,

$$\lim_{n \rightarrow \infty} d_X(x_n, x'_n) = \infty \iff \lim_{n \rightarrow \infty} d_Y(f(x_n), f(x'_n)) = \infty.$$

When the control functions ρ_{-}, ρ_{+} are affine, f is said to be a *quasi-isometric embedding*. The map f is said to be *large-scale Lipschitz* if it admits an affine upper control, without necessarily having a lower control.

Definition 2.1. A metric space (X, d) is *large-scale geodesic* if there exist constants $\lambda, c > 0$ and $b \geq 0$ such that, for every pair (x, x') of points of X , there exists a sequence $x_0 = x, x_1, \dots, x_n = x'$ of points in X such that $d(x_{i-1}, x_i) \leq c$ for $i = 1, \dots, n$ and $n \leq \lambda d(x, x') + b$.

If X is large-scale geodesic, then f admits a control function ρ_{+} which is affine.

Definition 2.2. A metric space X has *bounded geometry* if there exists $R_0 \geq 0$ such that, for every $R \geq 0$, there exists an integer N such that every ball of radius R in X can be covered by N balls of radius R_0 .

Remark 2.3. An example of a metric space that is not of bounded geometry is an infinite-dimensional Hilbert space, with the metric given by the norm.

Definition 2.4. Let X be a metric space with bounded geometry and let $\varepsilon > 0$. We define the ε -*volume* of a subset $A \subset X$, that we will denote $\text{Vol}_X^{\varepsilon}(A)$, as the minimal number of balls of radius ε needed to cover A .

We will denote by $B(x, r)$ the closed ball $B(x, r) := \{y \in X : d(y, x) \leq r\}$.

Definition 2.5. Let $\varepsilon > 0$. We define the ε -growth function of a metric space (X, d) with bounded geometry to be

$$\beta_X^\varepsilon(r) = \sup\{\text{Vol}_X^\varepsilon(B(x, r)) \mid x \in X\}.$$

We say that X has *polynomial growth* if there exist $\varepsilon > 0$ and $D \geq 0$ such that $\beta_X^\varepsilon(r) \preceq r^D$. We say that X has *exponential growth* if there exists $\varepsilon > 0$ such that $\beta_X^\varepsilon(r) \approx e^r$.

The comparison on functions $\mathbb{N} \rightarrow \mathbb{R}$ is defined as follows: $f \preceq g$ if there exists a constant $c > 0$ such that $f(n) \leq cg(cn + c) + c$ for all $n \in \mathbb{N}$, and $f \approx g$ if $f \preceq g$ and $g \preceq f$.

Remark 2.6. If X is large-scale geodesic and has bounded geometry, then it has at most exponential growth $\beta_X^\varepsilon(r) \preceq e^r$.

2.2. Coarse geometry of symmetric spaces and Euclidean buildings. Symmetric spaces and Euclidean buildings can be seen as leading examples of CAT(0) spaces. In fact, they seem to be the most rigid among all proper CAT(0) spaces [CM09][Lee]. We will recall their definition and basic properties.

2.2.1. Symmetric spaces. A *symmetric space* is a connected riemannian manifold M such that for all $x \in M$, there exists an isometry $\sigma_x \in \text{Isom}(M)$ such that $\sigma_x(x) = x$ and $T_x\sigma_x = -\text{Id}_{T_xM}$. It is said to be of *nonpositive curvature* if it has non-positive sectional curvature. If moreover it has no non-trivial Euclidean factor, M is called a *symmetric space of non-compact type*. All symmetric spaces of non-compact type can be obtained as coset spaces G/K , where G is a connected semi-simple Lie group with trivial center and no compact factors, and K is a maximal compact subgroup of G . The metric on G/K comes from the Killing form of $\text{Lie}(G)$. When M is a symmetric space of non-compact type, G is the identity component of the isometry group $\text{Isom}_0(M)$.

An important example of symmetric spaces of non-compact type is given by $\text{SL}_n(\mathbb{R})/\text{SO}_n(\mathbb{R})$. When $n = 2$, it corresponds to the hyperbolic plane \mathbb{H}^2 . More generally, if M is a symmetric space of non-compact type such that $\dim(\text{Isom}_0(M)) = n \geq 2$, then after rescaling the metrics of the de Rham factors of M by positive constants, M can be isometrically embedded in $\text{SL}_n(\mathbb{R})/\text{SO}_n(\mathbb{R})$ [Ebe96]. The symmetric space $\text{SL}_n(\mathbb{R})/\text{SO}_n(\mathbb{R})$ can also be seen as the collection of scalar products on \mathbb{R}^n , for which the unit ball has volume 1.

2.2.2. Euclidean buildings. Euclidean buildings are non-Archimedean analogues of symmetric spaces of non-compact type. In fact, to any semi-simple algebraic group over a local field, $\text{SL}_n(\mathbb{Q}_p)$ for example, we can associate a Euclidean building, called its Bruhat-Tits Building, on which the group acts isometrically in a very transitive way. Let us define Euclidean buildings in general. A general reference for buildings is [AB08], and for an introductory course see [Cap14].

Let $W < \text{Isom}(\mathbb{R}^n)$ be a discrete reflection group, i.e. a discrete subgroup of $\text{Isom}(\mathbb{R}^n)$ generated by orthogonal reflections through a collection \mathcal{H} of hyperplanes, called *walls*, and such that this collection is locally finite. The pattern determined by the set of walls defines a cellular decomposition Σ of \mathbb{R}^n , called a *Euclidean Coxeter complex*. The group W is called the *affine Weyl group* (or the *affine Coxeter group*) of the Euclidean Coxeter complex Σ . A *chamber* (or *alcove*) in that complex is defined as a connected component of $\mathbb{R}^n - \cup_{h \in \mathcal{H}} h$. The affine Weyl group W acts transitively on the set of chambers. The top-dimensional cells of Σ are the closures of chambers, also called *closed chambers*, which are compact if and only

if the group W acts cocompactly. A lower dimensional cell is the intersection of a closed chamber with a set of walls.

Definition 2.7. Let W be an affine Weyl group of \mathbb{R}^n , and Σ its associated Euclidean Coxeter complex. A *(discrete) Euclidean building* modeled on (Σ, W) is a cell complex B , which is covered by subcomplexes all isomorphic to Σ , called the *apartments* of B , and such that the following incidences properties hold:

- (1) Any two cells of B lie in some apartment.
- (2) Any two apartments, there is an isomorphism between them fixing their intersection pointwise.

Euclidean buildings are frequently called *Affine buildings* in the literature.

A Euclidean building is *thick* if each wall belongs to at least 3 half-apartments with disjoint interiors. If the affine Weyl group acts cocompactly on the apartments, then the chambers of the Euclidean Coxeter complex are polysimplices. This induces on the Euclidean building a structure of a polysimplicial complex. If the building is moreover irreducible, then this complex is a simplicial complex.

A Euclidean building always possesses a geometric realization, which is a complete CAT(0) metric space, and such that the restriction of its distance to each apartment is the Euclidean metric.

2.2.3. Model spaces and rank. Let S be a product of symmetric spaces of non-compact type and B a product of thick Euclidean buildings with cocompact affine Weyl group, with bounded geometry and no Euclidean factor. The spaces that we are going to consider are of the form $X = \mathbb{R}^n \times S \times B$. We will call such space a *model space*.

Definition 2.8. Let $X = \mathbb{R}^n \times S \times B$ be a model space. We define its *rank* as $\text{rank}(X) = \max\{k \in \mathbb{N} \mid \exists g : \mathbb{R}^k \rightarrow X \text{ an isometric embedding}\}$.

Example 2.9. For all $p \geq 1$, $\text{rank}(\mathbb{R}^p) = p$.

For all $n \geq 2$, $\text{rank}(\mathbb{H}^n) = 1$ and $\text{rank}(\text{SL}_n(\mathbb{R})/\text{SO}_n(\mathbb{R})) = \text{rank}(\text{BT}(\text{SL}_n(\mathbb{Q}_p))) = n - 1$, where $\text{BT}(\text{SL}_n(\mathbb{Q}_p))$ is the Bruhat-Tits building of $\text{SL}_n(\mathbb{Q}_p)$.

Remark 2.10. Note that the rank of a Euclidean building is equal to his dimension, and $\text{rank}(\mathbb{R}^n \times S \times B) = n + \text{rank}(S) + \dim(B)$.

2.2.4. Parallel sets. References for this subsection are [Ebe96],[Hel01] for symmetric spaces, and [KL97], [Lee] for Euclidean buildings.

Let X be a model space of rank k . A subspace of X is a *flat* if it is isometric to some Euclidean space. A *maximal flat* is a flat of dimension k , and flats of dimension 1 are the geodesics. A geodesic γ is said to be *regular* if it is contained in a unique maximal flat. Otherwise it is called *singular*. It is said to be *maximally singular* if it belongs to $(k - 1)$ half-flats of maximal dimension, with disjoint interiors. A *singular flat* is a flat which is the intersection of maximal flats.

Let X be a model space with no Euclidean factor, and let $F \subset X$ be a flat. If another flat F' is at finite Hausdorff distance from F then by the Flat Strip Theorem [BH13], since X is CAT(0), there exists a segment I such that the convex hull of the union $F \cup F'$ is

isometric to $F \times I$. In that case, the flats F and F' are called *parallel*. The flat F is a closed convex subset with extendible geodesics (i.e. each geodesic segment is contained in a bi-infinite geodesic in F), therefore we can consider its *parallel set* $P_X(F)$, which is the union of flats that are parallel to F . It is a closed convex subset of X isometric to a product

$$P_X(F) = F \times CS_X(F),$$

where $CS_X(F)$ is a model space. It is called the *cross section* of F . Moreover, if F is a singular flat then its cross section has no Euclidean factor and $\text{rank}(CS_X(F)) = \text{rank}(X) - \dim(F)$. The flats in X being just products of flats in its factors, and singular flats being products of singular flats in the factors, the cross section of a product is the product of cross sections (see [Lee] and [KL97]).

Example 2.11. *Let X be the Bruhat-Tits building associated to $\text{SL}_3(\mathbb{Q}_p)$ and let γ be a maximally singular geodesic. We then have that $CS_X(\gamma)$ is isometric to the $(p+1)$ -regular simplicial tree T_{p+1} , and*

$$P_X(\gamma) = \mathbb{R} \times T_{p+1}.$$

Example 2.12. *Let $Y = \text{SL}(3, \mathbb{R})/\text{SO}(3)$, and let γ be a maximally singular geodesic. We then have that $CS_Y(\gamma)$ is isometric to the hyperbolic plane \mathbb{H}^2 , and*

$$P_Y(\gamma) = \mathbb{R} \times \mathbb{H}^2.$$

In the proof of Theorem 1.10, we will need a uniform lower bound on the volume growth of cross sections of singular flats. It is obvious for symmetric spaces since cross sections of singular flats are symmetric spaces of lower rank, and there are only finitely many isometry classes. For Euclidean buildings we need the following result.

Proposition 2.13. *Let X be a thick Euclidean building with cocompact affine Weyl group. Then for every singular flat F , there exists $D = D(F) > 0$ such that the 3-regular tree of edge length D isometrically embeds into the cross section of F . This implies that there exists a uniform lower bound on the volume growth of cross sections of singular flats in X .*

Proof. Let X be such Euclidean building, and let $n = \dim X$. If $n = 1$, X is a tree and the singular flats are the vertices. Thus the cross sections are the whole building. Let $D > 0$ be the distance between vertices in an apartment. Since X is thick, it contains a 3-regular tree of edge length D .

Suppose that $n \geq 2$, and let F be a singular flat of dimension $k \leq n-1$. Let F' be a singular $(n-1)$ -flat (i.e. a wall) that contains F . Any $(n-1)$ -flat parallel to F' contains a k -flat parallel to F , therefore $P_X(F') \subset P_X(F)$, which implies that $\mathbb{R}^{n-1-k} \times CS_X(F') \subset CS_X(F)$. So it is enough to show it for $k = n-1$.

The affine Weyl group W is a semi-direct product $W = W_r \rtimes T$, where W_r is the associated finite Coxeter group and T is a group of translations generated by vectors orthogonal to some walls. Since it is cocompact, T is isomorphic to \mathbb{Z}^n . Let $F \subset X$ be a wall, and fix a generating set of T . Since the affine Weyl group is cocompact, there exists $D = D(F) > 0$ such that the walls parallel to F in an apartment are at distance an integer multiple of D , see also [KL97] Corollary 5.1.3. In particular, every wall parallel to F admits at least three parallel walls in X at distance D . Consider the graph G whose vertex set is the set of walls parallel to F , and whose edges are pairs of walls that differ by a translation by a generator of T in an apartment. If moreover the length of the edges is the distance in X between the

corresponding walls, then the graph G is isometric to the cross section of F , which is called the *wall tree* of F . It is a thick tree, see [Wei08] Chapter 10 and [KW14], and contains a subtree which is 3-regular of edge length D , obtained by only considering for each vertex two parallel walls at distance D .

Moreover, since there is only a finite number of singular flats up to translation, there is a uniform lower bound on the volume growth of such wall trees. \square

2.3. Metric currents. References for this section are [AK⁺00],[HKS22] and [Wen05].

Ambrosio and Kirchheim extended the classical theory of normal and integral currents developed by Federer and Fleming [FF60] to arbitrarily complete metric spaces.

Let (X, d) be a complete metric space and $k \geq 0$ and let $\mathcal{D}^k(X)$ denote the set of $(k + 1)$ -tuples (f, π_1, \dots, π_k) of Lipschitz functions on X with f bounded.

Definition 2.14. A k -dimensional metric current T on X is a multi-linear functional on $\mathcal{D}^k(X)$ satisfying the following properties:

- (1) If π_i^j converges point-wise to π_i as $j \rightarrow \infty$ and if $\sup_{i,j} \text{Lip}(\pi_i^j) < \infty$ then

$$T(f, \pi_1^j, \dots, \pi_k^j) \rightarrow T(f, \pi_1, \dots, \pi_k).$$

- (2) If $\{x \in X : f(x) \neq 0\}$ is contained in the union $\bigcup_{i=1}^k B_i$ of Borel sets B_i and if π_i is constant on B_i then

$$T(f, \pi_1, \dots, \pi_k) = 0.$$

- (3) There exists a finite Borel measure μ on X such that

$$(2.1) \quad |T(f, \pi_1, \dots, \pi_k)| \leq \prod_{i=1}^k \text{Lip}(\pi_i) \int_X |f| d\mu$$

for all $(f, \pi_1, \dots, \pi_k) \in \mathcal{D}^k(X)$.

The space of k -dimensional metric currents on X is denoted by $\mathbf{M}_k(X)$ and the minimal Borel measure μ satisfying (2.1) is called *mass* of T and written as $\|T\|$. Let us denote by $\mathbf{M}(T) = \|T\|(X)$ the *total mass* of T . The *support* of T is, by definition, the closed set $\text{spt}T$ of points $x \in X$ such that $\|T\|(B(x, r)) > 0$ for all $r > 0$.

Every function $g \in L^1_{loc}(\mathbb{R}^k)$ induces a current $\llbracket g \rrbracket \in \mathbf{M}_k(\mathbb{R}^k)$. Indeed, by Rademacher's theorem, Lipschitz functions are differentiable almost everywhere, so if $(f, \pi_1, \dots, \pi_k) \in \mathcal{D}^k(\mathbb{R}^k)$, $\llbracket g \rrbracket$ is defined by :

$$\llbracket g \rrbracket(f, \pi_1, \dots, \pi_k) := \int_{\mathbb{R}^k} g f \det \left(\frac{\partial \pi_i}{\partial x_j} \right) d\mathcal{L}^k$$

Which corresponds to the integration of the differential form $f d\pi_1 \wedge \dots \wedge d\pi_k$, weighted by f . Therefore, every Borel set $A \subset \mathbb{R}^k$ induces a current $\llbracket A \rrbracket := \llbracket \chi_A \rrbracket$.

The restriction of $T \in \mathbf{M}_k(X)$ to a Borel set $A \subset X$ is given by

$$(T \llcorner A)(f, \pi_1, \dots, \pi_k) := T(f \chi_A, \pi_1, \dots, \pi_k).$$

This expression is well-defined since T can be extended to a functional on tuples for which the first argument lies in $L^\infty(X, \|T\|)$.

The boundary of $T \in \mathcal{D}_k(X)$ is defined by analogy with Stokes formula:

$$\partial T(f, \pi_1, \dots, \pi_{k-1}) := T(1, f, \pi_1, \dots, \pi_{k-1}).$$

It is clear that ∂T satisfies conditions (1) and (2) in the above definition. If ∂T also has finite mass (condition (3)) then T is called a *normal current*. The space of normal currents is denoted by $\mathbf{N}_k(X)$.

The push-forward of $T \in \mathbf{M}_k(X)$ under a Lipschitz map φ from X to another complete metric space Y is given by

$$\varphi_{\#}T(g, \tau_1, \dots, \tau_k) := T(g \circ \varphi, \tau_1 \circ \varphi, \dots, \tau_k \circ \varphi),$$

for $(g, \tau_1, \dots, \tau_k) \in \mathcal{D}^k(Y)$. This defines a k -dimensional current on Y .

Let \mathcal{H}^k denote the Hausdorff k -dimensional measure. An \mathcal{H}^k -measurable set $A \subset X$ is said to be *countably \mathcal{H}^k -rectifiable* if there exist Lipschitz maps $f_i : B_i \rightarrow X$ from subsets $B_i \subset \mathbb{R}^k$ such that

$$\mathcal{H}^k(A - \bigcup f_i(B_i)) = 0.$$

Definition 2.15. A current $T \in \mathbf{M}_k(X)$ with $k \geq 1$ is said to be *rectifiable* if

- (1) $\|T\|$ is concentrated on a countably \mathcal{H}^k -rectifiable set and
- (2) $\|T\|$ vanishes on \mathcal{H}^k -negligible sets.

T is called *integer rectifiable* if, in addition, the following property holds:

- (3) For any Lipschitz map $\varphi : X \rightarrow \mathbb{R}^k$ and any open set $U \subset X$ there exists $\theta \in L^1(\mathbb{R}^k, \mathbb{Z})$ such that

$$\varphi_{\#}(T \llcorner U) = \llbracket \theta \rrbracket.$$

We denote the space of rectifiable currents by $\mathcal{R}_k(X)$, and $\mathcal{I}_k(X)$ for integer rectifiable currents. Integer rectifiable normal currents are called *integral currents*, denoted by $\mathbf{I}_k(X)$. An element $T \in \mathbf{I}_k(X)$ is called a *cycle* $\partial T = 0$.

The main motivation for using currents instead of Lipschitz chains is the following Slicing Theorem due to Ambrosio–Kirchheim [AK⁺00] that will play a crucial role in this paper.

Theorem 2.16. *Let be $T \in \mathbf{I}_k(X)$ and π a Lipschitz function on X . Then there exists for almost every $r \in \mathbb{R}$ an integral current $\langle T, \pi, r \rangle \in \mathbf{I}_{k-1}(X)$ with the following properties:*

- (1) $\langle T, \pi, r \rangle = \partial(T \llcorner \{\pi \leq r\}) - (\partial T) \llcorner \{\pi \leq r\}$,
- (2) $\|\langle T, \pi, r \rangle\|$ and $\|\partial \langle T, \pi, r \rangle\|$ are concentrated on $\pi^{-1}(\{r\})$,
- (3) $\mathbf{M}(\langle T, \pi, r \rangle) \leq \text{Lip}(\pi) \frac{d}{dr} \mathbf{M}(T \llcorner \{\pi \leq r\})$, which is just a reformulation of the co-area formula.

2.4. Filling functions. The basic idea of a filling function is to measure the difficulty of filling a boundary of a given size. There are several ways to make this rigorous, depending on the type of boundary and the type of filling. We will use the Homological filling, as in [Gro93][Leu14], which consists on filling Lipschitz cycles by Lipschitz chains, instead of the homotopical filling where we fill spheres by balls.

An *integral Lipschitz d -chain* in a complete metric space X is a finite linear combination $\Sigma = \sum_i a_i \sigma_i$, with $a_i \in \mathbb{Z}$, of Lipschitz maps $\sigma_i : \Delta^k \rightarrow X$ from the Euclidean k -dimensional simplex Δ^k to X . We will often call this simply a *Lipschitz k -chain*. The boundary operator is defined as in the case of singular chains.

Note that by the push-forward of metric currents, every Lipschitz chain in X induces an integral current. Indeed, if $\sigma_i : \Delta^k \rightarrow X$ is Lipschitz, then $f_{\#}(\llbracket \Delta^k \rrbracket)$ is an integral k -current in X . We define the *k -volume* of σ_i as the total mass of $f_{\#}(\llbracket \Delta^k \rrbracket)$:

$$\text{Vol}_k \sigma_i = \mathbf{M}(f_{\#}(\llbracket \Delta^k \rrbracket)).$$

Note that if X is a riemannian manifold then, by Rademacher's theorem, σ_i is differentiable almost everywhere and $\text{Vol}_k \sigma_i$ is also equal to the integral of the magnitude of its Jacobian. Note also that it is not necessarily equal to the volume of the image (its hausdorff measure), unless the map is injective, because the volume is counted with multiplicity.

We then define the *k -volume* of a Lipschitz k -chain as

$$\text{Vol}_k^X \Sigma := \sum_i |a_i| \text{vol}_k \sigma_i.$$

We wish to measure the difficulty to fill Lipschitz k -cycles by Lipschitz $(k+1)$ -chains.

To ensure that such a filling exists, the space considered must be k -connected, i.e. all homotopy groups $\pi_k(X)$ are trivial (which is the case for $\text{CAT}(0)$ spaces) hence the corresponding homology groups are also trivial.

More precisely, for an integral Lipschitz k -cycle Σ in k -connected space we define its *filling volume*

$$\text{FillVol}_{k+1}^X(\Sigma) := \inf\{\text{Vol}_{k+1}(\Omega) \mid \Omega \text{ is a Lipschitz } (k+1)\text{-chain with } \partial\Omega = \Sigma\}.$$

The *$(k+1)$ -dimensional filling function* of X is then given by

$$\text{FV}_{k+1}^X(l) := \sup\{\text{FillVol}_{k+1}^X(\Sigma) \mid \Sigma \text{ is a Lipschitz } k\text{-cycle in } X \text{ with } \text{Vol}_k^X(\Sigma) \leq l\}.$$

We are only interested in the asymptotic behaviour of these filling functions. We have that if X and Y are two k -connected manifolds or simplicial complexes which are quasi-isometric, then

$$\text{FV}_k^X(l) \sim \text{FV}_k^Y(l).$$

The equivalence relation on functions $\mathbb{R} \rightarrow \mathbb{R}$ is define as follows: we write $f \lesssim g$ if there is a constant $C > 0$ such that $f(x) \leq Cg(Cx + C) + Cx + C$. We write $f \sim g$ and say that they are *asymptotically equivalent* if $f \lesssim g$ and $g \lesssim f$.

Example 2.17.

- A hyperbolic space $X = \mathbb{H}^n$ satisfies for all $2 \leq k \leq n$, $\text{FV}_X^k(l) \sim l$.
- A Euclidean space $X = \mathbb{R}^n$ satisfies for all $2 \leq k \leq n$, $\text{FV}_X^k(l) \sim l^{\frac{k}{k-1}}$.
- More generally, Wenger [Wen05] showed that a complete, simply connected $\text{CAT}(0)$ space satisfies $\text{FV}_k^X(l) \lesssim l^{\frac{k}{k-1}}$ for all dimensions k .

A key theorem that we will use is due to Leuzinger and says that the filling functions can detect the rank of symmetric spaces:

Theorem 2.18 ([Leu14]).

Let $X = \mathbb{R}^d \times S \times B$ be a model space. Then

(i) X has Euclidean filling functions below the rank:

$$\text{FV}_k^X(l) \sim l^{\frac{k}{k-1}} \quad \text{if } 2 \leq k \leq \text{rank}(X);$$

(ii) X has linear filling functions above the rank:

$$\text{FV}_k^X(l) \sim l \quad \text{if } \text{rank}(X) < k \leq \dim(X).$$

When X is a symmetric space of non-compact type, this result has been correctly asserted by Gromov in [Gro93], section 5.D, and proposed a possible different proof for the upper bound by projecting the cycle to a maximal flat, see [Leu14] for more details. The proof of Leuzinger uses a different approach by projecting to a suitable horosphere.

Note that by the push-forward of metric currents, every Lipschitz chain in X induces an integral current. Moreover, if σ is a Lipschitz k -chain $\text{Vol}_X^k(\sigma) = \mathbf{M}(\sigma)$. Therefore, the chains that we will consider will be seen as currents, so it is convenient to define filling functions in this setting.

Definition 2.19. Let (X, d) be a complete metric space and $k \geq 0$ an integer. For a k -cycle $\Sigma \in \mathbf{I}_k(X)$, we define its $(k+1)$ -dimensional current-filling volume to be

$$\text{FillVol}_{k+1}^{X, \text{cr}}(\Sigma) := \inf\{\mathbf{M}(\Omega) \mid \Omega \in \mathbf{I}_{k+1} \text{ with } \partial\Omega = \Sigma\}.$$

Since every Lipschitz chain induces a metric current, it is clear that if Σ is a Lipschitz k -cycle seen in $\mathbf{I}_k(X)$, then $\text{FillVol}_{k+1}^{X, \text{cr}}(\Sigma) \leq \text{FillVol}_{k+1}^X(\Sigma)$.

2.5. Connect the dots argument.

Definition 2.20. A metric space X is *uniformly contractible* if there is a function $\phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that for any $x \in X$ and $r > 0$, the ball $B_X(x, r)$ is contractible in the ball $B_X(x, \phi(r))$.

Lemma 2.21. Let $f : X \rightarrow Y$ be a large-scale Lipschitz map, X a CW complex where the size of the cells is globally bounded, and Y is uniformly contractible. Then f is a bounded distance from a Lipschitz map $g : X \rightarrow Y$. That is, $\sup_{x \in X} d_Y(f(x), g(x)) \leq C$ for some constant $C > 0$.

Proof. Let us prove it by induction on the dimension of the cells of X , as suggested in [BW97]. Let $D = \sup_{\sigma} \text{diam}(\sigma)$ over all the cells of X , and let (λ, c) be the large-scale Lipschitz constants of f . For any vertex $v \in X^{(0)}$, define $g(v) := f(v)$. Given an edge $e = (u, v) \in X^{(1)}$, $f(\partial e) \subset B_Y(f(u), \lambda D + c)$, so $f(\partial e)$ can be filled inside of $B_Y(f(u), \phi(\lambda D + c))$. Let γ be such filling, and define $g(e) := \gamma$ in a continuous way that respects the boundary. Then g , defined at this stage on $X^{(1)}$, is uniformly close to f on $X^{(1)}$. Indeed, if $x \in X^{(1)}$ is in an edge $e = (u, v)$ so

$$\begin{aligned} d_Y(f(x), g(x)) &\leq d_Y(f(x), f(u)) + d_Y(g(u), g(x)) \\ &\leq \lambda D + c + \phi(\lambda D + c) \end{aligned}$$

Suppose that g is defined on $X^{(k)}$. Let σ be a $(k+1)$ -cell in X , and take $x \in \sigma$. $f(\partial\sigma) \subset B_Y(f(x), \lambda D + c)$, so it can be filled inside of $B_Y(f(x), \phi(\lambda D + c))$. Let Ω be such filling,

and define $g(\sigma) := \Omega$ in a continuous way that respects the boundary. Same as before, we can easily check that g is a bounded distance from f . \square

The domains that we will consider are always geodesic, so the coarse embeddings are large-scale Lipschitz. Since all our model spaces are $\text{CAT}(0)$, they are uniformly contractible. Moreover, they admit a CW structure where the size of the cells is globally bounded. Therefore we can always assume that our coarse embeddings are actually Lipschitz.

3. THE DOMAIN X IS A PRODUCT OF SPACES WITH EXPONENTIAL GROWTH

The goal of this section is to prove Theorem 1.9.

3.1. Preliminary lemmas. Let us give some useful lemmas first. One crucial property that we are going to use is that the ε -volume is coarsely preserved by coarse embeddings:

Lemma 3.1. *Let X and Y be metric spaces with bounded geometry, $f : X \rightarrow Y$ a coarse embedding, and let $\varepsilon > 0$. There exist $\alpha, \beta > 0$ such that for all $A \subset X$ with $\text{Vol}_X^\varepsilon(A) < \infty$,*

$$\alpha \text{Vol}_X^\varepsilon(A) \leq \text{Vol}_Y^\varepsilon(f(A)) \leq \beta \text{Vol}_X^\varepsilon(A).$$

Proof. Let f be such coarse embedding with control functions ρ_- and ρ_+ .

- Let $n \in \mathbb{N}$ and $x_1, \dots, x_n \in X$ such that

$$A \subset B_X(x_1, \varepsilon) \cup \dots \cup B_X(x_n, \varepsilon).$$

Then

$$f(A) \subset f(B_X(x_1, \varepsilon)) \cup \dots \cup f(B_X(x_n, \varepsilon)) \subset B_Y(f(x_1), \rho^+(\varepsilon)) \cup \dots \cup B_Y(f(x_n), \rho^+(\varepsilon)).$$

Since Y has bounded geometry, there exists $\beta = \text{Vol}_X^\varepsilon(B(\rho^+(\varepsilon))) \in \mathbb{N}$ such that every ball of radius $\rho^+(\varepsilon)$ is in the union of β balls of radius ε . Therefore $f(A)$ is in the union of $\beta \times n$ balls of radius ε . Since its volume is the minimum among such number of balls, we have

$$\text{Vol}_Y^\varepsilon(f(A)) \leq \beta n.$$

By taking n to be minimal, we have

$$\text{Vol}_Y^\varepsilon(f(B_X(x, R))) \leq \beta \text{Vol}_X^\varepsilon(B_X(x, R)).$$

- Now let $n \in \mathbb{N}$ and $y_1, \dots, y_n \in Y$ such that

$$f(B_X(x, R)) \subset B_Y(y_1, \varepsilon) \cup \dots \cup B_Y(y_n, \varepsilon).$$

Then

$$B_X(x, R) \subset f^{-1}(f(B_X(x, R))) \subset f^{-1}(B_Y(y_1, \varepsilon)) \cup \dots \cup f^{-1}(B_Y(y_n, \varepsilon)).$$

For any $i = 1, \dots, n$, $f^{-1}(B_Y(y_i, \varepsilon))$ has diameter less than $\rho_-^{-1}(2\varepsilon) : \forall a, b \in f^{-1}(B_Y(y_i, \varepsilon))$, $d_Y(f(a), f(b)) \leq 2\varepsilon$, so $d_X(a, b) \leq \rho_-^{-1}(2\varepsilon)$. For any $i = 1, \dots, n$ such that $f^{-1}(B_Y(y_i, \varepsilon))$ is non-empty, take some x_i in it. So $f^{-1}(B_Y(y_i, \varepsilon)) \subset B_X(x_i, \rho_-^{-1}(2\varepsilon))$.

Let us denote $D = \rho_-^{-1}(2\varepsilon)$. Therefore

$$B_X(x, R) \subset B_X(x_1, D) \cup \dots \cup B_X(x_n, D).$$

X has bounded geometry, so there exists $\gamma = \text{Vol}_X^\varepsilon(B(D)) \in \mathbb{N}$ such that every ball of radius D is in the union of γ balls of radius D . So $B_X(x, R)$ is in the union of $\gamma \times n$ balls of radius ε . Since its volume is the minimum among such number of balls, we have have

$$\text{Vol}_X^\varepsilon(B_X(x, R)) \leq \gamma n.$$

By taking n to be minimal and by denoting $\alpha = \gamma^{-1}$, we have

$$\text{Vol}_Y^\varepsilon(f(B_X(x, R))) \geq \alpha \text{Vol}_X^\varepsilon(B_X(x, R)).$$

□

Let us also prove the following lemma.

Lemma 3.2. *Let X be a metric space with bounded geometry, and let $\varepsilon, \delta > 0$. If $A \subset X$ with $\text{Vol}_X^\varepsilon(A) < \infty$, then*

$$\text{Vol}_X^\varepsilon(N_\delta(A)) \leq \text{Vol}_X^\varepsilon(B_X(\delta + \varepsilon)) \times \text{Vol}_X^\varepsilon(A).$$

Proof. X has bounded geometry, so there exist $n \in \mathbb{N}$ and $x_1, \dots, x_n \in X$ such that $A \subset B_X(x_1, \varepsilon) \cup \dots \cup B_X(x_n, \varepsilon)$.

Let $z \in N_\delta(A)$, i.e. there exists $a \in A$ such that $d_X(z, a) \leq \delta$. So there exists $i \in \{1, \dots, n\}$ such that $d_X(z, x_i) \leq \delta + \varepsilon$, therefore $N_\delta(A) \subset B_X(x_1, \varepsilon + \delta) \cup \dots \cup B_X(x_n, \varepsilon + \delta)$.

There exists $p = \text{Vol}_X^\varepsilon(B_X(\delta + \varepsilon))$ such that every ball of radius $\varepsilon + \delta$ can be covered by p balls of radius ε . Therefore $N_\delta(A)$ can be covered by $p \times n$ balls of radius ε , so

$$\text{Vol}_X^\varepsilon(N_\delta(A)) \leq \text{Vol}_X^\varepsilon(B_X(\delta + \varepsilon)) \times n.$$

By taking n to be minimal, we have the result. □

We will also need the following result.

Lemma 3.3. *Let X be a complete metric space, and $x, y \in X$. Let $T \in \mathbf{I}_1(X)$ such that $\partial T = \llbracket x \rrbracket - \llbracket y \rrbracket$, then*

$$\mathbf{M}(T) \geq d_X(x, y),$$

and

$$\text{FillVol}_1^{X, \text{cr}}(\llbracket x \rrbracket - \llbracket y \rrbracket) \geq d_X(x, y).$$

If moreover there is a geodesic segment from x to y , then

$$\text{FillVol}_1^{X, \text{cr}}(\llbracket x \rrbracket - \llbracket y \rrbracket) = d_X(x, y).$$

Proof. $\partial T = \llbracket x \rrbracket - \llbracket y \rrbracket$ means that for all $f : X \rightarrow \mathbb{R}$ Lipschitz and bounded, we have

$$\partial T(f) = f(x) - f(y).$$

By definition, we have $\partial T(f) = T(1, f)$. So $T(1, f) = f(x) - f(y)$.

T being a 1-current, there exists a finite Borel measure μ on X such that for all $g : X \rightarrow \mathbb{R}$ Lipschitz and bounded and for all $f : X \rightarrow \mathbb{R}$ Lipschitz

$$|T(g, f)| \leq \text{Lip}(f) \int_X |g(x)| d\mu(x).$$

Let $\mu = \|T\|$ be the minimal Borel measure. In particular, we have that $\mu(X) = \|T\|(X) = \mathbf{M}(T)$.

For $g = 1$ and for all $f : X \rightarrow \mathbb{R}$ Lipschitz, we have

$$|f(x) - f(y)| = |T(1, f)| \leq \text{Lip}(f) \int_X |1(x)| d\mu(x) = \text{Lip}(f) \mathbf{M}(T).$$

So $\mathbf{M}(T)$ satisfies : $\forall f : X \rightarrow \mathbb{R}$ Lipschitz, $|f(x) - f(y)| \leq \text{Lip}(f) \mathbf{M}(T)$. This implies that

$$\sup_{f \text{ 1-Lip}} \frac{|f(x) - f(y)|}{\text{Lip}(f)} \leq \mathbf{M}(T).$$

Consider the map $d_y : X \rightarrow \mathbb{R}$, $\forall z \in X$, $d_y(z) := d_X(y, z)$. It is a 1-Lipschitz map that satisfies:

$$\frac{|d_y(x) - d_y(y)|}{\text{Lip}(f)} = d_X(x, y).$$

Therefore

$$\sup_{f \text{ 1-Lip}} \frac{|f(x) - f(y)|}{\text{Lip}(f)} = d_X(x, y),$$

and

$$\mathbf{M}(T) \geq d_X(x, y).$$

By taking the infimum of the mass of all such 1-currents,

$$\text{FillVol}_1^{X, \text{cr}}(\llbracket x \rrbracket - \llbracket y \rrbracket) \geq d_X(x, y).$$

If there is a geodesic segment γ from y to x , consider the Lipschitz 1-chain realized by γ . Then $\partial\gamma = \llbracket x \rrbracket - \llbracket y \rrbracket$ and $\mathbf{M}(\gamma) = d_X(x, y)$, and the infimum is attained. \square

3.2. Proof of Theorem 1.9. Let us prove a more general result where we do not actually require the k -filling function of the target space to be linear, but we only need a weaker condition on the filling of the boundaries of some k -rectangles.

Theorem 3.4. *Let $X = X_1 \times \dots \times X_k$ be a product of geodesic metric spaces with bounded geometry, where $k \in \mathbb{N}^*$, and let Y be a uniformly contractible complete metric space with at most exponential growth. Let $f : X \rightarrow Y$ be a large-scale Lipschitz map. If*

- for all $i = 1, \dots, k$, X_i has exponential growth,
- there exists a sublinear function ϕ such that for every $d > 0$ big enough and every k -dimensional rectangle R_d , whose sides are geodesic segments in the X_i 's, with side lengths l_1, \dots, l_k that satisfies $l_1 = d, l_2 \leq d^{1/2}, \dots, l_k \leq d^{1/k}$, we have

$$\text{FillVol}_k^{Y, \text{cr}}(f(\partial R_d)) \leq \phi(d^{H_k}),$$

where $H_k = 1 + \frac{1}{2} + \dots + \frac{1}{k}$, then f is not a coarse embedding.

Remark 3.5. This condition means that the filling of the image of the boundary of such rectangles in Y is sublinear when compared to their filling in X .

Remark 3.6. This result can be seen as a generalization of the fact, for $k = 1$, that a space with exponential growth cannot be coarsely embedded with a sublinear ρ^+ in a space with at most exponential growth, because the second condition means that f admits a sublinear ρ^+ . Indeed, R_d is just a geodesic segment in X . Let x and y be its extremities. So the second condition can be rewritten as

$$\text{FillVol}_1^{Y, \text{cr}}(\llbracket f(x) \rrbracket - \llbracket f(y) \rrbracket) \leq \phi(d_X(x, y)).$$

By lemma 3.3, we have

$$d_Y(f(x), f(y)) \leq \text{FillVol}_1^{Y, \text{cr}}(\llbracket f(x) \rrbracket - \llbracket f(y) \rrbracket).$$

Therefore

$$d_Y(f(x), f(y)) \leq \phi(d_X(x, y)).$$

Proof. Let us prove the theorem by induction on k .

• For $k = 1$, let X be a metric space with exponential growth, and suppose that such a coarse embedding $f : X \rightarrow Y$ exists for a sublinear ϕ .

Let $x \in X$ and $R > 0$. Then

$$f(B_X(x, R)) \subset B_Y(f(x), \phi(R)).$$

Let $\varepsilon > 0$. By taking the ε -volumes :

$$\text{Vol}_Y^\varepsilon(f(B_X(x, R))) \leq \text{Vol}_Y^\varepsilon(B_Y(f(x), \phi(R))).$$

Y has at most exponential growth so there exists $\lambda \geq 0$ such that for all $y \in Y$ and for all $R > 0$,

$$\text{Vol}_Y^\varepsilon(B_Y(y, R)) \leq e^{\lambda R}.$$

But, on one hand f is a coarse embedding, so by lemma 3.1 it preserves the volume coarsely. In particular there exists $\alpha > 0$ such that $\forall x \in X$

$$\alpha \text{Vol}_X^\varepsilon(B_X(x, R)) \leq \text{Vol}_Y^\varepsilon(f(B_X(x, R))).$$

On the other hand, X has exponential growth, so there exists $\gamma > 0$ such that for all $x \in X$ and for all $R > 0$

$$\text{Vol}_X^\varepsilon(B_X(x, R)) \geq e^{\gamma R}.$$

So we finally get

$$\alpha e^{\lambda R} \leq \alpha \text{Vol}_X^\varepsilon(B_X(x, R)) \leq \text{Vol}_Y^\varepsilon(f(B_X(x, R))) \leq \text{Vol}_Y^\varepsilon(B_Y(f(x), \phi(R))) \leq e^{\lambda \phi(R)}.$$

Which is not possible for a sublinear function ϕ .

Note that for $k = 1$ we did not require X to be geodesic.

Before proving it for a $k \geq 2$, and just for the sake of clarity, let us first prove it for $k = 2$. Note that it already contains most of the conceptual difficulties of the general case.

• Let $X = X_1 \times X_2$ be a product of two geodesic metric spaces of exponential growth and Y be a metric space with at most exponential growth. Suppose that $f : X \rightarrow Y$ is a coarse embedding that satisfies the second condition, i.e. there exists a sublinear function ϕ such that for every $d > 0$ and every 2-dimensional rectangle R_d , whose sides are geodesic segments in the X_i 's, of side lengths l_1, l_2 with $l_1 = d$ and $l_2 \leq d^{1/2}$, we have

$$\text{FillVol}_2^{Y, \text{cr}}(f(\partial R_d)) \leq \phi(d\sqrt{d}).$$

Following the case $k = 1$, for all $x_2 \in X_2$, the copy $X_1 \times \{x_2\}$ cannot be sent sublinearly. Therefore, if we fix some $x_2 \in X_2$, there exist $\lambda > 0$ and two sequences $(a_n)_{n \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}}$ in $X_1 \times \{x_2\}$ such that

$$d_n := d_X(a_n, b_n) \xrightarrow{n \rightarrow +\infty} +\infty \quad \text{and} \quad d_Y(f(a_n), f(b_n)) \geq \lambda d_X(a_n, b_n).$$

Let $n \in \mathbb{N}$ and take $x'_2 \in X_2$ such that $d_{X_2}(x_2, x'_2) \leq \sqrt{d_n}$. Consider $a'_n = (\text{proj}_{X_1}(a_n), x'_2)$, and $b'_n = (\text{proj}_{X_1}(b_n), x'_2)$.

Let c_1 be a geodesic segment in X_1 going from $\text{proj}_{X_1}(a_n)$ to $\text{proj}_{X_1}(b_n)$, and c_2 be a geodesic segment in X_2 from x_2 to x'_2 .

Now consider the four following geodesic segments in X :

$$\gamma_1 = (c_1, x_2), \quad \gamma_2 = (\text{proj}_{X_1}(b_n), c_2), \quad \gamma_3 = (-c_1, x_2), \quad \gamma_4 = (\text{proj}_{X_1}(a_n), -c_2).$$

By concatenating them, i.e. taking their formal sum as Lipschitz chains, we get a Lipschitz 1-cycle that we will denote R_n .

The second condition on the filling volume implies that

$$\text{FillVol}_2^{Y, \text{cr}}(f(R_n)) \leq \phi \left(d_n \sqrt{d_n} \right).$$

This implies that there exists $S_n \in \mathbf{I}_2(Y)$ in Y such that $\partial S_n = f(R_n)$ and

$$\mathbf{M}(S_n) \leq \phi \left(d_n \sqrt{d_n} \right).$$

Now consider the 1-Lipschitz map $\pi : Y \rightarrow \mathbb{R}$, $\pi(y) = d_Y(y, f(\gamma_1))$.

By the Slicing Theorem, we have that for a.e. $t \in \mathbb{R}$, there exists $\langle S_n, \pi, t \rangle \in \mathbf{I}_1(Y)$ such that $\langle S_n, \pi, t \rangle = \partial(S_n \llcorner \{\pi \leq t\}) - (\partial S_n) \llcorner \{\pi \leq t\}$, and by integrating the co-area formula over the distance t , we have

$$\begin{aligned} \mathbf{M}(\langle S_n, \pi, t \rangle) &\leq \frac{d}{dt} \mathbf{M}(S_n \llcorner \{\pi \leq t\}), \\ \int_0^{+\infty} \mathbf{M}(\langle S_n, \pi, t \rangle) dt &\leq \int_0^{+\infty} \frac{d}{dt} \mathbf{M}(S_n \llcorner \{\pi \leq t\}), \\ \int_0^{+\infty} \mathbf{M}(\langle S_n, \pi, t \rangle) dt &\leq \mathbf{M}(S_n). \end{aligned}$$

Since $\mathbf{M}(S_n) \leq \phi \left(d_n \sqrt{d_n} \right)$, we get

$$(3.1) \quad \int_0^D \mathbf{M}(\langle S_n, \pi, t \rangle) dt \leq \int_0^{+\infty} \mathbf{M}(\langle S_n, \pi, t \rangle) dt \leq \phi \left(d_n \sqrt{d_n} \right),$$

where $D = d_Y(f(\gamma_1), f(\gamma_3))$. However, for a.e. $t \in]0, D[$, $\mathbf{M}(\langle S_n, \pi, t \rangle)$ cannot be too small since the current $\langle S_n, \pi, t \rangle$ almost gives a filling for the 0-cycle $\llbracket f(b_n) \rrbracket - \llbracket f(a_n) \rrbracket$.

Claim 3.7. For n big enough and for a.e. $t \in]0, D[$, $\mathbf{M}(\langle S_n, \pi, t \rangle) \geq \frac{\lambda}{2} d_n$.

Proof. For a.e. $t \in]0, D[$,

$$\begin{aligned} \partial(S_n) \llcorner \{\pi \leq t\} &= \partial(S_n) \llcorner \{\pi = 0\} + \partial(S_n) \llcorner \{0 < \pi \leq t\} \\ &= f(\gamma_1) + H_t. \end{aligned}$$

Where H_t is the 1-current $\partial(S_n) \llcorner \{0 < \pi \leq t\}$.

Since $\|\langle S_n, \pi, t \rangle\|$ is concentrated on $\pi^{-1}(\{t\})$,


$$\partial(S_n \llcorner \{\pi \leq t\}) = \langle S_n, \pi, t \rangle + f(\gamma_1) + H_t.$$

Which means that $\langle S_n, \pi, t \rangle + f(\gamma_1) + H_t$ is a 1-current that is actually a cycle.

So $-(\langle S_n, \pi, t \rangle + H_t)$ is a 1-chain that fills the 0-cycle $\partial f(\gamma_1) = f(\partial \gamma_1) = \llbracket f(b_n) \rrbracket - \llbracket f(a_n) \rrbracket$.

Therefore

$$\mathbf{M}(-(\langle S_n, \pi, t \rangle + H_t)) \geq \text{FillVol}_1^{Y, \text{cr}}(\llbracket f(b_n) \rrbracket - \llbracket f(a_n) \rrbracket).$$


 FIGURE 1. The cycle R_n in X , and a slice of the filling of its image in Y

Since, by the lemma 3.3 , $\text{FillVol}_1^{Y, \text{cr}}(\llbracket f(b_n) \rrbracket - \llbracket f(a_n) \rrbracket) \geq d_Y(f(a_n), f(b_n))$, we have

$$\mathbf{M}(-\langle S_n, \pi, t \rangle + H_t) \geq \lambda d_n.$$

So

$$\mathbf{M}(\langle S_n, \pi, t \rangle) + \mathbf{M}(H_t) \geq \lambda d_n.$$

Note that

$$\begin{aligned} H_t &= \partial S_n \llcorner \{0 < \pi \leq t\} \\ &= (f(\gamma_2) + f(\gamma_4)) \llcorner \{0 < \pi \leq t\}, \end{aligned}$$

because $0 < \pi < D$. So

$$H_t = f(\gamma_2) \llcorner \{0 < \pi \leq t\} + f(\gamma_4) \llcorner \{0 < \pi \leq t\}.$$

By taking the mass

$$\begin{aligned} \mathbf{M}(H_t) &\leq \mathbf{M}(f(\gamma_2) \llcorner \{0 < \pi \leq t\}) + \mathbf{M}(f(\gamma_4) \llcorner \{0 < \pi \leq t\}) \\ &\leq \mathbf{M}(f(\gamma_2)) + \mathbf{M}(f(\gamma_4)) \\ &\leq \text{Lip}(f)(\mathbf{M}(\gamma_2) + \mathbf{M}(\gamma_4)). \end{aligned}$$

Since the γ'_i 's are Lipschitz chains, their mass is equal to their volume. So $\mathbf{M}(\gamma_2) = \mathbf{M}(\gamma_4) \leq \sqrt{d_n}$, and $\mathbf{M}(H_t) \leq 2\text{Lip}(f)\sqrt{d_n}$.

Therefore there exists $N \in \mathbb{N}$ such that for all $n \geq N$ and for a.e. $t \in]0, D[$,

$$\mathbf{M}(\langle S_n, \pi, t \rangle) \geq \lambda d_n - 2\text{Lip}(f)\sqrt{d_n} \geq \frac{\lambda}{2}d_n. \quad \square$$

Thus by (3.1),

$$\begin{aligned} \phi(d_n\sqrt{d_n}) &\geq \int_0^D \mathbf{M}(\langle S_n, \pi, t \rangle) dt \\ &\geq D \frac{\lambda}{2}d_n. \end{aligned}$$

Which implies that

$$D \leq \frac{2\phi(d_n\sqrt{d_n})}{\lambda d_n}.$$

Let us denote $\psi(\sqrt{d_n}) = \frac{2\phi(d_n\sqrt{d_n})}{\lambda d_n}$. Note that ψ is sublinear.

Since $D = d_Y(f(\gamma_1), f(\gamma_3))$, the last inequality implies that there exists $z \in X$ such that $z \in \gamma_3$ and $d_Y(f(\gamma_1), f(z)) \leq \psi(\sqrt{d_n})$.

But $z \in \gamma_3$ implies that $d_X(z, \gamma_1) = d_X(x_2, x'_2)$.

By doing this process for all $n \geq N$ and all $x'_2 \in B_{X_2}(x_2, \sqrt{d_n})$, we get subsets $C_n \subset X$ that projects onto $B_{X_2}(x_2, \sqrt{d_n})$, i.e.

$$B_{X_2}(x_2, \sqrt{d_n}) \subset \text{proj}_{X_2}(C_n),$$

and such that

$$(3.2) \quad f(C_n) \subset N_{\psi(\sqrt{d_n})}(f(\gamma_1)),$$

since $\forall z \in C_n$, $d_Y(f(z), f(\gamma_1)) \leq \psi(\sqrt{d_n})$.




FIGURE 2. The subsets C_n in X that are highly compressed by f

The projection onto X_2 is 1-Lipschitz, so if we fix $\varepsilon > 0$,

$$\text{Vol}_X^\varepsilon(C_n) \geq \text{Vol}_X^\varepsilon(\text{proj}_{X_2}(C_n)) \geq \text{Vol}_{X_2}^\varepsilon\left(B_{X_2}\left(x_2, \sqrt{d_n}\right)\right).$$

X_2 has exponential growth, so there exists $\alpha > 0$ such that $\forall R > 0$, $\text{Vol}_{X_2}^\varepsilon(S_{X_2}(R)) \geq e^{\alpha R}$. So

$$\text{Vol}_X^\varepsilon(C_n) \geq e^{\alpha\sqrt{d_n}}.$$

While the equation (3.2) implies that

$$\text{Vol}_Y^\varepsilon(f(C_n)) \leq \text{Vol}_Y^\varepsilon(N_{\psi(\sqrt{d_n})}(f(\gamma_1))).$$

But, like we saw for the case $k = 1$, f coarsely preserves volumes, i.e. there exist $\delta, \delta' > 0$ such that

$$\delta \text{Vol}_X^\varepsilon(C_n) \leq \text{Vol}_Y^\varepsilon(f(C_n)) \leq \delta' \text{Vol}_X^\varepsilon(C_n).$$

Also, by lemma 3.2

$$\text{Vol}_Y^\varepsilon(N_{\psi(\sqrt{d_n})}(f(\gamma_1))) \leq \text{Vol}_Y^\varepsilon\left(B_Y(\varepsilon + \psi(\sqrt{d_n}))\right) \times \text{Vol}_Y^\varepsilon(f(\gamma_1)).$$

Y has at most exponential growth, so there exists $\beta > 0$ such that for all $R > 0$, $\text{Vol}_Y^\varepsilon(B_Y(R)) \leq e^{\beta R}$. Therefore, we have on one hand

$$\text{Vol}_Y^\varepsilon\left(B_Y(\varepsilon + \psi(\sqrt{d_n}))\right) \leq \exp\left(\varepsilon + \psi(\sqrt{d_n})\right).$$

On the other hand, by taking a partition of the geodesic segment γ_1 into sub-intervals of length ε , we get :

$$\text{Vol}_X^\varepsilon(\gamma_1) \leq \frac{d_n}{\varepsilon} + 1 \leq \frac{2d_n}{\varepsilon}.$$

Hence

$$\text{Vol}_Y^\varepsilon(f(\gamma_1)) \leq \delta' \text{Vol}_X^\varepsilon(\gamma_1) \leq \frac{2\delta'}{\varepsilon} d_n.$$

So we have

$$\text{Vol}_Y^\varepsilon(N_{\psi(\sqrt{d_n})}(f(\gamma_1))) \leq \exp\left(\varepsilon + \psi(\sqrt{d_n})\right) \times \frac{2\delta'}{\varepsilon} d_n.$$

We conclude from the previous inequalities that

$$\delta e^{\alpha\sqrt{d_n}} \leq \delta \text{Vol}_X^\varepsilon(C_n) \leq \text{Vol}_Y^\varepsilon(f(C_n)) \leq \exp\left(\varepsilon + \psi(\sqrt{d_n})\right) \times \frac{2\delta'}{\varepsilon} d_n.$$

Which implies finally that for all $n \geq N$

$$\delta \exp\left(\alpha\sqrt{d_n}\right) \leq \exp\left(\varepsilon + \psi(\sqrt{d_n})\right) \times \frac{2\delta'}{\varepsilon} d_n.$$

Which is not possible when $d_n \rightarrow \infty$ because ψ is sublinear. This completes the proof for $k = 2$.

- Suppose now that the result holds for some $k \geq 2$, and let us prove it for $k + 1$. Suppose that there is a coarse embedding $f : X = X_1 \times \cdots \times X_{k+1} \rightarrow Y$ that satisfies the second condition of the theorem, and let us fix some $x_{k+1} \in X_{k+1}$. The induction hypothesis implies that the restriction of f to the copy $X_1 \times \cdots \times X_k \times \{x_{k+1}\}$ of $X_1 \times \cdots \times X_k$ does not satisfy the second condition of the theorem. Which means that

there exists a sequence $(d_n)_n$ such that $d_n \rightarrow \infty$ and a sequence of k -dimensional rectangles R_{d_n} , whose sides are geodesic segments in the X_i 's, of lengths l_1, \dots, l_k , such that $l_1 = d_n, l_2 \leq d_n^{1/2}, \dots, l_k \leq d_n^{1/k}$ and there exists $\lambda > 0$ such that for all n

$$\text{FillVol}_k^{Y, \text{cr}}(f(\partial R_{d_n})) \geq \lambda d_n^{H_k}.$$

Let $n \in \mathbb{N}$ and let us take $x'_{k+1} \in X_{k+1}$ such that $d_{X_{k+1}}(x_{k+1}, x'_{k+1}) \leq d_n^{\frac{1}{k+1}}$, and consider the $(k+1)$ -dimensional rectangle $R'_{d_n} = R_{d_n} \times [x_{k+1}, x'_{k+1}]$.

Since f satisfies the second condition of the theorem, there exists a sublinear function ϕ such that

$$\text{FillVol}_{k+1}^{Y, \text{cr}}(f(\partial R'_{d_n})) \leq \phi(d_n^{H_{k+1}}).$$

This implies that there exists $V_n \in \mathbf{I}_{k+1}(Y)$ in Y such that $\partial V_n = f(\partial R'_{d_n})$ and

$$\mathbf{M}(V_n) \leq \phi(d_n^{H_{k+1}}).$$

Now consider the 1-Lipschitz map $\pi : Y \rightarrow \mathbb{R}$, $\pi(y) = d_Y(y, f(R_{d_n} \times \{x_{k+1}\}))$, i.e. the map that gives the distance from the image of the basis of the $(k+1)$ -dimensional rectangle R'_{d_n} . Again, by the Slicing Theorem, we have that for a.e. $t \in \mathbb{R}$, there exists $\langle V_n, \pi, t \rangle \in \mathbf{I}_k(Y)$ such that $\langle V_n, \pi, t \rangle = \partial(V_n \llcorner \{\pi \leq t\}) - (\partial V_n) \llcorner \{\pi \leq t\}$, and by integrating the co-area formula over the distance t , we have

$$\int_0^{+\infty} \mathbf{M}(\langle V_n, \pi, t \rangle) dt \leq \mathbf{M}(V_n).$$

Since $\mathbf{M}(V_n) \leq \phi(d_n^{H_{k+1}})$, we get

$$(3.3) \quad \int_0^D \mathbf{M}(\langle V_n, \pi, t \rangle) dt \leq \int_0^{+\infty} \mathbf{M}(\langle V_n, \pi, t \rangle) dt \leq \phi(d_n^{H_{k+1}}),$$

where $D = d_Y(f(R_{d_n} \times \{x_{k+1}\}), f(R_{d_n} \times \{x'_{k+1}\}))$.

However, for a.e. $t \in]0, D[$, $\mathbf{M}(\langle V_n, \pi, t \rangle)$ cannot be too small since $\langle V_n, \pi, t \rangle$ almost gives a filling of the $(k-1)$ -cycle $\partial f(R_{d_n} \times \{x_{k+1}\}) = f(\partial(R_{d_n} \times \{x_{k+1}\}))$, that we will just denote by $f(\partial R_{d_n})$. We will also denote $R_{d_n} \times \{x_{k+1}\}$ by R_{d_n} .

Claim 3.8. For n big enough and for a.e. $t \in]0, D[$, $\mathbf{M}(\langle V_n, \pi, t \rangle) \geq \frac{\lambda}{2} d_n^{H_k}$.

Proof. For a.e. $t \in]0, D[$,

$$\begin{aligned} \partial(V_n) \llcorner \{\pi \leq t\} &= \partial(V_n) \llcorner \{\pi = 0\} + \partial(V_n) \llcorner \{0 < \pi \leq t\} \\ &= f(R_{d_n}) + H_t. \end{aligned}$$

Where H_t is the k -current $\partial(V_n) \llcorner \{0 < \pi \leq t\}$.

Since $\|\langle V_n, \pi, t \rangle\|$ is concentrated on $\pi^{-1}(\{t\})$,

$$\partial(V_n \llcorner \{\pi \leq t\}) = \langle V_n, \pi, t \rangle + f(R_{d_n}) + H_t.$$

Which means that $\langle V_n, \pi, t \rangle + f(R_{d_n}) + H_t$ is actually a cycle.


So $-(\langle V_n, \pi, t \rangle + H_t)$ is a k -chain that fills the $(k-1)$ -cycle $\partial f(R_{d_n}) = f(\partial R_{d_n})$.

Therefore, by the induction hypothesis

$$\mathbf{M}(-(\langle V_n, \pi, t \rangle + H_t)) \geq \text{FillVol}_k^{Y, \text{cr}}(f(\partial R_{d_n})) \geq \lambda d_n^{H_k}.$$

So

$$\mathbf{M}(\langle V_n, \pi, t \rangle) + \mathbf{M}(H_t) \geq \lambda d_n^{H_k}.$$

FIGURE 3. A slice of V_n at distance t from $f(R_{d_n})$

Note that

$$\begin{aligned} H_t &= \partial(V_n) \llcorner \{0 < \pi \leq t\} \\ &= \left(\sum_{F \in \Delta} f(F) \right) \llcorner \{0 < \pi \leq t\}, \end{aligned}$$

where Δ is the set of side faces of R'_{d_n} , i.e. faces of R'_{d_n} except $R_{d_n} \times \{x_{k+1}\}$ and $R_{d_n} \times \{x'_{k+1}\}$, because $0 < \pi < D$. So by taking the mass

$$\begin{aligned} \mathbf{M}(H_t) &= \mathbf{M}\left(\left(\sum_{F \in \Delta} f(F)\right) \llcorner \{0 < \pi \leq t\}\right) \\ &\leq \sum_{F \in \Delta} \mathbf{M}\left((f(F)) \llcorner \{0 < \pi \leq t\}\right) \\ &\leq \sum_{F \in \Delta} \mathbf{M}(f(F)) \\ &\leq \text{Lip}(f) \sum_{F \in \Delta} \mathbf{M}(F). \end{aligned}$$

Since every side face F satisfies $\mathbf{M}(F) \leq d_n^{H_{k+1}-\frac{1}{s}}$, where $s \in \{1, \dots, k\}$, so every $F \in \Delta$ satisfies $\mathbf{M}(F) \leq d_n^{S_{k+1}-\frac{1}{k}}$. There are $2k$ side faces, So

$$\mathbf{M}(H_t) \leq 2k \operatorname{Lip}(f) d_n^{H_{k+1}-\frac{1}{k}}.$$

Hence

$$\begin{aligned} \mathbf{M}(\langle V_n, \pi, t \rangle) &\geq \lambda d_n^{H_k} - \mathbf{M}(H_t) \geq \lambda d_n^{H_k} - 2k \operatorname{Lip}(f) d_n^{H_{k+1}-\frac{1}{k}}. \\ d_n^{H_k} &\gg d_n^{H_{k+1}-\frac{1}{k}}, \text{ so by taking } n \text{ big enough we have } \lambda d_n^{H_k} - 2k \operatorname{Lip}(f) d_n^{H_{k+1}-\frac{1}{k}} \geq \frac{\lambda}{2} d_n^{H_k}. \end{aligned}$$

We conclude that for sufficiently big n and for a.e. $t \in]0, D[$

$$\mathbf{M}(\langle V_n, \pi, t \rangle) \geq \frac{\lambda}{2} d_n^{H_k}. \quad \square$$

Therefore, by (3.3)

$$\begin{aligned} \phi(d^{H_{k+1}}) &\geq \int_0^D \mathbf{M}(\langle V_n, \pi, t \rangle) dt \\ &\geq D \frac{\lambda}{2} d_n^{H_k}. \end{aligned}$$

This implies that

$$\frac{2 \phi(d_n^{H_{k+1}})}{\lambda d_n^{H_k}} \geq D.$$

Let us denote $\psi(d_n^{\frac{1}{k+1}}) = \frac{2 \phi(d_n^{H_{k+1}})}{\lambda d_n^{H_k}}$. Note that ψ is sublinear.

Since $D = d_Y(f(R_{d_n} \times \{x_{k+1}\}), f(R_{d_n} \times \{x'_{k+1}\}))$, the last inequality implies that $\exists z \in X$ such that $z \in R_{d_n} \times \{x'_{k+1}\}$ and $d_Y(f(R_{d_n} \times \{x_{k+1}\}), f(z)) \leq \psi(d_n^{\frac{1}{k+1}})$. But $z \in R_{d_n} \times \{x'_{k+1}\}$ implies that $d_X(z, R_{d_n} \times \{x_{k+1}\}) = d_{X_{k+1}}(x_{k+1}, x'_{k+1})$.

By doing this process for all $x'_{k+1} \in B_{X_{k+1}}(x_{k+1}, d_n^{\frac{1}{k+1}})$, we get a subset $C_n \subset X$ that projects onto $B_{X_{k+1}}(x_{k+1}, d_n^{\frac{1}{k+1}})$, i.e.

$$(3.4) \quad B_{X_{k+1}}(x_{k+1}, d_n^{\frac{1}{k+1}}) \subset \operatorname{proj}_{X_{k+1}}(C_n),$$

and such that

$$(3.5) \quad f(C_n) \subset N_{\psi(d_n^{\frac{1}{k+1}})}(f(R_{d_n} \times \{x_{k+1}\})),$$

since $\forall z \in C_n$, $d_Y(f(z), f(R_{d_n} \times \{x_{k+1}\})) \leq \psi(d_n^{\frac{1}{k+1}})$.

The projection onto X_{k+1} is 1-Lipschitz, so equation (3.4) implies that, if we fix $\varepsilon > 0$,

$$\operatorname{Vol}_X^\varepsilon(C_n) \geq \operatorname{Vol}_X^\varepsilon(\operatorname{proj}_{X_{k+1}}(C_n)) \geq \operatorname{Vol}_X^\varepsilon(B_{X_{k+1}}(x_{k+1}, d_n^{\frac{1}{k+1}})).$$

X_{k+1} has exponential growth, so $\exists \alpha > 0$ such that $\forall R > 0$, $\operatorname{Vol}_{X_{k+1}}^\varepsilon(B_{X_{k+1}}(R)) \geq e^{\alpha R}$. So

$$\operatorname{Vol}_X^\varepsilon(C_n) \geq \exp(\alpha d_n^{\frac{1}{k+1}}).$$

While the equation (3.5) implies that

$$\text{Vol}_Y^\varepsilon(f(C_n)) \leq \text{Vol}_Y^\varepsilon\left(N_{\psi(d_n^{\frac{1}{k+1}})}(f(R_{d_n} \times \{x_{k+1}\}))\right).$$

Since f coarsely preserves volumes, there exist $\delta, \delta' > 0$ such that

$$\delta \text{Vol}_X^\varepsilon(C_n) \leq \text{Vol}_Y^\varepsilon(f(C_n)) \leq \delta' \text{Vol}_X^\varepsilon(C_n).$$

By lemma 3.2, we have

$$\text{Vol}_Y^\varepsilon\left(N_{\psi(d_n^{\frac{1}{k+1}})}(f(R_{d_n} \times \{x_{k+1}\}))\right) \leq \text{Vol}_Y^\varepsilon(B_Y(\varepsilon + \psi(d_n^{\frac{1}{k+1}}))) \times \text{Vol}_Y^\varepsilon(f(R_{d_n} \times \{x_{k+1}\})).$$

Y has at most exponential growth, so there exists $\beta > 0$ such that for all $R > 0$,

$$\text{Vol}_Y^\varepsilon(B_Y(R)) \leq e^{\beta R}.$$

In particular, we have on hand that

$$\text{Vol}_Y^\varepsilon\left(B_Y(\varepsilon + \psi(d_n^{\frac{1}{k+1}}))\right) \leq \exp\left(\beta(\varepsilon + \psi(d_n^{\frac{1}{k+1}}))\right) \leq \exp\left(2\beta\psi(d_n^{\frac{1}{k+1}})\right).$$

On the other hand, by taking a partition of each side vector of R_{d_n} into sub-intervals of length ε , we get :

$$\text{Vol}_X^\varepsilon(R_{d_n} \times \{x_{k+1}\}) \leq \prod_{i=1}^k \left(\frac{d_n^{\frac{1}{i}}}{\varepsilon} + 1\right) \leq \prod_{i=1}^k \left(\frac{2d_n^{\frac{1}{i}}}{\varepsilon}\right) \leq (2/\varepsilon)^k d_n^{H_k}.$$

So, by denoting $A = (2/\varepsilon)^k$, we have

$$\begin{aligned} \text{Vol}_Y^\varepsilon(f(R_{d_n} \times \{x_{k+1}\})) &\leq \delta' \text{Vol}_X^\varepsilon(C^k(x_n; u_1^n, \dots, u_k^n)) \\ &\leq \delta' A d_n^{H_k}. \end{aligned}$$

Therefore

$$\text{Vol}_Y^\varepsilon\left(N_{\psi(d_n^{\frac{1}{k+1}})}(f(C^k(x_n; u_1^n, \dots, u_k^n)))\right) \leq \exp(2\beta\psi(d_n^{\frac{1}{k+1}})) \times \delta' A d_n^{S_k}.$$

We conclude from all the previous inequalities that

$$\delta \exp(\alpha d_n^{\frac{1}{k+1}}) \leq \delta \text{Vol}_X^\varepsilon(C_n) \leq \text{Vol}_Y^\varepsilon(f(C_n)) \leq \exp(2\beta\psi(d_n^{\frac{1}{k+1}})) \times \delta' A d_n^{H_k}.$$

Which implies finally that

$$\delta \exp(\alpha d_n^{\frac{1}{k+1}}) \leq \exp(2\beta\psi(d_n^{\frac{1}{k+1}})) \times \delta' A d_n^{H_k}.$$

Which is not possible when $d_n \rightarrow \infty$ because ψ is sublinear. This completes the proof. \square

Theorem 1.9 immediately follows from that: The k -dimensional filling function of Y being linear implies that there exists $C > 0$ such that for every $(k-1)$ -cycle Σ in Y $\text{FillVol}_k^Y(f(\Sigma)) \leq C \text{Vol}_{k-1}^Y(\Sigma)$. Therefore if R_d is k -dimensional rectangle in X , whose sides

are geodesic segments in the X_i 's, with side lengths l_1, \dots, l_k that satisfies $l_1 = d, l_2 \leq d^{1/2}, \dots, l_k \leq d^{1/k}$, Then

$$\begin{aligned} \text{FillVol}_k^Y(f(\partial R_d)) &\leq C \text{Vol}_{k-1}^Y(f(\partial R_d)) \\ &\leq C \text{Lip}(f) \text{Vol}_{k-1}^X(\partial R_d) \\ &\leq 2k C \text{Lip}(f) l_1 \times \dots \times l_{k-1} \\ &\leq 2k C \text{Lip}(f) d^{H_{k-1}}. \end{aligned}$$

Hence $\text{FillVol}_k^Y(f(\partial R_d)) = o(d^{H_k})$ and so the second condition of the theorem is satisfied.

4. THE DOMAIN X IS A MODEL SPACE WITH NO EUCLIDEAN FACTOR

4.1. Preliminary result on Weyl sectors. The goal of this subsection is to show a structure result for Weyl sectors in a model space. In both symmetric spaces and Euclidean buildings, singular hyperplanes in a maximal flat give rise to a root system. Let us first define the Weyl sectors associated to a root system.

Definition 4.1. Let Φ be a root system in a Euclidean space (E, \langle, \rangle) , see [Bou81] Chap.6, and let Δ be a basis of Φ . Consider the hyperplane perpendicular to each root in Δ . The complement of this finite set of hyperplanes is disconnected, and each connected component is called a *Weyl sector*. Equivalently, we may define Weyl sectors as the equivalence classes of the following relation:

$$u \sim v \quad \text{if} \quad \langle \alpha, u \rangle \cdot \langle \alpha, v \rangle > 0 \quad \text{for every } \alpha \in \Delta.$$

Remark 4.2. They are also called Weyl chambers in the literature. We prefer Weyl sectors to avoid any confusion with chambers in Euclidean buildings.

Let us now define the root system associated to a maximal flat in a model space.

- Let X be a Euclidean building of rank p . Let $x_0 \in X$ be a vertex, and let A be an apartment containing it. Let \mathcal{H} denote the set of walls in A that contains x_0 . Recall that \mathcal{H} is stable under reflections by walls in \mathcal{H} . So if we take for each wall in \mathcal{H} an orthogonal vector with a suitable size, we get a root system. The *Weyl sectors with tip at x_0 of the apartment A* are defined as the Weyl sectors of this root system. They are also the connected components of $A - \cup_{h \in \mathcal{H}} h$.

- Let M is a symmetric space of non-compact type of rank p . A reference for what follows is [Ebe96], Chap.2. Let $x_0 \in M$ and let F be a maximal flat containing x_0 . Let $G = \text{Isom}_0(M)$ be the identity component of the isometry group of M , \mathfrak{g} its Lie algebra and $K = \text{Stab}_G(x_0)$. Let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ be the Cartan decomposition determined by x_0 , and $\theta = \theta_{x_0}$ be the Cartan involution of \mathfrak{g} induced by x_0 .

Consider $\mathfrak{a} \subset \mathfrak{p}$ a maximal abelian subspace such that $F = \exp(\mathfrak{a}).x_0$, and the root space decomposition of \mathfrak{g} determined by \mathfrak{a}

$$\mathfrak{g} = \mathfrak{g}_0 \oplus \sum_{\alpha \in \Lambda} \mathfrak{g}_\alpha,$$

where $\Lambda \subset \mathfrak{a}^*$ is a finite subset called the set of roots determined by \mathfrak{a} , and

$$\mathfrak{g}_\alpha = \{X \in \mathfrak{g} \mid [A, X] = \alpha(A)X, \forall A \in \mathfrak{a}\}.$$

The bilinear form $\langle X, Y \rangle := -B(\theta(X), Y)$, where B is the Killing form of \mathfrak{g} , defines a positive definite inner product on \mathfrak{g} . This inner product induces an isomorphism between \mathfrak{a} and \mathfrak{a}^* , so for all $\alpha \in \Lambda$, let us denote h_α the unique vector in \mathfrak{a} such that we have

$$\alpha(A) = \langle h_\alpha, A \rangle, \quad \forall A \in \mathfrak{a}.$$

A *wall* in F , or equivalently in \mathfrak{a} , is a hyperplane of the form $\ker(\alpha)$. Note that we can identify \mathfrak{a} with F by the exponential map composed with the orbital map at x_0 , which is an isometry when restricted to \mathfrak{a} . For every $\alpha \in \Lambda$, consider the reflection S_α in the hyperplane $\mathfrak{a}_\alpha = \ker(\alpha)$, that is orthogonal to h_α . In particular, we have $S_\alpha(A) = A - 2\frac{\langle h_\alpha, A \rangle}{\langle h_\alpha, h_\alpha \rangle} h_\alpha$, $\forall A \in \mathfrak{a}$. The collection $\{h_\alpha, \alpha \in \Lambda\}$ form a root system. We define the *Weyl sectors with tip at x_0 of the maximal flat F* as the Weyl sectors of this root system. They are also the connected components of $\mathfrak{a} \setminus \cup_{\alpha \in \Lambda} \ker(\alpha)$.

Proposition 4.3. *Let X be a model space. Then its Weyl sectors are acute open simplicial cones generated by maximally singular vectors. In other words, if $x_0 \in X$ and F is a maximal flat containing x_0 , and if C is a Weyl sector of F with tip at x_0 , then after identifying x_0 with the origin of F :*

- There exist u_1, \dots, u_p maximally singular vectors in \overline{C} such that $\forall x \in C$, $\exists x_1, \dots, x_p \geq 0$ such that $x = \sum_{i=1}^p x_i u_i$, where $p = \text{rank } X$.
- For all $x, y \in C$, $\langle x, y \rangle > 0$.




FIGURE 4. A Weyl sector C generated by the maximal singular vectors u_1, u_2, u_3 .

Proof. Since maximal flats in a product space are products of maximal flats in the factors, Weyl sectors of a product space are products of Weyl sectors of the factors. Therefore, it is enough to show the result for an irreducible space, i.e. for a general root system in a Euclidean space.

- Let Φ be a root system in a Euclidean space $E = \mathbb{R}^p$. Let W_r be its Weyl group, i.e. the group generated by reflections through the walls associated to the roots of Φ .

Let C be a Weyl sector with tip at the origin. By [Bou81] Theorem 2 p.153 and section 6 p.64, there exist $\alpha_1, \dots, \alpha_p$ a basis of Φ^* (equivalently there exist h_1, \dots, h_p a basis of Φ) such that C is an open simplicial cone, whose walls are exactly the hyperplanes $\{\alpha_i = 0\}$

for $i = 1 \dots p$ (i.e. the subsets $\{x \in \mathbb{R}^p, \langle x, h_i \rangle = 0\}$ for $i = 1 \dots p$). Which means that $C = \{x = \sum_{i=1}^p x_i e_i \mid x_i \geq 0\}$, where

$$e_i \in \bigcap_{\substack{k=1 \\ k \neq i}}^p \ker(\alpha_k).$$

Since $\{\alpha_1, \dots, \alpha_p\}$ form a basis of Φ^* , they form a basis of $(\mathbb{R}^p)^*$, and the intersection is one dimensional. So the e_i 's are maximally singular vectors.

• Now let us show that C is acute. Again by [Bou81] Theorem 2 p.153, we can choose the basis $\{h_1, \dots, h_p\}$ in a unique way, up to permutation, such that the walls of C are again $\text{Ker}(\alpha_i)$ and such that h_i and C are both in the same side of the wall $\text{Ker}(\alpha_i)$. In that case, we have

$$C = \bigcap_{i=1}^p (\alpha_i > 0) = \{x \in E \mid \forall i \in \{1, \dots, p\}, \langle h_i, x \rangle > 0\}.$$

Since $\{h_1, \dots, h_p\}$ is a basis of the root system Λ , then $\langle h_i, h_j \rangle \leq 0$ for all $i \neq j$.

Therefore if $x, y \in C$, i.e. for every $i \in \{1, \dots, p\}$, $\langle x, h_i \rangle > 0$ and $\langle y, h_i \rangle > 0$, then by [Bou81] p.79 lemma 6, we have $\langle x, y \rangle > 0$. So C is acute. \square

Corollary 4.4. *Let C be a Weyl sector in a model space of rank p , and $u_1, \dots, u_p \in \overline{C}$ maximally singular vectors generating it. Then if $x = \sum_{i=1}^p x_i u_i \in C$, we have $\forall J \subset \{1, \dots, p\}$*

$$\|x\|^2 \geq \sum_{j \in J} \|x_j u_j\|^2.$$

Proof. Since $u_1, \dots, u_p \in \overline{C}$, by the proposition and the continuity of the inner product, we have $\forall i, j \in \{1, \dots, p\}$ $\langle u_i, u_j \rangle \geq 0$.

If $x = \sum_{i=1}^p x_i u_i \in C$, then

$$\|x\|^2 = \sum_{i=1}^p \|x_i u_i\|^2 + 2 \sum_{i,j=1}^p x_i x_j \langle u_i, u_j \rangle \geq \sum_{i=1}^p \|x_i u_i\|^2,$$

since for any $1 \leq i \leq p$, $x_i \geq 0$. \square

4.2. Parallelepipeds and Parallelograms. In the proof of Theorem 1.3, we considered rectangles in the product space. However, in the case of irreducible symmetric space of non-compact type or Euclidean buildings, the space does not factorize and we can no longer consider rectangles. More precisely, maximally singular geodesics are not necessarily orthogonal in a maximal flat. Therefore, we will have to work with cycles that are higher dimensional parallelograms in a maximal flat. We will define them as boundaries of higher dimensional parallelepipeds by induction, as sum of simplices.

Let E be a Euclidean space and $x_1, \dots, x_n \in E$. We denote by $[x_1 \dots x_n]$ the $(n-1)$ -simplex which is the convex hull of these points and the orientation is given by the order in which the vertices are listed. Its boundary $\partial[x_1 \dots x_n]$ is

$$\partial[x_1 \dots x_n] = \sum_{k=1}^n (-1)^{k+1} [x_1, \dots, \hat{x}_k, \dots, x_n],$$

where $[x_1, \dots, \hat{x}_k, \dots, x_n]$ is the $(n-2)$ -simplex given by omitting x_k .

Let us define the 0-dimensional parallelepiped in x , $C^0(x)$, as the 0-simplex $[x]$.

A 1-dimensional parallelepiped is defined from a point x and a vector $u \in E$ by

$$C^1(x; u) := [x, x + u].$$

Suppose that we defined $C^{n-1}(x; u_1, \dots, u_{n-1})$ as a sum of $(n-1)$ -simplices. To define the n -dimensional parallelepiped $C^n(x; u_1, \dots, u_n)$, we use its standard triangulation: x has n hyper-faces opposite to it. Each one of them is an $(n-1)$ -dimensional parallelepiped written as $C^{n-1}(x + u_k; u_1, \dots, \hat{u}_k, \dots, u_n)$, for $k \in \{1, \dots, n\}$, which is a sum of $(n-1)$ -simplices. We take its convex hull with x , that we can write as $[x; C^{n-1}(x + u_k; u_1, \dots, \hat{u}_k, \dots, u_n)]$, which is just the sum of the n -simplices that we get by taking the convex hull of x with the $(n-1)$ -simplices in $C^{n-1}(x + u_k; u_1, \dots, \hat{u}_k, \dots, u_n)$.

We define the n -dimensional parallelepiped $C^n(x; u_1, \dots, u_n)$ as the alternating sum of these terms:

$$C^n(x; u_1, \dots, u_n) := \sum_{k=1}^n (-1)^{k+1} [x, C^{n-1}(x + u_k; u_1, \dots, \hat{u}_k, \dots, u_n)].$$

We denote its boundary $P^{n-1}(x; u_1, \dots, u_n) := \partial C^n(x; u_1, \dots, u_n)$.




FIGURE 5. $P^1(x; u_1, u_2)$ as sum of two simplices

Lemma 4.5. *Let $y, u_1, \dots, u_n \in E$.*

• *We have*

$$C^m(y; u_1, \dots, u_k, \dots, u_n) = -C^m(y + u_k; u_1, \dots, -u_k, \dots, u_n),$$

therefore

$$C^m(y; u_1, \dots, u_n) = (-1)^n C^m(y + \sum_{k=1}^n u_k; -u_1, \dots, -u_n).$$

• *For all $x \in E$,*

$$\partial[x, C^m(y; u_1, \dots, u_{n-1})] = C^m(y; u_1, \dots, u_{n-1}) - [x, \partial C^m(y; u_1, \dots, u_{n-1})]$$

Proof. By the boundary formula and by linearity of the boundary operator. \square

Lemma 4.6. *Let $x, u_1, \dots, u_n \in E$. We can write the $(n-1)$ -dimensional parallelogram as a sum of $(n-1)$ -dimensional parallelepipeds :*

$$(4.1) \quad \begin{aligned} P^{n-1}(x; u_1, \dots, u_n) &= \sum_{k=1}^n (-1)^k C^{n-1}(x; u_1, \dots, \hat{u}_k, \dots, u_n) \\ &+ (-1)^n \sum_{k=1}^n (-1)^k C^{n-1}(x + \sum_{i=1}^n u_i; -u_1, \dots, -\hat{u}_k, \dots, -u_n). \end{aligned}$$

Proof. By the preceding lemma, we have

$$(4.2) \quad \begin{aligned} P^{n-1}(x; u_1, \dots, u_n) &= \partial C^m(x; u_1, \dots, u_n) \\ &= \partial \left(\sum_{k=1}^n (-1)^{k+1} [x, C^{m-1}(x + u_k; u_1, \dots, \hat{u}_k, \dots, u_n)] \right) \\ &= \sum_{k=1}^n (-1)^{k+1} C^{m-1}(x + u_k; u_1, \dots, \hat{u}_k, \dots, u_n) \\ &\quad - \sum_{k=1}^n (-1)^{k+1} [x, \partial C^{m-1}(x + u_k; u_1, \dots, \hat{u}_k, \dots, u_n)]. \end{aligned}$$

Also by lemma 4.5, we have

$$C^{m-1}(x + u_k; u_1, \dots, \hat{u}_k, \dots, u_n) = (-1)^{n-1} C^{m-1}(x + \sum_{i=1}^n u_i; -u_1, \dots, -\hat{u}_k, \dots, -u_n).$$

So

$$\begin{aligned} P^{n-1}(x; u_1, \dots, u_n) &= \sum_{k=1}^n (-1)^k (-1)^n C^{m-1}(x + \sum_{i=1}^n u_i; -u_1, \dots, -\hat{u}_k, \dots, -u_n) \\ &\quad - \sum_{k=1}^n (-1)^{k+1} [x, \partial C^{m-1}(x + u_k; u_1, \dots, \hat{u}_k, \dots, u_n)]. \end{aligned}$$

Let us denote

$$\begin{aligned} S_1 &= (-1)^n \sum_{k=1}^n (-1)^k C^{m-1}(x + \sum_{i=1}^n u_i; -u_1, \dots, -\hat{u}_k, \dots, -u_n), \\ S_2 &= \sum_{k=1}^n (-1)^{k+1} [x, \partial C^{m-1}(x + u_k; u_1, \dots, \hat{u}_k, \dots, u_n)], \end{aligned}$$

so that $P^{n-1}(x; u_1, \dots, u_n) = S_1 - S_2$. Let us first develop S_2 . We have

$$\begin{aligned} \partial C^{m-1}(x + u_k; u_1, \dots, \hat{u}_k, \dots, u_n) &= \sum_{\substack{p=1 \\ p \neq k}}^n (-1)^p \varepsilon(p, k) C^{m-2}(x + u_k; u_1, \dots, \hat{u}_p, \dots, \hat{u}_k, \dots, u_n) \\ &+ (-1)^{n-1} \sum_{\substack{p=1 \\ p \neq k}}^n (-1)^p \varepsilon(p, k) C^{m-2}(x + \sum_{i=1}^n u_i; -u_1, \dots, -\hat{u}_p, \dots, -\hat{u}_k, \dots, -u_n), \end{aligned}$$

where $\varepsilon(p, k) = 1$ if $p < k$ and $\varepsilon(p, k) = -1$ if $p > k$. Therefore

$$\begin{aligned}
S_2 &= \sum_{k=1}^n (-1)^{k+1} [x, \partial C^{n-1}(x + u_k; u_1, \dots, \hat{u}_k, \dots, u_n)] \\
&= \sum_{k=1}^n (-1)^{k+1} \left[x, \sum_{\substack{p=1 \\ p \neq k}}^n (-1)^p \varepsilon(p, k) C^{n-2}(x + u_k; u_1, \dots, \hat{u}_p, \dots, \hat{u}_k, \dots, u_n) \right] \\
&\quad + \sum_{k=1}^n (-1)^{k+1} \left[x, (-1)^{n-1} \sum_{\substack{p=1 \\ p \neq k}}^n (-1)^p \varepsilon(p, k) C^{n-2}(x + \sum_{i=1}^n u_i; -u_1, \dots, -\hat{u}_p, \dots, -\hat{u}_k, \dots, -u_n) \right] \\
&= \sum_{\substack{p=1, k=1 \\ p \neq k}}^n (-1)^{k+1} (-1)^p \varepsilon(p, k) \left[x, C^{n-2}(x + u_k; u_1, \dots, \hat{u}_p, \dots, \hat{u}_k, \dots, u_n) \right] \\
&\quad + (-1)^{n-1} \sum_{\substack{p=1, k=1 \\ p \neq k}}^n (-1)^{k+1} (-1)^p \varepsilon(p, k) \left[x, C^{n-2}(x + \sum_{i=1}^n u_i; -u_1, \dots, -\hat{u}_p, \dots, -\hat{u}_k, \dots, -u_n) \right] \\
&= \sum_{p=1}^n (-1)^p \sum_{\substack{k=1 \\ k \neq p}}^n (-1)^{k+1} \varepsilon(p, k) \left[x, C^{n-2}(x + u_k; u_1, \dots, \hat{u}_p, \dots, \hat{u}_k, \dots, u_n) \right] \\
&\quad + (-1)^{n-1} \sum_{\substack{p=1, k=1 \\ p \neq k}}^n (-1)^{p+k+1} \varepsilon(p, k) \left[x, C^{n-2}(x + \sum_{i=1}^n u_i; -u_1, \dots, -\hat{u}_p, \dots, -\hat{u}_k, \dots, -u_n) \right].
\end{aligned}$$

Remark that

$$\sum_{\substack{k=1 \\ k \neq p}}^n (-1)^{k+1} \varepsilon(p, k) \left[x, C^{n-2}(x + u_k; u_1, \dots, \hat{u}_p, \dots, \hat{u}_k, \dots, u_n) \right] = -C^{n-1}(x; u_1, \dots, \hat{u}_p, \dots, u_n).$$

So the first sum is just equal to $-\sum_{p=1}^n (-1)^p C^{n-1}(x; u_1, \dots, \hat{u}_p, \dots, u_n)$.

By denoting $E(p, k) := (-1)^{p+k+1} \left[x, C^{n-2}(x + \sum_{i=1}^n u_i; -u_1, \dots, -\hat{u}_p, \dots, -\hat{u}_k, \dots, -u_n) \right]$, the second sum can be written as

$$(-1)^{n-1} \sum_{\substack{p=1, k=1 \\ p \neq k}}^n \varepsilon(p, k) E(p, k).$$

Since $E(p, k) = E(k, p)$, and $\varepsilon(p, k) = -\varepsilon(k, p)$, this second sum is equal to zero. So

$$\begin{aligned}
P^{n-1}(x; u_1, \dots, u_n) &= S_1 - S_2 \\
&= (-1)^n \sum_{k=1}^n (-1)^k C^{n-1}(x + \sum_{i=1}^n u_i; -u_1, \dots, -\hat{u}_k, \dots, -u_n) \\
&\quad + \sum_{k=1}^n (-1)^k C^{n-1}(x; u_1, \dots, \hat{u}_k, \dots, u_n). \quad \square
\end{aligned}$$

Now, let us show the following decomposition result:

Proposition 4.7. *Let $x, u_1, \dots, u_{n+1} \in E$. If $u_{n+1} = \sum_{i=1}^p a_i$, with $a_1, \dots, a_p \in E$, Then $P^n(x; u_1, \dots, u_{n+1}) - \sum_{i=1}^p P^n(x + \sum_{k=1}^{i-1} a_k; u_1, \dots, u_n, a_i)$ is an n -cycle, sum of $2n(p+1)$ n -parallelepipeds which are of the form $C^n(y; w_1, \dots, w_n)$, where $w_n \in \{u_{n+1}, a_1, \dots, a_n\}$ and $(w_1, \dots, w_n) \in \{(u_1, \dots, \hat{u}_k, \dots, u_n) \mid k \in \{1, \dots, n\}\}$.*

In particular, if $\|u_1\| > \|u_2\| > \dots > \|u_{n+1}\|$, which is the case that we will be interested in, then the cycle obtained has small volume since all its parallelepipeds have vectors $\neq (u_1, \dots, u_n)$. In other words, we decompose the cycle $P^n(x; u_1, \dots, u_{n+1})$ along the vectors a_1, \dots, a_p , plus a residual cycle with a very small volume compared to the other cycles.




FIGURE 6. A decomposition of $P^1(x; u_1, u_2)$ along the vectors a_1 and a_2 . The residual cycle is the sum of the boundaries of the upper and lower triangles.

Proof. Let $i \in \{1, \dots, p\}$,

$$\begin{aligned}
P^n(x + \sum_{k=1}^{i-1} a_k; u_1, \dots, u_n, a_i) &= \sum_{s=1}^{n+1} (-1)^s C^n(x + \sum_{k=1}^{i-1} a_k; u_1, \dots, \hat{u}_s, \dots, u_n, a_i) \\
&\quad + (-1)^{n+1} \sum_{s=1}^{n+1} (-1)^s C^n(x + \sum_{k=1}^i a_k + \sum_{i=1}^n u_i; -u_1, \dots, -\hat{u}_s, \dots, -u_n, -a_i) \\
&= \sum_{s=1}^n (-1)^s C^n(x + \sum_{k=1}^{i-1} a_k; u_1, \dots, \hat{u}_s, \dots, u_n, a_i) \\
&\quad + (-1)^{n+1} \sum_{s=1}^n (-1)^s C^n(x + \sum_{k=1}^i a_k + \sum_{i=1}^n u_i; -u_1, \dots, -\hat{u}_s, \dots, -u_n, -a_i) \\
&\quad + (-1)^{n+1} C^n(x + \sum_{k=1}^{i-1} a_k; u_1, \dots, u_n) \\
&\quad + (-1)^{n+1} (-1)^{n+1} C^n(x + \sum_{k=1}^i a_k + \sum_{i=1}^n u_i; -u_1, \dots, -u_n).
\end{aligned}$$

By lemma 4.5, we have

$$C^n(x + \sum_{k=1}^i a_k + \sum_{i=1}^n u_i; -u_1, \dots, -u_n) = (-1)^n C^n(x + \sum_{k=1}^i a_k; u_1, \dots, u_n).$$

So

$$\begin{aligned}
&(-1)^{n+1} C^n(x + \sum_{k=1}^{i-1} a_k; u_1, \dots, u_n) \\
&\quad + (-1)^{n+1} (-1)^{n+1} C^n(x + \sum_{k=1}^i a_k + \sum_{i=1}^n u_i; -u_1, \dots, -u_n) \\
&= (-1)^{n+1} C^n(x + \sum_{k=1}^{i-1} a_k; u_1, \dots, u_n) \\
&\quad + (-1)^n C^n(x + \sum_{k=1}^i a_k; u_1, \dots, u_n).
\end{aligned}$$

By adding them, for $i \in \{1, \dots, p\}$, we get

$$\begin{aligned}
&\sum_{i=1}^p \left((-1)^{n+1} C^n(x + \sum_{k=1}^{i-1} a_k; u_1, \dots, u_n) + (-1)^n C^n(x + \sum_{k=1}^i a_k; u_1, \dots, u_n) \right) \\
&= (-1)^n \left(C^m(x + u_{n+1}; u_1, \dots, u_n) - C^m(x; u_1, \dots, u_n) \right).
\end{aligned}$$

So the sum of the p parallelograms $\sum_{i=1}^p P^n(x + \sum_{k=1}^{i-1} a_k; u_1, \dots, u_n, a_i)$ gives

$$\begin{aligned} & \sum_{i=1}^p \sum_{s=1}^n (-1)^s C^n(x + \sum_{k=1}^{i-1} a_k; u_1, \dots, \hat{u}_s, \dots, u_n, a_i) \\ & + (-1)^{n+1} \sum_{i=1}^p \sum_{s=1}^n (-1)^s C^n(x + \sum_{k=1}^i a_k + \sum_{i=1}^n u_i; -u_1, \dots, -\hat{u}_s, \dots, -u_n, -a_i) \\ & + (-1)^n \left(C^n(x + u_{n+1}; u_1, \dots, u_n) - C^n(x; u_1, \dots, u_n) \right). \end{aligned}$$

On the other hand, we have

$$\begin{aligned} P^n(x; u_1, \dots, u_{n+1}) &= \sum_{k=1}^{n+1} (-1)^k C^n(x; u_1, \dots, \hat{u}_k, \dots, u_{n+1}) \\ &+ (-1)^{n+1} \sum_{k=1}^{n+1} (-1)^k C^n(x + \sum_{i=1}^{n+1} u_i; -u_1, \dots, -\hat{u}_k, \dots, -u_{n+1}) \\ &= \sum_{k=1}^n (-1)^k C^n(x; u_1, \dots, \hat{u}_k, \dots, u_{n+1}) \\ &+ (-1)^{n+1} \sum_{k=1}^n (-1)^k C^n(x + \sum_{i=1}^n u_i; -u_1, \dots, -\hat{u}_k, \dots, -u_{n+1}) \\ &+ (-1)^{n+1} C^n(x; u_1, \dots, u_n) \\ &+ (-1)^n C^n(x + u_{n+1}; u_1, \dots, u_n). \end{aligned}$$

Since $(-1)^{n+1}(-1)^{n+1}C^n(x + \sum_{i=1}^{n+1} u_i; -u_1, \dots, -u_n) = (-1)^n C^n(x + u_{n+1}; u_1, \dots, u_n)$.
Finally, by taking the difference, all the chains whose vectors are (u_1, \dots, u_n) simplifies and we get :

$$\begin{aligned} & P^n(x; u_1, \dots, u_{n+1}) - \sum_{i=1}^p P^n(x + \sum_{k=1}^{i-1} a_k; u_1, \dots, u_n, a_i) \\ &= \sum_{k=1}^n (-1)^k C^n(x; u_1, \dots, \hat{u}_k, \dots, u_{n+1}) \\ &+ (-1)^{n+1} \sum_{k=1}^n (-1)^k C^n(x + \sum_{i=1}^{n+1} u_i; -u_1, \dots, -\hat{u}_k, \dots, -u_{n+1}) \\ &- \sum_{i=1}^p \sum_{s=1}^n (-1)^s C^n(x + \sum_{k=1}^{i-1} a_k; u_1, \dots, \hat{u}_s, \dots, u_n, a_i) \\ &- (-1)^{n+1} \sum_{i=1}^p \sum_{s=1}^n (-1)^s C^n(x + \sum_{k=1}^i a_k + \sum_{i=1}^n u_i; -u_1, \dots, -\hat{u}_s, \dots, -u_n, -a_i). \end{aligned}$$

This is a sum of $2n(p+1)$ n -parallelepipedes, whose last vector is either u_{n+1} or an a_i , and whose first $(n-1)$ -vectors are in $\{(u_1, \dots, \hat{u}_k, \dots, u_n) \mid k \in \{1, \dots, n\}\}$. \square

4.3. k -sequences.

In the proof of Theorem 1.3, we used k -dimensional rectangles of side lengths l_1, \dots, l_k that satisfy $l_1 = d, l_2 \leq d^{1/2}, \dots, l_k \leq d^{1/k}$. Parallelograms with the same side lengths will not work. We will need to replace the harmonic sequence by a decreasing positive sequence $(\alpha_n)_n$ that satisfies the additional property: $\frac{S_{n+1}}{n+1} < \alpha_n$ for all $n \in \mathbb{N}^*$, where $S_n = \sum_{k=1}^n \alpha_k$. However, it turns out that such sequence does not exist. We thank Mingkun Liu for having pointed it out to us. Indeed, such sequence must satisfy for all $n \in \mathbb{N}^*$

$$(\alpha_1 - \alpha_n) + \dots + (\alpha_{n-1} - \alpha_n) + 0 + (\alpha_{n+1} - \alpha_n) < 0.$$

Since the sequence is decreasing, for all $n \geq 2$

$$(\alpha_{n-1} - \alpha_n) + (\alpha_{n+1} - \alpha_n) < 0.$$

This implies that the sequence $(\alpha_n - \alpha_{n+1})_n$ is increasing, thus for all $n \geq 2$

$$\alpha_1 - \alpha_2 < \alpha_n - \alpha_{n+1}.$$

This is not possible since $\alpha_1 - \alpha_2 > 0$, and $(\alpha_n - \alpha_{n+1})_n$ converges to 0.

However, to bypass this problem, we can construct arbitrarily long finite sequences that satisfy the required properties. The condition $\frac{S_{n+1}}{n+1} < \alpha_n$ is equivalent to $\alpha_{n+1} < (n+1)\alpha_n - S_n$. We will use this condition to define our sequence by induction.

Let us consider the sequence $(\beta_n)_{n \in \mathbb{N}^*}$ defined by induction :

$$\beta_1 \in \mathbb{R}, \quad \text{and} \quad \forall n \in \mathbb{N}^*, \quad \beta_{n+1} = (n+1)\beta_n - S_n - 1,$$

where $S_n = \sum_{k=1}^n \beta_k$. For all $n \in \mathbb{N}^*$

$$\beta_{n+1} - \beta_n = (n+1)\beta_n - S_n - 1 - n\beta_{n-1} + S_{n-1} + 1 = n(\beta_n - \beta_{n-1}).$$

So

$$\beta_{n+1} - \beta_n = (\beta_2 - \beta_1) n!$$

Therefore

$$\beta_n = \beta_1 - (\beta_1 - \beta_2) \sum_{k=1}^{n-1} k!$$

Since $\beta_2 = \beta_1 - 1$, we get

$$\beta_n = \beta_1 - \sum_{k=1}^{n-1} k!$$

So $(\beta_n)_{n \in \mathbb{N}^*}$ is strictly decreasing and tends to $-\infty$. For any $k \in \mathbb{N}^*$, there exists $\beta_1 \in \mathbb{N}$ such that the first k -terms of the sequence are positive. We will denote by $(\alpha_n)_{n \in \mathbb{N}^*}$ the normalization of such sequence with respect to β_1 , i.e. for any $n \in \mathbb{N}^*$, $\alpha_n = \frac{\beta_n}{\beta_1}$.

Definition 4.8. For every $k \in \mathbb{N}^*$, we will call such sequence a k -sequence, i.e. a sequence $(\alpha_n)_{n \in \mathbb{N}^*}$ that satisfies: $\alpha_1 = 1$, $\alpha_k > 0$, and for all $n \in \mathbb{N}^*$ $\alpha_{n+1} < (n+1)\alpha_n - S_n$.

4.4. Proof of Theorem 1.10. We saw in section 2 that the cross section of a product is the product of cross sections. Therefore, to prove Theorem 1.10, it is enough to prove it when X is an irreducible symmetric space of non-compact type or a Euclidean building with no Euclidean factor. Let us prove the following more general result.

Theorem 4.9. *Let $k \geq 2$. Let X be a symmetric space of non-compact type of rank $\geq k$, or a thick Euclidean building with cocompact affine Weyl group, with bounded geometry and no Euclidean factor, of dimension $\geq k$. Let $\alpha_1, \dots, \alpha_k$ be the k -sequence defined before. Let Y be a uniformly contractible complete metric space with at most exponential growth, and let $f : X \rightarrow Y$ a large-scale Lipschitz map. If there exists a sublinear map ϕ such that $\forall x \in X$, for every F maximal flat that contains x , $\forall d > 0$, $\forall u_1, \dots, u_k \in T_x F \simeq F$ that satisfies*

- u_1, \dots, u_{k-1} are maximally singular,
 - $\|u_1\| = d^{\alpha_1}$ and $\forall i = 2, \dots, k$, $\|u_i\| \leq d^{\alpha_i}$,
- we have

$$\text{FillVol}_k^{Y, \text{cr}}(f(P^{k-1}(x; u_1, \dots, u_k))) \leq \phi(d^{S_k}),$$

then f is not a coarse embedding.

Proof. Let $r \geq 2$ be fixed. We will prove the theorem by induction on k , from $k = 2$ to $k = r$. To prove it for $r' > r$, we will need to change the r -sequence to an r' -sequence.

Let us start by fixing the r -sequence $\alpha_1 > \dots > \alpha_r > 0$, and we will show the theorem for $k = 2, \dots, r$.

- Let us prove it first for $k = 2$.

Let X be a symmetric space of non-compact type of rank $p \geq 2$, or a Euclidean building of rank $p \geq 2$. Suppose there exists a sublinear map ϕ such that $\forall x \in X$, for every F maximal flat that contains x , $\forall d > 0$, $\forall u_1, u_2 \in T_x F \simeq F$ that satisfies : u_1 is maximally singular, $\|u_1\| = d^{\alpha_1}$ and $\|u_2\| \leq d^{\alpha_2}$, we have

$$\text{FillVol}_2^{Y, \text{cr}}(f(P^1(x; u_1, u_2))) \leq \phi(d^{S_2}).$$

Since X has exponential growth and Y has at most exponential growth, the case $k = 1$ in Theorem 1.3 implies that X is not sent sublinearly, i.e. there exist $\lambda > 0$ and two sequences $(a_n)_{n \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}}$ in X such that $d_n := d_X(a_n, b_n) \rightarrow \infty$, and $\forall n \in \mathbb{N}$

$$d_Y(f(a_n), f(b_n)) \geq \lambda d_X(a_n, b_n).$$

For all $n \in \mathbb{N}$, let γ_n denote the geodesic segment in X going from a_n to b_n , which is unique since X is CAT(0). If γ_n is maximally singular, then there is nothing to do at this step. If not, consider a maximal flat F_n containing γ_n and we will denote $w_n := \frac{d_n}{\|\gamma_n'(0)\|} \gamma_n'(0)$ the directing vector of γ_n with magnitude d_n .

Let C be a Weyl sector containing γ_n , i.e. a Weyl sector of F_n with tip at a_n , such that $b_n \in \overline{C}$. So there exist u_1, \dots, u_p maximally singular vectors at a_n generating C , in particular there exist $\delta_1, \dots, \delta_p \geq 0$ such that $w_n = \sum_{i=1}^p \delta_i u_i$.

Now let us consider the geodesic segment γ_n as the 1-chain $C^1(a_n; w_n)$ in F_n , and its boundary $\partial C^1(a_n; w_n) = P^0(a_n; w_n)$. We have that

$$P^0(a_n; w_n) = P^0(a_n; \delta_1 u_1) + P^0(a_n + \delta_1 u_1; \delta_2 u_2) + \dots + P^0(a_n + \sum_{i=1}^{p-1} \delta_i u_i; \delta_p u_p).$$

So, by applying f which is functorial, we have

$$f(P^0(a_n; w_n)) = \sum_{i=1}^p f(P^0(a_n + \sum_{s=1}^{i-1} \delta_s u_s; \delta_i u_i)).$$

So

$$\begin{aligned} \text{FillVol}_1^{Y, \text{cr}}(f(P^0(a_n; w_n))) &= \text{FillVol}_1^{Y, \text{cr}}\left(\sum_{i=1}^p f(P^0(a_n + \sum_{s=1}^{i-1} \delta_s u_s; \delta_i u_i))\right) \\ &\leq \sum_{i=1}^p \text{FillVol}_1^{Y, \text{cr}}\left(f(P^0(a_n + \sum_{s=1}^{i-1} \delta_s u_s; \delta_i u_i))\right). \end{aligned}$$

Since $f(P^0(a_n; w_n)) = [f(b_n)] - [f(a_n)]$, by lemma 3.3

$$\text{FillVol}_1^{Y, \text{cr}}(f(P^0(a_n; w_n))) = d_Y(f(a_n), f(b_n)).$$

So there exists $i_0 \in \{1, \dots, p\}$ such that

$$\begin{aligned} \frac{1}{p} d_Y(f(a_n), f(b_n)) &\leq \text{FillVol}_1^{Y, \text{cr}}\left(f(P^0(a_n + \sum_{s=1}^{i_0-1} \delta_s u_s; \delta_{i_0} u_{i_0}))\right) \\ &= d_Y\left(f\left(a_n + \sum_{s=1}^{i_0-1} \delta_s u_s\right), f\left(a_n + \sum_{s=1}^{i_0} \delta_s u_s\right)\right). \end{aligned}$$

Let us denote $a'_n = a_n + \sum_{s=1}^{i_0-1} \delta_s u_s$ and $b'_n = a_n + \sum_{s=1}^{i_0} \delta_s u_s$. So we have

$$\frac{\lambda}{p} d_X(a_n, b_n) \leq \frac{1}{p} d_Y(f(a_n), f(b_n)) \leq d_Y(f(a'_n), f(b'_n)).$$

This implies in particular that $d_Y(f(a'_n), f(b'_n)) \rightarrow \infty$, which implies that $d_X(a'_n, b'_n) \rightarrow \infty$ because f is a coarse embedding.

Since C is an acute simplicial cone, we use corollary 4.4. We have

$$d_X(a_n, b_n) = \|w_n\| \geq \|\delta_{i_0} u_{i_0}\| = d_X(a'_n, b'_n).$$

We finally get

$$\frac{\lambda}{p} d_X(a'_n, b'_n) \leq d_Y(f(a'_n), f(b'_n)).$$

Now we have two sequences $(a'_n)_n$ and $(b'_n)_n$ in X that satisfies $d_X(a'_n, b'_n) \rightarrow \infty$, and $\forall n \in \mathbb{N}$

$$d_Y(f(a'_n), f(b'_n)) \geq \lambda' d_X(a'_n, b'_n).$$

Where $\lambda' = \frac{\lambda}{p}$, and more importantly, the geodesic between a'_n and b'_n is maximally singular.

To simplify the notations for the rest of the proof, we will just denote a'_n by a_n , b'_n by b_n and λ' by λ .

For every $n \in \mathbb{N}$, let us denote γ_n the bi-infinite geodesic in X that contains a_n and b_n , and consider $P_X(\gamma_n)$ the parallel set of γ_n . $P_X(\gamma_n)$ is a convex subset that splits metrically as

$$P_X(\gamma_n) = \mathbb{R} \times C_X(\gamma_n).$$

The cross section $C_X(\gamma_n)$ is either :

- A symmetric space of non-compact type of rank ≥ 1 , when X is a symmetric space,
- Or a Euclidean building with bounded geometry of dimension ≥ 1 , with no Euclidean

factor, when X is a Euclidean building.

Let us denote $C_X(\gamma_n)$ by X'_n . Note that all such cross sections have exponential growth, and by Proposition 2.13, there is a uniform lower bound on their growth. Therefore, if we fix $\varepsilon > 0$, there exists $\mu > 0$ such that for all $n \in \mathbb{N}$ and for all $R > 0$

$$\text{Vol}^\varepsilon(B_{X'_n}(R)) \geq \exp(\mu R).$$

Now we will work in this product space and apply the same strategy as when X is a product space.

$$P_X(\gamma_n) = \mathbb{R} \times X'_n.$$

Note that the slices $\mathbb{R} \times \{x'\}$, with $x' \in X'$, are the geodesics that are parallel to γ . Let $x'_n \in X'$ such that $\gamma_n = \mathbb{R} \times \{x'_n\}$.

For all $y \in B_{X'_n}(x'_n, d_n^{\alpha_2})$, consider the geodesic $[x'_n, y]$ in X'_n . Denote γ'_n the (unique) bi-infinite geodesic extension of $[x'_n, y]$, and $F''_n = F'_n \times \gamma'_n$ the 2-flat. Consider the 1-parallellogram $P^1(a_n; u_1^n, u_2^n)$ in F''_n , where u_1 is the (maximally singular) directing vector of γ_n that satisfies $\|u_1^n\| = d_X(a_n, b_n)$, and u_2^n is the unique vector in $T_{x_n} F''_n$ that is parallel to γ'_n and satisfies $\|u_2^n\| = d_{X'_n}(x'_n, y)$. This 1-parallellogram satisfies the induction hypothesis:

$$\text{FillVol}_2^{Y, \text{cr}}(f(P^1(a_n; u_1^n, u_2^n))) \leq \phi(d^{S_2}).$$

Which implies that there exists $V_n \in \mathbf{I}_2(Y)$ in Y such that $\partial V_n = f(P^1(a_n; u_1^n, u_2^n))$ and

$$\mathbf{M}(V_n) \leq \phi(d^{S_2}).$$

Now consider the 1-Lipschitz map $\pi : Y \rightarrow \mathbb{R}$, $\pi(z) = d_Y(z, f(C^1(a_n; u_1^n)))$.

By the Slicing Theorem, we have that for a.e. $t \in \mathbb{R}$, there exists $\langle V_n, \pi, t \rangle \in \mathbf{I}_1(Y)$ such that $\langle V_n, \pi, t \rangle = \partial(V_n \llcorner \{\pi \leq t\}) - (\partial V_n) \llcorner \{\pi \leq t\}$, and by integrating the co-area formula over the distance t , we have

$$\begin{aligned} \mathbf{M}(\langle V_n, \pi, t \rangle) &\leq \frac{d}{dt} \mathbf{M}(V_n \llcorner \{\pi \leq t\}), \\ \int_0^{+\infty} \mathbf{M}(\langle V_n, \pi, t \rangle) dt &\leq \int_0^{+\infty} \frac{d}{dt} \mathbf{M}(V_n \llcorner \{\pi \leq t\}), \\ \int_0^{+\infty} \mathbf{M}(\langle V_n, \pi, t \rangle) dt &\leq \mathbf{M}(V_n). \end{aligned}$$

Since $\mathbf{M}(V_n) \leq \phi(d^{S_2})$, we get

$$(4.3) \quad \int_0^D \mathbf{M}(\langle V_n, \pi, t \rangle) dt \leq \int_0^{+\infty} \mathbf{M}(\langle V_n, \pi, t \rangle) dt \leq \phi(d^{S_2})$$

Where $D = d_Y(f(C^1(a_n; u_1^n)), f(C^1(a_n + u_2^n; u_1^n)))$.

However, for a.e. $t \in]0, D[$, $\mathbf{M}(\langle V_n, \pi, t \rangle)$ cannot be too small since $\langle V_n, \pi, t \rangle$ almost gives a filling of the 0-cycle $\partial(f(C^1(a_n; u_1^n))) = \llbracket f(b_n) \rrbracket - \llbracket f(a_n) \rrbracket$.

Claim 4.10. For n big enough: for a.e. $t \in]0, D[$, $\mathbf{M}(\langle V_n, \pi, t \rangle) \geq \frac{\lambda}{2} d_n$.

Proof. For a.e. $t \in]0, D[$,

$$\begin{aligned} \partial(V_n) \llcorner \{\pi \leq t\} &= \partial(V_n) \llcorner \{\pi = 0\} + \partial(V_n) \llcorner \{0 < \pi \leq t\} \\ &= (f(b_n) - f(a_n)) + H_t. \end{aligned}$$

Where H_t is the 1-current $\partial(V_n) \llcorner \{0 < \pi \leq t\}$.

Since $\|\langle V_n, \pi, t \rangle\|$ is concentrated on $\pi^{-1}(\{t\})$,

$$\partial(V_n \llcorner \{\pi \leq t\}) = \langle V_n, \pi, t \rangle + (f(b_n) - f(a_n)) + H_t.$$

Which means that $\langle V_n, \pi, t \rangle + (f(b_n) - f(a_n)) + H_t$ is a 1-current that is actually a cycle. So $-(\langle V_n, \pi, t \rangle + H_t)$ is a 1-chain that fills the 0-cycle $[f(b_n)] - [f(a_n)]$. Therefore,

$$\mathbf{M}(-(\langle V_n, \pi, t \rangle + H_t)) \geq \text{FillVol}_k^{Y, \text{cr}}(f(b_n) - f(a_n)) \geq \lambda d_n.$$

So

$$\mathbf{M}(\langle V_n, \pi, t \rangle) + \mathbf{M}(H_t) \geq \lambda d_n.$$

Note that

$$\begin{aligned} H_t &= \partial(V_n) \llcorner \{0 < \pi \leq t\} \\ &= (f(C^1(a_n; u_2^n)) + f(C^1(b_n; u_2^n))) \llcorner \{0 < \pi \leq t\}. \end{aligned}$$

By taking the mass

$$\begin{aligned} \mathbf{M}(H_t) &= \mathbf{M}((f(C^1(a_n; u_2^n)) + f(C^1(b_n; u_2^n))) \llcorner \{0 < \pi \leq t\}) \\ &\leq \mathbf{M}(f(C^1(a_n; u_2^n)) \llcorner \{0 < \pi \leq t\}) + \mathbf{M}(f(C^1(b_n; u_2^n)) \llcorner \{0 < \pi \leq t\}) \\ &\leq \mathbf{M}(f(C^1(a_n; u_2^n))) + \mathbf{M}(f(C^1(b_n; u_2^n))) \\ &\leq \text{Lip}(f)(\mathbf{M}(C^1(a_n; u_2^n)) + \mathbf{M}(C^1(b_n; u_2^n))). \end{aligned}$$

Since $\mathbf{M}(C^1(a_n; u_2^n)) = \mathbf{M}(C^1(b_n; u_2^n)) \leq d_n^{\alpha_2}$, so

$$\mathbf{M}(H_t) \leq 2\text{Lip}(f)d_n^{\alpha_2}.$$

So

$$\mathbf{M}(\langle V_n, \pi, t \rangle) \geq \lambda d_n - \mathbf{M}(H_t) \geq \lambda d_n - 2\text{Lip}(f)d_n^{\alpha_2}.$$

Since $\alpha_2 < 1$, we have $d_n \gg d_n^{\alpha_2}$. So there exists $N \in \mathbb{N}$ such that for all $n \geq N$

$$\lambda d_n - 2\text{Lip}(f)d_n^{\alpha_2} \geq \frac{\lambda}{2} d_n.$$

So we conclude that for all $n \geq N$ and for a.e. $t \in]0, D[$,

$$\mathbf{M}(\langle V_n, \pi, t \rangle) \geq \frac{\lambda}{2} d_n. \quad \square$$

So we have

$$\begin{aligned} \phi(d_n^{S_2}) &\geq \int_0^D \mathbf{M}(\langle V_n, \pi, t \rangle) dt \\ &\geq D \frac{\lambda}{2} d_n. \end{aligned}$$

Which implies that

$$D \leq \frac{2}{\lambda} \frac{\phi(d_n^{S_2})}{d_n}.$$

Let us denote $\psi(d_n^{\alpha_2}) = \frac{2}{\lambda} \frac{\phi(d_n^{S_2})}{d_n}$. Note that ψ is sublinear.

Since $D = d_Y(f(C^1(a_n; u_1^n)), f(C^1(a_n + u_2^n; u_1^n)))$, the last inequality implies that there exists $z \in C^1(a_n + u_2^n; u_1^n)$ such that

$$d_Y(f(C^1(a_n; u_1^n)), f(z)) \leq \psi(d_n^{\alpha_2}).$$

But $z \in C^1(a_n + u_2^n; u_1^n)$ implies that $\text{proj}_{X'_n}(z) = y$.

If we choose another $y \in B_{X'_n}(x'_n, d_n^{\alpha_2})$, we get another z such that

$$\begin{aligned} \text{proj}_{X'_n}(z) &= y, \\ d_Y(f(C^1(a_n; u_1^n)), f(z)) &\leq \psi(d_n^{\alpha_2}). \end{aligned}$$

By doing this process $\forall y \in B_{X'_n}(x'_n, d_n^{\alpha_2})$, we get a subset $C_n \subset X$ that projects onto $B_{X'_n}(x'_n, d_n^{\alpha_2})$, i.e.

$$B_{X'_n}(x'_n, d_n^{\alpha_2}) \subset \text{proj}_{X'_n}(C_n),$$

and such that

$$f(C_n) \subset N_{\psi(d_n^{\alpha_2})}(f(C^1(a_n; u_1^n))).$$

Since the projection onto X'_n is 1-Lipschitz, it implies that

$$\text{Vol}_Y^\varepsilon(f(C_n)) \leq \text{Vol}_Y^\varepsilon\left(N_{\psi(d_n^{\alpha_2})}(f(C^1(a_n; u_1^n)))\right).$$

f coarsely preserves volumes, so there exist $\delta, \delta' > 0$ such that

$$\delta \text{Vol}_X^\varepsilon(C_n) \leq \text{Vol}_Y^\varepsilon(f(C_n)) \leq \delta' \text{Vol}_X^\varepsilon(C_n).$$

And

$$\text{Vol}_Y^\varepsilon\left(N_{\psi(d_n^{\alpha_2})}(f(C^1(a_n; u_1^n)))\right) \leq \text{Vol}_Y^\varepsilon(B_Y(\varepsilon + \psi(d_n^{\alpha_2})) \times \text{Vol}_Y^\varepsilon(f(C^1(a_n; u_1^n)))).$$

Y has at most exponential growth, so $\exists \beta > 0$ such that $\forall R > 0$, $\text{Vol}_Y^\varepsilon(B_Y(R)) \leq e^{\beta R}$. In particular, we have on hand that

$$\text{Vol}_Y^\varepsilon(B_Y(\varepsilon + \psi(d_n^{\alpha_2}))) \leq \exp(2\beta\psi(d_n^{\alpha_2})).$$

On the other hand, by taking a partition of the geodesic segment $[a_n, b_n]$ into sub-intervals of length ε , we get:

$$\text{Vol}_X^\varepsilon(C^1(a_n; u_1^n)) \leq \frac{\|u_1^n\|}{\varepsilon} + 1 \leq \frac{2\|u_1^n\|}{\varepsilon} \leq (2/\varepsilon)d_n.$$

So, by denoting $A = 2/\varepsilon$, we have

$$\begin{aligned} \text{Vol}_Y^\varepsilon(f(C^1(a_n; u_1^n))) &\leq \delta' \text{Vol}_X^\varepsilon(C^1(a_n; u_1^n)) \\ &\leq \delta' A d_n. \end{aligned}$$

Therefore

$$\text{Vol}_Y^\varepsilon\left(N_{\psi(d_n^{\alpha_2})}(f(C^1(a_n; u_1^n)))\right) \leq \exp(2\beta\psi(d_n^{\alpha_2})) \times \delta' A d_n.$$

We conclude from all the previous inequalities that

$$\delta \exp(\mu d_n^{\alpha_2}) \leq \delta \text{Vol}_X^\varepsilon(C_n) \leq \text{Vol}_Y^\varepsilon(f(C_n)) \leq \exp(2\beta\psi(d_n^{\alpha_2})) \times \delta' A d_n.$$

Which implies finally that for all $n \geq N$

$$\delta \exp(\mu d_n^{\alpha_2}) \leq \exp(2\beta\psi(d_n^{\alpha_2})) \times \delta' A d_n.$$

Which is not possible when $d_n \rightarrow \infty$ because ψ is sublinear.

This completes the proof of the case "rank $X \geq 2$ ".

Now suppose that it is true for rank $X \geq k$ for some $k \in \{2, \dots, r-1\}$, and let us prove it for $k+1$.

Let X be of rank $p \geq k+1$ and suppose that such a coarse embedding $f : X \rightarrow Y$ exists, i.e. there exists a sublinear function ϕ such that $\forall x \in X$, for every maximal flat F that contains x , $\forall d > 0$, $\forall u_1, \dots, u_{k+1} \in T_x F \simeq F$ that satisfies

- u_1, \dots, u_k are maximally singular,
- $\|u_1\| = d^{\alpha_1}$ and $\forall i = 2, \dots, k+1$, $\|u_i\| \leq d^{\alpha_i}$, we have

$$\text{FillVol}_{k+1}^{Y, \text{cr}}(f(P^k(x; u_1, \dots, u_{k+1}))) \leq \phi(d^{S_{k+1}}).$$

X is of rank $\geq k+1$, so it is of rank $\geq k$ and coarsely embeds into Y , therefore it does not satisfy the "rank $\geq k$ " case. This means that there exist a constant $\lambda > 0$, a sequence $(x_n)_n \in X$, maximal flats F_n containing x_n , a sequence d_n that goes to ∞ and $u_1^n, \dots, u_k^n \in T_{x_n} F_n$ such that

- u_1^n, \dots, u_{k-1}^n are maximally singular,
- $\|u_1^n\| = d_n^{\alpha_1}$ and $\forall i = 2, \dots, k$, $\|u_i^n\| \leq d_n^{\alpha_i}$, we have

$$\text{FillVol}_k^{Y, \text{cr}}(f(P^{k-1}(x_n; u_1^n, \dots, u_k^n))) \geq \lambda d_n^{S_k}.$$

The goal now is to find $\lambda' > 0$ and for every $n \in \mathbb{N}$: $x'_n \in F_n$ and maximally singular vectors $u_1^{n'}, \dots, u_k^{n'} \in F_n$ such that $\|u_1^{n'}\| = d_n^{\alpha_1}$ and $\forall i = 2, \dots, k$, $\|u_i^{n'}\| \leq d_n^{\alpha_i}$, and they all tends to ∞ , that satisfy

$$(4.4) \quad \text{FillVol}_k^{Y, \text{cr}}(f(P^{k-1}(x'_n; u_1^{n'}, \dots, u_k^{n'}))) \geq \lambda' d_n^{S_k}.$$

Let $n \in \mathbb{N}$. If u_k^n is maximally singular, there is nothing to do at this step. If not, take F_n a maximal flat at x_n that contains u_1^n, \dots, u_k^n , which is possible since rank $X = p \geq k+1$.

Let C be a Weyl sector of F_n with tip at x_n that contains u_k^n , i.e. such that $x_n + u_k^n \in \overline{C}$. Since C is a simplicial open cone generated by maximally singular vectors, there exist $e_1, \dots, e_p \in F_n$ maximally singular vectors generating C and $\delta_1, \dots, \delta_p \geq 0$ such that $u_k^n = \sum_{i=1}^p \delta_i e_i$.

Now let us consider the $(k-1)$ -parallelogram $P^{k-1}(x_n; u_1^n, \dots, u_k^n)$ seen as a $(k-1)$ -cycle in $F_n \simeq \mathbb{R}^p$. By the decomposition lemma 4.7, we decompose our parallelogram along the walls of the Weyl sector:

$$P^{k-1}(x_n; u_1^n, \dots, u_k^n) = \sum_{i=1}^p P^{k-1}(x_n + \sum_{s=1}^{i-1} \delta_s e_s; u_1^n, \dots, u_{k-1}^n, \delta_i e_i) + R_n^{k-1},$$

where R_n^{k-1} is a $(k-1)$ -cycle in F_n , sum of $2(k-1)(p+1)$ $(k-1)$ -chains, all of the form $C^{k-1}(y, w_1, \dots, w_{k-1})$, where $w_{k-1} \in \{u_k^n, \delta_1 e_1, \dots, \delta_p e_p\}$ and

$(w_1, \dots, w_{k-2}) \in \{(u_1^n, \dots, \hat{u}_j^n, \dots, u_{k-1}^n) \mid j \in \{1, \dots, k-1\}\}$.

So every $(k-1)$ -chain $\Sigma = C^{k-1}(y, w_1, \dots, w_{k-1})$ in R_n^{k-1} satisfy

$$\begin{aligned} \text{Vol}_{F_n}^{k-1}(\Sigma) &\leq \|w_1\| \cdots \|w_{k-2}\| \cdot \|w_{k-1}\| \\ &\leq \|u_1^n\| \cdots \|u_{k-2}^n\| \cdot \|u_k^n\| \\ &\leq d_n^{\alpha_1 + \dots + \alpha_{k-2} + \alpha_k} = d_n^{S_{k-2} + \alpha_k}. \end{aligned}$$

Therefore

$$\text{Vol}_{F_n}^{k-1}(R_n^{k-1}) \leq 2(k-1)(p+1) d_n^{S_{k-2} + \alpha_k}.$$

By applying f to the parallelogram decomposition:

$$f(P^{k-1}(x_n; u_1^n, \dots, u_k^n)) = \sum_{i=1}^p f(P^{k-1}(x_n + \sum_{s=1}^{i-1} \delta_s e_s; u_1^n, \dots, u_{k-1}^n, \delta_i e_i)) + f(R_n^{k-1}).$$

Since a filling of every cycle of the right-hand side gives a filling of the left-hand side, we have

$$\begin{aligned} \text{FillVol}_k^{Y, \text{cr}}(f(P^{k-1}(x_n; u_1^n, \dots, u_k^n))) &\leq \sum_{i=1}^p \text{FillVol}_k^{Y, \text{cr}}(f(P^{k-1}(x_n + \sum_{s=1}^{i-1} \delta_s e_s; u_1^n, \dots, u_{k-1}^n, \delta_i e_i))) \\ &\quad + \text{FillVol}_k^{Y, \text{cr}}(f(R_n^{k-1})). \end{aligned}$$

However, $\text{FillVol}_k^{Y, \text{cr}}(f(R_n^{k-1}))$ is very small compared to $\text{FillVol}_k^{Y, \text{cr}}(f(P^{k-1}(x_n; u_1^n, \dots, u_k^n)))$. Indeed, by considering the images by f of fillings of R_n^{k-1} in $F_n \subset X$, we have

$$\begin{aligned} \text{FillVol}_k^{Y, \text{cr}}(f(R_n^{k-1})) &\leq \text{Lip}(f) \text{FillVol}_k^{X, \text{cr}}(R_n^{k-1}) \\ &\leq \text{Lip}(f) \text{FillVol}_k^{F_n, \text{cr}}(R_n^{k-1}). \end{aligned}$$

By the Euclidean filling of R_n^{k-1} in the flat $F_n \simeq \mathbb{R}^p$, we have

$$\begin{aligned} \text{FillVol}_k^{F_n, \text{cr}}(R_n^{k-1}) &\leq (\text{Vol}_{F_n}^{k-1}(R_n^{k-1}))^{\frac{k}{k-1}} \\ &\leq [2(k-1)(p+1) d_n^{S_k - \alpha_{k-1}}]^{\frac{k}{k-1}}. \end{aligned}$$

So

$$\text{FillVol}_k^{Y, \text{cr}}(f(R_n^{k-1})) \leq \text{Lip}(f) [2(k-1)(p+1)]^{\frac{k}{k-1}} d_n^{\frac{k}{k-1}(S_k - \alpha_{k-1})}.$$

This is where the property of the k -sequence $(\alpha_n)_n$ comes in.

Let us show that $d_n^{\frac{k}{k-1}(S_k - \alpha_{k-1})} = o(d_n^{S_k})$:

$$\begin{aligned} S_k - \frac{k}{k-1}(S_k - \alpha_{k-1}) &= S_k - \left(1 + \frac{1}{k-1}\right)(S_k - \alpha_{k-1}) \\ &= -\frac{S_k}{k-1} + \alpha_{k-1} + \frac{\alpha_{k-1}}{k-1} \\ &= \frac{1}{k-1}(k \alpha_{k-1} - S_k). \end{aligned}$$

$k \alpha_{k-1} - S_k > 0$ because the sequence $(\alpha_n)_n$ satisfies $\forall i = 2, \dots, r : i \alpha_{i-1} > S_i$.

Therefore there exists $N \in \mathbb{N}$ such that for all $n \geq N$ we have:

$$\lambda d_n^{S_k} - \text{Lip}(f) [2(k-1)(p+1)]^{\frac{k}{k-1}} d_n^{\frac{k}{k-1}(S_k - \alpha_{k-1})} \geq \frac{\lambda}{2} d_n^{S_k}.$$

In particular, for $n \geq N$:

$$\text{FillVol}_k^{Y, \text{cr}}(f(P^{k-1}(x_n; u_1^n, \dots, u_k^n))) - \text{FillVol}_k^{Y, \text{cr}}(f(R_n^{k-1})) \geq \frac{\lambda}{2} d_n^{S_k}.$$

i.e. for $n \geq N$:

$$\sum_{i=1}^p \text{FillVol}_k^{Y, \text{cr}}(f(P^{k-1}(x_n + \sum_{s=1}^{i-1} \delta_s e_s; u_1^n, \dots, u_{k-1}^n, \delta_i e_i))) \geq \frac{\lambda}{2} d_n^{S_k}.$$

So there exists $i_0 \in \{1, \dots, p\}$ such that

$$\text{FillVol}_k^{Y,\text{cr}}\left(f\left(P^{k-1}\left(x_n + \sum_{s=1}^{i_0-1} \delta_s e_s; u_1^n, \dots, u_{k-1}^n, \delta_{i_0} e_{i_0}\right)\right)\right) \geq \frac{\lambda}{2p} d_n^{S_k}.$$

Let us denote $x'_n = x_n + \sum_{s=1}^{i_0-1} \delta_s e_s$, $u_1^{n'} = u_1^n, \dots, u_{k-1}^{n'} = u_{k-1}^n, u_k^{n'} = \delta_{i_0} e_{i_0}$, which are all maximally singular.

Remark that $\forall i \in \{2, \dots, k-1\}$, $\|u_i^{n'}\| \rightarrow \infty$ with a speed comparable to that of $d_n^{\alpha_i}$. Indeed, on one hand we have

$$\text{FillVol}_k^{Y,\text{cr}}\left(f\left(P^{k-1}\left(x'_n; u_1^{n'}, \dots, u_k^{n'}\right)\right)\right) \geq \frac{\lambda}{2p} d_n^{S_k}.$$

Note also that

$$\begin{aligned} \text{FillVol}_k^{Y,\text{cr}}\left(f\left(P^{k-1}\left(x'_n; u_1^{n'}, \dots, u_k^{n'}\right)\right)\right) &\leq \text{Lip}(f) \text{FillVol}_k^{X,\text{cr}}\left(P^{k-1}\left(x'_n; u_1^{n'}, \dots, u_k^{n'}\right)\right) \\ &\leq \text{Lip}(f) \text{FillVol}_k^{F_n,\text{cr}}\left(P^{k-1}\left(x'_n; u_1^{n'}, \dots, u_k^{n'}\right)\right) \\ &\leq \text{Lip}(f) \|u_1^{n'}\| \times \dots \times \|u_k^{n'}\|. \end{aligned}$$

Therefore

$$\frac{\lambda}{2p \text{Lip}(f)} d_n^{S_k} \leq \|u_1^{n'}\| \times \dots \times \|u_k^{n'}\|.$$

Since $\|u_1^{n'}\| = d_n^{\alpha_1}$ and $\forall i \in \{2, \dots, k\}$, $\|u_i^{n'}\| \leq d_n^{\alpha_i}$, we get $\forall i \in \{2, \dots, k\}$:

$$\frac{\lambda}{2p \text{Lip}(f)} d_n^{\alpha_i} \leq \|u_i^{n'}\| \leq d_n^{\alpha_i}.$$

In particular, $\|u_k^{n'}\| \gg d_n^{\alpha_{k+1}}$.

Now that we found the point $x'_n \in F_n$, the maximally singular vectors $u_1^{n'}, \dots, u_k^{n'} \in F_n$, and the positive constant $\lambda' = \frac{\lambda}{2p}$ that satisfy the following inequality, and to simplify the notations for the rest of the proof, we will just denote x'_n by x_n , λ' by λ and $u_i^{n'}$ by u_i^n . So that we have

$$\text{FillVol}_k^{Y,\text{cr}}\left(f\left(P^{k-1}\left(x_n; u_1^n, \dots, u_k^n\right)\right)\right) \geq \lambda d_n^{S_k},$$

with $\|u_1^n\| = d_n^{\alpha_1}$ and $\forall i = 2, \dots, k-1$, $\frac{\lambda}{\text{Lip}(f)} d_n^{\alpha_i} \leq \|u_i^n\| \leq d_n^{\alpha_i}$.

Let $n \geq N$. Consider the singular k -flat $F'_n \subset F_n$ at x_n generated by the maximally singular vectors u_1^n, \dots, u_k^n , and let $P_X(F'_n)$ be its parallel set. It splits metrically as

$$P_X(F'_n) = \mathbb{R}^k \times C_X(F'_n).$$

Where the cross section $C_X(F'_n)$, that we will denote X'_n , is either :

- a symmetric space of non-compact type of rank ≥ 1 , when X is a symmetric space,
- or a Euclidean building with bounded geometry of dimension ≥ 1 , with no Euclidean factor, when X is a Euclidean building.

By the same argument as before, if we fix $\varepsilon > 0$, there exists $\mu > 0$ such that for all $n \in \mathbb{N}$ and for all $R > 0$

$$(4.5) \quad \text{Vol}^\varepsilon(B_{X'_n}(R)) \geq \exp(\mu R).$$

Now we will work in this product space and apply the same strategy as when X is a product space. We have

$$P_X(F'_n) = \mathbb{R}^k \times X'_n.$$

Note that the slices $\mathbb{R}^k \times \{x'\}$, with $x' \in X'$, are the flats that are parallel to F'_n . So the k -flat F'_n is such a slice. Let $x'_n \in X'$ such that $F'_n = \mathbb{R}^k \times \{x'_n\}$.

For all $y \in B_{X'_n}(x'_n, d_n^{\alpha_{k+1}})$, consider the geodesic $[x'_n, y]$ in X'_n . Denote γ'_n the (unique) bi-infinite geodesic extension of $[x'_n, y]$, and $F''_n = F'_n \times \gamma'_n$ the $(k+1)$ -flat. Consider the k -parallelogram $P^k(x_n; u_1^n, \dots, u_k^n, u_{k+1}^n)$ in F''_n , where u_{k+1}^n is the unique vector in $T_{x_n} F''_n$ that is parallel to γ'_n and satisfies $\|u_{k+1}^n\| = d_{X'_n}(x'_n, y)$. This k -parallelogram satisfies the induction hypothesis:

$$\text{FillVol}_{k+1}^{Y, \text{cr}}(f(P^k(x_n; u_1^n, \dots, u_{k+1}^n))) \leq \phi(d^{S_{k+1}}).$$

Which implies that there exists $V_n \in \mathbf{I}_{k+1}(Y)$ in Y such that $\partial V_n = f(P^k(x_n; u_1^n, \dots, u_{k+1}^n))$ and

$$\mathbf{M}(V_n) \leq \phi(d^{S_{k+1}}).$$

Now consider the 1-Lipschitz map $\pi : Y \rightarrow \mathbb{R}$, $\pi(z) = d_Y(z, f(C^k(x_n; u_1^n, \dots, u_k^n)))$.

Again, by the Slicing Theorem, we have that for a.e. $t \in \mathbb{R}$, there exists $\langle V_n, \pi, t \rangle \in \mathbf{I}_k(Y)$ such that $\langle V_n, \pi, t \rangle = \partial(V_n \llcorner \{\pi \leq t\}) - (\partial V_n) \llcorner \{\pi \leq t\}$, and by integrating the co-area formula over the distance t , we have that

$$\int_0^{+\infty} \mathbf{M}(\langle V_n, \pi, t \rangle) dt \leq \mathbf{M}(V_n).$$

Since $\mathbf{M}(V_n) \leq \phi(d^{S_{k+1}})$, we get

$$(4.6) \quad \int_0^D \mathbf{M}(\langle V_n, \pi, t \rangle) dt \leq \int_0^{+\infty} \mathbf{M}(\langle V_n, \pi, t \rangle) dt \leq \phi(d^{S_{k+1}}),$$

where $D = d_Y(f(C^k(x_n; u_1^n, \dots, u_k^n)), f(C^k(x_n + u_{k+1}^n; u_1^n, \dots, u_k^n)))$.

However, for a.e. $t \in]0, D[$, $\mathbf{M}(\langle V_n, \pi, t \rangle)$ cannot be too small since it gives a filling of the $(k-1)$ -cycle $\partial f(C^k(x_n; u_1^n, \dots, u_k^n)) = f(P^{k-1}(x_n; u_1^n, \dots, u_k^n))$.

Claim 4.11. For n big enough: for a.e. $t \in]0, D[$, $\mathbf{M}(\langle V_n, \pi, t \rangle) \geq \frac{\lambda}{2} d_n^{S_k}$.

Proof. For every $t \in]0, D[$,

$$\begin{aligned} \partial(V_n) \llcorner \{\pi \leq t\} &= \partial(V_n) \llcorner \{\pi = 0\} + \partial(V_n) \llcorner \{0 < \pi \leq t\} \\ &= f(P^{k-1}(x_n; u_1^n, \dots, u_k^n)) + H_t \end{aligned}$$

Where H_t is the k -current $\partial(V_n) \llcorner \{0 < \pi \leq t\}$.

Since $\|\langle V_n, \pi, t \rangle\|$ is concentrated on $\pi^{-1}(\{t\})$,

$$\partial(V_n \llcorner \{\pi \leq t\}) = \langle V_n, \pi, t \rangle + f(P^{k-1}(x_n; u_1^n, \dots, u_k^n)) + H_t.$$

Which means that $\langle V_n, \pi, t \rangle + f(P^{k-1}(x_n; u_1^n, \dots, u_k^n)) + H_t$ is a k -current that is actually a cycle.

So $-(\langle V_n, \pi, t \rangle + H_t)$ is a k -chain that fills the $(k-1)$ -cycle $f(P^{k-1}(x_n; u_1^n, \dots, u_k^n))$. Therefore,

$$\mathbf{M}(-(\langle V_n, \pi, t \rangle + H_t)) \geq \text{FillVol}_k^{Y, \text{cr}}(f(P^{k-1}(x; u_1, \dots, u_k))) \geq \lambda d_n^{S_k}.$$

So

$$\mathbf{M}(\langle V_n, \pi, t \rangle) + \mathbf{M}(H_t) \geq \lambda d_n^{S_k}.$$

Note that

$$\begin{aligned} H_t &= \partial(V_n) \llcorner \{0 < \pi \leq t\} \\ &= \left(\sum_{F \in \Delta} f(F) \right) \llcorner \{0 < \pi \leq t\}, \end{aligned}$$

where Δ is the set of side faces of $P^k(x_n; u_1^n, \dots, u_{k+1}^n)$, i.e. faces whose vectors are not (u_1^n, \dots, u_k^n) , because $0 < \pi < D$. Indeed, by lemma 4.6

$$(4.7) \quad \begin{aligned} P^k(x_n; u_1^n, \dots, u_{k+1}^n) &= \sum_{s=1}^{k+1} (-1)^s C^k(x; u_1^n, \dots, \hat{u}_s^n, \dots, u_{k+1}^n) \\ &+ (-1)^{k+1} \sum_{k=1}^{k+1} (-1)^s C^k(x + \sum_{i=1}^{k+1} u_i^n; -u_1^n, \dots, -\hat{u}_s^n, \dots, -u_{k+1}^n). \end{aligned}$$

So $\Delta = \{C^k(x; u_1^n, \dots, \hat{u}_s^n, \dots, u_{k+1}^n), C^k(x + \sum_{i=1}^{k+1} u_i^n; -u_1^n, \dots, -\hat{u}_s^n, \dots, -u_{k+1}^n) \mid s \in \{1, \dots, k\}\}$. So by taking the mass

$$(4.8) \quad \begin{aligned} \mathbf{M}(H_t) &= \mathbf{M}\left(\left(\sum_{F \in \Delta} f(F)\right) \llcorner \{0 < \pi \leq t\}\right) \\ &\leq \sum_{F \in \Delta} \mathbf{M}\left((f(F)) \llcorner \{0 < \pi \leq t\}\right) \\ &\leq \sum_{F \in \Delta} \mathbf{M}(f(F)) \\ &\leq \text{Lip}(f) \sum_{F \in \Delta} \mathbf{M}(F). \end{aligned}$$

Since every side face F satisfies $\mathbf{M}(F) \leq d_n^{S_{k+1} - \alpha_s}$, where $s \in \{1, \dots, k\}$, so every $F \in \Delta$ satisfies $\mathbf{M}(F) \leq d_n^{S_{k+1} - \alpha_k}$. And there are $2k$ side faces, So

$$\mathbf{M}(H_t) \leq 2k \text{Lip}(f) d_n^{S_{k+1} - \alpha_k}.$$

So

$$\mathbf{M}(\langle V_n, \pi, t \rangle) \geq \lambda d_n^{S_k} - \mathbf{M}(H_t) \geq \lambda d_n^{S_k} - 2k \text{Lip}(f) d_n^{S_{k+1} - \alpha_k}.$$

The sequence $(\alpha_n)_n$ is decreasing, so $d_n^{S_k} \gg d_n^{S_{k+1} - \alpha_k}$. There exists $N \in \mathbb{N}$ such that for all $n \geq N$, $\lambda d_n^{S_k} - 2k \text{Lip}(f) d_n^{S_{k+1} - \alpha_k} \geq \frac{\lambda}{2} d_n^{S_k}$.

We conclude that for all $n \geq N$ and for a.e. $t \in]0, D[$,

$$\mathbf{M}(\langle V_n, \pi, t \rangle) \geq \frac{\lambda}{2} d_n^{S_k}. \quad \square$$

Therefore, we have

$$\begin{aligned} \phi(d_n^{S_{k+1}}) &\geq \int_0^D \mathbf{M}(\langle V_n, \pi, t \rangle) dt \\ &\geq D \frac{\lambda}{2} d_n^{S_k}. \end{aligned}$$

This implies that

$$D \leq \frac{2}{\lambda} \frac{\phi(d_n^{S_{k+1}})}{d_n^{S_k}}$$

Let us denote $\psi(d_n^{\alpha_{k+1}}) = \frac{2}{\lambda} \frac{\phi(d_n^{S_{k+1}})}{d_n^{S_k}}$. Note that ψ is sublinear.

Since $D = d_Y(f(C^k(x_n; u_1^n, \dots, u_k^n)), f(C^k(x_n + u_{k+1}^n; u_1^n, \dots, u_k^n)))$, the last inequality implies that there exists $z \in C^k(x_n + u_{k+1}^n; u_1^n, \dots, u_k^n)$ such that

$$d_Y(f(C^k(x_n; u_1^n, \dots, u_k^n)), f(z)) \leq \psi(d_n^{\alpha_{k+1}}).$$

But $z \in C^k(x_n + u_{k+1}^n; u_1^n, \dots, u_k^n)$ implies that $\text{proj}_{X'_n}(z) = y$.

If we choose another $y \in B_{X'_n}(x'_n, d_n^{\alpha_{k+1}})$, we get another $z \in C^k(x_n + u_{k+1}^n; u_1^n, \dots, u_k^n)$ such that

$$\begin{aligned} \text{proj}_{X'_n}(z) &= y, \\ d_Y(f(C^k(x_n; u_1^n, \dots, u_k^n)), f(z)) &\leq \psi(d_n^{\alpha_{k+1}}). \end{aligned}$$

By doing this process for all $y \in B_{X'_n}(x'_n, d_n^{\alpha_{k+1}})$, we get subsets $C_n \subset X$ that projects onto $B_{X'_n}(x'_n, d_n^{\alpha_{k+1}})$, i.e.

$$B_{X'_n}(x'_n, d_n^{\alpha_{k+1}}) \subset \text{proj}_{X'_n}(C_n),$$

and such that

$$f(C_n) \subset N_{\psi(d_n^{\alpha_{k+1}})}(f(C^k(x_n; u_1^n, \dots, u_k^n))).$$

Since the projection onto X'_n is 1-Lipschitz, if we fix $\varepsilon > 0$

$$\text{Vol}_X^\varepsilon(B_{X'_n}(x'_n, d_n^{\alpha_{k+1}})) \leq \text{Vol}_X^\varepsilon(\text{proj}_{X'_n}(C_n)) \leq \text{Vol}_X^\varepsilon(C_n).$$

By the uniform lower bound on the volume growth of all such cross sections (4.5), we get

$$(4.9) \quad \exp(\mu d_n^{\alpha_{k+1}}) \leq \text{Vol}_X^\varepsilon(B_{X'_n}(x'_n, d_n^{\alpha_{k+1}})) \leq \text{Vol}_X^\varepsilon(C_n).$$

On the other hand, we get

$$(4.10) \quad \text{Vol}_Y^\varepsilon(f(C_n)) \leq \text{Vol}_Y^\varepsilon(N_{\psi(d_n^{\alpha_{k+1}})}(f(C^k(x_n; u_1^n, \dots, u_k^n)))).$$

f coarsely preserves volumes, i.e. there exist $\delta, \delta' > 0$ such that

$$\delta \text{Vol}_X^\varepsilon(C_n) \leq \text{Vol}_Y^\varepsilon(f(C_n)) \leq \delta' \text{Vol}_X^\varepsilon(C_n)$$

By lemma 3.2, we have

$$\text{Vol}_Y^\varepsilon\left(N_{\psi(d_n^{\alpha_{k+1}})}(f(C^k(x_n; u_1^n, \dots, u_k^n)))\right) \leq \text{Vol}_Y^\varepsilon(B_Y(\varepsilon + \psi(d_n^{\alpha_{k+1}}))) \times \text{Vol}_Y^\varepsilon(f(C^k(x_n; u_1^n, \dots, u_k^n))).$$

Y has at most exponential growth, so there exists $\beta > 0$ such that $\forall R > 0$

$$\text{Vol}_Y^\varepsilon(B_Y(R)) \leq e^{\beta R}.$$

In particular, we have on hand that

$$\text{Vol}_Y^\varepsilon(B_Y(\varepsilon + \psi(d_n^{\alpha_{k+1}}))) \leq \exp(2\beta\psi(d_n^{\alpha_{k+1}})).$$

On the other hand, by taking a partition of each side vector into sub-intervals of length ε , we get a partition of the k -parallelepiped $C^k(x_n; u_1^n, \dots, u_k^n)$:

$$\text{Vol}_X^\varepsilon(C^k(x_n; u_1^n, \dots, u_k^n)) \leq \prod_{i=1}^k \left(\frac{\|u_i^n\|}{\varepsilon} + 1\right) \leq \prod_{i=1}^k \left(\frac{2\|u_i^n\|}{\varepsilon}\right) \leq (2/\varepsilon)^k d_n^{S_k}.$$

So, by denoting $A = (2/\varepsilon)^k$, we have

$$\begin{aligned} \text{Vol}_Y^\varepsilon(f(C^k(x_n; u_1^n, \dots, u_k^n))) &\leq \delta' \text{Vol}_X^\varepsilon(C^k(x_n; u_1^n, \dots, u_k^n)) \\ &\leq \delta' A d_n^{S_k}. \end{aligned}$$

Therefore

$$\text{Vol}_Y^\varepsilon\left(N_{\psi(d_n^{\alpha_{k+1}})}(f(C^k(x_n; u_1^n, \dots, u_k^n)))\right) \leq \exp(2\beta\psi(d_n^{\alpha_{k+1}})) \times \delta' A d_n^{S_k}.$$

We conclude from all the previous inequalities that

$$\delta \exp(\mu d_n^{\alpha_{k+1}}) \leq \delta \text{Vol}_X^\varepsilon(C_n) \leq \text{Vol}_Y^\varepsilon(f(C_n)) \leq \exp(2\beta\psi(d_n^{\alpha_{k+1}})) \times \delta' A d_n^{S_k}.$$

Which implies finally that for all $n \geq N$

$$\delta \exp(\mu d_n^{\alpha_{k+1}}) \leq \exp(2\beta\psi(d_n^{\alpha_{k+1}})) \times \delta' A d_n^{S_k}.$$

Which is not possible when $d_n \rightarrow \infty$ because ψ is sublinear. This completes the induction. \square

Similar to the previous section, Theorem 1.10 follows from that because a $(k-1)$ -parallelogram $P^{k-1}(x; u_1, \dots, u_k)$ in a maximal flat in X with $\|u_1\| = d^{\alpha_1}$ and $\forall i = 2, \dots, k$, $\|u_i\| \leq d^{\alpha_i}$ satisfies

$$\text{FillVol}_k^Y(f(P^{k-1}(x; u_1, \dots, u_k))) \leq 2k C \text{Lip}(f) d^{S_{k-1}} = o(d^{S_k}),$$

where C is the linear k -filling constant.

5. WHEN THE DOMAIN X HAS A ONE-DIMENSIONAL EUCLIDEAN FACTOR

The goal of this section is to prove Theorem 1.11, which implies Theorem 1.5.

5.1. Coarse embeddings of Euclidean spaces into lower rank. Let us prove the following result about coarse embeddings of Euclidean spaces.

Theorem 5.1. *Let $p \geq k \geq 2$ be integers. Let Y be a complete uniformly contractible metric space with linear k -dimensional filling function $\text{FV}_Y^k(l) \sim l$, and let $f : \mathbb{R}^p \rightarrow Y$ be a coarse embedding. Then for any $x \in \mathbb{R}^p$, and every linearly independent vectors $u_1, \dots, u_{k-1} \in \mathbb{R}^p$,*

$$\frac{\text{FillVol}_{k-1}^{Y, \text{cr}}(f(P^{k-2}(x; u_1, \dots, u_{k-1})))}{\|u_1\| \times \dots \times \|u_{k-1}\|} \rightarrow 0, \text{ when } \min\{\|u_i\|\} \text{ tends to } \infty.$$

Proof. Let $C > 0$ such that for any $(k-1)$ -cycle Σ we have $\text{FillVol}_k^{Y, \text{cr}}(\Sigma) \leq C \text{Vol}_Y^\varepsilon(\Sigma)$. Let $x \in \mathbb{R}^p$, and $u_1, \dots, u_{k-1} \in \mathbb{R}^p$ linearly independent vectors. Without loss of generality, we can assume that $\|u_1\| \geq \dots \geq \|u_{k-1}\|$. Let $u_k \in \mathbb{R}^p$ be a vector orthogonal to $\text{span}\{u_1, \dots, u_{k-1}\}$ such that $\|u_k\| = \|u_{k-1}\|^{1/2}$. Then the $(k-1)$ -parallelogram $P^{k-1}(x; u_1, \dots, u_k)$ is a $(k-1)$ -cycle. Therefore

$$\begin{aligned} \text{FillVol}_k^{Y, \text{cr}}(f(P^{k-1}(x; u_1, \dots, u_k))) &\leq C \text{Vol}_{k-1}^Y(f(P^{k-1}(x; u_1, \dots, u_k))) \\ &\leq C \text{Lip}(f) \text{Vol}_{k-1}^{\mathbb{R}^p}(P^{k-1}(x; u_1, \dots, u_k)) \\ &\leq C \text{Lip}(f) 2(k+1) \|u_1\| \times \dots \times \|u_{k-1}\|. \end{aligned}$$

Because $P^{k-1}(x; u_1, \dots, u_k)$ contains $2(k+1)$ faces, and their volume is at most $\|u_1\| \times \dots \times \|u_{k-1}\|$ since $\|u_1\| \geq \dots \geq \|u_{k-1}\|$.

Let $\Omega \in \mathbf{I}_k(Y)$ be a k -current in Y such that $\partial\Omega = f(P^{k-1}(x; u_1, \dots, u_k))$ and

$$\mathbf{M}(\Omega) \leq C \operatorname{Lip}(f) 2(k+1) \|u_1\| \times \dots \times \|u_{k-1}\|.$$

Consider the 1-Lipschitz map $\pi : Y \rightarrow \mathbb{R}$, $\pi(z) = d_Y(z, f(C^{k-1}(x; u_1, \dots, u_{k-1})))$.

By the Slicing Theorem, we have that for a.e. $t \in \mathbb{R}$, there exists $\langle \Omega, \pi, t \rangle \in \mathbf{I}_{k-1}(Y)$ such that $\langle \Omega, \pi, t \rangle = \partial(\Omega \llcorner \{\pi \leq t\}) - (\partial\Omega) \llcorner \{\pi \leq t\}$.

By integrating the co-area formula over the distance t , we have that

$$\int_0^{+\infty} \mathbf{M}(\langle \Omega, \pi, t \rangle) dt \leq \mathbf{M}(\Omega).$$

So

$$\int_0^D \mathbf{M}(\langle \Omega, \pi, t \rangle) dt \leq \int_0^{+\infty} \mathbf{M}(\langle \Omega, \pi, t \rangle) dt \leq C \operatorname{Lip}(f) 2(k+1) \|u_1\| \times \dots \times \|u_{k-1}\|.$$

Where $D = d_Y(f(C^{k-1}(x; u_1, \dots, u_{k-1})), f(C^{k-1}(x + u_k; u_1, \dots, u_{k-1})))$. However:

$$D \geq \rho^- (d_{\mathbb{R}^p}(C^{k-1}(x; u_1, \dots, u_{k-1}), C^{k-1}(x + u_k; u_1, \dots, u_{k-1}))).$$

u_k is orthogonal to $\operatorname{span}\{u_1, \dots, u_{k-1}\}$, so

$$d_{\mathbb{R}^p}(C^{k-1}(x; u_1, \dots, u_{k-1}), C^{k-1}(x + u_k; u_1, \dots, u_{k-1})) = \|u_k\| = \|u_{k-1}\|^{1/2}.$$

Therefore $D \geq \rho^- (\|u_{k-1}\|^{1/2})$. Let us denote $D' = \rho^- (\|u_{k-1}\|^{1/2})$. So

$$\int_0^{D'} \mathbf{M}(\langle \Omega, \pi, t \rangle) dt \leq C \operatorname{Lip}(f) 2(k+1) \|u_1\| \times \dots \times \|u_{k-1}\|.$$

So there exists $t_0 \in]0, D'[$ that satisfies the Slicing Theorem and such that

$$\frac{D'}{2} \mathbf{M}(\langle \Omega, \pi, t_0 \rangle) \leq C \operatorname{Lip}(f) 2(k+1) \|u_1\| \times \dots \times \|u_{k-1}\|.$$

However $\mathbf{M}(\langle \Omega, \pi, t_0 \rangle)$ almost gives a filling of the basis $f(P^{k-2}(x; u_1, \dots, u_{k-1}))$. Indeed, by (1) of the Slicing Theorem, we have

$$\partial(\Omega \llcorner \{\pi \leq t_0\}) = \langle \Omega, \pi, t_0 \rangle + f(P^{k-2}(x; u_1, \dots, u_{k-1})) + H_{t_0}.$$

Where H_{t_0} is the $(k-1)$ -current $\partial(\Omega) \llcorner \{0 < \pi < t_0\}$. Which means that $-(\langle \Omega, \pi, t_0 \rangle + H_{t_0})$ is a $(k-1)$ -chain that fills $f(P^{k-2}(x; u_1, \dots, u_{k-1}))$. Therefore,

$$\begin{aligned} \operatorname{FillVol}_{k-1}^{Y, \operatorname{cr}}(f(P^{k-2}(x; u_1, \dots, u_{k-1}))) &\leq \mathbf{M}(-(\langle \Omega, \pi, t_0 \rangle + H_{t_0})) \\ &\leq \mathbf{M}(\langle \Omega, \pi, t_0 \rangle) + \mathbf{M}(H_{t_0}). \end{aligned}$$

Note that

$$\begin{aligned} H_{t_0} &= \partial(\Omega) \llcorner \{0 < \pi < t_0\} \\ &= \left(\sum_{F \in \Delta} f(F) \right) \llcorner \{0 < \pi < t_0\}, \end{aligned}$$

where Δ is the set of side faces of $P^{k-1}(x_n; u_1^n, \dots, u_k^n)$, i.e. faces whose vectors are not $(u_1^n, \dots, u_{k-1}^n)$, because $0 < \pi < D$. Indeed, by lemma 4.6

$$\begin{aligned} P^{k-1}(x_n; u_1^n, \dots, u_k^n) &= \sum_{s=1}^k (-1)^s C^{k-1}(x; u_1^n, \dots, \hat{u}_s^n, \dots, u_k^n) \\ &\quad + (-1)^k \sum_{s=1}^k (-1)^s C^{k-1}(x + \sum_{i=1}^k u_i^n; -u_1^n, \dots, -\hat{u}_s^n, \dots, -u_k^n). \end{aligned}$$

So $\Delta = \{C^{k-1}(x; u_1^n, \dots, \hat{u}_s^n, \dots, u_k^n), C^{k-1}(x + \sum_{i=1}^k u_i^n; -u_1^n, \dots, -\hat{u}_s^n, \dots, -u_{k+1}^n) \mid s \in \{1, \dots, k-1\}\}$.

By taking the mass

$$\begin{aligned} \mathbf{M}(H_{t_0}) &= \mathbf{M}\left(\sum_{F \in \Delta} f(F)\right) \llcorner \{0 < \pi \leq t_0\} \\ &\leq \sum_{F \in \Delta} \mathbf{M}((f(F)) \llcorner \{0 < \pi \leq t_0\}) \\ &\leq \sum_{F \in \Delta} \mathbf{M}(f(F)) \\ &\leq \text{Lip}(f) \sum_{F \in \Delta} \mathbf{M}(F). \end{aligned}$$

Since every side face F satisfies $\mathbf{M}(F) \leq \frac{\|u_1\| \times \dots \times \|u_k\|}{\|u_s\|}$, where $s \in \{1, \dots, k-1\}$, every $F \in \Delta$ satisfies $\mathbf{M}(F) \leq \|u_1\| \times \dots \times \|u_{k-2}\| \|u_k\|$. There are $2k$ side faces, So

$$\mathbf{M}(H_{t_0}) \leq 2k \text{Lip}(f) \|u_1\| \times \dots \times \|u_{k-2}\| \|u_k\|.$$

So

$$\mathbf{M}(\langle \Omega, \pi, t_0 \rangle) + \mathbf{M}(H_{t_0}) \leq \|u_1\| \times \dots \times \|u_{k-1}\| \left(\frac{2C \text{Lip}(f) 2(k+1)}{D'} + \frac{2k \text{Lip}(f)}{\|u_{k-1}\|^{1/2}} \right).$$

Therefore

$$\frac{\text{FillVol}_{k-1}^{Y, \text{cr}}(f(P^{k-2}(x; u_1, \dots, u_{k-1})))}{\|u_1\| \times \dots \times \|u_{k-1}\|} \leq \left(\frac{2C \text{Lip}(f) 2(k+1)}{\rho^- (\|u_{k-1}\|^{1/2})} + \frac{2k \text{Lip}(f)}{\|u_{k-1}\|^{1/2}} \right).$$

and the right hand side clearly tends to ∞ when $\|u_{k-1}\|$ tends to ∞ . \square

Theorem 1.11 is an immediate consequence of this result.

Proof. Let $k \in \mathbb{N}$, X, Y be as in Theorem 1.11, and let $f : X \times \mathbb{R} \rightarrow Y$ be a coarse embedding.

Let $(\alpha_n)_n$ be a k -sequence. Let $x \in X$ and let F be a maximal flat in X containing x . Consider the maximal flat $E = F \times \mathbb{R}$ in $X \times \mathbb{R}$. By Theorem 5.1, there exists a sublinear function ϕ such that for all $d > 0$, for all u_1, \dots, u_k linearly independent vectors in $T_x E \simeq E$ such that $\|u_i\| = d^{\alpha_i}$,

$$\text{FillVol}_k^{Y, \text{cr}}(f(P^{k-1}(x; u_1, \dots, u_k))) \leq \phi(d^{S_k}).$$

In particular, u_1, \dots, u_k can be chosen in $T_x F \subset T_x E$. Therefore, when restricted to a copy $X \times \{z\}$ for some $z \in \mathbb{R}$, the coarse embedding $\tilde{f} := f \upharpoonright_X : X \rightarrow Y$ satisfies the condition

of Theorem 4.9, or Theorem 3.4, which is not possible. Therefore, a coarse embedding $f : X \times \mathbb{R} \rightarrow Y$ cannot exist. \square

Another consequence of this result is Theorem 1.6.

Proof. Let us suppose that there exists a subspace $E \simeq \mathbb{R}^k \subset \mathbb{R}^p$ that is sent quasi-isometrically with constants (λ, c) . By Lemma 2.21, we may assume that f is λ -Lipschitz. Let us also denote by f its restriction to E . So $f(E)$ is a maximal quasi-flat in Y . By the quasi-flats Theorem in [KL97] and [EF97], there exist $\delta > 0$ and maximal flats F_1, \dots, F_r in Y such that $f(E) \subset N_\delta(F_1 \cup \dots \cup F_r)$. We can assume that the union $F_1 \cup \dots \cup F_r$ is minimal in the sens that for all $i \in \{1, \dots, r\}$, $f(E)$ is not in a bounded neighborhood of $F_1 \cup \dots \cup F_r \setminus F_i$ (elsewhere we can just remove F_i and modify δ). Therefore by minimality of this union, there exists a sequence $(x_n)_n \in E$ such that $d_Y(f(x_n), F_1 \cup \dots \cup F_{r-1}) \geq n^2$, i.e. $(f(x_n))_n$ only stays in the δ -neighborhood of F_r .

Let $z \in B_E(x_n, n)$, so $d_Y(f(z), f(x_n)) \leq \lambda n$. On one hand we have

$$d_Y(f(x_n), F_1 \cup \dots \cup F_{r-1}) \leq d_Y(f(z), f(x_n)) + d_Y(f(z), F_1 \cup \dots \cup F_{r-1}).$$

So, for n big enough

$$\delta < n^2 - \lambda n \leq d_Y(f(z), F_1 \cup \dots \cup F_{r-1}).$$

On the other hand, $f(z) \in N_\delta(F_1 \cup \dots \cup F_r)$. So for n big enough, we have $f(z) \in N_\delta(F_r)$. Thus there exists $N \in \mathbb{N}$ such that for all $n \geq N$, $f(B_E(x_n, n)) \subset N_\delta(F_r)$.

Let us consider the projection map $\pi : Y \rightarrow F_r$, which is 1-Lipschitz since Y is CAT(0) and F_r is a closed convex subset. Let us denote $A := \cup_{n \geq N} B_E(x_n, n)$ and consider the map $g : A \rightarrow F_r$, $g(a) = \pi \circ f(a)$. It is a quasi-isometric embedding. Indeed, it is clearly λ -Lipschitz, and for $x, y \in A$ we have

$$\begin{aligned} d_Y(f(x), f(y)) &\leq d_Y(f(x), \pi(f(y))) + d_Y(\pi(f(x)), \pi(f(y))) + d_Y(\pi(f(y)), f(y)) \\ &\leq d_{F_r}(\pi(f(x)), \pi(f(y))) + 2\delta. \end{aligned}$$

Therefore, for all $x, y \in A$

$$\lambda d_A(x, y) - c - 2\delta \leq d_{F_r}(\pi(f(x)), \pi(f(y))) \leq \lambda d_A(f(x), f(y)).$$

Let us denote $c' = c + 2\delta$.

For all $n \geq N$, let $u_1^n, \dots, u_k^n \in E$ be orthogonal vectors such that $\|u_i\| = n^{1/i}$, and consider the $(k-1)$ -parallelograms $P^{k-1}(x_n; u_1^n, \dots, u_k^n)$ in E , which are actually boundaries of k -dimensional rectangles in this case. Note that for all $n \geq N$, $P^{k-1}(x_n; u_1^n, \dots, u_k^n)$ is in $B_E(x_n, n)$, so all the parallelograms are in A . Moreover, by the previous theorem, there exists a sublinear function ϕ such that for all $n \geq N$

$$\text{FillVol}_k^{Y, \text{cr}}(f(P^{k-1}(x_n; u_1^n, \dots, u_k^n))) \leq \phi(n^{H_k}).$$

π is 1-Lipschitz so

$$\text{FillVol}_k^{F_r, \text{cr}}(g(P^{k-1}(x_n; u_1^n, \dots, u_k^n))) \leq \phi(n^{H_k}).$$

Let $V_n \in \mathbf{I}_k(F_r)$ such that $\partial V_n = g(P^{k-1}(x_n; u_1^n, \dots, u_k^n))$ and $\mathbf{M}(V) \leq \phi(n^{H_k})$, and consider the 1-Lipschitz map $d_1 : F_r \rightarrow \mathbb{R}$, $d_1(z) = d_{F_r}(z, g(C^{k-1}(x_n; u_1^n, \dots, u_{k-1}^n)))$. By the Slicing Theorem,

$$\int_0^D \mathbf{M}(\langle V_n, \pi, t \rangle) dt \leq \int_0^{+\infty} \mathbf{M}(\langle V_n, \pi, t \rangle) dt \leq \mathbf{M}(V_n) \leq \phi(n^{H_k}).$$

Where $D = d_{F_r}(g(C^{k-1}(x_n; u_1^n, \dots, u_{k-1}^n)), g(C^{k-1}(x_n + u_k^n; u_1^n, \dots, u_{k-1}^n)))$. g is a (λ, c') quasi-isometric embedding and u_k^n is orthogonal to $\text{span}\{u_1^n, \dots, u_{k-1}^n\}$, so

$$D \geq \lambda d_X(C^{k-1}(x_n; u_1^n, \dots, u_{k-1}^n), C^{k-1}(x_n + u_k^n; u_1^n, \dots, u_{k-1}^n)) - c' \geq \lambda n^{1/k} - c'.$$

Up to taking bigger N , we can assume that for all $n \geq N$ and for all $i \in \{1 \dots k\}$, $\lambda n^{1/i} - c' \geq \frac{\lambda}{2} n^{1/i}$.

Therefore there exists $t_1 \in]0, \frac{\lambda}{2} n^{1/k}[$ such that

$$\frac{\lambda}{4} n^{1/k} \mathbf{M}(\langle V_n, \pi, t_1 \rangle) \leq \mathbf{M}(V_n).$$

Again, by adding a small current to $\langle V_n, \pi, t_1 \rangle$ of mass less than $2\lambda(k-1)n^{H_k - \frac{1}{k-1}}$, we get a filling of the cycle $g(P^{k-2}(x_n; u_1^n, \dots, u_{k-1}^n))$. So

$$\text{FillVol}_{k-1}^{F_r, \text{cr}}(g(P^{k-2}(x_n; u_1^n, \dots, u_{k-1}^n))) \leq \frac{4}{\lambda} \frac{\phi(n^{1/k})}{n^{1/k}} + 2\lambda(k-1)n^{H_k - \frac{1}{k-1}}.$$

Since $H_k - \frac{1}{k-1} < H_{k-1}$, the right hand side is $\ll n^{H_{k-1}}$. Let us denote by ψ the sublinear function such that

$$\text{FillVol}_{k-1}^{F_r, \text{cr}}(g(P^{k-2}(x_n; u_1^n, \dots, u_{k-1}^n))) \leq \psi(n^{H_{k-1}}).$$

We can apply the same strategy again by considering the 1-Lipschitz map $d_2 : F_r \rightarrow \mathbb{R}$, $d_2(z) = d_{F_r}(z, g(C^{k-2}(x_n; u_1^n, \dots, u_{k-2}^n)))$, and using the Slicing Theorem. By repeating this process $(k-1)$ times, we get a sublinear function φ such that for all $n \geq N$

$$\text{FillVol}_1^{F_r, \text{cr}}(g(P^0(x_n; u_1^n))) \leq \varphi(n).$$

If we denote $a_n = x_n$ and $b_n = x_n + u_1^n$, this implies that

$$\frac{\lambda}{2} n \leq d_{F_r}(g(a_n), g(b_n)) \leq \varphi(n).$$

Which contradicts the sublinearity of φ . □

Finally, let us give a proof of Corollary 1.7.

Proof. By a result of Bonk–Schramm [BS11], there exists $n \in \mathbb{N}$ such that Y quasi-isometrically embeds into \mathbb{H}^n . Let $g : Y \rightarrow \mathbb{H}^n$ be such an embedding with constants (λ, c) , and consider the coarse embedding $h = g \circ f : \mathbb{R}^p \rightarrow \mathbb{H}^n$. By Theorem 5.1, for all $x \in \mathbb{R}^p$, and every $u \in \mathbb{R}^p$

$$\frac{\text{FillVol}_1^{\mathbb{H}^n, \text{cr}}(h(P^0(x; u)))}{\|u\|} \rightarrow 0, \text{ when } \|u\| \text{ tends to } \infty.$$

In other words, and by lemma 3.3, there exists a sublinear function ϕ such that for all $x, y \in \mathbb{R}^p$, $d_{\mathbb{H}^n}(h(x), h(y)) \leq \phi(d_{\mathbb{R}^p}(x, y))$. However, g is a (λ, c) -quasi-isometric embedding so

$$\frac{1}{\lambda} d_Y(f(x), f(y)) - c \leq d_{\mathbb{H}^n}(h(x), h(y)).$$

Therefore, for all $x, y \in \mathbb{R}^p$

$$d_Y(f(x), f(y)) \leq \lambda \phi(d_{\mathbb{R}^p}(x, y)) + \lambda c,$$

and $\psi(t) := \lambda \phi(t) + \lambda c$ is sublinear. □

Remark 5.2. In higher rank, the Theorem 1.6 does not necessarily imply that the coarse embedding is uniformly compressing, as shown by the following example:

$$\begin{aligned} \psi : \quad \mathbb{R}^3 &\longrightarrow \mathbb{H}^3 \times \mathbb{R} \\ (x, y, z) &\longmapsto ((x, y, 1), z) \end{aligned}$$

Where \mathbb{H}^3 is the upper half-space model. ψ is the product of a horospherical embedding with the identity map on \mathbb{R} . It is clearly a coarse embedding, $\text{rank}(\mathbb{H}^3 \times \mathbb{R}) = 2$, but it is not uniformly compressing since the z -axis is sent isometrically.

6. FURTHER QUESTIONS

Question 6.1. Does every cocompact geodesic metric space with exponential growth admits a coarse embedding of a binary tree?

This question was already asked by Shalom in [Sha04]. Note that if the answer is positive, then Theorem 1.3 is just a consequence of Theorem 1.4. Indeed, Theorem 1.4 already treats the case when the domain X is a product of regular trees, since the $(p+1)$ -regular tree can be seen as the Bruhat-Tits building of $\text{SL}_2(\mathbb{Q}_p)$. Moreover, all regular trees of degree ≥ 3 are quasi-isometric. So if $X = X_1 \times \cdots \times X_k$ is a product of geodesic metric spaces of exponential growth, then X contains an isometric copy of a model space of rank k with no Euclidean factor. Therefore Theorem 1.3 follows from Theorem 1.4. This question in full generality is still open even for groups. In [DCT08], Cornuier and Tessera showed that if a group G is either a connected Lie group, or a finitely generated solvable group with exponential growth, then it contains a quasi-isometrically embedded free sub-semigroup on 2 generators. Thus it contains quasi-isometrically embedded binary tree. However, it is not known whether every locally compact compactly generated (or finitely generated) group with exponential growth contains a quasi-isometrically (or even coarsely) embedded copy of a binary tree.

Question 6.2. When there is no coarse embedding $X \rightarrow Y$, and $\text{rank}(X) = \text{rank}(Y) + 1$, can adding a Euclidean factor in the target make it possible? i.e. can we embed X into $Y \times \mathbb{R}^p$ for some $p \in \mathbb{N}$?

We answered this question negatively in Corollary 1.8 when $\text{rank}(X) > \text{rank}(Y) + 1$. But we do not know if the case $\text{rank}(X) = \text{rank}(Y) + 1$ is different. For example, there is no coarse embedding from the symmetric space $\text{SL}_3(\mathbb{R})/\text{SO}_3(\mathbb{R})$ of rank 2 into a hyperbolic space \mathbb{H}^n . Can we have a coarse embedding $\text{SL}_3(\mathbb{R})/\text{SO}_3(\mathbb{R}) \rightarrow \mathbb{H}^n \times \mathbb{R}^p$ for some $p \in \mathbb{N}$?

Question 6.3. What can we say about the limit set $\overline{f(X)} \cap \partial Y$ of a coarse embedding $f : X \rightarrow Y$ between model spaces, and in particular about coarse embeddings of \mathbb{R}^{n+1} into \mathbb{H}^n ? For instance, the limit sets of horospherical embeddings are reduced to a point. Is it the case for all coarse embeddings \mathbb{R}^{n+1} into \mathbb{H}^n ? Is the image of such embedding always contained in a horoball?

Note that Bowditch [Bow17] showed that when Y is Gromov-hyperbolic with bounded geometry and X is a geodesic metric space that has “fast growth”, in particular if it has exponential growth, then the limit set does not contain isolated points.

Our results extends to uniform lattices in symmetric spaces, since they are quasi-isometric to them. A natural question to ask is the following.

Question 6.4. What can be said about nonuniform lattices? For example, can we embed $SL_n(\mathbb{Z})$ into a symmetric space of rank $< n - 1$?

Fisher–Whyte [FW18] proved that $SL_2(\mathbb{R}) \times SL_2(\mathbb{R})$ quasi-isometrically embeds into $SL_3(\mathbb{R})$. Can we have a coarse embedding from $SL_2(\mathbb{Z}) \times SL_2(\mathbb{Z})$ into $SL_3(\mathbb{Z})$? Note that Leuzinger and Young managed recently to give higher filling functions of some nonuniform lattices [LY21].

Question 6.5. Can we extend these results to all cocompact CAT(0) spaces?

To do so, we need an analogue of Leuzinger’s result, i.e. is the filling of a cocompact CAT(0) space linear above the rank? This is an open question. Note that we know by [Wen11] that proper CAT(0) spaces admit a sub-Euclidean filling above the asymptotic rank.

REFERENCES

- [AB08] Peter Abramenko and Kenneth S Brown. *Buildings: theory and applications*, volume 248. Springer Science & Business Media, 2008.
- [AK⁺00] Luigi Ambrosio, Bernd Kirchheim, et al. Currents in metric spaces. *Acta Mathematica*, 185(1):1–80, 2000.
- [AS86] Michael T Anderson and Viktor Schroeder. Existence of flats in manifolds of nonpositive curvature. *Inventiones mathematicae*, 85(2):303–315, 1986.
- [Ass82] Patrice Assouad. Sur la distance de nagata. *CR Acad. Paris*, 294:31–34, 1982.
- [BDS07] Sergei Buyalo, Alexander Dranishnikov, and Viktor Schroeder. Embedding of hyperbolic groups into products of binary trees. *Inventiones mathematicae*, 169(1):153–192, 2007.
- [BF98] Noel Brady and Benson Farb. Filling-invariants at infinity for manifolds of nonpositive curvature. *Transactions of the American Mathematical Society*, 350(8):3393–3405, 1998.
- [BH13] Martin R Bridson and André Haefliger. *Metric spaces of non-positive curvature*, volume 319. Springer Science & Business Media, 2013.
- [BH21] Yves Benoist and Dominique Hulin. Harmonic quasi-isometric maps ii: negatively curved manifolds. *Journal of the European Mathematical Society*, 2021.
- [BHS21] Jason Behrstock, Mark F Hagen, and Alessandro Sisto. Quasiflats in hierarchically hyperbolic spaces. *Duke Mathematical Journal*, 170(5):909–996, 2021.
- [Bou81] Nicolas Bourbaki. *Groupes et algèbres de Lie: Chapitres 4, 5 et 6*. Masson, 1981.
- [Bow17] Brian H Bowditch. Limit sets of coarse embeddings. 2017.
- [BS11] Mario Bonk and Oded Schramm. Embeddings of gromov hyperbolic spaces. In *Selected Works of Oded Schramm*, pages 243–284. Springer, 2011.
- [BST12] Itai Benjamini, Oded Schramm, and Ádám Timár. On the separation profile of infinite graphs. *Groups, Geometry, and Dynamics*, 6(4):639–658, 2012.
- [BW92] Jonathan Block and Shmuel Weinberger. Aperiodic tilings, positive scalar curvature, and amenability of spaces. *Journal of the American Mathematical Society*, 5(4):907–918, 1992.
- [BW97] Jonathan Block and Shmuel Weinberger. Large scale homology theories and geometry. *AMS/IP Stud. in Adv. Math.*, 2:522–569, 1997.
- [Cap14] Pierre-Emmanuel Caprace. Lectures on CAT(0) spaces and their isometry groups. 2014.
- [CG19] Corentin Le Coz and Antoine Gournay. Separation profiles, isoperimetry, growth and compression. *arXiv preprint:1910.11733*, 2019.
- [CM09] Pierre-Emmanuel Caprace and Nicolas Monod. Isometry groups of non-positively curved spaces: structure theory. *Journal of topology*, 2(4):661–700, 2009.

- [DCT08] Yves De Cornulier and Romain Tessera. Quasi-isometrically embedded free sub-semigroups. *Geometry & Topology*, 12(1):461–473, 2008.
- [DKLMT20] Thiebaut Delabie, Juhani Koivisto, François Le Maître, and Romain Tessera. Quantitative measure equivalence between finitely generated groups. 2020.
- [Ebe96] Patrick Eberlein. *Geometry of nonpositively curved manifolds*. University of Chicago Press, 1996.
- [EF97] Alex Eskin and Benson Farb. Quasi-flats and rigidity in higher rank symmetric spaces. *Journal of the American Mathematical Society*, pages 653–692, 1997.
- [FF60] Herbert Federer and Wendell H Fleming. Normal and integral currents. *Annals of Mathematics*, pages 458–520, 1960.
- [Fra21] Charles Frances. Isometry group of lorentz manifolds: A coarse perspective. *arXiv preprint:2102.09213*, 2021.
- [FW18] David Fisher and Kevin Whyte. Quasi-isometric embeddings of symmetric spaces. *Geometry & Topology*, 22(5):3049–3082, 2018.
- [Gro88] Mikhaïl Gromov. Rigid transformation groups, in “géométrie différentielle”(d. bernard, choquet-bruhat, ed.). *Travaux encours*, 33, 1988.
- [Gro93] Mikhael Gromov. Asymptotic invariants of infinite groups. *Geometric group theory*, 2:1–295, 1993.
- [Hel01] Sigurdur Helgason. *Differential geometry and symmetric spaces*, volume 341. American Mathematical Soc., 2001.
- [HKS22] Jingyin Huang, Bruce Kleiner, and Stephan Stadler. Morse quasiflats i. *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 2022.
- [HMT20] David Hume, John M Mackay, and Romain Tessera. Poincaré profiles of groups and spaces. *Revista Matemática Iberoamericana*, 36(6):1835–1886, 2020.
- [HS17] David Hume and Alessandro Sisto. Groups with no coarse embeddings into hyperbolic groups. *New York Journal of Mathematics*, 23:1657–1670, 2017.
- [KL97] Bruce Kleiner and Bernhard Leeb. Rigidity of quasi-isometries for symmetric spaces and euclidean buildings. *Publications mathématiques de l’IHES*, 86:115–197, 1997.
- [Kle99] Bruce Kleiner. The local structure of length spaces with curvature bounded above. *Mathematische Zeitschrift*, 231(3):409–456, 1999.
- [KW14] Linus Kramer and Richard M Weiss. Coarse equivalences of euclidean buildings. *Advances in Mathematics*, 253:1–49, 2014.
- [KY06] Gennadi Kasparov and Guoliang Yu. The coarse geometric novikov conjecture and uniform convexity. *Advances in Mathematics*, 206(1):1–56, 2006.
- [Lee] Bernhard Leeb. A characterization of irreducible symmetric spaces and euclidean buildings of higher rank by their asymptotic geometry. *bonner mathematische schriften. Bonn Mathematical Publications*, 326:105–113.
- [Leu14] Enrico Leuzinger. Optimal higher-dimensional dehn functions for some cat (0) lattices. *Groups, Geometry, and Dynamics*, 8(2):441–466, 2014.
- [LY21] Enrico Leuzinger and Robert Young. Filling functions of arithmetic groups. *Annals of Mathematics*, 193(3):733–792, 2021.
- [MSW03] Lee Mosher, Michah Sageev, and Kevin Whyte. Quasi-actions on trees i. bounded valence. *Annals of mathematics*, pages 115–164, 2003.
- [Sha04] Yehuda Shalom. Harmonic analysis, cohomology, and the large-scale geometry of amenable groups. *Acta Mathematica*, 192(2):119, 2004.
- [STY02] Georges Skandalis, Jean-Louis Tu, and Guoliang Yu. The coarse Baum–Connes conjecture and groupoids. *Topology*, 41(4):807–834, 2002.
- [Tes20] Romain Tessera. On coarse embeddings of amenable groups into hyperbolic graphs. *arXiv preprint:2010.07205*, 2020.
- [Wei08] Richard M Weiss. *The Structure of Affine Buildings.(AM-168)*. Princeton University Press, 2008.
- [Wen05] Stefan Wenger. Isoperimetric inequalities of euclidean type in metric spaces. *GFAA, Geom. funct. anal.*, 5:534–554, 2005.

- [Wen11] Stefan Wenger. The asymptotic rank of metric spaces. *Commentarii Mathematici Helvetici*, 86(2):247–275, 2011.
- [Yu00] Guoliang Yu. The coarse Baum-Connes conjecture for spaces which admit a uniform embedding into hilbert space. , *Invent. Math*, 139, 2000.

UNIVERSITÉ DE PARIS, CNRS, IMJ-PRG
oussama.bensaid@imj-prg.fr