

Homophily in preferences or meetings? Identifying and estimating an iterative network formation model*

Luis Antonio Fantozzi Alvarez[†] Cristine Campos de Xavier Pinto[‡]
Vladimir Pinheiro Ponczek[§]

January 2022

Abstract

Is homophily in social and economic networks driven by a taste for homogeneity (preferences) or by a higher probability of meeting individuals with similar attributes (opportunity)? This paper studies identification and estimation of an iterative network game that distinguishes between these two mechanisms. Our approach enables us to assess the counterfactual effects across time of changing the meeting protocol between agents. As an application, we study the role of preferences and meetings in shaping classroom friendship networks in Brazil. We find that in a network structure in which homophily due to preferences is stronger than homophily due to meeting opportunities, tracking students may lead to welfare improvement, but the relative benefit of this policy diminishes over the school year.

Keywords: homophily; network formation; school tracking.

*We would like to thank Angelo Mele, Áureo de Paula, Braz Camargo, Bruno Ferman, Eduardo Mendes, Ricardo Masini and Sérgio Firpo for their useful comments and suggestions. We are also grateful to seminar participants at PUC-Rio, the 41st SBE meeting, the 2020 Econometric Society World Congress, Penn State Econometrics seminar, LAMES 2021, and the 2021 SBE seminar series. Luis Alvarez gratefully acknowledges financial support from Capes. Cristine Pinto and Vladimir Ponczek acknowledge financial support from CNPq.

[†]Doctoral candidate in Statistics, IME/USP: alvarez@ime.usp.br.

[‡]Associate Professor of Insper Institute of Education and Research: CristineCXP@insper.edu.br.

[§]Associate Professor of Economics, EESP/FGV: vponczek@fgv.br.

1 Introduction

Homophily, the observed tendency of agents with similar attributes to maintain relationships, is a salient feature in social and economic networks (Chandrasekhar, 2016; Jackson, 2010, and references therein). Inasmuch as it may drive network formation, homophily can produce relevant effects in outcomes as diverse as smoking behavior (Badev, 2021), test scores (Hsieh and Lee, 2016; Goldsmith-Pinkham and Imbens, 2013), the adoption of health innovations (Centola, 2011), and health outcomes (Kadelka and McCombs, 2021). As argued by Jackson (2021), such examples suggest that taking a proper account of homophily could improve the understanding and design of policies that aim to reduce social and economic inequality. It is therefore unsurprising that the appropriate modelling of homophily has been a focus in the recent push for estimable econometric models of network formation (Goldsmith-Pinkham and Imbens, 2013; Chandrasekhar and Jackson, 2016; Mele, 2017; Graham, 2016, 2017).

Homophily that is due to choice is distinct from homophily that is due to opportunity.¹ We shall label the former homophily in “preferences”; and the latter homophily in “meetings”. We provide an estimable econometric model that accounts for both types of homophily. This distinction has an important role: public policy may be able to alter the meeting technology between agents (say, by desegregating environments), but it may be less successful in changing preferences. So, the effect of a public policy that aims at changing individual connections (e.g. a tracking policy or a policy that induces people to move to a different neighborhood) depends on the type of homophily that is prevalent in that network. Theoretical models that distinguish between these mechanisms are offered by Currarini et al. (2009) and Bramoullé et al. (2012).² However, these models have some relevant limitations: first, they focus on steady-state or “long-run” behaviour, which may not be appropriate in settings where transitional dynamics may matter (as in our empirical application); second, they are either purely probabilistic³ (Bramoullé et al., 2012) or, in the case of Currarini et al. (2009), allow for only a restrictive set of pay-offs from rela-

¹Cf. Jackson (2010, p. 68) for a discussion.

²See also Currarini et al. (2010) for an attempt at estimation.

³In contrast to strategic models of network formation. See Jackson (2010) and de Paula (2017) for examples.

tionships.⁴ These limitations render these models unfit for some empirical analyses. This paper intends to fill this gap in the literature.

We study identification and estimation of a sequential network-formation algorithm originally developed by Mele (2017) (and by Christakis et al. (2020) and Badev (2021)), where agents meet sequentially in pairs in order to revise their relationship status. The model is well-grounded in the theoretical literature of strategic network formation (Jackson and Watts, 2002) and allows specifications that account for both “homophilies”. However, our approach differs from previous work in several aspects. While Mele (2017) discusses identification and estimation of utility parameters based on the model’s induced stationary distribution under large- and many-network asymptotics, we study identification and estimation of *both* preference- and meeting-related parameters under many-network asymptotics in a setting where networks are observed at *two* points of time. Given observation of several networks at two points in time, our results allow us to estimate *both* preference and meeting parameters. We use these to assess the counterfactual effects of changes in the meeting technology between agents – something that previous work has been unable to do.⁵ In addition, by identifying and estimating *both* meeting- and preference-related parameters, our methodology enables us to analyze the effects of policies *along the transition* to a new steady state, and not just at the model’s stationary (long-run) distribution. Our results also cover a larger – and arguably less restrictive – class of pay-offs and meeting processes than those of Mele (2017), who assumes that utilities admit a potential function and meeting probabilities do not depend on the existence of a link between agents in the current network. By studying identification and estimation of general classes of preference- and meeting-related parameters in possibly off-stationary-equilibrium settings, we hope to contribute to the model’s applicability and empirical usefulness, especially in conducting counterfactual analyses. We focus on estimation using EP-ABC (Barthelmé and Chopin, 2014; Barthelmé et al., 2018), a likelihood-free Bayesian algorithm that can deliver estimates relatively quickly.⁶ We believe this further enhances the usefulness of our

⁴Pay-offs of Currarini et al. (2009) depend only on the number of relationships with individuals of the same or different types. There is no role for indirect benefits.

⁵In fact, as we show in Section B of the Supplemental Appendix, meeting parameters are unidentified in the setting of Mele (2017).

⁶In a recent paper, Battaglini et al. (2021) use a variation of the standard ABC algorithm (see Sec-

approach.

As an application, we study how “homophilies” structure network formation in primary schools in Brazil (Pinto and Ponczek, 2017). We consider 30 municipal elementary schools in Recife, Pernambuco, for which baseline (early 2014) and follow-up (late 2014) data on 3rd- and 5th-grade intra-classroom friendship networks was collected. Using this information, we structurally estimate our model. We then assess how changes in the meeting technology between classmates impact homophily in friendships. Our results suggest that removing biases in meeting opportunities (shutting down homophily in the meeting process) does not decrease observed homophily patterns in students’ cognitive skills. By contrast, in a counterfactual scenario where the role of preferences is excluded from the network formation process, the probability that a student maintains a friendship with a classmate with a different level of cognitive skills increases. We provide evidence that both types of homophily are important determinants of the edges in our networks; although homophily due to preferences does appear to be more important. In this context, a tracking policy that reallocates students between classrooms according to their cognitive skills leads to welfare improvements – as students benefit from connecting with similar individuals – though this benefit appears to diminish (*vis-à-vis* leaving the network process unchanged) in the long run. Given the opportunity to connect with similar people, students have a positive jump in welfare in the short run. However, after they make their new friendships, they tend to keep these links, and the impact of the policy decreases in the long run – inasmuch that, by the end of the school year, current welfare is roughly the same in both the base and counterfactual scenarios.

In the next sections, we introduce the network formation game under consideration (Section 2); explore identification when information on the network structure is available at two distinct points of time (Section 3); and discuss estimation (Section 4). Section 5 presents the results of our application. Section 6 concludes.

tion 4.2) to estimate the parameters of a network formation game. The authors use the structure of the game to speed up their implementation. In contrast, we consider a variation of the ABC algorithm that uses the structure of the data to reduce the computational toll of estimation.

2 Setup

The setup expands upon the work of Mele (2017). We consider a network game with a finite set of agents $\mathcal{I} := \{1, 2, \dots, N\}$. Each agent $i \in \mathcal{I}$ is endowed with a $k \times 1$ vector of exogenous characteristics W_i . These vectors are stacked on matrix $X := \begin{bmatrix} W_1 & W_2 & \dots & W_N \end{bmatrix}'$. Agents' characteristics are drawn according to law \mathbb{P}_X before the game starts and remain *fixed* throughout. We denote the support of X by \mathcal{X} and a realization of X by an element $x \in \mathcal{X}$.

Time is discrete. At each round $t \in \mathbb{N}$ of the network formation process, agents' relations are described by a *directed network*. Information on the network is stored on an $N \times N$ *adjacency matrix*, with entry $g_{ij} = 1$ if i lists j as a friend and 0 otherwise.⁷ By assumption, $g_{ii} = 0$ for all $i \in \mathcal{I}$. We denote the set of all $2^{N(N-1)}$ possible adjacency matrices by \mathcal{G} .

Agent i 's utility from a network g when covariates are $X = x$ is described by a utility function $u_i : \mathcal{G} \times \mathcal{X} \mapsto \mathbb{R}$, $u_i(g, x)$. Utility may depend on the entire network and on the entire set of agents' covariates.

Agents are myopic, i.e. they form, maintain, or sever relationships based on the current utility these bring. In each round, a *matching process* m^t selects a pair of agents (i, j) . The matching process m^t is a stochastic process $\{m^t : t \in \mathbb{N}\}$ over $\mathcal{M} := \{(i, j) \in \mathcal{I} \times \mathcal{I} : i \neq j\}$. If the pair (i, j) is selected, agent i will get to choose whether to form/maintain or not form/sever a relationship with j . After the matching process selects a pair of agents, a pair of choice-specific idiosyncratic shocks $(\epsilon_{ij,t}(0), \epsilon_{ij,t}(1))$ are drawn, where $\epsilon_{ij,t}(1)$ corresponds to the taste shock in forming/maintaining a relationship with j at time t . These shocks are unobserved by the econometrician and enter additively in the utility of each choice.⁸ Given that choice is myopic, agent i forms/maintain a relation with j if, and only if

$$u_i([1, g_{-ij}], X) + \epsilon_{ij,t}(1) \geq u_i([0, g_{-ij}], X) + \epsilon_{ij,t}(0) \quad (1)$$

⁷Our model and main results are readily extended to the case of an *undirected* network where friendships are forcibly symmetric.

⁸Additive separability of unobserved shocks is a common assumption in the econometric literature on discrete choice and games (Aguirregabiria and Mira, 2010), though it is not innocuous. In our setting, it precludes factors unobserved by the econometrician from affecting the marginal effect of covariates and network characteristics on utility (homophily in preferences), as taste shocks act as pure location shifts.

where $[a, g_{-ij}]$ denotes an adjacency matrix with all entries equal to matrix g except for entry ij , which equals a .

The next two assumptions constrain the meeting process and the distribution of shocks.

Assumption 2.1. The meeting process $\{m^t : t \in \mathbb{N}\}$ is described by a *time-invariant* matching function $\rho : \mathcal{M} \times \mathcal{G} \times \mathcal{X} \mapsto [0, 1]$, where $\rho((i, j), g, x)$ is the probability that (i, j) is selected when covariates are $X = x$ and *the previous-round network was g* . Moreover, for all $g \in \mathcal{G}$, $x \in \mathcal{X}$, $(i, j) \in \mathcal{M}$, $\rho((i, j), g, x) > 0$.

Assumption 2.1 constrains the matching function to assign positive probability to all possible meetings under all possible values of covariates and previous-round networks. Note that this allows for dependence on the existence of previous-round links, which was not permitted by Mele (2017).⁹

Assumption 2.2. Shocks are iid draws across pairs and time, independent from X , from a *known* distribution $(\epsilon(0), \epsilon(1))' \sim F_\epsilon$ which is absolutely continuous with respect to the Lebesgue measure on \mathbb{R}^2 and which has positive density almost everywhere.

Conditional on $X = x$, we have that, under Assumptions 2.1 and 2.2 – and given an initial distribution $\mu_0(x) \in \Delta(\mathcal{G})$ ¹⁰ –, the network game just described induces a homogenous Markov chain $\{g^t : t \in \mathbb{N} \cup \{0\}\}$ on the set of networks \mathcal{G} . The $2^{N(N-1)} \times 2^{N(N-1)}$ transition matrix $\Pi(x)$ has entries $\Pi(x)_{gw}$, $g, w \in \mathcal{G}$, which specify the probability of transitioning to w given the current period network g .

For each $g \in \mathcal{G}$, define $N(g) := \{w \in \mathcal{G} \setminus \{g\} : \exists! (i, j) \in \mathcal{M}, g_{ij} \neq w_{ij}\}$ as the set of networks that differ from g in *exactly* one edge. Entries of $\Pi(x)$ take the form

$$\Pi(x)_{gw} = \begin{cases} \rho((i, j), g, x) F_{\epsilon(g_{ij}) - \epsilon(w_{ij})}(u_i(w, x) - u_i(g, x)) & \text{if } w \in N(g), g_{ij} \neq w_{ij} \\ \sum_{(i, j) \in \mathcal{M}} \rho((i, j), g, x) F_{\epsilon(1 - g_{ij}) - \epsilon(g_{ij})}(u_i(g, x) - u_i([1 - g_{ij}, g_{-ij}], x)) & \text{if } g = w \\ 0 & \text{elsewhere} \end{cases} \quad (2)$$

where $F_{\epsilon(1) - \epsilon(0)}$ and $F_{\epsilon(0) - \epsilon(1)}$ denote the distribution function of the difference in shocks.

⁹Do also note that, unlike Mele (2017), we do *not* assume that utilities admit a potential function. See Section 3.3 for a discussion.

¹⁰We denote by $\Delta(\mathcal{G})$ the set of all probability distributions on \mathcal{G} .

Remark 2.1. The transition matrix is irreducible and aperiodic. By Assumptions 2.1 and 2.2, the first and second cases in (2) are always positive for any $g \in \mathcal{G}$. We can thus always reach any other network w starting from any g with positive probability in finite time (irreducibility). Since the chain is irreducible and contains a self-loop ($\Pi(x)_{gg} > 0$), it is also aperiodic.

We next look for (conditional) stationary distributions. A stationary distribution is an element $\pi(x) \in \Delta(\mathcal{G})$ satisfying $\pi(x) = \Pi(x)' \pi(x)$.

Remark 2.2. The transition matrix $\Pi(x)$ admits a unique stationary distribution, which is a direct consequence of the Perron-Frobenius theorem for nonnegative irreducible matrices (Horn and Johnson, 2012, Theorem 8.4.4). Moreover, as the chain is irreducible and aperiodic, we have that, for any $\pi_0 \in \Delta(\mathcal{G})$, $\lim_{t \rightarrow \infty} (\Pi(x)^t)' \pi_0 = \pi(x)$ (Norris, 1997, Theorem 1.8.3), so we may interpret the invariant distribution as a “long-run” distribution (Mele, 2017).

Remark 2.3. The stationary distribution puts positive mass over all network configurations. Indeed, since $\pi(x)$ is a distribution, there exists some $g_0 \in \mathcal{G}$ such that $\pi(g_0|x) > 0$. Fix $w \in \mathcal{G}$. Since the chain is irreducible, there exists $k \in \mathbb{N}$ such that $(\Pi(x)^k)_{g_0,w} > 0$, where $(\Pi(x)^k)_{g_0,w} > 0$ denotes the (g_0, w) entry of $\Pi(x)^k$. However $\pi(x) = (\Pi(x)^k)' \pi(x) \implies \pi(w|x) > 0$.

3 Identification

In this section, we study identification under many-network asymptotics. In particular, we assume that we have access to a random sample (iid across c) of C networks $\{G_c^{T_0}, G_c^{T_1}, X_c\}_{c=1}^C$ stemming from the network formation game described in Section 2.¹¹ In this context, $G_c^{T_0}$ and $G_c^{T_1}$ are observations of network c over two (possibly nonconsecutive) periods¹² (labelled *first* and *second*); and X_c is the set of covariates associated with network

¹¹Under large-network asymptotics, we have access to a single (a few) network(s) with a large number of players.

¹²In Section B of the supplemental appendix, we briefly discuss (non)identification when only one period of data stemming from the stationary distribution of the network formation game is available.

c . We recall the law of X_c is \mathbb{P}_X , as X_c is a copy of X (i.e. a random variable with the same law as X). As in the previous section, realizations of X are denoted by lower-case letters, i.e. elements $x \in \mathcal{X}$.

Denote by $\Pi(X; \theta_0)$ the transition matrix under covariates X ; where $\theta_0 := ((u_i)_{i=1}^N, \rho)$ are the “true” parameters (functions). Further write τ_0 for the number of *rounds* of the network formation game that have taken place between the first and second period. We can identify $\Pi(X; \theta_0)^{\tau_0}$, the transition matrix to the power of the number of rounds of the network formation game which took place between the first and second period (τ_0), provided that the first-period conditional distribution, which we denote by $\pi_0(X)$, is such that $\pi_0(X) \gg 0$ \mathbb{P}_X -a.s. To see this more clearly, suppose X is empty. In this case, we could consistently estimate $(\Pi^{\tau_0})_{gw}$, $g, w \in \mathcal{G}$, by $(\widehat{\Pi}^{\tau_0})_{gw} = \sum_{c=1}^C \mathbb{1}\{G_c^{T_0} = g, G_c^{T_1} = w\} / \sum_{c=1}^C \mathbb{1}\{G_c^{T_0} = g\}$,¹³ provided $\mathbb{P}[G_c^{T_0} = g] > 0$.¹⁴ The next assumption summarizes this requirement.

Assumption 3.1 (Full support). $\pi_0(g|X) > 0$ for all $g \in \mathcal{G}$ \mathbb{P}_X -a.s.

Since $\Pi(X; \theta_0)^{\tau_0}$ is identified from the data under Assumption 3.1, the identification problem subsumes to (denoting by Θ the parameter space):¹⁵

$$\forall (\theta, \tau) \in \Theta \times \mathbb{N}, \quad (\theta, \tau) \neq (\theta_0, \tau_0) \implies \Pi(X; \theta)^\tau \neq \Pi(X; \theta_0)^{\tau_0}$$

where the inequality must hold with positive probability over the distribution of X (Newey and McFadden, 1994).

¹³In our setting, an agent is described by her set of exogenous characteristics W_i . When no such traits are included in the model, two adjacency matrices g and g' are deemed equal if they are equal up to relabeling of agents. When covariates are included, one compares adjacency matrices for a given labeling of agents – and such labeling will define the distance between the covariate matrices X and X' .

¹⁴The case where X has discrete support is similar to the case where no covariates are included: for each $x \in \mathcal{X}$, we consider observations c such that $X_c = x$ up to a relabeling of agents in c . We then use these observations to compute the transition probability given $X = x$. In the case where X contains continuous covariates, a consistent estimator is given by a kernel estimator (Li and Racine, 2006). Clearly, these estimators would behave poorly in most practical settings (especially if the number of players is moderate to large). We do not suggest to use them in practice, though; we rely on consistency only as an indirect argument for identification.

¹⁵Abstracting from measurability concerns, this is the set of utilities and matching functions that satisfy the assumptions in Section 2.

In our statement of the identification problem, the number of rounds in the network formation game is assumed to be unknown. There is no reason to expect τ_0 , the true number of rounds, to be known *a priori* by the researcher, unless the network formation algorithm has a clear empirical interpretation. Nonetheless, it is still possible to identify τ_0 under some assumptions. For all $\theta \in \Theta$ and $x \in \mathcal{X}$, $\Pi(x; \theta)$ is irreducible and has a strictly positive main diagonal. It clear, then, that the number of strictly positive entries in $\Pi(x; \theta)^\tau$ is nondecreasing in τ . Moreover, this number is strictly increasing for $\tau \leq N(N-1)$ ¹⁶ and does not depend on the choice of (x, θ) . Thus, provided that $\tau_0 \leq N(N-1)$, we can identify τ_0 by “counting”¹⁷ the number of positive entries in $\Pi(X; \theta_0)^{\tau_0}$.

Assumption 3.2 (Upper bound on τ_0). The number of rounds which took place in the network formation game between the first and second period (τ_0) is smaller than or equal to $N(N-1)$.

Remark 3.1. Under Assumption 3.2, τ_0 is identified.

A similar assumption is considered in Christakis et al. (2020), where the authors assume that $\tau_0 = N(N-1)/2$ (they work with an undirected network, so the set of available matches is divided by two) and that all meeting opportunities are played (though in an unknown order). In their setting, however, the assumption’s main purpose is to reduce the computational toll of evaluating the model likelihood (see Section 4.2 for a similar discussion). In our case, we require it for identification. We also emphasize that the bound in Assumption 3.2 is more or less restrictive, depending on the setting. Knowledge of the particular application in mind should help to assess its appropriability. Finally, we note that our main identification result (Proposition 3.1) holds irrespective of the bound, provided that τ_0 is identified or known *a priori*.

Under Assumption 3.2, we may thus assume, without loss, that τ_0 is known.

In the next subsections, we discuss identification of θ_0 .

¹⁶ $N(N-1)$ is the minimum number of rounds required for the probability of transitioning from a “fully empty” network ($g_{ij} = 0$ for all ij) to a “fully connected” network ($g_{ij} = 1$ for all ij) to be strictly positive.

¹⁷We provide a consistent estimator for τ_0 in Section 4.1.

3.1 Identification without restrictions

To illustrate the difficulty of identification without imposing further restrictions, let us briefly analyze identification of θ_0 from $\Pi(X; \theta_0)$, which is a necessary condition for identification of θ_0 from $\Pi(X; \theta_0)^{\tau_0}$. Indeed, knowledge of $\Pi(X; \theta_0)$ implies knowledge of $\Pi(X; \theta_0)^{\tau_0}$. Thus, in a sense, our analysis in this subsection provides a “best-case” scenario for achieving identification without additional restrictions.

As we do not impose further restrictions in the model, we essentially view X as non-stochastic throughout the remainder of this subsection and suppress dependence of $\Pi(X; \theta)$ on X by writing $\Pi(\theta)$. In order to make the identification problem clearer, define, for all $g \in \mathcal{G}$ and $w \in N(g)$ with $g_{ij} \neq w_{ij}$, $F_{ij}(g, w) := F_{\epsilon(g_{ij}) - \epsilon(w_{ij})}(u_i(w, X) - u_i(g, X))$. Observe that $F_{ij}(g, w) + F_{ij}(w, g) = 1$. Write $\rho_{ij}(g)$ for $\rho((i, j), g, X)$. Observe that $\sum_{(i,j) \in \mathcal{M}} \rho_{ij}(g) = 1$. Let $\gamma := ((\rho_{ij}(g))_{g \in \mathcal{G}, (i,j) \in \mathcal{M}}, (F_{ij}(g, w))_{g \in \mathcal{G}, w \in N(g), g_{ij} \neq w_{ij}})$ be a parameter vector, and γ_0 the “true” parameter. Observe that, under Assumptions 2.1 and 2.2, the parameter space, which we denote by Γ , is a subset of $\mathbb{R}_{++}^{\dim \gamma_0}$, an open set. Put another way, $\Gamma = \{\gamma \in \mathbb{R}_{++}^{\dim \gamma_0} : F_{ij}(g, w) + F_{ij}(w, g) = 1, \sum_{(k,l) \in \mathcal{M}} \rho_{k,l}(g) = 1 \text{ for all } g \in \mathcal{G}, w \in N(g) \text{ with } g_{ij} \neq w_{ij}\}$. Identification from the transition matrix thus requires us to show that, for all $\gamma' \in \Gamma$, $\gamma' \neq \gamma_0 \implies \Pi(\gamma') \neq \Pi(\gamma_0)$, where $\Pi(\gamma)$ is the matrix in (2) constructed under γ . If we can uniquely recover γ_0 from $\Pi(\gamma_0)$, then we can recover differences in utilities, $u_i(g, X) - u_i(w, X)$, for all $i \in \mathcal{I}$ and $g, w \in \mathcal{G}$, as $F_{\epsilon(1) - \epsilon(0)}$ is invertible under Assumption 2.2. Levels (and thus θ_0) are then identified under a location normalization on pay-offs (e.g. $u_i(g_0, X) = 0$ for all i and some g_0).

Observe that $\dim \gamma = N(N-1)2^{N(N-1)} + N(N-1)2^{N(N-1)}$. The first summand is the dimension of $(p_{ij}(g) : g \in \mathcal{G}, (i, j) \in \mathcal{M})$ and the second term is the dimension of $(F_{ij}(g, w) : (i, j) \in \mathcal{M}, g \in \mathcal{G}, w \in N(g), w_{ij} \neq g_{ij})$. Matrix $\Pi(\gamma)$ has $2^{N(N-1)}(N(N-1)+1)$ strictly positive entries. The parameter space imposes $2^{N(N-1)}$ restrictions of the type $\sum_{(i,j) \in \mathcal{M}} \rho_{i,j}(g) = 1$ and $N(N-1)2^{N(N-1)}/2$ restrictions of the type $F_{ij}(g, w) + F_{ij}(w, g) = 1$. A simple order condition would thus require

$$2^{N(N-1)} [2N(N-1)] \leq 2^{N(N-1)} [N(N-1) + 1 + N(N-1)/2] \quad (3)$$

which implies that N should be less than or equal to 2. The point is that the map $\gamma \mapsto \Pi(\gamma)$

is nonlinear, so the order condition is neither necessary nor sufficient for identification. Nonetheless, we are able to show directly that the model is identified when $N = 2$.

Claim 3.1. *If $N = 2$, then γ_0 is identified from $\Pi(\gamma_0)$ under Assumptions 2.1, 2.2, and 3.1.*

Proof. See Section C of the supplemental appendix. □

Extending such a direct argument to $N > 2$ is not feasible, as $\Pi(\gamma)$ is a $2^{N(N-1)} \times 2^{N(N-1)}$ matrix. Notice that (3) implies that the rank condition of Rothenberg (1971) for local identification is not satisfied. The problem is that, for this failure of the order condition to be sufficient for nonidentification, the Jacobian of $\gamma \mapsto \Pi(\gamma)$ must be rank-regular (i.e. it must have constant rank in a neighborhood of γ_0), which is not trivial to show. Of course, if that were the case, then we would know the model is nonidentified for $N > 2$.

Given the difficulty of establishing identification without imposing further restrictions, even when $\Pi(X; \theta_0)$ is known (or $\tau_0 = 1$), in the next subsections we explore the identifying power of restrictions on (i) how covariates affect utilities and the matching function; and (ii) how the network structure affects pay-offs and meetings. To make both the exposition and proofs clearer, in what follows we maintain the notation introduced in this section, and dependence of objects on covariates remains implicit whenever it does not cause confusion.

3.2 Identification with covariates

In this subsection, we explore the identifying power of restrictions on covariates. We work in the environment where we observe two periods of data and we assume τ_0 is identified or known *a priori*. We follow an identification at infinity approach in order to identify θ_0 (Tamer, 2003; Bajari et al., 2010).

To make the dependency in covariates explicit, we write $X_i^u(g)$ for the covariates that enter the utility of agent i under network g , i.e. we shall write $u_i(g, X) = u_i(g, X_i^u(g))$ for all $i \in \mathcal{I}$, $g \in \mathcal{G}$. We use $X^m(g)$ for the covariates that enter the matching function under network g , i.e. $\rho((i, j), g, X) = \rho((i, j), g, X^m(g))$ for all $(i, j) \in \mathcal{M}$ and $g \in \mathcal{G}$. We also define $u_i(w, X_i^u(w)) - u_i(g, X_i^u(g)) =: \delta_{ij}(g, w, X_{ij}^u(g, w))$, the gain in utility from each choice, for all $g \in \mathcal{G}$ and $w \in N(g)$ with $w_{ij} \neq g_{ij}$. In this case, $X_{ij}^u(g, w)$ is the subvector

of $[X_i^u(g), X_i^u(w)]$ with the covariates relevant in the marginal gain of i moving from g to w . We write $\Pi(X)^{\tau_0} = \Pi(X; \theta_0)^{\tau_0}$ for the observed transition matrix. In our case, we may take $X = [(X_{ij}^u(g, w))_{g \in \mathcal{G}, w \in N(g), w_{ij} \neq g_{ij}}, (X^m(g))_{g \in \mathcal{G}}]$. Finally, we use the notation $A \setminus B$ for the subvector of A such that, up to permutations, $A = [A \setminus B, B]$.

We next impose the following assumptions.

Assumption 3.3 (Location normalization). There exists some $g_0 \in \mathcal{G}$ such that $u_i(g_0, X_i^u(g_0)) = 0$ for all $i \in \mathcal{I}$.

Such a normalization is required in order to identify utilities in levels.

The next assumption states our exclusion restriction.

Assumption 3.4 (Large support exclusion restriction). For all $g \in \mathcal{G}$, $w \in N(g)$, $g_{ij} \neq w_{ij}$, there exists an $m \times 1$ subvector $Z_{ij}^u(g, w)$ of $X_{ij}^u(g, w)$, i.e. $X_{ij}^u(g, w) = [Z_{ij}^u(g, w), \tilde{X}_{ij}^u(g, w)]$, such that no covariate in $Z_{ij}^u(g, w)$ is an element of $X^m(g)$. Moreover, $Z_{ij}^u(g, w)$ admits a conditional Lebesgue density $f(Z_{ij}^u(g, w) | \tilde{X}_{ij}^u(g, w), X^m(g))$ that is positive a.e. (for almost all realizations of $[\tilde{X}_{ij}^u(g, w), X^m(g)]$); and there exists $\vec{r} \in \mathbb{R}^m$ s.t. $\lim_{t \rightarrow \infty} \delta_{ij}(g, w, [Z_{ij}^u(g, w) = t\vec{r}, \tilde{X}_{ij}^u(g, w)]) = \infty$.

Assumption 3.4 requires that, for each $g \in \mathcal{G}$, large support covariates be included in the marginal gain of each agent's choice under g , but excluded from the matching function under g . These covariates should admit, with positive probability, sufficiently "high" realizations such that the (conditional on X) probability of an agent selected by the matching process "accepting" a transition from g can be made arbitrarily close to unity.

The previous restrictions imply the next result:

Lemma 3.1. *Under Assumptions 2.1, 2.2, 3.1, 3.3 and 3.4, θ_0 is identified when $\tau_0 = 1$.*

Proof. See Appendix A.1. □

The corollary below provides a sufficient condition for identification of θ_0 for any known or identified τ_0 . It states a sufficient condition for recovery of $\Pi(X; \theta_0)^{\tau_0}$ from $\Pi(X; \theta_0)$. One can then apply Lemma 3.1 to achieve identification.

Corollary 3.1. *If $\Pi(X; \theta_0)$ is a.s. diagonalizable with the appropriate eigenvalue signs (nonnegative if τ_0 even), then, under the assumptions in Lemma 3.1, θ_0 is identified for any τ_0 known or identified.*

More generally, conditions for uniqueness of a stochastic τ_0 th root of a transition matrix are quite complicated. Higham and Lin (2011) give examples and sufficient conditions.

Remark 3.2. Lemma 3.1 would similarly hold if the large support variable were included in the matching function (but not in utilities). This may be more appropriate in some applied settings.

When $\tau_0 \geq 2$ and we do not know if $\Pi(X; \theta_0)$ is “appropriately” diagonalizable, we need stronger exclusion restrictions. We state a sufficient version (for all τ_0 identified or known) of this assumption below.

Assumption 3.5. The exclusion restriction in 3.4 holds as: no covariate in $Z_{ij}^u(g, w)$ is included in $[X^m(g), X^m(w), (X_{kl}^u(g, [1 - g_{kl}, g_{-kl}]), X_{kl}^u(w, [1 - w_{kl}, w_{-kl}]))_{(k,l) \neq (i,j)}]$; with $f(Z_{ij}^u(g, w) | X \setminus Z_{ij}^u(g, w))$ positive a.e. (for almost all realizations of $[X \setminus Z_{ij}^u(g, w)]$).

This stronger exclusion restriction requires that the large support covariates included in agent i 's marginal gain from transitioning from g to w be excluded not only from the matching function under g ; but also from the matching function under w and from all other pairs of agents' marginal gains from transitioning from g or w .

The next proposition presents the main identification result of the paper: identification of the vector of model parameters for any fixed τ_0 .

Proposition 3.1. *Suppose Assumptions 2.1, 2.2, 3.1, 3.3, and 3.5 hold. Then the model is identified for any τ_0 known or identified.*

Proof. See Appendix A.2. □

The intuition behind the main identification result of the paper is that when the instruments attain large values in their support, the probability of staying/returning to the same network after τ_0 rounds depends solely on the matching process, which remains unchanged by the exclusion restriction. Similarly, large values of the instruments for pairs $ms \neq ij$

imply that the probability of, starting from g_0 , arriving at network $w \in N(g)$, $w_{ij} \neq g_{ij}$ after τ_0 rounds, depends only on the meeting process **and** on the decisions of pair ij . These limiting probabilities then allow us to identify the desired parameters.

In Section D in the supplemental appendix, we show that identification would similarly hold if an analogous exclusion restriction on the matching function were true.

Remark 3.3 (A possible parametrization). If we take $\rho((i, j), g, X^m(g)) = \frac{\exp(\alpha'_g X_{ij}^m(g))}{\sum_{(k,l) \in \mathcal{M}} \exp(\alpha'_g X_{kl}^m(g))}$ and $u_i(g, X_i^u(g)) = \beta'_g X_i^u(g)$, where $X_i^u(g)$ may include other individuals' characteristics, then the model is identified under the previous restrictions, provided the usual rank conditions hold (cf. Amemiya (1985, p.286-292); also McFadden (1973)).¹⁸

Remark 3.4 (Additional parametric restrictions). While we focus on identification without parametric restrictions (besides those implied by the base model), it becomes clear from the proofs of our results that, if we are willing to impose further parametrizations, then it is possible to reduce the number of required instruments in the model. For example, if meeting probabilities are parameterized such that $\rho((i, j), g, X^m(g)) = \rho_0((i, j), X^m(g))$ for all $g \in \mathcal{G}$ and some base function ρ_0 , then it is possible, for each (i, j) , to identify $x \mapsto \rho_0((i, j), x)$ from the transitions starting on a single $g_0 \in \mathcal{G}$ and then *extrapolate it* to the remaining networks.

3.3 Identification via the network structure

Mele (2017) assumes that $\rho_{ij}(g, X) = \rho_{ij}([1 - g_{ij}, g_{-ij}], X)$ for all $g \in \mathcal{G}$. In other words, meeting probabilities do not depend on there being a link between ij . This constitutes an exclusion restriction with identifying power in our environment. Indeed, if $\tau_0 = 1$ and this hypothesis holds, $\Pi_{g, [1 - g_{ij}, g_{-ij}]} / \Pi_{[1 - g_{ij}, g_{-ij}], g} = F_{ij}(g, [1 - g_{ij}, g_{-ij}]) / (1 - F_{ij}(g, [1 - g_{ij}, g_{-ij}]))$, which establishes identification of utilities (and meeting probabilities thereupon) under a location normalization. For $\tau_0 > 1$, identification is less immediate,¹⁹ but we can rely on a

¹⁸Specifically. we would require $\mathbb{E}[(X_i^u(w) - X_i^u(g))(X_i^u(w) - X_i^u(g))']$ to have full rank for all $i \in \mathcal{I}$, $g \in \mathcal{G}$, $w \in N(g)$, with $X_i^u(g_0) = \vec{0}$ for all $i \in \mathcal{I}$; and $\mathbb{E}[\tilde{X}^m(g)' \tilde{X}^m(g)]$ to have full rank for all g , where $\tilde{X}^m(g)' = \left[(X_{1,1}^m(g) - \bar{X}^m(g))' \quad (X_{1,2}^m(g) - \bar{X}^m(g))' \quad \dots \quad (X_{N,(N-1)}^m(g) - \bar{X}^m(g))' \right]$ and $\bar{X}^m(g) = \sum_{(i,j) \in \mathcal{M}} \rho((i, j), g, X^m(g)) X_{ij}^m(g)$.

¹⁹We note, however, that, under such restriction on meeting probabilities, Equation (10) in Appendix A.2

sufficient condition such as “appropriate” diagonalizability, as in Corollary 3.1, to achieve identification.

If we further assume that (1) taste shocks are independent EV type 1; and (2) utility functions admit a potential function²⁰ $Q : \mathcal{G} \times \mathcal{X} \mapsto \mathbb{R}$; then the model’s (conditional on X) network stationary distribution is in the exponential family, i.e. $\pi(g|X) \propto \exp(Q(g, X))$ (Mele, 2017, Theorem 1). If we assume that $\{G_c^{T_0}\}_c$ are drawn from the model’s stationary distribution, then the model’s potential (and hence marginal utilities) is identified under standard assumptions (Newey and McFadden, 1994). Provided that utility functions in the game played before period T_0 remain unaltered in the game played between period T_0 and period T_1 , we can use the T_0 distribution to help identify the model.²¹ A necessary condition for this equality in utilities, provided that between period T_0 and period T_1 the matching function satisfies the restriction of Mele (2017), is that the period T_0 network distribution equals the period T_1 distribution.²² This is a testable condition.

More generally, we could try to achieve identification by restricting how payoffs are affected by the network structure. This approach is followed by de Paula et al. (2018) and Sheng (2020), who assume that network observations are pairwise-stable realizations of a (static) simultaneous-move complete-information game. In their setting, pairwise stability only enables partial identification. We recognize that further restrictions on how the network structure affects payoffs may enable point-identification in our setting, though we do not try to analyze these conditions in a general environment.

entails that $(\Pi^2)_{g, [1-g_{ij}, g_{-ij}]} / (\Pi^2)_{[1-g_{ij}, g_{-ij}], g} = F_{ij}(g, [1-g_{ij}, g_{-ij}]) / (1 - F_{ij}(g, [1-g_{ij}, g_{-ij}])),$ which identifies utilities when $\tau_0 = 2$. Identification of meeting probabilities is not immediate in this case, though.

²⁰A *potential function* is a map $Q : \mathcal{G} \times \mathcal{X} \mapsto \mathbb{R}$ satisfying $Q([1, g_{-ij}], X) - Q([0, g_{-ij}], X) = u_i([1, g_{-ij}], X) - u_i([0, g_{-ij}], X)$ for all $(i, j) \in \mathcal{M}, g \in \mathcal{G}$.

²¹We do not need to assume matching functions remain unaltered, just that before period T_0 they obey the restriction of Mele (2017).

²²If the equality of distributions is conditional on X , this is also a sufficient condition, as it is assumed that the potential function before period T_0 is identified from $\pi_0(g|X)$.

4 Estimation

In this section, we will analyze estimation. We have access to a sample of C networks, $\{G_c^{T_0}, G_c^{T_1}, X_c\}_{c=1}^C$, stemming from the network formation game previously described.

4.1 Estimating τ_0

We first propose to estimate τ_0 as follows:

$$\hat{\tau} = \max_c \{\|G_c^{T_1} - G_c^{T_0}\|_F\} \quad (4)$$

where $\|\cdot\|_F$ is the Froebenius norm. This estimator is intuitive: it amounts to “counting” in each network the number of differing edges between periods and then taking the maximum. It turns out that, under iid sampling and the bound in 3.2, $\hat{\tau} \xrightarrow{a.s.} \tau_0$.

Lemma 4.1. *Suppose $\{G_c^{T_0}, G_c^{T_1}, X_c\}_{c=1}^C$ is a random sample (iid across c). Under Assumptions 2.1, 2.2, and 3.2, $\hat{\tau} \xrightarrow{a.s.} \tau_0$.*

Proof. See Appendix A.3. □

As we argue in Appendix A.3, the previous Lemma can be extended to accommodate a setting where the sequence of observations is independently drawn from games with common parameters, but where the distribution of covariates and the number of players may vary with c .

4.2 Estimation of preference and meeting parameters

Let vector $\beta_0 \in \mathbb{B} \subseteq \mathbb{R}^l$ encompass a parametrization of preferences and meetings, i.e. $u_i(g, X) = u_i(g, X; \beta_0)$ and $\rho_{ij}(g, X) = \rho_{ij}(g, X; \beta_0)$ for all $(i, j) \in \mathcal{M}$, $g \in \mathcal{G}$. The network log-likelihood, conditional on X_c and $G_c^{T_0}$, is

$$l_c(G_c^{T_1} | G_c^{T_0}, X_c; \tau_0, \beta) = \sum_{g \in \mathcal{G}} \mathbb{1}\{G_c^{T_1} = g\} \ln \left((\Pi(X_c; \beta)^{\tau_0})_{G_c^{T_0}, g} \right)$$

and the sample log-likelihood, under an independent sequence of observations, is

$$\mathcal{L}(\{G_c^{T_1}\}_{c=1}^C | \{G_c^{T_0}\}_{c=1}^C; \tau_0, \beta) = \sum_{c=1}^C \sum_{g \in \mathcal{G}} \mathbb{1}\{G_c^{T_1} = g\} \ln \left((\Pi(X_c; \beta)^{\tau_0})_{G_c^{T_0} g} \right).$$

The second-step MLE estimator will thus be

$$\hat{\beta}_{\text{MLE}} \in \operatorname{argmax}_{\beta \in \mathbb{B}} \mathcal{L}(\{G_c^{T_1}\}_{c=1}^C | \{G_c^{T_0}\}_{c=1}^C; \hat{\tau}, \beta)$$

where $\hat{\tau}$ is the estimator discussed in the previous section. This formulation can be easily modified to accommodate for observations of networks with different numbers of players, provided they have common parameters.

Numerically, computation of the likelihood is complicated by the fact that we need to sum over all walks between $G_c^{T_0}$ and $G_c^{T_1}$.²³ For small τ_0 , this is feasible, but for higher values of τ_0 , it becomes impractical. For a given $\tau \in \mathbb{N}$, there are $[N(N-1)+1]^\tau$ walks starting from $G_c^{T_0}$ and ending in some network $g \in \mathcal{G}$. Evaluating the model likelihood requires summing over all walks ending in $G_c^{T_1}$. A “recursive” approach for evaluating the model likelihood would consist of, for each $c \in \{1, 2, \dots, C\}$, “writing down” the formula for each walk iteratively, i.e. starting from $G_c^{T_0}$, compute all possible $N(N-1)+1$ transitions in the first round; then, for each of these $N(N-1)+1$ possible transitions, compute the $N(N-1)+1$ transitions in the second round and multiply each of these probabilities by the probability of the associated transition in the first round, and so on; and then summing over all walks ending in $G_c^{T_1}$. Walks that “strand off” from $G_c^{T_1}$ in some round $r < \tau$ can be excluded from the next steps in the recursion,²⁴ which ameliorates the computational toll, but does not solve it.

Given the above difficulty, an interesting alternative is to work with simulation-based methods, which allows us to bypass direct evaluation of the model likelihood. We take a Bayesian perspective²⁵ and follow an approach known as likelihood-free estimation or

²³A walk between g and w in τ rounds is a sequence of networks g_1, \dots, g_τ such that $g_1 = g$, $g_\tau = w$ and $g_t \in N(g_{t-1}) \cup \{g_{t-1}\}$ for all $t = 2, \dots, \tau$.

²⁴By “strand off” we mean a path of realizations of the stochastic process g_c^t up to round r such that the probability of reaching $G_c^{T_1}$ in $\tau_0 - r$ rounds is 0.

²⁵From the frequentist point of view, a simulated method of moments estimator would be a possibility in our case, though the nonsmoothness of the objective function (which involves indicators of simulated network observations) as well as the poor properties of GMM estimators with many moment conditions

approximate Bayesian computation (ABC) (Sisson and Fan, 2011).²⁶ This method admits a close correspondence to nonparametric (frequentist) estimation (Blum, 2010) and indirect inference (Frazier et al., 2018). The methodology basically requires the researcher to be able to draw a sample (or statistics thereof) from the model given the parameters. Algorithm 1 outlines the simplest accept-reject ABC algorithm in our setting, where S is the maximum number of iterations and $p_0(\beta, \tau)$ is a prior distribution over $\mathbb{B} \times \mathbb{N}$.²⁷

Algorithm 1 Basic accept-reject ABC algorithm

define some tolerance $\epsilon > 0$

define a vector of m statistics $T : \mathcal{G}^C \mapsto \mathbb{R}^m$

compute the observed sample statistics $T_{\text{obs}} := T(\{G_c^{T_1}\}_{c=1}^C)$

for $s \in \{1, 2, \dots, S\}$ **do**

 draw $(\beta_s, \tau_s) \sim p_0$

 generate an artificial sample $\{\tilde{G}_c^{T_1}\}_{c=1}^C$ given $\{G_c^{T_0}, X_c\}_{c=1}^C$ and (β_s, τ_s)

 compute the simulated statistic $T_s := T(\{\tilde{G}_c^{T_1}\})$

 if $\|T_s - T_{\text{obs}}\| \leq \epsilon$, accept (β_s, τ_s)

end for

In this methodology, the researcher must make two crucial choices. One is the tolerance parameter. Here, we can use the recommendations of Li and Fearnhead (2018): we may (transition probabilities) in finite samples (Newey and Smith, 2004) are unappealing. An indirect inference approach is also unappealing, as low-dimensional sufficient statistics are unknown in our context.

²⁶From the Bayesian point of view, an alternative and well-known approach to simplify, but not bypass, a complicated model likelihood is data augmentation (Hobert, 2011). This alternative is not useful in our setting, though, due to the dimensionality of the support of the meeting process. Thus, an approach that conditions the likelihood on the (unobserved) matching process (reducing the number of walks starting at $G_c^{T_0}$ from $[N(N-1)+1]^\tau$ to 2^τ) is unworkable here in our case, since we still have to draw from the matching process distribution conditional on the data and the model parameters.

²⁷In practice, a few improvements can be made upon Algorithm 1 (Li and Fearnhead, 2018). First, we can use importance sampling: instead of drawing from the prior, we may draw from a proposal distribution $q_0(\beta, \tau)$ such that $\text{supp } p_0 \subseteq \text{supp } q_0$. Accepted draws should then be associated with weights $w_s := p_0(\beta_s, \tau_s)/q_0(\beta_s, \tau_s)$. Li and Fearnhead (2018) provide a data-driven method to select the proposal. Second, we can use a “smooth” rejection rule, i.e. we accept a draw with probability $K(\|T_s - T_{\text{obs}}\|/\epsilon)$, where $K(\cdot)$ is a rescaled univariate kernel such that $K(0) = 1$. See Section E of the supplemental appendix for details of this augmented algorithm.

choose ϵ so the algorithm produces a “reasonable” acceptable rate. The second important choice is the vector of statistics. This is closely related to identification: for a proper working of the ABC algorithm, the chosen vector of statistics should be informative of the model’s parameters (Li and Fearnhead, 2018; Frazier et al., 2018).²⁸ As an example, Battaglini et al. (2021) use a version of the ABC algorithm to estimate a network formation game. In their setting, the authors use the characterization of the model equilibrium in order to construct the summary statistics used to assess the quality of a parameter draw.

In Section E of the supplemental appendix, we show that, if we take the vector of summary statistics as the whole second-period network data, then, as $\epsilon \rightarrow 0$ as $S \rightarrow \infty$, the mean of the accepted draws $h(\theta_s)$ converges in probability to the expectation of $h(\cdot)$ with respect to the posterior, where $h(\cdot)$ is a function with finite moments (with respect to the prior distribution). This result motivates computation of approximations to the posterior mean and credible intervals. However, in order to ameliorate the curse of dimensionality associated with taking the whole second-period data as the vector of summary statistics,²⁹ we propose to use an alternative version of the ABC algorithm: the Expectation Propagation (EP) ABC (Barthelmé and Chopin, 2014; Barthelmé et al., 2018). In this method, we leverage the assumption of an independent sample of networks to factor the posterior distribution:

$$dF(\beta) \times \mathbb{P}[\mathbf{G}_1 | \beta, \mathbf{G}_0, \mathbf{X}] \stackrel{\text{independence}}{=} dF(\beta) \times \prod_{c=1}^C \mathbb{P}[G_c^{T_1} | \beta, G_c^{T_0}, X_c] \quad (5)$$

where $\mathbf{G}_d = (G_c^{T_d})_{c=1}^C$; $d \in \{0, 1\}$; $\mathbf{X} = (X_c)_{c=1}^C$; and the prior F only ranges over utility and meeting parameters because we assume that τ_0 is either fixed or estimated in a first step via (4) (see Section 5 for further discussion). EP-ABC assumes a Gaussian prior. The method considers a Gaussian approximation with mean μ_c and covariance matrix Ω_c to each network likelihood function $\beta \mapsto \mathbb{P}[G_c^{T_1} | \beta, G_c^{T_0}, X_c]$. It then proceeds by sequentially

²⁸Both authors study the frequentist properties of ABC algorithms. In their setting, where a finite dimensional vector of summary statistics is considered, it is crucial that the binding function $b(\beta_s, \tau_s) = \text{plim}_{C \rightarrow \infty} T_s$ identifies model parameters.

²⁹Intuitively, it will take many tries to get a draw that approximates the whole vector of observed data well. With a fixed number of simulations, this means that, to have a reasonably small Monte Carlo variance, we tolerate high levels of bias in the ABC approximation to the true posterior.

updating the approximations as follows. Suppose we are currently at network c . The goal is to find (μ_c, Ω_c) to minimize the Kullback-Leibler (KL) divergence between (i) a hybrid approximation that uses the true density at the c -th network and the Gaussian approximation at the remaining networks; and (ii) the “full” Gaussian approximation. Let us denote the prior distribution by $l_0(\beta)$ ($dF(\beta) =: l_0(\beta)$), the “true” likelihood function at each network by $l_c(\beta)$ ($\mathbb{P}[G_c^{T_1} | \beta, G_c^{T_0}, X_c] =: l_c(\beta)$), and let $f_c(\beta)$ be a Gaussian density with parameters (μ_c, Ω_c) . Barthelmé and Chopin (2014) show that the choice of (μ_c, Ω_c) that minimizes this KL divergence is

$$Z_c = \int l_0(\beta) \times \prod_{i \neq c}^C f_i(\beta) \times l_c(\beta) d\beta \quad (6)$$

$$\mu_c = \frac{1}{Z_c} \int l_0(\beta) \times \prod_{i \neq c}^C f_i(\beta) \times l_c(\beta) \times \beta d\beta \quad (7)$$

$$\Omega_c = \frac{1}{Z_c} \int l_0(\beta) \times \prod_{i \neq c}^C f_i(\beta) \times l_c(\beta) \times \beta \beta' d\beta - \mu_c \mu_c'. \quad (8)$$

These quantities are approximated by running a variant of the ABC algorithm (1) with prior $l_0(\beta) \times \prod_{i \neq c}^C f_i(\beta)$, which is Gaussian and can be evaluated; and likelihood $l_c(\beta) = \mathbb{P}[G_c^{T_1} | \beta, G_c^{T_0}, X_c]$, which involves simulating second period adjacency matrices **only** in network c . This ameliorates the curse of dimensionality, since at any given step a draw must be good at reproducing features of **the current network**. Starting from initial guesses for the (μ_c, Ω_c) equal to the prior parameters, the algorithm proceeds by sequentially going through all networks $c = 1, \dots, C$ and updating (μ_c, Ω_c) until meeting a convergence criterion or a full number of passes through all networks.³⁰

5 Application

In our application, we consider data on friendship networks from Pinto and Ponczek (2017). The dataset comprises information on 3rd- and 5th-graders from 30 elementary schools

³⁰In Section F of the supplemental appendix, we describe another method: “expectation propagation with ‘local’ summary statistics”, as an alternative to the EP-ABC described above. To further reduce dimensionality, this method replaces the vector of network edge indicators in the EP-ABC approach with data-driven “local” summary statistics.

in Recife, Brazil. Data on students’ traits and intraclassroom friendship networks was collected at the beginning (baseline) and at the end (followup) of the 2014 school year.³¹ Once missing observations are removed, our working sample comprises 161 classrooms (networks) totalling 1,589 students.³²

As a first step in our analysis, we attest that homophily is indeed a salient feature of our data. For that, we run the following regression model:

$$g_{ij,c,1} = \beta'W_{ij,c,0} + \alpha_i + \gamma_j + \epsilon_{ij,c} \quad (9)$$

where $g_{ij,c,1}$ equals 1 if, in classroom c , individual i nominates j as a friend *at the followup period*. Vector $W_{ij,c,0}$ consists of pairwise distances in gender, age (in years), and measures of cognitive and non-cognitive skills between i and j *at the baseline period*.³³ The specification controls for sender (α_i) and receiver (γ_j) fixed effects. We cluster standard errors at the classroom level.

Column 1 in Table 1 reports estimates obtained from running the above specification. Results indicate that homophily is pervasive, e.g. same-sex classmates are, on average, 19.8 pp more likely to be friends, compared to boy-and-girl pairs.

³¹More specifically, students could nominate up to 8 classmates in each of the three categories: classmates with whom they would (i) study, (ii) talk, or (iii) play. We consider an individual a friend if she appears on at least one of the three lists. No student exceeds 8 friends, since in most cases the three criteria coincide. In our raw dataset, only 1.01% of students reported 8 friends at the baseline, and this number falls to 0.3% at the followup. For a complete description of the dataset, see Pinto and Ponczek (2017).

³²Figure 1 in the supplemental appendix plots one such network.

³³Table 3 in the supplemental appendix presents summary statistics of our dyad-level covariates.

Table 1: Dyadic regressions

	<i>Dependent variable</i>	
	edge	
	(1)	(2)
distance on class list		−0.002*** (0.0005)
distance in age	−0.018*** (0.006)	0.002 (0.003)
distance in gender	−0.198*** (0.006)	−0.186*** (0.006)
distance in cognitive skills	−0.254*** (0.049)	−0.127*** (0.034)
distance in conscientiousness	−0.031*** (0.009)	−0.019*** (0.006)
distance in neuroticism	−0.017** (0.008)	−0.023*** (0.005)
Sender fixed effects?	Yes	No
Receiver fixed effects?	Yes	No
Time effect?	Yes	No
Observations	17,736	17,736
R ²	0.323	0.064
Adjusted R ²	0.185	0.063

Note: *p<0.1; **p<0.05; ***p<0.01
Standard errors clustered at the classroom level in parentheses.

As discussed in Section 3, nonparametric identification of our model requires an instrumental variable with large support. We use the distance between two students in the alphabetically-ordered class list, interacted with one minus an indicator of a link between the pair in the current network, as our “instrument”. This pair-level covariate has large support and enters the matching function, but is excluded from pairs’ marginal utilities.

The position in the class list should affect the odds of a pair meeting, e.g. through class activities in groups, but should not enter marginal utilities. We also assume this mechanism is important for pairs that are not currently friends.

Column 2 provides some reduced-form evidence of relevance of our class list distance variable. We run the specification in (9), but include our class list distance variable and exclude sender and receiver fixed effects. The covariate is statistically significant at the 1% level and with the expected sign: all else equal, classmates “one more student away” in the class list are 0.2 pp less likely to be friends.

In estimating our network formation model, we parameterize the matching function as follows. For individuals i and j in classroom c , we specify

$$\rho((i, j), g, X_c) \propto \exp(\beta'_m W_{ij,c,0} + \delta_0 g_{ij} + \delta_1(1 - g_{ij})Z_{ij,c,0})$$

where $W_{ij,c,0}$ and $Z_{ij,c,0}$ denote, respectively, our vector of pair-level covariates and class list distance variable *at the baseline*. Our specification allows for explicit dependence of meetings on a previous-period link between agents. The utility of individual i in classroom c follows the linear parametrization of Mele (2017), i.e.

$$\begin{aligned} u_i(g, X_c) = & \underbrace{\sum_{k \neq i} \beta'_{ud} \begin{pmatrix} 1 \\ W_{ik,c,0} \end{pmatrix} g_{ik}}_{\text{direct links}} + \underbrace{\sum_{k \neq i} \beta'_{ur} \begin{pmatrix} 1 \\ W_{ik,c,0} \end{pmatrix} g_{ki}}_{\text{mutual links}} + \\ & + \underbrace{\sum_{k \neq i} g_{ik} \sum_{\substack{l \neq i \\ l \neq k}} \beta'_{un} \begin{pmatrix} 1 \\ W_{il,c,0} \end{pmatrix} g_{kl}}_{\text{indirect links}} + \underbrace{\sum_{k \neq i} g_{ik} \sum_{\substack{l \neq i \\ l \neq k}} \beta'_{up} \begin{pmatrix} 1 \\ W_{kl,c,0} \end{pmatrix} g_{li}}_{\text{popularity}} \end{aligned}$$

where, as Mele (2017) does, we impose that $\beta_{un} = \beta_{up}$. This specification accounts for gains in direct links, reciprocity, indirect links, and “popularity” (individuals derive utility of serving as a “bridge” between agents). The restriction $\beta_{un} = \beta_{up}$ is an identification assumption in Mele’s setting, where networks are draws from the model’s stationary distribution. It is not required in our setup, though we enforce it for the sake of comparability. The difference in preference shocks is drawn from a logistic distribution.

We estimate our model using a two-step approach. In the first step, we estimate τ_0

through (4). Our first-step estimate of τ_0 leads to 76 rounds. ^{34,35}

In the second step, we use the EP-ABC method described in Section 4.2 to approximate for the posterior of preference- and matching-related parameters, where we take the first-step estimate of τ_0 as given.³⁶ We adopt independent zero-mean Gaussian distributions with standard deviation equal to two as priors, for both meeting and preference parameters.³⁷ We follow Barthelmé and Chopin’s (2014) method by making a single pass through the dataset, which appears to be sufficient for convergence. In the ABC step, we follow the suggestion of these authors and use a simple nonstochastic accept-reject rule for draws. We use the followup network data in the current classrom as the vector of statistics. Dimensionality in this case is smaller than using a standard ABC algorithm, since we consider just a single classroom at each step. For each classroom, we simulate 100,000 draws and aim for an acceptance rate of 1% (1,000 accepted draws). We use Halton random numbers in generating draws from the prior, as suggested by Barthelmé and Chopin (2014), in order to improve numerical stability.

Table 2 presents the posterior mean of the coefficients of the utility function and matching obtained by the Expectation Propagation ABC method described above. We also report posterior quantiles and the posterior probability of obtaining a negative parameter.

³⁴The estimator does not violate the bound $\hat{\tau} \leq N_c(N_c - 1)$ for 81 out of 161 networks in our sample, which is somewhat reassuring.

³⁵Identification of τ_0 in our setting extends immediately to the case where the number of agents in a network, N_c , is assumed to be random, provided that $\mathbb{P}[\tau_0 \leq N_c(N_c - 1)] > 0$.

³⁶From a frequentist perspective, we can motivate our estimator by appealing to some Bernstein-von-Mises theorem (Vaart, 1998, chapter 10) which ensures posterior asymptotic normality as $C \rightarrow \infty$ (see also the discussion of Frazier et al. (2018)). From a Bayesian point of view, our estimator imposes a degenerate prior on τ_0 at the frequentist estimator (4).

³⁷The value of 2 is chosen in accordance with the reduced-form patterns in Table 1, so as to cover with reasonable prior probability parameter values that enable either homophily in preferences by itself or homophily in meetings by itself to account for observed patterns.

Table 2: Posterior estimates - Expectation propagation

	Mean	Q 0.025	Q 0.975	Prob < 0
Meeting process				
distance in age	0.2920	0.1596	0.4245	0.0000
distance in gender	1.1646	0.8963	1.4328	0.0000
distance in cognitive skills	0.1770	-1.4080	1.7620	0.4134
distance in conscientiousness	0.1809	-0.1255	0.4873	0.1236
distance in neuroticism	-0.0473	-0.4188	0.3241	0.5986
g_ij	1.9856	1.5925	2.3787	0.0000
(1-g_ij)*distance in class list	-0.0966	-0.1392	-0.0539	1.0000
Utility – Direct Links				
intercept	1.0219	0.3494	1.6944	0.0014
distance in age	-0.1080	-0.3992	0.1832	0.7663
distance in gender	-2.6392	-3.3934	-1.8851	1.0000
distance in cognitive skills	-2.3364	-5.2264	0.5536	0.9435
distance in conscientiousness	-1.7369	-2.4102	-1.0637	1.0000
distance in neuroticism	-0.4752	-1.0809	0.1305	0.9379
Utility – Reciprocity Links				
intercept	1.1757	-0.0292	2.3806	0.0279
distance in age	0.1337	-0.6070	0.8744	0.3617
distance in gender	-0.7291	-2.4036	0.9455	0.8033
distance in cognitive skills	0.9659	-2.1432	4.0750	0.2713
distance in conscientiousness	4.2006	2.5797	5.8216	0.0000
distance in neuroticism	1.7257	0.5708	2.8805	0.0017
Utility – Indirect/Popularity				
intercept	0.4427	0.1463	0.7391	0.0017
distance in age	0.1295	-0.0867	0.3457	0.1202
distance in gender	-3.2515	-4.3049	-2.1981	1.0000
distance in cognitive skills	-0.9512	-2.7256	0.8232	0.8533
distance in conscientiousness	-0.5412	-0.9177	-0.1646	0.9976
distance in neuroticism	-0.8722	-1.3541	-0.3902	0.9998

The results indicate almost all mean estimates of utility parameters associated with covariates are negative. This suggests that homophily in preferences is pervasive and not restricted to direct links. We also find high posterior probabilities of a negative effect for the role of gender, not only for the direct links, but also for reciprocal links and popularity. For the matching parameters, the posterior probability of the coefficient associated with our instrument being negative is near 100%, which is reassuring. We also find high posterior probabilities of a negative effect for the role of cognitive skills and neuroticism in the direct and popularity components of utility.

Next, we proceed to counterfactual exercises. We consider the evolution of networks, starting from their baseline value, under four different sequences of matching parameters: (i) when these are kept at their estimated value (*base case*); (ii) when random unbiased matching is imposed across networks (*random meetings*); (iii) when, keeping grade and classroom size in schools fixed, we track students according to their cognitive skills (*tracking case*); and (iv) when, upon meeting, friendships are formed at random with probability 1/2 (*random friendships*).

Table 2 reports posterior means and 95% credible intervals of the projection coefficients of edge indicators at the followup period ($g_{ij,c,1}$) on an intercept and our main controls *at the baseline*. Compared with the observed data, magnitudes in the base case are broadly in line with the frequentist reduced form (Column (2) in Table 1 or Column (1) in Table 2). Nonetheless, our model appears to overstate the role of homophily in age; and we also understate the average value of $g_{ij,c,1}$ in the data. Comparing the second and third columns, we see that imposing random unbiased matching increases observed homophily patterns in gender, cognitive skills, and conscientiousness. This indicates that shutting down homophily in meetings does not lead to a decrease in observed homophily patterns. Moving on to the last column, when we close the channel of homophily due to preferences, the coefficients associated with homophily due to gender, cognitive skills, and conscientiousness decrease in magnitude compared to the base case. This is evidence that, in this example, homophily in preferences is more important than homophily in meetings. In the third column, we note that tracking leads to a weak reduced-form estimate of homophily in cognitive skills. This is expected as students now interact in homogeneous groups. It

also leads to a weaker pattern in the gender coefficient, which may be due to correlation of this attribute with cognitive skills at the baseline.³⁸

Table 3: Projection coefficients and edge statistics - Expectation propagation

	Data	Base case	Random matching	Tracking	Random friendship
Regression coefficients					
distance in class list	-0.0016 [-0.0028;-0.0004]	-0.0016 [-0.0023;-0.0009]	-0.0009 [-0.0017;-0.0003]	-0.0030 [-0.0041;-0.0020]	-0.0030 [-0.0045;-0.0016]
distance in age	0.0018 [-0.0048; 0.0083]	-0.0067 [-0.0143; 0.0010]	-0.0004 [-0.0074; 0.0071]	-0.0039 [-0.0104; 0.0029]	-0.0073 [-0.0126;-0.0017]
distance in gender	-0.1864 [-0.2072;-0.1657]	-0.1576 [-0.1701;-0.1456]	-0.2221 [-0.2383;-0.2061]	-0.1029 [-0.1144;-0.0918]	-0.0884 [-0.1001;-0.0780]
distance in cognitive skills	-0.1275 [-0.2172;-0.0378]	-0.1168 [-0.1805;-0.0486]	-0.1883 [-0.2502;-0.1260]	-0.0321 [-0.1076; 0.0426]	-0.0404 [-0.0968; 0.0171]
distance in conscientiousness	-0.0193 [-0.0332;-0.0053]	-0.0287 [-0.0397;-0.0181]	-0.0371 [-0.0493;-0.0238]	-0.0232 [-0.0325;-0.0140]	-0.0195 [-0.0295;-0.0097]
distance in neuroticism	-0.0231 [-0.0356;-0.0106]	-0.0102 [-0.0192;-0.0009]	-0.0232 [-0.0371;-0.0093]	-0.0131 [-0.0223;-0.0022]	-0.0049 [-0.0143; 0.0052]
Edge summary statistics (follow up)					
edge indicator mean	0.1788 [0.168;0.1896]	0.0837 [0.0775;0.0897]	0.1637 [0.1562;0.1717]	0.0573 [0.0521;0.0625]	0.1201 [0.1034;0.1395]
edge indicator stdev	0.3832 [0.374;0.3921]	0.2769 [0.2674;0.2858]	0.3700 [0.3630;0.3772]	0.2323 [0.2222;0.2421]	0.3248 [0.3045;0.3465]

Notes: In the first column, the output from a frequentist regression of the edge indicator g_{ij} on pair characteristics, along with 95% confidence intervals, are reported. Summary statistics on the edge indicator are also reported. Confidence intervals assume a Gaussian approximation and are constructed using standard errors clustered at the classroom level. Confidence intervals on the standard deviation of the edge indicator additionally use the delta method. Columns 2-4 report mean estimates and 95% credible intervals on the projection coefficients and summary statistics. These are obtained from 1,000 simulations of network data from draws of the posterior distribution under base and counterfactual modifications in parameter values.


Figures 1, 2, and 3 compare the evolution of aggregate utility in the base case with each of our counterfactual scenarios. We plot posterior means and 95% credible intervals of the aggregate utility index, $\sum_{c=1}^C \sum_{i=1}^{N_c} u_i(g_c^t, X_c)$, in the counterfactual scenario minus

³⁸Girls have, on average, 0.0185 more points in cognitive skills at the baseline than boys, and this difference is statistically significant at the 1% level.


the same index in the base case, for each round from the baseline to the followup period. We normalize the difference in indices by the number of students in our sample multiplied by minus the posterior mean estimate of the distance-in-gender direct utility parameter; so results can be interpreted as the number of direct same-sex links each student should receive in the base case so that they are indifferent between policies (without taking into account spillovers on the remaining components of utility).

Imposing random matching leads to a lower trajectory in aggregate utility over the school year. As expected from the previous analysis, random friendship formation leads to an even lower welfare than random matching. The decrease in welfare in both cases comes from a decrease in two components of the utility function: direct links and popularity. We also see that tracking leads to an improvement in welfare, though this benefit diminishes as time passes on. This indicates that a tracking policy in these schools has a positive effect on welfare in the short run, but this effect decreases over time, getting close to zero at the end of 76 rounds of iteration. The effect of tracking policies depends on the structure of the network, as pointed out by Jackson (2021). In our example, we have evidence that homophily due to preferences is stronger than homophily due to opportunities. Then, when we change the meeting opportunities and place individuals in homogeneous classrooms, their welfare increases substantially in the first rounds of the game. However, as soon as they make their new connections, the relative effect of the tracking policy starts to decrease *vis-à-vis* the base case. At the end of the game, the relative effect of the policy is close to zero.


Figure 1: Random matching vs. Base case




(a) Total



(b) Direct




(c) Mutual




(d) Indirect/Popularity


Figure 2: Tracking vs. Base case




(a) Total



(b) Direct




(c) Mutual



(d) Indirect/Popularity

Figure 3: Random friendships vs. Base case



6 Conclusion

In this article, we studied identification and estimation of a network formation model that distinguishes between homophily due to preferences and homophily due to meeting opportunities. The model builds upon the algorithm by Mele (2017), allowing for general classes of utilities and meeting processes. It is also well-grounded on the theoretical literature of network formation (Jackson and Watts, 2002; Jackson, 2010). We provided identification results in the case where a large-support “instrument” is included in preferences (meeting process) and excluded from the meeting process (preferences); and two periods of data from many networks are available. We also discussed a Bayesian estimation procedure that

bypasses direct evaluation of the model likelihood, a task which can be computationally unfeasible even for a moderate number of rounds of the network algorithm. Taken together, our approach enables users to estimate the counterfactual effects of changes in the meeting technology between agents across time, something that previous work was unable to do.

In the applied section of our article, we studied network formation in elementary schools in Northeastern Brazil. Our results suggest that tracking students according to their cognitive skills leads to improved welfare, though the benefits reduce over time. The effect of this policy can be associated with the structure of the networks. In these classroom networks, homophily due to preferences seems more salient than homophily due to meeting opportunities. This can explain the large positive short-run effect of the tracking policies, but the almost zero long run impact.

As emphasised in Section 3.3, analyzing how restrictions on the relationship between the network structure and payoffs may enable point-identification in our setting is an open question which may further enhance the model’s applicability, especially if it allows us to dispense with the exclusion restrictions currently required to identify the model. Another interesting topic for future research is the study of our model under single-network asymptotics, which may be more suitable in some applied settings. Finally, an interesting extension would be to couple our model with an outcome model (e.g. a linear-in-means model), which may enable practitioners to assess the effect of network policies on “hard” outcomes such as test scores.

References

- Aguirregabiria, V. and P. Mira (2010). Dynamic discrete choice structural models: A survey. *Journal of Econometrics* 156(1), 38 – 67. Structural Models of Optimization Behavior in Labor, Aging, and Health.
- Amemiya, T. (1985). *Advanced Econometrics*. Harvard University Press.
- Badev, A. (2021). Nash equilibria on (un)stable networks. *Econometrica* 89(3), 1179–1206.

- Bajari, P., H. Hong, and S. P. Ryan (2010). Identification and estimation of a discrete game of complete information. *Econometrica* 78(5), 1529–1568.
- Barthelmé, S., N. Chopin, and V. Cottet (2018). Divide and conquer in ABC: Expectation-propagation algorithms for likelihood-free inference. *Handbook of Approximate Bayesian Computation*, 415–34.
- Barthelmé, S. and N. Chopin (2014). Expectation propagation for likelihood-free inference. *Journal of the American Statistical Association* 109(505), 315–333.
- Battaglini, M., E. Patacchini, and E. Rainone (2021). Endogenous social interactions with unobserved networks. *The Review of Economic Studies*.
- Blum, M. G. B. (2010). Approximate bayesian computation: A nonparametric perspective. *Journal of the American Statistical Association* 105(491), 1178–1187.
- Bramoullé, Y., S. Currarini, M. O. Jackson, P. Pin, and B. W. Rogers (2012). Homophily and long-run integration in social networks. *Journal of Economic Theory* 147(5), 1754 – 1786.
- Centola, D. (2011). An experimental study of homophily in the adoption of health behavior. *Science* 334(6060), 1269–1272.
- Chandrasekhar, A. (2016). Econometrics of network formation. In *The Oxford Handbook of the Economics of Networks*, pp. 303–357. Oxford University Press.
- Chandrasekhar, A. and M. O. Jackson (2016). A network formation model based on sub-graphs. arXiv:1611.07658.
- Christakis, N., J. Fowler, G. W. Imbens, and K. Kalyanaraman (2020). An empirical model for strategic network formation. In B. Graham and A. de Paula (Eds.), *The Econometric Analysis of Network Data*, Chapter Chapter 6, pp. 123–148. Academic Press.
- Currarini, S., M. O. Jackson, and P. Pin (2009). An economic model of friendship: Homophily, minorities, and segregation. *Econometrica* 77(4), 1003–1045.

- Currarini, S., M. O. Jackson, and P. Pin (2010). Identifying the roles of race-based choice and chance in high school friendship network formation. *Proceedings of the National Academy of Sciences* 107(11), 4857–4861.
- de Paula, A. (2017). Econometrics of network models. In B. Honoré, A. Pakes, M. Piazzesi, and L. Samuelson (Eds.), *Advances in Economics and Econometrics: Eleventh World Congress*, Econometric Society Monographs, pp. 268–323. Cambridge University Press.
- de Paula, Á., S. Richards-Shubik, and E. Tamer (2018). Identifying preferences in networks with bounded degree. *Econometrica* 86(1), 263–288.
- Frazier, D. T., G. M. Martin, C. P. Robert, and J. Rousseau (2018). Asymptotic properties of approximate Bayesian computation. *Biometrika* 105(3), 593–607.
- Goldsmith-Pinkham, P. and G. W. Imbens (2013). Social networks and the identification of peer effects. *Journal of Business & Economic Statistics* 31(3), 253–264.
- Graham, B. S. (2016). Homophily and transitivity in dynamic network formation. Working Paper 22186, National Bureau of Economic Research.
- Graham, B. S. (2017). An econometric model of network formation with degree heterogeneity. *Econometrica* 85(4), 1033–1063.
- Higham, N. J. and L. Lin (2011). On p th roots of stochastic matrices. *Linear Algebra and its Applications* 435(3), 448 – 463. Special Issue: Dedication to Pete Stewart on the occasion of his 70th birthday.
- Hobert, J. P. (2011). Chapter 10 the data augmentation algorithm: Theory and methodology. In *Handbook of Markov Chain Monte Carlo*, pp. 253–294. CRC Press.
- Horn, R. A. and C. R. Johnson (2012). *Matrix Analysis*. Cambridge University Press.
- Hsieh, C.-S. and L. F. Lee (2016). A social interactions model with endogenous friendship formation and selectivity. *Journal of Applied Econometrics* 31(2), 301–319.
- Jackson, M. (2010). *Social and Economic Networks*. Princeton University Press.

- Jackson, M. O. (2021). Inequality's economic and social roots: The role of social networks and homophily. *SSRN Electronic Journal*.
- Jackson, M. O. and A. Watts (2002). The evolution of social and economic networks. *Journal of Economic Theory* 106(2), 265 – 295.
- Kadelka, C. and A. McCombs (2021). Effect of homophily and correlation of beliefs on COVID-19 and general infectious disease outbreaks. *PLOS ONE* 16(12).
- Li, Q. and J. S. Racine (2006). *Nonparametric Econometrics: Theory and Practice*. Princeton University Press.
- Li, W. and P. Fearnhead (2018). On the asymptotic efficiency of approximate Bayesian computation estimators. *Biometrika* 105(2), 285–299.
- McFadden, D. (1973). Conditional logit analysis of qualitative choice behaviour. In P. Zarembka (Ed.), *Frontiers in Econometrics*, pp. 105–142. New York, NY, USA: Academic Press New York.
- Mele, A. (2017). A structural model of dense network formation. *Econometrica* 85(3), 825–850.
- Newey, W. K. and D. McFadden (1994). Large sample estimation and hypothesis testing. In *Handbook of Econometrics*, Volume 4, Chapter 36, pp. 2111 – 2245. Elsevier.
- Newey, W. K. and R. J. Smith (2004). Higher order properties of gmm and generalized empirical likelihood estimators. *Econometrica* 72(1), 219–255.
- Norris, J. R. (1997). *Markov Chains*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press.
- Pinto, C. C. d. X. and V. P. Ponczek (2017). The building blocks of skill development. Unpublished.
- Rothenberg, T. J. (1971). Identification in parametric models. *Econometrica* 39(3), 577–591.

Sheng, S. (2020). A structural econometric analysis of network formation games through subnetworks. *Econometrica* 88(5), 1829–1858.

Sisson, S. A. and Y. Fan (2011). Chapter 12 likelihood-free mcmc. In *Handbook of Markov Chain Monte Carlo*, pp. 313–338. CRC Press.

Tamer, E. (2003). Incomplete simultaneous discrete response model with multiple equilibria. *The Review of Economic Studies* 70(1), 147–165.

Vaart, A. W. v. d. (1998). *Asymptotic Statistics*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press.

A Proofs of the main results

In the following, write $N^k(g)$, $k \in \mathbb{N}$ and $g \in \mathcal{G}$, for the set of networks that differ from g in *exactly* k edges.

A.1 Proof of Lemma 3.1

Starting from some $w \in N(g_0)$, $w_{ij} \neq g_{0ij}$, we can identify $\rho_{ij}(g_0, X^m(g)) = \lim_{t \rightarrow \infty} (\Pi(X \setminus Z_{ij}^u(g_0, w), Z_{ij}^u(g_0, w) = t\vec{r}_{g_0w}))_{g_0w}$, which is valid under the (conditional) large support assumption. We are then able to identify $u_i(w, X_i^u(w))$ thanks to the normalization on $u_i(g_0, X_i^u(g_0))$. Proceeding in a similar fashion iteratively on $w' \in N^2(g_0), N^3(g_0) \dots$, we identify all objects.

A.2 Proof of Proposition 3.1

Before presenting the proof of the main identification result in the paper, we present a lemma for the case in which $\tau_0 = 2$. This lemma provides the intuition for the proof of Proposition 3.1. We show identification when $\tau_0 = 2$ to get an idea of how the general case would look. As seen in the proof, the exclusion restriction given by Assumption 3.5 could be relaxed in this case, though the latter would then be insufficient for identification when $\tau_0 > 2$.

Lemma A.1. *Suppose Assumptions 2.1, 2.2, 3.1, 3.3, and 3.5 hold. Then, if $\tau_0 = 2$, the model is identified.*

Proof. First notice that $\Pi(X; \theta)^2$ takes the form

$$(\Pi(X; \theta)^2)_{gw} = \begin{cases} \rho_{ij}(g)F_{ij}(g, [w_{ij}, g])\rho_{kl}([w_{ij}, g])F_{kl}([w_{ij}, g], w) + \\ + \rho_{kl}(g)F_{kl}(g, [w_{kl}, g])\rho_{ij}([w_{kl}, g])F_{ij}([w_{kl}, g], w) & \text{if } w \in N^2(g), w_{ij} \neq g_{ij}, w_{kl} \neq g_{kl} \\ \Pi_{gg}\rho_{ij}(g)F_{ij}(g, w) + \rho_{ij}(g)F_{ij}(g, w)\Pi_{ww} & \text{if } w \in N(g), w_{ij} \neq g_{ij} \\ \Pi_{gg}\Pi_{gg} + \sum_{s \in N(g)} \Pi_{gs}\Pi_{sg} & \text{if } g = w \\ 0 & \text{otherwise} \end{cases} \quad (10)$$

Fix $g \in \mathcal{G}$, $w \in N(g)$, $g_{ij} \neq w_{ij}$. By driving $F_{ij}(g, w) \rightarrow 1$ and $F_{pq}(g, m) \rightarrow 0$ for all $m \in N(g) \setminus \{w\}$, $g_{pq} \neq m_{pq}$, the term $(\Pi^2)_{gg}$ identifies

$$\lim_{t^*} (\Pi^2)_{gg} = (1 - \rho_{ij}(g))^2$$

where \lim_{t^*} is shorthand for the appropriate limit.³⁹ Since $\lim_{t^*} (\Pi^2)_{gg} \in [0, 1]$, we can uniquely solve for $\rho_{ij}(g)$, thus establishing identification of $\rho_{ij}(g)$. Next, by taking $F_{pq}(g, m) \rightarrow 0$ for all $m \in N(g) \setminus \{w\}$, $F_{pq}(w, m) \rightarrow 0$ for all $m \in N(w) \setminus \{g\}$, the term Π_{gw} identifies

$$\begin{aligned} \lim_{t^{**}} \Pi_{gw} &= (1 - \rho_{ij}(g)F_{ij}(g, w))\rho_{ij}(g)F_{ij}(g, w) + \rho_{ij}(g)F_{ij}(g, w)(1 - \rho_{ij}(w)F_{ij}(w, g)) = \\ &\quad \rho_{ij}(g)F_{ij}(g, w)(2 - \rho_{ij}(w) + (\rho_{ij}(w) - \rho_{ij}(g))F_{ij}(g, w)) \end{aligned}$$

where $\lim_{t^{**}}$ is shorthand for the appropriate limit.⁴⁰ Since the right-hand term is strictly increasing in $F_{ij}(g, w)$, the “true” $F_{ij}(g, w)$ uniquely solves the equation, thus establishing identification. \square

Next we present the proof of Proposition 3.1:

Proof of Proposition 3.1. Fix $g \in \mathcal{G}$, $w \in N(g)$, $g_{ij} \neq w_{ij}$. Observe that

$$(\Pi^{\tau_0})_{gg} = \sum_{m \in N(g) \cup \{g\}} (\Pi^{\tau_0-1})_{gm} \Pi_{mg}.$$

We first prove the following claim:

³⁹I.e. a limit that drives $F_{ij}(g, w) \rightarrow 1$ and $F_{pq}(g, m) \rightarrow 0$ for all $m \in N(g) \setminus \{w\}$, $g_{pq} \neq m_{pq}$.

⁴⁰I.e. a limit that drives $F_{pq}(g, m) \rightarrow 0$ for all $m \in N(g) \setminus \{w\}$, $F_{pq}(w, m) \rightarrow 0$ for all $m \in N(w) \setminus \{g\}$.

Claim. Under a limit which drives $F_{pq}(w, m) \rightarrow 0$ for all $m \in N(w)$, $m_{pq} \neq w_{pq}$, and $F_{kl}(g, s) \rightarrow 0$ for all $s \in N(g) \setminus \{w\}$, $s_{kl} \neq g_{kl}$, we have $\lim_{t^*}(\Pi^{\tau_0})_{gg} = (1 - \rho_{ij}(g))^{\tau_0}$, where \lim_{t^*} is shorthand for the appropriate limit.

Proof. The case $\tau_0 = 1$ is readily verified by driving $F_{kl}(g, s) \rightarrow 0$ for all $s \in N(g) \setminus \{w\}$, $s_{kl} \neq g_{kl}$ and $F_{ij}(g, w) \rightarrow 1$. For $\tau_0 > 1$, we begin by noticing that we may drive $(\Pi^{\tau_0-1})_{gm} \rightarrow 0$ for all $m \in N(g) \setminus \{w\}$. Since m and g differ in exactly one edge (say, $m_{pq} \neq g_{pq}$), a transition in edge pq must appear in every summand in $(\Pi^{\tau_0-1})_{gm}$. Indeed, $(\Pi^{\tau_0-1})_{gm}$ sums over all possible transitions in edge pq from value g_{pq} to m_{pq} in $\tau_0 - 1$ rounds. Put another way, for each summand in $(\Pi^{\tau_0-1})_{gm}$, there exists $t \in \{0, 1 \dots \tau_0 - 2\}$, $g_{pq}^t = g_{pq}$ and $g_{pq}^{t+1} = m_{pq}$.⁴¹ Fix a summand in $(\Pi^{\tau_0-1})_{gm}$. We analyze the following cases:

1. There exists $t \in \{0, 1 \dots \tau_0 - 2\}$, $g_{pq}^t = g_{pq}$, $g_{pq}^{t+1} = m_{pq}$ **and** $g^t = g$. In this case, by taking $F_{pq}(g, m) \rightarrow 0$, we drive the summand to 0.
2. For all $t \in \{0, 1 \dots \tau_0 - 2\}$ such that $g_{pq}^t = g_{pq}$, $g_{pq}^{t+1} = m_{pq}$, we have $g^t \neq g$. Take t^* to be the smallest t satisfying the above. Observe that $t^* > 0$, as $g^0 = g$ (we always start at g). Since $g^{t^*} \neq g$, there exists $t' < t^*$, $g^{t'} = g$ and $g^{t'+1} = z$, $z \in N(g)$. If there exists some t' satisfying this property such that $z \neq w$ (with $g_{kl} \neq z_{kl}$), then driving $F_{kl}(g, z) \rightarrow 0$ vanishes the term. If, for all such t' , $z = w$, take t^{**} to be the maximum of t' . Observe that $t^{**} < t^*$. If $g^{t^{**}} = w$, we may safely drive $F_{pq}(w, [m_{pq}, w_{-pq}]) \rightarrow 0$. If not, then $t^{**} + 1 < t^*$ and there exists a transition from w to some element in $N(w)$ which we can safely drive to 0.

Since the above argument holds, irrespective of the summand (the common limit will vanish all terms), we conclude $(\Pi^{\tau_0-1})_{gm} \rightarrow 0$. Since $F_{ij}(g, w) \rightarrow 1$, $\lim_{t^*}(\Pi^{\tau_0-1})_{gw}\Pi_{wg} = 0$.⁴² The common limit in the statement of the claim thus leaves us with

$$\lim_{t^*}(\Pi^{\tau_0})_{gg} = \lim_{t^*}(\Pi^{\tau_0-1})_{gg} \lim_{t^*}(\Pi)_{gg} = \lim_{t^*}(\Pi^{\tau_0-1})_{gg}(1 - \rho_{ij}(g))$$

Induction then yields the desired result. □

⁴¹Recall g^t is the stochastic process on \mathcal{G} induced by the game.

⁴²In all previous arguments, we implicitly use the sandwich lemma to infer that, if one term of the product goes to zero, the whole product does. This is immediate, since we are working with products of probabilities.

Since $\lim_{t^*}(\Pi^{\tau_0})_{gg} \in [0, 1]$, we can uniquely solve for $\rho_{ij}(g)$, thus establishing identification.

Next, we proceed to identification of $F_{ij}(g, w)$. Note that

$$(\Pi^{\tau_0})_{gw} = \sum_{m \in N(w) \cup \{w\}} (\Pi^{\tau_0-1})_{gm} \Pi_{mw}.$$

We then prove the following claim:

Claim. *Under a limit which drives $F_{pq}(w, m) \rightarrow 0$ for all $m \in N(w) \setminus \{g\}$, $m_{pq} \neq w_{pq}$, and $F_{kl}(g, s) \rightarrow 0$ for all $s \in N(g) \setminus \{w\}$, $s_{kl} \neq g_{kl}$:*

$$\lim_{t^{**}}(\Pi^{\tau_0})_{gw} = \lim_{t^{**}}(\Pi^{\tau_0-1})_{gg} \rho_{ij}(g) F_{ij}(g, w) + \lim_{t^{**}}(\Pi^{\tau_0-1})_{gw} (1 - \rho_{ij}(w) F_{ij}(w, g))$$

where $\lim_{t^{**}}$ is shorthand for the appropriate limit. We also have that, under such a limit,

$$\lim_{t^{**}}(\Pi^{\tau_0})_{gw} + \lim_{t^{**}}(\Pi^{\tau_0})_{gg} = 1.$$

Proof. For $\tau_0 = 1$, we have

$$\begin{aligned} \lim_{t^{**}}(\Pi)_{gw} &= \rho_{ij}(g) F_{ij}(g, w) \\ \lim_{t^{**}}(\Pi)_{gg} &= (1 - \rho_{ij}(g) F_{ij}(g, w)). \end{aligned}$$

These expressions follow directly from the limit being taken and equation (2).

Consider next the case $\tau_0 > 1$. The limit in the statement of the lemma drives $(\Pi^{\tau_0-1})_{gm} \rightarrow 0$ for all $m \in N(w) \setminus \{g\}$. Indeed, if $m \in N(w) \setminus \{g\}$, then $m \in N^2(g)$. Recall that $(\Pi^{\tau_0-1})_{gm}$ sums over all possible transitions from g to m in $\tau_0 - 1$ rounds. Fix a summand in $(\Pi^{\tau_0-1})_{gm}$. If a transition from g occurs at pair $(a, b) \in \mathcal{M}$, $(a, b) \neq (i, j)$, then the limit vanishes the term. If all transitions from g occur at pair (i, j) (i.e. g only transitions to w), a transition from w must occur, since $m \in N^2(g)$. If w only transitions to g , then either $m = g$ or $m = w$, which is not true. Therefore, there exists a transition from w to some $z \in N(w) \setminus \{g\}$, so we can vanish the summand. The limit in the statement thus drives the term $(\Pi^{\tau_0-1})_{gm}$ to zero.

From the above discussion, we thus get

$$\begin{aligned} \lim_{t^{**}}(\Pi^{\tau_0})_{gw} &= \lim_{t^{**}}(\Pi^{\tau_0-1})_{gg}\Pi_{gw} + \lim_{t^{**}}(\Pi^{\tau_0-1})_{gw}\Pi_{ww} = \\ &= \lim_{t^{**}}(\Pi^{\tau_0-1})_{gg}\rho_{ij}(g)F_{ij}(g, w) + \lim_{t^{**}}(\Pi^{\tau_0-1})_{gw}(1 - \rho_{ij}(w)F_{ij}(w, g)) \end{aligned}$$

which establishes the first part of the claim.

Next, under the limit in the statement of the claim

$$\lim_{t^{**}}(\Pi^{\tau_0})_{gg} = \lim_{t^{**}}(\Pi^{\tau_0-1})_{gg}(1 - \rho_{ij}(g)F_{ij}(g, w)) + \lim_{t^{**}}(\Pi^{\tau_0-1})_{gw}\rho_{ij}(w)F_{ij}(w, g).$$

This follows from observation that, in the proof of the *previous* claim, we can still drive $(\Pi^{\tau_0-1})_{gm} \rightarrow 0$ for all $m \in N(g) \setminus \{w\}$ even though $F_{ij}(g, w)$ does not vanish.⁴³ We are thus left with the terms related to staying in g or transitioning to w in $\tau_0 - 1$ rounds.

Finally, the second part of the claim can be asserted by noticing that $\lim_{t^{**}}(\Pi)_{gw} + \lim_{t^{**}}(\Pi)_{gg} = 1$ and applying this fact inductively on the expression for $\lim_{t^{**}}(\Pi^{\tau_0})_{gw} + \lim_{t^{**}}(\Pi^{\tau_0})_{gg} = \lim_{t^{**}}(\Pi^{\tau_0-1})_{gw} + \lim_{t^{**}}(\Pi^{\tau_0-1})_{gg}$. \square

To establish identification of $F_{ij}(g, w)$, we need to show that the expression for $\lim_{t^{**}}(\Pi^{\tau_0})_{gw}$ is strictly increasing in $(0, 1)$ as a function of $F_{ij}(g, w)$. Denoting by $D_{F_{ij}(g, w)} \lim_{t^{**}}(\Pi^{\tau_0})_{gw}$ the derivative of $\lim_{t^{**}}(\Pi^{\tau_0})_{gw}$ as a function of $F_{ij}(g, w)$, we get

$$\begin{aligned} D_{F_{ij}(g, w)} \lim_{t^{**}}(\Pi^{\tau_0})_{gw} &= D_{F_{ij}(g, w)} \lim_{t^{**}}(\Pi^{\tau_0-1})_{gw}(1 - \rho_{ij}(w) + (\rho_{ij}(w) - \rho_{ij}(g))F_{ij}(g, w)) \\ &\quad + \rho_{ij}(g)(1 - \lim_{t^{**}}(\Pi^{\tau_0-1})_{gw}) + \rho_{ij}(w) \lim_{t^{**}}(\Pi^{\tau_0-1})_{gw} \end{aligned}$$

where we use that $\lim_{t^{**}}(\Pi^{\tau_0})_{gw} + \lim_{t^{**}}(\Pi^{\tau_0})_{gg} = 1$ and $F_{ij}(g, w) + F_{ij}(w, g) = 1$. By noticing $\rho_{ij}(g)(1 - \lim_{t^{**}}(\Pi^{\tau_0-1})_{gw}) + \rho_{ij}(w) \lim_{t^{**}}(\Pi^{\tau_0-1})_{gw} \geq \min\{\rho_{ij}(g), \rho_{ij}(w)\} > 0$ and applying induction on the fact that $D_{F_{ij}(g, w)} \lim_{t^{**}}(\Pi)_{gw} \geq 0$, we get that the derivative is strictly positive in $(0, 1)$, thus showing that the map is invertible and establishing identification of $F_{ij}(g, w)$. \square

⁴³Suppose g only transitions to w . Since $m \neq w$, w must transition to some other $z \in N(w)$. If w only transitions to g , then either $m = g$ or $m = w$, which is not true. Therefore, we can always vanish a summand in $(\Pi^{\tau_0-1})_{gm}$, even though $F_{ij}(g, w)$ does not vanish.

A.3 Proof of Lemma 4.1

Proof. Consider the event $\{\hat{\tau} \rightarrow \tau_0\}^{\mathcal{G}}$. Observe that $\{\hat{\tau} \rightarrow \tau_0\}^{\mathcal{G}} = \bigcap_{c \in \mathbb{N}} \{\|G_c^{T_1} - G_c^{T_0}\|_F < \tau_0\}$. Fix $k \in \mathbb{N}$ and notice that

$$\mathbb{P} \left[\bigcap_{c=1}^k \{\|G_c^{T_1} - G_c^{T_0}\|_F < \tau_0\} \right] = \phi^k$$

where ϕ is the ex-ante (unconditional) probability that $G_c^{T_1}$ differs from $G_c^{T_0}$ in strictly less than τ_0 edges. Since $\mathbb{P}[\|G_c^{T_1} - G_c^{T_0}\|_F = \tau_0] \in (0, 1)$ (which follows from 2.1, 2.2, and 3.2), we have $\phi \in (0, 1)$. Passing k to the limit and using continuity of \mathbb{P} from above, we conclude the desired result. \square

It is apparent from the proof above that it is possible extend the result of Lemma 4.1 in order to allow for a sequence of *independent* observations stemming from a game with common parameters, but allowing the distribution of covariates (and the number of players) to vary between networks. In this case, we must restrict the distribution of covariates to not shift excessively to high-probability regions. Formally, the proof would change as the ex-ante probability would now depend on c , i.e. we would have $\prod_{c=1}^k \phi_c$ in the formula. If $\limsup_{c \rightarrow \infty} \phi_c < 1$, we would get the same result.