

Conditional h -convexity with applications

Ismail Nikoufar and Davuod Saeedi

Received: date / Accepted: date

Abstract In this paper, we introduce the notion of conditional h -convex functions and we prove an operator version of the Jensen inequality for conditional h -convex functions. Using this type of functions, we give some refinements for Ky-Fan's inequality, arithmetic-geometric mean inequality, Chrystal inequality, and Hölder-McCarthy inequality. Many of the other inequalities can be refined by applying this new notion.

Keywords h -convex function · Jensen's inequality · Ky-Fan's inequality · arithmetic-geometric mean inequality · Hölder-McCarthy inequality.

Mathematics Subject Classification (2010) 26D15 · 15A39 · 47A63 · 46N10 · 47A60.

1 Introduction

Throughout this paper, we denote by \mathcal{H} a Hilbert space and by $B(\mathcal{H})$ the algebra of all bounded linear operators on \mathcal{H} . The subalgebra of all self-adjoint operators in $B(\mathcal{H})$ is denoted by $B_{sa}(\mathcal{H})$. An operator A in $B_{sa}(\mathcal{H})$ is positive whenever $\langle Ax, x \rangle \geq 0$ for all $x \in \mathcal{H}$ and we write $A \geq 0$. We denote by $Sp(A)$ the spectrum of an operator $A \in B(\mathcal{H})$.

One of the most important and most frequently applied in a diversity of mathematical fields, especially in mathematical analysis and statistics is Jensen's inequality. In this process, it has been improved, generalized and adjusted to various environments. It has been linked to the other important inequalities and often referred to as "king of inequalities".

I. Nikoufar (Corresponding Author)

Department of Mathematics, Payame Noor University, P.O. Box 19395-3697 Tehran, Iran

E-mail: nikoufar@pnu.ac.ir

D. Saeedi

Department of Mathematics, Payame Noor University, P.O. Box 19395-3697 Tehran, Iran

E-mail: dsaeedi3961@gmail.com

The convexity of functions is an important issue in many fields of science, for instance in economy and optimization. A function $f : \mathbb{I} \rightarrow \mathbb{R}$, $\mathbb{I} \subseteq \mathbb{R}$ is convex whenever the following inequality

$$f(\lambda u + (1 - \lambda)v) \leq \lambda f(u) + (1 - \lambda)f(v)$$

holds for all $u, v \in \mathbb{I}$ and for all $\lambda \in [0, 1]$ and the function $f : \mathbb{I} \rightarrow \mathbb{R}$ is concave whenever $-f$ is convex.

In 2007, Varošanec [18] introduced a wide class of functions the so called h -convex functions which generalizes convex, s -convex, Godunova-Levin, and P -class functions [10], [5, 13, 17, 14], [7], [3], [16], [4, 12].

A non-negative function $f : \mathbb{I} \rightarrow \mathbb{R}$ is an h -convex function on \mathbb{I} if

$$f(\lambda u + (1 - \lambda)v) \leq h(\lambda)f(u) + h(1 - \lambda)f(v), \quad (1)$$

where h is a non-negative function defined on the real interval \mathbb{J} , $u, v \in \mathbb{I}$ and $\lambda \in [0, 1] \subseteq \mathbb{J}$. For more results and generalizations regarding h -convexity, we refer the readers to see [1, 6, 8, 15].

Jensen's inequality for convex functions is one of the most important result in the theory of inequalities and for appropriate choices of the function many other famous inequalities are particular cases of this inequality. An operator version of the Jensen inequality for a convex function has been proved by Mond and Pečarić as follows ([11], [9]):

Theorem 1 *Let $f : [m, M] \rightarrow \mathbb{R}$ be a continuous convex function. Then,*

$$f(\langle Ax, x \rangle) \leq \langle f(A)x, x \rangle \quad (2)$$

for every self-adjoint operator A with $Sp(A) \subseteq [m, M]$ and every unit vector $x \in \mathcal{H}$.

In Section 2, we introduce the notion of conditional h -convex functions and we prove an operator version of the Jensen inequality for conditional h -convex functions. In Section 3, Using this type of functions, we give some refinements for Ky-Fan's inequality, arithmetic-geometric mean inequality, Chrystal inequality, and Hölder-McCarthy inequality. We claim that many of the other inequalities can be refined by applying the notion of conditional h -convex functions.

2 Mond and Pečarić inequality for conditional h -convex functions

In this section, we introduce the notion of conditional h -convex functions and we prove an operator version of the Jensen inequality for conditional h -convex functions. We indicate that an operator version of the Jensen inequality for conditional h -convex functions still holds similar to that Mond and Pečarić considered for convex functions.

Definition 1 A function $f : [m, M] \rightarrow \mathbb{R}$ is conditional convex at $v \in [m, M]$ if there exists a function $g : [m, M] \rightarrow \mathbb{R}$ such that $g(v) < v$ and

$$f(\lambda u + (1 - \lambda)v) \leq \lambda f(u) + (1 - \lambda)f(v) \quad (3)$$

for every $u \in [m, M]$ with $g(v) \leq u < v$ and $\lambda \in [0, 1]$.

It is clear that every convex function is conditional convex at every $v \in [m, M]$ by considering $g(t) = m$, but not vice versa.

Example 1 Define $f : [0, 2] \rightarrow \mathbb{R}$ by $f(t) = (t - 1)^3$ and $g(t) = \begin{cases} 1, & x = 2 \\ 2, & x \neq 2 \end{cases}$. The function f is not convex on $[0, 2]$, where f is conditional convex at $2 \in [0, 2]$. Consider $g(t) = \cos(\frac{4\pi}{3}x)$. Then f is conditional convex at $\frac{3}{2} \in [0, 2]$.

Definition 2 A non-negative function $f : [m, M] \rightarrow \mathbb{R}$ is conditional h -convex at $v \in [m, M]$ if there exists a function $g : [m, M] \rightarrow \mathbb{R}$ such that $g(v) < v$ and

$$f(\lambda u + (1 - \lambda)v) \leq h(\lambda)f(u) + h(1 - \lambda)f(v), \quad (4)$$

for every $u \in [m, M]$ with $g(v) \leq u < v$ and $\lambda \in [0, 1]$, where h is a non-negative function defined on the real interval \mathbb{J} , $[0, 1] \subseteq \mathbb{J}$. In this situation, we say the function f is g -conditional h -convex at $v \in [m, M]$.

It is clear that every h -convex function is g -conditional h -convex at every $v \in [m, M]$ by considering $g(t) = m$, but not vice versa.

For a non-negative and non-zero function $h : K \subset \mathbb{R} \rightarrow \mathbb{R}$, we define

$$M_K(h) = \inf_{t \in K} \frac{h(t)}{t}$$

and call it the Jensen's coefficient for h -convex functions on K . We consider all functions h such that the Jensen's coefficient $M_K(h)$ exists.

Theorem 2 Let $h : [0, 1] \rightarrow \mathbb{R}$ be a non-negative and non-zero function. If f is a continuous g -conditional h -convex function at $v \in [m, M]$, then

$$f(\langle Ax, x \rangle) \leq M_{(0,1)}(h) \langle f(A)x, x \rangle \quad (5)$$

for each $x \in \mathcal{H}$ with $\|x\| = 1$ and every self-adjoint operator A such that $Sp(A) \subseteq [g(v), v]$.

Proof It is clear that if $M_{(0,1)}(h) = +\infty$, then the inequality (5) holds. Assume that $M_{(0,1)}(h) < +\infty$. Since f is g -conditional h -convex at $v \in [m, M]$,

$$f(\lambda u + (1 - \lambda)v) - h(1 - \lambda)f(v) \leq h(\lambda)f(u) \quad (6)$$

for all $u \in [g(v), v]$ and for all $\lambda \in (0, 1)$. By dividing both sides of (6) with $\lambda \in (0, 1)$, one can reach

$$\frac{f(\lambda u + (1 - \lambda)v) - h(1 - \lambda)f(v)}{\lambda} \leq \frac{h(\lambda)}{\lambda} f(u) \quad (7)$$

for all $u \in [g(v), v]$. Define

$$\alpha := \min_{u \in [g(v), v]} \frac{f(\lambda u + (1 - \lambda)v) - h(1 - \lambda)f(v)}{\lambda(u - v)}. \quad (8)$$

The inequalities (7) and (8) entail that

$$\alpha(u - v) \leq \frac{h(\lambda)}{\lambda} f(u)$$

for all $u \in [g(v), v]$ and $\lambda \in (0, 1)$. Consider the linear function $l(t) := \alpha(t - v)$ and $\bar{g} = \langle Ax, x \rangle$ for every self-adjoint operator A with $Sp(A) \subseteq [g(v), v]$. This implies that $l(u) \leq \frac{h(\lambda)}{\lambda} f(u)$ for all $u \in [g(v), v]$ and $g(v) \leq \bar{g} \leq v$, respectively. Let $l'(t) := \alpha(t - \bar{g}) + f(\bar{g})$ be the straight line passing through the point $(\bar{g}, f(\bar{g}))$ and parallel to the line l . It follows from the continuity of the function f that

$$l'(\bar{g}) \geq f(\bar{g}) - \epsilon \quad (9)$$

for all $\epsilon > 0$. We now consider two cases:

(i) Assume that $l'(t) \leq \frac{h(\lambda)}{\lambda} f(t)$ for every $t \in [g(v), v]$. By using the functional calculus, one has $l'(A) \leq \frac{h(\lambda)}{\lambda} f(A)$ for every self-adjoint operator A with $Sp(A) \subseteq [g(v), v]$ and so

$$\langle l'(A)x, x \rangle \leq \frac{h(\lambda)}{\lambda} \langle f(A)x, x \rangle \quad (10)$$

for each $x \in \mathcal{H}$ with $\|x\| = 1$. Since l' is linear and ϵ is arbitrary the inequalities (9) and (10) imply

$$f(\langle Ax, x \rangle) \leq \frac{h(\lambda)}{\lambda} \langle f(A)x, x \rangle. \quad (11)$$

(ii) Assume that there exist some points $t \in [g(v), v]$ such that

$$l'(t) > \frac{h(\lambda)}{\lambda} f(t).$$

Define the sets T and S as follows:

$$T := \{t \in [g(v), \bar{g}] : l'(t) > \frac{h(\lambda)}{\lambda} f(t)\},$$

$$S := \{t \in [\bar{g}, v] : l'(t) > \frac{h(\lambda)}{\lambda} f(t)\}.$$

Consider $t_T := \max\{t : t \in T\}$ and $t_S := \min\{t : t \in S\}$. Let l_T be the line passing through the points $(t_T, 0)$ and $(\bar{g}, f(\bar{g}))$ and l_S the line passing through the points $(t_S, 0)$ and $(\bar{g}, f(\bar{g}))$. Define the function L on $[g(v), v]$ as follows:

$$L(t) := \begin{cases} l_T(t), & t \in [g(v), \bar{g}], \\ l_S(t), & t \in [\bar{g}, v]. \end{cases}$$

To show that $L(t) \leq \frac{h(\lambda)}{\lambda} f(t)$ holds for all $t \in [g(v), v]$, consider the partition $\{g(v), t_T, \bar{g}, t_S, v\}$ for the closed interval $[g(v), v]$. Note that $l_T(t) \leq 0$ for every

$t \in [g(v), t_T]$ and since $f(t) \geq 0$, we clearly observe that $l_T(t) \leq \frac{h(\lambda)}{\lambda}f(t)$ for every $t \in [g(v), t_T]$. On the other hand, we see that

$$l'(t) \leq \frac{h(\lambda)}{\lambda}f(t) \quad (12)$$

for every $t \in (t_T, \bar{g}]$, since if there exists $t_0 \in (t_T, \bar{g}]$ such that $l'(t_0) > \frac{h(\lambda)}{\lambda}f(t_0)$, then $t_0 \in T$ and $t_0 < t_T$, which is a contradiction. So, by letting t tends to t_T from right in (12), the inequality (12) indicates that

$$l'(t_T) \leq \frac{h(\lambda)}{\lambda}f(t_T). \quad (13)$$

Moreover, since $t_T \in \bar{T}$, the reverse inequality holds in (13) and hence l' is the line passing through the points $(t_T, \frac{h(\lambda)}{\lambda}f(t_T))$ and $(\bar{g}, f(\bar{g}))$ and its slope is $\alpha' = \frac{f(\bar{g}) - \frac{h(\lambda)}{\lambda}f(t_T)}{\bar{g} - t_T}$, where the slope of l_T is $\alpha_T = \frac{f(\bar{g})}{\bar{g} - t_T}$ and so $l_T(t) \leq l'(t)$ for every $t \in (t_T, \bar{g}]$. By the inequality (12) we observe that $l_T(t) \leq \frac{h(\lambda)}{\lambda}f(t)$ for every $t \in (t_T, \bar{g}]$. So, $L(t) = l_T(t) \leq \frac{h(\lambda)}{\lambda}f(t)$ for every $t \in [g(v), \bar{g}]$.

By the similar methods one can show that $L(t) = l_S(t) \leq \frac{h(\lambda)}{\lambda}f(t)$ for every $t \in [\bar{g}, v]$. Note that the lines l_T and l_S are joining at the point along the length of \bar{g} and so $l_T(\bar{g}) = l_S(\bar{g})$ and since f is continuous,

$$l_T(\bar{g}) = f(\bar{g}) \geq f(\bar{g}) - \epsilon \quad (14)$$

for arbitrary $\epsilon > 0$. For the case $Sp(A) \subseteq [g(v), \bar{g}]$, we have

$$f(\langle Ax, x \rangle) - \epsilon \leq l_T(\langle Ax, x \rangle) = \langle l_T(A)x, x \rangle \leq \frac{h(\lambda)}{\lambda} \langle f(A)x, x \rangle.$$

Moreover, for the case $Sp(A) \subseteq [\bar{g}, v]$, we have

$$\begin{aligned} f(\langle Ax, x \rangle) - \epsilon &\leq l_T(\langle Ax, x \rangle) = l_S(\langle Ax, x \rangle) = \langle l_S(A)x, x \rangle \\ &\leq \frac{h(\lambda)}{\lambda} \langle f(A)x, x \rangle \end{aligned}$$

and consequently we can deduce (11). Now, by taking the infimum over all $\lambda \in (0, 1)$ on both sides of the inequality (11), one can deduce (5).

Corollary 1 *Under the hypotheses of Theorem 2, if the function $\frac{h(\lambda)}{\lambda}$ is decreasing on $(0, 1)$, then $\lambda = \frac{1}{2}$ is the best possible in the inequality (11). Indeed, one has*

$$f(\langle Ax, x \rangle) \leq 2h\left(\frac{1}{2}\right) \langle f(A)x, x \rangle \quad (15)$$

for each $x \in \mathcal{H}$ with $\|x\| = 1$ and every self-adjoint operator A such that $Sp(A) \subseteq [g(v), v]$.

Proof We divide the interval $(0, 1)$ to two parts $(0, \frac{1}{2}]$ and $(\frac{1}{2}, 1)$, respectively. If $\lambda \in (0, \frac{1}{2}]$, then the infimum value of the function $\frac{h(\lambda)}{\lambda}$ occurs at the endpoint $\lambda = \frac{1}{2}$, since the function $\frac{h(t)}{t}$ is decreasing and so

$$M_{(0, \frac{1}{2}]}(h) = \frac{h(\frac{1}{2})}{\frac{1}{2}} = 2h(\frac{1}{2}).$$

When $\lambda \in (\frac{1}{2}, 1)$, we show that there is an h -convex function such that does not satisfy the inequality (11). Let $h(\lambda) = \lambda^\beta$, $0 < \beta < 1$, and $0 < \lambda < 1$. Then, $\frac{h(\lambda)}{\lambda}$ is decreasing on $(0, 1)$. Define $f : [0, \infty) \rightarrow \mathbb{R}$ by $f(t) = e^{-t}$. The function f is h -convex, since by using the convexity of f , we have

$$\begin{aligned} f(\lambda x + (1 - \lambda)y) &= e^{-(\alpha x + (1 - \alpha)y)} \\ &\leq \lambda e^{-x} + (1 - \lambda)e^{-y} \\ &\leq \lambda^\beta e^{-x} + (1 - \lambda)^\beta e^{-y} \\ &= h(\lambda)f(x) + h(1 - \lambda)f(y) \end{aligned}$$

for every $x, y \geq 0$ and $\lambda \in [0, 1]$. Consider $A = \begin{pmatrix} 0 & 0 \\ 0 & a \end{pmatrix}$, $a > 0$, and $x = (\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$. A simple calculation shows that $f(\langle Ax, x \rangle) = f(\frac{a}{2}) = e^{-\frac{a}{2}}$ and $\langle f(A)x, x \rangle = \frac{e^{-a}}{2}$. By applying the inequality (11), one can see that $e^{-\frac{a}{2}} \leq \lambda^{\beta-1} \frac{e^{-a}}{2}$ which implies $2 < 2e^{\frac{a}{2}} \leq \lambda^{\beta-1}$. On the other hand, for $\lambda \in (\frac{1}{2}, 1)$, we have $\lambda^{\beta-1} < \lambda^{-1} < 2$ which is a contradiction.

3 Some Applications

In this section, we provide some examples of conditional h -convex functions. Using this type of functions, we give some refinements for Ky-Fan's inequality, arithmetic-geometric mean inequality, Chrystal inequality, and Hölder-McCarthy inequality.

Lemma 1 Let $h : [0, 1] \rightarrow (0, \infty)$ defined by $h(t) = \frac{\alpha}{\beta} e^{t(1-t)}$ and let $g : (0, \frac{1}{2}] \rightarrow [0, \infty)$ defined by $g(t) = \frac{t^\alpha}{t^\alpha + (1-t)^\alpha}$ with $0 < \alpha < \beta \leq \alpha + 1$. Then, the function $f : (0, \frac{1}{2}] \rightarrow [0, \infty)$ defined by $f(t) = \ln(\frac{1-t}{t})$ is g -conditional h -convex at every $v \in (0, \frac{1}{2}]$.

Proof Let $v \in (0, \frac{1}{2}]$ and define the function F by

$$F(u, \lambda) = \frac{\alpha}{\beta} e^{t(1-t)} \ln \left(\frac{1-u}{u} \right) + \frac{\alpha}{\beta} e^{t(1-t)} \ln \left(\frac{1-v}{v} \right) - \ln \left(\frac{1 - (\lambda u + (1 - \lambda)v)}{\lambda u + (1 - \lambda)v} \right)$$

for every $u \in [g(v), v]$ and $\lambda \in [0, 1]$. It is enough to show that $F(u, \lambda) \geq 0$. We have

$$\begin{aligned} F(u, \lambda) &\geq \frac{\alpha}{\beta} \ln \left(\frac{1-u}{u} \right) + \frac{\alpha}{\beta} \ln \left(\frac{1-v}{v} \right) - \ln \left(\frac{1 - (\lambda u + (1-\lambda)v)}{\lambda u + (1-\lambda)v} \right) \\ &= \ln \left(\frac{1-u}{u} \right) - \ln \left(\frac{1 - (\lambda u + (1-\lambda)v)}{\lambda u + (1-\lambda)v} \right) \\ &\quad + \frac{\alpha}{\beta} \ln \left(\frac{1-v}{v} \right) - (1 - \frac{\alpha}{\beta}) \ln \left(\frac{1-u}{u} \right). \end{aligned} \quad (16)$$

It is obvious that $u < \lambda u + (1-\lambda)v$ for every $\lambda \in [0, 1]$. So, $\frac{1 - (\lambda u + (1-\lambda)v)}{\lambda u + (1-\lambda)v} < \frac{1-u}{u}$. It follows that

$$\ln \left(\frac{1-u}{u} \right) - \ln \left(\frac{1 - (\lambda u + (1-\lambda)v)}{\lambda u + (1-\lambda)v} \right) \geq 0. \quad (17)$$

On the other hand, we have $\frac{v^\alpha}{v^\alpha + (1-v)^\alpha} \leq u$ so that $\frac{1-u}{u} \leq (\frac{1-v}{v})^\alpha$. This ensures $\ln(\frac{1-u}{u}) \leq \alpha \ln(\frac{1-v}{v})$. Since $\beta \leq \alpha + 1$, this indicates that

$$(1 - \frac{\alpha}{\beta}) \ln \left(\frac{1-u}{u} \right) \leq \frac{1}{\beta} \ln \left(\frac{1-u}{u} \right) \leq \frac{\alpha}{\beta} \ln \left(\frac{1-v}{v} \right). \quad (18)$$

The inequalities (17) and (18) entail that $F(u, \lambda) \geq 0$.

Corollary 2 (*Refined Ky-Fan's inequality*) *If $v \in (0, \frac{1}{2}]$, $a_i \in [\frac{v^\alpha}{v^\alpha + (1-v)^\alpha}, v]$, $q_i \in (0, 1)$, and $\sum_{i=1}^n q_i = 1$ such that $0 < \alpha < \beta \leq \alpha + 1$, then*

$$\frac{\sum_{i=1}^n q_i (1-a_i)}{\sum_{i=1}^n q_i a_i} \leq \prod_{i=1}^n \left(\frac{1-a_i}{a_i} \right)^{\frac{\alpha}{\beta} q_i} < \prod_{i=1}^n \left(\frac{1-a_i}{a_i} \right)^{q_i}. \quad (19)$$

Proof Consider $f(t) = \ln(\frac{1-t}{t})$ on $(0, \frac{1}{2}]$. By Lemma 1, the function f is g -conditional h -convex at every $v \in (0, \frac{1}{2}]$. Define

$$A = \begin{pmatrix} a_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & a_n \end{pmatrix}, \quad x = \begin{pmatrix} \sqrt{q_1} \\ \vdots \\ \sqrt{q_n} \end{pmatrix}.$$

Then, $Sp(A) \subseteq [\frac{v^\alpha}{v^\alpha + (1-v)^\alpha}, v]$ and $\|x\| = 1$. According to Theorem 2 one knows

$$f(\langle Ax, x \rangle) \leq M_{(0,1)}(h) \langle f(A)x, x \rangle. \quad (20)$$

On the other hand, we have

$$\begin{aligned} \langle Ax, x \rangle &= \sum_{i=1}^n q_i a_i, \\ M_{(0,1)}(h) &= \inf_{\lambda \in [0,1]} \frac{h(\lambda)}{\lambda} = \frac{\alpha}{\beta}, \\ \langle f(A)x, x \rangle &= \sum_{i=1}^n q_i \ln \left(\frac{1-a_i}{a_i} \right). \end{aligned}$$

By replacing these quantities in (20), we get the first inequality in (19). The second inequality follows from the fact that $\frac{1-a_i}{a_i} \geq 1$ and $\frac{\alpha}{\beta} < 1$.

Lemma 2 *Let $h : [0, 1] \rightarrow (0, \infty)$ defined by $h(t) = \frac{\alpha}{\beta} e^{t(1-t)}$ and let $g : (0, 1] \rightarrow [0, \infty)$ defined by $g(t) = t^\alpha$ with $0 < \alpha < \beta \leq \alpha + 1$. Then, the function $f : (0, 1] \rightarrow [0, \infty)$ defined by $f(t) = -\ln(t)$ is g -conditional h -convex at every $v \in (0, 1]$.*

Proof Let $v \in (0, 1]$ and define the function F by

$$F(u, \lambda) = -\frac{\alpha}{\beta} e^{t(1-t)} \ln(u) - \frac{\alpha}{\beta} e^{t(1-t)} \ln(v) + \ln(\lambda u + (1-\lambda)v)$$

for every $u \in [v^\alpha, v]$ and $\lambda \in [0, 1]$. We demonstrate that $F(u, \lambda) \geq 0$. We have

$$\begin{aligned} F(u, \lambda) &\geq -\frac{\alpha}{\beta} \ln(u) - \frac{\alpha}{\beta} \ln(v) + \ln(\lambda u + (1-\lambda)v) \\ &= (1 - \frac{\alpha}{\beta}) \ln(u) - \frac{\alpha}{\beta} \ln(v) - \ln(u) + \ln(\lambda u + (1-\lambda)v). \end{aligned}$$

Since $u < \lambda u + (1-\lambda)v$ for every $\lambda \in [0, 1]$, $\ln(u) < \ln(\lambda u + (1-\lambda)v)$ and hence

$$-\ln(u) + \ln(\lambda u + (1-\lambda)v) > 0. \quad (21)$$

On the other hand, we have $v^\alpha \leq u$. Since $\beta - \alpha \leq 1$ and $0 < u < 1$, $u \leq u^{\beta-\alpha}$. These two inequalities ensure that $v^\alpha \leq u^{\beta-\alpha}$. So,

$$\frac{\alpha}{\beta} \ln(v) \leq \frac{\beta - \alpha}{\beta} \ln(u). \quad (22)$$

We realize from the inequalities (21) and (22) that $F(u, \lambda) \geq 0$.

Corollary 3 *(Refined Arithmetic-Geometric Mean inequality) If $v \in (0, 1]$, $a_i \in [v^\alpha, v]$, $q_i \in (0, 1)$, and $\sum_{i=1}^n q_i = 1$ such that $0 < \alpha < \beta \leq \alpha + 1$, then*

$$\prod_{i=1}^n a_i^{q_i} < \prod_{i=1}^n a_i^{\frac{\alpha}{\beta} q_i} \leq \sum_{i=1}^n q_i a_i. \quad (23)$$

Proof Consider $f(t) = -\ln(t)$ on $(0, 1]$. Due to Lemma 1, the function f is g -conditional h -convex at every $v \in (0, 1]$. Define the matrix A and the vector x as in the proof of Corollary 2. Then, $Sp(A) \subseteq [v^\alpha, v]$ and $\|x\| = 1$. Theorem 2 signifies the inequality (20) holds and by the assumptions we have

$$\begin{aligned} \langle Ax, x \rangle &= \sum_{i=1}^n q_i a_i, \\ M_{(0,1)}(h) &= \inf_{\lambda \in [0,1]} \frac{h(\lambda)}{\lambda} = \frac{\alpha}{\beta}, \\ \langle f(A)x, x \rangle &= -\sum_{i=1}^n q_i \ln(a_i). \end{aligned}$$

By replacing these quantities in (20), one can deduce

$$-\ln\left(\sum_{i=1}^n q_i a_i\right) \leq -\frac{\alpha}{\beta} \sum_{i=1}^n q_i \ln(a_i).$$

From this inequality, it follows that

$$\ln\left(\sum_{i=1}^n q_i a_i\right) \geq \sum_{i=1}^n \ln(a_i^{\frac{\alpha}{\beta} q_i}) = \ln\left(\prod_{i=1}^n a_i^{\frac{\alpha}{\beta} q_i}\right),$$

which is equivalent to the second inequality in (23). The first inequality comes from the fact that $0 < a_i \leq 1$ and $\frac{\alpha}{\beta} < 1$.

Lemma 3 *Let $h : [0, 1] \rightarrow (0, \infty)$ defined by $h(t) = \frac{\alpha}{\beta} e^{t(1-t)}$ and let $g : (0, \infty) \rightarrow [0, \infty)$ defined by $g(t) = \ln((1 + e^t)^{\frac{\beta}{\alpha} - 1} - 1)$ with $0 < \alpha < \beta \leq 2\alpha$. Then, the function $f : (0, \infty) \rightarrow [0, \infty)$ defined by $f(t) = \ln(1 + e^t)$ is g -conditional h -convex at every $v \in (0, \infty)$.*

Proof Let $v \in (0, \infty)$ and define the function F by

$$F(u, \lambda) = \frac{\alpha}{\beta} e^{t(1-t)} \ln(1 + e^u) + \frac{\alpha}{\beta} e^{t(1-t)} \ln(1 + e^v) - \ln(1 + e^{\lambda u + (1-\lambda)v})$$

for every $u \in [\ln((1 + e^v)^{\frac{\beta}{\alpha} - 1} - 1), v]$ and $\lambda \in [0, 1]$. We prove that $F(u, \lambda) \geq 0$. Note that

$$F(u, \lambda) \geq \frac{\alpha}{\beta} \ln(1 + e^u) + \frac{\alpha}{\beta} \ln(1 + e^v) - \ln(1 + e^{\lambda u + (1-\lambda)v}).$$

Since $\lambda u + (1 - \lambda)v < v$ for every $\lambda \in [0, 1]$,

$$\ln(1 + e^{\lambda u + (1-\lambda)v}) < \ln(1 + e^v). \quad (24)$$

Since $0 < \frac{\beta}{\alpha} - 1 \leq 1$, $\ln((1 + e^v)^{\frac{\beta}{\alpha} - 1} - 1) > 0$. On the other hand, we have $\ln((1 + e^v)^{\frac{\beta}{\alpha} - 1} - 1) \leq u$. So,

$$1 + e^v \leq (1 + e^u)^{\frac{\alpha}{\beta}} (1 + e^v)^{\frac{\alpha}{\beta}}. \quad (25)$$

In view of the inequalities (24) and (25), we find that

$$\ln(1 + e^{\lambda u + (1-\lambda)v}) < \ln((1 + e^u)^{\frac{\alpha}{\beta}} (1 + e^v)^{\frac{\alpha}{\beta}}),$$

which implies $F(u, \lambda) \geq 0$.

Corollary 4 (*Refined Chrystal's inequality*) *If $v > 0$, $q_i \in (0, 1)$, and $\sum_{i=1}^n q_i = 1$, $a_i, b_i \in [\ln((1 + e^v)^{\frac{\beta}{\alpha} - 1} - 1), v]$ such that $0 < \alpha < \beta \leq 2\alpha$, then*

$$\prod_{i=1}^n a_i^{q_i} + \prod_{i=1}^n b_i^{q_i} \leq \prod_{i=1}^n \left(\frac{(a_i + b_i)^{\frac{\alpha}{\beta}}}{b_i^{\frac{\alpha}{\beta} - 1}} \right)^{q_i} < \prod_{i=1}^n (a_i + b_i)^{q_i}. \quad (26)$$

Proof Consider $f(t) = \ln(1+e^t)$ on $(0, \infty)$. Then, the function f is g -conditional h -convex at every $v \in (0, \infty)$, by Lemma 3. Define the matrix A and the vector x as in the proof of Corollary 2. Then, $Sp(A) \subseteq [\ln((1+e^v)^{\frac{\beta}{\alpha}-1}-1), v]$ and $\|x\| = 1$. In view of Theorem 2 we know that the inequality (20) holds and by the assumptions we have

$$\begin{aligned}\langle Ax, x \rangle &= \sum_{i=1}^n q_i a_i, \\ M_{(0,1)}(h) &= \inf_{\lambda \in [0,1]} \frac{h(\lambda)}{\lambda} = \frac{\alpha}{\beta}, \\ \langle f(A)x, x \rangle &= \sum_{i=1}^n q_i \ln(1+e^{a_i}).\end{aligned}$$

By replacing these quantities in (20), we see that

$$\ln(1+e^{\sum_{i=1}^n q_i a_i}) \leq \frac{\alpha}{\beta} \sum_{i=1}^n q_i \ln(1+e^{a_i}). \quad (27)$$

On the other hand, since $\frac{\alpha}{\beta} < 1$,

$$\frac{\alpha}{\beta} \sum_{i=1}^n q_i \ln(1+e^{a_i}) < \sum_{i=1}^n q_i \ln(1+e^{a_i}). \quad (28)$$

Combining the inequalities (27) and (28), we get

$$\ln(1+e^{\sum_{i=1}^n q_i a_i}) \leq \frac{\alpha}{\beta} \sum_{i=1}^n q_i \ln(1+e^{a_i}) < \sum_{i=1}^n q_i \ln(1+e^{a_i}). \quad (29)$$

By replacing $\ln(\frac{a_i}{b_i})$ with a_i in (29), we observe that

$$\ln\left(1 + \prod_{i=1}^n \left(\frac{a_i}{b_i}\right)^{q_i}\right) \leq \ln\left(\prod_{i=1}^n \left(1 + \frac{a_i}{b_i}\right)^{\frac{\alpha}{\beta} q_i}\right) < \ln\left(\prod_{i=1}^n \left(1 + \frac{a_i}{b_i}\right)^{q_i}\right),$$

which implies

$$1 + \prod_{i=1}^n \left(\frac{a_i}{b_i}\right)^{q_i} \leq \prod_{i=1}^n \left(1 + \frac{a_i}{b_i}\right)^{\frac{\alpha}{\beta} q_i} < \prod_{i=1}^n \left(1 + \frac{a_i}{b_i}\right)^{q_i}.$$

Multiplying both sides by $\prod_{i=1}^n b_i^{q_i}$ gives (26).

Lemma 4 Let $h : [0, 1] \rightarrow (0, \infty)$ defined by $h(t) = \frac{\alpha}{\beta} e^{t(1-t)}$ and let $g : [0, \infty) \rightarrow [0, \infty)$ defined by $g(t) = (\sqrt[p]{\frac{\beta}{\alpha}} - 1)t$ with $p > 1$ and $0 < \alpha < \beta$. Then, the function $f : [0, \infty) \rightarrow [0, \infty)$ defined by $f(t) = t^p$ is g -conditional h -convex at every $v \in [0, \infty)$.

Proof Suppose $v \in [0, \infty)$ and let the function F be defined by

$$F(u, \lambda) = \frac{\alpha}{\beta} e^{t(1-t)} u^p + \frac{\alpha}{\beta} e^{t(1-t)} v^p - (\lambda u + (1-\lambda)v)^p$$

for every $u \in [(\sqrt[p]{\frac{\beta}{\alpha}} - 1)v, v]$ and $\lambda \in [0, 1]$. To show that $F(u, \lambda) \geq 0$, we observe that

$$\begin{aligned} F(u, \lambda) &\geq \frac{\alpha}{\beta} u^p + \frac{\alpha}{\beta} v^p - (\lambda u + (1-\lambda)v)^p \\ &= \frac{\alpha}{\beta} u^p - (1 - \frac{\alpha}{\beta})v^p + v^p - (\lambda u + (1-\lambda)v)^p. \end{aligned}$$

Since $\lambda u + (1-\lambda)v < v$ for every $\lambda \in [0, 1]$,

$$(\lambda u + (1-\lambda)v)^p < v^p. \quad (30)$$

Since $\frac{\beta}{\alpha} > 1$ and $(\sqrt[p]{\frac{\beta}{\alpha}} - 1)v \leq u$,

$$\left(\frac{\beta}{\alpha} - 1\right)v^p \leq u^p < \frac{\beta}{\alpha} u^p. \quad (31)$$

The inequalities (30) and (31) signify $F(u, \lambda) \geq 0$.

Corollary 5 (*Refined Hölder-McCarthy inequality*) *If $v > 0$, $p > 1$, and $0 < \alpha < \beta$, then*

$$\langle Ax, x \rangle^p \leq \frac{\alpha}{\beta} \langle A^p x, x \rangle < \langle A^p x, x \rangle$$

for every self-adjoint operator A with $Sp(A) \subseteq [(\sqrt[p]{\frac{\beta}{\alpha}} - 1)v, v]$ and every unit vector x .

Proof Define $f(t) = t^p$ on $(0, \infty)$. Due to Lemma 4 the function f is g -conditional h -convex at every $v \in (0, \infty)$. In account of Theorem 2 the inequality (20) holds and we have

$$\langle Ax, x \rangle^p \leq M_{(0,1)}(h) \langle A^p x, x \rangle,$$

where $M_{(0,1)}(h) = \inf_{\lambda \in [0,1]} \frac{h(\lambda)}{\lambda} = \frac{\alpha}{\beta}$. Since $\frac{\alpha}{\beta} < 1$, we get the desired result.

4 Declarations

We remark that the potential conflicts of interest and data sharing not applicable to this article and no data sets were generated during the current study.

References

1. M. Bombardelli and S. Varošanec, Properties of h -convex functions related to the Hermite-Hadamard-Fejér inequalities, *Compute. Math. Applic.* 58(9), 1869–1877 (2009).
2. S. N. Bernstein, Sur les fonctions absolument monotones, *Acta Math.* 52, 1–66 (1929).
3. S. S. Dragomir, J. Pečarić, and L. E. Persson, Some inequalities of Hadamard type, *Soochow J. Math.* 21(3), 335–341 (1995).
4. S. S. Dragomir and C. E. M. Pearce, Quasi-convex functions and Hadamard's inequality, *Bull. Aust. Math. Soc.* 57(3), 377–385 (1998).
5. S. S. Dragomir and Th. M. Rassias, *Ostrowski type inequalities and applications in numerical integration*, Kluwer Academic Publishers, Dordrecht, Boston, London, 2002.
6. S. S. Dragomir, Inequalities of Jensen type for h -convex functions on linear spaces, *Math. Moravica* 19(1), 107–121 (2015).
7. E. K. Godunova and V. I. Levin, Neravenstva dlja funkcii širokogo klassa, sodержaščego vypuklye, monotonnye i nekotorye drugie vidy funkcii, *Vyčislitel. Mat. i Mat. Fiz. Mežvuzov. Sb. Nauk. Trudov, MGPI, Moskva*, 138–142 (1985) (in Russian).
8. A. Háyzy, Bernstein–doetsch type results for h -convex functions, *Math. Inequal. Appl.* 14(3), 499–508 (2011).
9. T. Furuta, J. Mićić, J. Pečarić, Y. Seo, Mond–Pečarić method in operator inequalities, inequalities for bounded self-adjoint operators on a Hilbert Space, *Element*, Zagreb, 2005.
10. H. Hudzik and L. Maligranda, Some remarks on s -convex functions, *Aequationes Math.* 17, 100–111 (1994).
11. B. Mond, J. Pečarić, Convex inequalities in Hilbert space, *Houston J. Math.* 19, 405–420 (1993).
12. I. Nikoufar and D. Saeedi, Some Inequalities for P -Class Functions, *Filomat* 34(13), 4555–4566 (2020).
13. I. Nikoufar and D. Saeedi, An operator version of the Jensen inequality for s -convex functions, *Complex Anal. Oper. Theory* 15, 92 (2021). <https://doi.org/10.1007/s11785-021-01139-x>
14. I. Nikoufar and D. Saeedi, Operator inequalities associated with the Kantrovich type inequalities for s -convex functions, *arXiv preprint arXiv:2111.12837* (2021).
15. A. Olbryś, Representation theorems for h -convexity, *J. Math. Anal. Appl.* 426(2), 986–994 (2015).
16. C. E. M. Pearce and A. M. Rubinov, P -functions, quasi-convex functions and Hadamard-type inequalities, *J. Math. Anal. Appl.* 240, 92–104 (1999).
17. M. R. Pinheiro, Exploring the concept of s -convexity, *Aequationes Math.* 74, 201–209 (2007).
18. S. Varošanec, On h -convexity, *J. Math. Anal. Appl.* 326, 303–311 (2007).