

HARMONICALLY m -CONVEX SET-VALUED FUNCTION

GABRIEL SANTANA¹, MAIRA VALERA-LÓPEZ^{1*}

ABSTRACT. This research aimed to introduce the concept of harmonically m -convex set-valued functions, which is obtained from the combination of two definitions: harmonically m -convex functions and set-valued functions. In this work some properties and characteristics are developed, as well as a inequality of the Hermite-Hadamard type for such functions.

1. INTRODUCTION

Convexity is a fundamental concept in mathematics and has interesting applications in other areas of knowledge. The study of this definition has increased exponentially at the beginning of this century obtaining important results in this area (see [17]). In recent studies, new concepts of convexity have been introduced from the original definition, one of them is harmonic convexity. These concepts constitute the basis for the development of our work. Anderson *et al.* [1] and I. Iscan [8] have considered harmonic convex functions which can be viewed as an important class of convex functions. One can show that harmonic convex functions have some nice properties, which convex functions have. Noor *et al.* [22] introduced the class of harmonic h -convex functions with respect to an arbitrary nonnegative function $h(\cdot)$. This class is more general and contains several known and new classes of harmonic convex functions as special cases.

Several new convex sets have been introduced and investigated (see [2, 18]). However, the concept of m -convex set D was introduced in 1984 by G. Toader [25] and go on to presenting the definition of m -convex set value function and stating some properties and examples.

The notion of set-valued function arises at the beginning of the twentieth century, when C. Berge [4] introduced the concept of upper and lower limit of succession sets. Recently different researchers have been studied for different notions of set-valued convexity functions, as well as have been used to find the error in some inclusion problems with set restrictions, (see [5, 11, 15, 16, 19, 20, 21, 23, 24]). In particular, T. Lara *et al.* [10] introduced the notion of m -convex set-valued functions defined on a nonempty m -convex subset of a real linear space with values in the set of nonempty parts of an equality real linear space. In this research are given several characterizations as well as certain algebraic properties and examples.

Various integral inequalities for convex functions and their variant forms are being developed using novel ideas and techniques. Some recent developments demonstrate results of Hermite-Hadamard inequality for harmonically convex set-valued functions. Santana *et al.* [24] introduced the definition of harmonically convex and strongly harmonically convex set-valued functions, obtaining important results such as Hermite-Hadamard and Fejér inequalities, as well as a Benstein-Doetsch-type theorem.

The aim of this research is to introduce a new concept of harmonically m -convex set-valued function, in turn, this research explores some characteristics and properties that exhibit these. Also, a new Hermite-Hadamard type inequality for harmonically m -convex set-valued functions is developed. The results obtained in this investigation are being studied for stochastic processes, having as reference the following works [3, 6, 7, 12, 13, 14].

Date:

* Corresponding author: Gabriel Santana.

2020 *Mathematics Subject Classification.* Primary 26D15; Secondary 26D99, 26A51, 39B62, 46N10.

Key words and phrases. set-valued convex functions, Harmonically convex functions, Harmonically m -convex functions, Hermite-Hadamard type inequality, convex analysis.

2. PRELIMINARY

As part of our research it is necessary to provide the reader with some preliminary definitions used throughout this investigation in order to lay the foundations for the development of this work.

Definition 2.1. (see [9]) Let X a linear space and $m \in (0, 1]$. A nonempty subset D of X is said to be harmonic m -convex, if for all $x, y \in D$ and $t \in [0, 1]$, we have:

$$\frac{mxy}{tmx + (1-t)y} \in D.$$

In this case İ. İscan in [9] generalized the harmonically convex function definition introduced in [8] to harmonically (α, m) -convex function:

Definition 2.2. (see [9]) Let $f : D \subset (0, \infty) \rightarrow \mathbb{R}$ a function. Then, f is said to be harmonically (α, m) -convex function if for all $\alpha, t \in [0, 1]$, $m \in (0, 1]$ and $x, y \in D$, we have:

$$(2.1) \quad f\left(\frac{mxy}{tmx + (1-t)y}\right) \leq t^\alpha f(y) + m(1-t)^\alpha f(x).$$

Note that if we considered $\alpha = 1$ in (2.1), f is said to be a harmonically m -convex function and satisfies the following:

$$f\left(\frac{mxy}{tmx + (1-t)y}\right) \leq tf(y) + m(1-t)f(x).$$

For this kind of functions in [9] obtain the following result:

Theorem 2.3. (see [9]) Let $f : (0, \infty) \rightarrow \mathbb{R}$ be a harmonically m -convex function with $m \in (0, 1]$. If $0 < a < b < \infty$ and $f \in L[a, b]$, then one has the inequality:

$$\frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \leq \min \left\{ \frac{f(a) + f\left\{\frac{b}{m}\right\}}{2}, \frac{f(b) + f\left\{\frac{a}{m}\right\}}{2} \right\}.$$

In the other hand, G. Santana *et al.* [24] in 2018, introduced the definition of harmonically convex set-valued functions, extending the definition given by İ. İscan for real functions (see [8]).

Definition 2.4. (see [24]) Let X and Y linear spaces, D a harmonically convex subset of X and $F : D \subset X \rightarrow n(Y)$ a set-valued function. Then F is said harmonically convex function if for all $x, y \in D$ and $t \in [0, 1]$, we have:

$$tF(y) + (1-t)F(x) \subseteq F\left(\frac{xy}{tx + (1-t)y}\right)$$

Remark 2.5. Throughout this paper $n(Y)$ will denote the family of nonempty subsets of Y .

For this kind of functions they obtain many results as algebraic properties, Hermite-Hadamard and Fejer type inequalities and Bernstein-Doetsch type result.

To prove certain algebraic properties of the results in this research, we use two definitions established by T. Lara *et al.* in 2014 [10].

Definition 2.6. (see [10]) Let $F_1, F_2 : D \subseteq X \rightarrow n(Y)$ be two set-valued functions (or multifunctions) then:

- The union of F_1 and F_2 is a set-valued function $F_1 \cup F_2 : D \rightarrow n(Y)$ given by $(F_1 \cup F_2)(x) = F_1(x) \cup F_2(x)$ for each $x \in D$.
- The sum of F_1 and F_2 is the function $F_1 + F_2 : D \rightarrow n(Y)$ defined in its usual form $(F_1 + F_2)(x) = F_1(x) + F_2(x)$ for each $x \in D$.

Definition 2.7. (see [10]) Let X, Y, Z be linear spaces and D be a subset of X . Then:

- If $F_1 : D \rightarrow n(Y)$ and $F_2 : D \rightarrow n(Z)$, then the cartesian product function of F_1 and F_2 is the set-valued function $F_1 \times F_2 : D \rightarrow n(Y) \times n(Z)$ given by $(F_1 \times F_2)(x) = F_1(x) \times F_2(x)$ for each $x \in D$.
- If $F_1 : D \rightarrow n(Y)$ and $F_2 : n(Y) \rightarrow n(Z)$, then the composition function of F_1 and F_2 is the set-valued function $F_2 \circ F_1 : D \rightarrow n(Z)$ given by

$$(F_2 \circ F_1)(x) = F_2(F_1(x)) = \bigcup_{y \in F_1(x)} F_2(y),$$

for each $x \in D$.

Following the idea establish in [24], in this paper we extend that definition and introduce a new concept of convexity, combining the definitions of harmonically m -convex functions and set-valued functions. Then we define the following:

Definition 2.8. Let X and Y be linear spaces, D a harmonically m -convex subset of X and $F : D \subset X \rightarrow n(Y)$ a set-valued function. It said that F is said harmonically m -convex function if for all $x, y \in D$, $t \in [0, 1]$, and $m \in (0, 1]$, we have:

$$(2.2) \quad tF(y) + m(1-t)F(x) \subseteq F\left(\frac{mxy}{tmx + (1-t)y}\right).$$

We have some examples of this kind of function.

Example 2.9. Let $f_1, (-f_2) : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$ be harmonically m -convex functions with $f_1(x) \leq f_2(x)$ for all $x \in [a, b]$. Then, the set-valued function $F : D \subset X \rightarrow n(Y)$ defined by $F(x) = [f_1(x), f_2(x)]$ is harmonically m -convex.

In fact, since f_1 and $-f_2$ are harmonically m -convex functions, then for all $x, y \in D$, $t \in [0, 1]$ and $m \in (0, 1]$, we have:

$$f_1\left(\frac{mxy}{tmx + (1-t)y}\right) \leq tf_1(y) + m(1-t)f_1(x),$$

and

$$-f_2\left(\frac{mxy}{tmx + (1-t)y}\right) \leq -(tf_2(y) + m(1-t)f_2(x)),$$

if we multiply (-1) to both sides of the last inequality, we have:

$$f_2\left(\frac{mxy}{tmx + (1-t)y}\right) \geq tf_2(y) + m(1-t)f_2(x),$$

Then,

$$(2.3) \quad [tf_1(y) + m(1-t)f_1(x), tf_2(y) + m(1-t)f_2(x)] \subset \left[f_1\left(\frac{mxy}{tmx + (1-t)y}\right), f_2\left(\frac{mxy}{tmx + (1-t)y}\right) \right]$$

from (2.3) we obtain with a elementary calculus that

$$tF(y) + m(1-t)F(x) \subset F\left(\frac{mxy}{tmx + (1-t)y}\right).$$

Example 2.10. Let $H \in \mathbb{R}^3$ a harmonically m -convex subset, $F : \mathbb{R} \rightarrow n(\mathbb{R})$ a set-valued function defined by $F(x) = f(x)H$, where $f(x) = x^2$ is a harmonically m -convex function.

Then, since $f(x) = x^2$ is a harmonically m -convex function by definition, we have for all $x, y \in D$, $t \in [0, 1]$ and $m \in (0, 1]$ that

$$\left(\frac{mxy}{tmx + (1-t)y}\right)^2 \leq t(y)^2 + m(1-t)(x)^2,$$

using the harmonically m -convex properties of H we have to

$$(t(y)^2 + m(1-t)(x)^2)H \subseteq \left(\frac{mxy}{tmx + (1-t)y} \right)^2 H,$$

then,

$$\begin{aligned} (t(y)^2 + m(1-t)(x)^2)H &= t(y)^2H + m(1-t)(x)^2H \\ &= tF(y) + m(1-t)F(x) \\ &\subseteq F\left(\frac{mxy}{tmx + (1-t)y}\right). \end{aligned}$$

Thus F is a harmonically m -convex set-valued function.

3. MAIN RESULTS

The results obtained in this paper are based on the developments and ideas of İ. Iscan [9] and T. Lara *et al.* in [10]. The following proposition establishes a property over harmonic m -convex set.

Proposition 3.1. *Let harmonic m -convex ($m \neq 1$) subset D of X is said to be starshaped if, for all x in D and all t in the interval $(0, 1]$, the point tx also belongs to D . That is:*

$$tD \subseteq D.$$

Proof. Let D be a harmonically m -convex subset of X . If D is an empty set, there is nothing to prove. If, on the contrary, we consider D a nonempty set, let $x \in D$ then the point $x = \frac{mab}{tma + (1-t)b} \in D$ for everything $a, b \in D$ and $t \in [0, 1]$. Thus, $[m, 1]x = \{rx : m \leq r \leq 1\} \subset D$, in particular $mx \in D$. If $m = 0$, then $[0, 1]x \in D$, we got the desired result.

In the case $m > 0$, we similarly repeat the previous argument for mx (instead of x), in this case we have to $[m^2, m]x = [m, 1]mx \subseteq D$.

Inductively, we have that $[m^n, m^{n-1}]x \subseteq D$ for all $n \in \mathbb{N}$. Therefore $(0, 1]x = \bigcup_{n=1}^{\infty} [m^n, m^{n-1}]x \subseteq D$. Thus D satisfies $tD \subseteq D$ for $t \in (0, 1]$. \square

For harmonically m -convex set-valued functions we obtain the following results:

Proposition 3.2. *Let $F_1, F_2 : D \rightarrow n(Y)$ be harmonically m -convex set-valued functions with $F_1(x) \subseteq F_2(x)$ (or $F_2(x) \subseteq F_1(x)$) for each $x \in D$. Then the union of F_1 and F_2 ($F_1 \cup F_2$) is a harmonically m -convex set-valued function.*

Proof. Let F_1, F_2 be harmonically m -convex set-valued function with $x, y \in D$, $t \in [0, 1]$ and $m \in (0, 1]$. Let's assume $F_1(x) \subseteq F_2(x)$ (in the case $F_2(x) \subseteq F_1(x)$ is analogous) for each $x \in D$, then:

$$\begin{aligned} t(F_1 \cup F_2)(y) + m(1-t)(F_1 \cup F_2)(x) &= t(F_1(y) \cup F_2(y)) + m(1-t)(F_1(x) \cup F_2(x)) \\ &= tF_2(y) + m(1-t)F_2(x) \\ &\subseteq F_2\left(\frac{mxy}{tmx + (1-t)y}\right) \\ &= F_1\left(\frac{mxy}{tmx + (1-t)y}\right) \cup F_2\left(\frac{mxy}{tmx + (1-t)y}\right) \\ &= (F_1 \cup F_2)\left(\frac{mxy}{tmx + (1-t)y}\right). \end{aligned}$$

\square

Proposition 3.3. *If $F : D \subset X \rightarrow n(Y)$ is a harmonically m -convex set-valued function, then the image of F of any harmonically m -convex subset of D is a m -convex set of Y .*

Proof. Let A be a harmonically m -convex subset of $D \subset X$ and $a, b \in F(A) = \cup_{z \in A} F(z)$. Then $a \in F(x)$ and $b \in F(y)$ for some $x, y \in A$. Thus, for all $t \in [0, 1]$ and $m \in (0, 1]$, we have to:

$$tb + m(1-t)a \in tF(y) + m(1-t)F(x) \subseteq F\left(\frac{mxy}{tmx + (1-t)y}\right).$$

Since A is harmonically m -convex set, we have $\frac{mxy}{tmx + (1-t)y} \in A$ and $tb + m(1-t)a \in F(A)$ for all $t \in [0, 1]$. Which implies that $F(A)$ is m -convex set of Y . \square

Corollary 3.4. *If $F : D \subseteq X \rightarrow n(Y)$ is a harmonically m -convex set-valued function, then the range of F is m -convex set of Y .*

Proof. If we consider $A = D$ in Proposition 3.3 we get that $Rang(F) = F(D)$. \square

Proposition 3.5. *A set-valued function $F : D \rightarrow n(Y)$ is harmonically m -convex, if and only if,*

$$(3.1) \quad tF(B) + m(1-t)F(A) \subseteq F\left(\frac{mAB}{tmA + (1-t)B}\right),$$

for each $A, B \subseteq D$, $t \in [0, 1]$ and $m \in (0, 1]$.

Proof. (\Rightarrow) Let A, B be arbitrary subsets of D , $t \in [0, 1]$ and $m \in (0, 1]$. Let $x \in t(F(B) = \cup_{b \in B} F(b)) + m(1-t)F(A) = \cup_{a \in A} F(a)$, that is to say $x \in tF(b) + m(1-t)F(a)$ for some $a \in A$ and $b \in B$. Since F is harmonically m -convex and $a, b \in D$, it follows that:

$$tF(b) + m(1-t)F(a) \subseteq F\left(\frac{mab}{tma + (1-t)b}\right),$$

moreover, $\frac{mab}{tma + (1-t)b} \in \frac{mAB}{tmA + (1-t)B}$ and, in consequence,

$$F\left(\frac{mab}{tma + (1-t)b}\right) \subset F\left(\frac{mAB}{tmA + (1-t)B}\right).$$

Therefore, $x \in F\left(\frac{mAB}{tmA + (1-t)B}\right)$.

(\Leftarrow) For $x, y \in D$ $t \in [0, 1]$ and $m \in (0, 1]$, the result is obtained by replacing $A = \{x\}$ and $B = \{y\}$ in (3.1) and we obtain the desired result

$$tF(\{y\}) + m(1-t)F(\{x\}) \subseteq F\left(\frac{m\{x\}\{y\}}{tm\{x\} + (1-t)\{y\}}\right).$$

Then, F is a harmonically m -convex set-valued function. \square

In the following, consider that for nonempty linear space subsets A, B, C, D and α a scalar, the following properties are true:

- $\alpha(A \times B) = \alpha A \times \alpha B$,
- $(A \times C) + (B \times D) = (A + B) \times (C + D)$,
- Si $A \subseteq B$ y $C \subseteq D$ then $A \times C \subseteq B \times D$.

Proposition 3.6. *Let $F_1 : D \rightarrow n(Y)$ and $F_2 : D \rightarrow n(Z)$ harmonically m -convex set-valued functions. Then the cartesian product $F_1 \times F_2$ is a harmonically m -convex set-valued function.*

Proof. Let $x, y \in D$ and $t \in [0, 1]$, then:

$$\begin{aligned}
t(F_1 \times F_2)(y) + m(1-t)(F_1 \times F_2)(x) &= [tF_1(y) \times tF_2(y)] + [m(1-t)F_1(x) \times m(1-t)F_2(x)] \\
&= (tF_1(y) + m(1-t)F_1(x)) \times (tF_2(y) + m(1-t)F_2(x)) \\
&\subseteq F_1\left(\frac{mxy}{tmx + (1-t)y}\right) \times F_2\left(\frac{mxy}{tmx + (1-t)y}\right) \\
&= (F_1 \times F_2)\left(\frac{mxy}{tmx + (1-t)y}\right).
\end{aligned}$$

□

The following proposition establishes that the harmonically m -convex set-valued functions are closed under the sum and the product by a scalar.

Proposition 3.7. *Let X, Y be two linear spaces. If D is a harmonically m -convex subset of X and $F, G : D \subset X \rightarrow n(Y)$ two harmonically m -convex set-valued functions. Then $\lambda F + G$ is a harmonically m -convex set-valued function, for all λ .*

Proof. Let $x, y \in D \subset X$, $t \in [0, 1]$ and $m \in (0, 1]$. Since F and G are harmonically m -convex set-valued functions, we have:

$$tF(y) + m(1-t)F(x) \subseteq F\left(\frac{mxy}{tmx + (1-t)y}\right),$$

and,

$$tG(y) + m(1-t)G(x) \subseteq G\left(\frac{mxy}{tmx + (1-t)y}\right).$$

Thus,

$$\begin{aligned}
t(\lambda F + G)(y) + m(1-t)(\lambda F + G)(x) &= [t(\lambda F(y)) + m(1-t)(\lambda F(x))] + [tG(y) + m(1-t)G(x)] \\
&\subseteq \lambda F\left(\frac{mxy}{tmx + (1-t)y}\right) + G\left(\frac{mxy}{tmx + (1-t)y}\right) \\
&= (\lambda F + G)\left(\frac{mxy}{tmx + (1-t)y}\right).
\end{aligned}$$

□

Proposition 3.8. *Let x, Y be two linear spaces. If D is a harmonically m -convex subset of X and $F, G : D \subset X \rightarrow n(Y)$ two harmonically m -convex set-valued functions then $(F \cdot G)(x)$ is also harmonically m -convex set-valued function.*

Proof. First, from [24], we have that.

$$F(x_1)G(x_2) + F(x_2)G(x_1) \subset F(x_1)G(x_1) + F(x_2)G(x_2).$$

Then, for $t \in [0, 1]$ and $x_1, x_2 \in D$:

$$\begin{aligned}
(F \cdot G) \left(\frac{mx_1x_2}{mtx_1 + (1-t)x_2} \right) &= F \left(\frac{mx_1x_2}{mtx_1 + (1-t)x_2} \right) \cdot G \left(\frac{mx_1x_2}{mtx_1 + (1-t)x_2} \right) \\
&\supseteq [tF(x_1) + m(1-t)F(x_2)][tG(x_1) + m(1-t)G(x_2)] \\
&= t^2F(x_1)G(x_1) + mt(1-t)F(x_1)G(x_2) + mt(1-t)F(x_2)G(x_1) \\
&\quad + m^2(1-t)^2F(x_2)G(x_2) \\
&= t^2F(x_1)G(x_1) + mt(1-t)[F(x_1)G(x_2) + F(x_2)G(x_1)] \\
&\quad + m^2(1-t)^2F(x_2)G(x_2) \\
&\supseteq t^2F(x_1)G(x_1) + mt(1-t)[F(x_1)G(x_1) + F(x_2)G(x_2)] \\
&\quad + m^2(1-t)^2F(x_2)G(x_2) \\
&= t[t + m(1-t)]F(x_1)G(x_1) + [m(1-t)[t + m(1-t)]F(x_2)G(x_2)] \\
&\supseteq tF(x_1)G(x_1) + m(1-t)F(x_2)G(x_2).
\end{aligned}$$

This shows that the product of two harmonically m -convex set-valued functions is again harmonically m -convex set-valued function. \square

The following result follows the idea of İ. İscan in [9]. To integrate set-valued functions we use the definition given by R. J. Aumann, and if a function satisfies the requirements of being integrable under this integral definition given, we say that a set-valued function F is Aumann integrable under a certain domain (see [2]).

Theorem 3.9. *Let X, Y linear spaces, D be a harmonically m -convex subset of X and $F : D \subset X \rightarrow n(Y)$ a harmonically m -convex set-valued Aumann integrable function, then*

$$\min \left(\inf \left(\frac{F(a) + mF\left(\frac{b}{m}\right)}{2} \right), \inf \left(\frac{mF\left(\frac{a}{m}\right) + F(b)}{2} \right) \right) \subseteq \frac{ab}{b-a} \int_a^b \frac{F(x)}{x^2} dx.$$

Proof. Let $F : D \subset X \rightarrow n(Y)$ be a harmonically m -convex set-valued function, for every $x, y \in D$ we have to

$$tF(y) + m(1-t)F\left(\frac{x}{m}\right) \subset F\left(\frac{xy}{tx + (1-t)y}\right) = F\left(\frac{m\frac{x}{m}y}{tm\frac{x}{m} + (1-t)y}\right),$$

Then, we have the following:

$$(3.2) \quad tF(b) + m(1-t)F\left(\frac{a}{m}\right) \subset F\left(\frac{ab}{ta + (1-t)b}\right),$$

and

$$tF(a) + m(1-t)F\left(\frac{b}{m}\right) \subset F\left(\frac{ab}{tb + (1-t)a}\right),$$

for every $t \in [0, 1]$, $m \in (0, 1]$ and $a, b \in D$. Integrating both sides of (3.2) on $[0, 1]$ with respect to t , we get that

$$(3.3) \quad \int_0^1 tF(b) + m(1-t)F\left(\frac{a}{m}\right) dt \subset \int_0^1 F\left(\frac{ab}{ta + (1-t)b}\right) dt.$$

Integrating the left side of (3.3) we have:

$$\int_0^1 tF(b) + m(1-t)F\left(\frac{a}{m}\right) dt = \frac{F(b) + mF\left(\frac{a}{m}\right)}{2}.$$

By the Aumann integral definition we get, that integral on the right hand of (3.3) is defined as:

$$\int_0^1 F\left(\frac{ab}{ta + (1-t)b}\right) dt = \left\{ \int_0^1 f\left(\frac{ab}{ta + (1-t)b}\right) dt : f(x) \in F(x) \wedge t \in [0, 1] \right\}.$$

But,

$$\int_0^1 f\left(\frac{ab}{ta + (1-t)b}\right) dt = \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx,$$

then

$$\left\{ \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx : f(x) \in F(x) \wedge x \in [a, b] \right\} = \frac{ab}{b-a} \int_a^b \frac{F(x)}{x^2} dx.$$

In consequence:

$$\frac{F(b) + mF\left(\frac{a}{m}\right)}{2} \subseteq \frac{ab}{b-a} \int_a^b \frac{F(x)}{x^2} dx.$$

Similarly, we have that:

$$\frac{F(a) + mF\left(\frac{b}{m}\right)}{2} \subseteq \frac{ab}{b-a} \int_a^b \frac{F(x)}{x^2} dx,$$

so the required result is obtained.

$$\min \left(\inf \left(\frac{F(a) + mF\left(\frac{b}{m}\right)}{2} \right), \inf \left(\frac{mF\left(\frac{a}{m}\right) + F(b)}{2} \right) \right) \subseteq \frac{ab}{b-a} \int_a^b \frac{F(x)}{x^2} dx.$$

□

REFERENCES

- [1] Anderson, G. D., Vamanamurthy, M. K. and Vuorinen, M. (2007). Generalized convexity and inequalities. *Journal of Mathematical Analysis and Applications*, **335**(2), 1294-1308.
- [2] Aumann, R. J. (1965). *Integrals of set-valued functions*. *Journal of mathematical analysis and applications*, **12**(1), 1-12.
- [3] Barráez D. , González L., Merentes N., and Moros A. (2015). *On h-convex stochastic processes*, *Mathematica Aeterna*, **5**(4), 571-581.
- [4] Berge, C. (1963). *Topological Space: Including a treatment of Multi-Valued functions, vector spaces and convexity*, Dover Publications, INC. Mineola, New York.
- [5] Geletu, A. (2006). *Introduction to topological spaces and set-valued maps (Lecture notes)*, Department of Operations Research & Stochastics Ilmenau University of Technology. August 25.
- [6] González, L., Merentes, N., and Valera-López, M. (2015) *Some estimates on the Hermite-Hadamard inequality through convex and quasi-convex stochastic processes*, *Mathematica Aeterna*, **5**(5), 745-767.
- [7] González, L., Merentes, N., and Valera-López, M. (2016). *On (k, h)-convex stochastic process*, *Journal of New Theory*, (10), 19-29.
- [8] İscan, İ. (2014) *Hermite-Hadamard type inequalities for harmonically convex functions*, *Hacettepe Journal of Mathematics and statistics* , **43** (6), 935-942.
- [9] İscan, İ. (2016) *Hermite-Hadamard type inequalities for harmonically (α, m) -convex functions*, *Hacettepe Journal of Mathematics and Statistics*, **45** (2), 381-390.
- [10] Lara . T. , Merentes N., Quintero R. and Rosales E. (2019). *On m-convexity of set-valued functions*, *Advance in Operator Theory*, **4** (14), 767-783.
- [11] Leiva, H., Merentes, N., Nikodem, K., and Sánchez, J. L. (2013) *Strongly convex set-valued maps*, *Journal of Global Optimization*, **57**(3), 695-705.
- [12] Materano, J., Merentes, N., and Valera-López, M. (2015). *Some estimates on the Simpson's type inequalities through s-convex and quasi-convex stochastic processes*, *Mathematica Aeterna*, **5**(5), 673-705.
- [13] Materano, J., Merentes, N., and Valera-López, M. (2016). *On Ostrowski's type inequalities via convex, s-convex and quasi-convex stochastic processes*. *Mathematica Aeterna*, **6**(1), 47-85.
- [14] Materano, J. E., Merentes, N., and López-Valera, M. (2017). *On inequalities of Hermite-Hadamard type for stochastic processes whose third derivative absolute values are quasi-convex*, *Tamkang Journal of Mathematics*, **48**(2), 203-208.
- [15] Matkowski J. and Nikodem K. (1998), *Convex set-valued functions on $(0, +1)$ and their conjugate*, *Rocznik Nauk.-Dydakt. Prace Mat.* (15) , 103-107.
- [16] Mejia, O., Merentes, N., and Nikodem, K. (2014). *Strongly concavemapse set-valued*, *Mathematica Aeterna*, **4**, 477-487.
- [17] Merentes N., Ribas S. (2013). *El desarrollo del concepto de función convexa*, Ediciones IVIC, Caracas-Venezuela.
- [18] Narváez D. X., Restrepo G. (2011). *Funciones Multivaluadas*, Facultad de Ciencias Naturales y Exactas Universidad del Valle, Revista de Ciencias, septiembre 20.
- [19] Nikodem K. (1987), *On concave and midpoint concave set-valued functions*, *Glasnik Mat. Ser. III* **22**(42),(1), 69-76.
- [20] Nikodem, K. (1989). *A characterization of midconvex set-valued function*, *Acta Universitatis Carolinae. Mathematica et Physica*, 30(2), 125-129.

- [21] Nikodem, K., Sánchez, J. L., and Sánchez, L. (2014). *Jensen and Hermite-Hadamard inequalities for strongly convex set-valued maps*, *Mathematica Aeterna*, **4**(8), 979-987.
- [22] Noor, M. A., Noor, K. I., Awan, M. U., and Costache, S. (2015). *Some integral inequalities for harmonically h -convex functions*, *Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys*, **77**(1), 5-16.
- [23] Sadowska, E. (1996). *A sandwich with convexity for set-valued functions*, *Mathematica Pannonica*, **7**(1), 163-169.
- [24] Santana G., González L. and Merentes N. (2019) *Funciones conjunto valuadas armónicamente convexas*, *Divulgaciones Matemáticas*, **19**(1), 20-33.
- [25] Toader, G. H. (1984, October). *Some generalizations of the convexity*, Cluj-Napoca. In Proc. Colloq. Approx. Optim. **329**, 338.

¹ ESCUELA DE MATEMÁTICA, FACULTAD DE CIENCIAS, UNIVERSIDAD CENTRAL DE VENEZUELA, CARACAS 1010, VENEZUELA.

Email address: gaszsantana@gmail.com

Email address: maira.valera@ciens.ucv.ve