

# An error estimate for the Gauss-Jacobi-Lobatto quadrature rule

Concetta Laurita<sup>1</sup>

*Department of Mathematics, Computer Science and Economics, University of Basilicata, Viale dell'Ateneo Lucano n. 10, 85100 Potenza, ITALY*

---

## Abstract

An error estimate for the Gauss-Lobatto quadrature formula for integration over the interval  $[-1, 1]$ , relative to the Jacobi weight function  $w^{\alpha,\beta}(t) = (1-t)^\alpha(1+t)^\beta$ ,  $\alpha, \beta > -1$ , is obtained. This estimate holds true for functions belonging to some Sobolev-type subspaces of the weighted space  $L^1_{w^{\alpha,\beta}}([-1, 1])$ .

*Keywords:* Gauss-Lobatto formula; Jacobi weight function

*2020 MSC:* 65D30, 65D32

---

## 1. Introduction

For a Jacobi weight function  $w^{\alpha,\beta}(t) = (1-t)^\alpha(1+t)^\beta$ ,  $\alpha, \beta > -1$ , on the interval  $[-1, 1]$  we consider the  $(n+2)$ -point Gauss-Lobatto rule

$$\int_{-1}^1 f(t)w^{\alpha,\beta}(t)dt = w_0f(-1) + \sum_{k=1}^n w_kf(t_k) + w_{n+1}f(1) + e_n(f) \quad (1)$$

which is exact for polynomials of degree at most  $2n+1$ , i.e.

$$e_n(f) = 0, \quad \forall f \in \mathbb{P}_{2n+1}$$

being  $\mathbb{P}_n$  the set of all algebraic polynomials on  $[-1, 1]$  of degree at most  $n$ .

It is well known that the interior quadrature nodes  $t_k$ ,  $k = 1, \dots, n$ , are the zeros of the Jacobi polynomial of degree  $n$  orthonormal with respect to the Jacobi weight  $w^{\alpha+1,\beta+1}(t) = (1-t)^2w^{\alpha,\beta}(t)$ . The weights of the formula (1) are given by

$$w_k = \int_{-1}^1 l_k(t)w^{\alpha,\beta}(t)dt, \quad k = 0, 1, \dots, n+1 \quad (2)$$

where, setting from now on  $t_0 = -1$  and  $t_{n+1} = 1$ ,  $l_k(t)$  denotes the  $(k+1)$ -th Lagrange fundamental polynomial associated to the system of nodes  $\{t_0, t_1, \dots, t_n, t_{n+1}\}$ .

By standard arguments, it can be easily proved that the quadrature error  $e_n(f)$  satisfies the following estimate

$$|e_n(f)| \leq 2 \left( \int_{-1}^1 w^{\alpha,\beta}(t) dt \right) E_{2n+1}(f)_\infty, \quad \forall f \in C([-1, 1]),$$

where

$$E_n(f)_\infty = \inf_{P \in \mathbb{P}_n} \|f - P\|_\infty$$

---

*Email address:* concetta.laurita@unibas.it (Concetta Laurita)

<sup>1</sup>Tel: +390971205846

denotes the error of best approximation of a function  $f \in C([-1, 1])$  by means of polynomials of degree at most  $n$  with respect to the uniform norm.

The aim of the present paper is to provide a new error estimate for less regular functions belonging to some Sobolev-type subspaces of the weighted space  $L_{w^{\alpha, \beta}}^1([-1, 1])$ . A similar estimate is proved in [3] for the classical Gauss-Jacobi quadrature formula, in [1] for the Gauss-Lobatto rule with respect to the Legendre weight  $w^{0,0}$  and, more recently, in [2] for the Gauss-Radau formula with respect to a general Jacobi weight  $w^{\alpha, \beta}$ .

## 2. Notation and preliminary results

### 2.1. Notation

For a general weight function  $w(t)$  on  $[-1, 1]$  and  $1 \leq p < +\infty$ , let  $L_w^p$  denote the weighted space of all real-valued measurable functions  $f$  on  $[-1, 1]$  such that

$$\|f\|_{L_w^p} = \|fw\|_p = \left( \int_{-1}^1 |f(t)w(t)|^p dt \right)^{\frac{1}{p}} < +\infty,$$

and let  $W_r^p(w)$  be the following weighted Sobolev-type subspaces of  $L_w^p$

$$W_r^p(w) = \left\{ f \in L_w^p : f^{(r-1)} \in AC(-1, 1), \|f^{(r)}\varphi^r w\|_p < +\infty \right\},$$

where  $r \in \mathbb{N}$ ,  $r \geq 1$ ,  $\varphi(t) = \sqrt{1-t^2}$  and  $AC(-1, 1)$  is the collection of all functions which are absolutely continuous on every closed subset of  $(-1, 1)$ , equipped with the norm

$$\|f\|_{W_r^p(w)} = \|fw\|_p + \|f^{(r)}\varphi^r w\|_p.$$

For a function  $f \in L_w^p$ , the error of the best approximation of  $f$  in  $L_w^p$  by polynomials of degree at most  $n$  is defined as

$$E_n(f)_{w,p} = \inf_{P \in \mathbb{P}_n} \|f - P\|_{L_w^p}.$$

Fixed a Jacobi weight  $w^{\gamma, \delta}(t) = (1-t)^\gamma(1+t)^\delta$ ,  $\gamma, \delta > -1$ , we will denote by  $x_{n,k}^{\gamma, \delta}$  and  $\lambda_{n,k}^{\gamma, \delta}$ ,  $k = 1, \dots, n$ , the nodes and coefficients of the corresponding  $n$ -point Gauss-Jacobi quadrature rule on  $[-1, 1]$  and by

$$p_n^{\gamma, \delta}(t) = \gamma_n^{\gamma, \delta} t^n + \text{lower degree terms}, \quad \gamma_n^{\gamma, \delta} > 0,$$

the Jacobi polynomial of degree  $n$  orthonormal w.r.t.  $w^{\gamma, \delta}(t)$  having positive leading coefficient.

In the sequel  $\mathcal{C}$  will denote a positive constant which may assume different values in different formulas. We write  $\mathcal{C} = \mathcal{C}(a, b, \dots)$  to say that  $\mathcal{C}$  is dependent on the parameters  $a, b, \dots$  and  $\mathcal{C} \neq \mathcal{C}(a, b, \dots)$  to say that  $\mathcal{C}$  is independent of them. Moreover, we will write  $A \sim B$ , if there exists a positive constant  $\mathcal{C}$  independent of the parameters of  $A$  and  $B$  such that  $1/\mathcal{C} \leq A/B \leq \mathcal{C}$ .

### 2.2. Preliminary results

It is well known that for functions  $f$  belonging to  $W_1^p(w)$ , the following Favard inequality

$$E_n(f)_{w,p} \leq \frac{\mathcal{C}}{n} E_{n-1}(f')_{\varphi w,p}, \quad (3)$$

is fulfilled for a positive constant  $\mathcal{C}$  independent of  $n$  and  $f$  (see, for example, [3, (2.5.22), p. 172]). By iteration of inequality (3), it follows that, for  $f \in W_r^p(w)$ ,  $r \geq 1$ , the estimate

$$E_n(f)_{w,p} \leq \frac{\mathcal{C}}{n^r} E_{n-r}(f^{(r)})_{\varphi^r w,p}, \quad \mathcal{C} \neq \mathcal{C}(n, f) \quad (4)$$

holds true.

Let us recall that the knots  $x_{n,k}^{\gamma,\delta}$  and Christoffel numbers  $\lambda_{n,k}^{\gamma,\delta}$  of the Gaussian quadrature formula corresponding to the Jacobi weight  $w^{\gamma,\delta}$  satisfy the following properties (see, for instance, [3, (4.2.4), p. 249])

$$x_{n,k+1}^{\gamma,\delta} - x_{n,k}^{\gamma,\delta} \sim \frac{\sqrt{1-t^2}}{n}, \quad x_{n,k}^{\gamma,\delta} \leq t \leq x_{n,k+1}^{\gamma,\delta}, \quad (5)$$

and (see [6, (14), p.673])

$$\lambda_{n,k}^{\gamma,\delta} \sim \frac{\sqrt{1 - \left(x_{n,k}^{\gamma,\delta}\right)^2}}{n} w^{\gamma,\delta}(x_{n,k}^{\gamma,\delta}) \quad (6)$$

uniformly for  $1 \leq k \leq n$ ,  $n \in \mathbb{N}$ . Moreover, for the orthonormal polynomials  $\{p_n^{\gamma,\delta}(t)\}_n$  one has that (see [7, (12.7.2), p. 309])

$$\frac{\gamma_n^{\gamma,\delta}}{\gamma_{n-1}^{\gamma,\delta}} \sim 1, \quad \text{as } n \rightarrow \infty, \quad (7)$$

and (see, for instance, [5, p. 170]) the equality

$$\frac{1}{p_{n-1}^{\gamma,\delta}(x_{n,k}^{\gamma,\delta})} = \frac{\gamma_{n-1}^{\gamma,\delta}}{\gamma_n^{\gamma,\delta}} \lambda_{n,k}^{\gamma,\delta} (p_n^{\gamma,\delta})'(x_{n,k}^{\gamma,\delta}) \quad (8)$$

holds true. Furthermore (see [5, Corollary 9.34, p. 171])

$$p_n^{\gamma,\delta}(1) \sim n^{\alpha+\frac{1}{2}} \quad |p_n^{\gamma,\delta}(-1)| \sim n^{\beta+\frac{1}{2}}, \quad (9)$$

uniformly for  $n \in \mathbb{N}$  and, more generally, [3, (4.4.49)-(4.2.30), p. 255])

$$|p_n^{\gamma,\delta}(t)| \sim n^{\alpha+\frac{1}{2}}, \quad 1 - \frac{\mathcal{C}}{n^2} \leq t \leq 1, \quad (10)$$

$$|p_n^{\gamma,\delta}(t)| \sim n^{\beta+\frac{1}{2}}, \quad -1 \leq t \leq -1 + \frac{\mathcal{C}}{n^2}. \quad (11)$$

### 3. Main results

**Lemma 3.1.** *The nodes and the weights of the Gauss-Lobatto quadrature formula (1) satisfy the following relations*

$$w_0 \leq \mathcal{C} \Delta t_0 w^{\alpha,\beta}(t_1), \quad (12a)$$

$$w_k \sim \begin{cases} \Delta t_k w^{\alpha,\beta}(t_k), & k = 1, \dots, n-1 \\ \Delta t_{k-1} w^{\alpha,\beta}(t_k) & k = n \end{cases}, \quad (12b)$$

$$w_{n+1} \leq \mathcal{C} \Delta t_n w^{\alpha,\beta}(t_n), \quad (12c)$$

where  $\Delta t_k = t_{k+1} - t_k$ ,  $k = 0, 1, \dots, n$  and  $\mathcal{C} \neq \mathcal{C}(n)$ .

*Proof.* First, let us observe that the weights of the formula (1), given in (2), have the following alternative representation

$$w_0 = \frac{1}{1-t_0} \int_{-1}^1 \frac{p_n^{\alpha+1,\beta+1}(t)}{p_n^{\alpha+1,\beta+1}(t_0)} w^{\alpha+1,\beta}(t) dt, \quad (13a)$$

$$w_k = \frac{1}{(1-t_k^2)} \int_{-1}^1 \frac{p_n^{\alpha+1,\beta+1}(t)}{(t-t_k) (p_n^{\alpha+1,\beta+1})'(t_k)} w^{\alpha+1,\beta+1}(t) dt, \quad k = 1, \dots, n, \quad (13b)$$

$$w_{n+1} = \frac{1}{1+t_{n+1}} \int_{-1}^1 \frac{p_n^{\alpha+1,\beta+1}(t)}{p_n^{\alpha+1,\beta+1}(t_{n+1})} w^{\alpha,\beta+1}(t) dt. \quad (13c)$$

For  $k = 1, \dots, n$ , since

$$w_k = \frac{\lambda_{n,k}^{\alpha+1,\beta+1}}{1 - \left(x_{n,k}^{\alpha+1,\beta+1}\right)^2},$$

using (6) and (5) for  $\gamma = \alpha + 1$  and  $\delta = \beta + 1$ , we can deduce that

$$w_k \sim \begin{cases} w^{\alpha,\beta}(t_k)\Delta t_k & k = 1, \dots, n-1 \\ w^{\alpha,\beta}(t_k)\Delta t_{k-1} & k = n \end{cases}$$

i.e. (12b). Now, in order to prove (12a), let us observe that, since

$$\int_{-1}^1 \left[ \frac{p_n^{\alpha+1,\beta+1}(t)}{p_n^{\alpha+1,\beta+1}(t_0)} - \left( \frac{p_n^{\alpha+1,\beta+1}(t)}{p_n^{\alpha+1,\beta+1}(t_0)} \right)^2 \right] w^{\alpha+1,\beta}(t) dt = 0,$$

we can rewrite the first coefficient  $w_0$  in (13a) as follows

$$w_0 = \frac{1}{2} \left[ \frac{(p_n^{\alpha+1,\beta+1})'(t_1)}{p_n^{\alpha+1,\beta+1}(t_0)} \right]^2 \int_{-1}^1 \left[ \frac{p_n^{\alpha+1,\beta+1}(t)}{(t-t_1)(p_n^{\alpha+1,\beta+1})'(t_1)} \right]^2 \frac{(t-t_1)^2}{1+t} w^{\alpha+1,\beta+1}(t) dt.$$

Being  $(t-t_1)^2/(1+t) \leq \mathcal{C}$  and using (8) (with  $\gamma = \alpha + 1$  and  $\delta = \beta + 1$ ) for  $k = 1$  (we recall that in our notation  $x_{n,1}^{\alpha+1,\beta+1} \equiv t_1$ ), we get

$$\begin{aligned} w_0 &\leq \mathcal{C} \left[ \frac{(p_n^{\alpha+1,\beta+1})'(t_1)}{p_n^{\alpha+1,\beta+1}(t_0)} \right]^2 \lambda_{n,1}^{\alpha+1,\beta+1} \\ &= \mathcal{C} \frac{1}{[p_n^{\alpha+1,\beta+1}(t_0)]^2} \left( \frac{\gamma_n^{\alpha+1,\beta+1}}{\gamma_{n-1}^{\alpha+1,\beta+1}} \right)^2 \frac{1}{\lambda_{n,1}^{\alpha+1,\beta+1} [p_{n-1}^{\alpha+1,\beta+1}(t_1)]^2}. \end{aligned}$$

Now, in virtue of (6), it is

$$\lambda_{n,1}^{\alpha+1,\beta+1} \sim \frac{\left(1 - x_{n,1}^{\alpha+1,\beta+1}\right)^{\alpha+\frac{3}{2}} \left(1 + x_{n,1}^{\alpha+1,\beta+1}\right)^{\beta+\frac{3}{2}}}{n} \sim \frac{1}{n^{2\beta+4}}. \quad (14)$$

Taking into account (14), and (7), (9) and (11) (all applied with  $\gamma = \alpha + 1$  and  $\delta = \beta + 1$ ), we deduce the following estimate of  $w_0$

$$w_0 \leq \mathcal{C} \frac{n^{2\beta+4}}{n^{4\beta+6}} = \frac{\mathcal{C}}{n^{2\beta+2}}. \quad (15)$$

On the other hand,

$$\Delta t_0 w^{\alpha,\beta}(t_1) = \left(1 - x_{n,1}^{\alpha+1,\beta+1}\right)^\alpha \left(1 + x_{n,1}^{\alpha+1,\beta+1}\right)^{\beta+1} \sim \frac{1}{n^{2\beta+2}}$$

which, combined with (15), gives (12a). In order to prove (12c), proceeding in an analogous way, we start writing  $w_{m+1}$  as follows

$$w_{m+1} = \frac{1}{2} \left[ \frac{(p_n^{\alpha+1,\beta+1})'(t_n)}{p_n^{\alpha+1,\beta+1}(t_{n+1})} \right]^2 \int_{-1}^1 \left[ \frac{p_n^{\alpha+1,\beta+1}(t)}{(t-t_n)(p_n^{\alpha+1,\beta+1})'(t_n)} \right]^2 \frac{(t-t_n)^2}{1-t} w^{\alpha+1,\beta+1}(t) dt.$$

Then, using  $(t - t_n)^2/(1 - t) \leq \mathcal{C}$ , (6) and (8) for  $\gamma = \alpha + 1$ ,  $\delta = \beta + 1$  and  $k = n$ , (7), (9) and (10) also with  $\gamma = \alpha + 1$  and  $\delta = \beta + 1$ , we deduce the estimate

$$w_{n+1} \leq \mathcal{C} \left[ \frac{(p_n^{\alpha+1, \beta+1})'(t_n)}{p_n^{\alpha+1, \beta+1}(t_{n+1})} \right]^2 \lambda_{n,n}^{\alpha+1, \beta+1} \sim \frac{1}{n^{2\alpha+2}}$$

with  $\mathcal{C} \neq \mathcal{C}(n)$ . Since

$$\Delta t_n w^{\alpha, \beta}(t_n) \sim (1 - x_{n,n}^{\alpha+1, \beta+1})^{\alpha+1} (1 + x_{n,n}^{\alpha+1, \beta+1})^\beta \sim \frac{1}{n^{2\alpha+2}}$$

we can, finally, conclude that also (12c) holds true.  $\square$

Using the previous lemma we are able to prove our main result.

**Theorem 3.1.** *For  $f \in W_r^1(w^{\alpha, \beta})$ ,  $r \geq 1$ , the error of the Gauss-Lobatto quadrature formula (1) satisfies the following estimate*

$$|e_n(f)| \leq \frac{\mathcal{C}}{(2n)^r} E_{2n+1-r}(f^{(r)})_{\varphi^r w^{\alpha, \beta}, 1} \quad (16)$$

where  $\varphi(t) = \sqrt{1 - t^2}$  and  $\mathcal{C} \neq \mathcal{C}(f, n)$  is a positive constant.

*Proof.* We start by proving the estimate (16) in the case  $r = 1$ . First, we are going to show that

$$\sum_{k=0}^{n+1} w_k |f(t_k)| \leq \mathcal{C} \|f w^{\alpha, \beta}\|_1 + \frac{\mathcal{C}}{n} \int_{-1}^1 |f'(t)| \varphi(t) w^{\alpha, \beta}(t) dt. \quad (17)$$

Taking into account (12a), (12b) and (12c) we have

$$\begin{aligned} \sum_{k=0}^{n+1} w_k |f(t_k)| &\leq \mathcal{C} [ \Delta t_0 w^{\alpha, \beta}(t_1) |f(t_0)| + \sum_{k=1}^{n-1} \Delta t_k w^{\alpha, \beta}(t_k) |f(t_k)| + \\ &\quad + \Delta t_{n-1} w^{\alpha, \beta}(t_n) |f(t_n)| + \Delta t_n w^{\alpha, \beta}(t_n) |f(t_{n+1})| ]. \end{aligned}$$

In virtue of the following inequality

$$\left. \begin{aligned} (b-a)|f(a)| \\ (b-a)|f(b)| \end{aligned} \right\} \leq \int_a^b |f(t)| dt + (b-a) \int_a^b |f'(t)| dt,$$

it follows that

$$\begin{aligned} \sum_{k=0}^{n+1} w_k |f(t_k)| &\leq \mathcal{C} \left[ w^{\alpha, \beta}(t_1) \left( \int_{t_0}^{t_1} |f(t)| dt + \Delta t_0 \int_{t_0}^{t_1} |f'(t)| dt \right) \right. \\ &\quad + \sum_{k=1}^{n-1} w^{\alpha, \beta}(t_k) \left( \int_{t_k}^{t_{k+1}} |f(t)| dt + \Delta t_k \int_{t_k}^{t_{k+1}} |f'(t)| dt \right) \\ &\quad + w^{\alpha, \beta}(t_n) \left( \int_{t_{n-1}}^{t_n} |f(t)| dt + \Delta t_{n-1} \int_{t_{n-1}}^{t_n} |f'(t)| dt \right) \\ &\quad \left. + w^{\alpha, \beta}(t_n) \left( \int_{t_n}^{t_{n+1}} |f(t)| dt + \Delta t_n \int_{t_n}^{t_{n+1}} |f'(t)| dt \right) \right], \end{aligned}$$

from which, being for  $k = 0, 1, \dots, n$ ,

$$1 \pm t_k \sim 1 \pm t \sim 1 \pm t_{k+1}, \quad t_k \leq t \leq t_{k+1}, \quad (18)$$

and, taking into account (5) and also that

$$\begin{aligned}\Delta t_0 &\sim \frac{\sqrt{1-t^2}}{n}, & -1 < t \leq t_1, \\ \Delta t_n &\sim \frac{\sqrt{1-t^2}}{n}, & t_n \leq t < 1,\end{aligned}$$

we deduce

$$\begin{aligned}\sum_{k=0}^{n+1} w_k |f(t_k)| &\leq \mathcal{C} \left[ \int_{t_0}^{t_1} |f(t)| w^{\alpha,\beta}(t) dt + \frac{1}{n} \int_{t_0}^{t_1} |f'(t)| \varphi(t) w^{\alpha,\beta}(t) dt \right. \\ &+ \sum_{k=1}^{n-1} \left( \int_{t_k}^{t_{k+1}} |f(t)| w^{\alpha,\beta}(t) dt + \frac{1}{n} \int_{t_k}^{t_{k+1}} |f'(t)| \varphi(t) w^{\alpha,\beta}(t) dt \right) \\ &+ \left. \int_{t_n}^{t_{n+1}} |f(t)| w^{\alpha,\beta}(t) dt + \frac{1}{n} \int_{t_n}^{t_{n+1}} |f'(t)| \varphi(t) w^{\alpha,\beta}(t) dt \right] \\ &= \mathcal{C} \left[ \int_{-1}^1 |f(t)| w^{\alpha,\beta}(t) dt + \frac{1}{n} \int_{-1}^1 |f'(t)| \varphi(t) w^{\alpha,\beta}(t) dt \right]\end{aligned}$$

i.e. (17). Since the  $(n+1)$  quadrature formula (1) is exact for any polynomial of degree at most  $2n+1$ , for  $P \in \mathbb{P}_{2n+1}$  we have

$$\begin{aligned}|e_n(f)| &= |e_n(f-P)| \\ &\leq \left| \int_{-1}^1 (f-P)(t) w^{\alpha,\beta}(t) dt \right| + \left| \sum_{k=0}^{n+1} w_k (f-P)(t_k) \right| \\ &\leq \|(f-P)w^{\alpha,\beta}\|_1 + \sum_{k=0}^{n+1} w_k |(f-P)(t_k)|.\end{aligned}$$

Now, combining the previous inequality with (17) and the following relation (see, for instance, [3, p. 339], [4, p. 286])

$$\|(f-P)' \varphi w^{\alpha,\beta}\|_1 \leq \mathcal{C}(2n+2) \|(f-P)w^{\alpha,\beta}\|_1 + \mathcal{C}_1 E_{2n}(f')_{\varphi w^{\alpha,\beta},1},$$

we can deduce

$$\begin{aligned}|e_n(f)| &\leq \mathcal{C} \|(f-P)w^{\alpha,\beta}\|_1 + \frac{\mathcal{C}}{n} \|(f-P)' \varphi w^{\alpha,\beta}\|_1 \\ &\leq \mathcal{C} \|(f-P)w^{\alpha,\beta}\|_1 + \frac{\mathcal{C}_1}{n} E_{2n}(f')_{\varphi w^{\alpha,\beta},1}.\end{aligned}$$

Taking the infimum over  $P \in \mathbb{P}_{2n+1}$  and using the Favard inequality (4) we get

$$|e_n(f)| \leq \mathcal{C} E_{2n+1}(f)_{w^{\alpha,\beta},1} + \frac{\mathcal{C}_1}{n} E_{2n}(f')_{\varphi w^{\alpha,\beta},1} \leq \frac{\mathcal{C}}{2n} E_{2n}(f')_{\varphi w^{\alpha,\beta},1}. \quad (19)$$

The estimate (16) for  $r > 1$  can be deduced from (19) by iterating the application of the Favard inequality.  $\square$

## Acknowledgements

The author is partially supported by University of Basilicata (local funds). The author is member of the INdAM Research group GNCS and the TAA-UMI Research Group. This research has been accomplished within ‘‘Research Italian network on Approximation’’ (RITA).

## References

- [1] L. Fermo, C. Laurita, A Nyström method for a boundary integral equation related to the Dirichlet problem on domains with corners, *Numer. Math.* 130 (2015) 35–71.
- [2] C. Laurita, A new stable numerical method for Mellin integral equations in weighted spaces with uniform norm, *Calcolo* 57 (2020) 25.
- [3] G. Mastroianni, G.V. Milovanović, *Interpolation Processes Basic Theory and Applications*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, Heidelberg, 2009.
- [4] G. Mastroianni, M.G. Russo, Lagrange interpolation in weighted Besov spaces, *Constr. Approx.* 15 (1999) 257–289.
- [5] G.P. Nevai, *Orthogonal Polynomials*, Amer. Math. Soc., Providence, R.I., 1979.
- [6] G.P. Nevai, Mean convergence of Lagrange interpolation, III, *Trans. Amer. Math. Soc.* 282 (1984) 669–698.
- [7] G. Szegő, *Orthogonal polynomials*, volume 23 of *Amer. Math. Soc. Colloq. Publ.*, Amer. Math. Soc., Providence, R.I., 4th ed. edition, 1975.