

Hydrostatic equilibrium configurations of quark stars in a non-minimal geometry-matter coupling theory of gravity

G.A. Carvalho[‡], R. Lobato^{◊,‡}, P.H.R.S. Moraes[†], F. Rocha^{*}, D. Deb^{**} and M. Malheiro^{*}

[‡]*Universidade do Vale do Paraíba, Instituto de Pesquisa e*

Desenvolvimento (IP&D), 12244-000, São José dos Campos, SP, Brazil

^{*}*Instituto Tecnológico de Aeronáutica (ITA), 12228-900, São José dos Campos, SP, Brazil*

[◊]*Departamento de Física, Universidad de los Andes, Bogotá, Colombia*

[‡]*Department of Physics and Astronomy, Texas A&M University - Commerce, TX 75428, USA*

[†]*Universidade Federal do ABC (UFABC) - Centro de Ciências Naturais e Humanas (CCNH) - Avenida dos Estados 5001, 09210-580, Santo André, SP, Brazil*

^{**}*Department of Physics, Indian Institute of Science, Bangalore, 560012, India*

This work analyses the hydrostatic equilibrium configurations of strange stars in a non-minimal geometry-matter coupling (GMC) theory of gravity. Those stars are supposed to be made of strange quark matter, whose distribution is governed by the MIT equation of state. The non-minimal GMC theory is described by the following gravitational action: $f(R, L) = R/2 + L + \sigma RL$, where R represents the curvature scalar, L is the matter Lagrangian density, and σ is the coupling parameter. When considering this theory, the strange stars become larger and more massive. In particular, when $\sigma = 20$, the theory can achieve the 2.6 solar mass, suitable for describing the pulsars PSR J2215+5135 and PSR J1614-2230, and the mass of the secondary object in the GW190814 event. The 2.6 M_{\odot} is a value hardly achievable in General Relativity. The non-minimal GMC theory can also give feasible results to describe the macroscopical features of strange star candidates.

I. INTRODUCTION

Recent observations regarding type Ia supernovae [1–3] and cosmic microwave background radiation [4–6] indicate that, presently, our universe is going through an accelerated expanding phase. Within the General Theory of Relativity (GR) context, the inclusion of the cosmological constant Λ into the Einstein gravitational field equations is the standard way to explain the cosmic acceleration and provide a good agreement with the observed data. However, the inclusion of Λ faces the major setback due to a considerable mismatch of 120 orders of magnitude between its observational and theoretical values [7, 8].

This situation engaged different researchers on more sophisticated gravity theories by modifying the Einstein-Hilbert action, which gave rise to a new avenue known as modified/extended gravity theories. These theories offer a great opportunity to solve problems that still do not have convincing explanations within the GR framework. In this regard, T. Harko and F.S.N. Lobo generalized the well-known $f(R)$ -type gravity model [9, 10] by assuming that the gravitational Lagrangian is given by an arbitrary function of the Ricci scalar R and the matter Lagrangian L , in the so-called $f(R, L)$ theory [11]. The dynamics in such a theory can only exist in the presence of matter, which suggests a deeper link between spacetime and matter. In fact, the $f(R, L)$ gravity is a subclass of the geometry-matter coupling (GMC) theories [12–14], i.e., theories that allow geometry and matter scalars to be mixed in the gravitational action.

The viability of $f(R, L)$ gravity as an alternative explanation for the cosmic acceleration was analyzed from a dynamical system approach in [15]. Some constraints were obtained to $f(R, L)$ theories using the COBEFIRAS

measurement of the cosmic microwave background spectral radiance [16]. The application of the energy conditions in the $f(R, L)$ gravity can be seen in [17, 18]. Harko et al. have discussed the non-conservation of the energy-momentum tensor in $f(R, L)$ models in [19, 20] and related it to a mechanism responsible for gravitationally induced particle production. Very recently, the $f(R, L)$ gravity was studied from a thermodynamic point of view [21].

In reference [22], it was indicated that the $f(R, L)$ theories of gravity possibly be regarded as a subclass of the $f(R, T)$ gravity theories [23], in which T is the trace of the energy-momentum tensor, with the latter theory also allowing for the GMC. In addition, in [24] the $f(R, L)$ gravity action was generalized by inserting on it a scalar field. Moreover, a further model with GMC was proposed by Harko in [25]. In [26, 27] it was shown that GMC models can be candidates to solve fundamental issues of standard gravity, without considering dark energy [28, 29] and dark matter [30–32]. Note that it is generally believed that dark energy is the cause of the accelerated expansion. On the other hand, dark matter is an exotic kind of matter that does not interact with light, but interacts gravitationally and strongly affects the galactic and intergalactic dynamics. For a review of generalized GMC theories, one can also check [33].

A well-behaved extended or alternative theory of gravity must also show significant effect in the stellar astrophysics regime. In other words, a given alternative theory of gravity should exhibit significant effects in the cosmological and galactic scales, as well as predict the existence of observable stable, compact stellar objects, such as white dwarfs, neutron stars and black holes. In fact, the study and analysis of compact objects are of great importance in astrophysics because these objects provide

an excellent laboratory to study dense matter in extreme conditions, such as the strong gravity regime. In particular, neutron stars were already studied in $f(R, L)$ gravity, providing a remarkable increase in the maximum mass limit [34, 35]. In [34], the matter inside neutron stars was described by a relativistic polytropic equation of state (EoS), and also a Skyrme type EoS known as SLy4. It was showed in this theory that the mass of massive pulsars can be achieved, such as PSR J2215 + 5135, for both equations of state. It was pointed out that results for mass-radius relation in GMC gravity are strongly dependent on the stiffness of the EoS. A further investigation of neutron stars in this theory, considering several realistic nuclear matter equations of state was performed in [35]. NS masses and radii obtained were subject to observational constrains from massive pulsars, the gravitational wave event GW170817 and from the PSR J0030+0451 mass-radius from NASA's Neutron Star Interior Composition Explorer (NICER) data. It was shown that in this theory of gravity, the mass-radius results can accommodate massive pulsars. The mass-radius relation in GMC theory shows a sharp increase in the star mass around one solar mass up to $2.25 M_{\odot}$ with a very small change in the star radius, which is ~ 13 Km for a strong gravity matter coupling. These results are in agreement with the NS radius region constrained by PSR J0030+0451 and by the very massive PSR J0740+6620 obtained in NICER observations [36–38], and also in accordance with the GW170817 event [39].

Until now, hydrostatic equilibrium configurations of quark stars in a non-minimal GMC model have not been investigated. The strange quark matter, made up of approximately equal numbers of unconfined up, down and strange quarks, may be the absolute ground state of the strong interaction [40–43]. There may exist objects entirely made of strange matter [44, 45]. Note that three flavor strange quark matter is more stable than the two flavor non-strange ones [46, 47]. As a result, a new class of compact objects has been postulated to exist, namely quark stars, almost self-bounded systems with an energy density at the star surface - given in the MIT model by the confined Bag energy \mathcal{B} - in contrast with NSs that are bounded by gravity. Some constraints have been put to quark stars from gravitational waves [48, 49], particularly from the GW170817 event [39].

In the present work, we are particularly interested in analyzing quark stars in a non-minimal GMC model, which shall be presented in Section II A. In Section II B the hydrostatic equilibrium equations for the concerned theory will be reviewed. In Section III we will present the equation of state (EoS) that we shall consider for numerically solving the hydrostatic equilibrium equations in Section IV. In Section V we present our conclusions.

II. BASIC FORMALISM

A. A Non-minimal Geometry-Matter Coupling Theory

The concerned modified form of the Einstein-Hilbert action reads [11]:

$$S = \int d^4x \sqrt{-g} f(R, L), \quad (1)$$

being $f(R, L)$ an arbitrary function of R and L . The constants $8\pi G$ and c , with G being the Newtonian gravitational constant and c the speed of light, are taken as 1. One can observe from (1), that when $f(R, L) = R/2 + L$, the standard form of Einstein-Hilbert action is retrieved, which leads to the standard Einstein's field equations: $G_{\mu\nu} = T_{\mu\nu}$, where $T_{\mu\nu}$ is the energy-momentum tensor.

Following [50, 51], we consider a GMC model defined by $f(R, L) = R/2 + L(1 + \sigma R)$. Furthermore, we will assume $L = -p$ [52], where p is the pressure of the fluid. Applying the variational principle in (1), and taking into account the considerations above, it follows that the field equations become [11, 34]

$$(1 - 2\sigma p)G_{\mu\nu} + \frac{1}{3}Rg_{\mu\nu} - \frac{\sigma p}{3}Rg_{\mu\nu} = (1 + \sigma R) \left(T_{\mu\nu} - \frac{1}{3}Tg_{\mu\nu} \right) - 2\sigma \nabla_{\mu} \nabla_{\nu} p. \quad (2)$$

Furthermore, the covariant derivative of the energy-momentum tensor reads [11, 34]

$$\nabla^{\mu} T_{\mu\nu} = (-pg_{\mu\nu} - T_{\mu\nu}) \nabla^{\mu} \ln(\sigma R). \quad (3)$$

B. The Hydrostatic Equilibrium Equations in a Non-minimal Geometry-Matter Coupling Model

The hydrostatic equilibrium equations in the concerned GMC theory were previously derived by adopting a spherically symmetric metric in its canonical form,

$$ds^2 = e^{\alpha(r)} dt^2 - e^{\beta(r)} dr^2 - r^2 (d\theta^2 + \sin^2 \theta d\phi^2), \quad (4)$$

with $\alpha(r)$ and $\beta(r)$ being the potentials depending on r only. The energy-momentum tensor of a perfect fluid is defined as $T_{\mu\nu} = \text{diag}(e^{\alpha}\rho, e^{\beta}p, r^2p, r^2 \sin^2 \theta p)$, where ρ is the matter-energy density. From the substitution of (4) into (3) one can find that the energy-momentum tensor is covariantly conserved independently of the functional form assumed for $f(R, L)$. Detailed derivation is presented in [34].

The equilibrium configurations are obtained from the 00 and 11 components of the field equations,

$$\frac{(1 - 2\sigma p)}{r^2} [r(1 - e^{-\beta})]' + (1 - \sigma p) \frac{R}{3} = (1 + \sigma R) \left(\frac{2}{3}\rho + p \right) + \sigma e^{-\beta} \alpha' p', \quad (5)$$

$$\begin{aligned} & \frac{(1-2\sigma p)}{r^2} [e^{-\beta}(1-\alpha'r) - 1] + (\sigma p - 1)\frac{R}{3} \\ & = (1 + \sigma R)\frac{\rho}{3} - 2\sigma e^{-\beta} \left(p'' - \frac{\beta'}{2}p' \right), \end{aligned} \quad (6)$$

where prime represents the derivative with respect to radial coordinate r .

The conservation of the energy-momentum tensor leads to

$$p' = -(\rho + p)\frac{\alpha'}{2}. \quad (7)$$

Another equation can be derived from the trace of the field equations and reads

$$(1 + 2\sigma p)R = -(1 + \sigma R)T - 6\sigma \square p, \quad (8)$$

where \square is the D'Alembertian operator, defined as

$$\square = -e^{-\beta} \left(\frac{d^2}{dr^2} - \frac{\beta'}{2} \frac{d}{dr} + \frac{\alpha'}{2} \frac{d}{dr} + \frac{2}{r} \frac{d}{dr} \right). \quad (9)$$

In order to obtain the hydrostatic equilibrium configurations, (8) must be included in the system of differential equations (5), (6) and (7). The unknowns are R, α, β, ρ and p . An EoS must be provided to solve the set of differential equations. Details about the numerical procedure to solve the system of equations (5), (6), (7) and (8) are given in [34].

The stellar mass is obtained from

$$M = \int_0^{R_*} 4\pi r^2 \rho dr, \quad (10)$$

where R_* represents the stellar radius and by definition, pressure vanishes, $p(R_*) = 0$.

III. EQUATION OF STATE FOR NUCLEAR MATTER INSIDE QUARK STARS

The EoS for the matter inside the star is considered to be that for quark-gluon plasma, dubbed MIT bag model [53]. To consider hadronic masses in terms of their constituents, the MIT bag model considers that the quarks are inside a “bag” which reproduces the asymptotic freedom and confinement, i.e.,

$$p = \omega(\rho - 4\mathcal{B}), \quad (11)$$

where ω is the EoS parameter and \mathcal{B} is the bag constant, which we take as $\mathcal{B} = 60 \text{ MeV/fm}^3$ in allusion to [54], among many others. We adopt this bag constant value since the parameters of the maximum mass configurations for quark stars are similar to those for realistic neutron stars made of baryonic matter. The value of ω is related to the Quantum Chromodynamics coupling constant, and the strange quark mass [55]. For $\omega = 0.28$, the mass of the strange quark is $m_s = 250 \text{ MeV}$, and $\omega = 1/3$ for massless quarks. In this work, we adopt $\omega = 0.28$.




FIG. 1: Mass-radius relation of quark stars for different values of σ .




FIG. 2: Mass-central energy density relation of quark stars for different values of σ .

IV. RESULTS

In Figure 1, we present the mass-radius relation for quark stars in the GMC theory of gravitation for five values of σ , the GMC parameter. Blue circles mark the maximum mass points for each value of σ . The two horizontal lines denote the 2.3 and $2.6 M_\odot$ values of mass, which correspond to the PSR J2215+5135 [56], and the mass of the lighter object of the GW190814 event [65], respectively. When σ equals zero, GR results are retrieved. From the figure, it is possible to see that the radius of the star ranges from 7 to 11 km approximately. This radius interval lies within the expected values of compact star radii constraints from observed X-ray binaries, and GW170817 gravitational wave event [66, 67].

Figure 1 also shows that maximum mass points change according to the value of σ . For larger σ , more massive stars are found. Excluding the case where $\sigma = 20$, the

TABLE I: Physical parameters of observed strange star candidates derived using $\sigma = 10$ and $B = 60$ MeV/fm³.

SS candidate	Observed mass M/M_\odot	Predicted radius (km)	M/R_\star	Z_s	ρ_C (MeV/fm ³)	p_C (MeV/fm ³)
PSR J2215 + 5135	$2.27^{+0.17}_{-0.15}$ [56]	$10.64^{+0.38}$	0.301	0.586	1240	280
PSR J1614-2230	1.97 ± 0.04 [57]	$11.13^{+0.003}_{-0.09}$	0.261	0.448	715	133
Vela X-1	1.77 ± 0.08 [58]	11.13 ± 0.02	0.235	0.373	604	102
4U 1608-52	1.74 ± 0.14 [59]	$11.13^{+0.01}_{-0.06}$	0.231	0.363	591	98
PSR J1903+327	1.667 ± 0.021 [60]	$11.11^{+0.004}_{-0.01}$	0.221	0.340	562	90
4U 1820-30	1.58 ± 0.06 [61]	$11.07^{+0.02}_{-0.04}$	0.211	0.315	532	82
Cen X-3	1.49 ± 0.08 [58]	$11.01^{+0.05}_{-0.006}$	0.200	0.291	506	74
EXO 1785-248	1.3 ± 0.2 [62]	$10.85^{+0.17}_{-0.24}$	0.177	0.244	458	61
LMC X-4	1.29 ± 0.05 [58]	$10.84^{+0.05}_{-0.06}$	0.176	0.242	455	60
SMC X-1	1.04 ± 0.09 [58]	$10.52^{+0.13}_{-0.14}$	0.146	0.188	406	46
SAX J1808.4-3658	0.9 ± 0.3 [63]	$10.29^{+0.45}_{-0.71}$	0.130	0.161	383	40
4U 1538-52	0.87 ± 0.07 [58]	$10.23^{+0.11}_{-0.15}$	0.125	0.156	378	39
HER X-1	0.85 ± 0.15 [64]	$10.12^{+0.34}_{-0.27}$	0.124	0.153	375	38

TABLE II: Maximum masses with their correspondent radius, central energy density and central pressure for five values of σ .

σ	Maximum mass (M/M_\odot)	Radius (km)	ρ_C (MeV/fm ³)	p_C (MeV/fm ³)
0	1.97	10.10	1413	328
5	2.12	10.29	1413	328
10	2.28	10.43	1472	345
15	2.48	10.55	1559	369
20	2.71	10.51	1822	443

minimum radius also increases with increasing σ . This behavior is reversed for $\sigma = 20$. This is similar to the effects of charge on the stellar structure of compact objects [68–71], where increasing total charge initially increases the minimum radius, but at some point continued increasing in charge yields a minimum radius reduction. It is important to stress that in the GMC theory, the maximum star mass is not only a function of the bag constant, but also depends on σ , while in GR, the quark star mass increases with the star volume almost up to the maximum star mass that is only a function of the bag constant [72–74].

Increasing σ (for $\sigma > 15$) yields larger maximum mass stars with smaller radii, which means that the stars become more compact. This effect of increasing the maximum mass while diminishing minimum radii can lead to a violation of the Buchdahl limit if the star has a mass of $2.5 M_\odot$ and radius $R_\star < 8$ km. From Figure 1, it is clear that the maximum mass stars have radii between 9–10 km, indicating that the Buchdahl limit is maintained.

Figure 2 presents the mass-central density relation for the same five values of σ as in Figure 1. By increasing σ , maximum masses are always found for larger central energy densities. At smaller densities, the GMC results becomes closer to GR ones ($\sigma = 0$), which can be seen from both Figures 1 and 2. This indicates that GMC effects are more evident at a high density regime, so, in less compact systems, such as non-compact stellar objects or

in the solar system regime, GMC outcomes would be the same as the GR ones. Table II below is a list of the stellar parameters of the maximum mass quark stars in GMC theory, where we can see that the central energy densities of the maximum mass stars are between 3.0–3.6 times the nuclear saturation energy density ($\rho_0 \sim 140$ MeV/fm³).

We see that GMC theory can enhance maximum masses, allowing compact stars to sustain more mass against gravitational collapse, which indicates that the theory is capable of describing the data of massive compact stars, such as PSR J2215+5135 [56] and PSR J1614-2230 [57].

Recently, the LIGO (Laser Interferometer Gravitational wave Observatory)/VIRGO experiments have detected the GW190814 event [65], which is a binary merger of two compact objects, with one of them, a neutron star or a strange quark star, having a mass of $2.6 M_\odot$. This mass value can be described using $\sigma \approx 20$, which sets an observational upper limit to the GMC parameter.

In Figure 3, we present the compactness and surface redshift as a function of the central energy density of the stars. These quantities are given by

$$u = \frac{M}{R_\star} \quad \text{and} \quad Z_s = \frac{1}{\sqrt{1 - 2M/R_\star}} - 1, \quad (12)$$

respectively. It is observed that as the parameter σ increases, the compactification degree is higher. The sur-

TABLE III: Physical parameters of the strange star candidate LMC X-4 for different values of σ and $B = 60$ MeV/fm³.

σ	Predicted radius (km)	M/R_*	Z_s	ρ_C (MeV/fm ³)	p_C (MeV/fm ³)
0	10.54 ^{+0.11} _{-0.12}	0.181	0.251	502	73
5	10.69 ^{+0.05} _{-0.04}	0.178	0.247	476	66
10	10.84 ^{+0.05} _{-0.06}	0.176	0.242	455	60
15	10.97 ^{+0.06} _{-0.05}	0.174	0.238	439	56
20	11.10 ^{+0.06} _{-0.06}	0.172	0.234	424	52







FIG. 3: Compactness and surface red-shift as function of the star's central energy density.

face redshift is also shown to increase with central energy density and with σ .

In Figure 4, we show the energy density and pressure profiles for a strange quark star with a central pressure of 500 MeV/fm³ in the GMC theory. Pressure and energy density are shown to decrease as radial coordinate increases. In the cases where $\sigma \neq 0$, density and pressure are larger and take longer to diminish. This yields a larger mass according to increasing σ and a slightly larger radius. This exemplifies how increasing σ yields a weaker

FIG. 4: Profiles of pressure and energy density inside a bare quark star with central pressure of 500 MeV/fm³. This is an example to illustrate that the GMC theory yields to well-behaved scenarios, where it is straightforward to show that energy conditions are respected.

gravitational force, and hence how more mass is sustained against gravitational collapse. Moreover, the variations of pressure and energy density are always negative, indicating that the energy conditions [75] are respected.

In Table I, we have used $\sigma = 10$ and $B = 60$ MeV/fm³ to derive the radius, compactness, surface redshift, cen-

tral energy density and central pressure of strange star candidates. The table shows that the stars have high surface redshift (0.15-0.45) and compactness (0.12-0.26), which is reinforced by the results for central energy densities (3-3.6 ρ_0 , where ρ_0 represents an average value for nuclear saturation energy density). From Figure 1 and Table I, one can observe that most stars have radii within the range 10-11 km, which can be a good marker to analyze strange quark stars in GMC theories.

In Table III, we took as a test the mass of the strange star candidate LMC X-4 ($M = 1.29 \pm 0.05 M_\odot$) to predict its radius, compactness, surface redshift, central energy density and central pressure for different values of σ and $B = 60 \text{ MeV/fm}^3$. The LMC X-4 is an object extensively pointed as a strange star candidate due to its structure and characteristics [76–80].

V. CONCLUSIONS

In this work, we have studied strange stars in the background of a GMC theory, particularly, the one described by the gravitational action $f(R, L) = R/2 + L + \sigma RL$. The effects of the theory on strange star macroscopic properties are that the stars become larger and more massive, leading to larger compactness and surface redshift. One important feature of the theory is that it recovers GR for energy densities smaller than 250 MeV/fm^3 not depending on the value of σ .

In addition, the space-time metric outside the stars is described by the exterior Schwarzschild solution, which means that the space-time outside is neither AdS nor de Sitter, as it is in some $f(R)$ models. Also, the GMC theory explored here has, as a property, energy-momentum conservation, which is an advantage compared to various other modified theories of gravity.

In particular, when $\sigma = 20$, the GMC theory can achieve $2.6 M_\odot$, which can describe, e.g., the mass of

the pulsars PSR J2215+5135 and PSR J1614-2230, and of the secondary object in the GW190814 event, if those objects are considered to be strange stars. The theory can also describe strange quark star candidates with feasible physical properties, as detailed in Tables I and III.

When calculating the mass of the object for the GMC theory described here, we took into consideration that the mass is calculated according to (10). This means, in particular, that the gravitational mass is obtained in its conventional way as in GR calculations. This happens basically because, at the surface of the stars, the pressure vanishes and all the new terms coming from the GMC theory also go to zero, which in turn implies that the exterior Schwarzschild solution describes the exterior space-time. So, the interior space-time is smoothly connected to the exterior via junction condition with the Schwarzschild metric. Moreover, the energy-momentum tensor is covariantly conserved in the scenario of the present work, so there is not a spherical vacuum layer where the curvature scalar is non-zero [81].

Acknowledgments

GAC thanks Coordenação de Aperfeiçoamento Pessoal de Nível Superior (CAPES) for financial support under grant PNP/88887.368365/2019-00. FR would like to thank CAPES for financial support. R.V.L. is supported by U.S. Department of Energy (DOE) under grant DE-FG02-08ER41533 and to the LANL Collaborative Research Program by Texas A&M System National Laboratory Office and Los Alamos National Laboratory. MM acknowledge CAPES, CNPq and FAPESP thematic project 13/26258 – 4. PHRSM thanks CAPES for financial support. The research of DD is funded by the C.V. Raman Postdoctoral Fellowship (Reg. No. R(IA)CVR-PDF/2020/222) from the Department of Physics, Indian Institute of Science

-
- [1] S. Perlmutter et al., *Astrophys. J.* **517**, 565 (1999), ISSN 0004-637X.
 - [2] C. L. Bennett et al., *Astrophys. J. Suppl. Ser.* **148**, 97 (2003), ISSN 0067-0049.
 - [3] A. G. Riess et al., *Astron. J.* **116**, 1009 (1998), ISSN 0004-6256.
 - [4] D. N. Spergel et al., *Astrophys. J. Suppl. Ser.* **148**, 175 (2003), ISSN 0067-0049.
 - [5] D. N. Spergel et al., *Astrophys. J. Suppl. Ser.* **170**, 377 (2007), ISSN 0067-0049.
 - [6] P. A. R. Ade and others (BICEP2 Collaboration), *Phys. Rev. Lett.* **112**, 241101 (2014), ISSN 1079-7114.
 - [7] S. Weinberg, *Rev. Mod. Phys.* **61**, 1 (1989), ISSN 1539-0756.
 - [8] S. M. Carroll, *Living Rev. Relativ.* **4**, 1 (2001), ISSN 1433-8351.
 - [9] L. Amendola, R. Gannouji, D. Polarski, and S. Tsujikawa, *Phys. Rev. D* **75**, 083504 (2007), ISSN 2470-0029.
 - [10] J.-c. Hwang and H. Noh, *Phys. Lett. B* **506**, 13 (2001), ISSN 0370-2693.
 - [11] T. Harko and F. S. N. Lobo, *Eur. Phys. J. C* **70**, 373 (2010), ISSN 1434-6052.
 - [12] J. Wang, R. Gui, and W. Qiu, *Physics of the Dark Universe* **19**, 60 (2018).
 - [13] S. Nesseris, *Phys. Rev. D* **79**, 044015 (2009), ISSN 2470-0029.
 - [14] T. Delsate and J. Steinhoff, *Phys. Rev. Lett.* **109**, 021101 (2012), ISSN 1079-7114.
 - [15] R. P. L. Azevedo and J. Páramos, *Phys. Rev. D* **94**, 064036 (2016), ISSN 2470-0029.
 - [16] P. P. Avelino and R. P. L. Azevedo, *Phys. Rev. D* **97**, 064018 (2018), ISSN 2470-0029.
 - [17] J. Wang and K. Liao, *Classical Quantum Gravity* **29**, 215016 (2012), ISSN 0264-9381.
 - [18] Y.-B. Wu, Y.-Y. Zhao, Y.-Y. Jin, L.-L. Lin, J.-B. Lu, and X. Zhang, *Mod. Phys. Lett. A* **29** (2014).

- [19] T. Harko, F. S. N. Lobo, J. P. Mimoso, and D. Pavón, *Eur. Phys. J. C* **75**, 386 (2015), ISSN 1434-6052.
- [20] T. Harko, *Phys. Rev. D* **90**, 044067 (2014), ISSN 2470-0029.
- [21] B. Pourhassan and P. Rudra, *Phys. Rev. D* **101**, 084057 (2020), ISSN 2470-0029.
- [22] P. P. Avelino and L. Sousa, *Phys. Rev. D* **97**, 064019 (2018), ISSN 2470-0029.
- [23] T. Harko, F. S. N. Lobo, S. Nojiri, and S. D. Odintsov, *Phys. Rev. D* **84**, 024020 (2011), ISSN 2470-0029.
- [24] T. Harko, F. S. N. Lobo, and O. Minazzoli, *Phys. Rev. D* **87**, 047501 (2013), ISSN 2470-0029.
- [25] T. Harko, *Phys. Lett. B* **669**, 376 (2008), ISSN 0370-2693.
- [26] P. H. R. S. Moraes and P. K. Sahoo, *Eur. Phys. J. C* **77**, 480 (2017), ISSN 1434-6052.
- [27] T. Harko, *Phys. Rev. D* **81**, 044021 (2010), ISSN 2470-0029.
- [28] E. J. Copeland, M. Sami, and S. Tsujikawa, *Int. J. Mod. Phys. D* **15**, 1753 (2006), ISSN 0218-2718.
- [29] J. A. Frieman, M. S. Turner, and D. Huterer, *Annu. Rev. Astron. Astrophys.* **46**, 385 (2008), ISSN 0066-4146.
- [30] J. F. Navarro, C. S. Frenk, and S. D. M. White, *Astrophys. J.* **462**, 563 (1996), astro-ph/9508025.
- [31] B. Moore, S. Ghigna, F. Governato, G. Lake, T. Quinn, J. Stadel, and P. Tozzi, *Astrophys. J. Lett.* **524**, L19 (1999), astro-ph/9907411.
- [32] G. Bertone, D. Hooper, and J. Silk, *Phys. Rep.* **405**, 279 (2005), ISSN 0370-1573.
- [33] T. Harko and F. S. N. Lobo, *Galaxies* **2**, 410 (2014), ISSN 2075-4434.
- [34] G. A. Carvalho, P. H. R. S. Moraes, S. I. dos Santos, B. S. Gonçalves, and M. Malheiro, *Eur. Phys. J. C* **80**, 483 (2020), ISSN 1434-6052.
- [35] R. V. Lobato, G. A. Carvalho, and C. A. Bertulani, *Eur. Phys. J. C* **81**, 1 (2021), ISSN 1434-6052.
- [36] T. E. Riley, A. L. Watts, S. Bogdanov, P. S. Ray, R. M. Ludlam, S. Guillot, Z. Arzoumanian, C. L. Baker, A. V. Bilous, D. Chakrabarty, et al., *Astrophys. J. Lett.* **887**, L21 (2019), ISSN 2041-8213.
- [37] M. C. Miller, F. K. Lamb, A. J. Dittmann, S. Bogdanov, Z. Arzoumanian, K. C. Gendreau, S. Guillot, W. C. G. Ho, J. M. Lattimer, M. Loewenstein, et al., *Astrophys. J. Lett.* **918**, L28 (2021), ISSN 2041-8205.
- [38] T. E. Riley, A. L. Watts, P. S. Ray, S. Bogdanov, S. Guillot, S. M. Morsink, A. V. Bilous, Z. Arzoumanian, D. Choudhury, J. S. Deneva, et al., *Astrophys. J. Lett.* **918**, L27 (2021), ISSN 2041-8205.
- [39] B. P. Abbott et al., *Phys. Rev. Lett.* **119**, 161101 (2017), ISSN 1079-7114.
- [40] E. Farhi and R. L. Jaffe, *Phys. Rev. D* **30**, 2379 (1984), ISSN 2470-0029.
- [41] C. Alcock and A. Olinto, *Annu. Rev. Nucl. Part. Sci.* **38**, 161 (1988), ISSN 0163-8998.
- [42] A. R. Bodmer, *Phys. Rev. D* **4**, 1601 (1971), ISSN 2470-0029.
- [43] E. Witten, *Phys. Rev. D* **30**, 272 (1984), ISSN 2470-0029.
- [44] M. Malheiro, M. Fiolhais, and A. R. Taurines, *J. Phys. G: Nucl. Part. Phys.* **29**, 1045 (2003), ISSN 0954-3899.
- [45] P. H. R. S. Moraes and O. D. Miranda, *Mon. Not. R. Astron. Soc. Lett.* **445**, L11 (2014), ISSN 1745-3925.
- [46] F. Weber, *Prog. Part. Nucl. Phys.* **54**, 193 (2005), ISSN 0146-6410.
- [47] G. Panotopoulos and Á. Rincón, *Eur. Phys. J. C* **79**, 524 (2019), ISSN 1434-6052.
- [48] E.-P. Zhou, X. Zhou, and A. Li, *Phys. Rev. D* **97**, 083015 (2018), ISSN 2470-0029.
- [49] R. O. Gomes, P. Char, and S. Schramm, *Astrophys. J.* **877**, 139 (2019), ISSN 1538-4357.
- [50] N. M. Garcia and F. S. N. Lobo, *Phys. Rev. D* **82**, 104018 (2010), ISSN 2470-0029.
- [51] N. M. Garcia and F. S. N. Lobo, *Classical Quantum Gravity* **28**, 085018 (2011), ISSN 0264-9381.
- [52] T. Harko and F. S. N. Lobo, *Galaxies* **2**, 410 (2014), ISSN 2075-4434.
- [53] E. Farhi and R. L. Jaffe, *Phys. Rev. D* **30**, 2379 (1984), ISSN 2470-0029.
- [54] P. H. R. S. Moraes, J. D. V. Arbañil, and M. Malheiro, *J. Cosmol. Astropart. Phys.* **2016**, 005 (2016), 1511.06282.
- [55] R. L. Jaffe and F. E. Low, *Phys. Rev. D* **19**, 2105 (1979), ISSN 2470-0029.
- [56] M. Linares, T. Shahbaz, and J. Casares, *Astrophys. J.* **859**, 54 (2018), 1805.08799.
- [57] P. B. Demorest, T. Pennucci, S. M. Ransom, M. S. E. Roberts, and J. W. T. Hessels, *Nature* **467**, 1081 (2010), ISSN 1476-4687.
- [58] M. L. Rawls, J. A. Orosz, J. E. McClintock, M. A. P. Torres, C. D. Bailyn, and M. M. Buxton, *Astrophys. J.* **730**, 25 (2011), ISSN 0004-637X.
- [59] T. Güver, F. Özel, A. Cabrera-Lavers, and P. Wroblewski, *Astrophys. J.* **712**, 964 (2010), ISSN 0004-637X.
- [60] P. C. C. Freire, C. G. Bassa, N. Wex, I. H. Stairs, D. J. Champion, S. M. Ransom, P. Lazarus, V. M. Kaspi, J. W. T. Hessels, M. Kramer, et al., *Mon. Not. R. Astron. Soc.* **412**, 2763 (2011), ISSN 0035-8711.
- [61] T. Güver, P. Wroblewski, L. Camarota, and F. Özel, *Astrophys. J.* **719**, 1807 (2010), ISSN 0004-637X.
- [62] F. Özel, T. Güver, and D. Psaltis, *Astrophys. J.* **693**, 1775 (2009), ISSN 0004-637X.
- [63] P. Elebert, M. T. Reynolds, P. J. Callanan, D. J. Hurley, G. Ramsay, F. Lewis, D. M. Russell, B. Nord, S. R. Kane, D. L. DePoy, et al., *Mon. Not. R. Astron. Soc.* **395**, 884 (2009), ISSN 0035-8711.
- [64] M. K. Abubekurov, E. A. Antokhina, A. M. Cherepashchuk, and V. V. Shimanskii, *Astron. Rep.* **52**, 379 (2008), ISSN 1562-6881.
- [65] R. Abbott, T. D. Abbott, S. Abraham, F. Acernese, K. Ackley, C. Adams, R. X. Adhikari, V. B. Adya, C. Affeldt, M. Agathos, et al., *Astrophys. J. Lett.* **896**, L44 (2020), ISSN 2041-8205.
- [66] E. R. Most, L. R. Weih, L. Rezzolla, and J. Schaffner-Bielich, *Phys. Rev. Lett.* **120**, 261103 (2018), ISSN 1079-7114.
- [67] A. Marino, N. Degenaar, T. Di Salvo, R. Wijnands, L. Burderi, and R. Iaria, *Mon. Not. R. Astron. Soc.* **479**, 3634 (2018), ISSN 0035-8711.
- [68] H. Liu, X. Zhang, and D. Wen, *Phys. Rev. D* **89**, 104043 (2014), ISSN 2470-0029.
- [69] J. D. V. Arbañil and M. Malheiro, *Phys. Rev. D* **92**, 084009 (2015), ISSN 2470-0029.
- [70] G. A. Carvalho, J. D. V. Arbañil, R. M. Marinho, and M. Malheiro, *Eur. Phys. J. C* **78**, 1 (2018), ISSN 1434-6052.
- [71] F. Rocha, G. A. Carvalho, D. Deb, and M. Malheiro, *Phys. Rev. D* **101**, 104008 (2020), URL <https://link.aps.org/doi/10.1103/PhysRevD.101.104008>.
- [72] S. Banerjee, S. K. Ghosh, and S. Raha, *J. Phys. G: Nucl.*

- Part. Phys. **26**, L1 (2000), ISSN 0954-3899.
- [73] K. S. Cheng and T. Harko, Phys. Rev. D **62**, 083001 (2000), ISSN 2470-0029.
- [74] T. Harko and K. S. Cheng, Astron. Astrophys. **385**, 947 (2002), ISSN 0004-6361.
- [75] M. S. Morris, K. S. Thorne, and U. Yurtsever, Phys. Rev. Lett. **61**, 1446 (1988).
- [76] F. Weber, M. Orsaria, H. Rodrigues, and S.-H. Yang, Proc. Int. Astron. Union **8**, 61 (2012), ISSN 1743-9213.
- [77] D. Deb, F. Rahaman, S. Ray, and B. K. Guha, J. Cosmol. Astropart. Phys. **2018**, 044 (2018), ISSN 1475-7516.
- [78] G. A. Carvalho, S. I. Dos Santos, P. H. R. S. Moraes, and M. Malheiro, Int. J. Mod. Phys. D **29**, 2050075 (2020), ISSN 0218-2718.
- [79] M. Sharif and A. Waseem, Eur. Phys. J. C **78**, 1 (2018), ISSN 1434-6052.
- [80] A. Majid and M. Sharif, Universe **6**, 124 (2020), ISSN 2218-1997.
- [81] S. K. Maurya, A. Errehymy, D. Deb, F. Tello-Ortiz, and M. Daoud, Phys. Rev. D **100**, 044014 (2019), ISSN 2470-0029.