

# Causal-net category

Xuexing Lu

School of Mathematical Sciences, Zaozhuang University  
Zaozhuang, China

March 16, 2022

## Abstract

A causal-net is a finite acyclic directed graph. In this paper, we introduce a category, denoted as **Cau**, whose objects are causal-nets and morphisms are functors of path categories of causal-nets. It is called causal-net category and in fact the Kleisli category of the "free category on a causal-net" monad. We study several composition-closed classes of morphisms in **Cau**, which characterize interesting causal-net relations, such as coarse-graining, contraction, immersion-minor, topological minor, etc., and prove several useful decomposition theorems. In addition, we show that the notions of a coloring and a minor can be understood as a special kind of minimal-quotient and sub-quotient in **Cau**, respectively. Base on these results, we conclude that **Cau** is a natural setting for studying causal-nets, and the theory of **Cau** should shed new light on the category-theoretic understanding of graph theory.


## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Quotient</b>	<b>2</b>
2.1	Coarse-graining and coloring . . . . .	3
2.2	Vertex and edge coarse-graining . . . . .	4
2.3	Merging and contraction . . . . .	5
<b>3</b>	<b>Inclusion</b>	<b>7</b>
3.1	Immersion and strong-immersion . . . . .	8
3.2	Weak and topological embedding . . . . .	9
3.3	Embedding as sub-causal-net . . . . .	11
<b>4</b>	<b>Minor as sub-quotient</b>	<b>12</b>

## 1 Introduction

A **causal-net** is a finite acyclic directed graph. For each causal-net  $G$ , there is a **path category**  $\mathbf{P}(G)$  with vertices of  $G$  as objects, directed paths in  $G$  as morphisms, directed paths of length zero as identity morphisms and juxtaposition of directed paths as composition. We define a **morphism** between two causal-nets  $G_1$  and  $G_2$  to be a functor between their path categories, that is,  $Hom(G_1, G_2) \doteq Fun(\mathbf{P}(G_1), \mathbf{P}(G_2))$ . All causal-nets and their morphisms form a category, called **causal-net category**, and denoted as **Cau**. It is a full sub-category of the category **Cat** of small categories. The construction of path category is same as that of free category for general directed graphs (II.7, [8]) and **Cau** is in fact the Kleisli category of the "free category on a causal-net" monad (VI.5, [8]).

**Example 1.** The following figure shows an example of causal-net. The objects of its path category are  $v_1, v_2, v_3$ , and morphisms are listed as follows:  $Hom(v_1, v_1) = \{Id_{v_1}\}$ ,  $Hom(v_2, v_2) = \{Id_{v_2}\}$ ,  $Hom(v_3, v_3) = \{Id_{v_3}\}$ ,  $Hom(v_1, v_2) = \{e_1, e_2\}$ ,  $Hom(v_2, v_3) = \{e_3\}$ ,  $Hom(v_1, v_3) = \{e_3e_1, e_3e_2, e_4\}$ ,  $Hom(v_2, v_1) = Hom(v_3, v_2) = Hom(v_3, v_1) = \emptyset$ .



There are two main applications of **Cau**. One is as a basic language for a category-theoretic formulation of Baez's spin network construction for quantum gauge theories [2]. The details of this application will be shown elsewhere. In this paper, we will focus on the other application, that is, to demonstrate a new idea of bringing together graph theory and category theory, which is summarized as the slogan:

$$\text{Graph Theory} = \text{Kleisli Category.}$$

Concretely, we will show that some basic graph-theoretic operations on causal-nets can be naturally represented by morphisms of causal-nets (functors between path categories), and some common relations of causal-nets, such as coarse-graining, contraction, immersion-minor, topological minor, etc., can be characterized by composition-closed classes of morphisms. In addition, we interpret the notions of a coloring and a minor as a special kind of minimal-quotient and sub-quotient, respectively. We believe that these facts are sufficient to justify the fundamental role of **Cau** in the theory of causal-nets and give a new category-theoretic understanding of graph theory.

Evidently, the ideas in this paper can be directly applied to general directed graphs. As to undirected graphs, the ideas may be considered in the framework of dagger categories [7], and the general philosophy should be stated as follows.

$$\text{Ordinary Graph Theory} = \text{Dagger Kleisli Category}$$

This paper is organized as follows. In Section 2, we introduce several special quotient morphisms, including coarse-graining, coloring, vertex-coarse-graining, edge-coarse-graining, merging and contraction, explain their graph-theoretic means and show their relations by several decomposition theorems. We also interpret the notion of a coloring of a causal-net as a special kind of minimal-quotient, that is, as a minimal fusion. In Section 3, we introduce several special but natural inclusion morphisms, such as immersion, topological embedding, subdivision and embedding, etc., explain their graph-theoretic means and show several decomposition theorems. In Section 4, we interpret minor relation as a special kind of sub-quotient.

## 2 Quotient

In this section, we study various type of quotients, which are essential for understanding graphical calculi for kinds of monoidal categories and naturally appears in many physical theories. In our categorical setting, quotient has two different means: one is category-theoretical and the other is graph-theoretical. The later mean turns out to be a special case of the former one.

We start with the most general type, that is, the purely category-theoretic quotient.

**Definition 2.** A morphism of causal-nets is called a **quotient** if it is surjective on objects and full on morphisms.

For a graph-theoretic understanding of the category-theoretic quotient, we introduce some terminologies. Given a morphism  $\lambda : G_1 \rightarrow G_2$  of causal-nets. Edges of  $G_2$  are classified into three classes: (1)  $e \in E(G_2)$  is called a **null-edge** of  $\lambda$  if the preimage  $\lambda^{-1}(e)$  is an empty-set; (2)  $e \in E(G_2)$  is called a **simple-edge** of  $\lambda$  if  $\lambda^{-1}(e)$  has exactly one element; (3)  $e \in E(G_2)$  is called a **multiple-edge** of  $\lambda$  if  $\lambda^{-1}(e)$  has at least two elements.

Similarly, vertices of  $G_2$  are classified into three classes: (1)  $v \in V(G_2)$  is called a **null-vertex** of  $\lambda$  if the preimage  $\lambda^{-1}(v)$  is an empty-set; (2)  $v \in V(G_2)$  is called a **simple-vertex** of  $\lambda$  if  $\lambda^{-1}(v)$  has exactly one element; (3)  $v \in V(G_2)$  is called a **multiple-vertex** of  $\lambda$  if  $\lambda^{-1}(v)$  has at least two elements.

Since a morphism is just a functor between free categories, it is easy to see that a morphism is a quotient if and only if it has no null-vertex and no null-edge.

## 2.1 Coarse-graining and coloring


In this subsection, we introduce the notion of a coarse-graining, which characterizes the graph-theoretic notion of a quotient causal-net and naturally appears in the monadic description of a monoidal category [5, 4]. We also introduce the notion of a coloring of a causal-net, which is characterized as a special kind of minimal-quotient (as a minimal-fusion).

As before, given a morphism  $\lambda : G_1 \rightarrow G_2$ , edges of  $G_1$  are classified into three classes: (1)  $e \in E(G_1)$  is called a **contraction** of  $\lambda$  if  $l(\lambda(e)) = 0$ , that is, the length  $l(\lambda(e))$  of the directed path  $\lambda(e)$  is 0, which means that  $\lambda(e)$  is an identity morphism; (2)  $e \in E(G_1)$  is called a **segment** of  $\lambda$  if  $l(\lambda(e)) = 1$ , that is,  $\lambda(e)$  is an edge; (3)  $e \in E(G_1)$  is called a **subdivision** of  $\lambda$  if  $l(\lambda(e)) \geq 2$ , that is,  $\lambda(e)$  is a directed path of length no less than two.

**Definition 3.** A quotient is called a **coarse-graining** if it has no subdivision.

In other words, a coarse-graining is a morphism with no null-vertex, no null-edge and no subdivision. Clearly, both quotients and coarse-grainings are closed under composition.

**Example 4.** In the following figure,  $\lambda_1, \lambda_2$  show two examples of coarse-grainings of causal nets, where  $\lambda_1(v_1) = \lambda_1(v_2) = w_1$ ,  $\lambda_1(v_3) = w_2$ ,  $\lambda_1(e_1) = \lambda_1(e_2) = Id_{w_1}$ ,  $\lambda_1(e_3) = \lambda_1(e_4) = h$ ; and  $\lambda_2(v_1) = w_1$ ,  $\lambda_2(v_2) = \lambda_2(v_3) = w_2$ ,  $\lambda_2(e_1) = \lambda_2(e_2) = \lambda_2(e_4) = h$ ,  $\lambda_2(e_3) = Id_{w_2}$ .



The following theorem shows a simple graph-theoretic fact.

**Theorem 5.** Let  $H$  and  $G$  be two causal-nets.  $H$  is a quotient of  $G$  if and only if there is a coarse-graining  $\lambda : G \rightarrow H$ .

*Proof.* The proof is easy. Only things needed to know are that

$$V(H) = \frac{V(G)}{\sim_\lambda}, \quad E(H) = \frac{Seg(\lambda)}{\sim_\lambda},$$

where  $Seg(\lambda) \subseteq E(G_1)$  denotes the set of segments of  $\lambda$ , and  $\sim_f$  denotes the equivalence relation on a set  $X$  defined by a map  $f : X \rightarrow Y$  of sets in the way that  $x_1 \sim_f x_2 \iff f(x_1) = f(x_2)$ ,  $x_1, x_2 \in X$ .  $\square$

The following more restricted notion is useful to define the notion of a coloring of a causal-net.

**Definition 6.** A coarse-graining is called a **fusion** if it has no contraction.

We say causal-net  $H$  is a fusion of  $G$ , if there is a fusion  $\lambda : G \rightarrow H$ . Clearly, fusions are closed under composition and the fusion relation defines a partial order among causal-nets, which provides a natural background for understanding colorings of causal-nets.

**Definition 7.** A causal-net is called **complete** if it is minimal under the fusion relation, that is, all its fusions are isomorphisms.

The notion of a complete causal-net is an analog of that of a complete graph in ordinary graph theory. The relation of fusions with colorings is demonstrated as follows.

**Definition 8.** A fusion  $\lambda : G \rightarrow K$  is called a  **$K$ -coloring** of  $G$ , or simply, a **coloring** of  $G$ , if  $K$  is a complete causal-net.

Clearly, a coloring is just a minimal fusion and the relation of colorings of a causal-net with complete causal-nets is similar as the relation of vertex-colorings of a graph with complete graphs in ordinary graph theory. These notions will be further justified in Subsection 2.3.

**Remark 9.** The notion of a complete causal-net is totally parallel to that of a complete graph in ordinary graph. In prepare this paper, we do not know any non-trivial example.


## 2.2 Vertex and edge coarse-graining

In this subsection, we introduce two important sub-classes of coarse-grainings, both of which are closed under composition and naturally appear in some physical theories, such as loop quantum gravity [1], tensor network renormalization [3], etc.

**Definition 10.** A coarse-graining is called a **vertex-coarse-graining** if it has no multiple-edge.

Clearly, a vertex-coarse-graining can be characterized as a morphism with the property that its simple-edges and segments are in bijective with each other.


**Example 11.** The following vertex-coarse-graining is represented by the morphism  $\lambda$  with  $\lambda(v_1) = \lambda(v_2) = \lambda(v_3) = \lambda(v_4) = w$ ,  $\lambda(e_1) = \lambda(e_2) = \lambda(e_3) = Id_w$ .



**Definition 12.** A coarse-graining is called an **edge-coarse-graining** if it has no multiple-vertex and no contraction.

An edge-coarse-graining can be characterized as a morphism with only simple-vertex and without null-edge, contraction and subdivision.

**Example 13.** The following edge-coarse-graining is represented by the morphism  $\lambda$  with  $\lambda(e_1) = \lambda(e_2) = h_1$ ,  $\lambda(e_3) = \lambda(e_4) = h_2$ .



The following decomposition theorem is just a simple graph-theoretic fact and is useful for the description of the monad of the graphical calculus for monoidal categories [6, 4, 5].

**Theorem 14.** *Any coarse-graining  $\lambda$  can be uniquely represented as a composition of a vertex-coarse-graining  $\lambda_\nu$  and an edge-coarse-graining  $\lambda_\varepsilon$ .*

$$\begin{array}{ccc}
 G_1 & \xrightarrow[\lambda_\nu]{\lambda} & G_2 \\
 & \searrow \text{vertex-coarse-graining} & \nearrow \text{edge-coarse-graining} \\
 & & G_\nu
 \end{array}$$

$\lambda$  coarse-graining  
 $\lambda_\nu$  vertex-coarse-graining  
 $\lambda_\varepsilon$  edge-coarse-graining

*Proof.* The proof is easy. We only point out that

$$V(G_\nu) = \frac{V(G)}{\sim_\lambda}, \quad E(G_\nu) = \text{Seg}(\lambda),$$

with notations as before. □

**Remark 15.** *To abuse of terminologies, we are free to say that  $H$  is a coarse-graining, fusion, vertex-coarse-graining, edge-coarse-graining, or any other types of quotients of  $G$ , in case that there is a corresponding quotient morphism from  $G$  to  $H$ , respectively.*

### 2.3 Merging and contraction

In this subsection, we introduce several special classes of vertex-coarse-grainings, which are characterized by their fibers.

The following notion is parallel to the notion of a vertex-identification in ordinary graph theory.

**Definition 16.** *A vertex-coarse-graining is called a **merging** if it has no contraction.*

We say causal-net  $H$  is a merging of  $G$  if there is a merging  $\lambda : G \rightarrow H$ . Clearly, mergings are closed under composition.

Given a morphism  $\lambda : G_1 \rightarrow G_2$  and a vertex  $v$  of  $G_2$ , the **fiber**  $\mathcal{F}(v)$  of  $\lambda$  at  $v$  is defined as the sub-causal-net of  $G_1$  with  $\lambda^{-1}(v)$  as vertex set and  $\lambda^{-1}(Id_v)$  as edge set. If  $\lambda$  is a merging, then for any  $v \in V(G_2)$ ,  $\mathcal{F}(v)$  is discrete, i.e., a sub-causal-net with no edge, whose graph-theoretic mean is that  $\lambda^{-1}(v)$  is an independent subset of vertices of  $G_1$ .

Clearly, a merging can be characterized as a vertex-coarse-graining with all fibers discrete. It is well-known that in ordinary graph theory, a vertex-coloring of a graph corresponds to a partition of its vertex set into independent subsets. Similarly, a merging of a causal-net corresponds to a partition of its vertex set into independent subsets, and a merging can be viewed as a vertex-coloring of a causal-net with the additional property that: all directed edges between vertices of two different colors are in the same direction, more precisely, if vertices  $v$  and  $v'$  have the same color, and vertices  $w$  and  $w'$  have the same color, then the two facts that there is a directed edge from  $v$  to  $w$  and there is a directed edge from  $w'$  to  $v'$  can not hold at the same time.

To completely justify the notion of a complete causal-net and a coloring, we need the following result, which directly follows from Theorem 14.

**Corollary 17.** *Any fusion  $\lambda$  can be uniquely represented as a composition of a merging  $\lambda_m$  and an edge-coarse-graining  $\lambda_\varepsilon$ .*

$$\begin{array}{ccc}
 G & \xrightarrow[\lambda_m]{\lambda} & H \\
 & \searrow \text{merging} & \nearrow \text{edge-coarse-graining} \\
 & & G_m
 \end{array}$$

$\lambda$  fusion  
 $\lambda_m$  merging  
 $\lambda_\varepsilon$  edge-coarse-graining

Just as the former case of fusion relation, the merging relation and edge-coarse-graining relation defines two partial orders among causal-nets, respectively. The above corollary, then, provides us a good reason to view the partial order defined by the fusion relation as a kind of product of these two partial orders, and leads us to a conclusion that a fusion  $G$  is minimal if and only if  $G_m$  is minimal under merging and  $H$  is minimal under edge-coarse-graining.

It is not difficult to verify the following results: (1) a causal-net is simple (without a multi-edge) if and only if it is minimal under edge-coarse-graining; (2) a merging  $\lambda : G \rightarrow K$  defines a minimal vertex-coloring of  $G$  if and only if  $K$  is minimal under merging. The following theorem, following from these two results and Corollary 17, justifies Definition 7 and Definition 8.


**Theorem 18.** *A causal-net is complete if and only if it is simple and minimal under merging.*

Unlike mergings with all fibers discrete, in the opposite direction, we introduce the following notion.

**Definition 19.** *A vertex-coarse-graining is called a **contraction** if all its fibers are connected.*

Both mergings and contractions are closed under composition. The following theorem is a direct consequence of these definitions.

**Theorem 20.** *Any vertex-coarse-graining  $\lambda$  can be uniquely represented as a composition of a merging  $\lambda_m$  and a contraction  $\lambda_{con}$ .*




A causal-net is called a **causal-tree** or a **poly-tree**, if its underlying graph is a tree. The following generalization may be more useful.

**Definition 21.** *A causal-net is called a **causal-Tree** or a **poly-Tree**, if it has a causal-tree as an edge-coarse-graining.*

Straightforward, a causal-Tree is a causal-tree with multi-edges. The operation of contracting a multi-edge is called a **multi-edge-contraction**, which can be uniquely represented by a morphism with the unique nontrivial fiber being a multi-edge.

**Example 22.** *The following figure shows a causal-Tree and one of its multi-edge-contraction, which is represented by a morphism  $\lambda$ . The multi-edge  $\{e_1, e_2\}$  is contracted, which is represented by the condition  $\lambda(e_1) = \lambda(e_2) = Id_v$ .*



The following proposition gives a characterization of causal-Trees in terms of multi-edge-contractions, whose parallel result in ordinary graph theory is that a graph is a tree if and only if it can be transformed into a vertex by a sequence of edge-contractions.

**Proposition 23.** *A causal-net is a causal-Tree if and only if it can be transformed into a vertex through a sequence of multi-edge-contractions.*

The following notion is useful for understanding minor relation in Section 4.

**Definition 24.** A contraction is called a **tree-contraction** if all its fibers are causal-Trees.

We say that causal-net  $H$  is a tree-contraction of  $G$ , if there is a tree-contraction  $\lambda : G \rightarrow H$ . The following result directly follows from Proposition 23.

**Theorem 25.** Let  $H, G$  be two causal-nets.  $H$  is a tree-contraction of  $G$  if and only if  $H$  can be obtained from  $G$  by a sequence of multi-edge-contractions.

A causal-net is called a **directed multi-path** if it has a directed path as an edge-coarse-graining. Briefly, a directed multi-path is directed path with multi-edges. The following special case of tree-contraction will be used to characterize subdivisions in Section 3.2.

**Definition 26.** A vertex-coarse-graining is called a **path-contraction** if all its fibers are directed multi-paths.


### 3 Inclusion

In this section, we introduce various types of inclusions, whose parallel notions are of broad interest in graph theory. We start with the most general type—the purely category-theoretic inclusion.

**Definition 27.** A morphism of causal-nets is called an **inclusion** if it is injective on objects and fully faithful on morphisms.

An inclusion can be characterized as a morphism with no multiple-vertex, no multiple-edge and no contraction. However, as the following example shown, an inclusion may create **joint-vertices** and **joint-edges**. Clear, the class of inclusions is closed under composition.

**Example 28.** The following figure shows an example of an inclusion with  $\lambda(h_1) = e_3e_1$ ,  $\lambda(h_2) = e_3e_2$ . In this example, the inclusion creates a joint-edge  $e_3$  and a joint-vertex  $v$ .



The following theorem is a simple category-theoretic fact.

**Theorem 29.** Any morphism  $\lambda$  can be uniquely represented as a composition of a coarse-graining  $\lambda_c$  and an inclusion  $\lambda_{in}$ .

$$\begin{array}{ccc}
 G_1 & \xrightarrow[\text{coarse-graining}]{\lambda_c} & G_c \\
 & \xrightarrow[\text{inclusion}]{\lambda_{in}} & \\
 & \xrightarrow[\text{morphism}]{\lambda} & G_2
 \end{array}$$

*Proof.* The proof is easy, we leave it to readers. Only things we need to notice are that coarse-grainings have no subdivision, inclusions have no contraction, and that

$$V(G_c) = \frac{V(G_1)}{\sim_\lambda}, \quad E(G_c) = \frac{Seg(\lambda) \sqcup Subd(\lambda)}{\sim_\lambda},$$

where  $Subd(\lambda) \subseteq E(G_1)$  denotes the set of subdivisions of  $\lambda$  and the other notations are as before.  $\square$

### 3.1 Immersion and strong-immersion


In this subsection, we introduce two types of inclusions whose parallel notions in ordinary graph theory characterize those of an immersion-minor and a strong-immersion-minor, respectively.

Given a morphism  $\lambda : G_1 \rightarrow G_2$ , two subdivisions  $e_1, e_2$  of  $\lambda$  are called **edge-disjoint** if  $\lambda(e_1)$  and  $\lambda(e_2)$  has no common edge. We call  $\lambda$  **edge-disjoint** if any two subdivisions of  $\lambda$  are edge-disjoint.

**Definition 30.** An inclusion is called an *immersion* if it is edge-disjoint.

In other words, an immersion is an inclusion that does not create any joint-edges. But as shown in the following example, it may create joint-vertices.


**Example 31.** The following  $\lambda$  is an example of an immersion with  $\lambda(h_1) = e_3e_1, \lambda(h_2) = e_4e_2, \lambda(h) = e, \lambda(w) = v$ , and  $v$  is the unique joint-vertex created by  $\lambda$ .



A causal-net  $H$  is called an **immersion-minor** of  $G$ , if  $H$  can be obtained from  $G$  by a sequence of operations of deleting an isolated vertex, deleting an edge and an edge-lift (replacing a length two path  $e_2e_1$  by a length one path  $h$ , with the middle vertex of  $e_2e_1$  retained).

As the following example shown, an edge-lift can be view as a composition of a merging after an edge-subdivision, and clearly, an edge-lift does not change the number of vertices.

**Example 32.** The following figure shows an example of an edge-lift, which can be represented by a morphism in the opposite direction, with  $\lambda(h) = e_2e_1, \lambda(w) = v$ .



It is not difficult to verify that immersions are closed under composition. The following theorem is not difficult to verify.

**Theorem 33.** Let  $H, G$  be two causal-nets.  $H$  is an immersion-minor of  $G$  if and only if there is an immersion  $\lambda : H \rightarrow G$ .




A morphism  $\lambda$  is called **strong** if for any subdivision  $e$  of  $\lambda$ , all internal vertices of  $\lambda(e)$  are null-vertices of  $\lambda$ . In other words, an immersion is strong if all its subdividing-vertices are null-vertices. It is not difficult to see that strong immersions are closed under composition.

The following notion is parallel to that of a strong-immersion-minor in ordinary graph theory.

**Definition 34.** A causal-net  $H$  is a **strong-immersion-minor** of  $G$ , if there is a strong immersion  $\lambda : H \rightarrow G$ .


The immersion in Example 31 is not strong as the subdividing-vertex  $v$  is not a null-vertex (with  $w$  as a preimage). An immersion is called a **weak immersion** when it is not a strong one.

**Example 35.** We show an example of a strong-immersion-minor with  $\lambda(h_1) = e_3e_1$ ,  $\lambda(h_2) = e_4e_2$  and  $\lambda(h) = e$ . The unique subdividing-vertex  $v$  is a null-vertex.




We will end this subsection by listing several corollaries of Theorem 29.


**Corollary 36.** Any edge-disjoint morphism  $\lambda$  can be uniquely represented as a composition of a coarse-graining  $\lambda_c$  and an immersion  $\lambda_{im}$ .



**Corollary 37.** Any strong morphism  $\lambda$  can be uniquely represented as a composition of a coarse-graining  $\lambda_c$  and a strong inclusion  $\lambda_{str-in}$ .



**Corollary 38.** Any strong edge-disjoint morphism  $\lambda$  can be uniquely represented as a composition of a coarse-graining  $\lambda_c$  and a strong-immersion  $\lambda_{str-im}$ .



### 3.2 Weak and topological embedding


In this subsection, we introduce more restricted types of inclusions. We need more terminologies. Two subdivisions  $e_1, e_2$  of morphism  $\lambda$  are called **vertex-disjoint** if  $\lambda(e_1)$  and  $\lambda(e_2)$  has no common internal vertex. We call a morphism **vertex-disjoint** if any two of its subdivisions are vertex-disjoint. Vertex-disjoint is a more restricted property than edge-disjoint. Both Example 31 and 35 are not vertex-disjoint because  $\lambda(h_1)$  and  $\lambda(h_2)$  have a common internal vertex  $v$ .

The following notion is a special type of an immersion, which may be useful somewhere.

**Definition 39.** An inclusion is called a **weak-embedding** if it is vertex-disjoint.

As the following example shown, a weak-embedding may create joint-vertices, but none of these joint-vertices is a null-vertex.

**Example 40.** The following figure shows an example of a weak-embedding, with  $\lambda(h_1) = e_1$ ,  $\lambda(h_2) = e_3e_2$ . In this example, the weak-embedding  $\lambda$  creates a joint-vertex  $v$ , which is not a null-vertex, hence it is not a strong-immersion.




A single edge-lift, as shown in Example 32, can be reversely represented by a weak-embedding. As mentioned above, both Example 31 and 35 are not weak-embeddings, but they are immersions, which, by Theorem 33, implies that a composition of a sequence of edge-lifts in general may not be reversely represented by a weak-embedding. So weak-embeddings are not closed under composition.

The following notion is more useful.

**Definition 41.** An inclusion is called a **topological embedding** if it is strong and vertex-disjoint.

A causal-net  $H$  is called a **topological minor** of  $G$ , if  $G$  can be obtained from  $H$  by a sequence of operations of adding an isolated vertex, adding an edge, and subdividing an edge (inserting a middle vertex into an edge).

**Example 42.** The following morphisms  $\lambda_1$  and  $\lambda_2$  represent the topological minor relation of  $H$  and  $G$ , where  $\lambda_1(h_1) = e_2$ ,  $\lambda_1(h_3) = e_4e_3$  and  $\lambda_2(h_1) = e_3e_2$ ,  $\lambda_2(h_3) = e_4$ .



The graph-theoretic notion of a topological minor coincides with the category-theoretic notion of a topological embedding. The following theorem can be easily proved by induction on number of subdivisions of morphisms.

**Theorem 43.** Let  $H, G$  be two causal-nets.  $H$  is a topological-minor of  $G$  if and only if there is a topological embedding  $\lambda : H \rightarrow G$ .


Let  $\lambda : H \rightarrow G$  be a morphism,  $v \in V(G)$  and  $h \in E(H)$ , we say that  $h$  is a **cover** of  $v$  or  $v$  is **covered** by  $h$ , if  $v$  is a vertex of  $\lambda(h)$ .

**Definition 44.** A topological embedding  $\lambda : H \rightarrow G$  is called a **subdivision** if each non-isolated vertex of  $G$  has a cover and all isolated vertices of  $G$  are simple.

The following theorem characterizes subdivisions in terms of path-contractions.

**Theorem 45.** A morphism  $\lambda : H \rightarrow G$  is a subdivision if and only if it has a path-contraction  $\chi : G \rightarrow H$  as left inverse

**Example 46.** A subdivision  $\lambda$  may have different left inverses  $\chi_1, \chi_2$ , where  $\lambda(v_1) = w_1, \lambda(v_2) = w_3; \chi_1(w_1) = \chi_1(w_2) = v_1, \chi_1(w_3) = v_2; \chi_2(w_1) = v_1, \chi_2(w_2) = \chi_2(w_3) = v_2$ .



We mention that the inverse operation of an edge-subdivision is called a **smoothing**, as shown above, it can be represented by a path-contraction.

We say causal-net  $G$  is a **subdivision** of  $H$  if  $G$  can be obtained by subdividing a subset of edges of  $H$ . The following theorem gives a characterization of subdivision relation, which can be easily proved by induction on number of subdivisions of morphisms.

**Theorem 47.** Let  $H, G$  be two causal-nets.  $G$  is a subdivision of  $H$  if and only if there is a subdivision  $\lambda : H \rightarrow G$ .

It is easy to verify that topological embeddings and subdivisions are closed under composition.

### 3.3 Embedding as sub-causal-net

In this subsection, we consider the simplest type of inclusion.

**Definition 48.** A morphism is called an **embedding** if it is an inclusion with no subdivision.


It is straightforward to verify the following result.

**Proposition 49.** Let  $H, G$  be two causal-nets. Then the following conditions are equivalent:


- (1)  $H$  is a sub-causal-net of  $G$ ;
- (2) there is an embedding  $\lambda : H \rightarrow G$ ;
- (3)  $H$  can be obtained from  $G$  by a sequence of operations of deleting an edge, deleting an isolated vertex;
- (4)  $G$  can be obtained from  $H$  by a sequence of operations of adding an edge, adding an isolated vertex.

The following theorem is of particular importance for a category-theoretic understanding of Baez's spin network construction [2].


**Theorem 50.** Any morphism  $\lambda : G_1 \rightarrow G_2$  can be uniquely represented as a composition of a subdivision  $\lambda_s$ , a coarse-graining  $\lambda_c$  and an embedding  $\lambda_e$ .



*Proof.* The subdivision  $G_s$  is obtained from  $G$  by applying the operation that replacing each subdivision  $e \in \text{Subd}(\lambda)$  by a copy of its image  $\lambda(e)$ , which can be represented by the subdivision  $\lambda_s : G_1 \rightarrow G_s$  according to Theorem 47. Then  $\lambda$  can be naturally represented as a composition of  $\lambda_s$  and a morphism  $\lambda' : G_s \rightarrow G_2$  without subdivision, that is, we have the following commutative diagram




By Theorem 29,  $\lambda'$  can be decomposed as a composition of a coarse-graining  $\lambda_c : G_s \rightarrow G_c$  and an inclusion  $\lambda_l : G_c \rightarrow G_2$ . Since  $\lambda'$  has no subdivision,  $\lambda_l$  has no subdivision, which means that  $\lambda_l$  is an embedding. That is, we get the following diagram




Combining the above two commutative diagrams, we complete the proof.  $\square$

Without giving proofs, we end this subsection by listing several results, which may be helpful in understanding the combinatorial structure of **Cau**.


**Corollary 51.** *Any inclusion  $\lambda : G_1 \rightarrow G_2$  can be unique represented as a composition of a subdivision  $\lambda_s$ , a fusion  $\lambda_{fu}$  and an embedding  $\lambda_l$ .*



**Corollary 52.** *Any immersion  $\lambda : G_1 \rightarrow G_2$  can be unique represented as a composition of a subdivision  $\lambda_s$ , a merging  $\lambda_m$  and an embedding  $\lambda_l$ .*



**Corollary 53.** *Any topological embedding  $\lambda : G_1 \rightarrow G_2$  can be unique represented as a composition of a subdivision  $\lambda_s$  and an embedding  $\lambda_l$ .*




## 4 Minor as sub-quotient

In this section, we show a category-theoretic understanding of minor relation. Since in **Cau** multi-edges are allowed, we had better define a minor as follows.

**Definition 54.** *Let  $H, G$  be two causal-nets.  $H$  is called a **minor** of  $G$ , if  $H$  can be obtained from  $G$  by a sequence of operations of deleting an edge, deleting an isolated vertex, and contracting a multi-edge.*

Evidently, as usual, the minor relation is a partial order among causal-nets. But unlike the cases of topological minor and immersion-minor, minor relation in general can not be represented by morphisms of causal-nets, the reason is that the morphisms representing deleting and contraction are of different directions. However, minor relation can be viewed as a special kind of sub-quotient.


**Example 55.** The following figure shows that  $H$  is a minor of  $G$ .



The following result is a direct consequence of Theorem 25 and Proposition 49, which characterizes the minor relation as sub-tree-contraction.

**Theorem 56.** Let  $H, G$  be two causal-nets. Then the fact that  $H$  is a minor of  $G$  is equivalent to the following conditions:

- (1)  $H$  is a tree-contraction of a sub-causal-net  $K$  of  $G$ , that is, there is a causal-net  $K$  with an embedding  $\lambda_\iota : K \rightarrow G$  and a tree-contraction  $\lambda_{tr} : K \rightarrow H$ ;
- (2)  $H$  is a sub-causal-net of a tree-contraction  $L$  of  $G$ , that is, there is a causal-net  $L$  with an embedding  $\rho_\iota : H \rightarrow L$  and a tree-contraction  $\rho_{tr} : G \rightarrow L$ ;



Using the same paradigm, we can easily generalize the notion of a minor. Now we fix an arbitrary non-empty set  $\Sigma$  of connected causal-nets, and call an element of  $\Sigma$  a  $\Sigma$ -causal-net.

**Definition 57.** Let  $H$  and  $G$  be two causal-nets.  $H$  is called a  $\Sigma$ -minor of  $G$ , if  $H$  can be obtained from  $G$  by a sequence of operations of deleting an edge, deleting an isolated vertex and contracting a  $\Sigma$ -causal-net.

Clearly, when  $\Sigma$  is the set of multi-edges, a  $\Sigma$ -minor is just a minor. To characterize  $\Sigma$ -minors as sub-quotients, we introduce some notions. The first one is a parallel generalization of that of a causal-Tree (Definition 21).

**Definition 58.** A causal-net is called a  $\Sigma$ -tree if it has a causal-tree as a coarse-graining such that for each edge of the causal-tree, its pre-image is a  $\Sigma$ -causal-net.

The operation of contracting a  $\Sigma$ -causal-net is called a  $\Sigma$ -**contraction**, which can be uniquely represented by a morphism with the unique non-trivial fiber being a  $\Sigma$ -causal-net. The following proposition characterizes  $\Sigma$ -trees in terms of  $\Sigma$ -contractions, which can be view as a proper generalization of Proposition 23.

**Proposition 59.** *A causal-net is a  $\Sigma$ -tree if and only if it can be transformed into a vertex through a sequence of  $\Sigma$ -contractions.*

The notion of a tree-contraction is then generalized as follows.

**Definition 60.** *A contraction is called a  $\Sigma$ -tree-contraction if all its fibers are  $\Sigma$ -trees.*


As a generalization of Theorem 25, the following result implies that  $\Sigma$ -tree-contractions are closed under composition.

**Theorem 61.** *Let  $H, G$  be two causal-nets.  $H$  is a  $\Sigma$ -tree-contraction of  $G$  if and only if  $H$  can be obtained from  $G$  by a sequence of  $\Sigma$ -contractions.*

Now we can state the  $\Sigma$ -version of Theorem 62 as follows.

**Theorem 62.** *Let  $H, G$  be two causal-nets. Then the fact that  $H$  is a  $\Sigma$ -minor of  $G$  is equivalent to the following conditions:*


- (1)  $H$  is a  $\Sigma$ -tree-contraction of a sub-causal-net  $K$  of  $G$ , that is, there is a causal-net  $K$  with an embedding  $\lambda_\iota : K \rightarrow G$  and a  $\Sigma$ -tree-contraction  $\lambda_{\Sigma-tr} : K \rightarrow H$ ;
- (2)  $H$  is a sub-causal-net of a  $\Sigma$ -tree-contraction  $L$  of  $G$ , that is, there is a causal-net  $L$  with an embedding  $\rho_\iota : H \rightarrow L$  and a  $\Sigma$ -tree-contraction  $\rho_{\Sigma-tr} : G \rightarrow L$ ;



To see that  $\Sigma$ -minor defines a partial order, the only thing we need to notice is that  $\Sigma$ -minor, same as minor, is of operational type. Clearly we can go further to define a more general type of minor by choosing a set of composition-closed coarse-grainings.

At the end of this section, we mention that just as in ordinary graph theory, a topological minor is also a minor, which follows from the fact that subdivisions have path-contractions as left inverses (Theorem 45).

**Example 63.** *Topological minor relation implies minor relation.*



## References

- [1] S. Ariwahjoedi, J.S. Kosasih, C. Rovelli and F.P. Zen. How many quanta are there in a quantum spacetime? *Classical and Quantum Gravity*, 32(16), 2015.
- [2] J.C. Baez. Spin networks in gauge theory. *Advances in Mathematics*, 117:253-272, 1996.
- [3] G. Evenbly and G. Vidal. Tensor Network Renormalization. *Physical Review Letters*, 115, 2015.
- [4] S. Hu, X. Lu and Y. Ye. A graphical calculus for semi-groupal categories. *Applied Categorical Structures*, 27(2):163–197, April 2019, arXiv:1604.07276.
- [5] S. Hu, X. Lu and Y. Ye. Combinatorics and algebra of tensor calculus. *PhD Thesis*, 2015, arXiv:1501.01790.
- [6] A. Joyal and R. Street. The geometry of tensor calculus. I. *Advances in Mathematics*, 88(1):55–112, 1991.
- [7] M. Karvonen. The way of the dagger. *PhD Thesis*. 2018. arXiv:1904.10805.
- [8] S. Mac Lane. Categories for the Working Mathematician. *Graduate Texts in Mathematics*. 5 (Second ed.), Springer. 1998. ISBN 0-387-98403-8. Zbl 0906.18001.

**Xuexing Lu**

Email: xxlu@uzz.edu.cn