

---

# MATRIX BISPECTRALITY AND NONCOMMUTATIVE ALGEBRAS: BEYOND THE PROLATE SPHEROIDALS

---

**F. Alberto Grünbaum**  
Department of Mathematics  
University of California, Berkeley, CA  
grunbaum@math.berkeley.edu

**Brian D. Vasquez**  
IMPA, Rio de Janeiro  
bridava927@gmail.com

**Jorge P. Zubelli**  
Department of Mathematics, Khalifa University  
zubelli@gmail.com

DEDICATED TO VAUGHAN JONES, A TRULY INSPIRATIONAL FRIEND, WHO LEFT US SOME GREAT MEMORIES.

February 2, 2022

**Keywords** bispectral algebras · bispectral triple · presentations of finitely generated algebras · time-and-band limiting · integral and commuting differential operators

## 1 Preamble

The bispectral problem discussed in this paper has its origin in Duistermaat and Grünbaum (1986). It is motivated, as mentioned there, by an effort to understand and extend a remarkable phenomenon in Fourier analysis on the real line: the operator of time-and-band limiting is an integral operator admitting a second order differential operator with simple spectrum in its commutator. This property, which gives a good numerical way to compute the eigenfunctions of the integral operator, was put to good use in a series of papers by D. Slepian, H. Landau and H. Pollak at Bells Labs back in the 1960's, see Slepian and Pollak (1961); Landau and Pollak (1962, 1961, 1962); Slepian (1964, 1978, 1983, 1976) and is of interest in one other contribution in this issue, see Connes and Consani (2021); Connes and Marcolli (2008); Connes and Moscovici (2021). We are thankful to Luc Haine, of Louvain-la-Neuve, Belgium who alerted one of us (AG) on Nov 2021 about a talk by Alain Connes in the series Mathematical Picture Language. This talk delivered on Dec 7th. 2021 can be seen in Youtube and covers some of the contents of his work with various collaborators on the zeta function and its relations to the prolate spheroidal functions. For readers interested in this fascinating connection there is no better way of learning about this than watching the lecture. See Connes (2021).

If one tries to extend this property beyond the Fourier case by adding a potential  $V(x)$  to the operator  $-\left(\frac{d}{dx}\right)^2$  and replacing an expansion in exponentials by an expansion in term of the eigenfunctions  $\psi(x, z)$  of  $-\left(\frac{d}{dx}\right)^2 + V(x)$  it appears plausible that a certain property of these eigenfunctions  $\psi(x, z)$  could play a useful role. This property is now known as the bispectral property which will be formulated below. Its solution in the scalar case was the purpose of Duistermaat and Grünbaum (1986).

The version of the bispectral problem that we discuss in this paper is a noncommutative one, obtained by allowing all objects in the original formulation to be matrix valued. The details are given later.

Now for a bit of history: one of us (AG) gave a couple of talks, one at Vanderbilt in Nov 2013 and one in IMPA, Rio de Janeiro in March 2014. The first talk was at the invitation of Vaughan Jones and started by saying that the topic was most likely of no interest to him. At some point in the talk Vaughan said "all of this is about bi-modules and subfactors". The second talk had both Vaughan and another one of us (JZ) in the audience, there was some more mention of these

topics and then in Grünbaum (2014), written in April 2014, a reference is made to a future joint paper with Vaughan "On the bimodule structure of the bispectral problem, in preparation". The occasion of this second talk was a visit that Vaughan (and for part of it with Wendy) did to Argentina, Uruguay and Brazil. In Buenos Aires he delivered one lecture, in Rio he delivered a series of talks but in Montevideo besides giving a lecture he was received in a private audience by President Jose Mujica, described as "the world humblest head of state" by Wikipedia. Vaughan whose command of the spanish language was quite high lamented that he had troubles understanding President Mujica.

The examples in this short paper Grünbaum (2014) were discussed with Vaughan who showed some level of interest but we never managed to get him fully on board. Of course we assumed that we always had time to get him onto this project. The short paper Grünbaum (2014) makes three separate conjectures going with three different examples. These conjectures were proved to be correct in the thesis of the third of us (BVC) under the supervision of (JZ).

And now that sadly we have no chance of benefiting from the insights that Vaughan would have brought in, we present the problem and some results to a wider audience in the hope that someone may surmise what it was that Vaughan had in mind. Getting someone involved in this effort would be a nice way of honoring his memory.

## 2 Commuting integral and differential operators

The bispectral problem, introduced in the next section is motivated by the following very concrete problem in signal communication: a signal of support in the interval  $[-T, T]$  is transmitted over a channel that has bandwidth  $[-W, W]$  i.e., all frequencies in the signal beyond absolute value of  $W$  cannot be sent over. A mathematical formulation is as follows: an arbitrary signal in  $L^2(\mathbb{R})$  is chopped to the interval  $[-T, T]$  and then its Fourier transform is chopped to the interval  $[-W, W]$ . Denoting for simplicity these two chopping operations by  $T$  and  $W$  we are dealing with the operator

$$E = WFT$$

where  $\mathcal{F}$  stands for the Fourier transform. The spectral analysis of this operator, i.e., a look at its singular functions and singular values requires the consideration of the operator  $E^*E$ . It is easy to see that  $E^*E$  is an integral operator acting in  $L^2(-T, T)$  whose kernel is given by

$$\frac{\sin W(t-s)}{t-s}$$

and this bounded operator acts on a function in the space  $L^2(-T, T)$  by

$$(Kf)(s) = \int_{-T}^T \frac{\sin W(t-s)}{t-s} f(t) dt$$

for  $f \in L^2(-T, T)$  and  $s \in (-T, T)$ . This  $K$  commutes with the operator

$$(Lf)(x) = \left(-\frac{d}{dx}\right) (T^2 - x^2) \left(\frac{df}{dx}\right) + W^2 x^2 f(x)$$

defined on  $C^2$  functions.

One can show that this densely defined operator has a unique selfadjoint extension in  $L^2(-T, T)$  with eigenfunctions and eigenvalues that depend on the parameter  $W$ .

Its eigenfunctions are known as the prolate spheroidal wave functions, since this is one of the differential operators resulting in separating variables when solving for the eigenfunctions of the Laplacian on a prolate spheroid.

What other naturally appearing integral operators allow for commuting differential operators? Two other examples are the Bessel and Airy kernels, as in the work of Tracy and Widom, Tracy and Widom (1994a,b) in the context of **Random Matrix Theory**. For the Bessel case see also Slepian (1964); Grünbaum et al. (1982). There are other examples, but the search is nowhere close to finished. The **bispectral property**, to be formulated below was put forward in Duistermaat and Grünbaum (1986) as an important ingredient in the search for more examples of this commuting property.

## 3 The bispectral problem

The problem was posed and solved in Duistermaat and Grünbaum (1986). It is as follows:

Find all nontrivial instances where a function  $\psi(x, z)$  satisfies

$$L \left( x, \frac{d}{dx} \right) \psi(x, z) \equiv \left( - \left( \frac{d}{dx} \right)^2 + V(x) \right) \psi(x, z) = z\psi(x, z)$$

as well as

$$B \left( z, \frac{d}{dz} \right) \psi(x, z) \equiv \left( \sum_{i=0}^M b_i(z) \left( \frac{d}{dz} \right)^i \right) \psi(x, z) = \Theta(x)\psi(x, z).$$

All the functions  $V(x), b_i(k), \Theta(x)$  are, in principle, arbitrary except for smoothness assumptions. Notice that here  $M$  is arbitrary (finite). The operator  $L$  could be of higher order, but in Duistermaat and Grünbaum (1986) attention is restricted to order two.

The complete solution is given as follows:

**Theorem** *If  $M = 2$ , then  $V(x)$  is (except for translation) either  $\frac{c}{x^2}$  or  $ax$ , i.e. we have a Bessel or an Airy case. If  $M > 2$ , there are two families of solutions*

- a)  $L$  is obtained from  $L_0 = - \left( \frac{d}{dx} \right)^2$  by a finite number of Darboux transformations ( $L = AA^* \rightarrow \tilde{L} = A^*A$ ). In this case  $V$  is a rational solution of the Korteweg-deVries hierarchy of equations. Here  $A$  is a first order differential operator.
- b)  $L$  is obtained from  $L_0 = - \left( \frac{d}{dx} \right)^2 + \frac{1}{4x^2}$  after a finite number of (rational) Darboux transformations.

In all cases we have a solution of the ad-conditions, a complicated system of nonlinear differential equations. These conditions are necessary and sufficient.

Notice that the solutions organize themselves into nice manifolds.

The simplest example of case *a*) follows from  $L_0 = - \left( \frac{d}{dx} \right)^2$  by two Darboux transformations, one gets the operator

$$L_2 = - \left( \frac{d}{dx} \right)^2 + \frac{6(x^4 + 12t_3x)}{(x^3 - t_3)^2}.$$

In this case  $\Theta(x) = x^4 - 4t_3x$  and the differential operator in the spectral parameter is

$$B_2 \left( z, \frac{d}{dz} \right) = \left( - \left( \frac{d}{dz} \right)^2 + \frac{6}{z^2} \right)^2 + 4it_3 \left( \frac{d}{dz} \right)$$

The potential in the operator  $L_2 = - \left( \frac{d}{dx} \right)^2 + V(x, t)$  above satisfies the KdV equation.

It was later observed by Magri and Zubelli, see Zubelli and Magri (1991) that in case *b*) we are dealing with rational solutions of the Virasoro equations (i.e. master symmetries of KdV). The bigger picture became more apparent in the work Zubelli and Valerio Silva (2000) where it is shown that the generic rational potentials that decay at infinity and remain rational by all the flows of the master-symmetry KdV hierarchy are bispectral potentials for the Schrödinger operator.

In case *a*) the space of common solutions has dimension one and in case *b*) it has dimension two. One refers to these as the rank one and rank two situations.

Observe that the "trivial cases" when  $M = 2$  are self-dual in the sense that since the eigenfunctions  $\psi(x, z)$  are functions either of the product  $xz$  or of the sum  $x + z$ , one gets  $B$  by replacing  $z$  for  $x$  in  $L$ . The *bispectral involution* introduced in Wilson (1993) shows how this can be adapted in the "higher order cases".

## 4 The non-commutative version of the bispectral problem

A first noncommutative (or matrix) version of the bispectral problem was considered in J. Zubelli Ph.D. thesis at Berkeley, see also Zubelli (1990, 1992)Zubelli (1992) in the situation where both the physical space and spectral operators act on the same side of the eigenfunction and the eigenvalues are both scalar valued. Later on, several other versions were considered. See Sakhnovich and Zubelli (2001); Kasman (2015); Boyallian and Liberati (2008); Grünbaum and Iliev (2003); Grünbaum (2014); Geiger et al. (2017) and references therein. The noncommutative version of the bispectral problem displayed interesting connections with soliton equations as well. Indeed, in Zubelli (1992) it was shown that a large class of rational solutions to the AKNS hierarchy Ablowitz et al. (1974) led to matrix differential operators that displayed the bispectral property. Among the important equations in Mathematical Physics that are covered by the AKNS hierarchy one finds the modified KdV and the nonlinear Schrödinger equation. The matrix differential operator that appeared in this case was in turn related to Dirac operators. The connection between bispectrality and another important topic in Mathematical Physics, namely Huygens' principle in the strict sense Berest and Veselov (1994) turned out to appear also in the context of Matrix Bispectrality. Indeed, in Chalub and Zubelli (2006, 2005, 2004, 2001), it was shown that rational solutions to the AKNS hierarchy led to Dirac operators which satisfy Huygens' principle in the stric sense. In other words, the fundamental solutions of the perturbed Dirac equation in a suitably high space-time dimension had its support precisely on the surface of the light cone and not in its interior. Another interesting connection between Matrix bispectrality to soliton equations of Mathematical Physics was explored in Sakhnovich and Zubelli (2001).

In this present paper we take the **bi-module** structure of the problem into account and let the operators act on different sides as well as allow both eigenvalues to be matrix valued.

We consider triplets  $(L, \psi, B)$  satisfying the equations

$$L\psi(x, z) = \psi(x, z)F(z) \quad (\psi B)(x, z) = \theta(x)\psi(x, z) \quad (1)$$

with  $L = L(x, \frac{d}{dx})$ ,  $B = B(z, \frac{d}{dz})$  linear matrix differential operators, i.e.,  $L\psi = \sum_{i=0}^l a_i(x) \cdot (\frac{d}{dx})^i \psi$ ,  $\psi B = \sum_{j=0}^m (\frac{d}{dz})^j \psi \cdot b_j(z)$ . The functions  $a_i : U \subset \mathbb{C} \rightarrow M_N(\mathbb{C})$ ,  $b_j : V \subset \mathbb{C} \rightarrow M_N(\mathbb{C})$ ,  $F : V \subset \mathbb{C} \rightarrow M_N(\mathbb{C})$ ,  $\theta : U \subset \mathbb{C} \rightarrow M_N(\mathbb{C})$  and the nontrivial common eigenfunction  $\psi : U \times V \subset \mathbb{C}^2 \rightarrow M_N(\mathbb{C})$  are in principle compatible sized meromorphic matrix valued functions defined in suitable open subsets  $U, V \subset \mathbb{C}$ .

A triplet  $(L, \psi, B)$  satisfying (1) is called a bispectral triplet.

The study of the structure of the algebra of possible  $\theta(x)$  going with a fixed bispectral  $\psi(x, z)$  was first raised in Castro and Grünbaum (2006) and analyzed in Tirao (2011); Grünbaum and Tirao (2007). See also Casper (2018) and Zurrián (2017).

We consider now the examples and conjectures given in Grünbaum (2014) as well as their validation and further description given in Vasquez and Zubelli (2021).

For the benefit of the reader we give a few definitions before giving some explicit results in the next section.

**Definition 1.** Let  $\mathbb{K}$  be a field,  $C$  be a  $\mathbb{K}$ -algebra,  $A$  a subring of  $C$  and  $S \subset C$ . We define

$$A \cdot \langle S \rangle = \text{span}_{\mathbb{K}} \left\{ \prod_{j=1}^n s_j \mid s_1, \dots, s_n \in S \cup A, n \in \mathbb{N} \right\},$$

where the noncommutative product is understood from left to right, i.e.,  $\prod_{j=1}^{n+1} s_j := (\prod_{j=1}^n s_j) s_{n+1}$ , for  $n = 0, 1, 2, \dots$ . For completion,  $\prod_{j=1}^0 s_j := 1$ .

The set  $A \cdot \langle S \rangle$  is called the *subalgebra generated by  $S$  over  $A$*  and we call an element  $f \in A \cdot \langle S \rangle$  a noncommutative polynomial with coefficients in  $A$  and set of variables  $S$ .

**Definition 2.** Let  $C$  be a noncommutative ring and  $A$  a subring of  $C$ . We say that an element  $\alpha \in C$  is integral over  $A$  if there exists a noncommutative polynomial  $f$  with coefficients in  $A$  such that  $f(\alpha) = 0$ . Furthermore, we say that  $\beta \in C$  is integral over  $\alpha \in C$  if  $\beta$  is integral over  $A \cdot \langle \alpha \rangle$ . Finally,  $\alpha$  and  $\beta$  are associated integral if  $\alpha$  is integral over  $\beta$  and  $\beta$  is integral over  $\alpha$ .

In order to characterize the algebraic structure of bispectrality in the present noncommutative context, we start with the following definitions.

**Definition 3.** Let  $\mathbb{K}$  be a field, we denote by  $\mathbb{K}\langle x_\lambda \mid \lambda \in \Lambda \rangle$  the free algebra generated by the letters  $x_\lambda$ ,  $\lambda \in \Lambda$  i.e.,

$$\mathbb{K}\langle x_\lambda \mid \lambda \in \Lambda \rangle = \bigoplus_{F \subset \Lambda, F \text{ finite}} \bigoplus_{\lambda \in F} \mathbb{K} \cdot x_\lambda.$$

**Definition 4.** Let  $A$  be a  $\mathbb{K}$ -algebra. A presentation for an algebra  $A$  is a triple  $(\mathbb{K}\langle x_\lambda \mid \lambda \in \Lambda \rangle, f, I)$  such that  $I \subset A$  is an ideal and  $f : \mathbb{K}\langle x_\lambda \mid \lambda \in \Lambda \rangle / I \rightarrow A$  is an isomorphism. Furthermore, we say that  $A$  is finitely generated if there exists a presentation with  $\Lambda$  finite and finitely presented if there exists a presentation with  $\Lambda$  finite and the ideal  $I$  is generated by finitely many elements.

## 5 Three examples

Take for  $\Psi(x, z)$  the matrix valued function

$$\Psi(x, z) = e^{xz} \begin{pmatrix} z - 1/x & 1/x^2 \\ 0 & z - 1/x \end{pmatrix}$$

and consider all instances of matrix-valued polynomials  $\theta(x)$  and differential operators  $B$  (with matrix coefficients  $b_i(z)$ ) such that

$$\Psi B \equiv \sum_{i=1}^m \left( \frac{d}{dz} \right)^i \Psi b_i = \theta(x) \Psi(x, z).$$

In this case one has

$$L\Psi = -z^2\Psi$$

with

$$L = - \left( \frac{d}{dx} \right)^2 + 2 \begin{pmatrix} 1/x^2 & -2/x^3 \\ 0 & 1/x^2 \end{pmatrix}.$$

In other words for this specific differential operator in the variable  $x$  we are asking for all bispectral "partners" of  $L$ .

One finds that one such pair  $(B, \theta)$  is given by

$$B = \left( \frac{d}{dz} \right)^3 - 3 \left( \frac{d}{dz} \right)^2 \cdot \frac{1}{z} + 3 \left( \frac{d}{dz} \right) \cdot \frac{1}{z^2} + 3 \begin{pmatrix} 0 & 1/z^2 \\ 0 & 0 \end{pmatrix}$$

and  $\theta(x)$  the scalar-valued polynomial

$$\theta(x) = x^3.$$

The set of all possible  $\theta(x)$  is given by the following subalgebra  $\mathbb{A}$ . The complete statement is given by

**Theorem 1.** Let  $\Gamma$  be the sub-algebra of  $M_2(\mathbb{C})[x]$  of the form

$$\begin{pmatrix} r_0^{11} & r_0^{12} \\ 0 & r_0^{11} \end{pmatrix} + \begin{pmatrix} r_1^{11} & r_1^{12} \\ 0 & r_1^{11} \end{pmatrix} x + \begin{pmatrix} r_2^{11} & r_2^{12} \\ r_1^{11} & r_2^{22} \end{pmatrix} x^2 + \begin{pmatrix} r_3^{11} & r_3^{12} \\ r_2^{22} + r_2^{11} - r_1^{12} & r_3^{22} \end{pmatrix} x^3 + x^4 p(x),$$

where  $p \in M_2(\mathbb{C})[x]$  and all the variables  $r_0^{11}, r_0^{12}, r_1^{11}, r_1^{12}, r_2^{11}, r_2^{12}, r_2^{22}, r_3^{11}, r_3^{12}, r_3^{22} \in \mathbb{C}$ . Then  $\Gamma = \mathbb{A}$ . Moreover, for each  $\theta$  we have an explicit expression for the operator  $B$ .

Furthermore, we have the presentation  $\mathbb{A} = \mathbb{C} \cdot \langle \alpha_0, \alpha_1 \mid I = 0 \rangle$  with the ideal  $I$  given by

$$I := \langle \alpha_0^2, \alpha_1^3 + \alpha_0 \alpha_1 \alpha_0 - 3 \alpha_1 \alpha_0 \alpha_1 + \alpha_0 \alpha_1^2 + \alpha_1^2 \alpha_0 \rangle.$$

This is an example of an algebra with an integral element over a nilpotent one.

For the next example take for  $\psi(x, z)$  the matrix-valued function

$$\psi(x, z) = \left[ \frac{d}{dx} - \begin{pmatrix} 1/x & -1/x^2 & 1/x^3 \\ 0 & 1/x & -1/x^2 \\ 0 & 0 & 1/x \end{pmatrix} \right] e^{xz} I = e^{xz} \begin{pmatrix} z - 1/x & 1/x^2 & -1/x^3 \\ 0 & z - 1/x & 1/x^2 \\ 0 & 0 & z - 1/x \end{pmatrix}.$$

Here one can see that

$$L\psi = -z^2\psi$$

with

$$L = -\left(\frac{d}{dx}\right)^2 + 2\begin{pmatrix} 1/x^2 & -2/x^3 & 3/x^4 \\ 0 & 1/x^2 & -2/x^3 \\ 0 & 0 & 1/x^2 \end{pmatrix}.$$

The results in this case about the set of all possible  $\theta(x)$  are given below.

**Theorem 2.** *Let  $\Gamma$  the sub-algebra of  $M_3(\mathbb{C})[x]$  of the form*

$$\begin{aligned} & \begin{pmatrix} r_0^{11} & r_0^{12} & r_0^{13} \\ 0 & r_0^{22} & r_0^{23} \\ 0 & 0 & r_0^{11} \end{pmatrix} + \begin{pmatrix} r_1^{11} & r_1^{12} & r_1^{13} \\ r_0^{22} - r_0^{11} & r_1^{22} & r_1^{23} \\ 0 & r_0^{22} - r_0^{11} & r_1^{11} + r_0^{23} - r_0^{12} \end{pmatrix} x \\ & + \begin{pmatrix} r_1^{22} - r_1^{11} - r_0^{23} + r_0^{12} & r_2^{12} & r_2^{13} \\ r_0^{22} - r_0^{11} & r_2^{22} & r_2^{23} \\ r_1^{22} - r_1^{11} & r_2^{11} + r_1^{23} - r_1^{12} \end{pmatrix} x^2 + \begin{pmatrix} r_3^{11} & r_3^{12} & r_3^{13} \\ r_3^{21} & r_3^{22} & r_3^{23} \\ r_1^{22} - 2r_1^{11} - r_0^{23} + r_0^{12} & r_3^{32} & r_3^{33} \end{pmatrix} x^3 \\ & + \begin{pmatrix} r_4^{11} & r_4^{12} & r_4^{13} \\ r_4^{21} & r_4^{22} & r_4^{23} \\ r_3^{32} + r_3^{21} - r_2^{22} - r_2^{11} + r_1^{12} & r_4^{42} & r_4^{43} \end{pmatrix} x^4 \\ & + \begin{pmatrix} r_5^{11} & r_5^{12} & r_5^{13} \\ r_5^{21} & r_5^{22} & r_5^{23} \\ r_4^{32} + r_4^{21} - r_3^{33} - r_3^{22} - r_3^{11} + r_2^{23} + r_2^{12} - r_1^{13} & r_5^{42} & r_5^{43} \end{pmatrix} x^5 + x^6 p(x), \end{aligned}$$

where  $p \in M_3(\mathbb{C})[x]$  and all the variables  $r_0^{11}, r_0^{12}, \dots, r_5^{33} \in \mathbb{C}$  are arbitrary.

Then,  $\Gamma = \mathbb{A}$  and for each  $\theta$  we have an explicit expression for the operator  $B$ .

Furthermore, we have the presentation  $\mathbb{A} = \mathbb{C} \cdot \langle \alpha_2, \alpha_3 \mid I = 0 \rangle$  with

$$I = \langle \alpha_2^3, \alpha_3^2 - \alpha_3, (\alpha_3 \alpha_2)^2 \alpha_3 - 4\alpha_3 \alpha_2^2 \alpha_3 \rangle.$$

This is an example of an algebra with nilpotent and idempotent associated elements.

As the last example we consider a case when both "eigenvalues"  $F$  and  $\theta$  are matrix valued. Let

$$\psi(x, z) = \frac{e^{xz}}{(x-2)xz} \begin{pmatrix} \frac{x^3 z^2 - 2x^2 z^2 - 2x^2 z + 3xz + 2x - 2}{xz - 2} & \frac{1}{x} \\ \frac{xz - 2}{z} & x^2 z - 2xz - x + 1 \end{pmatrix}$$

and

$$L = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \cdot \left(\frac{d}{dx}\right)^2 + \begin{pmatrix} 0 & \frac{1}{(x-2)x^2} \\ -\frac{1}{x-2} & 0 \end{pmatrix} \cdot \left(\frac{d}{dx}\right) + \begin{pmatrix} -\frac{1}{x^2(x-2)^2} & \frac{x-1}{x^3(x-2)^2} \\ \frac{2x-1}{x(x-2)^2} & -\frac{2x^2-4x+3}{x^2(x-2)^2} \end{pmatrix},$$

then  $L\psi = \psi F$  with

$$F(z) = \begin{pmatrix} 0 & 0 \\ 0 & z^2 \end{pmatrix}.$$

It is easy to check that  $\psi B = \theta \psi$  for

$$B = \left(\frac{d}{dz}\right)^3 \cdot \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} + \left(\frac{d}{dz}\right)^2 \cdot \begin{pmatrix} 0 & 0 \\ -\frac{2z+1}{z} & 0 \end{pmatrix} + \left(\frac{d}{dz}\right) \cdot \begin{pmatrix} 1 & 0 \\ \frac{2(z-1)}{z^2} & 1 \end{pmatrix} + \begin{pmatrix} -z^{-1} & 0 \\ 6z^{-3} & z^{-1} \end{pmatrix}$$

and

$$\theta(x) = \begin{pmatrix} x & 0 \\ x^2(x-2) & x \end{pmatrix}.$$

In this case we characterize the algebra  $\mathbb{A}$  of all polynomial  $F$  such that there exist  $L = L(x, \frac{d}{dx})$  with  $L\psi = \psi F$  as follows

**Theorem 3.** *Let  $\Gamma$  be the sub-algebra of  $M_2(\mathbb{C})[z]$  of the form*

$$\begin{pmatrix} a & 0 \\ b-a & b \end{pmatrix} + \begin{pmatrix} c & c \\ a-b-c & -c \end{pmatrix} z + \begin{pmatrix} a-b-c & c+a-b \\ d & e \end{pmatrix} \frac{z^2}{2} + z^3 p(z),$$

where  $p \in M_2(\mathbb{C})[z]$  and all the variables  $a, b, c, d, e$  are arbitrary. Then  $\Gamma = \mathbb{A}$ .

Furthermore, we have the presentation  $\mathbb{A} = \mathbb{C} \cdot \langle \theta_1, \theta_3, \theta_4, \theta_5 \mid I = 0 \rangle$  with

$$\begin{aligned} I = & \langle \theta_1^2 - \theta_1, \theta_4^2, \theta_4\theta_5, \theta_4\theta_1 + \theta_4\theta_3 - 2\theta_4 - \theta_5\theta_4 - \theta_5^2, \theta_3^2 - \theta_3 + \theta_5 - 3\theta_3\theta_4\theta_3\theta_5 - \theta_1\theta_4 - \theta_5\theta_1, \\ & \theta_3\theta_1 - \theta_1 - \theta_4 - \frac{1}{2}\theta_4\theta_1 + \frac{1}{2}\theta_4\theta_3 + \theta_5\theta_1 - \frac{1}{2}\theta_5\theta_4 + \frac{1}{2}\theta_5^2 + \theta_3\theta_4 - \theta_1\theta_5 - \theta_3\theta_5, \\ & \theta_1\theta_3 - \theta_3 + \theta_4 + \theta_5 - \frac{3}{2}\theta_4\theta_1 + \frac{3}{2}\theta_4\theta_3 - 2\theta_5\theta_1 - \frac{3}{2}\theta_5\theta_4 + \frac{3}{2}\theta_5^2 + 3\theta_3\theta_4 + \theta_3\theta_5, \\ & \theta_5\theta_3 - \theta_4\theta_1 + \theta_4\theta_3 - \theta_5\theta_1 - \theta_5\theta_4 + \theta_5^2, \theta_5\theta_1\theta_5 - \theta_5^2\theta_1 - \theta_5\theta_4, \theta_5\theta_4\theta_1 - \theta_5^3 + \theta_5\theta_1\theta_4 + \theta_5^2\theta_1, \\ & \theta_4\theta_1\theta_5 + \theta_4\theta_3\theta_5 - \theta_3^3, \theta_5\theta_3\theta_4 + \theta_5\theta_1\theta_4 \rangle. \end{aligned}$$

This is an example of an algebra with two integer elements over one nilpotent and one idempotent. This is linked to the spin Calogero systems whose relation with bispectrality can be found in Bergvelt et al. (2009); Kasman (2015).

Theorems 1, 2 and 3 give positive answers to the Conjectures 1, 2 and 3 of Grünbaum (2014) about three bispectral full rank 1 algebras. Moreover, these algebras are Noetherian and finitely generated because they are contained in the  $N \times N$  matrix polynomial ring  $M_N(\mathbb{K}[x])$  (See Vasquez and Zubelli (2021)).

We close this section by remarking the important role played by Darboux transformations, which goes back to Duistermaat and Grünbaum (1986) and Zubelli and Magri (1991). Indeed, the three examples in this section are instances of rational Darboux transformations from the scalar matrix exponential functions. All such Darboux transformations were shown to be bispectral in Theorems 1.1 and 1.2 of Geiger et al. (2017).

## 6 A few extensions of the problems discussed above

Since one of the goals of this paper is to serve as an “invitation” to look at this problem extended to a wider audience, we give a road map with some selected references. Indeed, the bispectral problem has many different incarnations, and in our opinion we are still far from having a unified theory.

In the scalar case here are some early papers that should be mentioned are Kasman and Rothstein (1997); Wilson (1993, 1998); Bakalov et al. (1996, 1997, 1998a,b); Grünbaum and Yakimov (2002); Geiger et al. (2017).

One natural issue concerns the numerical aspects involving the prolate spheroidal functions. In this case, the reader may want to consult the book Osipov et al. (2013).

Another direction concerns, the purely discrete (actually finite) version of time-band-limiting. In an effort to better understand the commuting property in question, one of us looked at the case when the real line is replaced by the  $N$  roots of unity. See Grünbaum (1981) as well as Perlstadt (1984); Perline (1987). Expression (11) in Grünbaum (1981) contains a small typo: the  $r(2)$  in the denominator should be replaced by  $r(1)$ .

Moving on to the discrete-continuous version of the bispectral problem, we have that for the scalar case, involving orthogonal polynomials satisfying differential equations the problem had already been considered by S. Bochner (and even earlier). A very good introduction to this is given in Haine (2001) and its references. See also Grünbaum (2001); Grünbaum and Haine (1999); Grünbaum et al. (1997); Grünbaum et al. (1999).

For the matrix valued case there are two sources of early examples, one resulting from the theory of matrix valued spherical functions, see Grünbaum et al. (2001, 2002a,b, 2003); Grünbaum et al. (2004); Grünbaum et al. (2005), and another one see Durán and Grünbaum (2004, 2007); Durán and de la Iglesia (2008). See also Grünbaum and Iliev (2003) as well as Boyallian and Liberati (2008); Miranian (2005); Reach (1988).

Solutions of the bispectral problem can be used to obtain integral operators which reflect some ordinary differential operator in the sense of (for instance) Casper et al. (2019). This fact generalizes the commuting property in the scalar case. It would be interesting to see whether this could be extended to the matrix case.

In the (noncommutative) matrix case, by considering operators in the physical and spectral variables acting from opposite directions, we maintain the Ad-conditions that played a substantial role in Duistermaat and Grünbaum (1986).

---

In this case, this leads to the embedding of the bispectral algebras of eigenvalues into the matrix polynomial algebra  $M_N(\mathbb{C}[x])$ . See Vasquez and Zubelli (2021).

Another natural direction would be to look for a characterization for algebras relating to the spin-Calogero system for matrices of arbitrary size of matrix  $N$ . The examples  $N = 1$  and  $N = 2$  were generalized to arbitrary matrix size  $N$  and was characterized as a subalgebra of  $M_N(\mathbb{C}[x])$  using a family of maps  $\{P_k\}_{k \in \mathbb{N}}$  satisfying some nice properties such as translation and a product rule similar to the Leibniz rule. See Vasquez and Zubelli (2021).

Finally, let us go back to the original problem. We recall that both in the scalar and in the matrix case the motivation behind the bispectral problem was a desire to understand "what is behind" the remarkable commutativity property between the operator of time-and-band limiting in the Fourier, Bessel and Airy cases. It was hard to suspect that this problem would have connections with many of the recent developments in integrable systems. There has been some progress in connecting the bispectral problem with the commutativity property mentioned above, and here again- at least in the scalar case- there are some connections with Integrable systems, see for instance Grünbaum (2021); Casper and Yakimov (2020); Casper et al. (2019); Casper et al. (2021a,b). For connections between the bispectral problem and the commuting property see Castro and Grünbaum (2017); Grünbaum et al. (2017, 2018).

## Acknowledgments

BVDC was supported by CAPES grants 88882 332418/2019-01 as well as IMPA. JPZ was supported by CNPq grants 302161 and 47408, as well as by FAPERJ under the program *Cientistas do Nosso Estado* grant E-26/202.927/2017, Brazil. BDVC and JPZ acknowledge the support from the FSU-2020-09 grant from Khalifa University, UAE.

## References

- J.J. Duistermaat and F.A. Grünbaum. Differential equations in the spectral parameter. *Communications in Mathematical Physics*, 103(2):177–240, 1986.
- D. Slepian and H. O. Pollak. Prolate spheroidal wave functions, Fourier analysis and uncertainty. I. *Bell Syst. Tech. J.*, 40:43–63, 1961. ISSN 0005-8580. doi:10.1002/j.1538-7305.1961.tb03976.x.
- H. J. Landau and H. O. Pollak. Prolate spheroidal wave functions, Fourier analysis and uncertainty, III: The dimension of the space of essentially time- and band-limited signals. *Bell Syst. Tech. J.*, 41:1295–1336, 1962. ISSN 0005-8580. doi:10.1002/j.1538-7305.1962.tb03279.x.
- H. J. Landau and H. O. Pollak. Prolate spheroidal wave functions, Fourier analysis and uncertainty. II. *Bell Syst. Tech. J.*, 40:65–84, 1961. ISSN 0005-8580. doi:10.1002/j.1538-7305.1961.tb03977.x.
- D. Slepian. Prolate spheroidal wave functions, Fourier analysis and uncertainty. IV: Extensions to many dimensions; generalized prolate spheroidal functions. *Bell Syst. Tech. J.*, 43:3009–3057, 1964. ISSN 0005-8580. doi:10.1002/j.1538-7305.1964.tb01037.x.
- D. Slepian. Prolate spheroidal wave functions, Fourier analysis, and uncertainty - V: The discrete case. *Bell Syst. Tech. J.*, 57:1371–1430, 1978. ISSN 0005-8580. doi:10.1002/j.1538-7305.1978.tb02104.x.
- David Slepian. Some comments on Fourier analysis, uncertainty and modeling. *SIAM Review*, 25(3):379–393, 1983. ISSN 00361445. URL <http://www.jstor.org/stable/2029386>.
- David S. Slepian. On bandwidth. *Proceedings of the IEEE*, 64:292–300, 1976.
- Alain Connes and Caterina Consani. Spectral triples and zeta-cycles. *arXiv preprint:2106.01715*, 2021.
- Alain Connes and Matilde Marcolli. *Noncommutative geometry, quantum fields and motives*, volume 47. AMS Colloquium Publications, Volume 55, New Delhi: Hindustan Book Agency, 2008. ISBN 978-81-85931-80-7.
- Alain Connes and Henri Moscovici. Prolate spheroidal operator and zeta. *arXiv preprint:2112.05500*, 2021.
- Alain Connes. Prolate spheroidal functions and zeta, 2021. URL <https://www.youtube.com/watch?v=d8LgqGApv1o>.
- F. Alberto Grünbaum. Some noncommutative matrix algebras arising in the bispectral problem. *SIGMA, Symmetry Integrability Geom. Methods Appl.*, 10:paper 078, 9, 2014. ISSN 1815-0659. doi:10.3842/SIGMA.2014.078.
- Craig A. Tracy and Harold Widom. Fredholm determinants, differential equations and matrix models. *Commun. Math. Phys.*, 163(1):33–72, 1994a. ISSN 0010-3616. doi:10.1007/BF02101734.
- Craig A. Tracy and Harold Widom. Level-spacing distributions and the Airy kernel. *Commun. Math. Phys.*, 159(1): 151–174, 1994b. ISSN 0010-3616. doi:10.1007/BF02100489.

- 
- F. A. Grünbaum, L. Longhi, and M. Perlstadt. Differential operators commuting with finite convolution integral operators: Some non-Abelian examples. *SIAM J. Appl. Math.*, 42:941–955, 1982. ISSN 0036-1399. doi:10.1137/0142067.
- Jorge P. Zubelli and Franco Magri. Differential equations in the spectral parameter, Darboux transformations and a hierarchy of master symmetries for KdV. *Commun. Math. Phys.*, 141(2):329–351, 1991. ISSN 0010-3616. doi:10.1007/BF02101509.
- Jorge P. Zubelli and D. S. Valerio Silva. Rational solutions of the master symmetries of the KdV equation. *Commun. Math. Phys.*, 211(1):85–109, 2000. ISSN 0010-3616. doi:10.1007/s002200050803.
- George Wilson. Bispectral commutative ordinary differential operators. *J. Reine Angew. Math.*, 442:177–204, 1993. ISSN 0075-4102. doi:10.1515/crll.1993.442.177.
- J.P. Zubelli. Differential equations in the spectral parameter for matrix differential operators. *Physica D: Nonlinear Phenomena*, 43(2-3):269–287, 1990.
- Jorge P. Zubelli. On a zero curvature problem related to the ZS-AKNS operator. *J. Math. Phys.*, 33(11):3666–3675, 1992. ISSN 0022-2488.
- Jorge P. Zubelli. Rational solutions of nonlinear evolution equations, vertex operators, and bispectrality. *J. Differ. Equations*, 97(1):71–98, 1992. ISSN 0022-0396. doi:10.1016/0022-0396(92)90084-Z.
- Alexander Sakhnovich and Jorge P. Zubelli. Bundle bispectrality for matrix differential equations. *Integral Equations Oper. Theory*, 41(4):472–496, 2001. ISSN 0378-620X. doi:10.1007/BF01202105.
- Alex Kasman. Bispectrality of  $N$ -component KP wave functions: a study in non-commutativity. *SIGMA, Symmetry Integrability Geom. Methods Appl.*, 11:paper 087, 22, 2015. ISSN 1815-0659. doi:10.3842/SIGMA.2015.087.
- Carina Boyallian and Jose I. Liberati. Matrix-valued bispectral operators and quasideterminants. *J. Phys. A, Math. Theor.*, 41(36):11, 2008. ISSN 1751-8113. doi:10.1088/1751-8113/41/36/365209. Id/No 365209.
- F. Alberto Grünbaum and Plamen Iliev. A noncommutative version of the bispectral problem. *J. Comput. Appl. Math.*, 161(1):99–118, 2003. ISSN 0377-0427. doi:10.1016/S0377-0427(03)00548-X.
- Joel Geiger, Emil Horozov, and Milen Yakimov. Noncommutative bispectral Darboux transformations. *Trans. Am. Math. Soc.*, 369(8):5889–5919, 2017. ISSN 0002-9947. doi:10.1090/tran/6950.
- Mark J. Ablowitz, David J. Kaup, Alan C. Newell, and Harvey Segur. The inverse scattering transform–Fourier analysis for nonlinear problems. *Stud. Appl. Math.*, 53:249–315, 1974. ISSN 0022-2526. doi:10.1002/sapm1974534249.
- Yu Yu Berest and A P Veselov. Huygens’ principle and integrability. *Russian Mathematical Surveys*, 49(6):5, 1994. URL <http://stacks.iop.org/0036-0279/49/i=6/a=R02>.
- Fabio A. C. C. Chalub and Jorge P. Zubelli. Huygens’ principle for hyperbolic operators and integrable hierarchies. *Phys. D*, 213(2):231–245, 2006. ISSN 0167-2789. doi:10.1016/j.physd.2005.11.008. URL <http://dx.doi.org/10.1016/j.physd.2005.11.008>.
- Fabio A. C. C. Chalub and Jorge P. Zubelli. Huygens’ principle, Dirac operators, and rational solutions of the AKNS hierarchy. *Math. Phys. Anal. Geom.*, 8(3):187–210, 2005. ISSN 1385-0172. doi:10.1007/s11040-004-6683-3. URL <http://dx.doi.org/10.1007/s11040-004-6683-3>.
- Fabio A. C. C. Chalub and Jorge P. Zubelli. Matrix bispectrality and Huygens’ principle for Dirac operators. In *Partial differential equations and inverse problems*, volume 362 of *Contemp. Math.*, pages 89–112. Amer. Math. Soc., Providence, RI, 2004. doi:10.1090/conm/362/06607. URL <http://dx.doi.org/10.1090/conm/362/06607>.
- Fabio A. C. C. Chalub and Jorge P. Zubelli. On Huygens’ principle for Dirac operators and nonlinear evolution equations. *J. Nonlinear Math. Phys.*, 8(suppl.):62–68, 2001. ISSN 1402-9251. Nonlinear evolution equations and dynamical systems (Kolimbary, 1999).
- Mirta M. Castro and F. Alberto Grünbaum. The algebra of differential operators associated to a family of matrix-valued orthogonal polynomials: five instructive examples. *Int. Math. Res. Not.*, 2006(7):33, 2006. ISSN 1073-7928. doi:10.1155/IMRN/2006/47602. Id/No 47602.
- Juan Tirao. The algebra of differential operators associated to a weight matrix: a first example. In *Groups, algebras and applications. XVIII Latin American algebra colloquium, São Pedro, Brazil, August 3–8, 2009. Proceedings.*, pages 291–324. Providence, RI: American Mathematical Society (AMS), 2011. ISBN 978-0-8218-5239-2.
- F. Alberto Grünbaum and Juan Tirao. The algebra of differential operators associated to a weight matrix. *Integral Equations Oper. Theory*, 58(4):449–475, 2007. ISSN 0378-620X. doi:10.1007/s00020-007-1517-x.
- W. Riley Casper. Elementary examples of solutions to Bochner’s problem for matrix differential operators. *J. Approx. Theory*, 229:36–71, 2018. ISSN 0021-9045. doi:10.1016/j.jat.2018.02.004.

- 
- Ignacio Zurrián. The algebra of differential operators for a matrix weight: an ultraspherical example. *Int. Math. Res. Not.*, 2017(8):2402–2430, 2017. ISSN 1073-7928. doi:10.1093/imrn/rnw104.
- Brian D. Vasquez and Jorge P. Zubelli. Matrix bispectrality of full rank one algebras. *arXiv preprint:2108.06997*, 2021.
- Maarten Bergvelt, Michael Gekhtman, and Alex Kasman. Spin Calogero particles and bispectral solutions of the matrix KP hierarchy. *Math. Phys. Anal. Geom.*, 12(2):181–200, 2009. ISSN 1385-0172. doi:10.1007/s11040-009-9058-y.
- Jorge P. Zubelli and Franco Magri. Differential equations in the spectral parameter, Darboux transformations and a hierarchy of master symmetries for KdV. *Comm. Math. Phys.*, 141(2):329–351, 1991. ISSN 0010-3616.
- Alex Kasman and Mitchell Rothstein. Bispectral Darboux transformations: The generalized Airy case. *Physica D*, 102(3-4):159–176, 1997. ISSN 0167-2789. doi:10.1016/S0167-2789(96)00208-4.
- George Wilson. Collisions of Calogero-Moser particles and an adelic Grassmannian (with an appendix by I. G. Macdonald). *Invent. Math.*, 133(1):1–41, 1998. ISSN 0020-9910. doi:10.1007/s002220050237.
- B. Bakalov, E. Horozov, and M. Yakimov. General methods for constructing bispectral operators. *Phys. Lett., A*, 222(1-2):59–66, 1996. ISSN 0375-9601. doi:10.1016/0375-9601(96)00624-X.
- B. Bakalov, E. Horozov, and M. Yakimov. Bispectral algebras of commuting ordinary differential operators. *Commun. Math. Phys.*, 190(2):331–373, 1997. ISSN 0010-3616. doi:10.1007/s002200050244.
- B. Bakalov, E. Horozov, and M. Yakimov. Highest weight modules over the  $W_{1+\infty}$  algebra and the bispectral problem. *Duke Math. J.*, 93(1):41–72, 1998a. ISSN 0012-7094. doi:10.1215/S0012-7094-98-09302-4.
- Bojko Bakalov, Emil Horozov, and Milen Yakimov. Automorphisms of the Weyl algebra and bispectral operators. In *The bispectral problem. Proceedings of the CRM workshop, Montréal, Canada, March 1997*, pages 3–10. Providence, RI: American Mathematical Society, 1998b. ISBN 0-8218-0949-0.
- F. Alberto Grünbaum and Milen Yakimov. Discrete bispectral Darboux transformations from Jacobi operators. *Pac. J. Math.*, 204(2):395–431, 2002. ISSN 1945-5844. doi:10.2140/pjm.2002.204.395.
- Andrei Osipov, Vladimir Rokhlin, and Hong Xiao. *Prolate spheroidal wave functions of order zero. Mathematical tools for bandlimited approximation*, volume 187. New York, NY: Springer, 2013. ISBN 978-1-4614-8258-1; 978-1-4614-8259-8. doi:10.1007/978-1-4614-8259-8.
- F. Alberto Grünbaum. Eigenvectors of a Toeplitz matrix: Discrete version of the prolate spheroidal wave functions. *SIAM J. Algebraic Discrete Methods*, 2:136–141, 1981. ISSN 0196-5212. doi:10.1137/0602017.
- Marci Perlstadt. A property of orthogonal polynomial families with polynomial duals. *SIAM J. Math. Anal.*, 15:1043–1054, 1984. ISSN 0036-1410. doi:10.1137/0515081.
- Ronald Keith Perline. Discrete time-band limiting operators and commuting tridiagonal matrices. *SIAM J. Algebraic Discrete Methods*, 8:192–195, 1987. ISSN 0196-5212. doi:10.1137/0608016.
- Luc Haine. The Bochner-Krall problem: Some new perspectives. In *Special functions 2000: current perspective and future directions. Proceedings of the NATO Advanced Study Institute, Tempe, AZ, USA, May 29–June 9, 2000*, pages 141–178. Dordrecht: Kluwer Academic Publishers, 2001. ISBN 0-7923-7119-4.
- F. Alberto Grünbaum. The bispectral problem: An overview. In *Special functions 2000: current perspective and future directions. Proceedings of the NATO Advanced Study Institute, Tempe, AZ, USA, May 29–June 9, 2000*, pages 129–140. Dordrecht: Kluwer Academic Publishers, 2001. ISBN 0-7923-7119-4.
- F. Alberto Grünbaum and Luc Haine. Associated polynomials, spectral matrices and the bispectral problem. *Methods Appl. Anal.*, 6(2):209–224, 1999. ISSN 1073-2772. doi:10.4310/MAA.1999.v6.n2.a6.
- F. Alberto Grünbaum, Luc Haine, and Emil Horozov. On the Krall-Hermite and the Krall-Bessel polynomials. *Int. Math. Res. Not.*, 1997(19):953–966, 1997. ISSN 1073-7928. doi:10.1155/S1073792897000615.
- F. Alberto Grünbaum, Luc Haine, and Emil Horozov. Some functions that generalize the Krall-Laguerre polynomials. *J. Comput. Appl. Math.*, 106(2):271–297, 1999. ISSN 0377-0427.
- F. A. Grünbaum, I. Pacharoni, and J. Tirao. A matrix-valued solution to Bochner’s problem. *J. Phys. A, Math. Gen.*, 34(48):10647–10656, 2001. ISSN 0305-4470. doi:10.1088/0305-4470/34/48/328.
- F. A. Grünbaum, I. Pacharoni, and J. Tirao. Matrix valued spherical functions associated to the complex projective plane. *J. Funct. Anal.*, 188(2):350–441, 2002a. ISSN 0022-1236. doi:10.1006/jfan.2001.3840.
- F. A. Grünbaum, I. Pacharoni, and J. Tirao. Matrix valued spherical functions associated to the three dimensional hyperbolic space. *Int. J. Math.*, 13(7):727–784, 2002b. ISSN 0129-167X. doi:10.1142/S0129167X02001460.

- 
- F. A. Grünbaum, I. Pacharoni, and J. Tirao. Matrix valued orthogonal polynomials of the Jacobi type. *Indag. Math., New Ser.*, 14(3-4):353–366, 2003. ISSN 0019-3577. doi:10.1016/S0019-3577(03)90051-6.
- F. Alberto Grünbaum, Inés Pacharoni, and Juan Tirao. An invitation to matrix-valued spherical functions: Linearization of products in the case of complex projective space  $P_2(\mathbb{C})$ . In *Modern signal processing.*, pages 147–160. Cambridge: Cambridge University Press, 2004. ISBN 0-521-82706-X.
- F. Alberto Grünbaum, Inés Pacharoni, and Juan Alfredo Tirao. Matrix valued orthogonal polynomials of Jacobi type: the role of group representation theory. *Annales de l'Institut Fourier*, 55(6):2051–2068, 2005. doi:10.5802/aif.2151. URL <http://www.numdam.org/articles/10.5802/aif.2151/>.
- Antonio J. Durán and F. Alberto Grünbaum. Orthogonal matrix polynomials satisfying second-order differential equations. *Int. Math. Res. Not.*, 2004(10):461–484, 2004. ISSN 1073-7928. doi:10.1155/S1073792804132583.
- Antonio J. Durán and F. Alberto Grünbaum. Matrix orthogonal polynomials satisfying second-order differential equations: coping without help from group representation theory. *J. Approx. Theory*, 148(1):35–48, 2007. ISSN 0021-9045. doi:10.1016/j.jat.2007.02.003.
- Antonio J. Durán and Manuel D. de la Iglesia. Second-order differential operators having several families of orthogonal matrix polynomials as eigenfunctions. *Int. Math. Res. Not.*, 2008:24, 2008. ISSN 1073-7928. doi:10.1093/imrn/rnn084. Id/No rnn084.
- L. Miranian. Matrix-valued orthogonal polynomials on the real line: some extensions of the classical theory. *J. Phys. A, Math. Gen.*, 38(25):5731–5749, 2005. ISSN 0305-4470. doi:10.1088/0305-4470/38/25/009.
- Michael Reach. Generating difference equations with the Darboux transformation. *Communications in Mathematical Physics*, 119(3):385–402, September 1988. doi:10.1007/BF01218079.
- W. Riley Casper, F. Alberto Grünbaum, Milen Yakimov, and Ignacio Zurrián. Reflective prolate-spheroidal operators and the KP/KdV equations. *Proc. Natl. Acad. Sci. USA*, 116(37):18310–18315, 2019. ISSN 0027-8424. doi:10.1073/pnas.1906098116.
- F. Alberto Grünbaum. Commuting integral and differential operators and the master symmetries of the Korteweg-de Vries equation. *Inverse Probl.*, 37(8):21, 2021. ISSN 0266-5611. doi:10.1088/1361-6420/ac11c6. Id/No 085010.
- W. Riley Casper and Milen T. Yakimov. Integral operators, bispectrality and growth of Fourier algebras. *J. Reine Angew. Math.*, 766:151–194, 2020. ISSN 0075-4102. doi:10.1515/crelle-2019-0031.
- W. Riley Casper, F. Alberto Grünbaum, Milen Yakimov, and Ignacio Zurrián. Reflective prolate-spheroidal operators and the adelic grassmannian. *to appear in Comm. Pure and Applied Mathematics*, 2021a.
- W. R. Casper, F. A. Grünbaum, M. Yakimov, and I. Zurrián. Algebras of differential operators for kernels of Airy type. *in preparation*, 2021b.
- M. Castro and F. A. Grünbaum. Time-and-band limiting for matrix orthogonal polynomials of Jacobi type. *Random Matrices Theory Appl.*, 6(4):12, 2017. ISSN 2010-3263. doi:10.1142/S2010326317400019. Id/No 1740001.
- F. A. Grünbaum, I. Pacharoni, and I. Zurrián. Time and band limiting for matrix valued functions: an integral and a commuting differential operator. *Inverse Probl.*, 33(2):14, 2017. ISSN 0266-5611. doi:10.1088/1361-6420/aa53b8. Id/No 025005.
- F. Alberto Grünbaum, Luc Vinet, and Alexei Zhedanov. Algebraic Heun operator and band-time limiting. *Commun. Math. Phys.*, 364(3):1041–1068, 2018. ISSN 0010-3616. doi:10.1007/s00220-018-3190-0.