

A functional approach to a Gelfand-Tsetlin type base for \mathfrak{o}_5

D.V. Artamonov

A realization of representations of the Lie algebra \mathfrak{o}_5 in the space of functions on a group $Spin_5 \simeq Sp_4$ is considered. In a representation we take a Gelfand-Tsetlin type base associated with a restriction $\mathfrak{o}_5 \downarrow \mathfrak{o}_3$. Such a base is useful in problems appearing in quantum mechanics. We construct explicitly functions on the group that correspond to base vectors. As in the cases of Lie algebras \mathfrak{gl}_3 , \mathfrak{sp}_4 these functions can be expressed through A -hypergeometric functions (this does not hold for algebras of these series in higher dimensions). Using this realization formulas for the action of generators are obtained.

1 Introduction

In the paper [1] I.M. Gelfand and M.L. Tsetlin constructed a base in an irreducible finite dimensional representation of \mathfrak{o}_N . A construction of this base is based on investigation of branching of an irreducible representation of \mathfrak{o}_N under a restriction of algebras $\mathfrak{o}_N \downarrow \mathfrak{o}_{N-1}$. Later it was understood that it is reasonable to also consider a Gelfand-Tsetlin type base in representations of \mathfrak{o}_{2n+1} using restrictions $\mathfrak{o}_{2n+1} \downarrow \mathfrak{o}_{2n-1}$ i.e. restrictions inside the series B . Such a base was constructed by Molev in [2]. He also obtained formulas for the action of generators of the algebra \mathfrak{o}_{2n+1} in this base. Mention that construction of such a Gelfand-Tsetlin type base is not unique.

There are a lot of papers where authors construct a Gelfand-Tsetlin type base for \mathfrak{o}_5 by hand and use it to classify states of a five-dimensional quasi-spin in the nuclear shell model [3],[4],[5]. Such construction gives a base other than constructed in [2]. Let us mention also the paper [6] that is also motivated by problems coming from quantum mechanics. In these papers the Gelfand-Tsetlin type base for \mathfrak{o}_5 was constructed using a technique of the projection operator (see definition in [7]). This operator plays an important role in the theory of the Gelfand-Tsetlin bases (see a review [8]), and also in general constructions in [2].

In the present paper we construct the Gelfand-Tsetlin type base for \mathfrak{o}_5 , but we use new ideas (other than in [2]). A starting point is the paper [9], where in the case \mathfrak{gl}_3 the following observation is done. If one uses a functional realization then a function of the group GL_3 , corresponding to Gelfand-Tsetlin base vectors can be expressed through a Gauss hypergeometric function (see a modern approach in [14]). This fact was used in [15] to obtain an explicit formula for an arbitrary Clebsh-Gordan coefficient for \mathfrak{gl}_3 .

Explicit formulas for the Gelfand-Tsetlin base vectors for \mathfrak{sp}_4 (such base vectors were constructed in [19]) were obtained in [16]. In this paper the formulas for the action of generators were derived.

Mention that one has an isomorphism of Lie algebras $\mathfrak{sp}_4 \simeq \mathfrak{o}_5$, thus a representation V of the algebra \mathfrak{o}_5 is automatically a representation of \mathfrak{sp}_4 . But under this isomorphism the subalgebras $\mathfrak{sp}_2 \subset \mathfrak{sp}_4$ and $\mathfrak{o}_3 \subset \mathfrak{o}_5$ are not identified. Hence the Gelfand-Tsetlin type bases in V corresponding to these chains of subalgebras are different. In applications the base corresponding to the chain $\mathfrak{o}_3 \subset \mathfrak{o}_5$ is needed.

Thus in the present paper the following problem is considered. Take a Lie group with the Lie algebra \mathfrak{o}_5 . For example one can take SO_5 or $Spin_5 = Sp_4$. Consider functions on this group. This space of functions is a representation of \mathfrak{o}_5 , one has a canonical embedding of a arbitrary irreducible finite dimensional representation into this functional representation. It is natural to ask the following question: which functions correspond to the Gelfand-Tsetlin base vectors? What is the action of generators onto them? We obtain answers to both question in the present paper.

Surprisingly, the case of the algebra \mathfrak{o}_5 is much more difficult than the case of the algebra \mathfrak{sp}_4 . The first steps are similar to those in [16]. At the beginning it is more convenient to work with the space of functions on SO_5 . We establish a relation between restriction problems $\mathfrak{o}_5 \downarrow \mathfrak{o}_3$ and $\mathfrak{gl}_3 \downarrow \mathfrak{gl}_1^1$. Using this relation we construct a base in the space of \mathfrak{o}_3 -highest vectors with a fixed \mathfrak{o}_3 -weight in a given \mathfrak{o}_5 -representation. The function corresponding to basic \mathfrak{o}_3 -highest vectors are written as Γ -series.

To obtain formulas for functions corresponding to arbitrary vectors (which are not \mathfrak{o}_3 -highest), we need to apply a power of a lowering operator from \mathfrak{o}_3 . It is easy to check that one gets a difficult function which is not a Γ -series. Thus the situation differs from the situation in the paper [16].

To write the function in a simpler way we use an isomorphism $\mathfrak{sp}_4 \simeq \mathfrak{o}_5$ and realize our representation in the space of functions on $Spin_5 = Sp_4$. Then the functions corresponding to the Gelfand-Tsetlin base vectors are written as

¹A restriction problem $g \downarrow h$ (where $h \subset g$ are Lie algebras) is a problem of description of h -highest vectors in an irreducible representation of g

Γ -series.

But also there are similarities with the paper [16]. To derive formulas for the action of generators we use the Principle Lemma, formulated in Section 4, which gives a formula for a result of a multiplication of a Γ -series depending on minors of the matrix composed of matrix elements (considered as functions on the group) onto a minor. In the Lemma we claim that a lattice underlying the Γ -series is related to relations between minors.

The Gelfand-Tsetlin base in a representation of \mathfrak{o}_5 based on restrictions $\mathfrak{o}_5 \downarrow \mathfrak{o}_3$ is not unique. The base constructed in the present paper differs from bases constructed in papers cited above. A construction of the present paper establishes a remarkable relation with the theory of A -hypergeometric functions (see [20]). From one hand this is a beautiful demonstration of the unity of mathematics. From the other hand that makes possible to find explicit solution of problems that seem to have no easy solution in the traditional approach to the representation theory. Thus using a relation with the theory of A -hypergeometric functions in the case \mathfrak{gl}_3 one manages to obtain a really explicit formula for an arbitrary Clebsch-Gordan coefficient for \mathfrak{gl}_3 (see [15], and especially [18]).

2 Preliminary facts

2.1 Algebras \mathfrak{sp}_4 and \mathfrak{o}_5 . A functional realization of a representations

Take a complex linear space with coordinated indexed by $\pm 2, \pm 1$. Consider a Lie group Sp_4 , which consists of linear transformations preserving the skew-symmetric form $\omega = e_{-2} \wedge e_2 + e_{-1} \wedge e_1$.

It's Lie algebra \mathfrak{sp}_4 is spanned by

$$f_{i,j} = E_{i,j} - \text{sign}(i)\text{sign}(j)E_{-j,-i}, \quad i, j = \pm 2, \pm 1.$$

Take a complex linear space with coordinated indexed by $\pm 2, \pm 1, 0$. Consider a Lie group SO_5 , which consists of linear transformations preserving the scalar product whose matrix has ones on the second diagonal and zeros on other places.

It's Lie algebra \mathfrak{o}_5 is spanned by

$$F_{i,j} = E_{i,j} - E_{-j,-i}, \quad i, j = \pm 2, \pm 1, 0.$$

On the space of complex-valued functions on the group $G = Sp_4, SO_5$ the group itself acts by right shifts. Thus the space of functions is a representation of the group G and hence of it's Lie algebra. Let a_i^j be a function of a matrix

element where i is a column index and j is a row index. These indices are taken from $\{\pm 2, \pm 1, 0\}$. Compose determinants of size 1 and 2:

$$a_i = \det(a_i^{-2}), \quad a_{i_1, i_2} = \det(a_i^j)_{\substack{j=-2, -1 \\ i=i_1, i_2}} \quad (1)$$

They are also function on the group.

The Lie algebra acts onto (1) by the ruler

$$\begin{aligned} f_{i,j} a_{i_1} &= a_{i_1|_{j \rightarrow i}} - \text{sign}(i) \text{sign}(j) a_{i_1|_{-i \rightarrow -j}}, \\ F_{i,j} a_{i_1} &= a_{i_1|_{j \rightarrow i}} - a_{i_1|_{-i \rightarrow -j}}, \end{aligned}$$

where $i_1|_{j \rightarrow i}$ is a substitution of j instead of i , and in the case $j \neq i_1$ one gets 0. Analogously one defines an action onto a_{i_1, i_2} . And onto a product of determinants the action is defined by the Leibnitz ruler.

One can easily prove that

$$a_{-2}^{m-2-m-1} a_{-2, -1}^{m-1} \quad (2)$$

is a highest vector of weight $[m_{-2}, m_{-1}]$.

The functional realization has the following advantage: one can write conditions that define polynomials in (1) that form a representation with the highest vector (2) (in terms of the indicator system, see [19]).

2.2 A mapping $Sp_4 \rightarrow SO(5)$

There exists a covering $Sp_4 \rightarrow SO_5$ that induces an isomorphism $\mathfrak{sp}_4 = \mathfrak{o}_5$. Note that this isomorphism can be derived from the coincidence of root systems C_2 and B_2 .

The mapping is constructed as follows. Consider the space $V = \mathbb{C} \langle e_{-2}, e_{-1}, e_1, e_2 \rangle$ of the standard representation of Sp_4 . Then the action of Sp_4 onto $V \wedge V$ has a five-dimensional invariant subspace

$$\mathbb{C} \langle v_{-2} = e_{-2} \wedge e_{-1}, v_1 = e_{-2} \wedge e_1, v_{-1} = e_2 \wedge e_{-1}, v_2 = e_1 \wedge e_2, v_0 = \frac{1}{\sqrt{2}}(e_{-2} \wedge e_2 + e_{-1} \wedge e_1) \rangle.$$

The action of Sp_4 on this subspace has an invariant scalar product such that $\langle v_{-i}, v_i \rangle = 1$, and other scalar products vanish. Thus the elements of Sp_4 act on this space by orthogonal matrices 5×5 , hence one has $Sp_4 \rightarrow SO_5$. This mapping is an epimorphism, a preimage of each element of SO_5 consists of two elements (in particular a preimage of $E \in SO_5$ consists of E and $-E$ from Sp_4). Thus one has a mapping $Fun(SO_5) \rightarrow Fun(Sp_4)$.

The obtained mapping between Lie algebras look as follows. The positive root elements of \mathfrak{sp}_4 are mapped by the following ruler

$\frac{1}{\sqrt{2}}f_{-2,-1}$	$\frac{1}{2}f_{-2,2}$	$-\frac{1}{\sqrt{2}}f_{-2,1}$	$\frac{1}{2}f_{-1,1}$
$F_{-1,0}$	$F_{-2,1}$	$F_{-2,0}$	$F_{-2,-1}$

The negative root elements of \mathfrak{sp}_4 are mapped to corresponding negative root elements of \mathfrak{o}_5 .

For the Cartan elements one has

$\frac{1}{2}(f_{-2,-2} + f_{-1,-1})$	$\frac{1}{2}(f_{-2,-2} - f_{-1,-1})$
$F_{-2,-2}$	$F_{-1,-1}$

In particular a vector with a \mathfrak{sp}_4 -weight $[x, y]$ has a \mathfrak{o}_5 -weight $\frac{1}{2}[x + y, x - y]$.

From an explicit form of the mapping we conclude that a_i are transformed to b_{j_1, j_2} , more precise one has:

a_{-2}	a_{-1}	a_0	a_1	a_2
$b_{-2,-1}$	$b_{-2,1}$	$\frac{1}{\sqrt{2}}(b_{1,-1} + b_{-2,2})$	$b_{2,-1}$	$b_{1,2}$

(3)

Now let us find images of determinants $a_{i_1, i_2} \in Fun(SO_5)$. These determinants form a representation with the highest weight $[1, 1]$. Hence images of these functions in $Fun(Sp_4)$ form a representation with the highest weight $[2, 0]$, which is spanned by $b_{j_1} b_{j_2}$.

The highest vectors are $a_{-2,-1}$ and $(b_{-2})^2$. Let us identify them then using the identification of \mathfrak{o}_5 and \mathfrak{sp}_4 one obtains a correspondence

$(b_{-2})^2$	$\sqrt{2}b_{-2}b_{-1}$	$\sqrt{2}b_{-2}b_1$	$b_{-2}b_2 + b_{-1}b_1$	$b_{-2}b_2 - b_{-1}b_1$	$\sqrt{2}b_{-1}b_2$	$-\sqrt{2}b_1b_2$	$(b_{-1})^2$	$-(b_1)^2$	$(b_2)^2$
$a_{-2,-1}$	$a_{-2,0}$	$a_{-1,0}$	$a_{-2,2}$	$a_{-1,1}$	$a_{0,1}$	$a_{0,2}$	$a_{-2,1}$	$a_{2,-1}$	$a_{1,2}$

(4)

Obviously the correspondence respects products.

2.3 A-hypergeometric functions

Information about a Γ -series can be found in [20].

Let $\mathcal{B} \subset \mathbb{Z}^N$ be a lattice and let $\mu \in \mathbb{Z}^N$ be a fixed vector. We call it *the shift vector*. Define a *hypergeometric Γ -series* in variables z_1, \dots, z_N by the formula

$$\mathcal{F}_\gamma(z, \mathcal{B}) = \sum_{v \in \mathcal{B}} \frac{z^{v+\gamma}}{\Gamma(v+\gamma+1)}, \quad (5)$$

where $z = (z_1, \dots, z_N)$. We use a multi-index notation:

$$z^{v+\gamma} := \prod_{i=1}^N z_i^{v_i+\gamma_i}, \quad \Gamma(v+\gamma+1) := \prod_{i=1}^N \Gamma(v_i+\gamma_i+1).$$

Note that when at least one components of $v + \gamma$ is integer and negative that the corresponding summand in (5) vanishes. This is the reason, why the considered in the present paper Γ -series are finite sums. For simplicity we write factorials instead of Γ -functions.

We need the following ruler for differentiation of a Γ -series:

$$\frac{\partial}{\partial z_i} \mathcal{F}_\gamma = \mathcal{F}_{\gamma - e_i}, \quad (6)$$

where e_i is a standard base vector for the i -th coordinate.

An A -hypergeometric function satisfies the Gelfand-Kapranov-Zelevinsky system of PDE, which consists of equations of two types .

1. Let $\alpha = (\alpha_1, \dots, \alpha_N)$ be a vector orthogonal to \mathcal{B} , then

$$\alpha_1 z_1 \frac{\partial}{\partial z_1} \mathcal{F}_\gamma + \dots + \alpha_N z_N \frac{\partial}{\partial z_N} \mathcal{F}_\gamma = (\alpha_1 \gamma_1 + \dots + \alpha_N \gamma_N) \mathcal{F}_\gamma, \quad (7)$$

it is sufficient to consider base vectors of the orthogonal compliment to \mathcal{B} .

2. Let $v \in \mathcal{B}$ and $v = v_+ - v_-$, where vectors v_+ , v_- have non-negative coordinates. Take non-zero elements $v_+ = (\dots, v_{i_1}, \dots, v_{i_k}, \dots)$, $v_- = (\dots, v_{j_1}, \dots, v_{j_l}, \dots)$. Then

$$\left(\frac{\partial}{\partial z_{i_1}}\right)^{v_{i_1}} \dots \left(\frac{\partial}{\partial z_{i_k}}\right)^{v_{i_k}} \mathcal{F}_\gamma = \left(\frac{\partial}{\partial z_{j_1}}\right)^{v_{j_1}} \dots \left(\frac{\partial}{\partial z_{j_l}}\right)^{v_{j_l}} \mathcal{F}_\gamma. \quad (8)$$

It is sufficient to consider only base vectors $v \in \mathcal{B}$.

2.4 A Gelfand-Tsetlin type base for a chain $g \supset h$

Let us introduce a unified notation for Lie algebras: denote as g the algebra \mathfrak{o}_5 of \mathfrak{sp}_4 , and denote as h the subalgebra $\langle F_{0,-2}, F_{-2,-2}, F_{-2,0} \rangle$ or $\langle f_{-2,1}, f_{-2,-2} - f_{1,1}, f_{1,-2} \rangle$.

Let V be a standard representation of $g = \mathfrak{o}_5$ with the highest weight $\mu_2 = [m_{-2}, m_{-1}]$. Take a subalgebra $h = \mathfrak{o}_3 = \langle F_{0,-2}, F_{-2,-2}, F_{-2,0} \rangle$ and consider V as a representation of this algebra. Now it is not irreducible but it splits into a sum of irreducibles. Consider the space of \mathfrak{o}_3 -highest vectors with the highest weight $\mu_1 = [s_{-2}]$. Chose a base in this space and let μ'_2 be an index numerating the base vectors. In the space of a \mathfrak{o}_3 -representation generated by a chosen \mathfrak{o}_3 -highest vector chose a weight base and let μ'_1 be an index numerating the base vectors. Then one gets that in the space V there exists a base indexed by

$$\begin{pmatrix} \mu_2 \\ \mu'_2 \\ \mu_1 \\ \mu'_1 \end{pmatrix} \quad (9)$$

Such a base is called a Gelfand-Tsetlin type base. Since the choice of a base in the space of \mathfrak{o}_3 -highest vectors is not unique, the construction of such a base is not unique.

One can describe this construction on the language of $g = \mathfrak{sp}_4$. In this case one must consider a subalgebra $h = \langle f_{-2,1}, f_{-2,-2} - f_{1,1}, f_{1,-2} \rangle$, isomorphic to \mathfrak{sl}_2 .

3 A base in the space of h -highest vectors of a representation of g

Let us be given a representation V of the Lie algebra g . The first step in the construction of the Gelfand-Tsetlin type base in V for the chain $g \supset h$ is a construction of base in the space of h -highest vectors in g .

It is more natural to do it on the language of \mathfrak{o}_5 , however to finish the construction of the Gelfand-Tsetlin type base we need to pass to the language of \mathfrak{sp}_4 .

3.1 A base in the space of \mathfrak{o}_3 -highest vectors

In [19] an explicit description of conditions is given which define in the space of functions an irreducible representation with the highest vector (2). They look as follows

1. $L_- f = 0$, where L_- is a left infinitesimal shift by an element of SO_5 , corresponding to a negative root .
2. $L_{-i,-i} f = m_{-i} f$, where $L_{-i,-i}$, $i = 1, 2$ is a left infinitesimal shift by an element of SO_5 , corresponding to a Cartan element $F_{-i,-i}$.
3. f satisfies the indicator system

$$\begin{cases} L_{-2,-1}^{l_{-2}+1} f = 0, & l_{-2} = m_{-2} - m_{-1}, \\ L_{-1,0}^{l_{-1}+1} f = 0 & l_{-1} = 2m_{-1} \end{cases},$$

where $L_{i,j}$ is a left infinitesimal shift by an element $F_{i,j}$.

Consider the space of \mathfrak{o}_3 -highest vectors with a given \mathfrak{o}_3 -weight. A function is a \mathfrak{o}_3 -highest vector if it depends on

$$a_{-2}, a_{-1}, a_1, a_{-2,-1}, a_{-2,1}, a_{-1,1}. \quad (10)$$

Let us find among function depending on these variables the functions satisfying conditions 1-3. Let us introduce for shots time a new notation for determinants which includes upper indices of columns participating in the determinant (see (1))

$$a_i^{-2}, \quad a_{i_1, i_2}^{-2, -1}.$$

One can easily prove that $L_{-i, -j}$ acts as follows:

$$\begin{aligned} L_{-2, -2} a_{i_1}^{-2} &= a_{i_1}^{-2}, \quad L_{-1, -1} a_{i_1}^{-2} = 0, \quad L_{-i, -i} a_{i_1, i_2}^{-2, -1} = a_{i_1, i_2}^{-2, -1}, \\ L_{-2, -1} a_{i_1}^{-2} &= a_{i_1}^{-1}, \quad L_{-2, -1} a_{i_1}^{-1} = 0, \quad L_{-1, 0} a_{i_1}^{-2} = 0, \\ L_{-2, -1} a_{i_1, i_2}^{-2, -1} &= 0, \quad L_{-1, 0} a_{i_1, i_2}^{-2, -1} = a_{i_1, i_2}^{-2, 0}, \quad L_{-1, 0} a_{i_1, i_2}^{-2, 0} = a_{i_1, i_2}^{-2, 1}, \quad L_{-1, 0} a_{i_1, i_2}^{-2, 1} = 0 \end{aligned} \quad (11)$$

Among functions depending on (10), let us find functions satisfying the conditions 1-3.

Theorem 1. *A function depending on (10) is a \mathfrak{o}_3 -highest vector in an irreducible representation with the highest vector (2) if and only if the following condition holds.*

If the highest weight is integer then f is a polynomial in determinants and the sum of exponents of determinants of order i equals l_{-i} .

If the highest weight is half-integer then

$$f = (a_{-2, -1})^{\frac{1}{2}} f_1 + a_{-2, 0} (a_{-2, -1})^{-\frac{1}{2}} f_2, \quad (12)$$

where f_1 and f_2 are polynomials in determinants. The sum of exponents of determinants of order 1 in each summand of f equals l_{-2} , and the sum of exponents of determinants of order 2 equals l_{-1} .

Proof. Consider first the case of an integer highest weight, the vector (2) is a polynomial in determinants and under the action of elements of the algebra the space of polynomials is invariant. Thus an arbitrary vector of the representation is written as a polynomial in determinants. Then the statement of the Theorem follows from (11).

Now consider the case of the half-integer highest weight. First of all let us show that a functions satisfying the conditions 1-3, looks as follows (12).

Consider the case of the highest vector $(a_{-2, -1})^{\frac{1}{2}}$. An arbitrary vector of the representation is a linear combination of vectors

$$F_{0, -1}^{p-1} F_{-1, -2}^{p-2} (a_{-2, -1})^{\frac{1}{2}}.$$

Such a vector is nonzero only if $p_{-2} = 0$, and $p_{-1} = 0$ or 1 . Indeed if $p_{-1} = 1$ one gets the vector $\frac{1}{2}a_{-2,0}(a_{-2,-1})^{-\frac{1}{2}}$, and for $p_{-1} = 2$ one gets

$$\begin{aligned} & -\frac{1}{2}a_{-2,1}(a_{-2,-1})^{-\frac{1}{2}} - \frac{1}{4}(a_{-2,0})^2(a_{-2,-1})^{-\frac{3}{2}} = \\ & = -\frac{1}{2}(a_{-2,-1})^{-\frac{3}{2}}(a_{-2,1}a_{-2,-1} + \frac{1}{2}(a_{-2,0})^2) = 0, \end{aligned}$$

we have used a relations $a_{-2,0}^2 = -\frac{1}{2}a_{-2,1}a_{-2,-1}$.

Now consider the case of a half-integer highest weight. The highest vector (2) can be written as follows:

$$v_0 = v'_0(a_{-2,-1})^{\frac{1}{2}},$$

where v'_0 is a polynomial in determinants. An arbitrary vector f of the representation is obtained as a linear combination of vectors that are results of application to the highest vectors of the operators $F_{0,-1}^{p-1}F_{-1,-2}^{p-2}$. As a result one gets a vector of type (12).

Now we need to show that each vector of type (12), satisfying the conditions from the formulation of the Theorem satisfies conditions 1-3. Note that conditions 1,2 are satisfied automatically one needs to check only the condition 3.

The operator $L_{-2,-1}$ acts onto determinants of order 1 only. Such determinants are included only in f_1, f_2 , thus they occur in non-negative integer powers, the sum of powers of determinants of order 1 equals l_{-2} . Hence the condition $L_{-2,-1}^{l_{-2}+1}f = 0$ holds.

Now consider the equation $L_{-1,0}^{2m_{-1}+1}f = L_{-1,0}^{2[m_{-1}]+1+1}f = 0$, where $[m_{-1}]$ is an integer part. The operator $L_{-1,0}^{2[m_{-1}]+1+1}$ acts onto each summand in (12) by the Leibnitz ruler as follows.

Either $L_{-1,0}^{2[m_{-1}]+1+1}$ acts onto the second factor f_1 or f_2 only. Then one obtains 0, since in f_1 and f_2 the sum of powers of determinants of order 2 equals $[m_{-1}]$, but such polynomials are annihilated by $L_{-1,0}^{2[m_{-1}]+1}$.

Either $L_{-1,0}^{2[m_{-1}]+1}$ acts onto the second factor f_1 or f_2 , and $L_{-1,0}$ acts onto the first factor. Then one gets 0 by the same reason.

Either $L_{-1,0}^{2[m_{-1}]+2-k}$ acts onto the second factor, and the operator $L_{-1,0}^k$, where $k \geq 2$ acts onto the first factor. Since the first factor is a vector of representation with the highest weight $[\frac{1}{2}, \frac{1}{2}]$, then under the action $L_{-1,0}^2$ it vanishes.

Hence the vector of type (12) vanishes under the action of $L_{-1,0}^{2m_{-1}+1}$.

□

From the results of [2] it follows that in the space of \mathfrak{o}_3 -highest vectors of a \mathfrak{o}_5 -representation with a fixed \mathfrak{o}_3 -weight there is a base indexed by the number σ and an integer or half-integer diagram satisfying the betweenness conditions

$$\sigma, \begin{pmatrix} m_{-2} & & m_{-1} & & 0 \\ & k_{-2} & & k_{-1} & \\ & & s_{-2} & & \end{pmatrix}, \quad (13)$$

where $\sigma = 0, 1$. In the case $k_{-1} = 0$, one has $\sigma = 0$.

Below in the same space we construct another base whose elements are indexed by the same diagrams. For this new base we manage to write explicitly the functions corresponding to diagrams.

Consider the space \mathbb{C}^4 . Let us write the determinants in the following order:

$$a = (a_{-2}, a_{-1}, a_{-2,1}, a_{-1,1}).$$

To the diagram (13) the corresponds the vector

$$\gamma = (s_{-2} - m_{-1}, k_{-2} - s_{-2}, m_{-1} - k_{-1}, 0).$$

Introduce a lattice

$$\mathcal{B}^1 = \mathbb{Z} \langle (1, -1, -1, 1) \rangle.$$

Lemma 1. *Let us be given a representation of \mathfrak{o}_5 with the highest weight $[m_{-2}, m_{-1}]$. In the space of \mathfrak{o}_3 -highest vectors with weight s_{-2} there exists a base indexed by diagrams of type (13). In this diagram in the case of the integer highest weight if $\sigma = 1$ then $k_{-1} \geq 1$. To the diagram (13) there corresponds the function*

$$a_{-2,0}^\sigma a_1^{m_{-2}-k_{-2}} a_{-2,-1}^{k_{-1}-\sigma} \mathcal{F}_\gamma(a, \mathcal{B}^1). \quad (14)$$

The weights are calculated as follows: s_{-2} is a (-2) -component of the weight, a (-1) -component of the weight equals $-2(k_{-2} + k_{-1}) + (m_{-2} + m_{-1}) + s_{-2} + \sigma$.

Remark 1. *The function is written explicitly as follows*

$$a_{-2,0}^\sigma a_1^{m_{-2}-k_{-2}} a_{-2,-1}^{k_{-1}-\sigma} \sum \frac{1}{p_{-1}! p_{-1,1}! p_1! p_{-2,1}!} a_{-1}^{p_{-1}} a_{-1,1}^{p_{-1,1}} a_{-2}^{p_{-2}} a_{-2,1}^{p_{-2,1}}, \quad (15)$$

the summation is taken over all integer non-negative collections $p_{-1}, p_{-1,1}, p_1, p_{-2,1}$, such that

$$\begin{aligned} p_{-2} + p_{-1} &= k_{-2} - m_{-2}, & p_{-1} + p_{-2} &= k_{-2} - m_{-1}, \\ p_{-1,1} + p_{-2,1} &= m_{-1} - k_{-1}, & p_{-1} + p_{-1,1} &= k_{-2} - s_{-2}. \end{aligned}$$

Proof. Let us prove that such functions form a base in the considered linear space.

Consider first the case of the integer highest weight. The functions described in Theorem 1 are polynomials in determinants. Consider the admissible polynomial f in determinants satisfying the conditions of Theorem 1. Then f as an element of the representation with the highest vector v_0 can be written as a linear combination of polynomials of type $F_{-1,-2}^p F_{0,-1}^q v_0$, where $v_0 = a_{-2}^{m_2-m_1} a_{-2,-1}^{m_1}$. One has

$$F_{0,-1}^2 a_{-2,-1} = 2F_{0,-1} a_{-2,-1} a_{-2,0} = 2a_{-2,0}^2 - 2a_{-2,-1} a_{-2,1} = -4a_{-2,-1} a_{-2,1}, \quad (16)$$

using it in the case when $q = 2q'$ is even one gets

$$F_{-1,-2}^p F_{0,-1}^{2q'} v_0 = \text{const} E_{-1,-2}^p E_{1,-1}^{q'} v_0.$$

We formally define an action of $E_{i,j}$ onto symbols a_{i_1}, a_{i_1, i_2} by the ruler

$$E_{i,j} a_X = a_{X|_{j \rightarrow i}}.$$

The polynomial on the right satisfies the conditions of Theorem 1 for the algebra \mathfrak{gl}_3 and the highest weight $[m_{-2}, m_{-1}, 0]$. Thus one obtains an isomorphism between the span of vectors $F_{-1,-2}^p F_{0,-1}^{2q'} v_0$ the space of representation of \mathfrak{gl}_3 with the highest weight $[m_{-2}, m_{-1}, 0]$.

Let us use the Biedenharn-Baird Theorem (see [9], and its modern formulation in [14]). It states that in the space of representation \mathfrak{gl}_3 with the highest weight $[m_{-2}, m_{-1}, 0]$ in the functional realization the Gelfand-Tsetlin vector corresponding to the diagram (13) is written as the function $\mathcal{F}_\gamma(a, \mathcal{B}^1)$.

Thus in the span of vectors $F_{-1,-2}^p F_{0,-1}^{2q'} v_0$ there exists a base (14), given by diagrams (13), where $\sigma = 0$.

Note that in the considered case the eigenvalues of $F_{-2,-2}$ and $F_{-1,-1}$ correspond to eigenvalues of $E_{-2,-2}$ and $E_{-1,-1} - E_{1,1}$. They are equal to s_{-2} and $-2(k_{-2} + k_{-1}) + (m_{-2} + m_{-1}) + s_{-2}$.

In the case when $q = 2q' + 1$ is odd one has

$$F_{-1,-2}^p F_{0,-1}^{2q'+1} (a_{-2}^{m_2-m_1} a_{-2,-1}^{m_1}) = \text{const} (E_{-1,-2}^p E_{1,-1}^{q'} (a_{-2}^{m_2-m_1} a_{-2,-1}^{m_1-1})) a_{-2,0}.$$

If one removes $a_{-2,0}$, than the polynomial in determinants on the right satisfies the conditions of Theorem 1 for \mathfrak{gl}_3 and the highest weight $[m_{-2} - 1, m_{-1} - 1, 0]$. In the space of such polynomials there exists the Gelfand-Tsetlin base indexed by integer diagrams

$$\begin{pmatrix} m_{-2} - 1 & & m_{-1} - 1 & & 0 \\ & k_{-2} - 1 & & k_{-1} - 1 & \\ & & & & s_{-2} - 1 \end{pmatrix}. \quad (17)$$

Note that here $k_{-1} - 1 \geq 0$.

In the span $F_{-1,-2}^p F_{0,-1}^{2q'+1} v_0$ there exists a base (14), given by diagrams (13), where $\sigma = 1$ in the case $k_{-1} \geq 1$.

In the odd case the eigenvalues of $F_{-2,-2}$ and $F_{-1,-1}$ correspond to eigenvalues of $E_{-2,-2} - 1$ and $E_{-1,-1} - E_{1,1}$. They are equal to $s_{-2} - 1$ and $-2(k_{-2} - 1 + k_{-1} - 1) + (m_{-2} - 1 + m_{-1} - 1) + s_{-2} - 1 = -2(k_{-2} + k_{-1}) + (m_{-2} + m_{-1}) + s_{-2} + 1$.

Thus in the case of integer highest weight there exist a base (14), indexed by (13) where $\sigma = 0, 1$.

Now consider the case of a half-integer highest weight. One has

$$F_{0,-1} a_{-2,-1}^{1/2} = \frac{1}{2} a_{-2,0} a_{-2,-1}^{-1/2}, \quad F_{0,-1}^2 a_{-2,-1}^{1/2} = 0. \quad (18)$$

The highest vector looks as follows $v_0 = a_{-2}^{m_{-2}-m_{-1}} a_{-2,-1}^{[m_{-1}]+1/2}$, where $[m_{-1}]$ is an integer part. A vector of the representation can be written as a linear combination of vectors of type

$$F_{-1,-2}^p F_{0,-1}^q (a_{-2}^{m_{-2}-m_{-1}} a_{-2,-1}^{[m_{-1}]+1/2}).$$

If $q = 2q'$ is even then using (16), (18), one gets that the previous expression can be written as:

$$F_{-1,-2}^p F_{0,-1}^{2q'} (a_{-2}^{m_{-2}-m_{-1}} a_{-2,-1}^{[m_{-1}]+1/2}) = \text{const} E_{-1,-2}^p E_{1,-1}^{q'} (a_{-2}^{m_{-2}-m_{-1}} a_{-2,-1}^{[m_{-1}]}) a_{-2,-1}^{1/2}.$$

Thus there exists a natural isomorphism between the span of vectors of type $F_{-1,-2}^p F_{0,-1}^{2q'} v_0$ and the space of vectors in the representation \mathfrak{gl}_3 with the highest weight $[m_{-2} - \frac{1}{2}, m_{-1} - \frac{1}{2}, 0]$. In this space there exists a base indexed by integer diagrams of type

$$\begin{pmatrix} m_{-2} - \frac{1}{2} & & m_{-1} - \frac{1}{2} & & 0 \\ & k_{-2} - \frac{1}{2} & & k_{-1} - \frac{1}{2} & \\ & & & & s_{-2} - \frac{1}{2} \end{pmatrix}. \quad (19)$$

Using the formula for the vector corresponding to the diagram in the case \mathfrak{gl}_3 , one gets that in the span $F_{-1,-2}^p F_{0,-1}^{2q'} v_0$ there exists a base (14), indexed by half-integer diagrams of type (13) with $\sigma = 0$.

The eigenvalues of $F_{-2,-2}$ and $F_{-1,-1}$ correspond to eigenvalues of $E_{-2,-2} - \frac{1}{2}$ and $E_{-1,-1} - E_{1,1} - \frac{1}{2}$. One easily proves that s_{-2} is a (-2) -component of the

weight, and a (-1) -component of the weight equals $-2(k_{-2} + k_{-1}) + (m_{-2} + m_{-1}) + s_{-2}$.

Now let $q = 2q' + 1$ be odd, one has

$$F_{-1,-2}^p F_{0,-1}^{2q'+1} (a_{-2}^{m_{-2}-m_{-1}} a_{-2,-1}^{[m_{-1}]+1/2}) = \text{const}(E_{-1,-2}^p E_{1,-1}^{q'} a_{-2}^{m_{-2}-m_{-1}} a_{-2,-1}^{[m_{-1}]}) a_{-2,0}^{-1/2} a_{-2,-1}^{-1/2},$$

thus one gets an isomorphism between the span vectors $F_{-1,-2}^p F_{0,-1}^{2q'+1} v_0$ and the space of the representation of \mathfrak{gl}_3 with the highest weight $[m_{-2} - \frac{1}{2}, m_{-1} - \frac{1}{2}, 0]$. Thus in the span there exists the base (14), indexed by integer diagrams of type (13) with $\sigma = 1$. The eigenvalues of $F_{-2,-2}$ and $F_{-1,-1}$ correspond to eigenvalues of $E_{-2,-2} - \frac{1}{2}$ and $E_{-1,-1} - E_{1,1} + \frac{1}{2}$. One easily proves that s_{-2} is a (-2) -component of the weight, and a (-1) -component of the weight equals $-2(k_{-2} + k_{-1}) + (m_{-2} + m_{-1}) + s_{-2} + 1$.

Thus in the case of the half-integer highest weight these exists a base (14), indexed by diagrams (13), where $\sigma = 0, 1$.

Thus we have proved that in the space of functions defined by Theorem 1 there exists a base indexed by diagrams (13). Such functions define \mathfrak{o}_3 -highest vectors in a given irreducible representation of \mathfrak{o}_5 . According to [2], in this space there exist a base indexed by (13). Thus using the arguments of dimension we obtain that we have constructed a base in the space of all \mathfrak{o}_3 -highest vectors.

Let us formulate the ruler of calculation of the weight: s_{-2} is a (-2) -component of the weight, a (-1) -component of the weight equals $-2(k_{-2} + k_{-1}) + (m_{-2} + m_{-1}) + s_{-2} + \sigma$.

□

3.1.1 Further considerations

An arbitrary vector is obtained form a \mathfrak{o}_3 -highest vector corresponding to a diagram (13) by application of $\frac{F_{0,-2}^p}{p!}$, $p = 0, \dots, 2s_{-2}$. Such a vector is encoded by a diagram

$$\sigma, \begin{pmatrix} m_{-2} & m_{-1} & 0 \\ & k_{-2} & k_{-1} \\ & & s_{-2} & -s_{-2} \\ & & & s_{-1} \end{pmatrix} \quad (20)$$

But when one applies $\frac{F_{0,-2}^p}{p!}$ to $a_{-2,0}^\sigma a_1^{m_{-2}-k_{-2}} a_{-2,-1}^{k_{-2}-\sigma} \mathcal{F}_\gamma(a, \mathcal{B}_1)$ one gets not a Γ -series but a much more difficult functions.

It turns out that on the language of \mathfrak{sp}_4 one can construct the Gelfand-Tsetlin type base encoded by (20), such that each vector is presented by a function on Sp_4 , written as a Γ -series.

3.1.2 A base in the space of \mathfrak{h} -highest vectors

Let us write the results obtained above on the language of $g = \mathfrak{sp}_4$ and it's subalgebra h . Let us find an image of (14) in $Fun(Sp_4)$. The mapping $Fun(SO_5) \rightarrow Fun(Sp_4)$ is described explicitly in (3), (4). One gets that the image of the function (14) up to multiplication by a constant is written as

$$(b_{-2}b_{-1})^\sigma b_{-1,2}^{2(m_{-2}-k_{-2})} (b_{-2})^{k_{-1}} \cdot \mathcal{F}_\gamma(b_{-2,-1}, b_{-2,1}, (b_{-1})^2, -b_1b_{-1} + b_{-2}b_2, \mathcal{B}^1) \quad (21)$$

Let us show that this function is a Γ -series in *determinants* b_i, b_{i_1, i_2} . The function \mathcal{F} from (21) is written as

$$(b_{-2}b_{-1})^\sigma b_{-1,2}^{2(m_{-2}-k_{-2})} (b_{-2})^{k_{-1}} \cdot \sum_{n \in \mathbb{Z}} \frac{b_{-2,-1}^{\gamma_1+n}}{(\gamma_1+n)!} \frac{b_{-2,1}^{\gamma_2-n}}{(\gamma_2-n)!} \frac{b_{-1}^{2\gamma_3-2n}}{(2\gamma_3-2n)!} \frac{(-b_1b_{-1} + b_{-2}b_2)^n}{n!} \quad (22)$$

Consider in detail the sum in this expression. Let us open the brackets in the last factor. Then one gets an expression that is a Γ -series in determinants. It's shift vector δ has components

	$b_{-2,-1}$	$b_{-2,1}$	b_{-1}	$-b_1$	b_{-2}	b_2
δ	γ_1	γ_2	$2\gamma_3$	0	0	0

(23)

And the lattice is generated by two vectors (see row of the table below)

	$b_{-2,-1}$	$b_{-2,1}$	b_{-1}	b_1	b_{-2}	b_2
v_1	1	-1	-1	1	0	0
v_0	0	0	1	1	-1	-1

(24)

Put

$$\begin{aligned} \mathcal{B}^2 &= \mathbb{Z} \langle v_0, v_1 \rangle, \\ \delta &= (s_{-2} - m_{-1}, k_{-2} - s_{-2}, 2(m_{-1} - k_{-1}), 0, 0, 0) \end{aligned} \quad (25)$$

One obtains the Lemma.

Lemma 2. *On the language \mathfrak{sp}_4 a h -highest vector corresponding to a diagram is written as a function*

$$(b_{-2}b_{-1})^\sigma b_{-1,2}^{k_{-1}-\sigma} (b_{-2})^{2(m_{-2}-k_{-2})} \cdot \mathcal{F}_\delta(b_{-2,-1}, b_{-2,1}, b_1, b_{-1}, b_{-2}, b_2, \mathcal{B}^2). \quad (26)$$

4 The principle Lemma

In this section we formulate the main instrument that allows us to deal with Γ -series.

Lemma 3. *Let us be given a Γ -series in determinants b_i, b_{i_1, i_2} and the underlying lattice has generators $v_\alpha, \alpha = 1, \dots, K$, such that for $\alpha = 1, \dots, k, 1 \leq k \leq K$ the following holds*

1. *The generators v_α look as follows*

$$v_\alpha = e_{X_1} + e_{X_2} - e_{X_3} - e_{X_4},$$

where e_X is a unit vector corresponding to a coordinate $b_X, X = \{i\}$ or $\{i_1, i_2\}$.

2. *With each generator v_α a relation between determinants of the following type is associated*

$$b_{X_1} b_{X_2} + b_{X_3} b_{X_4} + b_{X_5} b_{X_6} = 0. \quad (27)$$

With each generator v_α let us associated a vector

$$r_\alpha = e_{X_1} + e_{X_2} - e_{X_5} - e_{X_6}. \quad (28)$$

Then modulo relations (27) one has

$$b_X \mathcal{F}_\gamma(b) = \sum_{s \in \mathbb{Z}_{\geq 0}^k} C_s^\gamma \mathcal{F}_{\gamma + e_X + sr}(b) \text{ mod } Pl, \quad (29)$$

here Pl is an ideal generated by (27).

The proof of this Lemma is similar to the proof of an analogous Lemma in the case \mathfrak{sp}_4 from [16] (see also [17]).

Let us give explicit formulas for the coefficients in (29). For $t, s \in \mathbb{Z}_{\geq 0}^k$ let us introduce functions (we use notations $tv := t_1 v_1 + \dots + t_k v_k, sr := s_1 r_1 + \dots + s_k r_k$ and a multi-index notation for factorials)

$$\mathcal{F}_\gamma^s(z) = \sum_{t \in \mathbb{Z}^k} \frac{(t+1) \dots (t+s) A^{\gamma+tv}}{s!(\gamma+tv)!} \quad (30)$$

Than as in [16], one obtains that

$$\begin{aligned}
C_s^\gamma &= \frac{\mathcal{F}_{\gamma+v}^s(1)}{\mathcal{F}_{\gamma+v+e_X+sr}^s(1)} - \sum_{p=0}^{s-1} \frac{\mathcal{F}_{\gamma+v}^p(1)\mathcal{F}_{\gamma+v+pr+e_X+(s-p)r}^{s-p}(1)}{\mathcal{F}_{\gamma+v+pr+e_X+(s-p)r}(1)\mathcal{F}_{\gamma+v+pr+e_X}(1)} = \\
&= \frac{\mathcal{F}_{\gamma+v}^s(1)}{\mathcal{F}_{\gamma+v+e_X+sr}^s(1)} - \sum_{p=0}^{s-1} \frac{\mathcal{F}_{\gamma+v}^p(1)\mathcal{F}_{\gamma+v+e_X+sr}^{s-p}(1)}{\mathcal{F}_{\gamma+v+e_X+sr}(1)\mathcal{F}_{\gamma+v+pr+e_X}(1)}
\end{aligned} \tag{31}$$

5 A function on Sp_4 , corresponding to a diagram

Let us find a function corresponding to a diagram (20) on the language \mathfrak{sp}_4 .

Note that a Γ -series (26) and the lattice (25) satisfy the conditions of the principle Lemma and a relation (27) corresponds only to v_1 .

5.1 A function corresponding to a h -highest diagram

In the space of h -highest vectors there exists a base indexed by (13), presented by functions (26), where δ is defined in (25).

Let us construct in this space another base also indexed by diagrams.

To the function (26) let us apply the ruler (29), one gets that the function (26) is presented as follows

$$const \cdot b_{2,-1}^{m-2} \mathcal{F}_\omega(b, \mathcal{B}^2) + l.o.t, \tag{32}$$

where

$$\omega = (s_{-2} - m_{-1}, k_{-2} - s_{-2}, 2(m_{-1} - k_{-1}) + \sigma, 0, 2k_{-2}) - \sigma, 0, \tag{33}$$

and $l.o.t.$ is a linear combination of functions of type $\mathcal{F}_{\omega+sr_1}$, $s \in \mathbb{Z}_{\geq 0}$.

Note that these functions also define h -highest vectors.

Introduce an ordering on the vectors δ

$$\delta' \prec \delta \Leftrightarrow \delta' = \delta + sr \text{ mod } \mathcal{B}^2$$

Then one gets that in the space of h -highest vectors one has two collections of functions indexed by diagrams: $\{b_{2,-1}^{m-2-k-2} \mathcal{F}_\omega(b, \mathcal{B}^2)\}$, and functions (26). The functions from the second collection are expressed from function of the first collection using a triangular operator. The second collection is a base in the space of h -highest vectors thus the first collection is also a base in this space.

Below in the space of h -highest vectors with a fixed h -weight the base $b_{2,-1}^{m-2-k-2} \mathcal{F}_\omega(b, \mathcal{B}^2)$ is considered.

5.2 A pseudo-action of \mathfrak{gl}_4

Determinants b_i, b_{i_1, i_2} satisfy some relations. These are the Plucker relations and also for example the relation $b_{-1,1} = -b_{-2,2}$. Consider the symbols B_i, B_{i_1, i_2} , that satisfy the Plucker relations *only*. Also they are supposed to be antisymmetric under permutations of i_1 and i_2 .

These symbols can be identified with function on GL_4 .

One has an action of \mathfrak{gl}_4 onto these symbols

$$E_{i,j}B_{i_1} = B_{i_1|j \rightarrow i}.$$

These formulas do not give an action of \mathfrak{gl}_4 onto determinants b_i, b_{i_1, i_2} . But the formula

$$f_{i,j} = E_{i,j} + \text{sign}(i)\text{sign}(j)E_{-j,-i}$$

gives an action of \mathfrak{sp}_4 onto B .

5.3 A function corresponding to a diagram

Consider an expression

$$B_{2,-1}^{m-2-k-2} \mathcal{F}_\omega(B, \mathcal{B}^2). \quad (34)$$

Apply the operator

$$\frac{f_{1,-2}^p}{p!} = \sum_{p_1+p_2=p} \frac{E_{2,-1}^{p_1}}{p_1!} \frac{E_{1,-2}^{p_2}}{p_2!}. \quad (35)$$

Here we use the fact that $E_{1,-2}$ and $E_{2,-1}$ commute and they do not act onto $B_{2,-1}^{m-2-k-2}$.

Remark 2. *It one tries to apply $\frac{F_{0,-2}^p}{p!}$ to the function (14), that it is not possible to write an analogous expression since the operators $E_{0,-2}$ and $E_{2,0}$ do not commute. This is the main obstacle in the case of the language \mathfrak{o}_5 .*

Let us write the operator $E_{1,-2}$ explicitly

$$B_1 \frac{\partial}{\partial B_{-2}} + B_{1,-1} \frac{\partial}{\partial B_{-2,-1}} + B_{1,2} \frac{\partial}{\partial B_{-2,2}}. \quad (36)$$

Analogously one writes $E_{2,-1}$ as follows:

$$B_2 \frac{\partial}{\partial B_{-1}} + B_{-2,2} \frac{\partial}{\partial B_{-2,-1}} + B_{1,2} \frac{\partial}{\partial B_{1,-1}}. \quad (37)$$

The summand in the expressions (36) and (37) commute.

Instead of the symbol B_1 let us introduce symbols B'_1 and B''_1 . In the function \mathcal{F}_δ , corresponding to a h -highest vector let us change B_1 to B'_1 . And in the operator (36) let us change B_1 to B''_1 .

Analogously one introduces B'_2 and B''_2 . One writes B'_2 instead of B_2 in \mathcal{F}_δ , and one writes B''_2 instead of B_2 in (37).

The operator $\frac{\partial}{\partial B_X}$ acts onto a Γ -series as a transformation $\omega \mapsto \omega - e_X$. Then we can write:

$$\frac{f_{1,-2}^p}{p!} \mathcal{F}_\omega = \sum_{p_1+p_2+q_1+q_2+q_3=p} \frac{(B''_1)^{p_1}}{p_1!} \frac{B_{1,-1}^{p_2-q_3}}{(p_2-q_3!)} \frac{(B''_2)^{q_1}}{q_1!} \frac{B_{-2,2}^{q_2}}{q_2!} \frac{B_{1,2}^{q_3}}{q_3!} \cdot \mathcal{F}_{\omega - p_1 e_{-2} - (p_2+q_2)e_{-2,-1} - q_1 e_{-1}}(B_{-2,-1}, B_{-2,1}, B_{-1}, -B'_1, B_{-2}, B'_2, B^2) \quad (38)$$

This expression is a Γ -series in variables

$$B''_1, B'_2, B''_2, B_{-1,1}, B_{-2,2}, B_{1,2}, B_{-2,-1}, B_{-2,1}, B_{-1}, -B'_1, B_{-2}, B'_1.$$

The shift vector looks as follows

$$(0, 0, 0, p, 0, 0, \delta - p e_{-2,-1}) \quad (39)$$

Let us write the generators of the lattice. These are the vectors involved in construction of \mathcal{F}_ω and also the vectors defining the shifts of exponents corresponding to the change in p_1, \dots, q_3 preserving the sum $p_1 + p_2 + p_3 + q_1 + q_2 + q_3 = p$. Such changes are generated by

	p_1	p_2	q_1	q_2	q_3
u_0	0	1	0	-1	0
u_1	0	0	0	-1	1
u_2	0	0	1	0	-1
u_3	1	-1	0	0	0

(40)

Using these vectors let us obtain vectors of changes of exponents of B_X . For the vector u_0 one obtains a vector

$$w_0 = e_{-1,1} - e_{-2,2}, \quad (41)$$

for vectors u_i , $i = 1, 2, 3$ one obtains vectors:

$$\begin{aligned} v_2 &= e_{-2,-1} + e_{1,2} - e_{-1,1} - e_{-2,2}, \\ v_3 &= e''_2 + e_{-1,1} - e_{-1} - e_{2,1}, \\ v_4 &= e''_1 + e_{-2,-1} - e_{-2} - e_{-1,1}. \end{aligned} \quad (42)$$

Thus one obtains the Theorem.

Theorem 2. *In the space of an irreducible representation of \mathfrak{g} there exists a Gelfand-Tsetlin type base corresponding to the chain $\mathfrak{g} \supset \mathfrak{h}$, indexed by diagrams (20), to a diagram there corresponds the function*

$$\begin{aligned}
& b_{2,-1}^{m_{-2}-k_{-2}} \mathcal{F}_\omega(b, \mathcal{B}_{GC}) \big|_{b'_1=b''_1=b_1, b'_2=b''_2=b_2}, \\
& b = (b''_1, b''_2, b_{-1,1}, b_{-2,2}, b_{1,2}, b_{-2,-1}, b_{-2,1}, b_{-1}, -b'_1, b_{-2}, b'_2), \\
& \omega = (0, 0, s_{-2} - s_{-1}, 0, 0, s_{-1} - m_{-1}, k_{-2} - s_{-2}, 2(m_{-1} - k_{-1}) + \sigma, 0, 2k_{-2} - \sigma, 0), \\
& \mathcal{B}_{GC} = \mathbb{Z} \langle v_0, w_0, v_1, v_2, v_3, v_4 \rangle.
\end{aligned} \tag{43}$$

Remark 3. *The vector ω satisfies the following selection rulers:*

$$\omega = (0, 0, \omega_{-1,1}, 0, 0, \omega_{-2,-1}, \omega_{-2,1}, \omega_{-1}, 0, \omega_{-2}, 0).$$

6 The action of generators

6.1 The action of an arbitrary $f_{i,j}$

Consider an arbitrary generator $f_{i,j}$ of the algebra \mathfrak{sp}_4 . Apply it to \mathcal{F}_ω , one gets

$$f_{i,j} \mathcal{F}_\omega = \sum_Y b_{i,Y} \frac{\partial}{\partial b_{j,Y}} \mathcal{F}_\omega - \text{sign}(i) \text{sign}(j) \sum_Z b_{-j,Z} \frac{\partial}{\partial b_{-i,Z}} \mathcal{F}_\omega$$

A summation is taken over all subsets $Y, X \subset \{\pm 1, \pm 2\}$, such that $i, j \notin Y$, $-i, -j \notin Z$. A differentiation acts onto a Γ -series by a change of a shift-vector, and a multiplication by a determinants acts according to (29). One has

$$\begin{aligned}
f_{i,j} \mathcal{F}_\omega &= \sum_Y \sum_{s^1 \in \mathbb{Z}_{\geq 0}^k} C_s^{\omega - e_Y, j} \mathcal{F}_{\omega + e_{Y,i} - e_{Y,j} + s^1 r} \\
&- \text{sign}(i) \text{sign}(j) \sum_Z \sum_{s^2 \in \mathbb{Z}_{\geq 0}^k} C_s^{\omega - e_Z, -i} \mathcal{F}_{\omega + e_{Z,-j} - e_{Z,-i} + s^2 r},
\end{aligned} \tag{44}$$

where C_s^ω is given by (31).

A transformation of the shift-vector

$$\omega \mapsto \omega + e_{Y,i} - e_{Y,j} + s^1 r, \quad \omega \mapsto \omega + e_{Z,-j} - e_{Z,-i} + s^2 r$$

corresponds to a transformation of the Gelfand-Tsetlin diagram. Let us derive it for some generators of \mathfrak{sp}_4 . It is sufficient to do it for $\frac{1}{2}(f_{-2,-2} + f_{-1,-1})$ and $\frac{1}{2}(f_{-2,-2} - f_{-1,-1})$, $f_{1,-2}$ and $f_{-2,1}$, $f_{-2,-1}$ and $f_{-1,-2}$.

6.2 The action of Cartan elements

The Cartan elements $\frac{1}{2}(f_{-2,-2} + f_{-1,-1})$ and $\frac{1}{2}(f_{-2,-2} - f_{-1,-1})$ on the language of \mathfrak{o}_5 turn into $F_{-2,-2}$ and $F_{-1,-1}$.

The Gelfand-Tsetlin type base consists of eigenvectors of $F_{-1,-1}$, $F_{-2,-2}$ and the eigenvalues equal to

$$\begin{aligned}\lambda_{-2} &= \sigma + 2(k_{-2} + k_{-1}) - (m_{-2} + m_{-1}) - s_{-2}, \\ \lambda_{-1} &= s_{-1}.\end{aligned}$$

This fact was derived in 3.1.

6.3 The action of $f_{1,-2}$ and $f_{-2,1}$

The function corresponding to a diagram is obtained from a function corresponding to a h -highest diagram by the action of $\frac{f_{1,-2}^{s_{-2}-s_{-1}}}{(s_{-2}-s_{-1})!}$

Hence the action of $f_{1,-2}$ gives a transformation $s_{-1} \mapsto s_{-1} - 1$ and a multiplication onto $s_{-2} - s_{-1} + 1$.

The action of $f_{-2,1}$ gives a transformation $s_{-1} \mapsto s_{-1} + 1$ and a multiplication onto $s_{-2} - s_{-1} + 1$.

6.4 The action of $f_{-1,1}$ and $f_{-2,2}$

Let us write $f_{-1,1}$ as

$$b_{-1} \frac{\partial}{\partial b'_1} + b_{-1} \frac{\partial}{\partial b''_1} + b_{-2,-1} \frac{\partial}{\partial b_{-2,1}} + b_{-1,2} \frac{\partial}{\partial b_{1,2}}$$

Also write $f_{-2,2}$ as

$$b_{-2} \frac{\partial}{\partial b'_2} + b_{-2} \frac{\partial}{\partial b''_2} + b_{-2,-1} \frac{\partial}{\partial b_{2,-1}} + b_{-2,1} \frac{\partial}{\partial b_{2,1}} +$$

The action of these expressions onto \mathcal{F}_δ is given by (44). Let us find a transformation of a diagram corresponding to the transformation $\omega \mapsto \omega - e_X + e_Y$ of the shift vector associated with the summand $b_Y \frac{\partial}{\partial b_X}$. Let us also describe a transformation of a diagram corresponding to the addition of r_α .

6.4.1 The addition of r_α

With each vector v_α one associates a vector r_α , defined in (28). Consider the vector

$$v_1 = e_{-2,-1} - e_{-2,1} - e_{-1} + e_1 \Rightarrow r_1 = e_{-2} + e_{-1,1} - e_{-2,1} - e_{-1}.$$

Addition of r_1 to ω from (43) is equivalent $\text{mod}\mathcal{B}_{GC}$ to the following transformation:

$$\begin{cases} s_{-1} \mapsto s_{-1} + 1, \\ \sigma = 0, k_{-1} \mapsto \sigma = 1, k_{-1} + 1 \\ \sigma = 1, k_{-1} \mapsto \sigma = 0, k_{-1} \end{cases}$$

Consider the vector

$$v_2 = e_{-2,-1} + e_{1,2} - e_{-1,1} - e_{-2,2} \Rightarrow r_2 = e_{-2,1} + e_{1,-2} - e_{-1,1} - e_{-2,2}$$

The addition of r_2 to ω from (43) is equivalent $\text{mod}\mathcal{B}_{GC}$ to the following transformation:

$$\begin{cases} k_{-2} \mapsto k_{-2} - 1, s_{-2} \mapsto s_{-2} - 2 \end{cases}$$

Consider the vector

$$v_3 = e_2'' + e_{-1,1} - e_{-1} - e_{1,2} \Rightarrow r_3 = e_1'' + e_{-1,2} - e_{-1} - e_{1,2}$$

The addition of r_3 to ω from (43) is equivalent $\text{mod}\mathcal{B}_{GC}$ to the following transformation:

$$\begin{cases} k_{-2} \mapsto k_{-2} - 1, \\ s_{-2} \mapsto s_{-2} - 1, \\ \sigma = 0, k_{-1} \mapsto \sigma = 1, k_{-1} + 1 \\ \sigma = 1, k_{-1} \mapsto \sigma = 0, k_{-1} \end{cases}$$

Consider the vector

$$v_5 = e_1'' + e_{-2,-1} - e_{-1,1} - e_{-2} \Rightarrow r_5 = e_{-1} + e_{-2,1} - e_{-1,1} - e_{-2}$$

The addition of r_4 to ω from (43) is equivalent $\text{mod}\mathcal{B}_{GC}$ to the following transformation:

$$\begin{cases} s_{-2} \mapsto s_{-2} - 1, \\ \sigma = 1, k_{-1} \mapsto \sigma = 0, k_{-1} - 1 \\ \sigma = 0, k_{-1} \mapsto \sigma = 1, k_{-1} \end{cases}$$

6.4.2 The transformation $\omega - e_{Y,1} + e_{Y,-1}$ associated with the action of $f_{-1,1}$

Consider the summand $b_{-1} \frac{\partial}{\partial b_1}$. The transformation of the shift vector looks as follows: $\omega \mapsto \omega + e_{-1} - e_1'$. It is equivalent $\text{mod}\mathcal{B}_{GC}$ to the following transformation of the diagram:

$$\left\{ s_{-2} \mapsto s_{-2} + 1, s_{-1} \mapsto s_{-1} + 1 \right.$$

Consider the summand $b_{-1} \frac{\partial}{\partial b_1}$. The transformation of the shift vector looks as follows: $\omega \mapsto \omega + e_{-1} - e'_1$. It is equivalent $\text{mod} \mathcal{B}_{GC}$ to the following transformation of the diagram:

$$\left\{ \begin{array}{l} s_{-2} \mapsto s_{-2} + 1 \\ \sigma = 1, k_{-1} \mapsto \sigma = 0, k_{-1} - 1 \\ \sigma = 0, k_{-1} \mapsto \sigma = 1, k_{-1} \end{array} \right.$$

Consider the summand $b_{-2,-1} \frac{\partial}{\partial b_{-2,1}}$. The transformation of the shift vector looks as follows: $\omega \mapsto \omega + e_{-2,-1} - e_{-2,1}$. It is equivalent $\text{mod} \mathcal{B}_{GC}$ to the following transformation of the diagram:

$$\left\{ s_{-2} \mapsto s_{-2} + 1, s_{-1} \mapsto s_{-1} + 1 \right.$$

Consider the summand $b_{-1,2} \frac{\partial}{\partial b_{1,2}}$. The transformation of the shift vector looks as follows: $\omega \mapsto \omega + e_{-1,2} - e_{1,2}$. It is equivalent $\text{mod} \mathcal{B}_{GC}$ to the following transformation of the diagram:

$$\left\{ k_{-2} \mapsto k_{-2} + 1, s_{-2} \mapsto s_{-2} + 1, s_{-1} \mapsto s_{-1} - 1 \right.$$

6.4.3 The transformation $\omega - e_{Y,2} + e_{Y,-2}$ associated with the action of $f_{-2,2}$

Consider the summand $b_{-2} \frac{\partial}{\partial b_2}$. The transformation of the shift vector looks as follows: $\omega \mapsto \omega + e_{-2} - e'_2$. It is equivalent $\text{mod} \mathcal{B}_{GC}$ to the following transformation of the diagram:

$$\left\{ k_{-1} \mapsto k_{-1} + 1, s_{-2} \mapsto s_{-2} + 1, s_{-1} \mapsto s_{-1} + 1 \right.$$

Consider the summand $b_{-2} \frac{\partial}{\partial b_2}$. The transformation of the shift vector looks as follows: $\omega \mapsto \omega + e_{-2} - e''_2$. It is equivalent $\text{mod} \mathcal{B}_{GC}$ to the following transformation of the diagram:

$$\left\{ \begin{array}{l} s_{-1} \mapsto s_{-1} - 1, \\ \sigma = 0, k_{-1} \mapsto \sigma = 1, k_{-1} + 1 \\ \sigma = 1, k_{-1} \mapsto \sigma = 0, k_{-1} \end{array} \right.$$

Consider the summand $b_{-2,-1} \frac{\partial}{\partial b_{2,-1}}$. The transformation of the shift vector looks as follows: $\omega \mapsto \omega + e_{-2,-1} - e_{2,-1}$. It is equivalent $\text{mod} \mathcal{B}_{GC}$ to the following transformation of the diagram:

$$\left\{ k_{-2} \mapsto k_{-2} + 1, s_{-2} \mapsto s_{-2} + 1, s_{-1} \mapsto s_{-1} + 1 \right.$$

Consider the summand $b_{-2,1} \frac{\partial}{\partial b_{2,1}}$. The transformation of the shift vector looks as follows: $\omega \mapsto \omega + e_{-2,1} - e_{2,1}$. It is equivalent $\text{mod } \mathcal{B}_{GC}$ to the following transformation of the diagram:

$$\left\{ s_{-2} \mapsto s_{-2} + 2, s_{-1} \mapsto s_{-1} + 2 \right.$$

References

- [1] Гельфанд И.М., Цетлин М.Л. Конечномерные представления группы ортогональных матриц, ДАН СССР, 1950, **51**, 825–828 (in Russian)
- [2] Molev A.I., Yangians and classical Lie algebras. NY : AMS, Mathematical Surveys and Monographs Vol 143, 2007.
- [3] Hecht K.T. , The vector coherent state method and its applications to problems of higher symmetries, Heidelberg: Springer, Lecture Notes in Phys., Vol 290, 1987
- [4] Helmers K., Symplectic invariants and Flowers' classification of shell model states, Nucl. Phys, 1961, **23**, 594–611
- [5] Artamonov D.V., Goloubeva V.A. Noncommutative Pfaffians and classification of states of five-dimensional quasi-spin, J.Math. Phys, 2012, **53**, 083504/1–083504/21
- [6] R. M. Asherova, Yu. F. Smirnov, V. N. Tolstoy Projected basis for many-phonon systems J. Phys. G: Nucl. Phys. vol. 4, 205 (1978)
- [7] R. M. Asherova, Yu. F. Smirnov, V. N. Tolstoy, “Description of a class of projection operators for semisimple complex Lie algebras”, Mat. Zametki, 26:1 (1979), 15–25; Math. Notes, 26:1 (1979), 499–504
- [8] V. N. Tolstoy, Fortieth anniversary of extremal projector method for Lie symmetries, Fuchs, J?rgen (ed.) et al., Noncommutative geometry and representation theory in mathematical physics. Satellite conference to the fourth European congress of mathematics, July 5–10, 2004, Karlstad, Sweden. Providence, RI: American Mathematical Society (AMS). Contemporary Mathematics 391, 371–384 (2005).
- [9] Biedenharn L. C., Baird G. E., *On the representations of semisimple Lie Groups II*, J. Math. Phys., **4**, (1963), 1449–1466.

- [10] Louck J. D., Biedenharn L. C., *A pattern calculus for tensor operators in the unitary groups*, Comm. Math. Phys. **8**, (1968), 89–131
- [11] Louck J. D., Biedenharn L. C., *Canonical adjoint tensor operators in $U(n)$* , J. Math. Phys., **11**, (1970), 2368–2411
- [12] Chacon E. , Ciftan M., Biedenharn L. C., *On the evaluation of the Multiplicity-Free Wigner coefficients of $U(n)$* , J. Math. Phys, **13**, (1972), 577–589
- [13] Louck J. D., Biedenharn L. C., *On the structure of the canonical tensor operators in the unitary groups II. An extension of the pattern calculus rules and the canonical splitting in $U(3)$* , J. Math. Phys, **13**, (1972), 1957–1984
- [14] D.V. Artamonov, *Formula for the Product of Gauss Hypergeometric Functions and Applications* ,J. Math. Sci. , 249, (2020), 817–826.
- [15] D.V. Artamonov , *The Clebsch–Gordan coefficients for the algebra \mathfrak{gl}_3 and hypergeometric functions* , Algebra i Analiz, 33:1 (2021), 1–29
- [16] D. V. Artamonov, “A Gelfand–Tsetlin-type basis for the algebra \mathfrak{sp}_4 and hypergeometric functions”, TMF, 206:3 (2021), 279–294; Theoret. and Math. Phys., 206:3 (2021), 243–257
- [17] D.V. Artamonov, *Antisymmetrization of the Gelfand–Kapranov–Zelevinskij Systems*, J. Math. Sci., 255, 535–542 (2021).
- [18] D.V. Artamonov, *3j-symbols for representation of the Lie algebra gl_3 in the Gelfand-Tselin base*, arXiv:2105.10761
- [19] D. P. Zhelobenko, *Compact Lie groups and their representations*. – American Mathematical Soc., 1973. – V. 40.
- [20] I. M. Gel’fand, M. I. Graev, V. S. Retakh, “General hypergeometric systems of equations and series of hypergeometric type”, Russian Math. Surveys, 47:4 (1992), 1–88