

Homotopic Policy Mirror Descent: Policy Convergence, Implicit Regularization, and Improved Sample Complexity

Yan Li [†]Tuo Zhao [‡]Guanghai Lan [§]

Abstract

We propose the homotopic policy mirror descent (HPMD) method for solving discounted, infinite horizon MDPs with finite state and action space, and study its policy convergence. We report three properties that seem to be new in the literature of policy gradient methods: (1) HPMD exhibits global linear convergence of the value optimality gap, and local superlinear convergence of the policy to the set of optimal policies with order γ^{-2} . The superlinear convergence of the policy takes effect after no more than $\mathcal{O}(\log(1/\Delta^*))$ number of iterations, where Δ^* is defined via a gap quantity associated with the optimal state-action value function; (2) HPMD also exhibits last-iterate convergence of the policy, with the limiting policy corresponding exactly to the optimal policy with the maximal entropy for every state. No regularization is added to the optimization objective and hence the second observation arises solely as an algorithmic property of the HPMD method. (3) For the stochastic HPMD method, we further demonstrate a better than $\mathcal{O}(|\mathcal{S}||\mathcal{A}|/\epsilon^2)$ sample complexity for small optimality gap ϵ , when assuming a generative model for policy evaluation.

1 Introduction

We consider a discrete time Markov decision process (MDP) denoted by the quintuple $\mathcal{M} = (\mathcal{S}, \mathcal{A}, \mathcal{P}, C, \gamma)$, where \mathcal{S} is the finite state space, \mathcal{A} is the finite action space, $\mathcal{P} : \mathcal{S} \times \mathcal{A} \rightarrow \Delta_{|\mathcal{S}|}$ is the transition kernel, $c : \mathcal{S} \times \mathcal{A} \rightarrow \mathbb{R}$ is the cost function satisfying $|c(s, a)| \leq C$ for all $(s, a) \in \mathcal{S} \times \mathcal{A}$, and $\gamma \in (0, 1)$ is the discount factor. Here $\Delta_{|\mathcal{X}|} = \left\{ p \in \mathbb{R}^{|\mathcal{X}|} : \sum_{i=1}^{|\mathcal{X}|} p_i = 1, p_i \geq 0 \right\}$ denotes the standard simplex defined over finite set \mathcal{X} .

A randomized, stationary policy is a mapping $\pi : \mathcal{S} \rightarrow \Delta_{|\mathcal{A}|}$, and we denote the set of all such policies by Π . At any timestep t , the policy explicitly governs what action to be made given the current state s_t , by $a_t \sim \pi(\cdot|s_t)$. Then a cost $c(s_t, a_t)$ is incurred, followed by the transition to the next state $s_{t+1} \sim \mathcal{P}(\cdot|s_t, a_t)$. The decision process is then repeated iteratively at future timesteps.

For a given policy π , we measure its performance by its value function $V^\pi : \mathcal{S} \rightarrow \mathbb{R}$,

$$V^\pi(s) = \mathbb{E} \left[\sum_{t=0}^{\infty} \gamma^t c(s_t, a_t) \mid s_0 = s, a_t \sim \pi(\cdot|s_t), s_{t+1} \sim \mathcal{P}(\cdot|s_t, a_t) \right].$$

We also define its state-action value function (Q-function) $Q^\pi : \mathcal{S} \times \mathcal{A} \rightarrow \mathbb{R}$, as

$$Q^\pi(s, a) = \mathbb{E} \left[\sum_{t=0}^{\infty} \gamma^t c(s_t, a_t) \mid s_0 = s, a_0 = a, a_t \sim \pi(\cdot|s_t), s_{t+1} \sim \mathcal{P}(\cdot|s_t, a_t) \right].$$

The planning objective of the MDP \mathcal{M} is to find an optimal policy π^* that minimizes the value for every state $s \in \mathcal{S}$ simultaneously,

$$V^{\pi^*}(s) \leq V^\pi(s), \quad \forall s \in \mathcal{S}, \quad \forall \pi \in \Pi. \quad (1)$$

Accordingly, we define the optimal value functions as

$$V^*(s) = \min_{\pi \in \Pi} V^\pi(s), \quad Q^*(s, a) = \min_{\pi \in \Pi} Q^\pi(s, a), \quad \forall s \in \mathcal{S}, a \in \mathcal{A}. \quad (2)$$

[†]H. Milton Stewart School of Industrial and Systems Engineering, Georgia Institute of Technology, Atlanta, GA, 30332. (E-mail: yli939@gatech.edu).

[‡]H. Milton Stewart School of Industrial and Systems Engineering, Georgia Institute of Technology, Atlanta, GA, 30332. (E-mail: touzhaog@gatech.edu).

[§]H. Milton Stewart School of Industrial and Systems Engineering, Georgia Institute of Technology, Atlanta, GA, 30332. (E-mail: george.lan@isye.gatech.edu).

It should be noted that the existence of an optimal policy π^* simultaneously minimizing the value for every state is well known in literature [21]. Hence we can succinctly reformulate (1) into a single-objective optimization problem

$$\min_{\pi} \mathbb{E}_{\rho} [V^{\pi}(s)], \quad \text{s.t. } \pi(\cdot|s) \in \Delta_{|\mathcal{A}|}, \forall s \in \mathcal{S}, \quad (3)$$

where ρ can be arbitrary distribution defined over \mathcal{S} . As pointed out by recent literature (e.g. [18]), setting ρ as the stationary state distribution induced by an optimal policy π^* of (1), denoted by ν^* , can simplify the analyses of various algorithms. In such a case, (3) becomes

$$\min_{\pi} \{f(\pi) := \mathbb{E}_{\nu^*} [V^{\pi}(s)]\}, \quad \text{s.t. } \pi(\cdot|s) \in \Delta_{|\mathcal{A}|}, \forall s \in \mathcal{S}. \quad (4)$$

It is also worth mentioning that the minimizers of (1) can be a proper subset of the solution set of (4) when ν^* does not have the full support on the state space. To assert equivalence, one needs the existence of an optimal policy π^* of (1) that has a fully supported steady state distribution (see Assumption 1). In addition, even though the objective $f(\pi)$ explicitly depends on the unknown distribution ν^* , the methods to be discussed in this paper do not require the information of ν^* .

Recently, there has been a rise of interests in designing first-order methods for searching the optimal policy of (1) and (4), which we term as the policy gradient methods in our ensuing discussions. Sublinear convergence of policy gradient methods with a constant stepsize has been discussed in [1], and strengthened to asymptotic linear convergence in [12]. Linear convergence with constant stepsize is shown in [4] for solving entropy-regularized MDPs, when applying the natural policy gradient method [11] with softmax policy parameterization. Until recently, linear convergence of policy gradient method is established in [13, 26] for MDPs without any regularization, by proposing an approximate policy mirror descent method. Policy gradient methods with a block update rule, which evaluate and update the policy for only a subset of states, have been also proposed in [14], with similar convergence properties. With softmax policy parameterization, linear convergence of natural policy gradient method is also established in [12] when using adaptive stepsizes. For large state or action space, there has also been fruitful developments on extending policy gradient methods to learning the optimal policy within a parameterized function class [23, 17]. The convergence of these methods, on the other hand, seems more subtle than their tabular counterparts, and often requires additional technical assumptions on the function class and the underlying MDP [1, 18, 25, 3].

This work complements existing developments of policy gradient methods with the following focuses:

- (1) *Convergence rate*: Linear convergence of policy gradient methods has been discussed in prior literature [14, 12, 26], while existing superlinear convergence result often requires strong algorithm-dependent assumptions [21, 12]. It is unclear whether policy gradient methods converge superlinearly in general settings.
- (2) *Convergence of the policy*: It is not clear whether policy gradient methods exhibits last-iterate convergence of the policy, as existing developments mostly establish convergence in optimality gap. To the best of our knowledge, it seems that only asymptotic subsequence convergence to an optimal policy can be claimed based on the value convergence [13, 26, 1]. More importantly, even when policy gradient methods exhibit last-iterate convergence, it is also unclear whether one can characterize the particular policy that the method converges to, as the number of optimal policies can be infinite.
- (3) *Sample complexity*: Prior developments have shown $\tilde{O}(|\mathcal{S}||\mathcal{A}|/\epsilon^2)$ sample complexity for policy gradient methods [13]. Superlinearly convergence, if holds true for policy gradient methods, naturally begs the question whether an improved sample complexity can be achieved as an implication of faster convergence.

Our contributions mainly exist in the following aspects. Firstly, we develop the homotopic policy mirror descent (HPMD) method and establish its global linear convergence, together with its local superlinear convergence in both the policy and the value with an order of γ^{-2} , provided the existence of an optimal policy with fully supported stationary state distribution. Moreover, our analysis shows that the superlinear convergence kicks in fairly quickly – after no later than $\log(1/\Delta^*)$ number of iterations, where Δ^* is defined via the gap value of the optimal Q-function.

Secondly, we show that the HPMD method exhibits last-iterate convergence of the policy while using exponentially increasing stepsizes. In addition, despite the existence of potentially infinitely many optimal policies, the policy converges exactly to the optimal policy $\pi_{\mathcal{U}}^*$ with the maximum entropy, given by

$$\pi_{\mathcal{U}}^*(a|s) := \begin{cases} 1/|\mathcal{A}^*(s)|, & a \in \mathcal{A}^*(s), \\ 0, & a \notin \mathcal{A}^*(s), \end{cases}$$

where $\mathcal{A}^*(s) = \operatorname{argmin}_{a \in \mathcal{A}} Q^*(s, a)$. This appears to be surprising since any Dirac measure δ_a with support $a \in \mathcal{A}^*(s)$ suffices to become the optimal strategy for every state s (see Lemma 3.1), and there is no regularization in the policy optimization objective (4) to enforce the limiting policy to explore every action $a \in \mathcal{A}^*(s)$ equally. We term such an observation as the implicit regularization of HPMD.

Thirdly, we develop the stochastic homotopic policy mirror descent (SHPMD) method and establish general conditions on the noise that ensures the similar convergence properties as deterministic HPMD. We further demonstrate that by exploiting superlinear convergence of SHPMD method, an improved sample complexity can be achieved when searching for the near-optimal policy with small optimality gap ϵ . Informally, with the gap quantity of the MDP denoted by $\Delta^*(\mathcal{M})$, the total number of sample complexity $N(\epsilon)$ can be bounded by

$$N(\epsilon) = \begin{cases} \tilde{\mathcal{O}}(|\mathcal{S}||\mathcal{A}|/\epsilon^2), & \epsilon \geq \epsilon_0 = \Theta(\Delta^*(\mathcal{M})^{3/2}), \\ \tilde{\mathcal{O}}(|\mathcal{S}||\mathcal{A}|/\epsilon_0^2 \left[1 + \log^2(1/\epsilon) + \log_{1/\gamma}(\log(\frac{1}{\epsilon}))\right]), & \epsilon < \epsilon_0. \end{cases}$$

To the best of our knowledge, our findings appear to be new in the literature of policy gradient methods. In particular, the superlinear convergence is established without prior strict assumptions [21, 12], and is established for both the policy and the value function. The last-iterate convergence of the policy has also not been established prior to our developments, neither does an exact characterization of the limiting policy (implicit regularization). Finally, the improved sample complexity by exploiting superlinear convergence of stochastic HPMD also appears to be the first of its kind.

To illustrate our theoretical findings, we apply HPMD to the benchmark GridWorld environment of size 10 (i.e., $|\mathcal{S}| = 100$)¹. The experiment findings are reported in Figure 1.




Figure 1: Convergence of optimality gap and policy parameters of the HPMD method on a randomly generated GridWorld MDP with grid size 10.

As can be seen from Figure 1a, the convergence of value optimality gap seems to consist of two phases. The first phase demonstrates linear convergence, as prescribed by its global convergence analysis. At the end of the first phase, the convergence clearly speeds up and the optimality gap improves much faster in the ensuing second phase. In addition, Figure 1b shows the convergence of HPMD to the set of optimal policies (see also Lemma 3.1 for a folklore characterization of optimal policies). Similar to the convergence in value, the convergence of the policy also demonstrates a phase transition, where a much faster convergence takes place in the second phase. Finally, Figure 1c shows that $\min_{a \in \mathcal{A}^*(s)} \pi_k(a|s)$ always converges to a value bounded away from zero, meaning that $\pi_k(a|s)$ is sufficiently exploring every action within $\mathcal{A}^*(s)$.

The rest of the paper is organized as follows. Section 2 introduces the deterministic HPMD method and establishes its global linear convergence. In Section 3, we establish the local superlinear convergence of both policy and value function, the last-iterate convergence of the policy, and the associated implicit regularization effect of the HPMD method. In Section 4, we extend our findings to the stochastic deterministic setting, and demonstrate that an improved sample complexity can be achieved by exploiting the local superlinear convergence. Discussions and concluding remarks are made in Section 5.

1.1 Notation and Terminology

Note that although the optimal value functions V^* and Q^* of (1) are unique, the optimal policies are not in general, and the MDP can have infinitely many optimal policies. We use Π^* to denote the set of optimal stationary policies

¹For more environment details, we refer readers to [14], which adopt the same experiment setup. Note that in Figure 1a, we vary the discount factor while keeping other defining parameters of the MDP unchanged.

Algorithm 1 The homotopic policy mirror descent (HPMD) method

Input: Initial policy π^0 , nonnegative parameters $\{\tau_k\}_{k \geq 0}$, and stepsizes $\{\eta_k\}_{k \geq 0}$.
for $k = 0, 1, \dots$ **do**
 Update policy:

$$\pi_{k+1}(\cdot|s) = \operatorname{argmin}_{p(\cdot|s) \in \Delta_{|\mathcal{A}|}} \eta_k \left[\langle Q^{\pi_k}(s, \cdot), p(\cdot|s) \rangle + \tau_k D_{\pi_0}^p(s) \right] + D_{\pi_k}^p(s), \quad \forall s \in \mathcal{S}$$

end for

of (1). We denote the stationary state distribution of a policy π by ν^π . In addition, we also define the discounted state visitation measure d_s^π induced by policy π as

$$d_s^\pi(s') = \sum_{t=0}^{\infty} \gamma^t \mathbb{P}^\pi(s_t = s' | s_0 = s), \quad \forall s \in \mathcal{S}, \quad (5)$$

where $\mathbb{P}^\pi(s_t = s' | s_0 = s)$ denotes the probability of reaching state s' at timestep t when starting at state s and following policy π .

For any distribution μ defined over set \mathcal{X} , we denote its support as $\operatorname{supp}(\mu) = \{x : \mu(x) > 0\}$, and its entropy as $\mathcal{H}(\mu) = -\sum_{x \in \operatorname{supp}(\mu)} \mu(x) \log \mu(x)$. For any function defined over a finite set \mathcal{X} , we denote its minimizers as $\operatorname{argmin}_{x \in \mathcal{X}} f(x) = \{x : f(x) = \min_{x' \in \mathcal{X}} f(x')\}$. For any pair of policy (π, π') , we define their ℓ_1 distance to be $\|\pi - \pi'\|_1 := \max_{s \in \mathcal{S}} \sum_{a \in \mathcal{A}} |\pi(a|s) - \pi'(a|s)|$. Accordingly, we define the ℓ_1 distance between a policy π and a policy set Π^* as $\operatorname{dist}_{\ell_1}(\pi, \Pi^*) := \inf_{\pi' \in \Pi^*} \|\pi - \pi'\|_1$.

2 Homotopic Policy Mirror Descent: Linear Convergence

We introduce the deterministic homotopic policy mirror descent (HPMD) method, and establish its global linear convergence. The HPMD method (Algorithm 1) minimizes the objective (4) by performing the following update at every iteration $k \geq 0$,

$$\pi_{k+1}(\cdot|s) = \operatorname{argmin}_{p(\cdot|s) \in \Delta_{|\mathcal{A}|}} \eta_k \left[\langle Q^{\pi_k}(s, \cdot), p(\cdot|s) \rangle + \tau_k D_{\pi_0}^p(s) \right] + D_{\pi_k}^p(s), \quad \forall s \in \mathcal{S}, \quad (6)$$

where $D_q^p(s) = \sum_{a \in \mathcal{A}} p(a|s) \log \left(\frac{p(a|s)}{q(a|s)} \right)$ denotes the Kullback–Leibler (KL) divergence, π^0 denotes the uniform policy.

In a nutshell, the HPMD method can be considered as a simplification of the approximate policy mirror descent (APMD) method proposed in [13] and extended in [26], by dropping the need for evaluating the perturbed state-action value function $Q_{\tau_k}^{\pi_k}$, defined by

$$Q_\tau^\pi(s, a) = \mathbb{E} \left[\sum_{t=0}^{\infty} \gamma^t (r(s_t, a_t) + \tau D_{\pi_0}^\pi(s_t)) \mid (s_0, a_0) = (s, a), a_t \sim \pi(\cdot|s_t), s_t \sim \mathcal{P}(\cdot|s_{t-1}, a_{t-1}) \right], \quad \tau \geq 0.$$

Thus aside from the conceptual simplicity, HPMD also allow an easier implementation without the requirement to compute the regularization term when evaluating the policy.

With proper specification of $\{\tau_k\}_{k \geq 0}$ and $\{\eta_k\}_{k \geq 0}$, we show that HPMD converges linearly to the minimum of the policy optimization objective (4).

Theorem 2.1. *By choosing $1 + \eta_k \tau_k = 1/\gamma$ and $\eta_k = \gamma^{-2(k+1)}$ in the HPMD method, then for any iteration $k \geq 1$, HPMD produces policy π_k with*

$$f(\pi_k) - f(\pi^*) \leq \gamma^k \left(f(\pi_0) - f(\pi^*) + \frac{4 \log |\mathcal{A}|}{1 - \gamma} \right). \quad (7)$$

Theorem 2.1 implies that to find an ϵ -optimal policy π_k such that $f(\pi_k) - f(\pi^*) \leq \epsilon$, HPMD requires at most $\log(1/\epsilon)$ number of iterations, which seems to describe the initial convergence behavior in Figure 1a faithfully. On the other hand, the linear convergence seems pessimistic when describing the convergence in the second stage observed in Figure 1a.

Moreover, unlike the scenario where the cost function includes a strongly convex regularization term [13, 4] and thus implying a unique optimal policy, the convergence of policy without strongly convex regularization seems less

investigated by the prior literature, as there can exist infinitely many optimal policies for the MDP. Meanwhile, as demonstrated Figure 1b, the convergence of policy seems fairly quick, and the limiting policy also demonstrated interesting behavior illustrated in Figure 1c – a nontrivial probability lower bound within $\mathcal{A}^*(s)$ for every state s .

In the next section, we will show that HPMD indeed converges faster than what linear convergence prescribes, and instead requires at most $\mathcal{O}(\min\{\log(1/\epsilon), \log(1/\Delta^*(\mathcal{M})) + \log\log(1/\epsilon)\})$ number of iterations, where $\Delta^*(\mathcal{M})$ is a gap quantity depending on the optimal state-action value function Q^* . The obtained iteration complexity naturally implies that the number of iterations first grows at the order of $\mathcal{O}(\log(1/\epsilon))$, as prescribed by the linear convergence. Then after the precision ϵ falls below $\mathcal{O}(\Delta^*(\mathcal{M}))$, the superlinear convergence kicks in and HPMD only needs an additional $\log\log(1/\epsilon)$ number of iterations. In addition, we also show that the policy converges to the set of optimal policies at a similar rate as the value function. More importantly, we will establish the last-iterate convergence of the policy, and conclude that the nontrivial probability lower bound observed in Figure 1c holds in general scenarios.

3 Local Superlinear Convergence and Implicit Regularization

Before stating our formal results, we make the following assumption throughout the remainder of the paper. As we have discussed in Section 1, Assumption 1 is needed to certify the single-objective policy optimization problem (4) being equivalent to the multi-objective planning problem of the MDP (1).

Assumption 1. *There exists $\pi^* \in \Pi^*$ such that ν^{π^*} has full support on \mathcal{S} . That is, $\text{supp}(\nu^{\pi^*}) = \mathcal{S}$. Accordingly, we define the distribution mismatch ratio $r := \gamma \max_{s \in \mathcal{S}, a \in \mathcal{A}} \max_{s' \in \mathcal{S}} \mathcal{P}(s'|s, a) / \nu^{\pi^*}(s')$.*

Since the state space \mathcal{S} is finite, one sufficient condition for Assumption (1) is that the Markov chain induced by π^* , with transition kernel $\mathbb{P}^{\pi^*}(s'|s) := \sum_{a \in \mathcal{A}} \mathbb{P}(s'|s, a)\pi^*(a|s)$, is irreducible. Such a condition is also satisfied by the GridWorld environment in Figure 1, since for any location on the 2D grid, there is always positive probability of reaching every neighboring gridpoint and thus any policy induces an irreducible Markov chain over \mathcal{S} .

It is also worth mentioning that Assumption 1 seems not strictly necessary for obtaining the results in our ensuing discussions, and can be removed with a slightly modified analysis. We refer interested readers to Section 5 for related discussions.

To proceed, we define $\Delta^*(s) = \min_{a \notin \mathcal{A}^*(s)} Q^*(s, a) - \min_{a \in \mathcal{A}} Q^*(s, a)$ if $\mathcal{A}^*(s) \neq \mathcal{A}$, and $\Delta^*(s) = \infty$ otherwise. Then the gap function of the MDP \mathcal{M} is defined as $\Delta^*(\mathcal{M}) = \min_{s \in \mathcal{S}} \Delta^*(s)$. Note that if $\Delta^*(\mathcal{M}) = \infty$, then any policy is an optimal policy (see also Lemma 3.1)

We begin by characterizing the parameter convergence of the HPMD method in the following theorem.

Theorem 3.1 (Policy Convergence). *Whenever $\Delta^*(\mathcal{M}) < \infty$, then with $1 + \eta_k \tau_k = 1/\gamma$ and $\eta_k = \gamma^{-2(k+1)}$, the iterate $\{\pi_k\}_{k \geq 0}$ generated by HPMD satisfies*

$$\text{dist}_{\ell_1}(\pi_k, \Pi^*) \leq 2C_\gamma |\mathcal{A}| \exp\left(-\frac{\Delta^*(\mathcal{M})}{2} \gamma^{-2k-1}\right),$$

for any iteration $k \geq K_1 := \lceil 3 \log_\gamma \left(\frac{\Delta^*(\mathcal{M})(1-\gamma)}{2r(4 \log |\mathcal{A}| + C)} \right) \rceil$, and $C_\gamma := \exp\left(\frac{2C\gamma^2}{(1-\gamma^3)(1-\gamma)}\right)$.

By having the superlinear convergence of policy parameters, the convergence in the value function V^π follows by invoking the performance difference lemma [10]. Specifically, we have the following corollary.

Corollary 3.1. *For any policy π , let $\pi^* = \text{argmin}_{\pi' \in \Pi^*} \|\pi' - \pi\|_1$ denotes the its projection onto Π^* in ℓ_1 -norm. Then we have $V^\pi(s) - V^{\pi^*}(s) \leq \frac{C}{(1-\gamma)^2} \|\pi - \pi^*\|_1$. In particular, for $k \geq K_1$ where K_1 is defined as in Theorem 3.1, we have*

$$V^{\pi_k}(s) - V^{\pi^*}(s) \leq \frac{2C|\mathcal{A}|C_\gamma}{(1-\gamma)^2} \exp\left(-\frac{\Delta^*(\mathcal{M})}{2} \gamma^{-2k-1}\right), \quad \forall s \in \mathcal{S}. \quad (8)$$

Proof. We start with the performance difference lemma [10]. Let π^* be the projection of π_k onto Π^* in ℓ_1 -norm. Then

$$\begin{aligned} V^{\pi_k}(s) - V^{\pi^*}(s) &= \frac{1}{1-\gamma} \mathbb{E}_{s' \sim d_s^{\pi_k}} \langle Q^*(s', \cdot), \pi_k(\cdot|s') - \pi^*(\cdot|s') \rangle \\ &\leq \frac{1}{1-\gamma} \mathbb{E}_{s' \sim d_s^{\pi_k}} \|Q^*(s', \cdot)\|_\infty \|\pi_k(\cdot|s') - \pi^*(\cdot|s')\|_1 \end{aligned}$$

$$\leq \frac{C}{(1-\gamma)^2} \|\pi_k - \pi^*\|_1,$$

where the last inequality uses the fact that $\|Q^*(\cdot, \cdot)\|_\infty \leq 1/(1-\gamma)$ and the definition of $\|\pi_k - \pi^*\|_1$. The claim then follows immediately after applying Theorem 3.1. \square

We briefly compare our result with related literature. Existing superlinear convergence for policy gradient methods have only been established for the value function, by exploiting the relations of policy gradient methods and policy iteration [12, 2], together with the superlinear convergence of the latter method [21]. However, such results assume strong assumptions on bounding the difference between the transition kernel induced from the generated policy π_k and the optimal policy, by the optimality gap of the policy π_k (see Theorem 6.4.8, [21]), which seems difficult to verify as the assumption is algorithm-dependent in nature.

On the other hand, our obtained result requires only the existence of an optimal policy with fully-supported stationary state distribution. The result establishes local superlinear convergence of policy parameters after $\log(1/\Delta^*(\mathcal{M}))$ number of iterations, which also parallels the algorithmic behavior of Newton's method in the optimization literature [19].

It is also worth mentioning that by extending similar arguments, the local superlinear convergence can also be established for the approximate policy mirror descent (APMD) method developed in [13, 26].

Our next result shows that the implicit exploration effect in Figure 1c is a general algorithmic property of HPMD. Before we proceed, we first recall a folklore result that characterizes the set of the optimal policies.

Lemma 3.1 (Characterization of Optimal Policies). *Let $Q^* \in \mathbb{R}^{|\mathcal{S}| \times |\mathcal{A}|}$ denote the optimal Q -function. Then the set of optimal policies Π^* is given by:*

$$\Pi^* = \left\{ \pi \in \Pi : \text{supp}(\pi(\cdot|s)) \subseteq \underset{a \in \mathcal{A}}{\text{argmin}} Q^*(s, a), \forall s \in \mathcal{S} \right\}.$$

From Lemma 3.1, an immediate consequence is that the optimal policy with the maximal entropy for every state, denoted by π_U^* , takes the form of

$$\pi_U^*(a|s) := \begin{cases} 1/|\mathcal{A}^*(s)|, & a \in \mathcal{A}^*(s), \\ 0, & a \notin \mathcal{A}^*(s). \end{cases}$$

We proceed to show that the HPMD converges exactly to this particular optimal policy, even though there is no regularization in its objective function (4).

Theorem 3.2 (Implicit Regularization). *We have the last-iterate convergence for the policy sequence $\{\pi_k\}_{k \geq 0}$ generated by HPMD, i.e., there exists $\pi^* \in \Pi$ such that*

$$\lim_{k \rightarrow \infty} \pi_k = \pi^*. \quad (9)$$

Furthermore, the converged policy equals to the optimal policy with the maximal entropy for every state $s \in \mathcal{S}$. That is

$$\lim_{k \rightarrow \infty} \pi_k(a|s) = \pi_U^*(a|s) := \begin{cases} 1/|\mathcal{A}^*(s)|, & a \in \mathcal{A}^*(s), \\ 0, & a \notin \mathcal{A}^*(s). \end{cases} \quad (10)$$

In particular, for any $0 < \epsilon < 1$, we have $\|\pi_k - \pi_U^*\|_\infty \leq \epsilon$, whenever

$$k \geq 2\bar{K}_1 + \log_\gamma \left(\frac{4r(4 \log |\mathcal{A}| + C)}{\epsilon(1-\gamma)(1-\gamma^2)\gamma} \right) + 2 \log_\gamma \left(\frac{8\gamma C |\mathcal{A}| C_\gamma}{\epsilon(1-\gamma^{1/2})(1-\gamma)\gamma} \right) + \log_\gamma \left(\frac{2 \log(C_\gamma |\mathcal{A}|/\epsilon)}{\Delta^*(\mathcal{M})} \right),$$

where $\bar{K}_1 = \min_k \left\{ k > K_1 : \Delta^*(\mathcal{M})\gamma^{-2k-1} \geq 5k \log(\frac{1}{\gamma}) \right\}$.

It is worth mentioning that aside from characterizing the limiting policy (10), we believe the last-iterate convergence of the policy (9) should be not considered trivial since (1) there can be infinitely many optimal policies; (2) the stepsize η_k is increasing in the HPMD method.

Theorem 3.2 is also related to [7], which shows that the actor-critic method produces policy iterate that has bounded Kullback–Leibler (KL) divergence to the optimal policy with maximal entropy π_U^* . Our differences mainly

exists in the following aspects: (1) We study general discounted MDP with finite state and action. On the other hand, [7] consider a slightly restrictive MDP class with a special linear transition kernel; (2) We establish exact convergence to the optimal policy with the maximal entropy π_U^* , instead of bounding the KL divergence $\text{KL}(\pi_k \|\pi_U^*)$ as in [7]; We refer interested readers to [7] for an excellent discussion on the benefits of maintaining a high-entropy policy throughout policy optimization process.

It is also worth pointing out that the implicit regularization of HPMD shares interesting connections with the homotopy method (i.e. regularization path) in statistics literature [27, 6, 20]. In particular, for an empirical risk minimization problem of the form $\min_{\theta \in \mathbb{R}^d} \mathcal{L}(\theta)$, where $\mathcal{L}(\theta)$ denotes the empirical risk on the training data, the regularization path $\{\Gamma(s) : s \geq 0\}$ is a path in \mathbb{R}^d , with $\Gamma(r)$ being a minimizer of the regularized problem $\min_{\theta \in \mathbb{R}^d} \mathcal{L}(\theta) + sR(\theta)$. Here $R(\cdot)$ denotes the regularization term to control the complexity. It is known that when training linear classifiers with linearly separable data and an exponentially-tailed loss, and $R(\theta)$ being ℓ_p -norm, then when taking $s \rightarrow 0$, the regularization path $\Gamma(r)$ converges in direction to the maximal margin (measured in ℓ_q -norm) linear classifier, where $1/p + 1/q = 1$ and $p \geq 1$ [22]. That is, the converged direction is given by

$$\min_{\theta} \|\theta\|_p, \text{ s.t. } y_i x_i^\top \theta \geq 1, \quad (11)$$

a minimal ℓ_p -norm solution satisfying separation the constraint (note that there are infinitely many feasible solutions).

Similarly, HPMD can be viewed as solving a sequence of regularized MDP problems with diminishing regularizations. In particular, define the regularized objective as

$$\min_{\pi} \{f_{\tau}(\pi) := \mathbb{E}_{\nu^*} [V_{\tau}^{\pi}(s)]\}, \text{ s.t. } \pi(\cdot|s) \in \Delta_{|\mathcal{A}|}, \forall s \in \mathcal{S},$$

where the regularized value function V_{τ}^{π} is defined as

$$V_{\tau}^{\pi}(s) = \mathbb{E} \left[\sum_{t=0}^{\infty} \gamma^t (c(s_t, a_t) + \tau D_{\pi_0}^{\pi}(s_t)) \mid s_0 = s, a_t \sim \pi(\cdot|s_t), s_{t+1} \sim \mathcal{P}(\cdot|s_t, a_t) \right].$$

Then HPMD can be viewed as solving the regularized objective f_{τ_n} *approximately* using only *one* step of policy gradient update. As Theorem 3.2 suggests, the HPMD method yields a solution π_U^* that is the solution of problem

$$\min_{\pi \in \Pi} \sum_{s \in \mathcal{S}} -\mathcal{H}(\pi(\cdot|s)), \text{ s.t. } V^{\pi}(s) \leq V^{\pi'}(s), \forall s \in \mathcal{S}, \forall \pi' \in \Pi. \quad (12)$$

Since the constraint in (12) states the feasible solution π to be the optimal policy of the original MDP (1), (12) in general also has infinitely many feasible solutions. By comparing (11) and (12), and note that $-\mathcal{H}(\pi(\cdot|s)) = D_{\pi_0}^{\pi}$, it should be clear that the implicit regularization of HPMD parallels the large-margin property of the homotopy method, in the sense that both seeks the minimal-complexity solution (measured by the negative entropy and ℓ_p -norm, respectively) among all feasible candidates.

Moreover, it is worth mentioning that implicit regularization effects of first order methods have also been extensively studied for empirical risk minimization problems [9, 24, 5, 15, 16], including their intimate connections to the regularization path [8]. To the best of our knowledge, Theorem 3.2 seems to be the first result on characterizing implicit regularization effects of first order methods in the RL literature.

4 Stochastic HPMD: Improved Sample Complexity

In this section, we introduce the stochastic HPMD (SHPMD) method, and show similar convergence properties as the deterministic HPMD method. In addition, we will establish that by exploiting the local superlinear convergence of SHPMD, one can improve upon the sample complexity of $\mathcal{O}(|\mathcal{S}||\mathcal{A}|/\epsilon^2)$, when searching for a near-optimal policy.

The SHPMD method (Algorithm 2) minimizes the objective (4) by performing the following update at every iteration $k \geq 0$,

$$\pi_{k+1}(\cdot|s) = \underset{p(\cdot|s) \in \Delta_{|\mathcal{A}|}}{\text{argmin}} \eta_k \left[\langle Q^{\pi_k, \xi_k}(s, \cdot), p(\cdot|s) \rangle + \tau_k D_{\pi_0}^p(s) \right] + D_{\pi_k}^p(s), \quad \forall s \in \mathcal{S}. \quad (13)$$

Different from the deterministic HPMD method, SHPMD uses a stochastic estimate Q^{π_k, ξ_k} of the state-action value function Q^{π_k} to perform the policy update. We impose the following conditions on the stochastic estimate $Q^{\pi_k, \xi_k} \in \mathbb{R}^{|\mathcal{S}| \times |\mathcal{A}|}$:

$$\mathbb{E}_{\xi_k} Q^{\pi_k, \xi_k} = \overline{Q}^{\pi_k}, \quad (14)$$

Algorithm 2 The stochastic homotopic policy mirror descent (SHPMD) method

Input: Initial policy π^0 , nonnegative parameters $\{\tau_k\}_{k \geq 0}$, and stepsizes $\{\eta_k\}_{k \geq 0}$.
for $k = 0, 1, \dots$ **do**
 Update policy:

$$\pi_{k+1}(\cdot|s) = \underset{p(\cdot|s) \in \Delta_{|\mathcal{A}|}}{\operatorname{argmin}} \eta_k \left[\langle Q^{\pi_k, \xi_k}(s, \cdot), p(\cdot|s) \rangle + \tau_k D_{\pi_0}^p(s) \right] + D_{\pi_k}^p(s), \quad \forall s \in \mathcal{S}$$

end for

$$\left\| \bar{Q}^{\pi_k} - Q^{\pi_k} \right\|_{\infty} \leq \varepsilon_k, \quad (15)$$

$$\mathbb{E} \left\| Q^{\pi_k, \xi_k} - Q^{\pi_k} \right\|_{\infty}^2 \leq \sigma_k^2, \quad (16)$$

To obtain such an estimate Q^{π_k, ξ_k} , we consider the simple scenario where we have the access to a generative model of the MDP. That is, for any state-action pair $(s, a) \in \mathcal{S} \times \mathcal{A}$, we can generate M_k independent trajectories, each with length T_k , by following policy π_k starting from (s, a) . The i -th trajectory, denoted by $\zeta_k^i(s, a)$, takes the form of $\zeta_k^i(s, a) = \{(s_0^i = s, a_0^i = a), (s_1^i, a_1^i), \dots, (s_{T_k-1}^i, a_{T_k-1}^i)\}$.

Let us denote the set of random variables for constructing the stochastic estimate as $\xi_k = \{\zeta_k^i(s, a), s \in \mathcal{S}, a \in \mathcal{A}, i \in [M_k]\}$. Then the stochastic estimator Q^{π_k, ξ_k} is defined by

$$Q^{\pi_k, \xi_k}(s, a) = \frac{1}{M_k} \sum_{i=1}^{M_k} \sum_{t=0}^{T_k-1} \gamma^t c(s_t^i, a_t^i), \quad s \in \mathcal{S}, a \in \mathcal{A}.$$

We first show that with certain conditions on $\{(\sigma_k, \varepsilon_k)\}_{k \geq 0}$, together with proper choices of $\{(\tau_k, \eta_k)\}_{k \geq 0}$, then SHPMD converges linearly to the minimum of objective (4).

Theorem 4.1. Take $1 + \eta_k \tau_k = 1/\gamma$ and $\eta_k = \gamma^{-(k+1)/2} \sqrt{\log|\mathcal{A}|}$ in the SHPMD method. Suppose

$$\sigma_k = \gamma^{(k+1)/2}, \quad \varepsilon_k = \gamma^{3(k+1)/4} \quad (17)$$

holds in (15), (16), then for any iteration $k \geq 1$, SHPMD produces policy π_k satisfying

$$\mathbb{E}[f(\pi_k) - f(\pi^*)] \leq \gamma^{k/2} \frac{6\sqrt{\log|\mathcal{A}|} + C}{(1-\gamma)(1-\gamma^{1/2})\gamma}, \quad \forall k \geq 1. \quad (18)$$

We then show that with proper choices of $\{(M_k, T_k)\}_{k \geq 0}$, condition (17) is satisfied and SHPMD converges linearly while achieving an $\mathcal{O}(|\mathcal{S}||\mathcal{A}|/\varepsilon^2)$ sample complexity for finding an ε -optimal policy.

Theorem 4.2. Take $1 + \eta_k \tau_k = 1/\gamma$ and $\eta_k = \gamma^{-(k+1)/2} \sqrt{\log|\mathcal{A}|}$ in the SHPMD method. Suppose at each iteration of SHPMD, the number of independent trajectories M_k and the trajectory length T_k satisfies

$$T_k \geq \frac{3(k+1)}{4} + \log_{\gamma} \left(\frac{1-\gamma}{2C} \right), \quad M_k \geq \frac{4C^2}{(1-\gamma)^2} \gamma^{-(k+1)}. \quad (19)$$

Then for any $\varepsilon > 0$, the SHPMD method finds a policy π_k with $\mathbb{E}[f(\pi_k) - f(\pi^*)] \leq \varepsilon$ in $k = 2 \log_{\gamma} \left(\frac{\varepsilon(1-\gamma)}{6\sqrt{\log|\mathcal{A}|+C}} \right)$, number of iterations. In addition, the total number of samples required by SHPMD can be bounded by

$$\mathcal{O} \left(\frac{|\mathcal{S}||\mathcal{A}| \log_{\gamma}(\varepsilon(1-\gamma))}{(1-\gamma)^5 (1-\gamma^{1/2})^2 \varepsilon^2} \right). \quad (20)$$

Similar to the deterministic HPMD method, the policy π_k in SHPMD also exhibits local superlinear convergence to the set of optimal policies, despite using noisy first order information with linearly decaying noise.

Theorem 4.3. Whenever $\Delta^*(\mathcal{M}) < \infty$, under the same setting as in Theorem 4.1, the iterate $\{\pi_k\}_{k \geq 0}$ generated by SHPMD satisfies

$$\mathbb{E}[\operatorname{dist}_{\ell_1}(\pi_k, \Pi^*)] \leq 2C_{\gamma}^n |\mathcal{A}| \exp \left(-\frac{\Delta^*(\mathcal{M}) \sqrt{\log|\mathcal{A}|}}{2} \gamma^{-k/2} \right), \quad \forall k \geq K_2,$$

for any iteration $k \geq K_2 := \lceil 3 \log_{\gamma} \left(\frac{\Delta^*(\mathcal{M})(1-\gamma)}{2r(6\log|\mathcal{A}|+C)} \right) \rceil$, and $C_{\gamma}^n := \exp \left(\frac{(2C+1)\gamma^2 \sqrt{\log|\mathcal{A}|}}{(1-\gamma)(1-\gamma^{3/4})} \right)$. On the other hand, if $\Delta^*(\mathcal{M}) = \infty$, then any policy is an optimal policy.

Similarly to Corollary 3.1, we can also establish the local superlinear convergence of the value function.

Corollary 4.1. *For any $k \geq K_2$ where K_2 is defined as in Theorem 4.3, then under the same setting as in Theorem 4.1, the iterate π_k in the SHPMD method satisfies*

$$\mathbb{E} \left[V^{\pi_k}(s) - V^{\pi^*}(s) \right] \leq \frac{2C|\mathcal{A}|C_\gamma^n}{(1-\gamma)^2} \exp \left(-\frac{\Delta^*(\mathcal{M})\sqrt{\log|\mathcal{A}|}}{2} \gamma^{-k/2} \right), \quad \forall s \in \mathcal{S}. \quad (21)$$

By exploiting the superlinear convergence above, we can further improve the sample complexity of the SHPMD method when searching for a policy with optimality gap smaller than $\mathcal{O}(\Delta^*(\mathcal{M})^{3/2})$.

Theorem 4.4 (Improved Sample Complexity). *By choosing $\{(\tau_k, \eta_k, M_k, T_k)\}_{k \geq 0}$ as prescribed in Theorem 4.2, for any precision $\epsilon > 0$, the total number of iterations $k(\epsilon)$ for finding an ϵ -optimal policy $\mathbb{E} [f(\pi_{k(\epsilon)}) - f(\pi^*)] \leq \epsilon$ can be bounded by*

$$k(\epsilon) = \begin{cases} \mathcal{O}(\log_\gamma(\epsilon(1-\gamma))), & \epsilon \geq \epsilon_0, \\ \mathcal{O}\left(\log_\gamma(\epsilon_0(1-\gamma)) + \log_\gamma\left(\frac{\Delta^*(\mathcal{M})\sqrt{\log|\mathcal{A}|}}{\log(C_\gamma^n/\epsilon)}\right)\right), & \epsilon < \epsilon_0. \end{cases}$$

In addition, the total number of sample complexity $N(\epsilon)$ can be bounded by

$$N(\epsilon) = \begin{cases} \mathcal{O}\left(\frac{|\mathcal{S}||\mathcal{A}|\log_\gamma(\epsilon(1-\gamma))}{(1-\gamma)^5(1-\gamma^{1/2})^2\epsilon^2}\right), & \epsilon \geq \epsilon_0, \\ \mathcal{O}\left(\frac{|\mathcal{S}||\mathcal{A}|\log_\gamma(\epsilon(1-\gamma))}{(1-\gamma)^5(1-\gamma^{1/2})^2\epsilon_0^2} \left[1 + \frac{\log^2(1/\epsilon)}{\log|\mathcal{A}|\Delta^*(\mathcal{M})^2} + \log_{1/\gamma}\left(\frac{\log(\frac{1}{\epsilon})}{\Delta^*(\mathcal{M})}\right)\right]\right), & \epsilon < \epsilon_0, \end{cases}$$

where $\epsilon_0 = \Delta^*(\mathcal{M})^{3/2}(1-\gamma)^{1/2} / \left[(1-\gamma^{1/2})\gamma\sqrt{6\log|\mathcal{A}| + C}\right]$.

By comparing Theorem 4.2 and Theorem 4.4, it is clear that the sample complexity of SHPMD is no worse than $\mathcal{O}(|\mathcal{S}||\mathcal{A}|/\epsilon^2)$ and strictly better when the optimality gap ϵ is small. Specifically, for $\epsilon \geq \epsilon_0 = \mathcal{O}(\Delta^*(\mathcal{M})^{3/2})$, the sample complexity of SHPMD grows at the order of $\mathcal{O}(|\mathcal{S}||\mathcal{A}|/\epsilon^2)$. Then for $\epsilon < \epsilon_0$, the sample complexity of SHPMD improves over $\mathcal{O}(|\mathcal{S}||\mathcal{A}|/\epsilon^2)$ by a factor of $\mathcal{O}(\epsilon_0^2 / [\epsilon^2(1 + \log^2(\frac{1}{\epsilon}) \log \log(\frac{1}{\epsilon}))])$.

We then proceed to show that with slightly smaller stepsizes, SHPMD also exhibits the last-iterate convergence of the policy almost surely. We state the formal result in an asymptotic fashion, while a non-asymptotic convergence characterization can also be obtained with essentially the same analysis.

Theorem 4.5 (Implicit Regularization). *If the parameter choice $\{(\eta_t, \tau_t, \epsilon_t, \sigma_t)\}_{t \geq 0}$ satisfies*

$$1 + \eta_t \tau_t = 1/\gamma, \sigma_t = \gamma^{t/2}, \eta_t \sigma_t = \gamma^{\beta t} \sqrt{\log|\mathcal{A}|}, \epsilon_t = \gamma^{\frac{3}{4}t}, \quad 0 < \beta < 1/2, \quad (22)$$

Then with probability 1, we have the last-iterate convergence for the policy sequence $\{\pi_k\}_{k \geq 0}$ generated by SHPMD, i.e., there exists $\pi^* \in \Pi$ such that $\lim_{k \rightarrow \infty} \pi_k = \pi^*$, a.s. Furthermore, the converged policy equals to the optimal policy with the maximal entropy for every state $s \in \mathcal{S}$. That is

$$\lim_{k \rightarrow \infty} \pi_k(a|s) = \pi_U^*(a|s) := \begin{cases} 1/|\mathcal{A}^*(s)|, & a \in \mathcal{A}^*(s), \\ 0, & a \notin \mathcal{A}^*(s), \end{cases} \quad a.s. \quad (23)$$

Finally, we show that for the particular choice of parameters in Theorem 4.5, the sample complexity of SHPMD with last-iterate convergence can be arbitrarily close to $\mathcal{O}(|\mathcal{S}||\mathcal{A}|/\epsilon^2)$ when taking $\beta \rightarrow 0$.

Theorem 4.6 (Sample Complexity with Last-iterate Convergence). *Let $1 + \eta_t \tau_t = 1/\gamma$ and $\eta_t = \gamma^{(\beta-1/2)t} \sqrt{\log|\mathcal{A}|}$ in the SHPMD method, with $0 < \beta < 1/2$. Suppose at each iteration, the number of independent trajectories M_k and the trajectory length T_k satisfies*

$$T_k \geq \frac{3(k+1)}{4} + \log_\gamma\left(\frac{1-\gamma}{2C}\right), M_k \geq \frac{4C^2}{(1-\gamma)^2} \gamma^{-(k+1)}.$$

Then for any $\epsilon > 0$, the SHPMD method finds an policy π_k with $\mathbb{E} [f(\pi_k) - f(\pi^*)] \leq \epsilon$ in $k = \frac{2}{1-2\beta} \log_\gamma\left(\frac{\epsilon(1-\gamma)(1-\gamma^{1/2})}{6\sqrt{\log|\mathcal{A}|+C}}\right)$, number of iterations. In addition, the total number of samples required by SHPMD can be bounded by

$$\mathcal{O}\left(\frac{|\mathcal{S}||\mathcal{A}|\log_\gamma(\epsilon(1-\gamma))}{(1-\gamma)^{3+2/(1-2\beta)}(1-\gamma^{1/2})^{2/(1-2\beta)}} \frac{1}{\epsilon^{2/(1-2\beta)}}\right).$$

We remark that similar to Theorem 4.4, the sample complexity for the SHPMD method with the last-iterate convergence of the policy can also be improved by exploiting its local superlinear convergence.

5 Discussions and Concluding Remarks

5.1 Dropping Assumption 1

We discuss the feasibility of retaining all obtained results without Assumption 1. As should be clear from our analysis in Appendix B, the key component for obtaining our results including the local superlinear convergence, last-iterate convergence of the policy, and the improved sample complexity, is the linear convergence of $V^{\pi_k}(s)$ to $V^*(s)$ for every state. To this end, it suffices to establish the linear convergence of the general weighted value objective

$$f_\rho(\pi) := \sum_{s \in \mathcal{S}} \rho(s) V^\pi(s)$$

for any distribution ρ defined over \mathcal{S} with full support. Clearly, $f_\rho(\pi)$ reduces to the policy optimization objective $f(\pi)$ defined in (4) by taking $\rho = \nu^*$. Establishing the global linear convergence for objective $f_\rho(\pi)$ follows the essentially the same analysis for the objective $f(\pi)$.

We illustrate such a procedure for the deterministic HPMD method. More specifically, in Appendix A.1, we take expectation w.r.t. $s \sim d_\rho^{\pi^*}$ on both sides of (31), instead of $s \sim \nu^*$, where $d_\rho^{\pi^*}(s) := \sum_{s' \in \mathcal{S}} \rho(s') d_{s'}^{\pi^*}(s)$. Then by reusing every other element developed in Appendix A.1, one can establish

$$f_\rho(\pi_k) - f_\rho(\pi^*) = \mathcal{O} \left(1 - (1 - \gamma) / \left\| d_\rho^{\pi^*} / \rho \right\|_\infty \right)^k,$$

where $\left\| d_\rho^{\pi^*} / \rho \right\|_\infty = \max_{s \in \mathcal{S}} d_\rho^{\pi^*}(s) / \rho(s)$. Note that the previous convergence recovers (7) when taking $\rho = \nu^*$. The same procedure can also be directly applied to adapt the analysis of the SHPMD method.

5.2 Concluding Remarks

In this paper, we introduce a new policy gradient method, named homotopic policy mirror descent (HPMD) and establish its global linear convergence property. We then show that the proposed HPMD can achieve local superlinear convergence under the condition that there is one optimal policy with fully supported stationary state distribution. In addition, HPMD exhibits last-iterate convergence of the policy, which is the first linearly converging policy gradient method with such a convergence property. We further demonstrate that the limiting policy of HPMD corresponds to the optimal policy with the maximal entropy, a property attributed to the implicit regularization effect of HPMD. Finally, we develop a stochastic version of HPMD and show similar convergence properties. By exploiting the local superlinear convergence, we further show that the stochastic HPMD method achieves a sample complexity that is strictly better than $\mathcal{O}(|\mathcal{S}||\mathcal{A}|/\epsilon^2)$ for small optimality gap ϵ .

References

- [1] Alekh Agarwal, Sham M Kakade, Jason D Lee, and Gaurav Mahajan. Optimality and approximation with policy gradient methods in markov decision processes. In *Conference on Learning Theory*, pages 64–66. PMLR, 2020.
- [2] Jalaj Bhandari and Daniel Russo. A note on the linear convergence of policy gradient methods. *arXiv preprint arXiv:2007.11120*, 2020.
- [3] Semih Cayci, Niao He, and R Srikant. Linear convergence of entropy-regularized natural policy gradient with linear function approximation. *arXiv preprint arXiv:2106.04096*, 2021.
- [4] Shicong Cen, Chen Cheng, Yuxin Chen, Yuting Wei, and Yuejie Chi. Fast global convergence of natural policy gradient methods with entropy regularization. *Operations Research*, 2021.
- [5] Suriya Gunasekar, Jason Lee, Daniel Soudry, and Nathan Srebro. Characterizing implicit bias in terms of optimization geometry. In *International Conference on Machine Learning*, pages 1832–1841. PMLR, 2018.
- [6] Trevor Hastie, Saharon Rosset, Robert Tibshirani, and Ji Zhu. The entire regularization path for the support vector machine. *Journal of Machine Learning Research*, 5(Oct):1391–1415, 2004.

- [7] Yuzheng Hu, Ziwei Ji, and Matus Telgarsky. Actor-critic is implicitly biased towards high entropy optimal policies. *arXiv preprint arXiv:2110.11280*, 2021.
- [8] Ziwei Ji, Miroslav Dudík, Robert E Schapire, and Matus Telgarsky. Gradient descent follows the regularization path for general losses. In *Conference on Learning Theory*, pages 2109–2136. PMLR, 2020.
- [9] Ziwei Ji and Matus Telgarsky. The implicit bias of gradient descent on nonseparable data. In *Conference on Learning Theory*, pages 1772–1798. PMLR, 2019.
- [10] Sham Kakade and John Langford. Approximately optimal approximate reinforcement learning. In *In Proc. 19th International Conference on Machine Learning*. Citeseer, 2002.
- [11] Sham M Kakade. A natural policy gradient. *Advances in neural information processing systems*, 14, 2001.
- [12] Sajad Khodadadian, Prakirt Raj Jhunjhunwala, Sushil Mahavir Varma, and Siva Theja Maguluri. On the linear convergence of natural policy gradient algorithm. *arXiv preprint arXiv:2105.01424*, 2021.
- [13] Guanghui Lan. Policy mirror descent for reinforcement learning: Linear convergence, new sampling complexity, and generalized problem classes. *arXiv preprint arXiv:2102.00135*, 2021.
- [14] Guanghui Lan, Yan Li, and Tuo Zhao. Block Policy Mirror Descent. *arXiv e-prints*, page arXiv:2201.05756, January 2022.
- [15] Yan Li, Ethan X Fang, Huan Xu, and Tuo Zhao. Implicit bias of gradient descent based adversarial training on separable data. 2020.
- [16] Yan Li, Caleb Ju, Ethan X Fang, and Tuo Zhao. Implicit regularization of bregman proximal point algorithm and mirror descent on separable data. *arXiv preprint arXiv:2108.06808*, 2021.
- [17] Timothy P Lillicrap, Jonathan J Hunt, Alexander Pritzel, Nicolas Heess, Tom Erez, Yuval Tassa, David Silver, and Daan Wierstra. Continuous control with deep reinforcement learning. *arXiv preprint arXiv:1509.02971*, 2015.
- [18] Boyi Liu, Qi Cai, Zhuoran Yang, and Zhaoran Wang. Neural trust region/proximal policy optimization attains globally optimal policy. 2019.
- [19] Jorge Nocedal and Stephen Wright. *Numerical optimization*. Springer Science & Business Media, 2006.
- [20] Mee Young Park and Trevor Hastie. L1-regularization path algorithm for generalized linear models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 69(4):659–677, 2007.
- [21] Martin L Puterman. *Markov decision processes: discrete stochastic dynamic programming*. John Wiley & Sons, 2005.
- [22] Saharon Rosset, Ji Zhu, and Trevor Hastie. Boosting as a regularized path to a maximum margin classifier. *The Journal of Machine Learning Research*, 5:941–973, 2004.
- [23] John Schulman, Sergey Levine, Pieter Abbeel, Michael Jordan, and Philipp Moritz. Trust region policy optimization. In *International conference on machine learning*, pages 1889–1897. PMLR, 2015.
- [24] Daniel Soudry, Elad Hoffer, Mor Shpigel Nacson, Suriya Gunasekar, and Nathan Srebro. The implicit bias of gradient descent on separable data. *The Journal of Machine Learning Research*, 19(1):2822–2878, 2018.
- [25] Lingxiao Wang, Qi Cai, Zhuoran Yang, and Zhaoran Wang. Neural policy gradient methods: Global optimality and rates of convergence. *arXiv preprint arXiv:1909.01150*, 2019.
- [26] Wenhao Zhan, Shicong Cen, Baihe Huang, Yuxin Chen, Jason D Lee, and Yuejie Chi. Policy mirror descent for regularized reinforcement learning: A generalized framework with linear convergence. *arXiv preprint arXiv:2105.11066*, 2021.
- [27] Peng Zhao and Bin Yu. Stagewise lasso. *The Journal of Machine Learning Research*, 8:2701–2726, 2007.

Supplementary Material for “Homotopic Policy Mirror Descent: Policy Convergence, Implicit Regularization, and Improved Sample Complexity”

We state the following folklore characterization of optimal policies of discounted MDP with finite state and action space, and present its proof for completeness.

Lemma (Restating Lemma 3.1). *Let $Q^* \in \mathbb{R}^{|\mathcal{S}| \times |\mathcal{A}|}$ denote the optimal Q -function. Then the set of optimal policies Π^* is given by:*

$$\Pi^* = \left\{ \pi \in \Pi : \text{supp}(\pi(\cdot|s)) \subseteq \underset{a \in \mathcal{A}}{\text{argmin}} Q^*(s, a), \forall s \in \mathcal{S} \right\}.$$

Proof. We first show that any $\pi^* \in \Pi^*$ is an optimal policy. Denote $V^* \in \mathbb{R}^{|\mathcal{S}|}$ as the optimal value function, then for such a π^* , we have for every state $s \in \mathcal{S}$,

$$\begin{aligned} V^*(s) &= \min_{a \in \mathcal{A}} Q^*(s, a) = \sum_{a \in \mathcal{A}^*(s)} \pi^*(a|s) Q^*(s, a) \\ &= \sum_{a \in \mathcal{A}} \pi^*(a|s) Q^*(s, a) \\ &= \sum_{a \in \mathcal{A}} \pi^*(a|s) [r(s, a) + \gamma \mathbb{E}_{s' \sim \mathcal{P}(s'|s, a)} V^*(s')] \\ &= \sum_{a \in \mathcal{A}} \pi^*(a|s) r(s, a) + \gamma \sum_{a \in \mathcal{A}} \sum_{s' \in \mathcal{S}} \pi^*(a|s) \mathcal{P}(s'|s, a) V^*(s'), \end{aligned} \quad (24)$$

where the first equality follows from the definition of Q^* and V^* , the third equality follows from the fact that $\pi^* \in \Pi^*$, Writing the previous relation in vector form, we have

$$r^{\pi^*} + \gamma \mathbb{P}^{\pi^*} V^* = V^*,$$

where $r^{\pi^*}(s) = \sum_{a \in \mathcal{A}} \pi^*(a|s) r(s, a)$, and $\mathbb{P}^{\pi^*}(s, s') = \sum_{a \in \mathcal{A}} \pi^*(a|s) \mathcal{P}(s'|s, a)$. Now since $(I - \gamma \mathbb{P}^{\pi^*})$ is invertible, we have equivalently

$$(I - \gamma \mathbb{P}^{\pi^*})^{-1} r^{\pi^*} = V^*,$$

Note that the left hand side is exactly the value function of π^* due to Bellman equation for the value vector: $V^\pi = r^\pi + \gamma \mathbb{P}^\pi V^\pi$ for any policy π . Hence $V^{\pi^*} = (I - \gamma \mathbb{P}^{\pi^*})^{-1} r^{\pi^*} = V^*$, which implies that π^* is an optimal policy.

We then show that any policy $\pi \notin \Pi^*$ can not be an optimal policy. Suppose there exists $s \in \mathcal{S}$, such that $\text{supp}(\pi(\cdot|s)) \not\subseteq \underset{a \in \mathcal{A}}{\text{argmin}} Q^*(s, a)$. Then repeating the previous argument in (24), we have

$$\begin{aligned} V^*(s) &= \min_{a \in \mathcal{A}} Q^*(s, a) < \sum_{a \in \mathcal{A}} \pi(a|s) Q^*(s, a) \\ &= \sum_{a \in \mathcal{A}} \pi(a|s) [r(s, a) + \gamma \mathbb{E}_{s' \sim \mathcal{P}(s'|s, a)} V^*(s')] \\ &= \sum_{a \in \mathcal{A}} \pi(a|s) r(s, a) + \gamma \sum_{a \in \mathcal{A}} \sum_{s' \in \mathcal{S}} \pi(a|s) \mathcal{P}(s'|s, a) V^*(s'), \end{aligned}$$

where the inequality comes from the fact that $\text{supp}(\pi(\cdot|s)) \not\subseteq \underset{a \in \mathcal{A}}{\text{argmin}} Q^*(s, a)$. Writing the previous relation in vector form, we have

$$r^\pi + \gamma \mathbb{P}^\pi V^* > V^*, \quad (25)$$

where $r^\pi(s) = \sum_{a \in \mathcal{A}} \pi(a|s) r(s, a)$, $\mathbb{P}^\pi(s, s') = \sum_{a \in \mathcal{A}} \pi(a|s) \mathcal{P}(s'|s, a)$, and the relation $>$ on $\mathbb{R}^{|\mathcal{S}|}$ is defined as: $a > b$ if $a_i - b_i \geq 0$ for all $i \in [|\mathcal{S}|]$ and $a_j > b_j$ for some $j \in [|\mathcal{S}|]$. Since $\|\gamma \mathbb{P}^\pi\| \leq \gamma < 1$, then $(I - \gamma \mathbb{P}^\pi)$ is invertible. Combined with the fact that \mathbb{P}^π has nonnegative entries, we also have

$$(I - \gamma \mathbb{P}^\pi)^{-1} z = \sum_{t=0}^{\infty} \gamma^t (\mathbb{P}^\pi)^t z \geq z > 0, \forall z \in \mathbb{R}^{|\mathcal{S}|}, z > 0.$$

Hence we obtain

$$a > b, a, b \in \mathbb{R}^{|\mathcal{S}|} \Rightarrow (I - \gamma \mathbb{P}^\pi)^{-1} a > (I - \gamma \mathbb{P}^\pi)^{-1} b.$$

By applying the previous relation to (25), we obtain

$$(I - \gamma \mathbb{P}^\pi)^{-1} r^\pi > V^*.$$

Note that the left hand side is exactly the value function of the policy π . That is, $V^\pi = (I - \gamma \mathbb{P}^\pi)^{-1} r^\pi > V^*$. Hence π is not an optimal policy. \square

A Global Linear Convergence of HPMD Methods

A.1 Convergence of Deterministic HPMD

Recall that we denote ν^* as the stationary state distribution generated by the optimal policy π^* . We will also define $\phi(\pi, \pi^*) = \mathbb{E}_{s \sim \nu^*} D_\pi^{\pi^*}(s)$. We first have the following lemma characterizing the update of the HPMD method.

Lemma A.1. *For any $p \in \Delta_{|\mathcal{A}|}$ and any $s \in \mathcal{S}$, we have*

$$\eta_k \langle Q^{\pi_k}(s, \cdot), \pi_{k+1}(\cdot|s) - p \rangle + \eta_k \tau_k (D_{\pi_0}^{\pi_{k+1}}(s) - D_{\pi_0}^p(s)) + D_{\pi_k}^{\pi_{k+1}}(s) \leq D_{\pi_k}^p(s) - (\tau_k \eta_k + 1) D_{\pi_{k+1}}^p(s). \quad (26)$$

Proof. From the optimality condition of the HPMD update (6), we have for any $p \in \Delta_{|\mathcal{A}|}$,

$$\eta_k \langle Q^{\pi_k}(s, \cdot), p - \pi_{k+1}(\cdot|s) \rangle + \eta_k \tau_k \langle \nabla D_{\pi_0}^{\pi_{k+1}}(s), p - \pi_{k+1}(\cdot|s) \rangle + \langle \nabla D_{\pi_k}^{\pi_{k+1}}(s), p - \pi_{k+1}(\cdot|s) \rangle \geq 0, \quad (27)$$

In addition, we note that $D_q^p = -\mathcal{H}(p) + \mathcal{H}(q) + \langle \nabla \mathcal{H}(q), p - q \rangle$, and hence D_q^p is a Bregman divergence with distance-generating function $-\mathcal{H}(p)$. Then we have the following identity

$$\begin{aligned} \langle \nabla D_{\pi_0}^{\pi_{k+1}}(s), p - \pi_{k+1}(\cdot|s) \rangle &= D_{\pi_0}^p(s) - D_{\pi_0}^{\pi_{k+1}}(s) - D_{\pi_{k+1}}^p(s), \\ \langle \nabla D_{\pi_k}^{\pi_{k+1}}(s), p - \pi_{k+1}(\cdot|s) \rangle &= D_{\pi_k}^p(s) - D_{\pi_k}^{\pi_{k+1}}(s) - D_{\pi_{k+1}}^p(s). \end{aligned}$$

Combining the previous two observations with (27), we immediately obtain the result. \square

We then establish the generic convergence property of the HPMD method.

Lemma A.2. *Suppose $1 + \eta_k \tau_k = 1/\gamma$, and $\alpha_k > 0$ satisfies*

$$\alpha_k \geq \alpha_{k+1} \gamma, \quad \frac{\alpha_k}{\gamma \eta_k} \geq \frac{\alpha_{k+1}}{\eta_{k+1}}, \quad (28)$$

then for any iteration k , HPMD satisfies

$$\alpha_k (f(\pi_{k+1}) - f(\pi^*)) + \frac{\alpha_k}{\gamma \eta_k} \phi(\pi_{k+1}, \pi^*) \leq \alpha_0 \gamma (f(\pi_0) - f(\pi^*)) + \frac{\alpha_0}{\eta_0} \phi(\pi_0, \pi^*) + \sum_{t=0}^k \frac{3\alpha_t}{\gamma \eta_t} \log |\mathcal{A}| \quad (29)$$

Proof. For each $s \in \mathcal{S}$, by plugging $p = \pi_k(\cdot|s)$ into (26), we obtain

$$\eta_k \langle Q^{\pi_k}(s, \cdot), \pi_{k+1}(\cdot|s) - \pi_k(\cdot|s) \rangle + \eta_k \tau_k (D_{\pi_0}^{\pi_{k+1}}(s) - D_{\pi_0}^{\pi_k}(s)) \leq -(\tau_k \eta_k + 1) D_{\pi_{k+1}}^{\pi_k}(s) \leq 0. \quad (30)$$

On the other hand, choosing $p = \pi^*(\cdot|s)$ in (26), we obtain

$$\begin{aligned} \eta_k \langle Q^{\pi_k}(s, \cdot), \pi_k(\cdot|s) - \pi^*(\cdot|s) \rangle + \eta_k \langle Q^{\pi_k}(s, \cdot), \pi_{k+1}(\cdot|s) - \pi_k(\cdot|s) \rangle + \eta_k \tau_k (D_{\pi_0}^{\pi_{k+1}}(s) - D_{\pi_0}^{\pi^*}(s)) \\ \leq D_{\pi_k}^{\pi^*}(s) - (\tau_k \eta_k + 1) D_{\pi_{k+1}}^{\pi^*}(s). \end{aligned} \quad (31)$$

We make the following observations,

$$\begin{aligned} \mathbb{E}_{s \sim \nu^*} \langle Q^{\pi_k}(s, \cdot), \pi_k(\cdot|s) - \pi^*(\cdot|s) \rangle &= \mathbb{E}_{s \sim \nu^*} \mathbb{E}_{s' \sim d_s^{\pi^*}} \langle Q^{\pi_k}(s', \cdot), \pi_k(\cdot|s') - \pi^*(\cdot|s') \rangle \\ &= -\mathbb{E}_{s \sim \nu^*} \mathbb{E}_{s' \sim d_s^{\pi^*}} \langle Q^{\pi_k}(s', \cdot), \pi^*(\cdot|s') - \pi_k(\cdot|s') \rangle \end{aligned}$$

$$\begin{aligned}
&= -(1-\gamma)\mathbb{E}_{s\sim\nu^*}(V^*(s) - V^{\pi^k}(s)) \\
&= (1-\gamma)(f(\pi_k) - f(\pi^*)), \tag{32}
\end{aligned}$$

where in the third equality we apply the performance difference lemma [10]. In addition, we also have

$$\begin{aligned}
&(1-\gamma)(V^{\pi_{k+1}}(s) - V^{\pi^k}(s)) + \mathbb{E}_{s'\sim d_s^{\pi_{k+1}}}[\tau_k(D_{\pi_0}^{\pi_{k+1}}(s') - D_{\pi_0}^{\pi^k}(s'))] \\
&= \mathbb{E}_{s'\sim d_s^{\pi_{k+1}}}[\langle Q^{\pi^k}(s', \cdot), \pi_{k+1}(\cdot|s') - \pi_k(\cdot|s') \rangle + \tau_k(D_{\pi_0}^{\pi_{k+1}}(s') - D_{\pi_0}^{\pi^k}(s'))] \\
&\leq (1-\gamma)[\langle Q^{\pi^k}(s, \cdot), \pi_{k+1}(\cdot|s) - \pi_k(\cdot|s) \rangle + \tau_k(D_{\pi_0}^{\pi_{k+1}}(s) - D_{\pi_0}^{\pi^k}(s))],
\end{aligned}$$

where the first equality uses again the performance difference lemma, and the inequality uses (30), and the fact that $d_s^{\pi_{k+1}}(s) \geq 1-\gamma$. Thus from the previous relation, we obtain

$$\langle Q^{\pi^k}(s, \cdot), \pi_{k+1}(\cdot|s) - \pi_k(\cdot|s) \rangle \geq V^{\pi_{k+1}}(s) - V^{\pi^k}(s) - \frac{2\tau_k}{1-\gamma} \max_{s\in\mathcal{S}} |D_{\pi_0}^{\pi_{k+1}}(s') - D_{\pi_0}^{\pi^k}(s')| \tag{33}$$

By taking expectation w.r.t. $s \sim \nu^*$ in (31), and combining (32), (33), we obtain

$$\begin{aligned}
&(1-\gamma)[f(\pi_k) - f(\pi^*)] + f(\pi_{k+1}) - f(\pi^*) \\
&\leq \frac{1}{\eta_k} \phi(\pi_k, \pi^*) - \left(\frac{1}{\eta_k} + \tau_k\right) \phi(\pi_{k+1}, \pi^*) + \frac{2\tau_k}{1-\gamma} \max_{s\in\mathcal{S}} |D_{\pi_0}^{\pi_{k+1}}(s') - D_{\pi_0}^{\pi^k}(s')| + \tau_k \max_{s\in\mathcal{S}} |D_{\pi_0}^{\pi_{k+1}}(s) - D_{\pi_0}^{\pi^k}(s)|.
\end{aligned}$$

After simple rearrangement, the previous relation becomes

$$\begin{aligned}
&f(\pi_{k+1}) - f(\pi^*) + \left(\frac{1}{\eta_k} + \tau_k\right) \phi(\pi_{k+1}, \pi^*) \\
&\leq \gamma(f(\pi_k) - f(\pi^*)) + \frac{1}{\eta_k} \phi(\pi_k, \pi^*) + \frac{2\tau_k}{1-\gamma} \max_{s\in\mathcal{S}} |D_{\pi_0}^{\pi_{k+1}}(s') - D_{\pi_0}^{\pi^k}(s')| + \tau_k \max_{s\in\mathcal{S}} |D_{\pi_0}^{\pi_{k+1}}(s) - D_{\pi_0}^{\pi^k}(s)|.
\end{aligned}$$

Suppose η_k, τ_k satisfies $1 + \eta_k \tau_k = 1/\gamma$, then the previous relation becomes

$$\begin{aligned}
&f(\pi_{k+1}) - f(\pi^*) + \frac{1}{\gamma\eta_k} \phi(\pi_{k+1}, \pi^*) \leq \gamma(f(\pi_k) - f(\pi^*)) + \frac{1}{\eta_k} \phi(\pi_k, \pi^*) + \frac{3\tau_k}{1-\gamma} \log|\mathcal{A}| \\
&= \gamma(f(\pi_k) - f(\pi^*)) + \frac{1}{\eta_k} \phi(\pi_k, \pi^*) + \frac{3}{\gamma\eta_k} \log|\mathcal{A}|, \tag{34}
\end{aligned}$$

where we use the fact that $0 \leq D_{\pi_0}^{\pi}(s) \leq \log|\mathcal{A}|$ since π_0 is a uniform policy. Now multiplying both sides of (34) by some positive constant α_k , where

$$\alpha_k \geq \alpha_{k+1}\gamma, \quad \frac{\alpha_k}{\gamma\eta_k} \geq \frac{\alpha_{k+1}}{\eta_{k+1}},$$

then summing up from $t = 0$ to k , we obtain

$$\alpha_k (f(\pi_{k+1}) - f(\pi^*)) + \frac{\alpha_k}{\gamma\eta_k} \phi(\pi_{k+1}, \pi^*) \leq \alpha_0 \gamma (f(\pi_0) - f(\pi^*)) + \frac{\alpha_0}{\eta_0} \phi(\pi_0, \pi^*) + \sum_{t=0}^k \frac{3\alpha_t}{\gamma\eta_t} \log|\mathcal{A}|.$$

□

Theorem (Restating Theorem 2.1). *By choosing $1 + \eta_k \tau_k = 1/\gamma$ and $\eta_k = \gamma^{-2(k+1)}$ in HPMD, then for any iteration $k \geq 1$, HPMD produces policy π_k with*

$$f(\pi_k) - f(\pi^*) \leq \gamma^k \left(f(\pi_0) - f(\pi^*) + \frac{4 \log|\mathcal{A}|}{1-\gamma} \right).$$

Proof of Theorem 2.1. We now specify the concrete choice of $\{(\eta_k, \tau_k, \alpha_k)\}_{k \geq 0}$ that yields the global linear convergence of the HPMD method. By choosing $\alpha_k = \gamma^{-(k+1)}$, $\eta_k = \gamma^{-2(k+1)}$, one can readily verify that condition (28) holds. Thus from the recursion (29), we obtain

$$\gamma^{-k} (f(\pi_k) - f(\pi^*)) + \gamma^{k-1} \phi(\pi_k, \pi^*) \leq f(\pi_0) - f(\pi^*) + \gamma \phi(\pi_0, \pi^*) + \sum_{t=0}^{k-1} \frac{3\alpha_t}{\gamma\eta_t} \log|\mathcal{A}|$$

$$\begin{aligned}
&\leq f(\pi_0) - f(\pi^*) + \gamma\phi(\pi_0, \pi^*) + \sum_{t=0}^{k-1} 3\gamma^t \log|\mathcal{A}| \\
&\leq f(\pi_0) - f(\pi^*) + \frac{4\log|\mathcal{A}|}{1-\gamma},
\end{aligned}$$

from which we immediately conclude

$$f(\pi_k) - f(\pi^*) \leq \gamma^k \left(f(\pi_0) - f(\pi^*) + \frac{4\log|\mathcal{A}|}{1-\gamma} \right).$$

□

A.2 Convergence of Stochastic HPMD

We establish convergence properties of the stochastic homotopic policy mirror descent (SHPMD) method. Recall that we denote ν^* as the stationary state distribution generated by the optimal policy π^* , and $\phi(\pi, \pi^*) = \mathbb{E}_{s \sim \nu^*} D_{\pi}^{\pi^*}(s)$.

We first have the following lemma characterizing the update of the SHPMD method.

Lemma A.3. *For any $p \in \Delta_{|\mathcal{A}|}$ and any $s \in \mathcal{S}$, we have*

$$\begin{aligned}
&\eta_k \langle Q^{\pi_k, \xi_k}(s, \cdot), \pi_{k+1}(\cdot|s) - p \rangle + \eta_k \tau_k (D_{\pi_0}^{\pi_{k+1}}(s) - D_{\pi_0}^p(s)) + D_{\pi_k}^{\pi_{k+1}}(s) \\
&\leq D_{\pi_k}^p(s) - (\tau_k \eta_k + 1) D_{\pi_{k+1}}^p(s).
\end{aligned} \tag{35}$$

The proof of Lemma A.3 follows exactly the same lines as in Lemma A.1. We then establish the generic convergence properties of the SHPMD method.

Lemma A.4. *Suppose $1 + \eta_k \tau_k = 1/\gamma$, and $\alpha_k > 0$ satisfies*

$$\alpha_k \geq \alpha_{k+1} \gamma, \quad \frac{\alpha_k}{\gamma \eta_k} \geq \frac{\alpha_{k+1}}{\eta_{k+1}}, \tag{36}$$

then for any iteration k , SHPMD satisfies

$$\begin{aligned}
&\mathbb{E} \left[\alpha_k (f(\pi_{k+1}) - f(\pi^*)) + \frac{\alpha_k}{\gamma \eta_k} \phi(\pi_{k+1}, \pi^*) \right] \\
&\leq \alpha_0 \gamma (f(\pi_0) - f(\pi^*)) + \frac{\alpha_0}{\eta_0} \phi(\pi_0, \pi^*) + \frac{2}{1-\gamma} \sum_{t=0}^k \alpha_t \varepsilon_t + \frac{2}{1-\gamma} \sum_{t=0}^k \alpha_t \eta_t \sigma_t^2 + \frac{3}{\gamma} \sum_{t=0}^k \frac{\alpha_t}{\eta_t} \log|\mathcal{A}|.
\end{aligned} \tag{37}$$

Proof. For each $s \in \mathcal{S}$, by plugging $p = \pi_k(\cdot|s)$ in (35), we obtain

$$\eta_k \langle Q^{\pi_k}(s, \cdot), \pi_{k+1}(\cdot|s) - \pi_k(\cdot|s) \rangle + \eta_k \tau_k (D_{\pi_0}^{\pi_{k+1}}(s) - D_{\pi_0}^{\pi_k}(s)) + D_{\pi_k}^{\pi_{k+1}}(s) \leq -(\tau_k \eta_k + 1) D_{\pi_{k+1}}^{\pi_k}(s) \leq 0. \tag{38}$$

On the other hand, choosing $p = \pi^*(\cdot|s)$ in (35), we obtain

$$\begin{aligned}
&\eta_k \langle Q^{\pi_k, \xi_k}(s, \cdot), \pi_k(\cdot|s) - \pi^*(\cdot|s) \rangle + \eta_k \langle Q^{\pi_k, \xi_k}(s, \cdot), \pi_{k+1}(\cdot|s) - \pi_k(\cdot|s) \rangle + \eta_k \tau_k (D_{\pi_0}^{\pi_{k+1}}(s) - D_{\pi_0}^{\pi^*}(s)) \\
&\leq D_{\pi_k}^{\pi^*}(s) - (\tau_k \eta_k + 1) D_{\pi_{k+1}}^{\pi^*}(s) - D_{\pi_k}^{\pi_{k+1}}(s).
\end{aligned} \tag{39}$$

Define $\delta_k = Q^{\pi_k, \xi_k} - Q^{\pi_k} \in \mathbb{R}^{|\mathcal{S}| \times |\mathcal{A}|}$, and denote $\delta_k(s, \cdot)$ as its s -th row, then for (39), we first have

$$\langle Q^{\pi_k, \xi_k}(s, \cdot), \pi_k(\cdot|s) - \pi^*(\cdot|s) \rangle = \langle Q^{\pi_k}(s, \cdot), \pi_k(\cdot|s) - \pi^*(\cdot|s) \rangle + \langle \delta_k(s, \cdot), \pi_k(\cdot|s) - \pi^*(\cdot|s) \rangle$$

Then from (32), we obtain

$$\mathbb{E}_{s \sim \nu^*} \langle Q^{\pi_k, \xi_k}(s, \cdot), \pi_k(\cdot|s) - \pi^*(\cdot|s) \rangle = (1-\gamma) (f(\pi_k) - f(\pi^*)) + \mathbb{E}_{s \sim \nu^*} \langle \delta_k(s, \cdot), \pi_k(\cdot|s) - \pi^*(\cdot|s) \rangle. \tag{40}$$

In addition, we also have

$$(1-\gamma) (V^{\pi_{k+1}}(s) - V^{\pi_k}(s)) + \mathbb{E}_{s' \sim d_s^{\pi_{k+1}}} [\tau_k (D_{\pi_0}^{\pi_{k+1}}(s') - D_{\pi_0}^{\pi_k}(s'))]$$

$$\begin{aligned}
&= \mathbb{E}_{s' \sim d_s^{\pi_{k+1}}} \left[\langle Q^{\pi_k, \xi_k}(s', \cdot), \pi_{k+1}(\cdot|s') - \pi_k(\cdot|s') \rangle + \tau_k (D_{\pi_0}^{\pi_{k+1}}(s') - D_{\pi_0}^{\pi_k}(s')) \right. \\
&\quad \left. - \langle \delta_k(s', \cdot), \pi_{k+1}(\cdot|s') - \pi_k(\cdot|s') \rangle \right] \\
&\leq \mathbb{E}_{s' \sim d_s^{\pi_{k+1}}} \left[\langle Q^{\pi_k, \xi_k}(s', \cdot), \pi_{k+1}(\cdot|s') - \pi_k(\cdot|s') \rangle + \tau_k (D_{\pi_0}^{\pi_{k+1}}(s') - D_{\pi_0}^{\pi_k}(s')) \right. \\
&\quad \left. + \|\delta_k(s', \cdot)\|_\infty \|\pi_{k+1}(\cdot|s') - \pi_k(\cdot|s')\|_1 \right] \\
&\leq \mathbb{E}_{s' \sim d_s^{\pi_{k+1}}} \left[\langle Q^{\pi_k, \xi_k}(s', \cdot), \pi_{k+1}(\cdot|s') - \pi_k(\cdot|s') \rangle + \tau_k (D_{\pi_0}^{\pi_{k+1}}(s') - D_{\pi_0}^{\pi_k}(s')) \right. \\
&\quad \left. + \frac{\eta_k}{2} \|\delta_k(s, \cdot)\|_\infty^2 + \frac{1}{2\eta_k} \|\pi_{k+1}(\cdot|s') - \pi_k(\cdot|s')\|_1^2 \right] \\
&\leq \mathbb{E}_{s' \sim d_s^{\pi_{k+1}}} \left[\langle Q^{\pi_k, \xi_k}(s', \cdot), \pi_{k+1}(\cdot|s') - \pi_k(\cdot|s') \rangle + \tau_k (D_{\pi_0}^{\pi_{k+1}}(s') - D_{\pi_0}^{\pi_k}(s')) \right. \\
&\quad \left. + \frac{\eta_k}{2} \|\delta_k(s', \cdot)\|_\infty^2 + \frac{1}{\eta_k} D_{\pi_k}^{\pi_{k+1}}(s') \right] \\
&\leq (1 - \gamma) \left[\langle Q^{\pi_k, \xi_k}(s, \cdot), \pi_{k+1}(\cdot|s) - \pi_k(\cdot|s) \rangle + \tau_k (D_{\pi_0}^{\pi_{k+1}}(s) - D_{\pi_0}^{\pi_k}(s)) + \frac{1}{\eta_k} D_{\pi_k}^{\pi_{k+1}}(s) \right] \\
&\quad + \frac{\eta_k}{2} \|\delta_k\|_\infty^2
\end{aligned}$$

where the first equality uses the performance difference lemma, and the last inequality uses (38) and the fact that $1 \geq d_s^{\pi_{k+1}}(s) \geq 1 - \gamma$. From the previous relation, we obtain

$$\begin{aligned}
\langle Q^{\pi_k, \xi_k}(s, \cdot), \pi_{k+1}(\cdot|s) - \pi_k(\cdot|s) \rangle &\geq (V^{\pi_{k+1}}(s) - V^{\pi_k}(s)) - \frac{2\tau_k}{1 - \gamma} \max_{s \in \mathcal{S}} |D_{\pi_0}^{\pi_{k+1}}(s) - D_{\pi_0}^{\pi_k}(s)| \\
&\quad - \frac{\eta_k}{2(1 - \gamma)} \|\delta_k\|_\infty^2 - \frac{1}{\eta_k} D_{\pi_k}^{\pi_{k+1}}(s).
\end{aligned} \tag{41}$$

Thus, by combining (39) and (41), we obtain

$$\begin{aligned}
&\eta_k \langle Q^{\pi_k, \xi_k}(s, \cdot), \pi_k(\cdot|s) - \pi^*(\cdot|s) \rangle + \eta_k \left[(V^{\pi_{k+1}}(s) - V^{\pi_k}(s)) - \frac{2\tau_k}{1 - \gamma} \max_{s \in \mathcal{S}} |D_{\pi_0}^{\pi_{k+1}}(s) - D_{\pi_0}^{\pi_k}(s)| \right. \\
&\quad \left. - \frac{\eta_k}{2(1 - \gamma)} \|\delta_k\|_\infty^2 \right] \\
&\leq D_{\pi_k}^{\pi^*}(s) - (\tau_k \eta_k + 1) D_{\pi_{k+1}}^{\pi^*}(s) + \tau_k \max_{s \in \mathcal{S}} |D_{\pi_0}^{\pi_{k+1}}(s) - D_{\pi_0}^{\pi^*}(s)|.
\end{aligned}$$

Taking expectation w.r.t. $s \sim \nu^*$ on both sides of the previous relation, then given the above relation, (40) and the definition of f , we obtain

$$\begin{aligned}
&(1 - \gamma) [f(\pi_k) - f(\pi^*)] + f(\pi_{k+1}) - f(\pi^*) \\
&\leq \frac{1}{\eta_k} \phi(\pi_k, \pi^*) - \left(\frac{1}{\eta_k} + \tau_k \right) \phi(\pi_{k+1}, \pi^*) + \frac{2\tau_k}{1 - \gamma} \max_{s \in \mathcal{S}} |D_{\pi_0}^{\pi_{k+1}}(s') - D_{\pi_0}^{\pi_k}(s')| \\
&\quad - \mathbb{E}_{s \sim \nu^*} \langle \delta_k(s, \cdot), \pi_k(\cdot|s) - \pi^*(\cdot|s) \rangle + \frac{\eta_k}{2(1 - \gamma)} \|\delta_k\|_\infty^2 + \tau_k \max_{s \in \mathcal{S}} |D_{\pi_0}^{\pi_{k+1}}(s) - D_{\pi_0}^{\pi^*}(s)|.
\end{aligned}$$

Simple rearrangement of the previous relation gives

$$\begin{aligned}
&f(\pi_{k+1}) - f(\pi^*) + \left(\frac{1}{\eta_k} + \tau_k \right) \phi(\pi_{k+1}, \pi^*) \\
&\leq \gamma (f(\pi_k) - f(\pi^*)) + \frac{1}{\eta_k} \phi(\pi_k, \pi^*) + \frac{2\tau_k}{1 - \gamma} \max_{s \in \mathcal{S}} |D_{\pi_0}^{\pi_{k+1}}(s') - D_{\pi_0}^{\pi_k}(s')| \\
&\quad - \mathbb{E}_{s \sim \nu^*} \langle \delta_k(s, \cdot), \pi_k(\cdot|s) - \pi^*(\cdot|s) \rangle + \frac{\eta_k}{2(1 - \gamma)} \|\delta_k\|_\infty^2 + \tau_k \max_{s \in \mathcal{S}} |D_{\pi_0}^{\pi_{k+1}}(s) - D_{\pi_0}^{\pi^*}(s)|.
\end{aligned}$$

Choosing $1 + \eta_k \tau_k = 1/\gamma$, then we obtain

$$\begin{aligned}
&f(\pi_{k+1}) - f(\pi^*) + \frac{1}{\gamma \eta_k} \phi(\pi_{k+1}, \pi^*) \\
&\leq \gamma (f(\pi_k) - f(\pi^*)) + \frac{1}{\eta_k} \phi(\pi_k, \pi^*) + \frac{3}{\gamma \eta_k} \log |\mathcal{A}| - \mathbb{E}_{s \sim \nu^*} \langle \delta_k(s, \cdot), \pi_k(\cdot|s) - \pi^*(\cdot|s) \rangle + \frac{\gamma \eta_k}{2(1 - \gamma)} \|\delta_k\|_\infty^2,
\end{aligned}$$

where we use the fact that $0 \leq D_{\pi_0}^\pi(s) \leq \log|\mathcal{A}|$ since π_0 is a uniform policy. Now multiplying both sides of previous relation with positive $\alpha_k > 0$, where

$$\alpha_k \geq \alpha_{k+1}\gamma, \quad \frac{\alpha_k}{\gamma\eta_k} \geq \frac{\alpha_{k+1}}{\eta_{k+1}},$$

and summing up from $t = 0$ to k , and taking expectation w.r.t $\{\xi_t\}_{t \geq 0}$, we obtain

$$\begin{aligned} & \mathbb{E} \left[\alpha_k (f(\pi_{k+1}) - f(\pi^*)) + \frac{\alpha_k}{\gamma\eta_k} \phi(\pi_{k+1}, \pi^*) \right] \\ & \leq \alpha_0 \gamma (f(\pi_0) - f(\pi^*)) + \frac{\alpha_0}{\eta_0} \phi(\pi_0, \pi^*) + 2 \sum_{t=0}^k \alpha_t \varepsilon_t + \frac{2}{1-\gamma} \sum_{t=0}^k \alpha_t \eta_t \sigma_t^2 + \frac{3}{\gamma} \sum_{t=0}^k \frac{\alpha_t}{\eta_t} \log|\mathcal{A}|, \end{aligned}$$

where we also uses condition (15), (16), together with $\langle \mathbb{E}_{\xi_k} \delta_k(s, \cdot), \pi^*(\cdot|s) - \pi_k(\cdot|s) \rangle \leq 2 \|\mathbb{E}_{\xi_k} \delta_k(s, \cdot)\|_\infty \leq 2\varepsilon_k$. \square

We now specify the concrete choice of $\{(\eta_k, \tau_k, \alpha_k)\}_{k \geq 0}$ and $\{(\varepsilon_k, \sigma_k)\}_{k \geq 0}$ that yields the global linear convergence of the SHPMD method.

Theorem (Restating Theorem 4.1). *Take $1 + \eta_k \tau_k = 1/\gamma$ and $\eta_k = \gamma^{-(k+1)/2} \sqrt{\log|\mathcal{A}|}$ in the SHPMD method. Suppose*

$$\sigma_k = \gamma^{(k+1)/2}, \quad \varepsilon_k = \gamma^{3(k+1)/4}$$

holds in (15), (16), then for any iteration $k \geq 1$, SHPMD produces policy π_k satisfying

$$\mathbb{E} [f(\pi_k) - f(\pi^*)] \leq \gamma^{k/2} \frac{6\sqrt{\log|\mathcal{A}|} + C}{(1-\gamma)(1-\gamma^{1/2})\gamma}.$$

Proof of Theorem 4.1. By taking $\sigma_t = \gamma^{(t+1)/2}$, $\eta_t = \gamma^{-(t+1)/2} \sqrt{\log|\mathcal{A}|}$, and $\alpha_t = \gamma^{-(t+1)}$, one can readily verify that

$$\frac{\alpha_t}{\alpha_{t+1}} = \gamma, \quad \frac{\alpha_t}{\gamma\eta_t} = \gamma^{-(t+1)/2-1} / \sqrt{\log|\mathcal{A}|}, \quad \frac{\alpha_{t+1}}{\eta_{t+1}} = \gamma^{-(t+2)/2} / \sqrt{\log|\mathcal{A}|}$$

and thus (36) is satisfied. In addition, we have

$$\frac{2}{1-\gamma} \sum_{t=0}^k \alpha_t \eta_t \sigma_t^2 + \frac{3}{\gamma} \sum_{t=0}^k \frac{\alpha_t}{\eta_t} \log|\mathcal{A}| \leq \frac{3\sqrt{\log|\mathcal{A}|}(\gamma^{-k/2} - 1)}{(1-\gamma)(\gamma^{-1/2} - 1)\gamma^{3/2}} \leq \frac{3\sqrt{\log|\mathcal{A}|}\gamma^{-k/2}}{(1-\gamma)(1-\gamma^{1/2})\gamma}.$$

On the other hand, choosing $\varepsilon_t = \gamma^{3(t+1)/4}$, we have

$$2 \sum_{t=0}^k \alpha_t \varepsilon_t \leq \frac{2\gamma^{-k/2}}{1-\gamma^{1/2}}.$$

Combining previous two observations and (37), we obtain

$$\begin{aligned} & \mathbb{E} \left[\gamma^{-k} (f(\pi_k) - f(\pi^*)) + \frac{\gamma^{-k/2-1}}{\sqrt{\log|\mathcal{A}|}} \phi(\pi_k, \pi^*) \right] \\ & \leq f(\pi_0) - f(\pi^*) + \frac{\gamma^{-1/2}}{\sqrt{\log|\mathcal{A}|}} \phi(\pi_0, \pi^*) + \frac{2\gamma^{-k/2}}{1-\gamma^{1/2}} + \frac{3\sqrt{\log|\mathcal{A}|}\gamma^{-k/2}}{(1-\gamma)(1-\gamma^{1/2})\gamma}, \quad \forall k \geq 1. \end{aligned}$$

which translates into

$$\begin{aligned} \mathbb{E} [f(\pi_k) - f(\pi^*)] & \leq \gamma^k (f(\pi_0) - f(\pi^*)) + \gamma^{k-1/2} \sqrt{\log|\mathcal{A}|} + 5\gamma^{k/2} \frac{\sqrt{\log|\mathcal{A}|}}{(1-\gamma^{1/2})(1-\gamma)\gamma} \\ & \leq \gamma^{k/2} \frac{6\sqrt{\log|\mathcal{A}|} + C}{(1-\gamma)(1-\gamma^{1/2})\gamma}, \quad \forall k \geq 1. \end{aligned}$$

\square

A.3 Sample Complexity of Stochastic HPMD

We assume the access to a generative model of the MDP. That is, for any state-action pair $(s, a) \in \mathcal{S} \times \mathcal{A}$, we can generate M_k independent trajectories, each with length T_k , by following policy π_k starting from (s, a) . The i -th trajectory, denoted by $\zeta_k^i(s, a)$, takes the form of $\zeta_k^i(s, a) = \{(s_0^i = s, a_0^i = a), (s_1^i, a_1^i), \dots, (s_{T_k-1}^i, a_{T_k-1}^i)\}$.

Let us denote the set of random variables for constructing the stochastic estimate as $\xi_k = \{\zeta_k^i(s, a), s \in \mathcal{S}, a \in \mathcal{A}, i \in [M_k]\}$. Then the stochastic estimator Q^{π_k, ξ_k} is defined by

$$Q^{\pi_k, \xi_k}(s, a) = \frac{1}{M_k} \sum_{i=1}^{M_k} \sum_{t=0}^{T_k-1} \gamma^t [c(s_t^i, a_t^i) + h^{\pi_k}(s_t^i)], \quad s \in \mathcal{S}, a \in \mathcal{A}.$$

One can readily verify that Q^{π_k, ξ_k, i_k} satisfies (15) and (16) with

$$\varepsilon_k = \frac{C\gamma^{T_k}}{1-\gamma}, \sigma_k^2 = 2\varepsilon^2 + \frac{2C^2}{(1-\gamma)^2 M_k} = \frac{2C^2}{(1-\gamma)^2} \left(\gamma^{2T_k} + \frac{1}{M_k} \right). \quad (42)$$

Theorem (Restating Theorem 4.2). *Suppose at each iteration of SHPMD, the number of independent trajectories M_k and the trajectory length T_k satisfies*

$$T_k \geq \frac{3(k+1)}{4} + \log_\gamma \left(\frac{1-\gamma}{2C} \right), M_k \geq \frac{4C^2}{(1-\gamma)^2} \gamma^{-(k+1)}.$$

Then for any $\epsilon > 0$, the SHPMD method finds an policy π_k with $\mathbb{E}[f(\pi_k) - f(\pi^)] \leq \epsilon$ in $k = 2 \log_\gamma \left(\frac{\epsilon(1-\gamma)(1-\gamma^{1/2})}{6\sqrt{\log|\mathcal{A}|+C}} \right)$, number of iterations. In addition, the total number of samples required by SHPMD can be bounded by*

$$\mathcal{O} \left(\frac{C^2(6\sqrt{\log|\mathcal{A}|}+C)^2|\mathcal{S}||\mathcal{A}|}{(1-\gamma)^5\epsilon^2} \log_\gamma \left(\frac{\epsilon(1-\gamma)(1-\gamma^{1/2})}{6\sqrt{\log|\mathcal{A}|+C}} \right) \right).$$

Proof of Theorem 4.2. By choosing M_k and T_k as in (19), one can readily verify that condition (17) is satisfied. To find an ϵ -optimal policy, from (18), one needs

$$k = 2 \log_\gamma \left(\frac{\epsilon(1-\gamma)(1-\gamma^{1/2})}{6\sqrt{\log|\mathcal{A}|+C}} \right),$$

number of iterations. Thus the total number of samples can be bounded by

$$\begin{aligned} |\mathcal{S}||\mathcal{A}| \sum_{t=0}^k M_t T_t &= |\mathcal{S}||\mathcal{A}| \sum_{t=0}^k \left[\frac{3(t+1)}{4} + \log_\gamma \left(\frac{1-\gamma}{2C} \right) \right] \frac{4C^2}{(1-\gamma)^2} \gamma^{-(t+1)} \\ &= \frac{4C^2|\mathcal{S}||\mathcal{A}|}{(1-\gamma)^2} \mathcal{O} \left(\frac{k\gamma^{-k}}{1-\gamma} \right) \\ &= \frac{4C^2|\mathcal{S}||\mathcal{A}|}{(1-\gamma)^3} \mathcal{O} \left(\log_\gamma \left(\frac{\epsilon(1-\gamma)}{6\sqrt{\log|\mathcal{A}|+C}} \right) \cdot \frac{(6\sqrt{\log|\mathcal{A}|}+C)^2}{(1-\gamma)^2(1-\gamma^{1/2})^2\epsilon^2} \right) \\ &= \mathcal{O} \left(\frac{4C^2(6\sqrt{\log|\mathcal{A}|}+C)^2|\mathcal{S}||\mathcal{A}|}{(1-\gamma)^5(1-\gamma^{1/2})^2\epsilon^2} \log_\gamma \left(\frac{\epsilon(1-\gamma)}{6\sqrt{\log|\mathcal{A}|+C}} \right) \right). \end{aligned}$$

□

B Policy Convergence

B.1 Deterministic HPMD

We proceed to show the policy convergence of the HPMD method.

Theorem (Restating Theorem 3.1). *We define $\Delta^*(s) = \min_{a \notin \mathcal{A}^*(s)} Q^*(s, a) - \min_{a \in \mathcal{A}} Q^*(s, a)$ if $\mathcal{A}^*(s) \neq \mathcal{A}$, and $\Delta^*(s) = \infty$ otherwise. Define the gap function of the MDP \mathcal{M} as $\Delta^*(\mathcal{M}) = \min_{s \in \mathcal{S}} \Delta^*(s)$. Then whenever $\Delta^*(\mathcal{M}) < \infty$, the iterate $\{\pi_k\}_{k \geq 0}$ generated by HPMD satisfies*

$$\text{dist}_{\ell_1}(\pi_k, \Pi^*) \leq 2C_\gamma |\mathcal{A}| \exp\left(-\frac{\Delta^*(\mathcal{M})}{2} \gamma^{-2k-1}\right), \quad \forall k \geq K_1,$$

for any iteration $k \geq K_1 := \lceil 3 \log_\gamma \left(\frac{\Delta^*(\mathcal{M})(1-\gamma)}{2r(4 \log |\mathcal{A}| + C)} \right) \rceil$, and $C_\gamma := \exp\left(\frac{2C\gamma^2}{(1-\gamma^3)(1-\gamma)}\right)$. On the other hand, if $\Delta^*(\mathcal{M}) = \infty$, then any policy is an optimal policy.

Proof of Theorem 3.1. Recall that from Assumption 1, there exists π^* such that ν^{π^*} has full support. From the definition of state-action value function, we have

$$Q^{\pi_k}(s, a) - Q^*(s, a) = \gamma \sum_{s' \in \mathcal{S}} \mathcal{P}(s'|s, a) [V^{\pi_k}(s') - V^*(s')]. \quad (43)$$

Hence we can conclude for $r := \gamma \max_{s \in \mathcal{S}, a \in \mathcal{A}} \max_{s' \in \mathcal{S}} \mathcal{P}(s'|s, a) / \nu^{\pi^*}(s')$,

$$\text{supp}(\mathcal{P}(\cdot|s, a)) \subseteq \nu^{\pi^*} \Rightarrow Q^{\pi_k}(s, a) - Q^*(s, a) \leq \gamma^k \frac{r(4 \log |\mathcal{A}| + C)}{1 - \gamma}, \quad (44)$$

where C denotes the upper bound of the cost.

We begin by inspecting the update rule of HPMD. Directly examining the optimality condition of the update (6), we obtain

$$\log \pi_{k+1}(a|s) = \frac{\log \pi_k(a|s) - \eta_k Q^{\pi_k}(s, a)}{\eta_k \tau_k + 1} + \Delta_k(s), \quad \forall (s, a) \in \mathcal{S} \times \mathcal{A}, \quad (45)$$

where $\Delta_k(s)$ denotes the normalization constant to make the right hand size of (45) the logarithm of a valid probability vector defined over \mathcal{A} . Combined the choice of $\{(\tau_k, \eta_k)\}_{k \geq 0}$ satisfying $1 + \tau_k \eta_k = 1/\gamma$, we have

$$\log \pi_{k+1}(a|s) = \gamma [\log \pi_k(a|s) - \eta_k Q^{\pi_k}(s, a)] + \Delta_k(s), \quad \forall (s, a) \in \mathcal{S} \times \mathcal{A}. \quad (46)$$

Define $z_i^k(s) = \log \pi_k(i|s)$ for any pair $(s, i) \in \mathcal{S} \times \mathcal{A}$. For the ease of presentation, let us fix a state $s \in \mathcal{S}$ and write z_i^k in short for $z_i^k(s)$, g_i^k in short for $Q^{\pi_k}(s, i)$, and Δ_k in short for $\Delta_k(s)$, when the context is clear. Then applying (46) recursively, we obtain that for any $i \in \mathcal{A}$,

$$z_i^{k+1} = \gamma^{k+1} z_i^0 - \sum_{t=0}^k \gamma^{k+1-t} \eta_t g_i^t + \sum_{t=0}^k \gamma^{k-t} \Delta_t.$$

Hence for any pair of actions $i, j \in \mathcal{A}$, we have

$$z_i^{k+1} - z_j^{k+1} = \gamma^{k+1} (z_i^0 - z_j^0) - \sum_{t=0}^k \gamma^{k+1-t} \eta_t (g_i^t - g_j^t) = - \sum_{t=0}^k \gamma^{k+1-t} \eta_t (g_i^t - g_j^t). \quad (47)$$

Recall that from Lemma 5.2, it suffices to characterize the convergence of $\pi_k(a|s)$ for $a \notin \mathcal{A}^*(s)$. It then suffices to consider any pair of action i, j satisfying $Q^*(s, i) < Q^*(s, j)$. Combining with the observation in (44) and the assumption that $\text{supp}(\nu^{\pi^*}) = \mathcal{S}$ for some $\pi^* \in \Pi^*$, we conclude that there exists $\underline{K}_1(s)$ such that

$$g_i^t - g_j^t \leq (Q^*(s, i) - Q^*(s, j))/2, \quad \forall t \geq \underline{K}_1(s). \quad (48)$$

Recall that we choose $\eta_k = \gamma^{-2(k+1)}$ in HPMD. Thus for any $k \geq \underline{K}_1(s)$, we have

$$\begin{aligned} z_i^{k+1} - z_j^{k+1} &= - \sum_{t=0}^{\underline{K}_1(s)} \gamma^{k+1-t} \eta_t (g_i^t - g_j^t) - \sum_{t=\underline{K}_1(s)+1}^k \gamma^{k+1-t} \eta_t (g_i^t - g_j^t) \\ &= - \sum_{t=0}^{\underline{K}_1(s)} \gamma^{k-3t-1} (g_i^t - g_j^t) - \sum_{t=\underline{K}_1(s)+1}^k \gamma^{k-3t-1} (g_i^t - g_j^t) \end{aligned}$$

$$\begin{aligned}
&\geq -\sum_{t=0}^{\underline{K}_1(s)} \gamma^{k-3t-1} \cdot \frac{2C}{1-\gamma} + \gamma^{-2k-1} [Q^*(s, j) - Q^*(s, i)] / 2 \\
&\geq -\frac{2C\gamma^{k-3\underline{K}_1(s)+2}}{(1-\gamma^3)(1-\gamma)} + \frac{\gamma^{-2k-1}}{2} [Q^*(s, j) - Q^*(s, i)]
\end{aligned}$$

Thus for any $k \geq 3\underline{K}_1(s)$, we obtain

$$z_i^{k+1} - z_j^{k+1} \geq -\frac{2C\gamma^2}{(1-\gamma^3)(1-\gamma)} + \frac{\gamma^{-2k-1}}{2} [Q^*(s, j) - Q^*(s, i)]$$

It remains to bound $\underline{K}_1(s)$ that satisfies (48). From the observation (44), to satisfy (48), it suffices to choose

$$\underline{K}_1(s) \geq \log_\gamma \left(\frac{[Q^*(s, j) - Q^*(s, i)](1-\gamma)}{2r(4\log|\mathcal{A}| + C)} \right). \quad (49)$$

For any $s \in \mathcal{S}$, define $\mathcal{A}^*(s) = \operatorname{argmin}_{a \in \mathcal{A}} Q^*(s, a)$, recall that from Lemma 5.2, we can characterize the set of optimal policy Π^* by

$$\Pi^* = \left\{ \pi \in \Pi : \operatorname{supp}(\pi(\cdot|s)) = \operatorname{argmin}_{a \in \mathcal{A}} Q^*(s, a), \forall s \in \mathcal{S} \right\}. \quad (50)$$

If $\mathcal{A}^*(s) \neq \mathcal{A}$, we denote $\Delta^*(s) = \min_{a \notin \mathcal{A}^*(s)} Q^*(s, a) - \min_{a \in \mathcal{A}^*(s)} Q^*(s, a)$, and $\Delta^*(s) = \infty$ otherwise. Define the gap function of MDP \mathcal{M} as $\Delta^*(\mathcal{M}) = \min_{s \in \mathcal{S}} \Delta^*(s)$, then whenever $\Delta^*(\mathcal{M}) < \infty$, from our previous discussion, we know that for $k \geq K_1 := \lceil 3 \log_\gamma \left(\frac{\Delta^*(\mathcal{M})(1-\gamma)}{2r(4\log|\mathcal{A}| + C)} \right) \rceil$, we have

$$z_i^{k+1}(s) - z_j^{k+1}(s) \geq -\frac{2C\gamma^2}{(1-\gamma^3)(1-\gamma)} + \frac{\gamma^{-2k-1}}{2} \Delta^*(\mathcal{M}), \forall s \in \mathcal{S}.$$

Thus from the definition of (z_i^k, z_j^k) , we obtain that for any $j \notin \mathcal{A}^*(s)$, for $k \geq K_1$,

$$\begin{aligned}
\pi_k(j|s) &\leq \pi_k(i|s) \exp \left(\frac{2C\gamma^2}{(1-\gamma^3)(1-\gamma)} - \frac{\Delta^*(\mathcal{M})}{2} \gamma^{-2k-1} \right) \\
&\leq C_\gamma \exp \left(-\frac{\Delta^*(\mathcal{M})}{2} \gamma^{-2k-1} \right), \forall s \in \mathcal{S},
\end{aligned} \quad (51)$$

where $C_\gamma := \exp \left(\frac{2C\gamma^2}{(1-\gamma^3)(1-\gamma)} \right)$. Hence from the characterization of the optimal policy in (50), we can conclude that

$$\operatorname{dist}_{\ell_1}(\pi_k, \Pi^*) \leq 2C_\gamma |\mathcal{A}| \exp \left\{ -\frac{\Delta^*(\mathcal{M})}{2} \gamma^{-2k-1} \right\}, \quad \forall k \geq K_1,$$

Thus the policy π_k converges superlinearly to the optimal policy set with order γ^{-2} , after K_1 number of iterations. \square

Theorem (Restating Theorem 3.2). *We have the last-iterate convergence for the policy sequence $\{\pi_k\}_{k \geq 0}$ generated by HPMD, i.e., there exists $\pi^* \in \Pi$ such that*

$$\lim_{k \rightarrow \infty} \pi_k = \pi^*.$$

Furthermore, the converged policy equals to the optimal policy with the maximal entropy for every state $s \in \mathcal{S}$. That is

$$\lim_{k \rightarrow \infty} \pi_k(a|s) = \pi_U^*(a|s) := \begin{cases} 1/|\mathcal{A}^*(s)|, & a \in \mathcal{A}^*(s), \\ 0, & a \notin \mathcal{A}^*(s). \end{cases}$$

In particular, for any $0 < \epsilon < 1$, we have $\|\pi - \pi_U^\|_\infty \leq \epsilon$, as long as*

$$k \geq 2\bar{K}_1 + \log_\gamma \left(\frac{4r(4\log|\mathcal{A}| + C)}{\epsilon(1-\gamma)(1-\gamma^2)\gamma} \right) + 2 \log_\gamma \left(\frac{8\gamma C|\mathcal{A}|C_\gamma}{\epsilon(1-\gamma^{1/2})(1-\gamma)\gamma} \right) + \log_\gamma \left(\frac{2\log(C_\gamma|\mathcal{A}|/\epsilon)}{\Delta^*(\mathcal{M})} \right),$$

where

$$\bar{K}_1 = \min_k \left\{ k > K_1 : \Delta^*(\mathcal{M})\gamma^{-2t-1} \geq 5t \log\left(\frac{1}{\gamma}\right) \right\}.$$

Proof of Theorem 3.2. We consider the scenario where $Q^*(s, i) = Q^*(s, j)$. Similar to the proof of Theorem 3.1, we have

$$z_i^{k+1} - z_j^{k+1} = \gamma^{k+1}(z_i^0 - z_j^0) - \sum_{t=0}^k \gamma^{k+1-t} \eta_t (g_i^t - g_j^t) = - \sum_{t=0}^k \gamma^{k+1-t} \eta_t (g_i^t - g_j^t). \quad (52)$$

Recall that from Corollary 3.1, we have

$$\begin{aligned} Q^{\pi^k}(s, a) - Q^*(s, a) &= \gamma \sum_{s' \in \mathcal{S}} \mathcal{P}(s'|s, a) (V^\pi(s) - V^*(s)) \leq \frac{\gamma C |\mathcal{A}|}{(1-\gamma)^2} \text{dist}_{\ell_\infty}(\pi, \Pi^*) \\ &\leq \frac{\gamma C |\mathcal{A}| C_\gamma}{(1-\gamma)^2} \exp\left(-\frac{\Delta^*(\mathcal{M})}{2} \gamma^{-2k-1}\right), \end{aligned} \quad (53)$$

whenever $k \geq K_1$. Thus for any $\bar{K}_1 \geq K_1$, we have

$$\sum_{t=0}^k \gamma^{k+1-t} \eta_t |g_i^t - g_j^t| \leq \underbrace{\sum_{t=0}^{\bar{K}_1} \gamma^{k+1-t} \eta_t |g_i^t - g_j^t|}_{(A)} + \underbrace{\sum_{t=\bar{K}_1+1}^k \gamma^{k+1-t} \eta_t |g_i^t - g_j^t|}_{(B)}. \quad (54)$$

We proceed to first bound term (A) in (54). Note that from (44) and the definition of $\eta_t = \gamma^{-2(t+1)}$,

$$\begin{aligned} \gamma^{k+1-t} \eta_t |g_i^t - g_j^t| &\leq \gamma^{k+1-t} \eta_t (|g_i^t - Q^*(s, i)| + |g_j^t - Q^*(s, j)|) \\ &\leq 2\gamma^{k-3t-1} \cdot r \gamma^t \frac{4 \log |\mathcal{A}| + C}{1-\gamma} = \gamma^{k-2t} \frac{2r(4 \log |\mathcal{A}| + C)}{(1-\gamma)\gamma}. \end{aligned}$$

Hence we can bound term (A) as

$$(A) = \sum_{t=0}^{\bar{K}_1} \gamma^{k+1-t} \eta_t |g_i^t - g_j^t| = \sum_{t=0}^{\bar{K}_1} \gamma^{k-2t} \frac{2r(4 \log |\mathcal{A}| + C)}{(1-\gamma)\gamma} = \gamma^{k-2\bar{K}_1+2} \frac{2r(4 \log |\mathcal{A}| + C)}{(1-\gamma)(1-\gamma^2)\gamma}. \quad (55)$$

We then proceed to bound term (B) in (54). From the definition of $\eta_t = \gamma^{-2(t+1)}$, the superlinear convergence of Q-function in (53), and the condition that $Q^*(s, i) = Q^*(s, j)$,

$$\begin{aligned} \sum_{t=\bar{K}_1}^k \gamma^{k+1-t} \eta_t |g_i^t - g_j^t| &\leq \sum_{t=\bar{K}_1}^k \gamma^{k+1-t} \eta_t (|g_i^t - Q^*(s, i)| + |g_j^t - Q^*(s, j)|) \\ &\leq \sum_{t=\bar{K}_1}^k \gamma^{k-3t} \frac{4\gamma C |\mathcal{A}| C_\gamma}{(1-\gamma)\gamma} \exp\left(-\frac{\Delta^*(\mathcal{M})}{2} \gamma^{-2t-1}\right) \\ &= \sum_{t=\bar{K}_1}^k \gamma^{k-t/2} \frac{4\gamma C |\mathcal{A}| C_\gamma}{(1-\gamma)\gamma} \exp\left(-\frac{\Delta^*(\mathcal{M})}{2} \gamma^{-2t-1} + \frac{5t}{2} \log\left(\frac{1}{\gamma}\right)\right) \end{aligned} \quad (56)$$

It suffices to choose \bar{K}_1 as

$$\bar{K}_1 = \min_k \left\{ k > K_1 : \Delta^*(\mathcal{M}) \gamma^{-2k-1} \geq 5k \log\left(\frac{1}{\gamma}\right) \right\},$$

Then for any $k \geq \bar{K}_1$, from (56), we obtain

$$(B) = \sum_{t=\bar{K}_1}^k \gamma^{k+1-t} \eta_t |g_i^t - g_j^t| \leq \sum_{t=\bar{K}_1}^k \gamma^{k-t/2} \frac{4\gamma C |\mathcal{A}| C_\gamma}{(1-\gamma)\gamma} \leq \gamma^{(k+1)/2} \frac{4\gamma C |\mathcal{A}| C_\gamma}{(1-\gamma^{1/2})(1-\gamma)\gamma} \quad (57)$$

Thus, by combing (55) and (57), we have for any $k \geq \bar{K}_1$,

$$\sum_{t=0}^k \gamma^{k+1-t} \eta_t |g_i^t - g_j^t| \leq \gamma^{k-2\bar{K}_1+2} \frac{2r(4 \log |\mathcal{A}| + C)}{(1-\gamma)(1-\gamma^2)\gamma} + \gamma^{(k+1)/2} \frac{4\gamma C |\mathcal{A}| C_\gamma}{(1-\gamma^{1/2})(1-\gamma)\gamma}. \quad (58)$$

Fixing \bar{K}_1 , and taking $k \rightarrow \infty$ in the previous relation, we obtain

$$\lim_{k \rightarrow \infty} \sum_{t=0}^k \gamma^{k+1-t} \eta_t |g_i^t - g_j^t| \rightarrow 0. \quad (59)$$

Since $\{\pi_k\}_{k \geq 0} \subset \Pi$ and Π is a compact set, hence $\{\pi_k\}_{k \geq 0} \subset \Pi$ has at least one limit point. Let π^* denote a limit point of the policy iterate $\{\pi_k\}_{k \geq 0}$, i.e., there exists subsequence $\{\pi_{k_l}\}_{l \geq 0}$ such that $\lim_{l \rightarrow \infty} \pi_{k_l} = \pi^*$. Denoting $z_i^*(s) = \log \pi^*(i|s)$ for any $i \in \mathcal{A}$, we will write z_i^* in short for $z_i^*(s)$ when the context is clear. Then from (52), (59) and the definition of (z_i^k, z_j^k) , we have

$$\frac{\pi^*(i|s)}{\pi^*(j|s)} = \exp(\log z_i^* - \log z_j^*) = \lim_{l \rightarrow \infty} \exp(\log z_i^{k_l} - \log z_j^{k_l}) = 1, \quad \forall i, j \in \mathcal{A}^*(s).$$

Combined with (51), we can characterize the limiting policy π^* as

$$\pi^*(a|s) = \pi_U^*(a|s) := \begin{cases} 1/|\mathcal{A}^*(s)|, & a \in \mathcal{A}^*(s), \\ 0, & a \notin \mathcal{A}^*(s). \end{cases} \quad (60)$$

Finally, note that since (60) characterizes all the limiting policies, we obtain the last-iterate convergence of the policy

$$\lim_{k \rightarrow \infty} \pi_k = \pi_U^*.$$

In addition, the finite-time convergence characterization can be readily obtained by combining (51) and (58). Specifically, to ensure $\|\pi_k - \pi_U^*\|_\infty \leq \epsilon$, we need $k \geq K_1$. Then for $k \geq \log_\gamma \left(\frac{2 \log(C_\gamma |\mathcal{A}|/\epsilon)}{\Delta^*(\mathcal{M})} \right)$, from (51), we have

$$\pi(j|s) \leq \frac{\epsilon}{|\mathcal{A}|}, \quad \forall s \in \mathcal{S}, j \notin \mathcal{A}^*(s).$$

On the other hand, for any $\epsilon < 1$, and any

$$k \geq 2\bar{K}_1 + \log_\gamma \left(\frac{4r(4 \log |\mathcal{A}| + C)}{\epsilon(1-\gamma)(1-\gamma^2)\gamma} \right) + 2 \log_\gamma \left(\frac{8\gamma C |\mathcal{A}| C_\gamma}{\epsilon(1-\gamma^{1/2})(1-\gamma)\gamma} \right),$$

then $|\pi(i|s) - 1/|\mathcal{A}^*(s)|| \leq \epsilon$ for any $i \in \mathcal{A}^*(s)$. Hence we can conclude that for any

$$k \geq 2\bar{K}_1 + \log_\gamma \left(\frac{4r(4 \log |\mathcal{A}| + C)}{\epsilon(1-\gamma)(1-\gamma^2)\gamma} \right) + 2 \log_\gamma \left(\frac{8\gamma C |\mathcal{A}| C_\gamma}{\epsilon(1-\gamma^{1/2})(1-\gamma)\gamma} \right) + \log_\gamma \left(\frac{2 \log(C_\gamma |\mathcal{A}|/\epsilon)}{\Delta^*(\mathcal{M})} \right),$$

then $\|\pi - \pi_U^*\|_\infty \leq \epsilon$, where

$$\bar{K}_1 = \min_k \left\{ k > K_1 : \Delta^*(\mathcal{M}) \gamma^{-2k-1} \geq 5t \log \left(\frac{1}{\gamma} \right) \right\}.$$

□

B.2 Stochastic HPMD

We proceed to show the policy convergence of the SHPMD method, which follows the similar analyses as the deterministic HPMD method.

Theorem (Restating Theorem 4.3). *We define $\Delta^*(s) = \min_{a \notin \mathcal{A}^*(s)} Q^*(s, a) - \min_{a \in \mathcal{A}^*(s)} Q^*(s, a)$ if $\mathcal{A}^*(s) \neq \mathcal{A}$, and $\Delta^*(s) = \infty$ otherwise. Define the gap function of the MDP \mathcal{M} as $\Delta^*(\mathcal{M}) = \min_{s \in \mathcal{S}} \Delta^*(s)$. Then whenever $\Delta^*(\mathcal{M}) < \infty$, the iterate π_k generated by SHPMD satisfies*

$$\mathbb{E} [\text{dist}_{\ell_1}(\pi_k, \Pi^*)] \leq 2C_\gamma^n |\mathcal{A}| \exp \left(-\frac{\Delta^*(\mathcal{M}) \sqrt{\log |\mathcal{A}|}}{2} \gamma^{-k/2} \right), \quad \forall k \geq K_2,$$

for any iteration $k \geq K_2 := \lceil 3 \log_\gamma \left(\frac{\Delta^*(\mathcal{M})(1-\gamma)}{2r(6 \log |\mathcal{A}| + C)} \right) \rceil$, and $C_\gamma^n := \exp \left(\frac{(2C+1)\gamma^2 \sqrt{\log |\mathcal{A}|}}{(1-\gamma)(1-\gamma^{3/4})} \right)$. On the other hand, if $\Delta^*(\mathcal{M}) = \infty$, then any policy is an optimal policy.

Proof of Theorem 4.3. Recall that from Assumption 1, there exists π^* such that ν^{π^*} has full support. From the definition of state-action value function, we have

$$Q^{\pi_k}(s, a) - Q^*(s, a) = \gamma \sum_{s' \in \mathcal{S}} \mathcal{P}(s'|s, a) [V^{\pi_k}(s') - V^*(s')].$$

Hence we can conclude for $r := \gamma \max_{s \in \mathcal{S}, a \in \mathcal{A}} \max_{s' \in \mathcal{S}} \mathcal{P}(s'|s, a) / \nu^{\pi^*}(s')$,

$$\text{supp}(\mathcal{P}(\cdot|s, a)) \subseteq \nu^{\pi^*} \Rightarrow \mathbb{E}[Q^{\pi_k}(s, a) - Q^*(s, a)] \leq \gamma^{k/2} \frac{r(5 \log|\mathcal{A}| + C)}{1 - \gamma}, \quad (61)$$

where C denotes the upper bound of the cost.

We begin by inspecting the update rule of SHPMD. Directly examining the optimality condition of the update (13), we obtain

$$\log \pi_{k+1}(a|s) = \frac{\log \pi_k(a|s) - \eta_k Q^{\pi_k, \xi_k}(s, a)}{\eta_k \tau_k + 1} + \Delta_k(s), \quad \forall (s, a) \in \mathcal{S} \times \mathcal{A}, \quad (62)$$

where $\Delta_k(s)$ denotes the normalization constant to make the right hand size of (45) the logarithm of a valid probability vector defined over \mathcal{A} . Combined the choice of $\{(\tau_k, \eta_k)\}_{k \geq 0}$ satisfying $1 + \tau_k \eta_k = 1/\gamma$, we have

$$\log \pi_{k+1}(a|s) = \gamma [\log \pi_k(a|s) - \eta_k Q^{\pi_k, \xi_k}(s, a)] + \Delta_k(s), \quad \forall (s, a) \in \mathcal{S} \times \mathcal{A}. \quad (63)$$

Define $z_i^k(s) = \log \pi_k(i|s)$ for any pair $(s, i) \in \mathcal{S} \times \mathcal{A}$. For the ease of presentation, let us fix a state $s \in \mathcal{S}$ and write z_i^k in short for $z_i^k(s)$, and Δ_k in short for $\Delta_k(s)$, when the context is clear. We will also denote g_i^k in short for $Q^{\pi_k}(s, i)$, \widehat{g}_i^k in short for $Q^{\pi_k, \xi_k}(s, i)$, and accordingly define $\delta_i^k = \widehat{g}_i^k - g_i^k$. Then applying (63) recursively, we obtain that for any $i \in \mathcal{A}$,

$$z_i^{k+1} = \gamma^{k+1} z_i^0 - \sum_{t=0}^k \gamma^{k+1-t} \eta_t \widehat{g}_i^t + \sum_{t=0}^k \gamma^{k-t} \Delta_t.$$

Hence for any pair of actions $i, j \in \mathcal{A}$, we have

$$z_i^{k+1} - z_j^{k+1} = \gamma^{k+1} (z_i^0 - z_j^0) - \sum_{t=0}^k \gamma^{k+1-t} \eta_t (\widehat{g}_i^t - \widehat{g}_j^t) = - \sum_{t=0}^k \gamma^{k+1-t} \eta_t (\widehat{g}_i^t - \widehat{g}_j^t). \quad (64)$$

Recall that from Lemma 5.2, it suffices to characterize the convergence of $\pi_k(a|s)$ for $a \notin \mathcal{A}^*(s)$. It then suffices to consider any pair of action i, j satisfying $Q^*(s, i) < Q^*(s, j)$. Combining with the observation in (61), we conclude that there exists $\underline{K}_2(s)$ such that

$$\mathbb{E}[g_i^t - g_j^t] \leq (Q^*(s, i) - Q^*(s, j))/2, \quad \forall t \geq \underline{K}_2(s). \quad (65)$$

Combined with the observation that $\mathbb{E}[\delta_i^t] \leq \|\mathbb{E}\delta_t\|_\infty = \varepsilon_t = \gamma^{3(k+1)/4}$, and $\eta_k = \gamma^{-(k+1)/2} \sqrt{\log|\mathcal{A}|}$ in SHPMD, for any $k \geq \underline{K}_2(s)$, we have

$$\begin{aligned} \mathbb{E}\pi_j^{k+1} &\leq \mathbb{E} \exp \left\{ \sum_{t=0}^k \gamma^{k+1-t} \eta_t (\widehat{g}_i^t - \widehat{g}_j^t) \right\} \\ &\leq \exp \left\{ \sum_{t=0}^k \gamma^{k+1-t} \mathbb{E}[g_i^t - g_j^t] + \sum_{t=0}^k \gamma^{k+1-t} \mathbb{E}[\delta_i^t - \delta_j^t] \right\} \\ &\leq \exp \left\{ \sum_{t=0}^{\underline{K}_2(s)} \gamma^{k+1-t} \eta_t \mathbb{E}[g_i^t - g_j^t] + \sum_{t=\underline{K}_2(s)+1}^k \gamma^{k+1-t} \eta_t \mathbb{E}[g_i^t - g_j^t] + \sum_{t=0}^k \gamma^{k+1-t} \eta_t (\|\mathbb{E}\delta_i^t\| + \|\mathbb{E}\delta_j^t\|) \right\} \\ &\leq \exp \left\{ \sum_{t=0}^{\underline{K}_2(s)} \gamma^{k-3t/2} \sqrt{\gamma \log|\mathcal{A}|} \mathbb{E}[g_i^t - g_j^t] + \sum_{t=\underline{K}_2(s)}^k \gamma^{k-3t/2} \sqrt{\gamma \log|\mathcal{A}|} \mathbb{E}[g_i^t - g_j^t] \right\} \end{aligned}$$

$$\begin{aligned}
& + 2 \sum_{t=0}^k \gamma^{k-\frac{3}{4}t} \sqrt{\log|\mathcal{A}|} \gamma^{5/4} \Big\} \\
& = \exp \left\{ \sum_{t=0}^{\underline{K}_2(s)} \gamma^{k-3t/2} \cdot \frac{2C\sqrt{\gamma\log|\mathcal{A}|}}{1-\gamma} - \gamma^{-k/2} \frac{\sqrt{\gamma\log|\mathcal{A}|}}{2} [Q^*(s,j) - Q^*(s,i)] - \gamma^{k/4+2} \frac{\sqrt{\log|\mathcal{A}|}}{1-\gamma^{3/4}} \right\} \\
& \leq \exp \left\{ \frac{2C\gamma^{k-\frac{3}{2}\underline{K}_2(s)}\gamma^2\sqrt{\log|\mathcal{A}|}}{(1-\gamma)(1-\gamma^{3/2})} - \gamma^{-k/2} \frac{\sqrt{\gamma\log|\mathcal{A}|}}{2} [Q^*(s,j) - Q^*(s,i)] + \gamma^{k/4+2} \frac{\sqrt{\log|\mathcal{A}|}}{1-\gamma^{3/4}} \right\}
\end{aligned}$$

Thus for any $k \geq 3\underline{K}_2(s)/2$, we obtain

$$\mathbb{E}[\pi_{k+1}(j|s)] \leq \exp \left\{ \frac{(2C+1)\gamma^2\sqrt{\log|\mathcal{A}|}}{(1-\gamma)(1-\gamma^{3/4})} - \gamma^{-k/2} \frac{\sqrt{\gamma\log|\mathcal{A}|}}{2} [Q^*(s,j) - Q^*(s,i)] \right\}.$$

It remains to bound $\underline{K}_2(s)$ that satisfies (65). From the observation (61), to satisfy (65), it suffices to choose

$$\underline{K}_2(s) \geq 2 \log_\gamma \left(\frac{[Q^*(s,j) - Q^*(s,i)](1-\gamma)}{2r(6\log|\mathcal{A}|+C)} \right). \quad (66)$$

For any $s \in \mathcal{S}$, define $\mathcal{A}^*(s) = \operatorname{argmin}_{a \in \mathcal{A}} Q^*(s, a)$, recall that from Lemma 5.2, we can characterize the set of optimal policy Π^* by

$$\Pi^* = \left\{ \pi \in \Pi : \operatorname{supp}(\pi(\cdot|s)) = \operatorname{argmin}_{a \in \mathcal{A}} Q^*(s, a), \forall s \in \mathcal{S} \right\}.$$

If $\mathcal{A}^*(s) \neq \mathcal{A}$, we denote $\Delta^*(s) = \min_{a \notin \mathcal{A}^*(s)} Q^*(s, a) - \min_{a \in \mathcal{A}^*(s)} Q^*(s, a)$, and $\Delta^*(s) = \infty$ otherwise. Define the gap function of MDP \mathcal{M} as $\Delta^*(\mathcal{M}) = \min_{s \in \mathcal{S}} \Delta^*(s)$, then whenever $\Delta^*(\mathcal{M}) < \infty$, from our previous discussion, we know that for $k \geq K_2 := \lceil 3 \log_\gamma \left(\frac{\Delta^*(\mathcal{M})(1-\gamma)}{2r(6\log|\mathcal{A}|+C)} \right) \rceil$ and any $j \notin \mathcal{A}^*(s)$,

$$\begin{aligned}
\mathbb{E}[\pi_{k+1}(j|s)] & \leq \exp \left(\frac{(2C+1)\gamma^2\sqrt{\log|\mathcal{A}|}}{(1-\gamma)(1-\gamma^{3/4})} - \frac{\gamma^{-k/2}\sqrt{\gamma\log|\mathcal{A}|}}{2} \Delta^*(\mathcal{M}) \right) \\
& = C_\gamma^n \exp \left(-\frac{\gamma^{-k/2}\sqrt{\gamma\log|\mathcal{A}|}}{2} \Delta^*(\mathcal{M}) \right), \forall s \in \mathcal{S}
\end{aligned}$$

where $C_\gamma^n := \exp \left(\frac{(2C+1)\gamma^2\sqrt{\log|\mathcal{A}|}}{(1-\gamma)(1-\gamma^{3/4})} \right)$. Hence from the characterization of the optimal policy in (50), we can conclude that

$$\mathbb{E}[\operatorname{dist}_{\ell_1}(\pi_k, \Pi^*)] \leq 2C_\gamma^n |\mathcal{A}| \exp \left(-\frac{\Delta^*(\mathcal{M})\sqrt{\log|\mathcal{A}|}}{2} \gamma^{-k/2} \right), \forall k \geq K_2,$$

Thus the policy π_k converges superlinearly to the optimal policy set with order $\gamma^{-1/2}$, after K_2 number of iterations. \square

Corollary (Restating Corollary 4.1). *For any policy π , let $\pi^* = \operatorname{argmin}_{\pi' \in \Pi} \|\pi' - \pi\|_\infty$ denotes the its projection onto Π^* in ℓ_∞ -norm. Then we have $V^\pi(s) - V^{\pi^*}(s) \leq \frac{C|\mathcal{A}|}{(1-\gamma)^2} \|\pi - \pi^*\|_\infty$. In particular, for $k \geq K_2$ where K_2 is defined as in Theorem 4.3, the iterate π_k in the SHPMD method satisfies*

$$\mathbb{E} \left[V^{\pi_k}(s) - V^{\pi^*}(s) \right] \leq \frac{2C|\mathcal{A}|C_\gamma^n}{(1-\gamma)^2} \exp \left(-\frac{\Delta^*(\mathcal{M})\sqrt{\log|\mathcal{A}|}}{2} \gamma^{-k/2} \right).$$

The proof of Corollary 4.1 follows from the exact same lines as in Corollary 3.1.

C Improved Sample Complexity

Theorem (Restating Theorem 4.4). *By choosing $\{(\tau_k, \eta_k, M_k, T_k)\}_{k \geq 0}$ the same as in Theorem 4.1 and Theorem 4.2, for any precision $\epsilon > 0$, the total number of iterations $k(\epsilon)$ for finding an ϵ -optimal policy $\mathbb{E} [f(\pi_{k(\epsilon)}) - f(\pi^*)] \leq \epsilon$ can be bounded by*

$$k(\epsilon) = \begin{cases} \mathcal{O}(\log_\gamma(\epsilon(1-\gamma))), & \epsilon \geq \epsilon_0, \\ \mathcal{O}\left(\log_\gamma(\epsilon_0(1-\gamma)) + \log_\gamma\left(\frac{\Delta^*(\mathcal{M})\sqrt{\log|\mathcal{A}|}}{\log(C_\gamma^n/\epsilon)}\right)\right), & \epsilon < \epsilon_0. \end{cases}$$

In addition, the total number of sample complexity $N(\epsilon)$ can be bounded by

$$N(\epsilon) = \begin{cases} \mathcal{O}\left(\frac{|\mathcal{S}||\mathcal{A}|\log_\gamma(\epsilon(1-\gamma))}{(1-\gamma)^5(1-\gamma^{1/2})^2\epsilon_0^2}\right), & \epsilon \geq \epsilon_0, \\ \mathcal{O}\left(\frac{|\mathcal{S}||\mathcal{A}|\log_\gamma(\epsilon(1-\gamma))}{(1-\gamma)^5(1-\gamma^{1/2})^2\epsilon_0^2} \left[1 + \frac{\log^2(1/\epsilon)}{\log|\mathcal{A}|\Delta^*(\mathcal{M})^2} + \log_{1/\gamma}\left(\frac{\log(\frac{1}{\epsilon})}{\Delta^*(\mathcal{M})}\right)\right]\right), & \epsilon < \epsilon_0, \end{cases}$$

where $\epsilon_0 = \Delta^*(\mathcal{M})^{3/2}(1-\gamma)^{1/2} / \left[(1-\gamma^{1/2})\gamma\sqrt{6\log|\mathcal{A}| + C}\right]$.

Proof of Theorem 4.4. Note that from the definition of ϵ_0, K_2 and the global linear convergence of SHPMD, for any $\epsilon < \epsilon_0$, the number of iterations $k(\epsilon)$ required can be larger than K_2 . From the local superlinear convergence in (21), it suffices to have

$$k(\epsilon) = K_2 + 2 \log_\gamma\left(\frac{\Delta^*(\mathcal{M})\sqrt{\log|\mathcal{A}|}}{2 \log(C|\mathcal{A}|C_\gamma^n/\epsilon(1-\gamma))}\right). \quad (67)$$

Thus we have the total number of samples can be bounded by

$$\begin{aligned} |\mathcal{S}||\mathcal{A}|\sum_{s=0}^{k(\epsilon)} M_s T_s &= |\mathcal{S}||\mathcal{A}|\left(\sum_{s=0}^{T_2} M_s T_s + \sum_{s=T_2+1}^{k(\epsilon)} M_s T_s\right) \\ &= \mathcal{O}\left(\frac{|\mathcal{S}||\mathcal{A}|\log_\gamma(\epsilon_0(1-\gamma))}{(1-\gamma)^5(1-\gamma^{1/2})^2\epsilon_0^2}\right) + \mathcal{O}\left(\frac{|\mathcal{S}||\mathcal{A}|}{(1-\gamma)^3}\gamma^{-k(\epsilon)}k(\epsilon)\right) \\ &= \mathcal{O}\left(\frac{|\mathcal{S}||\mathcal{A}|\log_\gamma(\epsilon_0(1-\gamma))}{(1-\gamma)^5(1-\gamma^{1/2})^2\epsilon_0^2}\right) \\ &\quad + \frac{|\mathcal{S}||\mathcal{A}|}{(1-\gamma)^3}\gamma^{-K_2}\mathcal{O}\left(K_2\gamma^{-(k(\epsilon)-K_2)} + (k(\epsilon) - K_2)\gamma^{-(k(\epsilon)-K_2)}\right) \end{aligned} \quad (68)$$

From the definition of K_2 , we have

$$\begin{aligned} \gamma^{-K_2} &= \mathcal{O}\left(\frac{1}{(1-\gamma)^2(1-\gamma^{1/2})^2\gamma^2\epsilon_0^2}\right), \\ K_2 &= \mathcal{O}\left(\log_\gamma(\epsilon_0(1-\gamma)(1-\gamma^{1/2})\gamma)\right), \\ \gamma^{-(k(\epsilon)-K_2)} &= 4 \log^2\left(\frac{C|\mathcal{A}|C_\gamma^n}{\epsilon(1-\gamma)}\right) / (\log|\mathcal{A}|\Delta^*(\mathcal{M})^2). \end{aligned}$$

By combining (67), (68) and the previous relation, we obtain

$$\begin{aligned} |\mathcal{S}||\mathcal{A}|\sum_{s=0}^{k(\epsilon)} M_s T_s &= \mathcal{O}\left(\frac{|\mathcal{S}||\mathcal{A}|\log_\gamma(\epsilon_0(1-\gamma))}{(1-\gamma)^5(1-\gamma^{1/2})^2\epsilon_0^2} \left[1 + \frac{4 \log^2\left(\frac{C|\mathcal{A}|C_\gamma^n}{\epsilon(1-\gamma)}\right)}{\log|\mathcal{A}|\Delta^*(\mathcal{M})^2} \log_\gamma\left(\frac{\Delta^*(\mathcal{M})\sqrt{\log|\mathcal{A}|}}{2 \log(C|\mathcal{A}|C_\gamma^n/\epsilon(1-\gamma))}\right)\right]\right) \\ &= \mathcal{O}\left(\frac{|\mathcal{S}||\mathcal{A}|\log_\gamma(\epsilon_0(1-\gamma))}{(1-\gamma)^5(1-\gamma^{1/2})^2\epsilon_0^2} \left[1 + \frac{\log^2(1/\epsilon)}{\log|\mathcal{A}|\Delta^*(\mathcal{M})^2} + \log_{1/\gamma}\left(\frac{\log(\frac{1}{\epsilon})}{\Delta^*(\mathcal{M})}\right)\right]\right). \end{aligned}$$

□

Theorem (Restating Theorem 4.5). *If the parameter choice $\{(\eta_t, \tau_t, \epsilon_t, \sigma_t)\}_{t \geq 0}$ satisfies*

$$1 + \eta_t \tau_t = 1/\gamma, \sigma_t = \gamma^{t/2}, \eta_t \sigma_t = \gamma^{\beta t} \sqrt{\log|\mathcal{A}|}, \epsilon_t = \gamma^{\frac{3}{4}t}, \quad 0 < \beta < 1/2,$$

Then with probability 1, we have the last-iterate convergence for the policy sequence $\{\pi_k\}_{k \geq 0}$ generated by SHPMD, i.e., there exists $\pi^ \in \Pi$ such that $\lim_{k \rightarrow \infty} \pi_k = \pi^*$, a.s. Furthermore, the converged policy equals to the optimal policy with the maximal entropy for every state $s \in \mathcal{S}$. That is*

$$\lim_{k \rightarrow \infty} \pi_k(a|s) = \pi_U^*(a|s) := \begin{cases} 1/|\mathcal{A}^*(s)|, & a \in \mathcal{A}^*(s), \\ 0, & a \notin \mathcal{A}^*(s), \end{cases} \quad a.s.$$

Proof of Theorem 4.5. We begin by establishing the convergence property of the SHPMD method with the specific choice of $\{(\eta_t, \tau_t, \epsilon_t, \sigma_t)\}_{t \geq 0}$ in (22). Specifically, from Lemma A.4, by choosing $\alpha_t = \gamma^{-t}$, one can readily verify that condition (36) is verified with

$$\gamma_t = \gamma \alpha_{t+1}, \quad \frac{\alpha_t}{\gamma \eta_t} > \frac{\alpha_{t+1}}{\eta_{t+1}}.$$

In addition, we have

$$\sum_{t=0}^k \frac{\alpha_t}{\eta_t} \log|\mathcal{A}| = \frac{\gamma^{-(1/2+\beta)k}}{1 - \gamma^{1/2+\beta}}, \quad \sum_{t=0}^k \alpha_t \eta_t \sigma_t^2 = \frac{\gamma^{-k/2}}{1 - \gamma^{1/2}} \sqrt{\log|\mathcal{A}|}, \quad \sum_{t=0}^k \alpha_t \epsilon_t = \frac{\gamma^{-k/4}}{1 - \gamma^{1/4}}.$$

Combining the above relation with Lemma (A.4), we obtain that

$$\mathbb{E}[f(\pi_k) - f(\pi^*)] \leq \gamma^{\frac{1}{2}-\beta k} \left[f(\pi_0) - f(\pi^*) + \frac{6 \log|\mathcal{A}|}{\gamma(1-\gamma)(1-\gamma^{1/2})} \right]. \quad (69)$$

Following similar lines as in the proof of Theorem 4.3, we can show that local superlinear convergence of SHPMD with the particular parameter choice. Specifically, for any $s \in \mathcal{S}$, and $i, j \in \mathcal{A}$ with $Q^*(s, i) < Q^*(s, j)$, we can show that there exists $\underline{K}_3(s)$ such that

$$\mathbb{E}[\pi_{k+1}(j|s)] \leq \exp \left\{ \frac{2C\gamma^{k-(3/2-\beta)\underline{K}_3(s)} \sqrt{\log|\mathcal{A}|}}{(1-\gamma)(1-\gamma^{3/2-\beta})} - \frac{\gamma^{-(1/2-\beta)k}}{2} [(Q^*(s, j) - Q^*(s, i))] \sqrt{\log|\mathcal{A}|} + \frac{\gamma^{k/4}}{1-\gamma^{3/4}} \sqrt{\log|\mathcal{A}|} \right\}.$$

where it suffices for $\underline{K}_3(s)$ to satisfy

$$\underline{K}_3(s) \geq \frac{2}{1-\beta} \log_\gamma \left(\frac{[Q^*(s, j) - Q^*(s, i)](1-\gamma)}{2r(6 \log|\mathcal{A}| + C)} \right).$$

Hence for any $k \geq (3/2 - \beta)\underline{K}_3(s)$, we obtain

$$\mathbb{E}[\pi_{k+1}(j|s)] \leq \exp \left\{ \frac{2C+1}{(1-\gamma)(1-\gamma^{3/4})} \sqrt{\log|\mathcal{A}|} - \frac{\gamma^{-(1/2-\beta)k}}{2} [(Q^*(s, j) - Q^*(s, i))] \sqrt{\log|\mathcal{A}|} \right\}.$$

For any $s \in \mathcal{S}$, define $\mathcal{A}^*(s) = \operatorname{argmin}_{a \in \mathcal{A}} Q^*(s, a)$, recall that from Lemma 5.2, we can characterize the set of optimal policy Π^* by $\Pi^* = \{\pi \in \Pi : \operatorname{supp}(\pi(\cdot|s)) = \operatorname{argmin}_{a \in \mathcal{A}} Q^*(s, a), \forall s \in \mathcal{S}\}$.

If $\mathcal{A}^*(s) \neq \mathcal{A}$, we denote $\Delta^*(s) = \min_{a \notin \mathcal{A}^*(s)} Q^*(s, a) - \min_{a \in \mathcal{A}^*(s)} Q^*(s, a)$, and $\Delta^*(s) = \infty$ otherwise. Define the gap function of MDP \mathcal{M} as $\Delta^*(\mathcal{M}) = \min_{s \in \mathcal{S}} \Delta^*(s)$, then whenever $\Delta^*(\mathcal{M}) < \infty$, from our previous discussion, we know that for $k \geq K_3 := \lceil \frac{2}{1-\beta} \log_\gamma \left(\frac{\Delta^*(\mathcal{M})(1-\gamma)}{2r(6 \log|\mathcal{A}| + C)} \right) \rceil$ and any $j \notin \mathcal{A}^*(s)$,

$$\begin{aligned} \mathbb{E}[\pi_{k+1}(j|s)] &\leq \exp \left(\frac{(2C+1)\sqrt{\log|\mathcal{A}|}}{(1-\gamma)(1-\gamma^{3/4})} - \frac{\gamma^{-(1/2-\beta)k} \sqrt{\gamma \log|\mathcal{A}|}}{2} \Delta^*(\mathcal{M}) \right) \\ &= C_\gamma^n \exp \left(-\frac{\gamma^{-(1/2-\beta)k} \sqrt{\log|\mathcal{A}|}}{2} \Delta^*(\mathcal{M}) \right), \quad \forall s \in \mathcal{S} \end{aligned} \quad (70)$$

where $C_\gamma^n := \exp\left(\frac{(2C+1)\sqrt{\log|\mathcal{A}|}}{(1-\gamma)(1-\gamma^{3/4})}\right)$. Hence from the characterization of the optimal policy in (50), we can conclude that

$$\mathbb{E}[\text{dist}_{\ell_1}(\pi_k, \Pi^*)] \leq 2C_\gamma^n |\mathcal{A}| \exp\left(-\frac{\Delta^*(\mathcal{M})\sqrt{\log|\mathcal{A}|}}{2}\gamma^{-k/2}\right), \quad \forall k \geq K_3.$$

Thus the policy π_k converges superlinearly to the optimal policy set with order $\gamma^{-1/2}$, after K_3 number of iterations.

Similar to the proof of Theorem 4.3, it suffices to consider scenario for $i, j \in \mathcal{A}$, where $Q^*(s, i) = Q^*(s, j)$. Note that for any $\bar{K}_3 < k$,

$$\begin{aligned} \mathbb{E}|z_j^{k+1} - z_i^{k+1}| &= \sum_{t=0}^k \gamma^{k+1-t} \mathbb{E}|g_i^t - g_j^t| + \sum_{t=0}^k \gamma^{k+1-t} \mathbb{E}|\delta_i^t - \delta_j^t| \\ &= \underbrace{\sum_{t=0}^{\bar{K}_3} \gamma^{k+1-t} \eta_t \mathbb{E}|g_i^t - g_j^t|}_{(A)} + \underbrace{\sum_{t=\bar{K}_3+1}^k \gamma^{k+1-t} \eta_t \mathbb{E}|g_i^t - g_j^t|}_{(B)} + \underbrace{\sum_{t=0}^k \gamma^{k+1-t} \eta_t \mathbb{E}[|\delta_i^t| + |\delta_j^t|]}_{(C)}. \end{aligned} \quad (71)$$

We proceed to bound each term in the previous relation. Note that from (61) and the definition of $\eta_t = \gamma^{(\beta-1/2)t} \sqrt{\log|\mathcal{A}|}$,

$$\begin{aligned} \gamma^{k+1-t} \eta_t |g_i^t - g_j^t| &\leq \gamma^{k+1-t} \eta_t (|g_i^t - Q^*(s, i)| + |g_j^t - Q^*(s, j)|) \\ &\leq \gamma^{k-(3/2-\beta)t} \frac{2r(6 \log|\mathcal{A}| + C) \sqrt{\log|\mathcal{A}|}}{1-\gamma}. \end{aligned}$$

Hence we can bound term (A) as

$$(A) = \sum_{t=0}^{\bar{K}_3} \gamma^{k+1-t} \eta_t |g_i^t - g_j^t| = \frac{2r(6s \log|\mathcal{A}| + C) \sqrt{\gamma \log|\mathcal{A}|}}{(1-\gamma)^2} \gamma^{k-(3/2-\beta)\bar{K}_3} \quad (72)$$

Recall that from Corollary 4.1, we have

$$\begin{aligned} \mathbb{E}[Q^{\pi_k}(s, a) - Q^*(s, a)] &= \gamma \sum_{s' \in \mathcal{S}} \mathcal{P}(s'|s, a) \mathbb{E}[V^\pi(s) - V^*(s)] \leq \frac{\gamma C}{1-\gamma} \mathbb{E}[\text{dist}_{\ell_1}(\pi, \Pi^*)] \\ &\leq \frac{2\gamma C |\mathcal{A}| C_\gamma^n}{1-\gamma} \exp\left(-\frac{\Delta^*(\mathcal{M})\sqrt{\log|\mathcal{A}|}}{2}\gamma^{-(1/2-\beta)k}\right), \end{aligned} \quad (73)$$

whenever $k \geq K_3$. Thus for any $\bar{K}_3 \geq K_3$, from the definition of $\eta_t = \gamma^{(\beta-1/2)t} \sqrt{\log|\mathcal{A}|}$, the superlinear convergence of Q-function in (53), and the condition that $Q^*(s, i) = Q^*(s, j)$,

$$\begin{aligned} (B) &\leq \sum_{t=\bar{K}_3}^k \gamma^{k+1-t} \eta_t |g_i^t - g_j^t| \leq \sum_{t=\bar{K}_3}^k \gamma^{k+1-t} \eta_t (|g_i^t - Q^*(s, i)| + |g_j^t - Q^*(s, j)|) \\ &\leq \sum_{t=\bar{K}_3}^k \gamma^{k-(\frac{3}{2}-\beta)t} \frac{2\gamma C |\mathcal{A}| C_\gamma^n}{1-\gamma} \exp\left(-\frac{\Delta^*(\mathcal{M})\sqrt{\log|\mathcal{A}|}}{2}\gamma^{-(1/2-\beta)t}\right) \\ &\leq \sum_{t=\bar{K}_3}^k \gamma^{k-t/2} \frac{2\gamma C |\mathcal{A}| C_\gamma^n}{1-\gamma} \exp\left(-\frac{\Delta^*(\mathcal{M})\sqrt{\log|\mathcal{A}|}}{2}\gamma^{-(1/2-\beta)t} + t \log\left(\frac{1}{\gamma}\right)\right) \end{aligned} \quad (74)$$

It suffices to choose \bar{K}_3 satisfying

$$\bar{K}_3 = \min_k \left\{ k > K_3 : \Delta^*(\mathcal{M})\sqrt{\log|\mathcal{A}|}\gamma^{-(1/2-\beta)k} \geq 2k \log\left(\frac{1}{\gamma}\right) \right\},$$

then for any $k \geq \bar{K}_3$, from (56), we obtain

$$(B) = \sum_{t=\bar{K}_3}^k \gamma^{k+1-t} \eta_t |g_i^t - g_j^t| \leq \sum_{t=\bar{K}_3}^k \gamma^{k-t/2} \frac{2\gamma C |\mathcal{A}| C_\gamma^n}{1-\gamma} \leq \gamma^{k/2} \frac{2\gamma C |\mathcal{A}| C_\gamma^n}{(1-\gamma)(1-\gamma^{1/2})}. \quad (75)$$

We then proceed to bound term (C). From the definition of $\eta_t \sigma_t = \gamma^{\beta t} \sqrt{\log |\mathcal{A}|}$ and the fact that $\mathbb{E} |\delta_i^t| \leq \mathbb{E} \|\delta_t\|_\infty \leq \left(\mathbb{E} \|\delta_t\|_\infty^2 \right)^{1/2} \leq \sigma_t$, we have

$$(C) \leq 2 \sum_{t=0}^k \gamma^{k+1-t} \eta_t \sigma_t \leq \frac{2\gamma^{\beta k} \sqrt{\log |\mathcal{A}|}}{1-\gamma^{1/2}}. \quad (76)$$

Thus by combining (72), (75), and (76), we obtain that

$$\mathbb{E} |z_j^{k+1} - z_i^{k+1}| \leq \frac{2r(6 \log |\mathcal{A}| + C) \sqrt{\gamma \log |\mathcal{A}|}}{(1-\gamma)^2} \gamma^{k-(3/2-\beta)\bar{K}_3} + \gamma^{k/2} \frac{2\gamma C |\mathcal{A}| C_\gamma^n}{(1-\gamma)(1-\gamma^{1/2})} + \frac{2\gamma^{\beta k} \sqrt{\log |\mathcal{A}|}}{1-\gamma^{1/2}}. \quad (77)$$

By choosing $v_k = \gamma^{\frac{\beta k}{2}}$, from Markov's inequality we obtain that for any $k \geq \bar{K}_3$,

$$\begin{aligned} & \mathbb{P} (|z_j^{k+1} - z_i^{k+1}| \geq v_k) \\ & \leq \frac{2r(6 \log |\mathcal{A}| + C) \sqrt{\gamma \log |\mathcal{A}|}}{(1-\gamma)^2} \gamma^{(1-\beta/2)k-(3/2-\beta)\bar{K}_3} + \gamma^{(1-\beta)/2} \frac{2\gamma C |\mathcal{A}| C_\gamma^n}{(1-\gamma)(1-\gamma^{1/2})} + \frac{2\gamma^{\beta k/2} \sqrt{\log |\mathcal{A}|}}{1-\gamma^{1/2}}, \end{aligned}$$

and hence $\sum_{k=1}^{\infty} \mathbb{P} (|z_j^{k+1} - z_i^{k+1}| \geq v_k) < \infty$. From Borel-Cantelli lemma, we conclude that with probability 1, $\lim_{k \rightarrow \infty} |z_j^k - z_i^k| = 0$. On the other hand, apply Borel-Cantelli lemma to (70), we also know that with probability 1, $\lim_{k \rightarrow \infty} \pi_k(j|s) = 0$ for any $j \notin \mathcal{A}^*(s)$. Combined previous two observations, we immediately conclude that $\lim_{k \rightarrow \infty} \pi_k(j|s) = 1/|\mathcal{A}^*(s)|$ for any $s \in \mathcal{S}$ and any $i \in \mathcal{A}^*(s)$.

Finally, we remark that a high-probability non-asymptotic convergence can be also obtained by using (70), (77) and the Markov inequality. \square

Theorem (Restating Theorem 4.6). *Let $1 + \eta_t \tau_t = 1/\gamma$ and $\eta_t = \gamma^{(\beta-1/2)t} \sqrt{\log |\mathcal{A}|}$ in the SHPMD method. Suppose at each iteration, the number of independent trajectories M_k and the trajectory length T_k satisfies*

$$T_k \geq \frac{3(k+1)}{4} + \log_\gamma \left(\frac{1-\gamma}{2C} \right), M_k \geq \frac{4C^2}{(1-\gamma)^2} \gamma^{-(k+1)}.$$

Then for any $\epsilon > 0$, the SHPMD method finds an policy π_k with $\mathbb{E} [f(\pi_k) - f(\pi^*)] \leq \epsilon$ in $k = \frac{2}{1-2\beta} \log_\gamma \left(\frac{\epsilon(1-\gamma)(1-\gamma^{1/2})}{6\sqrt{\log |\mathcal{A}| + C}} \right)$, number of iterations. In addition, the total number of samples required by SHPMD can be bounded by

$$\mathcal{O} \left(\frac{|\mathcal{S}| |\mathcal{A}| \log_\gamma (\epsilon(1-\gamma))}{(1-\gamma)^{3+2/(1-2\beta)} (1-\gamma^{1/2})^{2/(1-2\beta)}} \frac{1}{\epsilon^{2/(1-2\beta)}} \right).$$

Proof of Theorem 4.6. By choosing M_k and T_k as in (19), one can readily verify that condition (17) is satisfied. To find an ϵ -optimal policy, from (18), one needs

$$k = \frac{2}{1-2\beta} \log_\gamma \left(\frac{\epsilon(1-\gamma)(1-\gamma^{1/2})}{6\sqrt{\log |\mathcal{A}| + C}} \right),$$

number of iterations. Thus the total number of samples can be bounded by

$$\begin{aligned} |\mathcal{S}| |\mathcal{A}| \sum_{t=0}^k M_t T_t &= |\mathcal{S}| |\mathcal{A}| \sum_{t=0}^k \left[\frac{3(t+1)}{4} + \log_\gamma \left(\frac{1-\gamma}{2C} \right) \right] \frac{4C^2}{(1-\gamma)^2} \gamma^{-(t+1)} \\ &= \frac{4C^2 |\mathcal{S}| |\mathcal{A}|}{(1-\gamma)^2} \mathcal{O} \left(\frac{k \gamma^{-k}}{1-\gamma} \right) \end{aligned}$$

$$\begin{aligned}
&= \frac{4C^2|\mathcal{S}||\mathcal{A}|}{(1-\gamma)^3} \mathcal{O} \left(\frac{2}{1-2\beta} \log_\gamma \left(\frac{\epsilon(1-\gamma)}{6\sqrt{\log|\mathcal{A}|} + C} \right) \cdot \left[\frac{\sqrt{6\log|\mathcal{A}|} + C}{(1-\gamma)(1-\gamma^{1/2})\epsilon} \right]^{\frac{2}{1-2\beta}} \right) \\
&= \mathcal{O} \left(\frac{|\mathcal{S}||\mathcal{A}| \log_\gamma(\epsilon(1-\gamma))}{(1-\gamma)^{3+2/(1-2\beta)} (1-\gamma^{1/2})^{2/(1-2\beta)}} \frac{1}{\epsilon^{2/(1-2\beta)}} \right).
\end{aligned}$$

□