

# A systematic construction of integrable delay-difference and delay-differential analogues of soliton equations

Kenta Nakata<sup>1</sup> and Ken-ichi Maruno<sup>2</sup>

<sup>1</sup> Department of Pure and Applied Mathematics, School of Fundamental Science and Engineering, Waseda University, 3-4-1 Okubo, Shinjuku-ku, Tokyo 169-8555, Japan

<sup>2</sup> Department of Applied Mathematics, Faculty of Science and Engineering, Waseda University, 3-4-1 Okubo, Shinjuku-ku, Tokyo 169-8555, Japan

E-mail: kennakaxx@akane.waseda.jp and kmaruno@waseda.jp

25 January 2022

**Abstract.** We propose a systematic method for constructing integrable delay-difference and delay-differential analogues of known soliton equations such as the Lotka-Volterra, Toda lattice, and sine-Gordon equations and their multi-soliton solutions. It is carried out by applying a reduction and delay-differential limit to the discrete KP or discrete two-dimensional Toda lattice equations. Each of the delay-difference and delay-differential equations has the  $N$ -soliton solution, which depends on the delay parameter and converges to an  $N$ -soliton solution of a known soliton equation as the delay parameter approaches 0.

*Keywords:* delay-differential equations, delay-difference equations, soliton equations, integrable systems, multi-soliton solutions

Submitted to: *J. Phys. A: Math. Theor.*

## 1. Introduction

Delay differential equations have been used as mathematical models in various fields of science and engineering such as traffic flow, population dynamics, nonlinear optics, fluid mechanics and infectious disease [1, 2]. Because of their importance in various fields, the mathematical properties of delay differential equations have been actively studied. For example, exact solutions of delay differential equations play an important role in the study of traffic flow [3–6].

In recent years, there has been active research on integrability of delay differential equations. Quispel *et al* obtained a delay-differential equation which has a continuum limit to the first Painlevé equation from the viewpoint of a symmetry reduction to an integrable differential-difference equation which is the Lotka-Volterra (LV) equation (this equation is also called the Kac-van Moerbeke lattice or Volterra lattice) [7]. Levi and Winternitz also obtained delay-differential equations as reductions based on the symmetry group of the two-dimensional Toda (2DTL) lattice [8]. Grammaticos *et al* [9, 10] introduced delay-differential equations by applying the method blending the Painlevé analysis with the singularity confinement to

study delay-differential analogues of the Painlevé equations. After their monumental works, several important studies have revealed mathematical properties of integrable (ordinary) delay-differential equations [11–17].

On the other hand, there are few examples of integrable non-ordinary delay-differential equations. Villarroel and Ablowitz found the Lax pair for a delay-differential analogue of the 2DTL equation and established the inverse scattering formalism [18]. Recently, we constructed the  $N$ -soliton solution of the delay-differential analogues of the 2DTL equation [19]. Although there have been such works, research on integrable non-ordinary delay-differential equations is not fully developed.

In this paper, we propose a systematic method for constructing integrable delay-difference and delay-differential analogues of soliton equations and their multi-soliton solutions. Our construction starts from the discrete KP equation (or discrete 2DTL equation) and use reduction and delay-differential limit. As examples of our method, we present delay-difference and delay-differential analogues of the LV, Toda lattice (TL) and sine-Gordon (sG) equations and their exact  $N$ -soliton solutions.

This paper is organized as follows. In Section 2, we show a systematic method to construct integrable delay-difference and delay-differential analogues of soliton equations, and derive delay-difference and delay-differential analogues of the LV equation and their  $N$ -soliton solutions as an example. In Section 3 and 4, we construct delay-difference and delay-differential analogues of the TL and sG equations and their  $N$ -soliton solutions. Section 5 is devoted to conclusions.

## **2. A method to construct delay-difference and delay-differential analogues of soliton equations and construction of an integrable delay Lotka-Volterra equation**

In this section, we show how to construct integrable delay-difference and delay-differential analogues of soliton equations and their  $N$ -soliton solutions. The first step of our method is to obtain a discrete equation by applying a reduction including a free parameter  $\alpha$  to the discrete KP equation [20, 21]

$$a(b - c)f_{n+1,m,k}f_{n,m+1,k+1} + b(c - a)f_{n,m+1,k}f_{n+1,m,k+1} + c(a - b)f_{n,m,k+1}f_{n+1,m+1,k} = 0 \quad (1)$$

or discrete 2DTL equation [20, 22]

$$abf_{k+1,n+1,m}f_{k-1,n,m+1} + f_{k,n+1,m}f_{k,n,m+1} - (1 + ab)f_{k,n+1,m+1}f_{k,n,m} = 0, \quad (2)$$

where  $a, b, c$  are real constants. The equations obtained depend on the free parameter  $\alpha$ , the discrete variable  $n, m$  and the time-lattice parameter  $\delta$  which is defined by the parameters  $a, b, c$ . Note that the dependency on  $k$  is removed by reduction. Here  $\alpha$  is made to appear as shifts of  $m$  (such as  $f_{m+\alpha}$ ). Thus the equations obtained in this step are delay-difference analogues of soliton equations.

Then, we apply the delay-differential limit

$$\delta \rightarrow 0, \quad m\delta = t, \quad \alpha\delta = \tau = \text{const.} \quad (3)$$

to the equation we obtained by the reduction, where  $t$  is the continuous time variable. This limit yields a delay-differential equation which includes the delay parameter  $\tau$  as shifts of  $t$ . For example, we can actually check that the limit of  $f_{m+\alpha}$  is  $f(t + \tau)$ .

An important point of this process is that we can easily obtain explicit  $N$ -soliton solutions of delay-difference and delay-differential analogue of soliton equations. It is carried out by reduction and delay-differential limit to the  $N$ -soliton solution of the discrete KP or discrete 2DTL equations. In addition, we can obtain a known soliton equation and its  $N$ -soliton solution as  $\tau \rightarrow 0$  if we properly determine the reduction at the first step of our method. We remark that which soliton equation we obtain as  $\tau \rightarrow 0$  depends on the reduction. Therefore, considering different reductions at the first step of our method, we can construct different delay-difference and delay-differential analogues of soliton equations.

In this section, we show the construction of the delay-difference and delay-differential analogue of the LV equation and their  $N$ -soliton solutions as an example and explain the detail of our method.

First, we consider the discrete KP equation (1) and its  $N$ -soliton solution in the Gram determinant form [23]:

$$\begin{aligned} f_{n,m,k} &= \det \left( \delta_{ij} + \frac{\phi_i \psi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N} = \frac{\prod_{i=1}^N \phi_i}{\prod_{j=1}^N \phi_j} \det \left( \delta_{ij} \frac{\phi_j}{\phi_i} + \frac{\phi_j \psi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N} \\ &= \det \left( \delta_{ij} + \frac{\phi_j \psi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N}, \end{aligned} \quad (4)$$

$$\phi_i(n, m, k) = \beta_i (1 - ap_i)^{-n} (1 - bp_i)^{-m} (1 - cp_i)^{-k},$$

$$\psi_i(n, m, k) = \gamma_i (1 - aq_i)^n (1 - bq_i)^m (1 - cq_i)^k.$$

This Gram determinant form of the  $N$ -soliton solution can be rewritten as

$$\begin{aligned} f_{n,m,k} &= \sum_{I \subset \{1, \dots, N\}} \prod_{i \in I} \Phi_i \prod_{i < j, i, j \in I} \frac{(p_i - p_j)(q_i - q_j)}{(p_i - q_j)(q_i - p_j)}, \\ \Phi_i(n, m, k) &= \beta_i \gamma_i \left( \frac{1 - aq_i}{1 - ap_i} \right)^n \left( \frac{1 - bq_i}{1 - bp_i} \right)^m \left( \frac{1 - cq_i}{1 - cp_i} \right)^k. \end{aligned} \quad (5)$$

Here  $\sum_{I \subset \{1, \dots, N\}}$  is taken to be summed over all possible combinations in the set  $\{1, \dots, N\}$ .

Now, we apply the following reduction condition to the discrete KP equation (1):

$$f_{n,m,k+1} = f_{n+2,m+\alpha,k}, \quad (6)$$

where the free parameter  $\alpha$  is a fixed value considered as the delay. Setting  $c = -a$  and  $\delta = 2b/(a - b)$ , we obtain the following discrete bilinear equation by the reduction to the discrete KP equation (1):

$$(1 + \delta) f_n^{m+1+\alpha} f_{n-1}^m - \delta f_{n+1}^{m+\alpha} f_{n-2}^{m+1} - f_n^{m+\alpha} f_{n-1}^{m+1} = 0, \quad (7)$$

where  $f_n^m \equiv f_{n,m,k}$  and  $k$  can be omitted (thus we can set  $k = 0$ ). We can rewrite the bilinear equation (7) as follows by using Hirota's D-operators:

$$\begin{aligned} &\left( 2 \sinh \left( \frac{D_m}{2} \right) \sinh \left( \frac{D_n}{2} + \frac{\alpha D_m}{2} \right) - 2\delta \sinh \left( \frac{D_n - D_m}{2} \right) \sinh \left( D_n + \frac{\alpha D_m}{2} \right) \right) f_n^m \cdot f_n^m \\ &= 0. \end{aligned} \quad (8)$$

Here Hirota's D-operators are defined by

$$D_t^l g(t) \cdot h(t) = \left( \frac{\partial}{\partial t} - \frac{\partial}{\partial s} \right)^l g(t)h(s) \Big|_{s=t}, \quad e^{D^m} g_m \cdot h_m = g_{m+1} h_{m-1}. \quad (9)$$

Next we consider the reduction of the  $N$ -soliton solution. By applying the constraint

$$\left( \frac{1 - aq_i}{1 - ap_i} \right)^2 \left( \frac{1 - bq_i}{1 - bp_i} \right)^\alpha = \frac{1 - cq_i}{1 - cp_i} \quad (10)$$

to the  $N$ -soliton solution (4) and (5), we obtain

$$\begin{aligned} \Phi_i(n, m, k+1) &= \beta_i \gamma_i \left( \frac{1 - aq_i}{1 - ap_i} \right)^n \left( \frac{1 - bq_i}{1 - bp_i} \right)^m \left( \frac{1 - cq_i}{1 - cp_i} \right)^{k+1} \\ &= \beta_i \gamma_i \left( \frac{1 - aq_i}{1 - ap_i} \right)^{n+2} \left( \frac{1 - bq_i}{1 - bp_i} \right)^{m+\alpha} \left( \frac{1 - cq_i}{1 - cp_i} \right)^k \\ &= \Phi_i(n+2, m+\alpha, k), \end{aligned} \quad (11)$$

thus the reduction condition (6) is satisfied. Setting  $c = -a$ ,  $\delta = 2b/(a-b)$  and replacing  $p_i$  and  $q_i$  by  $(-2p_i - 1)/a$  and  $(-2q_i - 1)/a$  respectively, we obtain the  $N$ -soliton solution of the bilinear equation (7) by the reduction to (4) and (5):

$$f_n^m = \det \left( \delta_{ij} + \frac{\Phi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N} = \sum_{I \subset \{1, \dots, N\}} \prod_{i \in I} \Phi_i \prod_{i < j, i, j \in I} \frac{(p_i - p_j)(q_i - q_j)}{(p_i - q_j)(q_i - p_j)}, \quad (12)$$

$$\Phi_i(n, m) = \beta_i \gamma_i \left( \frac{1 + q_i}{1 + p_i} \right)^n \left( \frac{1 + \delta + \delta q_i}{1 + \delta + \delta p_i} \right)^m,$$

$$\frac{q_i}{p_i} = \left( \frac{1 + q_i}{1 + p_i} \right)^2 \left( \frac{1 + \delta + \delta q_i}{1 + \delta + \delta p_i} \right)^\alpha.$$

The delay parameter  $\alpha$  appears in the bilinear equation (7) as shifts of the discrete time variable  $m$ . We remark that the bilinear equation (7) and the  $N$ -soliton solution (12) in the case of  $\alpha = 0$  (thus  $p_i = 1/q_i$ ) are known as the bilinear equation of the discrete LV equation and its  $N$ -soliton solution [24, 25]. Thus the bilinear equation (7) can be seen as the bilinear equation of the delay-difference analogue of the LV equation.

By the dependent variable transformation

$$u_n^m = \frac{f_{n+1}^{m+\alpha} f_{n-2}^{m+1}}{f_n^{m+\alpha} f_{n-1}^{m+1}}, \quad (13)$$

the bilinear equation (7) is transformed into the nonlinear delay-difference equation

$$\frac{u_n^{m+1+\alpha} u_{n-1}^m}{u_n^{m+\alpha} u_{n-1}^{m+1}} = \frac{(1 + \delta u_{n+1}^{m+\alpha})(1 + \delta u_{n-2}^{m+1})}{(1 + \delta u_n^{m+\alpha})(1 + \delta u_{n-1}^{m+1})}, \quad (14)$$

which is the delay-difference analogue of the LV equation. If  $\alpha = 0$ , equation (14) is just a division of the discrete LV equation [24]

$$\frac{u_n^{m+1}}{u_n^m} = \frac{1 + \delta u_{n+1}^m}{1 + \delta u_{n-1}^{m+1}}, \quad \frac{u_{n-1}^{m+1}}{u_{n-1}^m} = \frac{1 + \delta u_n^m}{1 + \delta u_{n-2}^{m+1}}. \quad (15)$$

Now, we apply the delay-differential limit

$$\delta \rightarrow 0, \quad m\delta = t, \quad \alpha\delta = 2\tau, \quad (16)$$

where  $\tau$  is a constant value called the delay parameter. The delay-differential limits of (7) and (12) are calculated respectively as follows:

$$D_t f_n(t + \tau) \cdot f_{n-1}(t - \tau) - f_{n+1}(t + \tau) f_{n-2}(t - \tau) + f_n(t + \tau) f_{n-1}(t - \tau) = 0, \quad (17)$$

$$f_n(t) = \det \left( \delta_{ij} + \frac{\Phi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N} = \sum_{I \subset \{1, \dots, N\}} \prod_{i \in I} \Phi_i \prod_{i < j, i, j \in I} \frac{(p_i - p_j)(q_i - q_j)}{(p_i - q_j)(q_i - p_j)}, \quad (18)$$

$$\Phi_i(n, t) = \beta_i \gamma_i \left( \frac{1 + q_i}{1 + p_i} \right)^n e^{(q_i - p_i)t},$$

$$\frac{q_i}{p_i} = \left( \frac{1 + q_i}{1 + p_i} \right)^2 e^{2\tau(q_i - p_i)}.$$

Here the bilinear equation (17) can be rewritten as

$$\left( D_t \sinh \left( \frac{D_n}{2} + \tau D_t \right) - 2 \sinh \left( \frac{D_n}{2} \right) \sinh (D_n + \tau D_t) \right) f_n(t) \cdot f_n(t) = 0. \quad (19)$$

The bilinear equation (17) is a delay differential-difference equation which includes the delay  $\tau$  as shifts of the continuous time variable  $t$ . Putting  $\tau = 0$  (thus  $p_i = 1/q_i$ ), we can easily check that the bilinear equation (17) becomes the bilinear equation of the LV equation and (18) becomes the  $N$ -soliton solution of the LV equation [26]. Thus we claim that the bilinear equation (17) is the bilinear equation of the integrable delay LV equation and the bilinear equation (7) is the fully discrete analogue of (17).

**Remark 2.1.** The last relation of (18), which is the dispersion relation, is rewritten as

$$R(p_i) = R(q_i), \quad R(p) = \frac{pe^{-2\tau p}}{(1+p)^2}. \quad (20)$$

According to the graph of  $R(p)$ , it is easily seen that there exist  $p_i$  and  $q_i$  satisfying  $R(p_i) = R(q_i)$ .

**Remark 2.2.** We remark that soliton solutions of (17) are also obtained by Hirota's direct method [27], which does not require any reduction. We show a few examples of deriving soliton solutions without using the soliton solutions of the discrete KP equation. Let  $f_n(t)$  be 1-soliton or 2-soliton with the perturbation parameter  $\epsilon$ :

$$f_n(t) = 1 + \epsilon P_1^n e^{Q_1 t + \xi_{01}}, \quad (21)$$

$$f_n(t) = 1 + \epsilon P_1^n e^{Q_1 t + \xi_{01}} + \epsilon P_2^n e^{Q_2 t + \xi_{02}} + \epsilon^2 a_{12} (P_1 P_2)^n e^{(Q_1 + Q_2)t + \xi_{01} + \xi_{02}}. \quad (22)$$

Substituting them into the bilinear equations (19) and assuming that each of the orders of  $\epsilon$  vanishes, we have the following conditions.

$$F(P_i, Q_i) = 0 \quad (i = 1, 2), \quad a_{12} = -\frac{F(P_1/P_2, Q_1 - Q_2)}{F(P_1 P_2, Q_1 + Q_2)}, \quad (23)$$

where the function  $F$  is defined by

$$F(P, Q) = Q \left( \sqrt{P} e^{\tau Q} - \frac{1}{\sqrt{P}} e^{-\tau Q} \right) - \left( \sqrt{P} - \frac{1}{\sqrt{P}} \right) \left( P e^{\tau Q} - \frac{1}{P} e^{-\tau Q} \right). \quad (24)$$

By computations, we can check that the solutions (21) and (22) are equivalent to the case of  $N = 1, 2$  in the  $N$ -soliton solution (18). As we can see from this example, we can use Hirota's direct method even if the bilinear equation includes some delays.

Via the dependent variable transformation

$$u_n(t) = \frac{f_{n+1}(t+\tau)f_{n-2}(t-\tau)}{f_n(t+\tau)f_{n-1}(t-\tau)}, \quad (25)$$

the bilinear equation (17) is transformed into the nonlinear delay-differential equation

$$\frac{d}{dt} \log \frac{u_n(t+\tau)}{u_{n-1}(t-\tau)} = u_{n+1}(t+\tau) - u_n(t+\tau) - u_{n-1}(t-\tau) + u_{n-2}(t-\tau), \quad (26)$$

which is the nonlinear form of (17). If  $\tau = 0$ , equation (26) is just a subtraction of the following nonlinear forms of the LV equation [26].

$$\frac{d}{dt} \log u_n(t) = u_{n+1}(t) - u_{n-1}(t), \quad \frac{d}{dt} \log u_{n-1}(t) = u_n(t) - u_{n-2}(t). \quad (27)$$

Next, we derive the bilinear equation (17) and the  $N$ -soliton solution (18) directly from the semi-discrete KP equation [28]

$$acD_t f_{n+1}^k(t) \cdot f_n^{k+1}(t) - (a-c)(f_{n+1}^{k+1}(t)f_n^k(t) - f_{n+1}^k(t)f_n^{k+1}(t)) = 0. \quad (28)$$

We remark that the bilinear equation (28) and its solutions are derived by the continuum limit  $b \rightarrow 0$ ,  $mb = t$  of the discrete KP equation (1) and its solutions. The  $N$ -soliton solution in the Gram determinant form is given as

$$f_n^k(t) = \det \left( \delta_{ij} + \frac{\phi_i \psi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N} = \det \left( \delta_{ij} + \frac{\phi_j \psi_i}{p_i - q_j} \right)_{1 \leq i, j \leq N} \quad (29)$$

$$= \sum_{I \subset \{1, \dots, N\}} \prod_{i \in I} \Phi_i \prod_{i < j, i, j \in I} \frac{(p_i - p_j)(q_i - q_j)}{(p_i - q_j)(q_i - p_j)},$$

$$\phi_i(n, k, t) = \beta_i (1 - ap_i)^{-n} (1 - cp_i)^{-k} e^{p_i t}, \quad \psi_i(n, k, t) = \gamma_i (1 - aq_i)^n (1 - cq_i)^k e^{-q_i t},$$

$$\Phi_i(n, k, t) = \beta_i \gamma_i \left( \frac{1 - aq_i}{1 - ap_i} \right)^n \left( \frac{1 - cq_i}{1 - cp_i} \right)^k e^{(p_i - q_i)t}.$$

Now, we can easily obtain the bilinear equation (17) by applying the reduction condition

$$f_n^{k+1}(t) = f_{n+2}^k(t + 2\tau), \quad (30)$$

and setting  $a = 2$ ,  $c = -2$  to the semi-discrete KP equation (28). To realize this reduction condition for the  $N$ -soliton solution, we can apply the constraint

$$\frac{1 - cq_i}{1 - cp_i} = \left( \frac{1 - aq_i}{1 - ap_i} \right)^2 e^{2\tau(p_i - q_i)} \quad (31)$$

to (29). This constraint leads to

$$\begin{aligned} \Phi_i(n, k+1, t) &= \beta_i \gamma_i \left( \frac{1 - aq_i}{1 - ap_i} \right)^n \left( \frac{1 - cq_i}{1 - cp_i} \right)^{k+1} e^{(p_i - q_i)t} \\ &= \beta_i \gamma_i \left( \frac{1 - aq_i}{1 - ap_i} \right)^{n+2} \left( \frac{1 - cq_i}{1 - cp_i} \right)^k e^{(p_i - q_i)(t+2\tau)} \\ &= \Phi_i(n+2, k, t+2\tau), \end{aligned} \quad (32)$$

thus the reduction condition (30) is satisfied. Replacing  $p_i$  and  $q_i$  by  $(-2p_i - 1)/a$  and  $(-2q_i - 1)/a$  respectively and setting  $a = 2$ ,  $c = -2$ , we obtain the  $N$ -soliton solution (18).

This construction of the delay-differential analogue of the LV equation does not require the delay-differential limit.

The Wronskian solution of the semi-discrete KP equation (28) is given as follows [28]:

$$f_n^k(t) = \begin{vmatrix} \phi_1(n, k, t) & \phi_1^{(1)}(n, k, t) & \cdots & \phi_1^{(N-1)}(n, k, t) \\ \phi_2(n, k, t) & \phi_2^{(1)}(n, k, t) & \cdots & \phi_2^{(N-1)}(n, k, t) \\ \vdots & \vdots & \cdots & \vdots \\ \phi_N(n, k, t) & \phi_N^{(1)}(n, k, t) & \cdots & \phi_N^{(N-1)}(n, k, t) \end{vmatrix}, \quad (33)$$

$$\frac{\phi_i(n, k, t) - \phi_i(n-1, k, t)}{a} = \frac{\partial \phi_i(n, k, t)}{\partial t}, \quad \frac{\phi_i(n, k, t) - \phi_i(n, k-1, t)}{c} = \frac{\partial \phi_i(n, k, t)}{\partial t},$$

$$\phi_i^{(l)}(n, k, t) \equiv \frac{\partial^l \phi_i(n, k, t)}{\partial t^l}.$$

The  $N$ -soliton solution of the semi-discrete KP equation is given by choosing elements in the Wronskian solution as

$$\phi_i(n, k, t) = (1 - ap_i)^{-n} (1 - cp_i)^{-k} e^{p_i t + \zeta_{0i}} + (1 - aq_i)^{-n} (1 - cq_i)^{-k} e^{q_i t + \eta_{0i}}. \quad (34)$$

Now, we can easily obtain the bilinear equation (17) by applying the reduction condition

$$f_n^{k+1}(t) \approx f_{n+2}^k(t + 2\tau), \quad (35)$$

and setting  $a = 2$ ,  $c = -2$  to the semi-discrete KP equation (28). To realize this reduction condition for the  $N$ -soliton solutions, we can apply the constraint

$$(1 - ap_i)^{-2} (1 - cp_i) e^{2\tau p_i} = (1 - aq_i)^{-2} (1 - cq_i) e^{2\tau q_i} \quad (36)$$

to (34). Replacing  $p_i$  and  $q_i$  by  $(-2p_i - 1)/a$  and  $(-2q_i - 1)/a$  respectively and setting  $a = 2$ ,  $c = -2$ , we obtain the  $N$ -soliton solution in the Wronskian form

$$f_n^k(t) = \begin{vmatrix} \phi_1(n, k, t) & \phi_1^{(1)}(n, k, t) & \cdots & \phi_1^{(N-1)}(n, k, t) \\ \phi_2(n, k, t) & \phi_2^{(1)}(n, k, t) & \cdots & \phi_2^{(N-1)}(n, k, t) \\ \vdots & \vdots & \cdots & \vdots \\ \phi_N(n, k, t) & \phi_N^{(1)}(n, k, t) & \cdots & \phi_N^{(N-1)}(n, k, t) \end{vmatrix}, \quad (37)$$

$$\phi_i(n, k, t) = (1 + p_i)^{-n} e^{-p_i t + \tilde{\zeta}_{0i}} + (1 + q_i)^{-n} e^{-q_i t + \tilde{\eta}_{0i}}, \quad (i = 1, 2, \dots, N)$$

$$\frac{q_i}{p_i} = \left( \frac{1 + q_i}{1 + p_i} \right)^2 e^{2\tau(q_i - p_i)}.$$

**Remark 2.3.** We can also obtain the delay-differential analogue of the LV equation from the Bäcklund transformation (BT) of the 2DTL equation [27, 29]

$$\mu D_t f_n^{k-1}(t) \cdot f_{n+1}^k(t) - f_n^{k-1}(t) f_{n+1}^k(t) + f_{n+1}^{k-1}(t) f_n^k(t) = 0. \quad (38)$$

The following derivation is much easier than the above derivation from the semi-discrete KP equation. The  $N$ -soliton solution in the Gram determinant form is given as

$$f_n^k(t) = \det \left( \delta_{ij} + \frac{\phi_i \psi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N} = \det \left( \delta_{ij} + \frac{\phi_j \psi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N}, \quad (39)$$

$$\phi_i(n, k, t) = \beta_i p_i^k (1 - \mu p_i)^{-n} e^{-p_i t}, \quad \psi_i(n, k, t) = \gamma_i q_i^{-k} (1 - \mu q_i)^n e^{q_i t},$$

and the  $N$ -soliton solution in the Wronskian (Casorati determinant) form is given as

$$f_n^k(t) = \begin{vmatrix} \phi_1(n, k, t) & \phi_1^{(1)}(n, k, t) & \cdots & \phi_1^{(N-1)}(n, k, t) \\ \phi_2(n, k, t) & \phi_2^{(1)}(n, k, t) & \cdots & \phi_2^{(N-1)}(n, k, t) \\ \vdots & \vdots & \cdots & \vdots \\ \phi_N(n, k, t) & \phi_N^{(1)}(n, k, t) & \cdots & \phi_N^{(N-1)}(n, k, t) \end{vmatrix}, \quad (40)$$

$$\phi_i(n, k, t) = p_i^k (1 - \mu p_i)^{-n} e^{-p_i t + \zeta_0 i} + q_i^k (1 - \mu q_i)^{-n} e^{-q_i t + \eta_0 i}. \quad (i = 1, 2, \dots, N)$$

Applying the reduction condition

$$f_n^{k-1}(t) \simeq f_{n+2}^k(t + 2\tau), \quad (41)$$

and setting  $\mu = -1$ , we obtain the bilinear equation (17) and its  $N$ -soliton solution (18).

### 3. An integrable delay Toda lattice equation

In this section, we construct a delay differential-difference equation that should be called an integrable delay TL equation by using our method.

We first apply the reduction condition

$$f_{n,m,k+1} = f_{n+1,m+1+\alpha,k} \quad (42)$$

to the discrete KP equation (1) and its  $N$ -soliton solution (4). The independent variables  $n$  and  $m$  are considered to be the discrete space variable and discrete time variable respectively, and the parameter  $\alpha$  is a delay parameter. Replacing  $p_i$  and  $q_i$  by  $(1 - p_i)/c$  and  $(1 - q_i)/c$  respectively and setting  $\alpha\delta = (a - c)/a$ ,  $\delta = b/(b - c)$ , we have

$$f_n^{m+1+\alpha} f_n^{m-1} - \alpha\delta^2 f_{n+1}^{m+\alpha} f_{n-1}^m - (1 - \alpha\delta^2) f_n^{m+\alpha} f_n^m = 0, \quad (43)$$

$$f_n^m = \det \left( \delta_{ij} + \frac{\Phi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N} = \sum_{I \subset \{1, \dots, N\}} \prod_{i \in I} \Phi_i \prod_{i < j, i, j \in I} \frac{(p_i - p_j)(q_i - q_j)}{(p_i - q_j)(q_i - p_j)}, \quad (44)$$

$$\Phi_i = \beta_i \gamma_i \left( \frac{q_i - \alpha\delta}{p_i - \alpha\delta} \right)^n \left( \frac{1 - \delta q_i}{1 - \delta p_i} \right)^m,$$

$$\frac{q_i}{p_i} = \frac{q_i - \alpha\delta}{p_i - \alpha\delta} \left( \frac{1 - \delta q_i}{1 - \delta p_i} \right)^{1+\alpha}.$$

The bilinear equation (43) is rewritten as

$$\left( 2 \sinh \left( \frac{D_m}{2} \right) \sinh \left( \frac{D_m}{2} + \frac{\alpha D_m}{2} \right) - 2\alpha\delta^2 \sinh \left( \frac{D_n}{2} \right) \sinh \left( \frac{D_n}{2} + \frac{\alpha D_m}{2} \right) \right) f_n^{m+\alpha} \cdot f_n^m = 0. \quad (45)$$

We can consider that equation (43) is the bilinear equation of the delay discrete TL equation.

By the dependent variable transformation

$$1 + V_n^m = \frac{f_{n+1}^{m+\alpha} f_{n-1}^m}{f_n^{m+\alpha} f_n^m}, \quad (46)$$



we can transform (43) into the nonlinear delay-difference equation

$$\frac{(1 + V_n^{m+1+\alpha})(1 + V_n^{m-1})}{(1 + V_n^{m+\alpha})(1 + V_n^m)} = \frac{(1 + \alpha\delta^2 V_{n+1}^{m+\alpha})(1 + \alpha\delta^2 V_{n-1}^m)}{(1 + \alpha\delta^2 V_n^{m+\alpha})(1 + \alpha\delta^2 V_n^m)}, \quad (47)$$

which is the delay-difference analogue of the TL equation.

Now, we apply the delay-differential limit

$$\delta \rightarrow 0, \quad m\delta = t, \quad \alpha\delta = 2\tau \quad (48)$$

to (43) and (44), where  $t$  is the continuous time variable and  $\tau$  is the delay parameter. Then we obtain

$$D_t f_n(t + \tau) \cdot f_n(t - \tau) - 2\tau(f_{n+1}(t + \tau)f_{n-1}(t - \tau) - f_n(t + \tau)f_n(t - \tau)) = 0 \quad (49)$$

and

$$f_n(t) = \det \left( \delta_{ij} + \frac{\Phi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N} = \sum_{I \subset \{1, \dots, N\}} \prod_{i \in I} \Phi_i \prod_{i < j, i, j \in I} \frac{(p_i - p_j)(q_i - q_j)}{(p_i - q_j)(q_i - p_j)}, \quad (50)$$

$$\Phi_i = \beta_i \gamma_i \left( \frac{q_i - 2\tau}{p_i - 2\tau} \right)^n e^{(p_i - q_i)t},$$

$$\frac{q_i}{p_i} = \frac{q_i - 2\tau}{p_i - 2\tau} e^{2\tau(p_i - q_i)}.$$

Here the bilinear equation (49) is equivalent to

$$\left( D_t \sinh(\tau D_t) - 4\tau \sinh\left(\frac{D_n}{2}\right) \sinh\left(\frac{D_n}{2} + \tau D_t\right) \right) f_n(t) \cdot f_n(t) = 0. \quad (51)$$

Calculating the limit of this equation as  $\tau \rightarrow 0$ , we obtain the bilinear TL equation [30]

$$\left( D_t^2 - 4 \sinh^2\left(\frac{D_n}{2}\right) \right) f_n(t) \cdot f_n(t) = 0. \quad (52)$$

The last relation of (50) is described by  $(p_i + 1/p_i - q_i - 1/q_i)\tau + O(\tau^2) = 0$ , thus we have  $p_i = 1/q_i$  as  $\tau \rightarrow 0$ . Therefore, we can check that the limit of (50) as  $\tau \rightarrow 0$  is the  $N$ -soliton solution of the TL equation [27, 30, 31]. Thus we claim that the bilinear equation (49) is the bilinear equation of the delay TL equation.

**Remark 3.1.** If we use the relation  $\delta = (a - c)/a$  instead of the above one  $\alpha\delta = (a - c)/a$ , we obtain the bilinear equation

$$f_n^{m+1+\alpha} f_n^{m-1} - \delta^2 f_{n+1}^{m+\alpha} f_{n-1}^m - (1 - \delta^2) f_n^{m+\alpha} f_n^m = 0$$

instead of the delay discrete TL equation (43). This equation can be considered more natural than (43), because we can obtain the discrete TL equation from it by putting  $\alpha = 0$  [30]. However it does not yield a good delay-differential equation, because the order  $O(\delta^2)$  vanishes in the delay-differential limit. On the other hand, the bilinear equation (43) yields the good delay-differential equation (49), which should be called a delay TL equation.

We present the nonlinear form of the bilinear equation of the delay TL equation (49) under the dependent variable transformation

$$1 + V_n(t) = \frac{f_{n+1}(t + \tau)f_{n-1}(t - \tau)}{f_n(t + \tau)f_n(t - \tau)}. \quad (53)$$

By using this transformation, we can transform (49) into the delay-differential equation

$$\begin{aligned} \left(\frac{1}{2\tau}\right) \frac{d}{dt} \log \frac{1 + V_n(t + \tau)}{1 + V_n(t - \tau)} \\ = V_{n+1}(t + \tau) + V_{n-1}(t - \tau) - V_n(t + \tau) - V_n(t - \tau), \end{aligned} \quad (54)$$

which is the nonlinear form of (49). The limit of (54) as  $\tau \rightarrow 0$  is the nonlinear form of the TL equation [27, 30, 31]:

$$\frac{d^2}{dt^2} \log(1 + V_n(t)) = V_{n+1}(t) - 2V_n(t) + V_{n-1}(t). \quad (55)$$

We can also obtain the above result from the BT of the 2DTL equation [29]

$$\mu D_t f_n^{k-1}(t) \cdot f_{n+1}^k(t) - f_n^{k-1}(t) f_{n+1}^k(t) + f_{n+1}^{k-1}(t) f_n^k(t) = 0. \quad (56)$$

Applying the reduction condition

$$f_n^{k-1}(t) = f_{n+1}^k(t + 2\tau), \quad (57)$$

and setting  $\mu = -1/(2\tau)$ , we obtain the bilinear equation (43) and its  $N$ -soliton solution

$$\begin{aligned} f_n &= \det \left( \delta_{ij} + \frac{\Phi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N}, \\ \Phi_i &= \beta_i \gamma_i \left( \frac{2\tau + q_i}{2\tau + p_i} \right)^n e^{(q_i - p_i)t}, \\ \frac{q_i}{p_i} &= \frac{2\tau + q_i}{2\tau + p_i} e^{2\tau(q_i - p_i)}, \end{aligned} \quad (58)$$

which leads to (50) by replacing  $p_i$  and  $q_i$  by  $-p_i$  and  $-q_i$ . This construction of the delay TL equation does not require the delay-differential limit.

#### 4. An integrable delay sine-Gordon equation

In this section, we find an integrable delay sG equation by the process similarly to the previous sections. It is a delay partial differential equation which can be obtained simply.

We consider the bilinear equation of the discrete 2DTL equation [20, 22]

$$ab f_{k+1, n+1, m} f_{k-1, n, m+1} + f_{k, n+1, m} f_{k, n, m+1} - (1 + ab) f_{k, n+1, m+1} f_{k, n, m} = 0, \quad (59)$$

where  $k$  is the discrete space variable, and  $n, m$  are the discrete time variables. The  $N$ -soliton solution of (59) is given as follows [22]:

$$\begin{aligned} f_{k, n, m} &= \det \left( \delta_{ij} + \frac{\phi_i \psi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N} \\ &= \det \left( \delta_{ij} + \frac{\phi_j \psi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N} = \sum_{I \subset \{1, \dots, N\}} \prod_{i \in I} \Phi_i \prod_{i < j, i, j \in I} \frac{(p_i - p_j)(q_i - q_j)}{(p_i - q_j)(q_i - p_j)}, \\ \phi_i &= \beta_i p_i^k (1 - ap_i)^{-n} (1 + b/p_i)^{-m}, \quad \psi_i = \gamma_i q_i^{-k} (1 - aq_i)^n (1 + b/q_i)^m, \\ \Phi_i &= \beta_i \gamma_i \left( \frac{p_i}{q_i} \right)^k \left( \frac{1 - aq_i}{1 - ap_i} \right)^n \left( \frac{1 + b/q_i}{1 + b/p_i} \right)^m. \end{aligned} \quad (60)$$

Applying the reduction condition

$$f_{k+1,n-\alpha,m-\beta} = f_{k-1,n+\alpha,m+\beta}, \quad (61)$$

and setting

$$f_n^m \equiv f_{k,n,m} = f_{k-2,n+2\alpha,m+2\beta}, \quad g_n^m \equiv f_{k+1,n-\alpha,m-\beta} = f_{k-1,n+\alpha,m+\beta}, \quad a = b = \delta, \quad (62)$$

to (59) and (60) respectively, we obtain

$$(1 + \delta^2) f_{n+1}^{m+1} f_n^m - f_{n+1}^m f_n^{m+1} - \delta^2 g_{n+1+\alpha}^{m+\beta} g_{n-\alpha}^{m+1-\beta} = 0, \quad (63)$$

$$(1 + \delta^2) g_{n+1}^{m+1} g_n^m - g_{n+1}^m g_n^{m+1} - \delta^2 f_{n+1+\alpha}^{m+\beta} f_{n-\alpha}^{m+1-\beta} = 0 \quad (64)$$

and

$$f_n^m = \det \left( \delta_{ij} + \frac{\Phi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N}, \quad (65)$$

$$\Phi_i = \beta_i \gamma_i \left( \frac{1 - \delta q_i}{1 - \delta p_i} \right)^n \left( \frac{1 + \delta/q_i}{1 + \delta/p_i} \right)^m,$$

$$\left( \frac{p_i}{q_i} \right)^2 = \left( \frac{1 - \delta q_i}{1 - \delta p_i} \right)^{2\alpha} \left( \frac{1 + \delta/q_i}{1 + \delta/p_i} \right)^{2\beta}.$$

To construct a delay-difference analogue of the sG equation, we take  $g_n^m$  to be the complex conjugate of  $f_n^m$ . Considering the regularity conditions  $f_n^m \neq 0$ , the  $N$ -soliton solution is given as

$$f_n^m = \det \left( \delta_{ij} + \frac{\Phi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N}, \quad g_n^m = \det \left( \delta_{ij} - \frac{\Phi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N}, \quad (66)$$

$$\Phi_i = \sqrt{-1} \mu_i \left( \frac{1 - \delta q_i}{1 - \delta p_i} \right)^n \left( \frac{1 + \delta/q_i}{1 + \delta/p_i} \right)^m,$$

$$\frac{p_i}{q_i} = - \left( \frac{1 - \delta q_i}{1 - \delta p_i} \right)^\alpha \left( \frac{1 + \delta/q_i}{1 + \delta/p_i} \right)^\beta,$$

where  $\mu_i$  is a real constant.

Setting  $f_n^m = F_n^m + \sqrt{-1} G_n^m$  and  $g_n^m = F_n^m - \sqrt{-1} G_n^m$ , we can rewrite the bilinear equations (63) and (64) with

$$\left( 2 \sinh \left( \frac{D_n}{2} \right) \sinh \left( \frac{D_m}{2} \right) + 2\delta^2 \sinh \left( \frac{D_n}{2} + \frac{\alpha D_n}{2} + \frac{\beta D_m}{2} \right) \sinh \left( \frac{D_m}{2} - \frac{\alpha D_n}{2} - \frac{\beta D_m}{2} \right) \right) (F_n^m \cdot F_n^m - G_n^m \cdot G_n^m) = 0, \quad (67)$$

$$\left( 2 \sinh \left( \frac{D_n}{2} \right) \sinh \left( \frac{D_m}{2} \right) + 2\delta^2 \cosh \left( \frac{D_n}{2} + \frac{\alpha D_n}{2} + \frac{\beta D_m}{2} \right) \cosh \left( \frac{D_m}{2} - \frac{\alpha D_n}{2} - \frac{\beta D_m}{2} \right) \right) F_n^m \cdot G_n^m = 0. \quad (68)$$

When  $\alpha = \beta = 0$ , we can find that the bilinear equations (63), (64) (and also (67), (68)) and the  $N$ -soliton solution (66) are actually equivalent to the bilinear equations of the discrete sG equation and their  $N$ -soliton solution [32].

To construct a nonlinear form of the delay-difference analogue of the sG equation, we consider the dependent variable transformation

$$f_n^m = \exp \left( \frac{\rho_n^m}{4} + \sqrt{-1} \frac{\theta_n^m}{4} \right), \quad g_n^m = \exp \left( \frac{\rho_n^m}{4} - \sqrt{-1} \frac{\theta_n^m}{4} \right), \quad (69)$$

which is equivalent to

$$\theta_n^m = 2\sqrt{-1} \log \frac{g_n^m}{f_n^m} = 4 \tan^{-1} \frac{G_n^m}{F_n^m}, \quad \rho_n^m = 2 \log f_n^m g_n^m = 2 \log((F_n^m)^2 + (G_n^m)^2). \quad (70)$$

By using this transformation, we can transform the bilinear equations (63) and (64) into the nonlinear delay-difference equation

$$\begin{aligned} & \sin \left( \frac{\theta_{n+1}^m + \theta_n^{m+1} - \theta_{n+1}^{m+1} - \theta_n^m}{4} \right) \\ &= \delta^2 \exp \left( \frac{\rho_{n+1+\alpha}^{m+\beta} + \rho_{n-\alpha}^{m+1-\beta} - \rho_{n+1}^m - \rho_n^{m+1}}{4} \right) \sin \left( \frac{\theta_{n+1+\alpha}^{m+\beta} + \theta_{n-\alpha}^{m+1-\beta} + \theta_{n+1}^m + \theta_n^m}{4} \right), \end{aligned} \quad (71)$$

$$\begin{aligned} & \sinh \left( \frac{\rho_{n+1}^m + \rho_n^{m+1} - \rho_{n+1}^{m+1} - \rho_n^m}{4} \right) = \delta^2 \exp \left( \frac{-\rho_{n+1}^m - \rho_n^{m+1} + \rho_{n+1}^{m+1} + \rho_n^m}{4} \right) \\ & - \delta^2 \exp \left( \frac{\rho_{n+1+\alpha}^{m+\beta} + \rho_{n-\alpha}^{m+1-\beta} - \rho_{n+1}^{m+1} - \rho_n^m}{4} \right) \cos \left( \frac{\theta_{n+1+\alpha}^{m+\beta} + \theta_{n-\alpha}^{m+1-\beta} + \theta_{n+1}^m + \theta_n^{m+1}}{4} \right) \\ & - \delta^4 \exp \left( \frac{\rho_{n+1+\alpha}^{m+\beta} + \rho_{n-\alpha}^{m+1-\beta} - \rho_{n+1}^m - \rho_n^{m+1}}{4} \right) \sinh \left( \frac{\rho_{n+1+\alpha}^{m+\beta} + \rho_{n-\alpha}^{m+1-\beta} - \rho_{n+1}^{m+1} - \rho_n^m}{4} \right), \end{aligned} \quad (72)$$

which is the delay-difference analogue of the sG equation. Equation (72) in the case of  $\alpha = \beta = 0$  is the discrete sG equation [32]

$$\sin \left( \frac{\theta_{n+1}^m + \theta_n^{m+1} - \theta_{n+1}^{m+1} - \theta_n^m}{4} \right) = \delta^2 \sin \left( \frac{\theta_{n+1}^m + \theta_n^{m+1} + \theta_{n+1}^{m+1} + \theta_n^m}{4} \right). \quad (73)$$

Now, let us apply the delay-differential limit

$$\delta \rightarrow 0, \quad n\delta = x, \quad \alpha\delta = \xi, \quad m\delta = y, \quad \beta\delta = \eta \quad (74)$$

to the bilinear equations (63), (64) and the  $N$ -soliton solution (66). Here  $x, y$  are the continuous variables, and  $\xi, \eta$  are the delay parameters. Consequently we obtain the bilinear equations

$$D_x D_y f(x, y) \cdot f(x, y) + 2(f(x, y)f(x, y) - g(x + \xi, y + \eta)g(x - \xi, y - \eta)) = 0, \quad (75)$$

$$D_x D_y g(x, y) \cdot g(x, y) + 2(g(x, y)g(x, y) - f(x + \xi, y + \eta)f(x - \xi, y - \eta)) = 0, \quad (76)$$

and the  $N$ -soliton solution

$$f(x, y) = \det \left( \delta_{ij} + \frac{\Phi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N}, \quad g(x, y) = \det \left( \delta_{ij} - \frac{\Phi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N}, \quad (77)$$

$$\Phi_i = \sqrt{-1} \mu_i \exp \left( (p_i - q_i)x - \left( \frac{1}{p_i} - \frac{1}{q_i} \right) y \right),$$

$$\frac{p_i}{q_i} = - \exp \left( \xi(p_i - q_i) - \eta \left( \frac{1}{p_i} - \frac{1}{q_i} \right) \right).$$

Setting  $f(x, y) = F(x, y) + \sqrt{-1}G(x, y)$  and  $g(x, y) = F(x, y) - \sqrt{-1}G(x, y)$ , we can rewrite (75) and (76) with

$$\left( D_x D_y - 4 \sinh \left( \frac{\xi D_x}{2} + \frac{\eta D_y}{2} \right) \sinh \left( \frac{\xi D_x}{2} + \frac{\eta D_y}{2} \right) \right) (F \cdot F - G \cdot G) = 0, \quad (78)$$

$$\left( D_x D_y + 4 \cosh \left( \frac{\xi D_x}{2} + \frac{\eta D_y}{2} \right) \cosh \left( \frac{\xi D_x}{2} + \frac{\eta D_y}{2} \right) \right) F \cdot G = 0. \quad (79)$$

In the case of  $\xi = \eta = 0$ , the bilinear equations (75), (76) (and also (78), (79)) and the  $N$ -soliton solution (77) lead to the sG equation and their  $N$ -soliton solution [32–34].

To construct a nonlinear form of the delay-differential analogue of the sG equation, we consider the dependent variable transformation

$$f(x, y) = \exp \left( \frac{\rho(x, y)}{4} + \sqrt{-1} \frac{\theta(x, y)}{4} \right), \quad g(x, y) = \exp \left( \frac{\rho(x, y)}{4} - \sqrt{-1} \frac{\theta(x, y)}{4} \right), \quad (80)$$

which is equivalent to

$$\theta(x, y) = 2\sqrt{-1} \log \frac{g}{f} = 4 \tan^{-1} \frac{G}{F}, \quad \rho(x, y) = 2 \log fg = 2 \log(F^2 + G^2). \quad (81)$$

By using this transformation, we can transform the bilinear equations (75) and (76) into the nonlinear delay-differential equation

$$\begin{aligned} \frac{\partial^2}{\partial x \partial y} \theta(x, y) &= -4 \exp \left( \frac{\rho(x + \xi, y + \eta) + \rho(x - \xi, y - \eta) - 2\rho(x, y)}{4} \right) \\ &\quad \times \sin \left( \frac{\theta(x + \xi, y + \eta) + \theta(x - \xi, y - \eta) + 2\theta(x, y)}{4} \right), \end{aligned} \quad (82)$$

$$\begin{aligned} \frac{\partial^2}{\partial x \partial y} \rho(x, y) &= -4 + 4 \exp \left( \frac{\rho(x + \xi, y + \eta) + \rho(x - \xi, y - \eta) - 2\rho(x, y)}{4} \right) \\ &\quad \times \cos \left( \frac{\theta(x + \xi, y + \eta) + \theta(x - \xi, y - \eta) + 2\theta(x, y)}{4} \right). \end{aligned} \quad (83)$$

which is the delay-differential analogue of the sG equation. In the case of  $\xi = \eta = 0$ , the above nonlinear equation leads to the sG equation [32, 33]:

$$\frac{\partial^2}{\partial x \partial y} \theta(x, y) = -4 \sin \theta(x, y). \quad (84)$$

We can obtain the above delay-differential analogue of the sG equation from the 2DTL equation by a reduction. The bilinear equation of the 2DTL equation

$$D_x D_y f_k(x, y) \cdot f_k(x, y) + 2(f_k(x, y) f_k(x, y) - f_{k+1}(x, y) f_{k-1}(x, y)) = 0 \quad (85)$$

has the following  $N$ -soliton solution [22, 27]:

$$\begin{aligned} f_k(x, y) &= \det \left( \delta_{ij} + \frac{\phi_i \psi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N} \\ &= \det \left( \delta_{ij} + \frac{\phi_j \psi_j}{p_i - q_j} \right)_{1 \leq i, j \leq N} = \sum_{I \subset \{1, \dots, N\}} \prod_{i \in I} \Phi_i \prod_{i < j, i, j \in I} \frac{(p_i - p_j)(q_i - q_j)}{(p_i - q_j)(q_i - p_j)}, \end{aligned} \quad (86)$$

$$\phi_i = \beta_i p_i^k e^{p_i x - p_i^{-1} y}, \quad \psi_i = \gamma_i q_i^{-k} e^{-q_i x + q_i^{-1} y},$$

$$\Phi_i = \beta_i \gamma_i \exp \left( (p_i - q_i) x - \left( \frac{1}{p_i} - \frac{1}{q_i} \right) y \right),$$

where  $p_i, q_i, \beta_i, \gamma_i$  are constants. We apply the reduction condition

$$f_{k+1}(x - \xi, y - \eta) = f_{k-1}(x + \xi, y + \eta) \quad (87)$$

and set

$$f(x, y) \equiv f_k(x, y) = f_{k-2}(x + 2\xi, y + 2\eta), \quad (88)$$

$$g(x, y) \equiv f_{k+1}(x - \xi, y - \eta) = f_{k-1}(x + \xi, y + \eta). \quad (89)$$

Then we obtain the bilinear equations

$$D_x D_y f(x, y) \cdot f(x, y) + 2(f(x, y)f(x, y) - g(x + \xi, y + \eta)g(x - \xi, y - \eta)) = 0, \quad (90)$$

$$D_x D_y g(x, y) \cdot g(x, y) + 2(g(x, y)g(x, y) - f(x + \xi, y + \eta)f(x - \xi, y - \eta)) = 0. \quad (91)$$

For the  $N$ -soliton solution, the constraint

$$\left(\frac{p_i}{q_i}\right)^2 = \exp\left(2\xi(p_i - q_i) - 2\eta\left(\frac{1}{p_i} - \frac{1}{q_i}\right)\right) \quad (92)$$

provides the reduction condition (87). The constraint (92) leads to

$$\frac{p_i}{q_i} = \pm \exp\left(\xi(p_i - q_i) - \eta\left(\frac{1}{p_i} - \frac{1}{q_i}\right)\right). \quad (93)$$

To satisfy the reality and regularity conditions, we can choose

$$\frac{p_i}{q_i} = -\exp\left(\xi(p_i - q_i) - \eta\left(\frac{1}{p_i} - \frac{1}{q_i}\right)\right) \quad (94)$$

and  $\beta_i \gamma_i = \sqrt{-1} \mu_i$  where  $\mu_i$  are real constants. Thus we obtain the delay-differential analogue of the sG equation and its  $N$ -soliton solution from the 2DTL equation. This construction of the delay sG equation does not require the delay-differential limit.

## 5. Conclusions

We have presented the systematic method to construct delay-difference and delay-differential analogues of soliton equations and their  $N$ -soliton solutions.

Our construction starts from the discrete KP equation (or discrete 2DTL equation) and use reduction and delay-differential limit. As examples, we have presented the delay-difference and delay-differential analogues of the LV, TL and sG equations and their  $N$ -soliton solutions. We have also presented another construction of delay-differential analogues of soliton equations starting from the semi-discrete KP, BT of 2DTL and 2DTL equations without applying a delay-differential limit. In the construction of them, the important thing is to integrate discrete variables with continuous variables by reduction.

In this paper, we have not discussed Lax pairs and conserved quantities of the delay soliton equations and the relationship to the delay-differential Painlevé equations. These problems remains to be revealed in future studies.

This work was partially supported by JSPS KAKENHI Grant Numbers 18K03435, 17H02856 and JST/CREST.

## References

- [1] Erneux T 2009 *Applied Delay differential equations* (New York: Springer)
- [2] Rihan F A 2021 *Delay differential equations and Applications to Biology* (Singapore: Springer)
- [3] Nakanishi K, Itoh A, Igarashi Y and Bando M 1997 Solvable optimal velocity models and asymptotic trajectory *Phys. Rev. E* **55** 6519
- [4] Hasebe K, Nakayama A and Sugiyama Y 1999 Exact solutions of differential equations with delay for dissipative systems *Phys. Lett. A* **259** 135–139
- [5] Tutiya Y and Kanai M 2007 Exact Shock Solution of a Coupled System of Delay Differential Equations: A Car-Following Model *J. Phys. Soc. Jpn.* **76** 083002
- [6] Matsuya K and Kanai M 2015 Exact solution of a delay difference equation modeling traffic flow and their ultra-discrete limit arXiv:1509.0786
- [7] Quispel G R W, Capel H W and Sahadevan R 1992 Continuous symmetries of differential-difference equations: the Kac-van Moerbeke equation and Painlevé reduction *Phys. Lett. A* **170** 379–383
- [8] Levi D and Winternitz P 1993 Symmetries and conditional symmetries of differential-difference equations *J. Math. Phys.* **34** 3713–3730
- [9] Grammaticos B, Ramani A and Moreira I D C 1993 Delay-differential equations and the Painlevé transcendents *Physica A* **196** 574–590
- [10] Ramani A, Grammaticos B and Tamizhmani K M 1993 Painlevé analysis and singularity confinement: the ultimate conjecture *J. Phys. A: Math. Gen.* **26** L53–L58
- [11] Joshi N 2008 Direct ‘delay’ reductions of the Toda equation *J. Phys. A: Math. Theor.* **42** 022001–022009
- [12] Joshi N and Spicer P E 2009 Direct “delay” reductions of the Toda hierarchy *J. Phys. Soc. Jpn.* **78** 0940061–0940065
- [13] Carstea A S 2011 Bilinear approach to delay-Painlevé equations *J. Phys. A: Math. Theor.* **44** 105202
- [14] Viallet C M 2014 Algebraic entropy for differential-delay equations arXiv:1408.6161
- [15] Halburd R and Korhonen R 2017 Growth of meromorphic solutions of delay differential equations *Proc. Amer. Math. Soc.* **145** 2513–2526
- [16] Berntson B K 2018 Special solutions of bi-Riccati delay-differential equations *SIGMA* **14** 020
- [17] Stokes A 2020 Singularity confinement in delay-differential Painlevé equations *J. Phys. A: Math. Theor.* **53** 435201
- [18] Villarroel J and Ablowitz M 1993 On the method of solution of the differential-delay Toda equation *Phys. Lett. A* **180** 413–418
- [19] Tsunematsu A, Nakata K, Tanaka Y and Maruno K 2021 Delay reductions of the two-dimensional Toda lattice equation, submitted
- [20] Hirota R 1981 Discrete analogue of a generalized Toda equation *J. Phys. Soc. Jpn.* **50** 3785–3791
- [21] Miwa T 1982 On Hirota’s difference equations *Proc. Japan Acad. A* **58** 9–12
- [22] Hirota R, Ito M and Kako F 1988 Two-dimensional Toda lattice equation *Prog. Theor. Phys. Suppl.* **94** 42–58
- [23] Ohta Y, Hirota R, Tsujimoto S and Imai T 1993 Casorati and discrete Gram Type determinant representations of solutions to the discrete KP hierarchy *J. Phys. Soc. Jpn.* **62** 1872–1886
- [24] Hirota R and Tsujimoto S 1995 Conserved quantities of a class of nonlinear difference-difference equations *J. Phys. Soc. Jpn.* **64** 3125–3127
- [25] Maruno K, Kajiwara K, Nakao S and Oikawa M 1997 Bilinearization of discrete soliton equations and singularity confinement *Phys. Lett. A* **229** 173–182
- [26] Hirota R and Satsuma J 1976  $N$ -soliton solutions of nonlinear network equations describing a Volterra system *J. Phys. Soc. Jpn.* **40** 891–900
- [27] Hirota R 2004 *The Direct Method in Soliton Theory* (Cambridge: Cambridge University Press)
- [28] Li C X, Lafortune S and Shen S F 2016 A semi-discrete Kadomtsev-Petviashvili equation and its coupled integrable system *J. Math. Phys.* **57** 053503
- [29] Ohta Y, Kajiwara K, Matsukidaira J and Satsuma J 1993 Casorati determinant solution for the relativistic Toda lattice equation *J. Math. Phys.* **34** 5190–5204

- [30] Hirota R 1977 Nonlinear partial difference equations II. Discrete-time Toda equation *J. Phys. Soc. Jpn.* **43** 2074–2078
- [31] Hirota R and Satsuma J 1976 A Variety of Nonlinear Network Equations Generated from the Bäcklund Transformation for the Toda Lattice *Prog. Theor. Phys. Suppl.* **59** 64–100
- [32] Hirota R 1977 Nonlinear partial difference equations III; Discrete sine-Gordon equation *J. Phys. Soc. Jpn.* **43** 2079–2086
- [33] Hirota R 1972 Exact solution of the Sine-Gordon equation for multiple collisions of solitons *J. Phys. Soc. Jpn.* **33** 1459–1463
- [34] Ablowitz M J and Segur H 1981 *Solitons and the inverse scattering transform* (Philadelphia: SIAM)