

# Wiener Process of Fractals and Path-Integrals II: Emergent Einstein-Hilbert Action in Stochastic Process of Quantum Fields along with the Ricci Flow of the Space Geometry

Amir Abbass Varshovi\*

*Faculty of Mathematics and Statistics, Department of Applied Mathematics  
and Computer Science, University of Isfahan, Isfahan, IRAN.  
School of Mathematics, Institute for Research in Fundamental  
Sciences (IPM), P.O. Box: 19395-5746, Tehran, IRAN.*

**Abstract:** This is the second paper of a series of researches (that commenced with [79] and is followed by [80]) that aims to interpret the gravitational effects of nature within some consistent stochastic fractal-based (intrinsically conformal) path-integral formulation. In the present work, the Wiener stochastic process is employed to study the Brownian motion of fractal functions on a closed Riemannian manifold with dynamical geometry due to the Ricci flow. It has been shown that the Wiener measure automatically leads to the Einstein-Hilbert action and the path-integral formulation of scalar quantum field theory at the first local approximation. This would be interpreted as the more fundamental formulation of quantum field theory in presence of gravity. However, we establish that the emergence of Einstein-Hilbert action is independent of the matter field interactions and is a merely entropic effect stemming from the nature of the Ricci flow. We also extract an explicit formula for the cosmological constant in terms of the Ricci flow and Hamilton's theorem for 3-manifold. Then, we discuss the cosmological features of the FLRW solution in  $\Lambda$ CDM Model via the derived equations of the Ricci flow. We also argue the correlation between our formulations and the entropic aspects of gravity. Finally, we establish some theoretical evidence that proves that the second law of thermodynamics is the basic source of and more fundamental than gravity.

**Keywords:** Wiener Stochastic Process, Fractal Norm, Laplace-Beltrami Operator, Closed Riemannian Manifold, Einstein-Hilbert Action, Hamilton's Theorem, Ricci Flow, Entropic Force, Cosmological Constant,  $\Lambda$ CDM Model, Second Law of Thermodynamics.

## I. INTRODUCTION

Employing the Wiener stochastic process [92] for Wierstrass-like functions (the so called fractal functions) on a flat space manifold we have proved in [79] that the scalar quantum field theory stems from the Brownian motion of fractal functions along the time direction.<sup>1</sup> Actually, this consequence stands on a complex footing which consists of **i) Fractals** and **ii) Brownian motion**. In fact, fractal functions are continuous but nowhere differentiable functions on the whole (or a widespread subset of)

---

\*Electronic address: [ab.varshovi@sci.ui.ac.ir](mailto:ab.varshovi@sci.ui.ac.ir)/[amirabbassv@gmail.com](mailto:amirabbassv@gmail.com)/[amirabbassv@ipm.ir](mailto:amirabbassv@ipm.ir)

<sup>1</sup> See [30] for a nice presentation of the Wiener stochastic process and its applications as a probability measure.

of the underlying domain. We studied them according to the properties of the Weierstrass functions [89] and their asymptotic behavior in Fourier coefficients. Following the Hardy's results [47] we obtained a non-linear differential equation for such asymptotic behaviors of fractal functions. By solving the differential equation we established that  $f : S \rightarrow \mathbb{R}$ , with  $S = [-L, L]^D$ , is a fractal function if for each  $\sigma > 0$ ,  $\mu \geq 0$ , and  $\ell > 0$  there exists some Fourier mode  $n = (n_1, \dots, n_D) \in \mathbb{Z}^D$ , so that

$$\int_0^{|f_n|} e^{\sigma(\mu^2+n^2)x^2/2} dx > \ell, \quad (\text{I.1})$$

wherein  $f_n$  is the  $n$ -th Fourier coefficient of  $f$ , and  $n^2 = \sum_{i=1}^D n_i^2$ . In fact, if we set  $\Delta = -\sum_{i=1}^D \partial_i^2$  to be the Laplacian operator, then we could readily adjust (I.1) for the spectrum of the Laplacian operators. Let  $\{\psi_{\lambda,i}\}_{\lambda \in \text{spec} \Delta}^{i \in D_\lambda}$  be the complete set of orthonormal eigenfunctions of  $\Delta$  (i.e.  $\int_S \psi_{\lambda,i} \psi_{\lambda',i'} d^D x = \delta_{\lambda\lambda'} \delta_{ii'}$ ) for eigenvalues  $0 \leq \lambda s$  and some possible degeneracy space  $D_\lambda$  labeled by  $i$ . Then  $f : S \rightarrow \mathbb{R}$  is a fractal function if for each  $\sigma > 0$ ,  $\mu \geq 0$ , and  $\ell > 0$  there exists some Fourier mode  $(\lambda, i)$  so that

$$\int_0^{|f_{\lambda,i}|} e^{\sigma(\mu^2+\lambda)x^2/2} dx > \ell, \quad (\text{I.2})$$

wherein  $f_{\lambda,i} = \int_S \psi_{\lambda,i} f d^D x$ . In principle, the mechanism of which we employed in [79] for working out the fractality condition (I.3) on the space set  $S = [-L, L]^D$  (i.e. fractal periodic functions with period  $2L$  in each variable), could be generalized to any compact Riemannian manifold  $(M, g)$ . Actually, here we may concentrate on manifolds without boundary.<sup>2</sup> It is well-known that if  $(M, g)$  is a closed  $D$ -dimensional Riemannian manifold then the spectrum of the Laplacian  $\Delta$  is discrete and consists of an increasing sequence  $\{\lambda_k\}_{k=1}^\infty$  of eigenvalues (counted with degeneracies) with  $\lambda_1 = 0$  and  $\lambda_k \rightarrow \infty$  as  $k \rightarrow \infty$  [41].

On the other hand, the eigenfunctions (resp. the eigenforms) of the Laplacian operator provide a pure point spectrum, which means that  $L^2 M = L^2(\Lambda^0 M)$  (resp.  $L^2(\Lambda^n M)$ ,  $n > 0$ ) admits a complete orthonormal basis consisting of eigenfunctions  $\psi_{\lambda,i}$  of  $\Delta$  with associated eigenvalues  $\lambda s$ , i.e.  $\Delta \psi_{\lambda,i} = \lambda \psi_{\lambda,i}$ . This leads to a generalized Fourier expansion of continuous functions (resp. forms) on compact Riemannian manifolds. An extensive amount of investigations have been made concerning the eigenvalues and their relationship to the background Riemannian geometry.<sup>3</sup> These studies pertain to upper and lower bounds for eigenvalues and their asymptotics which are of the most interest in our study. There is a major conclusion that is valid for all compact Riemannian geometry; The Weyl's asymptotic law [90, 91] which asserts that  $\lambda_k$  asymptotically behaves as  $k^{2/D}$  as  $k \rightarrow \infty$ .

It is worth to remind that there are significant results for the first gap of the Laplacian eigenvalues:

- a)** The Yau's theorem [67, 94] which provides a lower bound for non-vanishing Laplacian eigenvalues on Riemannian manifolds with non-negative Ricci curvature as  $\lambda_1 \geq \frac{\pi^2}{d^2}$ , where  $d$  is the diameter of  $M$ .
- b)** The Lichnerowicz's theorem [36] for positive definite Ricci curvature, which says that  $\lambda_1 \geq KD$  whenever  $Ric_{\mu\nu} \geq (D-1)Kg_{\mu\nu}$ .
- c)** The Simon's lower bound for an Einstein's manifold of  $D \geq 3$  dimensions whose sectional curvature is bounded from below by a constant  $K_0$ , which states that  $M$

<sup>2</sup> So the Neumann and Dirichlet boundary conditions for extracting the eigenvalues and the eigenspaces of the Laplace-Beltrami operator would be irrelevant.

<sup>3</sup> See for example [3, 6, 8, 13, 14, 17, 22, 27, 36-39, 41, 49, 61, 62, 67, 69, 77, 90, 91, 94] and the references therein.

is either isometric to  $D$ -sphere  $S^D$  or else  $\lambda_1 > 2DK_0$  [77].<sup>4</sup> **d)** And finally, Cheeger's isoperimetric constant puts an important geometric lower bound for  $\lambda_1$ , mostly referred to as the Cheeger's inequality [14]. This inequality also gives rise to an upper bound for  $\lambda_1$  [8] which is essentially given in terms of the Ricci curvature of  $(M, g)$  [62].<sup>5</sup>

However, Weyl's asymptotic result guarantees that the asymptotic behavior of the Laplace-Beltrami operator on each  $D$ -dimensional closed Riemannian manifold equals that of flat  $D$ -torus  $\mathbb{T}^D$ , which we essentially considered in [79]. Moreover, the fractality condition is in particular a local property that can be studied by means of the harmonic analysis on Riemannian manifolds. Indeed, according to the Sobolev's theorem [40] any element of the Sobolev's space  $\mathcal{H}_t$  with  $t = [D/2] + 1 + s$ ,  $0 \leq s < 1$ , will uniformly converge to a continuous function whose derivative fails to converge on  $M$ , hence is a Weierstrass-like function.<sup>6</sup> Here, the Sobolev class of  $f : M \rightarrow \mathbb{R}$  is studied locally on  $M$  and the constant  $s$  could be considered as the self-similarity scale of the fractal function, as was pointed out by Hardy [47].

Since, the main criteria for studying the fractality in [79] stemmed from the asymptotic behavior of Fourier-Laplace coefficients (i.e. the expansion coefficients in terms of the eigenfunctions of the Laplacian operator on  $S = [-L, L]^D \cong \mathbb{T}^D$ ) of Weierstrass-like functions we readily obtain the same differential equations for such coefficients on closed Riemannian manifold  $M$ . Therefore, we have already obtained the generalized version of **Theorem 2** of [79]:<sup>7</sup>

**Theorem 1;** *Let  $(M, g)$  be a closed Riemannian manifold with Laplacian operator  $\Delta = dd^\dagger + d^\dagger d$ . Let  $\{\psi_{\lambda, i}\}_{\lambda \in \text{spec} \Delta}^{i \in D_\lambda}$  be the complete set of orthonormal eigenfunctions of  $\Delta$  for eigenvalues  $0 \leq \lambda s$  and some possible degeneracy space  $D_\lambda$  labeled by  $i$ . That is:  $\int_M \psi_{\lambda, i} \psi_{\lambda', i'} d\Omega_g = \delta_{\lambda\lambda'} \delta_{ii'}$ , for  $d\Omega_g$  the Riemannian volume form on  $M$ . Then,  $f : M \rightarrow \mathbb{R}$  is a fractal function if for each  $\sigma > 0$ ,  $\mu \geq 0$ , and  $\ell > 0$  there exists some Laplacian Fourier mode  $(\lambda, i)$  so that*

$$\int_0^{f_{\lambda, i}} e^{\sigma(\mu^2 + \lambda)x^2/2} dx > \ell, \quad (\text{I.3})$$

wherein  $f_{\lambda, i} = \int_M f \psi_{\lambda, i} d\Omega_g$ .

Employing the above theorem we can accomplish the Wiener stochastic process of fractal functions on general closed Riemannian manifold  $(M, g)$ . Through the next section, we follow the strategy of [79] on a given closed Riemannian manifold (as the manifold of space) and work out the generalized result of [79]. In fact, we will establish that a plain generalization of the Wiener Brownian process for propagating fractal functions on  $M$  will lead to the path-integral formulation of scalar quantum field theory on static curved space-time.

<sup>4</sup> See [18, 76] for more detailed review about the harmonic theory in Riemannian geometry and the lower bound of  $\lambda_1$  in Einstein-like and constant scalar curvature manifolds.

<sup>5</sup> See [61] for generalization of Cheeger's inequalities to the magnetic Laplacian  $\Delta^\alpha$  defined for  $d \mapsto d + \alpha$ ,  $\alpha \in \Omega^1 M$ , on closed Riemannian manifold  $M$ . See also [27] for more recent achievements about the eigenvalues of the magnetic Laplacian  $\Delta^\alpha$ .

<sup>6</sup> See chapter 6 of [86] for detailed applications of Sobolev's spaces on Riemannian manifolds.

<sup>7</sup> We omit the detailed arguments of the issue here and refer the reader to sections II and III of [79] for a more comprehensive discussion.

However, the main objective of this paper is accomplished in section III. Actually, in this section, the Wiener fractal measure is implemented for dynamical space-time with some time-dependent metric  $g_{ij}(t)$ . In fact, we will establish that the Einstein-Hilbert action can be rigorously worked out from an entropic evolution of the space geometry due to the Ricci flow. Moreover, we will establish that the emergence of gravity in this path-integral formulation is independent of the quantum field theory and is a merely entropic effect. Therefore, although the expectation value of the stress-energy tensor of the involved quantum matter fields will affect the metric of space  $M$  the time evolution of the metric is imposed by an entropic law comparable with the second law of thermodynamics. Hence, the Ricci flow could be assumed as a natural gauge fixing term for Einstein's field equation.

In section IV we study the cosmological results of the extracted stochastic-entropic theory of gravity, the so called *Wiener fractal gravity*, and work out the cosmological constant due to Hamilton's theorem for 3-manifolds [43]. Next, we will give a different interpretation for inflation and the dark energy. Comparison with the  $\Lambda$ CDM Model and the admitted FLRW solution in cosmology is concerned due in this section. Finally, in section V we study the entropic features of the Ricci flow according to Perelman's seminal paper [75], and we will conclude that the second law of thermodynamics could be regarded as the basic source of gravitational effects in nature. We then compare the Wiener fractal gravity with Verlinde's entropic gravity, stochastic gravity and Horava-Lifshitz gravity.

## II. WIENER FRACTAL MEASURE AND QUANTUM FIELDS ON CURVED SPACETIME

In fact, **Theorem 1** lets us to define the (*massive*) *fractal norm* on Riemannian manifold  $(M, g)$  as;

$$\ell_{\sigma, \mu}(f_{\lambda, i}) = \int_0^{|f_{\lambda, i}|} e^{\sigma(\mu^2 + \lambda)x^2/2} dx. \quad (\text{II.1})$$

Let  $\mathcal{N} \in \mathbb{N}$  and define  $C_{\mathcal{N}}$  be the collection of real functions  $f : M \rightarrow \mathbb{R}$  which are finite linear combinations of Laplacian eigenfunctions  $\psi_{\lambda, i}$  with eigenvalues  $0 \leq \lambda \leq \mathcal{N}$ . Each individual element of  $C_{\mathcal{N}}$  is conventionally called an  $\mathcal{N}$ -*bounded function (on  $M$ )*. Then, the fractal norm (II.1) leads to a *Lebesgue fractal measure*  $d\mu_{LF}$  on  $C_{\mathcal{N}}$  as:

$$d\mu_{LF} = \prod_{i=1}^{D_{\mathcal{N}}} dx_{\lambda, i}, \quad dx_{\lambda, i} = e^{\sigma(\mu^2 + \lambda)f_{\lambda, i}^2/2} df_{\lambda, i}, \quad (\text{II.2})$$

wherein  $D_{\mathcal{N}}$  is the dimension of  $C_{\mathcal{N}}$ . Since  $M$  is compact  $C_{\mathcal{N}}$  is finite-dimensional (i.e.  $D_{\mathcal{N}} < \infty$ ), hence  $d\mu_{LF}$  is well-defined. Now, we are ready to study the Brownian motion of  $\mathcal{N}$ -bounded functions by means of the Wiener stochastic process and the background Lebesgue fractal measure  $d\mu_{LF}$ . This Brownian motion is defined by means of an appropriate *Wiener fractal measure* on  $C_{\mathcal{N}}$  for Riemannian manifold  $(M, g)$ . Actually, the definition is straightforward and is given upon what we have obtained in [79]. We remember that the symmetric Wiener measure for a stochastic process in target space  $\mathbb{R}^D$  from the initial position  $x_I$  at  $t_0 = -T$  to the final position  $x_F$  at  $t_{N+1} = T$ , within the time slicing

$-T = t_0 < t_1 < \dots < t_N < t_{N+1} = T$  is:

$$\begin{aligned} dW(t_N, \dots, t_1) &= (4\pi T)^{D_N/2} \exp(|x_F - x_I|^2/4T) \\ &\times \frac{1}{(2\pi\tau_{N+1})^{D_N/2}} e^{-|x_F - x_N|^2/2\tau_{N+1}} d^D x_N \frac{1}{(2\pi\tau_N)^{D_N/2}} e^{-|x_N - x_{N-1}|^2/2\tau_N} \times \dots \\ &\times \dots \times \frac{1}{(2\pi\tau_2)^{D_N/2}} e^{-|x_2 - x_1|^2/2\tau_2} d^D x_1 \frac{1}{(2\pi\tau_1)^{D_N/2}} e^{-|x_1 - x_I|^2/2\tau_1}, \end{aligned} \quad (\text{II.3})$$

for  $x_i = x(t_i)$ ,  $\tau_i = t_i - t_{i-1}$ , and  $|x_i - x_{i-1}|^2 = \sum_{j=1}^D (x_i^j - x_{i-1}^j)^2$ . Therefore, the symmetric Wiener for Brownian motion of fractal functions on  $M$  restricted to  $C_N$  would be (assuming  $x_I = x_F = 0$ ):

$$dW(t_N, \dots, t_1) = (4\pi T)^{D_N/2} \left( \prod_{k=1}^{N+1} \Xi_k \right) \times \left( \prod_{k=1}^N d^{D_N} f_{\lambda,i}(t_k) \right), \quad (\text{II.4})$$

for

$$\begin{aligned} \Xi_k &= \frac{1}{(2\pi\tau_k)^{D_N/2}} \times \exp \left( -\frac{\tau_k}{2} \sum_{\lambda \in \text{spec}\Delta} \sum_{i \in D_\lambda} \left( \frac{1}{\tau_k} \int_{f_{\lambda,i}(t_{k-1})}^{f_{\lambda,i}(t_k)} e^{\sigma(\lambda+\mu^2)x^2/2} dx \right)^2 \right) \\ &\times \exp \left( \frac{\sigma}{2} \int_M (g^{ij} \partial_i f(t_{k-1}) \partial_j f(t_{k-1}) + \mu^2 f^2(t_{k-1})) d\Omega_g \right), \end{aligned} \quad (\text{II.5})$$

We refer to (II.4) as the *Wiener fractal measure*. Being of a nonlocal formula, this probability measure could be turned to a local structure by employing the following approximation as  $N \rightarrow \infty$ :

**Strategy A:** When  $\tau_k \rightarrow 0$  (i.e.  $N \rightarrow \infty$ ) we can simply apply the following replacement within the Wiener fractal measure:

$$\frac{1}{\tau_k} \int_{f_{\lambda,i}(t_{k-1})}^{f_{\lambda,i}(t_k)} e^{\sigma(\lambda+\mu^2)x^2/2} dx \quad \mapsto \quad \frac{\partial f_{\lambda,i}}{\partial t}(t_{k-1}), \quad (\text{II.6})$$

which leads to:

$$\Xi_k = \frac{1}{(2\pi\tau_k)^{D_N/2}} \exp \left( -\tau_k \int_M \frac{1}{2} \left\{ (\partial_t f(t_{k-1}))^2 - \frac{\sigma}{\tau_k} (g^{ij} \partial_i f(t_{k-1}) \partial_j f(t_{k-1}) + \mu^2 f^2(t_{k-1})) \right\} d\Omega_g \right). \quad (\text{II.7})$$

Actually, upon the strategy **A** we readily find:

$$\prod_{k=1}^{N+1} \Xi_k = \left( \frac{1}{(2\pi\theta)^{D_N/2}} \right)^{N+1} \exp \left( -\int_{\mathcal{M}} \frac{1}{2} \left\{ (\partial_t f)^2 - \frac{\sigma}{\theta} (g^{ij} \partial_i f \partial_j f + \mu^2 f^2) \right\} dt \wedge d\Omega_g \right), \quad (\text{II.8})$$

wherein  $\mathcal{M} = [-T, T] \times M$  is the space-time continuum,<sup>8</sup> and  $\theta = dt = 2T/N$  ( $\rightarrow 0$ ) is the unit of uniform time slicing. Set  $c = \sqrt{\sigma/\theta}$  to be the speed of light in vacuum, then we obtain:

$$\prod_{k=1}^{N+1} \Xi_k = \left( \frac{1}{(2\pi\theta)^{D_N/2}} \right)^{N+1} \exp \left( -c \int_{\mathcal{M}} \frac{1}{2} \left\{ \mathbf{g}^{\mu\nu} \partial_\mu f \partial_\nu f - \mu^2 f^2 \right\} d\Omega \right), \quad (\text{II.9})$$

<sup>8</sup> Indeed, as we pointed out in [79] we distinguish between terminologies *spacetime* and *space-time*, for which the former is realized to admit space and time dimensions substantially as an identical essence, whereas the latter embraces an inherent difference between them.

where  $d\Omega = dx^0 \wedge d\Omega_g$  is the Riemannian volume form on the space-time continuum  $\mathcal{M}$  for the induced time-independent metric  $\mathbf{g} = \mathbf{g}_{\mu\nu} dx^\mu \otimes dx^\nu = dx^0 \otimes dx^0 - g_{ij} dx^i \otimes dx^j$  with  $x^0 = ct$ . Note that  $\mathbf{g}$  is a Lorentzian metric with signature  $(+, -, -, -)$ , hence the theory is locally Lorentz invariant. If one defines  $f = \gamma^{-1}\phi$  for normalization factor  $\gamma = \sqrt{\hbar c}$  we read:

$$\prod_{k=1}^{N+1} \Xi_k = \left( \frac{1}{(2\pi\theta)^{D_{\mathcal{N}}/2}} \right)^{N+1} \exp \left( -\frac{1}{\hbar} \int_{\mathcal{M}} \frac{1}{2} \left\{ \mathbf{g}^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{m^2 c^2}{\hbar^2} \phi^2 \right\} d\Omega \right), \quad (\text{II.10})$$

for  $m = \hbar\mu/c$  to be the physical mass.<sup>9</sup> Hence, for the Feynman measure

$$\mathfrak{D}\phi = (4\pi T)^{D_{\mathcal{N}}/2} \left( \frac{1}{(2\pi\theta)^{D_{\mathcal{N}}/2}} \right)^{N+1} \gamma^{-ND_{\mathcal{N}}} \left( \prod_{k=1}^N d^{D_{\mathcal{N}}} \phi_{\lambda,i}(t_k) \right), \quad (\text{II.11})$$

the Wiener fractal measure with strategy **A** becomes:

$$dW_{\mathbf{A}}(t_N, \dots, t_1) = \exp \left( -\frac{1}{\hbar} \int_{\mathcal{M}} \frac{1}{2} \left\{ \mathbf{g}^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{m^2 c^2}{\hbar^2} \phi^2 \right\} d\Omega \right) \mathfrak{D}\phi. \quad (\text{II.12})$$

However, as we discussed in [79] the replacement (II.6) causes the heat kernel of the Gaussian term of the Wiener measure to be replaced with a quadratic function which is not always dominating. Therefore, the Wiener fractal measure (II.12) is not well-defined over  $C_{\mathcal{N}}$ . To overcome this subtlety, we should transfer to complex analysis. As we pointed out in [79] incorporating an infinitesimal heat kernel term after analytic continuation of the Wiener fractal measure would compensate some of the pathologies emerging from replacement (II.6) via the equation:

$$\lim_{\varepsilon \rightarrow 0} \left| \int_{\mathbb{R}} e^{(ia-\varepsilon)x^2} dx \right| = \int_{\mathbb{R}} e^{-|a|x^2} dx. \quad (\text{II.13})$$

Then, the next strategy is:

**Strategy B:** *To compensate the pathology of strategy A we should employ the analytic continuation of the Wiener fractal measure with an augmented heat kernel  $\varepsilon$ -term due to (II.13). Hence, we must obtain:*

$$\begin{aligned} dW(t_N, \dots, t_1) &\approx dW_{\mathbf{A},\mathbf{B}}(t_N, \dots, t_1) \\ &= \exp \left( \frac{i}{\hbar} \int_{\mathcal{M}} \frac{1}{2} \left\{ \mathbf{g}^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{(m^2 - i\varepsilon)c^2}{\hbar^2} \phi^2 \right\} d\Omega \right) \mathfrak{D}\phi, \\ &= \exp \left( \frac{i}{\hbar} \int_{\mathcal{M}} \frac{1}{2} \sqrt{|\det \mathbf{g}|} \left\{ \mathbf{g}^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{(m^2 - i\varepsilon)c^2}{\hbar^2} \phi^2 \right\} d^{D+1}x \right) \mathfrak{D}\phi, \end{aligned} \quad (\text{II.14})$$

within the limit of  $\varepsilon \rightarrow 0$ .

In particular, after employing strategies **A** and **B** we are effectively worked out the path-integral formulation of a scalar quantum field theory on a curved space-time. In principle, a well-defined (renormalizable) quantum field theory on curved space-time  $\mathcal{M}$  is defined for a sequence of Wiener integrable

<sup>9</sup> For more detailed discussion about the relations between fractal parameters  $\sigma$  and  $\mu^2$  and the physical constants  $\hbar$  and  $c$  see [79].

functions, say  $F^{(\mathcal{N})}(t_N, \dots, t_1) = \prod_{k=1}^N \exp(\tau_k \int_M V^{(\mathcal{N})}(\phi(t_{k-1})) d\Omega)$ , the so-called interaction term, each of which being defined over  $C_N$ , so that the Wiener path-integrals

$$\begin{aligned} I_N &= \int_{C_N} F^{(\mathcal{N})}(t_N, \dots, t_1) \{\phi(t_{i_1}) \cdots \phi(t_{i_n})\} dW(t_N, \dots, t_1) \\ &\approx \int_{C_N} F_{\mathbf{B}}^{(\mathcal{N})}(t_N, \dots, t_1) \{\phi(t_{i_1}) \cdots \phi(t_{i_n})\} dW_{\mathbf{A}, \mathbf{B}}(t_N, \dots, t_1) \end{aligned} \quad (\text{II.15})$$

result in finite amplitudes as  $\mathcal{N} \rightarrow \infty$  (and  $N \rightarrow \infty$  as an appropriate monotone increasing function of  $\mathcal{N}$  [79]) for each  $n$ , wherein  $F_{\mathbf{B}}^{(\mathcal{N})}(t_N, \dots, t_1) = \prod_{k=1}^N \exp(-i\tau_k \int_M V^{(\mathcal{N})}(\phi(t_{k-1})) d\Omega)$ . As we discussed in [79] employing negative renormalizable potential terms  $V^{(\mathcal{N})}$  always lead to finite and predictable expectation values  $I_N$  as  $\mathcal{N}, N \rightarrow \infty$ . Indeed, the former property ensures that the energy levels are bounded from below, and the latter one puts forward a reasonable correlation between  $V^{(\mathcal{N})}$ s (and the corresponding coupling constants) at different regimes of energy labeled with  $\mathcal{N}$ .<sup>10</sup>

In particular, a similar mechanism to the above machinery could be employed to work out the Maxwell and the Yang-Mills actions on curved space-time  $\mathcal{M}$  [80]. Actually, for such gauge theories, one should replace the Fourier expansion of the Laplacian operator with that of the generalized Bochner Laplacian defined on principal bundles over the space manifold  $M$ . Therefore, one may claim that the Standard Model, as the admitted prescription for the fundamental forces of nature, including gauge fields, fermions and the scalar Higgs boson, could be described and interpreted fascinatingly as a Brownian process of geometric structures over the space manifold, i.e. the universe. In the next section we show that the gravitation effect, within the framework of general relativity, is spectacularly worked out by evolving the cosmos geometry via an entropic equation.

### III. EINSTEIN-HILBERT ACTION AND THE RICCI FLOW OF THE SPACE METRIC

Let us revisit the symmetric Wiener measure (II.4) once again. Suppose that we need to perform some change of variables  $F_i : \mathbb{R}^D \rightarrow \mathbb{R}^D$  at each intermediate time section  $t_i$ ,  $1 \leq i \leq N$ .

$$\begin{aligned} dW(t_N, \dots, t_1) &= \mathfrak{C}_N \frac{1}{(2\pi\tau_N)^{D/2}} e^{-|\Delta x_N|^2/2\tau_N} d^D \Delta x_N \times \cdots \times \frac{1}{(2\pi\tau_1)^{D/2}} e^{-|\Delta x_1|^2/2\tau_1} d^D \Delta x_1 \\ &= \mathfrak{C}_N \frac{1}{(2\pi\tau_N)^{D/2}} e^{-|F_N(\Delta x_N)|^2/2\tau_N} d^D F_N(\Delta x_N) \times \cdots \times \frac{1}{(2\pi\tau_1)^{D/2}} e^{-|F_1(\Delta x_1)|^2/2\tau_1} d^D F_1(\Delta x_1) \quad (\text{III.1}) \\ &= \mathfrak{C}_N \frac{1}{(2\pi\tau_N)^{D/2}} e^{-|F_N(\Delta x_N)|^2/2\tau_N} J_N d^D \Delta x_N \times \cdots \times \frac{1}{(2\pi\tau_1)^{D/2}} e^{-|F_1(\Delta x_1)|^2/2\tau_1} J_1 d^D \Delta x_1, \end{aligned}$$

for  $\Delta x_i = x_i - x_{i-1}$ , and  $\mathfrak{C}_N = \left(\frac{2T}{\tau_{N+1}}\right)^{D/2} \exp(|x_F - x_I|^2/4T - |x_F - x_N|^2/2\tau_{N+1})$ ,<sup>11</sup> where  $J_i$  is the Jacobian determinant of transformation  $F_i$ . We can simply employ (III.1) for a dynamical geometry of space Riemannian manifold  $M$ . This geometric evolution is absolutely independent of

<sup>10</sup> For more details about the renormalizability of a scalar quantum field theory within the Wiener fractal mechanism see [79]. Also see [20, 51, 84, 85] for more discussions about quantization, regularization, renormalization, Casimir effect, Bogoliubov transformation and the  $S$ -matrix components of such theories. Moreover, we refer the readers to [55] for an interesting review on the topic.

<sup>11</sup> The contribution of  $x_N$  in  $\mathfrak{C}_N$  would provide an integration on the last time section within the total action, hence it is essentially a null set integral and can be easily removed from the calculations.

the topology of  $M$ , but rather it would have been related to the differential geometric structures due; hence the Riemannian metric  $g$ . Therefore, since the Wiener fractal measure was worked out for the Fourier expansion in terms of the Laplacian eigenfunctions on  $(M, g)$ , the time evolution of  $g$  would cause some change of variables within the measure as we considered in (III.1). The next theorem provides a consistent framework for such transformations versus infinitesimal variation of the metric.<sup>12</sup>

**Theorem 2 (ref. [4]);** *Let  $g(t) = g + th$ ,  $|t| \ll 1$ , be a small variation of the metric on Riemannian manifold  $(M, g)$ , for  $h$  a symmetric  $(0, 2)$ -tensor field on  $M$ . Set  $\Delta^{g(t)}$  and  $d\Omega_{g(t)}$  to be respectively the Laplacian operator and the Riemann volume form due to the metric  $g(t)$ . Let  $\lambda$  be an eigenvalue of  $\Delta = \Delta^{g(0)}$  with degeneracy space  $D_\lambda$  labeled by  $i$ . Then, there exist  $\psi_{\lambda,i}(t) \in C^\infty(M)$ , for each  $\lambda \in \text{spec}\Delta$ , and  $i \in D_\lambda$ , such that;*

- a)  $\Delta^{g(t)}\psi_{\lambda,i}(t) = \lambda_i(t)\psi_i(t)$ , for each  $\lambda \in \text{spec}\Delta$  and  $i \in D_\lambda$ , with  $t$ ,  $|t| \ll 1$ .
- b)  $\lambda_i(t)$  and  $\psi_{\lambda,i}(t)$  depend analytically on  $t$ ,  $|t| \ll 1$ , with  $\lambda_i(0) = \lambda$ , for each  $i \in D_\lambda$ .
- c)  $\{\psi_{\lambda,i}(t)\}_{\lambda \in \text{spec}\Delta}^{i \in D_\lambda}$  is orthonormal with respect to integration on  $M$  with  $d\Omega_{g(t)}$ .

Actually, we assume that the space metric would vary with the Ricci flow:

$$\frac{\partial}{\partial t}g_{ij}(t) = -2\text{Ric}_{ij}(t), \quad (\text{III.2})$$

for  $\text{Ric}(t)$  the Ricci curvature of evolving metric  $g(t)$ . In fact, as we will argue in the following, we regard the Ricci flow as an imposed natural entropic gauge fixing term for Einstein's field equation. One should note that although the Einstein equation is scale invariant, i.e. is symmetric with respect to transformation  $g_{ij} \rightarrow cg_{ij}$  for some  $c > 0$ , the Ricci flow disobeys this property. Hence, once the scale of time dimension  $t$  is fixed (see Eq. (III.3) in below where the time scale  $\zeta$  is introduced) the Ricci flow fixes the scale of  $g_{ij}$  and consequently the space volume. Recently, the Ricci flow has attracted a lot of attention in building a well-defined model of (topological) quantum gravity and several research have shown some theoretical evidence for an intimate correlation between gravity and the Ricci flow.<sup>13</sup>

Actually, the Ricci flow has been established to be a very fruitful mechanism to improve metrics in Riemannian geometry whenever  $M$  is compact. Indeed, it has been shown that in some specific conditions the Ricci flow converges to a canonical metric [43–45, 57]. For instance, in his seminal work, Hamilton has established that on each closed 3-manifold with positive Ricci curvature the Ricci flow converges to a canonical metric of positive constant sectional curvature [43]. In this sense, the Ricci flow may be regarded as a natural homotopy between a given metric of positive Ricci curvature and a canonical metric of constant sectional curvature. Moreover, Hamilton demonstrated that in this case the Ricci flow could be resolved within a normalized version of which the volume of the manifold is preserved along with the metric evolution. In principle, whenever the Ricci flow holds and admits a unique solution on  $(a, b) \subset \mathbb{R}$ , it converges to the canonical metric  $g_c$  as  $t \rightarrow b$ , where  $g_c$  is necessarily

<sup>12</sup> For the variation of eigenvalues and eigenfunctions of the Laplacian (and its generalized versions such as  $(p, q)$ -Laplacian and the Witten Laplacian) versus the metric evolution see for example [1, 4–6, 10, 21, 28, 29, 53, 58, 66] and the references therein

<sup>13</sup> See for example [23–26, 31–33, 42, 59, 68, 71, 93].

an Einstein metric, i.e.  $\lim_{t \rightarrow b} Ric_{ij}(t) = K g_{c_{ij}}$ , for some constant  $K$ .<sup>14</sup>

Without loss of generality we may assume that the Ricci flow exists on  $(-a, a) \subset \mathbb{R}$ , for some  $a > 0$ . Hence, by rescaling  $t \mapsto t/n$ ,  $n \in \mathbb{N}$ , the domain of the solution changes to  $(-na, na)$ , while on the right hand side of (III.3) one must replace  $-2$  with  $-2/n$ . In principle, since we concern about the stochastic process in  $(-T, T)$  for  $T \rightarrow \infty$ , the integer  $n$  must be chosen as large as possible. Generally, this Ricci flow could be rescaled as

$$\frac{\partial}{\partial t} g_{ij}(t) = -2\zeta Ric_{ij}(t), \quad (III.3)$$

for some small positive constant  $\zeta$ . In the following we will see that as a very tiny amplitude  $\zeta$  couples to negligible terms inserted in the Einstein-Hilbert Lagrangian density. More precisely, the Ricci flow scaling factor could be compared with the cosmological constant  $\Lambda$ .

However, it can be easily seen that the variation of the Riemannian volume form  $d\Omega_{g(t)}$  with respect to  $t$  is given in terms of the scalar curvature  $R(t)$ :

$$\frac{\partial}{\partial t} d\Omega_{g(t)} = -\zeta R(t) d\Omega_{g(t)}. \quad (III.4)$$

The above equation lets us to work out the explicit formulation of the Wiener fractal measure (III.1) along the Ricci flow time evolution. Initially, one should note that we have already considered a normalization condition as:

$$\int_M \psi_{\lambda,i}(t) \psi_{\lambda,j}(t) d\Omega_{g(t)} = \delta_{ij}. \quad (III.5)$$

Hence, by variation of (III.5) with respect to  $t = \delta t$  we obtain:

$$\begin{aligned} \epsilon_{\lambda,ii}(t) &= 1 - \frac{t\zeta}{2} \int_M R(t) \psi_{\lambda,i}^2(t) d\Omega_{g(t)}, \\ \epsilon_{\lambda,ij}(t) + \epsilon_{\lambda,ji}(t) &= -t\zeta \int_M R(t) \psi_{\lambda,i}(t) \psi_{\lambda,j}(t) d\Omega_{g(t)}, \quad (i \neq j) \end{aligned} \quad (III.6)$$

wherein  $\epsilon_{\lambda,ij}(t) = \int_M \psi_{\lambda,i}(0) \psi_{\lambda,j}(t) d\Omega_{g(t)}$ . Hence, we would readily set;

$$\epsilon_{\lambda,ij}(t) = \delta_{ij} - \frac{t\zeta}{2} \int_M R(t) \psi_{\lambda,i}(t) \psi_{\lambda,j}(t) d\Omega_{g(t)}. \quad (III.7)$$

Consequently, along the variation of  $g$  the basis of  $C_{\mathcal{N}}$  move linearly by the matrix  $\epsilon = \oplus_{\lambda} \epsilon_{\lambda}$ ,  $\lambda \in \text{spec}\Delta$ , which means that the Laplace-Fourier coefficients  $f_{\lambda,i}$ s must be transformed via  $\epsilon^{-1}$ . In fact, in (III.1) we must consider  $F_i$  to be block diagonal for the spectrum of the Laplacian operator  $\Delta^{g(t_i-1)}$  as:  $F_i = \oplus_{\lambda} (F_i)_{\lambda}$ , with;

$$(F_i)_{\lambda, mn} = \delta_{mn} + \frac{\tau_i \zeta}{2} \int_M R(t_i) \psi_{\lambda,m}(\tau_i) \psi_{\lambda,n}(\tau_i) d\Omega_{g(t_i)}, \quad (1 \leq i \leq N) \quad (III.8)$$

for  $m$  and  $n$  running through the degenerate eigenfunctions of  $\Delta^{g(t_i-1)}$  with common eigenvalue  $\lambda$ . Indeed, one can simply consider (III.8) as:  $F_i = 1 + r_i$ ,  $i = 1, \dots, N$ , where we easily gain;

$$\begin{aligned} J_i &= \det F_i = \exp \left( tr \left\{ \ln(1 + r_i) \right\} \right) \approx \exp \left( tr \{ r_i \} \right) \\ &= \exp \left( \sum_{\lambda \in \text{spec}\Delta^{g(t_i-1)}} \sum_{m \in D_{\lambda}} \frac{\tau_i \zeta}{2} \int_M R(t_i) \psi_{\lambda,m}^2(\tau_i) d\Omega_{g(t_i)} \right). \end{aligned} \quad (III.9)$$

<sup>14</sup> For:  $K = \frac{1}{D} \int_M \lim_{t \rightarrow b} R(t) d\Omega_{g_c} / \int_M d\Omega_{g_c}$ .

Actually, incorporating the full expression of the Jacobian determinant (III.9) in the Wiener fractal measure (III.1) leads to a highly complicated formulation for stochastic process of fractal functions on  $M$ . Therefore, we have to consider our first approximation of the Wiener fractal measure as:

**Approximation I:** *The trace operator of  $\sum_{\lambda \in \text{spec} \Delta^{g(t_{i-1})}} \sum_{m \in D_\lambda}$  in the exponent of  $J_i$  is in fact approximated by the dominating leading term  $\sum_{\lambda=0} \sum_{m \in D_0}$ . However, since  $M$  is connected we have  $\dim H_{dR}^1(M, \mathbb{R}) = 1$ , hence there is no degeneracy ( $\dim D_0 = 1$ ) and consequently the only considered eigenfunction is a constant. On the other hand, upon the normalization condition (III.5) we obtain  $\psi_0 = 1/\sqrt{V_i}$ , wherein  $V_i$  is the volume of  $M$  for metric  $g(t_i)$ ; i.e.  $V_i = \int_M d\Omega_{g(t_i)}$ . Therefore, we readily obtain:*

$$J_i = \exp \left( \frac{\tau_i \zeta}{2V_i} \int_M R(t_i) d\Omega_{g(t_i)} \right). \quad (\text{III.10})$$

The second approximation concerns about the volume of the space manifold  $M$ . According to (III.4) one can simply compute the derivation of this volume with respect to  $t$  as:

$$\frac{dV(t)}{dt} = -\zeta \int_M R(t) d\Omega_{g(t)}. \quad (\text{III.11})$$

Since  $\zeta \approx 0$  we could assume that within a specific period of time  $V(t)$  evolve slightly, hence is almost constant. For instance, for fixed  $t_0 \in [-T, T]$ , say the present age of the universe, we compute:

$$V_i = V - \zeta \int_{\mathcal{M}_{t_0}^{t_i}} R(t) dt \wedge d\Omega_{g(t)} \approx V - (t - t_0) \zeta \int_M R(t_0) d\Omega_{g(t_0)}, \quad (\text{III.12})$$

with  $V = V(t_0)$ , and  $\mathcal{M}_{t_0}^{t_i} = [t_0, t_i] \times M$  for  $t_i > t_0$  (resp.  $\mathcal{M}_{t_0}^{t_i} = [t_i, t_0] \times M$  for  $t_i < t_0$ ). Thus;

$$\frac{1}{V_i} \approx \frac{1}{V} + \frac{(t - t_0) \zeta}{V^2} \int_M R(t_0) d\Omega_{g(t_0)} + \mathcal{O}(\zeta^2/V^3) \approx \frac{1}{V}. \quad (\text{III.13})$$

Hence, the next approximation is reasonable.<sup>15</sup>

**Approximation II:** *We may also assume that the variation of  $V_i$  along the Ricci flow is negligible during the period of time in which we concern the Brownian stochastic evolution of quantum fields. Hence, we would simply consider  $V_i = V$ ,  $1 \leq i \leq N$ , and write the Jacobian determinant (III.10) as:*

$$J_i = \exp \left( \frac{\tau_i \zeta}{2V} \int_M R(t_i) d\Omega_{g(t_i)} \right). \quad (\text{III.14})$$

In particular, upon approximations **I** and **II** one can easily extract the contribution of the Jacobian determinants  $J_i$ s,  $1 \leq i \leq N$ , within the Wiener fractal measure (III.1) as:

$$\prod_{i=1}^N J_i = \exp \left( \frac{1}{2\xi} \sum \tau_i \int_M R(t_i) d\Omega_{g(t_i)} \right), \quad (\text{III.15})$$

<sup>15</sup> Indeed, as we see in the following the space volume  $V$  is effectively equivalent to the Newton's gravitation constant  $G$ . In principle, the dynamics of the space volume is formulated with a double integral on  $M \times M$ , hence is due to a non-local contribution of the scalar curvature of  $M$  within the Einstein-Hilbert action. However, the slight variation of  $G$  has already been formulated by means of a local interacting scalar field theory (i.e. dilaton) via the Brans-Dicke theory [7]. See also [87] for more discussions.

for  $\xi = V/\zeta$ , which in the limit of  $N \rightarrow \infty$  (i.e.  $\tau_i = 2T/N \rightarrow 0$ ) leads to:

$$\prod_{i=1}^N J_i = \exp \left( \frac{1}{2\xi c} \int_{\mathcal{M}} R d\Omega \right), \quad (\text{III.16})$$

wherein  $\mathcal{M} = [-T, T] \times M$  is the space-time continuum introduced above, and  $d\Omega = cdt \wedge d\Omega_{g(t)}$  is the Riemannian volume form for the Lorentzian metric  $\mathbf{g} = \mathbf{g}_{\mu\nu} dx^\mu \otimes dx^\nu = c^2 dt \otimes dt \oplus (-g_{ij}(t) dx^i \otimes dx^j)$  on  $\mathcal{M}$ . We should note that according to the rescaled Ricci flow (III.3) the scalar curvature  $R$  of  $g$  is in essential equal to the scalar curvature  $\mathcal{R}$  of  $\mathbf{g}$  up to an overall minus sign and some negligible terms of order  $\mathcal{O}(\zeta)$ . In fact, upon to (III.3) we see:

$$\mathcal{R} = -R + \frac{\zeta}{c^2} \left\{ g^{ij} \frac{\partial Ric_{ij}}{\partial t} + \frac{\partial R}{\partial t} \right\} - \frac{\zeta^2}{c^2} Ric_{ij} Ric^{ij} + \frac{\zeta^2}{c^2} R^2 = -R + \mathcal{O}(\zeta). \quad (\text{III.17})$$

Here, we may set our third approximation law:

**Approximation III:** *We would simply replace  $R$  by  $-\mathcal{R}$  in the Jacobian determinant  $J_i$  and ignore the remaining terms of order  $\mathcal{O}(\zeta)$ , i.e.:*

$$\prod_{i=1}^N J_i = \exp \left( -\frac{1}{2\xi c} \int_{\mathcal{M}} \mathcal{R} d\Omega \right). \quad (\text{III.18})$$

In principle, the above three approximations would provide a sufficient set of assumptions to work out the original formulation of general relativity via the Einstein-Hilbert action. Actually, in a local coordinate system on the space-time continuum  $\mathcal{M}$ , say  $(t, x^i)$ , we would obtain the following formula for the Jacobian term (III.18):

$$\prod_{i=1}^N J_i = \exp \left( -\frac{1}{2\xi c} \int_{\mathcal{M}} \sqrt{|\det \mathbf{g}|} \mathcal{R} dt \wedge dx^1 \wedge \dots \wedge dx^D \right). \quad (\text{III.19})$$

Hence, the Wiener stochastic process of fractals on Riemannian manifold  $M$  along with the Ricci flow automatically involves the Einstein-Hilbert action of general relativity within the approximations **I**, **II**, and **III**. In fact, the emergence of general relativity is merely due to the evolution of the cosmos geometry, hence having nothing to do with possible background interactions of the included matter fields. However, one should note that the analytic structure of the spatial metric  $g_{ij}$  fulfills Einstein's field equation via the Euler-Lagrange equations. Actually,  $\mathbf{g}_{\mu\nu}$  must obey the Ricci flow as an intrinsic property of the space-time geometry.

Following the machinery of the previous section, it can be easily seen that the rest parts of the Wiener fractal measure on the dynamical Riemannian manifold  $(M, g)$  would be:

$$\exp \left( -\frac{1}{\hbar} \int_{\mathcal{M}} \frac{1}{2} \left\{ \mathbf{g}^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{m^2 c^2}{\hbar^2} \phi^2 \right\} d\Omega \right) \mathfrak{D}\phi. \quad (\text{III.20})$$

wherein the strategy **A** is considered due. One should note that there is an essential difference between (II.14) and (III.20); In the former, the space-time metric  $\mathbf{g}$  is a static tensor, but for the latter it evolves via the Ricci flow. Moreover, in the latter the Feynman measure  $\mathfrak{D}\phi$  is given in terms of the Fourier expansion of the Laplacian operator  $\Delta^{g(t_{i-1})}$  at the  $i$ -th time section, while in the former the Feynman

measure is static. All in all, the Wiener fractal measure along with the Ricci flow and after the mentioned three approximations becomes:

$$dW(t_N, \dots, t_1) \approx \exp \left( -\frac{1}{\hbar} \int_{\mathcal{M}} \left\{ \frac{1}{2\kappa} \sqrt{|\det \mathbf{g}|} \mathcal{R} + \mathcal{L}_{K-G}(\phi) \right\} dx^{D+1} \right) \mathfrak{D}\phi, \quad (\text{III.21})$$

with  $\mathcal{L}_{K-G}(\phi) = \frac{\sqrt{|\det \mathbf{g}|}}{2} \{ \mathbf{g}^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{m^2 c^2}{\hbar^2} \phi^2 \}$ , the Klein-Gordon Lagrangian density, and  $\kappa = \xi c / \hbar$ . Hence, we readily obtain the Newton's gravitational constant in terms of fractal, geometric and physical parameters in  $D = 3$  dimensions:  $G = \xi c^4 / 8\pi \hbar$ . Therefore, if we use the standard values of  $G$ ,  $\hbar$  and  $c$  we obtain:  $\xi = V / \zeta = 2.1899700 \times 10^{-77}$  (m.s), which shows that the *cosmos entropic volume* (i.e. the cosmos volume for the metric involved in the Ricci flow) is extremely small.

One should note that this is not a problem at all, since the small amount of the cosmos entropic volume  $V$  is, in fact, equivalent to the large value of the Einstein-Hilbert coefficient (i.e.  $1/2\kappa = c^4 / 16\pi G$ ) in the SI units. Actually, the small amount of  $V$  stems from the smallness of the space metric entries  $g_{ij}$ s, and hence the huge amplitude of the scalar curvature  $\mathcal{R}$ . Hence, if we rescale the entropic metric  $g_{ij}$  with some large constant coefficient  $\zeta$  to gain an enormously large universe we readily obtain the standard form of Einstein-Hilbert action with a large coefficient as we expected before. In fact, in this case we have:  $g_{ij} \rightarrow \zeta g_{ij}$ ,  $\mathcal{R} \rightarrow \zeta^{-1} \mathcal{R}$ , and  $d\Omega \rightarrow \zeta^{D/2} d\Omega$ , and thus the Einstein-Hilbert action will be multiplied by  $\sqrt{\zeta}$  in  $D = 3$  dimensions. In particular, the extremely small amount of the cosmos entropic volume is an immediate consequence of the Ricci flow and could be regarded as the main reason of the hierarchy problem.<sup>16</sup>

To work out a well-defined theory from Wiener fractal measure (III.21) we must employ the strategy **B** once again. We readily obtain:

$$dW(t_N, \dots, t_1) \approx \exp \left( \frac{i}{\hbar} \int_{\mathcal{M}} \left\{ \frac{1}{2\kappa} \sqrt{|\det \mathbf{g}|} \mathcal{R} + \mathcal{L}_{K-G}(\phi) + \sqrt{|\det \mathbf{g}|} \frac{i\varepsilon}{2} \phi^2 \right\} dx^{D+1} \right) \mathfrak{D}\phi, \quad (\text{III.22})$$

for  $\varepsilon \rightarrow 0$ . Therefore, one can consider the Wiener fractal process (III.1) for the Lebesgue fractal measure (II.2) through with the Ricci flow as the more fundamental formulation of quantum field theory in dynamical curved spacetime, i.e. in presence of semi-classical gravity. Hence, we would claim that the basic source of the Einstein's theory of gravity stems from the dynamical Wiener fractal measure via the Ricci flow. We refer to this stochastic derivation of gravitational effect as the *Wiener fractal gravity*. As we discussed in last section, although including non-local terms the Wiener fractal gravity leads to finite predictable expectation values

$$\lim_{N, N \rightarrow \infty} \int_{C_N} F^{(N)}(t_N, \dots, t_1) \{ \phi(t_{i_1}) \cdots \phi(t_{i_n}) \} dW(t_N, \dots, t_1) \quad (\text{III.23})$$

for  $F^{(N)}(t_N, \dots, t_1) = \prod_{k=1}^N \exp \left( \tau_k \int_M V^{(N)}(\phi(t_{k-1})) d\Omega_{g(t_{k-1})} \right)$  with negative (bounded from above) and renormlizable interaction terms  $V^{(N)}(\phi(t_i))$ . This ensures the implication of the Wiener fractal gravity as a more fundamental formulation of quantum field theory in presence of semi-classical gravity. Through the next two sections, we try to study the theoretical and empirical properties of the Wiener fractal gravity.

<sup>16</sup> See [34] and the references therein for interesting discussions about the relation of hierarchy problem and the Bayesian statistics, which could be related to our achievements in this paper.

#### IV. RICCI FLOW, COSMOLOGICAL CONSTANT AND $\Lambda$ CDM MODEL

Upon the Hamilton's celebrated theorem we may consider the metric of the universe to be almost Einstein with  $Ric_{ij}(t) = Kg_{ij}(t) + \varepsilon_{ij}$ , for infinitesimal  $\varepsilon_{ij} \ll Kg_{ij}$ , and  $t > t_0$ , where  $t_0$  is considered as the current age of the universe.<sup>17</sup> With this assumption, the Ricci flow is;

$$\frac{\partial g_{ij}(t)}{\partial t} = -2\zeta K g_{ij}(t) - 2\zeta \varepsilon_{ij}, \quad t > t_0. \quad (\text{IV.1})$$

which has a unique solution as:<sup>18</sup>

$$g_{ij}(t) = e^{-2\zeta K(t-t_0)} g_{ij}(t_0) - \frac{\varepsilon_{ij}}{K} \quad \text{and}; \quad Ric_{ij}(t) = K e^{-2\zeta k(t-t_0)} g_{ij}(t_0). \quad (\text{IV.2})$$

Hence, the higher order terms of (III.17) would become;

$$\begin{aligned} & \frac{\zeta}{c^2} \left\{ g^{ij} \frac{\partial Ric_{ij}}{\partial t} + \frac{\partial R}{\partial t} \right\} - \frac{\zeta^2}{c^2} Ric_{ij} Ric^{ij} + \frac{\zeta^2}{c^2} R^2 \\ & = (D-3) D c^{-2} \zeta^2 K^2 + 2(D-3) \zeta^2 c^{-2} K e^{-2\zeta k(t-t_0)} tr\{\varepsilon g_0\} + 2\Lambda, \end{aligned} \quad (\text{IV.3})$$

for  $g_0 = g(t_0)$ , and the cosmological constant  $\Lambda$ ;<sup>19</sup>

$$\Lambda = \frac{1}{2} \zeta^2 c^{-2} e^{-4\zeta K(t-t_0)} \left\{ (tr\{\varepsilon g_0\})^2 - tr\{(\varepsilon g_0)^2\} \right\}. \quad (\text{IV.4})$$

Actually, upon Hamilton's theorem [43], the deviation matrix  $\varepsilon$  is positive definite for  $D = 3$  dimensions, hence the cosmological constant  $\Lambda$  is a positive number on closed 3-manifolds. On the other hand, we see surprisingly that in  $D = 3$  dimensions the constant component and the  $\varepsilon$ -linear term of (IV.3) vanish and we are left with only the cosmological constant  $\Lambda \propto e^{-4\zeta K t} \zeta^2 O(\varepsilon^2)$  which is in essential an extremely small amplitude. Thus, on three-dimensional space  $M$  we have:  $R = \mathcal{R} - 2\Lambda$ , and consequently the cosmological term  $-2\Lambda$  is simply inserted beside the scalar curvature  $\mathcal{R}$  within the Einstein-Hilbert Lagrangian density:

$$\prod_{i=1}^N J_i = \exp \left( -\frac{1}{2\xi c} \int_{\mathcal{M}} \sqrt{|\det \mathbf{g}|} (\mathcal{R} - 2\Lambda) dt \wedge dx^1 \wedge \dots \wedge dx^D \right). \quad (\text{IV.5})$$

Therefore, according to the above calculations the cosmological constant emerges only upon the metric evolution of the space manifold, hence independent from the matter fields and the zero-point energy fluctuations of quantum field theory. In principle, the cosmological constant  $\Lambda$  has an entropic source due to the Ricci flow of the universe three-dimensional geometry, but not an inevitable consequence of matter fields. Therefore, it seems that the Wiener fractal gravity would give a reliable consistent answer to the cosmological constant problem [50, 88] and the nature of dark energy.<sup>20</sup>

<sup>17</sup> In fact, according to experimental achievements  $K$  has a very tiny amplitude; i.e.  $K \approx 0$ . Indeed, the experimental data from different laboratories (for example due to WMAP, BOOMERanG, and Planck) confirm that the universe manifold  $M$  is flat with about 0.4 percent margin of deviation [95]. See also [16, 63].

<sup>18</sup> We may temporary assume that  $\varepsilon_{ij}$  is constant. Indeed, we suppose that  $\varepsilon_{ij}$  varies slightly through the cosmological time  $t$ .

<sup>19</sup> Actually,  $\Lambda$  is not a constant in general, but upon to Hamilton's theorem [43] for some appropriate interval of time it evolve slightly on  $M$ , hence could be considered as a constant at the first order of approximation.

<sup>20</sup> See [9] for another entropic viewpoint to the cosmological constant via the holographic approach in quantum gravity.

Moreover, for almost the canonical asymptotic (limit) Einstein metric  $g_c$  of the Ricci flow, i.e.  $Ric_{ij} \approx K g_{cij}$ , we readily obtain an inflating universe  $M$ . To see this one has to employ the Jacobi field equation for Einstein metric  $g_c$ . Let  $\gamma_s(t)$  be a continuous family of geodesics on  $\mathcal{M}$  labeled by  $s \in \mathbb{R}$ , each of which with initial condition  $\dot{\gamma}_s(0) = \partial/\partial t$ . Then, according to the Jacobi field equation we read for small enough  $t$ :

$$\frac{D^2}{dt^2} J(t) = \mathcal{R}(\dot{\gamma}_s(t), J(t)) \dot{\gamma}_s(t) \approx \mathcal{R}_{tit}{}^j J^i(t) \frac{\partial}{\partial x^j}, \quad (\text{IV.6})$$

wherein  $J(t) = \frac{\partial \gamma_s(t)}{\partial s}$  and  $D/dt$  is the covariant derivative along  $t$ , while  $\mathcal{R}_{\mu\nu\sigma}{}^\lambda$  is the space-time Riemannian curvature computed for the metric  $\mathbf{g} = c^2 dt \otimes dt \oplus (-g_c)$ .<sup>21</sup> Let us compute the relative acceleration of inertial celestial bodies due to the space-time metric  $\mathbf{g}$ . We simply set  $J(0) = r \frac{\partial}{\partial x^i}$ , assuming the observed galaxies are located along the  $i$ -th direction of the local coordinate system  $(x^\mu) = (x^0 = t, x^j = x^1, \dots, x^D)$ . We compute the relative acceleration as:<sup>22</sup>

$$a_r = a_r^\mu \frac{\partial}{\partial x^\mu} = \mathcal{R}_{0i0}{}^\mu \frac{\partial}{\partial x^\mu} = r \left\{ \frac{\partial}{\partial t} \Gamma_{ti}^\mu - \frac{\partial}{\partial x^i} \Gamma_{tt}^\mu + \Gamma_{t\nu}^\mu \Gamma_{it}^\nu - \Gamma_{i\nu}^\mu \Gamma_{tt}^\nu \right\} \frac{\partial}{\partial x^\mu} = \zeta^2 K^2 r \frac{\partial}{\partial x^i}, \quad (\text{IV.7})$$

which is in fact equal to:<sup>23</sup>

$$\ddot{r}(t) = H_\infty^2 r(t), \quad t > t', \quad (\text{IV.8})$$

for radial distance  $r(t)$  and *ultimate Hubble's constant*  $H_\infty = \zeta|K|$ .<sup>24</sup> In principal, (IV.8) has a unique dominating solution for  $t \rightarrow \infty$ :

$$r(t) = r_0 e^{H_\infty(t-t')} \quad \rightarrow \quad \dot{r}(t) = H_\infty r(t). \quad (\text{IV.9})$$

These conclusions could be slightly modified via the assumption of (IV.1) for the present age of the universe  $t_0$  and the upcoming future of the cosmological time  $t > t_0$ . In this case we obtain the accurate form of (IV.7) as:<sup>25</sup>

$$a_r = a_r^\mu \frac{\partial}{\partial x^\mu} = \zeta^2 K^2 r \frac{\partial}{\partial x^i} + 2\zeta^2 \varepsilon_{jm} \varepsilon_{mi} r \frac{\partial}{\partial x^j} + \zeta^2 e^{4K\zeta(t-t_0)} A_{0k}^j A_{0i}^k r \frac{\partial}{\partial x^j} + \mathcal{A}_i^j(\dot{\varepsilon}) r \frac{\partial}{\partial x^j} + \mathcal{O}(\varepsilon^3), \quad (\text{IV.10})$$

with  $A_{0i}^j = g_0^{jm} \varepsilon_{mi}$  and

$$\mathcal{A}_i^j(\dot{\varepsilon}) = -\zeta e^{2K\zeta(t-t_0)} g_0^{jm} \dot{\varepsilon}_{mi} + \frac{1}{K} (\dot{\varepsilon}_{jm} \varepsilon_{mi} + \varepsilon_{jm} \dot{\varepsilon}_{mi}). \quad (\text{IV.11})$$

Indeed, the dominating diagonal term of (IV.11) (for  $\varepsilon_{ij} \approx \varepsilon \delta_{ij}$  and  $g_0^{ij} \approx \delta^{ij}$ ) is:

$$a_r = a_r^\mu \frac{\partial}{\partial x^\mu} = r \left\{ \zeta^2 K^2 + \zeta^2 \varepsilon^2 \left( 2 + D e^{4K\zeta(t-t_0)} \right) - \zeta \dot{\varepsilon} \left( e^{2K\zeta(t-t_0)} + \frac{2\varepsilon}{K} \right) \right\} \frac{\partial}{\partial x^i}, \quad (\text{IV.12})$$

<sup>21</sup> Here, we assume for the coordinate index  $\mu$  to vary among  $\{t, 1, \dots, D\}$ .

<sup>22</sup> Upon the Ricci flow (III.3) we have:  $\Gamma_{ij}^0 = \zeta Ric_{ij}$ ,  $\Gamma_{0j}^i = \Gamma_{j0}^i = -\zeta g_c^{im} Ric_{mj}$ , and  $\Gamma_{0i}^0 = \Gamma_{i0}^0 = \Gamma_{00}^i = \Gamma_{00}^0 = 0$ .

<sup>23</sup> Here we assume  $t'$  to be close enough to the ultimate time for the Ricci flow to reach to the Einstein's metric  $g_c$  up to the extent of the experimental and observational accuracy.

<sup>24</sup> Although, according to the Hamilton's theorem [43]  $K$  is considered to be positive, here we prefer to employ its absolute value to involve Einstein manifolds with negative curvature as is considered in the FLRW metric.

<sup>25</sup> As we see once again the  $\varepsilon$ -linear term cancels out automatically.

which leads to a time-dependent Hubble's parameter;

$$H(t) = H_\infty \sqrt{1 + \frac{\varepsilon^2}{K^2} (2 + D e^{4K\zeta(t-t_0)}) + \frac{|\dot{\varepsilon}|}{\zeta K^2} \left( e^{2K\zeta(t-t_0)} + \frac{2\varepsilon}{K} \right)}, \quad (\text{IV.13})$$

since upon the Hamilton's theorem  $\dot{\varepsilon} < 0$  [43]. Set

$$\Omega_1(t) = \frac{\varepsilon^2}{K^2} \left( 2 + D e^{4K\zeta(t-t_0)} \right), \quad \Omega_2(t) = \frac{|\dot{\varepsilon}|}{\zeta K^2} \left( e^{2K\zeta(t-t_0)} + \frac{2\varepsilon}{K} \right). \quad (\text{IV.14})$$

Thus, we gain approximately:<sup>26</sup>

$$H(t) = H_0 \sqrt{1 + \Delta\Omega_1(t) + \Delta\Omega_2(t)}, \quad (\text{IV.15})$$

for  $\Delta\Omega_i(t) = \Omega_i(t) - \Omega_i(t_0)$ ,  $i = 1, 2$ , and the Hubble's constant  $H_0$ , given as;

$$H_0 = H_\infty \sqrt{1 + \Omega_1(t_0) + \Omega_2(t_0)}. \quad (\text{IV.16})$$

All in all, we consequently find out some major results:

**a)** *The universe is expanding radially, hence radial velocities must be observed among extra-galactic nebulae and the amplitudes of these velocities must be proportional to distances from the fiducial observer, just as Hubble pointed out in his seminal 1929 paper [56].*

**b)** *The acceleration is positive, hence galaxies flee from each other and the universe is inflating with an increasing rate of;*<sup>27</sup>

$$\frac{\dot{a}}{a} = H_0 \sqrt{1 + \Delta\Omega_1(t) + \Delta\Omega_2(t)}, \quad (\text{IV.17})$$

where  $a(t) = (1 + z)^{-1}$  is the scale factor for observed redshift  $z$ .<sup>28</sup>

**c)** *According to the arguments of the next section for similarities between the second law of thermodynamics and Wiener fractal gravity, hereby we conclude that the dark energy (and the negative pressure due) stems from an entropic source due to the Ricci flow.*<sup>29</sup>

Indeed, the fundamental assumptions of Wiener fractal gravity resembles the well-understood Friedmann-Lamitre-Robertson-Walker solution (besides the assumed properties of homogeneity and isotropy), for which the Einstein's field equations are only mandatory to work out the scale factor of the universe metric  $a(t)$  as a function of cosmological time  $t$ . This provides more common results between Wiener fractal gravity and the  $\Lambda$ CDM model. In particular, for the space metric of the FLRW

<sup>26</sup> We are working in a period of time  $[t_0, T]$  in which  $\Omega_i(t_0)\Delta\Omega_j(t) \ll \Omega_k(t)$ , with  $t_0 \leq t \leq T$ , for  $i, j, k = 1, 2$ .

<sup>27</sup> One should note that the inflation of the universe does not mean that the volume  $V(t)$  of  $M$  is increasing along the Ricci flow. Indeed, according to (III.4) the volume of  $M$  for Einstein metric  $g_c$  changes with the rate of  $dV(t)/dt = -\zeta KDV(t)$ . Hence, for positive  $K$  the volume is decreasing through the Ricci flow.

<sup>28</sup> See [70] as an interesting research which shows that the root-mean-square distance of two density volume terms  $\omega_1(t) \geq 0$  and  $\omega_2(t) \geq 0$  evolving with the heat (diffusion) equation  $\frac{\partial}{\partial t}\omega_i = \Delta^{g(t)}\omega_i$ ,  $i = 1, 2$ , is non-increasing along the backward Ricci flow  $\frac{\partial}{\partial t}g_{ij} = 2Ric_{ij}$ . Indeed, this would lead to an expansion theorem along the Ricci flow for density fields via transformations  $t \rightarrow -t$ . We regard this conclusion as an entropic force, which obviously include the ordinary pressure in thermodynamics. This result, in principle, establishes the entropic root of the Ricci flow in its own turn.

<sup>29</sup> See [9, 12] for more discussions about the entropic source of inflation.

solution we have:<sup>30</sup>  $Ric_{ij} = -2kg_{ij}$ ; hence  $g_{ij}$  is an Einstein metric which is the promised limit metric  $g_{cij}$  of the Ricci flow (for  $K = -2k$ ):

$$Ric_{ij} = Kg_{ij} \quad (IV.18)$$

Thus, upon the above arguments the FLRW metric would describe the universe with conclusions in an admissible agreement with the Wiener fractal gravity.

However, if one readily imposes the Ricci flow on the FLRW metric we simply obtain a first order differential equation (for  $k = 1$ ) for the scale factor  $a(t)$  as;  $\dot{a}(t) = 2\zeta a(t) = H_\infty a(t)$ , with the immediate solution  $a(t) = e^{H_\infty(t-t_0)}$ , which is in agreement with (IV.9). Nevertheless, upon the Hamilton's theorem for 3-manifolds [43], the Einstein metric is an ultimate (limit) solution of the Ricci flow, hence the FLRW metric is indeed an approximate solution for the Wiener fractal gravity. Anyway, when the Ricci flow is employed this approximation leads to simple equalities of  $\frac{\dot{a}}{a} = H_\infty$  and  $\frac{\ddot{a}}{a} = H_\infty^2$ , hence the Friedmann equations turn to:

$$H_\infty^2 + \frac{kc^2}{a^2} - \frac{\Lambda c^2}{3} = \frac{8\pi G}{3}\rho, \quad \text{and}; \quad H_\infty^2 + \frac{kc^2}{3a^2} - \frac{\Lambda c^2}{3} = -\frac{8\pi G}{3c^2}p, \quad (IV.19)$$

which result in an immediate approximate equality;

$$\rho + \frac{p}{c^2} = \frac{kc^2}{4\pi G}e^{-2H_\infty(t-t_0)}. \quad (IV.20)$$

Combining (IV.20) with<sup>31</sup>

$$\rho + \frac{3p}{c^2} = \frac{\Lambda c^2}{4\pi G} - \frac{3H_\infty^2}{4\pi G} \quad (IV.21)$$

leads to explicit albeit approximate solutions for the density  $\rho(t)$  and the pressure  $p(t)$  as:

$$\begin{aligned} \rho(t) &= \frac{1}{8\pi G} \left\{ 3H_\infty^2 + c^2 \left( 3ke^{-2H_\infty(t-t_0)} - \Lambda \right) \right\}, \\ p(t) &= \frac{c^2}{8\pi G} \left\{ c^2 \left( \Lambda - ke^{-2H_\infty(t-t_0)} \right) - 3H_\infty^2 \right\}. \end{aligned} \quad (IV.22)$$

Before closing this section we prefer to return to (IV.15) to have a brief look at the Hubble's parameter  $H(t)$ . It is well-known that the Friedmann equation is rewritten in terms of the density parameters as

$$H(t) = H_0 \sqrt{\Omega_{mat}a^{-3} + \Omega_{rad}a^{-4} + \Omega_k a^{-2} + \Omega_\Lambda a^{-3(1+\omega)}}, \quad (IV.23)$$

wherein  $\Omega_{mat}$  is the matter density including baryons and the cold dark matter,  $\Omega_{rad}$  is for radiation including neutrinos,<sup>32</sup>  $\Omega_k$  is due to the curvature and  $\Omega_\Lambda$  is the contribution of dark energy with the equation of state parameter  $\omega$ . To impose the Ricci flow on Friedmann equations for positive curvature  $k$  (i.e.  $K < 0$  in (IV.18)), one simply reads from (IV.14);  $\Delta\Omega_1 \propto \varepsilon^2 a^{-4}$  and  $\Delta\Omega_2 \propto \dot{\varepsilon} a^{-2}$ . We should note that the matter (including the cold dark matter) and the radiation contributions are encoded in  $\varepsilon$  due to Einstein's field equation. Therefore, the above argument leads to an agreement between the  $\Lambda$ CDM model formula (IV.23) and the equation (IV.15) which we obtained for the Wiener fractal gravity.

<sup>30</sup> For example one can read the equality for the FLRW spatial metric in reduced-circumference polar coordinates  $g_{ij} = a^2(t) \left( \frac{dr^2}{1-kr^2} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \right)$ .

<sup>31</sup> This equation is derived from the Ricci flow and Friedmann's equation:  $\frac{\ddot{a}}{a} = -\frac{4\pi G}{3} \left( \rho + \frac{3p}{c^2} \right) + \frac{\Lambda c^2}{3}$ .

<sup>32</sup> Actually, the radiation has a negligible amplitude [35]. See also [74] as a nice presentation of the issue.

## V. DISCUSSION: RICCI FLOW, THERMODYNAMICS AND GRAVITY

According to the Perelman's seminal paper [75] the Ricci flow is an entropic based differential equation. In fact, employing an appropriate diffeomorphism  $\varphi : M \rightarrow M$ , it can be seen that the Ricci flow is the gradient of Perelman's  $\mathcal{F}$ -entropy (i.e.  $\mathcal{F}(g_{ij}(t), f(t)) = \int_M (R + |\nabla f|^2) e^{-f} d\Omega_{g(t)}$ ) via a volume preserving variation of the metric  $g_{ij}(t)$ .<sup>33</sup> The entropy functional  $\mathcal{F}$  has a natural interpretation in terms of Bochner-Lichnerovicz formulas which we will use to formulate the Yang-Mills gauge theories within the Wiener fractal measure in an independent paper [80]. However, from the theoretical physics insights,  $\mathcal{F}$  and its first variation describes the low energy effective action in the string theory, wherein  $f$  is called dilaton field.<sup>34</sup>

On the other hand, Perelman's generalized entropy functional, the so called Perelman's  $\mathcal{W}$ -entropy (i.e.  $\mathcal{W}(g_{ij}(t), f(t), \tau) = \frac{1}{(4\pi\tau)^{n/2}} \int_M (\tau (R + |\nabla f|^2) + f - D) e^{-f} d\Omega_{g(t)}$ ) has a specific interpretation as the entropy of a canonical ensemble in statistical mechanics [75].<sup>35</sup> Thus, it seems that considering the Ricci flow for the evolution of the space geometry stem from an entropic source, which give rise to the reminiscence of thermodynamic interpretation of gravity. In principle, now a day there is no doubt that classical (and semi-classical) general relativity is intimately correlated to thermodynamics; there are already enough theoretical evidences which lead to a thermodynamical interpretation of the Einstein's theory of gravity [60, 73, 83].<sup>36</sup> Moreover, as we explained above, the Wiener fractal measure is based on the Brownian motion which is in its turn subject to an entropic force [72].

Thus, one may suspect that the Wiener fractal gravity has a lot of properties in common with Verlinde's entropic gravity [81, 82]. However, there is a profound distinction between these two approaches to gravitational force. In particular, the entropic source of the Wiener fractal gravity is due to the evolution of the space geometry via the Ricci flow,<sup>37</sup> regardless of the existence of matter fields involved in the space-time continuum  $\mathcal{M}$ , whereas the entropic source of the Verlinde's theory is solely due to the (quantum) matter fields interactions. Actually, the latter is based on a combination of Jacobson's thermodynamical viewpoint to gravity with 't Hooft's holographic principle and interpret gravity as a consequence of the information associated with the positions of material bodies. Hence, Verlinde's theory claims that gravity is not in fact a fundamental interaction, but an emergent phenomenon.

On the other hand, as we established by working out the Wiener fractal measure for evolving metric  $g_{ij}(t)$ , the Einstein-Hilbert action has an entropic origin. Nevertheless, this entropic source is independent of the interaction of matter fields and the information of material positions. Actually, although the space metric  $g_{ij}$  (or the space-time metric  $\mathbf{g}_{\mu\nu}$ ) fits with the distribution of matter fields and their interactions via the semi-classical solution of Einstein's field equation, its time evolution has an entropic essence due to the Ricci flow. Therefore, despite their seemingly common interpretation

<sup>33</sup> For more discussion about the appropriate diffeomorphism see [19] and [46].

<sup>34</sup> See for example [78], where the author shows that the generalization of the Perelman's  $\mathcal{F}$ -entropy to all loop orders equals to minus the central charge at the fixed points, in agreement with the general claim of the Zamolodchikov's c-theorem.

<sup>35</sup> Perelman's  $\mathcal{W}$ -entropy also describes the RG flow in quantum field theories. See also [64, 65] for more discussions due.

<sup>36</sup> The correlation between thermodynamics and general relativity was first put forward by Bekenstein and Hawking via studying the black hole entropy [2, 48].

<sup>37</sup> Note that we preserve the topology of the space manifold  $M$  along the Ricci flow.

based on entropic origin, the Wiener fractal gravity is fundamentally different from Verlinde's theory of entropic gravity.

Moreover, including squared terms of time derivation, the Wiener fractal measure is essentially symmetric with respect to transformation  $t \rightarrow -t$ , but, however, imposed for evolving the geometry of the background manifold  $M$ , the Ricci flow is in fact asymmetric versus reversing the time direction, hence is the only *time's arrow-dependent* term (i.e. entropic term) included in the Wiener fractal measure. Therefore, one may readily regard the second law of thermodynamics (hence all entropic forces) as an immediate aspect of the Ricci flow of the cosmos geometry.

Actually, the most significant strategy to work out the Einstein-Hilbert action in the Wiener fractal measure is due to the approximation **I** which let us restrict the trace of the Ricci transformations  $F_i$ s in (III.9) to the lowest eigenvalue of the Laplacian.<sup>38</sup> In principle, the emergence of the Einstein-Hilbert action is a mere consequence of evolving the geometric structure on  $M$  and has nothing to do with the quantum field theory (i.e. the interaction terms and their low and high energy scenarios) involved in the space-time continuum  $\mathcal{M}$ . Hence, the only fundamental law which we assumed within the Wiener Brownian process is due to the Ricci flow of the universe. Therefore, as we emphasized above, and demonstrated in [79, 80], one may conclude that the second law of thermodynamics is a fundamental law of nature, either more fundamental than gravitation or coming with gravity from the same basic source.

The stochastic base of the Wiener fractal gravity puts forward the idea of the existence of an intimate correlation between our path-integral formulation of general relativity and the theory of stochastic gravity [54, 55]. In fact, stochastic gravity stands on the same footing as the Wiener fractal gravity. Actually, in stochastic gravity, the semi-classical solutions are effectively modified by including background noise from quantum fields via the Langevin equation. For instance, the familiar moving charge quantum field system in the framework of stochastic gravity leads to the so called Abraham-Lorentz-Dirac-Langevine equation (a generalized version of ALD equation) which describes the stochastic dynamics of the moving charge within a correct and pathology-free setting [55].

Historically, stochastic (semi-classical) gravity commenced from the 1990s as the third step of a theoretical evolution of quantum gravity which its two first development stages was *quantum field theory in curved spacetime* (accomplished in the 1970s) and *semi-classical gravity* (established in the 1980s). Actually, the best introduction to stochastic gravity is found in [55] where the authors explain: *"While semi-classical gravity is based on the semi-classical Einstein equation with the source given by the expectation value of the stress-energy tensor of quantum fields, stochastic semi-classical gravity includes also its fluctuations in a new stochastic semi-classical Einstein-Langevin equation. If the centerpiece of semi-classical gravity is the vacuum expectation value of the stress-energy tensor of a quantum field, the centerpiece in stochastic semi-classical gravity is the symmetrized stress-energy bi-tensor and its expectation value known as the noise kernel. The mathematical properties of this quantity, its physical contents in relation to the behavior of fluctuations of quantum fields in curved spacetimes and their backreaction in the spacetime dynamics engendering induced metric fluctuations are the main focus of this theory."*

Indeed, stochastic gravity could be simply regarded as a rigorous mathematical technique to work

---

<sup>38</sup> As we show in [80], this is the main idea to obtain the same result for Yang-Mills fields in presence of gravity.

out the most accurate solution of Einstein's field equation extracted from the approximated Wiener fractal gravity. In fact, as we established above, according to the original form of the Wiener fractal measure subject to the dynamics of space geometry due to the Ricci flow, the quantum field fluctuations are basically non-local effects, admitting no definite Lagrangian density, whereas the gravitational features come from an explicit action which at the first approximation leads to the celebrated Einstein-Hilbert theory. In other words, stochastic gravity is in principle one of the best understood frameworks to work out the approximated expectation values of the Wiener fractal gravity as (III.23). However, computing these expectation values could be accomplished with some parallel methods of real analysis for static space-time and the ordinary Lebesgue measure [30]. Therefore, another parallel effort besides stochastic gravity to work out (III.23) is to develop the present analytical methods to dynamical space-time and the Lebesgue fractal measure.

The substantial distinguishing features between space and time dimensions and considering the space-time continuum as a product manifold  $\mathcal{M} = [-T, T] \times M$  (or considering the space manifolds as the foliation leaves of  $\mathcal{M}$ ), may also resemble the main ideas of Horava-Lifshitz gravity [52]. However, we should stress that this similarity only includes the fundamental viewpoints about the time and spatial dimensions and their distinction in physical equations.<sup>39</sup> In particular, there is a vast difference between Horava's theory and the Wiener fractal gravity in their basic technology strategies and the mathematics of formulation, while, however, it is apparent that the Wiener fractal gravity does not contain quantum fluctuations of the metric components in its present form, hence still being a semi-classical theory.

Actually, Horava's theory respects some significant admitted properties for quantum field theory such as the necessity for the Lagrangian density to only consist of local terms and to maintain the renormalizability of the theory by power-counting at the Lifshitz point  $z = 3$ , but the Wiener fractal gravity is a highly non-local theory at high energy levels which does not even admit a definite form of Lagrangian density in its original expression. Although both try to remove the problem of obtaining a consistent theory of gravity within the path-integral formulation, and also both address the problem to breaking the (local) Lorentz symmetry and general diffeomorphism invariance of the theory, but the Horava's strategy to overcome the subtlety is somehow active. He breaks the Lorentz invariance (and diffeomorphism invariance) by hand, puts forward the fixed Lifshitz point  $z = 3$  according to some evidence coming from condensed matter, and introduces and incorporates some foliation-preserving diffeomorphism invariant terms for bringing gravity into the quantum field theory in some renormalizable manner.

On the other hand, in Wiener fractal gravity we behave as passive. We do not impose any preferred geometric structure (such as some definite symmetries) and theoretical constraint into the theory. We only assume that the quantum fields are mostly distributed over the space  $M$  by fractal structures and their propagation along time is fundamentally a stochastic process that is partly subject to some dominating entropic force due to the Ricci flow. Therefore, the most geometric structures and theoretical constraints of the theory emerge naturally from the essential properties of the Wiener

---

<sup>39</sup> In other words, the separation of space and time dimensions is presupposed in both Horava-Lifshitz and Wiener fractal theories of gravity. However, it can be seen that some modified versions of Horava-Lifshitz gravity could give rise to dynamical mechanisms for breaking the Lorentz symmetry in the UV contribution. See for example [11, 15].

Brownian process and the entropic effect of the Ricci flow. In fact, the theory basically respects Heisenberg's uncertainty principle of quantum mechanics (due to its intrinsic stochastic property) and is fundamentally entropic (breaks the time-reversal symmetry).

Moreover, in Wiener fractal gravity we surprisingly showed that the source of gravity is intrinsically different from that of the three fundamental forces of nature which have been already formulated in the Standard Model, i.e. the electromagnetic and the weak and strong nuclear forces. Thus, the viewpoint of the Wiener fractal gravity to quantum interactions of nature may resolve and interpret the question of: "*Why we cannot unify gravity with other fundamental forces of nature?*" Indeed, upon what we have done above the answer is: *Because the quantum fundamental forces of nature stem from Brownian propagation of fractal quantum fields along time, whereas the gravity is basically rooted in entropic evolution of the universe.* Obviously, they address two fiercely distinct scales of nature, as we expected before.

## VI. SUMMARY AND CONCLUSION

In this paper we established a more fundamental formulation for quantum field theory in presence of gravity by employing three basic mathematical mechanisms:

- a) *The asymptotic properties of Fourier-Laplace coefficients of Weierstrass-like fractal functions,*
- b) *The Wiener stochastic process of fractal functions on a closed Riemannian manifold,*
- c) *The entropic based Ricci flow of the Riemannian metric on a closed Riemannian 3-manifold.*

We proved that the stochastic Brownian motion of fractal functions on a closed Riemannian manifold with dynamical geometry due to the Ricci flow causes the emergence of the Einstein-Hilbert action and the path-integral formulation of the corresponding quantum field theory within the Wiener probability measure. The resulting theory, the so called *Wiener fractal gravity*, could be regarded as a more fundamental formulation of gravity. We also established that the emergence of the Einstein-Hilbert Lagrangian density has nothing to do with the background quantum field theory and has a merely entropic source due to the Ricci flow.

We discussed the different sources of gravity and the other fundamental forces of nature based on the mentioned stochastic framework on dynamical Riemannian manifolds. Also, by computing the coupling constant of the Einstein-Hilbert action we interpreted the hierarchy problem in terms of the background geometric structures and the Ricci flow. Then, we worked out the cosmological constant according to Hamilton's theorem for 3-manifolds and showed that in  $D = 3$  dimensions this constant is positive and surprisingly extremely small. We also argued that the cosmological constant is a pure entropic/geometric term and is independent of the zero-point energy of the involved quantum field theories.

The correlation of the solutions of the corresponding Ricci flow and the achievements of the  $\Lambda$ CDM Model was discussed and we found some specific similarities to the FLRW metric and its cosmological features. Hubble's law and the acceleration of the universe expansion were worked out due to the Ricci flow and it was argued that the dark energy (hence the inflation) has an entropic source. Afterward, we compared our achievements with three important theories of gravity; **1)** Verlinde's entropic gravity, **2)**

Stochastic gravity, and **3**) Horava-Lifshitz gravity. We found both basic similarities and fundamental differences between these theories and the Wiener fractal gravity in some appropriate approximated versions.

Upon the above results, we are optimistic that the above formulation could provide a well-defined framework of quantum gravity. Indeed, within the mentioned formulation, the gravitational fields have no quantum (fractal) fluctuations and their effects are regarded in terms of expectation values via some semi-classical viewpoint. Actually, to insert these quantum effects within the theory one should localize the non-local fluctuations of the metric which give rise to appearance of Grassmannian superfields just similar to the mechanism of the Faddeev-Popov quantization method. Anyway, as we have shown in [80] obtaining a probability measure is only possible by employing Grassmannian spinor fields. This causes the emergence of fermionic Majorana quantum fields which accompany the bosonic gravitational fields within the interaction term of the Wiener path-integral. This could be regarded as a possible procedure to produce a path-integral formulation of supergravity by means of a probability measure.

## VII. ACKNOWLEDGMENTS

The author would like to dedicate this paper to Saya Ziaee for her birthday on January 22nd; This research would not have been possible without her emotional support. Moreover, the author would like to thank Shahroud Azami for his fruitful guidelines. Also, it should be noted that this research was in part supported by a grant from IPM (No. 1400810418).

- 
- [1] S. Azami, *Variation of the First Eigenvalue of  $(p, q)$ -Laplacian along the Ricci-Harmonic Flow*, Int. J. Nonlin. Anal. Appl. 12, No. 2: 193-204, 2021.
  - [2] A. Bekenstein, *Black Holes and the Second Law*, Lett. Nuovo Cimento. 4 (15): 99-104, 1972.
  - [3] M. Berger, *Sur le Spectre d'une Variété Riemannienne*. C.R. Acad. Sci. Paris 263: 13-16, 1963.
  - [4] M. Berger, *Sur les Premières Valeurs Propres des Variétés Riemanniennes*, Compositio Math. 26 (2): 129-149, 1973.
  - [5] M. Berger, P. Gauduchon, and E. Mazet, *Le Spectre d'une Variété Riemannienne*, Springer, 1971.
  - [6] A. L. Besse, *Einstein Manifolds*, Springer, 2008.
  - [7] C. H. Brans, and R. H. Dicke, *Mach's Principle and a Relativistic Theory of Gravitation*, Phys. Rev. 124 (3): 925-935, 1961.
  - [8] P. Buser, *A Note on the Isoperimetric Constant*, Ann. Sci. École Norm. Sup. 15, No. 2: 213-230, 1982.
  - [9] Y. F. Cai, J. Liu, and H. Li, *Entropic Cosmology: A Unified Model of Inflation and Late-Time Acceleration*, Phys. Lett. B. 690 (3): 213-219, 2010.
  - [10] X. D. Cao, *First Eigenvalues Geometric Operators under the Ricci Flow*, Proc. Amer. Math. Soc. 136: 4075-4078, 2008.
  - [11] A. Casalino, M. Rinaldi, L. Sebastiani, S. Vagnozzi, *Alive and Well: Mimetic Gravity and a Higher-Order Extension in Light of GW170817*, Class. Quant. Grav. 36: 017001, 2019.
  - [12] Z. Chang, M. H. Li, X. Li, *Unification of Dark Matter and Dark Energy in a Modified Entropic Force Model*, Commun. Theor. Phys. 56 (1): 184-192, 2011.
  - [13] I. Chavel, *Eigenvalues in Riemannian Geometry*, Academic Press, 1984.

- [14] J. Cheeger, *A Lower Bound for the Smallest Eigenvalue of the Laplacian*, In: Problems in Analysis (Papers dedicated to Salomon Bochner, 1969), Princeton University Press, Princeton, N. J., pp. 195–199, 1970.
- [15] G. Cognola, R. Myrzakulov, L. Sebastiani, S. Vagnozzi, S. Zerbini, *Covariant Horava-Like and Mimetic Horndeski Gravity: Cosmological Solutions and Perturbations*, *Class. Quant. Grav.* 33: 225014, 2016.
- [16] M. Demianski, N. Sanchez, and Y. N. Parijskij, *Topology of the Universe and the Cosmic Microwave Background Radiation*, Proceeding of the NATO Advanced Study Institute 130, Springer, 2003.
- [17] S. Deshmukh and A. Al-Eid, *Curvature Bounds for the Spectrum of a Compact Riemannian manifold of Constant Scalar Curvature*, *J. Geom. Anal.* 15(4): 589-606, 2005.
- [18] S. Deshmukh, *Eigenvalues of the Laplacian Operator on Compact Riemannian Manifolds*, *Arab. J. Math. Sc.* Vol. 13: 39-65, 2007.
- [19] D. DeTurck, *Deforming Metrics in the Direction of their Ricci Tensors*, In: H. D. Cao, B. Chow, S. C. Chu and S. T. Yau (Eds.), *Collected Papers on Ricci Flow*, Series in Geometry and Topology, 37, International Press, 2003.
- [20] B. S. DeWitt, *Quantum Field Theory in Curved Spacetime*, *Phys. Rep.* 19, No. 6: 295-357, 1975.
- [21] L. F. Di Cerbo, *Eigenvalues of the Laplacian under the Ricci Flow*, *Rend. Mat.* VII, Vol. 27, Rome, 2007.
- [22] H. Donnelly, *Eigenfunctions of the Laplacian on Compact Riemannian Manifolds*, *Asian J. Math.* Vol. 10, No. 1: 115–126, 2006.
- [23] F. Dowker, *Topology Change in Quantum Gravity*, 2002 [arXiv:gr-qc/0206020].
- [24] V. Dzhunushaliev, *Quantum Wormhole as a Ricci Flow*, *Int. J. Geom. Meth. Mod. Phys.* Vol. 06, No. 06: 1033-1046, 2009.
- [25] V. Dzhunushaliev, V. Folomeev, *Masking Singularities in Weyl Gravity and Ricci Flows*, *Eur. J. Phys. C* 81: 387-, 2021.
- [26] V. Dzhunushaliev, N. Serikbayev, and R. Myrzakulov, *Topology Change in Quantum Gravity and Ricci Flows*, 2010 [ arXiv:0912.5326 [gr-qc]].
- [27] M. Egidi, S. Liu, F. Müunch, and N. Peyerimhoff, *Ricci Curvature and Eigenvalue Estimates for the Magnetic Laplacian on Manifolds*, *Commun. Anal. Geom.* Vol. 29, No. 5, 2021.
- [28] A. El Soufi, and S. Ilias, *Laplacian Eigenvalue Functionals and Metric Deformations on Compact Manifolds*, *J. Geom. Phys.* 58: 89-104, 2008.
- [29] S. Fang, F. Yang, and P. Zhu, *Eigenvalues of Geometric Operators Related to the Witten Laplacian under the Ricci Flow*, *Glas. Math. J.*: 1-9, 2017.
- [30] G. B. Folland, *Real Analysis; Modern Techniques and Their Applications*, 2nd Ed., John Wiley and Sons, 1999.
- [31] A. Frenkel, P. Horava, and S. Randall, *Topological Quantum Gravity of the Ricci Flow*, preprint, 2020 [arXiv:2010.15369 [hep-th]].
- [32] A. Frenkel, P. Horava, and S. Randall, *The Geometry of Time in Topological Quantum Gravity of the Ricci Flow*, preprint, 2020 [arXiv:2011.06230 [hep-th]].
- [33] A. Frenkel, P. Horava, and S. Randall, *Perelman's Ricci Flow in Topological Quantum Gravity*, preprint, 2020 [arXiv:2011.11914 [hep-th]].
- [34] A. Fowlie, C. Balazs, G. White, L. Marzola, and M. Raidal, *Naturalness of the Relaxion Mechanism*, *JHEP* 2016 (8): 100, 2016.
- [35] J. A. Frieman, M. S. Turner, and D. Huterer, Dragan, *Dark Energy and the Accelerating Universe*, *Ann. Rev. Astr. Astrophys.* 46 (1): 385-432, 2008.
- [36] S. Gallot, D. Hulin, and J. Lafontaine, *Riemannian Geometry*, 2nd Ed., Springer-Verlag, 1990.
- [37] P. B. Gilkey, *Curvature and the Eigenvalues of the Laplacian for Elliptic Complexes*, *Adv. Math.* 10: 344-382, 1973.
- [38] P. B. Gilkey, *The Spectral Geometry of Riemannian Manifolds*, *J. Diff. Geom.* 10: 601-618, 1975.
- [39] P. B. Gilkey, *Recursion Relations and the Asymptotic Behavior of the Eigenvalues of the Laplacian*, *Comp.*

Math. Vol. 38, Fasc. 2: 201-240, 1979.

- [40] L. Giovanni, *A First Course in Sobolev Spaces*, Graduate Studies in Mathematics. 105. American Mathematical Society, 2009.
- [41] A. Grigor'yan, Y. Netrusov, and S. T. Yau, *Eigenvalues of Elliptic Operators and Geometric Applications*, Surv. Diff. Geom. IX: 147-217, 2004.
- [42] W. Graf, *Ricci Flow Gravity*, PMCPhys. A 1: 3, 2007.
- [43] R. S. Hamilton, *Three-Manifolds with Positive Ricci Curvature*, J. Diff. Geom. 17: 255-306, 1982.
- [44] R. S. Hamilton, *Four-Manifolds with Positive Curvature Operator*, J. Diff. Geom. 24: 153-179, 1986.
- [45] R. S. Hamilton, *The Ricci Flow on Surfaces*, Contemp. Math. 71, Amer. Math. Soc., Providence, RI: 237-262, 1988.
- [46] R. S. Hamilton, *Formation of Singularities in the Ricci Flow*, Surv. Diff. Geom. II: 7-136, 1995.
- [47] G. H. Hardy, *Weierstrass's Non-Differentiable Function*, Trans. Amer. Math. Soc. 17, 3: 301-325, 1916.
- [48] S. Hawking, *Particle Creation by Black Holes*, Commun. Math. Phys. 43 (3): 199-220, 1975.
- [49] A. Henrot, *Minimization Problems for Eigenvalues of the Laplacian*, J. Evol. Eqs. 3: 443-461, 2003.
- [50] M. P. Hobson, G. P. Efstathiou, and A. N. Lesenby, *General Relativity: An Introduction for Physicists*, Cambridge University Press, 2006.
- [51] S. Hollands, and R. M. Wald, *Quantum Fields in Curved Spacetime*, In: A. Ashtekar, B. Berger, J. Isenberg, and M. MacCallum (Eds.), *General Relativity and Gravitation: A Centennial Perspective*, pp. 513-552, Cambridge: Cambridge University Press, 2015.
- [52] P. Horava, *Quantum Gravity at a Lifshitz Point*, Phys. Rev. D. 79 (8): 084008, 2009.
- [53] S. Hou, and S. Yang, *Eigenvalues of the Laplace Operator with Potential under the Backward Ricci Flow on Locally Homogeneous 3-Manifolds*, Manus. Math. 2021.
- [54] B. L. Hu, and E. Verdaguer, *Stochastic Gravity: Theory and Applications*, Liv. Rev. Relativ. 7(1): 3, 2004.
- [55] B. L. Hu, and E. Verdaguer, *Semiclassical and Stochastic Gravity: Quantum Field Effects on Curved Spacetime*, Cambridge University Press, 2020.
- [56] E. Hubble, *A Relation between Distance and Radial Velocity among Extra-Galactic Nebulae*, Proc. Natl. Acad. Sci. USA 15(3):168-173, 1929.
- [57] G. Huisken, *Ricci Deformation of the Metric of a Riemannian Manifold*, J. Diff. Geom. 21: 47-62, 1985.
- [58] S. Hwang, J. Chang, and G. Yun, *Variational Characterizations of the Total Scalar Curvature and Eigenvalues of the Laplacian*, Pac. J. Math. 261, No. 2: 395-415, 2013.
- [59] J. M. Isidro, J. L. G. Santander, and P. Fernandez de Cordoba, *Ricci Flow, Quantum Mechanics and Gravity*, Geom. Meth. Mod. Phys. Vol. 06, No. 03: 505-512, 2009.
- [60] T. Jacobson, *Thermodynamics of Spacetime: The Einstein Equation of State*, Phys. Rev. Lett. 75: 1260-1263, 1995.
- [61] C. Lange, S. Liu, O. Post, and N. Peyerimhoff, *Frustration Index and Cheeger inequalities for Discrete and Continuous Magnetic Laplacians*, Calc. Var. Par. Diff. Eq. 54, No. 4: 4165-4196, 2015.
- [62] M. Ledoux, *Spectral Gap, Logarithmic Sobolev Constant, and Geometric Bounds*, Surv. Diff. Geom. IX: 219-240, 2004.
- [63] J. Levin, *Topology and the Cosmic Microwave Background*, Phys. Rept. 365: 251-333, 2002.
- [64] X. D. Li, *Perelman's Entropy Formula for the Witten Laplacian on Riemannian Manifolds via Bakry-Emery Ricci Curvature*, Math. Ann. 353:403-437, 2012.
- [65] X. D. Li, *From the Boltzmann H-Theorem to Perelman's W-Entropy Formula for the Ricci Flow*, Emer. Top. Diff. Equ. App.: 68-84, 2013.
- [66] A. Lichnerowicz, *Géométrie des Groupes de Transformations*, Trav. Rech. Math. 3, 1958.
- [67] P. Li, and S. T. Yau, *Eigenvalues of a Compact Riemannian Manifold*, AMS Proc. Symp. Pure Math. 36: 205-239, 1980.
- [68] M. Lulli, A. Marciano, and X. Shan, *Stochastic Quantization of General Relativity a la Ricci-Flow*, preprint,

2021 [ arXiv:2112.01490 [gr-qc]].

- [69] H. P. McKean, and I. M. Singer, *Curvature and the Eigenforms of the Laplacian*, J. Diff. Geo. 1: 43-69, 1967.
- [70] R. J. McCann, and P. M. Topping, *Ricci Flow, Entropy and Optimal Transportation*, Amer. J. Math. 132, No. 3: 711-130, 2010.
- [71] C. Morison, *Uniformisation as a Bridge Between Ricci Flow and General Relativity in Two Spatial Dimensions*, PhD. Thesis, University of Amsterdam, 2020.
- [72] R. M. Neumann, *Entropic Approach to Brownian Movement*, Amer. J. Phys. 48 (5): 354-357, 1980.
- [73] T. Padmanabhan, *Thermodynamical Aspects of Gravity: New Insights*, Rep. Prog. Phys. 73 (4): 6901, 2010.
- [74] P. J. E. Peebles, *Cosmology's Century; An Inside History of Our Modern Understanding of the Universe*, Princeton University Press, 2020.
- [75] G. Perelman, *The Entropy Formula for the Ricci Flow and its Geometric Applications*, preprint, 2002.
- [76] W. A. Poor, *Differential Geometric Structures*, Dover Publications, 1981.
- [77] U. Simon, *Curvature Bounds for the Spectrum of a Closed Einstein Spaces*, Can. J. Math. 4: 1087-1091, 1978.
- [78] A.A. Tseytlin, *On Sigma Model, RG flow, Central Charge Action and Perelman's Entropy*, Phys. Rev. D 75: 064024, 2007.
- [79] A. A. Varshovi, *Wiener Process of Fractals and Path-Integrals I: Emergent Lorentz Symmetry in Stochastic Process of Quantum Fields*, preprint, 2022 [arXiv:2201.098055 [hep-th]].
- [80] A. A. Varshovi, *Wiener Process of Fractals and Path-Integrals III*, preprint, 2022.
- [81] E. P. Verlinde, *On the Origin of Gravity and the Laws of Newton*, JHEP (4): 29, 2011.
- [82] E. P. Verlinde, *Emergent Gravity and the Dark Universe*, SciPost Phys. 2, 016, 2017.
- [83] M. Visser, *Conservative Entropic Forces*, JHEP (10): 140, 2011.
- [84] R. M. Wald, *The Backreaction Effect in Particle Creation in Curved Spacetime*, Commun. Math. Phys. 54:1-19, 1977.
- [85] R. M. Wald, *Trace Anomaly of a Conformally Invariant Quantum Field in Curved Space-Time*, Phys. Rev. D. 17:1477-1484, 1978.
- [86] F. W. Warner, *Foundations of Differentiable Manifolds and Lie Group*, Springer-Verlag, 1983.
- [87] S. Weinberg, *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity*, John Wiley and Sons, 1972.
- [88] S. Weinberg, *The Cosmological Constant Problem*, Rev. Mod. Phys. 61, No. 1: 1-23, 1989.
- [89] K. Weierstrass, *Über Continuirliche Functionen eines Reellen Arguments, die für Keinen Werth des Letzteren einen Bestimmten Differentialquotienten Besitzen*, Mathematische Werke von Karl Weierstrass, 2, Berlin, Germany: Mayer and Mller, pp. 71-74, 1895, English translation: G. Edgar, *On Continuous Functions of a Real Argument That Do not Possess a Well-Defined Derivative for Any Value of Their Argument*, *Classics on Fractals*, Studies in Nonlinearity, Addison-Wesley Publishing Company, pp. 3-9, 1993.
- [90] H. Weyl, *Über die Asymptotische Verteilung der Eigenwerte*, Nachrichten der Königlichen Gesellschaft der Wissenschaften zu Göttingen, pp. 110-117, 1911.
- [91] H. Weyl, *The Classical Groups*, Princeton University Press, 1946.
- [92] N. Wiener, *Norbert Wiener: Collected Works, Volume I; Mathematical Philosophy and Foundations; Potential Theory; Brownian Movement, Wiener Integrals, Ergodic and Chaos Theories, Turbulence and Statistical Mechanics*, P. Masani (Ed.), MIT Press, 1976
- [93] E. Woolgar, *Some Applications of Ricci Flow in Physics*, Can. J. Phys. 86: 645, 2008.
- [94] S.T. Yau, *Isoperimetric Constants and the First Eigenvalue of a Compact Riemannian Manifold*, Ann. Scient. Ec. Norm. Sup. 4: 487-507, 1985.
- [95] M. Y. Yoo, *Unexpected Connections*, Eng. Sc. LXXIV1: 30, 2011.