

Output regulation of infinite-dimensional nonlinear systems: a forwarding approach for contraction semigroups*

Nicolas Vanspranghe[†]Lucas Brivadis[‡]

Abstract

This paper deals with the problem of robust output regulation of systems governed by nonlinear contraction semigroups. After adding an integral action to the system, we design a feedback law based on the so-called forwarding approach. For small constant perturbations, we give sufficient conditions for the existence of a locally exponentially stable equilibrium at which the output coincides with the reference. Under additional assumptions, global asymptotic stability is achieved. All these conditions are investigated in the case of semilinear systems, and examples of application are given.

Key words. Output regulation, infinite-dimensional control systems, nonlinear contraction semigroups, forwarding design, semilinear systems.

AMS subject classifications. 93C20, 93C10, 93B52.

1 Introduction

Robust output regulation is one of the oldest problems in control theory. It consists in designing a feedback law which ensures that the output of a system tracks a given reference, even in the presence of external disturbances. If a feedback law can be designed to stabilize the system at some target point then a natural strategy to regulate the output robustly with respect to constant disturbances consists in adding an integral action to the system. Then, one must find a new feedback law that stabilizes the augmented system (i.e., the state and the output integrator) at some equilibrium point even in the presence of constant disturbances. The integrator guarantees that the output is at the reference when the full state is at the equilibrium.

While the theory of output regulation for linear finite-dimensional systems is well-understood since the seminal work of E. Davidson [9, 10], many problems remain open in the case of nonlinear and/or infinite-dimensional systems. In [23, 24], S. Pohjolainen proposed an extension of the finite-dimensional linear robust regulation theory to the infinite-dimensional context. In particular, a controller based on an output integrator was investigated. More recently, this work has been continued in [19, 20, 22] where an internal model principle was developed to reject a wider class of disturbances (and not only constant ones). This theory however struggles to deal with nonlinearities. In the finite-dimensional context, a strategy to take them into account is the so-called *forwarding* approach, devised by F. Mazenc and L. Praly in [18] and developed with different purposes in [1, 3, 13].

To the best of our knowledge, the present paper is a first attempt to tackle the robust regulation problem in a nonlinear infinite-dimensional abstract framework. We propose to approach the issue via

***Funding:** This work was partially funded by the French Grants ANR ODISSE (ANR-19-CE48-0004-01) and MIAI@Grenoble Alpes (ANR-19-P3IA-0003).

[†]Univ. Grenoble Alpes, CNRS, Grenoble INP, GIPSA-Lab, 38000 Grenoble, France. Email: nicolas.vanspranghe@gipsa-lab.fr.

[‡]Université Paris-Saclay, CNRS, CentraleSupélec, Laboratoire des Signaux et Systèmes, 91190, Gif-sur-Yvette, France. Email: lucas.brivadis@centralesupelec.fr.

the forwarding technique. Few extensions of this technique in the infinite-dimensional setting exist. In all of them [2, 16, 17, 27], the control system is essentially linear, except for a potential cone-bounded nonlinearity that may be applied to the input. In the recent work [11], the authors followed a contraction approach to achieve the output regulation of nonlinear finite-dimensional systems by means of an incremental forwarding approach. Inspired by this work, we focus on systems of contraction, namely, such that the distance between any pair of trajectories is exponentially converging towards zero. We particularly investigate the case of semilinear systems, that are Lipschitz perturbations of linear systems.

Organization of the paper In the first section of the paper, we state more precisely the class of systems under consideration and the output regulation problem. Then, we give some insights on the forwarding strategy we use to design a stabilizing feedback law. Main results are stated and discussed in Section 3, and proved in Section 4. Some examples of application are provided in Section 5.

Notation Let us introduce some notations and definitions. If H is a Hilbert space endowed with the norm $\|\cdot\|_H$, $\mathcal{B}_H(x, r)$ denotes the open ball of $(H, \|\cdot\|_H)$ centered at $x \in H$ of radius $r > 0$. Let Z be a Hilbert space. We denote by $\mathcal{C}(H, Z)$ the space of continuous maps from H to Z and by $\mathcal{C}^1(H, Z)$ the space of continuously Fréchet differentiable maps. We also denote by $\mathcal{L}(H, Z)$ the Banach algebra of bounded linear operators from H to Z . Given L in $\mathcal{L}(H, Z)$, the adjoint operator L^* is uniquely defined in $\mathcal{L}(Z, H)$ by $(Lx, y)_Z = (x, L^*y)_H$, where (\cdot, \cdot) denotes the scalar product. We shall say that a map $M : H \rightarrow Z$ is locally Lipschitz continuous if it is Lipschitz continuous on every bounded subset of H . Given a H -valued map \mathcal{A} defined on some subset $\mathcal{D}(\mathcal{A})$ of H , we say that w is a *strong* solution to the abstract evolution equation $dw/dt + \mathcal{A}(w) = 0$ on $(0, T)$ if w is absolutely continuous with respect to the topology of H , takes values in $\mathcal{D}(\mathcal{A})$, and solves the differential equation almost everywhere on $(0, T)$. A *weak* solution w is a limit of such strong solutions in $\mathcal{C}([0, T], H)$.

2 Problem statement

2.1 Systems under consideration

Consider a nonlinear infinite-dimensional controlled system with measured output of the form:

$$\frac{dw}{dt} + \mathcal{A}(w) = Bu(t), \quad (1a)$$

$$y = Cw, \quad (1b)$$

where w is the state of the system lying in some real Hilbert space H , y is the measured output lying in some real Hilbert space Z , u is the control input lying in some real Hilbert space U , $\mathcal{A} : \mathcal{D}(\mathcal{A}) \rightarrow H$ is a singled-valued maximal monotone operator defined on the dense subset $\mathcal{D}(\mathcal{A}) \subset H$ such that $\mathcal{A}(0) = 0$, $B : U \rightarrow H$ is a bounded linear map, and $C : \mathcal{D}(\mathcal{A}) \rightarrow Z$ is a (potentially unbounded) linear map defined on the (dense) subspace $\mathcal{D}(\mathcal{A})$. According to [25, Chapter 4, Proposition 3.1], $-\mathcal{A}$ is the generator of a nonlinear strongly continuous contraction semigroup over H , denoted by $\{\mathcal{T}_t\}$. In order to address the problem of output regulation, we assume that \mathcal{A} satisfies the following assumption.

Hypothesis 2.1 (Monotonicity). The nonlinear operator \mathcal{A} is strongly monotone, i.e., there exists $\alpha > 0$ such that

$$(\mathcal{A}(w_1) - \mathcal{A}(w_2), w_1 - w_2)_H \geq \alpha \|w_1 - w_2\|_H^2, \quad \forall w_1, w_2 \in \mathcal{D}(\mathcal{A}). \quad (2)$$

Hypothesis 2.1 implies that the semigroup $\{\mathcal{T}_t\}$ satisfies

$$\|\mathcal{T}_t w_1 - \mathcal{T}_t w_2\|_H \leq \exp(-\alpha t) \|w_1 - w_2\|_H, \quad \forall w_1, w_2 \in H, \forall t \in \mathbb{R}_+. \quad (3)$$

We address the problem of regulating the output y of (1) at some constant reference $y_{\text{ref}} \in Z$ in presence of constant disturbances $d \in H$ acting on the dynamics of the system. We proceed by adding an

integrator to the output of the system, so that the resulting full system may be written as follows:

$$\frac{dw}{dt} + \mathcal{A}(w) = Bu(t) + d, \quad (4a)$$

$$\frac{dz}{dt} = Cw - y_{\text{ref}}, \quad (4b)$$

where z lying in Z is the output integrator. The reason for that strategy is the following. Assume that a feedback law $\varphi : H \times Z \rightarrow U$ may be designed, so that (4) in closed-loop with $u(t) = \varphi(w(t), z(t))$ is asymptotically stable at $(0, 0) \in H \times Z$ when $[d, y_{\text{ref}}] = (0, 0)$. Then, formally, one can expect that for small perturbations of this system, i.e., for small values of $[d, y_{\text{ref}}]$, the closed-loop is still asymptotically stable at some new equilibrium point $[w^*, z^*]$ near $(0, 0) \in H \times Z$. Roughly speaking, this would imply that the dynamics of z , namely $dz/dt = Cw - y_{\text{ref}}$, still tend towards zero, that is, the output $y = Cw$ is regulated at y_{ref} . In this paper, we propose a design strategy of such a feedback law φ based on the so-called forwarding approach, for which we make precise this formal reasoning.

2.2 Forwarding design

Following the forwarding design [18], we consider a candidate Lyapunov function of the form:

$$V(w, z) = \frac{1}{2}\|w\|_H^2 + \frac{\rho}{2}\|z - \mathcal{M}(w)\|_Z^2 \quad (5)$$

where $\|w\|_H^2/2$ corresponds to the Lyapunov function associated to the contraction semigroup $\{\mathcal{T}_t\}$, $\mathcal{M} : H \rightarrow Z$ is some Fréchet differentiable nonlinear map that vanishes at zero to be tuned, and ρ is some positive constant to be fixed (large enough) later. Let us compute the derivative of V along a strong solution $[w, z]$ of the open-loop system (4) in the case where $[d, y_{\text{ref}}] = [0, 0]$:

$$\begin{aligned} \frac{d}{dt}V(w, z) &= -(w, \mathcal{A}(w))_H + (w, Bu(t))_H \\ &\quad + \rho(z - \mathcal{M}(w), d\mathcal{M}(w)\mathcal{A}(w) + Cw - d\mathcal{M}(w)Bu(t))_Z. \end{aligned} \quad (6)$$

Under Hypothesis 2.1, we have $(w, \mathcal{A}(w))_H \geq \alpha\|w\|_H^2$. The key idea of the forwarding design is to choose a nonlinear map \mathcal{M} satisfying the following functional equation, which guarantees that the graph of \mathcal{M} is positively invariant under the uncontrolled dynamics (4):

$$\mathcal{M}(0) = 0, \quad (7a)$$

$$d\mathcal{M}(w)\mathcal{A}(w) + Cw = 0, \quad \forall w \in \mathcal{D}(\mathcal{A}). \quad (7b)$$

In order to follow the forwarding approach, let us consider the next hypothesis.

Hypothesis 2.2. The functional equation (7) admits a continuously Fréchet differentiable solution $\mathcal{M} \in \mathcal{C}^1(H, Z)$.

For such a map \mathcal{M} , we obtain:

$$\frac{d}{dt}V(w, z) \leq -\alpha\|w\|_H^2 + (B^*w, u(t))_U - \rho(B^*d\mathcal{M}(w)^*[z - \mathcal{M}(w)], u(t))_U. \quad (8)$$

This suggests to set the feedback law as

$$u(t) = B^*d\mathcal{M}(w)^*[z - \mathcal{M}(w)]. \quad (9)$$

Then, applying Young's inequality, we get that for all $\varepsilon > 0$,

$$\frac{d}{dt}V(w, z) \leq -\alpha\|w\|_H^2 + \frac{\varepsilon}{2}\|B\|_{\mathcal{L}(U, H)}^2\|w\|_H^2 + \frac{1}{2\varepsilon}\|u\|_U^2 - \rho\|u\|_U^2. \quad (10)$$

In particular, setting $\rho = \|B\|_{\mathcal{L}(U, H)}^2/\alpha$ and $\varepsilon = 1/\rho$ yields

$$\frac{d}{dt}V(w, z) \leq -\frac{\alpha}{2}\|w\|_H^2 - \frac{\rho}{2}\|u\|_U^2 \leq 0, \quad (11)$$

so that V is indeed a Lyapunov function. These formal computations give an insight on the feedback law (9) to choose and on the Lyapunov function (5) to consider in order to stabilize (4). However, many important questions remain open, in particular concerning the well-posedness of the closed-loop, the stability properties in presence of $[d, y_{\text{ref}}]$, and the feasibility of the functional equation (7).

Remark 2.3 (On Hypothesis 2.2). Checking Hypothesis 2.2 on a general nonlinear system of the form (1) is difficult in general. Moreover, since \mathcal{M} is used in feedback law (9), an expression of \mathcal{M} in terms of parameters of the system should be given in order to implement the controller. For these two reasons, we investigate later in the paper the case of semilinear systems, for which we give sufficient conditions on \mathcal{A} for Hypothesis 2.2 to be satisfied, as well as an expression of \mathcal{M} in terms of \mathcal{A} and C .

3 Main results

We now state our results on the closed-loop system (4)-(9).

3.1 Well-posedness of the closed-loop dynamics

Our first result concerns the existence and uniqueness of solutions to the closed-loop equations (4)-(9).

Theorem 3.1 (Well-posedness). *Assume Hypotheses 2.1 and 2.2 hold and $d\mathcal{M}$ is locally Lipschitz continuous. Then,*

- (i) *For all $[d, y_{\text{ref}}] \in H \times Z$ and all $[w_0, z_0] \in H \times Z$, the closed-loop system (4)-(9) admits a unique maximal weak solution $[w, z]$ satisfying the initial condition $[w(0), z(0)] = [w_0, z_0]$ and defined on $[0, T_{\max}(w_0, z_0))$ for some $T_{\max}(w_0, z_0) \in (0, +\infty]$;*
- (ii) *Moreover, if $w_0 \in \mathcal{D}(\mathcal{A})$, then $[w, z]$ is a strong solution to (4)-(9);*
- (iii) *Furthermore, if $d = 0$ or $d\mathcal{M}$ is globally Lipschitz continuous, then (4)-(9) is forward complete, i.e., $T_{\max}(w_0, z_0) = +\infty$ for all $[w_0, z_0] \in H \times Z$.*

The proof of Theorem 3.1 is given in Section 4.1 and works around two technical difficulties (the possible unboundedness of C and the *a priori* non-monotonicity of the system with integral action) by an appropriate use of Hypothesis 2.2.

3.2 Sufficient conditions for output regulation

In view of the output regulation problem, we wish to investigate the existence of an attractive equilibrium. The following theorem is the main result of the paper.

Theorem 3.2 (Sufficient conditions for output regulation). *Assume Hypotheses 2.1 and 2.2 are satisfied. Assume that*

$$\text{Range } d\mathcal{M}(0)B = Z, \quad \text{i.e.,} \quad \exists \lambda > 0 \mid \forall z \in Z, \quad \|B^* d\mathcal{M}(0)^* z\|_Z^2 \geq \lambda \|z\|_Z^2. \quad (12)$$

If $d\mathcal{M}$ is globally (resp. locally) Lipschitz continuous, then there exist positive constants M , κ and r and a neighborhood \mathcal{N} of the origin in $H \times Z$ such that for any $[d, y_{\text{ref}}]$ in $\mathcal{B}_{H \times Z}(0, r)$ (resp. in $\{0\} \times \mathcal{B}_Z(0, r)$), the following results hold.

- *There exists an equilibrium point $[w^*, z^*] \in \mathcal{D}(\mathcal{A}) \times Z$ in \mathcal{N} of the closed-loop system (4)-(9) such that $Cw^* = y_{\text{ref}}$ and for all $[w_0, z_0]$ in \mathcal{N} , the corresponding solution $[w, z]$ of (4)-(9) satisfies, for all $t \in \mathbb{R}_+$,*

$$\|[w(t) - w^*, z(t) - z^*]\|_{H \times Z} \leq M \exp(-\kappa t) \|[w_0 - w^*, z_0 - z^*]\|_{H \times Z}. \quad (13)$$

- *Moreover, if*

$$\|B^* d\mathcal{M}(w)^* z\|_Z^2 \geq \lambda \|z\|_Z^2, \quad \forall z \in Z, \quad \forall w \in H, \quad (14)$$

then for all $[w_0, z_0]$ in $H \times Z$, the corresponding solution $[w, z]$ of (4)-(9) satisfies:

$$[w(t), z(t)] \rightarrow [w^*, z^*] \quad \text{in } H \times Z \quad \text{as } t \rightarrow +\infty. \quad (15)$$

The proof of Theorem 3.2 is given in Section 4.2. Equation (13) guarantees the local exponential stability of the equilibrium point $[w^*, z^*]$, while (15) provides global asymptotic stability under the additional condition (14). This condition is stronger than a global version of the range condition (12), in the sense that the constant λ must be independent of $w \in H$. We shall show in the proof how this uniformity implies the global attractivity of the local exponential basin of attraction \mathcal{N} .

Conditions (12) and (14) may be difficult to check in general. In Section 3.3, we show how to investigate them in the semilinear case. In the finite-dimensional context, (14) is equivalent to the incremental condition given in [11, Assumption 2]. In our work, it allows to obtain global asymptotic stability even in the presence of small perturbations. The local condition (12) is commonly used in the forwarding design (see, e.g., [1]) to obtain local exponential stability. Note that in the finite-dimensional context, (12) is sufficient to obtain local exponential stability and global asymptotic stability in absence of perturbations [1, Lemma 1].

It is clear that if the operator C is continuous, i.e., C belongs to $\mathcal{L}(H, Z)$, then the output $Cw(t)$ goes to y_{ref} in Z whenever $w(t)$ goes to w^* in H . If C is unbounded, the question is more delicate and is discussed in Section 3.4.

3.3 The semilinear case

Consider now the case where $\mathcal{A} = A + F$ with $\mathcal{D}(\mathcal{A}) = \mathcal{D}(A)$, where $-A$ is the infinitesimal generator of a strongly continuous semigroup $\{\mathcal{S}_t\}$ of linear operators on H , and F is a nonlinear mapping satisfying

$$F \in \mathcal{C}^1(H), \quad \text{d}F \text{ locally Lipschitz continuous}, \quad (16)$$

and without loss of generality, $\text{d}F(0) = 0$. Following [21], we shall say that \mathcal{A} is semilinear. In that context, let us introduce the following set of assumptions.

Hypothesis 3.3. The operator $\mathcal{A} = A + F$ is semilinear and satisfies the following properties:

- (i) C is A -bounded, i.e., there exist positive constants a and b such that

$$\|Cw\|_Z \leq a\|Aw\|_H + b\|w\|_H, \quad \forall w \in \mathcal{D}(A); \quad (17)$$

- (ii) 0 is in the resolvent set of A ;
- (iii) There exists $\alpha > 0$ such that

$$(Ah + \text{d}F(w)h, h)_H \geq \alpha\|h\|_H^2, \quad \forall w \in H, \forall h \in \mathcal{D}(A). \quad (18)$$

Remark that Hypothesis 3.3 implies both Hypothesis 2.1 and maximal monotonicity of \mathcal{A} . Indeed, writing $F(w_1) - F(w_2)$ as an integral of $\text{d}F$ along the line segment joining w_1 to w_2 , one can show that (18) implies (2). Then, standard results on Lipschitz perturbations of linear systems (see [21, Section 6.1]) together with (2) yield that $-\mathcal{A}$ generates a strongly continuous semigroup of contractions (denoted by $\{\mathcal{T}_t\}$) on H , which in turn implies that \mathcal{A} is maximal monotone by virtue of [14, Theorem 4].

Under Hypothesis 3.3, we show in the following theorem that (7) admits a solution that can be expressed in terms of C , A , and F .

Theorem 3.4 (Existence of \mathcal{M} in the semilinear case). *Assume Hypothesis 3.3 is satisfied. Then there exists a map $\mathcal{M} \in \mathcal{C}^1(H, Z)$ satisfying (7) with $\text{d}\mathcal{M}$ locally Lipschitz continuous and given for all $w \in H$ by*

$$\mathcal{M}(w) \triangleq -C \left\{ \lim_{\tau \rightarrow +\infty} \int_0^\tau \mathcal{T}_t w \, dt \right\} = -CA^{-1}w - CA^{-1} \int_0^{+\infty} F(\mathcal{T}_t w) \, dt. \quad (19)$$

For all $w, h \in H$,

$$\text{d}\mathcal{M}(w)h = -CA^{-1}h - CA^{-1} \int_0^{+\infty} \text{d}F(\mathcal{T}_t w) \text{d}\mathcal{T}_t(w)h \, dt, \quad (20)$$

where $\text{d}\mathcal{T}_t(w)$ denotes the differential of \mathcal{T}_t at w , which exists. Moreover, if F and $\text{d}F$ are globally Lipschitz continuous, then so is $\text{d}\mathcal{M}$.

The proof of Theorem 3.4 is given in Section 4.3. Its purpose is to guarantee the existence of the control law (9) for a wide class of semilinear systems and provide easy conditions under which requirements of Theorem 3.2 are met. A notable consequence of (20) is that the coercivity condition (12) in Theorem 3.2 simply reads as

$$\text{Range } CA^{-1}B = Z, \quad (21)$$

which, in the context of output regulation of finite-dimensional linear systems, corresponds to a *non-resonance* condition between A and the zero dynamics of the integrator via the Schur complement (see, e.g., [12]). On the other hand, Theorem 3.4 also states that $d\mathcal{M}$ inherits the Lipschitz properties of F and dF . Now, regarding Hypothesis 3.3, (18) is roughly speaking a sufficient condition under which solutions to the uncontrolled w -equation linearized around a given trajectory (also called *first variation equation*) uniformly converge to that trajectory. Equation (18) is easily verified in (at least) two situations of interest.

- The nonlinearity F contributes to the contraction behavior of the w -dynamics. This is the case for example if A is coercive and F is monotone (i.e. such that $(dF(w)h, h)_H$ is nonnegative for all w, h in H). More generally, the reader may refer to [15] in finite dimension or [26, Chapter V] for the contraction analysis of the system by means of its first variation equation.
- Variations of F are small with respect to linear dissipation brought by A . More precisely, if F is K -Lipschitz continuous, then $\|dF(\cdot)\|_{\mathcal{L}(H)}$ is bounded by K , hence (18) is satisfied if $(Ah, h)_H \geq \beta \|h\|_H^2$ for some constant $\beta > K$. In the same spirit, under (18) and (21), by observing¹ that for each w the integral map in (20) has operator norm bounded by K/α , one deduces from (20) that *global* uniform coercivity (14) holds whenever $K < \alpha$.

Remark 3.5 (Linear case). If \mathcal{A} is linear, i.e., $F = 0$, and 0 is in the resolvent set of A , then the solution \mathcal{M} to (7) is explicitly and uniquely determined as $\mathcal{M} = -CA^{-1}$. This choice corresponds to the linear forwarding approach followed by [17, 27].

3.4 The case of unbounded output

We conclude Section 3 with an informal discussion regarding the convergence of the output $Cw(t)$ towards the reference when the operator C is unbounded. To get some insight on the situation, assume for a moment that the original w -system is linear. Then, the solution \mathcal{M} to (7) provided by Theorem 3.4 is a bounded linear operator, and the closed-loop dynamics around the equilibrium are governed by a strongly continuous linear semigroup, which commutes with its generator. Recalling the notation from Section 3.3, it follows that for a *strong* solution $[w, z]$ to (4)-(9), $w(t) - w^*$ goes to 0 in $\mathcal{D}(A)$ endowed with the graph norm, hence $Cw(t)$ converges to y_{ref} provided that C is A -bounded. In the nonlinear case, the argument breaks down.

Alternatively, one may look for weaker notions of convergence. In many applications, unbounded output operators of interest enjoy an *admissibility* property with respect to the uncontrolled dynamics. In the linear theory, C is said to be A -admissible if C is A -bounded and there exist positive constants K and T such that

$$\int_0^T \|CS_t w_0\|_Z^2 dt \leq K \|w_0\|_H^2, \quad \forall w_0 \in \mathcal{D}(A). \quad (22)$$

In the semilinear case, one can deduce from (22) that, first of all, the output Cw is well-defined in $L_{\text{loc}}^2(0, +\infty; Z)$ even for weak solutions $[w, z]$ to the closed-loop equations (4)-(9), and secondly, that the output converges “in average” to the reference:

$$\lim_{\tau \rightarrow +\infty} \int_{\tau}^{T+\tau} \|Cw(t) - y_{\text{ref}}\|_Z^2 dt = 0 \quad (23)$$

for any (including weak) solution $[w, z]$ that converges to the equilibrium $[w^*, z^*]$ in $H \times Z$. When no semilinear structure is prescribed for the maximal monotone operator \mathcal{A} governing the w -dynamics, we

¹See Lemma 4.6 below for the bound K/α .

generalize (22) by assuming that for any w_i^0 in $\mathcal{D}(\mathcal{A})$ and f_i absolutely continuous with derivative in $L^2(0, T; H)$, the solution² w_i to $dw_i/dt + \mathcal{A}(w) = f_i$ with initial condition $w_i(0) = w_i^0$, $i \in \{1, 2\}$, satisfies

$$\int_0^T \|Cw_1(t) - Cw_2(t)\|_Z^2 dt \leq K \|w_1(0) - w_2(0)\|_H^2 + K \int_0^T \|f_1 - f_2\|_H^2 dt. \quad (24)$$

In that case, the same conclusions hold for closed-loop solutions.

4 Proofs of the main results

This section is devoted to the proofs of the results of the paper. In what follows, we will investigate well-posedness, global attractivity (15) and local exponential stability (13) of the closed-loop system (4)-(9) in the new coordinates $[w, \eta]$ where η is given by

$$\eta \triangleq z - \mathcal{M}(w). \quad (25)$$

In $[w, \eta]$ -coordinates, (4)-(9) may be equivalently rewritten as

$$\frac{dw}{dt} + \mathcal{A}(w) = BB^*d\mathcal{M}(w)^*\eta + d, \quad (26a)$$

$$\frac{d\eta}{dt} + d\mathcal{M}(w)BB^*d\mathcal{M}(w)^*\eta = -y_{\text{ref}} - d\mathcal{M}(w)d. \quad (26b)$$

Equation (26b) is obtained by differentiating (25) and using (7) combined with (26a). Since \mathcal{M} is continuous, (25) preserves the topology of the space: given $[w^*, z^*] \in H \times Z$, any solution $[w, z]$ of (4)-(9) converges towards some $[w^*, z^*] \in H \times Z$ if and only if $[w, \eta]$ converges towards $[w^*, \eta^*]$, where $\eta = z - \mathcal{M}(z)$ and $\eta^* = z^* - \mathcal{M}(w^*)$. Moreover, given a ball of H containing w^* , by local Lipschitz continuity of \mathcal{M} , there exist positive constants K_1 and K_2 such that

$$K_1 \| [w - w^*, z - z^*] \|_{H \times Z} \leq \| [w - w^*, \eta - \eta^*] \|_{H \times Z} \leq K_2 \| [w - w^*, z - z^*] \|_{H \times Z} \quad (27)$$

whenever w lies in that ball. Therefore, well-posedness, global attractivity (15), and local exponential stability (13), of (4)-(9) and (26) are actually equivalent. Finally, to alleviate notation when needed, we define

$$\mathcal{K}(w) \triangleq d\mathcal{M}(w)B \in \mathcal{L}(U, Z), \quad \forall w \in H. \quad (28)$$

4.1 Proof of Theorem 3.1 (well-posedness)

We start by proving that solutions to (26) exist at least on a finite time interval (Items (i) and (ii) of Theorem 3.1), and then we investigate forward completeness (Item (iii) of Theorem 3.1).

Step 1: Local well-posedness. First, we observe that (26) represents a locally Lipschitz perturbation of the following maximal monotone problem:

$$\frac{d}{dt}[w, \eta] + [\mathcal{A}(w), \eta] = 0. \quad (29)$$

Indeed, by letting

$$\mathcal{F}_{d, y_{\text{ref}}}[w, \eta] \triangleq \begin{bmatrix} -B\mathcal{K}(w)^*\eta - d \\ -\eta + \mathcal{K}(w)\mathcal{K}(w)^*\eta + y_{\text{ref}} + d\mathcal{M}(w)d \end{bmatrix}, \quad \forall [w, \eta] \in H \times Z, \quad (30)$$

we can rewrite (26) as follows:

$$\frac{d}{dt}[w, \eta] + [\mathcal{A}(w), \eta] + \mathcal{F}_{d, y_{\text{ref}}}[w, \eta] = 0. \quad (31)$$

²Existence and uniqueness of a strong solution is guaranteed by [25, Chapter 4, Theorem 4.1]. In particular, each w_i is absolutely continuous; hence $\mathcal{A}(w_i)$ is measurable, and so is Cw_i by (7).

The nonlinear map $\mathcal{F}_{d, y_{\text{ref}}}$ is locally Lipschitz continuous on $H \times Z$. Besides, since \mathcal{A} is maximal monotone on H , so is the mapping $[w, \eta] \mapsto [\mathcal{A}(w), \eta]$ on $H \times Z$ (with dense domain $\mathcal{D}(\mathcal{A}) \times Z$). Furthermore, $[\mathcal{A}(0), 0] = 0$. Thus, it follows from [6, Theorem 7.2] that for each initial condition $[w_0, \eta_0] \in H \times Z$, there exists a unique maximal weak solution $[w, \eta]$ to (26) defined on $[0, T_{\max}(w_0, \eta_0))$, with $T_{\max}(w_0, \eta_0) \in (0, +\infty]$. Moreover, if $w_0 \in \mathcal{D}(\mathcal{A})$, $[w, \eta]$ is actually a strong solution. Recalling that $z = \eta + \mathcal{M}(w)$ and $\mathcal{M} \in \mathcal{C}^1(H, Z)$, we obtain Items (i) and (ii). Finally, if $T_{\max}(w_0, \eta_0)$ is finite, then the norm of $[w(t), \eta(t)]$ must go to $+\infty$ as t approaches $T_{\max}(w_0, \eta_0)$.

Step 2: Sufficient conditions for forward completeness. Let us prove that if take $d = 0$ or $d\mathcal{M}$ is globally Lipschitz continuous, then $T_{\max}(w_0, \eta_0) = +\infty$ for any initial condition $[w_0, \eta_0] \in H \times Z$. Let $\rho > 0$ to be fixed (large enough) later on. Given a (strong) solution $[w, \eta]$ with initial data $[w_0, \eta_0] \in \mathcal{D}(\mathcal{A}) \times Z$, we have

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \{ \|w\|_H^2 + \rho \|\eta\|_Z^2 \} &= -(\mathcal{A}(w), w)_H + (BK(w)^* \eta, w)_H + (d, w)_H \\ &\quad - \rho \|\mathcal{K}(w)^* \eta\|_U^2 - \rho (y_{\text{ref}}, \eta)_Z - \rho (d\mathcal{M}(w)d, \eta)_Z \end{aligned} \quad (32)$$

holding almost everywhere (a.e.) on $(0, T_{\max}(w_0, \eta_0))$. Similarly as in our preliminary Lyapunov analysis (8)-(10), in order to deal with the term $BK(w)^* \eta$, we use (2) together with Young's inequality and choose $2\rho \geq \|B\|_{\mathcal{L}(U, H)}^2$ to obtain

$$\frac{d}{dt} \{ \|w\|_H^2 + \rho \|\eta\|_Z^2 \} \leq 2\|w\|_H^2 + 2\rho \|\eta\|_Z^2 + \rho \|y_{\text{ref}}\|_Z^2 + \|d\|_H^2 + \rho \|d\mathcal{M}(w)d\|_Z^2 \quad (33)$$

If $d = 0$ or $d\mathcal{M}$ is globally Lipschitz continuous, the following inequality holds:

$$\|d\mathcal{M}(w)d\|_Z^2 \leq K(1 + \|w\|_H^2) \|d\|_H^2 \quad (34)$$

for some $K > 0$ independent of w or d . Combining (33) and (34) yields

$$\frac{d}{dt} \{ \|w\|_H^2 + \rho \|\eta\|_Z^2 \} \leq K' \{ \|w\|_H^2 + \rho \|\eta\|_Z^2 \} + K' \quad \text{a.e.} \quad (35)$$

for some $K' > 0$ independent of the initial data. As a strong solution to (26), $[w, \eta]$ is absolutely continuous in $H \times Z$; therefore, we can deduce from (35) and Grönwall's inequality the following uniform estimate: for all $t \in [0, T_{\max}(w_0, \eta_0))$,

$$\|w(t)\|_H^2 + \rho \|\eta(t)\|_Z^2 \leq \exp(K't) \{ \|w_0\|_H^2 + \rho \|\eta_0\|_Z^2 + 1 \}. \quad (36)$$

We infer from (36) that the norm of $[w, \eta]$ cannot blow up in finite time; thus, $T_{\max}(w_0, \eta_0) = +\infty$. Furthermore, by passing to the limit, we see that (36) is satisfied for weak solutions as well, which means that the same conclusion holds for any initial data in $H \times Z$. This proves Item (iii) and ends the proof of Theorem 3.1.

4.2 Proof of Theorem 3.2 (output regulation)

Our main result concerning the output regulation problem is demonstrated here, along with some auxiliary results.

4.2.1 Outline of the proof and intermediate results

The proof of Theorem 3.2 relies on a series of lemmas that are given below. Let us first give some insight on the main strategy.

1. Under the range condition (12), the operators $\mathcal{K}(w)^* = B^*d\mathcal{M}(w)^*$ involved in the η -equation (26b) enjoy a coercivity property that is uniform with respect to w , provided that w is small. This is shown in Lemma 4.1.

2. This allows to prove that there exists a (not necessarily invariant) region around the origin of $H \times Z$ where the dynamics generated by (26) are strictly contractive. This is stated in Lemma 4.2.
3. Another consequence of the local coercivity property is the existence of suitable attracting sets for the closed-loop dynamics (26), as demonstrated in Lemma 4.3. Furthermore, if the reference y_{ref} and the perturbation d are sufficiently small, then those sets are contained in the contraction region of the previous step.
4. Finally, existence of a locally exponentially stable equilibrium for (26) is shown by using Banach fixed point arguments and properties of ω -limit sets associated with contraction semigroups.
5. Additionally, under the global coercivity condition (14), the attracting sets from Lemma 4.3 are in fact globally attractive, hence the equilibrium is also globally asymptotically stable.

Coming back to the $[w, z]$ -coordinates and thus to the original closed-loop system (4)-(9) is then straightforward. In what follows, we shall be careful regarding the dependence of the various neighborhoods involved in the analysis with respect to each parameter. Before proceeding further, we introduce some additional notation. Recalling (30) and (31), given $[d, y_{\text{ref}}]$, we denote by $\tilde{\mathcal{A}}_{d, y_{\text{ref}}}$ the nonlinear operator $[\mathcal{A}, \text{id}] + \mathcal{F}_{d, y_{\text{ref}}}$ defined on $\mathcal{D}(\mathcal{A}) \times Z$ and associated with the closed-loop system (26) in the $[w, \eta]$ -coordinates. Also, for $\rho > 0$, we denote by $\|\cdot\|_{H \times Z, \rho}$ the Hilbert norm on $H \times Z$ given by

$$\|[w, \eta]\|_{H \times Z, \rho}^2 \triangleq \|w\|_H^2 + \rho \|\eta\|_Z^2, \quad \forall [w, \eta] \in H \times Z. \quad (37)$$

We denote by $(\cdot, \cdot)_{H \times Z, \rho}$ the associated scalar product. All these norms are equivalent. Bearing in mind our Lyapunov analysis in the $[w, z]$ -coordinates in Section 2.2, the ρ -norm is connected to the Lyapunov function V by $V(w, z) = (1/2)\|[w, z - \mathcal{M}(w)]\|_{H \times Z, \rho}^2$. In the sequel, we either take $d = 0$ or assume that $d\mathcal{M}$ is globally Lipschitz continuous. Therefore, by virtue of Theorem 3.1, solutions to the closed-loop equations (26) are well-defined for all positive time and any initial condition in $H \times Z$. Recall from (12) in the statement of Theorem 3.2 that the following coercivity assumption is in force:

$$\|\mathcal{K}(0)^* z\|_U^2 \geq \lambda \|z\|_Z^2, \quad \forall z \in Z. \quad (38)$$

The first lemma is a consequence of (38) and continuity of $d\mathcal{M}$.

Lemma 4.1 (Local coercivity). *There exist positive constants $\tilde{\lambda}$ and \tilde{r} such that*

$$\|\mathcal{K}(w)^* z\|_U^2 \geq \tilde{\lambda} \|z\|_Z^2, \quad \forall z \in Z, \forall w \in \mathcal{B}_H(0, \tilde{r}). \quad (39)$$

Proof. Let $w \in H$ and $z \in Z$. Recall that $\mathcal{K}(w)^* z = B^* d\mathcal{M}(0)^* z + B^* [d\mathcal{M}(w) - d\mathcal{M}(0)]^* z$. Using Cauchy-Schwarz and Young inequalities together with (38), we get

$$\|\mathcal{K}(w)^* z\|_U^2 \geq \frac{2\lambda}{3} \|z\|_Z^2 - 2\|B\|_{\mathcal{L}(U, H)}^2 \| [d\mathcal{M}(w) - d\mathcal{M}(0)]^* \|_{\mathcal{L}(Z, U)}^2 \|z\|_Z^2. \quad (40)$$

By continuity of $d\mathcal{M}$ at 0, we can choose $\tilde{r} > 0$ such that

$$\|d\mathcal{M}(w) - d\mathcal{M}(0)\|_{\mathcal{L}(H, Z)}^2 \leq \frac{\lambda}{6\|B\|_{\mathcal{L}(U, H)}^2}, \quad \forall w \in \mathcal{B}_H(0, \tilde{r}). \quad (41)$$

Thus, by letting $\tilde{\lambda} \triangleq \lambda/3 > 0$, we obtain the desired inequality (39). \square

The following constant appears in the next two lemmas:

$$\kappa \triangleq \min\{\alpha/4, \tilde{\lambda}/4\}. \quad (42)$$

Those concern the contraction property of the dynamics governed by (26) around the origin and the existence of attractive sets depending on $[d, y_{\text{ref}}]$.

Lemma 4.2 (Local strong monotonicity). *There exists $\rho_0 > 0$ such that the following property holds: for all $\rho \geq \rho_0$, there exists a positive $r_{0,\rho} \leq \tilde{r}$ such that*

$$\begin{aligned} (\tilde{\mathcal{A}}_{d,y_{\text{ref}}}[w_1, \eta_1] - \tilde{\mathcal{A}}_{d,y_{\text{ref}}}[w_2, \eta_2], [w_1, \eta_1] - [w_2, \eta_2])_{H \times Z, \rho} \\ \geq \kappa \|[w_1, \eta_1] - [w_2, \eta_2]\|_{H \times Z, \rho}^2 \end{aligned} \quad (43)$$

for all $[w_1, \eta_1], [w_2, \eta_2] \in \mathcal{B}_{H \times Z}(0, r_{0,\rho}) \cap \mathcal{D}(\mathcal{A}) \times Z$, $d \in \mathcal{B}_H(0, r_{0,\rho})$ and $y_{\text{ref}} \in Z$.

Proof. Let $[w_1, \eta_2]$ and $[w_2, \eta_2]$ in $\mathcal{D}(\mathcal{A}) \times Z$. We write $\tilde{w} \triangleq w_1 - w_2$ and $\tilde{\eta} \triangleq \eta_1 - \eta_2$. Then, for any $\rho > 0$,

$$\begin{aligned} (\tilde{\mathcal{A}}_{d,y_{\text{ref}}}[w_1, \eta_1] - \tilde{\mathcal{A}}_{d,y_{\text{ref}}}[w_2, \eta_2], [\tilde{w}, \tilde{\eta}])_{H \times Z, \rho} \\ = (\mathcal{A}(w_1) - \mathcal{A}(w_2) - B[\mathcal{K}(w_1)^* \eta_1 - \mathcal{K}(w_2)^* \eta_2], \tilde{w})_H \\ + \rho(\mathcal{K}(w_1)\mathcal{K}(w_1)^* \eta_1 - \mathcal{K}(w_2)\mathcal{K}(w_2)^* \eta_2 + [d\mathcal{M}(w_1) - d\mathcal{M}(w_2)]d, \tilde{\eta})_Z. \end{aligned} \quad (44)$$

By adding and removing some terms, (44) can be rewritten as follows:

$$\begin{aligned} (\tilde{\mathcal{A}}_{d,y_{\text{ref}}}[w_1, \eta_1] - \tilde{\mathcal{A}}_{d,y_{\text{ref}}}[w_2, \eta_2], [\tilde{w}, \tilde{\eta}])_{H \times Z, \rho} = (\mathcal{A}(w_1) - \mathcal{A}(w_2), \tilde{w})_H \\ + \rho(\mathcal{K}(w_1)\mathcal{K}(w_1)^* \tilde{\eta}, \tilde{\eta})_Z + \rho([\mathcal{K}(w_1)\mathcal{K}(w_1)^* - \mathcal{K}(w_2)\mathcal{K}(w_2)^*] \eta_2, \tilde{\eta})_Z \\ - (B\mathcal{K}(w_1)^* \tilde{\eta}, \tilde{w})_H - (B[\mathcal{K}(w_1) - \mathcal{K}(w_2)]^* \eta_2, \tilde{w})_H + \rho([d\mathcal{M}(w_1) - d\mathcal{M}(w_2)]d, \tilde{\eta})_Z. \end{aligned} \quad (45)$$

Assume for a moment that $\|w_i\|_H \leq \tilde{r}/2$, where \tilde{r} is given by Lemma 4.1. Therefore, $\|\tilde{w}\|_H \leq \tilde{r}$; and using (2) and (39) together with the Lipschitz continuity of \mathcal{K} and $\mathcal{K}(\cdot)\mathcal{K}(\cdot)^*$ on $\mathcal{B}_H(0, \tilde{r})$, we infer from (45) that

$$\begin{aligned} (\tilde{\mathcal{A}}_{d,y_{\text{ref}}}[w_1, \eta_1] - \tilde{\mathcal{A}}_{d,y_{\text{ref}}}[w_2, \eta_2], [\tilde{w}, \tilde{\eta}])_{H \times Z, \rho} \geq \alpha \|\tilde{w}\|_H^2 + \rho \tilde{\lambda} \|\tilde{\eta}\|_Z^2 \\ - \rho K_1 \|\tilde{w}\|_H \|\eta_2\|_Z \|\tilde{\eta}\|_Z - K_2 \|\tilde{\eta}\|_Z \|\tilde{w}\|_H - K_3 \|\eta_2\|_Z \|\tilde{w}\|_H^2 - \rho K_4 \|\tilde{w}\|_H \|d\|_H \|\tilde{\eta}\|_Z. \end{aligned} \quad (46)$$

where the K_i are some positive constants independent of $[w_i, \eta_i]$, $[d, y_{\text{ref}}]$ and ρ . Given $\varepsilon > 0$, we employ Cauchy-Schwarz and Young inequalities to obtain

$$\begin{aligned} (\tilde{\mathcal{A}}_{d,y_{\text{ref}}}[w_1, \eta_1] - \tilde{\mathcal{A}}_{d,y_{\text{ref}}}[w_2, \eta_2], [\tilde{w}, \tilde{\eta}])_{H \times Z, \rho} \geq \alpha \|\tilde{w}\|_H^2 + \rho \tilde{\lambda} \|\tilde{\eta}\|_Z^2 \\ - \frac{\rho}{2} \{K_1 \|\eta_2\|_Z + K_4 \|d\|_H\} \{\|\tilde{w}\|_H^2 + \|\tilde{\eta}\|_Z^2\} - \frac{K_2}{2\varepsilon} \|\tilde{\eta}\|_Z^2 - \frac{\varepsilon K_2}{2} \|\tilde{w}\|_H^2 - K_3 \|\eta_2\|_Z \|\tilde{w}\|_H^2. \end{aligned} \quad (47)$$

Let $\varepsilon = \alpha/(2K_2)$ in (47) and define $\rho_0 \triangleq 2K_2/(\tilde{\lambda}\varepsilon)$. For all $\rho \geq \rho_0$,

$$\begin{aligned} (\tilde{\mathcal{A}}_{d,y_{\text{ref}}}[w_1, \eta_1] - \tilde{\mathcal{A}}_{d,y_{\text{ref}}}[w_2, \eta_2], [\tilde{w}, \tilde{\eta}])_{H \times Z, \rho} \geq \frac{\alpha}{2} \|\tilde{w}\|_H^2 + \frac{\rho \tilde{\lambda}}{2} \|\tilde{\eta}\|_Z^2 \\ - \frac{\rho}{2} \{K_1 \|\eta_2\|_Z + K_4 \|d\|_H\} \{\|\tilde{w}\|_H^2 + \|\tilde{\eta}\|_Z^2\} - K_3 \|\eta_2\|_Z \|\tilde{w}\|_H^2. \end{aligned} \quad (48)$$

Let $\rho \geq \rho_0$. We infer from (48) that there exists $r_{0,\rho} \leq \tilde{r}/2$ such that

$$(\tilde{\mathcal{A}}_{d,y_{\text{ref}}}[w_1, \eta_1] - \tilde{\mathcal{A}}_{d,y_{\text{ref}}}[w_2, \eta_2], [\tilde{w}, \tilde{\eta}])_{H \times Z, \rho} \geq \frac{\alpha}{4} \|\tilde{w}\|_H^2 + \frac{\rho \tilde{\lambda}}{4} \|\tilde{\eta}\|_Z^2 \quad (49)$$

as long as $[w_i, \eta_i] \in \mathcal{B}_{H \times Z}(0, r_{0,\rho})$ and $d \in \mathcal{B}_H(0, r_{0,\rho})$, which completes the proof. \square

Lemma 4.3 (Absorbing balls). *There exists $\rho_1 > 0$ such that the following property holds: for any $\rho \geq \rho_1$, there exist positive $r_{1,\rho}$ and K_ρ such that, if $[d, y_{\text{ref}}] \in \mathcal{B}_{H \times Z}(0, r_{1,\rho})$, then the estimate*

$$\begin{aligned} \|[w(t), \eta(t)]\|_{H \times Z, \rho}^2 \\ \leq \exp(-\kappa t) \|[w_0, \eta_0]\|_{H \times Z, \rho}^2 + K_\rho \{1 - \exp(-\kappa t)\} \|[d, y_{\text{ref}}]\|_{H \times Z}^2 \end{aligned} \quad (50)$$

holds for any solution $[w, \eta]$ to (26) with initial data $[w_0, \eta_0]$ in $\mathcal{B}_{H \times Z}(0, r_{1,\rho})$. Furthermore, if (39) holds globally, then (50) is true for all initial data in $H \times Z$.

Proof. All formal computations performed below are justified by considering appropriate sequence of strong solutions as provided by Theorem 3.1 and then passing to the limit at the very end. First, let

$$\rho_1 \triangleq \|B\|_{\mathcal{L}(U,H)}^2 \max\{1, 2\alpha^{-1}\}. \quad (51)$$

Let $\rho \geq \rho_1$ be fixed and let $[w, \eta]$ be a solution to (26) with initial condition $[w_0, \eta_0]$. As in the proof of Theorem 3.1, using (32) we obtain

$$\frac{d}{dt} \|[w, \eta]\|_{H \times Z, \rho}^2 \leq k_\rho \{ \|[w, \eta]\|_{H \times Z, \rho}^2 + \|y_{\text{ref}}\|_Z^2 + \|d\|_H^2 \} \quad (52)$$

for some $k_\rho > 0$ independent of $[w_0, \eta_0]$ and $[d, y_{\text{ref}}]$. It follows from (52) that

$$\|[w(t), \eta(t)]\|_{H \times Z, \rho}^2 \leq \exp(k_\rho t) \|[w_0, \eta_0]\|_{H \times Z, \rho}^2 + \exp(k_\rho t) \{ \|y_{\text{ref}}\|_Z^2 + \|d\|_H^2 \} \quad (53)$$

for all $t \in [0, 1]$. As a consequence of (53), there exists a positive constant r_ρ such that $\max\{\|w_0\|_H, \|\eta_0\|_Z, \|d\|_H, \|y_{\text{ref}}\|_Z\} \leq r_\rho$ implies $\|w(t)\|_H \leq \tilde{r}$ for all $t \in [0, 1]$, where \tilde{r} is defined in Lemma 4.1. Therefore, for such initial data, we can use (39) in (32) to refine our previous estimate: for all positive ε and μ , on $(0, 1)$ we have

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|[w, \eta]\|_{H \times Z, \rho}^2 &\leq -\alpha \|w\|_H^2 + \frac{1}{2\varepsilon} \|B\|_{\mathcal{L}(U,H)}^2 \|\mathcal{K}(w)^* \eta\|_U^2 + \varepsilon \|w\|_H^2 \\ &\quad + \frac{1}{2\varepsilon} \|d\|_H^2 - \rho \|\mathcal{K}(w)^* \eta\|_U^2 + \frac{\rho}{2\mu} \|y_{\text{ref}}\|_Z^2 + \frac{\rho}{2\mu} \|d\mathcal{M}(w)d\|_Z^2 + \rho\mu \|\eta\|_Z^2. \end{aligned} \quad (54)$$

First, recall that we have either $d\mathcal{M}$ globally Lipschitz continuous or $d = 0$. Hence, there exists $k > 0$ independent of $[w_0, \eta_0]$ and $[d, y_{\text{ref}}]$ such that $\|d\mathcal{M}(w)d\|_Z^2 \leq k \|d\|_H^2$. By choosing $\varepsilon = \alpha/2$ and $\mu = \tilde{\lambda}/4$, we deduce from (54) the following differential inequality, valid on $(0, 1)$:

$$\frac{1}{2} \frac{d}{dt} \|[w, \eta]\|_{H \times Z, \rho}^2 \leq -\frac{\alpha}{2} \|w\|_H^2 - \frac{\rho \tilde{\lambda}}{4} \|\eta\|_Z^2 + \frac{\rho}{2\mu} \|y_{\text{ref}}\|_Z^2 + \left\{ \frac{\rho k}{2\mu} + \frac{1}{2\varepsilon} \right\} \|d\|_H^2 \quad (55)$$

Applying Grönwall's inequality to (55) yields

$$\begin{aligned} \|[w(t), \eta(t)]\|_{H \times Z, \rho}^2 &\leq \exp(-\kappa t) \|[w_0, \eta_0]\|_{H \times Z, \rho}^2 + K_\rho (1 - \exp(-\kappa t)) \| [d, y_{\text{ref}}] \|_{H \times Z}^2 \end{aligned} \quad (56)$$

for all $t \in [0, 1]$, where κ is defined in (42) and $K_\rho \geq 1$ is some constant independent of $[w_0, \eta_0]$ and $[d, y_{\text{ref}}]$. Next, by norm equivalence, there exists $r_{1,\rho} > 0$ such that the following implication holds: if $[w_0, \eta_0]$ and $[d, y_{\text{ref}}]$ are in $\mathcal{B}_{H \times Z}(0, r_{1,\rho})$, then

$$\|[w_0, \eta_0]\|_{H \times Z, \rho} \leq 2^{-1/2} r_\rho, \quad \text{and} \quad \|[d, y_{\text{ref}}]\|_{H \times Z} \leq (2K_\rho)^{-1/2} r_\rho. \quad (57)$$

Now, we claim that the estimate (56) remains valid for all $t \geq 0$. Indeed, (56) shows that $\|[w(1), \eta(1)]\|_{H \times Z, \rho} \leq r_\rho$. Therefore, by definition of r_ρ , we infer from the estimate (53) applied to the initial data $[w(1), \eta(1)]$ that $\|w(t)\|_H \leq \tilde{r}$ for all $t \in [1, 2]$. As a consequence, the differential inequality (55) is valid on $(0, 2)$; hence, (56) holds on $[0, 2]$, with in particular $\|[w(2), \eta(2)]\|_{H \times Z, \rho} \leq r_\rho$, and so on. The conclusion readily follows by induction.

Moreover, if it is assumed that (39) holds for all $w \in H$, then (55) is valid on $(0, +\infty)$ whatever the initial condition, so that (53) immediately holds for all $t \geq 0$. In this case, no additional condition on $[d, y_{\text{ref}}]$ is required. \square

4.2.2 Proof of Theorem 3.2

We can now prove the main result.

Step 1: Setting all neighborhoods. Pick ρ such that $\rho \geq \max\{\rho_0, \rho_1\}$ as in Lemmas 4.2 and 4.3. In the sequel, $[d, y_{\text{ref}}]$ is assumed to lie in the intersection of $\mathcal{B}_H(0, r_{0,\rho}) \times Z$ and $\mathcal{B}_{H \times Z}(0, r_{1,\rho})$, so that the lemmas apply. Now that ρ is fixed, we will omit the dependence on ρ in further notation. Lemma 4.2

provides a neighborhood \mathfrak{K} of the origin in $H \times Z$, which we will refer to as the contraction region, where (43) holds. On the other hand, according to (50) in Lemma 4.3, the set

$$\mathfrak{Y}^{d, y_{\text{ref}}} \triangleq \mathcal{B}_{H \times Z, \rho}(0, K_\rho^{1/2} \|[d, y_{\text{ref}}]\|_{H \times Z}) \quad (58)$$

attracts all solutions to (26) originating from $\mathcal{B}_{H \times Z}(0, r_{1, \rho})$. By norm equivalence, any sufficiently small ball for the ρ -norm that is centered at the origin is contained in both \mathfrak{K} and $\mathcal{B}_{H \times Z}(0, r_{1, \rho})$. That being said, as a consequence of (50) and (58), there exists positive numbers δ and r such that, having let $\mathcal{B} \triangleq \mathcal{B}_{H \times Z, \rho}(0, \delta)$, the following properties hold for any $[d, y_{\text{ref}}] \in \mathcal{B}_{H \times Z}(0, r)$:

- The closure of the corresponding attracting set $\mathfrak{Y}_{d, y_{\text{ref}}}$ is contained in \mathcal{B} ;
- Solutions to (26) with initial data in \mathcal{B} remain in \mathfrak{K} .

If in addition we assume that (14) holds, i.e., (39) holds globally, then by Lemma 4.3,

- The set $\mathfrak{Y}_{d, y_{\text{ref}}}$ attracts all solutions to (26), whatever the initial data.

In what follows, we omit the dependence on $[d, y_{\text{ref}}]$ in the notation and we denote by $\{\tilde{\mathcal{T}}_t\}$ the strongly continuous semigroup associated with (26). Then, since solutions originating from (the non-empty open set) \mathcal{B} remain in the contraction region \mathfrak{K} , we infer from (43) together with a density argument that

$$\|\tilde{\mathcal{T}}_t[w_1, \eta_1] - \tilde{\mathcal{T}}_t[w_2, \eta_2]\|_{H \times Z, \rho} \leq \exp(-\kappa t) \|[w_1, \eta_1] - [w_2, \eta_2]\|_{H \times Z, \rho} \quad (59)$$

for all $t \geq 0$ and $[w_i, \eta_i] \in \mathcal{B}$, $i \in \{1, 2\}$.

Step 2: Existence of a fixed point. Pick an arbitrary $[w_0, \eta_0] \in \mathcal{B}$. By (59) and a usual contraction argument, we see that

$$\{\tilde{\mathcal{T}}_n[w_0, \eta_0]\}_{n \geq 0} \text{ is a Cauchy sequence in } H \times Z \quad (60)$$

and converges to a fixed point $[w_0^*, \eta_0^*]$ of the nonlinear operator $\tilde{\mathcal{T}}_1$. Now, consider the ω -limit set $\omega([w_0, \eta_0])$ of $[w_0, \eta_0]$ with respect to the evolution semigroup $\{\tilde{\mathcal{T}}_t\}$. By the sequential characterization of ω -limit sets (see [5, Lemma 2.1, p.19]), we observe that $[w_0^*, \eta_0^*] \in \omega([w_0, \eta_0])$, which means that $\omega([w_0, \eta_0])$ is non-empty. Moreover, it is positively invariant by definition. Finally, since $\mathfrak{Y}_{d, y_{\text{ref}}}$ attracts all solutions originating from \mathcal{B} , we must have

$$\omega([w_0, \eta_0]) \subset \overline{\mathfrak{Y}_{d, y_{\text{ref}}}} \subset \mathcal{B}. \quad (61)$$

By following *verbatim*³ the proof of [8, Theorem 1], we obtain that for each $t \geq 0$, $\tilde{\mathcal{T}}_t$ is an isometry on $\omega([w_0, \eta_0])$. On the other hand, for positive t , $\tilde{\mathcal{T}}_t$ is a strict contraction on $\omega([w_0, \eta_0])$; thus, $\omega([w_0, \eta_0])$ must be reduced to the singleton $\{[w_0^*, \eta_0^*]\}$. By invariance of the ω -limit set, $[w_0^*, \eta_0^*]$ is fixed by the semigroup $\{\tilde{\mathcal{T}}_t\}$. Moreover, it follows from (59) that $[w_0^*, \eta_0^*]$ is the unique fixed point of $\{\tilde{\mathcal{T}}_t\}$ in \mathcal{B} and is exponentially attractive in \mathcal{B} . Thus, we write $[w^*, \eta^*] \triangleq [w_0^*, \eta_0^*]$.

Step 3: The fixed point lies in the domain. We now prove that $w^* \in \mathcal{D}(\mathcal{A})$. Let $\varepsilon > 0$ sufficiently small and consider the ball $\mathcal{C} \triangleq \mathcal{B}_{H \times Z, \rho}([w^*, \eta^*], \varepsilon) \subset \mathfrak{K}$. Since $\mathcal{D}(\mathcal{A})$ is dense in H and \mathcal{C} has non-empty interior, we can pick some $[w_0, \eta_0] \in [\mathcal{D}(\mathcal{A}) \times Z] \cap \mathcal{C}$. Let $[w(t), \eta(t)] \triangleq \tilde{\mathcal{T}}_t[w_0, \eta_0]$. As a strong solution to (26), $[w, \eta]$ is differentiable a.e. in $H \times Z$ and

$$\frac{d}{dt}[w, \eta] + \tilde{\mathcal{A}}_{d, y_{\text{ref}}}[w, \eta] = 0 \quad \text{a.e.} \quad (62)$$

Besides, we infer from (59) that \mathcal{C} is positively invariant so that $\{\tilde{\mathcal{T}}_t\}$ restricted to \mathcal{C} is still a well-defined contraction semigroup. Thus, we can apply [7, Theorem 1.4] to obtain

$$\|\tilde{\mathcal{A}}_{d, y_{\text{ref}}}[w, \eta]\|_{H \times Z, \rho} \leq \|\tilde{\mathcal{A}}_{d, y_{\text{ref}}}[w(t_0), \eta(t_0)]\|_{H \times Z, \rho} \quad \text{a.e. on } (t_0, +\infty) \quad (63)$$

³The only difference with [8, Theorem 1] is that $\{\tilde{\mathcal{T}}_t\}$ is a contraction only on a region (containing the ω -limit set) that is not *a priori* positively invariant. However, in order to obtain the isometry property, contraction is only needed on the points of the ω -limit set.

for some $t_0 \geq 0$. In particular, $\|\mathcal{A}(w)\|_H$ is bounded a.e. on $(t_0, +\infty)$. On the other hand, $w(t)$ converges to w^* in H when t goes to $+\infty$. Therefore, it follows from \mathcal{A} being maximal monotone and [7, Lemma 2.3] that $w^* \in \mathcal{D}(\mathcal{A})$.

Step 4: Conclusion. Since $[w^*, \eta^*]$ belongs to $\mathcal{D}(\mathcal{A}) \times Z$ and is fixed by $\{\tilde{\mathcal{T}}_t\}$, $\tilde{\mathcal{A}}_{d, y_{\text{ref}}}[w^*, \eta^*] = 0$. We come back to the original $[w, z]$ coordinates by letting $z^* \triangleq \eta^* + \mathcal{M}(w^*)$ and $\mathcal{N} \triangleq \{[w, \eta + \mathcal{M}(w)], [w, \eta] \in \mathcal{B}\}$. Then, $[w^*, z^*]$ belongs to $[\mathcal{D}(\mathcal{A}) \times Z] \cap \mathcal{N}$ and is an equilibrium for (4)-(9); hence, $Cw^* = y_{\text{ref}}$. Because \mathcal{M} vanishes at 0 and is continuous, \mathcal{N} is indeed a neighborhood of 0. Local exponential stability of $[w^*, z^*]$ with decay rate κ and basin of attraction containing \mathcal{N} follows from (59) and our prior remarks regarding the change of coordinates; the constant M in (3) comes from (27) and equivalence with the ρ -norm. Additionally, under the stronger condition (14), $[w^*, z^*]$ is globally asymptotically stable. The proof is now complete.

4.3 Proof of Theorem 3.4 (semilinear case)

We first give some auxiliary results valid under the hypotheses of Section 3.3, and then we prove Theorem 3.4.

4.3.1 Preliminaries

We endow $\mathcal{D}(A)$ with the graph norm. First, we claim that *weak* solutions (in the sense of nonlinear semigroup theory) to

$$\frac{dw}{dt} + \mathcal{A}(w) = 0. \quad (64)$$

coincide with *mild* solutions (in the sense of perturbation of linear equation). In other words, solutions $t \mapsto \mathcal{T}_t w_0$ to (64) with initial data $w_0 \in H$ are characterized by

$$\mathcal{T}_t w_0 = \mathcal{S}_t w_0 + \int_0^t \mathcal{S}_{t-s} F(\mathcal{T}_s w_0) ds, \quad \forall t \geq 0. \quad (65)$$

Because F is continuously differentiable, *strong* solutions w to (64) enjoy the regularity $w \in \mathcal{C}^1(\mathbb{R}_+, H) \cap \mathcal{C}(\mathbb{R}_+, \mathcal{D}(A))$. Lemma 4.4 given below allows us to circumvent the possible unboundedness of the output operator C and, together with the A -boundeness of C , guarantees that \mathcal{M} given by the limit in (19) is a well-defined nonlinear mapping on the whole space H .

Lemma 4.4 (Integral formula). *For any $w_0 \in H$,*

$$\lim_{\tau \rightarrow +\infty} \int_0^\tau \mathcal{T}_t w_0 dt = A^{-1} w_0 + A^{-1} \int_0^{+\infty} F(\mathcal{T}_t w_0) dt. \quad (66)$$

In particular, $\lim_{\tau \rightarrow +\infty} \int_0^\tau \mathcal{T}_t(w_0) dt$ belongs to $\mathcal{D}(A)$.

Proof. Let $w_0 \in H$. Recall from standard linear semigroup theory that the vector-valued integral $\int_0^\tau \mathcal{S}_t w_0 dt$ belongs to $\mathcal{D}(A)$ and $A \int_0^\tau \mathcal{S}_t w_0 dt = w_0 - \mathcal{S}_\tau w_0$. Thus, by integrating the variation of the constant formula (65) over $(0, \tau)$, $\tau \geq 0$, we obtain

$$\int_0^\tau \mathcal{T}_t w_0 dt = A^{-1} w_0 - A^{-1} \mathcal{S}_\tau w_0 + \int_0^\tau \int_0^t \mathcal{S}_{t-s} F(\mathcal{T}_s w_0) ds dt. \quad (67)$$

Let $\tau \geq 0$. Then, by letting $I_\tau \triangleq \{(s, t) \in \mathbb{R}^2, 0 \leq s \leq t \leq \tau\}$, we can write the last term in (67) as an absolutely convergent integral over \mathbb{R}^2 :

$$\int_0^\tau \int_0^t \mathcal{S}_{t-s} F(\mathcal{T}_s w_0) ds dt = \iint_{\mathbb{R}^2} \mathbb{1}_{I_\tau}(s, t) \mathcal{S}_{t-s} F(\mathcal{T}_s w_0) ds dt, \quad (68)$$

where $\mathbb{1}_{I_\tau}$ denotes the indicator function of the set I_τ . Switching the order of integration in the right-hand side of (68) leads to

$$\int_0^\tau \int_0^t \mathcal{S}_{t-s} F(\mathcal{T}_s w_0) ds dt = \int_0^{+\infty} \left\{ \mathbb{1}_{I_\tau}(s, s) \int_0^{\tau-s} \mathcal{S}_t F(\mathcal{T}_s w_0) dt \right\} ds. \quad (69)$$

We shall apply Lebesgue's dominated convergence theorem to conclude, the parameter being τ . For fixed $s \geq 0$, $\lim_{\tau \rightarrow +\infty} \mathbb{1}_{I_\tau} = 1$, and $\lim_{\tau \rightarrow +\infty} \mathcal{S}_\tau F(\mathcal{T}_s w_0) = 0$ in H ; hence,

$$\lim_{\tau \rightarrow +\infty} \mathbb{1}_{I_\tau}(s, s) \int_0^{\tau-s} \mathcal{S}_t F(\mathcal{T}_s w_0) dt = A^{-1} F(\mathcal{T}_s w_0). \quad (70)$$

On the other hand, by α -exponential stability of $\{\mathcal{S}_t\}$, for all $\tau \geq 0$ and $s \geq 0$,

$$\left\| \mathbb{1}_{I_\tau}(s, s) \int_0^{\tau-s} \mathcal{S}_t F(\mathcal{T}_s w_0) dt \right\|_H \leq \frac{1}{\alpha} \|F(\mathcal{T}_s w_0)\|_H, \quad (71)$$

Equation (71) provides an integrable dominating function, which ends the proof. \square

Now, let us come back to the uncontrolled w -equation (64), which we linearize around a given trajectory $\{\mathcal{T}_t w_0, t \geq 0\}$:

$$\frac{dv}{dt} + Av + dF(\mathcal{T}_t w_0)v = 0. \quad (72)$$

Equation (72) is linear but non-autonomous in general. Given $h \in H$, (72) possesses a unique mild solution $t \mapsto v(t) = v(t; w_0, h)$ satisfying $v(0) = h$ and $v \in \mathcal{C}(\mathbb{R}_+, H)$ (see [21, Theorem 1.2, p.184]). Given w_0 in H , the next lemma states that the nonlinear operators \mathcal{T}_t are all differentiable at w_0 and their differentials coincide with the evolution family associated with (72). This is a classical fact for sufficiently smooth nonlinear dynamics; here, we give a proof that matches our particular set of hypotheses.

Lemma 4.5 (Differentiability of the semigroup). *Each operator \mathcal{T}_t is Fréchet differentiable. Furthermore, for any $t \geq 0$ and $w_0 \in H$, the differential $d\mathcal{T}_t(w_0)$ is given by*

$$d\mathcal{T}_t(w_0)h = v(t; w_0, h) \quad \text{for all } h \in H, \quad (73)$$

where $t \mapsto v(t; w_0, h)$ is the unique mild solution to (72) with initial data h .

Proof. Let $\tau \geq 0$ and $w_0 \in H$. It is clear that the mapping $h \mapsto v(\tau; w_0, h)$ is linear; it is also continuous by [21, Theorem 1.2, p.184]. First, since F is differentiable, the following Taylor formula holds:

$$F(a+b) - F(a) = dF(a)b + R(a, b), \quad \forall a, b \in H, \quad (74)$$

with $R(a, b) = o(\|b\|_H)$ when $\|b\|_H \rightarrow 0$ for fixed $a \in H$. Now, take a nonzero $h \in H$. Combining the variation of the constant formula (65) with (74) leads to

$$\begin{aligned} \mathcal{T}_t(w_0 + h) - \mathcal{T}_t w_0 &= \mathcal{S}_t h - \int_0^t \mathcal{S}_{t-s} dF(\mathcal{T}_s w_0) \{\mathcal{T}_s(w_0 + h) - \mathcal{T}_s w_0\} ds \\ &\quad - \int_0^t \mathcal{S}_{t-s} R(\mathcal{T}_s w_0, \mathcal{T}_s(w_0 + h) - \mathcal{T}_s w_0) ds. \end{aligned} \quad (75)$$

Omitting the dependence on w_0 and h for the moment, we write $v(t) \triangleq v(t; w_0, h)$ and $\mathcal{R}(t) \triangleq \mathcal{T}_t(w_0 + h) - \mathcal{T}_t w_0 - v(t)$. Now, taking the difference between (75) and the integral identity satisfied by v as a mild solution to (72), one obtains

$$\mathcal{R}(t) = \int_0^t \mathcal{S}_{t-s} dF(\mathcal{T}_s w_0) \mathcal{R}(s) ds - \int_0^t \mathcal{S}_{t-s} R(\mathcal{T}_s w_0, \mathcal{T}_s(w_0 + h) - \mathcal{T}_s w_0) ds \quad (76)$$

holding for all $0 \leq t \leq \tau$. Next, because dF is continuous on the compact set $\{\mathcal{T}_s w_0, 0 \leq s \leq \tau\}$, $\|dF(\mathcal{T}_s w_0)\|_{\mathcal{L}(H)}$ is bounded by some m independent of h . Besides, $\|\mathcal{S}_s\|_{\mathcal{L}(H)} \leq 1$ for all $s \geq 0$. Therefore, it follows from (76) that for all $0 \leq t \leq \tau$,

$$\|\mathcal{R}(t)\|_H \leq m \int_0^t \|\mathcal{R}(s)\|_H ds + \int_0^t \|R(\mathcal{T}_s w_0, \mathcal{T}_s(w_0 + h) - \mathcal{T}_s w_0)\|_H ds. \quad (77)$$

Using Grönwall's inequality in its integral form, we deduce from (77) that

$$\frac{\|\mathcal{R}(\tau)\|_H}{\|h\|_H} \leq \exp(m\tau) \int_0^\tau \frac{\|R(\mathcal{T}_s w_0, \mathcal{T}_s(w_0 + h) - \mathcal{T}_s w_0)\|_H}{\|h\|_H} ds. \quad (78)$$

To obtain the desired differentiability property, it suffices to show that the right-hand side of (78) converges to 0 as $\|h\|_H$ goes to 0. This is done using Lebesgue's dominated convergence theorem. Let $s \in [0, \tau]$. By the contraction property of $\{\mathcal{T}_i\}$, for any nonzero $h \in H$ such that $\mathcal{T}_s(w_0 + h) - \mathcal{T}_s w_0$ is nonzero, we have

$$\frac{\|R(\mathcal{T}_s w_0, \mathcal{T}_s(w_0 + h) - \mathcal{T}_s w_0)\|_H}{\|h\|_H} \leq \frac{\|R(\mathcal{T}_s w_0, \mathcal{T}_s(w_0 + h) - \mathcal{T}_s w_0)\|_H}{\|\mathcal{T}_s(w_0 + h) - \mathcal{T}_s w_0\|_H}, \quad (79)$$

and if $\mathcal{T}_s(w_0 + h) - \mathcal{T}_s w_0 = 0$, then $R(\mathcal{T}_s w_0, \mathcal{T}_s(w_0 + h) - \mathcal{T}_s w_0) = 0$. Either way, when $\|h\|_H \rightarrow 0$, $\|\mathcal{T}_s(w_0 + h) - \mathcal{T}_s w_0\|_H \rightarrow 0$, and by definition of the residual term R , the right-hand side of (79) must converge to 0 as well. To conclude the proof, let us estimate the left-hand side of (79) uniformly with respect to h . We observe that the set of all points $\mathcal{T}_s(w_0 + h)$, $0 \leq s \leq \tau$, $\|h\|_H \leq 1$, is contained in some open ball, on which F is K -Lipschitz continuous for some $K > 0$. Thus, using (74), one obtains that the left-hand side of (79) is smaller than $2K$. \square

We continue by establishing exponential decay of solutions to (72).

Lemma 4.6 (Stability of the linearized equation). *Let $w_0 \in H$. For any $h \in H$, the solution $t \mapsto v(t; w_0, h)$ to (72) with initial data h satisfies*

$$\|v(t; w_0, h)\|_H \leq \exp(-\alpha t) \|h\|_H, \quad \forall t \geq 0. \quad (80)$$

Proof. Let w_0 and h in H . Let $\tau \geq 0$ and $f \triangleq t \mapsto dF(\mathcal{T}_t w_0)v(t) \in \mathcal{C}([0, \tau], H)$. Then, pick sequences $\{h_n\} \subset \mathcal{D}(A)$ and $\{f_n\} \subset W^{1,2}(0, \tau; H)$ such that $h_n \rightarrow h$ in H and $f_n \rightarrow f$ in $L^2(0, \tau; H)$. For each n , there exists a unique strong solution v_n to $dv_n/dt + Av_n + f_n(t) = 0$ satisfying the initial condition $v_n(0) = h_n$. Furthermore,

$$\frac{1}{2} \frac{d}{dt} \|v_n\|_H^2 = -(Av_n, v_n)_H - (f_n, h)_H \quad \text{a.e. on } (0, \tau), \quad (81)$$

and v_n converges to v in $\mathcal{C}([0, \tau], H)$. Plugging (18) into (81) yields

$$\frac{1}{2} \frac{d}{dt} \|v_n\|_H^2 \leq -\alpha \|v_n\|_H^2 + (dF(\mathcal{T}_t w_0)v_n - f_n, v_n)_H \quad \text{a.e. on } (0, \tau). \quad (82)$$

We deduce from (82) that for all $0 \leq t \leq \tau$,

$$\|v_n(t)\|_H^2 \leq \exp(-2\alpha t) \|h_n\|_H^2 + \frac{1}{2\alpha} \int_0^\tau |(dF(\mathcal{T}_s w_0)v_n(s) - f_n(s), v_n(s))_H| ds. \quad (83)$$

As n goes to $+\infty$, the integral term in (83) tends to 0 and we obtain the desired result by passing to the limit. \square

4.3.2 Proof of Theorem 3.4

Now that we have established that all objects in the statement of the theorem are well-defined, we can give the proof of the result. Recall that CA^{-1} is a bounded linear operator. In view of the formula (66), the desired properties of \mathcal{M} and $d\mathcal{M}$ readily follow from those of the mappings $w \mapsto \int_0^{+\infty} F(\mathcal{T}_t w) dt$ and $w \mapsto \int_0^{+\infty} dF(\mathcal{T}_t w) d\mathcal{T}_t(w) dt$, which we investigate next.

Step 1: Differentiability. Let $w_0 \in H$. It suffices to prove that

$$\int_0^{+\infty} \frac{\|F(\mathcal{T}_t(w_0 + h)) - F(\mathcal{T}_t w_0) - dF(\mathcal{T}_t w_0) d\mathcal{T}_t(w_0)h\|_H}{\|h\|_H} dt \rightarrow 0 \quad (84)$$

when $\|h\|_H$ goes to 0. We use Lebesgue's dominated convergence theorem. Since F and \mathcal{T}_t are differentiable, by the chain rule, the integrand in (84) converges to 0 pointwise. Let us now find some integrable dominating function. As in the proof of Lemma 4.5, we can find some open ball where F is K -Lipschitz and which contains the set of all $\mathcal{T}_t(w_0 + h)$, $\|h\|_H \leq 1$, $t \geq 0$. Thus, it follows from (3) and (80) that the integrand in (84) is dominated by $t \mapsto 2K \exp(-\alpha t)$, which is integrable.

Step 2: Lipschitz continuity of the differential. Pick two elements w_1 and w_2 in H . We write $R \triangleq \max\{\|w_1\|_H, \|w_2\|_H\}$. Subsequent estimates are motivated by the following decomposition:

$$\begin{aligned} & dF(\mathcal{T}_t w_1) d\mathcal{T}_t(w_1) - dF(\mathcal{T}_t w_2) d\mathcal{T}_t(w_2) \\ &= dF(\mathcal{T}_t w_1) [d\mathcal{T}_t(w_1) - d\mathcal{T}_t(w_2)] + [dF(\mathcal{T}_t w_1) - dF(\mathcal{T}_t w_2)] d\mathcal{T}_t(w_2). \end{aligned} \quad (85)$$

First, $\mathcal{T}_t w_1$ and $\mathcal{T}_t w_2$ must remain in $\mathcal{B}_H(0, R)$, where dF is, say, K_R -Lipschitz continuous. Since $dF(0) = 0$, for all $t \geq 0$ we have

$$\|dF(\mathcal{T}_t w_1)\|_{\mathcal{L}(H)} \leq K_R \|\mathcal{T}_t w_1\|_H \leq K_R \|w_1\|_H. \quad (86)$$

Now, let us estimate $d\mathcal{T}_t(w_1) - d\mathcal{T}_t(w_2)$ in operator norm. Pick $h \in H$. In what follows, we denote by $v(t)$ the difference $v(t) \triangleq d\mathcal{T}_t(w_1)h - d\mathcal{T}_t(w_2)h$. Then, $v(0) = 0$ and v is a (mild) solution to the following non-autonomous equation:

$$\frac{dv}{dt} + Av + [dF(\mathcal{T}_t w_1) d\mathcal{T}_t(w_1) - dF(\mathcal{T}_t w_2) d\mathcal{T}_t(w_2)]h = 0, \quad (87)$$

which can be rewritten as

$$\frac{dv}{dt} + Av + dF(\mathcal{T}_t w_1)v + [dF(\mathcal{T}_t w_1) - dF(\mathcal{T}_t w_2)]d\mathcal{T}_t(w_2)h = 0. \quad (88)$$

Justifications for the formal computations performed below are similar to those in the proof of Lemma 4.6; they are omitted here. Taking the scalar product in H of (88) with v and using (18) along with Cauchy-Schwarz and Young inequalities leads to

$$\frac{1}{2} \frac{d}{dt} \|v\|_H^2 \leq -\frac{\alpha}{2} \|v\|_H^2 + \frac{1}{2\alpha} \|[dF(\mathcal{T}_t w_1) - dF(\mathcal{T}_t w_2)]d\mathcal{T}_t(w_2)\|_{\mathcal{L}(H)}^2 \|h\|_H^2. \quad (89)$$

Since $v(0) = 0$, we deduce from (89) multiplied by $\exp(\alpha t)$, (80) and (3) that

$$\|v(t)\|_H^2 \leq \frac{K_R^2 \exp(-\alpha t)}{3\alpha^2} \|w_1 - w_2\|_H^2 \|h\|_H^2, \quad \forall t \geq 0. \quad (90)$$

We infer from (86) and (90) that

$$\|dF(\mathcal{T}_t w_1) [d\mathcal{T}_t(w_1) - d\mathcal{T}_t(w_2)]\|_{\mathcal{L}(H)} \leq m R K_R^2 \exp(-\alpha t/2) \|w_1 - w_2\|_H \quad (91)$$

for all $t \geq 0$, where m is some constant independent of R ; on the other hand, coming back to the second term of (85), we also have

$$\|[dF(\mathcal{T}_t w_1) - dF(\mathcal{T}_t w_2)]d\mathcal{T}_t(w_2)\|_{\mathcal{L}(H)} \leq K_R \exp(-2\alpha t) \|w_1 - w_2\|_H. \quad (92)$$

Thus, the desired local Lipschitz continuity is obtained by applying the triangular inequality to (85) and integrating (91) and (92) over $(0, +\infty)$. Furthermore, if we assume that both F and dF are globally Lipschitz continuous, then for some K_F we can choose $K_R = K_F$ independent of R and replace (86) with $\|dF(\mathcal{T}_t w_1)\|_{\mathcal{L}(H)} \leq K_F$, thereby proving global Lipschitz continuity of the differential $d\mathcal{M}$.

Step 3: Conclusion. At this point, it remains to check that our candidate \mathcal{M} is a solution to (7). It is clear that $\mathcal{M}(0) = 0$. Take w_0 in $\mathcal{D}(\mathcal{A})$ and consider the associated strong solution $w \triangleq t \mapsto \mathcal{T}_t w_0 \in \mathcal{C}(\mathbb{R}_+, \mathcal{D}(\mathcal{A})) \cap \mathcal{C}^1(\mathbb{R}_+, H)$ to (64). Then,

$$\frac{\mathcal{M}(\mathcal{T}_t w_0) - \mathcal{M}(w_0)}{t} = C \left[\frac{1}{t} \int_0^t \mathcal{T}_s w_0 ds \right]. \quad (93)$$

As w is continuous in $\mathcal{D}(A)$, the term $t^{-1} \int_0^t \mathcal{T}_s w_0 ds$ converges to w_0 in $\mathcal{D}(A)$ when t approaches 0. Thus, C being A -bounded, the difference quotient in (93) converges to Cw_0 in Z . On the other hand, since $w \in \mathcal{C}^1(\mathbb{R}_+, H)$ and $\mathcal{M} \in \mathcal{C}^1(H, Z)$, by the chain rule, we have

$$\frac{d}{dt}[\mathcal{M}(\mathcal{T}_t w_0)] = -d\mathcal{M}(\mathcal{T}_t w_0)\mathcal{A}(\mathcal{T}_t w_0), \quad \forall t \geq 0. \quad (94)$$

Evaluating (94) at $t = 0$ yields $d\mathcal{M}(w_0) + Cw_0 = 0$ by uniqueness of the limit.

5 Examples

We provide two examples for which our previous results apply. In what follows, we use standard notation for real-valued Lebesgue and Sobolev spaces.

5.1 Sine-Gordon equation

Let ξ , γ , and L be positive constants. Let \mathcal{O} be a non-empty open subset of $\Omega \triangleq (0, L)$. Consider the following damped sine-Gordon equation with control acting on \mathcal{O} and homogeneous Dirichlet boundary conditions:

$$\partial_{tt}\theta + \xi \partial_t \theta - \partial_{xx}\theta + \gamma \sin(\theta) = \mathbf{1}_{\mathcal{O}}(x)u(x, t) \quad \text{in } (0, L) \times (0, +\infty), \quad (95a)$$

$$\theta(0, t) = \theta(L, t) = 0 \quad \text{for all } t \geq 0. \quad (95b)$$

In this example inspired by [26, Chapter IV], (95) may represent the voltage dynamics of the continuous limit case for coupled Josephson junctions, with the control u being proportional to the applied current. The uncontrolled dynamics generated by (95) are well-posed on the energy space $H \triangleq H_0^1(\Omega) \times L^2(\Omega)$. Using the state variable $w = [\theta, \partial_t \theta]$, we can recast (95) into a semilinear evolution problem on H as in Section 3.3. Letting $\mathcal{D}(A) \triangleq [H^2(\Omega) \times H_0^1(\Omega)] \cap H$ and $U \triangleq L^2(\mathcal{O})$, we define the unbounded linear operator A , the input operator B , and the nonlinear mapping F by

$$A[\theta, \zeta] \triangleq [-\zeta, -\partial_{xx}\theta + \xi\zeta + \gamma\theta] \quad \forall [\theta, \zeta] \in \mathcal{D}(A), \quad (96a)$$

$$Bu = [0, \mathbf{1}_{\mathcal{O}}u] \quad \forall u \in U, \quad (96b)$$

$$F[\theta, \zeta] \triangleq [0, \gamma \sin(\theta) - \gamma\theta] \quad \forall [\theta, \zeta] \in H. \quad (96c)$$

As an output, consider the Neumann trace at, say, $x = 0$:

$$y(t) = Cw(t) = \partial_x \theta(0, t), \quad (97)$$

which is modeled by an unbounded (but A -bounded) scalar-valued operator. That A^{-1} exists in $\mathcal{L}(H)$ can be proved using Riesz representation theorem in $H_0^1(\Omega)$. Besides, F is in $\mathcal{C}^1(H)$ and both F and dF are globally Lipschitz continuous, dF being given by

$$dF(\theta, \zeta)[h_1, h_2] = [0, \gamma \cos(\theta)h_1 - \gamma h_1], \quad \forall [\theta, \zeta], [h_1, h_2] \in H. \quad (98)$$

We equip H with a scalar product that is equivalent to the usual one:

$$([\theta_1, \zeta_1], [\theta_2, \zeta_2])_{H, \varepsilon} \triangleq \int_{\Omega} \partial_x \theta_1 \partial_x \theta_2 dx + \int_{\Omega} (\zeta_1 + \varepsilon \theta_1)(\zeta_2 + \varepsilon \theta_2) dx \quad (99)$$

where $\varepsilon \triangleq \min\{\xi/4, \lambda_1/(2\xi)\}$, with λ_1 being the optimal Poincaré inequality constant. Then, after some computations similar to [26, Section IV.1.2], we obtain

$$(Ah + dF(\theta, \eta)h, h)_{H, \varepsilon} \geq \frac{\varepsilon}{2} \|h\|_{H, \varepsilon}^2 - \gamma \lambda_1 \|h\|_{H, \varepsilon}^2, \quad \forall h \in \mathcal{D}(A), \forall [\theta, \zeta] \in H. \quad (100)$$

Therefore, Hypothesis 3.3 is satisfied as long as $\gamma < \varepsilon/(2\lambda_1)$. In that case, Theorem 3.4 provides a suitable solution \mathcal{M} to (7) with $d\mathcal{M}$ globally Lipschitz continuous, upon which a forwarding control

law can be built for the output regulation problem. Since the range of $CA^{-1}B$ is non-zero⁴ and hence \mathbb{R} , Theorem 3.2 guarantees the existence of a locally exponentially stable equilibrium for the closed-loop system with small reference and disturbance. Furthermore, following the discussion subsequent to Theorem 3.2, the global coercivity condition (14) holds whenever $\varepsilon/2(1 + \lambda_1) > \gamma$, in which case the theorem provides a globally asymptotically stable equilibrium.

5.2 A pre-stabilized Wilson-Cowan equation

The following example is inspired by the study of neural fields (see for instance [4]). Let Ω be a bounded domain in \mathbb{R}^n and \mathcal{O} be an open subset of Ω . Given a positive gain α , a kernel $k \in L^\infty(\Omega \times \Omega)$, and a smooth scalar nonlinearity s that has bounded derivative and vanishes at 0, consider the following non-local evolution equation:

$$\partial_t w(x, t) + \alpha w(x, t) + \int_{\Omega} k(x, \nu) s(w(\nu, t)) d\nu = \mathbb{1}_{\mathcal{O}}(x) u(x, t) \quad \text{on } \Omega \times (0, +\infty). \quad (101)$$

We look at the (vector-valued) output given by

$$y(t) = Cw(t) = w|_{\mathcal{O}}(t). \quad (102)$$

Having set $H \triangleq L^2(\Omega)$, we define two mappings K and F on H by $[Kw](x) \triangleq s'(0) \int_{\Omega} k(x, \nu) w(\nu) d\nu$ and $[F(w)](x) \triangleq \int_{\Omega} k(x, \nu) \{s(w(\nu)) - s'(0)w(\nu)\} d\nu$ for all $w \in H$. Then, K is a bounded linear operator, and F is continuously differentiable with F and dF globally Lipschitz continuous. We also let $A \triangleq \alpha \text{id} + K$ and $Bu \triangleq \mathbb{1}_{\mathcal{O}}u$ with $U \triangleq L^2(\mathcal{O})$. As an integral operator, K is compact; thus, 0 lies in the resolvent set of A except for a bounded and countable set of values for α . Here, A is continuous; therefore, we can choose $Z \triangleq \text{Range } A^{-1}B$, which as a closed subspace of H is a Hilbert space as well. By letting $C \triangleq \text{id}$, the condition (21) is automatically satisfied. Furthermore, $Z = H$ if and only if $\mathcal{O} = \Omega$. As for condition (18), we have

$$(Ah + dF(w)h, h)_H \geq (\alpha - M_{k,s}) \|h\|_H^2, \quad \forall w, h \in H, \quad (103)$$

where $M_{k,s} \triangleq \iint_{\Omega \times \Omega} |k(x, \nu) s'(\nu)|^2 dx d\nu$. Hypothesis 3.3 is satisfied whenever $\alpha > M_{k,s}$, while global uniform coercivity (14) holds provided that $\alpha > 2M_{k,s}$.

6 Conclusions

We finish our article with some comments on our results. We wish to point out two notable byproducts of our approach in terms of *robustness*.

In order to take into account nonlinear behavior of actuators (e.g., saturation) in the feedback loop, one may investigate stability properties of (4)-(9) when a nonlinearity g is applied to the input u . This is also relevant in applications where the control signal must satisfy some prescribed bound in norm (see e.g. [17]). Consider a Lipschitz continuous map g that vanishes at 0 and is *strongly* monotone in some neighborhood of 0. Then, the local (strict) contraction property of the $[w, \eta]$ -dynamics is preserved. This leaves room for a possible adaptation of Theorem 3.2 in the case of saturated or *a priori* bounded control.

We also believe that our framework provides tools to analyse the behavior of the closed-loop (4)-(9) under certain time-varying disturbances. Indeed, given $[d_0, y_{\text{ref}}]$ as in Theorem 3.2, to which we associate an equilibrium $[w^*, z^*]$, consider a disturbance of the form $d(t) = d_0 + d_1(t)$ with d_1 small in $L^2(0, +\infty; H)$. It can then be deduced from (43) that the system (4)-(9) is *incrementally* input-to-state stable in a neighborhood of $[w^*, z^*]$, allowing us to quantify the deviation from equilibrium due to the exogenous signal d_1 in terms of its L^2 -norm.

Finally, putting aside the problem of output regulation and following [16,17], we might be interested in stabilizing the cascade composed of the (actuated) w -subsystem and a more general z -subsystem

⁴Consider for instance the image by $CA^{-1}B$ of $\mathbb{1}_I$ where I is some segment contained in \mathcal{O} .

governed by $dz/dt = Sz + Cw$, where S is a skew-adjoint operator on Z . This would require to investigate a nonlinear Sylvester equation of the form

$$d\mathcal{M}(w)\mathcal{A}(w) + S\mathcal{M}(w) + Cw = 0, \quad \forall w \in \mathcal{D}(\mathcal{A}). \quad (104)$$

Under the condition that a sufficiently regular solution \mathcal{M} to (104) exists, the Lyapunov analysis performed in Section 2.2 remains valid, which is a good starting point for analysing stability of the new closed-loop with control law given by (9).

Acknowledgments

The authors would like to thank Vincent Andrieu, Daniele Astolfi and Christophe Prieur for many fruitful discussions.

References

- [1] Daniele Astolfi and Laurent Praly. Integral action in output feedback for multi-input multi-output nonlinear systems. *IEEE Transactions on Automatic Control*, 62(4):1559–1574, 2017.
- [2] Ismaila Balogoun, Swann Marx, and Daniele Astolfi. ISS Lyapunov strictification via observer design and integral action control for a Korteweg-de Vries equation. working paper or preprint, July 2021.
- [3] S. Benachour, V. Andrieu, L. Praly, and H. Hammouri. Forwarding design with prescribed local behavior. *IEEE Transactions on Automatic Control*, 58(12):3011–3023, 2013.
- [4] Ugo Boscain, Dario Prandi, Ludovic Sacchelli, and Giuseppina Turco. A bio-inspired geometric model for sound reconstruction. *The Journal of Mathematical Neuroscience*, 11(1):1–18, 2021.
- [5] Igor Chueshov. *Introduction to the theory of infinite-dimensional dissipative systems*. University lectures in contemporary mathematics. Acta Scientific Publ. House, Kharkiv, 2002.
- [6] Igor Chueshov, Matthias Eller, and Irena Lasiecka. On the attractor for a semilinear wave equation with critical exponent and nonlinear boundary dissipation. *Communications in Partial Differential Equations*, 27(9-10):1901–1951, January 2002.
- [7] Michael G. Crandall and Amnon Pazy. Semi-groups of nonlinear contractions and dissipative sets. *Journal of functional analysis*, 3(3):376–418, 1969.
- [8] C.M. Dafermos and M. Slemrod. Asymptotic behavior of nonlinear contraction semigroups. *Journal of Functional Analysis*, 13(1):97–106, May 1973.
- [9] E. Davison. A generalization of the output control of linear multivariable systems with unmeasurable arbitrary disturbances. *IEEE Transactions on Automatic Control*, 20(6):788–792, 1975.
- [10] E. Davison. The robust control of a servomechanism problem for linear time-invariant multivariable systems. *IEEE Transactions on Automatic Control*, 21(1):25–34, 1976.
- [11] Mattia Giaccagli, Daniele Astolfi, Vincent Andrieu, and Lorenzo Marconi. Sufficient conditions for global integral action via incremental forwarding for input-affine nonlinear systems. *IEEE Transactions on Automatic Control*, pages 1–1, 2021.
- [12] Alberto Isidori, Lorenzo Marconi, and Andrea Serrani. *Robust autonomous guidance: an internal model approach*. Springer Science & Business Media, 2003.
- [13] G. Kaliora and A. Astolfi. On the stabilization of feedforward systems with bounded control. *Systems & control letters*, 54(3):263–270, 2005.
- [14] Yukio Komura. Differentiability of nonlinear semigroups. *Journal of the Mathematical Society of Japan*, 21(3):375–402, 1969.

- [15] Winfried Lohmiller and Jean-Jacques E. Slotine. On contraction analysis for non-linear systems. *Automatica*, 34(6):683–696, 1998.
- [16] Swann Marx, Daniele Astolfi, and Vincent Andrieu. Forwarding-Lyapunov design for the stabilization of coupled ODEs and exponentially stable PDEs. working paper or preprint, November 2021.
- [17] Swann Marx, Lucas Brivadis, and Daniele Astolfi. Forwarding techniques for the global stabilization of dissipative infinite-dimensional systems coupled with an ODE. *Mathematics of Control, Signals, and Systems*, 33, July 2021.
- [18] Frederic Mazenc and Laurent Praly. Adding integrations, saturated controls, and stabilization for feedforward systems. *IEEE Transactions on Automatic Control*, 41(11):1559–1578, 1996.
- [19] L. Paunonen and S. Pohjolainen. Internal model theory for distributed parameter systems. *SIAM Journal on Control and Optimization*, 48(7):4753–4775, 2010.
- [20] Lassi Paunonen. Stability and robust regulation of passive linear systems. *SIAM Journal on Control and Optimization*, 57(6):3827–3856, 2019.
- [21] Amnon Pazy. *Semigroups of linear operators and applications to partial differential equations*, volume 44. Springer Science & Business Media, 2012.
- [22] Duy Phan and Lassi Paunonen. Finite-dimensional controllers for robust regulation of boundary control systems. *Mathematical Control & Related Fields*, 11(1):95–117, 2021.
- [23] S. Pohjolainen. Robust multivariable PI-controller for infinite dimensional systems. *IEEE Transactions on Automatic Control*, 27(1):17–30, 1982.
- [24] Seppo Pohjolainen. Robust controller for systems with exponentially stable strongly continuous semigroups. *Journal of Mathematical Analysis and Applications*, 111(2):622–636, 1985.
- [25] Ralph Edwin Showalter. *Monotone operators in Banach space and nonlinear partial differential equations*, volume 49. American Mathematical Society, 2013.
- [26] Roger Temam. *Infinite-dimensional dynamical systems in mechanics and physics*, volume 68. Springer Science & Business Media, 2012.
- [27] A. Terrand-Jeanne, V. Andrieu, V. D. S. Martins, and C.-Z. Xu. Adding integral action for open-loop exponentially stable semigroups and application to boundary control of PDE systems. *IEEE Transactions on Automatic Control*, 2019.