

Weighted biharmonic equations involving continuous potential under exponential nonlinear growth

Brahim Dridi ⁽¹⁾ and Rached Jaidane ⁽²⁾

⁽¹⁾ Department of Mathematics, El Manar Preparatory Institute for Engineering Studies Tunis, University of Tunis El Manar, Tunisia. Address e-mail: dridibr@gmail.com

⁽²⁾ Department of Mathematics, Faculty of Science of Tunis, University of Tunis El Manar, Tunisia. Address e-mail: rachedjaidane@gmail.com

Abstract. We deal with the following weighted biharmonic problem

$$\begin{cases} \Delta(w(x)\Delta u) + V(x)u = f(x, u) & \text{in } B \\ u > 0 & \text{in } B \\ u = \frac{\partial u}{\partial n} = 0 & \text{on } \partial B, \end{cases}$$

where B is the unit ball of \mathbb{R}^4 , the non-linearity is assumed to have critical exponential growth in view of Adams's type inequalities. The weight $w(x)$ is of logarithm type and the potential V is a positive continuous function on \overline{B} . It is proved that there is a nontrivial positive weak solution to this equation by the mountain Pass Theorem. We avoid the loss of compactness by proving a concentration compactness result and by a suitable asymptotic condition.

Keywords: Adams inequality, Moser-Trudinger's inequality, Nonlinearity of exponential growth, Mountain pass method, Compactness level.

2010 **Mathematics Subject classification:** 35J20, 35J30, 35K57, 35J60.

1 Introduction

In this paper, we study the following weighted problem

$$\begin{cases} L := \Delta(w(x)\Delta u) + V(x)u = f(x, u) & \text{in } B \\ u > 0 & \text{in } B \\ u = \frac{\partial u}{\partial n} = 0 & \text{on } \partial B, \end{cases} \quad (1.1)$$

where B is the unitary disk in \mathbb{R}^4 , $f(x, t)$ is continuous in $B \times \mathbb{R}$ and behaves like $\exp\{t^{\frac{2}{1-\beta}}\}$, $\beta \in (0, 1)$ as $t \rightarrow +\infty$, for some $\alpha > 0$. The weight $w(x)$ is given by

$$w(x) = \left(\log \frac{e}{|x|}\right)^{1-\beta}, \beta \in (0, 1). \quad (1.2)$$

The potential $V : \overline{B} \rightarrow \mathbb{R}$ is a positive continuous function satisfying some conditions.

Problems of critical exponential growth in second order elliptic equations without weight in dimension $N = 2$

$$-\Delta u = f(x, u) \text{ in } \Omega \subset \mathbb{R}^2.$$

have been studied considerably [2, 21, 25, 29].

In dimension $N \geq 2$, the critical exponential growth is given by the well known Trudinger-Moser inequality [31, 38]

$$\sup_{\int_{\Omega} |\nabla u|^N \leq 1} \int_{\Omega} e^{\alpha|u|^{\frac{N}{N-1}}} dx < +\infty \text{ if and only if } \alpha \leq \alpha_N,$$

where $\alpha_N = \omega_{N-1}^{\frac{1}{N-1}}$ with ω_{N-1} is the area of the unit sphere S^{N-1} in \mathbb{R}^N .

After that, a logarithmic Trudinger-Moser inequality was used in a crucial way in [27] to study the Liouville equation of the form

$$\begin{cases} -\Delta u = \lambda \frac{e^u}{\int_{\Omega} e^u} & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.3)$$

where Ω an open domain of \mathbb{R}^N , $N \geq 2$ and λ a positive parameter.

The equation (1.3) has a long history and has been derived in the study of multiple condensate solution in the Chern-Simons-Higgs theory[36, 37]. It also appeared in the study of Euler Flow [8, 9, 15, 23].

Later, the Trudinger-Moser inequality was improved to weighted inequalities [11, 12]. The influence of the weight in the Sobolev norm was studied as the compact embedding [24].

When the weight is of logarithmic type, Calanchi and Ruf [13] extend the Trudinger-Moser inequality and proved the following results in the weighted Sobolev space

$$W_{0,rad}^{1,N}(B, \rho) = cl\{u \in C_{0,rad}^{\infty}(B) \mid \int_B |\nabla u|^N \rho(x) dx < \infty\} :$$

Theorem 1.1 [12]

(i) Let $\beta \in [0, 1)$ and let ρ given by $\rho(x) = (\log \frac{1}{|x|})^{\beta}$, then

$$\int_B e^{|u|^{\gamma}} dx < +\infty, \quad \forall u \in W_{0,rad}^{1,N}(B, \rho), \text{ if and only if } \gamma \leq \gamma_{N,\beta} = \frac{N}{(N-1)(1-\beta)} = \frac{N'}{1-\beta}$$

and

$$\sup_{\substack{u \in W_{0,rad}^{1,N}(B, \rho) \\ \int_B |\nabla u|^N w(x) dx \leq 1}} \int_B e^{\alpha|u|^{\gamma_{N,\beta}}} dx < +\infty \Leftrightarrow \alpha \leq \alpha_{N,\beta} = N[\omega_{N-1}^{\frac{1}{N-1}}(1-\beta)]^{\frac{1}{1-\beta}}$$

where ω_{N-1} is the area of the unit sphere S^{N-1} in \mathbb{R}^N and N' is the Hölder conjugate of N .

(ii) Let ρ given by $\rho(x) = (\log \frac{e}{|x|})^{N-1}$, then

$$\int_B \exp\{e^{|u|^{\frac{N}{N-1}}}\} dx < +\infty, \quad \forall u \in W_{0,rad}^{1,N}(B, \rho)$$

and

$$\sup_{\substack{u \in W_{0,rad}^{1,N}(B,\rho) \\ \|u\|_\rho \leq 1}} \int_B \exp\{\beta e^{\omega \frac{1}{N-1} |u|^{\frac{N}{N-1}}}\} dx < +\infty \Leftrightarrow \beta \leq N,$$

where ω_{N-1} is the area of the unit sphere S^{N-1} in \mathbb{R}^N and N' is the Hölder conjugate of N .

These results opened the way to study second order weighted elliptic problems in dimension $N \geq 2$. We cite the work of Calanchi et al [14], ie the following problem

$$\begin{cases} -\nabla \cdot (\nu(x) \nabla u) = f(x, u) & \text{in } B \\ u > 0 & \text{in } B \\ u = 0 & \text{on } \partial B, \end{cases}$$

with the weight $\nu(x) = \log(\frac{e}{|x|})$ and where the function $f(x, t)$ is continuous in $B \times \mathbb{R}$ and behaves like $\exp\{e^{\alpha t^2}\}$ as $t \rightarrow +\infty$, for some $\alpha > 0$. Also, recently, Deng et al [17] and Zhang[40] studied the following problem

$$\begin{cases} -\operatorname{div}(\rho(x) |\nabla u|^{N-2} \nabla u) = f(x, u) & \text{in } B \\ u = 0 & \text{on } \partial B, \end{cases}$$

where $N \geq 2$, the function $f(x, t)$ is continuous in $B \times \mathbb{R}$ and behaves like $\exp\{e^{\alpha t^{\frac{N}{N-1}}}\}$ as $t \rightarrow +\infty$, for some $\alpha > 0$. The authors proved that there is a non-trivial solution to this problem using Mountain Pass theorem.

Also, we mention that Baraket et al [6] studied the following non-autonomous weighted elliptic equations

$$\begin{cases} -\operatorname{div}(\rho(x) |\nabla u|^{N-2} \nabla u) + \xi(x) |u|^{N-2} u = f(x, u) & \text{in } B \\ u > 0 & \text{in } B \\ u = 0 & \text{on } \partial B, \end{cases}$$

where B is the unit ball of \mathbb{R}^N , $N > 2$, $f(x, t)$ is continuous in $B \times \mathbb{R}$ and behaves like $\exp\{e^{\alpha t^{\frac{N}{N-1}}}\}$ as $t \rightarrow +\infty$, for some $\alpha > 0$. $\xi : B \rightarrow \mathbb{R}$ is a positive continuous function satisfying some conditions. The weight $\rho(x)$ is given by $\rho(x) = (\log \frac{e}{|x|})^{N-1}$.

The biharmonic equation in dimension $N > 4$

$$\Delta^2 u = f(x, u) \text{ in } \Omega \subset \mathbb{R}^N,$$

where the nonlinearity f has subcritical and critical polynomial growth of power less than $\frac{N+4}{N-4}$, have been extensively studied [7, 19, 22, 34].

For bounded domains $\Omega \subset \mathbb{R}^4$, in [1, 32] the authors proved the following Adams' inequality

$$\sup_{u \in S} \int_\Omega (e^{\alpha u^2} - 1) dx < +\infty \Leftrightarrow \alpha \leq 32\pi^2$$

where $S = \{u \in W_0^{2,2}(\Omega) \mid (\int_\Omega |\Delta u|^2 dx)^{\frac{1}{2}} \leq 1\}$, and for unbounded domains, $\Omega \subset \mathbb{R}^4$,

$$S = \{u \in W_0^{2,2}(\Omega) \mid (\int_B u^2 dx + 2 \int_B |\nabla u|^2 dx + \int_B |\Delta u|^2 w(x) dx)^{\frac{1}{2}} \leq 1\}.$$

This last result opened the way to study fourth-order problems with subcritical or critical nonlinearity. We cite the work of Sani [33]

$$\Delta^2 u + V(x)u = f(x, u) \text{ in } H^2(\mathbb{R}^4).$$

We study the existence of the nontrivial solutions when the nonlinear terms have the critical exponential growth in the sense of Adams' inequalities [39]. Our approach is variational methods such as the Mountain Pass Theorem with Palais-Smale condition combining with a concentration compactness result.

Let $\Omega \subset \mathbb{R}^4$, be a bounded domain and $w \in L^1(\Omega)$ be a nonnegative function. We introduce the Sobolev spaces

$$W_0^{2,2}(\Omega, w) = cl\{u \in C_0^\infty(\Omega) \mid \int_B |\Delta u|^2 w(x) dx < \infty\}.$$

We will focus on radial functions and consider the subspace

$$W_0^{2,2}(\Omega, w) = cl\{u \in C_{0,rad}^\infty(\Omega) \mid \int_B |\Delta u|^2 w(x) dx < \infty\}.$$

The choice of the weight and the space $W_{0,rad}^{2,2}(B, w)$ are motivated by the following exponential inequality.

Theorem 1.2 [39] *Let $\beta \in (0, 1)$ and let w given by (1.2), then*

$$\sup_{\substack{u \in W_{0,rad}^{2,2}(B, w) \\ \int_B |\Delta u|^2 w(x) dx \leq 1}} \int_B e^{\alpha |u|^{\frac{2}{1-\beta}}} dx < +\infty \iff \alpha \leq \alpha_\beta = 4[8\pi^2(1-\beta)]^{\frac{1}{1-\beta}} \quad (1.4)$$

Let $\gamma := \frac{2}{1-\beta}$, in view of inequality (1.4), we say that f has critical growth at $+\infty$ if there exists some $0 < \alpha_0 \leq \alpha_\beta$,

$$\lim_{s \rightarrow +\infty} \frac{|f(x, s)|}{e^{\alpha s^\gamma}} = 0, \quad \forall \alpha \leq \alpha_\beta \text{ such that } \alpha_0 < \alpha \leq \alpha_\beta \quad \text{and} \quad \lim_{s \rightarrow +\infty} \frac{|f(x, s)|}{e^{\alpha s^\gamma}} = +\infty, \quad \forall \alpha < \alpha_0. \quad (1.5)$$

To study the solvability of the problem (1), consider the space

$$E = \{u \in W_{0,rad}^{2,2}(B, w) \mid \int_B V(x)|u|^2 dx < +\infty\},$$

endowed with the norm

$$\|u\| = \left(\int_B w(x)|\Delta u|^2 dx + \int_B V(x)|u|^2 dx \right)^{\frac{1}{2}}. \quad (1.6)$$

We note that this norm is issued from the product scalar

$$\langle u, v \rangle = \int_B (\Delta u \cdot \Delta v w(x) + V(x)uv) dx.$$

Let us now state our results. In this paper, we always assume that the nonlinearities $f(x, t)$ satisfies these conditions:

(H₁) $f : B \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous, radial in x , and $f(x, t) = 0$, for $t \leq 0$.

(H₂) There exist $t_0 > 0, M > 0$ such that $0 < F(x, t) = \int_0^t f(x, s)ds \leq M|f(x, t)|$,
 $\forall t > t_0, \forall x \in B$.

(H₃) $0 < F(x, t) \leq \frac{1}{2}f(x, t)t, \forall t > 0, \forall x \in B$.

(H₄) $\limsup_{t \rightarrow 0} \frac{2F(x, t)}{t^2} < \lambda_1$ uniformly in x , where λ_1 is the first eigenvalue of the operator with Dirichlet boundary condition that is

$$\lambda_1 = \inf_{u \in E, u \neq 0} \frac{\|u\|^2}{\int_B |u|^2 dx}. \quad (1.7)$$

This eigenvalue λ_1 exists and the corresponding eigen function ϕ_1 is positive and belongs to $L^\infty(B)$ [18].

and the potential V is continuous on \overline{B} and verifies

(V₁) $V(x) \geq V_0 > 0$ in B for some $V_0 > 0$.

We say that u is a solution to the problem (1), if u is a weak solution in the following sense.

Definition 1.1 A function u is called a solution to (1) if $u \in E$ and

$$\int_B (w(x) \Delta u \Delta \varphi + V u \varphi) dx = \int_B f(x, u) \varphi dx, \quad \text{for all } \varphi \in E. \quad (1.8)$$

It is easy to see that seeking weak solutions of the problem (1) is equivalent to find nonzero critical points of the following functional on E :

$$J(u) = \frac{1}{2} \int_B |\Delta u|^2 w(x) dx + \frac{1}{2} \int_B V(x) u^2 dx - \int_B F(x, u) dx, \quad (1.9)$$

where $F(x, u) = \int_0^u f(x, t) dt$.

We prove the following result.

Theorem 1.3 *Assume that V is continuous and verifies (V_1) . Assume that the function f has a critical growth at $+\infty$ and satisfies the conditions (H_1) , (H_2) , (H_3) and (H_4) . If in addition f verifies the asymptotic condition*

$$(H5) \quad \lim_{t \rightarrow \infty} \frac{f(x, t)t}{e^{\alpha_0 t^\gamma}} \geq \gamma_0 \quad \text{uniformly in } x, \quad \text{with } \gamma_0 > \frac{1024(1-\beta)}{\alpha_0^{1-\beta}},$$

then the problem (1) has a nontrivial solution.

In general the study of fourth order partial differential equations is considered an interesting topic. The interest in studying such equations was stimulated by their applications in micro-electro-mechanical systems, phase field models of multi-phase systems, thin film theory, surface diffusion on solids, interface dynamics, flow in Hele-Shaw cells, see [16, 20, 30]. However many applications are generated by the weighted elliptic problems, such as the study of traveling waves in suspension bridges, radar imaging (see, for example [4, 26]).

In this article, in spite of a possible failure of the Palais-Smale compactness condition, we apply mini-max methods and construct suitable sequence of Adam's type to avoid this compactness loss.

This paper is organized as follows. In Section 2, we present some necessary preliminary knowledge about functional space. In section 3, we give some useful lemmas for the compactness analysis. In section 4, we prove that the energy J satisfies the two geometric properties, and the compactness condition but under a given level. Finally, we conclude with the proofs of the main results in section 5.

Through this paper, the constant C may change from line to another and we sometimes index the constants in order to show how they change.

2 Weighted Lebesgue and Sobolev Spaces setting

Let $\Omega \subset \mathbb{R}^N$, $N \geq 2$, be a bounded domain in \mathbb{R}^N and let $w \in L^1(\Omega)$ be a nonnegative function. To handle with weighted operator, we need to introduce some functional spaces $L^p(\Omega, w)$, $W^{m,p}(\Omega, w)$, $W_0^{m,p}(\Omega, w)$ and some of their properties that will be used later. Let $S(\Omega)$ be the set of all measurable real-valued functions defined on Ω and two measurable functions are considered as the same element if they are equal almost everywhere.

Following Drabek et al. and Kufner in [18, 24], the weighted Lebesgue space $L^p(\Omega, w)$ is defined as follows:

$$L^p(\Omega, w) = \left\{ u : \Omega \rightarrow \mathbb{R} \text{ measurable; } \int_{\Omega} w(x)|u|^p dx < \infty \right\},$$

for any real number $1 \leq p < \infty$.

This is a normed vector space equipped with the norm

$$\|u\|_{p,w} = \left(\int_{\Omega} w(x)|u|^p dx \right)^{\frac{1}{p}}.$$

For $w(x) = 1$, we find the standard Lebesgue space $L^p(\Omega)$ and its norm

$$\|u\|_p = \left(\int_{\Omega} |u|^p dx \right)^{\frac{1}{p}}.$$

For $m \geq 2$, let w be a given family of weight functions w_{τ} , $|\tau| \leq m$, $w = \{w_{\tau}(x) \mid x \in \Omega, |\tau| \leq m\}$. In [18], the corresponding weighted Sobolev space was defined as

$$W^{m,p}(\Omega, w) = \{u \in L^p(\Omega) \text{ such that } D^{\tau}u \in L^p(\Omega, w) \text{ for all } 1 \leq |\tau| \leq m\}$$

endowed with the following norm:

$$\|u\|_{W^{m,p}(\Omega, w)} = \left(\sum_{|\alpha| \leq m-1} \int_{\Omega} |D^{\alpha}u|^p dx + \sum_{|\alpha|=m} \int_{\Omega} |D^{\alpha}u|^p w(x) dx \right)^{\frac{1}{p}}.$$

If we also assume that $w(x) \in L^1_{loc}(\Omega)$, then $C_0^{\infty}(\Omega)$ is a subset of $W^{m,p}(\Omega, w)$ and we can introduce the space

$$W_0^{m,p}(\Omega, w)$$

as the closure of $C_0^{\infty}(\Omega)$ in $W^{m,p}(\Omega, w)$.

$L^p(\Omega, w)$ and $W^{m,p}(\Omega, w)$ are separable, reflexive Banach spaces provided that $w(x)^{-\frac{1}{p-1}} \in L^1_{loc}(\Omega)$. For $w(x) = 1$, one finds the standard Sobolev spaces $W^{m,p}(\Omega)$ and $W_0^{m,p}(\Omega)$.

Then the space $E = \{u \in W_{0,rad}^{2,2}(B, w) \mid \int_B V(x)|u|^2 dx < +\infty\}$ is a Banach and reflexive space provided (V_1) is satisfied. The space E is endowed with the norm

$$\|u\| = \left(\int_B w(x)|\Delta u|^2 dx + \int_B V(x)|u|^2 dx \right)^{\frac{1}{2}}$$

which is equivalent to the following norm (see lemma 1)

$$\|u\|_{W_{0,rad}^{2,2}(B, w)} = \left(\int_B u^2 dx + \int_B |\nabla u|^2 dx + \int_B |\Delta u|^2 w(x) dx \right)^{\frac{1}{2}}.$$

3 Preliminary for the compactness analysis

In this section, we will derive several technical lemmas for our use later. First we begin by the radial lemma.

Lemma 1 *Assume that V is continuous and verifies (V_1) .*

(i) *Let u be a radially symmetric function in $C_0^2(B)$. Then, we have*

(i) [39]

$$|u(x)| \leq \frac{1}{2\sqrt{2\pi}} \frac{|\log(\frac{e}{|x|})|^{1-\beta} - 1|^{\frac{1}{2}}}{\sqrt{1-\beta}} \int_B w(x) |\Delta u|^2 dx \leq \frac{1}{2\sqrt{2\pi}} \frac{|\log(\frac{e}{|x|})|^{1-\beta} - 1|^{\frac{1}{2}}}{\sqrt{1-\beta}} \|u\|^2.$$

(ii) $\int_B e^{|u|^\gamma} dx < +\infty, \forall u \in E.$

(iii) There exists a positive constant C such that for all $u \in E$

$$\int_B V|u|^2 dx \leq C\|u\|^2$$

and then the norms $\|\cdot\|$ and $\|u\|_{W_{0,rad}^{2,2}(B,w)} = \left(\int_B u^2 dx + \int_B |\nabla u|^2 dx + \int_B |\Delta u|^2 w(x) dx \right)^{\frac{1}{2}}$ are equivalents.

(iv) The following embedding is continuous

$$E \hookrightarrow L^q(B) \text{ for all } q \geq 1.$$

(v) E is compactly embedded in $L^q(B)$ for all $q \geq 1$.

Proof

(i) see [39]

(ii) Follows from (i) and by density.

(iii) By Poincaré inequality, for all $u \in W_{0,rad}^{1,2}(B)$

$$\int_B |u|^2 \leq C \int_B |\nabla u|^2.$$

Using the Green formula, we get

$$\int_B |\nabla u|^2 = \int_B \nabla u \nabla u = - \int_B u \Delta u + \underbrace{\int_{\partial B} u \frac{\partial u}{\partial n}}_{=0} \leq \left| \int_B u \Delta u \right|,$$

By Young inequality, we get for all $\varepsilon > 0$

$$\left| \int_B u \Delta u \right| \leq \frac{1}{2\varepsilon} \int_B |\Delta u|^2 + \frac{\varepsilon}{2} \int_B |u|^2 \leq \frac{1}{2\varepsilon} \int_B w(x) |\Delta u|^2 + \frac{\varepsilon}{2} \int_B |u|^2,$$

Hence

$$\left(1 - \frac{\varepsilon}{2} C^2\right) \int_B |\nabla u|^2 \leq \frac{1}{2\varepsilon} \int_B |\Delta u|^2 w(x),$$

Then,

$$\int_B u^2 dx + \int_B |\nabla u|^2 dx + \int_B |\Delta u|^2 w(x) dx)^{\frac{1}{2}} \leq C \int_B |\Delta u|^2 w(x) dx \leq C \|u\|^2$$

On the other hand, from (i) we have for all $u \in E$,

$$\int_B V|u|^2 dx \leq m \int_B |u|^2 dx \leq \frac{m}{8\pi^2} \|u\|^2 \int_B \frac{|\log(\frac{e}{|x|})|^{1-\beta} - 1}{1-\beta} dx \leq \frac{m}{4} \|u\|^2 \int_0^1 r^3 \frac{|\log(\frac{e}{r})|^{1-\beta} - 1}{1-\beta} dr \leq C \|u\|^2,$$

where $m = \max_{x \in \overline{B}} V(x)$. Then (ii) follows.

(iv) Since $w(x) \geq 1$, then following embedding are continuous

$$E \hookrightarrow W_{0,rad}^{2,2}(B, w) \hookrightarrow W_{0,rad}^{2,2}(B) \hookrightarrow L^q(B) \quad \forall q \geq 2.$$

We also have by the Hölder inequality,

$$\int_B |u| dx \leq \left(\int_B \frac{1}{V} dx \right)^{\frac{1}{2}} \left(\int_B V|u|^2 dx \right)^{\frac{1}{2}} \leq \left(\int_B \frac{1}{V} dx \right)^{\frac{1}{2}} \|u\|.$$

For any $1 < \beta_0 < 2$, there holds

$$\int_B |u|^{\beta_0} dx \leq \int_B (|u| + |u|^2) dx \leq \left(\int_B \frac{1}{V} dx \right)^{\frac{1}{2}} \|u\| + \frac{1}{V_0} \|u\|^2.$$

Thus, we get the continuous embedding $E \hookrightarrow L^q(B)$ for all $q \geq 1$.

(v) The above embedding is also compact. Indeed, let $u_k \subset E$ be a sequence such that $\|u_k\| \leq C$ for all k . Then $\|u_k\|_{W_{0,rad}^{2,2}(B, w)} \leq C$, for all k . On the other hand, we have the following compact embedding [18] $W_{0,rad}^{2,2}(B, w) \hookrightarrow W_{0,rad}^{1,2}(B) \hookrightarrow L^q$ for all q such that $1 \leq q < 4$, then up to a subsequence, there exists some $u \in W_{0,rad}^{2,2}(B, w)$, such that u_k convergent to u strongly in $L^q(B)$ for all q such that $1 \leq q < 4$. Without loss of generality, we may assume that

$$\begin{cases} u_k & \rightharpoonup u & \text{weakly in } E \\ u_k & \rightarrow u & \text{strongly in } L^1(B) \\ u_k(x) & \rightarrow u(x) & \text{almost everywhere in } B. \end{cases} \quad (3.1)$$

For $q > 1$, it follows from (3.1) and the continuous embedding $E \hookrightarrow L^p(B)$ ($p \geq 1$) that

$$\begin{aligned} \int_B |u_k - u|^q dx &= \int_B |u_k - u|^{\frac{1}{2}} |u_k - u|^{q-\frac{1}{2}} dx \\ &\leq \left(\int_B |u_k - u| dx \right)^{\frac{1}{2}} \left(\int_B |u_k - u|^{2q-1} dx \right)^{\frac{1}{2}} \\ &\leq C \left(\int_B |u_k - u| dx \right)^{\frac{1}{2}} \rightarrow 0. \end{aligned}$$

This concludes the lemma. □

Second, we give the following useful lemma.

Lemma 2 [21] Let $\Omega \subset \mathbb{R}^N$ be a bounded domain and $f : \overline{\Omega} \times \mathbb{R}$ a continuous function. Let $\{u_n\}_n$ be a sequence in $L^1(\Omega)$ converging to u in $L^1(\Omega)$. Assume that $f(x, u_n)$ and $f(x, u)$ are also in $L^1(\Omega)$. If

$$\int_{\Omega} |f(x, u_n)u_n| dx \leq C,$$

where C is a positive constant, then

$$f(x, u_n) \rightarrow f(x, u) \text{ in } L^1(\Omega).$$

In the sequel, we prove a concentration compactness result of Lions type.

Lemma 3 Let $(u_k)_k$ be a sequence in E . Suppose that, $\|u_k\| = 1$, $u_k \rightharpoonup u$ weakly in E , $u_k(x) \rightarrow u(x)$ a.e $x \in B$, and $u \neq 0$. Then

$$\sup_k \int_B e^{p \alpha_\beta |u_k|^\gamma} dx < +\infty, \text{ where } \alpha_\beta = 4[8\pi^2(1 - \beta)]^{\frac{1}{1-\beta}},$$

for all $1 < p < U(u)$ where $U(u)$ is given by:

$$U(u) := \begin{cases} \frac{1}{(1 - \|u\|^2)^{\frac{1}{2}}} & \text{if } \|u\| < 1 \\ +\infty & \text{if } \|u\| = 1 \end{cases}$$

Proof For $a, b \in \mathbb{R}$, $q > 1$. If q' its conjugate i.e. $\frac{1}{q} + \frac{1}{q'} = 1$ we have, by young inequality, that

$$e^{a+b} \leq \frac{1}{q} e^{qa} + \frac{1}{q'} e^{q'b}.$$

Also, we have

$$(1 + a)^q \leq (1 + \varepsilon)a^q + \left(1 - \frac{1}{(1 + \varepsilon)^{\frac{1}{q-1}}}\right)^{1-q}, \quad \forall a \geq 0, \quad \forall \varepsilon > 0 \quad \forall q > 1. \quad (3.2)$$

So, using (3.2), we get

$$\begin{aligned} |u_k|^\gamma &= |u_k - u + u|^\gamma \\ &\leq (|u_k - u| + |u|)^\gamma \\ &\leq (1 + \varepsilon)|u_k - u|^\gamma + \left(1 - \frac{1}{(1 + \varepsilon)^{\frac{1}{\gamma-1}}}\right)^{1-\gamma} |u|^\gamma \end{aligned}$$

which implies that

$$\begin{aligned} \int_B e^{p \alpha_\beta |u_k|^\gamma} dx &\leq \frac{1}{q} \int_B e^{pq \alpha_\beta (1 + \varepsilon) |u_k - u|^\gamma} dx \\ &\quad + \frac{1}{q'} \int_B e^{pq' \alpha_\beta \left(1 - \frac{1}{(1 + \varepsilon)^{\frac{1}{\gamma-1}}}\right)^{1-\gamma} |u|^\gamma} dx, \end{aligned}$$

for any $p > 1$. Since $(1 - \frac{1}{(1+\varepsilon)^{\frac{1}{\gamma-1}}})^{1-\gamma} \leq 1$, then

$$\frac{1}{q'} \int_B e^{pq' \alpha_\beta (1 - \frac{1}{(1+\varepsilon)^{\frac{1}{\gamma-1}}})^{1-\gamma} |u|^\gamma} dx \leq \frac{1}{q'} \int_B e^{pq' \alpha_\beta |u|^\gamma} dx = \frac{1}{q'} \int_B e^{((pq' \alpha_\beta)^{\frac{1}{\gamma}} |u|)^\gamma} dx.$$

From Lemma 1 (ii), the last integral is finite.

To complete the evidence, we have to prove that for every p such that $1 < p < U(u)$,

$$\sup_k \int_B e^{pq \alpha_\beta (1+\varepsilon) |u_k - u|^\gamma} dx < +\infty, \quad (3.3)$$

for some $\varepsilon > 0$ and $q > 1$.

In the following, we suppose that $\|u\| < 1$ and in the case of $\|u\| = 1$, the proof is similar.

When

$$\|u\| < 1$$

and

$$p < \frac{1}{(1 - \|u\|^2)^{\frac{\gamma}{2}}},$$

there exists $\nu > 0$ such that

$$p(1 - \|u\|^2)^{\frac{\gamma}{2}}(1 + \nu) < 1.$$

On the other hand, we have

$$\|u_k - u\|^2 = \|u_k\|^2 - \|u\|^2 + o(1) \text{ where } o(1) \rightarrow 0 \text{ as } k \rightarrow +\infty.$$

Then,

$$\|u_k - u\|^2 = 1 - \|u\|^2 + o(1),$$

so,

$$\lim_{k \rightarrow +\infty} \|u_k - u\|^\gamma = (1 - \|u\|^2)^{\frac{\gamma}{2}}.$$

Therefore, for every $\varepsilon > 0$, there exists $k_\varepsilon \geq 1$ such that

$$\|u_k - u\|^\gamma \leq (1 + \varepsilon)(1 - \|u\|^2)^{\frac{\gamma}{2}}, \quad \forall k \geq k_\varepsilon.$$

If we take $q = 1 + \varepsilon$ with $\varepsilon = \sqrt[3]{1 + \nu} - 1$, then $\forall k \geq k_\varepsilon$, we have

$$pq(1 + \varepsilon) \|u_k - u\|^\gamma \leq 1.$$

Consequently,

$$\begin{aligned} \int_B e^{pq \alpha_\beta (1+\varepsilon) |u_k - u|^\gamma} dx &\leq \int_B e^{(1+\varepsilon)pq \alpha_\beta (\frac{|u_k - u|}{\|u_k - u\|})^\gamma \|u_k - u\|^\gamma} dx \\ &\leq \int_B e^{\alpha_\beta (\frac{|u_k - u|}{\|u_k - u\|})^\gamma} dx \\ &\leq \sup_{\|u\| \leq 1} \int_B e^{\alpha_\beta |u|^\gamma} dx < +\infty. \end{aligned}$$

Now, (3.3) follows from (1.4). This complete the proof and lemma 3 is proved.

4 The variational formulation

Since the reaction term f is of critical exponential growth, there exist positive constants c_1 and c_2 such that

$$|f(x, t)| \leq c_1 \exp\{e^{c_2 t^2}\}, \quad \forall x \in B, \quad \forall t \in \mathbb{R} \quad (4.1)$$

and so, by using (H_1) , the functional J given by (1.9) is C^1 .

4.1 The mountain pass geometry of the energy

In the sequel, we prove that the functional J has a mountain pass geometry.

Proposition 4.1 *Assume that the hypothesis (H_1) , (H_2) , (H_3) , (H_4) , and (V_1) hold. Then*

- (i) *there exist $\rho, \beta_0 > 0$ such that $\mathcal{J}(u) \geq \beta_0$ for all $u \in E$ with $\|u\| = \rho$.*
- (ii) *Let φ_1 be a normalized eigenfunction associated to λ_1 in E . Then, $\mathcal{J}(t\varphi_1) \rightarrow -\infty$, as $t \rightarrow +\infty$.*

Proof It follows from the hypothesis (H_4) that there exists $t_0 > 0$ and there exists $\varepsilon \in (0, 1)$ such that

$$F(x, t) \leq \frac{1}{2} \lambda_1 (1 - \varepsilon_0) |t|^2, \quad \text{for } |t| < t_0. \quad (4.2)$$

Indeed,

$$\limsup_{t \rightarrow 0} \frac{2F(x, t)}{t^2} < \lambda_1$$

or

$$\inf_{\tau > 0} \sup\left\{ \frac{2F(x, t)}{t^2}, \quad 0 < t < \tau \right\} < \lambda_1$$

Since this inequality is strict, then there exists $\varepsilon_0 > 0$ such that

$$\inf_{\tau > 0} \sup\left\{ \frac{2F(x, t)}{t^2}, \quad 0 < t < \tau \right\} < \lambda_1 - \varepsilon_0,$$

hence, there exists $t_0 > 0$ such that

$$\sup\left\{ \frac{2F(x, t)}{t^2}, \quad 0 < t < t_0 \right\} < \lambda_1 - \varepsilon_0.$$

Hence

$$\forall |t| < t_0 \quad F(x, t) \leq \frac{1}{2} \lambda_1 (1 - \varepsilon_0) t^2.$$

From (H3) and (4.1) and for all $q > 2$, there exist a constant $C > 0$ such that

$$F(x, t) \leq C|t|^q e^{a t^\gamma}, \quad \forall |t| > t_1. \quad (4.3)$$

So

$$F(x, t) \leq \frac{1}{2}\lambda_1(1 - \varepsilon_0)|t|^2 + C|t|^q e^{a t^\gamma}, \quad \text{for all } t \in \mathbb{R}. \quad (4.4)$$

Since

$$\mathcal{J}(u) = \frac{1}{2}\|u\|^2 - \int_B F(x, u)dx,$$

we get

$$\mathcal{J}(u) \geq \frac{1}{2}\|u\|^2 - \frac{1}{2}\lambda_1(1 - \varepsilon_0)|t|^2 - C \int_B |u|^q e^{a u^\gamma} dx.$$

But $\lambda_1\|u\|_2^2 \leq \|u\|^2$ and from the Hölder inequality, we obtain

$$\mathcal{J}(u) \geq \frac{\varepsilon_0}{2}\|u\|^2 - C\left(\int_B e^{2a|u|^\gamma} dx\right)^{\frac{1}{2}}\|u\|_{2q}^q. \quad (4.5)$$

From the Theorem 1.2, if we choose $u \in E$ such that

$$2a\|u\|^\gamma \leq \alpha_\beta, \quad (4.6)$$

we get

$$\int_B e^{2a|u|^\gamma} dx = \int_B e^{2a\|u\|^\gamma \left(\frac{|u|}{\|u\|}\right)^\gamma} dx < +\infty.$$

On the other hand $\|u\|_{2q} \leq C\|u\|$ (Lemma 1), so

$$\mathcal{J}(u) \geq \frac{\varepsilon_0}{2}\|u\|^2 - C\|u\|^q,$$

for all $u \in E$ satisfying (4.6). Since $2 < q$, we can choose $\rho = \|u\| > 0$ as the maximum point of the function $g(\sigma) = \frac{\varepsilon_0}{2}\sigma^2 - C\sigma^q$ on the interval $[0, (\frac{\alpha_\beta}{2a})^{\frac{1}{\gamma}}]$ and $\beta_0 = g(\rho)$, $\mathcal{J}(u) \geq \beta_0 > 0$.

(ii) Let $\phi_1 \in E \cap L^\infty(B)$ be the normalized eigen function associated to the eigen-value defined by (1.7) ie such that $\|\phi_1\| = 1$. We define the function

$$\varphi(t) = J(t\phi_1) = \frac{t^2}{2}\|\phi_1\|^2 - \int_B F(x, t\phi_1)dx.$$

Then using (H₁), (H₂) and integrating, we get the existence of a constant $C > 0$ such that

$$F(x, t) \geq C e^{\frac{1}{M}t}, \quad \forall |t| \geq t_0.$$

In particular, for $p > 2$, there exists C such that

$$F(x, t) \geq C|t|^p - C \geq C|t|^p - C, \quad \forall t \in \mathbb{R}, \quad x \in B.$$

Hence,

$$\varphi(t) = J(t\phi_1) \leq \frac{t^2}{2} \|\phi_1\|^2 - |t|^p \|\phi_1\|_p - c_5 \rightarrow -\infty, \text{ as } t \rightarrow +\infty,$$

and it's easy to conclude.

4.2 The compactness level of the energy

The main difficulty in the variational approach to the critical growth problem is the lack of compactness. Precisely the global Palais-Smale condition does not hold. Eventually, some partial Palais-Smale condition still holds under a given level. In the following proposition, we identify the first compactness level of the energy.

Proposition 4.2 *Let J be the energy associated to the problem (1) defined by (1.9), and suppose that the conditions (V_1) , (H_1) , (H_2) , (H_3) and (H_4) are satisfied. If the function $f(x, t)$ satisfies the condition (1.6) for some $\alpha_0 > 0$, then the functional J satisfies the Palais-Smale condition $(PS)_c$ for any*

$$c < \frac{1}{2} \left(\frac{\alpha_\beta}{\alpha_0} \right)^{\frac{2}{\gamma}},$$

where $\alpha_\beta = 4[8\pi^2(1 - \beta)]^{\frac{1}{1-\beta}}$.

Proof Consider a $(PS)_c$ sequence in E , for some $c \in \mathbb{R}$, that is

$$J(u_n) = \frac{1}{2} \|u_n\|^2 - \int_B F(x, u_n) dx \rightarrow c, \quad n \rightarrow +\infty \quad (4.7)$$

and

$$|\langle J'(u_n), \varphi \rangle| = \left| \int_B w(x) \Delta u_n \cdot \Delta \varphi dx + \int_B V u_n \varphi dx - \int_B f(x, u_n) \varphi dx \right| \leq \varepsilon_n \|\varphi\|, \quad (4.8)$$

for all $\varphi \in E$, where $\varepsilon_n \rightarrow 0$, when $n \rightarrow +\infty$.

Also, inspired by [14], it follows from (H_2) that for all $\varepsilon > 0$ there exists $t_\varepsilon > 0$ such that

$$F(x, t) \leq \varepsilon t f(x, t), \quad \text{for all } |t| > t_\varepsilon \text{ and uniformly in } x \in B, \quad (4.9)$$

and so, by (4.7), for all $\varepsilon > 0$ there exists a constant $C > 0$

$$\frac{1}{2} \|u_n\|^2 \leq C + \int_B F(x, u_n) dx,$$

hence

$$\frac{1}{2} \|u_n\|^2 \leq C + \int_{|u_n| \leq t_\varepsilon} F(x, u_n) dx + \varepsilon \int_B f(x, u_n) u_n dx$$

and so, from (4.8), we get

$$\frac{1}{2}\|u_n\|^2 \leq C_1 + \varepsilon\varepsilon_n\|u_n\| + \varepsilon\|u_n\|^2,$$

for some constant $C_1 > 0$. Since

$$\left(\frac{1}{2} - \varepsilon\right)\|u_n\|^2 \leq C_1 + \varepsilon\varepsilon_n\|u_n\|, \quad (4.10)$$

we deduce that the sequence (u_n) is bounded in E . As consequence, there exists $u \in E$ such that, up to subsequence, $u_n \rightharpoonup u$ weakly in E , $u_n \rightarrow u$ strongly in $L^q(B)$, for all $q \geq 1$ and $u_n(x) \rightarrow u(x)$ a.e. in B .

Furthermore, we have, from (4.7) and (4.8), that

$$0 < \int_B f(x, u_n)u_n \leq C, \quad (4.11)$$

and

$$0 < \int_B F(x, u_n) \leq C. \quad (4.12)$$

Since by Lemma 2.1 in [21], we have

$$f(x, u_n) \rightarrow f(x, u) \text{ in } L^1(B) \text{ as } n \rightarrow +\infty, \quad (4.13)$$

then, it follows from (H_2) and the generalized Lebesgue dominated convergence theorem that

$$F(x, u_n) \rightarrow F(x, u) \text{ in } L^1(B) \text{ as } n \rightarrow +\infty. \quad (4.14)$$

So,

$$\lim_{n \rightarrow +\infty} \|u_n\|^2 = 2\left(c + \int_B F(x, u)dx\right). \quad (4.15)$$

Using (4.4), we have

$$\lim_{n \rightarrow \infty} \int_B f(x, u_n)u_n dx = 2\left(c + \int_B F(x, u)dx\right). \quad (4.16)$$

Then by (H_3) and (4.8), we get

$$\lim_{n \rightarrow +\infty} 2 \int_B F(x, u_n)dx \leq \lim_{n \rightarrow +\infty} \int_B f(x, u_n)u_n dx = 2\left(d + \int_B F(x, u)dx\right). \quad (4.17)$$

Also, it follows from (4.8), (4.9), (4.13) and (4.14), by passing to the limit, we obtain that u is a weak solution of the problem (1). Taking $\varphi = u$ as a test function, we get

$$\int_B |\Delta u|^2 w(x)dx + \int_B V u^2 dx = \int_B f(x, u)u dx \geq 2 \int_B F(x, u)dx.$$

Hence $J(u) \geq 0$. We also have by the Fatou's lemma and (4.14)

$$0 \leq J(u) \leq \frac{1}{2} \liminf_{n \rightarrow \infty} \|u_n\|^2 - \int_B F(x, u)dx = c.$$

So, we will finish the proof by considering three cases for the level c .

Case 1. $c = 0$. In this case

$$0 \leq J(u) \leq \liminf_{n \rightarrow +\infty} J(u_n) = 0.$$

So,

$$J(u) = 0$$

and then by (4.14)

$$\lim_{n \rightarrow +\infty} \frac{1}{2} \|u_n\|^2 = \int_B F(x, u) dx = \frac{1}{2} \|u\|^2.$$

It follows that $u_n \rightarrow u$ in E .

Case 2. $c > 0$ and $u = 0$. We prove that this case cannot happen.

From (4.7) and (4.8) with $v = u_n$, we have

$$\lim_{n \rightarrow +\infty} \|u_n\|^2 = 2c \quad \text{and} \quad \lim_{n \rightarrow +\infty} \int_B f(x, u_n) u_n dx = 2c.$$

Again by (4.11) we have

$$\left| \|u_n\|^2 - \int_B f(x, u_n) u_n dx \right| \leq C\varepsilon_n.$$

First we claim that there exists $q > 1$ such that

$$\int_B |f(x, u_n)|^q dx \leq C, \tag{4.18}$$

so

$$\|u_n\|^2 \leq C\varepsilon_n + \left(\int_B |f(x, u_n)|^q dx \right)^{\frac{1}{q}} \left(\int_B |u_n|^{q'} dx \right)^{\frac{1}{q'}}$$

where q' the conjugate of q . Since (u_n) converge to $u = 0$ in $L^{q'}(B)$,

$$\lim_{n \rightarrow +\infty} \|u_n\|^2 = 0$$

which in contradiction with $c > 0$.

For the proof of the claim, since f has subcritical or critical growth, for every $\varepsilon > 0$ and $q > 1$ there exists $t_\varepsilon > 0$ and $C > 0$ such that for all $|t| \geq t_\varepsilon$, we have

$$|f(x, t)|^q \leq C e^{\alpha_0(\varepsilon+1)t^\gamma}. \tag{4.19}$$

Consequently,

$$\begin{aligned} \int_B |f(x, u_n)|^q dx &= \int_{\{|u_n| \leq t_\varepsilon\}} |f(x, u_n)|^q dx + \int_{\{|u_n| > t_\varepsilon\}} |f(x, u_n)|^q dx \\ &\leq 2\pi^2 \max_{B \times [-t_\varepsilon, t_\varepsilon]} |f(x, t)|^q + C \int_B e^{\alpha_0(\varepsilon+1)|u_n|^\gamma} dx. \end{aligned}$$

Since $2c < (\frac{\alpha\beta}{\alpha_0})^{\frac{2}{\gamma}}$, there exists $\eta \in (0, \frac{1}{2})$ such that $2c = (1 - 2\eta)(\frac{\alpha\beta}{\alpha_0})^{\frac{2}{\gamma}}$. On the other hand, $\|u_n\|^\gamma \rightarrow (2c)^{\frac{\gamma}{2}}$, so there exists $n_\eta > 0$ such that for all $n \geq n_\eta$, we get $\|u_n\|^\gamma \leq (1 - \eta)\frac{\alpha\beta}{\alpha_0}$. Therefore,

$$\alpha_0(1 + \varepsilon)\left(\frac{|u_n|}{\|u_n\|}\right)^\gamma \|u_n\|^\gamma \leq (1 + \varepsilon)(1 - \eta)\alpha\beta.$$

We choose $\varepsilon > 0$ small enough to get

$$\alpha_0(1 + \varepsilon)\|u_n\|^\gamma \leq \alpha\beta.$$

Therefore, the second integral is uniformly bounded in view of (1.4) and the claim is proved.

Case 3. $c > 0$ and $u \neq 0$. In this case, we claim that $J(u) = c$ and therefore, we get

$$\lim_{n \rightarrow +\infty} \|u_n\|^2 = 2\left(c + \int_B F(x, u)dx\right) = 2(J(u) + \int_B F(x, u)dx) = \|u\|^2.$$

Do not forget that

$$J(u) \leq \frac{1}{2} \liminf_{n \rightarrow +\infty} \|u_n\|^2 - \int_B F(x, u)dx = c.$$

Suppose that $J(u) < c$. Then,

$$\|u\|^\gamma < \left(2\left(c + \int_B F(x, u)dx\right)\right)^{\frac{\gamma}{2}}. \quad (4.20)$$

Set

$$v_n = \frac{u_n}{\|u_n\|}$$

and

$$v = \frac{u}{\left(2\left(c + \int_B F(x, u)dx\right)\right)^{\frac{1}{2}}}.$$

We have $\|v_n\| = 1$, $v_n \rightharpoonup v$ in E , $v \neq 0$ and $\|v\| < 1$. So, by Lemma 3, we get

$$\sup_n \int_B e^{p\alpha\beta|v_n|^\gamma} dx < \infty,$$

for $1 < p < U(v) = (1 - \|v\|^2)^{-\frac{\gamma}{2}}$.

As in the case (2), we are going to estimate $\int_B |f(x, u_n)|^q dx$.

For $\varepsilon > 0$, one has

$$\begin{aligned} \int_B |f(x, u_n)|^q dx &= \int_{\{|u_n| \leq t_\varepsilon\}} |f(x, u_n)|^q dx + \int_{\{|u_n| > t_\varepsilon\}} |f(x, u_n)|^q dx \\ &\leq 2\pi^2 \max_{B \times [-t_\varepsilon, t_\varepsilon]} |f(x, t)|^q + C \int_B e^{\alpha_0(1+\varepsilon)|u_n|^\gamma} dx \\ &\leq C_\varepsilon + C \int_B e^{\alpha_0(1+\varepsilon)\|u_n\|^\gamma |v_2|^\gamma} dx \leq C, \end{aligned}$$

provided that $\alpha_0(1 + \varepsilon)\|u_n\|^\gamma \leq p \alpha_\beta$ and $1 < p < U(v) = (1 - \|v\|^2)^{\frac{-\gamma}{2}}$.

Since

$$(1 - \|v\|^2)^{\frac{-\gamma}{2}} = \left(\frac{2(c + \int_B F(x, u)dx)}{2(c + \int_B F(x, u)dx) - \|u\|^2} \right)^{\frac{\gamma}{2}} = \left(\frac{c + \int_B F(x, u)dx}{c - J(u)} \right)^{\frac{\gamma}{2}}$$

and

$$\lim_{n \rightarrow +\infty} \|u_n\|^\gamma = (2(c + \int_B F(x, u)dx))^{\frac{\gamma}{2}},$$

then,

$$\alpha_0(1 + \varepsilon)\|u_n\|^\gamma \leq \alpha_0(1 + 2\varepsilon)(2(c + \int_B F(x, u)dx))^{\frac{\gamma}{2}}.$$

But $J(u) \geq 0$ and $c < \frac{1}{2}(\frac{\alpha_{2,\beta}}{\alpha_0})^{\frac{2}{\gamma}}$, then if we choose $\varepsilon > 0$ small enough such that

$$\frac{\alpha_0}{\alpha_\beta}(1 + 2\varepsilon) < \left(\frac{1}{2(c - J(u))} \right)^{\frac{\gamma}{2}},$$

we get,

$$(1 + 2\varepsilon)((c - J(u))^{\frac{\gamma}{2}}) < \frac{\alpha_\beta}{2^{\frac{\gamma}{2}}\alpha_0}.$$

So, the sequence $(f(x, u_n))$ is bounded in L^q , $q > 1$.

Since $\langle J'(u_n), (u_n - u) \rangle = o(1)$, we have from the boundedness of $\{f(x, u_n)\}$ in $L^q(B)$ for $q > 1$, we can prove that $u_n \rightarrow u$ strongly in E . Indeed, we have

$$\|u_n - u\|^2 = \langle u_n, u_n - u \rangle - \langle u, u_n - u \rangle = \langle u_n, u_n - u \rangle + o_n(1) \rightarrow 0 \text{ as } n \rightarrow +\infty.$$

From (4.5) and using the Hölder inequality, we get

$$\begin{aligned} |\langle u_n, u_n - u \rangle| &\leq \varepsilon_n \|u_n - u\| + \left| \int_B f(x, u_n)(u_n - u)dx \right| \\ &\leq C\varepsilon_n + \left(\int_B |f(x, u_n)|^q dx \right)^{\frac{1}{q}} \left(\int_B |u_n - u|^{q'} dx \right)^{\frac{1}{q'}} \rightarrow 0 \text{ as } n \rightarrow +\infty. \end{aligned}$$

Hence,

$$\lim_{n \rightarrow +\infty} \|u_n\|^2 = 2(c + \int_B F(x, u)dx) = \|u\|^2$$

and this contradicts (4.20). So, $J(u) = c$ and consequently, $u_n \rightarrow u$.

5 Proof of the main results

In the sequel, we will estimate the minimax level of the energy J . We will prove that the mountain pass level c satisfies

$$c < \frac{1}{2} \left(\frac{\alpha_\beta}{\alpha_0} \right)^{\frac{2}{\gamma}}.$$

For this purpose, we will prove that there exists $z \in E$ such

$$\max_{t \geq 0} J(tz) < \frac{1}{2} \left(\frac{\alpha_\beta}{\alpha_0} \right)^{\frac{2}{\gamma}}. \quad (5.1)$$

5.1 Adams functions

Now, we will construct particular functions, namely the Adams functions. We consider the sequence defined for all $n \geq 3$ by

$$(5.2) \quad w_n(x) = \begin{cases} \left(\frac{\log(e\sqrt[4]{n})}{\alpha_\beta} \right)^{\frac{1}{\gamma}} - \frac{|x|^{2(1-\beta)}}{2\left(\frac{\alpha_\beta}{4n}\right)^{\frac{1}{\gamma}} (\log(e\sqrt[4]{n}))^{\frac{\gamma-1}{\gamma}}} + \frac{1}{2\left(\frac{\alpha_\beta}{4}\right)^{\frac{1}{\gamma}} (\log(e\sqrt[4]{n}))^{\frac{\gamma-1}{\gamma}}} & \text{if } 0 \leq |x| \leq \frac{1}{\sqrt[4]{n}} \\ \frac{\left(\log\left(\frac{e}{|x|}\right) \right)^{1-\beta}}{\left(\frac{\alpha_\beta}{4} \log(e\sqrt[4]{n}) \right)^{\frac{1}{\gamma}}} & \text{if } \frac{1}{\sqrt[4]{n}} \leq |x| \leq \frac{1}{2} \\ \zeta_n & \text{if } \frac{1}{2} \leq |x| \leq 1 \end{cases}$$

where $\zeta_n \in C_0^\infty(B)$ is such that

$$\zeta_n|_{x=\frac{1}{2}} = \frac{1}{\left(\frac{\alpha_\beta}{16} \log(e^4 n)\right)^{\frac{1}{\gamma}}} (\log 2e)^{1-\beta}, \quad \frac{\partial \zeta_n}{\partial x}|_{x=\frac{1}{2}} = \frac{-2(1-\beta)}{\left(\frac{\alpha_\beta}{4} \log(e\sqrt[4]{n})\right)^{\frac{1}{\gamma}}} (\log(2e))^{-\beta}$$

$$\zeta_n|_{\partial B} = \frac{\partial \zeta_n}{\partial x}|_{\partial B} = 0 \text{ and } \xi_n, \nabla \xi_n, \Delta \xi_n \text{ are all } o\left(\frac{1}{\log(e\sqrt[4]{n})}\right).$$

Let $v_n(x) = \frac{w_n}{\|w_n\|}$. We have, $v_n \in E$, $\|v_n\|^2 = 1$.

We compute $\Delta w_n(x)$, we get

$$\Delta w_n(x) = \begin{cases} \frac{-(1-\beta)(4-2\beta)|x|^{-2\beta}}{\left(\frac{\alpha_\beta}{4n}\right)^{\frac{1}{\gamma}} (\log(e\sqrt[4]{n}))^{\frac{\gamma-1}{\gamma}}} & \text{if } 0 \leq |x| \leq \frac{1}{\sqrt[4]{n}} \\ \frac{-(1-\beta) \left(\log\left(\frac{e}{|x|}\right) \right)^{-\beta} \left(2 + \beta \left(\log\left(\frac{e}{|x|}\right) \right)^{-1} \right)}{\left(\frac{\alpha_\beta}{4} \log(e\sqrt[4]{n}) \right)^{\frac{1}{\gamma}}} & \text{if } \frac{1}{\sqrt[4]{n}} \leq |x| \leq \frac{1}{2} \\ \Delta \zeta_n & \text{if } \frac{1}{2} \leq |x| \leq 1 \end{cases}$$

So,

$$\|\Delta w_n\|_{2,w}^2 = 2\pi^2 \underbrace{\int_0^{\frac{1}{\sqrt[4]{n}}} r^3 |\Delta w_n(x)|^2 \left(\log \frac{e}{r}\right)^\beta dr}_{I_1} + 2\pi^2 \underbrace{\int_{\frac{1}{\sqrt[4]{n}}}^{\frac{1}{2}} r^3 |\Delta w_n(x)|^2 \left(\log \frac{e}{r}\right)^\beta dr}_{I_2} + 2\pi^2 \underbrace{\int_{\frac{1}{2}}^1 r^3 |\Delta w_n(x)|^2 \left(\log \frac{e}{r}\right)^\beta dr}_{I_3}$$

we have,

$$\begin{aligned}
I_1 &= 2\pi^2 \frac{(1-\beta)^2(4-2\beta)^2}{\left(\frac{\alpha\beta}{4n}\right)^{\frac{2}{\gamma}} (\log(e\sqrt[4]{n}))^{\frac{2(\gamma-1)}{\gamma}}} \int_0^{\frac{1}{\sqrt[4]{n}}} r^{3-4\beta} \left(\log \frac{e}{r}\right)^\beta dr \\
&= 2\pi^2 \frac{(1-\beta)^2(4-2\beta)^2}{\left(\frac{\alpha\beta}{4n}\right)^{\frac{2}{\gamma}} (\log(e\sqrt[4]{n}))^{\frac{2(\gamma-1)}{\gamma}}} \left[\frac{r^{4-4\beta} (\log \frac{e}{r})^\beta}{4-4\beta} \right]_0^{\frac{1}{\sqrt[4]{n}}} \\
&+ 2\pi^2 \frac{\beta(1-\beta)^2(4-2\beta)^2}{\left(\frac{\alpha\beta}{4n}\right)^{\frac{2}{\gamma}} (\log(e\sqrt[4]{n}))^{\frac{2(\gamma-1)}{\gamma}}} \int_0^{\frac{1}{\sqrt[4]{n}}} \frac{r^{4-4\beta}}{4-4\beta} \left(\log \frac{e}{r}\right)^{\beta-1} dr \\
&= o\left(\frac{1}{\log e\sqrt[4]{n}}\right).
\end{aligned}$$

Also,

$$\begin{aligned}
I_2 &= 2\pi^2 \frac{(1-\beta)^2}{\left(\frac{\alpha\beta}{4}\right)^{\frac{2}{\gamma}} (\log(e\sqrt[4]{n}))^{\frac{2}{\gamma}}} \int_{\frac{1}{\sqrt[4]{n}}}^{\frac{1}{2}} \frac{1}{r} \left(\log \frac{e}{r}\right)^{-\beta} (2 + \beta (\log \frac{e}{r})^{-1})^2 dr \\
&= -2\pi^2 \frac{(1-\beta)^2}{\left(\frac{\alpha\beta}{4}\right)^{\frac{2}{\gamma}} (\log(e\sqrt[4]{n}))^{\frac{2}{\gamma}}} \left[\frac{\beta^2}{-1-\beta} \left(\log \frac{e}{r}\right)^{-\beta-1} + 4 \left(\log \frac{e}{r}\right)^{-\beta} + \frac{4}{1-\beta} \left(\log \frac{e}{r}\right)^{1-\beta} \right]_{\frac{1}{\sqrt[4]{n}}}^{\frac{1}{2}} \\
&= 1 + o\left(\frac{1}{(\log e\sqrt[4]{n})^{\frac{2}{\gamma}}}\right).
\end{aligned}$$

and $I_3 = o\left(\frac{1}{(\log e\sqrt[4]{n})^{\frac{2}{\gamma}}}\right)$. Then $\|\Delta w_n\|_{2,w}^2 = 1 + o\left(\frac{1}{(\log e\sqrt[4]{n})^{\frac{2}{\gamma}}}\right)$. Also,

$$\text{for } 0 \leq |x| \leq \frac{1}{\sqrt[4]{n}}, v_n^\gamma(x) \geq \left(\frac{\log(e\sqrt[4]{n})}{\alpha\beta}\right) + o(1).$$

5.2 Key lemmas

Lemma 4 Assume $V(x)$ is continuous and (V_1) is satisfied. Then there holds $\lim_{n \rightarrow +\infty} \|w_n\|^2 = 1$

Proof We have

$$\begin{aligned}
\|w_n\|^2 &= \int_B |\Delta w_n|^2 w(x) dx + \int_B V w_n^2 dx \\
&= 1 + o\left(\frac{1}{(\log e\sqrt[4]{n})^{\frac{2}{\gamma}}}\right) + \int_{0 \leq |x| \leq \frac{1}{\sqrt[4]{n}}} V w_n^2 dx + \int_{\frac{1}{\sqrt[4]{n}} \leq |x| \leq \frac{1}{2}} V w_n^2 dx + \int_{|x| \geq \frac{1}{2}} V \zeta_n^2 dx.
\end{aligned}$$

For $|x| \leq \frac{1}{\sqrt[4]{n}}$, $w_n^2 \leq \left(\left(\frac{\log(e\sqrt[4]{n})}{\alpha\beta}\right)^{\frac{1}{\gamma}} + \frac{1}{2\left(\frac{\alpha\beta}{4}\right)^{\frac{1}{\gamma}} (\log(e\sqrt[4]{n}))^{\frac{\gamma-1}{\gamma}}}\right)^2$. Then,

$$\int_{0 \leq |x| \leq \frac{1}{\sqrt[4]{n}}} V w_n^2 dx \leq 2\pi^2 m \left(\left(\frac{\log(e\sqrt[4]{n})}{\alpha\beta}\right)^{\frac{1}{\gamma}} + \frac{1}{2\left(\frac{\alpha\beta}{4}\right)^{\frac{1}{\gamma}} (\log(e\sqrt[4]{n}))^{\frac{\gamma-1}{\gamma}}}\right)^2 \int_0^{\frac{1}{\sqrt[4]{n}}} r^3 dr = o_n(1)$$

Also,

$$\int_{\frac{1}{\sqrt[4]{n}} \leq |x| \leq \frac{1}{2}} V w_n^2 dx \leq 2\pi^2 m \frac{1}{\left(\frac{\alpha_\beta}{4} \log(e\sqrt[4]{n})\right)^{\frac{1}{\gamma}}} \int_{\frac{1}{\sqrt[4]{n}}}^{\frac{1}{2}} r^3 \left(\log\left(\frac{e}{r}\right)\right)^2 dr$$

So,

$$\int_{\frac{1}{\sqrt[4]{n}} \leq |x| \leq \frac{1}{2}} V w_n^2 dx \leq 2\pi^2 m \frac{1}{\left(\frac{\alpha_\beta}{4} \log(e\sqrt[4]{n})\right)^{\frac{1}{\gamma}}} \int_{\frac{1}{\sqrt[4]{n}}}^{\frac{1}{2}} e^2 r dr = o_n(1).$$

Finally,

$$\int_{\frac{1}{\sqrt[4]{n}}}^{\frac{1}{2}} V w_n^2 dx \leq m \int_{|x| \geq \frac{1}{2}} \zeta_n^2 dx = o_n(1).$$

Hence, $\|w_n\|^2 \leq 1 + o_n(1)$ and consequently $\|w_n\|^\gamma \leq 1 + o_n(1)$.

In the same way, using the fact that for all $0 \leq |x| \leq \frac{1}{\sqrt[4]{n}}$, $w_n^2 \geq \left(\frac{\log(e\sqrt[4]{n})}{\alpha_\beta}\right)^{\frac{2}{\gamma}}$, we get

$$\int_{0 \leq |x| \leq \frac{1}{\sqrt[4]{n}}} V w_n^2 dx \geq V_0 2\pi^2 \left(\frac{\log(e\sqrt[4]{n})}{\alpha_\beta}\right)^{\frac{2}{\gamma}} \int_0^{\frac{1}{\sqrt[4]{n}}} r^3 dr = o_n(1)$$

and

$$\int_{\frac{1}{\sqrt[4]{n}} \leq |x| \leq \frac{1}{2}} V w_n^2 dx \geq 2\pi^2 V_0 \frac{1}{\left(\frac{\alpha_\beta}{4} \log(e\sqrt[4]{n})\right)^{\frac{2}{\gamma}}} \int_{\frac{1}{\sqrt[4]{n}}}^{\frac{1}{2}} r^3 \left(\log\left(\frac{e}{r}\right)\right)^2 dr = o_n(1)$$

Consequently, $1 + o_n(1) \leq \|w_n\|^2 \leq 1 + o_n(1)$.

5.3 Min-Max level estimate

We are going to the desired estimate.

Lemma 5 *For the sequence (v_n) identified by (5.2), there exists $n \geq 1$ such that*

$$\max_{t \geq 0} J(tv_n) < \frac{1}{2} \left(\frac{\alpha_\beta}{\alpha_0}\right)^{\frac{2}{\gamma}}. \quad (5.3)$$

Proof By contradiction, suppose that for all $n \geq 1$,

$$\max_{t \geq 0} J(tv_n) \geq \frac{1}{2} \left(\frac{\alpha_\beta}{\alpha_0}\right)^{\frac{2}{\gamma}}.$$

Therefore, for any $n \geq 1$, there exists $t_n > 0$ such that

$$\max_{t \geq 0} J(tv_n) = J(t_n v_n) \geq \frac{1}{2} \left(\frac{\alpha_\beta}{\alpha_0}\right)^{\frac{2}{\gamma}}$$

and so,

$$\frac{1}{2}t_n^2 - \int_B F(x, t_n v_n) dx \geq \frac{1}{2} \left(\frac{\alpha_\beta}{\alpha_0} \right)^{\frac{2}{\gamma}}.$$

Then, by using (H1)

$$t_n^2 \geq \left(\frac{\alpha_\beta}{\alpha_0} \right)^{\frac{2}{\gamma}}. \quad (5.4)$$

On the other hand,

$$\frac{d}{dt} J(tv_n) \Big|_{t=t_n} = t_n - \int_B f(x, t_n v_n) v_n dx = 0,$$

that is

$$t_n^2 = \int_B f(x, t_n v_n) t_n v_n dx. \quad (5.5)$$

Now, we claim that the sequence (t_n) is bounded in $(0, +\infty)$.

Indeed, it follows from (H5) that for all $\varepsilon > 0$, there exists $t_\varepsilon > 0$ such that

$$f(x, t)t \geq (\gamma_0 - \varepsilon)e^{\alpha_0 t^\gamma} \quad \forall |t| \geq t_\varepsilon, \quad \text{uniformly in } x \in B. \quad (5.6)$$

Using (5.4) and (5.5), we get

$$t_n^2 = \int_B f(x, t_n v_n) t_n v_n dx \geq \int_{0 \leq |x| \leq \frac{1}{\sqrt[4]{n}}} f(x, t_n v_n) t_n v_n dx.$$

Since

$$\frac{t_n}{\|w_n\|} \left(\frac{\log e^{\sqrt[4]{n}}}{\alpha_\beta} \right)^{\frac{1}{\gamma}} \rightarrow \infty \quad \text{as } n \rightarrow +\infty,$$

then it follows from (5.6) that for all $\varepsilon > 0$, there exists n_0 such that for all $n \geq n_0$

$$\begin{aligned} t_n^2 &\geq (\gamma_0 - \varepsilon) \int_{0 \leq |x| \leq \frac{1}{\sqrt[4]{n}}} e^{\alpha_0 t_n^\gamma v_n^\gamma} dx \\ t_n^2 &\geq 2\pi^2(\gamma_0 - \varepsilon) \int_0^{\frac{1}{\sqrt[4]{n}}} r^3 e^{\alpha_0 t_n^\gamma \left(\frac{\log(e^{\sqrt[4]{n}})}{\alpha_\beta} \right) + o(1)} dr \end{aligned} \quad (5.7)$$

Hence,

$$1 \geq 2\pi^2(\gamma_0 - \varepsilon) e^{\alpha_0 t_n^\gamma \left(\frac{\log(e^{\sqrt[4]{n}})}{\alpha_\beta} \right) + o(1)} - 3 \log n - 2 \log t_n.$$

Therefore (t_n) is bounded. Also, we have from the formula (5.5) that

$$\lim_{n \rightarrow +\infty} t_n^2 \geq \left(\frac{\alpha_\beta}{\alpha_0} \right)^{\frac{2}{\gamma}}.$$

Now, suppose that

$$\lim_{n \rightarrow +\infty} t_n^2 > \left(\frac{\alpha_\beta}{\alpha_0} \right)^{\frac{2}{\gamma}},$$

then for n large enough, there exists some $\delta > 0$ such that $t_n^\gamma \geq \frac{\alpha\beta}{\alpha_0} + \delta$. Consequently the right hand side of (5.7) tends to infinity and this contradicts the boundedness of (t_n) . Since (t_n) is bounded, we get

$$\lim_{n \rightarrow +\infty} t_n^2 = \left(\frac{\alpha\beta}{\alpha_0}\right)^\frac{2}{\gamma}. \quad (5.8)$$

Let

$$\begin{aligned} \mathcal{A}_n &= \{x \in B \mid t_n v_n \geq t_\varepsilon\} \text{ and } \mathcal{C}_n = B \setminus \mathcal{A}_n, \\ t_n^2 &= \int_B f(x, t_n v_n) t_n v_n dx = \int_{\mathcal{A}_n} f(x, t_n v_n) t_n v_n dx + \int_{\mathcal{C}_n} f(x, t_n v_n) t_n v_n dx \\ &\geq (\gamma_0 - \varepsilon) \int_{\mathcal{A}_n} e^{\alpha_0 t_n^\gamma v_n^\gamma} dx + \int_{\mathcal{C}_n} f(x, t_n v_n) t_n v_n dx \\ &= (\gamma_0 - \varepsilon) \int_B e^{\alpha_0 t_n^\gamma v_n^\gamma} dx - (\gamma_0 - \varepsilon) \int_{\mathcal{C}_n} e^{\alpha_0 t_n^\gamma v_n^\gamma} dx \\ &\quad + \int_{\mathcal{C}_n} f(x, t_n v_n) t_n v_n dx. \end{aligned}$$

Since $v_n \rightarrow 0$ a.e in B , $\chi_{\mathcal{C}_n} \rightarrow 1$ a.e in B , therefore using the dominated convergence theorem, we get

$$\int_{\mathcal{C}_n} f(x, t_n v_n) t_n v_n dx \rightarrow 0 \text{ and } \int_{\mathcal{C}_n} e^{\alpha_0 t_n^\gamma v_n^\gamma} dx \rightarrow \frac{\pi^2}{2}.$$

Then,

$$\lim_{n \rightarrow +\infty} t_n^2 = \left(\frac{\alpha\beta}{\alpha_0}\right)^\frac{2}{\gamma} \geq (\gamma_0 - \varepsilon) \lim_{n \rightarrow +\infty} \int_B e^{\alpha_0 t_n^\gamma v_n^\gamma} dx - (\gamma_0 - \varepsilon) \frac{\pi^2}{2}.$$

On the other hand,

$$\int_B e^{\alpha_0 t_n^\gamma v_n^\gamma} dx \geq \int_{\frac{1}{\sqrt[3]{n}} \leq |x| \leq \frac{1}{2}} e^{\alpha_0 t_n^\gamma v_n^\gamma} dx + \int_{\mathcal{C}_n} e^{\alpha_0 t_n^\gamma v_n^\gamma} dx.$$

Then, using (5.4)

$$\lim_{n \rightarrow +\infty} t_n^2 \geq \lim_{n \rightarrow +\infty} (\gamma_0 - \varepsilon) \int_B e^{\alpha_0 t_n^\gamma v_n^\gamma} dx \geq \lim_{n \rightarrow +\infty} (\gamma_0 - \varepsilon) 2\pi^2 \int_{\frac{1}{\sqrt[3]{n}}}^{\frac{1}{2}} r^3 e^{\frac{4 \left(\log \frac{\varepsilon}{r}\right)^2}{\log(e^{\frac{1}{\sqrt[3]{n}}}) \|w_n\|^\gamma}} dr.$$

Therefore, we get

$$\begin{aligned}
\lim_{n \rightarrow +\infty} t_n^2 &\geq \lim_{n \rightarrow +\infty} (\gamma_0 - \varepsilon) \int_B e^{\alpha_0 t_n^\gamma v_n^\gamma} dx \\
&\geq \lim_{n \rightarrow +\infty} 2\pi^2 (\gamma_0 - \varepsilon) \frac{\|w_n\|^\gamma \log(e \sqrt[4]{n})}{4} e^4 \int_{\frac{4 \log 2e}{\|w_n\|^\gamma \log(e \sqrt[4]{n})}}^{\frac{4}{\|w_n\|^\gamma}} e^{\frac{\|w_n\|^\gamma \log(e \sqrt[4]{n})}{4} (s^2 - 4s)} ds \\
&\geq \lim_{n \rightarrow +\infty} 2\pi^2 (\gamma_0 - \varepsilon) \frac{\|w_n\|^\gamma \log(e \sqrt[4]{n})}{4} e^4 \int_{\frac{4 \log 2e}{\|w_n\|^\gamma \log(e \sqrt[4]{n})}}^{\frac{4}{\|w_n\|^\gamma}} e^{-\frac{\|w_n\|^\gamma \log(e \sqrt[4]{n})}{4} 4s} ds \\
&= \lim_{n \rightarrow +\infty} (\gamma_0 - \varepsilon) \frac{\pi^2}{2} e^4 (-e^{-4 \log e \sqrt[4]{n}} + e^{-4 \log(2e)}) \\
&= (\gamma_0 - \varepsilon) \frac{\pi^2 e^{4(1 - \log 2e)}}{2} = (\gamma_0 - \varepsilon) \frac{\pi^2}{32}.
\end{aligned}$$

It follows that

$$\left(\frac{\alpha_\beta}{\alpha_0}\right)^{\frac{2}{\gamma}} \geq (\gamma_0 - \varepsilon) \frac{\pi^2}{32}$$

for all $\varepsilon > 0$. So,

$$\gamma_0 \leq \frac{1024(1 - \beta)}{\alpha_0^{1-\beta}},$$

which is in contradiction with the condition (H5).

Now by Proposition 4.2, the functional \mathcal{J} satisfies the (PS) condition at a level $c < \frac{1}{2} \left(\frac{\alpha_\beta}{\alpha_0}\right)^{\frac{2}{\gamma}}$. Also, by Proposition 4.1, we deduce that the functional J has a nonzero critical point u in \mathbf{W} . From maximum principle, the solution u of the problem (1) is positive. The Theorem 1.3 is proved.

References

- [1] D. R. Adams, *A sharp inequality of J. Moser for higher order derivatives*, Annals of Math., 128 (1988), 385-398.
- [2] Adimurthi, *Positive solutions of the semilinear Dirichlet problem with critical growth in the unit disc in R^2* , Proc. Indian Acad. Sci. Math. Sci., 99 (1989), 49-73.
- [3] A. Adimurthi, *Existence results for the semilinear Dirichlet problem with critical growth for the n -Laplacian*, Houst. J. Math. 7 (1991), 285-298. MR 1079983 — Zbl 0732.35028
- [4] L.E. Andersson, T. Elfving and G.H. Golub, *Solution of biharmonic equations with application to radar imaging*, J. Comp. and Appl. Math. 94 (2) (1998), 153-180.
- [5] A. Ambrosetti and P. H. Rabinowitz, *Dual variational methods in critical points theory and applications*, J. Funct. Anal. 14 (1973), 349-381.

- [6] S. Baraket, R. Jaidane, *Non-autonomous weighted elliptic equations with double exponential growth*, An. Șt. Univ. Ovidius Constanța, Vol. 29(3), 2021, 33-66.
- [7] Elvise Berchio, Filippo Gazzola and Tobias Weth, *Critical growth biharmonic elliptic problems under Steklov-type boundary conditions*, Adv. Differential Equations, 12 (2007), 381-406.
- [8] E. Caglioti, P.L. Lions, C. Marchioro and M. Pulvirenti, *A Special Class of Stationary Flows for Two-Dimensional Euler Equations: a Statistical Mechanics Description*, Communications in Mathematical Physics, 1992, vol. 143, no. 3, 501-525. DOI: 10.1007/BF02099262
- [9] E. Caglioti, P.L. Lions, C. Marchioro and M. Pulvirenti, *A Special Class of Stationary Flows for Two-Dimensional Euler Equations: a Statistical Mechanics Description. II*, Communications in Mathematical Physics, 1995, vol. 174, no. 2, pp. 229-260. DOI: 10.1007/BF02099602
- [10] M. Calanchi, *"Some weighted inequalities of Trudinger - Moser Type" in Analysis and Topology in Nonlinear Differential Equations*, Progress in Nonlinear Differential Equations and Appl., Birkhauser vol 85 (2014), 163-174. MR3330728
- [11] M. Calanchi and B. Ruf, *On a Trudinger-Moser type inequalities with logarithmic weights*, Journal of Differential Equations no. 3 (2015), 258-263. Doi: 10.1016/j.jde.2014.11.019
- [12] M. Calanchi and B. Ruf, *Trudinger-Moser type inequalities with logarithmic weights in dimension N* , Nonlinear Analysis, Series A; Theory Methods and Applications 121 (2015), 403-411. DOI: 10.1016/j.na.2015.02.001
- [13] M. Calanchi and B. Ruf, *Weighted Trudinger-Moser inequalities and Applications*, Bulletin of the South Ural State University. Ser. Mathematical Modelling, programming and Computer Software vol. 8 no. 3 (2015), 42-55. DOI: 10.14529/mmp150303
- [14] M. Calanchi, B. Ruf and F. Sani, *Elliptic equations in dimension 2 with double exponential nonlinearities*, NoDea Nonlinear Differ. Equ. Appl., **24** (2017), Art. 29.
- [15] S. Chanillo and M. Kiessling, *Rotational Symmetry of Solutions of Some Nonlinear Problems in Statistical Mechanics and in Geometry*, Communications in Mathematical Physics, 1994, vol. 160, no. 2, 217-238. DOI: 10.1007/BF02103274
- [16] C.P. Dănet, *Two maximum principles for a nonlinear fourth order equation from thin plate theory*, Electron. J. Qual. Theory Differ. Equ. 31 (2014), 1-9.
- [17] S. Deng, T. Hu and C. Tang, *N -laplacian problems with critical double exponential nonlinearities*, DISCRETE AND CONTINUOUS DYNAMICAL SYSTEMS, **41** (2021), 987-1003.
- [18] P. Drabek, A. Kufner and F. Nicolosi, *Quasilinear Elliptic Equations with Degenerations and Singularities*, Walter de Gruyter, Berlin (1997). DOI:10.1515/9783110804775
- [19] D. E. Edmunds, D. Fortunato and E. Jannelli, *Critical exponents, critical dimensions and the biharmonic operator*, Arch. Rational Mech. Anal., 112 (1990), 269-289.

- [20] A. Ferrero and G. Warnault, *On a solutions of second and fourth order elliptic with power type nonlinearities*, Nonlinear Anal. TMA 70 (8) (2009),2889-2902.
- [21] D.G. de Figueiredo, O.H. Miyagaki and B. Ruf, *Elliptic equations in \mathbb{R}^2 with nonlinearities in the critical growth range*, Calc. Var. Partial Differential Equations 3 (2) (1995), 139-153. DOI: 10.1007/BF01205003.
- [22] Filippo Gazzola, Hans-Christoph Grunau and Marco Squassina, *Existence and nonexistence results for critical growth biharmonic elliptic equations*, Calc. Var. Partial Differential Equations, 18 (2003), 117-143.
- [23] M.K.-H. Kiessling, *Statistical Mechanics of Classical Particles with Logarithmic Interactions*, Communications on Pure and Applied Mathematics, 1993, vol. 46, 27-56. DOI:10.1002/cpa.3160460103
- [24] A. Kufner, *Weighted Sobolev spaces*, John Wiley and Sons Ltd, 1985. Doi: 10.1112/blms/18.2.220
- [25] Nguyen Lam and Guozhen Lu, *Elliptic equations and systems with subcritical and critical exponential growth without the Ambrosetti-Rabinowitz condition*, J GeomAnal DOI 10.1007/s12220-012-9330-4
- [26] N. Lam and G. Lu, *Existence and multiplicity of solutions to equations of N -Laplacian type with critical exponential growth in \mathbb{R}^N* , J. Funct. Anal. 262 no. 3 (2012), 1132-1165.
- [27] J. Liouville, *Sur l'équation aux dérivées partielles*, Journal de Mathématiques Pures et Appliquées **18** (1853), 71-72.
- [28] P.L. Lions, *The Concentration-compactness principle in the Calculus of Variations*, Part 1, Revista Iberoamericana 11 (1985), 185-201. MR 778970 — Zbl 0541.49009
- [29] O. H. Miyagaki and M. A. S. Souto, *Superlinear problems without Ambrosetti and Rabinowitz growth condition*, J. Differential Equations, 245 (2008), 3628-3638.
- [30] T.G. Myers, *Thin films with high surface tension*, SIAM Rev. 40 (3) (1998), 441-462.
- [31] J. Moser, *A sharp form of an inequality by N. Trudinger*, Indiana Univ. Math. J., 20 (1970/71), 1077-1092.
- [32] B. Ruf and F. Sani, *Sharp Adams-type inequalities in \mathbb{R}^N* , Trans. Amer. Math. Soc., in press, doi.org/10.1090/S0002-9947-2012-05561-9
- [33] F. Sani, *A biharmonic equation in \mathbb{R}^4 involving nonlinearities with critical exponential growth*, COMMUNICATIONS ON PURE AND APPLIED ANALYSIS Volume 12, Number 1 405-428 doi:10.3934/cpaa.2013.12.405
- [34] Wolfgang Reichel and Tobias Weth, *Existence of solutions to nonlinear, subcritical higher order elliptic Dirichlet problems*, J. Differential Equations, 248 (2010), 1866-1878.

- [35] R.E. Volker, *Nonlinear flow in porous media by finite elements*, Journal of the Hydraulics Division, Vol. 95 (6) (1969), 2093-2114.
- [36] G. Tarantello, *Condensate Solutions for the Chern - Simons - Higgs Theory*, Journal of Mathematical Physics vol. 37 (1996), 3769-3796. DOI: 10.1063/1.531601
- [37] G. Tarantello, *Analytical Aspects of Liouville-Type Equations with Singular Sources*, Handbook of Differential Equations (M. Chipot and P. Quittner, eds.), Elsevier, North Holland 2004, 491-592.
- [38] N. S. Trudinger, *On imbeddings into Orlicz spaces and some applications*, J. Math. Mech., 17 (1967), 473-483.
- [39] L. Wang, M. Zhu , *Adams' inequality with logarithm weight in \mathbb{R}^4* , PROCEEDINGS OF THE AMERICAN MATHEMATICAL SOCIETY, Volume 149, Number 8, August 2021, 3463–3472. doi: org/10.1090/proc/15488
- [40] C. Zhang, *Concentration-Compactness principle for Trudinger–Moser inequalities with logarithmic weights and their applications*, Nonlinear Anal. **197** (2020), 1-22.