

ON THE DYNAMICAL BOGOMOLOV CONJECTURE FOR FAMILIES OF SPLIT RATIONAL MAPS

NIKI MYRTO MAVRAKI AND HARRY SCHMIDT

ABSTRACT. We prove that Zhang’s dynamical Bogomolov conjecture (corrected appropriately) holds uniformly along 1-parameter families of rational split maps and curves. This provides dynamical analogues of recent results of Dimitrov-Gao-Habegger [DGH1, DGH3] and Kühne [Ku2]. In fact, we prove a stronger Bogomolov-type result valid for families of split maps in the spirit of the relative Bogomolov conjecture [DGH2, Conjecture 1.2]. We thus provide first instances of a generalization of Baker-DeMarco’s conjecture [BD1] to higher dimensions. Our proof contains both arithmetic and analytic ingredients. We establish a characterization of curves that are preperiodic under the action of a non-exceptional split rational endomorphism (f, g) of $(\mathbb{P}_{\mathbb{C}}^1)^2$ with respect to the measures of maximal entropy of f and g , extending a previous result of Levin-Przytycki [LP]. We further establish a height inequality for families of split maps and varieties comparing the values of a fiberwise Call-Silverman canonical height with a height on the base and valid for most points of a non-preperiodic variety. This provides a dynamical generalization of [Ha1, Theorem 1.3] and generalizes results of Call-Silverman [CS] and Baker [Ba] to higher dimensions. In particular, we establish a geometric Bogomolov theorem for split rational maps and varieties of arbitrary dimension.

1. INTRODUCTION

Motivated by the Manin-Mumford conjecture, proved in [Ra], and its strengthening as in Bogomolov’s conjecture, proved in [Ul, Zh3], Zhang conjectured that analogous statements [Zh4, Conjectures 1.2.1 and 4.1.7] hold in a more general dynamical setting. Our goal in this paper is to prove uniform and relative versions of the dynamical Bogomolov conjecture for 1-parameter families of split maps.

Our main results are stated as Theorem 1.2, Theorem 1.6, Theorem 1.8 and Theorem 1.9 in what follows. To ease the reader into our work, we begin with an application that may be of independent interest in complex dynamics. DeMarco, Krieger and Ye [DKY2, Conjecture 1.4] conjectured that there is a constant $M > 0$ such that for any two rational functions f

Date: January 26, 2022.

and g of degree $d \geq 2$ defined over \mathbb{C} , we have that

$$(1.1) \quad \text{either } \#\text{Prep}(f) \cap \text{Prep}(g) \leq M \text{ or } \text{Prep}(f) = \text{Prep}(g),$$

where $\text{Prep}(\phi)$ denotes the set of preperiodic points of a self-map of a set ϕ . As evidence, they show [DKY2, Theorem 1.1] that there is a constant $M > 0$ such that

$$\#\text{Prep}(z^2 + t_1) \cap \text{Prep}(z^2 + t_2) < M,$$

for all complex $t_1 \neq t_2$ or equivalently $\text{Prep}(z^2 + t_1) \neq \text{Prep}(z^2 + t_2)$; see [Bea, Theorem 1]. They thus establish that their conjecture holds along the surface in $\text{Rat}_d \times \text{Rat}_d$ given by unicritical quadratic polynomials. Here Rat_d denotes the space of rational maps of degree d .

A first application of our results shows that [DKY2, Conjecture 1.4] holds along arbitrary curves in $\text{Rat}_d \times \text{Rat}_d$ defined over the algebraic numbers.

Theorem 1.1. *Let $C \subset \text{Rat}_d \times \text{Rat}_d$ be a curve defined over $\overline{\mathbb{Q}}$ and let $p_i : C \rightarrow \text{Rat}_d$ be its projection to the i -th copy of Rat_d . For $\underline{\mu} \in C(\mathbb{C})$ let $f_{p_i(\underline{\mu})}$ denote the rational map associated to $p_i(\underline{\mu}) \in \text{Rat}_d(\mathbb{C})$ for $i = 1, 2$. Then, there exists $M = M(C) > 0$ such that*

$$\text{either } \#\text{Prep}(f_{p_1(\underline{\mu})}) \cap \text{Prep}(f_{p_2(\underline{\mu})}) < M \text{ or } \text{Prep}(f_{p_1(\underline{\mu})}) = \text{Prep}(f_{p_2(\underline{\mu})}).$$

Note here that the common preperiodic points of f and g in (1.1) can be seen as the preperiodic points under the action of the split map $(f, g) : \mathbb{P}_1^2 \rightarrow \mathbb{P}_1^2$, with $(f, g)(x, y) = (f(x), g(y))$, which lie on the diagonal in \mathbb{P}_1^2 . Replacing the diagonal with a different irreducible curve C , Ghioca, Nguyen and Ye [GNY1] have established that

$$(1.2) \quad \#\text{Prep}(f, g) \cap C(\mathbb{C}) < +\infty,$$

unless C is preperiodic under the action of (f, g) ; that is if f or g is non-exceptional. The latter assumption ensures that counterexamples to Zhang's original conjectures are avoided; see [GTZ, GT].

Also in view of recent breakthroughs of Dimitrov, Gao and Habegger [DGH3] and Kühne [Ku2], who established a uniform version of the classical Manin-Mumford and Bogomolov conjectures, one may wonder when is $\#\text{Prep}(f, g) \cap C(\mathbb{C})$ bounded uniformly. In Theorem 1.2 we show that this holds if the maps and the curve vary along a 1-parameter family.

Kühne has further established a stronger relative version of Bogomolov's conjecture for varieties in fibered products of elliptic families [Ku1]. In Theorem 1.7 we prove instances of an analogous statement in the setting of families of split maps.

1.1. **Setup.** In order to state our results we introduce some notation and terminology. Throughout we let B be a projective, regular, irreducible curve over $\overline{\mathbb{Q}}$ and write $K = \overline{\mathbb{Q}}(B)$. We let $\Phi : \mathbb{P}_1^\ell \rightarrow \mathbb{P}_1^\ell$ be an endomorphism defined over K for $\ell \in \mathbb{N}_{\geq 1}$. We say that Φ is a *split polarized endomorphism* of polarization degree $d \geq 2$ over K if it is defined as $\Phi(z_1, \dots, z_\ell) = (\mathbf{f}_1(z_1), \dots, \mathbf{f}_\ell(z_\ell))$, where $\mathbf{f}_1, \dots, \mathbf{f}_\ell$ are rational functions of fixed degree $d \geq 2$ defined over K . Recall that $\mathbf{f} \in K(z)$ is isotrivial if there is a Möbius transformation $\mathbf{M} \in \overline{K}(z)$ such that $\mathbf{M} \circ \mathbf{f} \circ \mathbf{M}^{-1} \in \overline{\mathbb{Q}}(z)$. If all \mathbf{f}_i are non-isotrivial for all $i = 1, \dots, \ell$, then we say that Φ is *isotrivial-free*. The map Φ defines an endomorphism of a quasiprojective variety $\Phi : B_0 \times \mathbb{P}_1^\ell \rightarrow B_0 \times \mathbb{P}_1^\ell$, where B_0 is a Zariski open and dense subset of B and Φ is defined over $\overline{\mathbb{Q}}$. Similarly a (geometrically) irreducible subvariety $\mathbf{X} \subset \mathbb{P}_1^\ell$ over K gives a subvariety $\mathcal{X} \subset B_0 \times \mathbb{P}_1^\ell$ over $\overline{\mathbb{Q}}$.

We denote by $\pi : B_0 \times \mathbb{P}_1^\ell \rightarrow B_0$ the projection map and for each $t \in B_0(\mathbb{C})$ we write $\Phi_t = \Phi|_{\pi^{-1}(\{t\})}$ and $\mathcal{X}_t = \mathcal{X} \cap \pi^{-1}(\{t\})$. We then have a fiber-wise Call-Silverman canonical height [CS] associated to Φ , assigning to each point $Z = (t, z_1, \dots, z_\ell) \in (B_0 \times \mathbb{P}_1^\ell)(\overline{\mathbb{Q}})$ the value of a canonical height in its fiber as

$$(1.3) \quad \hat{h}_\Phi(Z) = \hat{h}_{\Phi_{\pi(Z)}}(Z) = \sum_{i=1}^{\ell} \hat{h}_{f_{i,t}}(z_i) \in \mathbb{R}_{\geq 0}.$$

We note here that the canonical height $\hat{h}_\Phi(Z)$ vanishes exactly when $Z \in \text{Prep}(\Phi_{\pi(Z)})$. More generally it can be thought of as a measure of distance between Z and the preperiodic points of the specialization $\Phi_{\pi(Z)}$ on its fiber.

In what follows we fix B_0 to be the largest open subset of B such that Φ_t is a polarized endomorphism of polarization degree $d \geq 2$ and \mathcal{X}_t is irreducible for each $t \in B_0(\overline{\mathbb{Q}})$. Throughout the article, bold letters refer to geometric objects defined over K .

1.2. **Instances of uniformity in the dynamical Bogomolov conjecture.** Our first main result establishes the first instance of uniformity in the dynamical Bogomolov and Manin-Mumford conjectures [Zh4, Conjectures 1.2.1 and 4.1.7] or [GT, Conjecture 1.2] for varying non-polynomial maps or along a family of varying curves. For the statement to hold, one requires the notion of a weakly special curve (see Definition 1.5), as we shall explain shortly.

Theorem 1.2. *Let $\Phi : \mathbb{P}_1^2 \rightarrow \mathbb{P}_1^2$ be a split polarized endomorphism over K with polarization degree ≥ 2 . Let $\mathbf{C} \subset \mathbb{P}_1^2$ be an irreducible curve defined over K that is not weakly Φ -special.*

Then there exist $\epsilon > 0$ and $M > 0$ depending only on Φ and \mathbf{C} such that

$$(1.4) \quad \#\{z \in \mathcal{C}_t(\overline{\mathbb{Q}}) : \hat{h}_{\Phi_t}(z) < \epsilon\} \leq M,$$

for all but finitely many $t \in B_0(\overline{\mathbb{Q}})$. In particular,

$$\#\mathcal{C}_t(\overline{\mathbb{Q}}) \cap \text{Prep}(\Phi_t) \leq M,$$

for all but finitely many $t \in B_0(\overline{\mathbb{Q}})$.

That one should exclude certain $t \in B_0(\overline{\mathbb{Q}})$ in the conclusion of the theorem is clear. For instance if \mathcal{C}_t is Φ_t -preperiodic, then it contains infinitely many $\text{Prep}(\Phi_t)$ -preperiodic points. That there are at most finitely many such exceptional t is not a priori clear but follows from our theorem. More generally, we can show that for all but finitely many t the curve \mathcal{C}_t stays a distance away from being Φ_t -preperiodic. The said distance is measured by a real valued Arakelov-Zhang height function

$$t \mapsto \hat{h}_{\Phi_t}(\mathcal{C}_t);$$

taking non-negative values for $t \in B_0(\overline{\mathbb{Q}})$ (see §2.) In view of Zhang's fundamental inequality, characterizing its zeros amounts to proving the dynamical Manin-Mumford conjecture [GNY1]. More precisely, as a consequence of Theorem 1.2, we obtain the following.

Corollary 1.3. *Let $\Phi : \mathbb{P}_1^2 \rightarrow \mathbb{P}_1^2$ be a split polarized endomorphism over K with polarization degree ≥ 2 . Let $\mathbf{C} \subset \mathbb{P}_1^2$ be an irreducible curve defined over K that is not weakly Φ -special. Then the following hold.*

- (1) *There are at most finitely many $t \in B(\mathbb{C})$ such that \mathcal{C}_t is Φ_t -preperiodic.*
- (2) *There is $\epsilon > 0$ such that the set $\{t \in B_0(\overline{\mathbb{Q}}) : \hat{h}_{\Phi_t}(\mathcal{C}_t) < \epsilon\}$ is finite.*

Let us now explain the necessity of a condition on \mathbf{C} in the statement of Theorem 1.2. First, recalling that preperiodic curves contain infinitely many preperiodic points, we need to exclude the possibility that \mathbf{C} is Φ -preperiodic for the conclusion of Theorem 1.2 to remain true. If Φ is given by a pair of Lattès maps, which are associated to algebraic groups, then we further have to assume that \mathbf{C} is not the projection of a curve with genus 1. Moreover, if the curve is vertical $\mathbf{C} = \mathbf{c} \times \mathbb{P}_1$ for a point $\mathbf{c} \in K$ assuming that it is not Φ -preperiodic does not suffice for the conclusion to hold. Indeed, by an application of Montel's theorem there might still be infinitely many $t \in B_0(\overline{\mathbb{Q}})$ such that \mathcal{C}_t is Φ_t -preperiodic; see [De1]. To make matters worse, even excluding all such t one cannot hope to find uniform $\epsilon, M > 0$

satisfying (3.5). If for instance \mathbf{f}_1 is not isotrivial, one can solve the equations $f_{1,t}^n(c(t)) = 1$ for varying n and apply [CS, Theorem 1.4].

Our notion of a weakly Φ -special curve aims to summarize all such necessary assumptions. As our examples reveal, we need to consider maps that are associated to groups separately. To this end, we give here the notion of an exceptional map. Our vocabulary might not completely agree with some terms in the established literature. However a consensus does not seem to exist.

Definition 1.4. For this definition we assume that all objects are defined over some algebraically closed field. We say that a rational map $f : \mathbb{P}_1 \rightarrow \mathbb{P}_1$ of degree $\deg(f) \geq 2$ is *exceptional* if there exists an algebraic group G_f (which is either an elliptic curve or the multiplicative group), an isogeny $\Psi_f : G_f \rightarrow G_f$ and a dominant map $\varphi_f : G_f \rightarrow \mathbb{P}_1$ such that

$$\varphi_f \circ \Psi_f = f \circ \varphi_f.$$

If f is not exceptional, then we say that f is *ordinary*.

Equipped with Definition 1.4 we can now define the notion of a weakly special curve for a split map.

Definition 1.5. For this definition we assume that all objects are defined over some algebraically closed field. Let $C \subset \mathbb{P}_1^2$ be an irreducible curve and denote its projection into the i -th coordinate of \mathbb{P}_1 by $\pi_i : C \rightarrow \mathbb{P}_1$ for $i = 1, 2$. Let $\phi = (f, g) : \mathbb{P}_1^2 \rightarrow \mathbb{P}_1^2$ be a split polarized endomorphism of polarization degree ≥ 2 . We say that C is *weakly ϕ -special* if one of the following hold:

- (1) π_i is not dominant for some $i = 1, 2$; or
- (2) C is ϕ -preperiodic; or
- (3) both f and g are exceptional maps with associated algebraic groups G_f, G_g and maps Ψ_f, Ψ_g and there exists a coset $H \subset G_f \times G_g$ such that $C = (\Psi_f, \Psi_g)(H)$.

We say that C is *ϕ -special* if it is ϕ -preperiodic or both f and g are exceptional maps with associated algebraic groups G_f, G_g and maps Ψ_f, Ψ_g and there exists a torsion coset $H \subset G_f \times G_g$ such that $C = (\Psi_f, \Psi_g)(H)$.

Determining the equations that define preperiodic curves is a notoriously difficult problem. That said, a precise description of curves that are invariant under the action of split maps acting by a polynomial on each coordinate has been given in the celebrated work of Medvedev

and Scanlon [MS]. That at hand one can write several applications of Theorem 1.2. For instance, given $(n, m) \in \mathbb{Z} \setminus \{\pm(1, -1)\}$ and an ordinary polynomial $f(z) \in \overline{\mathbb{Q}}[z]$, we can bound the number of solutions

$$x^n y^m = t, \text{ with } x, y \in \text{Prep}(f)$$

independently of t .

1.3. Towards a relative dynamical Bogomolov conjecture. Theorem 1.2 is in fact a corollary of our next result, which gives instances of dynamical analogues of the relative Bogomolov theorem of Kühne [Ku1]. In particular, it provides a better description of the geometric distribution of the sets in (1.4).

Theorem 1.6. *Let $\Phi : \mathbb{P}_1^2 \rightarrow \mathbb{P}_1^2$ and $\Psi : \mathbb{P}_1^\ell \rightarrow \mathbb{P}_1^\ell$ be isotrivial-free split polarized endomorphisms over K both of polarization degree $d \geq 2$ for $\ell \geq 2$. Let $\mathbf{C} \subset \mathbb{P}_1^2$ be an irreducible curve that is not weakly Φ -special and let $\mathbf{X} \subset \mathbb{P}_1^\ell$ be a irreducible subvariety that is not Ψ -preperiodic, both defined over K . Then there exists $\epsilon = \epsilon(\Phi, \mathbf{C}, \Psi, \mathbf{X}) > 0$ such that the set*

$$\{(P, Q) \in (\mathcal{C} \times_{B_0} \mathcal{X})(\overline{\mathbb{Q}}) : \hat{h}_\Phi(P) + \hat{h}_\Psi(Q) < \epsilon\}$$

is not Zariski dense in $\mathcal{C} \times_{B_0} \mathcal{X}$. Here $\hat{h}_\Phi(\cdot)$, $\hat{h}_\Psi(\cdot)$ denote the corresponding fiber-wise heights from (1.3).

This result fits in the theme of unlikely intersections and is inspired by the Pink-Zilber conjectures and by the classical relative Bogomolov conjecture [DGH2, Conjecture 1.2]; see [Za] for an overview of related problems in the classical setting. More generally, the Pink-Zilber conjectures suggest that the following holds.

Conjecture 1.7. *Let $\ell \geq 2$ and $\mathbf{F} : \mathbb{P}_1^\ell \rightarrow \mathbb{P}_1^\ell$ be an isotrivial-free split polarized endomorphism over K with polarization degree ≥ 2 . Let $\mathbf{V} \subset \mathbb{P}_1^\ell$ be an irreducible projective subvariety defined over K and such that $\dim \mathbf{V} < \ell - 1$. Assume that \mathbf{V} is not contained in an \mathbf{F} -preperiodic subvariety other than \mathbb{P}_1^ℓ . Then there exists $\epsilon = \epsilon(\mathbf{F}, \mathbf{V}) > 0$ such that the set*

$$\{P \in \mathcal{V}(\overline{\mathbb{Q}}) : \hat{h}_F(P) \leq \epsilon\}$$

is not Zariski dense in \mathcal{V} , where $\hat{h}_F(\cdot)$ is the fiber-wise height as in (1.3).

Theorem 1.6 is the first instance of Conjecture 1.7 for \mathbf{V} with positive dimension and in a strictly dynamical setting. The only known case of Conjecture 1.7 (and of [DGH2, Conjecture 1.2]) in the case that $\dim \mathbf{V} \geq 1$ is Kühne’s result [Ku1, Theorem 1], who established the conjecture when all components of \mathbf{F} are Lattès maps, thereby extending [DM1, Theorem 1.4]. For related results in the case that $\epsilon = 0$ see also [MZ1, MZ2, MZ3, MZ4].

That said, there has been a lot of work concerning cases of the conjecture for points \mathbf{V} . Then it follows from Baker-DeMarco’s conjecture [BD1, Conjecture 1.10], [De1, Conjecture 6.1]. Its first case is established in the breakthrough article of Baker-DeMarco [BD2]. A lot of work has centered around other special cases – see [DWY1, GHT1, GHT2, BD1, DWY2, FG1, FG2, GKN, GKY, GY]. Most notably, recently Favre-Gauthier [FG3] established the conjecture for points \mathbf{V} and split maps \mathbf{F} acting by a polynomial on each coordinate.

Our next results provide with necessary ingredients towards our proof of Theorems 1.6 and 1.2.

1.4. Weakly special curves and measures of maximal entropy. For each rational map $f \in \mathbb{C}(z)$ with $\deg f \geq 2$ there is an invariant measure μ_f supported on the Julia set of f which was constructed by Lyubich [Ly] and Freire-Lopes-Mañé [FLM]. It is the unique measure with no point masses and such that $f^*\mu_f = (\deg f)\mu_f$. Rigidity questions such as when do two rational maps share the same invariant measure have been studied extensively; see [Ye] and the references therein for a relatively recent account.

Levin and Przytycki [LP, Theorem A] have shown that the measure μ_f determines the rational map f up to certain operations. Our next result can be seen as a generalization of [LP, Theorem A]. It allows us to characterize weakly special curves by looking at measures μ_f .

Theorem 1.8. *Let $f_i \in \mathbb{C}(z)$ be rational maps of degree $d \geq 2$ for $i = 1, 2$. Let $C \subset \mathbb{P}_1^2$ be an irreducible complex algebraic curve with dominant projections $\pi_i : C \rightarrow \mathbb{P}_1$ for $i = 1, 2$. Then $\deg(\pi_2)\pi_1^*\mu_{f_1} = \deg(\pi_1)\pi_2^*\mu_{f_2}$ if and only if $\text{supp}(\pi_1^*\mu_{f_1}) \cap \text{supp}(\pi_2^*\mu_{f_2}) \neq \emptyset$ and C is (f_1, f_2) -weakly special.*

In fact we establish a more general statement, not assuming that f_1 and f_2 have the same degree; see Theorem 3.1. Assuming further that there exists a point $\lambda \in C(\mathbb{C})$ such that $\pi_1(\lambda)$ is a repelling f_1 -periodic point and $\pi_2(\lambda)$ is f_2 -preperiodic, a related result is obtained in [GNY1, Theorem 5.1] and was used to establish the dynamical Bogomolov conjecture. Showing that such a point exists is non-trivial. In [GNY1] the authors infer it from Mimar’s

main theorem [Mi] in view of their assumption that C contains infinitely many (f_1, f_2) -preperiodic points. But in order to establish the uniform statement in Theorem 1.2 we cannot operate under this helpful assumption.

1.5. A height inequality and a geometric dynamical Bogomolov theorem. Finally, in Theorem 4.1, we establish a geometric Bogomolov theorem for split maps over 1-dimensional function fields of characteristic zero. This leads us to an inequality comparing the fiber-wise Call-Silverman canonical height with a height on the base and valid for most points of a non-preperiodic subvariety. For a split polarized endomorphism Φ over K , we let

$$\mathcal{X}^{\Phi, \star} = \mathcal{X} \setminus \bigcup_{\mathcal{Z}} \mathcal{Z},$$

where the union runs over all subvarieties \mathcal{Z} of \mathcal{X} that are irreducible components of Φ -special subvarieties projecting dominantly onto the base B .

Theorem 1.9. *Let $\ell \geq 2$ and $\Phi : \mathbb{P}_1^\ell \rightarrow \mathbb{P}_1^\ell$ be an isotrivial-free split polarized endomorphism over K with polarization degree at least equal to 2. Let $\mathbf{X} \subset \mathbb{P}_1^\ell$ be an irreducible projective subvariety defined over K and let \mathcal{N} be ample line bundle on B . Then exactly one of the following holds:*

- (1) *Either $\mathcal{X}^{\Phi, \star} = \emptyset$ and \mathbf{X} is Φ -preperiodic ; or*
- (2) *the set $\mathcal{X}^{\Phi, \star}$ is Zariski open and dense in \mathcal{X} and there are constants $c_1, c_2 > 0$ depending on Φ, \mathbf{X} and \mathcal{N} and such that*

$$(1.5) \quad \hat{h}_\Phi(P) \geq c_1 h_{\mathcal{N}}(\pi(P)) - c_2, \text{ for all } P \in \mathcal{X}^{\Phi, \star}(\overline{\mathbb{Q}}).$$

As an immediate corollary, we obtain uniformity in the dynamical Bogomolov conjecture across 1-parameter families of isotrivial-free split rational maps and varieties; albeit under the assumption that their height is large enough.

Corollary 1.10. *Let $\ell \geq 2$ and $\Phi : \mathbb{P}_1^\ell \rightarrow \mathbb{P}_1^\ell$ be an isotrivial-free split polarized endomorphism. Let $\mathbf{X} \subset \mathbb{P}_1^\ell$ be an irreducible subvariety that is not Φ -preperiodic. Then there are constants $c_3 = c_3(\Phi, \mathbf{X}), c_4 = c_4(\Phi, \mathbf{X})$ such that the set*

$$(1.6) \quad \left\{ z \in \mathcal{X}_t(\overline{\mathbb{Q}}) : (t, z) \in \mathcal{X}^{\Phi, \star}(\overline{\mathbb{Q}}) \text{ and } \hat{h}_{\Phi_t}(z) \leq c_4 \max\{h(t), 1\} \right\},$$

is empty for all $t \in B_0(\overline{\mathbb{Q}})$ with $h(t) > c_3$.

We expect that the techniques in this article can be generalized to allow for proving that the set in (1.6) has uniformly bounded cardinality without the assumption on the height of

t. However there are several technical obstacles, such as generalizing Theorem 1.8 to higher dimensions. We defer this investigation to a future article.

Theorem 1.9 is a dynamical generalization of Habegger’s result [Ha1, Theorem 1.3] which dealt with the case that all components \mathbf{f}_i of Φ are Lattès maps associated to elliptic curves. It should be noted that a first conditional result in the direction of Theorem 1.9, dealing with the case that \mathbf{X} is a point, is established by Call-Silverman [CS]. Under the assumption that the geometric (over the function field K) canonical height of \mathbf{X} under Φ is non-zero, they derive an inequality such as (1.5). In order to then get the result in Theorem 1.9 in this case, it is challenging to characterize the points of non-zero geometric canonical height as non-preperiodic. This issue has been clarified by Baker [Ba] following work of Benedetto [Ben]; see also [De1]. Therefore, in the case that \mathbf{X} is a point, Theorem 1.9 follows when combining the results in [CS] and [Ba]. And to prove Theorem 1.9 in general, we need to generalize the results in both [CS] and [Ba] to higher dimensions. To this end we provide a proof generalizing Call-Silverman’s result in the appendix; see Theorem A.1. A related result sufficient for our purposes has also been established recently by Yuan and Zhang [YZ2]. To generalize Baker’s result to higher dimensional subvarieties, we establish a geometric dynamical Bogomolov theorem for split maps; see Theorem 4.1.

It is worth pointing out that Habegger’s proof of Theorem 1.9 for maps coming from products of elliptic curves in [Ha1, Theorem 1.3] did not rely on the analogous geometric Bogomolov theorem, though the latter had at the time already been established by Gubler [Gu2]. Instead, Habegger used local monodromy and the Betti map and as a corollary, he then deduced a new proof of the geometric Bogomolov conjecture for products of elliptic curves over 1-dimensional function fields. This approach was later generalized to 1-parameter families of abelian varieties by Gao and Habegger [GH] who thus established the geometric Bogomolov conjecture over 1-dimensional function fields. In [DGH3] the authors generalized their work to families of abelian varieties over a base of arbitrary dimension. This inequality, coupled with the results of Gao on the mixed Ax-Schanuel conjecture [Ga1, Ga2], played a key role in their proof of uniformity in the Mordell-Lang conjecture and for curves of large height.

1.6. A discussion of the proof of Theorems 1.2 and 1.6. Let us now briefly describe the strategy of our proof of Theorems 1.2 and 1.6. First we point out that an application of Bézout’s theorem allows us to deduce Theorem 1.2 from a special case of Theorem 1.6 and its complimentary Theorem 5.6 in the case that Φ has an isotrivial component. It

is worth pointing out here that Dimitrov, Gao and Habegger [DGH2] combined the full strength of the relative Bogomolov conjecture [DGH2, Conjecture 1.2] with their work in [DGH3] to establish uniformity in the classical Mordell-Lang and Bogomolov problems. To prove Theorem 1.6, we first use Theorem 1.9. This allows us to apply the recent arithmetic equidistribution theorem of Yuan and Zhang [YZ2]. And we do so multiple times – a very fruitful idea in Bogomolov-type problems that was first employed by Ullmo [U1]. To leverage the output of this equidistribution result, we use the theory of currents as well as a local Hodge index theorem [YZ1]. Perhaps surprisingly, these tools allow us to reduce Theorem 1.6 to the complex analytic Theorem 1.8. This comes in stark contrast with Kühne’s proofs in [Ku1, Ku2] and with [DGH1, DGH3] which relied on a far stronger ‘functional transcendence’ result, known as the mixed Ax-Schanuel conjecture [Ga2]. In particular, we get a new, simplified proof of special cases of Kühne’s aforementioned results which relies solely on Ax’s theorem [Ax].

It is worth pointing out that the proof of [DKY2], which built on the strategy from [DKY1], requires careful estimates of a certain energy pairing at each Berkovich space corresponding to a place of \mathbb{Q} . Such estimates are non-trivial, especially in the setting of non-polynomial maps. Our proof bypasses these issues. Indeed, it only requires the complex analytic Theorem 1.8, which ensures that certain archimedean energy pairings are non-zero; see Proposition 3.6.

1.7. Structure of the paper. The structure of this paper is as follows. In §2 we briefly recall some background and terminology that we use throughout this article. In §3 we establish Theorem 1.8 in the stronger form of Theorem 3.1. In §4 we establish Theorem 1.9 as well as an analogous result for curves \mathbf{X} and maps Φ having an isotrivial component; see Theorem 4.9. We do so by establishing the geometric dynamical Bogomolov theorems 4.1 and 4.3. In §5 we combine our results to establish Theorem 1.6 and its complimentary Theorem 5.6. We deduce Theorem 1.2 as a consequence of Theorem 5.6 and of a special case of Theorem 1.6. We further deduce Corollary 1.3. In §6 we establish Theorem 1.1. Finally, in the appendix we generalize Call-Silverman’s specialization result [CS, Theorem 4.1] to higher dimensional subvarieties.

Acknowledgements: The first-named author would like to thank Laura DeMarco and Lars Kühne for many interesting conversations about topics related to this article over the last few years. The second-named author heartily thanks Robert Wilms for many enlightening discussions surrounding adelic line bundles. Both authors thank Laura DeMarco and Dragos Ghioca for valuable comments and suggestions on a previous version of this paper that greatly

improved the current presentation. We would also like to thank the University of Basel for the great working environment it provided during the initial stages of preparation of this article.

2. PRELIMINARIES

A notion of adelic metrized line bundles was introduced by Zhang [Zh1] in the number field case and by Gubler [Gu3] in the function field case. We begin by briefly summarizing the relevant terms for our article. For a more exhaustive introduction we direct the readers to the survey articles of Chambert-Loir [CL2] and Yuan [Yu2] as well as to the recent book by Yuan and Zhang [YZ2] and the references therein. An excellent source is also the appendices of [YZ1].

First let \mathfrak{K} be an algebraically closed field, complete with respect to a non-trivial archimedean or non-archimedean absolute value $|\cdot|$. Let Y and L be a projective variety, respectively a line bundle on Y both defined over \mathfrak{K} . Let Y^{an} denote the Berkovich analytification of Y and L^{an} to be the analytification of L as a line bundle of Y^{an} . For $x \in Y^{\text{an}}$, we let $\mathfrak{K}(x)$ be its residue field. A *continuous metric* $\|\cdot\|$ on Y is defined to be a collection of $\mathfrak{K}(x)$ -norms $\|\cdot\|_x$ one on each fiber $L^{\text{an}}(x)$ for $x \in Y^{\text{an}}$ such that for each rational section s of L the map $x \mapsto \|s(x)\|_x := \|s(x)\|$ is continuous on Y^{an} away from the poles of s . The pair $(L, \|\cdot\|)$ is called a (continuous) metrized line bundle on Y . A metric is called *semipositive* if it is a uniform limit of semipositive model metrics as in [YZ1, Definition 5.1]. Limits of metrics are taken with respect to the topology induced by the supremum norm. A metrized line bundle L on Y is *integrable* if it can be written as $\bar{L} = \bar{L}_1 - \bar{L}_2$, where \bar{L}_i are semipositive metrized line bundles on Y .

Associated to a semipositive metrized line bundle \bar{L} , there is a curvature form $c_1(\bar{L})$, defined on the Berkovich analytification Y^{an} . One can then define a positive measure $c_1(\bar{L})^{\dim(Y)}$ on Y^{an} . In the non-archimedean places the curvature forms and associated measures are constructed by Chambert-Loir [CL1]. We refer the reader to [CL2] for a more detailed exposition.

By Chambert-Loir [CL1], Gubler [Gu1] and Chambert-Loir-Thuillier [CLT] we can define local heights of subvarieties Z of Y , though not uniquely, which can be computed through integration against measures obtained from the aforementioned curvature forms. More precisely, if $n = \dim Z$, \bar{L}_i are integrable metrized line bundles and we choose sections ℓ_i of L_i with disjoint support for $i = 1, \dots, n$, then the local height of Z with respect to (ℓ_1, \dots, ℓ_n)

can be defined inductively through

$$(2.1) \quad \widehat{\operatorname{div}(\ell_0)} \cdots \widehat{\operatorname{div}(\ell_n)} \cdot [Z] := \widehat{\operatorname{div}(\ell_1)} \cdots \widehat{\operatorname{div}(\ell_n)} \cdot [\operatorname{div}(\ell_0) \cdot Z] - \int_{Y^{\text{an}}} \log \|\ell_0\|_{c_1(\bar{L}_1)} \cdots c_1(\bar{L}_n) \delta_Z.$$

A summary for intersection numbers sufficient for our purposes can be found in [YZ1]. We point out here that in the special case that $Y = C$ is a curve and \bar{L}_0, \bar{L}_1 are defined on trivial line bundles $L_0 = L_1 = \mathcal{O}_C$, we may use the regular sections 1_{L_0} and 1_{L_1} to define the local intersection number of the metrics of L_0 and L_1 , as follows

$$(2.2) \quad \bar{L}_0 \cdot \bar{L}_1 := \widehat{\operatorname{div}(1_{L_0})} \cdot \widehat{\operatorname{div}(1_{L_1})} \cdot [C].$$

Now let \mathcal{F} be either a number field or the function field of a smooth projective curve defined over $\bar{\mathbb{Q}}$. Denote by $M_{\mathcal{F}}$ the set of places v of \mathcal{F} normalized so that the product formula

$$(2.3) \quad \prod_{v \in M_{\mathcal{F}}} |x|_v^{n_v} = 1$$

holds for each nonzero $x \in \mathcal{F}$. Here $n_v = [\mathcal{F}_v : \mathbb{Q}_v]$ if \mathcal{F} is a number field, whereas $n_v = 1$ in the function field case. For each $v \in M_{\mathcal{F}}$ we denote by \mathcal{F}_v a completion of \mathcal{F} with respect to v and write $\bar{\mathcal{F}}_v$ for its algebraic closure and \mathbb{C}_v for its completion.

Let Y be an irreducible projective variety over \mathcal{F} and L a line bundle on Y . A (continuous) *semipositive adelic metric* on L is a collection $\{\|\cdot\|_v\}$ of continuous semipositive \mathcal{F}_v -metrics $\|\cdot\|_v$ on L over all places v of \mathcal{F} satisfying certain coherence conditions; [Yu2, Gu3]. We denote the pair of our line bundle and the family of metrics $\|\cdot\|_v$ by

$$\bar{L} := (L, \{\|\cdot\|_v\})$$

and call \bar{L} an *adelic (semipositive) metrized line bundle*. An adelic metric on L is called *integrable* if it is the difference of two semipositive adelic metrics. Following [YZ1], we denote the space of integrable adelic metrized line bundles on Y by $\widehat{\operatorname{Pic}}(Y)_{\text{int}}$.

For this article, the most important example of an adelic semipositive metrized line bundle is the so-called *ϕ -invariant metric* \bar{L}_{ϕ} associated to an endomorphism $\phi : Y \rightarrow Y$ defined over \mathcal{F} with $\phi^*L \simeq L^{\otimes d}$ for some $d \geq 2$. This metric was defined by Zhang [Zh1] following Tate's limiting argument.

Zhang [Zh1, Theorem (1.4)], Gubler [Gu1, Gu2, Gu3] and Chambert-Loir-Thuillier [CLT] developed a height theory for subvarieties and an intersection theory for integrable adelic metrized line bundles for number fields and function fields respectively. So denoting the set of $n \geq 0$ dimensional cycles of Y by $Z_n(Y)$, their respective intersection numbers are written as

$$(2.4) \quad \begin{aligned} \widehat{\text{Pic}}(Y)_{\text{int}}^{n+1} \times Z_n(Y) &\rightarrow \mathbb{R} \\ (\bar{L}_1, \dots, \bar{L}_n, D) &\mapsto \bar{L}_1 \cdots \bar{L}_n \cdot D = \bar{L}_1|_D \cdots \bar{L}_n|_D. \end{aligned}$$

When $D = Y$ we omit it from the above notation.

Using the intersection numbers in (2.4) one can give a notion for the height of a subvariety associated to any $\bar{L} \in \widehat{\text{Pic}}(Y)_{\text{int}}$. More specifically, if Z is an irreducible closed subvariety of Y defined over $\bar{\mathcal{F}}$, we denote by Z_{gal} the closed \mathcal{F} -subvariety of Y corresponding to Z and write

$$(2.5) \quad h_{\bar{L}}(Z) = \frac{\bar{L}^{\dim Z+1} \cdot Z_{\text{gal}}}{(\dim Z + 1) \deg_L(Z_{\text{gal}})} \in \mathbb{R}.$$

In the case that $Z = \{x\}$ is just a point in $Y(\bar{\mathcal{F}})$, one has

$$(2.6) \quad h_{\bar{L}}(x) = -\frac{1}{|\text{Gal}(\bar{\mathcal{F}}/\mathcal{F}) \cdot x|} \sum_{v \in M_{\mathcal{F}}} \sum_{z \in \text{Gal}(\bar{\mathcal{F}}_v/\mathcal{F}_v)x} n_v \log \|s(x)\|_v,$$

where s is meromorphic section of L that is regular and non-vanishing at x .

As an example, in the case of the ϕ -invariant metric \bar{L}_ϕ , the height $h_{\bar{L}_\phi}$ is the canonical height $\hat{h}_{L,\phi}$ of ϕ introduced by Call-Silverman [CS] taken with respect to the line bundle L (multiplied by the degree $[\mathcal{F} : \mathbb{Q}]$ in the number field case.)

For $i = 1, \dots, \dim Y + 1$ we have the i th successive minimum

$$(2.7) \quad e_i(Y, \bar{L}) = \sup_Z \inf_{P \in Y(\bar{\mathcal{F}}) \setminus Z} h_{\bar{L}}(P),$$

where Z ranges over all closed subsets of codimension i in Y . Zhang [Zh1] and Gubler [Gu2, Lemma 4.1, Proposition 4.3] in the setting of number fields and function fields respectively proved a fundamental inequality establishing a relationship between these successive minima and the height of Y .

Lemma 2.1 (Zhang-Gubler fundamental inequality). *Assume that \bar{L} is a semipositive continuous adelicly metrized line bundle on an irreducible projective d -dimensional variety Y*

defined over \mathcal{F} . Then

$$\frac{e_1(Y, \overline{L}) + \cdots + e_{d+1}(Y, \overline{L})}{\dim Y + 1} \leq h_{\overline{L}}(Y) \leq e_1(Y, \overline{L}).$$

We finish this section with a convention. Throughout this article when $\phi = (f_1, \dots, f_\ell) : \mathbb{P}_1^\ell \rightarrow \mathbb{P}_1^\ell$ is a split polarized endomorphism over \mathcal{F} with polarization degree $d \geq 2$ we will be working with the line bundle

$$L_0 = \{\infty\} \times \mathbb{P}_1 \cdots \times \mathbb{P}_1 + \cdots + \mathbb{P}_1 \times \cdots \times \mathbb{P}_1 \times \{\infty\},$$

giving a polarization. We will abbreviate the height from (2.5) in this case by

$$(2.8) \quad \hat{h}_\phi(\cdot) := h_{(\overline{L}_0)_\phi}(\cdot).$$

3. FROM MEASURES TO WEAKLY SPECIAL CURVES

Our goal in this section is to prove Theorem 1.8. In fact we will prove a more general result applicable to non-polarized endomorphisms.

Recall that for a rational map $f \in \mathbb{C}(z)$ with $\deg f \geq 2$ there is an invariant measure μ_f supported on the Julia set of f , denoted by J_f . This measure was constructed by Lyubich [Ly] and Freire-Lopes-Mañé [FLM]. It is the unique measure with no point masses and such that $f^*\mu_f = (\deg f)\mu_f$. We will show that the following holds.

Theorem 3.1. *Let $f_i \in \mathbb{C}(z)$ be rational maps of (not necessarily equal) degree $\deg(f_i) \geq 2$ for $i = 1, 2$. Let $C \subset \mathbb{P}_1^2$ be an irreducible complex algebraic curve with dominant projections $\pi_i : C \rightarrow \mathbb{P}_1$ for $i = 1, 2$. Then $\deg(\pi_2)\pi_1^*\mu_{f_1} = \deg(\pi_1)\pi_2^*\mu_{f_2}$ if and only if $\text{supp}(\pi_1^*\mu_{f_1}) \cap \text{supp}(\pi_2^*\mu_{f_2}) \neq \emptyset$ and there exist positive integers ℓ, k such that $\deg(f_1)^\ell = \deg(f_2)^k$ and C is weakly (f_1^ℓ, f_2^k) -special.*

Our proof relies on results from complex dynamics. For basic notions and results we refer the reader to [Mi]. We will use Koebe's 1/4-theorem, which we recall here for the reader's convenience. For $a \in \mathbb{C}$ and $r \in \mathbb{R}$ we write

$$B(a, r) = \{z \in \mathbb{C} : |z - a| < r\}.$$

Theorem 3.2 (Koebe 1/4-theorem). *Suppose that $f : B(a, r) \rightarrow \mathbb{C}$ is a univalent function. Then $f(B(a, r)) \supseteq B(f(a), \frac{r}{4}|f'(a)|)$.*

Let J_f be the Julia set of a rational function $f \in \mathbb{C}(w)$ with $\deg f \geq 2$. A holomorphic function $h : U \rightarrow \mathbb{C}$ defined on a neighborhood U of $z_0 \in J_f$ is called a *symmetry* of J_f if

- $h(U \cap J_f) = h(U) \cap J_f$; and
- if J_f is a circle, the entire Riemann sphere or a line segment, there is a constant $\alpha > 0$ such that for any Borel set A on which $h|_A$ is injective we have $\mu_f(h(A)) = \alpha\mu_f(A)$.

A family \mathfrak{S} of symmetries of J_f is called *non-trivial* if \mathfrak{S} is normal **and** no limit function of \mathfrak{S} is constant. The following result is classical and will be central in our proof. It combines Levin's results [Le] as well as the work of Douady and Hubbard [DH], which relied on ideas by Thurston.

Theorem 3.3 ([DH],[Le]). *If the Julia set of a rational map f has an infinite non-trivial family of symmetries, then f is exceptional.*

Next we record here the following lemma, which was essential also in Levin-Przytycki's proof of Theorem 3.1 for the case of the diagonal curve [LP].

Lemma 3.4. *Let f be an ordinary rational function of degree ≥ 2 defined over \mathbb{C} . Then there exists a set $E_f \subset J_f$ with $\mu_f(E_f) > \frac{1}{2}$ and numbers $r_f > 0$ and $\kappa_f > 1$, so that for each $x \in E_f$ one can find increasing sequences of natural numbers $t_n \rightarrow \infty$ and $k_n \rightarrow \infty$ (as $n \rightarrow \infty$) such that for each $n \in \mathbb{N}$ the map $f^{k_n} : B(x, e^{-t_n}) \rightarrow \mathbb{P}_1$ is injective and we have*

$$B(f^{k_n}(x), r_f/\kappa_f) \subset f^{k_n}(B(x, e^{-t_n})) \subset B(f^{k_n}(x), r_f).$$

Proof. The map f has positive Lyapunov exponent and μ_f is an ergodic invariant measure (see [Ly, FLM, Ru]), so the result follows immediately from [LP, Lemma 1]. \square

We are now ready to prove Theorem 3.1.

Proof of Theorem 3.1: Let f_i , C and π_i be as in the statement of the theorem. We will first show that (1) implies (2). We have

$$(3.1) \quad \mu_C := \frac{\pi_1^* \mu_{f_1}}{\deg(\pi_1)} = \frac{\pi_2^* \mu_{f_2}}{\deg(\pi_2)}$$

and in particular

$$(3.2) \quad \pi_1^{-1}(J_{f_1}) = \pi_2^{-1}(J_{f_2}).$$

By the proof of [Mi, Theorem 4.2] equation (3.2) implies that J_{f_1} is a smooth set (a circle, a line segment or the entire sphere) if and only if J_{f_2} is. Moreover by [DH] a rational function is exceptional if and only if its Julia set is smooth (a circle, a line segment or the entire sphere) and has a smooth measure of maximal entropy. Therefore, by (3.1) we infer that f_1 is ordinary if and only if f_2 is.

Assume first that both f_i are ordinary and let $E_i := E_{f_i} \subset J_{f_i}$, $r_i := r_{f_i} > 0$ and $\kappa_i := \kappa_{f_i} > 1$ be the quantities associated to f_i defined in Lemma 3.4 for $i = 1, 2$. Since $\mu_C(\pi_i^{-1}(E_i)) = \mu_{f_i}(E_i) > \frac{1}{2}$ for $i = 1, 2$ we have

$$(3.3) \quad \mu_C(\pi_1^{-1}(E_1) \cap \pi_2^{-1}(E_2)) > 0.$$

Let $\mathcal{R} \subset C(\mathbb{C})$ be the (finite) set of ramification points of π_1 and π_2 . As μ_C does not charge points equation (3.3) ensures that there is $p_0 \in \pi_1^{-1}(E_1) \cap \pi_2^{-1}(E_2) \setminus \mathcal{R}$.

We set s to be a branch of $\pi_2 \circ \pi_1^{-1}$ near $\pi_1(p_0)$ and we may assume without loss of generality that

$$(3.4) \quad |s'(\pi_1(p_0))| < 1.$$

Indeed, either $|(\pi_2 \circ \pi_1^{-1})'(\pi_1(p_0))| < 1$ or $|(\pi_1 \circ \pi_2^{-1})'(\pi_2(p_0))| < 1$ and if the latter happens we may just reverse the roles of f_1, f_2 in what follows.

By (3.5) and applying Lemma 3.4 to $\pi_i(p_0) \in E_i \setminus \pi_i(\mathcal{R})$ we can find strictly increasing sequences $t_n \rightarrow \infty$ and $k_n^i \rightarrow \infty$ as $n \rightarrow \infty$ so that for each $n \in \mathbb{N}$, we have

$$(3.5) \quad \sup_{n \in \mathbb{N}} \sup \{|s'(z)| : z \in B(\pi_1(p_0), e^{-t_n})\} \leq 1,$$

the map

$$(3.6) \quad f_i^{k_n^i} : B(\pi_i(p_0), e^{-t_n}) \rightarrow \mathbb{P}_1$$

is injective, $B(\pi_i(p_0), e^{-t_n}) \cap \pi_i(\mathcal{R}) = \emptyset$ and

$$(3.7) \quad B(f_i^{k_n^i}(\pi_i(p_0)), r_i/\kappa) \subset f_i^{k_n^i}(B(\pi_i(p_0), e^{-t_n})) \subset B(f_i^{k_n^i}(\pi_i(p_0)), r_i),$$

where $\kappa := \max\{\kappa_1, \kappa_2\} > 1$. As the Julia set J_{f_i} is compact (in the topology induced by the chordal distance on $\mathbb{P}_1(\mathbb{C})$) passing to a subsequence we may also assume that

$$(3.8) \quad f_i^{k_n^i}(\pi_i(p_0)) \rightarrow p_i,$$

for some $p_i \in J_{f_i}$ as $n \rightarrow \infty$. Passing to a further subsequence we can thus infer that

$$(3.9) \quad B(f_1^{k_n^1}(\pi_1(p_0)), c_2 r_1) \subset B(p_1, c_1 r_1) \subset f_1^{k_n^1}(B(\pi_1(p_0), e^{-t_n})),$$

for positive constants $c_1, c_2 > 0$ and all $n \in \mathbb{N}$. From (3.9) and (3.5) we infer that

$$(3.10) \quad s \circ f_1^{-k_n^1}(B(p_1, c_1 r_1)) \subset B(\pi_2(p_0), e^{-t_n}),$$

so that by (3.7) and (3.8) after perhaps passing to another subsequence, it also holds that

$$(3.11) \quad f_2^{k_n^2} \circ s \circ f_1^{-k_n^1} B(p_1, c_1 r_1) \subset B(p_2, 2r_2).$$

By Montel's theorem we thus infer that the functions

$$\tilde{h}_n := f_2^{k_n^2} \circ s \circ f_1^{-k_n^1} : B(p_1, c_1 r_1) \rightarrow \mathbb{P}_1(\mathbb{C})$$

form a normal family.

Next, we will show that $\{\tilde{h}_n\}_n$ has no subsequence that converges to a constant function. Applying the Cauchy integral formula to $f_1^{k_n^1}$ at $\pi_1(p_0)$ and using (3.7) and (3.8) we deduce

$$(3.12) \quad |(f_1^{-k_n^1})'(f_1^{k_n^1}(\pi_1(p_0)))| \geq c_3 r_1^{-1} e^{-t_n},$$

for an constant $c_3 > 0$ and all $n \in \mathbb{N}$. From (3.9), inequality (3.12) and Koebe's 1/4-theorem we deduce

$$(3.13) \quad B(\pi_1(p_0), c_4 e^{-t_n}) \subset f^{-k_n^1}(B(p_1, c_1 r_1))$$

for some $c_4 > 0$. By Koebe's 1/4-theorem and the fact that $s'(\pi_1(p_0)) \neq 0$ we also have that

$$(3.14) \quad B(\pi_2(p_0), c_5 e^{-t_n}) \subset s(B(\pi_1(p_0), c_4 e^{-t_n}))$$

for some $c_5 > 0$ and all $n \in \mathbb{N}$. Similarly, applying the Cauchy integral formula to $f_2^{-k_n^2}$ at $f_2^{k_n^2}(\pi_2(p_0))$ and using (3.7) and (3.8) we get

$$(3.15) \quad |(f_2^{k_n^2})'(\pi_2(p_0))| \geq c_6 r_2 e^{t_n},$$

for $c_6 > 0$ and all $n \in \mathbb{N}$. In view of (3.15), Koebe's 1/4-theorem combined with (3.13) and (3.14) implies that

$$(3.16) \quad B(f_2^{k_n^2}(\pi_2(p_0)), c_7 r_2) \subset f_2^{k_n^2} \circ s \circ f_1^{-k_n^1}(B(p_1, c_1 r_1))$$

for a positive constant c_7 . This proves our claim that $\{\tilde{h}_n\}_{n \in \mathbb{N}}$ has no subsequence which converges to a constant function.

Our next task is to construct a non-trivial family of symmetries of J_{f_1} . To this end let h^* be a limit function of the family $\{\tilde{h}_n\}_{n \in \mathbb{N}}$. We can choose a ball $B^* := B(p^*, r^*) \subset B(p_1, c_1 r_1)$ centered at $p^* \in J_{f_1}$ such that $h^*(B(p^*, 2r^*)) \cap \pi_2(\mathcal{R}) = \emptyset$ (indeed recalling that $\pi_1(p_0) \in J_{f_1}$ and that (3.8) holds we can choose $p^* = f_1^{k_n^1}(\pi_1(p_0))$ for some large n). As restrictions of sequences of functions converge to the restriction of the limit function, passing to a subsequence of \tilde{h}_n we may assume that $\tilde{h}_n(B^*) \cap \pi_2(\mathcal{R}) = \emptyset$ for all $n \in \mathbb{N}$ and $\tilde{h}_n \rightarrow h^*$

(which is non-constant). Now consider the functions defined by

$$(3.17) \quad h_n := s^{-1} \circ \tilde{h}_n : B^* \rightarrow \mathbb{P}_1(\mathbb{C}).$$

As we have seen, they form a normal family with no subsequence that converges to a constant function. Moreover if $A \subset B^*$ is a Borel measurable set (on which h_n is injective), then because of our assumption (3.1) and our construction we have

$$(3.18) \quad \mu_{f_1}(h_n(A)) = \frac{\deg(f_2)^{k_n^2}}{\deg(f_1)^{k_n^1}} \mu_{f_1}(A).$$

Thus we have a non-trivial family of symmetries of J_{f_1} . As f_1 is ordinary, by Theorem 3.3 we infer that $\{h_n\}$ is eventually a constant sequence. Thus there exist positive integers k, ℓ and non-negative integers N, M such that

$$(3.19) \quad f_2^{M+k} \circ s \circ f_1^{-(N+\ell)} = f_2^M \circ s \circ f_1^{-N}.$$

From (3.18) it follows directly that $\deg(f_2)^k = \deg(f_1)^\ell$. We are now going to show that

$$(3.20) \quad (f_1^{N+\ell}, f_2^{M+k})(C) = (f_1^N, f_2^M)(C).$$

Assuming (3.20) holds, as in [GNy1, Lemma 6.1] it follows that if $\pi_1(p)$ is preperiodic by f_1 for a point $p \in C(\mathbb{C})$ then $\pi_2(p)$ is preperiodic by f_2 . Thus we can choose p such that $\pi_1(p)$ is a repelling periodic point of f_1^ℓ that is neither a ramification point of π_1 nor of any iterate of f_1 . Then $\pi_2(p)$ is also preperiodic by f_2 . Taking $L \in \mathbb{N}$ such that $f_2^{Lk}(\pi_2(p))$ is fixed by f_2^{Lk} and $\pi_1(p)$ is fixed by $f_1^{L\ell}$, we see that $(f_1^{L\ell}, f_2^{Lk})(C)$ satisfies the conditions of [GNy1, Proposition 5.1] and the conclusion of our theorem follows.

To prove (3.20) consider the image of an arc

$$\mathcal{I} := \{(f_1^{-(N+\ell)}(x), s \circ f_1^{-(N+\ell)}(x)) : x \in B^*\}$$

which is well-defined by our construction. Since the inverse f^{-N} is well-defined it holds that

$$(3.21) \quad \begin{aligned} (f_1^{N+\ell}, f_2^{M+k})(\mathcal{I}) &= \{(x, f_2^M \circ s \circ f_1^{-N}(x)) : x \in B^*\} \\ &= \{(f_1^N \circ f_1^{-N}(x), f_2^M \circ s \circ f_1^{-N}(x)) : x \in B^*\} \\ &= \{(f_1^N(y), f_2^M \circ s(y)) : y \in f_1^{-N}(B^*)\}. \end{aligned}$$

We infer that $(f_1^{N+\ell}, f_2^{M+k})(\mathcal{I}) \subset (f_1^N, f_2^M)(C)$. It is also evident by (3.20) that $(f_1^{N+\ell}, f_2^{M+k})(\mathcal{I}) \subset (f_1^{N+\ell}, f_2^{M+k})(C)$, so

$$(f_1^{N+\ell}, f_2^{M+k})(\mathcal{I}) \subset (f_1^N, f_2^M)(C) \cap (f_1^{N+\ell}, f_2^{M+k})(C)$$

and by Bézout's theorem and the fact that C is irreducible our claim (3.20) follows. This finishes the proof in the case that both f_1, f_2 are ordinary.

If now both f_1 and f_2 are exceptional then the statement becomes more classical and should be known. We have not found a proof in the literature and thus provide a quick sketch here. First note that if one of the groups, say G_{f_1} is an elliptic curve then the other has to be as well as the support of μ_{f_1} is $\mathbb{P}_1(\mathbb{C})$ and if $G_{f_2} = \mathbb{G}_m$ then the support is a topological circle or an interval. If both groups G_{f_1}, G_{f_2} are elliptic curves then we can employ the arguments in [DM2]. In place of Bertrand's theorem [Ber, Théorém 3] we can use the main theorem in [Ax]. Assume now that $G_{f_1} = G_{f_2} = \mathbb{G}_m$. Then $\Psi_{f_i}^* \mu_{f_i}$ is a rational multiple of the Haar measure on the circle S^1 , denoted by μ_{S^1} for $i = 1, 2$. Let H be an irreducible component of $(\Psi_{f_1}, \Psi_{f_2})^{-1}(C)$ and let $\tilde{\pi}_1, \tilde{\pi}_2$ be the two canonical projections from $H \subset \mathbb{G}_m^2$ to \mathbb{G}_m . The first assumption of the theorem then implies that $\tilde{\pi}_1^* \mu_{S^1} = \alpha \tilde{\pi}_2^* \mu_{S^1}$ for $\alpha \in \mathbb{Q}^*$. It follows that there exists a non-constant analytic curve $\gamma : (0, 1) \rightarrow H$ such that $\pi_i \circ \gamma \in S^1$ for $i = 1, 2$ and such that $\tilde{\pi}_1, \tilde{\pi}_2$ are injective on the support of γ and that

$$\int_{1/2}^t dg_1 = \alpha \int_{1/2}^t dg_2$$

where $g_i = \log \tilde{\pi}_i \circ \gamma, i = 1, 2$ for all t in a neighborhood of $1/2$. We infer that $\log \pi_i \circ \gamma(t), i = 1, 2$ are linearly related over \mathbb{Q} modulo a constant function. By Ax's theorem [Ax] we infer that H is a coset.

We will now show that (2) implies (1). Assume that the curve C , which projects dominantly into both coordinates, is weakly (f_1, f_2) -special. We will show that (3.1) holds. First assume that both maps f_1 and f_2 are ordinary so that $\deg(f_1^k) = \deg(f_2^\ell)$ and C is (f_1^k, f_2^ℓ) -preperiodic for $k, \ell \geq 1$. If both maps f_i and C are defined over $\overline{\mathbb{Q}}$, then the result is known to hold. Indeed by [GNY1, Lemma 6.1] the curve C contains infinitely many (f_1^k, f_2^ℓ) -preperiodic points. The (3.1) follows by [GNY1, Proposition 4.1]. Otherwise, assume that f_1, f_2 and C are defined over the function field $K = \overline{\mathbb{Q}}(V)$ of a variety V defined over $\overline{\mathbb{Q}}$ with positive dimension. We can spread the f_i and C out to get endomorphisms $f_{i,U} : U \times \mathbb{P}_1 \rightarrow U \times \mathbb{P}_1$ and a family of curves $\mathcal{C} \subset U \times \mathbb{P}_1^2$ on a Zariski open and dense subset U of V with the property that fiberwise $f_{i,t}$ have degree $\deg f_i$ and \mathcal{C}_t is irreducible for $t \in U$ and $i = 1, 2$. Then for each $t \in U(\overline{\mathbb{Q}})$, the maps $f_{i,t}$ and the curve \mathcal{C}_t are defined

over $\overline{\mathbb{Q}}$ and by [GNy1, Lemma 6.1] and [GNy1, Proposition 4.1] we infer that

$$(3.22) \quad \frac{\pi_{1,t}^* \mu_{f_{1,t}}}{\deg(\pi_{1,t})} = \frac{\pi_{2,t}^* \mu_{f_{2,t}}}{\deg(\pi_{2,t})}.$$

Since $U(\overline{\mathbb{Q}})$ is dense in $V(\mathbb{C})$ in the euclidean topology and the measures above vary continuously, we infer that (3.22) holds for all $t \in V(\mathbb{C})$. In particular, (3.22) holds for the point t giving the embedding of $\overline{\mathbb{Q}}(V)$ into \mathbb{C} and the conclusion follows. Assume now that f_1 and f_2 are exceptional. The existence of a positive dimensional coset dominating both factors implies that G_{f_1}, G_{f_2} are either both elliptic curves or both equal to \mathbb{G}_m . If they are both elliptic curves, then it is clear from the definition of the Betti coordinates that they are linearly related and equation 3.1 follows once we express the measures in terms of the Betti coordinates as in [CDMZ, §4 Addendum]. If $G_{f_1} = G_{f_2} = \mathbb{G}_m$ then by our assumption we have a coset H in \mathbb{G}_m^2 with $C = (\Psi_{f_1}, \Psi_{f_2})(H)$. Denoting its two projections by $\tilde{\pi}_i : H \rightarrow \mathbb{G}_m$, for $i = 1, 2$, it suffices now to show that $\tilde{\pi}_i^* \mu_{S^1}$ are linearly related for $i = 1, 2$. Our assumption that $\text{supp}(\pi_1^* \mu_{f_1}) \cap \text{supp}(\pi_2^* \mu_{f_2}) \neq \emptyset$ ensures that on H it holds that $\tilde{\pi}_1(z)^{k_1} \tilde{\pi}_2(z)^{k_2} = \theta$ for two non-zero rational integers k_1, k_2 and $\theta \in S^1$. Then our claim follows taking logarithms. This completes the proof of our theorem. \square

3.1. Applications of Theorem 1.8. We proceed to discuss the consequences of Theorem 1.8 which will be crucial in our proof of Theorem 1.6 (hence also Theorem 1.2.)

For a curve $\mathbf{C} \subset \mathbb{P}_1 \times \mathbb{P}_1$ we let $\pi_i : \mathbf{C} \rightarrow \mathbb{P}_1$ for $i = 1, 2$ denote its projections into the first and second coordinate. For each i and $t \in B_0(\mathbb{C})$ we write $\pi_{i,t} : \mathcal{C}_t \rightarrow \mathbb{P}_1$ for the corresponding specialization in the fiber \mathcal{C}_t . We denote by $|\cdot|$ the standard absolute value in \mathbb{C} and for $x, y \in \mathbb{C}$ we write

$$\|(x, y)\| = \max\{|x|, |y|\}.$$

We first record the following easy lemma.

Lemma 3.5. *Let $\Phi : \mathbb{P}_1^2 \rightarrow \mathbb{P}_1^2$ be a split endomorphism and let $\mathbf{C} \subset \mathbb{P}_1^2$ be an irreducible curve, both defined over K . Then for all $t \in B_0(\mathbb{C})$ outside of a finite set $E_{\Phi, \mathbf{C}} \subset B_0(\overline{\mathbb{Q}})$ the following hold:*

- (1) \mathcal{C}_t is an irreducible curve of genus equal to the genus of \mathbf{C} ;
- (2) $\deg(\pi_{i,t}) = \deg(\pi_i)$, for $i = 1, 2$;
- (3) either \mathbf{f}_i is exceptional or the map $f_{i,t}$ is ordinary, for $i = 1, 2$;

(4) The homogeneous lifts $F_{i,t}$ of $f_{i,t}(X) = \frac{a_d(t)X^d + \dots + a_0(t)}{b_d(t)X^d + \dots + b_0(t)}$, given by

$$F_{i,t}(x, y) = (a_d(t)x^d + \dots + a_0(t)y^d, b_d(t)x^d + \dots + b_0(t)y^d),$$

define a function holomorphic in t , for $i = 1, 2$.

Proof. There are several ways to prove Lemma 3.5 and we sketch one. For (1) one can choose for example a planar smooth model for \mathbf{C} and use the genus degree formula to see that the fibres \mathcal{C}_t where the genus drops or \mathcal{C}_t is reducible is finite and consists of $\overline{\mathbb{Q}}$ -points. For (3) we note that an equation for \mathbf{f}_i induces a map $B \rightarrow \mathcal{M}_d$ into the moduli space of rational maps of degree $d = \deg(\mathbf{f}_i)$. The exceptional maps correspond to subvarieties of \mathcal{M}_d (defined over $\overline{\mathbb{Q}}$) and thus the claim follows. The conditions (2) and (4) are clearly satisfied outside a finite set. \square

In what follows we will often refer to the set $E_{\Phi, \mathbf{C}}$ as it is defined in Lemma 3.5. The following consequence of Theorem 1.8 will be crucial in our proof of Theorem 1.6.

Proposition 3.6. *Let $\Phi : \mathbb{P}_1^2 \rightarrow \mathbb{P}_1^2$ be a split polarized endomorphism and let $\mathbf{C} \subset \mathbb{P}_1^2$ be an irreducible curve that is not weakly Φ -special, both defined over K . Then for each compact set $\mathcal{D} \subset B_0 \setminus E_{\Phi, \mathbf{C}}$, there a continuous function $\psi_{\mathcal{D}} : \pi|_{\mathcal{C}}^{-1}(\mathcal{D}) \rightarrow \mathbb{R}$ with the property that*

$$(3.23) \quad \int_{\mathcal{C}_t} \psi_{\mathcal{D}}|_{\mathcal{C}_t} ((\deg \pi_2)\pi_{1,t}^* \mu_{f_{1,t}} - (\deg \pi_1)\pi_{2,t}^* \mu_{f_{2,t}}) < 0,$$

for each $t \in \mathcal{D}$ such that \mathcal{C}_t is not weakly Φ_t -special.

Proof. Recall first that both projections π_i are dominant since \mathbf{C} is not weakly ϕ -special. For $i = 1, 2$ we let $L_i = \pi_i^*(\infty)$ and denote the specializations at $t \in B_0(\mathbb{C})$ by $L_{i,t} = \pi_{i,t}^*(\infty)$. We write $d_i := \deg(\pi_i)$. Assume that both Φ and \mathcal{C} are defined over a number field \mathcal{K} . We fix an embedding $\mathcal{K} \subset \mathbb{C}$ and in what follows we work locally at the fixed archimedean place corresponding to the standard absolute value. Since our place is fixed we omit the index v from the subsequent notations.

Let $t \in B_0(\mathbb{C}) \setminus E_{\Phi, \mathbf{C}}$. As recalled in §2 and defined by Zhang [Zh1] (see also [GNY1]), we have an invariant local metric on $L_{i,t}$, which we denote by $\overline{L}_{i,t} = (L_{i,t}, \|\cdot\|_{i,t})$. In particular outside of the support of the canonical meromorphic section 1 of $L_{i,t}$ we have

$$-\log \|1\|_{i,t}(z) = G_{F_{i,t}}(\pi_{i,t}(z), 1) = \lim_{n \rightarrow \infty} \frac{\log \|F_{i,t}^n(\pi_{i,t}(z), 1)\|}{d^n},$$

where $F_{i,t}$ is the lift from Lemma 3.5 and we view $\pi_{i,t}(z)$ as a point in \mathbb{C} (identified with its image on \mathbb{P}_1 .) Here $G_{F_{i,t}} : \mathbb{C}^2 \setminus \{(0,0)\} \rightarrow \mathbb{R}$ is the escape rate given by

$$G_{F_{i,t}}(x, y) = \lim_{n \rightarrow \infty} \frac{\log \|F_{i,t}^n(x, y)\|}{d^n}.$$

Recall that by construction and since $t \notin E_{\Phi, \mathbf{C}}$ we have $c_1(\bar{L}_{i,t}) = \pi_{i,t}^* \mu_{f_{i,t}}$. Since the line bundle $M_t := d_2 L_{1,t} - d_1 L_{2,t}$ has degree 0, it has an associated flat metric, which we denote by $\bar{M}_t^0 = (M_t, \log \theta_t)$ satisfying $c_1(\bar{M}_t^0) \equiv 0$ as in [YZ1, Definition 5.10]. Here $\log \theta_t$ is the Green function of M_t . Consider the (local) metrized line bundle

$$(3.24) \quad \bar{N}_t := \overline{d_2 L_{1,t} - d_1 L_{2,t} - M_t^0} = d_2 \bar{L}_{1,t} - d_1 \bar{L}_{2,t} - \bar{M}_t^0.$$

Since \bar{M}_t^0 is flat, we have

$$c_1(\bar{N}_t) = d_2 c_1(\bar{L}_{1,t}) - d_1 c_1(\bar{L}_{2,t}) = d_2 \pi_{1,t}^* \mu_{f_{1,t}} - d_1 \pi_{2,t}^* \mu_{f_{2,t}}.$$

Since \bar{N}_t is supported on the trivial line bundle $\mathcal{O}_{\mathcal{C}_t}$, the local Hodge-index [YZ, Theorem 2.1] states that the local pairing satisfies $(\bar{N}_t \cdot \bar{N}_t) \leq 0$ with equality if and only if \bar{N}_t is a trivially metrized line bundle. But if the latter happens, then we have $d_2 c_1(\bar{L}_{1,t}) = d_1 c_1(\bar{L}_{2,t})$, which by Theorem 1.8 implies that \mathcal{C}_t is weakly Φ_t -special. Thus for each $t \in B_0(\mathbb{C}) \setminus E_{\Phi, \mathbf{C}}$ such that \mathcal{C}_t is not weakly Φ_t -special, the local pairing is strictly negative; that is

$$\int_{\mathcal{C}_t} (d_2 G_{F_{1,t}}(\pi_{1,t}(z), 1) - d_1 G_{F_{2,t}}(\pi_{2,t}(z), 1) - \log \theta_t(z)) (d_2 d\pi_{1,t}^* \mu_{f_{1,t}}(z) - d_1 d\pi_{2,t}^* \mu_{f_{2,t}}(z)) < 0.$$

Now let $\mathcal{D} \subset B_0 \setminus E_{\Phi, \mathbf{C}_1}$ be a compact set. Choose a smooth function $\chi_{\mathcal{D}}$ so that $\chi_{\mathcal{D}} \equiv 1$ on \mathcal{D} and $\chi_{\mathcal{D}} \equiv 0$ outside a small neighborhood of \mathcal{D} and define

$$(3.25) \quad \psi_{\mathcal{D}}(t, z) = \chi_{\mathcal{D}}(t) (d_2 G_{F_{1,t}}(\pi_{1,t}(z), 1) - d_1 G_{F_{2,t}}(\pi_{2,t}(z), 1) - \log \theta_t(z)),$$

for $t \in \mathcal{D}$ and $z \in \mathcal{C}_t(\mathbb{C})$. We have shown that this function satisfies (3.23). It remains to show that it is continuous.

Since by Lemma 3.5 the lift $F_{i,t}$ is holomorphic for t in the compact \mathcal{D} , from [HP, FS] or the proof of [BH, Proposition 1.2], we know that the escape rate $G_{F_{i,t}}(x, y)$ is continuous as a function of $(t, (x, y)) \in \mathcal{D} \times \mathbb{C}^2 \setminus \{(0,0)\}$. Moreover, for each $t \in \mathcal{D}$ the function $\pi_{i,t}(z)$ is continuous outside of its set of poles in $L_{i,t}$ and since the escape rate scales logarithmically (i.e. $G_{F_{i,t}}(\alpha x, \alpha y) = G_{F_{i,t}}(x, y) + \log |\alpha|$ for each $\alpha \in \mathbb{C} \setminus \{0\}$) we infer that

$$d_2 G_{F_{1,t}}(\pi_{1,t}(z), 1) - d_1 G_{F_{2,t}}(\pi_{2,t}(z), 1)$$

is continuous as a function of $z \in \mathcal{C}_t$ with logarithmic singularities at M_t (with multiplicities as prescribed there.) The function θ_t has been chosen exactly so that these logarithmic singularities ‘cancel out’. Indeed recall that for each t , the $\log \theta_t$ may be expressed as $\sum_{P \in M_t} \log h \circ \iota_P$ (taking multiplicities into account), where ι_P is the embedding of \mathcal{C}_t into its Jacobian $\text{Jac}(\mathcal{C}_t)$ via a base point P and h is given as in [GH, p.304]. By construction $\log \theta_t$ is smooth on \mathcal{C}_t outside of the support of M_t and has logarithmic singularities with multiplicities given by M_t . Moreover it depends locally smoothly on the period matrix of $\text{Jac}(\mathcal{C}_t)$ for $t \in B_0 \setminus E_{\Phi, \mathbf{c}}$ because of the first condition in Lemma 3.5 and as M_t has degree 0 also globally smoothly on t . Therefore $\psi_{\mathcal{D}}$ is continuous. This completes the proof of our proposition. \square

4. A GEOMETRIC BOGOMOLOV THEOREM: PROOF OF THEOREM 1.9 AND APPLICATIONS

Let $\ell \geq 2$ and $\Phi : \mathbb{P}_1^\ell \rightarrow \mathbb{P}_1^\ell$ be an isotrivial-free split polarized endomorphism and let $\mathbf{X} \subset \mathbb{P}_1^\ell$ be an irreducible projective subvariety of \mathbb{P}_1^ℓ defined over K . In this section we aim to prove Theorem 1.9. As it turns out, this amounts to establishing the dynamical Bogomolov conjecture for split maps over the function field K , which is the content of Theorem 4.1.

Throughout this section the line bundle

$$L = \{\infty\} \times \mathbb{P}_1 \cdots \times \mathbb{P}_1 + \cdots + \mathbb{P}_1 \times \cdots \times \mathbb{P}_1 \times \{\infty\}$$

is fixed and we henceforth write $h_{\overline{L}_\Phi} = \hat{h}_\Phi$ for canonical height from (2.5).

Theorem 4.1. *Let $\ell \geq 2$ and $\Phi = (\mathbf{f}_1, \dots, \mathbf{f}_\ell) : \mathbb{P}_1^\ell \rightarrow \mathbb{P}_1^\ell$ be an isotrivial-free split polarized endomorphism such that each \mathbf{f}_i has degree $d \geq 2$ and is defined over K . Let $\mathbf{X} \subset \mathbb{P}_1^\ell$ be an irreducible subvariety defined over K with dimension ≥ 1 . Then the following are equivalent:*

- (1) $\hat{h}_\Phi(\mathbf{X}) = 0$;
- (2) \mathbf{X} contains a generic¹ sequence $\{a_n\} \subset \mathbf{X}(\overline{K})$ of points with $\hat{h}_\Phi(a_n) \rightarrow 0$ ²;
- (3) \mathbf{X} is Φ -preperiodic.

Moreover, we prove an analogous geometric Bogomolov result in the case that \mathbf{X} is a curve and Φ is not isotrivial-free. To state it, we need to define the notion of an isotrivial curve.

¹The sequence $\{a_n\} \subset \mathbf{X}(\overline{K})$ is generic if every infinite subsequence of $\{a_n\}$ is Zariski dense in \mathbf{X} .

²We say that $\{a_n\}$ is a \hat{h}_Φ -small sequence.

Definition 4.2. We say that $\Phi = (\mathbf{f}_1, \mathbf{f}_2)$ is isotrivial if both \mathbf{f}_1 and \mathbf{f}_2 are isotrivial. If so, let \mathbf{M}_i be Möbius transformations defined over \overline{K} such that $\mathbf{M}_i \circ \mathbf{f}_i \circ \mathbf{M}_i^{-1}$ is defined over $\overline{\mathbb{Q}}$ for $i = 1, 2$. A curve $\mathbf{C} \subset \mathbb{P}_1 \times \mathbb{P}_1$ is Φ -isotrivial if $(\mathbf{M}_1, \mathbf{M}_2)(\mathbf{C})$ is defined over $\overline{\mathbb{Q}}$.

Note that this is a field theoretic condition that does not reflect deep dynamical properties. The analogous geometric Bogomolov statement is then as follows.

Theorem 4.3. *Let $\mathbf{C} \subset \mathbb{P}_1^2$ be a curve defined over K whose projection to each factor \mathbb{P}_1 is dominant. Let $\Phi = (\mathbf{f}_1, \mathbf{f}_2) : \mathbb{P}_1^2 \rightarrow \mathbb{P}_1^2$ be a split polarized endomorphism of polarization degree $d \geq 2$ defined over K . If Φ is isotrivial, then \mathbf{C} contains a generic \hat{h}_Φ -small sequence if and only if \mathbf{C} is Φ -isotrivial. Moreover, if exactly one of \mathbf{f}_1 or \mathbf{f}_2 is isotrivial, then \mathbf{C} does not contain a generic \hat{h}_Φ -small sequence.*

As a corollary, we obtain the following statement which will be useful in the proof of Theorem 1.2 in the non-isotrivial-free case.

Corollary 4.4. *Let $\phi = (\mathbf{f}_1, \mathbf{f}_2) : \mathbb{P}_1^2 \rightarrow \mathbb{P}_1^2$ be a split polarized endomorphism of polarization degree $d \geq 2$ defined over K such that at least one of $\mathbf{f}_1, \mathbf{f}_2$ is isotrivial and let \mathbf{f} an endomorphism of $\mathbb{P}_{1,K}$ of degree d . If ϕ is isotrivial assume that \mathbf{C} is not ϕ -isotrivial. Let (\mathbf{X}, Φ) be either $(\mathbb{P}_1 \times \mathbf{C}, (\mathbf{f}, \phi))$ or $(\mathbf{C} \times \mathbf{C}, (\phi, \phi))$. Then $\hat{h}_\Phi(\mathbf{X}) > 0$.*

Proof. The proof follows by Theorem 2.1 in view of Gubler's inequality in Lemma 2.1. \square

To place these results in the context of the work of Dimitrov-Gao-Habegger [DGH1, DGH3], Kühne [Ku2] and Yuan-Zhang [YZ2] we make the following remark.

Remark 4.5. Subvarieties with $\hat{h}_\Phi(\mathbf{X}) \neq 0$ coincide with non-degenerate subvarieties in the sense of [YZ2]. This follows by Lemma A.3. One can also infer that the two conditions coincide a posteriori by comparing [YZ2, Lemma 5.22] and Gauthier-Vigny's result [GV, Theorem B]. Thus, in the language of [YZ2], Theorem 4.1 characterizes the non-degenerate subvarieties.

Our proof is inspired by the proof of the analogous statement over number fields in Mimar's thesis [Mi] and in [GNY1] for the case of curves. For the higher dimensional case we are inspired by [GNY2], which itself relied on an idea used in the proof of [YZ1, Theorem 4.13]. It should be mentioned, that we do not use an equidistribution theorem as in [GNY1, GNY2]. Instead we solely rely on the Hodge index theorem which has recently been established in

the function field setting by [Ca], Gubler's inequality as in Lemma 2.1 and a result of Baker [Ba] characterizing points with geometric canonical height equal to zero.

In the case that Φ is isotrivial-free, these ingredients allow us to reduce the geometric Bogomolov statement of Theorem 4.1 to the dynamical Manin-Mumford over \mathbb{C} and consequently we use the results in [GNY1, GNY2, GT, Gu2]. In the setting of Theorem 4.3, as we shall see, the mathematical content of the proof is slightly different given that the role of preperiodic curves is replaced by that of isotrivial curves.

Before we begin with our proof we point out that the following holds.

Remark 4.6. If $\mathbf{f} : \mathbb{P}_1 \rightarrow \mathbb{P}_1$ is defined over K and is not isotrivial then \mathbf{f} is exceptional if and only if \mathbf{f} is a flexible Lattès map so that the group $G_{\mathbf{f}}$ from Definition 1.4 is an elliptic curve and $\Psi_{\mathbf{f}}$ is given by multiplication by an integer.

We start by proving Theorem 4.1 in the case of hypersurfaces.

Proposition 4.7. *Let $\ell \geq 2$ and $\Phi = (\mathbf{f}_1, \dots, \mathbf{f}_\ell) : \mathbb{P}_1^\ell \rightarrow \mathbb{P}_1^\ell$ be an isotrivial-free split polarized endomorphism defined over K . Let $\mathbf{H} \subset \mathbb{P}_1^\ell$ be an irreducible hypersurface defined over K which projects dominantly onto any subset of $\ell - 1$ coordinate axis. Assume that \mathbf{H} contains a generic sequence $\{a_n\} \subset \mathbf{H}(\overline{K})^3$ with $\hat{h}_\Phi(a_n) \rightarrow 0$ as $n \rightarrow \infty$. Then the following hold:*

(1) *There exists $c_1, \dots, c_\ell \in \mathbb{R}$ not all equal to zero such that*

$$(4.1) \quad \sum_{j=1}^{\ell} c_j \hat{h}_{\mathbf{f}_j}(\mathbf{x}_j) = 0,$$

for all $\mathbf{x} = (\mathbf{x}_1, \dots, \mathbf{x}_\ell) \in \mathbf{H}(\overline{K})$; and

(2) *\mathbf{H} is Φ -preperiodic. Moreover, either $l = 2$ or all maps $\mathbf{f}_i, i = 1, \dots, \ell$ are exceptional.*

Proof. First we show that (1) implies (2). Permuting the \mathbf{f}_j if necessary we may assume that $c_\ell \neq 0$ in (4.1). Let $\mathbf{x} = (\mathbf{x}_1, \dots, \mathbf{x}_\ell) \in \mathbf{H}(\overline{K})$ be such that $\mathbf{x}_i \in \text{Prep}(\mathbf{f}_i)$ and so $\hat{h}_{\mathbf{f}_i}(\mathbf{x}_i) = 0$ for all $i = 1, \dots, \ell - 1$. Then (4.1) yields that $\hat{h}_{\mathbf{f}_\ell}(\mathbf{x}_\ell) = 0$. Since \mathbf{f}_ℓ is not isotrivial, [Ba, Theorem 1.6] implies that $\mathbf{x}_\ell \in \text{Prep}(\mathbf{f}_\ell)$ as well. Henceforth we embed $\overline{K} \hookrightarrow \mathbb{C}$ and abusing the notation slightly we view Φ and \mathbf{H} as being defined over $\overline{K} \subset \mathbb{C}$. Let $\pi_\ell : \mathbf{H} \rightarrow \mathbb{P}_1^{\ell-1}$ be the projection to the first $\ell - 1$ coordinates and note that it is a dominant map by our

³the term generic here means that every infinite subsequence of $\{a_n\}_{n \in \mathbb{N}}$ is Zariski dense in \mathbf{H} .

assumption. Let $\Pi := \text{Prep}(\mathbf{f}_1) \times \cdots \times \text{Prep}(\mathbf{f}_{\ell-1}) \subset \mathbb{P}_1^{\ell-1}$. We have established that

$$(4.2) \quad \pi_\ell^{-1}(\Pi) \subset \text{Prep}(\Phi) \cap \mathbf{H}.$$

Moreover the Zariski closure $\overline{\Pi}$ clearly contains a generic sequence of $\Phi_\ell := (\mathbf{f}_1, \dots, \mathbf{f}_{\ell-1})$ -preperiodic points (since each \mathbf{f}_i has infinitely many preperiodic points.) Hence from (4.2) we see that \mathbf{H} contains a generic sequence of Φ -preperiodic points. Now if $\ell > 2$, then by [GNY2, Theorem 1.2] and in view of Remark 4.6 we infer that all components \mathbf{f}_i of Φ are Lattés maps corresponding to integer-multiplications on elliptic curves. By Gubler's result [Gu2] (or by [Ha1, Theorem 1.4]) it follows that \mathbf{H} is Φ -preperiodic. If $\ell = 2$, then the conclusion follows by [GNY1, Theorem 1.1] if one of the \mathbf{f}_i is ordinary or by [Gu2] otherwise.

It thus suffices to prove that (1) holds. Let $\Phi_i = (\mathbf{f}_1, \dots, \mathbf{f}_{i-1}, \mathbf{f}_{i+1}, \dots, \mathbf{f}_\ell) : \mathbb{P}_1^{\ell-1} \rightarrow \mathbb{P}_1^{\ell-1}$. As in §2 to each Φ_i we can associate a Φ_i -invariant adelic metrized line bundle \overline{L}_{Φ_i} (with underlying line bundle chosen so that $h_{\overline{L}_{\Phi_i}} = \hat{h}_{\Phi_i}$.) Now let $\pi_i : \mathbf{H} \rightarrow \mathbb{P}_1^{\ell-1}$ be the projection forgetting the i -th factor and restricted to \mathbf{H} , and define $\overline{L}_i = \pi_i^*(\overline{L}_{\Phi_i})$, for $i = 1, \dots, \ell$.

Using additive notation for our line bundles we let $\overline{M} = \sum_{i=1}^{\ell} \overline{L}_i$. For each vector $\vec{c} = (c_1, \dots, c_\ell) \in \mathbb{Q}^\ell \setminus \vec{0}$ we write $L_{\vec{c}} = \sum_{i=1}^{\ell} c_i L_i$ so that $\overline{L}_{\vec{c}} = \sum_{i=1}^{\ell} c_i \overline{L}_i$ is the associated adelically metrized line bundle on \mathbf{H} . We proceed, using an argument inspired by [YZ1, Theorem 4.13] as in [GNY2, Lemma 5.3], to show that there is a choice of $\vec{c} \neq \vec{0}$ such that the height associated to $\overline{L}_{\vec{c}}$ is identically equal to zero as in (1). To this end, we will use the Hodge index theorem [Ca, Theorem 3.1] applied to appropriate metrized line bundles. We first note that exactly as in [GNY2, Claim 5.4] there is $\vec{c} \neq 0$ which can be chosen to have rational entries so that

$$(4.3) \quad \begin{aligned} L_{\vec{c}} \cdot M^{\ell-2} &= 0 \\ L_{\vec{c}}^2 \cdot M^{\ell-3} &= 0. \end{aligned}$$

From now on we fix such \vec{c} . Then by the first Lefschetz-type theorem [YZ, Theorem 5.20], $L_{\vec{c}}$ is numerically trivial. By our assumption that \mathbf{H} contains a generic sequence of \hat{h}_Φ -small points, we have

$$e_i(\mathbf{H}, \overline{M} + m_1 \overline{L}_{i_1} + m_2 \overline{L}_{i_2}) = 0,$$

for all $i = 1, \dots, \ell$, $m_1, m_2 \in \mathbb{N}$ and $i_1, i_2 = 1, \dots, \ell$ (recall the definition of successive minima from (2.7)). Therefore, using Gubler's inequality as in Theorem 2.1 we infer that $(\overline{M} + m_1 \overline{L}_{i_1} + m_2 \overline{L}_{i_2})^\ell = 0$ as well. Thus, by the multilinearity of the intersection numbers

as in [GNY2, equation (5.4.4)] we have

$$(4.4) \quad \overline{L}_{\mathcal{E}}^2 \cdot \overline{M}^{\ell-2} = 0.$$

Moreover if m is large enough so that all coefficients of the \overline{L}_i in both $m\overline{M} + \overline{L}_{\mathcal{E}}$ and $m\overline{M} - \overline{L}_{\mathcal{E}}$ are non-negative, we have that both metrized line bundles $m\overline{M} + \overline{L}_{\mathcal{E}}$ and $m\overline{M} - \overline{L}_{\mathcal{E}}$ are semipositive. In other words, $\overline{L}_{\mathcal{E}}$ is \overline{M} -bounded. Recalling that (4.3) also hold, we are almost in the right setting to apply [Ca, Theorem 3.1] except that \overline{M} may not be arithmetically positive (as in [Ca, Definition 2.4])⁴. To fix this we modify \overline{M} by a small factor; see [YZ, Theorem 4.13] and [Ca, page 97] for this argument in the number field and function field setting respectively. Let $\pi : \mathbf{H} \rightarrow \text{Spec}(K)$ be the structure morphism and let \overline{D} be an adelic metrized line bundle on $S := \text{Spec}(K)$ such that $\overline{D} \cdot S > 0$. The modified $\overline{M}' := \overline{M} + \pi^*\overline{D}$ is now arithmetically positive and we still have that $\overline{L}_{\mathcal{E}}$ is \overline{M}' -bounded. Since $L_{\mathcal{E}}$ is numerically trivial, we see that still $L_{\mathcal{E}} \cdot M^{\ell-2} = 0$ and further the addition of the term $\pi^*\overline{D}$, leaves the intersection number $\overline{L}_{\mathcal{E}}^2 \cdot \overline{M}'^{\ell-2}$ equal to zero. We can therefore apply the Hodge index [Ca, Theorem 3.1] in conjunction with [Ca, Theorem 3.2] to get that $h_{\overline{L}}$ is identically zero on \mathbf{H} as claimed. \square

We are now ready to prove Theorem 4.1.

Proof of Theorem 4.1: That (1) and (2) are equivalent follows by Gubler's fundamental inequalities as in Lemma 2.1. In detail, first notice that the height \hat{h}_{Φ} takes non-negative values at points and as a result all the successive minima associated to \overline{L}_{Φ} are non-negative. It then follows by [Gu2, Lemma 4.1] that $\hat{h}_{\Phi}(\mathbf{X}) = 0$ if and only if the first successive minimum of $\overline{L}_{\Phi}|_{\mathbf{X}}$ is equal to zero or equivalently if (2) holds. That (3) implies (1) is well-known, see e.g [Fab, Theorem 3.2 (iv)]. It remains to see that (2) implies (3). But as in [GNY2, §2, Proposition 2.1] (see also [GT]), the situation reduces to that of Proposition 4.7. This completes our proof. \square

To prove Theorem 4.3 we also need the following result that is implicitly given in [Ba].

Theorem 4.8. [Ba] *Let $\mathbf{f} \in K(z)$ be a non-isotrivial rational map. For a preperiodic point $z_0 \in \mathbb{P}_1(\overline{K})$ denote by $\text{prep}(z_0)$ the cardinality of its orbit. Then the following holds.*

$$[K(z_0) : K] \rightarrow \infty, \text{ as } \text{prep}(z_0) \rightarrow \infty.$$

⁴Let $\pi : \mathbf{H} \rightarrow \text{Spec}(K)$ be the structure morphism. The adelic metrized line bundle \overline{M} is arithmetically positive if M is ample and $\overline{M} - \pi^*\overline{D}$ is nef for some adelic metrized line bundle \overline{D} on $\text{Spec}(K)$ with $\overline{D} \cdot \text{Spec}(K) > 0$.

Proof. This follows from the proof of [Ba, Theorem 1.6]. In fact the constant C'_2 in [Ba] scales linearly with $[K(z_0) : K]$ passing from K to the field extension $K(z_0)$. The other constants stay equal and thus using the fact that the orbit of z_0 by \mathbf{f} stays in $\mathbb{P}_1(K(z_0))$ the bound M gives an upper bound for $\text{prep}(z_0)$ and is bounded from above by a linear function in $[K(z_0) : K]$. \square

We can now prove Theorem 4.3.

Proof of Theorem 4.3: Recall that \mathbf{C} projects dominantly to both factors \mathbb{P}_1 . As in the proof of Theorem 4.1, using the Hodge index theorem [Ca, Theorem 3.1] and Gubler's fundamental inequalities in Lemma 2.1, we infer that if $\mathbf{C} \subset \mathbb{P}_1^2$ contains a generic \hat{h}_Φ -small sequence, then there exist constants $c_1, c_2 \in \mathbb{R}$, not both equal to zero, such that

$$(4.5) \quad c_1 \hat{h}_{\mathbf{f}_1}(z_1) + c_2 \hat{h}_{\mathbf{f}_2}(z_2) = 0,$$

for all $(z_1, z_2) \in \mathbf{C}(\overline{K})$. If one of c_1 or c_2 is zero then one of the coordinates of \mathbf{C} is constant which contradicts our assumption that \mathbf{C} projects dominantly onto each factor. Thus $c_1 c_2 \neq 0$ and we can deduce, as in the proof of Theorem 4.1, that \mathbf{C} contains a Zariski-dense set of points of height 0. If Φ is isotrivial and $(\mathbf{M}_1, \mathbf{M}_2)$ is a pair of Möbius transformations such that $(\mathbf{M}_1, \mathbf{M}_2) \circ \Phi \circ (\mathbf{M}_1^{-1}, \mathbf{M}_2^{-1})$ is defined over $\overline{\mathbb{Q}}$, then in view of [De1, Theorem 1.1] this implies that $(\mathbf{M}_1, \mathbf{M}_2)(\mathbf{C})$ contains infinitely many algebraic points (defined over $\overline{\mathbb{Q}}$). Therefore, $(\mathbf{M}_1, \mathbf{M}_2)(\mathbf{C})$ is in fact defined over $\overline{\mathbb{Q}}$ as claimed.

For the converse implication note that if $\overline{\mathbf{M}}_1, \overline{\mathbf{M}}_2$ are Möbius maps such that $(\mathbf{M}_1, \mathbf{M}_2) \circ \Phi \circ (\mathbf{M}_1^{-1}, \mathbf{M}_2^{-1})$ is defined over $\overline{\mathbb{Q}}$ and $(\mathbf{M}_1, \mathbf{M}_2)(\mathbf{C})$ is defined over $\overline{\mathbb{Q}}$, then \mathbf{C} contains infinitely many points P with $\hat{h}_\Phi(P) = 0$.

Assume now that \mathbf{f}_1 is isotrivial and \mathbf{f}_2 is not. In view of [Ba, Corollary 1.8], our assumptions imply that the curve $(\mathbf{M}_1, \text{id})(\mathbf{C})$ contains infinitely many points such that the first coordinate is in $\mathbb{P}_1(\overline{\mathbb{Q}})$ (and thus in particular in $\mathbb{P}_1(K)$) and the second coordinate is an \mathbf{f}_2 -preperiodic point. However by Theorem 4.8 it holds that $[K(z_n) : K] \rightarrow \infty$ as $n \rightarrow \infty$, for any infinite sequence of distinct points z_n that are \mathbf{f}_2 -preperiodic. This is a contradiction to the assumption that \mathbf{C} projects dominantly to each factor \mathbb{P}_1 . This completes the proof of this theorem. \square

We are now ready to deduce Theorem 1.9.

Proof of Theorem 1.9: If \mathbf{X} has dimension equal to 0 (and is therefore a point) then the result follows combining Call-Silverman's result [CS, Theorem 4.1] and Baker's [Ba, Theorem 1.6], since Φ is isotrivial-free. We thus let $\ell \geq 2$ and $\mathbf{X} \subset \mathbb{P}_1^\ell$ be a positive dimensional irreducible

subvariety over K . First note that by Theorem 4.1 we have that $\hat{h}_{\Phi}(\mathbf{X}) = 0$ if and only if \mathbf{X} is an irreducible component of a Φ -preperiodic subvariety or equivalently $\mathcal{X}^{\Phi,*} = \emptyset$. Assume now that \mathbf{X} is not Φ -preperiodic, so that $\mathcal{X}^{\Phi,*} \neq \emptyset$. We need to show that $\mathcal{X}^{\Phi,*} \neq \emptyset$ is Zariski open and dense in \mathcal{X} . To this end, it suffices to show that there are at most finitely many $\mathcal{Z} \subset \mathcal{X}$ that are irreducible components of Φ -special subvarieties projecting dominantly onto B and are maximal with this property. Equivalently, we have to show that \mathbf{X} (which is not Φ -preperiodic) contains at most finitely many maximal Φ -preperiodic subvarieties. This in turn follows from our geometric Bogomolov Theorem 4.1 (a Manin-Mumford version would suffice). Now since \mathbf{X} is not Φ -preperiodic, by Theorem 4.1 we infer that $\hat{h}_{\Phi}(\mathbf{X}) > 0$. Thus by Theorem A.1 or by [YZ2, Theorem 6.5] (recalling Remark 4.5) we deduce that there is a proper Zariski closed $\mathcal{V} \subset \mathcal{X}$ so that our desired inequality (1.5) holds for all $P \in (\mathcal{X}^{\Phi,*} \setminus \mathcal{V})(\overline{\mathbb{Q}})$. We will show that our inequality holds for all $\overline{\mathbb{Q}}$ -points of $\mathcal{X}^{\Phi,*}$ by induction of the dimension on \mathcal{X} . If \mathcal{X} has dimension equal to 1, then the exceptional set \mathcal{V} is merely a finite collection of points which we can incorporate by adjusting the constants in the inequality. Thus the base case follows. Now assume that \mathcal{X} has dimension at least 2. We decompose \mathcal{V} into finitely many irreducible components \mathcal{V}_i for $i = 1, \dots, s$. So it suffices to show that the inequality holds at all points in $\mathcal{V}_i \cap \mathcal{X}^{\Phi,*}$ for $i = 1, \dots, s$. Notice that \mathcal{V}_i has codimension at least one in \mathcal{X} , so we may induct on its dimension and apply our theorem to each \mathcal{V}_i . Adjusting the constants, we conclude that the inequality (1.5) holds for all $\overline{\mathbb{Q}}$ -points in $(\mathcal{X}^{\Phi,*} \setminus \mathcal{V}) \cup \mathcal{V}_1^{\Phi,*} \cup \dots \cup \mathcal{V}_s^{\Phi,*} = \mathcal{X}^{\Phi,*}$. This completes our proof. \square

In fact, arguing exactly as in the proof above and using Corollary 4.4 in place of Theorem 4.1 we infer that the following holds.

Theorem 4.9. *Let $\mathbf{C} \subset \mathbb{P}_1^2$ be a curve over K and $\Phi : \mathbb{P}_1^2 \rightarrow \mathbb{P}_1^2$ be a polarized split map $\Phi = (\mathbf{f}_1, \mathbf{f}_2)$ such that at least one of $\mathbf{f}_1, \mathbf{f}_2$ is isotrivial. If Φ is isotrivial assume that \mathbf{C} is not Φ -isotrivial. Then $\mathcal{C}^{\Phi,*}$ is Zariski open in \mathcal{C} and there are constants $c_1 = c_1(\Phi, \mathbf{C}) > 0$ and $c_2 = c_2(\Phi, \mathbf{C})$ such that*

$$(4.6) \quad \hat{h}_{\Phi}(P) \geq c_1 h_{\mathcal{N}}(\pi(P)) - c_2, \text{ for all } P \in \mathcal{C}^{\Phi,*}(\overline{\mathbb{Q}}).$$

4.1. Applications of Theorem 1.9: equidistribution and non-triviality of measures. Towards the proof of Theorems 1.6 and 1.2 we will need an arithmetic equidistribution result. To this end, we rely on recent work of Yuan and Zhang [YZ2], who generalized Kühne's equidistribution theorem [Ku2, Theorem 1], by building a theory of adelic metrized

line bundles on essentially quasiprojective varieties. Using this theory Yuan [Yu1] has recently provided a new proof of [Ku2, Theorem 2 and Theorem 3].

Let Φ and \mathbf{X} be as in §4. Associated to the pair (Φ, \mathbf{X}) we have a pair (Φ, \mathcal{X}) defined over a number field \mathcal{K} , where $\Phi : B_0 \times \mathbb{P}_1^\ell \rightarrow B_0 \times \mathbb{P}_1^\ell$ and $\mathcal{X} \subset B_0 \times \mathbb{P}_1^\ell$. In what follows we fix an embedding $\mathcal{K} \subset \mathbb{C}$ and as a result an archimedean place $|\cdot|_\infty$ of \mathcal{K} extending the classical absolute value in \mathbb{Q} . By [YZ2, §6.1.1, Theorem 6.1] we can associate an invariant adelic line bundle \overline{L}_Φ to the endomorphism Φ which has the property that $\Phi^* \overline{L}_\Phi \simeq \overline{L}_\Phi^{\otimes d}$. We further have a canonical measure $\mu_{\overline{L}_\Phi|_{\mathcal{X}}}$ defined in $\mathcal{X}(\mathbb{C})$ as in [YZ2, §3.6.6].

Definition 4.10. We say that $Z := \{x_n\}_n \subset \mathcal{X}(\overline{\mathbb{Q}})$ is a (Φ, \mathcal{X}) -generic small sequence if every infinite subsequence of Z is Zariski dense in \mathcal{X} and moreover $\hat{h}_\Phi(x_n) \rightarrow 0$ as $n \rightarrow \infty$.

We can now state the equidistribution theorem that will be crucial for our proof.

Proposition 4.11. *Let $\Phi : \mathbb{P}_1^\ell \rightarrow \mathbb{P}_1^\ell$ be an isotrivial-free, split polarized endomorphism and $\mathbf{X} \subset \mathbb{P}_1^\ell$ an irreducible subvariety that is not Φ -preperiodic, both defined over K . Let $\{x_n\}_n \subset \mathcal{X}(\overline{\mathbb{Q}})$ be a (Φ, \mathcal{X}) -generic small sequence. Then the Galois orbit of $\{x_n\}_n$ is equidistributed over $\mathcal{X}(\mathbb{C})$ with respect to the canonical measure $\mu_{\overline{L}_\Phi|_{\mathcal{X}}}$.*

Proof. Recalling Remark 4.5, this is an immediate consequence of our geometric Bogomolov Theorem 4.1 coupled with [YZ2, Theorem 1.6]. \square

We point out here that [YZ2, Theorem 1.6] yields a similar equidistribution statement for each place $v \in M_{\mathcal{K}}$. We shall only need their result at the fixed archimedean place. Kühne has pointed out that his proof of [Ku2, Theorem 1] adapts to this dynamical setting and it would be sufficient for us. Gauthier [Gau] has also recently established a related equidistribution result. It is likely that the assumptions in his theorem are satisfied in our setting.

We conclude this section by observing the following simple but crucial for us consequence of Theorem 1.9 and (a very special case of) Proposition 4.11.

Proposition 4.12. *Under the same assumptions as in Proposition 4.11, assume further that there is a (Φ, \mathcal{X}) -generic small sequence. Let $F \subset B_0(\overline{\mathbb{Q}})$ be a finite set. Then there is a compact set $\mathcal{D} = \mathcal{D}(F) \subset B_0 \setminus F$ such that $\mu_{\overline{L}_\Phi|_{\mathcal{X}}}(\pi|_{\mathcal{X}}^{-1}(\mathcal{D})) > 0$.*

Proof. Let $F \subset B_0(\mathbb{C})$ be a finite set. Let (t_n, z_n) be a generic (Φ, \mathcal{X}) -small sequence. By Theorem 1.9 we know that $\mathcal{X}^{\Phi, \star}$ is Zariski dense in \mathcal{X} , so we may assume passing to a

subsequence that $(t_n, z_n) \in \mathcal{X}^{\Phi, *(\overline{\mathbb{Q}})}$ for all $n \in \mathbb{N}$. Let h be a height on B associated to an ample divisor. Invoking Theorem 1.9 we infer that $h(t_n) \leq M$ for a constant M depending only on (Φ, \mathbf{X}) . This in turn prevents a positive proportion of Galois conjugates of t_n from accumulating near F ; see Masser-Zannier's [MZ2, Lemma 8.2]. More precisely, there is a compact set $\mathcal{D} = \mathcal{D}(F, \Phi, \mathbf{X}) \subset B_0(\mathbb{C}) \setminus F$ and a constant $a > 0$ depending only on Φ, \mathbf{X} and F such that

$$(4.7) \quad \frac{1}{[\mathcal{K}(t_n) : \mathcal{K}]} \sum_{\sigma \in \text{Gal}(\mathcal{K}(t_n)/\mathcal{K})} \chi_{\mathcal{D}}(\sigma(t_n)) \geq a,$$

for all $n \in \mathbb{N}$, where $\chi_{\mathcal{D}}$ denotes the characteristic function of \mathcal{D} . Now we can find a decreasing non-negative sequence of continuous functions $\chi_{\mathcal{D}, \epsilon} : B^0 \times \mathbb{P}_1^{\ell} \rightarrow \mathbb{R}_{\geq 0}$ for $\epsilon > 0$ such that $\chi_{\mathcal{D}, \epsilon}(Z) = \chi_{\mathcal{D}}(\pi(Z)) = 1$, for Z with $\pi(Z) \in \mathcal{D}$ and $\chi_{\mathcal{D}, \epsilon}(Z) = 0$ for Z with $\pi(Z)$ away from a compact ϵ -neighborhood of \mathcal{D} . In particular $\chi_{\mathcal{D}, \epsilon}(Z) \geq \chi_{\mathcal{D}}(\pi(Z))$ for all Z . Applying Proposition 4.11 with test function $\chi_{\mathcal{D}, \epsilon}$ along our sequence of small points we infer

$$\frac{1}{[\mathcal{K}(t_n, z_n) : \mathcal{K}]} \sum_{\sigma \in \text{Gal}(\mathcal{K}(t_n)/\mathcal{K})} \sum_{\rho \in \text{Gal}(\mathcal{K}(t_n, z_n)/\mathcal{K}) : \rho|_{\mathcal{K}(t_n)} = \sigma} \chi_{\mathcal{D}, \epsilon}(\sigma(t_n), \rho(z_n)) \rightarrow \int \chi_{\mathcal{D}, \epsilon} d\mu_{\overline{L}_{\Phi}|_{\mathcal{C}}}.$$

The definition of $\chi_{\mathcal{D}, \epsilon}$ along with inequality (4.7) then allow us to conclude that

$$\int \chi_{\mathcal{D}, \epsilon} d\mu_{\overline{L}_{\Phi}|_{\mathcal{C}}} \geq a,$$

for each ϵ . Letting $\epsilon \rightarrow 0$ the lemma follows by monotone convergence. \square

In the case of non isotrivial-free maps, in view of Theorem 4.3, the above argument yields that the following holds.

Proposition 4.13. *With assumptions as in Theorem 4.9, assume further that there is a (Φ, \mathcal{C}) -generic small sequence. Let $F \subset B_0(\overline{\mathbb{Q}})$ be a finite set. Then there is a compact set $\mathcal{D} = \mathcal{D}(F) \subset B_0 \setminus F$ such that $\mu_{\overline{L}_{\Phi}|_{\mathcal{C}}}(\pi|_{\mathcal{C}}^{-1}(\mathcal{D})) > 0$.*

5. INSTANCES OF RELATIVE BOGOMOLOV FOR SPLIT MAPS

In this section we prove Theorem 1.6 and subsequently deduce Theorem 1.2 as well as Corollary 1.3. To this end we will need the notion of a fiber-wise Green current associated to a family of polarized endomorphisms, which we shall first introduce. For the basics on currents we refer the reader to [Aea].

Let Λ be a smooth quasi-projective complex variety. Let $\pi : \mathcal{A} \rightarrow \Lambda$ be a family of irreducible complex projective varieties of dimension $k \geq 1$, so that $\mathcal{A}_\lambda = \pi^{-1}(\{\lambda\})$ is an irreducible complex variety of dimension k for each $\lambda \in \Lambda$. Let $f : \mathcal{A} \rightarrow \mathcal{A}$ be complex family of endomorphisms, i.e. f is analytic and for each $\lambda \in \Lambda$ the map $f_\lambda : \mathcal{A}_\lambda \rightarrow \mathcal{A}_\lambda$ is a morphism. Assume that $f^*\mathcal{L} \simeq \mathcal{L}^{\otimes d}$ for a line bundle \mathcal{L} on \mathcal{A} and that for each $\lambda \in \Lambda$ we have $f_\lambda^*\mathcal{L}_\lambda \simeq \mathcal{L}_\lambda^{\otimes d}$, where $\mathcal{L}_\lambda = \mathcal{L}|_{\mathcal{A}_\lambda}$. We encode the above data by $(\Lambda, \mathcal{A}, f, \mathcal{L})$. We let $\widehat{\omega}$ be a continuous positive $(1, 1)$ -form on \mathcal{A} cohomologous to a multiple of \mathcal{L} such that $\omega_\lambda := \widehat{\omega}|_{\mathcal{A}_\lambda}$ is a Kähler form on $(\mathcal{A}_\lambda)_{\text{reg}}$. We assume that $\widehat{\omega}$ is **normalized** so that

$$(5.1) \quad \int_{\mathcal{A}_\lambda} \omega_\lambda^k = 1,$$

for each $\lambda \in \Lambda$. The sequence $d^{-n}(f^n)^*(\widehat{\omega})$ converges in the sense of currents to a closed positive $(1, 1)$ -current \widehat{T}_f with continuous potentials on \mathcal{A} . We refer to \widehat{T}_f as the *fiber-wise Green current* of f . It is closed and positive and has continuous potentials. Moreover, for all $\lambda \in \Lambda$ the slice $\widehat{T}_f|_{\mathcal{A}_\lambda}$ is well-defined and we have $f_\lambda^*\widehat{T}_f|_{\mathcal{A}_\lambda} = d\widehat{T}_f|_{\mathcal{A}_\lambda}$. For these facts we refer the reader to [GV, Proposition 9].

Remark 5.1. In the setting of Proposition 4.11, we have $\mu_{\overline{\mathcal{T}}_\Phi|\mathcal{X}} = \frac{1}{v}\widehat{T}_\Phi^{\dim \mathcal{X}} \wedge [\mathcal{X}]$, where $v > 0$ is a constant such that the right-hand side is a probability measure. This transpires from the construction of the measure in [YZ2]. We remark that by [GV, Theorem B] or [YZ2] we know that $v = \hat{h}_\Phi(\mathbf{X})$.

In what follows we work in the setting of Theorem 1.6. So $\Phi = (\mathbf{f}_1, \mathbf{f}_2) : \mathbb{P}_1^2 \rightarrow \mathbb{P}_1^2$ and $\Psi : \mathbb{P}_1^\ell \rightarrow \mathbb{P}_1^\ell$ for $\ell \geq 2$ are isotrivial-free split polarizable endomorphisms both of polarization degree $d \geq 2$, $\mathbf{C} \subset \mathbb{P}_1^2$ is an irreducible curve that is not weakly Φ -special and $\mathbf{X} \subset \mathbb{P}_1^\ell$ is an irreducible subvariety that is not Ψ -preperiodic all defined over K . These induce endomorphisms of quasiprojective varieties $\Phi : B_0 \times \mathbb{P}_1^2 \rightarrow B_0 \times \mathbb{P}_1^2$, $f_i : B_0 \times \mathbb{P}_1 \rightarrow B_0 \times \mathbb{P}_1$ for $i = 1, 2$ and $\Psi : B_0 \times \mathbb{P}_1^\ell \rightarrow B_0 \times \mathbb{P}_1^\ell$, as well as subvarieties $\mathcal{C} \subset B_0 \times \mathbb{P}_1^2$ and $\mathcal{X} \subset B_0 \times \mathbb{P}_1^\ell$ defined over $\overline{\mathbb{Q}}$. For $i = 1, 2$ we write

$$\begin{aligned} f_i \times_{B_0} \Psi : B_0 \times \mathbb{P}_1^\ell &\rightarrow B_0 \times \mathbb{P}_1^\ell \\ (t, z_1, z_2, \dots, z_\ell) &\mapsto (t, f_{i,t}(z_1), \Psi_t(z_2, \dots, z_\ell)). \end{aligned}$$

Finally, we fix several projection maps as in the following diagram.

$$(5.2) \quad \begin{array}{ccccc} \mathcal{C} \times_{B_0} \mathcal{X} & \xrightarrow{\tilde{\pi}_1} & \mathbb{P}_1 \times \mathcal{X} & \xrightarrow{q} & \mathcal{X} \\ & \searrow^{\tilde{\pi}_2} & \downarrow p & & \downarrow q_B \\ \mathcal{C} & \xrightarrow[\pi_2]{} & B_0 \times \mathbb{P}_1 & \xrightarrow{p_B} & B_0 \end{array}$$

Here π_i and $\tilde{\pi}_i$ denote the projections to the i -th copy of \mathbb{P}_1 for $i = 1, 2$.

Our first step to proving Theorem 1.6 is to apply the equidistribution result in Proposition 4.11 (in fact 3 times) and deduce the following equality of measures.

Lemma 5.2. *Under the assumptions in Theorem 1.6 assume further that there is a $(\Phi \times_{B_0} \Psi, \mathcal{C} \times_{B_0} \mathcal{X})$ -generic small sequence. Then there exists $\alpha \in \mathbb{R}_{>0}$ such that*

$$(5.3) \quad \pi_1^* p_* (\widehat{T}_{f_1 \times_{B_0} \Psi}^{\wedge(1+\dim \mathcal{X})}) \wedge [\mathbb{P}_1 \times \mathcal{X}] = \alpha \pi_2^* p_* (\widehat{T}_{f_2 \times_{B_0} \Psi}^{\wedge(1+\dim \mathcal{X})}) \wedge [\mathbb{P}_1 \times \mathcal{X}].$$

Proof. Let $z := \{z_n\}_{n \in \mathbb{N}}$ be the generic small sequence as in the statement. We will trace its journey through the diagram (5.2). Since $\mathbf{C} \times \mathbf{X}$ is not (Φ, Ψ) -preperiodic, Proposition 4.11 (see also Remark 5.1) yields that the Galois orbits of z equidistribute with respect to

$$P := \frac{1}{v} \widehat{T}_{\Phi \times_{B_0} \Psi}^{\wedge(1+\dim \mathcal{X})} \wedge [\mathcal{C} \times_{B_0} \mathcal{X}],$$

where $v \in \mathbb{R}_{>0}$. Moreover, for each $i = 1, 2$ the sequence $\tilde{\pi}_i(z) := \{\tilde{\pi}_i(z_n)\}_n$ is $(f_i \times_{B_0} \Psi, \mathbb{P}_1 \times \mathcal{X})$ -generic and small. Since $\mathbb{P}_1 \times \mathbf{X}$ is not (f_i, Ψ) -preperiodic, Proposition 4.11 yields that its Galois conjugates equidistribute with respect to $\frac{1}{v_i} \widehat{T}_{f_i \times_{B_0} \Psi}^{\wedge(1+\dim \mathcal{X})} \wedge [\mathbb{P}_1 \times \mathcal{X}]$ for $v_i > 0$. We infer

$$(5.4) \quad \tilde{\pi}_{i*} P = \frac{1}{v_i} \widehat{T}_{f_i \times_{B_0} \Psi}^{\wedge(1+\dim \mathcal{X})} \wedge [\mathbb{P}_1 \times \mathcal{X}],$$

for each $i = 1, 2$. The aforementioned equidistribution statements further imply that the Galois conjugates of $p \circ \tilde{\pi}_i(z) = \pi_i \circ \pi_{\mathcal{C}}(z)$ equidistribute with respect to

$$(5.5) \quad p_* \tilde{\pi}_{i*} P = \pi_{i*} \pi_{\mathcal{C}*} P.$$

Since π_i is a finite map, we have $\pi_i^* \pi_{i*} \pi_{\mathcal{C}*} P = \deg(\pi_i) \pi_{\mathcal{C}*} P$. Recalling (5.4) and pulling back (5.5) by π_i we thus infer

$$\frac{\deg(\pi_2)}{v_1} \pi_1^* p_* (\widehat{T}_{f_1 \times_{B_0} \Psi}^{\wedge(1+\dim \mathcal{X})}) \wedge [\mathbb{P}_1 \times \mathcal{X}] = \frac{\deg(\pi_1)}{v_2} \pi_2^* p_* (\widehat{T}_{f_2 \times_{B_0} \Psi}^{\wedge(1+\dim \mathcal{X})}) \wedge [\mathbb{P}_1 \times \mathcal{X}].$$

The lemma follows. \square

Next our goal is to express the measures in Lemma 5.2 fiberwise. First, we will reinterpret their definition so that ultimately the characterization of slicing of currents applies. Our end goal is to infer that (5.3) contradicts Proposition 3.6 due to our assumption that \mathbf{C} is not weakly Φ -special. We first record a couple of lemmata.

Lemma 5.3. *With notation as in diagram (5.2), there is a constant $\kappa_i > 0$ such that $\widehat{T}_{f_i \times_{B_0} \Psi}^{\wedge(1+\dim \mathcal{X})} \wedge [\mathbb{P}_1 \times \mathcal{X}] = \kappa_i(p^*\widehat{T}_{f_i} \wedge q^*(\widehat{T}_{\Psi}^{\wedge(\dim \mathcal{X})} \wedge [\mathcal{X}]))$, for $i = 1, 2$.*

Proof. We fix projections $\Pi_1 : (B_0 \times \mathbb{P}_1) \times_{B_0} (B_0 \times \mathbb{P}_1^\ell) \rightarrow B_0 \times \mathbb{P}_1$ and $\Pi_2 : (B_0 \times \mathbb{P}_1) \times_{B_0} (B_0 \times \mathbb{P}_1^\ell) \rightarrow B_0 \times \mathbb{P}_1^\ell$. The map $f_i \times_{B_0} \Phi$ is polarized with respect to $\mathcal{L} = \Pi_1^* \mathcal{L}_1 \otimes \Pi_2^* \mathcal{L}_2$, where $\mathcal{L}_1 = B_0 \times \{\infty\}$ on $B_0 \times \mathbb{P}_1$ and $\mathcal{L}_2 = B_0 \times (\{\infty\} \times \mathbb{P}_1 \cdots \times \mathbb{P}_1 + \cdots + \mathbb{P}_1 \times \cdots \times \mathbb{P}_1 \times \{\infty\})$ on $B_0 \times \mathbb{P}_1^\ell$. Let $\widehat{\omega}_i$ be a continuous positive $(1, 1)$ -form cohomologous to a multiple of \mathcal{L}_i and normalized as in (5.1). Then $\Pi_1^* \widehat{\omega}_1 + \Pi_2^* \widehat{\omega}_2$ is cohomologous to a multiple of \mathcal{L} and by the limiting construction of the fiber-wise Green current as in e.g. [GV, Proposition 9] we infer

$$(5.6) \quad \widehat{T}_{f_i \times_{B_0} \Psi} = c(\Pi_1^* \widehat{T}_{f_i} + \Pi_2^* \widehat{T}_{\Psi}).$$

for a constant $c > 0$ (to ensure that both sides are normalized as in (5.1)), so that

$$(5.7) \quad \widehat{T}_{f_i \times_{B_0} \Psi} \wedge [\mathbb{P}_1 \times \mathcal{X}] = c(p^* \widehat{T}_{f_i} + q^*(\widehat{T}_{\Psi} \wedge [\mathcal{X}]))$$

Notice now that since $B_0 \times \mathbb{P}^1$ is 2-dimensional we have $(p^* \widehat{T}_{f_i})^{\wedge k} = p^*(\widehat{T}_{f_i})^{\wedge k} = 0$ (in the sense of currents) for $k \geq 3$ and similarly looking at the dimensions we have $q^*(\widehat{T}_{\Psi} \wedge [\mathcal{X}])^{\wedge \dim \mathcal{X} + 1} = 0$. Moreover \mathbb{P}_1 is \mathbf{f}_i -preperiodic, so that $\hat{h}_{\mathbf{f}_i}(\mathbb{P}_1) = 0$. By [GV, Theorem B] (or [YZ, Lemma 5.22]) and since \widehat{T}_{f_i} is a positive current we also have $(p^* \widehat{T}_{f_i})^{\wedge 2} = 0$. Hence passing to the $(\dim \mathcal{X} + 1)$ -th power in (5.7) we get

$$(5.8) \quad \widehat{T}_{f_i \times_{B_0} \Psi}^{\wedge(1+\dim \mathcal{X})} \wedge [\mathbb{P}_1 \times \mathcal{X}] = c^{\dim \mathcal{X} + 1} (\dim \mathcal{X} + 1) (p^* \widehat{T}_{f_i} \wedge q^*(\widehat{T}_{\Psi}^{\wedge(\dim \mathcal{X})} \wedge [\mathcal{X}])).$$

The lemma follows. \square

Lemma 5.4. *With notation as in diagram 5.2 we have*

$$p_* q^*(\widehat{T}_{\Psi}^{\wedge(\dim \mathcal{X})} \wedge [\mathcal{X}]) = p_B^* q_{B^*}(\widehat{T}_{\Psi}^{\wedge(\dim \mathcal{X})} \wedge [\mathcal{X}]).$$

Proof. Chasing the diagram 5.2 we observe that if $g_1 : B_0 \rightarrow \mathbb{R}$ and $g_2 : \mathbb{P}_1 \rightarrow \mathbb{R}$ are smooth and compactly supported functions and we let $g'_2 : B_0 \times \mathbb{P}_1 \rightarrow \mathbb{R}$ be given by $g'_2(t, z) = g_2(z)$,

then

$$\begin{aligned}
 (5.9) \quad \langle p_* q^* (\widehat{T}_\Psi^{\wedge(\dim \mathcal{X})} \wedge [\mathcal{X}]), (g_1 \circ p_B) g'_2 \rangle &= \langle q^* (\widehat{T}_\Psi^{\wedge(\dim \mathcal{X})} \wedge [\mathcal{X}]), (g_1 \circ p_B \circ p)(g'_2 \circ p) \rangle \\
 &= \langle \widehat{T}_\Psi^{\wedge(\dim \mathcal{X})} \wedge [\mathcal{X}], g_1 \circ q_B \rangle \langle [\mathbb{P}_1], g_2 \rangle \\
 &= \langle p_B^* q_{B*} (\widehat{T}_\Psi^{\wedge(\dim \mathcal{X})} \wedge [\mathcal{X}]), (g_1 \circ p_B) g'_2 \rangle.
 \end{aligned}$$

Since by [DS, Proposition 2.2.5] the space of functions generated by $g_1 g_2$ is dense in the space of smooth and compactly supported functions $B_0 \times \mathbb{P}_1 \rightarrow \mathbb{R}$, the lemma follows. \square

Finally we record here the following approximation lemma.

Lemma 5.5. *With notation as in diagram (5.2), there exists a sequence of smooth currents $\{R_n\}_{n \in \mathbb{N}}$ on B_0 such that*

$$R_n \rightarrow q_{B*} (\widehat{T}_\Psi^{\wedge(\dim \mathcal{X})} \wedge [\mathcal{X}]); \quad \pi_i^* \widehat{T}_{f_i} \wedge \pi_i^* p_B^* R_n \rightarrow \pi_i^* \widehat{T}_{f_i} \wedge \pi_i^* p_B^* q_{B*} (\widehat{T}_\Psi^{\wedge(\dim \mathcal{X})} \wedge [\mathcal{X}]),$$

for $n \rightarrow \infty$ and $i = 1, 2$.

Proof. Since the pushforward and pullback are continuous operators in the space of currents, it is enough to prove that there exists a sequence of smooth currents \tilde{R}_n on \mathcal{X} such that

$$\begin{aligned}
 (5.10) \quad \tilde{R}_n &\rightarrow \widehat{T}_\Psi^{\wedge(\dim \mathcal{X})} \wedge [\mathcal{X}]; \text{ and} \\
 p^* \widehat{T}_{f_i} \wedge q^* \tilde{R}_n &\rightarrow p^* \widehat{T}_{f_i} \wedge q^* (\widehat{T}_\Psi^{\wedge(\dim \mathcal{X})} \wedge [\mathcal{X}]).
 \end{aligned}$$

To show (5.10) note that by [Aea, Proposition A.21] each plurisubharmonic (psh) function can be approximated by a decreasing sequence of smooth psh functions on each affine open set and by [BT, Theorem 2.1] wedging and taking the limit of decreasing psh functions commutes. Since currents act on forms with compact support it suffices to work on open affine sets and the lemma follows. \square

We are now ready to prove Theorem 1.6.

Proof of Theorem 1.6: Under the assumptions in Theorem 1.6, assume that there is a $(\Phi \times_{B_0} \Psi, \mathcal{C} \times_{B_0} \mathcal{X})$ -generic small sequence to end in a contradiction. Combining Lemmata 5.2, 5.3 and 5.3 and using the projection formula for currents we infer that

$$(5.11) \quad \pi_1^* \widehat{T}_{f_1} \wedge \pi_1^* p_B^* (q_{B*} (\widehat{T}_\Psi^{\wedge(\dim \mathcal{X})} \wedge [\mathcal{X}])) = \beta \pi_2^* \widehat{T}_{f_2} \wedge \pi_2^* p_B^* (q_{B*} (\widehat{T}_\Psi^{\wedge(\dim \mathcal{X})} \wedge [\mathcal{X}])),$$

for $\beta \in \mathbb{R}_{>0}$. To simplify our notation we let $\Pi_{\mathcal{C}} := p_B \circ \pi_i : \mathcal{C} \rightarrow B_0$ and write

$$(5.12) \quad R := q_{B*} (\widehat{T}_\Psi^{\wedge(\dim \mathcal{X})} \wedge [\mathcal{X}])$$

Let $E_{\Phi, \mathcal{C}}$ be the finite set as in Lemma 3.5. Since \mathbf{X} is not Ψ -preperiodic, by Proposition 4.12 we can find a compact set $\mathcal{D} \subset B_0 \setminus E_{\Phi, \mathcal{C}}$ such that

$$(5.13) \quad R \wedge [\mathcal{D}] > 0.$$

We will first show that $\beta = \deg(\pi_1)/\deg(\pi_2)$. Let $\chi_{\mathcal{D}} : B_0 \rightarrow \mathbb{R}$ be a non-negative smooth function approximating the characteristic function of \mathcal{D} , so that (5.11) yields

$$(5.14) \quad \int_{\mathcal{C}} \chi_{\mathcal{D}} \circ \Pi_{\mathcal{C}} \wedge \pi_1^* \widehat{T}_{f_1} \wedge \Pi_{\mathcal{C}}^* R = \beta \int_{\mathcal{C}} \chi_{\mathcal{D}} \circ \Pi_{\mathcal{C}} \wedge \pi_2^* \widehat{T}_{f_2} \wedge \Pi_{\mathcal{C}}^* R.$$

Let us pretend for a moment that R is smooth. Then by the characterization of slicing as in [BB, Proposition 4.3] equation (5.14) would yield

$$(5.15) \quad \int_{B_0} \left(\chi_{\mathcal{D}}(t) \int_{\mathcal{C}_t} \pi_{1,t}^* T_{f_{1,t}} \right) R = \beta \int_{B_0} \left(\chi_{\mathcal{D}}(t) \int_{\mathcal{C}_t} \pi_{2,t}^* T_{f_{2,t}} \right) R,$$

where $T_{f_{i,t}}$ denotes the slice of \widehat{T}_{f_i} above t . We recall that this slice is well-defined and furthermore $T_{f_{i,t}} = \mu_{f_{i,t}}$; see e.g. [GV, Propositions 4 and 9]. By Wirtinger's formula (see e.g. [CL2]) we further have $\int_{\mathcal{C}_t} \pi_{1,t}^* T_{f_{1,t}} = \deg(\pi_{1,t}) = \deg \pi_1$ for $t \in \mathcal{D}$, so that recalling (5.13) equation (5.15) yields that $\beta = \deg(\pi_1)/\deg(\pi_2)$ as claimed.

Of course, we have been falsely assuming that R is smooth. To reconcile this we will use Lemma 5.5 to reduce to the smooth case. We let $\{R_n\}_{n \in \mathbb{N}}$ be the sequence of smooth currents from Lemma 5.5. From the definition of slicing [BB, Proposition 4.3] we have that

$$\int_{\mathcal{C}} u \pi_i^* \widehat{T}_{f_i} \wedge \pi_{\mathcal{C}}^* R_n = \int_{B_0} \left(\int_{\mathcal{C}_t} u|_{\mathcal{C}_t} T_{f_{i,t}} \right) R_n,$$

for every smooth compactly supported function u on \mathcal{C} and all $n \in \mathbb{N}$. From Lemma 5.5 we infer that the same equality holds with R_n replaced by R . The operator

$$\mathfrak{D}(u) = \int_{B_0} \left(\int_{\mathcal{C}_t} u|_{\mathcal{C}_t} T_{f_{i,t}} \right) R$$

is defined on smooth compactly supported functions and we established that its value equals $\int_{\mathcal{C}} u \pi_i^* \widehat{T}_{f_i} \wedge \pi_{\mathcal{C}}^* R$. Following the proof of [BB, Proof of Proposition 4.3, Appendix] we see that \mathfrak{D} can be continuously extended to continuous compactly supported functions. From the continuity of the current $\pi_i^* \widehat{T}_{f_i} \wedge \pi_{\mathcal{C}}^* R$ it follows that

$$(5.16) \quad \int_{\mathcal{C}} u \pi_i^* \widehat{T}_{f_i} \wedge \pi_{\mathcal{C}}^* R = \int_{B_0} \left(\int_{\mathcal{C}_t} u|_{\mathcal{C}_t} T_{f_{i,t}} \right) R,$$

for all continuous and compactly supported functions u and $i = 1, 2$. Setting $u = \chi_{\mathcal{D}}$ we arrive at the equality (5.15) and thus we have (5.11) with $\beta = \deg(\pi_1)/\deg(\pi_2)$.

Similarly, using (5.16) with u being the continuous function $\psi_{\mathcal{D}}$ corresponding to \mathcal{D} from Proposition 3.6 we infer from (5.11) that

$$(5.17) \quad \int_{\mathcal{D}} \left(\int_{\mathcal{C}_t} \psi_{\mathcal{D}}|_{\mathcal{C}_t} \deg(\pi_2) d\pi_{1,t}^* \mu_{f_{1,t}}(z) \right) R = \int_{\mathcal{D}} \left(\int_{\mathcal{C}_t} \psi_{\mathcal{D}}|_{\mathcal{C}_t} \deg(\pi_1) d\pi_{2,t}^* \mu_{f_{2,t}}(z) \right) R,$$

Recalling that \mathcal{D} is chosen as in (5.13) and $\psi_{\mathcal{D}}$ satisfies the statement of Proposition 3.6, equation (5.17) implies that the set $\mathcal{P} := \{t \in \mathcal{D} : \mathcal{C}_t \text{ is weakly } \Phi_t\text{-special}\}$ has positive R measure. In particular since R doesn't charge points (e.g. by Lemma 5.5), the set \mathcal{P} is uncountable and must therefore contain a transcendental parameter $t_0 \in B_0(\mathbb{C}) \setminus B_0(\overline{\mathbb{Q}})$. But since B is 1-dimensional this contradicts our assumption that \mathbf{C} is not weakly Φ -special. The theorem follows. \square

We also record here the following result.

Theorem 5.6. *Let $\Phi = (\mathbf{f}_1, \mathbf{f}_2) : \mathbb{P}_1^2 \rightarrow \mathbb{P}_1^2$ be a split polarized endomorphism of polarization degree $d \geq 2$ and assume that \mathbf{f}_i is isotrivial for some $i \in \{1, 2\}$. Let $\mathbf{C} \subset \mathbb{P}_1^2$ be an irreducible curve that is not weakly Φ -special. Then there exists $\epsilon = \epsilon(\Phi, \mathbf{C}) > 0$ such that*

$$\{(P_1, P_2) \in \mathcal{C} \times_{B_0} \mathcal{C}(\overline{\mathbb{Q}}) : \hat{h}_{\Phi_t}(P_1) + \hat{h}_{\Phi_t}(P_2) < \epsilon\}$$

is not Zariski dense in $\mathcal{C} \times_{B_0} \mathcal{C}$.

Proof. If \mathbf{C} is not Φ -isotrivial then with Theorems 4.3, 4.9 and 4.4 and Propositions 4.13, 4.4 we can follow exactly the proof of Theorem 1.6. Note that by the equidistribution theorem in [YZ1], Corollary 4.4 is precisely the input needed to obtain an equidistribution statement as in Proposition 4.11 and thus to infer a statement as in Lemma 5.2. If \mathbf{C} is Φ -isotrivial then the statement follows from [GNY2, Theorem 2.2], [Zh2] and [Zh3]. \square

We can now deduce Theorem 1.2.

Proof of Theorem 1.2: Let Φ and \mathbf{C} be as in the statement of Theorem 1.2. Assume that Theorem 1.6 (or Theorem 5.6 in the non-isotrivial-free case) holds for $\Phi = \Psi : \mathbb{P}_1^2 \rightarrow \mathbb{P}_1^2$, for $\mathbf{X} = \mathbf{C}$ and with constant $\epsilon > 0$. So we know that the Zariski closure \mathcal{M} of the set $\{(P, Q) \in (\mathcal{C} \times_B \mathcal{C})(\overline{\mathbb{Q}}) : \hat{h}_{\Phi}(P) + \hat{h}_{\Phi}(Q) < \epsilon\}$ has dimension at most equal to 2. For each $t \in B_0(\overline{\mathbb{Q}})$ we let

$$\Sigma_t := \{z \in \mathcal{C}_t(\overline{\mathbb{Q}}) : \hat{h}_{\Phi_t}(z) < \epsilon/2\}.$$

Clearly we have that

$$\Sigma_t \times \Sigma_t \subset \mathcal{M}_t \subset \mathcal{C}_t \times \mathcal{C}_t.$$

Let $\mathcal{M} = \mathcal{M}_1 \cup \mathcal{M}_2$, where \mathcal{M}_1 denotes the finite union of the irreducible components of \mathcal{M} that are fibral or 1-dimensional and \mathcal{M}_2 those that are 2-dimensional and project dominantly to the base B . Since \mathbf{C} is not weakly Φ -special the quantity $\#(\Sigma_t \times \Sigma_t \cap \mathcal{M}_{1,t})$ is uniformly bounded for all but finitely many $t \in B_0(\overline{\mathbb{Q}}) \setminus E_{\Phi, \mathbf{C}}$. So it suffices to restrict our attention to \mathcal{M}_2 and we may assume further that it is irreducible, passing to an irreducible component. For each $t \in B_0(\overline{\mathbb{Q}}) \setminus E_{\Phi, \mathbf{C}}$ we have a projection map (in the first or second coordinate) between two curves $\pi_t : \mathcal{M}_{2,t} \rightarrow \mathcal{C}_t$ which is dominant and so also surjective. In particular for each $P \in \Sigma_t$ we have $\#\pi_t^{-1}(P) \leq D$ for a constant $D = D(\mathcal{M}_2)$. Therefore,

$$(5.18) \quad \#\Sigma_t \times \Sigma_t \leq D\#\Sigma_t.$$

Moreover, since \mathcal{M} has dimension at most 2 we infer that for all $t \in B_0(\overline{\mathbb{Q}})$ outside a finite set $E'_{\Phi, \mathbf{C}}$ we have $\#\Sigma_t < \infty$. Therefore, the theorem follows by (5.18). \square

Finally we deduce Corollary 1.3.

Proof of Corollary 1.3: Recall the definition of the essential minima from (2.7). Let $t \in B_0(\overline{\mathbb{Q}})$ and write \overline{L}_{Φ_t} for the invariant adelic metrized line bundle extending $L = \infty \times \mathbb{P}_1 + \mathbb{P}_1 \times \{\infty\}$. Clearly $e_0(\mathcal{C}_t, \overline{L}_{\Phi_t}) \geq 0$. Moreover, by Theorem 1.2 there is $\epsilon > 0$ such that $e_1(\mathcal{C}_t, \overline{L}_{\Phi_t}) \geq \epsilon$ for all but finitely many $t \in B_0(\overline{\mathbb{Q}})$. By Zhang's inequality in Lemma 2.1 we deduce that $\hat{h}_{\overline{L}_{\Phi_t}}(\mathcal{C}_t) = \hat{h}_{\Phi_t}(\mathcal{C}_t) \geq \epsilon$ for all but finitely many t . This proves (2).

For (1), note that if $t \in B_0(\overline{\mathbb{Q}})$ is such that \mathcal{C}_t is Φ_t -preperiodic, then by Zhang's inequality an [GNY1, Lemma 6.1] we have $\hat{h}_{\Phi_t}(\mathcal{C}_t) = 0$. On the other hand if there is $t \in B_0(\mathbb{C}) \setminus B_0(\overline{\mathbb{Q}})$ such that \mathcal{C}_t is Φ_t -preperiodic then since B is 1-dimensional, \mathbf{C} would be Φ -preperiodic, contradicting our assumption that it not weakly Φ -special. This completes the proof of the corollary. \square

6. COMMON PREPERIODIC POINTS

In this section we establish Theorem 1.1.

Proof of Theorem 1.1: Let $C \subset \text{Rat}_d \times \text{Rat}_d$ be a curve defined over $\overline{\mathbb{Q}}$ with projections p_1 and p_2 . We set $B = C$ and recall that $K = \overline{\mathbb{Q}}(B)$. Let $\Phi = (f_{p_1(\mathbf{c})}, f_{p_2(\mathbf{c})})$ where \mathbf{c} is the generic point of C .

We first prove that the statement holds if we restrict to transcendental points $\underline{\mu} \in C(\mathbb{C}) \setminus C(\overline{\mathbb{Q}})$. In fact, then $\underline{\mu}$ is the embedded image of the generic point of C into \mathbb{C} (so it induces an embedding $K \hookrightarrow \mathbb{C}$) and Theorem 1.1 reduces to the dynamical Manin-Mumford conjecture over \mathbb{C} [GNY1] if one of the maps $(f_{p_1(\underline{c})}, f_{p_2(\underline{c})})$ is ordinary or if both are exceptional to the classical Manin-Mumford conjecture [Hi]. Note that if Δ is Φ -special then $\text{Preper}(f_{p_1(\underline{\mu})}) = \text{Preper}(f_{p_2(\underline{\mu})})$ for all $\underline{\mu} \in C(\mathbb{C})$.

We may thus restrict our attention to algebraic points in $C(\overline{\mathbb{Q}})$ and we will consider two cases depending on whether Φ is isotrivial or not. Assume first that Φ is not isotrivial. Then by Theorem 1.2 we infer that there exists a bound M for all algebraic points $C(\overline{\mathbb{Q}})$ as in Theorem 1.1 outside a finite set of exceptions $E \subset C(\overline{\mathbb{Q}})$ unless Δ is weakly Φ -special. As Φ is not isotrivial Δ is weakly Φ -special if and only if it is Φ -preperiodic. If for some $\underline{\mu} \in E$, the diagonal Δ is not $(f_{p_1(\underline{\mu})}, f_{p_2(\underline{\mu})})$ -special we can apply the main theorem of [GNY1] or [Hi] and adjust the bound M accordingly. If on the other hand, Δ is $(f_{p_1(\underline{\mu})}, f_{p_2(\underline{\mu})})$ -special then $\text{Preper}(f_{p_1(\underline{\mu})}) = \text{Preper}(f_{p_2(\underline{\mu})})$. Thus Theorem 1.1 is proven if Φ is not isotrivial.

Now assume that Φ is isotrivial. Then there exists a pair of Möbius transformation $(\mathbf{M}_1, \mathbf{M}_2)$ such that $\mathbf{M}_i \circ f_{p_i(\underline{c})} \circ \mathbf{M}_i^{-1}$ is defined over $\overline{\mathbb{Q}}$ for $i = 1, 2$. As at least one p_1, p_2 is non-constant, at least one of $\mathbf{M}_1, \mathbf{M}_2$ is not defined over $\overline{\mathbb{Q}}$. Thus $(\mathbf{M}_1, \mathbf{M}_2)(\Delta)$ is not defined over $\overline{\mathbb{Q}}$ and Δ is not Φ -isotrivial. If Δ is not weakly Φ -special, then the result follows by Theorem 1.2. If on the other hand Δ is weakly Φ -special, then $(\mathbf{M}_1, \mathbf{M}_2)(\Delta)$ is the image of a coset in a fixed algebraic group as in Definition 1.5. If for $\underline{\mu} \in C(\overline{\mathbb{Q}})$ the corresponding coset is a torsion coset (in its fiber) then $\text{Preper}(f_{p_1(\underline{\mu})}) = \text{Preper}(f_{p_2(\underline{\mu})})$. If it is not a torsion coset then $\text{Preper}(f_{p_1(\underline{\mu})}) \cap \text{Preper}(f_{p_2(\underline{\mu})}) = \emptyset$. This concludes the proof. \square

APPENDIX A. A SPECIALIZATION RESULT

In this appendix we establish a dynamical analogue of [GH, Theorem 1.4] for 1-parameter families of polarized endomorphisms (not necessarily split). More precisely, in Theorem A.1 we provide a relation between the fiber-wise Call-Silverman canonical height and a height on the base, generalizing Call-Silverman's result [CS] to higher dimensions. Yuan and Zhang [YZ, Theorem 6.5] have recently obtained a more general result using their new theory of adelicly metrized line bundles on quasi-projective varieties. Here we rely on simpler machinery. Our approach is adapted from the strategy in [DGH1, DGH3], that in turn built on [Hal].

To fix notation, let B be a projective, regular, irreducible curve over $\overline{\mathbb{Q}}$ and write $K = \overline{\mathbb{Q}}(B)$. Let \mathbf{A} be a projective normal variety and L be a very ample line bundle on \mathbf{A} , both defined over K . Let $\Phi : \mathbf{A} \rightarrow \mathbf{A}$ be an endomorphism defined over K such that $\Phi^*(L) \simeq L^{\otimes d}$ for $d \in \mathbb{Z}_{\geq 2}$. We call the triple (\mathbf{A}, Φ, L) a *polarized endomorphism* over K with polarization degree d .

We note that Φ can be viewed as a 1-parameter family of endomorphisms each defined over $\overline{\mathbb{Q}}$. To this end, consider a B_0 -model $(\mathcal{A}, \Phi, \mathcal{L})$ of (\mathbf{A}, Φ, L) , where $B_0 \subset B$ is Zariski open. More specifically, \mathcal{A} is a variety defined over $\overline{\mathbb{Q}}$, $\pi : \mathcal{A} \rightarrow B_0$ is normal and flat and has generic fiber A , $\Phi : \mathcal{A} \dashrightarrow \mathcal{A}$ is the rational map associated to Φ , the generic fiber of \mathcal{L} is L and for each $t \in B_0$ the restriction of \mathcal{L} on the fiber of \mathcal{A} over t is a very ample line bundle such that $(\Phi|_{\pi^{-1}(t)})^*(\mathcal{L}|_{\pi^{-1}(t)}) \simeq (\mathcal{L}|_{\pi^{-1}(t)})^{\otimes d}$. For simplicity we write $\Phi_t := \Phi|_{\pi^{-1}(t)}$, $\mathcal{A}_t := \mathcal{A}|_{\pi^{-1}(t)}$ and $\mathcal{L}_t := \mathcal{L}|_{\pi^{-1}(t)}$. Thus for each $t \in B_0(\overline{\mathbb{Q}})$ the triple $(\mathcal{A}_t, \Phi_t, \mathcal{L}_t)$ is a polarized endomorphism of polarization degree d defined over a number field. In particular we have a fiberwise Call-Silverman canonical height defined for algebraic points $P \in \mathcal{A}^0 = \pi^{-1}(B_0)$ by

$$\hat{h}_{\Phi, \mathcal{L}}(P) = \hat{h}_{\Phi_{\pi(P)}, \mathcal{L}_{\pi(P)}}(P),$$

where $\hat{h}_{\Phi_t, \mathcal{L}_t}$ is the canonical height defined in [CS] associated to the endomorphism $\Phi_t : \mathcal{A}_t \rightarrow \mathcal{A}_t$ for $t \in B_0(\overline{\mathbb{Q}})$ with a polarization given by \mathcal{L}_t .

Recall that for an irreducible subvariety \mathbf{X} of \mathbf{A} defined over K and write $h_{\overline{L}_{\Phi}}(\mathbf{X}) \geq 0$ for the height of \mathbf{X} with respect to the canonical metric $\overline{L}_{\Phi} := (L, \{\|\cdot\|_{\Phi, L, v}\}_v)$ as in §2. We write $\mathcal{X} \subset \mathcal{A}$ for a B_0 -model of \mathbf{X} . We can now state our theorem.

Theorem A.1. *Let $(\mathcal{A}, \Phi, \mathcal{L})$ be a B_0 -model of a polarized endomorphism (\mathbf{A}, Φ, L) over K with polarization degree $d \geq 2$ and let $\mathbf{X} \subset \mathbf{A}$ be an irreducible subvariety. Let also \mathcal{N} be an ample line bundle on B . Assume that $h_{\overline{L}_{\Phi}}(\mathbf{X}) > 0$. Then there is a constant $\chi := \chi(\mathbf{X}, \Phi, L) > 0$ satisfying the following. For each $\epsilon > 0$, there is a Zariski open subset $\mathcal{U} := \mathcal{U}(\Phi, \mathbf{X}, \mathcal{N}, \epsilon) \subset \mathcal{X}$ and a constant $m := m(\mathbf{X}, \Phi, L, \mathcal{N}, \epsilon) > 0$ such that*

$$(A.1) \quad \hat{h}_{\Phi, \mathcal{L}}(P) \geq (\chi - \epsilon) h_{\mathcal{N}}(\pi(P)) - m,$$

for all $P \in \mathcal{U}(\overline{\mathbb{Q}})$.

Remark A.2. The constant χ in Theorem A.1 is given as

$$\chi = \frac{h_{\overline{L}_{\Phi}}(\mathbf{X})}{c},$$

where $c := c(\Phi, L, \deg_L(\mathbf{X})) > 0$ is described in (A.10).

To prove this theorem we will need to work on projective varieties, on which we can employ the height machine. To this end, following [Fab], we define inductively a sequence $(\mathcal{A}_N, \mathcal{L}_N)$ of B -models of $(\mathbf{A}, L^{\otimes d^{N-1}})$ such that \mathcal{L}_N is nef for each N as follows: For $N = 1$ note that \mathbf{A} is projective and L is very ample so we may embed \mathbf{A} into a projective space over K by L and subsequently identify \mathbb{P}_K^n with the generic fiber of $\mathbb{P}^n \times B$ to get

$$\mathbf{A} \hookrightarrow \mathbb{P}_K^n \hookrightarrow \mathbb{P}^n \times B.$$

We let \mathcal{A}_1 be the Zariski closure of \mathbf{A} in $\mathbb{P}^n \times B$ and write $\mathcal{L} = \mathcal{O}_{\mathcal{A}_1}(1, 0)$. Letting $\pi : \mathcal{A}_1 \rightarrow B$ denote the projection to the second factor, we may choose a very ample line bundle \mathcal{M} on B such that $\mathcal{L} \otimes \pi^* \mathcal{M}$ is nef; see the proof of [Fab, Lemma 3.1]. We let $\mathcal{L}_1 = \mathcal{L} \otimes \pi^* \mathcal{M}$ and have a B -model of (\mathbf{A}, L) given by $(\mathcal{A}_1, \mathcal{L}_1)$.

For the inductive step suppose that $(\mathcal{A}_N, \mathcal{L}_N)$ is a B -model of $(\mathbf{A}, L^{\otimes d^{N-1}})$ such that \mathcal{L}_N is nef and let $j_N : \mathbf{A} \hookrightarrow \mathcal{A}_N$ be the inclusion of the generic fiber. Let

$$\Gamma_{\Phi} = (i_d, \Phi) : \mathbf{A} \rightarrow \mathbf{A} \times_K \mathbf{A}$$

be the graph morphism on the generic fiber and write

$$\tilde{\Gamma}_{\Phi, N} = (j_N, j_N) \circ \Gamma_{\Phi} : \mathbf{A} \rightarrow \mathcal{A}_N \times_B \mathcal{A}_N.$$

Then define \mathcal{A}_{N+1} as the Zariski closure of $\tilde{\Gamma}_{\Phi, N}(\mathbf{A})$ in $\mathcal{A}_N \times_B \mathcal{A}_N$ and let $\mathcal{L}_{N+1} = (\text{pr}_2^* \mathcal{L}_N)|_{\mathcal{A}_{N+1}}$, where $\text{pr}_2 : \mathcal{A}_N \times_B \mathcal{A}_N \rightarrow \mathcal{A}_N$ is the projection to the second factor. Note that \mathcal{L}_{N+1} is nef as the pullback of a nef line bundle. If \mathbf{X} is a subvariety of \mathbf{A} over K , we denote by \mathcal{X}_N its Zariski closure in \mathcal{A}_N . We can now describe the $h_{\overline{\mathcal{L}}_{\Phi}}$ -height of our subvariety as a limit of intersection numbers in these models.

Lemma A.3. *Let (\mathbf{A}, Φ, L) be a polarized endomorphism over K of polarization degree $d \geq 2$. Let \mathbf{X} be an irreducible subvariety of \mathbf{A} of dimension $d_X \geq 0$. Then we have*

$$h_{\overline{\mathcal{L}}_{\Phi}}(\mathbf{X}) = \lim_{N \rightarrow \infty} \frac{(\mathcal{L}_N^{d_X+1} \cdot \mathcal{X}_N)}{(d_X + 1)d^{(N-1)(d_X+1)}}.$$

Proof. The lemma follows from the work in [Fab, Gu3]. For each $N \geq 0$ we have constructed B -models $(\mathcal{A}_N, \mathcal{L}_N)$ of $(\mathbf{A}, L^{\otimes d^{N-1}})$. These induce semipositive continuous adelic metrics $\{\|\cdot\|_{\mathcal{L}_N, v}\}_v$ on $L^{\otimes d^{N-1}}$ as in [Fab, page 351], which we denote by $\overline{\mathcal{L}}_N$. As shown in [Fab, §3.1] the sequence of metrized line bundles $\{\overline{\mathcal{L}}_N^{\otimes 1/d^{N-1}}\}_N$ converges uniformly to the Φ -invariant metric $\overline{\mathcal{L}}_{\Phi}$ supported on L and moreover outside of finitely many places v of K we have

$\|\cdot\|_{\mathcal{L}_N, v}^{1/d^{N-1}} = \|\cdot\|_{\mathcal{L}_\Phi, v}$ for all N . The lemma then follows in view of [Gu3, Theorem 3.5 (c)] or [Fab, Theorem 2.1 (ii)] (see also the discussion around equation (3) in [Fab]). \square

We proceed to clarify the construction of our models and describe the heights associated to them. Let $\iota : \mathbf{A} \hookrightarrow \mathbb{P}_K^n$ be the embedding over K induced by L . We first note that by [Fak, Corollary 2.2] we can find an endomorphism $\mathbf{F} : \mathbb{P}^n \rightarrow \mathbb{P}^n$ defined over K and such that $\mathbf{F} \circ \iota = \iota \circ \Phi$. Spreading it out and shrinking B_0 if necessary we have for each $t \in B_0$ an embedding $\iota_t : \mathcal{A}_t \hookrightarrow \mathbb{P}^n$ given by \mathcal{L}_t and an endomorphism $F_t : \mathbb{P}^n \rightarrow \mathbb{P}^n$ such that $F_t \circ \iota_t = \iota_t \circ \Phi_t$. Note here that $\mathcal{L} = \iota^*(\mathcal{L})$. We write $\mathcal{A}^0 = \pi^{-1}(B_0)$ to be the part of \mathcal{A} on which Φ is an endomorphism. For $k \in \mathbb{N}$ we let $(\mathbb{P}^n \times B_0)^{\times k}$ denote the k th fibered power over B_0 and write $F^{\times k} : (\mathbb{P}^n \times B_0)^{\times k} \rightarrow (\mathbb{P}^n \times B_0)^{\times k}$ to be the endomorphism acting as F_t on each fiber \mathbb{P}^n while fixing $t \in B_0$. By our definition $\mathcal{A}_N \subset (\mathbb{P}^n \times B)^{\times 2^{N-1}}$ is the Zariski closure of \mathcal{A}_N^0 , where \mathcal{A}_N^0 is defined inductively as follows:

$$(A.2) \quad \begin{aligned} \mathcal{A}_1^0 &= \{(\iota_t(z), t) : (z, t) \in \mathcal{A}^0\}, \\ \mathcal{A}_{N+1}^0 &= \{(Q, F^{\times 2^{N-1}}(Q)) : Q \in \mathcal{A}_N^0\}. \end{aligned}$$

Recalling that $\mathcal{L}_{N+1} = (\text{pr}_2^* \mathcal{L}_N)|_{\mathcal{A}_{N+1}}$, the functoriality of heights allows us to choose the heights associated to \mathcal{L}_{N+1} inductively so that

$$h_{\mathcal{L}_{N+1}}(Q, F^{\times 2^{N-1}}(Q)) = h_{\mathcal{L}_N}(F^{\times 2^{N-1}}(Q)) + O_N(1),$$

for $Q \in \mathcal{A}_N^0(\overline{\mathbb{Q}})$ with a bounded error term depending on N but independent of Q . More specifically if for $Z := (z, t) \in \mathcal{A}_t(\overline{\mathbb{Q}}) \times B_0(\overline{\mathbb{Q}})$ we define $Z_{N+1} \in \mathcal{A}_{N+1}^0(\overline{\mathbb{Q}})$ as the (unique) point in $\mathcal{A}_{N+1}^0(\overline{\mathbb{Q}})$ determined by Z (its ‘first’ projective coordinate is $\iota_t(z)$ and ‘last’ projective coordinate is given by $F_t^N(\iota_t(z))$), then we have

$$(A.3) \quad \begin{aligned} h_{\mathcal{L}_{N+1}}(Z_{N+1}) &= h_{\iota^* \mathcal{L}_1}(\Phi^N(Z)) + O_N(1) = h_{\iota_t^* \mathcal{L}_t}(\Phi_t^N(z)) + h_{\mathcal{M}}(t) + O_N(1) \\ &= h_{\mathcal{L}_t}(\Phi_t^N(z)) + h_{\mathcal{M}}(t) + O_N(1). \end{aligned}$$

It also follows from our construction that $\mathcal{X}_N \subset \mathcal{A}_N$ defined earlier is the Zariski closure of \mathcal{X}_N^0 , where for each N the set \mathcal{X}_{N+1}^0 is the ‘graph’ of \mathcal{X}_N^0 under $F^{\times 2^{N-1}}$.

Finally, we write $f_N : \mathcal{A}_N \rightarrow B$ and define $\mathcal{M}_N = (f_N)^*(\mathcal{M})$. We will need the following bound on intersection numbers.

Lemma A.4. *We have $(\mathcal{M}_N \cdot \mathcal{L}_N^{\dim \mathbf{X}} \cdot \mathcal{X}_N) \leq cd^{(N-1)\dim \mathbf{X}}$ for a constant $c := c(\Phi, L, \mathbf{X}) > 0$.*

Proof. Write $d_X := \dim \mathbf{X}$. We may use \mathcal{M} to embed $B \hookrightarrow \mathbb{P}^m$ for some m . Abusing notation slightly we continue to denote by F the map induced by F defined earlier which is now an endomorphism of $\mathbb{P}^n \times (\mathbb{P}^m \setminus Z)$ for a Zariski closed subset $Z \subset \mathbb{P}^m$ corresponding to the fibers above $B \setminus B_0$. We also identify $\mathcal{X}_N^0, \mathcal{A}_N^0$ and their Zariski closures $\mathcal{X}_N, \mathcal{A}_N$ with their images in $(\mathbb{P}^n \times \mathbb{P}^m)^{2^{N-1}}$. Letting $x = [x_0, \dots, x_n]$ and $s = [s_0, \dots, s_m]$ denote the projective coordinates on \mathbb{P}^n and \mathbb{P}^m respectively, we write

$$F(x, s) = ([f_0(x, s) : \dots : f_n(x, s)], s)$$

for bihomogeneous $f_i(x, s)$ of bidegree (d, D_1) in (x, s) for each $i = 0, \dots, n$. Since $\mathcal{X}_1 \subset \mathbb{P}^n \times \mathbb{P}^m$ dominates the base but Z does not, we may henceforth assume that f_0 is not identically zero on $(x, s) \in \mathcal{X}_1$. The ℓ -th iterate of F is denoted by

$$F^\ell(x, s) = ([f_0^\ell(x, s) : \dots : f_n^\ell(x, s)], s).$$

It is easy to see that the $f_i^\ell(x, s)$ are bihomogeneous of bidegree (d^ℓ, D_ℓ) in (x, s) for each $i = 0, \dots, n$ and we have $D_{\ell+1} = D_1 + dD_\ell$, so that $D_\ell = \frac{d^\ell - 1}{d - 1} D_1 \leq d^\ell D_1$.

If we denote by

$$\text{pr}_N : (\mathbb{P}^n \times \mathbb{P}^m)^{2^{N-1}} \rightarrow \mathbb{P}^n \times \mathbb{P}^n \times \mathbb{P}^m$$

the projection to the first and last two factors in the product of the 2^N projective spaces above, then we have

$$\begin{aligned} \text{(A.4)} \quad \text{pr}_N^* \mathcal{O}(0, 1, 1) &= \mathcal{L}_N \\ \text{pr}_N^* \mathcal{O}(0, 0, 1) &= \mathcal{M}_N. \end{aligned}$$

Since pr_N has degree equal to 1, by the projection formula we get

$$\text{(A.5)} \quad (\mathcal{L}_N^{d_X} \cdot \mathcal{M}_N \cdot \mathcal{X}_N) = (\mathcal{O}(0, 1, 1)^{d_X} \cdot \mathcal{O}(0, 0, 1) \cdot \text{pr}_{N*} \mathcal{X}_N).$$

We may now follow closely the arguments in [DGH3, Proposition 4.3]. Writing $[\Gamma]$ for the cycle in $\mathbb{P}^n \times \mathbb{P}^n \times \mathbb{P}^m$ representing the subvariety $\{(x, y, s) : y \in \mathbb{P}^n, (x, s) \in \mathcal{X}_1\}$, we see that $\text{pr}_{N*}(\mathcal{X}_N) \subset \mathbb{P}^n \times \mathbb{P}^n \times \mathbb{P}^m$ is an irreducible component of the intersection of $V_i := \{y_i f_0^{N-1}(x, s) - f_i^{N-1}(x, s) y_0 = 0\}$ and Γ over all $i = 1, \dots, n$. In fact, as in [DGH3, Proposition 4.3 before equation (4.12)], using Fulton's positivity result [Ful98, Corollary 12.2.(a)], we get

$$\text{(A.6)} \quad (\mathcal{L}_N^{d_X} \cdot \mathcal{M}_N \cdot \mathcal{X}_N) \leq (\mathcal{O}(0, 1, 1)^{d_X} \cdot \mathcal{O}(0, 0, 1) \cdot \mathcal{O}(d^{N-1}, 1, D_{N-1}) \cdot [\Gamma]).$$

Continuing as in [DGH3] we observe that the positive cycle $[\Gamma] \subset \mathbb{P}^n \times \mathbb{P}^n \times \mathbb{P}^m$ is linearly equivalent with

$$\sum_{i+p=n+m-d_X-1} a_{ip} H_1^i H_2^p,$$

where $a_{ip} \in \mathbb{Z}_{\geq 0}$ and H_i are the pullbacks of the factors $\mathcal{X}_1 \subset \mathbb{P}^n \times \mathbb{P}^m$. We thus infer that the right hand side of (A.6) is smaller than

$$(A.7) \quad \sum_{i+p=n+m-d_X-1} a_{ip} (\mathcal{O}(0, 1, 1)^{d_X} \cdot \mathcal{O}(0, 0, 1) \cdot \mathcal{O}(d^{N-1}, 1, D_{N-1})^n \cdot \mathcal{O}(1, 0, 0)^i \cdot \mathcal{O}(0, 0, 1)^j).$$

Using the linearity of intersection numbers and their vanishing properties exactly as in [DGH3] it is then easy to infer that

$$(A.8) \quad (\mathcal{L}_N^{d_X} \cdot \mathcal{M}_N \cdot \mathcal{X}_N) \leq \sum_{\substack{i+p=n+m-d_X-1, i+i''=n \\ j'+p'=d_X, j'+j''=n \\ i''+j''+p''=n, p+p'+p''=m-1}} a_{ip} \binom{d_X}{j', p'} \binom{n}{i'', j'', p''} d^{(N-1)i''} D_{N-1}^{p''}.$$

Noting that $p'' + i'' = n - j'' = j' = d_X - p' \leq d_X$ and that $D_\ell \leq d^\ell D_1$, we thus infer that

$$(A.9) \quad (\mathcal{L}_N^{d_X} \cdot \mathcal{M}_N \cdot \mathcal{X}_N) \leq d^{(N-1)d_X} D_1^{d_X} 2^{d_X} 3^n \sum_{i+p=n+m-d_X-1} a_{ip}.$$

The proposition then follows with

$$(A.10) \quad c = D_1^{d_X} 2^{d_X} 3^n \sum_{i+p=n+m-d_X-1} a_{ip}.$$

□

We are now ready to prove Theorem A.1.

Proof of Theorem A.1: For simplicity we write $d_X = \dim \mathbf{X}$. Let $\epsilon \in (0, h_{\overline{\mathcal{L}}_\Phi}(\mathbf{X})/3)$. By Lemma A.3, we can find $N_\epsilon \in \mathbb{N}$ so that

$$(h_{\overline{\mathcal{L}}_\Phi}(\mathbf{X}) + \epsilon)(d_X + 1)d^{(N-1)(d_X+1)} \geq (\mathcal{L}_N^{d_X+1} \cdot \mathcal{X}_N) \geq (h_{\overline{\mathcal{L}}_\Phi}(\mathbf{X}) - \epsilon)(d_X + 1)d^{(N-1)(d_X+1)},$$

for all $N \geq N_\epsilon$. Let $N \geq N_\epsilon$ be arbitrary. For simplicity we write $\tilde{\mathcal{L}}_N := \mathcal{L}_N|_{\mathcal{X}_N}$ and $\tilde{\mathcal{M}}_N := \mathcal{M}_N|_{\mathcal{X}_N}$. Note that \mathcal{X}_{N+1} is a projective variety of dimension $d_X + 1$ and its line bundles $\tilde{\mathcal{M}}_{N+1}$ and $\tilde{\mathcal{L}}_{N+1}$ are nef. Let $c > 0$ be as in Lemma A.4 and $y_\epsilon \in \mathbb{Q}_{>0}$ be such that

$$(A.11) \quad 0 < \frac{(h_{\overline{\mathcal{L}}_\Phi}(\mathbf{X}) - 2\epsilon)}{c} < y_\epsilon \leq \frac{(h_{\overline{\mathcal{L}}_\Phi}(\mathbf{X}) - \epsilon)}{c}.$$

Because of the bounds on the intersection numbers in Lemmata A.3 and A.4, Siu's theorem [La3, Theorem 2.2.15] yields that the line bundle $\tilde{\mathcal{L}}_N - y_\epsilon d^{N-1} \tilde{\mathcal{M}}_N$ is big. In particular, there is $c_1(N, \epsilon) > 0$ so that

$$h_{\tilde{\mathcal{L}}_N}(Q) > y_\epsilon d^{N-1} h_{\tilde{\mathcal{M}}_N}(Q) - c_1(N, \epsilon),$$

for all algebraic points Q in a Zariski open subset of \mathcal{X}_N . Our considerations for the heights in (A.3) then yield that we can find a constant $c_2(N, \epsilon)$ with

$$(A.12) \quad h_{\mathfrak{L}_t}(\Phi_t^{N-1}(z)) + h_{\mathcal{M}}(t) > y_\epsilon d^{N-1} h_{\mathcal{M}}(t) - c_2(N, \epsilon),$$

for all (z, t) in a Zariski open subset \mathcal{U}_N of \mathcal{X} . By [CS, Theorem 3.1] we also have

$$(A.13) \quad \hat{h}_{\Phi_t, \mathfrak{L}_t}(\Phi_t^{N-1}(z)) - h_{\mathfrak{L}_t}(\Phi_t^{N-1}(z)) \geq -\kappa(h_{\mathcal{M}}(t) + 1),$$

for all (z, t) in a possibly smaller open $\mathcal{U}_N \subset \mathcal{X}$ and for positive constant κ which can be chosen independently of N .

We now proceed to fix $N = N'_\epsilon \geq N_\epsilon$ such that $\frac{\kappa+1}{d^{N-1}} < \frac{\epsilon}{c}$. Combining the inequalities (A.12) and (A.13) we get that for all (z, t) in a (fixed) Zariski open subset of \mathcal{X} the following holds

$$(A.14) \quad \begin{aligned} d^{N-1} \hat{h}_{\Phi_t, \mathfrak{L}_t}(z) &= \hat{h}_{\Phi_t, \mathfrak{L}_t}(\Phi_t^{N-1}(z)) \geq h_{\mathfrak{L}_t}(\Phi_t^{N-1}(z)) - \kappa(h_{\mathcal{M}}(t) + 1) \\ &\geq (y_\epsilon d^{N-1} - 1 - \kappa) h_{\mathcal{M}}(t) - c_2 - \kappa. \end{aligned}$$

Dividing by d^N and noting our choice for N and y_ϵ and that $h_{\mathcal{M}}$ is non-negative, we get

$$\begin{aligned} \hat{h}_{\Phi_t, \mathfrak{L}_t}(z) &\geq \left(y_\epsilon - \frac{\kappa+1}{d^{N-1}} \right) h_{\mathcal{M}}(t) - \frac{c_2 + \kappa}{d^{N-1}} \\ &\geq \left(\frac{h_{\tilde{\mathcal{L}}_\Phi}(\mathbf{X}) - 3\epsilon}{c} \right) h_{\mathcal{M}}(t) - \frac{c_2 + \kappa}{d^{N-1}}. \end{aligned}$$

The theorem follows. □

REFERENCES

- [Aea] M. Abate, E. Bedford, M. Brunella, T.-C. Dinh, D.-Schleicher and N. Sibony, Holomorphic dynamical systems, *Lecture Notes in Mathematics*, **1998**, Lectures given at the C.I.M.E. Summer School held in Cetraro, July 7–12, 2008, Edited by Graziano Gentili, Jacques Guenot and Giorgio Patrizio, Springer-Verlag, Berlin; Fondazione C.I.M.E., Florence, 2010, xiv+342.
- [Ax] J. Ax, Some topics in differential algebraic geometry. I. Analytic subgroups of algebraic groups, *Amer. J. Math.*, **94**, (1972), 1195–1204.

- [Ba] M. Baker. A finiteness theorem for canonical heights attached to rational maps over function fields, *J. Reine Angew. Math.*, **626**, (2009), 205–233.
- [BD1] M. Baker and L. DeMarco. Special curves and postcritically-finite polynomials. *Forum of Math. Pi* **1** (2013), 35 pp.
- [BD2] M. Baker and L. DeMarco. Preperiodic points and unlikely intersections. *Duke Math. J.* **159** (2011), 1–29.
- [BH] M. Baker and L.-C. Hsia. Canonical heights, transfinite diameters, and polynomial dynamics. *J. Reine Angew. Math.* **585** (2005), 61–92.
- [BB] G. Bassanelli and F. Berteloot. Bifurcation currents in holomorphic dynamics on \mathbb{P}^k , *J. Reine Angew. Math.*, **608**(2007), 201–235.
- [Bea] A. F. Beardon. Symmetries of Julia sets, *Bull. London Math. Soc.*, **22** (1990), no. 6, 576–582.
- [BT] E. Bedford and B.A. Taylor. A new capacity for plurisubharmonic functions. *Acta Math.* **149**(1982), no.1-2, 1–40.
- [Ben] R. L. Benedetto. Preperiodic points of polynomials over global fields, *J. Reine Angew. Math.*, **608** (2007), 123–153.
- [Ber] D. Bertrand. Extensions de D-modules et groupes de Galois différentiels, *p-adic analysis (Trento, 1989)*, volume 1454 of *Lecture Notes in Math.*, pages 125–141. Springer, Berlin, 1990.
- [BH] B. Branner and J. H. Hubbard. The iteration of cubic polynomials. I. The global topology of parameter space. *Acta Math.* **160**(1988), 143–206.
- [CS] G. S. Call and J. H. Silverman. Canonical heights on varieties with morphisms. *Compositio Math.* **89**(1993), 163–205.
- [Ca] A. Carney. The arithmetic Hodge index theorem and rigidity of dynamical systems over function fields, *Pacific J. Math.*, **309**, 2020, no.1, 71–102.
- [CL1] A. Chambert-Loir. Mesures et équidistribution sur les espaces de Berkovich. *J. Reine Angew. Math.* **595**(2006), 215–235.
- [CL2] A. Chambert-Loir. Heights and measures on analytic spaces. A survey of recent results, and some remarks. In *Motivic integration and its interactions with model theory and non-Archimedean geometry. Volume II*, volume 384 of *London Math. Soc. Lecture Note Ser.*, pages 1–50. Cambridge Univ. Press, Cambridge, 2011.
- [CLT] A. Chambert-Loir, A. Thuillier. Mesures de Mahler et équidistribution logarithmique, *Ann. Inst. Fourier (Grenoble)*, **59**(2009), no.3, 977–1014.
- [CDMZ] P. Corvaja, J. Demeio, D. Masser, U. Zannier. On the torsion values for sections of an elliptic scheme *J. Reine Angew. Math.*, **782**(2022), 1–41.
- [De1] L. DeMarco. Bifurcations, intersections, and heights *Algebra Number Theory* **10**(2016), 1031–1056.
- [DKY1] L. DeMarco, H. Krieger and H. Ye. Uniform Manin-Mumford for a family of genus 2 curves, *Ann. of Math. (2)* **191**, 2020, no.3, 949–1001.
- [DKY2] L. DeMarco, H. Krieger and H. Ye. Common preperiodic points for quadratic polynomials. *arXiv:1911.02458*.

- [DGH1] V. Dimitrov, Z. Gao, and P. Habegger. Uniform bound for the number of rational points on a pencil of curves. *Int. Math. Res. Not. IMRN*, 2021, no.2, 1138–1159.
- [DGH2] V. Dimitrov, Z. Gao, and P. Habegger. A consequence of the relative Bogomolov conjecture. *J. Number Theory*, **230**(2022), 146–160.
- [DGH3] V. Dimitrov, Z. Gao, and P. Habegger. Uniformity in Mordell–Lang for curves. *Ann. of Math.*, **194**, 2021, no.1, 237–298.
- [DM1] L. DeMarco and N.M. Mavraki. Variation of canonical height and equidistribution. *Amer. J. Math.*, **142** (2020), no.2, 443–473.
- [DM2] L. DeMarco and N.M. Mavraki. Elliptic surfaces and arithmetic equidistribution for \mathbb{R} -divisors on curves. arXiv:2012.14529.
- [DWY1] L. DeMarco, X. Wang and H. Ye. Torsion points and the Lattès family. *Amer. J. Math.* **138** (2016), no. 3, 697–732.
- [DWY2] L. DeMarco, X. Wang and H. Ye. Bifurcation measures and quadratic rational maps. *Proc. Lond. Math. Soc.*, (3) **111** (2015), no. 1, 149–180.
- [DS] T.-C. Dinh and N. Sibony. Introduction to the theory of currents notes dated September 21, 2005
- [DH] A. Douady and J. Hubbard. *A proof of Thurston’s topological characterization of rational functions*. In *Acta Math.* 171, 263–297 (1993)
- [Fab] X. W. C Faber. Equidistribution of dynamically small subvarieties over the function field of a curve, *Acta Arith.*, **137**, (2009), no.4, 345–389.
- [Fak] N. Fakhruddin. Questions on self maps of algebraic varieties, *J. Ramanujan Math. Soc.*, **18** (2003), no.2, 109–122.
- [FG1] C. Favre and T. Gauthier. Distribution of postcritically finite polynomials. *Israel J. Math.*, **209** (2015), 235–292.
- [FG2] C. Favre and T. Gauthier. Classification of special curves in the space of cubic polynomials. *Int. Math. Res. Not. IMRN*, (2018), no.2, 362–411.
- [FG3] Ch. Favre and Th. Gauthier. The arithmetic of polynomial dynamical pairs, *volume 214 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 2022*
- [FLM] A. Freire, A. Lopes and R. Mañé. *An invariant measure for rational maps*, *Bol. Soc. Brasil. Mat.* **14**, 1983, no.1, 45–62.
- [FS] J. E. Fornæss and N. Sibony. Complex dynamics in higher dimensions. In *Complex Potential Theory (Montreal, PQ, 1993)*, pages 131–186. Kluwer Acad. Publ., Dordrecht, 1994.
- [Ful98] W. Fulton. *Intersection theory*, volume 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, second edition, 1998.
- [Ga1] Z. Gao. Generic rank of Betti map and unlikely intersections, *Compos. Math.*, **156**(2020), no.12, 2469–2509.
- [Ga2] Z. Gao. Mixed Ax-Schanuel for the universal abelian varieties and some applications, *Compos. Math.*, **156**(2020), no.11, 2263–2297.

- [GH] Z. Gao and Ph. Habegger. Heights in families of abelian varieties and the geometric Bogomolov conjecture, *Ann. of Math. (2)*, **189**(2019), no.2, 527–604.
- [Gau] T. Gauthier. Good height functions on quasi-projective varieties: equidistribution and applications in dynamics *arXiv:2105.02479*, May 2021
- [GV] T. Gauthier, G. Vigny. The geometric dynamical Northcott and Bogomolov properties *arXiv:1912.07907*
- [GHT1] D. Ghioca, L.-C. Hsia and T. Tucker. Preperiodic points for families of polynomials. *Algebra and Number Theory*. **7** (2013), no. 3, 701–732.
- [GHT2] D. Ghioca, L.-C. Hsia and T. Tucker. Preperiodic points for families of rational maps. *Proc. London Math. Soc.* **110** (2015), 395–427.
- [GKN] D. Ghioca, H. Krieger and K. Nguyen. A case of the Dynamical André-Oort Conjecture. *Int. Math. Res. Not. IMRN* vol. **2016** (2016), 738–758.
- [GKY] D. Ghioca, H. Krieger, K. Nguyen and H. Ye. The Dynamical André-Oort Conjecture: unicritical polynomials. *Duke Math. Journal* **166** (2017), no.1, 1–25.
- [GNY1] D. Ghioca, K. Nguyen and H. Ye. The dynamical Manin-Mumford conjecture and the dynamical Bogomolov conjecture for split rational maps, *J. Eur. Math. Soc. (JEMS)*, **21**(2019), no.5, 1571–1594.
- [GNY2] D. Ghioca, K. Nguyen and H. Ye. The dynamical Manin-Mumford conjecture and the dynamical Bogomolov conjecture for endomorphisms of $(\mathbb{P}^1)^n$, *Compos. Math.*, **154**, no.7, 1441–1472.
- [GT] D. Ghioca and T. J. Tucker. *Proof of a Dynamical Bogomolov Conjecture for lines under polynomial actions*. *Proc. Amer. Math. Soc.* **138** (2010), 937–942.
- [GT] D. Ghioca and T. J. Tucker, A reformulation of the dynamical Manin-Mumford conjecture, *Bull. Aust. Math. Soc.*, **103** (2021), no.1, 154–161.
- [GTZ] D. Ghioca, T. Tucker and Sh. W. Zhang, Towards a Dynamical Manin-Mumford Conjecture *Int. Math. Res. Not.* (2011), 5109–5122.
- [GY] D. Ghioca and H. Ye. A dynamical variant of the André-Oort Conjecture. To appear *Int. Math. Res. Not.*.
- [GH] Ph. Griffiths and J. Harris. Principles of algebraic geometry, *Wiley Classics Library*, Reprint of the 1978 original, *John Wiley & Sons, Inc., New York*, (1994), xiv+813
- [Gu1] W. Gubler. Local heights of subvarieties over non-Archimedean fields, *J. Reine Angew. Math.*, **498**, (1998), 61–113.
- [Gu2] W. Gubler. The Bogomolov conjecture for totally degenerate abelian varieties. *Invent. Math.*, **169**, (2007), no. 2, 377–400.
- [Gu3] W. Gubler. Equidistribution over function fields. *Manuscripta Math.*, **127**, (2008), no. 4, 485–510.
- [Ha1] Ph. Habegger. Special points on fibered powers of elliptic surfaces, *J. Reine Angew. Math.*, **685**(2013), 143–179.
- [Hi] M. Hindry. Autour d’une conjecture de Serge Lang *Invent. Math.* **94** (1988), 3, 575–603.
- [HP] J. Hubbard and P. Papadopol. Superattractive fixed points in \mathbf{C}^n . *Indiana Univ. Math. J.* **43**(1994), 321–365.

- [Ku1] L. Kühne, The Relative Bogomolov Conjecture for Fibered Products of Elliptic Surfaces, arXiv:2103.06203.
- [Ku2] L. Kühne, Equidistribution in Families of Abelian Varieties and Uniformity, arXiv:2101.10272.
- [La3] R. Lazarsfeld. *Positivity in algebraic geometry. I*, volume 48 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics*. Springer-Verlag, Berlin, 2004.
- [Le] G. Levin. Symmetries on Julia sets. *Mat. Zametki* **48**, 72–79, 159 (1990), (in Russian).
- [LP] G. Levin and F. Przytycki. When do two rational functions have the same Julia set?, *Proc. Amer. Math. Soc.*, **125** (1997), no.7, 2179–2190.
- [Ly] M. Lyubich. Entropy properties of rational endomorphisms of the Riemann sphere. *Ergodic Theory Dynamical Systems* **3**(1983), 351–385.
- [MS] A. Medvedev and Th. Scanlon. Invariant varieties for polynomial dynamical systems, *Ann. of Math. (2)*, **179**(2014), no.1, 81–177.
- [MZ1] D. Masser and U. Zannier. Torsion anomalous points and families of elliptic curves. *Amer. J. Math.* **132**(2010), 1677–1691.
- [MZ2] D. Masser and U. Zannier. Torsion points on families of squares of elliptic curves. *Math. Ann.* **352**(2012), 453–484.
- [MZ3] D. Masser and U. Zannier. Torsion points on families of products of elliptic curves. *Adv. Math.* **259**(2014), 116–133.
- [MZ4] D. Masser and U. Zannier. Torsion points on families of simple abelian surfaces and Pell’s equation over polynomial rings. *J. Eur. Math. Soc. (JEMS)* **17**(2015), 2379–2416. With an appendix by E. V. Flynn.
- [Mi] J. Milnor. Dynamics in one complex variable. *arXiv*, 1990
- [Mi] A. Mimar. On the preperiodic points of an endomorphism of $\mathbb{P}^1 \times \mathbb{P}^1$ which lie on a curve, *Thesis (Ph.D.)—Columbia University, ProQuest LLC, Ann Arbor, MI* (1997)
- [Ra] M. Raynaud. Courbes sur une variété abélienne et points de torsion. *Invent. Math.* **71**(1983), 207–233.
- [Ru] D. Ruelle. An inequality for the entropy of differentiable maps, *Bol. Soc. Brasil. Mat.*, **9**(1978), no.1, 83–87.
- [Ul] E. Ullmo. Positivité et discrétion des points algébriques des courbes. *Ann. of Math. (2)* **147**(1998), 167–179.
- [Ye] H. Ye. Rational functions with identical measure of maximal entropy. *Adv. Math.*, 268:373–395, 2015.
- [Yu1] X. Yuan. Arithmetic bigness and a uniform Bogomolov-type result, arXiv:2108.05625.
- [Yu2] X. Yuan. Algebraic dynamics, canonical heights and Arakelov geometry. Algebraic dynamics, canonical heights and Arakelov geometry, *Fifth International Congress of Chinese Mathematicians. Part 1, 2, AMS/IP Stud. Adv. Math.*, 51, pt. 1 (2012)**2**, 893–929
- [YZ] X. Yuan and Sh.-W. Zhang. The arithmetic Hodge index theorem for adelic line bundles, *Math. Ann.*, **367**, (2017), no.3-4, 1123–1171.

- [YZ1] X. Yuan and S. W. Zhang. The arithmetic Hodge index theorem for adelic line bundles, *Math. Ann.* **367**(2017), no.3-4, 1123–1171.
- [YZ2] X. Yuan and S. W. Zhang. Adelic line bundles over quasi-projective varieties. *preprint*
- [Za] U. Zannier. Some problems of unlikely intersections in arithmetic and geometry, With appendixes by David Masser. *Annals of Mathematics Studies*. Volume 181, Princeton University Press, Princeton, NJ, 2012.
- [Zh1] S.-W. Zhang. Small points and adelic metrics, *J. Algebraic Geom.*, **4**, (1995), no.2, 281–300.
- [Zh2] S.-W. Zhang. Positive line bundles on arithmetic surfaces, *Ann. of Math. (2)*, **136**, (1992), no. 3, 569–587.
- [Zh3] S.-W. Zhang. Equidistribution of small points on abelian varieties, *Ann. of Math.* **147** (1998), no. 1, 159–165.
- [Zh4] S.-W. Zhang. Distributions in algebraic dynamics. *Surveys in differential geometry.*, **10** (2006), 381–430.

Email address: mavraki@math.harvard.edu

Email address: harry.schmidt@unibas.ch