

# RÉNYI ENTROPIES OF THE FREE FERMI GAS IN MULTI-DIMENSIONAL SPACE AT HIGH TEMPERATURE

HAJO LESCHKE, ALEXANDER V. SOBOLEV, WOLFGANG SPITZER

*To the memory of Harold Widom (1932–2021)*

ABSTRACT. We study the local and (bipartite) entanglement Rényi entropies of the free Fermi gas in multi-dimensional Euclidean space in thermal equilibrium. We prove the positivity of the entanglement entropies at any temperature for Rényi index  $\gamma \leq 1$ . Furthermore, for general  $\gamma > 0$  we establish the asymptotics of the entropies at high temperature and at large scaling parameter for two different regimes – for fixed chemical potential and also for fixed particle density. In particular, we thereby provide the last remaining building block for a complete proof of our low- and high-temperature results presented (for  $\gamma = 1$ ) in *J. Phys. A: Math. Theor.* **49**, 30LT04 (2016) (Corrigendum: **50**, 129501 (2017)) but being supported there only by the basic proof ideas.

## CONTENTS

1. Introduction	2
1.1. Physics background	2
1.2. Pseudo-differential operators and entropies	3
1.3. Description of results	4
2. Basic definitions and basic results	6
2.1. Conditions on the truncating region $\Lambda$	6
2.2. The asymptotic coefficient $\mathcal{B}_d$ and its properties	6
2.3. Asymptotics of $\text{tr } D_\alpha(a, \Lambda; f)$	7
2.4. Multi-scale symbols	9
3. On positivity of Rényi entropies	9
3.1. Concavity of the entropy function $\eta_\gamma$	9
3.2. Operator concavity of $\eta_\gamma$	11
4. Quasi-commutator bounds	12
5. Preparations for high temperature asymptotics: the model symbol	14
5.1. Conditions on the single-particle Hamiltonian $h$	15
5.2. The model symbol	15
6. Main results on the high-temperature asymptotics	19
6.1. Case of a fixed chemical potential $\mu$	19
6.2. Case of a fixed particle density $\rho$	20

---

*Date:* 5 January 2022.

*2020 Mathematics Subject Classification.* Primary 47G30, 35S05; Secondary 47B10, 47B35.

*Key words and phrases.* Non-smooth functions of Wiener–Hopf operators, asymptotic trace formulas, Rényi (entanglement) entropy of fermionic equilibrium states.

6.3. Concluding remarks	21
Appendix A. The Fermi symbol at high temperature $T$ and fixed particle density $\rho$	22
References	23

## 1. INTRODUCTION

**1.1. Physics background.** During the last decades the so-called entanglement entropy (EE) has become a useful single number quantifier of non-classical correlations between subsystems of a composite quantum-mechanical system [1, 13]. For example, one may imagine a macroscopically large system consisting of a huge number of particles in the state of thermal equilibrium at some temperature  $T \geq 0$ . All the particles inside some bounded spatial region  $\Lambda$  may then be considered to constitute one subsystem and the particles outside of  $\Lambda$  another one. The corresponding EE, more precisely the spatially bipartite thermal EE, now quantifies, to some extent, how strongly these two subsystems are correlated “across the interface” between  $\Lambda$  and its complement.

In the simplified situation where the particles do not dynamically interact with each other, such as in the ideal gas or, slightly more general, in the free gas, all possible correlations are entirely due to either the Bose–Einstein or the (Pauli–)Fermi–Dirac statistics due to the assumed indistinguishability of the particles. The present study is devoted to the latter case. Accordingly, we consider the free Fermi gas [3, 10] infinitely extended in the Euclidean space  $\mathbb{R}^d$  of an arbitrary dimension  $d \geq 1$ . Although the fermions neither interact with each other nor with any external field, their EE remains a complicated function of the region  $\Lambda \subset \mathbb{R}^d$  which is difficult to study by analytical methods. In general one can only hope for estimates and asymptotic results for its (physically interesting) growth when  $\Lambda$  is replaced with  $\alpha\Lambda$  with a parameter  $\alpha > 0$  and  $\alpha$  becomes large. A decisive progress towards the understanding of the growth of the EE at  $T = 0$ , in other words of the ground-state EE, is due to Gioev and Klich [11, 12]. They observed, remarkably enough, that this growth is related to a conjecture of Harold Widom [27, 30] about the quasi-classical Szegő-type asymptotics for traces of (smooth) functions of multi-dimensional versions of truncated Wiener–Hopf operators with discontinuous symbols. After Widom’s conjecture had been proved by one of us [20, 22] the gate stood open to confirm [15] the precise (leading) large-scale growth conjectured in [12] and, in addition, to establish its extension from the von Neumann EE to the one-parameter family of (quantum) Rényi EE’s.

In the present study we only consider the case of a true thermal state characterized by a *strictly* positive temperature  $T > 0$  (and a chemical potential  $\mu \in \mathbb{R}$  or, equivalently, a particle density  $\rho > 0$ ). On the one hand, the positive temperature case is simpler, because the *Fermi function*  $E \mapsto 1/(1 + \exp(E/T))$  on the real line  $\mathbb{R}$  is smooth in contrast to its zero-temperature limit, the Heaviside unit-step function. On the other hand, a reasonable definition of the EE (see (1.9)) is more complicated, because the thermal state is not a pure state. Nevertheless, due to the presence of the “smoothing parameter”  $T > 0$  the leading asymptotic growth of the EE as  $\alpha \rightarrow \infty$ , is determined by an asymptotic coefficient again going back to Widom [26, 29], see also [18, 8]. We denote this coefficient by  $\mathcal{B}_d$ , and it is introduced in (2.4).

It is interesting from a physics point of view to study the scaling asymptotics  $\alpha \rightarrow \infty$  as the temperature  $T$  varies. The double asymptotics of the EE and of its coefficient  $\mathcal{B}_d$  are not

simple to analyze and hard to guess by heuristic arguments. For low temperatures, that is small  $T > 0$ , this analysis has been performed in [17] for  $d = 1$  and in [24] for  $d \geq 2$  yielding a result consistent with that for  $T = 0$  in [12, 15].

For high temperatures quantum effects become weaker and the free Fermi gas should exhibit the properties of the corresponding classical free gas without correlations (for fixed particle density). In particular, the ideal Fermi gas should behave like the time-honored Maxwell–Boltzmann gas. Hence one purpose of our study is to determine the precise two-parameter asymptotics of the EE as  $\alpha \rightarrow \infty$ ,  $T \rightarrow \infty$ .

**1.2. Pseudo-differential operators and entropies.** In order to define the EE we first introduce the pseudo-differential operator

$$(1.1) \quad (\text{Op}_\alpha(a)u)(\mathbf{x}) := \frac{\alpha^d}{(2\pi)^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} e^{i\alpha \boldsymbol{\xi} \cdot (\mathbf{x} - \mathbf{y})} a(\boldsymbol{\xi}) u(\mathbf{y}) \, d\mathbf{y} d\boldsymbol{\xi}, \quad \mathbf{x} \in \mathbb{R}^d, \quad \alpha > 0,$$

with a smooth complex-valued function  $a$  that we call symbol, where  $u$  is a Schwartz function. The number  $\alpha > 0$  is a scaling parameter. In our large-scale applications it is either  $\alpha$  itself or a certain combination of  $\alpha$  with the temperature  $T$  that will become large.

The main role will be played by the *truncated Wiener–Hopf operator*

$$(1.2) \quad W_\alpha(a, \Lambda) := \chi_\Lambda \text{Op}_\alpha(a) \chi_\Lambda,$$

where  $\chi_\Lambda$  is the indicator function of a truncating Borel set  $\Lambda \subset \mathbb{R}^d$ . Clearly, for a bounded symbol  $a$  the operators  $\text{Op}_\alpha(a)$  and  $W_\alpha(a; \Lambda)$  are bounded on the Hilbert space  $L^2(\mathbb{R}^d)$ . Given a *test* function  $f : \mathbb{R} \rightarrow \mathbb{R}$ , we are interested in the operator  $f(W_\alpha(a, \Lambda))$  and in the operator difference

$$(1.3) \quad D_\alpha(a, \Lambda; f) := \chi_\Lambda f(W_\alpha(a, \Lambda)) \chi_\Lambda - W_\alpha(f \circ a, \Lambda),$$

where the symbol  $f \circ a$  is the composition of  $f$  and  $a$ .

For a bounded  $\Lambda$ , under appropriate conditions on  $a$  and  $f$  both operators in (1.3) are trace class. Nevertheless the operator (1.3) is trace class for a large class of unbounded sets  $\Lambda$  (see Condition 2.1 below). If there is no risk of confusion, we sometimes omit the dependence on the set  $\Lambda$  and write simply  $D_\alpha(a; f)$ . Our analysis of the scaling behavior of the entropies is based on the asymptotics of the trace  $\text{tr} D_\alpha(a; f)$  as  $\alpha \rightarrow \infty$ . The reciprocal parameter  $\alpha^{-1}$  can be naturally viewed as Planck’s constant, and hence the limit  $\alpha \rightarrow \infty$  can be regarded as the quasi-classical limit. By a straightforward change of variables the operator (1.3) is unitarily equivalent to  $D_1(a, \alpha\Lambda; f)$ , so that  $\alpha \rightarrow \infty$  can be also interpreted as a large-scale limit.

The macroscopic thermal equilibrium state of the free Fermi gas depends, first of all, on its classical single-particle Hamiltonian  $h = h(\boldsymbol{\xi})$ , sometimes also called the energy-momentum (dispersion) relation. The minimal condition on the function  $h$  that we impose are as follows. We assume that  $h \in C^\infty(\mathbb{R}^d)$  and for some  $m > 0$  the bounds are satisfied:

$$(1.4) \quad |\partial_{\boldsymbol{\xi}}^n h(\boldsymbol{\xi})| \lesssim \langle \boldsymbol{\xi} \rangle^{2m}, \quad \text{for all } n \in \mathbb{N}_0^d, \quad \text{and} \quad h(\boldsymbol{\xi}) \gtrsim |\boldsymbol{\xi}|^{2m} \quad \text{for sufficiently large } \boldsymbol{\xi} \in \mathbb{R}^d.$$

Here  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ , and the notations “ $\lesssim$ ”, “ $\gtrsim$ ”, “ $\langle \boldsymbol{\xi} \rangle$ ” are defined at the end of the introduction.

For a given  $h$ , (absolute) temperature  $T > 0$  and chemical potential  $\mu \in \mathbb{R}$  we introduce the *Fermi symbol*

$$(1.5) \quad a_{T, \mu}(\boldsymbol{\xi}) := \frac{1}{1 + \exp\left(\frac{h(\boldsymbol{\xi}) - \mu}{T}\right)}, \quad \boldsymbol{\xi} \in \mathbb{R}^d.$$

Next, we define the Rényi *entropy functions*  $\eta_\gamma : \mathbb{R} \rightarrow [0, \ln(2)]$  for each  $\gamma > 0$ . For  $\gamma \neq 1$ ,  $\eta_\gamma$  is given by

$$(1.6) \quad \eta_\gamma(t) := \begin{cases} \frac{1}{1-\gamma} \ln [t^\gamma + (1-t)^\gamma] & \text{for } t \in (0, 1) \\ 0 & \text{for } t \notin (0, 1) \end{cases},$$

and for  $\gamma = 1$  (the von Neumann case) it is defined by the limit

$$(1.7) \quad \eta_1(t) := \lim_{\gamma \rightarrow 1} \eta_\gamma(t) = \begin{cases} -t \ln(t) - (1-t) \ln(1-t) & \text{for } t \in (0, 1) \\ 0 & \text{for } t \notin (0, 1) \end{cases}.$$

Finally, we define the  $\gamma$ -Rényi *local entropy* associated with a bounded region  $\Lambda$  by

$$(1.8) \quad S_\gamma(T, \mu; \Lambda) := \text{tr } \eta_\gamma(W_1(a_{T, \mu}, \Lambda)),$$

and the  $\gamma$ -Rényi *entanglement entropy* (EE) with respect to the bipartition  $\mathbb{R}^d = \Lambda \cup (\mathbb{R}^d \setminus \Lambda)$  by

$$(1.9) \quad H_\gamma(T, \mu; \Lambda) := \text{tr } D_1(a_{T, \mu}, \Lambda; \eta_\gamma) + \text{tr } D_1(a_{T, \mu}, \mathbb{R}^d \setminus \Lambda; \eta_\gamma).$$

Conditions (1.4) guarantee that the above entropies are well-defined, see [17]. In the formula (1.9) either  $\Lambda$  or its complement  $\mathbb{R}^d \setminus \Lambda$  is assumed to be bounded, see Sect. 2 for details.

It is useful to observe that (for bounded  $\Lambda$ ) the local entropy (1.8) can be expressed as follows:

$$(1.10) \quad S_\gamma(T, \mu; \Lambda) = s_\gamma(T, \mu) |\Lambda| + \text{tr } D_1(a_{T, \mu}, \Lambda; \eta_\gamma),$$

where the function

$$(1.11) \quad s_\gamma(T, \mu) := \frac{\text{tr } W_1(\eta_\gamma \circ a_{T, \mu}, \Lambda)}{|\Lambda|} = (2\pi)^{-d} \int_{\mathbb{R}^d} \eta_\gamma(a_{T, \mu}(\boldsymbol{\xi})) \, d\boldsymbol{\xi},$$

is called the *entropy density*. Here  $|\Lambda|$  denotes the volume (Lebesgue measure) of the region  $\Lambda$ .

**1.3. Description of results.** Our first result is of non-asymptotic nature. In Sect. 3 we explore concavity properties of the function  $\eta_\gamma$ . First we notice that  $\eta_\gamma$  is concave on the interval  $[0, 1]$  for all  $\gamma \in (0, 2]$ , so in the case of bounded  $\Lambda$  one can use the Berezin–Lieb inequality to establish a lower bound for the local entropy (1.8) via its density (1.11), see Corollary 3.3. For the EE (1.9) this argument is not applicable, and instead we observe that  $\eta_\gamma$  is *operator concave* for  $\gamma \in (0, 1]$ . Next, instead of the Berezin–Lieb inequality we use the Davis operator inequality (3.5), which implies that the EE is non-negative for all  $\gamma \in (0, 1]$  (including the most important case  $\gamma = 1$  corresponding to the von Neumann EE), see Theorem 3.5. We do not know whether the EE is positive for  $\gamma \in (1, 2]$ . Later on however, in Sect. 6 we see that the EE becomes positive at large temperature for all  $\gamma \in (0, 2]$ , see remark 1 in Subsect. 6.3.

The main objective of the present paper is to study the asymptotics of  $H_\gamma(T, \mu; \alpha\Lambda)$  as  $\alpha \rightarrow \infty$  and  $T \rightarrow \infty$ . For this, we have to impose stronger conditions on the Hamiltonian  $h$  compared to (1.4), see Subsect. 5.1. In particular,  $h$  should be asymptotically homogeneous as  $\boldsymbol{\xi} \rightarrow \infty$ . We distinguish between two cases: the regime of a constant chemical potential  $\mu$  (i.e.  $\mu$  is constant as  $T \rightarrow \infty$ ), and the regime of a constant mean particle density

$$(1.12) \quad \varrho(T, \mu) := \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} a_{T, \mu}(\boldsymbol{\xi}) \, d\boldsymbol{\xi}.$$

To be precise, in the latter regime we consider the chemical potential  $\mu = \mu(T)$  as a function of temperature  $T$  such that  $\varrho(T, \mu)$  is kept (asymptotically) constant as  $T$  becomes large, i.e.

$$\varrho(T, \mu(T)) \rightarrow \rho \quad \text{as } T \rightarrow \infty,$$

with a constant  $\rho > 0$ . Under this condition the limit  $T \rightarrow \infty$  corresponds to the classical free gas limit. Indeed, the integrated density of states

$$(1.13) \quad \mathcal{N}(T) := (2\pi)^{-d} \int_{h(\boldsymbol{\xi}) < T} d\boldsymbol{\xi}$$

of the Hamiltonian  $h$  evaluated at temperature  $T$ , tends to infinity as  $T \rightarrow \infty$ . As a consequence,  $\rho/\mathcal{N}(T) \rightarrow 0$ . In physical terms the latter means that the number of particles per unit volume is small compared to the number of available states at temperature  $T$  per unit volume, which indicates the passage of the quantum gas to its classical regime.

The results on high temperature scaling asymptotics are presented in Theorem 6.1 (for constant  $\mu$ ) and in Theorem 6.3 (for the constant mean particle density). We postpone the discussion of these theorems until Section 6. Here we only mention two important facts: a) in both regimes the asymptotics still hold if  $T \rightarrow \infty$  and the scaling parameter  $\alpha$  is fixed, e.g.  $\alpha = 1$ , b) the EE with Rényi index  $\gamma > 2$ , becomes *negative* at fixed particle density and high temperature. This suggests that the case  $\gamma > 2$  is only of limited physical interest.

The main asymptotic orders as  $T \rightarrow \infty$  and  $\alpha = 1$  for both regimes have been announced in [16] for  $\gamma = 1$  for all dimensions  $d \geq 1$ . Here we provide proofs for all  $\gamma > 0$  and concentrate on the multi-dimensional case  $d \geq 2$ . The case  $d = 1$  is not discussed for lack of space.

The paper is organized as follows. In Sect. 2 we present the basic information such as conventions on the truncation regions and definition of the asymptotic coefficient  $\mathcal{B}_d$ . Apart from this, Sect. 2 also contains the results borrowed mostly from [25], that are used throughout the paper. In Sect. 3 we study the concavity of the functions  $\eta_\gamma$ ,  $\gamma > 0$  and discuss positivity of the entanglement entropy. In Sect. 4 we derive elementary trace class bounds for some abstract self-adjoint bounded operators. These bounds are based on estimates for *quasi-commutators* of the form  $f(A)J - Jf(B)$  with self-adjoint bounded  $A, B$ , bounded  $J$ , and suitable functions  $f$ . With the help of the bounds from Sect. 4 in Sect. 5 we obtain the large  $\alpha$  and  $T$  asymptotics for the operator  $D_\alpha(p_T, \Lambda; f)$  for a symbol  $p_T$  that models the Fermi symbol (1.5) in the fixed  $\mu$  or fixed  $\rho$  regimes. In Sect. 6 we collect our results on the high temperature asymptotics for the EE (1.9), see Theorems 6.1, 6.3. Their proofs are directly based on the formulas obtained in Sect. 5. In Subsect. 6.3 we also comment on the asymptotics of the local entropy (1.8). The Appendix contains a short calculation clarifying the structure of the Fermi symbol when the mean particle density is fixed as  $T \rightarrow \infty$ .

Throughout the paper we adopt the following conventions. For two non-negative numbers (or functions)  $X$  and  $Y$  depending on some parameters, we write  $X \lesssim Y$  (or  $Y \gtrsim X$ ) if  $X \leq CY$  with some positive constant  $C$  independent of those parameters. If  $X \lesssim Y$  and  $X \gtrsim Y$ , then we write  $X \asymp Y$ . To avoid confusion we often make explicit comments on the nature of (implicit) constants in the bounds.

For partial derivatives we use the standard notation  $\partial_{\boldsymbol{\xi}}^n := \partial_{\xi_1}^{n_1} \partial_{\xi_2}^{n_2} \cdots \partial_{\xi_d}^{n_d}$  for  $\boldsymbol{\xi} = (\xi_1, \xi_2, \dots, \xi_d) \in \mathbb{R}^d$  and  $n = (n_1, n_2, \dots, n_d) \in \mathbb{N}_0^d$ .

The notation  $B(\mathbf{z}, R) \subset \mathbb{R}^d$ ,  $\mathbf{z} \in \mathbb{R}^d$ ,  $R > 0$ , is used for the open ball of radius  $R$ , centered at the point  $\mathbf{z}$ . For any vector  $\mathbf{v} \in \mathbb{R}^d$ , we use the standard notation  $\langle \mathbf{v} \rangle := \sqrt{1 + |\mathbf{v}|^2}$ .

We dedicate this paper to the memory of Harold Widom (1932–2021). His ground-breaking results on the asymptotic expansions for traces of pseudo-differential operators have been highly influential to many researchers including us. Without his results our present and also previous contributions to the study of fermionic entanglement entropy would have been unthinkable. We are deeply indebted to Widom's ingenious insights. All three of us had the honor and pleasure of meeting him at a memorable workshop in March 2017 hosted by the American Institute of Mathematics in San Jose, California, USA.

## 2. BASIC DEFINITIONS AND BASIC RESULTS

In this section we collect some definitions and facts from [25] that are instrumental in the proof of our main results. We always assume that  $d \geq 2$ .

**2.1. Conditions on the truncating region  $\Lambda$ .** We call a domain (an open, connected set) Lipschitz, if it can be described locally as an epigraph of a Lipschitz function, see [21] for details. We call  $\Lambda$  a *Lipschitz region* if  $\Lambda$  is a union of finitely many Lipschitz domains such that their closures are pair-wise disjoint. We always assume that  $\Lambda$  satisfies the following condition.

- Condition 2.1.** (1) *The set  $\Lambda \subset \mathbb{R}^d$ ,  $d \geq 2$ , is a Lipschitz region, and either  $\Lambda$  or  $\mathbb{R}^d \setminus \Lambda$  is bounded.*  
 (2) *The boundary  $\partial\Lambda$  is piece-wise  $C^1$ -smooth.*

Note that  $\Lambda$  and  $\mathbb{R}^d \setminus \bar{\Lambda}$  satisfy Condition 2.1 simultaneously.

**2.2. The asymptotic coefficient  $\mathcal{B}_d$  and its properties.** Suppose that  $a \in C^\infty(\mathbb{R}^d)$  satisfies the condition

$$(2.1) \quad |\partial_{\boldsymbol{\xi}}^n a(\boldsymbol{\xi})| \lesssim \langle \boldsymbol{\xi} \rangle^{-\beta} \quad \text{with some } \beta > d,$$

for all  $\boldsymbol{\xi} \in \mathbb{R}^d$ , all  $n \in \mathbb{N}_0^d$  and some implicit constants that may depend on  $n$ . Here we use the notations  $\lesssim$ , and  $\langle \boldsymbol{\xi} \rangle$  both defined at the end of the Introduction.

In order to state the asymptotic formula for  $\text{tr } D_\alpha(a, \Lambda; f)$  we need to introduce first the asymptotic coefficient. For a function  $f : \mathbb{R} \rightarrow \mathbb{C}$  and any  $u, v \in \mathbb{R}$  we consider the integral

$$(2.2) \quad U(u, v; f) := \int_0^1 \frac{f((1-t)u + tv) - [(1-t)f(u) + tf(v)]}{t(1-t)} dt.$$

It is well-defined for any Hölder continuous function  $f$  and positive if  $f$  is concave.

For  $d \geq 2$  and every unit vector  $\mathbf{e} \in \mathbb{S}^{d-1}$  we define a functional of the symbol  $a$  by the principal-value integral:

$$(2.3) \quad \mathcal{A}_d(a, \mathbf{e}; f) := \frac{1}{8\pi^2} \lim_{\epsilon \downarrow 0} \int \int_{\mathbb{R}^d, |t| > \epsilon} \frac{U(a(\boldsymbol{\xi}), a(\boldsymbol{\xi} + t\mathbf{e}); f)}{t^2} dt d\boldsymbol{\xi}.$$

Finally we define the main asymptotic coefficient by

$$(2.4) \quad \mathcal{B}_d(a, \partial\Lambda; f) := \frac{1}{(2\pi)^{d-1}} \int_{\partial\Lambda} \mathcal{A}_d(a, \mathbf{n}_{\mathbf{x}}; f) dS_{\mathbf{x}},$$

where  $\mathbf{n}_x$  is the (unit outward) normal vector at the point  $x \in \partial\Lambda$  and  $dS_x$  is the canonical  $(d-1)$ -dimensional area measure on the boundary surface  $\partial\Lambda$ .

For functions  $f \in C^2(\mathbb{R})$  under the condition (2.1) the integral (2.3) exists in the usual sense, is bounded uniformly in  $\mathbf{e} \in \mathbb{S}^{d-1}$ , and hence (2.4) is trivially finite, see [25, Section 3]. However, in order to accommodate the Rényi functions  $\eta_\gamma$  we allow non-smooth functions satisfying the following condition.

**Condition 2.2.** *The function  $f$  is in  $C^2(\mathbb{R} \setminus X) \cap C(\mathbb{R})$  where  $X = \{x_1, x_2, \dots, x_N\} \subset \mathbb{R}$  is some finite set. The function  $f$  satisfies for all  $x \in \mathbb{R} \setminus X$  the bound*

$$(2.5) \quad |f^{(k)}(x)| \lesssim \sum_{z \in X} |x - z|^{\delta - k}, \quad k = 0, 1, 2,$$

with some  $\delta > 0$ .

Under this condition  $\mathcal{B}_d(a, \partial\Lambda; f)$  is finite:

**Proposition 2.3.** [25, Corollary 3.4] *Assume that  $f$  satisfies Condition 2.2 with some  $\delta > 0$ , and that  $\Lambda$  satisfies Condition 2.1. Let  $a$  be a real-valued symbol satisfying (2.1) with  $\beta > \max\{d, d\delta^{-1}\}$ . Then the coefficient  $\mathcal{B}_d(a, \partial\Lambda; f)$  is finite.*

Let us point out a few simple properties of this coefficient.

**Remark 2.4.** (1) Since  $\mathcal{A}_d(a, \mathbf{e}; f) = \mathcal{A}_d(a, -\mathbf{e}; f)$ , the coefficients  $\mathcal{B}_d$  for the regions  $\Lambda$  and  $\mathbb{R}^d \setminus \bar{\Lambda}$  coincide.

(2) If the symbol  $a$  is spherically symmetric (for example, if the Hamiltonian  $h$  in the definition of the Fermi symbol  $a_{T,\mu}$  is spherically symmetric) then the surface area  $|\partial\Lambda|$  factors out of  $\mathcal{B}_d(a, \partial\Lambda; f)$ . The remaining integral  $\mathcal{A}_d(a, \mathbf{e}; f)$  (for any choice of unit vector  $\mathbf{e}$ ) is still very complicated and hard to compute explicitly for general  $f$ .

(3) Because of the definition (2.2), the coefficient (2.4) is non-negative for concave functions  $f$ , and non-positive for convex ones. For example, the function  $\eta_\gamma$  with  $0 < \gamma \leq 2$  is concave on the interval  $(0, 1)$  (see Lemma 3.1), and hence for symbols  $a$  such that  $0 \leq a \leq 1$ , we have  $\mathcal{B}_d(a, \partial\Lambda; \eta_\gamma) \geq 0$ .

One other, less obvious important property of the coefficient  $\mathcal{B}_d(a, \partial\Lambda; f)$  is its continuity as a function of the symbol  $a$ . To state the result, in the next and subsequent assertions we consider a family of symbols  $\{a_0, a_\lambda\}$ ,  $\lambda > 0$ , all of them satisfying (2.1) with some  $\beta > \max\{d, d\delta^{-1}\}$ , uniformly in  $\lambda$ , and is such that  $a_\lambda \rightarrow a_0$  pointwise as  $\lambda \downarrow 0$ .

**Proposition 2.5.** [25, Corollary 3.5] *Let the function  $f$  and region  $\Lambda$  be as in Proposition 2.3. Then*

$$(2.6) \quad \mathcal{B}_d(a_\lambda, \partial\Lambda; f) \rightarrow \mathcal{B}_d(a_0, \partial\Lambda; f), \quad \lambda \downarrow 0.$$

**2.3. Asymptotics of  $\text{tr } D_\alpha(a, \Lambda; f)$ .** Now we are in a position to state the required asymptotic facts.

**Proposition 2.6.** [25, Theorem 2.3] *Assume that  $f$  satisfies Condition 2.2 with some  $\delta > 0$ , and that  $\Lambda$  is a region satisfying Condition 2.1. Suppose that  $a \in C^\infty(\mathbb{R}^d)$ ,  $d \geq 2$ , is a real-valued function that satisfies (2.1) with  $\beta > \max\{d, d\delta^{-1}\}$ . Then the operator  $D_\alpha(a, \Lambda; f)$  is trace class and*

$$(2.7) \quad \lim_{\alpha \rightarrow \infty} \alpha^{1-d} \text{tr } D_\alpha(a, \Lambda; f) = \mathcal{B}_d(a, \partial\Lambda; f).$$

This limit is uniform in the class of symbols  $a$  that satisfy (2.1) with the same implicit constants.

Proposition 2.6 was used in [25] to find the scaling asymptotics for the entanglement entropy at fixed temperature. To study the high-temperature regime, we need the continuity of this result in the symbol  $a$ :

**Corollary 2.7.** *Let the function  $f$  and the region  $\Lambda$  be as in Proposition 2.6. Then*

$$(2.8) \quad \lim \alpha^{1-d} \operatorname{tr} D_\alpha(a_\lambda, \Lambda; f) = \mathcal{B}_d(a_0, \partial\Lambda; f),$$

where the limit is taken as  $\alpha \rightarrow \infty$  and  $\lambda \rightarrow 0$  at the same time.

*Proof.* According to Proposition 2.6,

$$\lim_{\alpha \rightarrow \infty} \alpha^{1-d} \operatorname{tr} D_\alpha(a_\lambda, \Lambda; f) = \mathcal{B}_d(a_\lambda, \partial\Lambda; f),$$

uniformly in  $\lambda$ . Together with (2.6) this leads to (2.8).  $\square$

The next two lemmata describe the asymptotics for “small” symbols.

**Proposition 2.8.** [25, Theorem 2.5] *Let the region  $\Lambda$  be as in Proposition 2.6. Let  $f_0(t) := M|t|^\delta$  with some complex  $M$  and  $\delta > 0$ . Suppose that the function  $f \in \mathcal{C}^2(\mathbb{R} \setminus \{0\})$  satisfies the condition*

$$(2.9) \quad \lim_{t \rightarrow 0} |t|^{n-\delta} \frac{d^n}{dt^n} (f(t) - f_0(t)) = 0, \quad n = 0, 1, 2.$$

Then

$$(2.10) \quad \lim_{\alpha \rightarrow \infty} \lim_{\lambda \rightarrow 0} (\alpha^{1-d} \lambda^{-\delta} \operatorname{tr} D_\alpha(\lambda a_\lambda, \Lambda; f)) = \mathcal{B}_d(a_0, \partial\Lambda; f_0).$$

In the next proposition instead of the homogeneous function  $t \mapsto |t|^\delta$  we consider the function

$$(2.11) \quad \eta(t) := -t \ln |t|, \quad t \in \mathbb{R},$$

which still leads to a homogeneous asymptotic behavior.

**Proposition 2.9.** [25, Theorem 2.6] *Let the region  $\Lambda$  be as Proposition 2.6. Suppose that the function  $f \in \mathcal{C}^2(\mathbb{R} \setminus \{0\})$  satisfies the condition*

$$(2.12) \quad \lim_{t \rightarrow 0} |t|^{n-1} \frac{d^n}{dt^n} (f(t) - \eta(t)) = 0, \quad n = 0, 1, 2.$$

Then

$$(2.13) \quad \lim_{\alpha \rightarrow \infty} \lim_{\lambda \rightarrow 0} (\alpha^{1-d} \lambda^{-1} \operatorname{tr} D_\alpha(\lambda a_\lambda, \Lambda; f)) = \mathcal{B}_d(a_0, \partial\Lambda; \eta).$$

The asymptotic results listed above are useful but not sufficient for our study. Given that our symbol of interest (1.5) depends on two parameters, i.e.  $T$  and  $\mu$ , in the course of the proof we come across symbols that do not satisfy (2.1) uniformly in those parameters, which makes the above results inapplicable. Next we consider a wider class of symbols that we call *multi-scale* symbols, see [17, Sect. 3], which allows us to circumvent this difficulty.



**2.4. Multi-scale symbols.** Consider  $a \in C^\infty(\mathbb{R}^d)$  for which there exist positive continuous functions  $v$  and  $\tau$  such that  $v$  is bounded and

$$(2.14) \quad |\partial_{\xi}^k a(\xi)| \lesssim \tau(\xi)^{-|k|} v(\xi), \quad k \in \mathbb{N}_0^d, \quad \xi \in \mathbb{R}^d,$$

with constants independent of  $\xi$ . It is natural to call  $\tau$  the *scale (function)* and  $v$  the *amplitude (function)*. The scale  $\tau$  is assumed to be globally Lipschitz with Lipschitz constant  $\nu < 1$ , that is,

$$(2.15) \quad |\tau(\xi) - \tau(\eta)| \leq \nu |\xi - \eta|, \quad \text{for all } \xi, \eta \in \mathbb{R}^d.$$

Under this assumption the amplitude  $v$  is assumed to satisfy the bounds

$$(2.16) \quad v(\eta) \asymp v(\xi), \quad \text{for all } \eta \in B(\xi, \tau(\xi)),$$

with implicit constants independent of  $\xi$  and  $\eta$ . It is useful to think of  $v$  and  $\tau$  as (functional) parameters. They, in turn, can depend on other parameters, e.g. numerical parameters like  $\alpha$  and  $T$ . For example, all the results in the previous subsections assumed that  $a$  satisfied (2.1), which translates into (2.14) with  $v(\xi) = \langle \xi \rangle^{-\beta}$  and  $\tau(\xi) = 1$ . On the other hand, in Section 5 we encounter amplitudes and scales depending on the temperature  $T$ .

We need only one result involving multi-scale symbols.

**Proposition 2.10.** [17, Lemma 3.4] *Suppose that the domain  $\Lambda$  satisfies Condition 2.1, and let the functions  $\tau$  and  $v$  be as described above. Suppose that the symbol  $a$  satisfies (2.14), and that the conditions*

$$(2.17) \quad \alpha \tau_{\inf} \gtrsim 1, \quad \tau_{\inf} := \inf_{\xi \in \mathbb{R}^d} \tau(\xi) > 0,$$

hold. Then for any  $q \in (0, 1]$  we have

$$(2.18) \quad \|[\text{Op}_\alpha(a), \chi_\Lambda]\|_q^q \lesssim \alpha^{d-1} \int_{\mathbb{R}^d} \frac{v(\xi)^q}{\tau(\xi)} d\xi.$$

This bound is uniform in the symbols  $a$  satisfying (2.14) with the same implicit constants.

### 3. ON POSITIVITY OF RÉNYI ENTROPIES

In this section we are concerned with the positivity of the entanglement entropy (1.9). In this question the central issue is concavity of the functions  $\eta_\gamma$  for various values of  $\gamma$ . Since these results are not of asymptotic character, we assume that  $\alpha = 1$ .

**3.1. Concavity of the entropy function  $\eta_\gamma$ .** First we establish concavity of  $\eta_\gamma = \eta_\gamma(t)$  as a function of the variable  $t \in (0, 1)$ , and establish its consequences for the local and entanglement entropies. The next lemma is elementary.

**Lemma 3.1.** *The Rényi entropy function  $\eta_\gamma$  is concave on the interval  $(0, 1)$  if  $0 < \gamma \leq 2$ , and neither convex nor concave if  $\gamma > 2$ .*

*Proof.* We simply compute the second derivative of  $\eta_\gamma$  on  $(0, 1)$ . If  $\gamma = 1$ , then  $\eta_1''(t) = -t^{-1}(1-t)^{-1} < 0$ , and hence  $\eta_1$  is concave.

Now assume that  $\gamma \neq 1$ . Then a direct differentiation gives the formula

$$(3.1) \quad \eta_\gamma''(t)(t^\gamma + (1-t)^\gamma)^2 = -\gamma(t(1-t))^{\gamma-2} - \frac{\gamma}{1-\gamma}(t^{\gamma-1} - (1-t)^{\gamma-1})^2.$$

If  $\gamma < 1$ , then the left-hand side is clearly negative for all  $t \in (0, 1)$ , and hence  $\eta_\gamma$  is concave. If  $\gamma = 2$ , then the right-hand side equals  $-8t(1-t) < 0$ , and hence  $\eta_2$  is concave. If  $\gamma > 2$ , then it follows from (3.1) that

$$\eta_\gamma''(0) = \frac{\gamma}{\gamma-1} > 0, \quad \text{and} \quad \eta_\gamma''(1/2) = -\gamma < 0,$$

and hence the function  $\eta_\gamma$  is neither concave nor convex.

It remains to consider the case  $\gamma \in (1, 2)$ . Rewrite (3.1) as

$$\begin{aligned} \eta_\gamma''(t)(t^\gamma + (1-t)^\gamma)^2 &= -\frac{\gamma}{\gamma-1}g_{\gamma-1}(t), \\ g_p(t) &:= p(t(1-t))^{p-1} - (t^p - (1-t)^p)^2, \end{aligned}$$

and let us prove that  $g_p(t) \geq 0, t \in (0, 1)$ , for all  $p \in (0, 1)$ . The inequality  $g_p(t) \geq 0$  is equivalent to

$$(3.2) \quad (t(1-t))^{1-p}(t^{2p} + (1-t)^{2p}) \leq 2t(1-t) + p.$$

Denote

$$A = 2^{p-1} \max_{t \in [0,1]} (t^{2p} + (1-t)^{2p}) = \begin{cases} 2^{-p} & \text{if } 0 < p < 1/2 \\ 2^{p-1} & \text{if } 1/2 \leq p < 1 \end{cases}.$$

Thus, by the Young inequality

$$ab \leq \frac{a^{q_1}}{q_1} + \frac{b^{q_2}}{q_2}, \quad a, b \geq 0, \quad q_1, q_2 > 1, \quad \frac{1}{q_1} + \frac{1}{q_2} = 1,$$

used with  $a = (t(1-t))^{1-p}, q_1 = (1-p)^{-1}$  and  $b = 1, q_2 = p^{-1}$ , we get

$$\begin{aligned} (t(1-t))^{1-p}(t^{2p} + (1-t)^{2p}) &\leq A(2t(1-t))^{1-p} \\ &\leq A(1-p)(2t(1-t)) + Ap \leq 2t(1-t) + p. \end{aligned}$$

This coincides with (3.2) and the proof of the lemma is now complete.  $\square$

The concavity property, established above is useful for studying the local entropy (1.8), which is well-defined for bounded domains  $\Lambda$ . To this end, we recall the following abstract fact known as the Berezin (or Berezin–Lieb) inequality, see [5]. Let  $\mathcal{H}$  be a (separable) Hilbert space,  $A$  a self-adjoint operator  $A$  on  $\mathcal{H}$  and an orthogonal projection  $P$  on  $\mathcal{H}$ .

**Proposition 3.2.** *Let  $f : I \rightarrow \mathbb{R}$  be a concave function on the interval  $I \subset \mathbb{R}$ . Suppose that the spectrum of  $A$  is contained in  $I$ ,  $PAP$  is compact and that the operator*

$$(3.3) \quad \mathcal{D}(A, P; f) := Pf(PAP)P - Pf(A)P$$

*is trace class (If  $0 \notin I$ , then the function  $f(PAP)$  is understood as an operator on the subspace  $P\mathcal{H}$ ). Then  $\text{tr } \mathcal{D}(A, P; f) \geq 0$ .*

The above proposition can be found, e.g. in [14, Theorem A.1], although it is stated there for convex rather than concave functions.

Now, in order to study the entropies, assume that  $\Lambda$  satisfies Condition 2.1, and the function  $h$  in (1.5) is as specified in (1.4). Thus the function (1.5) satisfies (2.1) for each  $T > 0$  and  $\mu \in \mathbb{R}$ . We observe that the function  $\eta_\gamma$  satisfies Condition 2.2 with arbitrary  $\delta < \min\{1, \gamma\}$  for all  $\gamma > 0$ , and the set  $X = \{0, 1\}$ . Thus, due to Proposition 2.6, the operators  $D_1(a_{T,\mu}, \Lambda; \eta_\gamma)$

and  $D_1(a_{T,\mu}, \mathbb{R}^d \setminus \Lambda; \eta_\gamma)$  are trace class, so that the EE (1.9) is finite. If we assume that  $\Lambda$  is bounded, then by (1.10) the local entropy is also finite, and it possesses the following property.

**Corollary 3.3.** *Suppose that the region  $\Lambda$  satisfies Condition 2.1 and is bounded. Assume that the function  $h$  satisfies (1.4), and that  $0 < \gamma \leq 2$ . Then  $\text{tr } D_1(a_{T,\mu}, \Lambda; \eta_\gamma) \geq 0$  and the local entropy (1.8) satisfies the inequality*

$$(3.4) \quad S_\gamma(T, \mu; \Lambda) \geq s_\gamma(T, \mu)|\Lambda|,$$

where  $s_\gamma(T, \mu)$  is the entropy density defined in (1.11).

*Proof.* We use Proposition 3.2 with  $A = \text{Op}_1(a_{T,\mu})$ ,  $P = \chi_\Lambda$  and the concave function  $f = \eta_\gamma$ . Since  $0 \leq A \leq \mathbb{1}$  and  $PAP$  is compact, Proposition 3.2 is indeed applicable, and hence  $\text{tr } D_1(a, \Lambda; \eta_\gamma) \geq 0$ , which by (1.10) entails (3.4).  $\square$

We stress that Proposition 3.2 cannot be applied if the set  $\Lambda$  is unbounded, since in this case the operator  $\chi_\Lambda \text{Op}_\alpha(a_{T,\mu}) \chi_\Lambda$  does not have to be compact. Thus Corollary (3.3) cannot be used to determine the sign of the EE(1.9). To rectify this we use the *operator concavity* of  $\eta_\gamma$ .

**3.2. Operator concavity of  $\eta_\gamma$ .** Let us recall the basic definitions, see e.g. [19].

Let  $I \subseteq \mathbb{R}$  be an interval. A function  $f : I \rightarrow \mathbb{R}$  is called *operator monotone* if for all bounded self-adjoint operators  $A$  and  $B$  on a Hilbert space  $\mathcal{H}$  with spectra in  $I$  and with  $A \leq B$  the (operator) inequality  $f(A) \leq f(B)$  on  $\mathcal{H}$  holds.

A function  $f : I \rightarrow \mathbb{R}$  on the interval  $I$  is called *operator concave* if for all bounded self-adjoint operators  $A$  and  $B$  on  $\mathcal{H}$  with spectra in  $I$  and for all  $\lambda \in [0, 1]$  the inequality  $f(\lambda A + (1-\lambda)B) \geq \lambda f(A) + (1-\lambda)f(B)$  holds. If the inequality sign is reversed, then the function  $f$  is said to be *operator convex*.

Every operator monotone (concave) function is monotone (concave).

We use the following standard facts.

- (1) The function  $t \mapsto t^p$ ,  $t \in [0, \infty)$ , is operator concave and operator monotone if  $0 < p \leq 1$ .
- (2) The function  $t \mapsto \ln(t)$ ,  $t \in (0, \infty)$ , is operator concave and operator monotone.
- (3) The function  $t \mapsto -t \ln(t)$ ,  $t \in [0, \infty)$ , is operator concave.

All these examples can be found in [2]. It was proved in [9] that any operator concave function  $f$  satisfies the following *Davis operator inequality* for all orthogonal projections  $P$  on  $\mathcal{H}$  and all bounded self-adjoint operators  $A$  with spectrum in  $I$ :

$$(3.5) \quad Pf(PAP)P \geq Pf(A)P.$$

If  $0 \notin I$ , then the function  $f(PAP)$  is understood as an operator on the subspace  $P\mathcal{H}$ . If  $0 \in I$  and  $f(0) = 0$ , then  $f(PAP) \geq Pf(A)P$ .

**Theorem 3.4.** *If  $0 < \gamma \leq 1$  then  $\eta_\gamma$  is operator concave on  $I = [0, 1]$ .*

*Proof.* Assume first that  $0 < \gamma < 1$ . Let  $A$  and  $B$  be self-adjoint operators, such that  $0 \leq A, B \leq \mathbb{1}$ . The function  $f_\gamma(t) = t^\gamma + (1-t)^\gamma$  is operator concave on the interval  $I = [0, 1]$  (example (1) above) and the logarithm is operator monotone on  $(0, \infty)$  (example (2)). Thus for all  $\lambda \in [0, 1]$  we have

$$\eta_\gamma(\lambda A + (1-\lambda)B) = \frac{1}{1-\gamma} \ln[f_\gamma(\lambda A + (1-\lambda)B)] \geq \frac{1}{1-\gamma} \ln[\lambda f_\gamma(A) + (1-\lambda)f_\gamma(B)].$$

Now, since the logarithm is operator concave on  $(0, \infty)$  (again example (2)), the right-hand side is greater than or equal to

$$\frac{1}{1-\gamma} [\lambda \ln(f_\gamma(A)) + (1-\lambda) \ln(f_\gamma(B))] = \lambda \eta_\gamma(A) + (1-\lambda) \eta_\gamma(B),$$

as required.

If  $\gamma = 1$  then we proceed more directly and use that  $\eta(t) := -t \ln(t)$  is an operator concave function (see example (3)). This immediately implies that  $\eta_1(t) = \eta(t) + \eta(1-t)$  is also operator concave.  $\square$

Now we can use the inequality (3.5) with  $f = \eta_\gamma$ ,  $0 < \gamma \leq 1$ . Setting  $P = \chi_\Lambda$ ,  $A = \text{Op}_1(a_{T,\mu})$ , and using Proposition 2.6 yields the following theorem.

**Theorem 3.5.** *Suppose that the region  $\Lambda$  satisfies Condition 2.1. Assume that the function  $h$  is as in (1.4), and that  $0 < \gamma \leq 1$ . Then the operator  $D_1(a_{T,\mu}, \Lambda; \eta_\gamma)$  is trace class and positive, and hence the  $\gamma$ -Rényi entanglement entropy (1.9) is non-negative.*

This method however cannot be used for the  $\gamma$ -Rényi EE with  $\gamma > 1$  because of the following negative result.

**Theorem 3.6.** *If  $\gamma > 1$  then  $\eta_\gamma$  is not operator concave on  $I = [0, 1]$ .*

*Proof.* For convenience instead of  $\eta_\gamma$  we consider the function

$$f_\gamma(s) := -\eta_\gamma(s + \frac{1}{2}) = \frac{1}{\gamma - 1} \ln \left[ \left(\frac{1}{2} - s\right)^\gamma + \left(\frac{1}{2} + s\right)^\gamma \right], \quad s \in \left[-\frac{1}{2}, \frac{1}{2}\right].$$

Our objective is to show that  $f_\gamma$  is not operator convex on  $[-1/2, 1/2]$ . If  $f_\gamma$  were operator convex, then by [4, Corollary 1],  $f_\gamma$  would be analytic on the complex plane with cuts along the half-lines  $(-\infty, -1/2)$  and  $(1/2, \infty)$ . Let us prove that such an analytic continuation of  $f_\gamma$  is impossible. Let  $s = iy/2$  with  $y > 0$ . Then

$$\frac{1}{2} \pm s = \frac{1}{2} \sqrt{1 + y^2} \exp[\pm i \tan^{-1}(y)]$$

so that

$$(3.6) \quad \left(\frac{1}{2} - s\right)^\gamma + \left(\frac{1}{2} + s\right)^\gamma = 2^{1-\gamma} (1 + y^2)^{\gamma/2} \cos[\gamma \tan^{-1}(y)].$$

Since  $\gamma > 1$ , there exists a finite  $y_0 > 0$  such that  $\gamma \tan^{-1}(y_0) = \pi/2$ , so that the right-hand side of (3.6) changes sign at  $y = y_0$ . This implies that the function  $f_\gamma$  has a branching point at  $s_0 = iy_0/2$ , and hence it cannot be analytic in the whole of upper half-plane. This proves that  $f_\gamma$  is not operator convex, as claimed.  $\square$

#### 4. QUASI-COMMUTATOR BOUNDS

Here we collect some bounds in the Schatten–von Neumann classes  $\mathfrak{S}_p$ ,  $p > 0$ , on a (separable) Hilbert space  $\mathcal{H}$ , see e.g. [6, Chapter 11] for the definition and properties of  $\mathfrak{S}_p$ . We recall that the functional  $\|A\|_p := (\text{tr } |A|^p)^{1/p}$  defines on  $\mathfrak{S}_p$  a norm if  $p \geq 1$  and a quasi-norm if  $0 < p < 1$ . However, for  $0 < p < 1$  the quasi-norm is actually a  $p$ -norm, that is, it satisfies the following “triangle inequality” for all  $A, B \in \mathfrak{S}_p$ :

$$(4.1) \quad \|A + B\|_p^p \leq \|A\|_p^p + \|B\|_p^p.$$

This inequality is used systematically in what follows.

In this section the main part is played by estimates for *quasi-commutators*  $f(A)J - Jf(B)$  with self-adjoint  $A, B$  and bounded  $J$ . The following assertion is adapted from [23, Theorem 2.4].

**Proposition 4.1.** *Suppose that  $f$  satisfies Condition 2.2 with some  $\delta > 0$ . Let  $A, B$  be two bounded self-adjoint operators and let  $J$  be a bounded operator. Suppose that  $AJ - JB \in \mathfrak{S}_q$  where  $q$  satisfies  $0 < q < \min\{1, \delta\}$ . Then*

$$(4.2) \quad \|f(A)J - Jf(B)\|_1 \lesssim \|J\|^{1-q} (1 + \|A\|^{\delta-q} + \|B\|^{\delta-q}) \|AJ - JB\|_q^q,$$

with a constant independent of the operators  $A, B, J$ . The constant in (4.2) may depend on the set  $X$  in Condition 2.2, and is uniform in the functions satisfying (2.5) with the same implicit constants.

One should say that Theorem 2.4 from [23] provides estimates of the type (4.2) in arbitrary normed or quasi-normed operator ideals of compact operators and gives a more precise dependence of the constant on the function  $f$ . In this paper the bound (4.2) is sufficient.

All subsequent bounds involving the function  $f$  are uniform in  $f$  in the sense specified in Proposition 4.1. We apply Proposition 4.1 to obtain various bounds for the operator  $\mathcal{D}(A, P; f) = Pf(PAP)P - Pf(A)P$  with an orthogonal projection  $P$ , as defined above in (3.3).

**Corollary 4.2.** *Let the function  $f$  and the parameter  $q$  be as in Proposition 4.1. Let  $A, B$  be bounded self-adjoint operators and let  $J$  be a bounded operator such that*

$$(4.3) \quad [A, J] = [B, J] = 0, \quad (A - B)J = 0.$$

Then

$$(4.4) \quad \|\mathcal{D}(A, P; f)J\|_1 + \|J\mathcal{D}(A, P; f)\|_1 \lesssim \|[P, J]\|_q^q + \|[JA, P]\|_q^q,$$

and

$$(4.5) \quad \|\mathcal{D}(A, P; f)J - J\mathcal{D}(B, P; f)\|_1 \lesssim \|[J, P]\|_q^q + \|[J, P]\|_1.$$

The implicit constants in the above bounds depend on the norms  $\|A\|, \|B\|$ , and  $\|J\|$ , but they are uniform in the operators  $A, B, J$  whose norms are bounded by the same constants. They are also uniform in the functions  $f$  in the sense specified in Proposition 4.1.

*Proof.* The proof is based mainly on the bound (4.2). The assumption (4.3) considerably simplifies the calculations, and we use it without further comments.

We prove (4.4) for the first term on the left-hand side only, as the second one is done in the same way. We rewrite

$$(4.6) \quad \mathcal{D}(A, P; f)J = P(f(PAP)PJ - PJf(A)) - P[f(A), PJ].$$

Then we use (4.2) to estimate the first term on the right-hand side,

$$\begin{aligned} \|Pf(PAP)PJ - PJf(A)\|_1 &\leq \|f(PAP)PJ - PJf(A)\|_1 \\ &\lesssim \|PAPJ - PJA\|_q^q \lesssim \|P(AJ - JA)P\|_q^q + \|[P, J]\|_q^q + \|[JA, P]\|_q^q \\ &= \|[P, J]\|_q^q + \|[JA, P]\|_q^q. \end{aligned}$$

For the second term on the right-hand side of (4.6) we also use (4.2),

$$\begin{aligned} \|P[f(A), PJ]\|_1 &\lesssim \|APJ - PJA\|_q^q \lesssim \|(AJ - JA)P\|_q^q + \|[P, J]\|_q^q + \|[JA, P]\|_q^q \\ &= \|[P, J]\|_q^q + \|[JA, P]\|_q^q. \end{aligned}$$

Adding up the above two bounds we get (4.4).

In order to prove (4.5) we first estimate the difference

$$(4.7) \quad \begin{aligned} Pf(PAP)PJ - JPf(PBP)P &= P(f(PAP)J - Jf(PBP))P \\ &\quad + Pf(PAP)[P, J] - [J, P]f(PBP)P. \end{aligned}$$

Then we use (4.2) to estimate the first term on the right-hand side,

$$\|P(f(PAP)J - Jf(PBP))P\|_1 \lesssim \|PAPJ - JPBP\|_q^q \leq \|[J, P]\|_q^q.$$

To estimate the second and the third term on the right-hand side of (4.7) notice that  $\|f(A)\| \lesssim 1$ ,  $\|f(B)\| \lesssim 1$  uniformly in the operators  $A, B$  and in the function  $f$ . Consequently,

$$\|Pf(PAP)[P, J] - [J, P]f(PBP)P\|_1 \leq 2\|[J, P]\|_1 \|f\|_{L^\infty} \lesssim \|[J, P]\|_1.$$

This ensures (4.5) for (4.7).

Now we estimate

$$(4.8) \quad Pf(A)PJ - JPf(B)P = P(f(A)J - Jf(B))P + Pf(A)[P, J] - [J, P]f(B)P.$$

In view of (4.3), the first term on the right-hand side vanishes, and hence

$$\|Pf(A)PJ - JPf(B)P\|_1 \leq 2\|[P, J]\|_1 \|f\|_{L^\infty} \lesssim \|[J, P]\|_1.$$

Together with the bound (4.5) for (4.7) this completes the proof of (4.5).  $\square$

**Corollary 4.3.** *Let  $A, B, J$  be arbitrary bounded operators that satisfy (4.3), and let  $\mathbb{1}$  be the identity operator. Then*

$$(4.9) \quad \begin{aligned} \|\mathcal{D}(A, P; f) - \mathcal{D}(B, P; f)\|_1 &\lesssim \|[J, P]\|_q^q + \|[(\mathbb{1} - J)A, P]\|_q^q \\ &\quad + \|[(\mathbb{1} - J)B, P]\|_q^q + \|[J, P]\|_1. \end{aligned}$$

*This bound is uniform in  $A, B, J$  and in the function  $f$  in the same sense as in Corollary 4.2.*

*Proof.* We represent

$$\begin{aligned} \mathcal{D}(A, P; f) - \mathcal{D}(B, P; f) &= \mathcal{D}(A, P; f)J - J\mathcal{D}(B, P; f) \\ &\quad + \mathcal{D}(A, P; f)(\mathbb{1} - J) - (\mathbb{1} - J)\mathcal{D}(B, P; f), \end{aligned}$$

and apply Corollary 4.2.  $\square$

## 5. PREPARATIONS FOR HIGH TEMPERATURE ASYMPTOTICS: THE MODEL SYMBOL

The purpose of this section is to obtain the large  $\alpha$  and large  $T$  asymptotics for the operator  $D_\alpha(p_T, \Lambda; f)$  with the symbol  $p_T$  (see (5.4)) modeling the original Fermi symbol (1.5). We recall that  $d \geq 2$  throughout.

**5.1. Conditions on the single-particle Hamiltonian  $h$ .** First we need to specify the conditions on the Hamiltonian  $h$  in (1.5) in addition to (1.4).

We assume that  $h \in C^\infty(\mathbb{R}^d)$  and that for some  $m \in \mathbb{N}$ , the following bound holds:

$$(5.1) \quad |\partial_{\boldsymbol{\xi}}^n h(\boldsymbol{\xi})| \lesssim \langle \boldsymbol{\xi} \rangle^{2m-|n|}, \quad n \in \mathbb{N}_0^d, \quad \text{for all } \boldsymbol{\xi} \in \mathbb{R}^d.$$

Furthermore, we assume that there exists a real-valued homogeneous function  $h_0$  of order  $2m$  (that is,  $h_0(t\boldsymbol{\xi}) = t^{2m}h_0(\boldsymbol{\xi})$  for all  $\boldsymbol{\xi} \in \mathbb{R}^d$  and all  $t > 0$ ), such that

$$(5.2) \quad |\boldsymbol{\xi}|^{-2m} |h(\boldsymbol{\xi}) - h_0(\boldsymbol{\xi})| \rightarrow 0 \quad \text{as } |\boldsymbol{\xi}| \rightarrow \infty.$$

We suppose that  $h_0$  is non-degenerate in the sense that

$$(5.3) \quad 2\nu := \min_{|\boldsymbol{\xi}|=1} h_0(\boldsymbol{\xi}) > 0.$$

The condition (5.2) implies that  $h$  is bounded from below.

**5.2. The model symbol.** Starting from two continuous functions  $\phi_T$  and  $\omega_T$  of the temperature  $T \in [1, \infty)$  we consider the symbol  $p_T$  defined as

$$(5.4) \quad p_T(\boldsymbol{\xi}) := \frac{1}{\phi_T + \omega_T \exp(\frac{h(\boldsymbol{\xi})}{T})}, \quad \boldsymbol{\xi} \in \mathbb{R}^d.$$

We assume that

$$(5.5) \quad \phi_T \rightarrow \phi_\infty \geq 0, \quad \text{and} \quad \omega_T \rightarrow \omega_\infty > 0 \quad \text{as } T \rightarrow \infty.$$

We also introduce the ‘‘limit’’ symbol  $p_\infty$  defined naturally as

$$(5.6) \quad p_\infty(\boldsymbol{\xi}) := \frac{1}{\phi_\infty + \omega_\infty \exp(h_0(\boldsymbol{\xi}))}.$$

Throughout this section we assume that the function  $f$  satisfies Condition 2.2 with some  $\delta > 0$  and that the region  $\Lambda$  satisfies Condition 2.1. Since  $\Lambda$  is always fixed, we omit it from the notation and write  $D_\alpha(p_T; f)$  and  $\mathcal{B}_d(p_T; f)$ .

**Theorem 5.1.** *Let  $p_T$  be the symbol defined in (5.4). Then*

$$(5.7) \quad \lim \alpha^{-(d-1)} T^{-\frac{d-1}{2m}} \text{tr} D_\alpha(p_T; f) = \mathcal{B}_d(p_\infty; f),$$

as  $\alpha T^{1/2m} \rightarrow \infty$  and  $T \rightarrow \infty$ .

We also consider the operator  $D_\alpha(\lambda_T p_T; f)$  with the symbol  $\lambda_T p_T$ , where  $\lambda_T$  is a positive function of  $T$  that tends to zero as  $T \rightarrow \infty$ . In this case we assume that the function  $f$  has only one singular point,  $z = 0$ .

**Theorem 5.2.** *Let  $f_0$  be the function  $f_0(t) = M|t|^\delta$ ,  $t \in \mathbb{R}$  with some (complex) number  $M$  and some  $\delta > 0$ . Let  $\eta(t) = -t \ln |t|$ . Assume that  $\alpha T^{1/2m} \rightarrow \infty$  and  $T \rightarrow \infty$ . Then*

(1) *If  $f \in C^2(\mathbb{R} \setminus \{0\}) \cap C(\mathbb{R})$  satisfies (2.9), then*

$$(5.8) \quad \lim \alpha^{-(d-1)} T^{-\frac{d-1}{2m}} \lambda_T^{-\delta} \text{tr} D_\alpha(\lambda_T p_T; f) = \mathcal{B}_d(p_\infty; f_0).$$

(2) *If  $f \in C^2(\mathbb{R} \setminus \{0\}) \cap C(\mathbb{R})$  satisfies (2.12), then*

$$(5.9) \quad \lim \alpha^{-(d-1)} T^{-\frac{d-1}{2m}} \lambda_T^{-1} \text{tr} D_\alpha(\lambda_T p_T; f) = \mathcal{B}_d(p_\infty; \eta).$$

To prove these theorems we compare the operators  $D_\alpha$  for two different symbols defined as follows. Let the function  $\zeta \in C^\infty(\mathbb{R}^d)$  be such that  $\zeta(\boldsymbol{\xi}) = 0$  if  $|\boldsymbol{\xi}| \leq 1$  and  $\zeta(\boldsymbol{\xi}) = 1$  if  $|\boldsymbol{\xi}| \geq 2$ . We introduce the scaled functions

$$(5.10) \quad \zeta_T(\boldsymbol{\xi}) := \zeta(\boldsymbol{\xi}T^{-\frac{1}{2m}}), \quad \tilde{\zeta}_T(\boldsymbol{\xi}) := \zeta_T(\boldsymbol{\xi}/2),$$

so that  $\zeta_T \tilde{\zeta}_T = \tilde{\zeta}_T$ . Let us fix a number  $r \in (0, 1]$  and define the operators

$$A := \text{Op}_\alpha(p_T), \quad B := \text{Op}_\alpha(\zeta_{rT} p_T), \quad P := \chi_\Lambda, \quad J := \text{Op}_\alpha(\tilde{\zeta}_{rT}).$$

The norms of these operators are bounded uniformly in  $T$  and  $(A - B)J = 0$ . Thus we can use Corollary 4.3.

**Lemma 5.3.** *Let us assume that  $T \gtrsim 1$  and  $\alpha(rT)^{\frac{1}{2m}} \gtrsim 1$ . Then with the above notations we have*

$$(5.11) \quad \|D_\alpha(p_T; f) - D_\alpha(\zeta_{rT} p_T; f)\|_1 \lesssim \alpha^{d-1} (rT)^{\frac{d-1}{2m}},$$

with an implicit constant independent of  $\alpha, T$ , and  $r$ .

*Proof.* Let us estimate the right-hand side of (4.9) and let us start with a bound for  $\|[J, P]\|_q$ ,  $q \leq 1$ . Since  $[J, P] = -[\mathbb{1} - J, P]$ , we use Proposition 2.10 with  $a = \mathbb{1} - \tilde{\zeta}_{rT}$ . This symbol satisfies (2.14) with scale and amplitude functions

$$\tau(\boldsymbol{\xi}) = (rT)^{\frac{1}{2m}}, \quad v(\boldsymbol{\xi}) = \langle \boldsymbol{\xi}(rT)^{-\frac{1}{2m}} \rangle^{-\beta}, \quad \boldsymbol{\xi} \in \mathbb{R}^d,$$

with an arbitrary  $\beta > 0$ . We assume that  $\beta q > d$ . The conditions (2.15) and (2.16) are trivially satisfied, and hence Proposition 2.10 is applicable. Estimate the integral on the right-hand side of (2.18):

$$\int_{\mathbb{R}^d} \frac{v(\boldsymbol{\xi})^q}{\tau(\boldsymbol{\xi})} d\boldsymbol{\xi} = (rT)^{-\frac{1}{2m}} \int_{\mathbb{R}^d} \langle \boldsymbol{\xi}(rT)^{-\frac{1}{2m}} \rangle^{-\beta q} d\boldsymbol{\xi} \lesssim (rT)^{\frac{d-1}{2m}}.$$

Thus, under the existing assumptions on  $\alpha, T$ , and  $r$  the condition (2.17) is satisfied, and hence, by (2.18), we have

$$\|[J, P]\|_q^q = \|[\text{Op}_\alpha(a), \chi_\Lambda]\|_q^q \lesssim \alpha^{d-1} (rT)^{\frac{d-1}{2m}}.$$

Estimating  $\|[(\mathbb{1} - J)A, P]\|_q$  and  $\|[(\mathbb{1} - J)B, P]\|_q$  is somewhat trickier. For the first commutator we use Proposition 3.2 with  $a = (\mathbb{1} - \tilde{\zeta}_{rT})p_T$ . First estimate the derivatives of  $e^{h(\boldsymbol{\xi})/T}$  for  $|\boldsymbol{\xi}| \leq 4(rT)^{\frac{1}{2m}}$  using (5.1):

$$|\partial_{\boldsymbol{\xi}}^k e^{\frac{h(\boldsymbol{\xi})}{T}}| \lesssim \langle \boldsymbol{\xi} \rangle^{-|k|} e^{\frac{h(\boldsymbol{\xi})}{T}} \lesssim \langle \boldsymbol{\xi} \rangle^{-|k|}, \quad k \in \mathbb{N}_0^d.$$

Furthermore,

$$|\partial_{\boldsymbol{\xi}}^k (1 - \tilde{\zeta}_r(\boldsymbol{\xi}))| \lesssim (rT)^{-\frac{|k|}{2m}} \chi_{\{|\boldsymbol{\xi}| \leq 4(rT)^{\frac{1}{2m}}\}}(\boldsymbol{\xi}) \lesssim \langle \boldsymbol{\xi} \rangle^{-|k|} \chi_{\{|\boldsymbol{\xi}| \leq 4(rT)^{\frac{1}{2m}}\}}(\boldsymbol{\xi}).$$

Therefore, we obtain from (5.4) that

$$|\partial_{\boldsymbol{\xi}}^k a(\boldsymbol{\xi})| \lesssim \langle \boldsymbol{\xi} \rangle^{-|k|} \langle \boldsymbol{\xi}(rT)^{-\frac{1}{2m}} \rangle^{-\beta}.$$



with an arbitrary  $\beta > d/q$ . Consequently, the symbol  $a$  satisfies (2.14) with the scale and amplitude

$$\tau(\boldsymbol{\xi}) = \frac{1}{2}\langle \boldsymbol{\xi} \rangle, \quad v(\boldsymbol{\xi}) = \langle \boldsymbol{\xi}(rT)^{-\frac{1}{2m}} \rangle^{-\beta}, \quad \boldsymbol{\xi} \in \mathbb{R}^d.$$

Again the conditions (2.15), (2.16), and (2.17) are satisfied, and we can use Proposition 2.10, which produces the bound

$$\int_{\mathbb{R}^d} \frac{v(\boldsymbol{\xi})^q}{\tau(\boldsymbol{\xi})} d\boldsymbol{\xi} \leq 2 \int_{\mathbb{R}^d} |\boldsymbol{\xi}|^{-1} \langle \boldsymbol{\xi}(rT)^{-\frac{1}{2m}} \rangle^{-\beta q} d\boldsymbol{\xi} \lesssim (rT)^{\frac{d-1}{2m}}.$$

Thus by (2.18),

$$\|[(\mathbb{1} - J)A, P]\|_q^q = \|[\text{Op}_\alpha(a), \chi_\Lambda]\|_q^q \lesssim \alpha^{d-1} (rT)^{\frac{d-1}{2m}}.$$

The bound for the commutator  $[(\mathbb{1} - J)B, P]$  is proved in the same way.

Substituting the above bounds into (4.9), we get the required estimate (5.11).  $\square$

A useful consequence of this fact is the following continuity statement:

**Corollary 5.4.** *With the function  $\zeta_r$  defined in (5.10) and the symbol  $p_\infty$  defined in (5.6) we have*

$$(5.12) \quad \lim_{r \rightarrow 0} \mathcal{B}_d(\zeta_r p_\infty; f) = \mathcal{B}_d(p_\infty; f).$$

*Proof.* We apply Lemma 5.3 with  $h = h_0$ ,  $T = 1$ , and the constant functions  $\omega \equiv \omega_\infty$  and  $\phi \equiv \phi_\infty$ . Then, for  $\alpha r \gtrsim 1$ ,

$$\|D_\alpha(p_\infty; f) - D_\alpha(\zeta_r p_\infty; f)\|_1 \lesssim \alpha^{d-1} r^{\frac{d-1}{2m}}.$$

Therefore,

$$|\alpha^{1-d} \text{tr} D_\alpha(p_\infty; f) - \alpha^{1-d} \text{tr} D_\alpha(\zeta_r p_\infty; f)| \leq r^{\frac{d-1}{2m}}.$$

Now, we use Proposition 2.6 to obtain the estimate

$$|\mathcal{B}_d(p_\infty; f) - \mathcal{B}_d(\zeta_r p_\infty; f)| \lesssim r^{\frac{d-1}{2m}}.$$

This leads to (5.12), as required.  $\square$

We already have a result on the continuity of the asymptotic coefficient, Proposition 2.5. However, this proposition is not applicable to the symbol  $\zeta_r p_T$ , since its derivatives are not bounded uniformly in  $r > 0$ . This explains the necessity of Corollary 5.4.

The next lemma provides the same asymptotics as in Theorems 5.1 and 5.2, but for the truncated symbol  $p_T$ . Recall that  $\lambda_T$  is a function such that  $\lambda_T \rightarrow 0$  as  $T \rightarrow \infty$ .

**Lemma 5.5.** *Assume that  $r \in (0, 1]$  is fixed and that  $\alpha T^{\frac{1}{2m}} \rightarrow \infty, T \rightarrow \infty$ . Then*

$$(5.13) \quad \lim \alpha^{-(d-1)} T^{-\frac{d-1}{2m}} \text{tr} D_\alpha(\zeta_{rT} p_T; f) = \mathcal{B}_d(\zeta_r p_\infty; f).$$

*If  $f$  satisfies (2.9), then*

$$(5.14) \quad \lim \alpha^{-(d-1)} T^{-\frac{d-1}{2m}} \lambda_T^{-\delta} \text{tr} D_\alpha(\lambda_T \zeta_{rT} p_T; f) = \mathcal{B}_d(\zeta_r p_\infty; f_0).$$

*If  $f$  satisfies (2.12), then*

$$(5.15) \quad \lim \alpha^{-(d-1)} T^{-\frac{d-1}{2m}} \lambda_T^{-1} \text{tr} D_\alpha(\lambda_T \zeta_{rT} p_T; f) = \mathcal{B}_d(\zeta_r p_\infty; \eta).$$

*Proof.* By a straightforward change of variables in the definition (1.1), we obtain

$$\text{Op}_\alpha(\zeta_{rT} p_T) = \text{Op}_L(b_T) \quad \text{and} \quad D_\alpha(\zeta_{rT} p_T; f) = D_L(b_T; f), \quad L := \alpha T^{\frac{1}{2m}},$$

where for  $\xi \in \mathbb{R}^d$

$$b_T(\xi) := \zeta_r(\xi) p_T(T^{\frac{1}{2m}} \xi).$$

Thanks to the condition (5.2), for all  $\xi \neq 0$ , we have

$$(5.16) \quad T^{-1} h(T^{\frac{1}{2m}} \xi) \rightarrow h_0(\xi) \quad \text{and hence} \quad b_T(\xi) \rightarrow \zeta_r(\xi) p_\infty(\xi), \quad \text{as } T \rightarrow \infty.$$

Assuming that  $T \gtrsim 1$ , an elementary calculation using (5.1) leads to the bounds

$$(5.17) \quad |\partial_\xi^n b_T(\xi)| + |\partial_\xi^n (\zeta_r(\xi) p_\infty(\xi))| \lesssim e^{-\nu|\xi|^{2m}}, \quad n \in \mathbb{N}_0^d, \xi \in \mathbb{R}^d,$$

with implicit constants that depend on the parameter  $r$ . We recall that  $\nu$  is defined in (5.3). Since  $b_T$  satisfies (5.17) uniformly in  $T \gtrsim 1$ , we obtain by Corollary 2.7 that

$$\lim_{L \rightarrow \infty, T \rightarrow \infty} L^{1-d} D_L(b_T; f) = \mathcal{B}_d(\zeta_r p_\infty; f).$$

By the above change of variables, this leads to (5.13).

Formulas (5.14) and (5.15) follow along the same lines from Propositions 2.8 and 2.9.  $\square$

*Proof of Theorem 5.1.* Estimate:

$$\begin{aligned} |\alpha^{-(d-1)} T^{-\frac{d-1}{2m}} \text{tr} D_\alpha(p_T; f) - \mathcal{B}_d(p_\infty; f)| &\leq (\alpha T^{\frac{1}{2m}})^{1-d} \|D_\alpha(p_T; f) - D_\alpha(\zeta_{rT} p_T; f)\|_1 \\ &\quad + |(\alpha T^{\frac{1}{2m}})^{1-d} \text{tr} D_\alpha(\zeta_{rT} p_T; f) - \mathcal{B}_d(\zeta_r p_\infty; f)| \\ &\quad + |\mathcal{B}_d(\zeta_r p_\infty; f) - \mathcal{B}_d(p_\infty; f)|. \end{aligned}$$

By (5.11), (5.13) we obtain

$$\limsup |\alpha^{-(d-1)} T^{-\frac{d-1}{2m}} \text{tr} D_\alpha(p_T; f) - \mathcal{B}_d(p_\infty; f)| \leq r^{\frac{d-1}{2m}} + |\mathcal{B}_d(\zeta_r p_\infty; f) - \mathcal{B}_d(p_\infty; f)|,$$

where  $\limsup$  is taken as  $\alpha T^{\frac{1}{2m}} \rightarrow \infty$ ,  $T \rightarrow \infty$ . Taking  $r \rightarrow 0$  and using (5.12) we obtain (5.7).  $\square$

*Proof of Theorem 5.2.* We recall that the only singular point of the function  $f$  is  $z = 0$ . We assume that  $f$  satisfies (2.9), so that for all  $t \neq 0$ ,

$$(5.18) \quad |f^{(n)}(t)| \lesssim |t|^{\delta-n}, \quad n = 0, 1, 2.$$

Consequently, the function  $\tilde{f}_T(t) := \lambda_T^{-\delta} f(\lambda_T t)$ ,  $t \in \mathbb{R}$ , satisfies the same inequalities with the same constants. Now we can apply the argument in the previous proof to the operator

$$D_\alpha(p_T; \tilde{f}_T) = \lambda_T^{-\delta} D_\alpha(\lambda_T p_T; f).$$

More precisely, we estimate

$$\begin{aligned} |\alpha^{-(d-1)} T^{-\frac{d-1}{2m}} \lambda_T^{-\delta} \text{tr} D_\alpha(\lambda_T p_T; f) - \mathcal{B}_d(p_\infty; f_0)| &\leq (\alpha T^{\frac{1}{2m}})^{1-d} \|D_\alpha(p_T; \tilde{f}_T) - D_\alpha(\zeta_{rT} p_T; \tilde{f}_T)\|_1 \\ &\quad + |(\alpha T^{\frac{1}{2m}})^{1-d} \lambda_T^{-\delta} \text{tr} D_\alpha(\lambda_T \zeta_{rT} p_T; f) - \mathcal{B}_d(\zeta_r p_\infty; f_0)| \\ (5.19) \quad &\quad + |\mathcal{B}_d(\zeta_r p_\infty; f_0) - \mathcal{B}_d(p_\infty; f_0)|. \end{aligned}$$

By (5.11), (5.14) we obtain

$$\begin{aligned} \limsup \left| \alpha^{-(d-1)} T^{-\frac{d-1}{2m}} \lambda^{-\delta} \operatorname{tr} D_\alpha(\lambda p_T; f) - \mathcal{B}_d(p_\infty; f_0) \right| \\ \leq r^{\frac{d-1}{2m}} + |\mathcal{B}_d(\zeta_r p_\infty; f_0) - \mathcal{B}_d(p_\infty; f_0)|, \end{aligned}$$

where  $\limsup$  is taken as  $\alpha T^{\frac{1}{2m}} \rightarrow \infty$ ,  $T \rightarrow \infty$ . Taking  $r \rightarrow 0$  and using (5.12) we obtain (5.8).

Now we assume that  $f$  satisfies (2.12). We use (5.19) with  $\delta = 1$  and  $f_0$  replaced by  $\eta$ . Then

$$\begin{aligned} \left| \alpha^{-(d-1)} T^{-\frac{d-1}{2m}} \lambda_T^{-1} \operatorname{tr} D_\alpha(\lambda_T p_T; f) - \mathcal{B}_d(p_\infty; \eta) \right| \\ \leq (\alpha T^{\frac{1}{2m}})^{1-d} \|D_\alpha(p_T; \tilde{f}_T) - D_\alpha(\zeta_{rT} p_T; \tilde{f}_T)\|_1 \\ + \left| (\alpha T^{\frac{1}{2m}})^{1-d} \lambda_T^{-1} \operatorname{tr} D_\alpha(\lambda_T \zeta_{rT} p_T; f) - \mathcal{B}_d(\zeta_r p_\infty; \eta) \right| \\ + |\mathcal{B}_d(\zeta_r p_\infty; \eta) - \mathcal{B}_d(p_\infty; \eta)|. \end{aligned}$$

As before, the last term on the right-hand side tends to zero due to (5.12). The second term vanishes as  $\alpha T^{\frac{1}{2m}} \rightarrow \infty$ ,  $T \rightarrow \infty$  due to (5.15). To estimate the first term we represent  $f = \eta + g$ , so that  $g$  satisfies (5.18) with  $\delta = 1$ . Therefore,

$$\begin{aligned} D_\alpha(p_T; \tilde{f}_T) - D_\alpha(\zeta_{rT} p_T; \tilde{f}_T) &= [D_\alpha(p_T; \tilde{\eta}_T) - D_\alpha(\zeta_{rT} p_T; \tilde{\eta}_T)] \\ &\quad + [D_\alpha(p_T; \tilde{g}_T) - D_\alpha(\zeta_{rT} p_T; \tilde{g}_T)], \end{aligned}$$

where  $\tilde{\eta}_T(t) = \lambda_T^{-1} \eta(\lambda_T t)$  and  $\tilde{g}_T(t) = \lambda_T^{-1} g(\lambda_T t)$ . As in the previous calculation, the second term is estimated with the help of (5.11) by  $r^{\frac{d-1}{2m}}$ . Since  $\tilde{\eta}_T(t) = \eta(t) - t \ln(\lambda_T)$  and the functional (1.3) equals zero for linear functions, we have

$$D_\alpha(p_T; \tilde{\eta}_T) - D_\alpha(\zeta_{rT} p_T; \tilde{\eta}_T) = D_\alpha(p_T; \eta) - D_\alpha(\zeta_{rT} p_T; \eta).$$

The function  $\eta$  satisfies (5.18) for all  $\delta < 1$ , and hence by (5.11) the above difference is again estimated by  $r^{\frac{d-1}{2m}}$ . This entails (5.9). Now the proof of Theorem 5.2 is complete.  $\square$

## 6. MAIN RESULTS ON THE HIGH-TEMPERATURE ASYMPTOTICS

In this section we adapt Theorems 5.1 and 5.2 to two different asymptotic regimes when the temperature becomes large. In the first regime we fix the chemical potential  $\mu$ , which is reasonable since we work in the grand-canonical formalism [3, 7]. In the second regime we fix the particle density. This is often the physically more relevant case and this will be discussed in more detail after presenting the results. As above,  $d \geq 2$ .

**6.1. Case of a fixed chemical potential  $\mu$ .** In this case the symbol is given by the Fermi function  $a_{T,\mu}$  defined in (1.5). Clearly, this symbol coincides with  $p_T$  for  $\phi_T := 1$  and  $\omega_T := \exp(-\mu/T)$ , so that the limit symbol is given by

$$a_\infty(\boldsymbol{\xi}) := p_\infty(\boldsymbol{\xi}) = \frac{1}{1 + \exp(h_0(\boldsymbol{\xi}))}, \quad \boldsymbol{\xi} \in \mathbb{R}^d.$$

For the function  $f$  we take  $\eta_\gamma$  which satisfies Condition 2.2 with  $X = \{0, 1\}$ , for an arbitrary  $\delta < \min\{1, \gamma\}$ . Thus, remembering definition (1.9), Remark 2.4(1) and applying Theorem 5.1 we obtain the following result.

**Theorem 6.1.** *Let  $\Lambda$  be a region satisfying Condition 2.1. Then*

$$\lim \alpha^{-(d-1)} T^{-\frac{d-1}{2m}} \mathbb{H}_\gamma(T, \mu; \alpha\Lambda) = 2\mathcal{B}_d(a_\infty, \partial\Lambda; \eta_\gamma),$$

as  $\alpha T^{\frac{1}{2m}} \rightarrow \infty$  and  $T \rightarrow \infty$ .

**6.2. Case of a fixed particle density  $\rho$ .** Here we have to consider the chemical potential  $\mu = \mu(T)$  as a function of the temperature  $T$  such that the mean particle density  $\varrho(T, \mu)$  (see (1.12) for the definition) is kept asymptotically constant as  $T$  becomes large. As shown in the appendix, to ensure that

$$(6.1) \quad \varrho(T, \mu(T)) \rightarrow \rho \quad \text{as } T \rightarrow \infty,$$

with a constant  $\rho > 0$ , one assumes that the function  $z_\rho(T) = \exp(-\mu(T)/T)$  satisfies (A.5). In this case we use for the Fermi symbol the notation (A.6):

$$F_{T,\rho}(\boldsymbol{\xi}) = \frac{1}{1 + z_\rho(T) \exp\left(\frac{h(\boldsymbol{\xi})}{T}\right)}, \quad \boldsymbol{\xi} \in \mathbb{R}^d.$$

This implies that

$$(6.2) \quad F_{T,\rho} = \lambda_T p_T, \quad \lambda_T = \rho \varkappa^{-1} T^{-\frac{d}{2m}}, \quad \text{where}$$

$$p_T \quad \text{is given by (5.4) with } \phi_T = \lambda_T, \quad \omega_T = \rho \varkappa^{-1} z_\rho(T) T^{-\frac{d}{2m}},$$

and the constant  $\varkappa$  is defined in (A.4). Clearly,  $\phi_T \rightarrow 0$ , and according to (A.5)  $\omega_T \rightarrow 1$  as  $T \rightarrow \infty$ . Hence the limit symbol is defined as

$$(6.3) \quad F_\infty(\boldsymbol{\xi}) := p_\infty(\boldsymbol{\xi}) = e^{-h_0(\boldsymbol{\xi})}, \quad \boldsymbol{\xi} \in \mathbb{R}^d.$$

To study the symbol  $F_{T,\rho}$  we use Theorem 5.2 with  $f = \eta_\gamma$ . For a start we need to understand the small  $t$  behavior of  $\eta_\gamma(t)$ . This behavior depends on the value of  $\gamma$  and we distinguish five different regimes. To state the result in a unified way for all  $\gamma$ , we define the functions  $f_\gamma, \eta_\gamma^{\text{eff}}$ , and the parameter  $\delta = \delta_\gamma$  in Table 1. We recall that the function  $\eta$  is defined in (2.11).

	$\delta$	$f_\gamma(t)$	$\eta_\gamma^{\text{eff}}(t)$
$0 < \gamma < 1$	$\gamma$	$\eta_\gamma(t)$	$\frac{1}{1-\gamma} t^\gamma$
$\gamma = 1$	1	$\eta_1(t) - t$	$\eta(t)$
$1 < \gamma < 2$	$\gamma$	$\eta_\gamma(t) - \frac{\gamma}{\gamma-1} t$	$\frac{1}{1-\gamma} t^\gamma$
$\gamma = 2$	3	$\eta_2(t) - 2t$	$-\frac{4}{3} t^3$
$\gamma > 2$	2	$\eta_\gamma(t) - \frac{\gamma}{\gamma-1} t$	$\frac{\gamma}{2(\gamma-1)} t^2$

TABLE 1. Functions  $f_\gamma$  and  $\eta_\gamma^{\text{eff}}$

The next lemma shows that the function  $\eta_\gamma^{\text{eff}}$  describes the effective asymptotic contribution of  $f_\gamma$  as  $t \rightarrow 0$ .

**Lemma 6.2.** *Let the parameter  $\delta = \delta_\gamma$  and the functions  $f_\gamma$  and  $\eta_\gamma^{\text{eff}}$  on the interval  $(0, 1)$  be as defined in Table 1. Then the following asymptotic relation is satisfied:*

$$(6.4) \quad \lim_{t \rightarrow 0} t^{n-\delta} \frac{d^n}{dt^n} (f_\gamma(t) - \eta_\gamma^{\text{eff}}(t)) = 0, \quad n = 0, 1, 2.$$

*Proof.* If we assume that  $\gamma \neq 1$  and expand  $\eta_\gamma$  at  $t = 0$ , then we obtain

$$(6.5) \quad (1 - \gamma)\eta_\gamma(t) = t^\gamma - \gamma t - \frac{\gamma}{2}t^2 + O(t^3) + O(t^{2\gamma}) + O(t^{1+\gamma}).$$

The notation  $m(t) = O(t^\alpha)$  means that  $|m^{(k)}(t)| \lesssim t^{\alpha-k}$ , for all  $k = 0, 1, \dots$ . This expansion proves the statement of the lemma for all  $\gamma \neq 1$ ,  $\gamma \neq 2$ .

If  $\gamma = 2$ , then (6.5) is insufficient since the terms with  $t^2$  cancel out. Instead we use the expansion  $\eta_\gamma(t) = 2t - \frac{4}{3}t^3 + O(t^4)$ , which gives the required result.

For  $\gamma = 1$  we have  $\eta_1(t) = \eta(t) + t + O(t^2)$ . This leads to (6.4) again.  $\square$

Now we are in a position to state and prove our second main result for the EE.

**Theorem 6.3.** *Suppose that the mean particle density satisfies (6.1) with some  $\rho > 0$ , so that the Fermi symbol is given by (A.6). Let  $\Lambda$  be a region satisfying Condition 2.1. Let  $\eta_\gamma^{\text{eff}}$  and  $\delta = \delta_\gamma$  be as defined in Table 1. Then*

$$(6.6) \quad \lim \alpha^{1-d} T^{\frac{d(\delta-1)+1}{2m}} H_\gamma(T, \mu; \alpha\Lambda) = 2\rho^\delta \varkappa^{-\delta} \mathcal{B}_d(F_\infty, \partial\Lambda; \eta_\gamma^{\text{eff}}),$$

as  $\alpha T^{\frac{1}{2m}} \rightarrow \infty, T \rightarrow \infty$ .

*Proof.* We replace the symbol  $F_{T,\rho}$  with  $\lambda_T p_T$  as specified in (6.2). Furthermore, since the functional (1.3) equals zero for linear functions  $f$ , we have

$$D_\alpha(\lambda_T p_T, \Lambda; \eta_\gamma) = D_\alpha(\lambda_T p_T, \Lambda; f_\gamma).$$

As the function  $f_\gamma$  satisfies the condition (6.4) we can use Theorem 5.2 which gives

$$\lim \alpha^{1-d} T^{-\frac{d-1}{2m}} \lambda_T^{-\delta} \text{tr} D_\alpha(\lambda_T p_T, \Lambda; f_\gamma) = \mathcal{B}_d(p_\infty, \partial\Lambda; \eta_\gamma^{\text{eff}}),$$

as  $\alpha T^{\frac{1}{2m}} \rightarrow \infty, T \rightarrow \infty$ . The same formula holds for the region  $\mathbb{R}^d \setminus \bar{\Lambda}$ . Since  $\lambda_T = \rho \varkappa^{-1} T^{-\frac{d}{2m}}$  and  $p_\infty = F_\infty$  (see (6.3)), the required formula (6.6) follows due to the definition (1.9) and Remark 2.4(1).  $\square$

### 6.3. Concluding remarks.

- (1) We have proved positivity of the Rényi EE if  $\gamma \leq 1$  (see Theorem 3.5) and positivity of the asymptotic coefficient  $\mathcal{B}_d$  if  $\gamma \leq 2$  (see Remark 2.4(3)). It is an open question whether positivity of EE extends to all  $\gamma \leq 2$  for all temperatures.
- (2) In the fixed particle density case, for  $\gamma \leq 2$ , the test function  $\eta_\gamma^{\text{eff}}$  in Table 1 is (strictly) concave so that the coefficient  $\mathcal{B}_d(F_\infty, \partial\Lambda; \eta_\gamma^{\text{eff}}) > 0$ . On the other hand, if  $\gamma > 2$ , then  $\eta_\gamma^{\text{eff}}$  is convex so that this coefficient is (strictly) *negative*. This unexpected change of sign suggests that the definition of EE that we use is somewhat unphysical for  $\gamma > 2$ .

As a side remark, the coefficient  $\mathcal{B}_d(F_\infty, \partial\Lambda; \eta_\gamma^{\text{eff}})$ ,  $\gamma > 2$ , can be computed rather explicitly. Indeed, it is easy to see that  $U(u, v; t^2) = -(u - v)^2$ , so that for  $\gamma > 2$  we have

$$U(u, v; \eta_\gamma^{\text{eff}}) = -\frac{\gamma}{2(\gamma - 1)}(u - v)^2.$$

This observation and the use of Parseval's identity, as in the proof of Proposition 1 in [28], allow one to find the coefficient (2.3) and hence (2.4) explicitly. In particular, for the Hamiltonian  $h_0(\boldsymbol{\xi}) = |\boldsymbol{\xi}|^2/2$  (which includes the ideal Fermi gas) we obtain the formula

$$\mathcal{B}_d(F_\infty, \partial\Lambda; \eta_\gamma^{\text{eff}}) = -\frac{\gamma}{\gamma - 1} 2^{-d-3} \pi^{-(d+1)/2} |\partial\Lambda|.$$

- (3) For  $\gamma \in (0, 2)$  the test function  $\eta_\gamma^{\text{eff}}$  is exactly the classical entropy function of the Maxwell–Boltzmann gas. This confirms our expectations stated in the Introduction. This conclusion does not hold for  $\gamma \geq 2$ .
- (4) In both high temperature regimes (fixed chemical potential or fixed particle density) the quasi-classical parameter  $\alpha$  may be fixed to, say,  $\alpha = 1$ . Thus the truncating set  $\alpha\Lambda$  is fixed and only the temperature tends to infinity.
- (5) Using the relation (1.10) and Theorem 5.1 one can also derive appropriate asymptotic formulas for the local entropy (1.8). For example, assuming that  $\mu$  is fixed, as in Theorem 6.1 we easily obtain the relation

$$\lim \alpha^{-(d-1)} T^{-\frac{d-1}{2m}} \left[ \mathcal{S}_\gamma(T, \mu; \alpha\Lambda) - \alpha^d s_\gamma(T, \mu) |\Lambda| \right] = \mathcal{B}_d(a_\infty, \partial\Lambda; \eta_\gamma),$$

as  $T \rightarrow \infty$ ,  $\alpha T^{\frac{1}{2m}} \rightarrow \infty$ . However, in order to obtain from this formula a proper asymptotic expansion for the local entropy, one needs to find an expansion for the entropy density  $s_\gamma(T, \mu)$  as  $T \rightarrow \infty$ . A convenient starting point for the derivation of such an expansion could be, for example, the representation (10.8) in [17].

A similar formula can be written in the case when the particle density is fixed. It is worth noting that the negativity of the asymptotic coefficient  $\mathcal{B}_d(F_\infty, \partial\Lambda; \eta_\gamma^{\text{eff}})$  for  $\gamma > 2$  implies that at sufficiently high temperature the local entropy (1.8) satisfies the bound  $\mathcal{S}_\gamma(T, \mu(T); \Lambda) < s_\gamma(T, \mu(T)) |\Lambda|$ , instead of (3.4) which holds for  $\gamma \leq 2$  and all  $T > 0$ .

#### APPENDIX A. THE FERMI SYMBOL AT HIGH TEMPERATURE $T$ AND FIXED PARTICLE DENSITY $\rho$

Let the Hamilton function  $h$  be as in Subsection 5.1. Recall the formula (1.12) for the mean particle density  $\varrho = \varrho(T, \mu)$ :

$$(A.1) \quad \varrho(T, \mu) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} a_{T, \mu}(\boldsymbol{\xi}) \, d\boldsymbol{\xi} = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} \frac{1}{1 + \exp(-\mu/T) \exp(h(\boldsymbol{\xi})/T)} \, d\boldsymbol{\xi}.$$

We consider the chemical potential as a function of  $T$ , that is,  $\mu = \mu(T)$ . Our aim is to find conditions on  $z(T) := \exp(-\mu(T)/T)$  that ensure that

$$(A.2) \quad \varrho(T, \mu(T)) \rightarrow \rho \quad \text{as } T \rightarrow \infty,$$

where  $\rho$  on the right-hand side is a given strictly positive constant. To reflect the dependence on  $\rho$  we write  $z(T) = z_\rho(T)$ , and note in passing that the quantity  $\exp(\mu/T)$  is usually called the fugacity so that  $z_\rho$  is its reciprocal. A change of variables in the integral (A.1) gives

$$(A.3) \quad \varrho(T, \mu) = \frac{T^{\frac{d}{2m}}}{(2\pi)^d} \int_{\mathbb{R}^d} \frac{1}{1 + z_\rho(T) \exp(T^{-1}h(\boldsymbol{\xi}T^{\frac{1}{2m}}))} d\boldsymbol{\xi}.$$

Since  $T^{-1}h(T^{\frac{1}{2m}}\boldsymbol{\xi}) \rightarrow h_0(\boldsymbol{\xi})$  as  $T \rightarrow \infty$ , for each  $\boldsymbol{\xi} \neq 0$ , it is clear that the condition (A.2) requires that  $z_\rho(T) \rightarrow \infty$ . Consequently,

$$(A.4) \quad \varrho(T, \mu) = \frac{T^{\frac{d}{2m}}}{z_\rho(T)} \varkappa(1 + o(1)), \quad \varkappa := \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{-h_0(\boldsymbol{\xi})} d\boldsymbol{\xi}.$$

Thus according to (A.2),

$$(A.5) \quad z_\rho(T) = T^{\frac{d}{2m}} \varkappa \rho^{-1} (1 + o(1)) \quad \text{as } T \rightarrow \infty.$$

To reflect the dependence on  $\rho$  (instead of  $\mu$ ) we use for the Fermi symbol (1.5) the notation

$$(A.6) \quad F_{T,\rho}(\boldsymbol{\xi}) := \frac{1}{1 + z_\rho(T) \exp\left(\frac{h(\boldsymbol{\xi})}{T}\right)}, \quad \boldsymbol{\xi} \in \mathbb{R}^d.$$

As mentioned in the Introduction, the high temperature limit under the condition (A.2) corresponds to the classical Maxwell–Boltzmann gas limit. This fact can be conveniently restated in terms of the function  $z_\rho$  as follows. Because of (5.2), the integrated density of states  $\mathcal{N}(T)$  defined in (1.13), for large  $T$  satisfies the relation  $\mathcal{N}(T) \asymp T^{\frac{d}{2m}}$ , so (A.5) implies that  $z_\rho(T)^{-1} \asymp \rho \mathcal{N}(T)^{-1} \rightarrow 0$ .

## REFERENCES

- [1] L. Amico, R. Fazio, A. Osterloh, and V. Vedral, *Entanglement in many-body systems*. Rev. Mod. Phys. **80**: 517–576, 2008.
- [2] T. Ando, *Concavity of certain maps on positive definite matrices and applications to Hadamard products*. Linear Algebra Appl. **26**: 203–241, 1979.
- [3] R. Balian, *From Microphysics to Macrophysics II*. Springer, Berlin, 1992.
- [4] J. Benda and S. Sherman, *Monotone and convex operator functions*. Trans. Amer. Math. Soc. **79**: 58–71, 1955.
- [5] F. A. Berezin, *Convex operator functions*. Math. USSR Sb. **17**: 269–277, 1972.
- [6] M. S. Birman and M. Z. Solomjak, *Spectral Theory of Self-Adjoint Operators in Hilbert Space*. Reidel, Dordrecht, 1987. [Translated from the 1980 Russian original by S. Khrushchëv and V. Peller]
- [7] O. Bratteli and D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics 2*. 2nd edn., Springer, Berlin, 1997.
- [8] A. M. Budylin and V. S. Buslaev, *On the asymptotic behaviour of the spectral characteristics of an integral operator with a difference kernel on expanding domains*, pp. 16–60 in: *Differential Equations, Spectral Theory, Wave Propagation*, Probl. Mat. Fiz. vol. **13**. Leningrad Univ., Leningrad, 1991. [in Russian]
- [9] C. Davis, *A Schwarz inequality for convex operator functions*. Proc. Amer. Math. Soc. **8**: 42–44, 1957.
- [10] E. Fermi, *Zur Quantelung des idealen einatomigen Gases*. Z. Physik **36**: 902–912, 1926. [in German]
- [11] D. Gioev, *Szegő limit theorem for operators with discontinuous symbols and applications to entanglement entropy*. Int. Math. Res. Not. **2006**: 95181, 2006.
- [12] D. Gioev and I. Klich, *Entanglement entropy of fermions in any dimension and the Widom conjecture*. Phys. Rev. Lett. **96**: 100503, 2006.

- [13] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, *Quantum entanglement*. Rev. Mod. Phys. **81**: 865–942, 2009.
- [14] A. Laptev and Y. Safarov, *Szegő type limit theorems*. J. Funct. Anal. **138**: 544–559, 1996.
- [15] H. Leschke, A. V. Sobolev, and W. Spitzer, *Scaling of Rényi entanglement entropies of the free Fermi-gas ground state: a rigorous proof*. Phys. Rev. Lett. **112**: 160403, 2014.
- [16] H. Leschke, A. V. Sobolev, and W. Spitzer, *Large-scale behaviour of local and entanglement entropy of the free Fermi gas at any temperature*. Journal of Physics A: Math.Theor. **49**: 30LT04, 2016; Corrigendum. **50**: 129501, 2017.
- [17] H. Leschke, A. V. Sobolev, and W. Spitzer, *Trace formulas for Wiener–Hopf operators with applications to entropies of free fermionic equilibrium states*. J. Funct. Anal. **273**: 1049–1094, 2017.
- [18] R. Roccaforte, *Asymptotic expansions of traces for certain convolution operators*. Trans. Amer. Math. Soc. **285**: 581–602, 1984.
- [19] B. Simon, *Loewner’s Theorem on Monotone Matrix Functions*. Springer, Cham, 2019.
- [20] A. V. Sobolev, *Pseudo-Differential Operators with Discontinuous Symbols: Widom’s Conjecture*. Mem. Amer. Math. Soc. **222**(1043). AMS, Providence, Rhode Island, 2013.
- [21] A. V. Sobolev, *On the Schatten-von Neumann properties of some pseudo-differential operators*. J. Funct. Anal. **266**: 5886–5911, 2014.
- [22] A. V. Sobolev, *Wiener-Hopf operators in higher dimensions: the Widom conjecture for piece-wise smooth domains*. Integr. Equ. Oper. Theory **81**: 435–449, 2015.
- [23] A. V. Sobolev, *Functions of self-adjoint operators in ideals of compact operators*. J. London Math. Soc. (2) **95**: 157–176, 2017.
- [24] A. V. Sobolev, *Quasi-classical asymptotics for functions of Wiener–Hopf operators: smooth versus non-smooth symbols*. Geom. Funct. Anal. **27**: 676–725, 2017.
- [25] A. V. Sobolev, *On Szegő formulas for truncated Wiener-Hopf operators*. Integr. Equ. Oper. Theory **91**: 1, 2019.
- [26] H. Widom, *A theorem on translation kernels in  $n$  dimensions*. Trans. Amer. Math. Soc. **94**: 170–180, 1960.
- [27] H. Widom, *On a class of integral operators with discontinuous symbol*, pp. 477–500 in: *Toeplitz Centennial*, ed. by I. Gohberg, *Operator Theory: Advances and Applications* vol. **4**. Springer, Basel, 1982.
- [28] H. Widom, *A trace formula for Wiener-Hopf operators*. J. Oper. Theory: **8**: 279–298, 1982.
- [29] H. Widom, *Asymptotic Expansions for Pseudodifferential Operators on Bounded Domains, Lecture Notes in Mathematics* vol. **1152**, Springer, Berlin, 1985.
- [30] H. Widom, *On a class of integral operators on a half-space with discontinuous symbol*. J. Funct. Anal. **88**: 166–193, 1990.

INSTITUT FÜR THEORETISCHE PHYSIK, UNIVERSITÄT ERLANGEN-NÜRNBERG, STAUDTSTRASSE 7, 91058 ERLANGEN, GERMANY

*Email address:* [hajo.leschke@physik.uni-erlangen.de](mailto:hajo.leschke@physik.uni-erlangen.de)

DEPARTMENT OF MATHEMATICS, UNIVERSITY COLLEGE LONDON, GOWER STREET, LONDON, WC1E 6BT UK

*Email address:* [a.sobolev@ucl.ac.uk](mailto:a.sobolev@ucl.ac.uk)

FAKULTÄT FÜR MATHEMATIK UND INFORMATIK, FERNUNIVERSITÄT IN HAGEN, UNIVERSITÄTSSTRASSE 1, 58097 HAGEN, GERMANY

*Email address:* [wolfgang.spitzer@fernuni-hagen.de](mailto:wolfgang.spitzer@fernuni-hagen.de)