

SEMICLASSICAL ANALYSIS OF A NONLOCAL BOUNDARY VALUE PROBLEM RELATED TO MAGNITUDE

HEIKO GIMPERLEIN, MAGNUS GOFFENG, NIKOLETTA LOUCA

ABSTRACT. We study a Dirichlet boundary problem related to the fractional Laplacian in a manifold. Its variational formulation arises in the study of magnitude, an invariant of compact metric spaces given by the reciprocal of the ground state energy. Using recent techniques developed for pseudodifferential boundary problems we discuss the structure of the solution operator and resulting properties of the magnitude. In a semiclassical limit we obtain an asymptotic expansion of the magnitude in terms of curvature invariants of the manifold and the boundary, similar to the invariants arising in short-time expansions for heat kernels.

CONTENTS

1. Introduction	1
2. The symbolic structure near the diagonal	7
3. Global behavior of \mathcal{Z} on compact manifolds	26
4. The operator \mathcal{Z} on Sobolev spaces for a manifold with boundary	33
5. Structure of the inverse operator in the presence of a boundary	37
6. Conditional expectations of Q_X^{-1} and \mathcal{Z}_X^{-1}	47
Appendix A. Overview of conormal distributions	54
Appendix B. Parameter dependent pseudodifferential operators	60
Appendix C. Inverting Q in an extended Boutet de Monvel calculus	65
Appendix D. The meromorphic Fredholm theorem	68
Appendix E. Partial fraction decompositions of symbols	68
Appendix F. Evaluation of some boundary symbols at zero	70
References	75

1. INTRODUCTION

The analysis of boundary problems for nonlocal operators has attracted much interest in recent years, including Dirichlet and Neumann problems for fractional Laplacians in a Euclidean domain. In this article we motivate and initiate the semiclassical analysis of related boundary problems, with applications to the Leinster-Willerton conjecture for the magnitude invariant of compact metric spaces.

To be specific, we consider the integral equation with parameter $R > 0$

$$(1) \quad \int_X e^{-Rd(x,y)} u(y) dy = f(x).$$

Here (X, d) is a compact metric space, and we focus on when X is a manifold with boundary and d is a distance function satisfying additional regularity assumptions. Already when $X \subseteq \mathbb{R}^2$ is the unit disc, close to nothing was known for the solutions to (1). We shall prove in this paper that for X a compact n -dimensional manifold with boundary, Equation (1) is well posed for f in the Sobolev space $\overline{H}^{(n+1)/2}(X)$ and admits a unique solution $u_R \in \dot{H}^{-(n+1)/2}(X)$ for sufficiently large $R \gg 0$ (for notation, see page 33). We relate the integral equation (1) to a pseudodifferential boundary value problem which is elliptic with parameter.

Key words and phrases. 51F99 (primary), 28A75, 58J40, 58J50 (secondary).

Our main results concern structural properties of solutions to Equation (1) such as asymptotic behavior as $R \rightarrow \infty$ and meromorphic extensions in the parameter R to sectors $\Gamma \subseteq \mathbb{C}$. The methods for pseudodifferential boundary value problems that we use date far back, see the work of Gregory Eskin [9] and Lars Hörmander [21], but have in recent years seen much development in work of Gerd Grubb [14, 15, 16, 17, 18].

The solution to (1) for the right hand side $f = 1$ enters in the so called magnitude function of (X, d) , studied extensively in for instance [2, 11, 25, 26, 28, 32, 33, 43, 44]. The empirical properties of the solution to (1) have recently found applications in data science, leading to precise conjectures for its structural properties [6, 7].

The case $f = 1$ can also be considered as a minimizing problem. Equation (1) relates to the ground state energy

(2)

$$\begin{aligned} \mathcal{E}(R; X, d) &:= \inf \left\{ \int_X \int_X e^{-Rd(x,y)} d\mu(x) d\mu(y) : \mu \text{ a signed Borel measure with } \mu(X) = 1 \right\} = \\ &= \inf \left\{ \sum_{x,y \in X} e^{-Rd(x,y)} c(x)c(y) : c : X \rightarrow \mathbb{R} \text{ has finite support and } \sum_{x \in X} c(x) = 1 \right\}. \end{aligned}$$

More precisely, if R is such that (X, Rd) is positive definite (i.e. the matrix $(e^{-Rd(x,y)})_{x,y \in F}$ is positive definite for any finite $F \subseteq X$), then by [33] a solution u_R to Equation (1) with $f = 1$ satisfies

$$\int_X u_R(x) dx = \frac{1}{\mathcal{E}(R; X, d)}.$$

Let us digest on the problem of finding $\mathcal{E}(R; X, d)$ and studying its semiclassical limit, as it has been broadly studied in various mathematical communities. The ground state energy $\mathcal{E}(R; X, d)$ is that of a signed distribution of finitely many particles on X where a particle in x interacts with a particle in y under the potential $e^{-Rd(x,y)}$. As such, the scaling parameter $R > 0$ should be thought of as an order parameter with $R \rightarrow \infty$ corresponding to a semiclassical limit. The non-locality of Equation (1) and the ground state energy $\mathcal{E}(R; X, d)$ makes the problem of explicit computation an impossibility, however in the semiclassical limit the problem localizes and is asymptotically described in terms of geometric invariants. Related problems concerning ground state energies with nonlocal interaction potentials arise in the mean field description of interacting particle systems, such as [10]. Specifically for log gases the dependence of the ground state energy on the geometry has been of interest [46]. In complex geometry, Berman has studied similar minimization problems from which geometric structures emerged [3, 4]. Related operators also appear in image processing [1].

The integral equation (1) is, as mentioned above, related to magnitude – an invariant that has been extensively studied since it was introduced by Tom Leinster [25]. We presume no prerequisites from the reader on magnitude in this paper, but for the convenience of the reader we summarize the implications to magnitude here and expand on this relation in the follow up paper [12]. From its category-theoretic origin, magnitude has found unexpected applications from algebraic topology [13, 27, 40] and applied category theory [8, 35] to data science [6, 7] and mathematical biology [24].

For a metric space (X, d) this invariant leads to a function $\mathcal{M}_X : (0, \infty) \rightarrow \mathbb{R} \cup \{\infty\}$. When the metric space (X, Rd) is positive definite, and in particular for compact sets $X \subset \mathbb{R}^n$ [32], $\mathcal{M}_X(R)$ is defined as $\mathcal{M}_X(R) = \int_X u_R(x) dx$, where u_R satisfies the magnitude equation

$$(3) \quad \int_X e^{-Rd(x,y)} u(y) dy = 1.$$

The work [33] provides an abstract Hilbert space framework in which to pose this equation.

In the case of compact sets $X \subset \mathbb{R}^n$, Meckes [33] gives an interpretation of the magnitude in potential theoretic terms, as a generalized capacity:

$$(4) \quad \mathcal{M}_X(R) = \frac{1}{Rn!\omega_n} \inf \left\{ \|(R^2 + \Delta)^{(n+1)/4} h_R\|_{L^2(\mathbb{R}^n)}^2 : h_R \in H^{(n+1)/2}(\mathbb{R}^n), h_R = 1 \text{ on } X \right\}.$$

Here ω_n denotes the volume of the unit ball in \mathbb{R}^n . The minimizer of (4) is attained at a function $h_R \in H^{(n+1)/2}(\mathbb{R}^n)$ solving the non-local exterior problem

$$(5) \quad \begin{cases} (R^2 + \Delta)^{(n+1)/2} h_R = 0, & \text{in } \mathbb{R}^n \setminus X, \\ h_R = 1, & \text{in } X. \end{cases}$$

If n is odd, this was studied as a boundary value problem for the five-dimensional unit ball in [2], and extended to odd-dimensional unit balls in [45]. Few explicit computations of magnitude are known outside the realm of compact domains in odd-dimensional Euclidean space, and even there the state-of-the art [11] can only provide asymptotic results in the semiclassical limit and ensure existence of meromorphic extensions. In particular, nothing was previously known about magnitude even in such a simple case as the unit disk $X = B_2 \subseteq \mathbb{R}^2$.

We provide a framework for a refined analysis and explicit computations for solutions to Equation (1) when X is a smooth, n -dimensional, compact manifold with boundary, independent of the parity of n . The framework relies on recent advances for pseudodifferential boundary problems and initiates their semiclassical analysis. We work under certain regularity assumptions on the distance function d , firstly that its square is *regular at the diagonal* (see Definition 2.2) and secondly that it has *property (MR)* (see Definition 3.3 and 4.2). Our first assumption ensures that the distance function behaves to leading term as a Euclidean distance, and is satisfied by any geodesic distance function or a pullback thereof under an embedding. The first assumption ensures that the diagonal behavior in Equation (1) is governed by a pseudodifferential operator of order $-n - 1$ which is elliptic with parameter. Our second assumption – property (MR) – is a technical condition to ensure that the off-diagonal behavior in Equation (1) is negligible. Property (MR) is satisfied for subspace distances in manifolds whose distance functions squared are smooth, but it in fact fails for higher dimensional tori and real projective spaces.

1.1. Main results. Let us summarize the main results of this paper. The results all circle around the family of integral operators

$$\mathcal{Z}_X(R)u(x) := \frac{1}{R} \int_X e^{-Rd(x,y)} u(y) dy, \quad R \in \mathbb{C} \setminus \{0\}.$$

Here X is a compact manifold with boundary equipped with a distance function d and a volume density dy . We assume that d^2 is smooth in a small neighborhood of the diagonal $x = y$ and there in local coordinates admits a Taylor expansion (for any $N > 0$)

$$(6) \quad d(x, y)^2 = H_{d^2, x}(v) + \sum_{j=3}^{\infty} C_{d^2}^j(x; v) + O(|v|^{N+1}).$$

Here $v = x - y$, and where H_{d^2} is a Riemannian metric on X and $C_{d^2}^j$ in local coordinates is a symmetric j -form in v . This condition can be summarized in the terminology that d^2 is *regular at the diagonal*, see Definition 2.2 and for more details on the Taylor expansion, see Equation (10). We fix a function $\chi \in C^\infty(X \times X)$ such that $\chi = 1$ on a neighborhood of the diagonal $x = y$ and d^2 is smooth on the support of χ . The localization of \mathcal{Z} to near the diagonal is the integral operator

$$Q_X(R)u(x) := \frac{1}{R} \int_X \chi(x, y) e^{-Rd(x,y)} u(y) dy, \quad R \in \mathbb{C} \setminus \{0\}.$$

We remark that if d^2 is smooth on all of $X \times X$, e.g. for a domain or a submanifold in \mathbb{R}^n with the induced metric, it holds that $\mathcal{Z}_X - Q_X$ is smoothing with parameter on any sector $\Gamma \subseteq \mathbb{C}_+$ with opening angle $< \pi/2$.

Theorem 1.1. *Let X be a compact n -dimensional manifold with boundary and d such that d^2 is regular at the diagonal (see Definition 2.2). The family of integral operators Q_X is an elliptic pseudodifferential operator with parameter $R \in \mathbb{C}_+$ of order $-n - 1$, and its principal symbol is*

$$\bar{\sigma}_{-n-1}(Q_X)(x, \xi, R) = n! \omega_n (R^2 + g_{d^2}(\xi, \xi))^{-(n+1)/2},$$

where g_{d^2} is the dual metric to H_{d^2} from the Taylor expansion (6). The properties of Q_X can be summarized as follows:

- (1) In each coordinate chart, the full symbol of Q_X can be computed by an iterative scheme as in Theorem 2.9.
- (2) There exists an R_0 such that

$$Q_X(R) : \dot{H}^{-\frac{n+1}{2}}(X) \rightarrow \bar{H}^{\frac{n+1}{2}}(X),$$

is invertible for $\operatorname{Re}(R) > R_0$ and $\arg(R) < \pi/(n+1)$. Here $\dot{H}^{-\frac{n+1}{2}}(X)$, respectively $\bar{H}^{\frac{n+1}{2}}(X)$, denote the Sobolev spaces of supported, respectively extendable distributions in X (see Section 4).

- (3) If $\partial X = \emptyset$, then $Q_X(R)^{-1}$ is an elliptic pseudodifferential operator of order $n+1$ whenever it exists. The full symbol of Q_X^{-1} can be computed by an iterative scheme as in Theorem 2.21.
- (4) If $\partial X \neq \emptyset$, Q_X is elliptic with parameter of type $-(n+1)/2$ and factorization index $-\infty$ in an extended Boutet de Monvel calculus. In particular, there is a classical parameter dependent parametrrix in an extended Boutet de Monvel calculus of order $n+1$, type $(n+1)/2$ and factorization index $-\infty$.

Moreover, if d^2 is smooth then all the properties above hold also for \mathcal{Z}_X .

Theorem 1.1 is found in the bulk of the text as follows. The first statement and item (1) is found in Theorem 2.9. Item (2) is proven in Theorem 4.1, see also Corollary 2.22 of Theorem 2.23 for the simpler case of $\partial X = \emptyset$. Item (3) follows from Theorem 2.21 and Corollary 2.22. Item (4) is contained in Theorem 4.9.

The operator Q_X is generally more well behaved than \mathcal{Z}_X ; the off-diagonal singularities of d can create problems in considering \mathcal{Z}_X as a map between Sobolev spaces. For examples of such phenomena, see Subsection 3.4. We impose one of two conditions on d ; property (MR) and property (SMR) respectively to ensure that Q_X and \mathcal{Z}_X share common functional analytic features as operators between Sobolev space. The reader can find the precise definition of property (MR) and property (SMR) in Definition 3.3 (for $\partial X = \emptyset$) and Definition 4.2 (for $\partial X \neq \emptyset$). We note that property (MR) and property (SMR) hold on any sector $\Gamma \subseteq \mathbb{C} \setminus \{0\}$ as soon as d^2 is smooth on all of $X \times X$, e.g. for a domain in \mathbb{R}^n or more generally for a compact submanifold with boundary in a manifold with d^2 smooth.

Theorem 1.2. *Let X be a compact n -dimensional manifold with boundary and let d be a distance function such that d^2 is regular at the diagonal (see Definition 2.2). The family of operators*

$$Q_X(R) : \dot{H}^{-\frac{n+1}{2}}(X) \rightarrow \bar{H}^{\frac{n+1}{2}}(X), \quad R \in \mathbb{C} \setminus \{0\},$$

is a holomorphic family of Fredholm operators that are invertible on a sector. The inverse

$$Q_X(R)^{-1} : \bar{H}^{\frac{n+1}{2}}(X) \rightarrow \dot{H}^{-\frac{n+1}{2}}(X), \quad R \in \mathbb{C} \setminus \{0\},$$

is a meromorphic family of Fredholm operators.

If d satisfies property (SMR) on a sector Γ , then also

$$\mathcal{Z}_X(R) : \dot{H}^{-\frac{n+1}{2}}(X) \rightarrow \bar{H}^{\frac{n+1}{2}}(X), \quad R \in \Gamma,$$

is a holomorphic family of Fredholm operators whose inverse family

$$\mathcal{Z}_X(R)^{-1} : \bar{H}^{\frac{n+1}{2}}(X) \rightarrow \dot{H}^{-\frac{n+1}{2}}(X), \quad R \in \Gamma,$$

is a meromorphic family of Fredholm operators.

For the purposes described above, we are interested in precise asymptotic information about solutions to $\mathcal{Z}_X(R)u = f$. To this end, we were not able to utilize the extended Boutet de Monvel calculus appearing in Theorem 1.1, item (4), but rather we describe the inverse operator \mathcal{Z}_X^{-1} via Wiener-Hopf factorizations.

Theorem 1.3. *Let X be a compact n -dimensional manifold with boundary and d a distance function whose square is regular at the diagonal. For some $R_0 \geq 0$ and any $R \in \Gamma_{\pi/(n+1)}(R_0)$, we can write*

$$Q_X^{-1} = \chi_1 A \chi_1' + \chi_2 (\varphi^{-1})^* W_+ W_- \varphi^* \chi_2' + S,$$

where $\chi_1, \chi'_1 \in C_c^\infty(X^\circ)$, and $\chi_2, \chi'_2 \in C^\infty(X)$ are functions supported in a collar neighborhood U_0 of ∂X in X such that

$$\chi_1 + \chi_2 = 1 \quad \text{and} \quad \chi'_j|_{\text{supp}(\chi_j)} = 1, \quad j = 1, 2,$$

$\varphi : \partial X \times [0, 1) \rightarrow U_0$ is a collar identification, and the operators S , W_- and W_+ satisfy the following:

(1) $S : \overline{H}^\mu(X) \rightarrow \dot{H}^{-\mu}(X)$ is a continuous operator with

$$\|S\|_{\overline{H}^\mu(X) \rightarrow \dot{H}^{-\mu}(X)} = O(R^{-\infty}), \quad \text{as } R \rightarrow \infty.$$

(2) $W_+ : L^2(\partial X \times [0, \infty)) \rightarrow \dot{H}^{-\mu}(\partial X \times [0, \infty))$ is the properly supported pseudodifferential operator of mixed-regularity $(\mu, 0)$ from Definitions 5.18 which is invertible for large $R > 0$ and in local coordinates has an asymptotic expansion modulo $S^{\mu, -\infty}$ as in Lemma 5.21 and preserves support in $\partial X \times [0, \infty) \subseteq \partial X \times \mathbb{R}$. Moreover, for $\chi, \chi' \in C + C_c^\infty(\partial X \times [0, \infty))$ with $\chi\chi' = 0$, it holds that

$$\|\chi W_+ \chi'\|_{L^2(\partial X \times [0, \infty)) \rightarrow H^{-\mu}(\partial X \times \mathbb{R})} = O(R^{-\infty}), \quad \text{as } R \rightarrow \infty.$$

(3) $W_- : \overline{H}^\mu(\partial X \times [0, \infty)) \rightarrow L^2(\partial X \times [0, \infty))$ is the properly supported pseudodifferential operator of mixed-regularity $(\mu, 0)$ from Definitions 5.18 which is invertible for large $R > 0$ and in local coordinates has an asymptotic expansion modulo $S^{\mu, -\infty}$ as in Lemma 5.21 and preserves support in $\partial X \times (-\infty, 0] \subseteq \partial X \times \mathbb{R}$. Moreover, for $\chi, \chi' \in C + C_c^\infty(\partial X \times \mathbb{R})$ with $\chi\chi' = 0$, it holds that

$$\|\chi W_- \chi'\|_{H^\mu(\partial X \times \mathbb{R}) \rightarrow L^2(\partial X \times \mathbb{R})} = O(R^{-\infty}), \quad \text{as } R \rightarrow \infty.$$

The reader can find Theorem 1.3 stated as Theorem 5.22 in the body of the text. A key feature of the construction in Theorem 1.3 is that it provides us with a method to compute the symbolic structure of the inverse Q_X^{-1} .

Theorem 1.4. *Let X be a compact n -dimensional manifold with boundary and d a distance function whose square is regular at the diagonal. In the sector $\text{Re}(R) > R_0$ and $\arg(R) < \pi/(n+1)$, we can write*

$$\langle Q(R)^{-1} 1_X, 1_X \rangle = \sum_{k=0}^{\infty} c_k(X, d) R^{n-k} + O(\text{Re}(R)^{-\infty}), \quad \text{as } \text{Re}(R) \rightarrow \infty \text{ in } \Gamma,$$

where the coefficients $c_k(X, d)$ are given as

$$c_k(X, d) = \int_X a_{k,0}(x, 1) dx + \int_{\partial X} B_{d^2, k}(x) dx',$$

where

- (1) $a_{k,0}(\cdot, 1) \in C^\infty(X)$ is an invariant polynomial in the entries of the Taylor expansion (6) as described in Theorem 2.27 and can be computed inductively using Lemma 2.25, with $a_{k,0} = 0$ if k is odd; and
- (2) $B_{d^2, k} \in C^\infty(\partial X)$ is an invariant polynomial in the entries of the Taylor coefficients of d^2 at the diagonal in X near ∂X as described in Proposition 6.9 and can be inductively computed using Lemma 5.19.

In particular, we have that

$$\begin{aligned} c_0(X, d) &= \frac{\text{vol}(X)}{n! \omega_n} \\ c_1(X, d) &= \frac{(n+1) \text{vol}(\partial X)}{2n! \omega_n}, \\ c_2(X, d) &= \frac{n+1}{6 \cdot n! \omega_n} \int_X s_{d^2} dx + \frac{(n-1)(n+1)^2}{8 \cdot n! \omega_n} \int_{\partial X} H_{d^2} dx'. \end{aligned}$$

where the scalar curvature s_{d^2} is defined as in Theorem 6.1 and the mean curvature H_{d^2} of the distance function is defined as in Theorem 6.13.

The reader can find Theorem 1.4 stated as Theorem 6.13 in the body of the text.

Corollary 1.5. *Let X be a compact n -dimensional manifold with boundary and d a distance function whose square is regular at the diagonal. Write $\mathcal{E}(R; X, d)$ for the ground state energy from Equation (2).*

- (1) *If d has property (MR) on a sector Γ , the ground state energy function $\mathcal{E}(R; X, d)$ is a well defined meromorphic function of $R \in \Gamma$.*
- (2) *If d has property (SMR) on a sector Γ , the ground state energy function $\mathcal{E}(R; X, d)$ has the semiclassical limit*

$$\mathcal{E}(R; X, d) = \sum_{k=0}^{\infty} \varepsilon_k(X, d) R^{-n-k} + O(\operatorname{Re}(R)^{-\infty}), \quad \text{as } \operatorname{Re}(R) \rightarrow \infty \text{ in } \Gamma,$$

where

$$\begin{aligned} \varepsilon_0(X, d) &= \frac{n! \omega_n}{\operatorname{vol}_n(X)}, \\ \varepsilon_1(X, d) &= -\frac{(n+1) \operatorname{vol}_{n-1}(X)}{2 \operatorname{vol}_n(X)}, \\ \varepsilon_2(X, d) &= \frac{(n+1)^2 \operatorname{vol}_{n-1}(X)^2}{4 \operatorname{vol}_n(X)^2} - \frac{n! \omega_n c_2(X, d)}{\operatorname{vol}_n(X)}, \end{aligned}$$

and more generally

$$\varepsilon_k(X, d) = p_{k,n} \left(\frac{c_1(X, d)}{\operatorname{vol}_n(X)}, \dots, \frac{c_k(X, d)}{\operatorname{vol}_n(X)} \right),$$

for a universal polynomial $p_{k,n}$ of total degree k (where each $c_j(X, d)$ is declared to be of degree j).

1.2. Notational conventions. To avoid confusion, we will use the terms Riemannian metrics and distance function to separate the notions of metrics that appear in Riemannian geometry and metric geometry, respectively.

The Fourier transform on \mathbb{R}^n is denoted by \mathcal{F} . We use the convention that

$$D_x = -i \frac{\partial}{\partial x}.$$

For $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$, we write $|\alpha| = \sum_j \alpha_j$, $D_x^\alpha = D_{x_1}^{\alpha_1} \cdots D_{x_n}^{\alpha_n}$ and $x^\alpha = x_1^{\alpha_1} \cdots x_n^{\alpha_n}$. In this convention, for a Schwartz function f on \mathbb{R}^n ,

$$\mathcal{F}(D_x^\alpha f)(\xi) = \xi^\alpha \mathcal{F}f(\xi) \quad \text{and} \quad \mathcal{F}(x^\alpha f)(\xi) = (-D_\xi)^\alpha \mathcal{F}f(\xi),$$

and the product of pseudodifferential symbols is up to smoothing operators defined from a symbol of the form

$$p \# q \sim \sum_{\alpha} \frac{1}{\alpha!} \partial_\xi^\alpha p D_x^\alpha q = \sum_{\alpha} \frac{1}{\alpha!} D_\xi^\alpha p \partial_x^\alpha q.$$

We write M for a manifold and X for a compact manifold with boundary, or occasionally a general compact metric space. We let n denote the dimension of M or X and use the notation

$$\mu := \frac{n+1}{2}.$$

We write $\operatorname{Diag}_M := \{(x, x) : x \in M\}$ for the diagonal in $M \times M$. If (X, d) is a compact metric space such that the matrix $(e^{-Rd(x,y)})_{x,y \in F}$ is positive definite for any finite subset $F \subseteq X$, we say that (X, d) is positive definite. If (X, Rd) is positive definite for all $R > 0$, we say that (X, d) is stably positive definite. This terminology follows [32].

For a manifold M , we write $C_c^\infty(M)$ for the Fréchet space of compactly supported smooth functions and $\mathcal{D}'(M)$ for its topological dual – the distributions on M . If X is a compact manifold with boundary, it can always be embedded as a domain in a manifold M and we write $C^\infty(X) \subseteq C(X)$ for the restrictions to X of elements in $C^\infty(M)$.

A sector $\Gamma \subseteq \mathbb{C}$ is a conical subset, i.e., $\lambda\Gamma \subset \Gamma$ for all $\lambda > 1$. Standard examples we use throughout the paper are $\mathbb{C}_+ = \{z \in \mathbb{C} : \operatorname{Re}(z) > 0\}$ and

$$\Gamma_\alpha(R_0) := \{z \in \mathbb{C} : |\operatorname{Arg}(z)| < \alpha, \operatorname{Re}(z) > R_0\}.$$

If we make a claim concerning $R \rightarrow +\infty$, it is implicitly assumed to be a limit along the real line. We also note that for sectors $\Gamma \subseteq \mathbb{C}_+$ of opening angle $\alpha < \pi/2$, there is a $C_\alpha > 0$ with $C_\alpha^{-1}|R| \leq \operatorname{Re}(R) \leq C_\alpha|R|$.

For two Banach spaces V_1 and V_2 , we write $\mathbb{B}(V_1, V_2)$ for the space of bounded operators $V_1 \rightarrow V_2$ and $\mathbb{K}(V_1, V_2)$ for the space of compact operators $V_1 \rightarrow V_2$. Both form Banach spaces in the norm topology.

We write $\mathbb{N} = \{0, 1, 2, 3, \dots\}$ for the set of natural numbers.

1.3. Acknowledgments. We thank Tony Carbery, Gerd Grubb, Tom Leinster, Rafe Mazzeo, Mark Meckes, Niels Martin Møller, Grigori Rozenblum, Jan-Philip Solovej and Simon Willerton for fruitful and encouraging discussions.

MG was supported by the Swedish Research Council Grant VR 2018-0350. NL was supported by The Maxwell Institute Graduate School in Analysis and its Applications, a Centre for Doctoral Training funded by the UK Engineering and Physical Sciences Research Council (Grant EP/L016508/01), the Scottish Funding Council, Heriot-Watt University and the University of Edinburgh.

2. THE SYMBOLIC STRUCTURE NEAR THE DIAGONAL

To better understand the operator \mathcal{Z} we first consider the case of a manifold M . This analysis describes compact manifolds (see Subsection 2.4 and Section 3) and we carry it over to the interior of a compact manifold with boundary below in Section 4 and 5. We formulate our results in terms of the operator

$$(7) \quad \mathcal{Z}(R)f(x) := \frac{1}{R} \int_M e^{-Rd(x,y)} f(y) dy,$$

whose dependence on $R \neq 0$ is holomorphic under suitable assumptions studied in Section 3 below. Here we are implicitly using a volume density on M , and the operator depends on this choice. We shall later fix a certain choice adapted to the distance function. We shall specify the domain and codomain of this operator more precisely later on. For now, we can consider \mathcal{Z} an operator $C_c^\infty(M) \rightarrow \mathcal{D}'(M)$ by setting

$$\langle \mathcal{Z}\varphi, \psi \rangle = \frac{1}{R} \int_{M \times M} e^{-Rd(x,y)} \varphi(y) \psi(x) dx dy, \quad \text{for } \varphi, \psi \in C_c^\infty(M).$$

2.1. On a class of pseudodifferential operators with parameter. We pick a function $\chi \in C^\infty(M \times M)$ such that $\chi = 1$ near Diag_M and is supported in a small neighborhood of Diag_M . The precise choice of χ will not play an important role, but we shall later specify conditions on its support. The operator \mathcal{Z} decomposes as

$$(8) \quad \mathcal{Z} = Q + L, \quad \text{where } Q(R)f(x) := \frac{1}{R} \int_M \chi(x, y) e^{-Rd(x,y)} f(y) dy.$$

We call the operator Q the *localization of \mathcal{Z} near the diagonal*. In this section we focus our attention to Q . Distance functions might be non-smooth away from the diagonal despite being quite regular at the diagonal and this off-diagonal behavior of the distance function dictates whether or not L is negligible. The remainder L will be studied further in Section 3.

This subsection is devoted to proving that the localized part Q of \mathcal{Z} is a parameter-dependent pseudodifferential operator, for a brief overview thereof, see Appendix B. We will treat a slightly more general class of operators than Q . Consider a family of operators that take the form

$$(9) \quad Q_{G,\chi}(R)f(x) := \frac{1}{|R|_{\text{Mc}}} \int_M \chi(x, y) e^{-|R|_{\text{Mc}} \sqrt{G(x,y)}} f(y) dV(y).$$

Here we have written

$$|R|_{\text{Mc}} := \begin{cases} R, & \operatorname{Re}(R) > 0, \\ -R, & \operatorname{Re}(R) < 0, \end{cases}$$

for the McIntosh modulus which extends the absolute value to a holomorphic function in $\mathbb{C} \setminus i\mathbb{R}$. We shall mainly be concerned with the cases $R \in \mathbb{R}$ and $R \in \mathbb{C}_+$. The cut-off function χ is as above with the additional constraint that G is smooth on $\operatorname{supp}(\chi)$. The function $G : M \times M \rightarrow [0, \infty)$ should be *regular at the diagonal* as made precise in Definition 2.2 below: to define this

notion we first introduce further terminology. Note that $T(M \times M) = p_1^*TM \oplus p_2^*TM$ where $p_j : M \times M \rightarrow M$, $j = 1, 2$, denotes the projection onto the j :th factor. Over the diagonal Diag_M , the map $Dp_1 \oplus Dp_2 : T(M \times M)|_{\text{Diag}_M} \rightarrow TM \oplus TM$ is an isomorphism. We define the transversal tangent bundle to the diagonal to be

$$T_{\text{tra}}\text{Diag}_M := \ker(Dp_1 + Dp_2 : T(M \times M)|_{\text{Diag}_M} \rightarrow TM) \subseteq T(M \times M)|_{\text{Diag}_M}.$$

The restriction $Dp_1| : T_{\text{tra}}\text{Diag}_M \rightarrow TM$ is an isomorphism.

Definition 2.1. For a smooth function G defined in a neighborhood of Diag_M , we define the transversal Hessian H_G as the quadratic form on $T_{\text{tra}}\text{Diag}_M$ obtained from restricting the Hessian of G over the diagonal to the transversal tangent bundle.

Definition 2.2. A function $G : M \times M \rightarrow [0, \infty)$ is said to be *regular at the diagonal* if there is a tubular neighborhood U of the diagonal Diag_M such that $G|_U \in C^\infty(U)$ is a smooth function satisfying that

- $G|_U$ and $dG|_U$ vanish on $\text{Diag}_M \subseteq U$;
- $G(x) > 0$ for $x \in U \setminus \text{Diag}_M$; and
- the transversal Hessian H_G is positive definite in all points of Diag_M .

Remark 2.3. The prototypical example of a function G regular at the diagonal is $G = d^2$ for suitable distance functions d . The function d^2 is regular at the diagonal for the Euclidean distance, or when d is the geodesic distance on a Riemannian manifold (see Example 2.18 below) or more generally a distance function induced from pulling back a geodesic distance along an embedding of M (see Example 2.17 below for an example).

To show that $Q_{G,\chi}(R)$ is an elliptic pseudodifferential operator with parameter, and to describe its full symbol, we shall use a slight detour. The basic idea used in computing the full symbol of $Q_{G,\chi}(R)$ is to do an inverse Fourier transform in R , and then Fourier transform all conormal variables. When we Fourier transform in the R -variable the Schwartz kernel – depending on (x, y, R) – transforms to a conormal distribution on $U \times \mathbb{R}$ (conormal to $\text{Diag}_M \times \{0\}$) – depending on (x, y, η) – that we then Fourier transform in all the transversal directions (v, η) where $v = x - y$, thus producing the full symbol depending on (x, ξ, R) . To compute the Fourier transform in the R -direction, we use the following elementary lemma.

Proposition 2.4. For a parameter $a \geq 0$, and $F_a(R) := \text{F.P.} \frac{e^{-a|R|}}{|R|}$, we have that

$$\mathcal{F}F_a(\eta) = -\log(\eta^2 + a^2) + \log(2) - 2\gamma,$$

where γ is the Euler-Mascheroni constant.

Proof. By taking a derivative in the parameter a and using that the Fourier transform of $R \mapsto e^{-a|R|}$ is $2a(\eta^2 + a^2)^{-1}$, we see that

$$\frac{\partial}{\partial a} \mathcal{F}F_a(\eta) = -2a(\eta^2 + a^2)^{-1}.$$

As such, $\mathcal{F}F_a(\eta) = -\log(\eta^2 + a^2) + c_0(\eta)$ for some tempered distribution c_0 . Setting $a = 0$ and using Proposition A.8, we see that c_0 is a constant. By Proposition A.6 we have that

$$c_0 = \beta_{0,1} = 2\log(2) + \frac{1}{2}\psi(1/2) - \gamma = \log(2) - 2\gamma.$$

□

Proposition 2.5. Let $G : M \times M \rightarrow [0, \infty)$ be a function which is regular at the diagonal, see Definition 2.2. For a neighborhood U of Diag_M on which G is smooth, define $\tilde{G} \in C^\infty(U \times \mathbb{R})$ by

$$\tilde{G}(x, y, \eta) := \eta^2 + G(x, y),$$

and the conormal distribution $\log(\tilde{G}) \in I^{-n-1}(U \times \mathbb{R}; \text{Diag}_M \times \{0\})$ as in Proposition A.10. Then there exists a canonical metric g_G on T^*M such that

$$\sigma_{-n-1}(\log(\tilde{G}))(x, \xi, R) = -2\pi n! \omega_n(R^2 + g_G(\xi, \xi))^{-(n+1)/2}.$$

The canonical metric g_G is dual to the transversal Hessian of G under the isomorphism $Dp_1| : T_{\text{tra}}\text{Diag}_M \rightarrow TM$.

In the following, unless specified otherwise we shall consider M endowed with the Riemannian metric g_G . This allows, in particular, to define a Laplace operator on M .

Proof. Since G is regular at the diagonal, the function \tilde{G} satisfies the assumptions of Proposition A.10 and the result follows therefrom. \square

To compute the full symbol of $Q_{G,\chi}$ we use a Taylor expansion in the direction transversal to the diagonal. Consider a function $G : M \times M \rightarrow [0, \infty)$ which is regular at the diagonal, see Definition 2.2. Consider a coordinate chart $U_0 \subseteq M$. The coordinates on U_0 induce coordinates (x, y) on $U_0 \times U_0$ and we can identify a neighborhood of $\text{Diag}_M \cap (U_0 \times U_0)$ with a neighborhood of the zero section in $T_{\text{tra}}\text{Diag}_M|_{U_0}$ via the map $(x, y) \mapsto (x, x - y)$. If the coordinate chart U_0 on M satisfies that G is smooth on $U_0 \times U_0$, Taylor's theorem implies that for any $N \in \mathbb{N}$ we can on $U_0 \times U_0$ write

$$(10) \quad G(x, y) = H_{G,x}(x - y) + \sum_{j=3}^N C_G^{(j)}(x; x - y) + r_N(x, x - y),$$

for $|x - y|$ small enough, where r_N is a smooth function with $r_N(x, v) = O(|v|^{N+1})$ as $v \rightarrow 0$, H_G is the transversal Hessian of G , and $C_G^{(j)} : U_0 \rightarrow \text{Sym}^j(T_{\text{tra}}\text{Diag}_M|_{U_0})$ takes values in the symmetric j -forms on the transversal tangent bundle $T_{\text{tra}}\text{Diag}_M|_{U_0}$. A short computation shows that H_G indeed is a Riemannian metric on M under the isomorphism $Dp_1| : T_{\text{tra}}\text{Diag}_M \rightarrow TM$. However, each $C_G^{(j)}$ depends on the choice of coordinates, we nevertheless suppress this dependence in the notation.

Since there is a canonical isomorphism $T_{\text{tra}}\text{Diag}_M|_{U_0} \cong TM|_{U_0}$, the symmetric j -form $C_G^{(j)} : U_0 \rightarrow \text{Sym}^j(T_{\text{tra}}\text{Diag}_M|_{U_0})$ appearing in the Taylor expansion (10) of G defines a j :th order differential operator

$$C_G^{(j)}(x, -D_\xi) : C^\infty(T^*M|_{U_0}) \rightarrow C^\infty(T^*M|_{U_0}),$$

obtained by quantizing the coordinate functions, i.e. $C_G^{(j)}(x, -D_\xi)$ acts as multiplication operators by $C_G^{(j)}(x, v)$ under the fiberwise inverse Fourier transform (in the v -direction). For a $k \in \mathbb{N}_+$ and a multiindex $\gamma \in \mathbb{N}_{\geq 3}^k$, we can define a differential operator $C_G^{(\gamma)}(x, -D_\xi)$ on $T^*M|_{U_0}$ by

$$C_G^{(\gamma)}(x, -D_\xi) := \prod_{l=1}^k C_G^{(\gamma_l)}(x, -D_\xi).$$

Since each $C_G^{(\gamma_l)}(x, -D_\xi)$ acts as multiplication operators under the inverse Fourier transform, the differential operators $C_G^{(\gamma_l)}(x, -D_\xi)$, $l = 1, \dots, k$, commute. The order of $C_G^{(\gamma)}(x, -D_\xi)$ is $|\gamma| := \sum_{l=1}^k \gamma_l$. For $j \in \mathbb{N}$, define the finite set

$$I_j := \{\gamma \in \cup_{k=1}^\infty \mathbb{N}_{\geq 3}^k : |\gamma| = j + 2k\}.$$

For instance, we have that

$$I_1 = \{3\}, \quad I_2 = \{(3, 3), 4\}, \quad \text{and} \quad I_3 = \{(3, 3, 3), (4, 3), (3, 4), 5\}.$$

The role of I_j will become clear in Theorem 2.9 below describing the full symbol of $Q_{G,\chi}$ from Equation (9) in a coordinate chart. For $\gamma \in \mathbb{N}^k$, we set $\text{rk}(\gamma) := k$. In other words, $\gamma \in \cup_{k=1}^\infty \mathbb{N}_{\geq 3}^k$ belongs to I_j if and only if $j = |\gamma| - 2\text{rk}(\gamma)$. We remark that $|\gamma| \geq 3$ and $\text{rk}(\gamma) > 0$ is implicit for $\gamma \in I_j$ since $I_j \subseteq \cup_{k=1}^\infty \mathbb{N}_{\geq 3}^k$. The number of elements in $I_j \cap \mathbb{N}_{\geq 3}^k$ is the same as the number of ways to write $j - k$ as a sum of k natural numbers, and so

$$\#(I_j \cap \mathbb{N}_{\geq 3}^k) = \binom{j-1}{k-1}.$$

The following properties of I_j follows.

Proposition 2.6. *Let $j > 0$. The set $I_j \subseteq \cup_{k>0} \mathbb{N}_{\geq 3}^k$ satisfies the following*

- $\max\{|\gamma| : \gamma \in I_j\} = 3j$ and is attained at $\gamma = \vec{3} \in \mathbb{N}^j$.
- $\max\{\gamma_i : \gamma \in I_j\} = j + 2$ and is attained at $\gamma = j + 2 \in \mathbb{N}^1$.
- $\#I_j = 2^{j-1}$.

For notational purposes, we introduce the following notation.

Definition 2.7. For an integer $n \in \mathbb{N} \cup -2\mathbb{N} - 1$, we introduce the notation

$$\omega_n := \frac{\pi^{n/2}}{\Gamma\left(\frac{n}{2} + 1\right)}.$$

If $n > 0$, then ω_n is the volume of the unit ball in n -dimensions.

To simplify the computations in the subsequent theorem, we note the following relations for the Γ -function.

Proposition 2.8. For natural numbers $n, k \in \mathbb{N}$ such that $n > 2k + 1$, we have that

$$\Gamma\left(\frac{n+1}{2} - k\right) \Gamma\left(\frac{n-2k}{2} + 1\right) = \sqrt{\pi} \frac{\Gamma(n-2k+1)}{2^{n-2k}} = \sqrt{\pi} \frac{(n-2k)!}{2^{n-2k}}.$$

Moreover, we have the identities

$$\begin{aligned} \frac{(-1)^{k+1}}{k!} \pi^{(n-1)/2} 2^{n-2k} \Gamma\left(\frac{n+1}{2} - k\right) &= (-1)^{k+1} (n-2k)! \omega_{n-2k} \omega_{2k}, & \text{for } 2k < n \\ \frac{(-1)^{k+1}}{k!} \pi^{(n-1)/2} 2^{n-2k} \Gamma\left(\frac{n+1}{2} - k\right) &= \frac{(-1)^{n/2+1} \omega_{2k}}{(2k-n)! \omega_{2k-n}} = & \text{for } 2k - n \in 2\mathbb{N} \\ &= \frac{(-1)^{n/2+1} (2k-n+1)}{2(2\pi)^{2k-n}} \omega_{2k-n+1} \omega_{2k}, \\ \frac{(-1)^{\frac{n+1}{2}} \pi^{(n-1)/2}}{2^{2k-n} (k - \frac{n+1}{2})! k!} &= \frac{(-1)^{\frac{n+1}{2}}}{(2\pi)^{2k-n}} \omega_{2k} \omega_{2k-n-1}, & \text{for } 2k - n \in 2\mathbb{N} + 1 \end{aligned}$$

Proof. The Legendre duplication formula $\Gamma(\zeta)\Gamma(\zeta+1/2) = 2^{1-2\zeta} \sqrt{\pi} \Gamma(2\zeta)$ applied to $\zeta = n-2k+1$ implies the first stated identity, and the second one follows from the identity $\Gamma(1/2 - m) = (-4)^m m! / (2m)!$. Combining these identities with the definition of ω_n produces the first and second identities. The third identity follows from the definition of ω_n . \square

We now arrive at the main result of this subsection, describing the full symbol of $Q_{G,\chi}$. The reader should keep in mind that we are primarily interested in the function $G(x, y) := d(x, y)^2$ for a distance function d such that d^2 is regular at the diagonal. In this case $Q_{G,\chi}(R) = Q(R)$ is the localization of \mathcal{Z} at the diagonal for $\text{Re}(R) > 0$. We therefore formulate our results on the sector \mathbb{C}_+ , albeit for $Q_{G,\chi}$ they hold in the sector $\mathbb{C} \setminus i\mathbb{R}$.

Theorem 2.9. Let M be an n -dimensional manifold and $G : M \times M \rightarrow [0, \infty)$ a function which is regular at the diagonal. We denote the Riemannian metric on T^*M dual to the transversal Hessian H_G by g_G , as in Proposition 2.5. Consider the operator

$$Q_{G,\chi}(R)f := \frac{1}{|R|_{\text{Mc}}} \int_M \chi(x, y) e^{-|R|_{\text{Mc}} \sqrt{G(x,y)}} f(y) dy,$$

where $\chi \in C^\infty(M \times M)$ a function with $\chi = 1$ near Diag_M and supported only where G is smooth, and we use the Riemannian volume density defined from g_G .

We have that $Q_{G,\chi} \in \Psi_{\text{cl}}^{-n-1}(M; \mathbb{C}_+)$ is a classical elliptic pseudodifferential operator with parameter of order $-n-1$ with principal symbol

$$\bar{\sigma}_{-n-1}(Q_{G,\chi})(x, \xi, R) = n! \omega_n (R^2 + g_G(\xi, \xi))^{-(n+1)/2}.$$

In a coordinate chart U_0 on M , the full symbol q of $Q_{G,\chi}$ has a classical asymptotic expansion $q \sim \sum_{j=0}^{\infty} q_j$ computed from the Taylor expansion (10) and each q_j is the homogeneous symbol of degree $-n-1-j$ which for $j > 0$ and for n odd is given by

$$\begin{aligned} q_j(x, \xi, R) &= \sum_{\gamma \in I_j, \text{rk}(\gamma) < (n+1)/2} \mathbf{c}_{\text{rk}(\gamma), n} C_G^{(\gamma)}(x, -D_\xi) (R^2 + g_G(\xi, \xi))^{-(n+1)/2 + \text{rk}(\gamma)} - \\ &- \sum_{\gamma \in I_j, \text{rk}(\gamma) \geq (n+1)/2} \mathbf{c}_{\text{rk}(\gamma), n} C_G^{(\gamma)}(x, -D_\xi) \left[(R^2 + g_G(\xi, \xi))^{-(n+1)/2 + \text{rk}(\gamma)} \log(R^2 + g_G(\xi, \xi)) \right], \end{aligned}$$

and for n even given as

$$q_j(x, \xi, R) = \sum_{\gamma \in I_j} \mathbf{c}_{\text{rk}(\gamma), n} C_G^{(\gamma)}(x, -D_\xi)(R^2 + g_G(\xi, \xi))^{-(n+1)/2 + \text{rk}(\gamma)}$$

Here the coefficients are computed as

$$\mathbf{c}_{k, n} := \begin{cases} (-1)^k (n-2k)! \omega_{n-2k} \omega_{2k}, & \text{for } 2k < n \\ \frac{(-1)^{1-n/2} \omega_{2k}}{(2k-n)! \omega_{2k-n}}, & \text{for } 2k - n \in 2\mathbb{N} \\ \frac{(-1)^{\frac{n+1}{2}}}{(2\pi)^{2k-n}} \omega_{2k} \omega_{2k-n-1}, & \text{for } 2k - n \in 2\mathbb{N} + 1 \end{cases}$$

Remark 2.10. Exact expressions for q_1 and q_2 are given below in Proposition 2.14 and 2.15, respectively.

Remark 2.11. From the expression for q_j in Theorem 2.9, it is not clear that q_j is homogeneous of degree $-n-1-j$ when n is odd. We shall see in the proof that this is in fact the case.

Proof of Theorem 2.9. By Proposition 2.4, the Schwartz kernel $\frac{1}{|R|} \chi(x, y) e^{-|R| \sqrt{G(x, y)}}$ of $Q_{G, \chi}$ is the Fourier transform in the η -direction of

$$K(x, y, \eta) = -\frac{1}{2\pi} \chi(x, y) \log(\eta^2 + G(x, y)).$$

The extra 2π is coming from Fourier inversion in one dimension.

Let U denote a neighborhood of the diagonal Diag_M on which G is smooth. It follows from Proposition 2.5 (cf. Proposition A.10) that $K \in I^{-n-1}(U \times \mathbb{R}; \text{Diag}_M \times \{0\})$ with principal symbol

$$\bar{\sigma}_{-n-1}(K)(x, \xi, R) = n! \omega_n (R^2 + g_G(\xi, \xi))^{-(n+1)/2}.$$

Define $K_0(x, y, \eta) := \log(\eta^2 + G(x, y)) \in I^{-n-1}(U \times \mathbb{R}; \text{Diag}_M \times \{0\})$. We compute in a coordinate chart U_0 that $K_0 \in CI^{-n-1}(U \times \mathbb{R}; \text{Diag}_M \times \{0\})$ and using a uniform asymptotic expansion we use Proposition B.14 to show that the Fourier transform in the η -direction of K is the Schwartz kernel of a pseudodifferential operator with parameter.

In a coordinate chart U_0 , we introduce the coordinates $(x, v) = (x, x - y)$ on $U_0 \times U_0$. Using Equation (10), we can write

$$K_0(x, y, \eta) = -\log(\eta^2 + H_G(v, v)) - \log\left(1 + \frac{\sum_{j=3}^N C_G^{(j)}(v, v) + r_N(x, v)}{\eta^2 + H_G(v, v)}\right).$$

For small v , we can Taylor expand

$$K_0(x, y, \eta) = -\log(\eta^2 + H_G(v, v)) - \sum_{j=1}^N \sum_{\gamma \in I_j} \frac{(-1)^{\text{rk}(\gamma)}}{\text{rk}(\gamma)} \frac{C_G^{(\gamma)}(v)}{(\eta^2 + H_G(v, v))^{\text{rk}(\gamma)}} + \tilde{r}_N(x, v, \eta).$$

We note that, by the definition of I_j , each term in the second sum $\sum_{\gamma \in I_j} \frac{(-1)^{\text{rk}(\gamma)}}{\text{rk}(\gamma)} \frac{C_G^{(\gamma)}(v)}{(\eta^2 + H_G(v, v))^{\text{rk}(\gamma)}}$ is homogeneous of degree j . We also note that $\tilde{r}_N(x, v, \eta) = O((|\eta| + |v|)^{N+1})$ and a short computation gives that $\partial_x^\alpha \partial_v^\beta \partial_\eta^k \tilde{r}_N = O((|\eta| + |v|)^{N+1-|\beta|-k})$ for any multiindices α, β and k . As such, we have a uniform asymptotic expansion $K \sim \sum_{j=0}^\infty K_j$ (cf. Definition B.13) where

$$(11) \quad K_j(x, v, \eta) = \begin{cases} -\frac{1}{2\pi} \log(\eta^2 + H_G(v, v)), & j = 0, \\ \frac{1}{2\pi} \sum_{\gamma \in I_j} \frac{(-1)^{\text{rk}(\gamma)+1}}{\text{rk}(\gamma)} \frac{C_G^{(\gamma)}(v)}{(\eta^2 + H_G(v, v))^{\text{rk}(\gamma)}}, & j > 0. \end{cases}$$

We conclude from Proposition B.14 that $Q_{G, \chi} \in \Psi_{\text{cl}}^{-n-1}(M; \mathbb{R})$ is a pseudodifferential operator with parameter. Proposition B.14 implies that

$$\bar{\sigma}_{-n-1}(Q_{G, \chi})(x, \xi, R) = \bar{\sigma}_{-n-1}(K)(x, \xi, R) = n! \omega_n (R^2 + g_G(\xi, \xi))^{-(n+1)/2}.$$

This is invertible in $C^\infty(S(T^*M \oplus \mathbb{R}))$ so $Q_{G, \chi}$ is elliptic with parameter. It is readily seen that $\bar{\sigma}_{-n-1}(Q_{G, \chi})(x, \xi, R) \neq 0$ also for $R \in \mathbb{C}_+$, and the following symbol computation shows that $Q_{G, \chi} \in \Psi_{\text{cl}}^{-n-1}(M; \mathbb{C}_+)$ is an elliptic pseudodifferential operator with parameter.

Let us turn to the full symbol of $Q_{G,\chi}$. We will compute it from Equation (11) and the Fourier transform computations of Subsection A.1 of the appendix. For $k > 0$, denote the Fourier transform of $(\eta^2 + H_G(v, v))^{-k}$ in the (v, η) -direction by $F_k(x, \xi, R)$, that is

$$F_k(x, \xi, R) := \int_{T_x M \oplus \mathbb{R}} \frac{e^{-i\xi \cdot v - iR\eta}}{(\eta^2 + H_G(v, v))^k} dv d\eta.$$

Using homogeneity and rotational symmetries, F_k can be computed as in Proposition A.7 to be

$$\frac{\pi^{(n+1)/2} 2^{-2k+n+1} \Gamma\left(\frac{n+1}{2} - k\right)}{(k-1)!} (R^2 + g_G(\xi, \xi))^{-(n+1)/2+k},$$

for $2k - n - 1 \notin 2\mathbb{N}$, and

(12)

$$\frac{(-1)^{k-\frac{n+1}{2}} \pi^{(n+1)/2}}{2^{2k-n-1} \left(k - \frac{n+1}{2}\right)! (k-1)!} \left[(R^2 + g_G(\xi, \xi))^{-(n+1)/2+k} (-\log(R^2 + g_G(\xi, \xi)) + \beta_{k-(n+1)/2, n+1}) \right],$$

for $2k - n - 1 \in 2\mathbb{N}$. It follows that for $\gamma \in I_j$, the symbol of the term

$$\frac{C_G^{(\gamma)}(v)}{(\eta^2 + H_G(v, v))^{\text{rk}(\gamma)}},$$

is given by $C_G^{(\gamma)}(x, -D_\xi) F_{\text{rk}(\gamma)}(x, \xi, R)$ which is computed to be

$$\frac{\pi^{(n+1)/2} 2^{-2\text{rk}(\gamma)+n+1} \Gamma\left(\frac{n+1}{2} - \text{rk}(\gamma)\right)}{(\text{rk}(\gamma) - 1)!} C_G^{(\gamma)}(x, -D_\xi) (R^2 + g_G(\xi, \xi))^{-(n+1)/2+\text{rk}(\gamma)},$$

for $2\text{rk}(\gamma) - n - 1 \notin 2\mathbb{N}$, and

$$\frac{(-1)^{\text{rk}(\gamma) - \frac{n+1}{2}} \pi^{(n+1)/2}}{2^{2\text{rk}(\gamma) - n - 1}} \left(\text{rk}(\gamma) - \frac{n+1}{2} \right)! (\text{rk}(\gamma) - 1)!.$$

$$(13) \quad C_G^{(\gamma)}(x, -D_\xi) \left[(R^2 + g_G(\xi, \xi))^{-(n+1)/2+\text{rk}(\gamma)} (-\log(R^2 + g_G(\xi, \xi)) + \beta_{\text{rk}(\gamma) - (n+1)/2, n+1}) \right],$$

for $2\text{rk}(\gamma) - n - 1 \in 2\mathbb{N}$. We conclude that the symbol q_j of K_j , for $j > 0$, is (up to the term $\beta_{\text{rk}(\gamma), n+1}$) given by the formula in the statement of the theorem for the pre-factors $\mathbf{c}_{k,n}$:

$$\mathbf{c}_{k,n} = \begin{cases} \frac{(-1)^{k+1}}{k!} \pi^{(n-1)/2} 2^{n-2k} \Gamma\left(\frac{n+1}{2} - k\right), & \text{for } 2k - n - 1 \notin 2\mathbb{N}, \\ \frac{(-1)^{\frac{n+1}{2}} \pi^{(n-1)/2}}{2^{2k-n} \left(k - \frac{n+1}{2}\right)! k!}, & \text{for } 2k - n - 1 \in 2\mathbb{N}. \end{cases}$$

Therefore $\mathbf{c}_{k,n}$ takes the form prescribed in the theorem by Proposition 2.8.

We finish the proof by showing that $\beta_{\text{rk}(\gamma) - (n+1)/2, n+1}$ does not contribute in Equation (13) and that there is no logarithmic term when expanding the ξ -derivatives in Equation (13). If $2\text{rk}(\gamma) - n - 1 \in 2\mathbb{N}$, then $-(n+1)/2 + \text{rk}(\gamma) \in \mathbb{N}$ and so $(R^2 + g_d(\xi, \xi))^{-(n+1)/2+\text{rk}(\gamma)}$ is a polynomial of degree $2\text{rk}(\gamma) - n - 1 \in 2\mathbb{N}$ in ξ . For $\gamma \in I_j \subseteq \cup_k \mathbb{N}_{\geq 3}^k$, we have that

$$j + 2\text{rk}(\gamma) = |\gamma| \geq 3\text{rk}(\gamma).$$

Therefore, if $2\text{rk}(\gamma) - n - 1 \in 2\mathbb{N}$ then $C_d^{(\gamma)}(x, D_\xi) (R^2 + g_d(\xi, \xi))^{-(n+1)/2+\text{rk}(\gamma)}$ is a polynomial of degree $2\text{rk}(\gamma) - n - 1 - |\gamma| \leq -\text{rk}(\gamma) - n - 1 < 0$ and therefore it must be identically zero. In other words, we have the equality

$$(14) \quad C_G^{(\gamma)}(x, -D_\xi) (R^2 + g_G(\xi, \xi))^{-(n+1)/2+\text{rk}(\gamma)} = 0,$$

for $2\text{rk}(\gamma) - n - 1 \in 2\mathbb{N}$. This equality proves that q_j can not contain a term with a logarithmic factor, and as such q_j is homogeneous of degree $-n - 1 - j$ for all j , and

$$\begin{aligned} & C_G^{(\gamma)}(x, -D_\xi) \left[(R^2 + g_G(\xi, \xi))^{-(n+1)/2+\text{rk}(\gamma)} (-\log(R^2 + g_G(\xi, \xi)) + \beta_{\text{rk}(\gamma) - (n+1)/2, n+1}) \right] = \\ & = -C_G^{(\gamma)}(x, -D_\xi) \left[(R^2 + g_G(\xi, \xi))^{-(n+1)/2+\text{rk}(\gamma)} \log(R^2 + g_G(\xi, \xi)) \right]. \end{aligned}$$

□

We note the following corollary that will play a role later in the paper when we consider manifolds with boundary, and study the inverse of Q in the Boutet de Monvel calculus. See more in Theorem 4.9 and Appendix C. Recall that a complete classical parameter dependent symbol $p \sim \sum_l p_l$ of order m on a compact manifold M is said to satisfy the μ -transmission condition at a hypersurface $Y \subseteq M$ if in local coordinates $x = (x', x_n)$, with Y being defined by $x_n = 0$, satisfies that

$$D_x^\beta D_\xi^\alpha D_R^j p_l((x', 0), (0, \xi_n), 0) = e^{\pi i(m-l-2\mu-|\alpha|-j)} D_x^\beta D_\xi^\alpha D_R^j p_l((x', 0), (0, -\xi_n), 0).$$

One sometimes in short says that p has type μ . A classical pseudodifferential operator with parameter is said to be of type μ if its symbol is of type μ . See more in for instance [15, Proposition 1]. Item (b) of Theorem 3.7 implies the following.

Corollary 2.12. *Let M be a manifold and $G : M \times M \rightarrow [0, \infty)$ a function which is regular at the diagonal. Set $\mu := -(n+1)/2$. The pseudodifferential operator with parameter $Q_{G,x}$ satisfies the μ -transmission condition along any hypersurface in M . In fact, the asymptotic expansion $\sum_{l=0}^\infty q_l$ of the full symbol of Q from Theorem 3.7 satisfies that*

$$D_x^\beta D_\xi^\alpha D_R^j q_l(x, \xi, R) = (-1)^{|\alpha|+j+l} D_x^\beta D_\xi^\alpha D_R^j q_l(x, -\xi, -R),$$

for any multiindices $\alpha, \beta \in \mathbb{N}^n$, $x \in M$ and $(\xi, R) \neq 0$.

Remark 2.13. The computation that

$$q_0(x, \xi, R) = n! \omega_n (R^2 + g_G(\xi, \xi))^{-(n+1)/2},$$

is compatible with known symbol computations in \mathbb{R}^n for $G(x, y) = |x - y|^2$ in which case $q = q_0$ is the full symbol expansion when using Euclidean coordinates. Indeed, this statement follows from the fact that $e^{-R|v|}$ is the Fourier transform of $n! \omega_n (R^2 + |\xi|^2)^{-(n+1)/2}$, see for instance [2, Equation (3)]. We remark that the coordinate dependent symbol computations of Theorem 2.9 will be used also in \mathbb{R}^n . The reason for using Theorem 2.9 in \mathbb{R}^n is that to describe the operator near the boundary of a domain with the Wiener-Hopf factorization techniques of Section 5 below we need to “straighten out the boundary”, i.e. choose coordinates in which the boundary locally looks like a half-space. We make such computations more precise in Subsection 2.2 below.

Let us give some further details in computing the symbols q_1 and q_2 . The precise information contained in q_1 and q_2 will be used later to compute the first terms in the inverse of Q and the asymptotics of its expectation values in Section 6. Before entering into the symbol computations of q_1 and q_2 , let us introduce some notation. Since g is a metric on T^*M , it can be viewed as a symmetric tensor in $TM \otimes TM$ and for a covector ξ , the contraction $\iota_\xi g_G$ takes values in TM . In the coordinate chart, each C_G^j takes values in the symmetric j -forms on $TM|_{U_0}$. As such, expressions such as $C_G^3(x, g_G \otimes \iota_\xi g_G)$ or $C_G^4(x, g_G \otimes g_G)$, for instance, make sense. The reader should be aware that such expressions are individually not coordinate invariant, the transformation rules for these expressions can be deduced either from the Taylor expansion (10) or from the transformation rules for pseudodifferential operators. For computational purposes, we also note that

$$d_\xi g_G(\xi, \xi) = 2 \iota_\xi g_G.$$

Here d_ξ denotes the fiberwise exterior differential and we are implicitly using the canonical identification $T^*(T^*M) = \pi^* TM \oplus \pi^* T^*M$ and that the TM -summand is where d_ξ maps to. A useful tool in formulating the computations is the Pochhammer k -symbol. For $x \in \mathbb{R}$, $n \in \mathbb{N}$ and $k \in \mathbb{Z}$, we write

$$(x)_{n,k} := \underbrace{x(x+k)(x+2k) \cdots (x+(n-1)k)}_{n \text{ factors}},$$

with the convention that $(x)_{0,k} = 1$ for any k and $(0)_{n,k} = 1$ for any n and k .

A combinatorial argument shows that

$$\begin{aligned} C_G^3(x, -D_\xi)(R^2 + g_G(\xi, \xi))^{-\nu} &= -3 \cdot 2^2 i(\nu)_{2,1} C_G^3(x, g_G \otimes \iota_\xi g_G)(R^2 + g_G(\xi, \xi))^{-\nu-2} + \\ &\quad + 2^3 i(\nu)_{3,1} C_G^3(x, \iota_\xi g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G)(R^2 + g_G(\xi, \xi))^{-\nu-3}. \end{aligned}$$

$$\begin{aligned} C_G^4(x, -D_\xi)(R^2 + g_G(\xi, \xi))^{-\nu} &= 3 \cdot 2^2(\nu)_{2,1} C_G^4(x, g_G \otimes g_G)(R^2 + g_G(\xi, \xi))^{-\nu-2} - \\ &\quad - 6 \cdot 2^3(\nu)_{3,1} C_G^4(x, g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G)(R^2 + g_G(\xi, \xi))^{-\nu-3} + \\ &\quad + 2^4(\nu)_{4,1} C_G^4(x, \iota_\xi g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G)(R^2 + g_G(\xi, \xi))^{-\nu-4}. \end{aligned}$$

$$\begin{aligned} C_G^3(x, -D_\xi)^2(R^2 + g_G(\xi, \xi))^{-\nu} &= -24 \cdot 2^4(\nu)_{4,1} C_G^3(x, g_G \otimes \iota_\xi g_G)^2(R^2 + g_G(\xi, \xi))^{-\nu-4} + \\ &\quad + 6 \cdot 2^5(\nu)_{5,1} C_G^3(x, \iota_\xi g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G) C_G^3(x, g_G \otimes \iota_\xi g_G)(R^2 + g_G(\xi, \xi))^{-\nu-5} - \\ &\quad - 2^6(\nu)_{6,1} C_G^3(x, \iota_\xi g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G)^2(R^2 + g_G(\xi, \xi))^{-\nu-6} + \\ &\quad + 3 \cdot 2^3(\nu)_{3,1} (C_G^3 \otimes C_G^3)(x, g_G \otimes g_G \otimes g_G)(R^2 + g_G(\xi, \xi))^{-\nu-3}. \end{aligned}$$

Proposition 2.14. *Let M and $Q_{G,\chi}$ be as in Theorem 2.9. In a coordinate chart on M , the term q_1 of degree $-n-2$ appearing in the full symbol q of $Q_{G,\chi}$ is given by*

$$\begin{aligned} q_1(x, \xi, R) &= -i \left(6C_G^3(x, g_G \otimes \iota_\xi g_G)(R^2 + g_G(\xi, \xi))^{-2} - \right. \\ &\quad \left. - 8C_G^3(x, \iota_\xi g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G)(R^2 + g_G(\xi, \xi))^{-3} \right), \end{aligned}$$

if $n = 1$, and if $n > 1$ we have that

$$\begin{aligned} q_1(x, \xi, R) &= -i(n^2 - 1)\mathfrak{c}_{1,n} \left(3C_G^3(x, g_G \otimes \iota_\xi g_G)(R^2 + g_G(\xi, \xi))^{-(n+1)/2-1} - \right. \\ &\quad \left. - (n+3)C_G^3(x, \iota_\xi g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G)(R^2 + g_G(\xi, \xi))^{-(n+1)/2-2} \right), \end{aligned}$$

where $\iota_\xi g_G$ denotes the contraction of the metric g_G on T^*M along the covector ξ .

Proof. We note that $I_1 = \{\gamma \in \cup_{k=1}^\infty \mathbb{N}_{\geq 3}^k : |\gamma| = 1 + 2\text{rk}(\gamma)\} = \{3\}$. For $n = 1$, we have $\mathfrak{c}_{1,1} = -1/2$ and since we are in the critical case we compute as follows

$$\begin{aligned} q_1(x, \xi, R) &= -i\mathfrak{c}_{1,1} C^3(x, 1) \partial_\xi^3 \log(R^2 + g_G(\xi, \xi)) = iC^3(x, 1) \partial_\xi^2 \frac{\iota_\xi g_G}{R^2 + g_G(\xi, \xi)} = \\ &= iC_G^3(x, 1) \left(-\frac{6g_G \iota_\xi g_G}{(R^2 + g_G(\xi, \xi))^2} + \frac{8(\iota_\xi g_G)^3}{(R^2 + g_G(\xi, \xi))^3} \right), \end{aligned}$$

which proves the case $n = 1$.

For $n > 1$, we compute using the identities above that

$$\begin{aligned} q_1(x, \xi, R) &= i(n+1)(n-1)\mathfrak{c}_{1,n} \left(-3C_G^3(x, g_G \otimes \iota_\xi g_G)(R^2 + g_G(\xi, \xi))^{-(n+1)/2-1} + \right. \\ &\quad \left. + (n+3)C_G^3(x, \iota_\xi g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G)(R^2 + g_G(\xi, \xi))^{-(n+1)/2-2} \right). \end{aligned}$$

□

The next proposition follows from a similar computation.

Proposition 2.15. *Let M and $Q_{G,\chi}$ be as in Theorem 2.9. In a coordinate chart on M , the term q_2 of degree $-n-3$ appearing in the full symbol q of $Q_{G,\chi}$ is given as follows; for $n = 1$ we have that*

$$\begin{aligned} q_2(x, \xi, R) &= \mathfrak{c}_{2,1} C^3(x, 1)^2 \partial_\xi^6 ((R^2 + g_G(\xi, \xi)) \log(R^2 + g_G(\xi, \xi))) + \mathfrak{c}_{1,1} C^4(x, 1) \partial_\xi^4 \log(R^2 + g_G(\xi, \xi)) = \\ &= -\frac{C^3(x, 1)^2}{16} \left(-\frac{2^3 \cdot 15g_G^3}{(R^2 + g_G)^2} + \frac{2^4 \cdot 90(\iota_\xi g_G)^2 g_G^2}{(R^2 + g_G)^3} - \frac{2^4 \cdot 80(\iota_\xi g_G)^4 g_G}{(R^2 + g_G)^4} + \frac{2^6 \cdot 24(\iota_\xi g_G)^6}{(R^2 + g_G)^5} \right) \\ &\quad - \frac{C^4(x, 1)}{2} \left(-\frac{2^2 \cdot 3g_G^2}{(R^2 + g_G)^2} + \frac{2^3 \cdot 12(\iota_\xi g_G)^2 g_G}{(R^2 + g_G)^3} - \frac{2^4 \cdot 6(\iota_\xi g_G)^4}{(R^2 + g_G)^4} \right), \end{aligned}$$

and for $n = 3$, we have that

$$\begin{aligned}
q_2(x, \xi, R) &= \mathbf{c}_{2,3} C^3(x, \partial_\xi)^2 (\log(R^2 + g_G(\xi, \xi))) + \mathbf{c}_{1,3} C^4(x, \partial_\xi) (R^2 + g_G(\xi, \xi))^{-1} = \\
&= \mathbf{c}_{2,3} 240 (C_G^3 \otimes C_G^3) (g_G \otimes g_G \otimes g_G) (R^2 + g_G(\xi, \xi))^{-3} - \\
&\quad - \mathbf{c}_{2,3} 45 \cdot 96 C_G^3(x, g_G \otimes \iota_\xi g_G)^2 (R^2 + g_G(\xi, \xi))^{-4} + \\
&\quad + \mathbf{c}_{2,3} 120 \cdot 96 C_G^3(x, \iota_\xi g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G) C_G^3(x, g_G \otimes \iota_\xi g_G) (R^2 + g_G(\xi, \xi))^{-5} - \\
&\quad - \mathbf{c}_{2,3} 80 \cdot 96 C_G^3(x, g_G \otimes \iota_\xi g_G)^2 (R^2 + g_G(\xi, \xi))^{-6} + \\
&\quad + \mathbf{c}_{1,3} 24 C^4(x, g_G \otimes g_G) (R^2 + g_G(\xi, \xi))^{-3} - \\
&\quad - \mathbf{c}_{1,3} 288 C^4(x, g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G) (R^2 + g_G(\xi, \xi))^{-4} + \\
&\quad + \mathbf{c}_{1,3} 192 C^4(x, \iota_\xi g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G) (R^2 + g_G(\xi, \xi))^{-5}
\end{aligned}$$

and finally for $n \neq 1, 3$, we have that

$$\begin{aligned}
q_2(x, \xi, R) &= -\mathbf{c}_{2,n} C^3(x, \partial_\xi)^2 (R^2 + g_G(\xi, \xi))^{-(n-3)/2} + \mathbf{c}_{1,n} C^4(x, \partial_\xi) (R^2 + g_G(\xi, \xi))^{-(n-1)/2} = \\
&= -24 \mathbf{c}_{2,n} (n+3)_{4,-2} C^3(x, g_G \otimes \iota_\xi g_G)^2 (R^2 + g_G(\xi, \xi))^{-(n+1)/2-2} + \\
&\quad + 6 \mathbf{c}_{2,n} (n+5)_{5,-2} C^3(x, \iota_\xi g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G) C^3(x, g_G \otimes \iota_\xi g_G) (R^2 + g_G(\xi, \xi))^{-(n+1)/2-3} - \\
&\quad - \mathbf{c}_{2,n} (n+7)_{6,-2} C^3(x, \iota_\xi g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G)^2 (R^2 + g_G(\xi, \xi))^{-(n+1)/2-4} + \\
&\quad + 3 \mathbf{c}_{2,n} (n+5)_{5,-2} (C^3 \otimes C^3)(x, g_G \otimes g_G \otimes g_G) (R^2 + g_G(\xi, \xi))^{-(n+1)/2-1} + \\
&\quad + 3 \mathbf{c}_{1,n} (n^2 - 1) C^4(x, g_G \otimes g_G) (R^2 + g_G(\xi, \xi))^{-(n+1)/2-1} - \\
&\quad - 6 \mathbf{c}_{1,n} (n+3)_{3,-2} C^4(x, g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G) (R^2 + g_G(\xi, \xi))^{-(n+1)/2-2} + \\
&\quad + \mathbf{c}_{1,n} (n+5)_{4,-2} C^4(x, \iota_\xi g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G) (R^2 + g_G(\xi, \xi))^{-(n+1)/2-3},
\end{aligned}$$

where $\iota_\xi g_G$ denotes the contraction of the metric g_G on T^*M along the covector ξ .

2.2. Examples and further structure in the symbol computations. In the preceding subsection we saw a detailed computation of the full symbol of $Q_{\chi, G}$ in terms of the Taylor expansion. Let us consider a few important special cases where further structures can be visible in the symbol expansion, and proceed with a structural statement of for the entries in the full symbol in the general case.

Example 2.16 (Symbol computations near a boundary in Euclidean space). We consider the manifold $M = \mathbb{R}^n$ and the function $G(x, y) = |x - y|^2$ which is regular at the diagonal. This example fits into the bigger picture of the paper seeing that $G(x, y) = d(x, y)^2$. It will later in the paper be crucial to describe the symbol of $Q = Q_{G, \chi}$ near the boundary of a domain $X \subseteq \mathbb{R}^n$ with smooth boundary. Fix a point $x_0 \in \partial X$. Up to a rigid motion, we can assume that $x_0 = 0$ and that the normal vector of ∂X in x_0 is orthogonal to the plane $x_n = 0$, where we write $x = (x', x_n)$ for $x' \in \mathbb{R}^{n-1}$ and $x_n \in \mathbb{R}$. There is a neighborhood $U_0 = U_{00} \times (-\varepsilon, \varepsilon)$ of $0 \in \mathbb{R}^n$ and a smooth function $\varphi \in C^\infty(U_{00})$ such that $X \cap U_0 = \{x = (x', x_n) \in U_0 : \varphi(x') < x_n\}$. Since the normal vector of ∂X in x_0 is orthogonal to the plane $x_n = 0$, $\nabla_{x'} \varphi(0) = 0$. Near $x_0 = 0$, we use the coordinates

$$(x', x_n) \mapsto (x', x_n - \varphi(x')).$$

In these new coordinates, the domain X looks like the half-space $\{(x', x_n) : x_n > 0\}$ locally near $x_0 = 0$. We compute that in these coordinates $G = G(x, y)$ can be written as

$$\begin{aligned}
|(x', x_n - \varphi(x')) - (y', y_n - \varphi(y'))|^2 &= |x' - y'|^2 + (x_n - y_n - (\varphi(x') - \varphi(y')))^2 = \\
&= |x - y|^2 - 2(x_n - y_n)(\varphi(x') - \varphi(y')) + (\varphi(x') - \varphi(y'))^2 = \\
&= |v|^2 + \sum_{j=2}^N \sum_{|\alpha'|=j-1} \frac{-2\partial_{x'}^{\alpha'} \varphi(x')}{\alpha'!} v_n (v')^{\alpha'} + \\
&\quad + \sum_{j=2}^N \sum_{\substack{|\alpha'|+|\beta'|=j, \\ |\alpha'|, |\beta'|>0}} \frac{\partial_{x'}^{\alpha'} \varphi(x') \partial_{y'}^{\beta'} \varphi(y')}{\alpha'! \beta'!} (v')^{\alpha'+\beta'} + O(|v|^{N+1}),
\end{aligned}$$

where the sums over α' and β' ranges over $\alpha', \beta' \in \mathbb{N}^{n-1}$ and we have written $v = x - y$, $v' = x' - y'$ and $v_n = x_n - y_n$. Introducing the notation

$$\nabla^j \varphi(x'; v') := \sum_{|\alpha'|=j} \frac{\partial_{x'}^{\alpha'} \varphi(x')}{\alpha'!} (v')^{\alpha'},$$

we have in these coordinates that

$$\begin{aligned} G(x, y) &= |v|^2 - 2v_n \nabla \varphi(x') \cdot v' + (\nabla \varphi(x') \cdot v')^2 + \sum_{j=3}^N -2v_n \nabla^{j-1} \varphi(x'; v') \\ &+ \sum_{j=3}^N \sum_{\substack{k+l=j, \\ k, l > 0}} \nabla^k \varphi(x'; v') \nabla^l \varphi(x'; v') + O(|v|^{N+1}) \end{aligned}$$

In particular, we can conclude that

$$H_G(v) = |v|^2 - 2v_n \nabla \varphi(x') \cdot v' + (\nabla \varphi(x') \cdot v')^2,$$

so H_G is represented by the $n \times n$ -matrix

$$H_G = \begin{pmatrix} 1_{n-1} + \nabla \varphi(x') \nabla \varphi(x')^T & -\nabla \varphi(x') \\ -\nabla \varphi(x')^T & 1 \end{pmatrix}.$$

Therefore, in the same basis we have that

$$g_G = H_G^{-1} = \begin{pmatrix} 1_{n-1} & \nabla \varphi(x') \\ \nabla \varphi(x')^T & 1 + |\nabla \varphi(x')|^2 \end{pmatrix}.$$

We note that $H_G|_{T\partial X}$ is the Riemannian metric on ∂X induced from the Euclidean metric on \mathbb{R}^n and the inclusion $\partial X \hookrightarrow \mathbb{R}^n$. We also conclude that

$$(15) \quad C_G^j(x; v) = v_n C_G^{j,1}(x; v') + C_G^{j,0}(x; v')$$

where

$$C_G^{j,1}(x; v') := -2\nabla^{j-1} \varphi(x'; v') \quad \text{and} \quad C_G^{j,0}(x; v') := \sum_{k+l=j, k, l > 0} \nabla^k \varphi(x'; v) \nabla^l \varphi(x'; v).$$

Therefore, in these coordinates near the boundary of a domain in \mathbb{R}^n , we have for $\gamma \in I_j$ that

$$C_G^{(\gamma)}(x, -D_\xi) = (-1)^{|\gamma|} \prod_{l=1}^{\text{rk}(\gamma)} \left[\sum_{|\alpha'|=|\gamma_l|-1} \frac{-2\partial_{x'}^{\alpha'} \varphi(x')}{\alpha'!} D_{\xi'}^{\alpha'} D_{\xi_n} + \sum_{\substack{|\alpha'|+|\beta'|=|\gamma_l|, \\ |\alpha'|, |\beta'| > 0}} \frac{\partial_{x'}^{\alpha'} \varphi(x') \partial_{x'}^{\beta'} \varphi(x')}{\alpha'! \beta'!} D_{\xi'}^{\alpha'+\beta'} \right].$$

This gives a method for computing the homogeneous symbol q_j of degree $-n-1-j$ for any j following Theorem 2.9.

The principal symbol is computed as in Theorem 2.9. By Proposition 2.14 we compute for $n > 1$ that

$$\begin{aligned} q_1(x, \xi, R) &= -i(n^2 - 1) \mathbf{e}_{1,n} \left(6g_G(\nabla^2 \varphi) g_G(\xi, \nabla \varphi - e_n) (R^2 + g_G(\xi, \xi))^{-(n+1)/2-1} - \right. \\ &\quad \left. - 2(n+3) \nabla^2 \varphi(\iota_\xi g_G, \iota_\xi g_G) g_G(\xi, \nabla \varphi - e_n) (R^2 + g_G(\xi, \xi))^{-(n+1)/2-2} \right) = \\ &= i(n^2 - 1) \mathbf{e}_{1,n} \left(6\xi_n g_G(\nabla^2 \varphi) (R^2 + g_G(\xi, \xi))^{-(n+1)/2-1} - \right. \\ &\quad \left. - 2(n+3) \xi_n \nabla^2 \varphi((\xi' + \xi_n \nabla \varphi)^{\otimes 2}) (R^2 + g_G(\xi, \xi))^{-(n+1)/2-2} \right), \end{aligned}$$

By Proposition 2.15 we compute for $n \neq 1, 3$ that

$$\begin{aligned}
q_2(x, \xi, R) &= -24c_{2,n}(n+3)_{4,-2}4(g_G(\nabla^2\varphi))^2(g_G(\xi, \nabla\varphi - e_n))^2(R^2 + g_G(\xi, \xi))^{-(n+1)/2-2}+ \\
&\quad + 6c_{2,n}(n+5)_{5,-2}4g_G(\nabla^2\varphi)(g_G(\xi, \nabla\varphi - e_n))^2\nabla^2\varphi(\iota_\xi g_G, \iota_\xi g_G)(R^2 + g_G(\xi, \xi))^{-(n+1)/2-3}- \\
&\quad - c_{2,n}(n+7)_{6,-2}4(\nabla^2\varphi(\iota_\xi g_G, \iota_\xi g_G))^2(g_G(\xi, \nabla\varphi - e_n))^2(R^2 + g_G(\xi, \xi))^{-(n+1)/2-4}+ \\
&\quad + 3c_{2,n}(n+5)_{5,-2} \cdot 4(g_G(\nabla^2\varphi))^2g_G(\nabla\varphi - e_n, \nabla\varphi - e_n)(R^2 + g_G(\xi, \xi))^{-(n+1)/2-1}+ \\
&\quad + 3c_{1,n}(n^2 - 1)\left(2(g_G \otimes g_G)(\nabla^3\varphi \otimes (\nabla\varphi - e_n)) + (g_G(\nabla^2\varphi))^2\right)(R^2 + g_G(\xi, \xi))^{-(n+1)/2-1}- \\
&\quad - 6c_{1,n}(n+3)_{3,-2}\left(2(g_G \otimes \iota_\xi g_G)(\nabla^3\varphi)g_G(\xi, \nabla\varphi - e_n) + \right. \\
&\quad \quad \left. + (g_G(\nabla^2\varphi))(\nabla^2\varphi(\iota_\xi g_G, \iota_\xi g_G))\right)(R^2 + g_G(\xi, \xi))^{-(n+1)/2-2}+ \\
&\quad + c_{1,n}(n+5)_{4,-2}\left(2(\iota_\xi g_G)^{\otimes 3}(\nabla^3\varphi)g_G(\xi, \nabla\varphi - e_n) + \right. \\
&\quad \quad \left. + (\nabla^2\varphi(\iota_\xi g_G, \iota_\xi g_G))^2\right)(R^2 + g_G(\xi, \xi))^{-(n+1)/2-3} = \\
&= -48c_{2,n}(n+3)_{4,-2}(g_G(\nabla^2\varphi))^2\xi_n^2(R^2 + g_G(\xi, \xi))^{-(n+1)/2-2}+ \\
&\quad + 24c_{2,n}(n+5)_{5,-2}g_G(\nabla^2\varphi)\xi_n^2\nabla^2\varphi((\xi' + \xi_n\nabla\varphi)^{\otimes 2})(R^2 + g_G(\xi, \xi))^{-(n+1)/2-3}- \\
&\quad - 4c_{2,n}(n+7)_{6,-2}(\nabla^2\varphi((\xi' + \xi_n\nabla\varphi)^{\otimes 2}))^2\xi_n^2(R^2 + g_G(\xi, \xi))^{-(n+1)/2-4}+ \\
&\quad + (12c_{2,n}(n+5)_{5,-2} + 3c_{1,n}(n^2 - 1))(g_G(\nabla^2\varphi))^2(R^2 + g_G(\xi, \xi))^{-(n+1)/2-1}+ \\
&\quad - 6c_{1,n}(n+3)_{3,-2}\left(-2\xi_n\nabla^3\varphi(1_{n-1} \otimes (\xi' + \xi_n\nabla\varphi)) + \right. \\
&\quad \quad \left. + (g_G(\nabla^2\varphi))(\nabla^2\varphi((\xi' + \xi_n\nabla\varphi)^{\otimes 2}))\right)(R^2 + g_G(\xi, \xi))^{-(n+1)/2-2}+ \\
&\quad + c_{1,n}(n+5)_{4,-2}\left(-2\xi_n(\nabla^3\varphi)((\xi' + \xi_n\nabla\varphi)^{\otimes 3}) + \right. \\
&\quad \quad \left. + (\nabla^2\varphi((\xi' + \xi_n\nabla\varphi)^{\otimes 2}))^2\right)(R^2 + g_G(\xi, \xi))^{-(n+1)/2-3},
\end{aligned}$$

and for $n = 3$ that

$$\begin{aligned}
q_2(x, \xi, R) &= (c_{2,3}240 \cdot 4 + c_{1,3}24)(g_G(\nabla^2\varphi))^2(R^2 + g_G(\xi, \xi))^{-3}- \\
&\quad - c_{2,3}45 \cdot 96 \cdot 4(g_G(\nabla^2\varphi))^2\xi_n^2(R^2 + g_G(\xi, \xi))^{-4}+ \\
&\quad + c_{2,3}120 \cdot 96 \cdot 4g_G(\nabla^2\varphi)\xi_n^2\nabla^2\varphi((\xi' + \xi_n\nabla\varphi)^{\otimes 2})(R^2 + g_G(\xi, \xi))^{-5}- \\
&\quad - c_{2,3}80 \cdot 96 \cdot 4(g_G(\nabla^2\varphi))^2\xi_n^2(R^2 + g_G(\xi, \xi))^{-6}+ \\
&\quad - c_{1,3}288\left(-2\xi_n(\nabla^3\varphi)(1_{n-1} \otimes (\xi' + \xi_n\nabla\varphi)) + \right. \\
&\quad \quad \left. (g_G(\nabla^2\varphi))(\nabla^2\varphi((\xi' + \xi_n\nabla\varphi)^{\otimes 2}))\right)(R^2 + g_G(\xi, \xi))^{-4}+ \\
&\quad + c_{1,3}192\left(-2\xi_n(\nabla^3\varphi)((\xi' + \xi_n\nabla\varphi)^{\otimes 3}) + (\nabla^2\varphi((\xi' + \xi_n\nabla\varphi)^{\otimes 2}))^2\right)(R^2 + g_G(\xi, \xi))^{-5}.
\end{aligned}$$

We note that $g_G(\nabla^2\varphi)(x_0)$ is $(n-1)/2$ times the mean curvature in x_0 .

Example 2.17 (Symbol computations for a submanifold of Euclidean space). We consider a submanifold $M \subseteq \mathbb{R}^N$ and the function $G(x, y) = |x - y|^2$ which is regular at the diagonal. This example fits into the bigger picture of the paper seeing that $G(x, y) = d(x, y)^2$ where d is the distance function on M making the inclusion $M \subseteq \mathbb{R}^N$ isometric. To Taylor expand G as in (10), we take coordinates around a point $x_0 \in M$ such that M near x_0 is parametrized by

$$\begin{cases} x_l = x_l, & l = 1, \dots, n \\ x_l = \varphi_l(x_1, \dots, x_n), & l = n+1, \dots, N \end{cases}$$

for some functions $\varphi_{n+1}, \dots, \varphi_N$. Write $x = (x_1, \dots, x_n)$. In these coordinates,

$$\begin{aligned} G(x, y) &= |x - y|^2 + \sum_{l=n+1}^N |\varphi_l(x) - \varphi_l(y)|^2 = \\ &= |v|^2 + \sum_{j=2}^{N_0} \sum_{l=n+1}^N \sum_{\substack{|\alpha|+|\beta|=j, \\ |\alpha|, |\beta| > 0}} \frac{\partial_x^\alpha \varphi_l(x) \partial_x^\beta \varphi_l(x)}{\alpha! \beta!} v^{\alpha+\beta} + O(|v|^{N_0+1}), \end{aligned}$$

where $v = x - y$. Therefore, we conclude that

$$H_G(v) = |v|^2 + \sum_{l=n+1}^N (\nabla \varphi_l(x) \cdot v)^2,$$

and for $j > 2$,

$$C_G^j(x, v) = \sum_{l=n+1}^N \sum_{\substack{|\alpha|+|\beta|=j, \\ |\alpha|, |\beta| > 0}} \frac{\partial_x^\alpha \varphi_l(x) \partial_x^\beta \varphi_l(x)}{\alpha! \beta!} v^{\alpha+\beta} = \sum_{l=n+1}^N \sum_{\substack{i+k=j, \\ i, k > 0}} (\nabla^i \varphi_l \otimes \nabla^k \varphi_l)(v).$$

Computations similar to those in Example 2.16 can be carried out also for submanifolds. To preserve the reader's sanity, we spare the details.

Example 2.18 (Symbol computations for geodesic distances). Consider a manifold M equipped with a Riemannian metric g_M . To avoid having to prescribe the distance between different components, we assume that M is connected. The geodesic distance $d_{\text{geo}} : M \times M \rightarrow [0, \infty)$ is defined by

$$d_{\text{geo}}(x, y) := \inf \{L(c) : c \text{ is a smooth path from } x \text{ to } y\},$$

where the length $L(c)$ of a path $c : [0, 1] \rightarrow M$ is defined by

$$L(c) := \int_0^1 \sqrt{g_{M, c(t)}(\dot{c}(t), \dot{c}(t))} dt.$$

Here we write $\dot{c} : [0, 1] \rightarrow TM$ for the derivative of the path, so $\dot{c}(t) \in T_{c(t)}M$. For a suitable neighborhood $U \subseteq TM$ of the zero section, the Riemannian metric defines an exponential map $\exp : U \rightarrow M$. More precisely, for a small enough $v \in T_x M$, $\exp_x(v) = \exp(x, v) \in M$ is defined in local coordinates as $\exp_x(v) = w_x(v; 1)$ where $w_x(v; \cdot) : [0, 1] \rightarrow TM$ is the solution to the second order ordinary differential equation

$$(16) \quad \begin{cases} \ddot{w}_x(v, t) + \Gamma_{w_x(v; t)}(\dot{w}_x(v, t), \dot{w}_x(v, t)) = 0, \\ w(v; 0) = x, \\ \dot{w}(v, 0) = -v, \end{cases}$$

where Γ is the affine connection defined from g , which in local coordinates is a vector valued symmetric bilinear form on the tangent bundle. In these local coordinates, for x and y close enough we have that

$$d_{\text{geo}}(x, y) = |\exp_x^{-1}(y)|_g^2.$$

In particular, d_{geo}^2 is smooth in a neighborhood of the diagonal.

Let us compute the Taylor expansion of d_{geo}^2 as in Equation (10) and prove that d_{geo}^2 is regular at the diagonal. We use a coordinate neighborhood as above, and write $v = x - y$. We are looking for the Taylor expansion in v of the coordinate function $X_x(v) = \exp_x^{-1}(x - v)$. We Taylor expand

$$w_x(v, t) = x - vt + \sum_{k=2}^N \frac{w_x^{(k)}(v; 0)}{k!} t^k + O(|tv|^{N+1}).$$

We note that $v \mapsto w_x^{(k)}(v; 0)$ is a homogeneous polynomial of degree k , we denote this by $W_k(x; v)$. It follows from Equation (16) that

$$(17) \quad \begin{cases} W_1(v) = -v, \\ W_2(v) = -\Gamma(v, v), \\ W_3(v) = -d\Gamma(v, v, v) - 2\Gamma(v, \Gamma(v, v)). \end{cases}$$

The higher order terms W_4, W_5, \dots can be computed inductively from Equation (16). Write the Taylor expansion of the unknown function X_x in v as

$$X_x(v) = \sum_{k=1}^N X^{(k)}(x; v) + O(|v|^{N+1}),$$

where $X^{(k)}(x; \cdot)$ is a homogeneous polynomial of degree k in v . The identity $X_x(v) = \exp_x^{-1}(x-v)$ is equivalent to $\exp_x(X_x(v)) = x - v$ which implies that

$$(18) \quad -\sum_{k=1}^N X^{(k)}(x; v) + \sum_{k=2}^N \frac{W_k(x, \sum_{l=1}^N X^{(l)}(x; v))}{k!} = -v + O(|v|^{N+1}).$$

Considering the first order term, we see that $X^{(1)}(v) = v$. The higher order terms can be inductively determined by considering each homogeneous term separately:

$$(19) \quad X^{(k)}(v) = \left[\sum_{j=2}^k \frac{W_j(x, \sum_{l=1}^{k-1} X^{(l)}(x; v))}{j!} \right]_{(k)},$$

where $[\cdot]_{(k)}$ denotes the homogeneous term of degree k . Using Equation (17) and (18), we compute the first terms to be

$$\begin{cases} X^{(1)}(v) = v, \\ X^{(2)}(v) = \frac{1}{2}W_2(x; v) = -\frac{1}{2}\Gamma(v, v), \\ X^{(3)}(v) = \frac{1}{6}W_3(x; v) + \frac{1}{2} [W_2(x; v - \frac{1}{2}\Gamma(v, v))]_{(3)} = \\ = -\frac{1}{6}d\Gamma(v, v, v) + \frac{1}{6}\Gamma(v, \Gamma(v, v)). \end{cases}$$

We summarize these computations in a proposition.

Proposition 2.19. *Let M be a manifold equipped with a Riemannian metric g_M and let d_{geo} denote the geodesic distance. Then d_{geo}^2 is regular at the diagonal and in a coordinate neighborhood the Taylor expansion as in Equation (10) takes the form*

$$d_{\text{geo}}^2(x, y) = |v|_{g_M}^2 + C_{d_{\text{geo}}^2}^3(x; v) + C_{d_{\text{geo}}^2}^4(x; v) + O(|v|_{g_M}^5),$$

where

$$\begin{aligned} C_{d_{\text{geo}}^2}^3(x; v) &= -g_M(v, \Gamma(v, v)), \\ C_{d_{\text{geo}}^2}^4(x; v) &= \frac{1}{4}|\Gamma(v, v)|_{g_M}^2 + \frac{1}{3}(g(v, d\Gamma(v, v, v)) - g(v, \Gamma(v, \Gamma(v, v)))). \end{aligned}$$

The higher order terms $C_{d_{\text{geo}}^2}^j$ can be computed inductively from Equation (19).

Let us return to the general case and describe the overall structure of the terms appearing in the full symbol expansion of $Q_{G, \chi}$.

Lemma 2.20. *Let M and G be as in Theorem 2.9. The entries in the full symbol expansion $q \sim \sum_{j=0}^{\infty} q_j$ of $Q_{G, \chi} \in \Psi_{\text{cl}}^{-n-1}(M; \mathbb{C}_+)$ takes the form*

$$q_j(x, \xi, R) = \sum_{k=0}^{3j} P_{k,j}(x, \xi) (R^2 + g_G(\xi, \xi))^{-\frac{n+1+j+k}{2}},$$

where $P_{k,j}$ are of the form

(1) $P_{k,j}$ is a homogeneous polynomial of degree k in ξ and

$$P_{k,j} \equiv 0 \quad \text{if } j - k \notin 2\mathbb{Z}.$$

(2) If $j - k \in 2\mathbb{Z}$, the polynomial $P_{k,j}$ takes the form

$$P_{k,j}(x, \xi) = \sum_{\gamma \in I_j} \rho_{\gamma,k,j} C_G^{(\gamma)}(x, \underbrace{\iota_\xi g_G \otimes \iota_\xi g_G}_{k \text{ times}} \otimes \underbrace{g_G \otimes \cdots \otimes g_G}_{\frac{|\gamma|-k}{2} \text{ times}}),$$

for some coefficients $\rho_{\gamma,k,j} \in \mathbb{Q}\pi^{n/2} + \mathbb{Q}\pi^{(n+1)/2} + \mathbb{Q}\pi^{(n-1)/2}$.

In the last item, we note that if $j - k \in 2\mathbb{Z}$, then $|\gamma| - k \in 2\mathbb{Z}$ for all $\gamma \in I_j$.

Proof. It follows from the computations in Theorem 2.9 that q_j can be written as a finite sum

$$q_j(x, \xi, R) = \sum_{k \geq 0} P_{k,j}(x, \xi) (R^2 + g_G(\xi, \xi))^{-\frac{n+1+j+k}{2}},$$

where $P_{k,j}$ is a polynomial in ξ whose coefficients (as a polynomial in ξ) are all polynomials in the Taylor coefficients of G at the diagonal. For the degrees to match, $P_{k,j}$ must be of degree k . Since the powers $R^2 + g_G(\xi, \xi)$ must differ from $-(n+1)/2$ by an integer, $P_{k,j} = 0$ unless $j - k \notin 2\mathbb{Z}$. Finally, the largest possible degree k for which $\xi^\alpha (R^2 + g_G(\xi, \xi))^{-\frac{n+1+j+k}{2}}$ (with $|\alpha| = k$) can be a summand in q_j is if the derivative $C_G^{(\gamma)}(x, D_\xi)$ acts only on $(R^2 + g_G(\xi, \xi))^{-(n+1)/2 + \text{rk}(\gamma)}$ (or $(R^2 + g_G(\xi, \xi))^{-(n+1)/2 + \text{rk}(\gamma)} \log(R^2 + g_G(\xi, \xi))$ if $2\text{rk}(\gamma) - n - 1 \in 2\mathbb{N}$) and in that case $\alpha = \gamma$, so the maximal degree of $P_{k,j}$ is the size of the largest index in I_j , i.e. $3j$. This proves item (1). To prove item (2), one notices that by Theorem 2.9, all possible entries in q_j consists of terms of the form

$$C_G^{(\gamma)}(x, \underbrace{\iota_\xi g_G \otimes \iota_\xi g_G}_{k \text{ times}} \otimes \underbrace{g_G \otimes \cdots \otimes g_G}_{\frac{|\gamma|-k}{2} \text{ times}}) (R^2 + g_G(\xi, \xi))^{-\frac{n+1+j+k}{2}},$$

with a coefficient being a rational number times either $\pi^{n/2}$, $\pi^{(n+1)/2}$ or $\pi^{(n-1)/2}$. \square

2.3. The symbol structure of the parametrix. The operator $Q_{G,\chi} \in \Psi_{\text{cl}}^{-n-1}(M; \mathbb{C}_+)$ considered in the previous subsection is elliptic with parameter by Theorem 2.9. In particular, it admits a parametrix $A_{G,\chi} \in \Psi_{\text{cl}}^{n+1}(M; \mathbb{C}_+)$, that is an operator with parameter so that $A_{G,\chi} Q_{G,\chi} - 1, Q_{G,\chi} A_{G,\chi} - 1 \in \Psi_{\text{cl}}^{-\infty}(M; \mathbb{C}_+)$.

Theorem 2.21. *Let M be an n -dimensional manifold and $G : M \times M \rightarrow [0, \infty)$ a function which is regular at the diagonal. The full symbol a in local coordinates of the parametrix $A_{G,\chi} \in \Psi_{\text{cl}}^{n+1}(M; \mathbb{C}_+)$ of $Q_{G,\chi} \in \Psi_{\text{cl}}^{-n-1}(M; \mathbb{C}_+)$ has an asymptotic expansion $a \sim \sum_{j=0}^{\infty} a_j$ where a_j is constructed inductively from the symbol expansion $q \sim \sum_j q_j$ of Theorem 2.9 by*

$$a_j = -a_0 \sum_{k+l+|\alpha|=j, l < j} \frac{1}{\alpha!} \partial_\xi^\alpha q_k D_x^\alpha a_l.$$

For $n = 1$ the first two terms are given by

$$\begin{aligned} a_0(x, \xi, R) &= \frac{1}{2} (R^2 + g_G(\xi, \xi)) \\ a_1(x, \xi, R) &= -\frac{i}{2} (\partial_\xi g_G)(\partial_x g_G) (R^2 + g_G(\xi, \xi))^{-1} + \frac{3i}{2} C_G^3(x, g_G \otimes \iota_\xi g_G) - \\ &\quad - 2i C_G^3(x, \iota_\xi g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G) (R^2 + g_G(\xi, \xi))^{-1} \end{aligned}$$

and for $n > 1$

$$\begin{aligned} a_0(x, \xi, R) &= (q_0(x, \xi, R))^{-1} = \frac{1}{n! \omega_n} (R^2 + g_G(\xi, \xi))^{(n+1)/2}, \\ a_1(x, \xi, R) &= -\frac{(n+1)^2 i}{n! \omega_n} g_G(d_x g_G(\xi, \xi), \xi) (R^2 + g_G(\xi, \xi))^{(n+1)/2-2} + \\ &\quad + \frac{3i \mathfrak{c}_{1,n}(n^2-1)}{(n!)^2 \omega_n^2} C_G^3(x, g_G \otimes \iota_\xi g_G) (R^2 + g_G(\xi, \xi))^{(n+1)/2-1} - \\ &\quad - \frac{i \mathfrak{c}_{1,n}(n+3)_{3,-2}}{(n!)^2 \omega_n^2} C_G^3(x, \iota_\xi g_G \otimes \iota_\xi g_G \otimes \iota_\xi g_G) (R^2 + g_G(\xi, \xi))^{(n+1)/2-2}. \end{aligned}$$

The homogeneous terms a_j each takes the form

$$a_j(x, \xi, R) = \sum_{k=0}^{3j} \tilde{P}_{k,j}(x, \xi) (R^2 + g_G(\xi, \xi))^{\frac{n+1-j-k}{2}},$$

where $\tilde{P}_{k,j}$ are homogeneous polynomials of degree k in ξ and

$$\tilde{P}_{k,j} \equiv 0 \quad \text{if } j - k \notin 2\mathbb{Z}.$$

The coefficients of $\tilde{P}_{k,j}$ as a polynomial in ξ are all polynomials in derivatives of the Taylor coefficients $(C_G^{(\gamma)})_{\gamma \in \cup_{k \leq j} I_k}$ of G at the diagonal (from Equation (10)) of total degree j .

In the structural statement at the end of the proposition, the total degree refers to the degree of the polynomial when counting an order i derivative of a Taylor coefficient $C_G^{(\gamma)}$ for $\gamma \in I_k$ to have degree $k + i$.

Proof. It follows from the parametrix construction for elliptic operators (see e.g. [38, Chapter I.5]) that $a \sim \sum_{j=0}^{\infty} a_j$ where $a_0 = q_0^{-1}$ and $a_j := -a_0 \sum_{k+l+|\alpha|=j, l < j} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} q_k D_x^{\alpha} a_l$ for $j > 0$.

Let us prove the structural statement about a_j by induction on j . It is clear for $j = 0$. Assume that the structural statement holds for $l < j + 1$. We have that

$$\begin{aligned} a_{j+1} &= -a_0 \sum_{k+l+|\alpha|=j+1, l < j+1} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} q_k D_x^{\alpha} a_l = \\ &= -\frac{1}{n! \omega_n} (R^2 + g_G(\xi, \xi))^{(n+1)/2} \sum_{k+l+|\alpha|=j+1, l < j+1} \sum_{i_1=0}^{N(k)} \sum_{i_2=0}^{\tilde{N}(l)} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} \left[P_{i_1, k}(x, \xi) (R^2 + g_G(\xi, \xi))^{-\frac{n+1+k+i_1}{2}} \right] \\ &\quad \cdot D_x^{\alpha} \left[\tilde{P}_{i_2, l}(x, \xi) (R^2 + g_G(\xi, \xi))^{\frac{n+1-l-i_2}{2}} \right], \end{aligned}$$

which proves that a_{j+1} has the claimed structure.

What remains is to compute a_1 . We write

$$\begin{aligned} a_1(x, \xi, R) &= -a_0 \sum_{|\alpha|=1} \partial_{\xi}^{\alpha} q_0 D_x^{\alpha} a_0 - a_0^2 q_1 = \\ &= -\frac{1}{n! \omega_n} (R^2 + g_G(\xi, \xi))^{(n+1)/2} \sum_{|\alpha|=1} \partial_{\xi}^{\alpha} (R^2 + g_G(\xi, \xi))^{-(n+1)/2} D_x^{\alpha} (R^2 + g_G(\xi, \xi))^{(n+1)/2} \\ &\quad + \frac{i(n^2 - 1) \mathfrak{c}_{1, n}}{(n!)^2 \omega_n^2} (R^2 + g_G(\xi, \xi))^{n+1} \left(3C_G^3(x, g_G \otimes \iota_{\xi} g_G) (R^2 + g_G(\xi, \xi))^{-(n+1)/2-1} - \right. \\ &\quad \left. - (n+3) C_G^3(x, \iota_{\xi} g_G \otimes \iota_{\xi} g_G \otimes \iota_{\xi} g_G) (R^2 + g_G(\xi, \xi))^{-(n+1)/2-2} \right) = \\ &= -\frac{(n+1)^2 i}{n! \omega_n} g_G(d_x g_G(\xi, \xi), \xi) (R^2 + g_G(\xi, \xi))^{(n+1)/2-2} + \\ &\quad + \frac{3i \mathfrak{c}_{1, n} (n^2 - 1)}{(n!)^2 \omega_n^2} C_G^3(x, g_G \otimes \iota_{\xi} g_G) (R^2 + g_G(\xi, \xi))^{(n+1)/2-1} - \\ &\quad - \frac{i \mathfrak{c}_{1, n} (n+3)_{3, -2}}{(n!)^2 \omega_n^2} C_G^3(x, \iota_{\xi} g_G \otimes \iota_{\xi} g_G \otimes \iota_{\xi} g_G) (R^2 + g_G(\xi, \xi))^{(n+1)/2-2} \end{aligned}$$

□

We use the notation $\Gamma_{\alpha}(R_0) := \{z \in \mathbb{C} : |\text{Arg}(z)| < \alpha, \text{Re}(z) > R_0\}$.

Corollary 2.22. *Let M be a compact n -dimensional manifold and $G : M \times M \rightarrow [0, \infty)$ a function which is regular at the diagonal. For some $R_0 > 0$, the operator $Q_{G, \chi}(R) \in \Psi_{\text{cl}}^{-n-1}(M)$ is invertible as an operator $H^{-(n+1)/2}(M) \rightarrow H^{(n+1)/2}(M)$ for any $R \in \Gamma_{\pi/(n+1)}(R_0)$ and*

$$Q_{G, \chi}^{-1} - A_{G, \chi} \in \Psi_{\text{cl}}^{-\infty}(M; \Gamma_{\pi/(n+1)}(R_0)).$$

In particular, $Q_{G, \chi}^{-1}$ is a pseudodifferential with parameter in $\Gamma_{\pi/(n+1)}(R_0)$ whose full symbol in local coordinates has an asymptotic expansion $a \sim \sum_{j=0}^{\infty} a_j$ where a_j is as in Theorem 2.21.

Corollary 2.22 follows from Theorem 2.9 and 2.21 using standard techniques for pseudodifferential with parameter. For the convenience of the reader, we include its proof.

Proof. Let Δ denote a positive Laplace operator on M whose principal symbol coincides with the metric g_G dual to the transversal Hessian of G . It follows from Theorem 2.9 that $\bar{\sigma}_{-n-1}((R^2 + \Delta)^{-\mu}) = \bar{\sigma}_{-n-1}(Q)$. Hence, the operator $\mathfrak{r}(R) := 1 - (R^2 + \Delta)^{\mu/2}Q(R^2 + \Delta)^{\mu/2}$ is a parameter-dependent pseudodifferential operator of order -1 . We write

$$Q_{G,\chi} = (R^2 + \Delta)^{-\mu/2}(1 - \mathfrak{r})(R^2 + \Delta)^{-\mu/2}.$$

The order of \mathfrak{r} is -1 , so by Theorem B.11 (see page 63) we have that $\|\mathfrak{r}(R)\|_{L^2(M) \rightarrow L^2(M)} = O(\operatorname{Re}(R)^{-1})$ as $\operatorname{Re}(R) \rightarrow \infty$. We conclude that there exists an $R_0 > 1$ such that $\|\mathfrak{r}(R)\|_{L^2(M) \rightarrow L^2(M)} \leq \frac{1}{2}$ for $R \in \Gamma_{\pi/(n+1)}(R_0)$. Since $(R^2 + \Delta)^{-\mu/2} : H^s(M) \rightarrow H^{s+\mu}(M)$ is invertible for $R \in \Gamma_{\pi/(n+1)}(0)$ and any $s \in \mathbb{R}$, we can for $R \in \Gamma_{\pi/(n+1)}(R_0)$ invert $Q_{G,\chi}$ as the absolutely convergent series of operators $H^\mu(M) \rightarrow H^{-\mu}(M)$ given by

$$Q_{G,\chi}^{-1} = \sum_{k=0}^{\infty} (R^2 + \Delta)^{\mu/2} \mathfrak{r}(R)^k (R^2 + \Delta)^{\mu/2}.$$

It remains to prove that $Q_{G,\chi}^{-1} - A_{G,\chi} \in \Psi_{\text{cl}}^{-\infty}(M; \Gamma_{\pi/(n+1)}(R_0))$. By the construction above, $Q_{G,\chi}^{-1} \in \Psi_{\text{cl}}^{n+1}(M; \Gamma_{\pi/(n+1)}(R_0))$. Therefore the classes $[Q_{G,\chi}^{-1}]$ and $[A_{G,\chi}]$ in the formal symbol algebra $\Psi_{\text{cl}}^{n+1}(M; \Gamma)/\Psi_{\text{cl}}^{-\infty}(M; \Gamma_{\pi/(n+1)}(R_0))$ are both inverses to

$$[Q_{G,\chi}] \in \cup_{k \in \mathbb{Z}} \Psi_{\text{cl}}^k(M; \Gamma_{\pi/(n+1)}(R_0))/\Psi_{\text{cl}}^{-\infty}(M; \Gamma_{\pi/(n+1)}(R_0)).$$

By the uniqueness of inverses, $[Q_{G,\chi}^{-1}] = [A_{G,\chi}] \in \Psi_{\text{cl}}^{n+1}(M; \Gamma_{\pi/(n+1)}(R_0))/\Psi_{\text{cl}}^{-\infty}(M; \Gamma_{\pi/(n+1)}(R_0))$. \square

2.4. Analytic results for Q on compact manifolds. The results of the previous subsections have analytic implications in the case that M is a compact manifold. We consider the scale of Hilbert spaces $H_R^s(M) := (R^2 + \Delta)^{-s/2}L^2(M)$ defined for $R \in \mathbb{R} \setminus \{0\}$ and $s \in \mathbb{R}$ with the Hilbert space structure making $(R^2 + \Delta)^{-s/2} : L^2(M) \rightarrow H_R^s(M)$ unitary. Here Δ could be any choice of Laplacian, but for the sake of simplicity we fix the Laplacian associated with the Riemannian metric associated with a function regular at the diagonal. By elliptic regularity, $H^s(M) = H_R^s(M)$ as vector spaces with equivalent norms independently of the choice of Laplacian, but the Hilbert space structure differs in a non-uniform way as R varies. At this stage, we shall start to concern ourselves with extensions of Q to the complex numbers, so we phrase our results in terms of the operator

$$(20) \quad Q(R)f(x) := \frac{1}{R} \int_M \chi(x, y) e^{-Rd(x, y)} f(y) dy,$$

where d is a distance function whose square is regular at the diagonal and χ a function being 1 near the diagonal such that d^2 is smooth on the support of χ . We note that $Q(R) = Q_{d^2, \chi}(R)$ for $\operatorname{Re}(R) > 0$.

Theorem 2.23. *Let M be a compact n -dimensional manifold and $d : M \times M \rightarrow [0, \infty)$ a distance function whose square is regular at the diagonal. Set $\mu := (n + 1)/2$. The operator*

$$Q(R) : H^{-\mu}(M) \rightarrow H^\mu(M),$$

defined from the expression (20) is a well defined Fredholm operator for all $R \in \mathbb{C} \setminus \{0\}$ and there is an R_0 such that $Q(R)$ is invertible for all $R \in \Gamma_{\pi/(n+1)}(R_0)$. Moreover, the following holds:

- a) *For each $R \in \mathbb{C} \setminus \{0\}$, $Q(R) \in \Psi_{\text{cl}}^{-n-1}(M)$ is an elliptic pseudodifferential operator and the family of operators*

$$(Q(R) : H^{-\mu}(M) \rightarrow H^\mu(M))_{R \in \mathbb{C} \setminus \{0\}}$$

depends holomorphically on $R \in \mathbb{C} \setminus \{0\}$. Moreover, the holomorphic family

$$(Q(R)^{-1} : H^\mu(M) \rightarrow H^{-\mu}(M))_{R \in \Gamma_{\pi/(n+1)}(R_0)},$$

extends meromorphically to $R \in \mathbb{C} \setminus \{0\}$.

b) There is a $C > 0$ such that

$$C^{-1}\|f\|_{H_{|R|}^{-\mu}(M)}^2 \leq \operatorname{Re}\langle f, Q(R)f \rangle_{L^2} \leq C\|f\|_{H_{|R|}^{-\mu}(M)}^2,$$

for $R \in \Gamma_{\pi/(n+1)}(R_0)$ and $f \in H^{-\mu}(M)$. In particular, for $R \in \Gamma_{\pi/(n+1)}(R_0)$, $Q(R)$ is coercive in form sense on $L^2(M)$ and the sesquilinear form $\operatorname{Re}\langle \cdot, Q(R)\cdot \rangle$ is uniformly equivalent to the inner product of $H_{|R|}^{-\mu}(M)$.

Proof. The first statements of the theorem follows from Corollary 2.22.

Part a) follows from the meromorphic Fredholm theorem (see Appendix D) upon proving that $R \mapsto Q(R) \in \mathbb{B}(H^{-\mu}(M), H^{\mu}(M))$ is holomorphic with values in the set of Fredholm operators. We can write $Q(R)$ as the integral operator with Schwartz kernel

$$(21) \quad \frac{1}{R}\chi(x, y)e^{-Rd(x, y)} = \frac{\chi(x, y)}{R} + \sum_{k=0}^{\infty} \frac{R^k}{(k+1)!}\chi(x, y)d(x, y)^{k+1}.$$

Using the fact that d^2 is regular at the diagonal, an argument as in the proof of Theorem 2.9 shows that $Q(R)$ is an elliptic pseudodifferential operator of order $-n-1$. By Proposition A.7, the principal symbol of $Q(R)$ (for fixed R) is given by

$$\sigma_{-n-1}(Q(R))(x, \xi) = \pi^{(n-1)/2}2^n \Gamma\left(\frac{n+1}{2}\right) |\xi|_{g_G}^{-n-1} = n!\omega_n |\xi|_{g_{d^2}}^{-n-1}.$$

Therefore $R \mapsto Q(R) \in \mathbb{B}(H^{-\mu}(M), H^{\mu}(M))$ takes values in the set of Fredholm operators. The expression (21), and again an argument as in the proof of Theorem 2.9, shows that $R \mapsto Q(R) \in \mathbb{B}(H^{-\mu}(M), H^{\mu}(M))$ depends holomorphically on $R \in \mathbb{C} \setminus \{0\}$.

Part b) follows from the Gårding inequality (see Corollary B.12 on page 64) using that $\operatorname{Re}(Q)$ has positive principal symbol as a pseudodifferential operator with parameter by Theorem 2.9 on page 10). \square

2.5. Evaluation at $\xi = 0$ of some symbols. For later purposes, we will be interested in knowing the value of the homogeneous component of the full symbol of the parametrix of Q at $\xi = 0$, constructed as in Theorem 2.21. The following lemma shows that the evaluations of symbols of operators with parameter provides coordinate invariant expressions, therefore containing invariants of a pseudodifferential operator with parameter.

Lemma 2.24. *Assume that M is a manifold and that $A \in \Psi_{\text{cl}}^m(M; \Gamma)$ is a properly supported pseudodifferential operator with parameter. Then there exists a sequence $(a_{j,0})_{j \in \mathbb{N}} \subseteq C^\infty(M \times \Gamma)$ such that*

- i) Each $a_{j,0} = a_{j,0}(x, R)$ is homogeneous of degree $m-j$ in R .
- ii) In each local coordinate chart,

$$a_{j,0}(x, R) = a_j(x, 0, R),$$

where $a \sim \sum_j a_j$ is a homogeneous expansion of the full symbol of A in that chart.

Moreover, for any $N \in \mathbb{N}$, we have that

$$(22) \quad [A(R)1](x) = \sum_{j=0}^N a_j(x, R) + r_N(x, R) = \sum_{j=0}^N a_j(x, 1)R^{m-j} + r_N(x, R),$$

where $r_N \in C^\infty(M \times \Gamma)$ is a function such that for any compact $K \subseteq M$ it holds that

$$\sup_{x \in K} |\partial_x^\alpha \partial_R^k r_N(x, R)| = O(\operatorname{Re}(R)^{m-N+|\alpha|+k}), \quad \text{as } \operatorname{Re}(R) \rightarrow +\infty.$$

Proof. Choose a partition of unity $(\chi_j)_j \subseteq C_c^\infty(M)$ subordinate to a locally finite covering by coordinate charts, and choose $(\tilde{\chi}_j)_j \subseteq C_c^\infty(M)$ such that $\tilde{\chi}_j$ is supported in a coordinate chart and $\tilde{\chi}_j = 1$ on $\operatorname{supp}(\chi_j)$. We have that $\sum_j \chi_j A \tilde{\chi}_j$ converges in weak sense to a properly supported operator, and $A - \sum_j \chi_j A \tilde{\chi}_j \in \Psi_{\text{cl}}^{-\infty}(M; \Gamma)$. Therefore, we can assume that A is supported in a coordinate chart. In a coordinate chart, and a homogeneous expansion $a \sim \sum_j a_j$ of the full symbol of A , the method of stationary phase (see for instance [22, Chapter VII.7]) implies that $[A(R)1](x) = \sum_{j=1}^N a_j(x, 0, R) + r_N(x, R)$ as in Equation (22). It is clear that

that the function $A(R)1 \in C^\infty(M)$ and its asymptotic expansion is independent of choice of coordinates, and the lemma follows. \square

Lemma 2.25. *Let M be an n -dimensional manifold, $G : M \times M \rightarrow [0, \infty)$ be a function regular at the diagonal, and $(a_{j,0})_{j \in \mathbb{N}} \subseteq C^\infty(M \times \mathbb{C}_+)$ the sequence defined as in Lemma 2.24 from $A_{G,\chi} \in \Psi_{\text{cl}}^{n+1}(M; \mathbb{C}_+)$ (cf. Theorem 2.21). The functions $a_{j,0}$ are determined in local coordinates by the properties that*

$$a_{j,0}(x, R) = 0, \quad \text{for all } j \text{ odd,}$$

and for even j , $a_{j,0}(x, R)$ is determined inductively from $a_{0,0}(x, R) = \frac{1}{n!\omega_n} R^{n+1}$ and for $j > 0$,

$$(23) \quad a_{j,0}(x, R) = -\frac{1}{n!\omega_n} R^{n+1} \sum_{\substack{k+2l+p=j \\ 2l < j, 2|k+p}} i^p q_{k,p}(x, R) \cdot \nabla_x^p a_{j-2k,0}(x, R),$$

where $q_{k,p}(x, R)$ denotes the p -linear form

$$q_{k,p}(x, R) \cdot v := \sum_{|\alpha|=p} \partial_\xi^\alpha q_k(x, \xi, R) v^\alpha |_{\xi=0} = \partial_t^p q_k(x, tv, R) |_{t=0}.$$

Proof. By Lemma 2.24, we can perform all computations in a coordinate chart. The structural description of a_j from Lemma 2.21 implies that for any (x, ξ, R) and $j \in \mathbb{N}$ it holds that

$$a_j(x, -\xi, R) = (-1)^j a_j(x, \xi, R).$$

We conclude that $a_j(x, 0, R) = 0$, and even that $\nabla_x^p a_{j,0}(x, R) = 0$ for any p , when j is odd. To compute a_j for even j , we note that since $a_j = -a_0 \sum_{k+l+|\alpha|=j, l < j} \frac{1}{\alpha!} \partial_\xi^\alpha q_k D_x^\alpha a_l$, the formula (23) follows using that the only contributions are for even l , and evenness of j implies that $2|k+p$. \square

Remark 2.26. The formulas in Theorem 2.9 shows that for n odd, and j even,

$$\begin{aligned} q_{j,0}(x, R) = & R^{-n-1-j} \sum_{\gamma \in I_j, \text{rk}(\gamma) < (n+1)/2} \mathbf{c}_{\text{rk}(\gamma), n} (-1)^{\text{rk}(\gamma)-1} (|\gamma|/2)! (n+1-2\text{rk}(\gamma))_{|\gamma|/2, -2} C_G^{(\gamma)}(x, g_G^{\otimes |\gamma|/2}) + \\ & - R^{-n-1-j} \sum_{\gamma \in I_j, \text{rk}(\gamma) \geq (n+1)/2} \mathbf{c}_{\text{rk}(\gamma), n} (-1)^{\text{rk}(\gamma)-1} (|\gamma|/2)! (n+1-2\text{rk}(\gamma))_{|\gamma|/2-1, -2} C_G^{(\gamma)}(x, g_G^{\otimes |\gamma|/2}), \end{aligned}$$

and for n even, and j even,

$$q_{j,0}(x, R) = R^{-n-1-j} \sum_{\gamma \in I_j} \mathbf{c}_{\text{rk}(\gamma), n} (-1)^{\text{rk}(\gamma)-1} (|\gamma|/2)! (n+1-2\text{rk}(\gamma))_{|\gamma|/2, -2} C_G^{(\gamma)}(x, g_G^{\otimes |\gamma|/2}),$$

and $\mathbf{c}_{\text{rk}(\gamma), n}$ is as in Theorem 2.9. Note that by definition of the set I_j , $\gamma \in I_j$ satisfies that $|\gamma|$ is even if and only if j is even.

Theorem 2.27. *Let M be an n -dimensional manifold, $G : M \times M \rightarrow [0, \infty)$ be a function regular at the diagonal, and $(a_{j,0})_{j \in \mathbb{N}} \subseteq C^\infty(M \times \mathbb{C}_+)$ the restriction to 0 of the full symbol of $A_{G,\chi}$. Then for each $j > 0$, $a_{j,0}$ is a polynomial in $(C_G^{(\gamma)})_{\gamma \in \cup_{k \leq j} I_k}$ and its derivatives contracted by the metric g_G and its derivatives of total degree j where each $C_G^{(\gamma)}$, $\gamma \in I_k$, has degree k , the metric has degree zero and x -derivatives increase the order by 1.*

In the special case $j = 0$ we have

$$a_{0,0}(x, R) = \frac{1}{n!\omega_n} R^{n+1},$$

and for $j = 2$ and $n = 3$, we have that

$$a_{2,0}(x, R) = -\frac{24R^2}{(3!\omega_3)^2} \left(10\mathbf{c}_{1,3} C_G^4(x, g_G \otimes g_G) - \mathbf{c}_{2,3} (C_G^3 \otimes C_G^3)(x, g_G \otimes g_G \otimes g_G) \right),$$

while for $j = 2$ and $n \neq 1, 3$, we have that

$$(24) \quad a_{2,0}(x, R) = -\frac{3R^{n-1}}{(n!\omega_n)^2} \left(\mathbf{c}_{1,n} (n^2 - 1) C_G^4(x, g_G \otimes g_G) - \mathbf{c}_{2,n} (n+5)_{5,-2} (C_G^3 \otimes C_G^3)(x, g_G \otimes g_G \otimes g_G) \right),$$

and for $j = 4$ we have the identity

$$(25) \quad \begin{aligned} a_{4,0}(x, R) &= -\frac{R^{2n+2}}{(n!\omega_n)^2} q_{4,0}(x, R) - \frac{n+1}{R^2} g_G(\nabla_x^2 a_{2,0}(x, R)) - \\ &\quad - \frac{\mathbf{c}_{1,n}(n^2-1)}{R^2 n! \omega_n} (C_G^3 \otimes \nabla_x a_{2,0})(g_G \otimes g_G) - \frac{R^{n+1}}{n! \omega_n} q_{2,0}(x, R) a_{2,0}(x, R). \end{aligned}$$

Proof. The structural statement about $a_{j,0}(x, R)$ is readily deduced from Lemma 2.25 and induction by using the property that if $\gamma \in I_j$ and $\gamma' \in I_{j'}$ then $(\gamma, \gamma') \in I_{j+j'}$.

The equation for $a_{0,0}$ is immediate from Lemma 2.21. Using Lemma 2.25, we have that $a_{2,0}(x, R) = -a_{0,0}^2 q_{2,0}$ and the formula (24) follows from that

$$q_{2,0}(x, R) = 3\mathbf{c}_{1,n}(n^2-1)C_G^4(x, g_G \otimes g_G)R^{-n-3} + 3\mathbf{c}_{2,n}(n+5)_{5,-2}(C_G^3 \otimes C_G^3)(x, g_G \otimes g_G \otimes g_G)R^{-n-3},$$

for $n \neq 1, 3$ and a similar expression for $n = 3$.

Let us compute $a_{4,0}(x, R)$. By Lemma 2.25 we have that

$$\begin{aligned} a_{4,0}(x, R) &= -\frac{R^{n+1}}{n! \omega_n} \left(\sum_{k=0}^4 i^{-k} q_{k,4-k}(x, R) \cdot \nabla_x^{4-k} \frac{R^{n+1}}{n! \omega_n} - \sum_{k=0}^2 i^{-k} q_{k,2-k}(x, R) \cdot \nabla_x^{2-k} a_{2,0}(x, R) \right) = \\ &= \frac{R^{n+1}}{n! \omega_n} \left(q_{0,2}(x, R) \cdot \nabla_x^2 a_{2,0}(x, R) - i q_{1,1}(x, R) \cdot \nabla_x a_{2,0}(x, R) - q_{2,0}(x, R) a_{2,0}(x, R) \right) - \\ &\quad - \frac{R^{2n+2}}{(n! \omega_n)^2} q_{4,0}(x, R), \end{aligned}$$

and the computation is complete upon using Proposition 2.14. \square

Remark 2.28. The full expression for $a_{4,0}$ can be computed from Equation (25). We omit the full details, but let us note an expression for $q_{4,0}$. Since

$$I_4 = \{6, (3, 5), (4, 4), (5, 3), (3, 3, 4), (3, 4, 3), (4, 3, 3), (3, 3, 3, 3)\},$$

we have that

$$\begin{aligned} q_{4,0}(x, R) &= R^{-n-5} \mathbf{c}_{1,n} 3!(n-1)_{3,-2} C_G^6(x, g_G^{\otimes 3}) - R^{-n-5} \mathbf{c}_{2,n} 4!(n-3)_{4,-2} C_G^{(3,5)}(x, g_G^{\otimes 4}) - \\ &\quad - R^{-n-5} \mathbf{c}_{2,n} 4!(n-3)_{4,-2} C_G^{(4,4)}(x, g_G^{\otimes 4}) - R^{-n-5} \mathbf{c}_{2,n} 4!(n-3)_{4,-2} C_G^{(5,3)}(x, g_G^{\otimes 4}) + \\ &\quad + R^{-n-5} \mathbf{c}_{3,n} 5!(n-5)_{5,-2} C_G^{(3,3,4)}(x, g_G^{\otimes 5}) + R^{-n-5} \mathbf{c}_{3,n} 5!(n-5)_{5,-2} C_G^{(3,4,3)}(x, g_G^{\otimes 5}) + \\ &\quad + R^{-n-5} \mathbf{c}_{3,n} 5!(n-5)_{5,-2} C_G^{(4,3,3)}(x, g_G^{\otimes 5}) - R^{-n-5} \mathbf{c}_{4,n} 6!(n-7)_{6,-2} C_G^{(3,3,3,3)}(x, g_G^{\otimes 6}). \end{aligned}$$

Example 2.29 (Evaluations of symbols for domains in Euclidean space). Let us return to the computations on Euclidean space from Example 2.16. We consider $G(x, y) = |x - y|^2$ – the square of the Euclidean distance. Since $q(x, \xi, R) = n! \omega_n (R^2 + |\xi|^2)^{-(n+1)/2}$ is a full symbol of $Q_{G,\chi}$ in Euclidean coordinates, $a(x, \xi, R) = \frac{1}{n! \omega_n} (R^2 + |\xi|^2)^{(n+1)/2}$ is a full symbol of $A_{G,\chi}$. Therefore

$$(26) \quad a_{j,0}(x, R) = \begin{cases} \frac{1}{n! \omega_n} R^{n+1}, & j = 0, \\ 0, & j > 0. \end{cases}$$

By Lemma 2.25 this holds in any coordinate system on Euclidean space. We remark that the bulk of computations carried out in Example 2.16 will mainly be of interest when inverting Q near the boundary of a domain, while the computation (26) relates to interior terms.

Example 2.30 (Evaluations of symbols for submanifolds of Euclidean space). We return to submanifolds $M \subseteq \mathbb{R}^N$ and the function $G(x, y) = |x - y|^2$ which is regular at the diagonal as in Example 2.17 above. We take coordinates as in Example 2.17. By Theorem 2.27 we have that

$$\begin{aligned}
a_{0,0}(x, R) &= \frac{1}{n! \omega_n} R^{n+1} \text{ and} \\
a_{2,0}(x, R) &= -\frac{3\mathfrak{c}_{1,n}(n^2-1)}{(n! \omega_n)^2} \sum_{l=n+1}^N \sum_{\substack{i+k=4, \\ i,k>0}} (\nabla^i \varphi_l \otimes \nabla^k \varphi_l) (g_G^{\otimes 2}) R^{n-1} - \\
&\quad - \frac{3\mathfrak{c}_{2,n}(n+5)_{3,-2}}{(n! \omega_n)^2} \sum_{l_1, l_2=n+1}^N \sum_{\substack{i_1+k_1=i_2+k_2=3, \\ i_1, i_2, k_1, k_2>0}} (\nabla^{i_1} \varphi_{l_1} \otimes \nabla^{k_1} \varphi_{l_1} \otimes \nabla^{i_2} \varphi_{l_2} \otimes \nabla^{k_2} \varphi_{l_2})(x, g_G^{\otimes 3}) R^{n-1},
\end{aligned}$$

where g_G is the dual metric to the transversal Hessian:

$$H_G(v) = |v|^2 + \sum_{l=n+1}^N (\nabla \varphi_l(x) \cdot v)^2.$$

Example 2.31 (Evaluations of symbols for geodesic distances). Consider the geodesic distance on a Riemannian manifold M as in Example 2.18. Let g_M denote the Riemannian metric, and recall from Example 2.18 that g_M coincides with the transversal Hessian of d_{geo}^2 at the diagonal. We can compute $a_{2,0}$ in this case by means of known Riemannian curvatures. We fix a point x and choose coordinates so that Γ vanishes in that point (normal coordinates). In these coordinates, $C_{d_{\text{geo}}^2}^3(x, v) = 0$ and $C_{d_{\text{geo}}^2}^4(x, \cdot)$ is a third of the Riemannian curvature in x by Proposition 2.19. By Theorem 2.27 we conclude that for $n = 3$, we have that

$$a_{2,0}(x, R) = -\frac{80R^2}{(3! \omega_3)^2} s_g(x),$$

while for $n \neq 1, 3$, we have that

$$a_{2,0}(x, R) = -\frac{R^{n-1}}{(n! \omega_n)^2} \mathfrak{c}_{1,n}(n^2-1) s_g(x),$$

where s_g denotes the scalar curvature.

3. GLOBAL BEHAVIOR OF \mathcal{Z} ON COMPACT MANIFOLDS

In the previous section we studied the localization Q of \mathcal{Z} near the diagonal. Here \mathcal{Z} is the operator with parameter R defined from a distance function as in Equation (7). We now turn to study \mathcal{Z} by considering distance functions for which $L := \mathcal{Z} - Q$ in a certain sense is negligible so that Q dominates.

3.1. Controlling the off-diagonal part of \mathcal{Z} . In this subsection we shall study the remainder $L = \mathcal{Z} - Q$. We note that

$$L(R)f(x) = \frac{1}{R} \int_M (1 - \chi(x, y)) e^{-Rd(x, y)} f(y) dy.$$

We start making two initial observations concerning the remainder term L . The first observation concerns the behavior of the remainder term in L^2 .

Proposition 3.1. *Let M be a compact manifold and $d : M \times M \rightarrow [0, \infty)$ a distance function on M . Then*

$$\mathbb{C} \setminus \{0\} \ni R \mapsto L(R) \in \mathcal{L}^2(L^2(M)),$$

defines a holomorphic Hilbert-Schmidt valued function. Moreover, L satisfies that

$$\|L(R)\|_{\mathcal{L}^2(L^2(M))} = O(\text{Re}(R)^{-\infty}) \quad \text{as } \text{Re}(R) \rightarrow +\infty.$$

By the standard norm estimate $\|K\|_{\mathbb{B}} \leq \|K\|_{\mathcal{L}^2}$, the statement in the proposition also holds in the operator norm. We remark that by Theorem 2.9 and B.11, the localization to the diagonal satisfies

$$\|Q(R)\|_{\mathcal{L}^p(L^2(M))} = O(\text{Re}(R)^{\frac{n}{p} - n - 1}) \quad \text{as } \text{Re}(R) \rightarrow +\infty,$$

for any $p > 1 + 1/n$ (the bound is not uniform in p).

Proof. The function $\mathbb{C} \setminus \{0\} \ni R \mapsto \frac{1}{R}(1 - \chi(\cdot, \cdot))e^{-Rd(\cdot, \cdot)} \in C(M \times M)$ is clearly holomorphic in the norm on $C(M \times M)$. Since $\chi = 1$ in a neighborhood of the diagonal, compactness of M implies that $\|\frac{1}{R}(1 - \chi(\cdot, \cdot))e^{-Rd(\cdot, \cdot)}\|_{C(M \times M)} = O(\operatorname{Re}(R)^{-\infty})$ as $\operatorname{Re}(R) \rightarrow +\infty$. Since M is compact, an integral operator K with kernel $k \in C(M \times M)$ satisfies the estimate for the Hilbert-Schmidt norm $\|K\|_{\mathcal{L}^2(L^2(M))} \leq \operatorname{vol}(M)\|k\|_{C(M \times M)}$. The proposition follows. \square

The second observation concerns how an a priori estimate with respect to the off-diagonal remainder L affects distributional solutions to the magnitude equation $\mathcal{Z}u = 1$.

Proposition 3.2. *Let M be a compact manifold and $d : M \times M \rightarrow [0, \infty)$ a distance function on M such that d^2 is regular at the diagonal. Let $N \in \mathbb{N}$ and Γ be a sector. Assume that $f \in C^\infty(M)$ and that $(u_R)_{R \in \Gamma} \subseteq \mathcal{D}'(M)$ is a family of solutions to*

$$\mathcal{Z}(R)u_R = f$$

satisfying that $\langle L(R)u_R, \psi \rangle = O(\operatorname{Re}(R)^{-N})$ as $\operatorname{Re}(R) \rightarrow +\infty$ in Γ for all $\psi \in C^\infty(M)$. Then for large enough $\operatorname{Re}(R)$,

$$u_R = Q(R)^{-1}f + v_R,$$

where $(v_R)_{R \in \Gamma} \subseteq \mathcal{D}'(M)$ is a family satisfying that $\langle v_R, \psi \rangle = O(\operatorname{Re}(R)^{-N+n+1})$ as $\operatorname{Re}(R) \rightarrow +\infty$ in Γ for all $\psi \in C^\infty(M)$. In particular, for any $\psi \in C^\infty(M)$,

$$\langle u_R, \psi \rangle = \langle Q(R)^{-1}f, \psi \rangle + O(\operatorname{Re}(R)^{-N+n+1}), \quad \text{as } \operatorname{Re}(R) \rightarrow +\infty \text{ in } \Gamma.$$

Proof. The equation $\mathcal{Z}(R)u_R = f$ implies that

$$u_R = Q(R)^{-1}f - Q(R)^{-1}L(R)u_R.$$

Consider the distribution $v_R := -Q(R)^{-1}L(R)u_R \equiv u_R - Q(R)^{-1}f$. For $\psi \in C^\infty(M)$, we have that

$$\langle v_R, \psi \rangle = \langle L(R)u_R, Q(R)^{-1}\psi \rangle = O(\operatorname{Re}(R)^{-N+n+1}),$$

because of the assumption on $L(R)u_R = O(\operatorname{Re}(R)^{-N})$ in a weak sense and the fact that $Q(R)^{-1}$ is a pseudodifferential operator with parameter of order $n+1$ preserving $C^\infty(M)$ with uniform norm estimates $\|Q(R)^{-1}f\|_{C^k} \leq C_k(1 + \operatorname{Re}(R))^{n+1}\|f\|_{C^{n+k+1}}$. \square

Proposition 3.1 implies that L is small as an operator on L^2 , and Proposition 3.2 implies that the resulting remainder term will not alter the asymptotic properties of solutions given an a priori estimate. However, as Q is of negative order and acts compactly on L^2 , well-posedness of the problem in L^2 is not assured. We shall circumvent this problem by imposing a regularity assumption on the distance function that forces the magnitude equation naturally into a Sobolev space framework. We discuss examples satisfying this assumption, as well as counterexamples, below in the Subsections 3.2 and 3.4, respectively.

Definition 3.3 (Property (MR) of distance functions). Let M be an n -dimensional compact manifold and $d : M \times M \rightarrow [0, \infty)$ a distance function on M such that d^2 is regular at the diagonal. Set $\mu := (n+1)/2$. For a sector $[1, \infty) \subseteq \Gamma \subseteq \mathbb{C}$, we say that d has *property (MR)* on Γ if for any $R \in \Gamma$, $L(R)$ extends to a continuous mapping $H^{-\mu}(M) \rightarrow H^\mu(M)$ with

$$\|L(R)\|_{H^{-\mu}(M) \rightarrow H^\mu(M)} = O(\operatorname{Re}(R)^{-\infty}), \quad \text{as } \operatorname{Re}(R) \rightarrow +\infty \text{ in } \Gamma.$$

If $L(R) : H^{-\mu}(M) \rightarrow H^\mu(M)$ is a compact operator for $R \in \Gamma$ and

$$\Gamma \ni R \mapsto L(R) \in \mathbb{K}(H^{-\mu}(M), H^\mu(M)),$$

is holomorphic in norm sense, we say that d has *property (SMR)* on Γ .

The acronyms MR and SMR stand for magnitude regularity and strong magnitude regularity, respectively. Assuming these properties, the operator \mathcal{Z} inherits relevant analytic and geometric properties from Q . Property (MR) will be used to compute asymptotic solutions to the magnitude equation $R\mathcal{Z}u = 1$, while property (SMR) will be used for constructing meromorphic extensions of \mathcal{Z}^{-1} . If Γ is a sector on which a distance function has property (MR), it is clear that $0 \notin \Gamma$. We consider such results for compact manifolds below in Subsection 3.3.

3.2. Examples of distance functions satisfying property (MR) and (SMR). Let us give a method to produce distance functions with property (SMR):

Proposition 3.4. *Let M be a compact manifold embedded into a Riemannian manifold $i : M \rightarrow W$ such that the square of its geodesic distance $d_{\text{geo},W}^2$ is smooth, for instance $W = \mathbb{R}^N$, $W = \mathbb{H}_{N,\mathbb{R}}$ or $W = \mathbb{H}_{N,\mathbb{C}}$, for some $N \in \mathbb{N}$.*

The distance function $d : M \times M \rightarrow [0, \infty)$, $d(x, y) := d_{\text{geo},W}(i(x), i(y))$ satisfies

- i) d^2 is smooth on $M \times M$ and regular at the diagonal.
- ii) $L \in \Psi^{-\infty}(M; \mathbb{C}_+)$ and $L(R) \in \Psi^{-\infty}(M)$ for any $R \in \mathbb{C} \setminus \{0\}$.
- iii) d has property (SMR) on $\mathbb{C} \setminus \{0\}$.

Proof. Since $d_{\text{geo},W}^2$ is smooth, it is clear that d^2 is smooth on $M \times M$. It follows from Proposition 2.19 that d^2 is regular at the diagonal, and in fact g_{d^2} is the pullback metric i^*g_W from the Riemannian metric g_W on W . Therefore the singular support of d is the diagonal, and the integral kernel of $L(R)$ is smooth so $L(R)$ extends to a continuous mapping $L(R) : H^s(M) \rightarrow H^t(M)$ for any $s, t \in \mathbb{R}$. Moreover, for any vector fields X_1, \dots, X_m on $M \times M$ we can estimate

$$|X_1 \cdots X_m ((1 - \chi)e^{-Rd})| \leq C_m |R|^{m-1} e^{-\varepsilon \text{Re}(R)}.$$

Again, we write $\varepsilon := \inf\{d(x, y) : (x, y) \in \chi^{-1}(1)\} > 0$. In particular, we readily can deduce that $L \in \Psi^{-\infty}(M; \mathbb{C}_+)$, and for any $s, t \in \mathbb{R}$,

$$\|L(R)\|_{H^s(M) \rightarrow H^t(M)} = O(\text{Re}(R)^{-\infty}), \quad \text{as } \text{Re}(R) \rightarrow +\infty.$$

Therefore d has property (SMR) on $\mathbb{C} \setminus \{0\}$. □

In light of Proposition 3.4, we note the following corollary of Theorem 2.23.

Corollary 3.5. *Let M be a compact manifold equipped with a distance function $d : M \times M \rightarrow [0, \infty)$ such that d^2 is smooth, e.g. a subspace distance as in Proposition 3.4. Then $\mathcal{Z}(R) \in \Psi_{\text{cl}}^{-n-1}(M)$ is an elliptic pseudodifferential operator for any $R \in \mathbb{C} \setminus \{0\}$. Furthermore, $\mathcal{Z} \in \Psi_{\text{cl}}^{-n-1}(M; \mathbb{C}_+)$ is elliptic with parameter and its full symbol coincides with that of Q as given in Theorem 2.9.*

Another example of distance functions with property (MR) arises on spheres.

Proposition 3.6. *Let d denote the geodesic distance on a sphere S^n in its round metric. The distance function d has property (MR) on $\mathbb{C} \setminus \{0\}$ but fails to satisfy property (SMR) on any sector.*

Proof. The square of the geodesic distance is regular at the diagonal by Proposition 2.19. We note that d is smooth on $\{(x, y) \in S^n \times S^n : x \neq \pm y\}$. Consider the operator $Uf(x) := f(-x)$. The operator U acts via pullback along the antipodal mapping $\varphi(x) := -x$ which acts isometrically due to $O(n)$ -invariance of the Riemannian metric on S^n . The operator U extends to a unitary on all Sobolev spaces $H^s(S^n)$, $s \in \mathbb{R}$. Since it holds that $d(x, y) = \pi - d(x, \varphi(y))$ we can conclude that the integral kernel of $L(R)U$ is given by $\chi_\varphi e^{-Rd} e^{-\pi R}$, where $\chi_\varphi(x, y) := 1 - \chi(x, \varphi(y))$. In particular, since χ_φ satisfies that $\chi_\varphi = 1$ on the diagonal and $\chi_\varphi = 0$ on a neighborhood of the off-diagonal singularities of d , the operator $L(R)Ue^{\pi R}$ is an elliptic pseudodifferential operator with parameter of order $-n - 1$ by the same argument as in Theorem 2.9. It follows that property (SMR) fails on any sector. Using that U is unitary, $L(R)$ extends to a continuous operator $H^{-\mu}(S^n) \rightarrow H^\mu(S^n)$ with

$$\|L(R)Ue^{\pi R}\|_{H^{-\mu}(S^n) \rightarrow H^\mu(S^n)} = O(1),$$

as $\text{Re}(R) \rightarrow \infty$. Since U is unitary, we deduce that

$$\|L(R)\|_{H^{-\mu}(S^n) \rightarrow H^\mu(S^n)} = O(e^{-\pi \text{Re}(R)}) = O(\text{Re}(R)^{-\infty}) \quad \text{as } \text{Re}(R) \rightarrow \infty.$$

□

3.3. Analytic results for the operator \mathcal{Z} . We are now ready to extend the results of Section 2 for Q to results on the operator \mathcal{Z} for compact manifolds with a distance function satisfying property (MR) as defined in Definition 3.3.

Theorem 3.7. *Let M be a compact n -dimensional manifold and $d : M \times M \rightarrow [0, \infty)$ a distance function on M with property (MR) on the sector Γ . Set $\mu := (n+1)/2$. Then there is an $R_0 \geq 0$ such that*

$$\mathcal{Z}(R) : H^{-\mu}(M) \rightarrow H^{\mu}(M),$$

is invertible for all $R \in \Gamma \cap \Gamma_{\pi/(n+1)}(R_0)$. Moreover, for $R \in \Gamma \cap \Gamma_{\pi/(n+1)}(R_0)$

$$\mathcal{Z}^{-1} = Q^{-1} + \mathcal{R},$$

where $Q^{-1} \in \Psi_{\text{cl}}^{n+1}(M; \Gamma_{\pi/(n+1)}(R_0))$ is the elliptic pseudodifferential operator with parameter constructed in Corollary 2.22 and $\mathcal{R} : H^{\mu}(M) \rightarrow H^{-\mu}(M)$ is a family of operators parametrized by $R \in \Gamma \cap \Gamma_{\pi/(n+1)}(R_0)$ such that

$$\|\mathcal{R}\|_{H^{\mu}(M) \rightarrow H^{-\mu}(M)} = O(\text{Re}(R)^{-\infty}), \quad \text{as } \text{Re}(R) \rightarrow +\infty \text{ in } \Gamma \cap \Gamma_{\pi/(n+1)}(R_0).$$

Moreover, there is a constant $C > 0$ such that

$$(27) \quad C^{-1} \|f\|_{H_{|R|}^{-\mu}(M)}^2 \leq \text{Re}\langle f, \mathcal{Z}(R)f \rangle_{L^2} \leq C \|f\|_{H_{|R|}^{-\mu}(M)}^2,$$

for $R \in \Gamma \cap \Gamma_{\pi/(n+1)}(R_0)$ and $f \in H^{-\mu}(M)$. In particular, for $R \in \Gamma \cap \Gamma_{\pi/(n+1)}(R_0)$, $\mathcal{Z}(R)$ is coercive in form sense on $L^2(M)$ for the $H^{-\mu}$ -norm.

Proof. The operator Q is invertible by Corollary 2.22. We can therefore write

$$Q^{-1}\mathcal{Z} = 1 + Q^{-1}L,$$

as operators on $H^{-\mu}(M)$. Since Q^{-1} is a pseudodifferential operator with parameter, we have that

$$(28) \quad \begin{aligned} \|Q(R)^{-1}L(R)\|_{H^{-\frac{n+1}{2}}(M) \rightarrow H^{-\frac{n+1}{2}}(M)} &= O(\text{Re}(R)^{-\infty}), \\ &\text{as } \text{Re}(R) \rightarrow +\infty \text{ in } \Gamma \cap \Gamma_{\pi/(n+1)}(R_0). \end{aligned}$$

Therefore, for $\text{Re}(R) \gg 0$ the operator $(1 + Q^{-1}L)^{-1}Q^{-1}$ exists and is a left inverse to \mathcal{Z} . By an analogous argument,

$$(29) \quad \begin{aligned} \|L(R)Q(R)^{-1}\|_{H^{\frac{n+1}{2}}(M) \rightarrow H^{\frac{n+1}{2}}(M)} &= O(\text{Re}(R)^{-\infty}), \\ &\text{as } \text{Re}(R) \rightarrow +\infty \text{ in } \Gamma \cap \Gamma_{\pi/(n+1)}(R_0), \end{aligned}$$

and \mathcal{Z} has the right inverse $Q^{-1}(1 + LQ^{-1})^{-1}$ for $\text{Re}(R) \gg 0$. Therefore $(1 + Q^{-1}L)^{-1}Q^{-1} = Q^{-1}(1 + LQ^{-1})^{-1}$ and this operator is an inverse to \mathcal{Z} . By the estimates (28) and (29), it follows that

$$\mathcal{R} = \mathcal{Z}^{-1} - Q^{-1} = Q^{-1} \left((1 + LQ^{-1})^{-1} - 1 \right) = \sum_{k=0}^{\infty} (-1)^k Q^{-1} (LQ^{-1})^k,$$

as a norm convergent sum and has the required decay property as $\text{Re}(R) \rightarrow +\infty$.

The estimate (27) follows from the decay property of \mathcal{R} and Theorem 2.23. \square

The following result is immediate from Lemma 2.24 and Theorem 3.7.

Corollary 3.8. *Let M be a compact n -dimensional manifold and $d : M \times M \rightarrow [0, \infty)$ a distance function on M with property (MR) on the sector Γ . Take the sequence of homogeneous functions $(a_{j,0})_{j \in \mathbb{N}} \subseteq C^{\infty}(M \times \Gamma \cap \mathbb{C}_+)$ as in Lemma 2.25. Then, for any $N \in \mathbb{N}$, we have that*

$$[\mathcal{Z}(R)^{-1}1](x) = \sum_{j=0}^N a_j(x, R) + r_N(x, R),$$

where $r_N \in C(\Gamma \cap \mathbb{C}_+, H^{-\mu}(M))$, for $\mu = (n+1)/2$, is a function such that

$$\|r_N(\cdot, R)\|_{H^{-\mu}(M)} = O(\text{Re}(R)^{n+1-N}), \quad \text{as } \text{Re}(R) \rightarrow +\infty \text{ in } \Gamma.$$

Remark 3.9. We remark that the role that property (MR) plays in Corollary 3.8 is to ensure existence of a distributional solution to $\mathcal{Z}(R)u_R = 1$. For *computing* the integrated asymptotics $\langle 1, \mathcal{Z}(R)^{-1}1 \rangle$, property (MR) is not necessary. Indeed, with no assumptions of property (MR), but assuming that $d : M \times M \rightarrow [0, \infty)$ is a distance function on M such that d^2 is regular at the diagonal and that $(u_R)_{R > R_0} \subseteq \mathcal{D}'(M)$ is a family of solutions to

$$\mathcal{Z}(R)u_R = 1$$

satisfying that $\langle L(R)u_R, \psi \rangle = O(\operatorname{Re}(R)^{-N})$ as $\operatorname{Re}(R) \rightarrow +\infty$ for all $\psi \in C^\infty(M)$, we have that

$$\langle u_R, 1 \rangle = \sum_{j=1}^N \int_M a_j(x, R) dx + O(\operatorname{Re}(R)^{n+1-N}),$$

for any N . This follows from Lemma 2.24 and Proposition 3.2.

One instance where solutions to $\mathcal{Z}(R)u_R = 1$ exist, yet the distance function need not satisfy property (MR), is the geodesic distance on a compact symmetric space $M = G/H$. See Proposition 3.17 below for examples of symmetric spaces failing to satisfy property (MR). The problem $\mathcal{Z}(R)u_R = 1$ was studied for compact symmetric spaces in [44]. For a compact symmetric space $M = G/H$, we use the normalized G -invariant measure induced by the Haar measure. By symmetry, the function

$$u_R(x) = \frac{1}{\int_{G/H} e^{-Rd(x,y)} dy},$$

is constant and therefore solves $\mathcal{Z}(R)u_R = 1$. It is readily verified that $L(R)u_R = O(R^{-\infty})$ in distributional sense. We conclude that each $a_j(x, R)$ is constant and that

$$u_R(x) = u_R(eH) = \sum_{j=0}^N a_j(eH, R) + O(\operatorname{Re}(R)^{n+1-N}),$$

for any N . For examples of computations of u_R for compact symmetric spaces, see [44].

The next result poses an obstruction to property (MR) for distance functions and should be viewed as complementary to Theorem 3.16. Recall the following terminology from [32]: a compact metric space (X, d) is said to be positive definite if for any finite subset $F \subseteq X$, the matrix $(e^{-d(x,y)})_{x,y \in F}$ is positive definite. If (X, Rd) is positive definite for all $R > 0$, we say that (X, d) is stably positive definite.

Corollary 3.10. *Assume that d is a distance function on a compact manifold M with property (MR) on $[1, \infty)$. Then there exists an $R_0 \geq 0$ such that (M, Rd) is positive definite for all $R > R_0$.*

Proof. Consider the quadratic form $\mathfrak{q}_R(u) = \langle u, \mathcal{Z}(R)u \rangle_{L^2}$, $u \in H^{-\mu}(M)$. By Theorem 3.7, \mathfrak{q}_R is positive definite for $R > R_0$, for some $R_0 \geq 0$. In particular, for any subspace $V \subseteq H^{-\mu}(M)$ the restriction of \mathfrak{q}_R to V is also positive definite for $R > R_0$. For a finite subset $F \subseteq M$, consider the subspace $V_F \subseteq H^{-\mu}(M)$ spanned by $\{\delta_x : x \in F\}$. In the basis $(\delta_x)_{x \in F}$, the quadratic form $\mathfrak{q}_R|_{V_F}$ is represented by the $|F| \times |F|$ -matrix $(e^{-Rd(x,y)})_{x,y \in F}$ and so it is positive definite for $R > R_0$. \square

Remark 3.11. It was proven in [32, Subsection 3.2] that if M is a compact Riemannian manifold with $\pi_1(M) \neq 0$, the geodesic distance is not stably positive definite, i.e. there exists an $R > 0$ and a finite subset $F \subseteq M$ such that $(e^{-Rd_{\text{geo}}(x,y)})_{x,y \in F}$ fails to be positive definite. The manifold $M = S^1$ is not simply connected, and therefore fails to be stably positive definite. but nevertheless Proposition 3.6 implies that $M = S^1$ with its geodesic distance has property (MR). Therefore, property (MR) does not imply stably positive definiteness of the metric space but only an asymptotic version thereof.

Theorem 3.12. *Let M be an n -dimensional compact manifold with a distance function d having property (SMR) on Γ and set $\mu = (n+1)/2$. Then the operator*

$$\mathcal{Z}(R) : H^{-\mu}(M) \rightarrow H^\mu(M),$$

is a well defined Fredholm operator for all $R \in \Gamma$ invertible for $R \in \Gamma \cap \Gamma_{\pi/(n+1)}(R_0)$. Moreover, the operator $\mathcal{Z}(R) : H^{-\mu}(M) \rightarrow H^{\mu}(M)$ depends holomorphically on R in Γ and $\mathcal{Z}(R)^{-1} : H^{\mu}(M) \rightarrow H^{-\mu}(M)$ depends holomorphically on $R \in \Gamma \cap \Gamma_{\pi/(n+1)}(R_0)$ and admits a meromorphic extension to Γ .

Proof. By Theorem 2.23, Q is a holomorphic function on Γ with values in the Fredholm operators and by property (SMR), L is a compact valued holomorphic function on Γ . Therefore \mathcal{Z} defines a holomorphic function on Γ with values in the Fredholm operators. Since \mathcal{Z} is invertible for a large enough R , see Theorem 3.7, the theorem follows from the meromorphic Fredholm theorem, see Appendix D. \square

Remark 3.13. To ensure holomorphicity of \mathcal{Z} and \mathcal{Z}^{-1} on sectors, the full property (SMR) is not needed. Indeed, if d has property (MR) on a sector Γ and $\Gamma \ni R \mapsto L(R) \in \mathbb{B}(H^{-\mu}(M), H^{\mu}(M))$ is additionally holomorphic (in norm sense) then by Theorem 3.7, for some $R_0 \geq 0$, the mapping

$$\Gamma \cap \Gamma_{\pi/(n+1)}(R_0) \ni R \mapsto \mathcal{Z}(R) \in \mathbb{B}(H^{-\mu}(M), H^{\mu}(M))$$

is a holomorphic family of invertible operators. These properties are inherited by its inverse,

$$\Gamma \cap \Gamma_{\pi/(n+1)}(R_0) \ni R \mapsto \mathcal{Z}(R)^{-1} \in \mathbb{B}(H^{\mu}(M), H^{-\mu}(M)).$$

By the proof of Proposition 3.6, this discussion applies to S^n showing that for some R_0 , $\mathcal{Z}(R)^{-1}$ is holomorphic for $R \in \Gamma \cap \Gamma_{\pi/(n+1)}(R_0)$ in this case.

The following result follows from Corollary 3.5 and Theorem 3.12.

Theorem 3.14. *Let M be a compact manifold with a distance function d such that d^2 is smooth on $M \times M$ and regular at the diagonal (cf. Proposition 3.4), then $\mathcal{Z}(R) : H^{-\mu}(M) \rightarrow H^{\mu}(M)$ depends holomorphically on $R \in \mathbb{C} \setminus \{0\}$ and the operator $\mathcal{Z}(R)^{-1} : H^{\mu}(M) \rightarrow H^{-\mu}(M)$ extends meromorphically to $\mathbb{C} \setminus \{0\}$.*

3.4. Examples of distance functions that fail to satisfy property (MR). Property (MR) of a distance function, as defined in Definition 3.3, is a notable restriction on the singular support and singularity structure of the distance function. We remark here that by a singular point, we mean any point in the singular support, i.e. one in which the function is not C^∞ . To better understand how these singularities affect the operator theoretic properties of L , the reader is encouraged to review the proof of Proposition 3.6 where a crucial feature used in the proof is that the geodesic distance on spheres near an off-diagonal singularity has the same singular features as it has near the antipode of the singularity. An important property used there can be stated as having control of the dimension of the off-diagonal singular support of the metric. We make an elementary observation that follows from the smoothness of the function $(0, \infty) \ni t \mapsto \sqrt{t} \in (0, \infty)$.

Proposition 3.15. *Let d be a distance function on a manifold M . Then it holds that*

$$\text{singsupp}(d) \setminus \text{Diag}_M = \text{singsupp}(d^2) \setminus \text{Diag}_M.$$

Let us give a sufficient condition for a distance function (regular at the diagonal) *not* to satisfy property (MR). We note that an additional obstruction was provided above in Proposition 3.10.

Theorem 3.16. *Let M be a compact manifold and d a distance function such that d^2 is regular at the diagonal. Assume that there exists a submanifold $N \subseteq M \times M$ with $N \subseteq \text{singsupp}(d^2) \setminus \text{Diag}_M$ such that any point $z_0 \in N$ admits a neighborhood U_0 in $M \times M$ and a coordinate chart*

$$\varphi : \mathbb{R}_t^{\dim(N)} \times \mathbb{R}_s^{\dim(M) - \dim(N)} \rightarrow U_0,$$

with $\varphi(0) = x_0$, $\varphi^{-1}(U_0 \cap N) = \mathbb{R}_t^{\dim(N)} \times \{0\}$ and $(t, s) \mapsto \varphi^*d(t, s) - |s|$ being smooth in a neighborhood of 0. Then $L(R)$ does not extend to a continuous operator

$$H^{-\mu}(M) \rightarrow H^s(M),$$

for any $s > (3 \dim(M) + 1)/2 - \dim(N)$ where $\mu = (\dim(M) + 1)/2$. In particular, if $\dim(N) > \dim(M)$ then d does not have property (MR) (see Definition 3.3) on any sector containing a half-ray $[R_0, \infty)$.

Proof. We pick a point $(x_0, y_0) \in N$ and neighborhoods U_1 and U_2 of x_0 and y_0 respectively, such that there exists a coordinate chart φ as in item ii) above on $U_0 = U_1 \times U_2$. Pick functions $\chi_1 \in C_c^\infty(U_1)$ and $\chi_2 \in C_c^\infty(U_2)$ such that $\chi_1 = 1$ near x_0 , $\chi_2 = 1$ near y_0 and such that $(t, s) \mapsto \varphi^*d(t, s) - |s|$ is smooth on $\varphi^{-1}(\text{supp}(\chi_1) \times \text{supp}(\chi_2))$. Clearly, it suffices to prove that for any $R > 0$, the operator $\chi_1 L(R) \chi_2$ does not extend to a continuous operator $H^{-\mu}(M) \rightarrow H^s(M)$ for any $s > (3 \dim(M) + 1)/2 - \dim(N)$.

Set $k = \dim(N)$ and $n = \dim(M)$. The Schwartz kernel of $\chi_1 L(R) \chi_2$ is a distribution on $M \times M$ supported in $U_0 = U_1 \times U_2$ and pulling this kernel back along φ , we arrive at the distribution

$$K(t, s) = \chi_0(s, t) e^{-R|s|},$$

where $\chi_0 = \varphi^*[(\chi_1 \otimes \chi_2)(1 - \chi)] e^{-R\psi} \in C_c^\infty(\mathbb{R}^{2n})$ for $\psi(t, s) = \varphi^*d(t, s) - |s|$. It follows from Proposition A.7 and combining a Taylor expansion with asymptotic completeness, that $K \in C I^{k-2n-1}(\mathbb{R}^{2n}, \mathbb{R}^k)$. We conclude that $\chi_1 L(R) \chi_2$ is a Fourier integral operator of order $k-2n-1$ and this operator is elliptic in a neighborhood of (x_0, y_0) . Therefore, since $\chi_1 L(R) \chi_2$ is elliptic near (x_0, y_0) of order $k-2n-1$ it does not extend to a continuous operator $H^{-(n+1)/2}(M) \rightarrow H^s(M)$ for any

$$s > -(n+1)/2 - (k-2n-1) = (3n+1)/2 - k.$$

□

Proposition 3.17. *Let $n > 1$. The geodesic distance on the n -dimensional torus $M = \mathbb{T}^n$ or the real projective space $M = \mathbb{R}P^n$ fails to satisfy property (MR) (see Definition 3.3) on any sector containing a half-ray $[R_0, \infty)$. In fact, $L(R)$ does not extend to a continuous map $H^{-\mu}(M) \rightarrow H^s(M)$ for $s > -n/2 + 3/2$.*

Proof. The proofs for both cases follow the same lines and rely on Theorem 3.16. For $\mathbb{R}P^n$ we give a more geometric argument, and for \mathbb{T}^n we give a coordinate oriented argument.

We first prove the result for real projective space. The projective space $\mathbb{R}P^n$ is the quotient of S^n by the antipodal map $x \mapsto -x$. This quotient map is a covering map, and it is locally isometric with respect to the geodesic distance. For $x \in \mathbb{R}P^n$, the equator $\tilde{E}(x) \subseteq S^n$ is defined by

$$\tilde{E}(x) = \{v \in S^n : v \cdot x = 0\}.$$

The condition $v \cdot x = 0$ being invariant under the antipodal map, $\tilde{E}(x)$ only depends on $x \in \mathbb{R}P^n$ and not on a choice of pre-image of x in S^n . We let $E(x) \subseteq \mathbb{R}P^n$ denote the image under the quotient map. The off-diagonal singular support of the geodesic distance on $\mathbb{R}P^n$ is the set

$$\{(x, y) \in \mathbb{R}P^n \times \mathbb{R}P^n : y \in E(x)\}.$$

Indeed, any geodesic in $\mathbb{R}P^n$ from a point x lifts uniquely up to a geodesic on S^n up until the point it crosses $E(x)$ where the geodesic distance $d_{\text{geo}}(x, \cdot)$ has a kink. The projection mapping $p_1 : \text{singsupp}(d_{\text{geo}}) \setminus \text{Diag}_{\mathbb{R}P^n} \rightarrow \mathbb{R}P^n$ is a locally trivial $\mathbb{R}P^{n-1}$ -bundle on $\mathbb{R}P^n$. Therefore, $N = \text{singsupp}(d_{\text{geo}}) \setminus \text{Diag}_{\mathbb{R}P^n} \rightarrow \mathbb{R}P^n$ is a manifold of dimension $2n-1 > n$ and suitable local trivializations of $p_1 : N \rightarrow \mathbb{R}P^n$ satisfy the assumptions of item ii) in Theorem 3.16. We conclude from Theorem 3.16 that $L(R)$ has no bounded extension to an operator $H^{-\mu}(\mathbb{R}P^n) \rightarrow H^s(\mathbb{R}P^n)$ for any $s > -n/2 + 3/2$.

We now prove the result for the n -dimensional torus \mathbb{T}^n . Write $\mathbb{T}^n = \mathbb{R}^n/\mathbb{Z}^n$. While \mathbb{T}^n is a symmetric space, i.e. $\mathbb{T}^n = G/H$ for $G = \mathbb{T}^n$ and $H = 1$, it is instructive to consider $d(x, y)$ for $x = 0$. We have that $d(0, y) = |y|$ where we represent y by as an element of the fundamental domain $[-1/2, 1/2]^n$, and $\mathbb{T}^n \ni y \mapsto d(0, y)$ as a function on \mathbb{T}^n is the \mathbb{Z}^n -periodic extension of $[-1/2, 1/2]^n \ni y \mapsto |y|$. Therefore $y \mapsto d(0, y)$ has kinks on the image of $\partial([-1/2, 1/2]^n)$ in \mathbb{T}^n . In particular, we see that the off-diagonal singular support of d is the set

$$\{(x, y) \in \mathbb{T}^n \times \mathbb{T}^n : x - y \in \partial([-1/2, 1/2]^n) + \mathbb{Z}^n\}.$$

Consider the submanifold $N_0 := \{1/2\} \times (-1/4, 1/4)^{n-1} \subseteq \mathbb{R}^n$ and define the $2n-1$ -dimensional submanifold

$$N := \{(x, y) \in \mathbb{T}^n \times \mathbb{T}^n : x - y \in N_0 + \mathbb{Z}^n\} \subseteq \text{singsupp}(d^2) \setminus \text{Diag}_{\mathbb{T}^n}.$$

Consider a point

$$x_0 = (x, 1/2, y') \in N.$$

On the open ball of radius $1/4$ centred at x_0 , we introduce the coordinates $u = x_1 - y_1 - 1/2$ and $t = (t', y)$ where $t' = x' - y'$ in terms of standard coordinates $x = (x_1, x')$ and $y = (y_1, y')$. In these coordinates, we have that

$$d(x, y) = \sqrt{\left(\frac{1}{2} - |u|\right)^2 + |t'|^2} = \sqrt{-|u| + |t'|^2 + \frac{1}{4} + |u|^2}.$$

By shrinking the neighborhood of x_0 , we can Taylor expand

$$\begin{aligned} \sqrt{-|u| + |t'|^2 + \frac{1}{4} + |u|^2} &= \sqrt{|t'|^2 + \frac{1}{4} + |u|^2} \sqrt{1 - \frac{|u|}{|t'|^2 + \frac{1}{4} + |u|^2}} = \sum_{k=0}^{\infty} \alpha_k \frac{|u|^k}{(|t'|^2 + \frac{1}{4} + |u|^2)^{k-1}} = \\ &= |u| \underbrace{\sum_{k=0}^{\infty} \alpha_{2k+1} \frac{|u|^{2k}}{(|t'|^2 + \frac{1}{4} + |u|^2)^{2k}}}_{g(u, t')} + \sum_{k=0}^{\infty} \alpha_{2k} \frac{|u|^{2k}}{(|t'|^2 + \frac{1}{4} + |u|^2)^{2k-1}}. \end{aligned}$$

The functions g and \tilde{g} are smooth near 0. Since $g(0, 0) \neq 0$, we can define the new coordinate $s := u\tilde{g}(u, t')$. We conclude that in these coordinates $d(x, y) - |s|$ is smooth. We conclude from Theorem 3.16 that $L(R)$ has no bounded extension to an operator $H^{-\mu}(\mathbb{T}^n) \rightarrow H^s(\mathbb{T}^n)$ for any $s > -n/2 + 3/2$. \square

Remark 3.18. The analytical issues arising from the remainder term L reflect fundamental problems in Riemannian geometry. The singular support of the geodesic distance is akin to the conjugate locus of the Riemannian metric, which in general is hard to describe, see [5, 42] and for related technical issues arising in the X -ray transform on a Riemannian manifold see [20]. Furthermore, Theorem 3.16 shows that even when the singular support is a tractable set, i.e. when it looks like a submanifold near some point, dimensional obstructions to property (MR) appear. This gives rise to the analytic problem that the operator L in the decomposition $\mathcal{Z} = Q + L$ is in general of order higher than $-n - 1$ in the Sobolev order, while it is infinitely decaying in the parameter R . In this case L is of higher order than Q , which is elliptic with parameter of order $-n - 1$ and which determines the analytic and geometric properties of \mathcal{Z} in this article.

4. THE OPERATOR \mathcal{Z} ON SOBOLEV SPACES FOR A MANIFOLD WITH BOUNDARY

We now turn to compact manifolds with boundary. For simplicity, we tacitly assume that X is a compact domain in a manifold M and, for the purposes of this section, it suffices to assume that X has a C^0 -boundary. Recall that a domain is said to have C^0 -boundary if its boundary can be realized as the graph of a continuous function. We call such spaces X a compact manifold with C^0 -boundary. We may then study the operators \mathcal{Z} and Q in M and deduce results in X by restriction to distributions supported in X . In this section we study analytic properties and meromorphic extensions. Asymptotic properties are studied in the following section under additional regularity assumptions on the boundary. For notational clarity, we indicate the manifold on which an operator is defined by a subscript, e.g. \mathcal{Z}_X and \mathcal{Z}_M for the corresponding operator on X and M , respectively.

We shall make use of the following scales of Sobolev spaces. For $s \in \mathbb{R}$ and $R > 0$, write

$$\dot{H}_R^s(X) := \{u \in H_R^s(M) : \text{supp}(u) \subseteq X\}, \quad \text{and} \quad \overline{H}_R^s(X) := H_R^s(M) / \dot{H}_R^s(M \setminus X).$$

An approximation argument shows that $C_c^\infty(X^\circ) \subseteq \dot{H}_R^s(X)$ is dense for any $s \in \mathbb{R}$, see [31, Theorem 3.29]. We equip these Sobolev spaces with Hilbert space structure induced from $H_R^s(M)$, i.e. $\dot{H}_R^s(X) \subseteq H_R^s(M)$ as a subspace and $\overline{H}_R^s(X)$ as a quotient. We note that for $s = 0$, $\dot{H}_R^0(X) = \overline{H}_R^0(X) = L^2(X)$. The L^2 -pairing between $\dot{H}_R^s(X)$ and $\overline{H}_R^{-s}(X)$ is a perfect pairing and induces an isomorphism $\dot{H}_R^s(X)^* \cong \overline{H}_R^{-s}(X)$ (uniformly in R). For $R = 1$, we omit R from the notation. We remark that any pseudodifferential operator with parameter $A \in \Psi_{\text{cl}}^m(M; \Gamma)$ induces a continuous operator

$$A_X : \dot{H}^s(X) \rightarrow \overline{H}^{s-m}(X),$$

defined by the composition

$$\dot{H}^s(X) \hookrightarrow H_c^s(M) \xrightarrow{A} H_{\text{loc}}^{s-m}(M) \rightarrow \overline{H}^{s-m}(X).$$

Here the last map is the quotient map, $H_c^s(M)$ denotes the space of compactly supported distributions that are s -Sobolev regular and $H_{\text{loc}}^{s-m}(M)$ denotes the space of distributions that are locally $s-m$ -Sobolev regular.

Let us make two remarks regarding the operator $A_X : \dot{H}^s(X) \rightarrow \overline{H}^{s-m}(X)$. Firstly, for any $s \in \mathbb{R}$ density ensures that $A_X : \dot{H}^s(X) \rightarrow \overline{H}^{s-m}(X)$ is determined by continuity and the restriction $A_X : C_c^\infty(X^\circ) \rightarrow C^\infty(X)$. Secondly, if $s = -m/2$ and A is formally self-adjoint, then A_X is determined from the polarization identity by the continuous quadratic form

$$q_{A_X}(u) := \langle u, Au \rangle_{L^2}, \quad u \in \dot{H}^{-m/2}(X),$$

defined from A and the perfect L^2 -pairing $\dot{H}^{m/2}(X) \times \overline{H}^{-m/2}(X) \rightarrow \mathbb{C}$.

The analogue of Theorem 2.23 for Q_X is the following theorem.

Theorem 4.1. *Let X be a compact n -dimensional manifold with C^0 -boundary and d a distance function on X such that d^2 is regular at the diagonal. Set $\mu = (n+1)/2$. Then the family of operators*

$$Q_X := Q_M|_X : \dot{H}^{-\mu}(X) \rightarrow \overline{H}^\mu(X),$$

is a well defined family of Fredholm operators for all $R \in \mathbb{C} \setminus \{0\}$. For some $R_0 \geq 0$, $Q_X(R)$ is invertible for all $R \in \Gamma_{\pi/(n+1)}(R_0)$. Moreover, the following holds:

a) *The family of operators*

$$(Q_X(R) : \dot{H}^{-\mu}(X) \rightarrow \overline{H}^\mu(X))_{R \in \mathbb{C} \setminus \{0\}},$$

depends holomorphically on $R \in \mathbb{C} \setminus \{0\}$. Moreover, the holomorphic family

$$(Q_X(R)^{-1} : \overline{H}^\mu(X) \rightarrow \dot{H}^{-\mu}(X))_{R \in \Gamma_{\pi/(n+1)}(R_0)},$$

extends meromorphically to $R \in \mathbb{C} \setminus \{0\}$.

b) *There are $C, R_0 > 0$ such that*

$$C^{-1} \|f\|_{\dot{H}_{|R|}^{-\mu}(X)}^2 \leq \text{Re} \langle f, Q_X(R)f \rangle_{L^2} \leq C \|f\|_{\dot{H}_{|R|}^{-\mu}(X)}^2,$$

for $R \in \Gamma_{\pi/(n+1)}(R_0)$ and $f \in \dot{H}^{-\mu}(X)$. In particular, for $R \in \Gamma_{\pi/(n+1)}(R_0)$, the operator $Q_X(R)$ is coercive on L^2 in the form sense and the sesquilinear form $\langle \cdot, Q_X(R) \cdot \rangle_{L^2}$ is uniformly equivalent to the inner product of $\dot{H}_R^{-\mu}(X)$.

Proof. We first prove part b). This is a direct consequence of adapting Theorem 2.23, part b), to compactly supported distributions that are $-\mu$ -Sobolev regular and using that $\dot{H}_{|R|}^s(X) \subseteq H_{|R|}^s(M)$ is an isometric inclusion.

To prove part a), we note that for $R \in \mathbb{C} \setminus \{0\}$, $Q_M(R)$ is a lower-order perturbation of $Q_M(R_0)$ for any $R_0 \gg 0$. Therefore, the Rellich lemma implies that the quadratic form

$$q_{Q,R}(u) := \langle u, Q(R)u \rangle_{L^2} \equiv \langle u, Q_X(R)u \rangle_{L^2}, \quad u \in H^{-\mu}(X),$$

is a compact perturbation of q_{Q,R_0} . Therefore, the difference

$$Q_X(R) - Q_X(R_0) : \dot{H}^{-\mu}(X) \rightarrow \overline{H}^\mu(X),$$

is a compact operator. Since part b) implies that $Q_X(R_0)$ is invertible for a large enough $R_0 \gg 0$, we conclude that $(Q_X(R) : \dot{H}^{-\mu}(X) \rightarrow \overline{H}^\mu(X))_{R \in \mathbb{C} \setminus \{0\}}$ is a Fredholm family.

It remains to prove the assertion for the inverse of $Q_X(R)$. The family of operators $(Q_X(R) : \dot{H}^{-\mu}(X) \rightarrow \overline{H}^\mu(X))_{R \in \mathbb{C} \setminus \{0\}}$ is obtained from the holomorphic family of operators $(Q_M(R) : H_c^{-\mu}(M) \rightarrow H_{\text{loc}}^\mu(M))_{R \in \mathbb{C} \setminus \{0\}}$ (see Theorem 2.23, part a) via inclusions and projections, so it is also holomorphic. As such, $(Q_X(R)^{-1} : \overline{H}^\mu(X) \rightarrow \dot{H}^{-\mu}(X))_{R \in \Gamma_{\pi/(n+1)}(R_0)}$ extends meromorphically to $\mathbb{C} \setminus \{0\}$ by the meromorphic Fredholm theorem (see Appendix D). \square

Similarly to the ideas in Section 3, we shall transfer the results of Theorem 4.1 to the operator \mathcal{Z}_X using property (MR). For a domain, let us make the notion of property (MR) more precise.

Definition 4.2. Let X be a compact manifold with C^0 -boundary and d a distance function on X such that d^2 is regular at the diagonal. For a sector $[1, \infty) \subseteq \Gamma \subseteq \mathbb{C}$, we say that d has *property (MR)* on Γ if (X, d) is isometrically embedded as a domain with smooth boundary in a manifold M equipped with a distance function d_M , such that d_M^2 is regular at the diagonal and and for any $R \in \Gamma$, $L(R)$ extends to a continuous mapping $H_c^{-\mu}(M) \rightarrow H_{\text{loc}}^\mu(M)$ with

$$\|L(R)\|_{\dot{H}^{-\mu}(K) \rightarrow \overline{H}^\mu(K')} = O(\text{Re}(R)^{-\infty}), \quad \text{as } \text{Re}(R) \rightarrow +\infty \text{ in } \Gamma,$$

for any compact subsets $K, K' \subseteq M$.

If, for any compact subsets $K, K' \subseteq M$, the operator $L(R) : \dot{H}^{-\mu}(K) \rightarrow \overline{H}^\mu(K')$ is compact for $R \in \Gamma$ and

$$\Gamma \ni R \mapsto L(R) \in \mathbb{K}(\dot{H}^{-\mu}(K), \overline{H}^\mu(K)),$$

is holomorphic in norm sense, we say that d has *property (SMR)* on Γ .

In the absence of a boundary, the definition of property (MR) for a manifold with boundary (Definition 4.2) is readily seen to be equivalent to property (MR) for a compact manifold (Definition 3.3). The reader is encouraged to think of the definition of property (MR) for a manifold with boundary as the distance function having ‘‘property (MR) on a neighborhood of the manifold with boundary’’. Let us consider two examples of distance functions with property (SMR).

Example 4.3 (Domains in Riemannian manifolds with small diameter). Assume that $X \subseteq M$ is a compact domain with C^0 -boundary in a Riemannian manifold with geodesic distance $d_{\text{geo}, M}$. If the diameter of X is strictly smaller than the injectivity radius of M , $d_{\text{geo}, M}^2$ is smooth on a neighborhood of X . The same argument as in Proposition 3.4 shows that the distance function $d_{\text{geo}} := d_{\text{geo}, M}|_X$ on X has property (SMR) on $\mathbb{C} \setminus \{0\}$. In fact, in this case, $L \in \Psi^{-\infty}(X; \mathbb{C}_+)$ and $L(R) \in \Psi^{-\infty}(X)$ for any $R \in \mathbb{C} \setminus \{0\}$.

Example 4.4 (Submanifolds with boundary). Assume that X is a compact manifold with C^0 -boundary embedded in a manifold $i : X \rightarrow W$ and d_W is a distance function on W such that the square d_W^2 is smooth on $W \times W$ and regular at the diagonal. This arises for instance for $W = \mathbb{R}^N$, $W = \mathbb{H}_{N, \mathbb{R}}$ or $W = \mathbb{H}_{N, \mathbb{C}}$, for some $N \in \mathbb{N}$, with their geodesic distance.

The subspace distance function $d : X \times X \rightarrow [0, \infty)$, $d(x, y) := d_W(i(x), i(y))$ will then satisfy

- i) d^2 is smooth on $X \times X$ and regular at the diagonal.
- ii) $L \in \Psi^{-\infty}(X; \mathbb{C}_+)$ and $L(R) \in \Psi^{-\infty}(X)$ for any $R \in \mathbb{C} \setminus \{0\}$.
- iii) d has property (SMR) on $\mathbb{C} \setminus \{0\}$.

This follows by the same arguments as in Proposition 3.4 by choosing a submanifold $M \subseteq W$ in which $i(X)$ is a compact domain (with C^0 -boundary).

For a distance function with property (MR), we tacitly assume that the manifold with C^0 -boundary is embedded into the manifold M implementing property (MR). The next result is proven exactly as Theorem 3.7 but using Theorem 4.1 instead of Corollary 2.22.

Theorem 4.5. *Let X be a compact n -dimensional manifold with C^0 -boundary and d a distance function on X with property (MR) on the sector Γ . Then there is an $R_0 \geq 0$ such that*

$$\mathcal{Z}_X(R) : \dot{H}^{-\mu}(X) \rightarrow \overline{H}^\mu(X),$$

is invertible for all $R \in \Gamma \cap \Gamma_{\pi/(n+1)}(R_0)$. Moreover,

$$\mathcal{Z}_X^{-1} = Q_X^{-1} + \mathcal{R}_X,$$

where Q_X^{-1} is the inverse of Q_X (existing by Theorem 4.1) and $\mathcal{R} : \overline{H}^\mu(X) \rightarrow \dot{H}^{-\mu}(X)$ is a family of operators such that

$$\|\mathcal{R}\|_{\overline{H}^\mu(X) \rightarrow \dot{H}^{-\mu}(X)} = O(\text{Re}(R)^{-\infty}), \quad \text{as } \text{Re}(R) \rightarrow +\infty \text{ in } \Gamma \cap \Gamma_{\pi/(n+1)}(R_0).$$

Moreover, there is a $C > 0$ such that

$$(30) \quad C^{-1} \|f\|_{\dot{H}_{|R|}^{-\mu}(X)}^2 \leq \text{Re}\langle f, \mathcal{Z}_X(R)f \rangle_{L^2} \leq C \|f\|_{\dot{H}_{|R|}^{-\mu}(X)}^2,$$

for $R \in \Gamma \cap \Gamma_{\pi/(n+1)}(R_0)$ and $f \in \dot{H}^{-\mu}(X)$. In particular, for $R \in \Gamma \cap \Gamma_{\pi/(n+1)}(R_0)$, $\text{Re}\mathcal{Z}_X(R)$ is positive in form sense on $L^2(X)$ for the $H^{-\mu}$ -norm.

The next result poses an obstruction to property (MR) for distance functions on manifolds with boundary. It is proven in the same way as Corollary 3.10 but using Theorem 4.5 instead of Theorem 3.7.

Corollary 4.6. *Assume that d is a distance function on a manifold with C^0 -boundary X with property (MR) on $[1, \infty)$. Then there exists an $R_0 \geq 0$ such that for all finite subsets $F \subseteq X$ the $|F| \times |F|$ -matrix $(e^{-Rd(x,y)})_{x,y \in F}$ is positive definite for all $R > R_0$.*

Using Theorem 4.1 instead of Theorem 3.7, the next result is proven ad verbatim as Theorem 3.12.

Theorem 4.7. *Let X be an n -dimensional compact manifold with C^0 -boundary and assume that distance function d has property (SMR) on Γ . There is an $R_0 \geq 0$ such that the operator*

$$\mathcal{Z}_X(R) : \dot{H}^{-\mu}(X) \rightarrow \overline{H}^\mu(X),$$

is a well defined Fredholm operator for all $R \in \Gamma$ and invertible for $R \in \Gamma \cap \Gamma_{\pi/(n+1)}(R_0)$. Moreover, the operator $\mathcal{Z}_X(R) : \dot{H}^{-\mu}(X) \rightarrow \overline{H}^\mu(X)$ depends holomorphically on R in Γ and $\mathcal{Z}_X(R)^{-1} : \overline{H}^\mu(X) \rightarrow \dot{H}^{-\mu}(X)$ depends meromorphically on $R \in \Gamma$.

Similarly to Corollary 3.5, we deduce the following special case of Theorem 4.7.

Corollary 4.8. *Let X be an n -dimensional compact manifold with C^0 -boundary and assume that d is a distance function satisfying that d^2 is smooth on $X \times X$ and regular at the diagonal (e.g. as in Example 4.3 or 4.4). Then the family of operators*

$$(\mathcal{Z}_X(R) : \dot{H}^{-\mu}(X) \rightarrow \overline{H}^\mu(X))_{R \in \mathbb{C} \setminus \{0\}},$$

depends holomorphically on $R \in \mathbb{C} \setminus \{0\}$ and for some R_0 the family

$$(\mathcal{Z}_X(R)^{-1} : \overline{H}^\mu(X) \rightarrow \dot{H}^{-\mu}(X))_{R \in \Gamma_{\pi/(n+1)}(R_0)},$$

is holomorphic and extends meromorphically to $R \in \mathbb{C} \setminus \{0\}$.

The above results in this subsection provide a functional analytic description of \mathcal{Z} and its dependence on R . The following result gives a slightly more precise description of \mathcal{Z} in terms of an extended Boutet de Monvel calculus; for its proof we refer to Appendix C. This result requires the boundary to be smooth. Let us remark that while this result describes the inverse of \mathcal{Z} in terms of a well studied pseudodifferential calculus, we merely include it as an observation as it is not used for computations below. The next section is devoted to an even more direct asymptotic construction of the inverse of \mathcal{Z} that is put to use in the subsequent section.

Theorem 4.9. *Let X be an n -dimensional compact manifold with boundary and consider a distance function d such that d^2 is regular at the diagonal. Write $\mu := (n+1)/2$. The operator*

$$Q_X : \dot{H}^{-\mu}(X) \rightarrow \overline{H}^\mu(X),$$

is an elliptic pseudodifferential operator with parameter R of order $-n-1$, type $-(n+1)/2$ and factorization index $-\infty$ in the Boutet de Monvel calculus. In particular, Q_X admits a classical parameter dependent parametrrix in an extended Boutet de Monvel calculus (made precise in Appendix C).

If moreover d has property (MR) on a sector Γ , then for $R \in \Gamma \cap \Gamma_{\pi/(n+1)}(R_0)$, \mathcal{Z}^{-1} differs from a classical parameter dependent operator in an extended Boutet de Monvel calculus by an operator $\mathcal{R} : \overline{H}^\mu(X) \rightarrow \dot{H}^{-\mu}(X)$ of infinitely low order in R in the sense that

$$\|\mathcal{R}\|_{\overline{H}^\mu(X) \rightarrow \dot{H}^{-\mu}(X)} = O(\operatorname{Re}(R)^{-\infty}), \quad \text{as } \operatorname{Re}(R) \rightarrow +\infty \text{ in } \Gamma \cap \Gamma_{\pi/(n+1)}(R_0).$$

Remark 4.10. If d is the distance function pulled back along an embedding $X \hookrightarrow W$, where the distance function on W is smooth off-diagonally, then also \mathcal{Z} is an elliptic pseudodifferential operator with parameter R of order $-n-1$, type $-(n+1)/2$ and factorization index $-\infty$ in the Boutet de Monvel calculus and admits a classical parameter dependent parametrrix in an extended Boutet de Monvel calculus. In particular, \mathcal{Z}_X^{-1} is a classical parameter dependent operator in an extended Boutet de Monvel calculus if X is a domain in Euclidean space or real/complex hyperbolic space.

5. STRUCTURE OF THE INVERSE OPERATOR IN THE PRESENCE OF A BOUNDARY

Consider a compact manifold with boundary X equipped with a distance function d whose square is regular at the diagonal. As above, we set $\mu := (n+1)/2$ where $n := \dim(X)$. If d has property (MR), Theorem 4.5 ensures that computations for Q_X relate to computations for \mathcal{Z}_X up to a term of infinitely low order in R (as $\operatorname{Re}(R) \rightarrow \infty$), so we focus on the operator Q_X . As proved in Theorem 4.1 above, the localized operator $Q_X : \dot{H}^{-\mu}(X) \rightarrow \overline{H}^\mu(X)$ is an isomorphism for large enough R in the sector $\Gamma_{\pi/(n+1)}$. We shall now describe the inverse of Q_X in more precise terms under the assumption that the boundary is smooth. The inverse $Q_X^{-1} : \overline{H}^\mu(X) \rightarrow \dot{H}^{-\mu}(X)$ will be computed as a sum of

- a pseudodifferential operator (with parameter) in the interior;
- a composition of two *mixed-regularity* pseudodifferential operator near the boundary, where the two factors are obtained from inverting a Wiener-Hopf factorization of the magnitude operator at the boundary; as well as
- an error term that acts as order 2μ , mapping $\overline{H}^\mu(X) \rightarrow \dot{H}^{-\mu+1}(X)$, whose norm is $O(|R|^{-\infty})$ as $R \rightarrow \infty$.

Asymptotically, only the two first terms play a role, and in the next section we compute the asymptotics of conditional expectations from the symbols of these first two terms. To describe the decomposition, we shall need further terminology.

5.1. Mixed-regularity symbols and Sobolev spaces.

Definition 5.1. Let $s, t \in \mathbb{R}$ and $n \in \mathbb{N}_{>0}$. Write coordinates in \mathbb{R}^n as $\xi = (\xi', \xi_n) \in \mathbb{R}^{n-1} \times \mathbb{R}$. We define the Sobolev space of mixed-regularity (s, t) as

$$H^{s,t}(\mathbb{R}^n) := \left\{ f \in \mathcal{S}'(\mathbb{R}^n) : \int_{\mathbb{R}^n} \langle \xi \rangle^s \langle \xi' \rangle^t |\hat{f}(\xi)|^2 d\xi < \infty \right\}.$$

If $\Omega \subseteq \mathbb{R}^n$ is a domain (so $\Omega = \overline{\Omega^\circ}$), we define

$$\dot{H}^{s,t}(\Omega) := \{f \in H^{s,t}(\mathbb{R}^n) : \operatorname{supp}(f) \subseteq \Omega\}, \quad \text{and} \quad \overline{H}^{s,t}(\Omega) := H^{s,t}(\mathbb{R}^n)/H^{s,t}(\overline{\Omega^c}).$$

We also define $\dot{H}_c^{s,t}(\Omega)$ and $\overline{H}_c^{s,t}(\Omega)$ as the elements with compact support. The local Sobolev spaces of mixed-regularity are defined as

$$\begin{aligned} \dot{H}_{\text{loc}}^{s,t}(\Omega) &:= \{f \in \mathcal{S}'(\mathbb{R}^n) : \chi f \in \dot{H}^{s,t}(\Omega) \forall \chi \in C_c^\infty(\mathbb{R}^n)\}, \\ \overline{H}_{\text{loc}}^{s,t}(\Omega) &:= H_{\text{loc}}^{s,t}(\mathbb{R}^n)/H_{\text{loc}}^{s,t}(\overline{\Omega^c}). \end{aligned}$$

We note that $\dot{H}^{s,t}(\Omega) \subseteq H^{s,t}(\mathbb{R}^n)$ is a closed subspace. We call the quotient mapping $\dot{H}^{s,t}(\mathbb{R}^n) \rightarrow \overline{H}^{s,t}(\Omega)$ the restriction mapping. For large enough $s > 0$, we can identify $\dot{H}^{s,t}(\Omega)$ with a subspace of $\overline{H}^{s,t}(\Omega)$ (but not for small $s < 0$). A standard computations with Fourier transforms shows that the the identity operator induces continuous mappings $\overline{H}^{s,t}(\Omega) \rightarrow \overline{H}^{s',t'}(\Omega)$ and $\dot{H}^{s,t}(\Omega) \rightarrow \dot{H}^{s',t'}(\Omega)$ if and only if $s \geq s'$ and $t \geq t'$ (which is locally compact if and only if $s > s'$ and $t > t'$).

Definition 5.2. Let $u, m \in \mathbb{R}$ and $n \in \mathbb{N}_{>0}$. Let $\Gamma \subseteq \mathbb{C}$ be a sector and $U \subseteq \mathbb{R}^n$ an open subset. Write coordinates in \mathbb{R}^n as $\xi = (\xi', \xi_n) \in \mathbb{R}^{n-1} \times \mathbb{R}$. We say that $a \in C^\infty(U \times \mathbb{R}^n \times \Gamma)$ is a symbol with parameter of mixed-regularity (u, m) if for any compact $K \subseteq U$, $k \in \mathbb{N}$, $\alpha \in \mathbb{N}^n$ and $\beta \in \mathbb{N}^n$ there is a constant $C > 0$ such that

$$\sup_{x \in K} \left| \partial_{\xi_n}^k \partial_{(\xi', R)}^\beta \partial_x^\alpha a(x, \xi', \xi_n, R) \right| \leq C \langle (\xi, R) \rangle^{u-k} \langle (\xi', R) \rangle^{m-|\beta|},$$

for all $(\xi, R) \in \mathbb{R}^{n+1}$. We let $S^{u,m}(U; \Gamma)$ denote the space of symbols with parameter of mixed-regularity (u, m) . We set $S^{u,-\infty}(U; \Gamma) := \bigcap_{m \in \mathbb{R}} S^{u,m}(U; \Gamma)$.

For $a \in S^{u,m}(U; \Gamma)$ we define

$$\operatorname{Op}(a) : C_c^\infty(U) \rightarrow C^\infty(U), \quad \operatorname{Op}(a)f(x) := \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} a(x, \xi, R) \hat{f}(\xi) d\xi.$$

Let $\Psi^{u,m}(U; \Gamma)$ denote the linear space of operators $C_c^\infty(U) \rightarrow C^\infty(U)$ spanned by $\{\operatorname{Op}(a) : a \in S^{u,m}(U; \Gamma)\}$ and smoothing operators with parameter. We set $\Psi^{u,-\infty}(U; \Gamma) := \bigcap_{m \in \mathbb{R}} \Psi^{u,m}(U; \Gamma)$.

Example 5.3. Assume that $0 \notin \Gamma$. Suppose that $a(x, \xi, R) = b(x, \xi', R)(\xi_n - h(x, \xi', R))^u$ where b is a homogeneous symbol with parameter of order m and h is a homogeneous symbol with parameter of order 1. A short computation shows that $\partial_{\xi_n}^k \partial_{(\xi', R)}^\beta \partial_x^\alpha a$ is a sum of terms of the form $\tilde{b}(x, \xi', R)(\xi_n - h(x, \xi', R))^{u-k-l}$ where \tilde{b} is homogeneous of order $m - |\beta| + l$. Therefore $a \in S^{u, m}(U; \Gamma)$, and in fact $a \in S^{u_0, m+u-u_0}(U; \Gamma)$ for any $u_0 \leq u$.

We note that since differentiation in the (ξ', R) -direction improves the order of decay, we have for $a \in S^{u, m}(U; \Gamma)$ and $f \in C_c^\infty(U)$ that

$$Op(a)f(x) := \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n \times \mathbb{R}^{n-1}} a(x, \xi, R) \mathcal{F}_{y_n \rightarrow \xi_n} f(y', \xi_n) e^{i\xi_n x_n + i\xi'(x'-y)'} dy' d\xi,$$

as an oscillatory integral. Moreover, we can conclude the following result from standard techniques of oscillatory integrals (cf. [38, Chapter I.1]).

Proposition 5.4. *Assume that $A \in \Psi^{u, m}(U; \Gamma)$. Then for any $\chi, \chi' \in C^\infty(U)$ with $\chi\chi' = 0$ it holds that $\chi A \chi' \in \Psi^{u, -\infty}(U; \Gamma)$. In particular, if $\chi, \chi' \in C_c^\infty(U)$ satisfies $\chi\chi' = 0$ then $\|\chi A \chi'\|_{H^s \rightarrow H^{s'}} = O(|R|^{-\infty})$ as $R \rightarrow \infty$ for all $s \geq s' + u$.*

The last statement of the proposition follows from the next theorem.

Theorem 5.5. *Let $u, m, s, t, s', t' \in \mathbb{R}$ and $n \in \mathbb{N}_{>0}$. Let $A \in \Psi^{u, m}(\mathbb{R}^n; \Gamma)$. Then A extends to a continuous operator*

$$A : H_c^{s, t}(\mathbb{R}^n) \rightarrow H_{\text{loc}}^{s', t'}(\mathbb{R}^n),$$

if $s \geq s' + u$ and $t \geq t' + m$. In this case, we have for any $\chi, \chi' \in C_c^\infty(\mathbb{R}^n)$ that there exists a $C = C(s, s', A, \chi, \chi') > 0$

$$\|\chi A \chi'\|_{H^{s, t}(\mathbb{R}^n) \rightarrow H^{s', t'}(\mathbb{R}^n)} \leq C(1 + |R|)^{t-t+m}.$$

Proof. The first part follows from the Calderón-Vaillancourt theorem. The second part follows from noting that Calderón-Vaillancourt's theorem proves the case $t' + m = t$ and for $A \in \Psi^{u, m}(\mathbb{R}^n; \Gamma)$ compactly supported, then $(R^2 + \Delta')^{(t-t'-m)/2} A \in \Psi^{u, t-t'}(\mathbb{R}^n; \Gamma)$ (where Δ' denotes the Laplacian in the x' -direction). \square

Theorem 5.6. *Let $u, m, s, t, s', t' \in \mathbb{R}$ and $n \in \mathbb{N}_{>0}$. Let $a \in S^{u, m}(\mathbb{R}^n; \Gamma)$ be compactly supported in the x -direction and assume that a extends to a holomorphic function in $\text{Im}(\xi_n) < 0$. Set $A := Op(a)$. Then A restricts to and induces, respectively, continuous operators*

$$\dot{A} : \dot{H}^{s, t}(\mathbb{R}_+^n) \rightarrow \dot{H}^{s', t'}(\mathbb{R}_+^n) \quad \text{and} \quad \bar{A} : \bar{H}^{s, t}(\mathbb{R}_-^n) \rightarrow \bar{H}^{s', t'}(\mathbb{R}_-^n),$$

if $s \geq s' + u$ and $t \geq t' + m$.

Proof. Note that \bar{A} is well defined as the operator induced from $A : H^{s, t}(\mathbb{R}^n) \rightarrow H^{s', t'}(\mathbb{R}^n)$ since \dot{A} preserves supports in \mathbb{R}_+^n . The result follows from the Paley-Wiener theorem. \square

Remark 5.7. Under the assumptions of Theorem 5.6, Theorem 5.5 implies that for any s and s' with $s \geq s' + u$ there is a $C > 0$ such that for $t \geq t' + m$

$$\|A\|_{\dot{H}^{s, t}(\mathbb{R}_+^n) \rightarrow \dot{H}^{s', t'}(\mathbb{R}_+^n)} \leq C(1 + |R|)^{t-t+m} \quad \text{and} \quad \|A\|_{\bar{H}^{s, t}(\mathbb{R}_-^n) \rightarrow \bar{H}^{s', t'}(\mathbb{R}_-^n)} \leq C(1 + |R|)^{t-t+m}.$$

Definition 5.8. Let $u \in \mathbb{R}$. Consider a sequence $(a_j)_{j \in \mathbb{N}}$ of mixed-regularity symbols with $a_j \in S^{u, m_j}(U; \Gamma)$ for a sequence $m_j \rightarrow -\infty$. Set $m := \max_j m_j$. If $a \in S^{u, m}(U; \Gamma)$ satisfies that for any N , there is an M such that $a - \sum_{j=0}^M a_j \in S^{u, -N}(U; \Gamma)$, we write

$$a \sim \sum_{j=0}^{\infty} a_j,$$

and call a the asymptotic sum of $(a_j)_{j \in \mathbb{N}}$.

Proposition 5.9. *Let $u \in \mathbb{R}$. For any sequence $(a_j)_{j \in \mathbb{N}}$ of mixed symbols with $a_j \in S^{u, m_j}(U; \Gamma)$ for a sequence $m_j \rightarrow -\infty$, the asymptotic sum $a \sim \sum_{j=0}^{\infty} a_j$ exists in $S^{u, m}(U; \Gamma)$, where $m := \max_j m_j$. The asymptotic sum is uniquely determined modulo $S^{u, -\infty}(U; \Gamma)$.*

Proof. The proof of this proposition is carried out ad verbatim as in [23, Proposition 18.1.3] upon replacing the ξ in [23] with (ξ', R) . \square

Again, using that differentiation in the (ξ, R) -direction improves the order of decay, we can conclude several results from the standard situation (cf. [38, Chapter I]). For instance, the analogue of [38, Theorem 3.1, Chapter I.3] extends modulo $\Psi^{u, -\infty}$ to $\Psi^{u, m}$ which implies asymptotic expansions of products and adjoints.

Proposition 5.10. *Let $A \in \Psi^{u, m}(U; \Gamma)$ and $B \in \Psi^{u', m'}(U; \Gamma)$ be mixed-regularity pseudodifferential operators out of which at least one is properly supported. Then $AB \in \Psi^{u+u', m+m'}(U; \Gamma)$ is a mixed-regularity pseudodifferential operator. Moreover, if $a \in S^{u, m}(U; \Gamma)$ and $b \in S^{u', m'}(U; \Gamma)$ are symbols with $A - \text{Op}(a)$ and $B - \text{Op}(b)$ being smoothing with parameter, then $AB - \text{Op}(c)$ is smoothing with parameter where $c \in S^{u+u', m+m'}(U; \Gamma)$ is uniquely determined modulo $S^{u+u', -\infty}(U; \Gamma)$ as the asymptotic sum*

$$c \sim \sum_{\alpha \in \mathbb{N}^n} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} a D_x^{\alpha} b.$$

5.2. Wiener-Hopf factorization of Q_X near the boundary. Using the machinery of the previous subsection, we shall now factorize the operator Q_X near the boundary into factors that extend holomorphically into the upper respectively lower half-plane, and use Theorem 5.6 to (near the boundary) invert these individual factors as operators $\dot{H}^{-\mu}(X) \rightarrow L^2(X)$ and $L^2(X) \rightarrow \overline{H}^{\mu}(X)$, respectively. The reader should recall the structure of the full symbol of Q from Theorem 2.9. We shorten the notation and write C^j for $C_{d^2}^j$, where $C_{d^2}^j$ are the Taylor coefficients of d^2 as in Equation (10).

As above, we consider a compact manifold with boundary X a distance function d on X whose square is regular at the diagonal (cf. Definition 2.2). We tacitly fix a manifold M containing X as a smooth compact domain to which d extends as a distance function whose square is regular at the diagonal. In particular, we can Taylor expand d^2 near any point in the diagonal as in Equation (10) and its Taylor coefficients enter the full symbol of Q as in Theorem 2.9. To study the behavior at the boundary, we first reduce to the model case that $X = \partial X \times [0, \infty)$, as a domain in $\partial X \times \mathbb{R}$. We remark that $\partial X \times [0, \infty)$ is not compact, but we shall later on only use the constructed operators in a form localized to near the compact boundary.

Proposition 5.11. *Let X be a compact manifold with boundary with a distance function d whose square is regular at the diagonal, embedded into a manifold M as in the preceding paragraph. Consider the compact manifold $Y = \partial X$ and choose a tubular neighborhood $U \subseteq M$ of ∂X and a diffeomorphism $\varphi : U \rightarrow Y \times (-1, 1)$, with $\partial X = \varphi^{-1}(Y \times \{0\})$. Then there exists a classical elliptic pseudodifferential operator with parameter $Q^{\partial} \in \Psi_{\text{cl}}^{-n-1}(Y \times \mathbb{R}; \mathbb{C}_+)$ and a number $\varepsilon > 0$ such that*

- Q^{∂} is translation invariant outside a compact subset in the sense that there exists a $t_0 > 0$ such that if $f \in C_c^{\infty}(Y \times \mathbb{R})$ is supported in $\{(y, t) : \pm t > t_0\}$ then $[Q^{\partial} f](\cdot \mp s) = Q^{\partial}[f(\cdot \mp s)]$ for all $s > 0$.
- For all $f \in C_c^{\infty}(Y \times (-\varepsilon, \varepsilon))$ it holds that $Q(\varphi^* f)$ is supported in U and

$$Q(\varphi^* f) = \varphi^*(Q^{\partial} f).$$

- the principal symbol of Q^{∂} is given by

$$\sigma_{-n-1}(x, \xi, R) = c(R^2 + g(\xi, \xi))^{-\mu},$$

where g is a Riemannian metric on $Y \times \mathbb{R}$ which is translation invariant in the \mathbb{R} -direction outside a compact and coincides with $\varphi^* g_{d^2}$ on $Y \times (-1, 1)$.

Proof. Construct g from interpolating between g_{d^2} near 0 and something at infinity. Construct Q^{∂} from interpolation along the real line between Q near 0 and $(R^2 + \Delta_g)^{-\mu}$ at infinity. The second part follows from that Q by construction has small propagation. \square

Henceforth, we shall fix a choice of Q^{∂} and g as in Proposition 5.11.

Remark 5.12. To fix a choice of a diffeomorphism $\varphi : U \rightarrow Y \times (-1, 1)$ is (up to a self-diffeomorphism of Y) equivalent to choosing a vector field defined near $Y = \partial X$ which is transversal to the boundary. This choice of vector field, or equivalently, the last entry of $\varphi : U \rightarrow Y \times (-1, 1)$, gives rise to a coordinate that we denote by $x_n : U \rightarrow (-1, 1)$. We remark that it is always possible to use the transversal vector field to be the metric normal to the boundary, in which case we have that on ∂X :

$$g_{d^2} = dx_n^2 + g_{\partial X, d^2},$$

where $g_{\partial X, d^2}$ is the induced Riemannian metric on ∂X . For computational purposes, it becomes clumsy to restrict to the case when the transversal vector field is orthogonal to the boundary but for some considerations it simplifies the formulas.

Following the notation of Subsection 5.1, we write x_n for this transversal coordinate and x' for coordinates on ∂X . Similarly, ξ_n denotes the cotangent variable in the transversal direction and ξ' denotes the cotangent variables along ∂X .

We let $q^\partial \in S_{\text{cl}}^{-2\mu}(Y \times \mathbb{R}; \Gamma)$ denote the full symbol of Q^∂ . We note that $q^\partial \sim \sum_{j=0}^{\infty} q_j^\partial$ in $S^{-2\mu}(Y \times \mathbb{R}; \mathbb{C})$ where each $q_j^\partial \in S^{-2\mu-j}(Y \times \mathbb{R}; \mathbb{C})$ is a homogeneous symbol in (ξ, R) of order $-2\mu - j = -n - 1 - j$ and near $t = 0$, we have in any local coordinates on Y that $q_j^\partial = q_j$ where q_j is computed as in Theorem 2.9 using the coordinates induced from Y and φ .

Proposition 5.13. *There are unique homogeneous degree 1 symbols*

$$h_\pm = h_\pm(x, \xi', R) \in S^1(T^*Y \times \mathbb{R}, Y \times \mathbb{R}; \mathbb{C}),$$

that determine the complex solutions to the equation $R^2 + g_x((\xi', \xi_n), (\xi', \xi_n)) = 0$ for fixed (x, ξ', R) with $\pm \text{Im}(\xi_n) > 0$. Furthermore, there is a unique function $h_0 \in C^\infty(Y \times \mathbb{R}, \mathbb{R}_{>0})$ such that

$$R^2 + g(\xi, \xi) = h_0(x)(\xi_n - h_+(x, \xi', R))(\xi_n - h_-(x, \xi', R)).$$

Moreover, we have that

$$h_\pm(x, \xi', R) = -\frac{\xi'(b(x))}{h_0(x)} \pm i \frac{\sqrt{R^2 + g_Y(\xi', \xi') - (\xi'(b))^2}}{\sqrt{h_0(x)}},$$

for suitable b and g_Y determined from the metric.

Proof. It is not hard to see that h_0 and h_\pm are well defined and unique, but let us construct them explicitly. We can decompose

$$(31) \quad g = \begin{pmatrix} h_0 & b \\ b^T & g_Y \end{pmatrix},$$

where g_Y is a metric on Y on each slice, and $b \in C^\infty(Y \times \mathbb{R}, TY)$. Since g is translation invariant outside a compact, h_0 , b and g_Y are translation invariant outside a compact. We have that

$$R^2 + g(\xi, \xi) = R^2 + h_0 \xi_n^2 + 2\xi'(b)\xi_n + g_Y(\xi', \xi').$$

We see that h_0 in the above definition is the h_0 in Equation (31) and that the complex roots are as prescribed. The proposition follows. \square

Theorem 5.14. *Let $q^\partial \in S_{\text{cl}}^{-2\mu}(Y \times \mathbb{R}; \Gamma)$ be as above. Then there exists $q_\pm^\partial \in S^{-\mu, 0}(Y \times \mathbb{R}; \Gamma)$ that are translation invariant outside a compact such that*

- (1) $q_\pm^\partial \in S^{-\mu, 0}(Y \times \mathbb{R}; \Gamma)$ admits asymptotic expansions (in $S^{-\mu, 0}$ in the sense of Definition 5.8 on page 38)

$$q_\pm^\partial \sim \sum_{j=0}^{\infty} q_{\pm, j}^\partial,$$

where $q_{\pm, 0}^\partial \in S^{-\mu, 0}$ and for $j > 0$,

$$q_{\pm, j}^\partial(x, \xi, R) = \sum_{k=-1}^{j-1} b_{\pm, j, k}(x, \xi', R)(\xi_n - h_\pm(x, \xi', R))^{-\mu-j+k} \in S^{-\mu-1, -j+1},$$

where $b_{\pm,j,k}$ is homogeneous of degree $-k$ in (ξ', R) and can be computed by an iterative scheme of partial fraction decompositions as a homogeneous rational function in derivatives of h_0 , h_+ and h_- . The first terms are given by

$$(32) \quad \begin{aligned} q_{+,0}^\partial &= n! \omega_n (\xi_n - h_+)^{-\mu}, \\ q_{-,0}^\partial &= h_0^{-\mu} (\xi_n - h_-)^{-\mu}, \end{aligned}$$

and $q_{\pm,1}$ are computed in Proposition 5.16 below.

- (2) The mixed-regularity symbols $q_\pm^\partial \in S^{-\mu,0}(Y \times \mathbb{R}; \Gamma)$ and $q_{\pm,j}^\partial \in S^{-\mu-1,-j+1}$ admit holomorphic extensions to $\mp \text{Im}(\xi_n) > 0$.
- (3) It holds that

$$q^\partial = \sum_\alpha \frac{1}{\alpha!} \partial_\xi^\alpha q_-^\partial D_x^\alpha q_+^\partial \quad \text{mod } S^{-2\mu,-\infty}.$$

Proof. Let us first massage the statements of the theorem. We want to construct $q_\pm^\partial \sim \sum_{j=0}^\infty q_{\pm,j}^\partial \in S^{-\mu,0}$ admitting holomorphic extensions to $\mp \text{Im}(\xi_n) > 0$ and satisfying $q^\partial = \sum_\alpha \frac{1}{\alpha!} \partial_\xi^\alpha q_-^\partial D_x^\alpha q_+^\partial \text{ mod } S^{-2\mu,-\infty}$. We remark that to ensure item (2), i.e. the holomorphic extension of q_\pm^∂ , it suffices to construct each $q_{\pm,j}^\partial$ so that it admits a holomorphic extension to $\mp \text{Im}(\xi_n) > 0$. We also note that the requirement on the composition is equivalent to

$$(33) \quad q_j^\partial = \sum_{k+l+|\alpha|=j} \frac{1}{\alpha!} \partial_\xi^\alpha q_{-,k}^\partial D_x^\alpha q_{+,l}^\partial.$$

We take the formula (32) as a definition, and note that $q_{\pm,0}^\partial$ satisfy the structural statement in item (1), extends holomorphically to $\mp \text{Im}(\xi_n) > 0$ and $q_0^\partial = q_{-,0}^\partial q_{+,0}^\partial$. Using an idea described in [21], Equation (33) is for $j > 0$ equivalent to

$$\frac{q_{+,j}^\partial}{q_{+,0}^\partial} + \frac{q_{-,j}^\partial}{q_{-,0}^\partial} = \frac{q_j^\partial}{q_0^\partial} - \frac{1}{q_0^\partial} \sum_{\substack{k+l+|\alpha|=j \\ k,l < j}} \frac{1}{\alpha!} \partial_\xi^\alpha q_{-,k}^\partial D_x^\alpha q_{+,l}^\partial$$

We proceed by induction. Assume that we have constructed $q_{\pm,k}$ for $k < j$ satisfying the statements of items (1), (2), and (3) in the relevant degrees. Using Lemma 2.20 and item (1) for $q_{\pm,k}$ for $k < j$, we can use Lemma E.1 to uniquely partial fraction decompose

$$\frac{q_j^\partial}{q_0^\partial} - \frac{1}{q_0^\partial} \sum_{\substack{k+l+|\alpha|=j \\ k,l < j}} \frac{1}{\alpha!} \partial_\xi^\alpha q_{-,k}^\partial D_x^\alpha q_{+,l}^\partial = \mathfrak{q}_{+,j} + \mathfrak{q}_{-,j},$$

where $\mathfrak{q}_{\pm,j}$ by Proposition 2.6 and Theorem 2.9 takes the form

$$\mathfrak{q}_{\pm,j}(x, \xi, R) = \sum_{k=-1}^{j-1} \mathfrak{b}_{\pm,j,k}(x, \xi', R) (\xi_n - h_\pm(x, \xi', R))^{-j+k} \in S^{-1,-j+1},$$

where $\mathfrak{b}_{\pm,j,k}$ is homogeneous of degree $-k$ in (ξ', R) and can be explicitly computed from the results of Appendix E and Theorem 2.9. We now define

$$q_{\pm,j}^\partial := q_{\pm,0}^\partial \mathfrak{q}_{\pm,j},$$

and note that it by construction satisfies items (1), (2), and (3) in the relevant degrees. \square

Remark 5.15. As noted in Proposition 5.13, the symbols h_+ and h_- are directly determined from the metric and the choice of transversal to the boundary. Moreover, as the proof of Theorem 5.14 shows, each of the symbols $b_{\pm,j,k} = b_{\pm,j,k}(x, \xi', R)$ depends

- (1) polynomially on $(C_G^{(\gamma)})_{\gamma \in \cup_{k \leq j} I_k}$ and its derivatives contracted by g_G and $\iota_n g_G$ and its derivatives
- (2) polynomially on h_+ , h_- and rationally on $h_+ - h_-$ and $h_0^{-1/2}$,

The total degree is j where each $C_G^{(\gamma)}$, $\gamma \in I_k$, has degree k , the metric and h_0 have degree zero, h_+ and h_- have degree 1 and x -derivatives increase the order by 1.

Proposition 5.16. *For $n > 1$, the terms $q_{\pm,1}$ appearing in the expansion of q_{\pm} in Theorem 5.14 are given by:*

$$\begin{aligned} q_{+,1}^{\partial} &= n! \omega_n \mathbf{a}_{0,+}(x, \xi', R)(\xi_n - h_+)^{-\mu-1} + n! \omega_n \mathbf{a}_{1,+}(x, \xi', R)(\xi_n - h_+)^{-\mu-2}, \\ q_{-,1}^{\partial} &= \mathbf{a}_{0,-}(x, \xi', R) h_0^{-\mu} (\xi_n - h_-)^{-\mu-1} + \mathbf{a}_{1,-}(x, \xi', R) h_0^{-\mu} (\xi_n - h_-)^{-\mu-2}, \end{aligned}$$

where the homogeneous symbols $\mathbf{a}_{+,0}$ (of degree 0) and $\mathbf{a}_{+,1}$ (of degree 1) are explicitly given in Equation (34) and (35) below, and the homogeneous symbols $\mathbf{a}_{-,0}$ (of degree 0) and $\mathbf{a}_{-,1}$ (of degree 1) are explicitly given in Equation (36) and (37) below.

Proof. Let us first make some preliminary computations with the decomposition from Lemma E.1 using the explicit forms in Corollary E.4. We let ι_n denote contraction by the unit normal. We can write

$$C^3(x, g \otimes \iota_{\xi} g) = C^3(x, g \otimes \iota_n g) \xi_n + C^3(x, g \otimes \iota_{\xi'} g).$$

Using Corollary E.4 we decompose

$$\begin{aligned} C^3(x, g \otimes \iota_{\xi} g) (R^2 + g_d(\xi, \xi))^{-1} &= \frac{h_0^{-1} C^3(x, g \otimes \iota_n g)}{h_+ - h_-} [h_+(\xi_n - h_+)^{-1} - h_-(\xi_n - h_-)^{-1}] \\ &\quad + C^3(x, g \otimes \iota_{\xi'} g) \frac{h_0^{-1}}{h_+ - h_-} [(\xi_n - h_+)^{-1} - (\xi_n - h_-)^{-1}]. \end{aligned}$$

We can write

$$\begin{aligned} C^3(x, \iota_{\xi} g \otimes \iota_{\xi} g \otimes \iota_{\xi} g) &= C^3(x, \iota_n g \otimes \iota_n g \otimes \iota_n g) \xi_n^3 + 3C^3(x, \iota_{\xi'} g \otimes \iota_n g \otimes \iota_n g) \xi_n^2 + \\ &\quad + 3C^3(x, \iota_{\xi'} g \otimes \iota_{\xi'} g \otimes \iota_n g) \xi_n + C^3(x, \iota_{\xi'} g \otimes \iota_{\xi'} g \otimes \iota_{\xi'} g). \end{aligned}$$

Using Corollary E.4 we decompose

$$\begin{aligned} C^3(x, \iota_{\xi} g \otimes \iota_{\xi} g \otimes \iota_{\xi} g) (R^2 + g_d(\xi, \xi))^{-2} &= \\ \frac{C^3(x, \iota_n g \otimes \iota_n g \otimes \iota_n g) h_0^{-2}}{(h_+ - h_-)^2} &\left[h_+^3 (\xi_n - h_+)^{-2} + h_-^3 (\xi_n - h_-)^{-2} + \right. \\ &\quad \left. + \frac{h_+^2 (h_+ - 3h_-)}{h_+ - h_-} (\xi_n - h_+)^{-1} - \frac{h_-^2 (h_- - 3h_+)}{h_+ - h_-} (\xi_n - h_-)^{-1} \right] \\ + \frac{3h_0^{-2} C^3(x, \iota_{\xi'} g \otimes \iota_n g \otimes \iota_n g)}{(h_+ - h_-)^2} &\left[h_+^2 (\xi_n - h_+)^{-2} + h_-^2 (\xi_n - h_-)^{-2} - \right. \\ &\quad \left. - \frac{2h_+ h_-}{h_+ - h_-} ((\xi_n - h_+)^{-1} - (\xi_n - h_-)^{-1}) \right] \\ + \frac{3h_0^{-2} C^3(x, \iota_{\xi'} g \otimes \iota_{\xi'} g \otimes \iota_n g)}{(h_+ - h_-)^2} &\left[h_+ (\xi_n - h_+)^{-2} + h_- (\xi_n - h_-)^{-2} - \right. \\ &\quad \left. - \frac{h_+ + h_-}{h_+ - h_-} ((\xi_n - h_+)^{-1} - (\xi_n - h_-)^{-1}) \right] \\ + \frac{h_0^{-2} C^3(x, \iota_{\xi'} g \otimes \iota_{\xi'} g \otimes \iota_{\xi'} g)}{(h_+ - h_-)^2} &\left[(\xi_n - h_+)^{-2} + (\xi_n - h_-)^{-2} - \right. \\ &\quad \left. - \frac{2}{h_+ - h_-} ((\xi_n - h_+)^{-1} - (\xi_n - h_-)^{-1}) \right]. \end{aligned}$$

Using Proposition 2.14, we compute that

$$\begin{aligned} \frac{q_1^{\partial}}{q_0^{\partial}} - \frac{1}{q_0^{\partial}} \sum_{|\alpha|=1} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} q_{-,0}^{\partial} D_x^{\alpha} q_{+,0}^{\partial} &= \\ = - \frac{3i \mathbf{c}_{1,n} (n^2 - 1)}{n! \omega_n} C^3(x, g \otimes \iota_{\xi} g) (R^2 + g_d(\xi, \xi))^{-1} &+ \\ + \frac{i \mathbf{c}_{1,n} (n+3) 3_{-2}}{n! \omega_n} C^3(x, \iota_{\xi} g \otimes \iota_{\xi} g \otimes \iota_{\xi} g) (R^2 + g_d(\xi, \xi))^{-2} &+ \\ + \frac{i(n+1)^2 h_0}{4} (\nabla_{\xi'} h_- \cdot \nabla_{x'} h_+ - \partial_{x_n} h_+) (R^2 + g_d(\xi, \xi))^{-1}. & \end{aligned}$$

Decomposing as in the paragraphs above, we obtain

$$q_{+,1} = n! \omega_n \mathbf{a}_{0,+}(x, \xi', R) (\xi_n - h_+)^{-\mu-1} + n! \omega_n \mathbf{a}_{1,+}(x, \xi', R) (\xi_n - h_+)^{-\mu-2},$$

where

$$(34) \quad \begin{aligned} \mathbf{a}_{0,+}(x, \xi', R) = & -\frac{3i\mathbf{c}_{1,n}(n^2-1)}{n!\omega_n} (C^3(x, g \otimes \iota_n g)h_+ + C^3(x, g \otimes \iota_{\xi'} g)) \frac{h_0^{-1}}{h_+ - h_-} + \\ & + \frac{i\mathbf{c}_{1,n}(n+3)_{3,-2}}{n!\omega_n} \frac{h_0^{-2}}{(h_+ - h_-)^3} \\ & \left[C^3(x, \iota_n g \otimes \iota_n g \otimes \iota_n g)h_+^2(h_+ - 3h_-) - 6h_+h_-C^3(x, \iota_n g \otimes \iota_n g \otimes \iota_{\xi'} g) - \right. \\ & \left. - 3(h_+ + h_-)C^3(x, \iota_n g \otimes \iota_{\xi'} g \otimes \iota_{\xi'} g) - 2C^3(x, \iota_{\xi'} g \otimes \iota_{\xi'} g \otimes \iota_{\xi'} g) \right] - \\ & + \frac{i(n+1)^2}{4} \frac{(\nabla_{\xi'} h_- \cdot \nabla_{x'} h_+ - \partial_{x_n} h_+)}{h_+ - h_-}, \end{aligned}$$

and

$$(35) \quad \begin{aligned} \mathbf{a}_{1,+}(x, \xi', R) = & \frac{i\mathbf{c}_{1,n}(n+3)_{3,-2}}{n!\omega_n} \frac{h_0^{-2}}{(h_+ - h_-)^2} \\ & \left[C^3(x, \iota_n g \otimes \iota_n g \otimes \iota_n g)h_+^3 + 3h_+^2 C^3(x, \iota_{\xi'} g \otimes \iota_n g \otimes \iota_n g) + \right. \\ & \left. + 3h_+ C^3(x, \iota_{\xi'} g \otimes \iota_{\xi'} g \otimes \iota_n g) + C^3(x, \iota_n g \otimes \iota_{\xi'} g \otimes \iota_{\xi'} g) \right]. \end{aligned}$$

We also obtain

$$q_{-,1} = \mathbf{a}_{0,-}(x, \xi', R) h_0^{-\mu} (\xi_n - h_-)^{-\mu-1} + \mathbf{a}_{1,-}(x, \xi', R) h_0^{-\mu} (\xi_n - h_-)^{-\mu-2},$$

where

$$(36) \quad \begin{aligned} \mathbf{a}_{0,-}(x, \xi', R) = & \frac{3i\mathbf{c}_{1,n}(n^2-1)}{n!\omega_n} (C^3(x, g \otimes \iota_n g)h_- + C^3(x, g \otimes \iota_{\xi'} g)) \frac{h_0^{-1}}{h_+ - h_-} - \\ & - \frac{i\mathbf{c}_{1,n}(n+3)_{3,-2}}{n!\omega_n} \frac{h_0^{-2}}{(h_+ - h_-)^3} \\ & \left[C^3(x, \iota_n g \otimes \iota_n g \otimes \iota_n g)h_-^2(h_- - 3h_+) - 6h_+h_-C^3(x, \iota_n g \otimes \iota_n g \otimes \iota_{\xi'} g) - \right. \\ & \left. - 3(h_+ + h_-)C^3(x, \iota_n g \otimes \iota_{\xi'} g \otimes \iota_{\xi'} g) - 2C^3(x, \iota_{\xi'} g \otimes \iota_{\xi'} g \otimes \iota_{\xi'} g) \right] + \\ & - \frac{i(n+1)^2}{4} \frac{(\nabla_{\xi'} h_- \cdot \nabla_{x'} h_+ - \partial_{x_n} h_+)}{h_+ - h_-}, \end{aligned}$$

and

$$(37) \quad \begin{aligned} \mathbf{a}_{1,-}(x, \xi', R) = & \frac{i\mathbf{c}_{1,n}(n+3)_{3,-2}}{n!\omega_n} \frac{h_0^{-2}}{(h_+ - h_-)^2} \\ & \left[C^3(x, \iota_n g \otimes \iota_n g \otimes \iota_n g)h_-^3 + 3h_-^2 C^3(x, \iota_{\xi'} g \otimes \iota_n g \otimes \iota_n g) + \right. \\ & \left. + 3h_- C^3(x, \iota_{\xi'} g \otimes \iota_{\xi'} g \otimes \iota_n g) + C^3(x, \iota_n g \otimes \iota_{\xi'} g \otimes \iota_{\xi'} g) \right]. \end{aligned}$$

The stated formulas follow. \square

Lemma 5.17. *The operators $Q_{\pm}^{\partial} := Op(q_{\pm}^{\partial}) \in \Psi^{-\mu,0}(Y \times \mathbb{R}; \Gamma)$ satisfy that*

$$(1) \quad Q - Q_- Q_+ \in \Psi^{-2\mu, -\infty}(Y \times \mathbb{R}; \Gamma);$$

- (2) Q_+ restricts to a well defined operator $\dot{H}^{-\mu}(Y \times [0, \infty)) \rightarrow L^2(Y \times [0, \infty))$ which is invertible for large R ;
- (3) Q_- restricts to a well defined operator $L^2(Y \times [0, \infty)) \rightarrow \overline{H}^\mu(Y \times [0, \infty))$ which is invertible for large R .

Proof. Part (1) follows from Proposition 5.10 and Theorem 5.14. Parts (2) and (3) follow from Theorem 5.5 and Theorem 5.6. \square

Definition 5.18. Define $w_{\pm, j} \in S^{\mu, -j}$ inductively by

$$w_{\pm, 0}(x, \xi, R) := (q_{\pm, 0}^\partial)^{-1} = \begin{cases} \frac{1}{n! \omega_n} (\xi_n - h_+(x, \xi', R))^\mu, & \text{for } +, \\ h_0(x)^\mu (\xi_n - h_-(x, \xi', R))^\mu, & \text{for } -, \end{cases}$$

and then

$$w_{\pm, j} := -w_{\pm, 0} \sum_{k+l+|\alpha|=j, l < j} \frac{1}{\alpha!} \partial_\xi^\alpha q_{\pm, k}^\partial D_x^\alpha w_{\pm, l}.$$

We also define $w_\pm := \sum_j w_{\pm, j} \in S^{\mu, 0}(Y \times \mathbb{R}; \Gamma)$ and $W_\pm \in \Psi^{\mu, 0}(Y \times \mathbb{R}; \Gamma)$ is defined as a properly supported modification of $Op(w_\pm)$ with the same full symbol which is translation invariant outside a compact subset.

Lemma 5.19. *Let $w_\pm \in S^{\mu, 0}(Y \times \mathbb{R}; \Gamma)$ be as above. Then*

- (1) *The asymptotic expansion (in $S^{\mu, 0}$ in the sense of Definition 5.8 on page 38) of $w_\pm \in S^{\mu, 0}(Y \times \mathbb{R}; \Gamma)$*

$$w_\pm \sim \sum_{j=0}^{\infty} w_{\pm, j},$$

can for $j > 0$ be expanded in a finite sum

$$w_{\pm, j}(x, \xi, R) = \sum_{k=0}^{j-1} \mathfrak{w}_{\pm, j, k}(x, \xi', R) (\xi_n - h_\pm(x, \xi', R))^{\mu-j+k} \in S^{\mu-1, -j+1},$$

where $\mathfrak{w}_{\pm, j, k}$ is homogeneous of degree $-k$ in (ξ', R) and can be computed by an iterative scheme as a rational function of derivatives of h_0 , h_+ and h_- .

- (2) *The mixed-regularity symbols $w_\pm \in S^{\mu, 0}(Y \times \mathbb{R}; \Gamma)$ and $w_{\pm, j} \in S^{\mu-1, -j+1}$ admit holomorphic extensions to $\mp \text{Im}(\xi_n) > 0$.*
- (3) *The symbols $\mathfrak{w}_{\pm, j, k} = \mathfrak{w}_{\pm, j, k}(x, \xi', R)$ depends*
- polynomially on $(C_G^{(\gamma)})_{\gamma \in \cup_{k \leq j} I_k}$ and its derivatives contracted by g_G and $\iota_n g_G$ and its derivatives*
 - polynomially on h_+ , h_- and rationally on $h_+ - h_-$ and $h_0^{-1/2}$,*
- The total degree is j where each $C_G^{(\gamma)}$, $\gamma \in I_k$, has degree k , the metric and h_0 have degree zero, h_+ and h_- have degree 1 and x -derivatives increase the order by 1.*

Proof. Items (1) and (2) follow from a short induction argument with the construction (in Definition 5.18) and Theorem 5.14. \square

We now compute $w_{\pm, 1}$. By definition, we have that

$$w_{\pm, 1} = -w_{\pm, 0}^2 q_{\pm, 1}^\partial - w_{\pm, 0} \sum_{|\alpha|=1} \partial_\xi^\alpha q_{\pm, 0}^\partial D_x^\alpha w_{\pm, 0}.$$

A short algebraic manipulation with the computation of $q_{\pm, 1}$ from Proposition 5.16 gives the following formulas.

Proposition 5.20. *For $n > 1$, the terms $w_{\pm,1}$ appearing in the expansion of w_{\pm} in Lemma 5.19 are given by:*

$$\begin{aligned} w_{+,1}(x, \xi', \xi_n, R) &= -\frac{1}{n! \omega_n} \mathbf{a}_{0,+}(x, \xi', R) (\xi_n - h_+)^{\mu-1} - \frac{1}{n! \omega_n} \mathbf{a}_{1,+}(x, \xi', R) (\xi_n - h_+)^{\mu-2} - \\ &\quad - \frac{i(n+1)^2}{4 \cdot n! \omega_n} (\partial_{x_n} h_+ - \nabla_{\xi'} h_+ \cdot \nabla_{x'} h_+) (\xi_n - h_+)^{\mu-2}, \\ w_{-,1}(x, \xi', \xi_n, R) &= -\mathbf{a}_{0,-}(x, \xi', R) h_0^\mu (\xi_n - h_-)^{\mu-1} - \mathbf{a}_{1,-}(x, \xi', R) h_0^\mu (\xi_n - h_-)^{\mu-2} - \\ &\quad - \frac{i(n+1)^2}{4} (\partial_{x_n} h_- - \nabla_{\xi'} h_- \cdot \nabla_{x'} h_-) h_0^\mu (\xi_n - h_-)^{\mu-2} - \\ &\quad - \frac{i(n+1)^2}{4} (\partial_{x_n} h_0 - \nabla_{\xi'} h_0 \cdot \nabla_{x'} h_0) h_0^{\mu-1} (\xi_n - h_-)^{\mu-1}, \end{aligned}$$

where the homogeneous symbols $\mathbf{a}_{+,0}$ (of degree 0) and $\mathbf{a}_{+,1}$ (of degree 1) were explicitly given in Equation (34) and (35) above, and the homogeneous symbols $\mathbf{a}_{-,0}$ (of degree 0) and $\mathbf{a}_{-,1}$ (of degree 1) were explicitly given in Equation (36) and (37) above.

Lemma 5.21. *The operators $W_{\pm} \in \Psi^{\mu,0}(Y \times \mathbb{R}; \Gamma)$ satisfy that*

- (1) $1 - W_{\pm} Q_{\pm}^{\partial}, 1 - Q_{\pm}^{\partial} W_{\pm} \in \Psi^{0,-\infty}(Y \times \mathbb{R}; \Gamma)$;
- (2) W_- preserves supports in $Y \times (-\infty, 0]$ and restricts to a well defined operator $\overline{H}^{\mu}(Y \times [0, \infty)) \rightarrow L^2(Y \times [0, \infty))$ which is invertible for large R ;
- (3) W_+ preserves supports in $Y \times [0, \infty)$ and restricts to a well defined operator $L^2(Y \times [0, \infty)) \rightarrow \dot{H}^{-\mu}(Y \times [0, \infty))$ which is invertible for large R .

In particular, the operators

$$\begin{aligned} S_0 &:= 1 - W_+ W_- Q^{\partial} : \dot{H}^{-\mu}(Y \times [0, \infty)) \rightarrow \dot{H}^{-\mu}(Y \times [0, \infty)), \quad \text{and} \\ S_1 &:= 1 - Q^{\partial} W_+ W_- : \overline{H}^{\mu}(Y \times [0, \infty)) \rightarrow \overline{H}^{\mu}(Y \times [0, \infty)), \end{aligned}$$

are normbounded by $O(R^{-\infty})$.

Proof. Part (1) follows from Proposition 5.10 and Lemma 5.19. Parts (2) and (3) follow from Theorem 5.5 and Theorem 5.6. \square

5.3. Decomposition of the inverse magnitude operator.

Theorem 5.22. *Let X be an n -dimensional compact manifold with boundary and d a distance function whose square is regular at the diagonal. Set $\mu := (n+1)/2$. Let $Q_X : \dot{H}^{-\mu}(X) \rightarrow \overline{H}^{\mu}(X)$ denote the restriction of Q_{χ, d^2} to X and let $A \in \Psi_{\text{cl}}^{n+1}(X; \mathbb{C}_+)$ denote a parametrix of Q_{χ, d^2} . For some $R_0 \geq 0$ and any $R \in \Gamma_{\pi/(n+1)}(R_0)$, we can write*

$$Q_X^{-1} = \chi_1 A \chi'_1 + \chi_2 (\varphi^{-1})^* W_+ W_- \varphi^* \chi'_2 + S,$$

where $\chi_1, \chi'_1 \in C_c^\infty(X^\circ)$, and $\chi_2, \chi'_2 \in C^\infty(X)$ are functions supported in a collar neighborhood U_0 of ∂X in X such that

$$\chi_1 + \chi_2 = 1 \quad \text{and} \quad \chi'_j|_{\text{supp}(\chi_j)} = 1, \quad j = 1, 2,$$

$\varphi : \partial X \times [0, 1) \rightarrow U_0$ is a collar identification, and the operators S , W_- and W_+ satisfy the following:

- (1) $S : \overline{H}^{\mu}(X) \rightarrow \dot{H}^{-\mu}(X)$ is a continuous operator with

$$\|S\|_{\overline{H}^{\mu}(X) \rightarrow \dot{H}^{-\mu}(X)} = O(R^{-\infty}), \quad \text{as } R \rightarrow \infty.$$

- (2) $W_+ : L^2(\partial X \times [0, \infty)) \rightarrow \dot{H}^{-\mu}(\partial X \times [0, \infty))$ is the properly supported pseudodifferential operator of mixed-regularity $(\mu, 0)$ from Definitions 5.18 which is invertible for large $R > 0$ and in local coordinates has an asymptotic expansion modulo $S^{\mu, -\infty}$ as in Lemma 5.21 and preserves support in $\partial X \times [0, \infty) \subseteq \partial X \times \mathbb{R}$. Moreover, for $\chi, \chi' \in \mathbb{C} + C_c^\infty(\partial X \times [0, \infty))$ with $\chi \chi' = 0$, it holds that

$$\|\chi W_+ \chi'\|_{L^2(\partial X \times [0, \infty)) \rightarrow \dot{H}^{-\mu}(\partial X \times \mathbb{R})} = O(R^{-\infty}), \quad \text{as } R \rightarrow \infty.$$

- (3) $W_- : \overline{H}^\mu(\partial X \times [0, \infty)) \rightarrow L^2(\partial X \times [0, \infty))$ is the properly supported pseudodifferential operator of mixed-regularity $(\mu, 0)$ from Definitions 5.18 which is invertible for large $R > 0$ and in local coordinates has an asymptotic expansion modulo $S^{\mu, -\infty}$ as in Lemma 5.21 and preserves support in $\partial X \times (-\infty, 0] \subseteq \partial X \times \mathbb{R}$. Moreover, for $\chi, \chi' \in \mathcal{C} + C_c^\infty(\partial X \times \mathbb{R})$ with $\chi\chi' = 0$, it holds that

$$\|\chi W_- \chi'\|_{H^\mu(\partial X \times \mathbb{R}) \rightarrow L^2(\partial X \times \mathbb{R})} = O(R^{-\infty}), \quad \text{as } R \rightarrow \infty.$$

Proof. The properties of W_+ and W_- listed in items (2) and (3) follow from the results of Subsection 5.1. We note that it follows from the previous subsection that

$$\chi_1 Q_M^{-1} \chi'_1 + \chi_2 (\varphi^{-1})^* W_+ W_- \varphi^* \chi'_2 : \overline{H}^\mu(X) \rightarrow \dot{H}^{-\mu}(X),$$

is a well defined continuous operator. Pick a χ_3 supported close to ∂X with $\chi_3 = 1$ on $\text{supp}(\chi'_2)$. We compute that

$$\begin{aligned} & (\chi_1 Q_M^{-1} \chi'_1 + \chi_2 (\varphi^{-1})^* W_+ W_- \varphi^* \chi'_2) Q_X = \\ &= \chi_1 + \chi_1 Q_M^{-1} (\chi'_1 - 1) Q_X + \chi_2 (\varphi^{-1})^* W_+ W_- \varphi^* Q_X \chi_3 + \\ & \quad + \chi_2 (\varphi^{-1})^* W_+ W_- \varphi^* (\chi'_2 - 1) Q_X \chi_3 + \chi_2 (\varphi^{-1})^* W_+ W_- \varphi^* \chi'_2 Q_X (1 - \chi_3) = \\ &= \chi_1 + \chi_1 Q_M^{-1} (\chi'_1 - 1) Q_X + \chi_2 (\varphi^{-1})^* W_+ W_- Q^\partial \varphi^* \chi_3 + \\ & \quad + \chi_2 (\varphi^{-1})^* W_+ W_- \varphi^* (\chi'_2 - 1) Q_X \chi_3 + \chi_2 (\varphi^{-1})^* W_+ W_- \varphi^* \chi'_2 Q_X (1 - \chi_3) = \\ &= \chi_1 + \chi_1 Q_M^{-1} (\chi'_1 - 1) Q_X + \chi_2 + (\varphi^{-1})^* S_0 \varphi^* \chi_3 + \\ & \quad + \chi_2 (\varphi^{-1})^* W_+ W_- \varphi^* (\chi'_2 - 1) Q_X \chi_3 + \chi_2 (\varphi^{-1})^* W_+ W_- \varphi^* \chi'_2 Q_X (1 - \chi_3) = \\ &= 1 + \underbrace{\chi_1 Q_M^{-1} (\chi'_1 - 1) Q_X}_{=: S_2} + (\varphi^{-1})^* S_0 \varphi^* \chi_3 + \\ & \quad + \underbrace{\chi_2 (\varphi^{-1})^* W_+ W_- \varphi^* (\chi'_2 - 1) Q_X \chi_3}_{=: S_3} + \underbrace{\chi_2 (\varphi^{-1})^* W_+ W_- \varphi^* \chi'_2 Q_X (1 - \chi_3)}_{=: S_4}. \end{aligned}$$

Since $\chi_1 (\chi'_1 - 1) = 0$, S_2 is a smoothing operator with parameter. Similarly, since $\chi'_2 (1 - \chi_3) = 0$, S_4 is a smoothing operator with parameter. Using Proposition 5.4 and Lemma 5.21, respectively, we conclude that $S_3 : \dot{H}^{-\mu}(X) \rightarrow \dot{H}^{-\mu}(X)$ and $(\varphi^{-1})^* S_0 \varphi^* \chi_3 : \dot{H}^{-\mu}(X) \rightarrow \dot{H}^{-\mu}(X)$ are continuous with norms bounded by $O(R^{-\infty})$ as $R \rightarrow \infty$. In particular,

$$S_5 := (\chi_1 Q_M^{-1} \chi'_1 + \chi_2 (\varphi^{-1})^* W_+ W_- \varphi^* \chi'_2) Q_X - 1,$$

satisfies that $S_5 : \dot{H}^{-\mu}(X) \rightarrow \dot{H}^{-\mu}(X)$ is continuous and $\|S_5\|_{\dot{H}^{-\mu}(X) \rightarrow \dot{H}^{-\mu}(X)} = O(R^{-\infty})$ as $R \rightarrow \infty$. We conclude that $(1 + S_5)^{-1}$ exists for large R and $\|1 - (1 + S_5)^{-1}\|_{\dot{H}^{-\mu}(X) \rightarrow \dot{H}^{-\mu}(X)} = O(R^{-\infty})$ as $R \rightarrow \infty$. We therefore have that

$$Q_X^{-1} = (1 + S_5)^{-1} (\chi_1 Q_M^{-1} \chi'_1 + \chi_2 (\varphi^{-1})^* W_+ W_- \varphi^* \chi'_2) = \chi_1 Q_M^{-1} \chi'_1 + \chi_2 (\varphi^{-1})^* W_+ W_- \varphi^* \chi'_2 + S,$$

where

$$S := (1 - (1 + S_5)^{-1}) (\chi_1 Q_M^{-1} \chi'_1 + \chi_2 (\varphi^{-1})^* W_+ W_- \varphi^* \chi'_2).$$

Since $\|1 - (1 + S_5)^{-1}\|_{\dot{H}^{-\mu}(X) \rightarrow \dot{H}^{-\mu}(X)} = O(R^{-\infty})$ as $R \rightarrow \infty$, the same holds for S and the proof is complete. \square

By combining Theorem 4.5 with Theorem 5.22, we arrive at the following corollary.

Corollary 5.23. *Let X be an n -dimensional compact manifold with boundary and d a distance function with property (MR) on Γ . For some $R_0 \geq 0$ and any $R \in \Gamma_{\pi/(n+1)}(R_0) \cap \Gamma$, we can write*

$$\mathcal{Z}_X^{-1} = \chi_1 A \chi'_1 + \chi_2 (\varphi^{-1})^* W_+ W_- \varphi^* \chi'_2 + \tilde{S},$$

where A , W_+ , W_- and $\chi_1, \chi'_1, \chi_2, \chi'_2 \in C^\infty(X)$ are as in Theorem 5.22 and $\tilde{S} : \overline{H}^\mu(X) \rightarrow \dot{H}^{-\mu}(X)$ is a continuous operator with

$$\|\tilde{S}\|_{\overline{H}^\mu(X) \rightarrow \dot{H}^{-\mu}(X)} = O(\text{Re}(R)^{-\infty}), \quad \text{as } \text{Re}(R) \rightarrow \infty \text{ in } \Gamma.$$

6. CONDITIONAL EXPECTATIONS OF Q_X^{-1} AND \mathcal{Z}_X^{-1}

A large motivation for this paper is the relation of the operator \mathcal{Z}_X with magnitude. For that purpose, we shall be interested in computing conditional expectations of Q_X^{-1} and \mathcal{Z}_X^{-1} against the constant function 1. In an accompanying paper [12], we prove that this conditional expectation of $(R\mathcal{Z}_X(R))^{-1}$ coincides with the magnitude function. The section is divided into three subsections: firstly, we study the case of no boundary, secondly we proceed to compute the asymptotic expansion of the conditional expectation of Q_X^{-1} and finally we produce explicit formulas for the asymptotic expansion and consider examples. As in the previous sections, we perform computations for Q that later translates into results for \mathcal{Z} under assumptions of property (MR).

6.1. Asymptotic expansions for compact manifolds. Let us consider the case that $X = M$ is a compact manifold. Starting from Lemma 2.24 we compute the asymptotics of $\langle Q(R)^{-1}1, 1 \rangle$ as $R \rightarrow \infty$ for a pseudodifferential operator with parameter R . Let $\text{vol}_d(M)$ denote the volume of M in the Riemannian metric defined from the transversal Hessian of d^2 at the diagonal.

Theorem 6.1. *Let M be an n -dimensional compact manifold with a distance function d whose square is regular at the diagonal. Let $(a_{j,0})_{j \in \mathbb{N}} \subseteq C^\infty(M; \mathbb{C}_+)$ denote the sequence of homogeneous functions obtained from restriction to $\xi = 0$ of the full symbol of Q_M^{-1} , as in Lemma 2.24. It holds that*

$$\langle 1, Q_M(R)^{-1}1 \rangle \sim \sum_{k=0}^{\infty} c_k(M, d) R^{n+1-k} + O(\text{Re}(R)^{-\infty}), \quad \text{as } \text{Re}(R) \rightarrow +\infty,$$

where

$$c_k(M, d) = \int_M a_{k,0}(x, 1) dx.$$

Here dx is the Riemannian volume density defined from g_{d^2} . The functions $a_{k,0}(x, 1)$ depend on the Taylor expansion (10) as described in Theorem 2.27 and can be computed inductively using Lemma 2.25. In particular,

$$c_k(M, d) = \begin{cases} 0, & \text{when } k \text{ is odd,} \\ \frac{\text{vol}_d(M)}{n! \omega_n}, & \text{when } k = 0, \\ \frac{n+1}{6 \cdot n! \omega_n} \int_X s_{d^2} dx, & \text{when } k = 2, \end{cases}$$

where s_{d^2} in local coordinates is computed as the polynomial in the Taylor coefficients of d^2 at the diagonal given as

$$s_{d^2}(x) := 3C^4(x, g \otimes g) - 3 \frac{\mathfrak{c}_{2,n}(n+5)(n^2-9)}{\mathfrak{c}_{1,n}} (C^3 \otimes C^3)(x, g \otimes g \otimes g), \quad \text{if } n \neq 1, 3$$

$$s_{d^2}(x) := 3 \left(10C_G^4(x, g_G \otimes g_G) - \frac{\mathfrak{c}_{2,3}}{\mathfrak{c}_{1,3}} (C_G^3 \otimes C_G^3)(x, g_G \otimes g_G \otimes g_G) \right), \quad \text{if } n = 3$$

Proof. The asymptotic expansion

$$\langle Q_M^{-1}1, 1 \rangle = \sum_{k=0}^{\infty} c_k(M, d) R^{n+1-k} + O(R^{-\infty}),$$

where $c_k(M, d) = \int_M a_{k,0}(x, 1) dx$ follows directly from Lemma 2.24 and the fact that Q_M^{-1} is of order $n+1$. It follows from Lemma 2.25 that $a_k(x, 0, 1) = 0$ for odd k . It follows from Theorem 2.27 that c_0 and c_2 take the prescribed form. \square

The justification for the notation s_{d^2} in Theorem 6.1 comes from Example 2.31 which shows that for the geodesic distance on a Riemannian manifold, s_{d^2} is the scalar curvature. We further conclude the following corollary.

Corollary 6.2. *Let M be a compact Riemannian manifold equipped with its geodesic distance. Then*

$$\langle 1, Q_M(R)^{-1}1 \rangle = \frac{\text{vol}_d(M)}{n! \omega_n} R^{n+1} + \frac{n+1}{6 \cdot n! \omega_n} \int_M s dx R^{n-1} + O(\text{Re}(R)^{n-3}), \quad \text{as } \text{Re}(R) \rightarrow +\infty,$$

where s denotes the scalar curvature of M .

Combining Theorem 3.7 with Theorem 6.1 we arrive at the following corollary.

Corollary 6.3. *Let M be an n -dimensional compact manifold with a distance function d with property (MR) on Γ . It holds that*

$$\langle 1, \mathcal{Z}_M(R)^{-1}1 \rangle \sim \sum_{k=0}^{\infty} c_k(M, d) R^{n+1-k} + O(\text{Re}(R)^{-\infty}), \quad \text{as } \text{Re}(R) \rightarrow +\infty \text{ in } \Gamma,$$

where $c_k(M, d)$ is as in Theorem 6.1.

6.2. A lengthy exercise in integration by parts. To study the asymptotic expansions of $\langle 1, Q_X^{-1}1 \rangle$ in the presence of a boundary, we need a series of smaller lemmas. The reader should recall the notation from Theorem 5.22.

Lemma 6.4. *Let X be an n -dimensional compact manifold with boundary and d a distance function whose square is regular at the diagonal. It holds that*

$$\langle Q_X^{-1}1, 1 \rangle_{L^2(X)} = \langle A1, \chi_1 \rangle_{L^2(X)} + \langle W_- 1, (W_+)^*(\chi_2 \circ \varphi) \rangle_{L^2(\partial X \times [0, \infty))} + O(R^{-\infty}).$$

Proof. We first note that W_- preserves supports in $\partial X \times (-\infty, 0] \subseteq \partial X \times \mathbb{R}$ by Lemma 5.21 and $(W_+)^*$ preserves supports in $\partial X \times (-\infty, 0] \subseteq \partial X \times \mathbb{R}$ since W_+ preserves supports in $\partial X \times [0, \infty)$ by Lemma 5.21. Therefore, viewing 1 as an element of $\overline{H}_{\text{loc}}^{\mu}(\partial X \times [0, \infty))$ and $\chi_2 \circ \varphi$ as an element of $\overline{H}_c^{\mu}(\partial X \times [0, \infty))$, the images $W_- 1 \in L_{\text{loc}}^2(\partial X \times [0, \infty))$ and $(W_+)^*(\chi_2 \circ \varphi) \in L_c^2(\partial X \times [0, \infty))$ are well defined and $\langle W_- 1, (W_+)^*(\chi_2 \circ \varphi) \rangle_{L^2(\partial X \times [0, \infty))}$ is well defined. By the same token, $\langle A1, \chi_1 \rangle_{L^2(X)}$ is defined as the inner product of $\chi_1 \in L^2(X)$ with the restriction of $A1_M \in L^2(M)$ to X .

Since $\chi_j'|_{\text{supp}(\chi_j)} = 1$, for $j = 1, 2$, it follows that

$$\begin{aligned} \langle A1, \chi_1 \rangle_{L^2(X)} &= \langle A\chi_1', \chi_1 \rangle_{L^2(X)} + O(R^{-\infty}) \quad \text{and} \\ \langle W_- 1, (W_+)^*(\chi_2 \circ \varphi) \rangle_{L^2(\partial X \times [0, \infty))} &= \langle W_- (\chi_2' \circ \varphi), (W_+)^*(\chi_2 \circ \varphi) \rangle_{L^2(\partial X \times [0, \infty))} + O(R^{-\infty}). \end{aligned}$$

The last equality follows from Proposition 5.4. Therefore, Theorem 5.22 reduces the statement of the theorem to the property that $\langle S1, 1 \rangle_{L^2(X)} = O(R^{-\infty})$. This is clear from the property of S that $\|S\|_{\overline{H}^{\mu} \rightarrow \dot{H}^{-\mu}} = O(R^{-\infty})$. \square

Lemma 6.5. *Let X be an n -dimensional compact manifold with boundary and d a distance function whose square is regular at the diagonal. Let $(a_{j,0})_{j \in \mathbb{N}} \subseteq C^{\infty}(M; \mathbb{C}_+)$ denote the sequence of homogeneous functions obtained from restriction to $\xi = 0$ of the full symbol of A , as in Lemma 2.24. It holds that*

$$\langle A1, \chi_1 \rangle = \sum_{k=0}^{\infty} c_{k, \chi_1}(X, d) R^{n+1-k} + O(R^{-\infty}),$$

where

$$c_{k, \chi_1}(M, d) = \int_X \chi_1(x) a_{k,0}(x, 1) dx.$$

Here dx is the Riemannian volume density defined from g_{d^2} .

Proof. The lemma follows immediately from Lemma 2.24 since χ_1 has compact support in X° . \square

Lemma 6.6. *Let $a = a(x', \xi) \in C^\infty(\mathbb{R}^{n-1} \times \mathbb{R}^n)$ be a polynomially bounded smooth function with compact support in x' , and $\chi \in \mathcal{S}(\mathbb{R}^n)$ a real even Schwartz function. Then as $R \rightarrow +\infty$,*

$$\begin{aligned} \frac{1}{(2\pi)^n} \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}^n} a(x', \xi) R^n \hat{\chi}(-R\xi) e^{-Rix'\xi'} d\xi dx &= \\ &= \sum_{\alpha \in \mathbb{N}^n} \frac{D_x^\alpha \chi(0)}{\alpha!} \int_{\mathbb{R}^{n-1}} D_{\xi=0}^\alpha \left(a(x', \xi) e^{-ix'\xi'} \right) dx R^{-|\alpha|} + O(R^{-\infty}). \end{aligned}$$

In particular, if χ is locally constant near 0, then

$$\frac{1}{(2\pi)^n} \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}^n} a(x', \xi) R^n \hat{\chi}(-R\xi) e^{-Rix'\xi'} d\xi dx = \chi(0) \int_{\mathbb{R}^{n-1}} a(x', 0) dx + O(R^{-\infty}).$$

Proof. Consider the distribution $u_R(\xi) := R^n \hat{\chi}(-R\xi)$. For any test function $\varphi \in \mathcal{S}(\mathbb{R}^n)$, we compute that

$$\begin{aligned} (u_R, \varphi) &= \int_{\mathbb{R}^n} \hat{\chi}(-\xi) \varphi(\xi/R) d\xi = \sum_{\alpha \in \mathbb{N}^n} \frac{D_x^\alpha \varphi(0)}{\alpha!} \int_{\mathbb{R}^n} \hat{\chi}(-\xi) \xi^\alpha d\xi R^{-|\alpha|} + O(R^{-\infty}) = \\ &= (2\pi)^n \sum_{\alpha \in \mathbb{N}^n} \frac{D_x^\alpha \chi(0)}{\alpha!} R^{-|\alpha|} (\delta^\alpha, \varphi) + O(R^{-\infty}) \end{aligned}$$

We conclude that in $\mathcal{S}'(\mathbb{R}^n)$, we have an asymptotic expansion

$$u_R = (2\pi)^n \sum_{\alpha \in \mathbb{N}^n} \frac{D_x^\alpha \chi(0)}{\alpha!} R^{-|\alpha|} \delta^\alpha + O(R^{-\infty}).$$

Using standard methods for oscillatory integrals, we see that the same expansion holds also for in the weak topology against polynomially bounded smooth functions.

We compute that

$$\begin{aligned} \frac{1}{(2\pi)^n} \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}^n} a(x', \xi) R^n \hat{\chi}(-R\xi) e^{-Rix'\xi'} d\xi dx &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}^n} a(x', \xi) u_R(\xi) e^{-Rix'\xi'} d\xi dx \\ &= \sum_{\alpha \in \mathbb{N}^n} \frac{D_x^\alpha \chi(0)}{\alpha!} R^{-|\alpha|} \int_{\mathbb{R}^{n-1}} (\delta_\xi^\alpha, a(x', \xi) e^{-ix'\xi'}) dx + O(R^{-\infty}) = \\ &= \sum_{\alpha \in \mathbb{N}^n} \frac{D_x^\alpha \chi(0)}{\alpha!} \int_{\mathbb{R}^{n-1}} D_{\xi=0}^\alpha \left(a(x', \xi) e^{-ix'\xi'} \right) dx R^{-|\alpha|} + O(R^{-\infty}). \end{aligned}$$

□

Lemma 6.7. *Let X be an n -dimensional compact manifold with boundary and d a distance function whose square is regular at the diagonal. We denote the symbols of W_\pm by w_\pm (as in Definition 5.18 and Lemma 5.18). Then it holds that*

$$\langle W_- 1, W_+^* \chi_2 \rangle = \sum_{k=0}^{\infty} c_{k, \chi_2}(X, d) R^{n+1-k} + O(R^{-\infty}),$$

where

$$\begin{aligned} c_{k, \chi_2}(X, d) &= \int_X \chi_2(x) a_{k,0}(x, 1) dx + \\ &+ \sum_{\substack{k=|\beta|+\gamma_n+j+l \\ \gamma_n > 0}} \frac{i^{|\beta|+|\gamma_n|} (-1)^{|\beta|+1}}{\beta! (\beta_n + \gamma_n)!} \int_{\partial X} \partial_x^\beta w_{-,j}(x', 0, 0, 1) \partial_{x_n}^{\gamma_n-1} \partial_\xi^{\beta+(0, \gamma_n)} w_{+,l}(x', 0, 0, 1) dx'. \end{aligned}$$

Here dx is the Riemannian volume density on X defined from g_{d^2} and dx' the induced Riemannian volume density on ∂X .

Proof. The computation can be reduced to one in local coordinates, so we can assume that w_+ and w_- are symbols of mixed-regularity $(-\mu, 0)$ in \mathbb{R}^n , and up to $O(R^{-\infty})$ we can treat W_+ and

W_- as compactly based. As such, we replace M by \mathbb{R}^n and X by \mathbb{R}_+^n in all computations. Let w_+^* denote the symbol of W_+^* . By the same arguments as in [38, Chapter I.3], we have that

$$(38) \quad w_+^*(x, \xi, R) \sim \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} D_x^{\alpha} \overline{w_+(x, \xi, R)},$$

in the sense of Definition 5.8. We note that Equation (38) only identifies w_+^* up to $S^{\mu, -\infty}$ but this suffices as symbols from $S^{\mu, -\infty}$ will only contribute to the conditional expectation with $O(R^{-\infty})$.

Using that W_- and W_+^* preserves supports in \mathbb{R}_-^n , we can consider χ_2 as an element of $C_c^{\infty}(\mathbb{R}^n)$, and write

$$\langle W_- 1, W_+^* \chi_2 \rangle_{L^2(\mathbb{R}_+^n)} = \int_{\mathbb{R}_+^n} [W_- 1](x) \overline{[W_+^* \chi_2](x)} dx,$$

where

$$W_+^* \chi_2(x) := \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} w_+^*(x, \xi, R) \hat{\chi}_2(\xi) d\xi,$$

is computed from the action of $(Q_+^{-1})^*$ on $\chi \in C_c^{\infty}(\mathbb{R}^n)$. We compute that

$$\begin{aligned} \langle W_- 1, W_+^* \chi_2 \rangle_{L^2(\mathbb{R}_+^n)} &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}_+^n} \int_{\mathbb{R}^n} w_-(x, 0, R) \overline{w_+^*(x, \xi, R) \hat{\chi}_2(\xi) e^{ix\xi}} d\xi dx = \\ &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}_+^n} \int_{\mathbb{R}^n} w_-(x, 0, R) \overline{w_+^*(x, \xi, R) \hat{\chi}_2(-\xi) e^{-ix\xi}} d\xi dx = \\ &= \sum_{\alpha} \frac{(-1)^{|\alpha|}}{(2\pi)^n \alpha!} \int_{\mathbb{R}_+^n} \int_{\mathbb{R}^n} w_-(x, 0, R) \partial_{\xi}^{\alpha} D_x^{\alpha} w_+(x, \xi, R) \hat{\chi}_2(-\xi) e^{-ix\xi} d\xi dx + O(R^{-\infty}) = \\ &= \sum_{\alpha} \frac{(-1)^{\alpha_n}}{(2\pi)^n \alpha!} \int_{\mathbb{R}_+^n} \int_{\mathbb{R}^n} D_{x'}^{\alpha'} (w_-(x, 0, R) e^{-ix\xi}) D_{x_n}^{\alpha_n} \partial_{\xi}^{\alpha} w_+(x, \xi, R) \hat{\chi}_2(-\xi) d\xi dx + O(R^{-\infty}) = \\ &= \sum_{\alpha} \sum_{\gamma' + \beta' = \alpha'} \frac{(-1)^{\alpha_n}}{(2\pi)^n \beta'! \gamma'! \alpha_n!} \int_{\mathbb{R}_+^n} \int_{\mathbb{R}^n} D_{x'}^{\beta'} w_-(x, 0, R) D_{x_n}^{\alpha_n} \partial_{\xi}^{\alpha} w_+(x, \xi, R) (-i\xi')^{\gamma'} \hat{\chi}_2(-\xi) e^{-ix\xi} d\xi dx + O(R^{-\infty}) = \\ &= \sum_{\gamma, \beta} \frac{1}{(2\pi)^n \beta! \gamma!} \int_{\mathbb{R}_+^n} \int_{\mathbb{R}^n} D_x^{\beta} w_-(x, 0, R) \partial_{\xi}^{\gamma + \beta} w_+(x, \xi, R) (-i\xi)^{\gamma} \hat{\chi}_2(-\xi) e^{-ix\xi} d\xi dx + \\ &+ \sum_{\substack{\gamma, \beta, \\ \gamma_n > 0}} \sum_{k=0}^{\beta_n} b_{\gamma, \beta, k} \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}^n} D_x^{\beta - (0, k)} w_-(x', 0, 0, R) D_{x_n}^{\gamma_n - 1} \partial_{\xi}^{\beta + \gamma} w_+(x', 0, \xi, R) (i\xi)^{(\gamma', k)} \hat{\chi}_2(-\xi) e^{-ix'\xi'} d\xi dx + \\ &+ O(R^{-\infty}). \end{aligned}$$

where

$$b_{\gamma, \beta, k} = \frac{i(-1)^{|\gamma_n|+1} \beta_n!}{(2\pi)^n \beta'! \gamma'! (\beta_n + \gamma_n)! k! (\beta_n - k)!}$$

By the composition formula for pseudodifferential operators (see [38, Chapter I.3]), we have that

$$\begin{aligned} \sum_{\gamma, \beta} \frac{1}{(2\pi)^n \beta! \gamma!} \int_{\mathbb{R}_+^n} \int_{\mathbb{R}^n} D_x^{\beta} w_-(x, 0, R) \partial_{\xi}^{\gamma + \beta} w_+(x, \xi, R) (-i\xi)^{\gamma} \hat{\chi}_2(-\xi) e^{-ix\xi} d\xi dx = \\ = \int_{\mathbb{R}_+^n} \chi_2(x) a(x, 0, R) dx + O(R^{-\infty}) = \langle Q_M^{-1} 1, \chi_2 \rangle_{L^2(\mathbb{R}_+^n)} + O(R^{-\infty}). \end{aligned}$$

We can therefore continue our calculation

$$\begin{aligned} \langle W_- 1, W_+^* \chi_2 \rangle_{L^2(\mathbb{R}_+^n)} &= \langle Q_M^{-1} 1, \chi_2 \rangle_{L^2(\mathbb{R}_+^n)} \\ &+ \sum_{\substack{\gamma, \beta, \\ \gamma_n > 0}} \sum_{k=0}^{\beta_n} b_{\gamma, \beta, k} \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}^n} D_x^{\beta - (0, k)} w_-(x', 0, 0, R) D_{x_n}^{\gamma_n - 1} \partial_{\xi}^{\beta + \gamma} w_+(x', 0, \xi, R) (i\xi)^{(\gamma', k)} \hat{\chi}_2(-\xi) e^{-ix'\xi'} d\xi dx + \\ &+ O(R^{-\infty}) \end{aligned}$$

To obtain an asymptotic expansion, we expand w_{\pm} in its defining homogeneous expansion $w_{\pm} \sim \sum w_{\pm,l}$ from Definition 5.18 (see also Lemma 5.21). We see that

$$\begin{aligned} & \sum_{\substack{\gamma,\beta, \\ \gamma_n > 0}} \sum_{k=0}^{\beta_n} b_{\gamma,\beta,k} \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}^n} D_x^{\beta-(0,k)} w_{-}(x', 0, 0, R) D_{x_n}^{\gamma_n-1} \partial_{\xi}^{\beta+\gamma} w_{+}(x', 0, \xi, R) (i\xi)^{(\gamma',k)} \hat{\chi}_2(-\xi) e^{-ix\xi} d\xi dx = \\ & = \sum_{i=0}^{\infty} \sum_{\substack{i=|\beta|+|\gamma|+j+l, \\ \gamma_n > 0}} \sum_{k=0}^{\beta_n} b_{\gamma,\beta,k} \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}^n} D_x^{\beta-(0,k)} w_{-,j}(x', 0, 0, R) D_{x_n}^{\gamma_n-1} \partial_{\xi}^{\beta+\gamma} w_{+,l}(x', 0, \xi, R) (i\xi)^{(\gamma',k)} \hat{\chi}_2(-\xi) e^{-ix'\xi'} d\xi dx \end{aligned}$$

Let us consider each of the terms

$$\begin{aligned} & \sum_{\substack{i=|\beta|+|\gamma|+j+l, \\ \gamma_n > 0}} \sum_{k=0}^{\beta_n} b_{\gamma,\beta,k} \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}^n} D_x^{\beta-(0,k)} w_{-,j}(x', 0, 0, R) D_{x_n}^{\gamma_n-1} \partial_{\xi}^{\beta+\gamma} w_{+,l}(x', 0, \xi, R) (i\xi)^{(\gamma',k)} \hat{\chi}_2(-\xi) e^{-ix'\xi'} d\xi dx = \\ & = R^{2\mu-i} \sum_{\substack{i=|\beta|+|\gamma|+j+l, \\ \gamma_n > 0}} \sum_{k=0}^{\beta_n} b_{\gamma,\beta,k} \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}^n} D_x^{\beta-(0,k)} w_{-,j}(x', 0, 0, 1) D_{x_n}^{\gamma_n-1} \partial_{\xi}^{\beta+\gamma} w_{+,l}(x', 0, \frac{\xi}{R}, 1) (i\xi)^{(\gamma',k)} \hat{\chi}_2(-\xi) e^{-ix'\xi'} d\xi dx. \end{aligned}$$

We can compute each of the terms using Lemma 6.6 which implies that

$$\begin{aligned} & \frac{1}{(2\pi)^n} \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}^n} D_x^{\beta-(0,k)} w_{-,j}(x', 0, 0, 1) D_{x_n}^{\gamma_n-1} \partial_{\xi}^{\beta+\gamma} w_{+,l}(x', 0, \xi, 1) (i\xi)^{(\gamma',k)} R^n \hat{\chi}_2(-R\xi) e^{-Rix'\xi'} d\xi dx = \\ & = \begin{cases} O(R^{-\infty}), & \text{if } (\gamma', k) \neq 0, \\ \int_{\mathbb{R}^{n-1}} D_x^{\beta} w_{-,j}(x', 0, 0, 1) D_{x_n}^{\gamma_n-1} \partial_{\xi}^{\beta+(0,\gamma_n)} w_{+,l}(x', 0, 0, 1) dx + O(R^{-\infty}), & \text{if } (\gamma', k) = 0 \end{cases} \end{aligned}$$

We conclude that

$$\begin{aligned} & \langle Q^{-1}1, (Q_+^{-1})^* \chi_2 \rangle_{L^2(\mathbb{R}_+^n)} - \langle Q_M^{-1}1, \chi_2 \rangle_{L^2(\mathbb{R}_+^n)} = \\ & = R^{2\mu} \sum_{i=0}^{\infty} \sum_{\substack{i=|\beta|+\gamma_n+j+l, \\ \gamma_n > 0}} \frac{i(-1)^{|\gamma_n|+1} R^{-i}}{\beta'!(\beta_n + \gamma_n)!} \int_{\mathbb{R}^{n-1}} D_x^{\beta} w_{-,j}(x', 0, 0, 1) D_{x_n}^{\gamma_n-1} \partial_{\xi}^{\beta+(0,\gamma_n)} w_{+,l}(x', 0, 0, 1) dx + O(R^{-\infty}). \end{aligned}$$

After using $D = -i\partial$, the boundary contributions have been computed.

The lemma now follows from Lemma 2.24 giving the asymptotic expansion

$$\langle Q_M^{-1}1, \chi_2 \rangle_{L^2(\mathbb{R}_+^n)} = \sum_k R^{n+1-k} \int_{\mathbb{R}_+^n} \chi_2(x) a_k(x, 0, 1) dx.$$

□

6.3. Asymptotic expansions for compact manifolds with boundary. We now study asymptotic expansions of $\langle 1, Q_X^{-1}1 \rangle$ for a compact manifold with boundary, and give a procedure to compute the coefficients. An important difference to the case of empty boundary is the boundary contributions: we identify the from Lemma 6.7 as follows.

Definition 6.8. If X is an n -dimensional compact manifold with boundary and d a distance function whose square is regular at the diagonal, then we define the sequence of functions $(B_{d^2,k})_{k>0} \subseteq C^\infty(\partial X)$ by

$$B_{d^2,k}(x') := \sum_{\substack{k=|\beta|+\gamma_n+j+l, \\ \gamma_n > 0}} \frac{i^{|\beta|+|\gamma_n|} (-1)^{|\beta|+1}}{\beta'!(\beta_n + \gamma_n)!} \partial_x^{\beta} w_{-,j}(x', 0, 0, 1) \partial_{x_n}^{\gamma_n-1} \partial_{\xi}^{\beta+(0,\gamma_n)} w_{+,l}(x', 0, 0, 1).$$

For notational simplicity, we set $B_0 := 0$.

Proposition 6.9. *Let X be an n -dimensional manifold with boundary, d a distance function whose square is regular at the diagonal, and $(B_{d^2,k})_{k>0} \subseteq C^\infty(\partial X)$ as in Definition 6.8. Then for each $k > 0$, $B_{d^2,k}$ is a polynomial in $(C_G^{(\gamma)})_{\gamma \in \cup_{k \leq j} I_k}$ and its derivatives contracted by the metric g_G , its contraction along the normal to the boundary, and its derivatives of total degree j where each $C_G^{(\gamma)}$, $\gamma \in I_k$, has degree k , the metric has degree zero and x -derivatives increase the order by 1.*

Computing from the results of Appendix F, we can describe the special cases $k = 1$ and $k = 2$.

Proposition 6.10. *Let (X, d) be as in Proposition 6.9. Then*

$$B_{d^2,1}(x') = \frac{(n+1)}{2 \cdot n! \omega_n \sqrt{h_0(x')}}.$$

In particular, if x_n is the transversal coordinate defined from the unit normal to ∂X (in g_{d^2}), then

$$B_{d^2,1}(x') = \frac{(n+1)}{2 \cdot n! \omega_n}.$$

Proof. By definition, we have that

$$\begin{aligned} B_{d^2,1}(x') &= -iw_{-,0}(x', 0, 0, 1) \partial_{\xi_n} w_{+,0}(x', 0, 0, 1) = \\ &= -\frac{i}{n! \omega_n} h_0(x')^\mu (-h_-(x', 0, 1))^\mu \partial_{\xi_n} [(\xi_n - h_+(x, 0, 1))^\mu] |_{\xi_n=0} = \\ &= -\frac{i(n+1)}{2 \cdot n! \omega_n} h_0(x')^\mu (-h_-(x', 0, 1))^\mu (-h_+(x, 0, 1))^{\mu-1} = \\ &= -\frac{i(n+1)}{2 \cdot n! \omega_n} h_0(x') (-h_-(x', 0, 1)) = \frac{(n+1)}{2 \cdot n! \omega_n \sqrt{h_0(x')}}. \end{aligned}$$

Here we have used the identities of Lemma F.1. If x_n is the transversal coordinate defined from the unit normal to ∂X , then $h_0 = 1$. \square

Proposition 6.11. *Let (X, d) be as in Proposition 6.9 and assume that x_n is the transversal coordinate defined from the unit normal ∂_n to ∂X (in g_{d^2}). Then there are universal polynomials α_1 and α_2 with rational coefficients such that*

$$B_{d^2,2}(x') = \frac{\alpha_1(n)}{n! \omega_n} C^3(x', \partial_n \otimes \partial_n \otimes \partial_n) + \frac{\alpha_2(n)}{n! \omega_n} C^3(x', g \otimes \partial_n).$$

Proof. By definition, we have that

$$\begin{aligned} B_{d^2,2}(x) &= \frac{1}{2} w_{-,0}(x', 0, 0, 1) \partial_{x_n} \partial_{\xi_n}^2 w_{+,0}(x', 0, 0, 1) - \frac{1}{2} \partial_{x_n} w_{-,0}(x', 0, 0, 1) \partial_{\xi_n}^2 w_{+,0}(x', 0, 0, 1) - \\ &\quad - iw_{-,1}(x', 0, 0, 1) \partial_{\xi_n} w_{+,0}(x', 0, 0, 1) - iw_{-,0}(x', 0, 0, 1) \partial_{\xi_n} w_{+,1}(x', 0, 0, 1) + \\ &\quad + \nabla_{x'} w_{-,0}(x', 0, 0, 1) \cdot \nabla_{\xi'} \partial_{\xi_n} w_{+,0}(x', 0, 0, 1). \end{aligned}$$

These expressions were computed in Section F of the appendix. If x_n is the transversal coordinate defined from the unit normal to ∂X , then $h_0 = 1$ and $b = 0$. The lemmas of Section F, for $h_0 = 1$ and $b = 0$, shows that $B_{d^2,2}(x')$ is in the linear span of $C^3(x', \partial_n \otimes \partial_n \otimes \partial_n)$ and $C^3(x', g \otimes \partial_n)$, and carefully inspecting the computations imply the existence of the universal polynomials α_1 and α_2 . \square

Proposition 6.12. *Let $X \subseteq \mathbb{R}^n$ be a domain with smooth boundary equipped with the Euclidean distance. Then for a universal polynomial β , it holds that*

$$B_{d^2,2} = \frac{\beta(n)}{n! \omega_n} H,$$

where H denotes the mean curvature of the boundary.

Proof. We compute that

$$\frac{c_{1,n}(n^2 - 1)}{2(n! \omega_n)^2} = -\frac{(n-2)! \omega_{n-2} \omega_2 (n^2 - 1)}{2(n! \omega_n)^2} = -\frac{(n-1)! 2\pi \omega_{n-2}}{(n-1)! n \omega_n} \frac{n+1}{4 \cdot n! \omega_n} = -\frac{n+1}{4 \cdot n! \omega_n}.$$

Fix a point $x_0 \in \partial X$ and choose coordinates as in Example 2.16; in other words we write ∂X as the graph of a function $\varphi = \varphi(x')$ with $\nabla \varphi(x_0) = 0$. The computations in Example 2.16 show that $h_0(x') = 1 + |\nabla \varphi(x')|^2$ is x_n -independent and that $b(x_0) = \nabla \varphi(x_0) = 0$. Moreover, $C^3(x, v)$ is a first order polynomial in v_n , so $C^3(x, \iota_n g \otimes \iota_n g \otimes \iota_n g) = 0$. A short computation using Equation (15) gives us that

$$C^3(x_0, g \otimes \iota_n g) = -2g_{x_0}(\nabla^2 \varphi(x_0)) = -(n-1)H(x_0),$$

where H denotes the mean curvature. We conclude that

$$B_{d^2,2}(x_0) = \frac{n^2 - 1}{4 \cdot n! \omega_n} H(x_0).$$

□

Combining Lemmas 6.4, 6.5 and 6.7 we arrive at the following theorem:

Theorem 6.13. *Let X be an n -dimensional compact manifold with boundary and d a distance function whose square is regular at the diagonal. Denote the Riemannian volume density on X defined from g_{d^2} by dx and the induced Riemannian volume density on ∂X by dx' . It holds that*

$$(39) \quad \langle 1, Q_X^{-1} 1 \rangle_{L^2(X)} = \sum_{k=0}^{\infty} c_k(X, d) R^{n+1-k} + O(R^{-\infty}), \quad \text{as } R \rightarrow +\infty,$$

where the coefficients $c_k(X, d)$ are given as

$$c_k(X, d) = \int_X a_{k,0}(x, 1) dx + \int_{\partial X} B_{d^2,k}(x) dx',$$

where

- (1) $a_{k,0}(\cdot, 1) \in C^\infty(X)$ is an invariant polynomial in the entries of the Taylor expansion (10) as described in Theorem 2.27 and can be computed inductively using Lemma 2.25, with $a_{k,0} = 0$ if k is odd; and
- (2) $B_{d^2,k} \in C^\infty(\partial X)$ is an invariant polynomial in the entries of the Taylor coefficients of d^2 at the diagonal in X near ∂X as described in Proposition 6.9 and can be inductively computed using Lemma 5.19.

In particular, we have that

$$(40) \quad c_0(X, d) = \frac{\text{vol}(X)}{n! \omega_n}$$

$$(41) \quad c_1(X, d) = \frac{(n+1) \text{vol}(\partial X)}{2n! \omega_n},$$

$$(42) \quad c_2(X, d) = \frac{n+1}{6 \cdot n! \omega_n} \int_X s_{d^2} dx + \frac{(n-1)(n+1)^2}{8 \cdot n! \omega_n} \int_{\partial X} H_{d^2} dx'.$$

where the scalar curvature s_{d^2} is defined as in Theorem 6.1 and the mean curvature H_{d^2} is an explicit function in the linear span of $C^3(x', \partial_n \otimes \partial_n \otimes \partial_n)$ and $C^3(x', g \otimes \partial_n)$ that coincides with the usual mean curvature for domains in \mathbb{R}^n .

Our notation H_{d^2} in Theorem 6.13 is justified by Proposition 6.12 showing that $H_{d^2} = H$ is the mean curvature if X is a domain in Euclidean space.

Proof. The expression in Equation (39) follows from Lemma 6.4 by adding together the computation of Lemma 6.5 with that in Lemma 6.7.

The computation (40) follows from the fact that $a_{0,0}(x, 1) = \frac{1}{n! \omega_n}$ (see Theorem 2.27). The computation (41) follows from Proposition 6.10. The computation (42) is a consequence of Theorem 2.27 (computing the interior contribution) and Proposition 6.11 (computing the boundary contribution). □

Combining Theorem 4.5 with Theorem 6.13 we arrive at the following corollary.

Corollary 6.14. *Let X be an n -dimensional compact manifold with boundary and d a distance function with property (MR) on $[R_0, \infty)$, for some $R_0 \geq 0$. It holds that*

$$\langle 1, \mathcal{Z}_X^{-1} 1 \rangle_{L^2(X)} = \sum_{k=0}^{\infty} c_k(X, d) R^{n+1-k} + O(R^{-\infty}), \quad \text{as } R \rightarrow +\infty,$$

where the coefficients $c_k(X, d)$ are as in Theorem 6.13.

Corollary 6.15. *If $X \subseteq \mathbb{R}^n$ is a compact domain with smooth boundary, then*

$$\begin{aligned} c_0(X, d) &= \frac{\text{vol}(X)}{n! \omega_n} \\ c_1(X, d) &= \frac{\mu \text{vol}(\partial X)}{n! \omega_n}, \\ c_2(X, d) &= \frac{\mu^2 (n-1)}{2 \cdot n! \omega_n} \int_{\partial X} H dS. \end{aligned}$$

The computation of Corollary 6.15 is compatible with the computations of [11] for $\mu \in \mathbb{N}$. We note that the precise proportionality constant in c_2 follows from the computation from [11] for $\mu \in \mathbb{N}$ since the pre-factor by Proposition 6.12 is determined as a universal polynomial in n .

APPENDIX A. OVERVIEW OF CONORMAL DISTRIBUTIONS

Pseudodifferential operators with parameters will play an important role in our study of the operators Q and \mathcal{Z} , both to prove meromorphic extensions and to compute asymptotic expansions. We will use an approach to parameter dependence described in terms of conormal distributions to which the operator Q is susceptible.

First we recall the basics of conormal distributions. We follow the approach of [23, Chapter 18.2]. For a tempered distribution $u \in \mathcal{S}'(\mathbb{R}^N)$ we write $\mathcal{F}u$ or sometimes \hat{u} for its Fourier transform. We define the following subsets of \mathbb{R}^N as $B_1 := B(0, 1)$ and $B_j := B(0, 2^j) \setminus \overline{B(0, 2^{j-1})}$ for $j > 1$. For a function $u \in \mathcal{S}'(\mathbb{R}^N)$ we write $u = \sum_{j=1}^{\infty} u_j$ in a distributional sense where u_j is determined by $\hat{u}_j = \hat{u} \chi_{B_j}$. Following the notation of [23, Appendix B], for $s \in \mathbb{R}$, we set

$${}^{\infty}H_s(\mathbb{R}^N) := \{u \in \mathcal{S}'(\mathbb{R}^N) : \sup_{j \in \mathbb{N}_+} 2^{sj} \|u_j\|_{L^2(\mathbb{R}^N)} < \infty\}.$$

Written in terms of the standard notation for Besov spaces, ${}^{\infty}H_s(\mathbb{R}^N) = B_{\infty, 2}^s(\mathbb{R}^N)$. By [23, Theorem 18.2.9], the notion of belonging to ${}^{\infty}H_s(\mathbb{R}^N)$ is coordinate invariant for compactly supported distributions.

If Z is a smooth N -dimensional manifold, and $u \in \mathcal{D}'(Z)$ satisfies that for each coordinate chart $\kappa : U \rightarrow U' \subseteq \mathbb{R}^N$ and $\chi \in C_c^\infty(U)$, $(\kappa^{-1})^*(\chi u) \in {}^{\infty}H_s(\mathbb{R}^N)$ we say that u is locally ${}^{\infty}H_s$. We define

$${}^{\infty}H_s^{\text{loc}}(Z) := \{u \in \mathcal{D}'(Z) : u \text{ is locally } {}^{\infty}H_s\}.$$

Due to coordinate invariance on compacts of the space ${}^{\infty}H_s(\mathbb{R}^N)$, it holds that

$${}^{\infty}H_s^{\text{loc}}(\mathbb{R}^N) = \{u \in \mathcal{D}'(\mathbb{R}^N) : \chi u \in {}^{\infty}H_s(\mathbb{R}^N) \forall \chi \in C_c^\infty(\mathbb{R}^N)\}.$$

Due to the locality of the definition, if $(\kappa_\alpha)_{\alpha \in I}$ is a locally finite atlas on a smooth manifold Z and $(\chi_\alpha)_{\alpha \in I}$ a subordinate smooth partition of unity, $u \in {}^{\infty}H_s^{\text{loc}}(Z)$ if and only if $(\kappa_\alpha^{-1})^*(\chi_\alpha u) \in {}^{\infty}H_s(\mathbb{R}^N)$ for all $\alpha \in I$.

Definition A.1 (Definition 18.2.6 of [23]). Let Z be a smooth N -dimensional manifold and $Y \subseteq Z$ a smooth submanifold which is closed in the topology of Z . We let $I^m(Z; Y)$ denote the space of conormal distributions of order m . We say that $u \in \mathcal{D}'(Z)$ is a distribution conormal to Y of order m if for each collection of first order differential operator L_1, \dots, L_p (with C^∞ -coefficients) that are tangential to Y , it holds that

$$L_1 \cdots L_p u \in {}^{\infty}H_{-m - (N-k)/2}^{\text{loc}}(Z).$$

We let $I^m(Z, Y)$ denote the space of distributions conormal to Y of order m .

We here define the order of conormal distributions in such a way that the order of the conormal distribution matches the order of the operator it will define. Our convention differs from that in [23] but is consistent with that in [39].

The reader should note that for $u \in I^m(Z, Y)$, it holds that $u|_{Z \setminus Y} \in C^\infty(Z \setminus Y)$. Moreover, if L is an arbitrary differential operator (with C^∞ -coefficients) of order p then $Lu \in I^{m+p}(Z, Y)$ whenever $u \in I^m(Z, Y)$. However, if $u \in I^m(Z, Y)$ then for any collection of first order differential

operator L_1, \dots, L_p (with C^∞ -coefficients) that are tangential to Y , then $L_1 \cdots L_p u \in I^m(Z, Y)$. By elliptic regularity, we have that

$$I^{-\infty}(Z, Y) := \bigcap_{m \in \mathbb{R}} I^m(Z, Y) = C^\infty(Z).$$

Conormality of a distribution to Y encompasses being a smooth function away from Y and a prescribed type of singularity as one approaches Y . The singularity at Y is characterized by a symbol in the direction normal to Y .

Definition A.2. Let $E \rightarrow Y$ be a real vector bundle over a k -dimensional smooth manifold. We say that $a \in C^\infty(E)$ is a symbol of order $m \in \mathbb{R}$ if for each vector bundle trivialization $E|_U \rightarrow U \times \mathbb{R}^p$ over an open set $U \subseteq Y$ every compact $K \subseteq U$ and multiindices $\beta \in \mathbb{N}^k$, $\gamma \in \mathbb{N}^p$ there is a constant $C > 0$ such that

$$|\partial_x^\beta \partial_\xi^\gamma a(x, \xi)| \leq C(1 + |\xi|^2)^{(m-|\gamma|)/2}, \text{ for } (x, \xi) \in K \times \mathbb{R}^p.$$

Here we have identified a with a function on $U \times \mathbb{R}^p$ using the trivialization $E|_U \rightarrow U \times \mathbb{R}^p$.

We let $S^m(E)$ denote the space of symbols of order m on $E \rightarrow Y$. If $Y = \mathbb{R}^k$ and $E = Y \times \mathbb{R}^p$, we write $S^m(\mathbb{R}^k \times \mathbb{R}^p)$ instead of $S^m(E)$.

Let $Z = \mathbb{R}^N$ and $Y = \mathbb{R}^k$ embedded into the first factor and write the coordinates on \mathbb{R}^N as $x = (y, z)$ where $y \in \mathbb{R}^k$ and $z \in \mathbb{R}^{N-k}$. It was proven in [23, Theorem 18.2.8] that the map $a \mapsto u$, where

$$(43) \quad u(x) := \int_{\mathbb{R}^{N-k}} a(y, z, \xi) e^{iz \cdot \xi} d\xi,$$

defines a surjection $S^m(\mathbb{R}^N \times \mathbb{R}^{N-k}) \rightarrow I^m(Z, Y)$. Moreover, it was also proven in [23, Lemma 18.2.1] that the map $\tilde{a} \mapsto u$, where

$$(44) \quad u(x) := \int_{\mathbb{R}^{N-k}} \tilde{a}(y, \xi) e^{iz \cdot \xi} d\xi,$$

defines an isomorphism $S^m(\mathbb{R}^k \times \mathbb{R}^{N-k}) \rightarrow I^m(Z, Y)$. The relationship between the a appearing in Equation (43) and the \tilde{a} appearing in Equation (44) is stated in [23, Lemma 18.2.1] to be

$$\tilde{a}(y, \xi) \sim \sum_{\alpha \in \mathbb{N}^{N-k}} \frac{1}{\alpha!} \partial_z^\alpha D_\xi^\alpha a(y, z, \xi)|_{z=0}.$$

The right hand side is an asymptotic sum, for more details see [23], and is only well defined up to smooth functions. Here \sim denotes equality up to smooth errors.

In the case of a general manifold, we need some further set up. Consider a smooth manifold Z and a smooth submanifold $Y \subseteq Z$ closed in the topology of Z . We define the conormal bundle $N^*Y \rightarrow Y$ as the kernel of the restriction mapping $T^*Z|_Y \rightarrow T^*Y$ and the normal bundle $NY \rightarrow Y$ as the vector bundle $NY := TZ|_Y/TY$. By construction, $N^*Y = (NY)^*$ as vector bundles on Y . We equip the normal bundle with a fixed fiberwise volume density. After making a choice of metric, one can find a tubular neighborhood U_Y of $Y \subseteq Z$ and construct a fiber preserving diffeomorphism $\psi : U_Y \rightarrow NY$ mapping Y onto the zero section of $N(Y)$. Here we consider the tubular neighborhood to be a ball bundle over Y . We pick a $\chi_Y \in C^\infty(U_Y)$ with $\chi_Y = 1$ in a neighborhood of Y such that the projection mapping $\text{supp}(\chi_Y) \xrightarrow{\psi} NY \rightarrow Y$ is proper. We define the mapping

$$q : S^m(N^*Y) \rightarrow \mathcal{D}'(Z),$$

as the composition of the fiberwise Fourier transform $S^m(N^*Y) \rightarrow \mathcal{D}'(NY)$ (with respect to the volume density), pullback along ψ and multiplication by χ_Y . We can also define the mapping

$$\sigma : I^m(Z, Y) \rightarrow \mathcal{D}'(N^*Y),$$

as the composition of multiplication by χ_Y , pullback along ψ and Fourier transform in the fiber direction (with respect to the volume density).

Theorem A.3. *The maps q and σ give well defined maps*

$$q : S^m(N^*Y) \rightarrow I^m(Z, Y) \quad \text{and} \quad \sigma : I^m(Z, Y) \rightarrow S^m(N^*Y),$$

that induce isomorphisms

$$\begin{aligned}\bar{q} : S^m(N^*Y)/S^{-\infty}(N^*Y) &\rightarrow I^m(Z, Y)/C^\infty(Z) \quad \text{and} \\ \bar{\sigma} : I^m(Z, Y)/C^\infty(Z) &\rightarrow S^m(N^*Y)/S^{-\infty}(N^*Y).\end{aligned}$$

The quotient mappings

$$\begin{aligned}\bar{q}_m : S^m(N^*Y)/S^{m-1}(N^*Y) &\rightarrow I^m(Z, Y)/I^{m-1}(Z, Y) \quad \text{and} \\ \bar{\sigma}_m : I^m(Z, Y)/I^{m-1}(Z, Y) &\rightarrow S^m(N^*Y)/S^{m-1}(N^*Y),\end{aligned}$$

are mutual inverses and independent of the choice of ψ and χ_Y .

Proof. The details of the proof can be found in [23, Chapter XVIII.2]. The reader can also consult [39, Chapter 4], see Theorem 4.3.2 and Theorem 4.3.16. \square

To digest this theorem, let us consider what this means in practice. Fix a tubular neighborhood U_Y of Y inside Z and a bundle isomorphism $\psi : U_Y \rightarrow NY$. It says that $u \in I^m(Z, Y)$ if and only if u is C^∞ outside Y and we for any $\chi \in C_c^\infty(Z)$ supported in a small enough open set $U \subseteq U_Y$ can write

$$\chi u(y, z) = \int_{\mathbb{R}^{N-k}} a(y, z, \xi) e^{iz \cdot \xi} d\xi + r(x),$$

in local coordinates on U for a symbol $a \in S^m(N^*Y)$ and an $r \in C^\infty(U)$. We note that the order of u is determined by the order of the symbol a .

An important subclass of conormal distributions that we will later make use of are the classical ones. Classically simplifies computations and are by definition given by asymptotic sums, making them highly suitable for studying asymptotics of the magnitude with. Let us define classicality in the context of both symbols and conormal distributions. For a vector bundle $E \rightarrow Y$ and $t > 0$, we let λ_t denote the dilation of functions and distributions in the fiber direction. We say that a function or distribution u is homogeneous of degree $m \in \mathbb{C}$ if $\lambda_t u = t^m u$ for all $t > 0$.

Definition A.4. Consider a smooth N -dimensional manifold Z and a smooth k -dimensional submanifold $Y \subseteq Z$ closed in the topology of Z . Fix a tubular neighborhood U_Y of Y inside Z , a bundle isomorphism $\psi : U_Y \rightarrow NY$ and a \cdot . Let $m \in \mathbb{C}$.

- (1) We say that $a \in C^\infty(N^*Y)$ is a classical symbol of order m if there is a collection $(a_j)_{j \in \mathbb{N}} \subseteq C^\infty(N^*Y \setminus Y)$, with a_j being homogeneous of degree $m - j$, and a smooth function $\chi \in C^\infty(N^*Y)$ which is 1 outside a fiberwise precompact neighborhood of Y , such that

$$a \sim \sum_{j=0}^{\infty} \chi a_j.$$

We let $CS^m(N^*Y)$ denote the space of classical symbols of order m .

- (2) For $m \notin \mathbb{Z}$, we say that $u \in \mathcal{D}'(Z)$ is a classical conormal distribution of order m if $u|_{Z \setminus Y}$ is smooth and there is a collection $(u_j)_{j \in \mathbb{N}} \in \mathcal{D}'(NY)$, with u_j being homogeneous of degree $j - m - N + k$, and a smooth function $\chi \in C^\infty(Z)$ which is 1 in a neighborhood of Y such that for each l there is an L with

$$\chi u - \sum_{j=0}^L \psi^* u_j \in C^l(Z).$$

We let $CI^m(Z, Y)$ denote the space of classical conormal distributions of order m .

- (3) For $m \in \mathbb{Z}$, we say that $u \in \mathcal{D}'(Z)$ is a classical conormal distribution of order m if $u|_{Z \setminus Y}$ is smooth and there is a collection $(u_j)_{j \in \mathbb{N}} \in \mathcal{D}'(NY)$, with u_j being homogeneous of degree $j - m - N + k$, a collection $(p_j)_{j \in \mathbb{N}} \in C^\infty(NY)$ of fiber wise homogeneous polynomials, with each p_j being of degree $j - m - N + k$, and a smooth function $\chi \in C^\infty(Z)$ which is 1 in a neighborhood of Y such that for each l there is an L with

$$\chi u - \sum_{j=0}^L \psi^*(u_j + p_j \log |\cdot|) \in C^l(Z).$$

We let $CI^m(Z, Y)$ denote the space of classical conormal distributions of order m .

Let us make some remarks on the case $m \in \mathbb{Z}$. Here $\log|\cdot| \in C^\infty(NY \setminus Y)$ denotes the fiberwise logarithm in some choice of Riemannian metric. The reader should note that $p_j = 0$ if $j < m + N - k$. The appearance of logarithms is due to the fact that despite symbols of the form $|\xi|^{-m}$ being classical and homogeneous outside $\xi \neq 0$, their Fourier transforms are not homogeneous if $m \in -N + k - \mathbb{N}$ and contains logarithms. We shall explore this further in our main example below where the Fourier transform of $|\xi|^{k-N}$ is precisely the logarithm. This is of relevance to us since the conormal distribution defined from the operator Q is a logarithm (see Proposition 2.5).

Proposition A.5. *It holds that $CS^m(N^*Y) \subseteq S^{\text{Re}(m)}(N^*Y)$ and $CI^m(Z, Y) \subseteq I^{\text{Re}(m)}(Z, Y)$. Moreover, the maps from Theorem A.3 induce mappings*

$$q : CS^m(N^*Y) \rightarrow CI^m(Z, Y) \quad \text{and} \quad \sigma : CI^m(Z, Y) \rightarrow CS^m(N^*Y),$$

that induce isomorphisms

$$\begin{aligned} \bar{q} : CS^m(N^*Y)/S^{-\infty}(N^*Y) &\rightarrow CI^m(Z, Y)/C^\infty(Z) \quad \text{and} \\ \bar{\sigma} : CI^m(Z, Y)/C^\infty(Z) &\rightarrow CS^m(N^*Y)/S^{-\infty}(N^*Y), \\ \bar{q}_m : CS^m(N^*Y)/CS^{m-1}(N^*Y) &\rightarrow CI^m(Z, Y)/CI^{m-1}(Z, Y) \quad \text{and} \\ \bar{\sigma}_m : CI^m(Z, Y)/CI^{m-1}(Z, Y) &\rightarrow CS^m(N^*Y)/CS^{m-1}(N^*Y) \end{aligned}$$

The last two mappings are independent of the choices of ψ and χ_Y and are mutual inverses.

The reader should note that if we pick a Riemannian metric on N^*Y , the restriction mapping to the associated sphere bundle $SN^*Y \subseteq N^*Y$ of the leading term induces an isomorphism $CS^m(N^*Y)/CS^{m-1}(N^*Y) \cong C^\infty(SN^*Y)$. We can as such view $\bar{\sigma}_m$ as an isomorphism $CI^m(Z, Y)/CI^{m-1}(Z, Y) \rightarrow C^\infty(SN^*Y)$.

We shall not give the details of the proof of Theorem A.5, but merely note that it is a trivial consequence of Theorem A.3 in conjunction with homogeneity properties of the Fourier transform. We shall give the details for the case of interest to us below in Subsection A.1.

Conormal distributions often arise from the Schwartz kernels of pseudodifferential operators. To study the operator Q , we will need to apply conormal distributions to pseudodifferential operators with parameters. But let us nevertheless provide the reader with some context first by reviewing pseudodifferential operators from the point of view of conormal distributions.

Consider a manifold M and set $Z := M \times M$. We let $Y := \text{Diag}_M \subseteq Z$ denote the diagonal. Consider a continuous operator $A : C_c^\infty(M) \rightarrow C^\infty(M)$ with Schwartz kernel $K_A \in \mathcal{D}'(M \times M)$. In terms of conormal distributions, A is called a pseudodifferential operator of order $m \in \mathbb{R}$ if $K_A \in I^m(M \times M, \text{Diag}_M)$. The standard definition of a pseudodifferential operator is equivalent by Theorem A.3. In the case at hand $Y \cong M$ via the projection map and $N^*Y = T^*M$. As such, the symbol mapping induces isomorphisms $I^m(M \times M, \text{Diag}_M)/C^\infty(M \times M) \cong S^m(T^*M)/S^{-\infty}(T^*M)$ and analogous mappings in the classical setting.

A.1. Examples of conormal distributions. In this subsection we consider two examples coming from powers and logarithms of distance functions in Euclidean space. Both examples will play a pivotal role in our study of the operator Q . In lack of finding a good reference we have included a large amount of details.

We consider $Z = \mathbb{R}^N$ and $Y = \mathbb{R}^k$ viewed as a submanifold of Z . We write the Euclidean coordinates as $x = (y, z)$ for $y \in Y = \mathbb{R}^k$ and $z \in \mathbb{R}^{N-k}$. For $t \in \mathbb{R}_+$, define the automorphism λ_t of $C_c^\infty(Z)$ by $\lambda_t \varphi(y, z) := \varphi(y, tz)$. We define λ_t on $\mathcal{D}'(Z)$ by

$$\langle \lambda_t u, \varphi \rangle := t^{-k} \langle u, \lambda_{1/t} \varphi \rangle.$$

Take an $\alpha \in \mathbb{C}$ and define $u_{\alpha,0} \in C^\infty(Z \setminus Y)$ by $u_{\alpha,0}(y, z) := |z|^{-\alpha}$. If $\alpha \notin k - N + \mathbb{N}$, [22, Theorem 3.2.3] guarantees that $u_{\alpha,0}$ has a unique extension to a distribution $u_\alpha \in \mathcal{D}'(Z)$ such that $\lambda_t u_\alpha = t^{-\alpha} u_\alpha$ for all $t > 0$. A short computation shows that u_α depends holomorphically on $\alpha \in \mathbb{C} \setminus (k - N + \mathbb{N})$. If $f = f(w)$ is a function depending holomorphically on w in a punctured

neighborhood of α , we write $\text{F.P.}_{w=\alpha}f$ for the constant coefficient in the Laurent expansion of f around $w = \alpha$. For $\alpha \in \mathbb{C}$, we use the notation $\text{F.P.}|z|^{-\alpha}$ to denote the distribution

$$\langle \text{F.P.}|z|^{-\alpha}, \varphi \rangle := \text{F.P.}_{w=\alpha} \langle u_w, \varphi \rangle.$$

It follows from the construction that if $\alpha \in \mathbb{C} \setminus (k - N + \mathbb{N})$, then $\text{F.P.}|z|^{-\alpha}$ is homogeneous of degree $-\alpha$, i.e. $\lambda_t \text{F.P.}|z|^{-\alpha} = t^{-\alpha} \text{F.P.}|z|^{-\alpha}$ for all $t > 0$. By [22, Theorem 3.2.3], $\text{F.P.}|z|^{-\alpha} = u_\alpha$ for $\alpha \in \mathbb{C} \setminus (k - N + \mathbb{N})$. The computation [22, Equation (3.2.24), page 76] shows that if $\alpha \in k - N + \mathbb{N}$ we have for all $t > 0$ that

$$\lambda_t \text{F.P.}|z|^{-\alpha} - t^{-\alpha} \text{F.P.}|z|^{-\alpha} = \log(t) \sum_{\beta \in \mathbb{N}^k: |\beta| = \alpha + N - k} c_\beta \partial_z^\beta \delta_Y,$$

for some coefficients c_β and $\delta_Y \in \mathcal{D}'(Z)$ is the distribution defined by

$$\langle \delta_Y, \varphi \rangle = \int_{\mathbb{R}^k} \varphi(y, 0) dy.$$

Proposition A.6. *Let $\alpha \in \mathbb{C}$. Consider the distribution $\text{F.P.}|z|^{-\alpha}$ on \mathbb{R}^{N-k} as constructed in the preceding paragraph (for a fixed y). Then $\text{F.P.}|z|^{-\alpha}$ is a tempered distribution on \mathbb{R}^{N-k} and for $\xi \neq 0$, we have that*

$$\mathcal{F}\text{F.P.}|z|^{-\alpha} = \begin{cases} \pi^{(N-k)/2} 2^{\alpha+N-k} \frac{\Gamma(\frac{\alpha+N-k}{2})}{\Gamma(-\frac{\alpha}{2})} |\xi|^{-N+k-\alpha}, & \alpha \in \mathbb{C} \setminus (-N+k-2\mathbb{N}), \\ \frac{\pi^{(N-k)/2} (-1)^l}{2^{2l} l! \Gamma(\frac{N-k}{2} + l)} |\xi|^{2l} (-\log|\xi|^2 + \beta_{l, N-k}), & \alpha = -N+k-2l, l \in \mathbb{N}, \end{cases}$$

where

$$\beta_{l, N-k} := 2 \log(2) + \frac{1}{2} \psi((N-k)/2 + l) - H_l - \gamma,$$

and $\psi(z) := \frac{\Gamma'(z)}{\Gamma(z)}$, $H_0 = 0$ and $H_l := \sum_{j=1}^l \frac{1}{j}$ for $l > 0$, and γ is the Euler-Mascheroni constant.

We note that in the case $\alpha \in 2\mathbb{N}$, the right hand side of the equality has a zero which is compatible with that $\mathcal{F}\text{F.P.}|z|^{-\alpha}$ is a sum of distributions supported at $\xi = 0$. A closely related computation of $\langle \mathcal{F}\text{F.P.}|z|^{-\alpha}, \varphi \rangle$ for test functions φ with vanishing moments may be found in [36], Lemma 25.2. It implies the above result except for the value of $\beta_{l, N-k}$.

Proof. The distribution $\text{F.P.}|z|^{-\alpha}$ is polynomially bounded and therefore a tempered distribution. To compute its Fourier transform, we first assume that $\alpha \notin -N+k-2\mathbb{N}$. In this case, $\text{F.P.}|z|^{-\alpha}$ is homogeneous of degree α , so its Fourier transform must be homogeneous of degree $-N+k-\alpha$. Moreover, $\text{F.P.}|z|^{-\alpha}$ is $O(N-k)$ -invariant, so its Fourier transform must be $O(N-k)$ -invariant. The space of $O(N-k)$ -invariant tempered distributions of degree $\alpha \notin -N+k-2\mathbb{N}$ is one-dimensional and therefore there is a constant c_α such that $\mathcal{F}\text{F.P.}|z|^{-\alpha} = c_\alpha \text{F.P.}|\xi|^{-N+k-\alpha}$ if $\alpha \notin (-N+k-2\mathbb{N}) \cup 2\mathbb{N}$ and $\mathcal{F}\text{F.P.}|z|^{-\alpha}$ is supported at $\xi = 0$ if $\alpha \in 2\mathbb{N}$. To verify that $c_\alpha = \pi^{(N-k)/2} 2^{\alpha+N-k} \frac{\Gamma(\frac{\alpha+N-k}{2})}{\Gamma(-\frac{\alpha}{2})}$ one restricts to α in the range $(-N+k, -(N-k)/2)$ and integrates against the standard Gaussian (see [37, Exemple 5, Chapitre VII.7] but beware of the non-standard convention with 2π in the Fourier transform of [37]).

The case that $\alpha \in -N+k-2\mathbb{N}$ follows from that the Fourier transform is continuous and therefore commutes with taking finite part values. Let us compute the details. Near $\alpha_l = -N+k-2l$ we have that

$$c_\alpha = \frac{2a_l}{\alpha - \alpha_l} + b_l + O(\alpha - \alpha_l),$$

where

$$a_l = \frac{\pi^{(N-k)/2} (-1)^l}{2^{2l} l! \Gamma(\frac{N-k}{2} + l)} \quad \text{and} \quad b_l = \frac{\pi^{(N-k)/2} (-1)^l}{2^{2l} l! \Gamma(\frac{N-k}{2} + l)} \beta_{l, N-k}.$$

The expressions for a_l and b_l follows from the fact that the Γ -function has no zeroes and its poles are situation in $\zeta \in -\mathbb{N}$ where

$$\text{Res}_{\zeta=-k} \Gamma(\zeta) = \frac{(-1)^k}{k!} \quad \text{and} \quad \text{F.P.}_{\zeta=-k} \Gamma(\zeta) = \frac{(-1)^{k+1}}{k!} (\gamma + H_k).$$

Taking the finite part of $\mathcal{F}\text{F.P.}|z|^\alpha = c_\alpha \text{F.P.}|\xi|^{-N+k-\alpha}$ at $\alpha = \alpha_l$, gives us for $\xi \neq 0$ that

$$\mathcal{F}\text{F.P.}|z|^{-N+k-2l} = b_l |\xi|^{2l} - 2a_l |\xi|^{2l} \log |\xi| = b_l |\xi|^{2l} - a_l |\xi|^{2l} \log |\xi|^2,$$

which produces the expression in the statement of the proposition. \square

Write $CS^{m,k}$ for the class of log-classical symbols and $CI^{m,k}$ for the corresponding class of log-classical conormal distributions, see more in [30].

Proposition A.7. *Let $Z = \mathbb{R}^N$, $Y = \mathbb{R}^k$ and $\alpha \in \mathbb{C} \setminus \{0\}$. The distribution $\text{F.P.}|z|^{-\alpha}$ constructed in the preceding paragraph satisfies that $\text{F.P.}|z|^{-\alpha} \in CI^{\alpha-N+k,j}(Z, Y)$, where $j \in \{0, 1\}$ and $j = 1$ if and only if $\alpha \in N - k + 2\mathbb{N}$, and*

$$\bar{\sigma}_{\alpha-N+k}(\text{F.P.}|z|^{-\alpha})(y, \xi) = \begin{cases} \pi^{(N-k)/2} 2^{-\alpha+N-k} \frac{\Gamma(\frac{-\alpha+N-k}{2})}{\Gamma(\frac{\alpha}{2})} |\xi|^{-N+k+\alpha}, & \alpha \in \mathbb{C} \setminus (N-k+2\mathbb{N}), \\ \frac{\pi^{(N-k)/2} (-1)^l}{2^{2l} l! \Gamma(\frac{N-k}{2} + l)} |\xi|^{2l} (-\log |\xi|^2 + \beta_{l, N-k}), & \alpha = N - k + 2l, l \in \mathbb{N}. \end{cases}$$

Proof. For notational simplicity, set $u := \text{F.P.}|z|^{-\alpha}$. Pick a compactly supported $\chi_0 \in C_c^\infty(\mathbb{R}^{N-k})$ with $\chi = 1$ near $z = 0$. Write $u = u_1 + u_2$ where $\hat{u}_1 = \chi \hat{u}$ and $u_2 = u - u_1$. The distribution u_1 is constant in the y -direction. By Proposition A.6, u_1 is the Fourier transform in the z -direction of the compactly supported distribution $c_\alpha \chi \text{F.P.}|\xi|^{-N+k+\alpha}$, so u_1 is smooth. It therefore suffices to prove that $u_2 \in CI^{\alpha-N+k}(Z, Y)$. Consider the classical symbol $a(\xi) := (1 - \chi(\xi)) \hat{u}(\xi)$. By viewing a as a constant function of $x = (y, z) \in \mathbb{R}^n$, we have that $a \in CS^{-N+k+\alpha, j}(\mathbb{R}^N \times \mathbb{R}^{N-k})$ by the computation of Proposition A.6. Moreover,

$$u_2(y, z) = \int_{\mathbb{R}^{N-k}} a(y, z, \xi) e^{iz \cdot \xi} d\xi,$$

so $u_2 \in CI^{\alpha-N+k}(Z, Y)$ by Proposition A.5. \square

For $Z = \mathbb{R}^N$ and $Y = \mathbb{R}^k$, with coordinates $x = (y, z)$ for $y \in Y = \mathbb{R}^k$ and $z \in \mathbb{R}^{N-k}$, we define the function $u(y, z) := \log |z|^2$. This function is locally integrable and defines a distribution. Let us turn to studying its singularity at Y . First we need an elementary Fourier transform computation that follows from Proposition A.6.

Proposition A.8. *The function $\tilde{u}(z) := \log |z|^2$ on \mathbb{R}^{N-k} is a tempered distribution and for $\xi \neq 0$,*

$$\mathcal{F}\tilde{u} = -2\pi(N-k-1)! \omega_{N-k-1} |\xi|^{k-N},$$

where ω_{N-k-1} denotes the volume of the unit ball in dimension $N-k-1$.

Proof. Since \tilde{u} is locally integrable and polynomially bounded, it is a tempered distribution. To compute its Fourier transform, we note that $\lambda_t \tilde{u} - \tilde{u} = \log(t)$. In other words, \tilde{u} is homogeneous of degree 0 up to a constant function. In particular, $\mathcal{F}\tilde{u}$ must therefore be homogeneous of degree $-n+k$ up to a multiple of δ_0 . Moreover, \tilde{u} is $O(N-k)$ -invariant, so $\mathcal{F}\tilde{u}$ must be $O(N-k)$ -invariant. The space of $O(N-k)$ -invariant elements of $\mathcal{S}'(\mathbb{R}^{N-k})/\mathcal{C}\delta_0$ that are homogeneous of degree $-N+k$ is one dimensional and is spanned by the equivalence class defined from $\text{F.P.} \frac{1}{|\xi|^{N-k}}$. It is clear that $\mathcal{F}\tilde{u}$ is not compactly supported, so its class in $\mathcal{S}'(\mathbb{R}^{N-k})/\mathcal{C}\delta_0$ is non-zero. It follows from the computation in Proposition A.7 that for $\xi \neq 0$,

$$\mathcal{F}\tilde{u} = -2^{N-k} \pi^{(N-k)/2} \Gamma\left(\frac{N-k}{2}\right) |\xi|^{k-N}.$$

A short computation, as in Proposition 2.8, gives that

$$\Gamma\left(\frac{N-k}{2}\right) \Gamma\left(\frac{N-k-1}{2} + 1\right) = \sqrt{\pi} \frac{(N-k-1)!}{2^{N-k-1}}.$$

Therefore we have that

$$2^{N-k} \pi^{(N-k)/2} \Gamma\left(\frac{N-k}{2}\right) = 2\pi(N-k-1)! \frac{\pi^{(N-k-1)/2}}{\Gamma\left(\frac{N-k-1}{2} + 1\right)} = 2\pi(N-k-1)! \omega_{N-k-1}.$$

\square

Proposition A.9. *Let $Z = \mathbb{R}^N$ and $Y = \mathbb{R}^k$. The distribution $\log|z|$ satisfies that $\log|z| \in CI^{-N+k}(Z, Y)$ and*

$$\bar{\sigma}_{-N+k}(\log|z|)(y, \xi) = -2\pi(N-k-1)!\omega_{N-k-1}|\xi|^{k-N}, \quad \xi \neq 0.$$

The proof is analogous to that of Proposition A.7 and is therefore omitted.

For $g \in C^\infty(Z)$ and a submanifold $Y \subseteq Z$ with a prescribed tubular neighborhood $U \cong NY$, we define the transversal Hessian of g in $y \in Y$ as the symmetric bilinear form on $(NY)_y$ defined from the Hessian at the zero section of g restricted to $(NY)_y$ along the tubular neighborhood.

Proposition A.10. *Assume that $Y \subseteq Z$ is an k -dimensional smooth submanifold of an N -dimensional smooth manifold. Let $\tilde{G} \in C^\infty(Z)$ be a smooth function such that*

- \tilde{G} and $d\tilde{G}$ vanishes on Y ;
- $\tilde{G}(x) > 0$ for $x \notin Y$; and
- for each $y \in Y$, the transversal Hessian $H_{\tilde{G}}$ of \tilde{G} (defined as in Definition 2.1) is a positive definite quadratic form on the transversal tangent bundle of $Y \subseteq Z$.

Then $\log \tilde{G} \in CI^{-N+k}(Z; Y)$ and its principal symbol $\bar{\sigma}_{-N+k}(\log \tilde{G}) \in C^\infty(N^*Y \setminus Y)$ is given by

$$\bar{\sigma}_{-N+k}(\log \tilde{G})(y, \xi) = -2\pi(N-k-1)!\omega_{N-k-1}|g_{\tilde{G}}(\xi, \xi)|^{-(N-k)/2}, \quad \xi \neq 0,$$

where $g_{\tilde{G}}$ is the metric dual to the transversal Hessian $H_{\tilde{G}}$.

Proof. The statement is local, so we can assume that there is an open set $U \subseteq \mathbb{R}^n$ containing 0 such that $Z = U$ and $Y = U \cap \mathbb{R}^k$. Since dg vanishes on TY , we can consider its restriction to Y to be a section $dg|_Y : Y \rightarrow N^*Y$. Under the assumption that the transversal Hessian of g is non-degenerate in all points of Y , we can assume that U is taken small enough to be able to choose coordinates (\tilde{y}, \tilde{z}) on U such that $g(\tilde{y}, \tilde{z}) = |\tilde{z}|^2$ for $\tilde{z} \neq 0$. For notational simplicity, we assume that $g(y, z) = |z|^2$ and that $Z = U = \mathbb{R}^N$ and $Y = \mathbb{R}^k$. The proposition now follows from Proposition A.9. \square

APPENDIX B. PARAMETER DEPENDENT PSEUDODIFFERENTIAL OPERATORS

The reason for the above adventure through the seven circles of conormal distributions is that it simplifies our description of the magnitude operator as a pseudodifferential operator with parameter. We shall make heavy use of parameter dependent pseudodifferential operators, so let us recall their definition and describe their kernel structure. In this subsection we only consider classical symbols and operators.

Definition B.1. Let $\Gamma \subseteq \mathbb{C}$ be a conical subset, $m \in \mathbb{C}$ and $n, p \in \mathbb{N}$. A function $a = a(x, \xi, R) \in C^\infty(\mathbb{R}^p \times \mathbb{R}^n \times \Gamma)$ is said to be a classical symbol with parameter of order m if there exists a sequence $(a_j)_{j \in \mathbb{N}} \subseteq C^\infty(\mathbb{R}^p \times ((\mathbb{R}^n \times \Gamma) \setminus \{0\}))$ of functions homogeneous of degree $m - j$ in the sense that

$$a_j(x, t\xi, tR) = t^{m-j}a_j(x, \xi, R), \quad \forall t > 0,$$

and a $\chi \in C_c^\infty(\mathbb{R}^n \times \Gamma)$ such that

$$a \sim \sum_j (1 - \chi)a_j.$$

We write $CS^m(\mathbb{R}^p \times \mathbb{R}^n; \Gamma)$ for the space of classical symbols with parameter of order m .

For an n -dimensional manifold M , we define $CS^m(M; \Gamma) \subseteq C^\infty(T^*M \oplus \Gamma)$ as the space of functions $a \in C^\infty(T^*M \oplus \Gamma)$ such that in each local chart $\phi : U \subseteq M \rightarrow \mathbb{R}^n$ and $\chi \in C_c^\infty(U)$, $(D\phi)^*(\chi a) \in CS^m(\mathbb{R}^n \times \mathbb{R}^n; \Gamma)$.

Remark B.2. Let us clarify what we mean by $a \sim \sum_j (1 - \chi)a_j$. We write $a \sim \sum_j (1 - \chi)a_j$ when for any $\alpha \in \mathbb{N}^p$, $\beta \in \mathbb{N}^n$, $k, N \in \mathbb{N}$ and compact $K \subseteq \mathbb{R}^p$ there is a constant $C > 0$ and an $N_0 \in \mathbb{N}$ such that

$$\left| \partial_x^\alpha \partial_\xi^\beta \partial_R^k \left(a - \sum_{j=0}^{N_0} (1 - \chi)a_j \right) (x, \xi, R) \right| \leq C(1 + |\xi| + |R|)^{-N}, \quad x \in K, \xi \in \mathbb{R}^n, R \in \Gamma.$$

If this condition is satisfied, one can in fact take N_0 to be any natural number $\geq N - |\beta| - k + \operatorname{Re}(m)$. Conversely, if $(a_j)_{j \in \mathbb{N}} \subseteq C^\infty(\mathbb{R}^p \times ((\mathbb{R}^n \times \Gamma) \setminus \{0\}))$ is a sequence of functions homogeneous of degree $m - j$ then for any $\chi \in C_c^\infty(\mathbb{R}^p \times \Gamma)$ which equals 1 in a neighborhood of $0 \in \mathbb{R}^n \times \Gamma$ there exists an $a \in CS^m(\mathbb{R}^p \times \mathbb{R}^n; \Gamma)$ with $a \sim \sum_j (1 - \chi)a_j$.

Remark B.3. We use the notation $S^{-\infty}(\mathbb{R}^p \times \mathbb{R}^n; \Gamma)$ for the space of smoothing symbols, i.e. the space of $a \in C^\infty(\mathbb{R}^p \times \mathbb{R}^n \times \Gamma)$ satisfying that for any $\alpha \in \mathbb{N}^p$, $\beta \in \mathbb{N}^n$, $k, N \in \mathbb{N}$ and compact $K \subseteq \mathbb{R}^p$ there is a constant $C > 0$ such that

$$\left| \partial_x^\alpha \partial_\xi^\beta \partial_R^k a \right| \leq C(1 + |\xi| + |R|)^{-N}.$$

Note that $S^{-\infty}(\mathbb{R}^p \times \mathbb{R}^n; \Gamma) = \bigcap_{m \in \mathbb{R}} CS^m(\mathbb{R}^p \times \mathbb{R}^n; \Gamma)$. For $a, b \in C^\infty(\mathbb{R}^p \times \mathbb{R}^n \times \Gamma)$ we write $a \sim b$ if $a - b$ is a smoothing symbol.

Remark B.4. For an n -dimensional manifold M , we shall implicitly use a Riemannian metric on M and write $S(T^*M \oplus \Gamma) \rightarrow M$ to denote fiber bundle of elements $(\xi, R) \in T^*M \oplus \Gamma$ such that $|\xi|^2 + |R|^2 = 1$. An element $a \in CS^m(M; \Gamma)$ is determined modulo lower order terms (i.e. $CS^{m-1}(M; \Gamma)$) by its principal symbol. In other words, if $a \sim \sum_j (1 - \chi)a_j$ then $a - a_0 \in CS^{m-1}(M; \Gamma)$. Since a_0 is homogeneous of degree m , it is determined by its restriction to $S(T^*M \oplus \Gamma)$. We deduce that $a \mapsto a_0|_{S(T^*M \oplus \Gamma)}$ defines an isomorphism

$$CS^m(M; \Gamma)/CS^{m-1}(M; \Gamma) \cong C^\infty(S(T^*M \oplus \Gamma)).$$

We will mainly be interested in the case $p = 2n$ or $p = n$ where n is the dimension of the underlying manifold. If $\Gamma = \mathbb{R}$ our definition coincides with that of classical symbols on $\mathbb{R}^p \times \mathbb{R}^{n+1}$.

Let us turn to the quantization of symbols to operators. We can consider an element $K \in S^{-\infty}(\mathbb{R}^{2n} \times \{0\}; \Gamma) \subseteq C^\infty(\mathbb{R}^{2n} \times \Gamma)$ as a family of operators $K = K(R) : C_c^\infty(\mathbb{R}^n) \rightarrow C^\infty(\mathbb{R}^n)$ defined by

$$K(R)f(x) := \int_{\mathbb{R}^n} K(x, y, R)f(y)dy.$$

Such a family of operators is called a smoothing operator with parameter. We write $\Psi^{-\infty}(\mathbb{R}^n; \Gamma)$ for the space of smoothing operators with parameters. If $a = a(x, y, \xi, R) \in CS^m(\mathbb{R}^{2n} \times \mathbb{R}^n; \Gamma)$ we can associate a family of operators $Op(a) = Op(a)(R) : C_c^\infty(\mathbb{R}^n) \rightarrow C^\infty(\mathbb{R}^n)$ defined by

$$Op(a)(R)f(x) := \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} a(x, y, \xi, R)f(y)e^{i\xi \cdot (x-y)}dyd\xi.$$

We note that in this order of integration, both integrals are well defined for $f \in C_c^\infty(\mathbb{R}^n)$. Alternatively, we can consider this as an oscillatory integral in which the order of integration is irrelevant. A family of operators $A = A(R) : C_c^\infty(\mathbb{R}^n) \rightarrow C^\infty(\mathbb{R}^n)$ is called a classical pseudodifferential operator of order m with parameter if there is a classical symbol $a \in CS^m(\mathbb{R}^{2n} \times \mathbb{R}^n; \Gamma)$ and a smoothing operator $K \in \Psi^{-\infty}(\mathbb{R}^n; \Gamma)$ such that

$$A = Op(a) + K.$$

We write $\Psi_{\text{cl}}^m(\mathbb{R}^n; \Gamma)$ for the space of classical pseudodifferential operators of order m with parameter. For two operators $A_1, A_2 : C_c^\infty(\mathbb{R}^n) \rightarrow C^\infty(\mathbb{R}^n)$ we write $A_1 \sim A_2$ if $A_1 - A_2$ is smoothing with parameter. If $A \in \Psi_{\text{cl}}^m(\mathbb{R}^n; \Gamma)$ satisfies $A = \chi A \chi'$ for some $\chi, \chi' \in C_r^\infty(\mathbb{R}^n)$, we say that A is compactly supported.

Proposition B.5. *Let $m \in \mathbb{C}$ and consider a conical subset $\Gamma \subseteq \mathbb{C}$. Then the following holds:*

- (1) *Assume that $\Gamma = \mathbb{R}$ and let $A \in \Psi_{\text{cl}}^m(\mathbb{R}^n; \mathbb{R})$ be a classical pseudodifferential operators of order m with parameter. There is an $A' \in \Psi_{\text{cl}}^m(\mathbb{R}^n; \mathbb{R})$ with $A \sim A'$ such that the family of Schwartz kernels $K_{A'}(R) \in \mathcal{D}'(\mathbb{R}^n \times \mathbb{R}^n)$ satisfies that $\mathcal{F}_{R \rightarrow \eta} K_{A'} \in CI^m(\mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}; \mathbb{R}^n \times \{0\})$, where $\mathcal{F}_{R \rightarrow \eta} K_{A'}$ denotes the Fourier transform in the R -direction and we identify $\mathbb{R}^n \times \{0\} \subseteq \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}$ as a submanifold via the diagonal inclusion $\mathbb{R}^n \rightarrow \mathbb{R}^n \times \mathbb{R}^n$.*

(2) Let $A = Op(a) + K \in \Psi_{\text{cl}}^m(\mathbb{R}^n; \Gamma)$ for an $a \in CS^m(\mathbb{R}^{2n} \times \mathbb{R}^n; \Gamma)$. For

$$\tilde{a}(x, \xi, R) := \sum_{\alpha \in \mathbb{N}^n} \frac{1}{\alpha!} \partial_y^\alpha D_\xi^\alpha a(x, y, \xi, R)|_{x=y} \in CS^m(\mathbb{R}^n \times \mathbb{R}^n; \Gamma),$$

we have that $Op(a) \sim Op(\tilde{a})$. In particular, we can write $A = Op(\tilde{a}) + \tilde{K}$ for an $\tilde{a} \in CS^m(\mathbb{R}^n \times \mathbb{R}^n; \Gamma)$ and a smoothing operator with parameter \tilde{K} .

(3) An element A as in (2) is uniquely determined up to smoothing operators by \tilde{a} modulo smoothing symbols. In particular, the mapping

$$\Psi_{\text{cl}}^m(\mathbb{R}^n; \Gamma) / \Psi^{-\infty}(\mathbb{R}^n; \Gamma) \rightarrow CS^m(\mathbb{R}^n; \Gamma) / S^{-\infty}(\mathbb{R}^n; \Gamma), \quad A \mapsto \tilde{a},$$

is a well defined isomorphism.

(4) If $A = Op(\tilde{a}) \in \Psi_{\text{cl}}^m(\mathbb{R}^n; \Gamma)$ and $B = Op(\tilde{b}) \in \Psi_{\text{cl}}^{m'}(\mathbb{R}^n; \Gamma)$ for $\tilde{a} \in CS^m(\mathbb{R}^n; \Gamma)$ and $\tilde{b} \in CS^{m'}(\mathbb{R}^n; \Gamma)$ then $AB \in \Psi_{\text{cl}}^{m+m'}(\mathbb{R}^n; \Gamma)$ and $AB \sim Op(a\#b)$ where

$$a\#b(x, \xi) \sim \sum_{\alpha} \frac{1}{\alpha!} D_\xi^\alpha a(x, \xi, R) \partial_x^\alpha b(x, \xi, R) \in CS^{m+m'}(\mathbb{R}^n \times \mathbb{R}^n; \Gamma).$$

Proof. For (1) we take A' to be the classical pseudodifferential operator defined from that $K_{A'} = \chi K_A$ where $\chi \in C^\infty(\mathbb{R}^n \times \mathbb{R}^n)$ is defined from $\chi(x, y) = \chi_0(|x - y|^2)$ for some $\chi_0 \in C_c^\infty(\mathbb{R})$ with $\chi_0 = 1$ in a neighborhood of 0. It is clear from standard arguments with oscillatory integrals that $K_{A'} - K_A \in \Psi^{-\infty}(\mathbb{R}^n; \mathbb{R})$. Since $K_{A'}$ is properly supported, we can write $A' = Op(a)$ for an $a \in CS^m(\mathbb{R}^{2n} \times \mathbb{R}^n; \mathbb{R})$. It follows from the arguments around Equation (43) that

$$[\mathcal{F}_{R \rightarrow \eta} K_{A'}](x, y, \eta) = \int_{\mathbb{R}^{n+1}} a(x, y, \xi, R) e^{i(x-y)\xi - i\eta R} dR d\xi,$$

defines an element of $CI^m(\mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}; \mathbb{R}^n \times \{0\})$.

Items (2), (3) and (4) follow as in the case of no parameters, see [38, Theorem 3.1], [39, Theorem 4.3.2] (with $\rho = 1$ and $\delta = 0$), and [38, Theorem 3.4], respectively. \square

Remark B.6. With regards to item (1) in Proposition B.5, we refer to $a \in CS^m(\mathbb{R}^{2n} \times \mathbb{R}^n; \Gamma)$ as a “two-variable symbol” and $\tilde{a} \in CS^m(\mathbb{R}^n \times \mathbb{R}^n; \Gamma)$ as a “one-variable symbol”. In the applications we are concerned with, the two-variable symbol is easily obtained. Due to item (3), i.e. that the two variable symbol is not determined by the operator, and item (4) we shall need to use the transition from two-variable symbols to one-variable symbols. The isomorphism in item (3) of Proposition B.5 is called the full symbol isomorphism.

For the remainder of this subsection let M denote an n -dimensional smooth manifold. For simplicity we assume that M is compact. For $m \in \mathbb{C}$ and a conical subset $\Gamma \subseteq \mathbb{C}$, an family of operators $A = A(R) : C^\infty(M) \rightarrow C^\infty(M)$ is a classical pseudodifferential operators of order m with parameter if for any coordinate chart $\phi : U \subseteq M \rightarrow U' \subseteq \mathbb{R}^n$, and any $\chi, \chi' \in C_c^\infty(U')$ the composition of operators defined on $f \in C_c^\infty(\mathbb{R}^n)$ as

$$f \mapsto \phi^*(\chi f) \mapsto A\phi^*(\chi f) \mapsto (\phi^{-1})^*(\chi' A\phi^*(\chi f)) \in C^\infty(\mathbb{R}^n),$$

is a classical pseudodifferential operators of order m with parameter on \mathbb{R}^n . We write $\Psi_{\text{cl}}^m(M; \Gamma)$ for the space of classical pseudodifferential operators of order m with parameter on M . The space of smoothing operators $\Psi^{-\infty}(M; \Gamma)$ is by definition isomorphic to the space $\mathcal{S}(\Gamma; C^\infty(M \times M))$ consisting of Schwartz functions from Γ into the Fréchet space of smoothing operators $C^\infty(M \times M)$.

Since the full symbol of a pseudodifferential operator with parameter is uniquely determined modulo smoothing symbols in local coordinates, we obtain a full symbol isomorphism

$$\Psi_{\text{cl}}^m(M; \Gamma) / \Psi^{-\infty}(M; \Gamma) \cong CS^m(M; \Gamma) / S^{-\infty}(M; \Gamma).$$

The next theorem contains well known statements that summarize the properties of pseudodifferential operators with parameter we shall make use of. First, let us introduce some notation. The $2n + 1$ -dimensional manifold $M \times M \times \mathbb{R}$ carries an \mathbb{R} -action by translation in the third factor. We shall assume that $Z \subseteq M \times M \times \mathbb{R}$ is an \mathbb{R} -invariant open neighborhood of $\text{Diag}_M \times \{0\} \subseteq M \times M \times \mathbb{R}$, i.e. that there is an open neighborhood $U \subseteq M \times M$ of

Diag_M such that $Z = U \times \mathbb{R}$. From Proposition B.5 and the coordinate transformation laws of pseudodifferential operators we deduce the following theorem.

Theorem B.7. *Let $m \in \mathbb{C}$ and let $\Gamma \subseteq \mathbb{C}$ be a sector.*

- (1) *Assume that $\Gamma = \mathbb{R}$. Item (1) of Proposition B.5 and the full symbol mapping, respectively, induce isomorphisms*

$$\begin{aligned} CI^m(Z; \text{Diag}_M \times \{0\})/C^\infty(Z) &\cong \Psi_{\text{cl}}^m(M; \Gamma)/\Psi^{-\infty}(M; \Gamma) \cong \\ &\cong CS^m(M; \Gamma)/S^{-\infty}(M; \Gamma), \end{aligned}$$

and

$$\begin{aligned} CI^m(Z; \text{Diag}_M \times \{0\})/CI^{m-1}(Z; \text{Diag}_M \times \{0\}) &\cong \\ &\cong CS^m(M; \Gamma)/CS^{m-1}(M; \Gamma) \cong C^\infty(S(T^*M \oplus \Gamma)). \end{aligned}$$

- (2) *The principal mapping that maps a pseudodifferential operator with parameter to its leading homogeneous term defines an isomorphism*

$$\bar{\sigma}_m : \Psi_{\text{cl}}^m(M; \Gamma)/\Psi_{\text{cl}}^{m-1}(M; \Gamma) \rightarrow CS^m(M; \Gamma)/CS^{m-1}(M; \Gamma) \cong C^\infty(S(T^*M \oplus \Gamma)).$$

- (3) *If $A \in \Psi_{\text{cl}}^m(M; \Gamma)$ and $B \in \Psi_{\text{cl}}^{m'}(M; \Gamma)$, then $AB \in \Psi_{\text{cl}}^{m+m'}(M; \Gamma)$. The full symbol $a\#b$ of AB is computed in local coordinates from the full symbols a and b of A and B , respectively, and the formula*

$$a\#b(x, \xi) \sim \sum_{\alpha} \frac{1}{\alpha!} D_{\xi}^{\alpha} a(x, \xi, R) \partial_x^{\alpha} b(x, \xi, R) \in CS^{m+m'}(M; \Gamma).$$

Definition B.8. Let $m \in \mathbb{C}$, M be a compact manifold and $A \in \Psi_{\text{cl}}^m(M; \Gamma)$. We say that A is elliptic if $\bar{\sigma}_m(A) \in C^\infty(S(T^*M \oplus \Gamma))$ is an invertible element.

Remark B.9. The reader should beware that in item (3) of Theorem B.7, the full symbol depends on the choice of coordinates.

The following proposition follows from item (2) of Theorem B.7 and asymptotic completeness of the space of pseudodifferential operators with parameter.

Proposition B.10. *Let $A \in \Psi_{\text{cl}}^m(M; \Gamma)$ be an operator with full symbol $a \in CS^m(M; \Gamma)$. The following are equivalent:*

- *A is elliptic*
- *There exists $b \in CS^{-m}(M; \Gamma)$ with $ab = 1$ outside a compact subset of $T^*M \oplus \Gamma$.*
- *There exists $B \in \Psi_{\text{cl}}^{-m}(M; \Gamma)$ such that $AB - 1, BA - 1 \in \Psi^{-\infty}(M; \Gamma)$.*

We recall that we have implicitly used a Riemannian metric on M . Using this metric, we also define the scale of Hilbert spaces $H^s(M) := (1 + \Delta)^{-s/2} L^2(M)$ for $s \in \mathbb{R}$, where Δ is the Laplace operator defined from the Riemannian metric. We also consider the scale $H_R^s(M) := (R^2 + \Delta)^{-s/2} L^2(M)$ defined for $R \in \mathbb{R} \setminus \{0\}$ and $s \in \mathbb{R}$. We note that elliptic regularity implies that $H^s(M) = H_R^s(M)$ as vector spaces with equivalent norms, but their Hilbert space structure differs in a non-uniform way as R varies. The difference in Hilbert space structure is best seen from the mapping properties of pseudodifferential operators with parameters, which is made precise in the following theorem.

Theorem B.11. *Let $\Gamma \subseteq \Gamma_{\alpha}(0) \cup -\Gamma_{\alpha}(0)$ be a bisector with opening angle $< \alpha \in [0, \pi/2)$, and $A \in \Psi_{\text{cl}}^m(M; \Gamma)$. For any $s, t \in \mathbb{R}$ with $t \leq s - \text{Re}(m)$ there is a constant $C_{s,t} > 0$ such that for $R \in \Gamma \setminus \{0\}$*

$$\|Af\|_{H_{|R|}^t} \leq C_{s,t} (1 + |R|)^{t-s+\text{Re}(m)} \|f\|_{H_{|R|}^s}, \forall f \in H^s(M).$$

Moreover, $A : H_{|R|}^s(M) \rightarrow H_{|R|}^t(M)$ is compact if $t < s - \text{Re}(m)$. In particular, if A is elliptic then there are for any $s \in \mathbb{R}$ constants $C_s, R_0 > 0$ such that for $|R| > R_0$

$$\frac{1}{C_s} \|f\|_{H_{|R|}^s} \leq \|Af\|_{H_{|R|}^{s-\text{Re}(m)}} \leq C_s \|f\|_{H_{|R|}^s}, \forall f \in H^s(M).$$

Proof. It is clear that $A : H_{|R|}^s(M) \rightarrow H_{|R|}^t(M)$ is compact if $t < s - \text{Re}(M)$ by the corresponding statement without parameters. We start by showing the norm estimate $\|Af\|_{H_{|R|}^t} \leq C_{s,t}(1 + |R|)^{t-s+\text{Re}(m)}\|f\|_{H_{|R|}^s}$. Since Γ does not contain the imaginary axis, the operator $(R^2 + \Delta)^{m/2} \in \Psi_{\text{cl}}^m(M; \Gamma)$ is an invertible elliptic element. Assume that $\alpha \in [\pi/4, \pi/2)$. Since there is a positive angle between Γ and the imaginary axis we have a lower quadratic form estimate

$$(R^2 + \Delta)^*(R^2 + \Delta) \geq (1 - \cos^2(2\alpha))|R|^4, \quad \text{for } R \in \Gamma.$$

It therefore follows by interpolation that

$$(45) \quad \|(R^2 + \Delta)^{m/2}f\|_{H_{|R|}^t} \leq C_{s,t}(1 + |R|)^{t-s+\text{Re}(m)}\|f\|_{H_{|R|}^s}, \quad f \in H^s(M).$$

By combining Equation (45) with item (4) of Theorem B.7 we can reduce to $t = s = m = 0$. To prove the theorem in the case $t = s = m = 0$, we need to show that point evaluation in R defines a continuous representation $\Psi_{\text{cl}}^0(M; \Gamma) \rightarrow \mathcal{B}(L^2(M))$ with uniform seminorm bounds. This statement follows from the fact that point evaluations in R defines a continuous mapping $\Psi_{\text{cl}}^0(M; \Gamma) \rightarrow \Psi_{\text{cl}}^0(M)$ in the respective Fréchet topologies with uniform seminorm bounds and the Calderón-Vaillancourt theorem implies that the action of zero order pseudodifferential operators on $L^2(M)$ defines a continuous representation $\Psi_{\text{cl}}^0(M) \rightarrow \mathcal{B}(L^2(M))$.

Fix an $s \in \mathbb{R}$. We want to show that for some R_0 and C_s , we have the estimates $\frac{1}{C_s}\|f\|_{H_{|R|}^s} \leq \|Af\|_{H_{|R|}^{s-\text{Re}(m)}} \leq C_s\|f\|_{H_{|R|}^s}$ for $|R| > R_0$. By the preceding paragraph, the upper estimate holds. To show the lower estimate we note that if $A \in \Psi_{\text{cl}}^m(M; \Gamma)$ is elliptic, then there is a $B \in \Psi_{\text{cl}}^{-m}(M; \Gamma)$ such that $AB - 1, BA - 1 \in \Psi^{-\infty}(M; \Gamma)$. Write A_s for the continuous operator $H_{|R|}^s(M) \rightarrow H_{|R|}^{s-\text{Re}(m)}(M)$ defined from A and B_s for the continuous operator $H_{|R|}^{s-\text{Re}(m)}(M) \rightarrow H_{|R|}^s(M)$ defined from B . From the preceding paragraph, we have that $\|1 - A_s B_s\|, \|1 - B_s A_s\| = O(R^{-\infty})$ so there is an R_0 such that $B_s(R) - A_s(R)^{-1}$ is a smoothing operator for $|R| > R_0$. We conclude from the preceding paragraph that there is a C' such that $\|B_s g\|_{H_{|R|}^s} \leq C'\|g\|_{H_{|R|}^{s-\text{Re}(m)}}$ for all $g \in H_{|R|}^{s-\text{Re}(m)}(M)$. This shows that $\frac{1}{C'}\|f\|_{H_{|R|}^s} \leq \|Af\|_{H_{|R|}^{s-\text{Re}(m)}}$ for all $f \in H_{|R|}^s(M)$ and $|R| > R_0$, and we conclude the lower estimate. \square

Corollary B.12 (Gårding inequality). *Let $\Gamma \subseteq \Gamma_\alpha(0) \cup -\Gamma_\alpha(0)$ be a bisector with opening angle $< \alpha \in [0, \pi/2)$, and $A \in \Psi_{\text{cl}}^m(M; \Gamma)$ a formally self-adjoint operator with strictly positive principal symbol, i.e. for some $\varepsilon > 0$, $\bar{\sigma}(A)(x, \xi, R) \geq \varepsilon$ for all $|\xi|^2 + |R|^2 = 1$ and $x \in M$.*

Then for large R , the quadratic form $f \mapsto \langle Af, f \rangle_{L^2}$ is continuous, positive and coercive on $H_{|R|}^{\text{Re}(m)/2}$. To be precise, there is an $R_0 > 0$ and a $C > 0$ such that

$$\frac{1}{C}\|f\|_{H_{|R|}^{\text{Re}(m)/2}} \leq \langle Af, f \rangle_{L^2} \leq C\|f\|_{H_{|R|}^{\text{Re}(m)}}, \quad \forall f \in H^s(M), \quad |R| > R_0.$$

Proof. It follows in the same way as in [38, Proposition 6.1], that there is a $B \in \Psi_{\text{cl}}^{m/2}(M; \Gamma)$ such that $r := A - B^*B \in \Psi^{-\infty}(M; \Gamma)$. It is clear that B is elliptic and that R is self-adjoint, so the previous theorem implies that for some $C_0 > 0$,

$$\frac{1}{C}\|f\|_{H_{|R|}^{\text{Re}(m)/2}} - \langle rf, f \rangle_{L^2} \leq \langle Af, f \rangle_{L^2} \leq C\|f\|_{H_{|R|}^{\text{Re}(m)}} + \langle rf, f \rangle_{L^2}, \quad \forall f \in H^s(M).$$

Since $r \in \Psi^{-\infty}(M; \Gamma)$, we have that $\|r(R)\|_{L^2 \rightarrow L^2} = O(|R|^{-\infty})$ and the corollary follows. \square

One of the reasons for introducing conormal distributions above was that it will give us a direct way of verifying that the magnitude operator is an elliptic pseudodifferential operator with parameter. We will next present the relevant technical result needed for such an endeavor. For a compact smooth n -dimensional manifold, identified with its diagonal in $M \times M$, we let $Z = U \times \mathbb{R} \subseteq M \times M \times \mathbb{R}$ denote an \mathbb{R} -invariant tubular neighborhood of the diagonal of M . After fixing some metric on M that we for notational simplicity assume to have injectivity radius > 1 , we can assume that we have exponential coordinates on Z identifying Z with the ball bundle

$$BM \oplus \mathbb{R} = \{(x, v, \eta) \in TM \oplus \mathbb{R} : |v| < 1\}.$$

This is the ball bundle in the normal bundle of $\text{Diag}_M \times \{0\} \subseteq Z$ under the identification of M with its diagonal. We denote the coordinate in the \mathbb{R} -direction by η since for conormal distributions it plays the role of a dual variable (cf. item (1) of Proposition B.5). In the situation at hand, the conormal bundle of $\text{Diag}_M \times \{0\} \subseteq Z$ is under the identification of M with its diagonal given by $T^*M \oplus \mathbb{R} \rightarrow M$. We denote coordinates in the fiber of the conormal bundle by (ξ, R) . We note that under these identifications, we have a canonical isomorphism

$$(46) \quad CS^m(N^*(\text{Diag}_M \times \{0\} \subseteq Z)) \cong CS^m(M; \mathbb{R}).$$

Compare this to Theorem B.7. A problem we need to address is that the correspondence between operators and symbols is only well behaved modulo smoothing terms, and the notions of smoothing symbols and operators in the setting of conormal distributions differ from that in the setting of the parameter dependent calculus. To address the issue at the level of operators, instead of equivalence classes, we introduce a uniform notion of asymptotic expansions.

Assume that $K \in CI^m(Z; M \times \{0\})$ in exponential coordinates has a classical asymptotic expansion of the form $K \sim \sum_{j=0}^{\infty} \chi K_j$ as in Definition A.4. Here $\chi \in C_c^\infty(U)$ is a function with $\chi = 1$ near the diagonal. In the above choice of coordinates, K_j is a smooth function on $TM \oplus \mathbb{R} \setminus (M \times \{0\})$. If $m \notin \mathbb{Z}$, K_j is homogeneous of degree $-m - j - n - 1$. If $m \in \mathbb{Z}$, $K_j = u_j + p_j \log(|v|^2 + \eta^2)$ where u_j is homogeneous of degree $-m - j - n - 1$ and p_j is a homogeneous polynomial in (v, η) of degree $-m - j - n - 1$ (in particular $p_j = 0$ if $j < -m - n - 1$).

Definition B.13. We shall say that $K \in CI^m(Z; M \times \{0\})$ has a **uniform asymptotic expansion** if for any $\alpha \in \mathbb{N}^p$, $\beta \in \mathbb{N}^n$, $k, N \in \mathbb{N}$ there is a constant $C > 0$ and an $N_0 \in \mathbb{N}$ such that for $R \neq 0$

$$\left| \partial_x^\alpha \partial_v^\beta \partial_\eta^k \left(K - \sum_{j=0}^{N_0} \chi K_j \right) \right| \leq C(1 + |v| + |\eta|)^{-N}.$$

For $x \in M$, we write $\exp_x : T_x M \rightarrow M$ for the exponential map. Note that under our assumption on the injectivity radius, $\exp_x : B_x M \rightarrow M$ is a diffeomorphism onto its range.

Proposition B.14. *Let M be a smooth n -dimensional manifold and Z is as in the preceding paragraphs. Assume that $K \in CI^m(Z; M \times \{0\})$ admits a uniform asymptotic expansion. Then there is a pseudodifferential operator with parameter $A \in \Psi^m(M; \mathbb{R})$ such that for any R , the Schwartz kernel of $A(R)$ is given by $\mathcal{F}_{\eta \rightarrow R} K$. In other words, for $f \in C^\infty(M)$, $A(R)f$ is defined as the oscillatory integral*

$$A(R)f(x) := \int_{B_x M \oplus \mathbb{R}} K(x, v, \eta) f(\exp_x(v)) e^{i\eta R} dv d\eta.$$

In particular, the full symbol of A in $CS^m(M; \mathbb{R})/S^{-\infty}(M; \mathbb{R})$ coincides with the full symbol of K in $CS^m(N^(\text{Diag}_M \times \{0\} \subseteq Z))/S^{-\infty}(N^*(\text{Diag}_M \times \{0\} \subseteq Z))$ under the isomorphism (46).*

Proof. If the uniform asymptotic expansion of K only contains one term, i.e. $K = K_0$, the statement of the proposition holds by homogeneity properties of the Fourier transform. As such, the proposition is in fact a statement concerning the asymptotic completeness of the space of pseudodifferential operators with parameters. This is proven just as in the usual setting (see [23, Proposition 18.1.3]). \square

APPENDIX C. INVERTING Q IN AN EXTENDED BOUTET DE MONVEL CALCULUS

In this Appendix we prove Theorem 4.9. More generally, we discuss the magnitude operator $\mathcal{Z}_X(R)$ in the context of general techniques which have been developed for elliptic pseudodifferential boundary problems. While $\mathcal{Z}_X(R)$ is not always a pseudodifferential operator, for distance functions having property (MR) Theorem 4.5 shows that $\mathcal{Z}_X(R)$ closely relates to the pseudodifferential operator $Q_X(R)$. While there is an extensive theory for boundary problems for elliptic differential operators, developed over many decades, boundary problems for pseudodifferential operators pose severe difficulties since the operators are nonlocal. Much of the recent study was motivated by fractional powers of the Laplacian. In particular, for a larger class of operators pseudodifferential methods were developed starting with [19], under the assumption that the operator satisfies a μ -transmission condition at the boundary. They allowed to show, for example,

regularity results for solutions of elliptic Dirichlet problems and integration by parts formulas. A basic observation is that such problems could be reduced to problems in the Boutet de Monvel calculus, originally developed for differential boundary problems [19]. While the results in this Appendix are not directly used in this article, they complement the specific analysis of $\mathcal{Z}(R)$ by results and challenges for a large class of related problems. We refer to [15, 19] for further context and notation.

As the techniques are local, it is sufficient to prove Theorem 4.9 for $X \subset \subset \mathbb{R}^n$. Consider the Dirichlet problem

$$(47) \quad \begin{aligned} r^+ P_R u_R &= f && \text{in } X \\ u_R &= 0 && \text{in } X^c. \end{aligned}$$

Generalizing $Q_X(R)$, here P_R is a parameter dependent, classical pseudodifferential operator on X which is elliptic of order $-m = -n - 1$, of type $-m/2$ and has factorization index $-m/2$. We make the further assumption that it is of infinite regularity.

We reduce (47) to a problem of type and order 0 in the Boutet de Monvel calculus, where a parametrix can be constructed using classical techniques. The reduction relies on the following operators to raise the order of P_R , while preserving support in X , see [19, Section 2.5]:

Definition C.1. For $l \in \frac{1}{2}\mathbb{Z}$, define $\lambda_{\pm}^l \in \Psi_{\text{cl}}^l(\mathbb{R}^n; \Gamma)$

$$\lambda_-^l(\xi, R) = \left(\langle \xi', R \rangle \psi \left(\frac{\xi_n}{b \langle \xi', R \rangle} \right) - i \xi_n \right)^l, \quad \lambda_+^l(\xi, R) = \overline{\lambda_-^l}(\xi, R).$$

Here $\psi \in \mathcal{S}(\mathbb{R})$ with $\text{supp } \mathcal{F}^{-1}\psi \subset \overline{\mathbb{R}}_-$ and $\psi(0) = 1$, and $b \geq 2 \sup |\partial_t \psi(t)|$. We set $\Lambda_{\pm, R}^l := \text{Op}(\lambda_{\pm}^l(\xi, R))$.

The symbols λ_{\pm}^l are uniformly parameter-elliptic on $\mathbb{R}^n \times \mathbb{R}_+^{n+1}$ of order l , and they are of factorization index and type l , respectively 0.

Composition of (47) on the left with $r_+ \Lambda_{-, R}^{\frac{m}{2}} e_+$ gives

$$r_+ \Lambda_{-, R}^{\frac{m}{2}} e_+ r_+ P_R u_R = r_+ \Lambda_{-, R}^{\frac{m}{2}} P_R u_R = r_+ \Lambda_{-, R}^{\frac{m}{2}} e_+ f.$$

Letting $v_R := r_+ \Lambda_{+, R}^{-\frac{m}{2}} u_R$, equivalently $u_R = \Lambda_{+, R}^{\frac{m}{2}} e_+ v_R$, and denoting $\mathcal{A}_R := \Lambda_{-, R}^{\frac{m}{2}} P_R \Lambda_{+, R}^{\frac{m}{2}}$ gives the following problem for v_R :

$$(48) \quad \begin{aligned} r_+ \mathcal{A}_R e_+ v_R &= r_+ \Lambda_{-, R}^{\frac{m}{2}} e_+ f && \text{in } X \\ v_R &= 0 && \text{in } M \setminus \overline{X}. \end{aligned}$$

We next show that (48) is a problem in the Boutet de Monvel calculus, based on arguments in [19, Section 2.5]:

Proposition C.2. *The operator \mathcal{A}_R is an elliptic parameter-dependent pseudodifferential operator of order, type, and factorization index 0.*

Using a parametrix constructed in this calculus, one recovers the solution to the original problem (47) by the transformation $u_R = \Lambda_{+, R}^{\frac{m}{2}} e_+ v_R$. In particular, this implies the assertions about $Q_X(R)$ in Theorem 4.9. Unfortunately, for the purposes of this paper the general properties of the parametrix after composition with $\Lambda_{+, R}^{\frac{m}{2}}$ are not currently understood sufficiently to directly yield results for the magnitude problem.

The second part of Theorem 4.9, about $\mathcal{Z}_X(R)$, then follows from the decomposition $\mathcal{Z}_X(R)^{-1} = Q_X(R)^{-1} + \mathcal{R}_X(R)$ in Theorem 4.5.

Proof. That it is a pseudodifferential operator follows from the fact that it is a composition of honest pseudodifferential operators. It is immediate that it is of order $\frac{m}{2} - m + \frac{m}{2} = 0$ and factorization index $-\frac{m}{2} + \frac{m}{2} = 0$.

All that is left to show is that the composition $\lambda_{-, R}^{m/2} \# \sigma \# \lambda_{+, R}^{m/2}$ is of type 0, where by σ we denote the symbol of \mathcal{A}_R .

By our assumption on P_R and due to the nature of the Λ operators, we see that we can regard $\tilde{\mathcal{A}}(x, (\xi', R), \xi_n) = \mathcal{A}_R(x, \xi, R)$ as a classical elliptic ψ do, with the parameter R absorbed in the

ξ' variable. For easier notation, we still write the new variable as ξ' , and use v and w to denote the symbols $\lambda_{-,R}^{\frac{m}{2}}$ and $\lambda_{+,R}^{\frac{m}{2}}$ respectively.

We first show that the composition $\sigma\#w$ is of type 0.

Since σ is of type $-\frac{m}{2}$, we have that

$$(49) \quad D_x^\beta D_\xi^\alpha \sigma_{-m-j}(x, \xi', -N) = (-1)^{(-m-2(-\frac{m}{2})-j-|\alpha|)} D_x^\beta D_\xi^\alpha \sigma_{-m-j}(x, \xi', N)$$

for all multiindices β, α , where σ_j are the terms in the polyhomogeneous expansion. Similarly, since w is of type $\frac{m}{2}$ it satisfies

$$(50) \quad D_x^\beta D_\xi^\alpha w_{\frac{m}{2}-j}(x, \xi', -N) = (-1)^{(\frac{m}{2}-2(\frac{m}{2})-j-|\alpha|)} D_x^\beta D_\xi^\alpha w_{\frac{m}{2}-j}(x, \xi', N)$$

for all multiindices β, α , where w_j are the terms in the polyhomogeneous expansion.

Similarly to above, denoting the terms in the polyhomogeneous expansion of $\sigma\#w$ by $[\sigma\#w]_l$, we have that

$$\begin{aligned} [\sigma\#w]_{-\frac{m}{2}-l} &= \left[\sum_{\alpha} \frac{1}{\alpha!} (D_\xi^\alpha \sigma_{-m-j})(D_x^\alpha w_{\frac{m}{2}-k}) \right]_{-\frac{m}{2}-l} \\ &= \sum_{\alpha:|\alpha|=l-k-j} \frac{1}{\alpha!} (D_\xi^\alpha \sigma_{-m-j})(D_x^\alpha w_{\frac{m}{2}-k}) \end{aligned}$$

and thus

$$\begin{aligned} D_x^\beta D_\xi^\gamma [\sigma\#w]_{-\frac{m}{2}-l} &= \sum_{\alpha:|\alpha|=l-k-j} \frac{1}{\alpha!} D_x^\beta D_\xi^\gamma [(D_\xi^\alpha \sigma_{-m-j})(D_x^\alpha w_{\frac{m}{2}-k})] \\ &= \sum_{\alpha:|\alpha|=l-k-j} \frac{1}{\alpha!} D_x^\beta \left[\sum_{\nu:\nu \leq \gamma} \binom{\gamma}{\nu} (D_\xi^{\alpha+\nu} \sigma_{-m-j})(D_\xi^{\gamma-\nu} D_x^\alpha w_{\frac{m}{2}-k}) \right] \\ &= \sum_{\alpha:|\alpha|=l-k-j} \frac{1}{\alpha!} \sum_{\substack{\nu:\nu \leq \gamma \\ \nu':\nu' \leq \beta}} \binom{\beta}{\nu'} \binom{\gamma}{\nu} (D_x^{\beta-\nu'} D_\xi^{\alpha+\nu} \sigma_{-m-j})(D_x^{\alpha+\nu'} D_\xi^{\gamma-\nu} w_{\frac{m}{2}-k}). \end{aligned}$$

By (49) and (50), we have that

$$D_x^{\beta-\nu'} D_\xi^{\alpha+\nu} \sigma_{-m-j}(x, \xi', -N) = (-1)^{(-j-|\alpha+\nu|)} D_x^{\beta-\nu'} D_\xi^{\alpha+\nu} \sigma_{-m-j}(x, \xi', N)$$

and

$$D_x^{\alpha+\nu'} D_\xi^{\gamma-\nu} w_{\frac{m}{2}-k}(x, \xi', -N) = (-1)^{(-\frac{m}{2}-k-|\gamma-m|)} w_{\frac{m}{2}-k}(x, \xi', N).$$

We thus have that

$$\begin{aligned} &D_x^\beta D_\xi^\gamma [\sigma\#w]_{-\frac{m}{2}-l}(x, \xi', -N) \\ &= \sum_{\alpha:|\alpha|=l-k-j} \frac{1}{\alpha!} \sum_{\substack{\nu:\nu \leq \gamma \\ \nu':\nu' \leq \beta}} \binom{\beta}{\nu'} \binom{\gamma}{\nu} (-1)^{(-\frac{m}{2}-(j+k)-(|\alpha+\gamma|))} \\ &\quad (D_x^{\beta-\nu'} D_\xi^{\alpha+\nu} \sigma_{-m-j}(x, \xi', N))(D_x^{\alpha+\nu'} D_\xi^{\gamma-\nu} w_{\frac{m}{2}-k}(x, \xi', N)) \\ &= (-1)^{(-\frac{m}{2}-l-\gamma)} \sum_{\alpha:|\alpha|=l-k-j} \frac{1}{\alpha!} D_x^\beta D_\xi^\gamma [D_\xi^\alpha \sigma_{-m-j}(x, \xi', N) D_x^\alpha w_{\frac{m}{2}-k}(x, \xi', N)] \\ &= (-1)^{(-\frac{m}{2}-l-\gamma)} D_x^\beta D_\xi^\gamma [\sigma\#w]_{-\frac{m}{2}-l}(x, \xi', N) \end{aligned}$$

and we see that this symbol is of type 0.

We now want to show that $v\#\sigma\#w$ is of type 0. We let $s := \sigma\#w$ and thus all we want to show is that $v\#s$ is of type 0, and the calculation is analogous to the one above. Both v and s are of type 0 and in addition the composition is of order 0.

As above,

$$\begin{aligned} [v\#s]_{-l} &= \left[\sum_{\alpha} \frac{1}{\alpha!} (D_{\xi}^{\alpha} v_{-j}) (D_x^{\alpha} s_{-k}) \right]_{-l} \\ &= \sum_{\alpha: |\alpha|=l-k-j} \frac{1}{\alpha!} (D_{\xi}^{\alpha} v_{-j}) (D_x^{\alpha} s_{-k}) \end{aligned}$$

and similarly

$$D_x^{\beta} D_{\xi}^{\gamma} [v\#s]_{-l} = \sum_{\alpha: |\alpha|=l-k-j} \frac{1}{\alpha!} \sum_{\substack{\nu: \nu \leq \gamma \\ \nu': \nu' \leq \beta}} \binom{\beta}{\nu'} \binom{\gamma}{\nu} (D_x^{\beta-\nu'} D_{\xi}^{\alpha+\nu} v_{-j}) (D_x^{\alpha+\nu'} D_{\xi}^{\gamma-\nu} s_{-k}).$$

Since v is such that

$$D_x^{\beta} D_{\xi}^{\alpha} v_{-j}(x, \xi', -N) = (-1)^{(-j-|\alpha|)} D_x^{\beta} D_{\xi}^{\alpha} v_{-j}(x, \xi', N)$$

and the same for s the calculation follows from above. \square

APPENDIX D. THE MEROMORPHIC FREDHOLM THEOREM

The meromorphic Fredholm theorem describes the inverse of a holomorphic family of Fredholm operators. The version we consider can be found in [29, Proposition 1.1.8]. For two Hilbert spaces \mathcal{H}_1 and \mathcal{H}_2 let $\text{Fred}(\mathcal{H}_1, \mathcal{H}_2) \subseteq \mathbb{B}(\mathcal{H}_1, \mathcal{H}_2)$ denote the set of Fredholm operators, which is open in the operator norm.

Theorem D.1. *Let $D \subseteq \mathbb{C}$ be a connected domain and $T : D \rightarrow \text{Fred}(\mathcal{H}_1, \mathcal{H}_2)$ a holomorphic function. Assume that $T(\lambda)$ is invertible for at least one $\lambda \in D$. Then the set*

$$Z := \{z \in D : 0 \in \text{Spec}(T(z))\}$$

is a discrete subset of D , and the function $\lambda \mapsto T(\lambda)^{-1}$ exists as a meromorphic function $D \rightarrow \text{Fred}(\mathcal{H}_1, \mathcal{H}_2)$. For λ near any $z \in Z$, $\lambda \mapsto T(\lambda)^{-1}$ has a pointwise norm-convergent Laurent expansion

$$(51) \quad T(\lambda)^{-1} = \sum_{k=-N}^{\infty} T_k (\lambda - z)^k,$$

where T_k are finite rank operators whenever $k < 0$.

APPENDIX E. PARTIAL FRACTION DECOMPOSITIONS OF SYMBOLS

We note the following structural result from basic calculus:

Lemma E.1. *For all $l, m \in \mathbb{N}$ with $m < 2l$, there exists homogeneous rational functions (with rational coefficients) of (h_+, h_-) denoted by $\beta_{l,m,0,\pm}, \beta_{l,m,1,\pm}, \dots, \beta_{l,m,l-1,\pm}$, where each $\beta_{l,m,j,\pm} = b_{l,m,j,\pm}(h_+, h_-)$ has homogeneous degree $m - j - l$ in (h_+, h_-) , such that*

$$\xi_n^m (\xi_n - h_+)^{-l} (\xi_n - h_-)^{-l} = \sum_{j=0}^{l-1} \beta_{l,m,j,+}(h_+, h_-) (\xi_n - h_+)^{j-l} + \sum_{j=0}^{l-1} \beta_{l,m,j,-}(h_+, h_-) (\xi_n - h_-)^{j-l}.$$

For explicit computations of the first few terms, we require the precise form of $\beta_{l,m,j,\pm}$:

Proposition E.2. *Let $h_+, h_- \in \mathbb{C}$ be two distinct complex numbers and $l \in \mathbb{N}_{>0}$. Then for $\xi_n \in \mathbb{C} \setminus \{h_+, h_-\}$, we have that*

$$\begin{aligned} &(\xi_n - h_+)^{-l} (\xi_n - h_-)^{-l} \\ &= \frac{1}{(h_+ - h_-)^l} \sum_{j=0}^{l-1} \frac{(-1)^j}{(h_+ - h_-)^j} \binom{l+j-1}{j} ((\xi_n - h_+)^{-l+j} + (-1)^{l-j} (\xi_n - h_-)^{-l+j}) \end{aligned}$$

Proof. We show the assertion by induction in l . The case $l=1$ is easily checked:

$$(\xi_n - h_+)^{-1}(\xi_n - h_-)^{-1} = \frac{1}{h_+ - h_-} \left((\xi_n - h_+)^{-1} - (\xi_n - h_-)^{-1} \right).$$

Assume now that the assertion holds for l . To show it for $l+1$, we take derivatives $\frac{\partial^2}{\partial h_+ \partial h_-}$ on both sides:

$$\frac{\partial^2}{\partial h_+ \partial h_-} (\xi_n - h_+)^{-l} (\xi_n - h_-)^{-l} = l^2 (\xi_n - h_+)^{-l-1} (\xi_n - h_-)^{-l-1},$$

respectively,

$$\begin{aligned} & \frac{\partial^2}{\partial h_+ \partial h_-} (\xi_n - h_+)^{-l} (\xi_n - h_-)^{-l} \\ &= \sum_{j=0}^{l-1} (-1)^j \binom{l+j-1}{j} \frac{\partial^2}{\partial h_+ \partial h_-} \left\{ \frac{1}{(h_+ - h_-)^{l+j}} \left((\xi_n - h_+)^{-l+j} + (-1)^{l-j} (\xi_n - h_-)^{-l+j} \right) \right\}. \end{aligned}$$

Consider

$$\begin{aligned} & \frac{\partial^2}{\partial h_+ \partial h_-} \left\{ \frac{1}{(h_+ - h_-)^{l+j}} \left((\xi_n - h_+)^{-l+j} + (-1)^{l-j} (\xi_n - h_-)^{-l+j} \right) \right\} \\ &= \left(\partial_{h_+} \partial_{h_-} \frac{1}{(h_+ - h_-)^{l+j}} \right) \left((\xi_n - h_+)^{-l+j} + (-1)^{l-j} (\xi_n - h_-)^{-l+j} \right) \\ &+ \left(\partial_{h_+} \frac{1}{(h_+ - h_-)^{l+j}} \right) \left((-1)^{l-j} \partial_{h_-} (\xi_n - h_-)^{-l+j} \right) \\ &+ \left(\partial_{h_-} \frac{1}{(h_+ - h_-)^{l+j}} \right) \left(\partial_{h_+} (\xi_n - h_+)^{-l+j} \right). \end{aligned}$$

Using that

$$\begin{aligned} \partial_{h_{\pm}} (\xi_n - h_{\pm})^{-l} &= l (\xi_n - h_{\pm})^{-l-1}, \\ \partial_{h_+} (h_+ - h_-)^{-l} &= -l (h_+ - h_-)^{-l-1}, \quad \partial_{h_-} (h_+ - h_-)^{-l} = l (h_+ - h_-)^{-l-1}, \end{aligned}$$

as well as

$$\partial_{h_+} \partial_{h_-} (h_+ - h_-)^{-l} = -l(l+1) (h_+ - h_-)^{-l-2},$$

we conclude

$$\begin{aligned} & \frac{\partial^2}{\partial h_+ \partial h_-} (\xi_n - h_+)^{-l} (\xi_n - h_-)^{-l} \\ &= \sum_{j=0}^{l-1} (-1)^j \binom{l+j-1}{j} \left\{ -\frac{(l+j)(l+j+1)}{(h_+ - h_-)^{l+j+2}} \left((\xi_n - h_+)^{-l+j} + (-1)^{l-j} (\xi_n - h_-)^{-l+j} \right) \right. \\ &+ \left. \frac{(l-j)(l+j)}{(h_+ - h_-)^{l+j+1}} (\xi_n - h_+)^{-l+j-1} + (-1)^{l-j+1} \frac{(l-j)(l+j)}{(h_+ - h_-)^{l+j+1}} (\xi_n - h_-)^{-l+j-1} \right\}. \end{aligned}$$

Note that the coefficient of $(\xi_n - h_+)^{-l+k} = (\xi_n - h_+)^{-(l+1)+(k+1)}$ is given by

$$\begin{aligned} & (-1)^k \binom{l+k-1}{k} \left(-(l+k)(l+k+1) \right) + (-1)^{k+1} \binom{l+k}{k+1} (l-k-1)(l+k+1) \\ &= (-1)^{k+1} \left(\binom{l+k-1}{k} (l+k) + \binom{l+k}{k+1} (l-k-1) \right) (l+k+1) \\ &= (-1)^{k+1} \left(\frac{(l+k-1)!}{k!(l-1)!} (l+k) + \frac{(l+k)!}{(k+1)!(l-1)!} (l-k-1) \right) (l+k+1) \\ &= (-1)^{k+1} l^2 \binom{l+k+1}{k+1}. \end{aligned}$$

A similar computation applies to the coefficient of $(\xi_n - h_-)^{-l+k} = (\xi_n - h_-)^{-(l+1)+(k+1)}$, so that

$$\begin{aligned} l^2(\xi_n - h_+)^{-l-1}(\xi_n - h_-)^{-l-1} &= \frac{\partial^2}{\partial h_+ \partial h_-} (\xi_n - h_+)^{-l} (\xi_n - h_-)^{-l} \\ &= \frac{1}{(h_+ - h_-)^{l+1}} \sum_{j=0}^l \frac{(-1)^j l^2}{(h_+ - h_-)^j} \binom{l+j}{j} ((\xi_n - h_+)^{-l-1+j} + (-1)^{l+1-j} (\xi_n - h_-)^{-l-1+j}) . \end{aligned}$$

The asserted formula follows for exponent $l+1$ after dividing by l^2 . \square

More generally, by differentiating with respect to h_{\pm} we obtain:

Lemma E.3. *For all $l, m \in \mathbb{N}$ and two distinct complex numbers $h_+, h_- \in \mathbb{C}$, consider the rational function:*

$$K_{m,l}(\xi_n) := (\xi_n - h_+)^{-m} (\xi_n - h_-)^{-l}.$$

This rational function can be decomposed as

$$\begin{aligned} &K_{m,l}(\xi_n) \\ &= \sum_{j=0}^{m-1} \frac{(-1)^j}{(h_+ - h_-)^{l+j}} \binom{l+j-1}{j} (\xi_n - h_+)^{-m+j} + \sum_{j=0}^{l-1} \frac{(-1)^m}{(h_+ - h_-)^{m+j}} \binom{m+j-1}{j} (\xi_n - h_-)^{-l+j} . \end{aligned}$$

These following formulas allow us to obtain explicit partial fraction decompositions for the terms relevant to the factorization of \mathcal{Z}_R .

Corollary E.4. *For all $l \in \mathbb{N}$ and two distinct complex numbers $h_+, h_- \in \mathbb{C}$,*

$$\begin{aligned} \xi_n (\xi_n - h_+)^{-l} (\xi_n - h_-)^{-l} &= K_{l-1,l}(\xi_n) + h_+ K_{l,l}(\xi_n) , \\ \xi_n^2 (\xi_n - h_+)^{-l} (\xi_n - h_-)^{-l} &= K_{l-1,l-1}(\xi_n) + h_- K_{l-1,l}(\xi_n) + h_+ K_{l,l-1}(\xi_n) + h_+ h_- K_{l,l}(\xi_n) , \\ \xi_n^3 (\xi_n - h_+)^{-l} (\xi_n - h_-)^{-l} &= K_{l-2,l-1}(\xi_n) + (2h_+ + h_-) K_{l-1,l-1}(\xi_n) + \\ &\quad + (h_+^2 + h_-^2 + h_+ h_-) K_{l-1,l}(\xi_n) + h_+^2 h_- K_{l,l}(\xi_n) \end{aligned}$$

In particular, we have the formulas

$$\begin{aligned} (h_+ - h_-) \xi_n (\xi_n - h_+)^{-1} (\xi_n - h_-)^{-1} &= h_+ (\xi_n - h_+)^{-1} - h_- (\xi_n - h_-)^{-1} , \\ (h_+ - h_-)^2 \xi_n (\xi_n - h_+)^{-2} (\xi_n - h_-)^{-2} &= h_+ (\xi_n - h_+)^{-2} + h_- (\xi_n - h_-)^{-2} - \\ &\quad - \frac{h_+ + h_-}{h_+ - h_-} ((\xi_n - h_+)^{-1} - (\xi_n - h_-)^{-1}) , \\ (h_+ - h_-)^2 \xi_n^2 (\xi_n - h_+)^{-2} (\xi_n - h_-)^{-2} &= h_+^2 (\xi_n - h_+)^{-2} + h_-^2 (\xi_n - h_-)^{-2} - \\ &\quad - \frac{2h_+ h_-}{h_+ - h_-} ((\xi_n - h_+)^{-1} - (\xi_n - h_-)^{-1}) , \\ (h_+ - h_-)^2 \xi_n^3 (\xi_n - h_+)^{-2} (\xi_n - h_-)^{-2} &= h_+^3 (\xi_n - h_+)^{-2} + h_-^3 (\xi_n - h_-)^{-2} + \\ &\quad + \frac{h_+^2 (h_+ - 3h_-)}{h_+ - h_-} (\xi_n - h_+)^{-1} - \frac{h_-^2 (h_- - 3h_+)}{h_+ - h_-} (\xi_n - h_-)^{-1} . \end{aligned}$$

APPENDIX F. EVALUATION OF SOME BOUNDARY SYMBOLS AT ZERO

For the purpose of computations in Subsection 6.3, we are interested in evaluating some symbols at $\xi = 0$ and $R = 1$. We will use the notations from the Sections 5 and 6 freely. We tacitly assume that $n > 1$ to avoid limit cases.

Recall from Proposition 5.13 that

$$R^2 + g(\xi, \xi) = h_0(\xi_n - h_+)(\xi_n - h_-),$$

where $h_{\pm} = h_{\pm}(x, \xi', R) \in S^1(T^*Y \times \mathbb{R}, Y \times \mathbb{R}; \mathbb{C})$ are of the form

$$h_{\pm}(x, \xi', R) = -\frac{\xi'(b(x))}{h_0(x)} \pm i \frac{\sqrt{R^2 + g_Y(\xi', \xi') - (\xi'(b))^2}}{\sqrt{h_0(x)}}.$$

Here we use the splitting of the metric

$$g = \begin{pmatrix} h_0 & b \\ b^T & g_Y \end{pmatrix},$$

For $x_n = 0$, i.e. on ∂X , we write x' instead of $(x', 0)$. We conclude the following lemma from elementary computations.

Lemma F.1. *The following identities hold on ∂X :*

$$(52) \quad h_0(x')^\nu (-h_+(x', 0, 1))^\nu (-h_-(x', 0, 1))^\nu = 1, \quad \text{for all } \nu \in \mathbb{R}.$$

$$(53) \quad h_\pm(x', 0, 1) = \pm i h_0(x')^{-1/2}.$$

$$(54) \quad \partial_{x_n} h_\pm(x', 0, 1) = \mp \frac{i \partial_{x_n} h_0(x')}{2 h_0(x')^{3/2}}.$$

$$(55) \quad \nabla_{x'} h_\pm(x', 0, 1) = \mp \frac{i \nabla_{x'} h_0(x')}{2 h_0(x')^{3/2}}.$$

$$(56) \quad \nabla_{\xi'} h_\pm(x', 0, 1) = -\frac{b(x')}{h_0(x')}.$$

From Lemma F.1 we deduce the following series of lemmas. We use the notation $\mu = \frac{n+1}{2}$.

Lemma F.2. *The following identity holds on ∂X :*

$$w_{-,0}(x', 0, 0, 1) \partial_{x_n} \partial_{\xi_n}^2 w_{+,0}(x', 0, 0, 1) = \frac{\mu(\mu-1)(\mu-2)}{2n!\omega_n} \partial_{x_n} h_0(x').$$

Proof. We compute that

$$\begin{aligned} & w_{-,0}(x', 0, 0, 1) \partial_{x_n} \partial_{\xi_n}^2 w_{+,0}(x', 0, 0, 1) = \\ &= -\frac{\mu(\mu-1)(\mu-2)}{n!\omega_n} \partial_{x_n} h_+(x', 0, 1) h_0(x)^\mu (-h_-(x', 0, 1))^\mu (-h_-(x', 0, 1))^{\mu-3} = \\ &= -\frac{\mu(\mu-1)(\mu-2)}{n!\omega_n} \partial_{x_n} h_+(x', 0, 1) h_0(x')^3 (-h_-(x', 0, 1))^3 = \\ &= -\frac{\mu(\mu-1)(\mu-2)}{n!\omega_n} \left(-\frac{i \partial_{x_n} h_0(x')}{2 h_0(x')^{3/2}} \right) h_0(x')^3 (i h_0(x')^{-1/2})^3 = \\ &= \frac{\mu(\mu-1)(\mu-2)}{2n!\omega_n} \partial_{x_n} h_0(x') \end{aligned}$$

□

Lemma F.3. *The following identity holds on ∂X :*

$$\partial_{x_n} w_{-,0}(x', 0, 0, 1) \partial_{\xi_n}^2 w_{+,0}(x', 0, 0, 1) = -\frac{\mu^2(\mu-1)}{2 \cdot n!\omega_n} \partial_{x_n} h_0(x').$$

Proof. We compute that

$$\begin{aligned} & \partial_{x_n} w_{-,0}(x', 0, 0, 1) \partial_{\xi_n}^2 w_{+,0}(x', 0, 0, 1) = \\ &= \frac{\mu^2(\mu-1)}{n!\omega_n} \left(\partial_{x_n} h_0(x') h_0(x')^{\mu-1} (-h_-(x', 0, 1))^{\mu-} \right. \\ & \quad \left. - \partial_{x_n} h_-(x', 0, 1) h_0(x')^\mu (-h_-(x', 0, 1))^{\mu-1} \right) (-h_+(x', 0, 1))^{\mu-2} = \\ &= \frac{\mu^2(\mu-1)}{n!\omega_n} \left(\partial_{x_n} h_0(x') \frac{h_-(x', 0, 1)}{h_+(x', 0, 1)} + \partial_{x_n} h_-(x', 0, 1) \frac{h_0(x')}{h_+(x', 0, 1)} \right) = \\ &= \frac{\mu^2(\mu-1)}{n!\omega_n} \left(-\partial_{x_n} h_0(x') + \frac{i \partial_{x_n} h_0(x')}{2 h_0(x')^{3/2}} \frac{h_0(x')}{i h_0(x')^{-1/2}} \right) = -\frac{\mu^2(\mu-1)}{2 \cdot n!\omega_n} \partial_{x_n} h_0(x') \end{aligned}$$

□

Lemma F.4. *The following identity holds on ∂X :*

$$\begin{aligned} w_{-,1}(x', 0, 0, 1) \partial_{\xi_n} w_{+,0}(x', 0, 0, 1) &= \\ &= \frac{i\mu \mathbf{c}_{1,n}(n^2 - 1)}{(n! \omega_n)^2} \left(\frac{3}{2} C^3(x', g \otimes \iota_n g) + \frac{17(n+3)}{4h_0(x)} C^3(x', \iota_n g \otimes \iota_n g \otimes \iota_n g) \right) - \\ &\quad - \frac{7i\mu^3}{4 \cdot n! \omega_n} \left(\partial_{x_n} h_0(x') + \frac{b(x')}{h_0(x')} \cdot \nabla_{x'} h_0(x') \right) \end{aligned}$$

Proof. Recall the computation of $w_{-,1}$ from Proposition 5.20, and the homogeneous symbols $\mathbf{a}_{-,0}$ (of degree 0) and $\mathbf{a}_{-,1}$ (of degree 1) computed in Proposition 5.16 and explicitly given in Equation (36) and (37), respectively. We compute that

$$\begin{aligned} w_{-,1}(x', 0, 0, 1) \partial_{\xi_n} w_{+,0}(x', 0, 0, 1) &= \\ &= \frac{\mu}{n! \omega_n} \left(-\mathbf{a}_{-,0}(x', 0, 1) h_0(x)^\mu (-h_-(x', 0, 1))^{\mu-1} - \mathbf{a}_{-,1}(x', 0, 1) h_0(x)^\mu (-h_-(x', 0, 1))^{\mu-2} - \right. \\ &\quad - i\mu^2 (\partial_{x_n} h_-(x', 0, 1) - \nabla_{\xi'} h_-(x', 0, 1) \cdot \nabla_{x'} h_-(x', 0, 1)) h_0(x)^\mu (-h_-(x', 0, 1))^{\mu-2} - \\ &\quad - i\mu^2 (\partial_{x_n} h_0(x') - \nabla_{\xi'} h_-(x', 0, 1) \cdot \nabla_{x'} h_0(x')) h_0(x')^{\mu-1} (-h_-(x', 0, 1))^{\mu-1} \left. \right) \cdot \\ &\quad \cdot (-h_+(x', 0, 1))^{\mu-1} = \\ &= \frac{\mu}{n! \omega_n} \left(-\mathbf{a}_{-,0}(x', 0, 1) h_0(x) - \mathbf{a}_{-,1}(x', 0, 1) h_0(x) (-h_+(x', 0, 1)) - \right. \\ &\quad - i\mu^2 (\partial_{x_n} h_-(x', 0, 1) - \nabla_{\xi'} h_-(x', 0, 1) \cdot \nabla_{x'} h_-(x', 0, 1)) h_0(x)^2 (-h_+(x', 0, 1)) - \\ &\quad \left. - i\mu^2 (\partial_{x_n} h_0(x') - \nabla_{\xi'} h_-(x', 0, 1) \cdot \nabla_{x'} h_0(x')) \right) = \\ &= \frac{\mu}{n! \omega_n} \left(-\mathbf{a}_{-,0}(x', 0, 1) h_0(x) + i\mathbf{a}_{-,1}(x', 0, 1) h_0(x)^{3/2} - \right. \\ &\quad - i\mu^2 \left(\frac{i}{2} \frac{\partial_{x_n} h_0(x')}{h_0(x')^{3/2}} + \frac{i}{2} \frac{b(x')}{h_0(x')} \cdot \frac{\nabla_{x'} h_0(x')}{h_0(x)^{3/2}} \right) h_0(x)^2 (-ih_0(x')^{-1/2}) - \\ &\quad \left. - i\mu^2 \left(\partial_{x_n} h_0(x') + \frac{b(x')}{h_0(x')} \cdot \nabla_{x'} h_0(x') \right) \right) = \\ &= \frac{\mu}{n! \omega_n} \left(-\mathbf{a}_{-,0}(x', 0, 1) h_0(x) + i\mathbf{a}_{-,1}(x', 0, 1) h_0(x)^{3/2} - \right. \\ &\quad \left. - \frac{3i}{2} \mu^2 \left(\partial_{x_n} h_0(x') + \frac{b(x')}{h_0(x')} \cdot \nabla_{x'} h_0(x') \right) \right). \end{aligned} \tag{57}$$

Computing with Equation (36) and (37) gives us

$$\begin{aligned} \mathbf{a}_{0,-}(x, 0, R) &= \frac{3i\mathbf{c}_{1,n}(n^2 - 1)}{n! \omega_n} C^3(x', g \otimes \iota_n g) \frac{h_0(x')^{-1} h_-(x', 0, 1)}{h_+(x', 0, 1) - h_-(x', 0, 1)} - \\ &\quad - \frac{i\mathbf{c}_{1,n}(n+3)_{3,-2}}{n! \omega_n} \frac{h_0(x')^{-2} h_-(x', 0, 1)^2 (h_-(x', 0, 1) - 3h_+(x', 0, 1))}{(h_+(x', 0, 1) - h_-(x', 0, 1))^3} C^3(x', \iota_n g \otimes \iota_n g \otimes \iota_n g) + \\ &\quad - \frac{i(n+1)^2 (\nabla_{\xi'} h_-(x', 0, 1) \cdot \nabla_{x'} h_+(x', 0, 1) - \partial_{x_n} h_+(x', 0, 1))}{4 (h_+(x', 0, 1) - h_-(x', 0, 1))} = \\ &= -\frac{i\mathbf{c}_{1,n}(n^2 - 1)}{n! \omega_n} \left(\frac{3}{2h_0(x')} C^3(x', g \otimes \iota_n g) + \frac{4(n+3)}{h_0(x)^2} C^3(x', \iota_n g \otimes \iota_n g \otimes \iota_n g) \right) + \\ &\quad + \frac{i\mu^2}{4h_0(x')} \left(\partial_{x_n} h_0(x') + \frac{b(x')}{h_0(x')} \cdot \nabla_{x'} h_0(x') \right), \end{aligned}$$

and

$$\begin{aligned} \mathbf{a}_{1,-}(x, \xi', R) &= \frac{i\mathbf{c}_{1,n}(n+3)_{3,-2}}{n!\omega_n} \frac{h_-(x', 0, 1)^3}{h_0(x)^2(h_+(x', 0, 1) - h_-(x', 0, 1))^2} C^3(x', \iota_n g \otimes \iota_n g \otimes \iota_n g) = \\ &= \frac{\mathbf{c}_{1,n}(n^2-1)(n+3)}{4 \cdot n!\omega_n h_0(x')^{5/2}} C^3(x', \iota_n g \otimes \iota_n g \otimes \iota_n g) \end{aligned}$$

Continuing from Equation (57), we have that

$$\begin{aligned} w_{-,1}(x', 0, 0, 1) \partial_{\xi_n} w_{+,0}(x', 0, 0, 1) &= \\ &= \frac{\mu}{n!\omega_n} \left(\frac{i\mathbf{c}_{1,n}(n^2-1)}{n!\omega_n} \left(\frac{3}{2} C^3(x', g \otimes \iota_n g) + \frac{4(n+3)}{h_0(x)} C^3(x', \iota_n g \otimes \iota_n g \otimes \iota_n g) \right) - \right. \\ &\quad \left. - \frac{i\mu^2}{4} \left(\partial_{x_n} h_0(x') + \frac{b(x')}{h_0(x')} \cdot \nabla_{x'} h_0(x') \right) + i \frac{\mathbf{c}_{1,n}(n^2-1)(n+3)}{4 \cdot n!\omega_n h_0(x')} C^3(x', \iota_n g \otimes \iota_n g \otimes \iota_n g) - \right. \\ &\quad \left. - \frac{3i}{2} \mu^2 \left(\partial_{x_n} h_0(x') + \frac{b(x')}{h_0(x')} \cdot \nabla_{x'} h_0(x') \right) \right) = \\ &= \frac{i\mu\mathbf{c}_{1,n}(n^2-1)}{(n!\omega_n)^2} \left(\frac{3}{2} C^3(x', g \otimes \iota_n g) + \frac{17(n+3)}{4h_0(x)} C^3(x', \iota_n g \otimes \iota_n g \otimes \iota_n g) \right) - \\ &\quad - \frac{7i\mu^3}{4 \cdot n!\omega_n} \left(\partial_{x_n} h_0(x') + \frac{b(x')}{h_0(x')} \cdot \nabla_{x'} h_0(x') \right). \end{aligned}$$

□

Lemma F.5. *The following identity holds on ∂X :*

$$\begin{aligned} w_{-,0}(x', 0, 0, 1) \partial_{\xi_n} w_{+,1}(x', 0, 0, 1) &= \\ &= \frac{i\mathbf{c}_{1,n}(n^2-1)}{(n!\omega_n)^2} \left(-\frac{3(\mu-1)}{2} C^3(x', g \otimes \iota_n g) + \frac{\mu(n+3)}{4h_0(x')} C^3(x', \iota_n g \otimes \iota_n g \otimes \iota_n g) \right) + \\ &\quad + \frac{i\mu^2(3\mu-5)}{4 \cdot n!\omega_n} \left(\partial_{x_n} h_0(x') + \frac{b(x')}{h_0(x')} \cdot \nabla_{x'} h_0(x') \right). \end{aligned}$$

Proof. Recall the computation of $w_{+,1}$ from Proposition 5.20, and the homogeneous symbols $\mathbf{a}_{+,0}$ (of degree 0) and $\mathbf{a}_{+,1}$ (of degree 1) computed in Proposition 5.16 and explicitly given in Equation (34) and (35), respectively. Note that $\mathbf{a}_{+,0}$, $\mathbf{a}_{+,1}$, h_0 and h_{\pm} are independent of ξ_n . We compute that

$$\begin{aligned} w_{-,0}(x', 0, 0, 1) \partial_{\xi_n} w_{+,1}(x', 0, 0, 1) &= \\ &= -\frac{1}{n!\omega_n} h_0(x')^{\mu} (-h_-(x', 0, 1))^{\mu} \cdot \\ &\quad \cdot \left((\mu-1)\mathbf{a}_{+,0}(x', 0, 1)(-h_+(x', 0, 1))^{\mu-2} + (\mu-2)\mathbf{a}_{+,1}(x', 0, 1)(-h_+(x', 0, 1))^{\mu-3} + \right. \\ &\quad \left. + i\mu^2(\mu-2)(\partial_{x_n} h_+(x', 0, 1) - \nabla_{\xi'} h_+(x', 0, 1) \cdot \nabla_{x'} h_+(x', 0, 1))(-h_+(x', 0, 1))^{\mu-3} \right) = \\ &= -\frac{1}{n!\omega_n} \left((\mu-1)\mathbf{a}_{+,0}(x', 0, 1)h_0(x')^2(-h_-(x', 0, 1))^2 + \right. \\ &\quad \left. + (\mu-2)\mathbf{a}_{+,1}(x', 0, 1)h_0(x')^3(-h_-(x', 0, 1))^3 + \right. \\ &\quad \left. + i\mu^2(\mu-2)(\partial_{x_n} h_+(x', 0, 1) - \nabla_{\xi'} h_+(x', 0, 1) \cdot \nabla_{x'} h_+(x', 0, 1))h_0(x')^3(-h_-(x', 0, 1))^3 \right) = \\ &= \frac{1}{n!\omega_n} \left((\mu-1)\mathbf{a}_{+,0}(x', 0, 1)h_0(x') + i(\mu-2)\mathbf{a}_{+,1}(x', 0, 1)h_0(x')^{3/2} + \right. \\ &\quad \left. + \frac{i\mu^2(\mu-2)}{2} \left(\partial_{x_n} h_0(x') + \frac{b(x')}{h_0(x')} \cdot \nabla_{x'} h_0(x') \right) \right). \end{aligned} \tag{58}$$

Computing with Equation (34) and (35) gives us

$$\begin{aligned}
\mathfrak{a}_{0,+}(x', 0, 1) &= -\frac{3i\mathfrak{c}_{1,n}(n^2-1)}{n!\omega_n} C^3(x', g \otimes \iota_n g) \frac{h_+(x', 0, 1)}{h_0(x')(h_+(x', 0, 1) - h_-(x', 0, 1))} + \\
&+ \frac{i\mathfrak{c}_{1,n}(n+3)_{3,-2}}{n!\omega_n} \frac{h_+(x', 0, 1)^2(h_+(x', 0, 1) - 3h_-(x', 0, 1))}{h_0(x)^2(h_+(x', 0, 1) - h_-(x', 0, 1))^3} C^3(x, \iota_n g \otimes \iota_n g \otimes \iota_n g) + \\
&+ i\mu^2 \frac{\nabla_{\xi'} h_-(x', 0, 1) \cdot \nabla_{x'} h_+(x', 0, 1) - \partial_{x_n} h_+(x', 0, 1)}{h_+(x', 0, 1) - h_-(x', 0, 1)} = \\
&= \frac{i\mathfrak{c}_{1,n}(n^2-1)}{n!\omega_n} \left(-3C^3(x', g \otimes \iota_n g) \frac{ih_0(x)^{-1/2}}{h_0(x')2ih_0(x)^{-1/2}} + \right. \\
&+ (n+3) \frac{(ih_0(x')^{-1/2})^2 4ih_0(x)^{-1/2}}{h_0(x)^2(2ih_0(x')^{-1/2})^3} C^3(x, \iota_n g \otimes \iota_n g \otimes \iota_n g) \left. \right) + \\
&+ i\mu^2 \frac{\frac{b(x')}{h_0(x')} \cdot \frac{i\nabla_{x'} h_0(x')}{2h_0(x)^{3/2}} + \frac{i}{2} \frac{\partial_{x_n} h_0(x')}{h_0(x)^{3/2}}}{2ih_0(x')^{-1/2}} = \\
&= \frac{i\mathfrak{c}_{1,n}(n^2-1)}{n!\omega_n} \left(-\frac{3}{2h_0(x')} C^3(x', g \otimes \iota_n g) + \frac{n+3}{2h_0(x')^2} C^3(x, \iota_n g \otimes \iota_n g \otimes \iota_n g) \right) + \\
&+ \frac{i\mu^2}{4h_0(x')} \left(\partial_{x_n} h_0(x') + \frac{b(x')}{h_0(x')} \cdot \nabla_{x'} h_0(x') \right)
\end{aligned}$$

and

$$\begin{aligned}
\mathfrak{a}_{1,+}(x', 0, 1) &= \frac{i\mathfrak{c}_{1,n}(n+3)_{3,-2}}{n!\omega_n} \frac{h_+(x', 0, 1)^3}{h_0(x)^2(h_+(x', 0, 1) - h_-(x', 0, 1))^2} C^3(x, \iota_n g \otimes \iota_n g \otimes \iota_n g) = \\
&= -\frac{\mathfrak{c}_{1,n}(n^2-1)(n+3)}{4 \cdot n!\omega_n h_0(x')^{5/2}} C^3(x, \iota_n g \otimes \iota_n g \otimes \iota_n g).
\end{aligned}$$

Continuing from Equation (58), we have that

$$\begin{aligned}
w_{-,0}(x', 0, 0, 1) \partial_{\xi_n} w_{+,1}(x', 0, 0, 1) &= \\
&= \frac{1}{n!\omega_n} \left((\mu-1) \frac{i\mathfrak{c}_{1,n}(n^2-1)}{n!\omega_n} \left(-\frac{3}{2} C^3(x', g \otimes \iota_n g) + \frac{n+3}{2h_0(x')} C^3(x, \iota_n g \otimes \iota_n g \otimes \iota_n g) \right) + \right. \\
&+ \frac{i\mu^2(\mu-1)}{4} \left(\partial_{x_n} h_0(x') + \frac{b(x')}{h_0(x')} \cdot \nabla_{x'} h_0(x') \right) + \\
&- \frac{i\mathfrak{c}_{1,n}(n^2-1)(n+3)(\mu-2)}{4 \cdot n!\omega_n h_0(x')} C^3(x, \iota_n g \otimes \iota_n g \otimes \iota_n g) + \\
&\quad \left. + \frac{i\mu^2(\mu-2)}{2} \left(\partial_{x_n} h_0(x') + \frac{b(x')}{h_0(x')} \cdot \nabla_{x'} h_0(x') \right) \right) = \\
&= \frac{i\mathfrak{c}_{1,n}(n^2-1)}{(n!\omega_n)^2} \left(-\frac{3(\mu-1)}{2} C^3(x', g \otimes \iota_n g) + \frac{\mu(n+3)}{4h_0(x')} C^3(x, \iota_n g \otimes \iota_n g \otimes \iota_n g) \right) + \\
&+ \frac{i\mu^2(3\mu-5)}{4 \cdot n!\omega_n} \left(\partial_{x_n} h_0(x') + \frac{b(x')}{h_0(x')} \cdot \nabla_{x'} h_0(x') \right).
\end{aligned}$$

□

Lemma F.6. *The following identity holds on ∂X :*

$$\nabla_{x'} w_{-,0}(x', 0, 0, 1) \cdot \nabla_{\xi'} \partial_{\xi_n} w_{+,0}(x', 0, 0, 1) = -\frac{\mu^2(\mu-1)}{2 \cdot n!\omega_n} \frac{b(x')}{h_0(x')} \cdot \nabla_{x'} h_0(x').$$

Proof. We compute that

$$\begin{aligned} & \nabla_{x'} w_{-,0}(x', 0, 0, 1) \cdot \nabla_{\xi'} \partial_{\xi_n} w_{+,0}(x', 0, 0, 1) = \\ &= -\frac{\mu^2(\mu-1)}{n! \omega_n} \left(\nabla_{x'} h_0(x') h_0(x')^{\mu-1} (-h_-(x', 0, 1))^\mu - \nabla_{x'} h_-(x', 0, 1) h_0(x')^\mu (-h_-(x', 0, 1))^{\mu-1} \right) \\ & \quad \cdot \nabla_{\xi'} h_+(x', 0, 1) (-h_+(x', 0, 1))^{\mu-2} = \\ &= -\frac{\mu^2(\mu-1)}{n! \omega_n} \left(\nabla_{x'} h_0(x') \cdot \nabla_{\xi'} h_+(x', 0, 1) \frac{h_-(x', 0, 1)}{h_+(x', 0, 1)} + \right. \\ & \quad \left. + \nabla_{x'} h_-(x', 0, 1) \cdot \nabla_{\xi'} h_+(x', 0, 1) \frac{h_0(x')}{h_+(x', 0, 1)} \right) = \\ &= -\frac{\mu^2(\mu-1)}{n! \omega_n} \left(\frac{b(x')}{h_0(x')} \cdot \nabla_{x'} h_0(x') - \frac{b(x')}{2h_0(x')} \cdot \nabla_{x'} h_0(x') \right) = -\frac{\mu^2(\mu-1)}{2 \cdot n! \omega_n} \frac{b(x')}{h_0(x')} \cdot \nabla_{x'} h_0(x') \end{aligned}$$

□

REFERENCES

- [1] H. Antil, T. Berry, J. Harlim, *Fractional diffusion maps*, Applied and Computational Harmonic Analysis 54 (2021), pp. 145–175.
- [2] J. A. Barceló, and A. Carbery, *On the magnitudes of compact sets in Euclidean spaces*, Amer. J. Math. 140 (2018), pp. 449–494.
- [3] R. Berman, S. Boucksom, D. Witt Nyström, *Fekete points and convergence towards equilibrium measures on complex manifolds*, Acta Math. 207 (2011), no. 1, pp. 1–27.
- [4] R. J. Berman, *Determinantal point processes and fermions on polarized complex manifolds: bulk universality*, Algebraic and analytic microlocal analysis, pp. 341–393, Springer Proc. Math. Stat., 269, Springer, Cham, 2018.
- [5] R. L. Bishop, *Decomposition of cut loci*, Proc. Amer. Math. Soc. 65 (1977), pp. 133–136.
- [6] E. Bunch, J. Kline, D. Dickinson, S. Bhat, G. Fung, *Weighting vectors for machine learning: numerical harmonic analysis applied to boundary detection*, arXiv:2106.00827.
- [7] E. Bunch, J. Kline, D. Dickinson, G. Fung, *Practical applications of metric space magnitude and weighting vectors*, arXiv:2006.14063.
- [8] S. Cho, *Quantaes, persistence, and magnitude homology*, arXiv:1910.02905.
- [9] G. I. Eskin, *Boundary value problems for elliptic pseudodifferential equations*, Translations of Mathematical Monographs, 52. American Mathematical Society, Providence, R.I., 1981.
- [10] R. L. Frank, E. H. Lieb, *A ‘liquid-solid’ phase transition in a simple model for swarming, based on the ‘no flat-spots’ theorem for subharmonic functions*, Indiana Univ. Math. J. 67 (2018), pp. 1547–1569.
- [11] H. Gimperlein, M. Goffeng, *On the magnitude function of domains in Euclidean space*, Amer. J. Math. 143 (2021), pp. 939–967.
- [12] H. Gimperlein, M. Goffeng, N. Louca, *The magnitude and spectral geometry*, preprint.
- [13] D. Govc, R. Hepworth, *Persistent magnitude*, J. Pure Appl. Algebra 225 (2021), no. 3, Paper No. 106517, 40 pp.
- [14] G. Grubb, *Spectral results for mixed problems and fractional elliptic operators*, J. Math. Anal. Appl. 421 (2015), pp. 1616–1634.
- [15] G. Grubb, *Fractional Laplacians on domains, a development of Hörmander’s theory of μ -transmission pseudodifferential operators*, Adv. Math. 268 (2015), pp. 478–528.
- [16] G. Grubb, *Local and nonlocal boundary conditions for μ -transmission and fractional elliptic pseudodifferential operators*, Anal. PDE 7 (2014), pp. 1649–1682.
- [17] G. Grubb, *Green’s formula and a Dirichlet-to-Neumann operator for fractional-order pseudodifferential operators*, Comm. Partial Differential Equations 43 (2018), pp. 750–789.
- [18] G. Grubb, *Integration by parts and Pohozaev identities for space-dependent fractional-order operators*, J. Differential Equations 261 (2016), pp. 1835–1879.
- [19] G. Grubb, *Functional Calculus of Pseudodifferential Boundary Problems*, Birkhäuser Boston, 1996.
- [20] S. Holman, G. Uhlmann, *On the microlocal analysis of the geodesic X-ray transform with conjugate points*, J. Differential Geom. 108 (2018), no. 3, pp. 459–494.
- [21] L. Hörmander, *Seminar notes on pseudo-differential operators and boundary problems*, Lectures at IAS Princeton 1965-66, available from Lund University, <https://lup.lub.lu.se/record/7c9d8485-d7d6-4d47-80a5-a5533685c4ca>.
- [22] L. Hörmander, *The analysis of linear partial differential operators. I. Distribution theory and Fourier analysis*, Reprint of the 1994 edition. Classics in Mathematics. Springer, Berlin, 2007.
- [23] L. Hörmander, *The analysis of linear partial differential operators. III. Pseudo-differential operators*, Reprint of the 1994 edition. Classics in Mathematics. Springer, Berlin, 2007.
- [24] T. Leinster, and C. A. Cobbold, *Measuring diversity: the importance of species similarity*, Ecology 93 (2012), pp. 477–489.

- [25] T. Leinster, *The magnitude of metric spaces*, Doc. Math. 18 (2013), pp. 857–905.
- [26] T. Leinster, M. Meckes, *The magnitude of a metric space: from category theory to geometric measure theory*, in: N. Gigli (ed.), *Measure theory in non-smooth spaces*, pp. 156–193, Partial Differ. Equ. Meas. Theory, De Gruyter Open, Warsaw, 2017.
- [27] T. Leinster, M. Shulman, *Magnitude homology of enriched categories and metric spaces*, Algebr. Geom. Topol. 21 (2021), pp. 2175–2221.
- [28] T. Leinster, S. Willerton, *On the asymptotic magnitude of subsets of Euclidean space*, Geom. Dedicata 164(1) (2013), pp. 287–310.
- [29] M. Lesch, *Operators of Fuchs type, conical singularities, and asymptotic methods*, Teubner-Texte zur Mathematik, 136. B. G. Teubner Verlagsgesellschaft mbH, Stuttgart, 1997. arXiv:dg-ga/9607005.
- [30] M. Lesch, *On the noncommutative residue for pseudodifferential operators with log-polyhomogeneous symbols*, Ann. Global Anal. Geom. 17, 2 (1999), pp. 151–187.
- [31] W. McLean, *Strongly elliptic systems and boundary integral equations*, Cambridge University Press, Cambridge, 2000.
- [32] M. Meckes, *Positive definite metric spaces*, Positivity 17 (2013), pp. 733–757.
- [33] M. Meckes, *Magnitude, diversity, capacities and dimensions of metric spaces*, Potential Anal. 42 (2015), pp. 549–572.
- [34] M. Meckes, *On the magnitude and intrinsic volumes of a convex body in Euclidean space*, Mathematika 66 (2020), pp. 343–355.
- [35] N. Otter, *Magnitude meets persistence. Homology theories for filtered simplicial sets*, arXiv:1807.01540.
- [36] S. G. Samko, A. A. Kilbas, O. I. Marichev, *Fractional Integrals and Derivatives: Theory and Applications*, Gordon and Breach Science Publishers, Amsterdam, 1993.
- [37] L. Schwartz, *Théorie des distributions*, Hermann, Paris, 1997.
- [38] M. Shubin, *Pseudodifferential operators and spectral theory*, Springer-Verlag, Berlin, second edition, 2001.
- [39] S. R. Simanca, *Pseudo-differential operators*, Pitman Research Notes in Mathematics Series, 236. Longman Scientific & Technical, Harlow; copublished in the United States with John Wiley & Sons, Inc., New York, 1990.
- [40] V. W. Summer, *Torsion in the Khovanov homology of links and the magnitude homology of graphs*, Ph.D. thesis, North Carolina State University, 2019, available at <https://repository.lib.ncsu.edu/handle/1840.20/36588>
- [41] H. Triebel, D. Yang, *Spectral theory of Riesz potentials on quasi-metric spaces*, Math. Nachr. 238 (2002), pp. 160–184.
- [42] F. W. Warner, *The conjugate locus of a Riemannian manifold*, Amer. J. Math. 87 (1965), pp. 575–604.
- [43] S. Willerton, *Heuristic and computer calculations for the magnitude of metric spaces*, arXiv:0910.5500.
- [44] S. Willerton, *On the magnitude of spheres, surfaces and other homogeneous spaces*, Geom. Dedicata 168 (2014), pp. 291–310.
- [45] S. Willerton, *The magnitude of odd balls via Hankel determinants of reverse Bessel polynomials*, Discrete Anal. 2020, Paper No. 5, 42 pp.
- [46] A. Zabrodin, P. Wiegmann, *Large- N expansion for the 2D Dyson gas*, Journal of Physics A 39 (2006), pp. 8933–8963.

HEIKO GIMPERLEIN, NIKOLETTA LOUCA
 MAXWELL INSTITUTE FOR MATHEMATICAL SCIENCES AND
 DEPARTMENT OF MATHEMATICS, HERIOT-WATT UNIVERSITY
 EDINBURGH EH14 4AS
 UNITED KINGDOM

MAGNUS GOFFENG,
 CENTRE FOR MATHEMATICAL SCIENCES
 UNIVERSITY OF LUND
 BOX 118, SE-221 00 LUND
 SWEDEN

Email address: h.gimperlein@hw.ac.uk, n124@hw.ac.uk, magnus.goffeng@math.lth.se