

On generally covariant mathematical formulation of Feynman integral in Lorentz signature

András László

Wigner Research Centre for Physics, Budapest

E-mail: laszlo.andras@wigner.hu

Abstract. It is widely accepted that the Feynman integral is one of the most promising methodologies for defining a generally covariant formulation of nonperturbative interacting quantum field theories (QFTs) without a fixed prearranged causal background. Recent literature suggests that if the spacetime metric is not fixed, e.g. because it is to be quantized along with the other fields, one may not be able to avoid considering the Feynman integral in the original Lorentz signature, without Wick rotation. Several mathematical phenomena are known, however, which are at some point showstoppers to a mathematically sound definition of Feynman integral in Lorentz signature. The Feynman integral formulation, however, is known to have a differential reformulation, called to be the master Dyson–Schwinger (MDS) equation for the field correlators. In this paper it is shown that a particular presentation of the MDS equation can be cast into a mathematically rigorously defined form: the involved function spaces and operators can be strictly defined and their properties can be established. Therefore, MDS equation can serve as a substitute for the Feynman integral, in a mathematically sound formulation of constructive QFT, in arbitrary signature, without a fixed background causal structure. It is also shown that even in such a generally covariant setting, there is a canonical way to define the Wilsonian regularization of the MDS equation. The main result of the paper is a necessary and sufficient condition for the regularized MDS solution space to be nonempty, for conformally invariant Lagrangians. This theorem also provides an iterative approximation algorithm for obtaining regularized MDS solutions, and is guaranteed to be convergent whenever the solution space is nonempty. The algorithm could eventually serve as a method for putting Lorentz signature QFTs onto lattice, in the original metric signature.

Keywords: Feynman integral, master Dyson–Schwinger equation, generally covariant, constructive field theory

1. Introduction

By now, a lot is known about the mathematically sound formulation of interacting quantum field theory (QFT), using perturbation theory [1]. However, still until now, there is no widely accepted concise mathematical formulation known for nonperturbative interacting QFT. Strictly speaking, as of now, it is only conjectured that eventually one could well-define an interacting QFT model in a nonperturbative manner, in a constructive way, e.g. as specified by a Lagrangian. A well known promising attempt for the nonperturbative approach is the algebraic quantum field theory (AQFT) [2, 3]. AQFT is known to capture several important qualitative aspects of the QFT formalism in physics, such as the spin-statistics theorem, but there are no known concrete AQFT constructions in the complexity of e.g. a 3+1 dimensional full quantum electrodynamics. Concrete AQFT models, as of now, are only known for free particles in arbitrary dimensions, or for simple systems, such as discrete Ising models in 1+1 dimensional and discrete spacetimes, or for particular simple systems in spacetime dimensions typically lower than 3+1. There are also recent advances of perturbative AQFT on causal sets, in which framework concrete interacting models are constructed by now [4], assuming a finite system of causal sets. Due to the difficulties of nonperturbative formulation, the perturbative rigorous formulation of constructive QFT (pQFT) was seriously considered by a number of authors [5, 6, 7, 8, 9]. In particular, [10] explicitly proves the renormalizability of Yang–Mills interactions over globally hyperbolic spacetimes. Moreover, a generally covariant framework was already developed [11]. However, it is generally thought that the only promising framework, which could be capable of formalizing *nonperturbative* interacting generally covariant QFT models in the continuum limit, is likely to be the Feynman integral formulation [12].

A lot is known about Feynman integrals [13, 14], but in Lorentz signature, without taking a Wick rotation, it seems to be still not a completely understood mathematical construction, although the modern literature seems to tighten the noose on the measure theoretically well defined Feynman integral [15, 16]. Other authors [17] argue, that Feynman integral should not be, strictly speaking, understood in the measure theoretical sense, i.e. in the sense of infinitesimal summation, but in a more generalized sense. That kind of picture is indeed supported by the fact that e.g. for a fermionic system, the Feynman integral is defined as a Berezin integral, which indeed has little link with integration in terms of infinitesimal summation. To complicate the picture, recent literature suggests [10, 18, 19, 20] that in order to apply Feynman integral formalism to a generally covariant setting, in which case the a background spacetime metric is not fixed, the applicability of the usual Wick rotation from Lorentzian to Euclidean signature can be problematic.

The above issues with the Lorentz signature Feynman integral formulation can be circumvented via using the well known differential reformulation of Feynman integral formalism, called to be the master Dyson–Schwinger (MDS) equations for the field correlators (see e.g. [21] for a didactic review). From the usually presented form of the

MDS equation in the QFT literature, it is not immediately evident that the function spaces and operators defining the MDS equation are well defined. In this paper, however, it is argued that they are, and as such, the MDS equation can be used to substitute the Feynman integral for a mathematically sound definition of constructive nonperturbative generally covariant QFT. It will be also argued, that the Wilsonian regularized version of the MDS equation can also be canonically defined in a generally covariant setting. The main result of the paper is a theorem about a necessary and sufficient condition for the regularized MDS equation to have nonempty solution space, for theories with classically conformally invariant Lagrangians. The pertinent theorem is constructive in the sense that it provides a (probably slowly converging) iterative algorithm for approximating MDS solutions, which is guaranteed to be convergent whenever the solution space is nonempty. This method can eventually be also employed for doing lattice QFT-like calculations in arbitrary signatures, in particular, in the original Lorentz signature.

The structure of the paper is as follows. In Section 2 the heuristic form of the MDS equation is recalled, as derived from the heuristic Feynman integral formulation in usual QFT. The rest of the paper intends to keep mathematical rigor. In Section 3 the mathematically strict definition of a classical field theory is recalled, in order to pin down the precise topology of the involved function spaces and the continuity properties of a typical Euler–Lagrange functional, which is key in the construction. In Section 4 the function spaces and operators needed to define the (unregularized) MDS operator are presented. In Section 5, these are generalized in the distributional sense, and the Wilsonian regularized version of the MDS equation is invoked and justified. Section 6 is dedicated for the main theorem of the paper about a necessary and sufficient condition for the existence of solutions for the regularized MDS equation, for conformally invariant Lagrangians. The paper heavily relies on the theory of non-normable topological vector spaces (TVS), and therefore an entire Appendix is dedicated to a recollection of important and sometimes counterintuitive theorems on these.

2. Feynman integral and the heuristic form of the MDS equation

We briefly recall the justification of the MDS equation in the Feynman integral formulation of QFT. Let F denote the space of all, i.e. off-shell classical field configurations. As expanded in Section 3, in most models it is safe to assume that F is a topological affine space, carrying a nuclear Fréchet topology. Its underlying vector space \mathbb{F} is the topological vector space of field variations, and \mathbb{F}^* will denote its topological dual understood with the standard strong dual topology. In the Feynman integral formulation of QFT, then it is postulated that the evaluation method for Feynman type, i.e. causally ordered quantum vacuum expectation value of observables in a (not necessarily unique) vacuum state ρ is the following. Given a fixed reference field $\psi_0 \in F$ and test functionals $J_1, \dots, J_n \in \mathbb{F}^*$, the causally ordered quantum vacuum expectation value of the real valued homogeneous multipolynomial functional $(J_1 | \cdot - \psi_0) \cdots (J_n | \cdot - \psi_0) : F \rightarrow \mathbb{R}$ is

declared to be

$$\int_{\psi \in F} (J_1 | \psi - \psi_0) \cdots (J_n | \psi - \psi_0) e^{i\frac{1}{\hbar} S(\psi)} d\rho(\psi) \Big/ \int_{\psi \in F} e^{i\frac{1}{\hbar} S(\psi)} d\rho(\psi), \quad (1)$$

where the symbol $d\rho(\cdot)$ denotes the hypothetical Feynman measure corresponding to a vacuum state ρ , $(\cdot | \cdot)$ denotes the duality pairing form between \mathbb{F}^* and \mathbb{F} , whereas $S : F \rightarrow \mathbb{R}$ is the action functional of the underlying classical field theory. In the heuristic calculations, $d\rho(\cdot)$ is handled as if it were a Lebesgue measure on F , and as if $e^{i\frac{1}{\hbar} S(\cdot)} d\rho(\cdot)$ were a finite measure, having finite moments and analytic Fourier transform. A sign change $\hbar \mapsto -\hbar$ would correspond to a reversal in the causal ordering, if there were any *a priori* causal structure over the spacetime manifold (which in fact, is not needed to be assumed at this point). The hypothetical *partition function* condenses all these information about the state ρ , and would be a mapping

$$Z_{\hbar, \psi_0} : \mathbb{F}^* \rightarrow \mathbb{C}, \quad J \mapsto Z_{\hbar, \psi_0}(J) := \int_{\psi \in F} e^{i(J | \psi - \psi_0)} e^{i\frac{1}{\hbar} S(\psi)} d\rho(\psi), \quad (2)$$

i.e. the formal Fourier transform of the hypothetical measure $e^{i\frac{1}{\hbar} S(\cdot)} d\rho(\cdot)$. The collection of *n-field correlators*

$$G_{\hbar, \psi_0}^{(n)} := \left((-i)^n \frac{1}{Z_{\hbar, \psi_0}(J)} D^{(n)} Z_{\hbar, \psi_0}(J) \right) \Big|_{J=0} \quad (3)$$

is an other means to rephrase these information about the state ρ , and also can be used to evaluate the quantum expectation values Eq.(1) by simple duality pairing, like $(J_1 \otimes \dots \otimes J_n | G_{\hbar, \psi_0}^{(n)})$. Here $D^{(n)} Z_{\hbar, \psi_0}$ is assumed to behave like the n -th Fréchet derivative of the partition function $J \mapsto Z_{\hbar, \psi_0}(J)$, implicitly assuming that Z_{\hbar, ψ_0} is n -times continuously Fréchet differentiable (and for fermion fields, this differentiation is assumed to be a graded differentiation). Since the partition function would be a map $Z_{\hbar, \psi_0} : \mathbb{F}^* \rightarrow \mathbb{C}$, the collection of field correlators $G_{\hbar, \psi_0} := (G_{\hbar, \psi_0}^{(0)}, G_{\hbar, \psi_0}^{(1)}, \dots, G_{\hbar, \psi_0}^{(n)}, \dots)$ would sit in $\mathcal{T}(\mathbb{F}) := \bigoplus_{n \in \mathbb{N}_0} \otimes^n \mathbb{F}$, i.e. in the tensor algebra of \mathbb{F} .

Let $E(\psi) := D_F S(\psi)$ denote the Euler–Lagrange functional, i.e. the derivative of the action functional S , evaluated at the classical field configuration $\psi \in F$. It would be a map $E : F \times \mathbb{F} \rightarrow \mathbb{R}$, $(\psi, \delta\psi) \mapsto (E(\psi) | \delta\psi) := (D_F S(\psi) | \delta\psi)$, being linear in its second variable, since it is a derivative. In the usual QFT protocol it is assumed that the EL functional E is multipolynomial, and thus so is the real valued map $\psi \mapsto (E(\psi) | \delta\psi)$ for any fixed field variation $\delta\psi \in \mathbb{F}$. Let $\mathbf{E}((-i)D_{\mathbb{F}^*} + \psi_0)$ be the multipolynomial differential operator defined by the polynomial coefficients of the Euler-Lagrange functional E . Applying the usual rules of formal Fourier transform, a function $Z : \mathbb{F}^* \rightarrow \mathbb{C}$ is of the form Eq.(2), up to a complex multiplier, if and only if it satisfies the master Dyson–Schwinger (MDS) equation

$$\left(\mathbf{E}((-i)D_{\mathbb{F}^*} + \psi_0) Z \right) \Big|_J = -\hbar J Z(J) \quad (\forall J \in \mathbb{F}^*), \quad (4)$$

see e.g. [21] for a didactic derivation. The operational meaning of this usual presentation of the MDS equation might not seem immediately evident. However, expressing Z_{\hbar, ψ_0}

via its formal Taylor series, encoded by the collection of field correlators $G_{\hbar, \psi_0} \in \mathcal{T}(\mathbb{F})$, the MDS equation Eq.(4) is seen to be equivalent to

$$\text{we search for } G \in \mathcal{T}(\mathbb{F}) \text{ such that :} \\ G^{(0)} = 1 \quad \text{and} \quad \mathcal{L}_{(\mathbf{E}_{\psi_0}|\delta\psi)} G = i \hbar L_{\delta\psi} G \quad (\forall \delta\psi \in \mathbb{F}). \quad (5)$$

Here, the symbol $L_{\delta\psi}$ denotes the left-multiplication operator in the tensor algebra $\mathcal{T}(\mathbb{F})$ by the one-vector $\delta\psi \in \mathbb{F}$. The symbol \mathcal{L}_p denotes the left-insertion operator by some element p from the topological dual space of $\mathcal{T}(\mathbb{F})$. The map $E_{\psi_0} : \mathbb{F} \rightarrow \mathbb{F}^*$ is defined via $E_{\psi_0} := E \circ (\mathbf{I}_{\mathbb{F}} + \psi_0)$ from the original Euler–Lagrange functional $E : F \rightarrow \mathbb{F}^*$, i.e. it is the Euler–Lagrange functional with respect to a fixed reference field $\psi_0 \in F$, expressed on the space of field variations \mathbb{F} . Since it was assumed to be multipolynomial, it can eventually be regarded as a linear map $E_{\psi_0} : \mathcal{T}(\mathbb{F}) \rightarrow \mathbb{F}^*$. As such, it may be identified with an element $\mathbf{E}_{\psi_0} \in (\mathcal{T}(\mathbb{F}))^* \otimes \mathbb{F}^*$, and correspondingly $(E_{\psi_0}|\delta\psi)$ with $(\mathbf{E}_{\psi_0}|\delta\psi) \in (\mathcal{T}(\mathbb{F}))^* \otimes \mathbb{F}^*$ ($\forall \delta\psi \in \mathbb{F}$), which then has the corresponding left-insertion operator $\mathcal{L}_{(\mathbf{E}_{\psi_0}|\delta\psi)}$ acting over $\mathcal{T}(\mathbb{F})$. The spaces and operators involved in Eq.(5) would be perfectly meaningful if the space of fields F were finite dimensional, and could be used as a substitute for Feynman integral formulation Eq.(1), regardless of e.g. a metric signature or other auxiliary information on the details of the underlying classical theory described by the Euler–Lagrange functional E . In Section 3 and 4 it shall be shown that the pertinent objects can be made well-defined even when F is indeed the infinite dimensional space of off-shell field configurations in a realistic field theory. The Eq.(5) presentation of the MDS equation does not seem to be described in the literature.

In QFT, is also necessary to consider the Wilsonian regularized version of the Feynman integral. The Wilsonian regularization is understood as performing the Feynman integral Eq.(1) on a subspace of off-shell fields with their high frequency modes suppressed. In a generally covariant setting the meaning of this might not seem immediately evident, but Wilsonian regularized Feynman integral can be postulated as

$$\int_{\delta\psi \in \mathcal{R}[\mathbb{F}]} O(\delta\psi + \psi_0) \, d\mathcal{R}_* \mu_{\psi_0}(\delta\psi) \bigg/ \int_{\delta\psi \in \mathcal{R}[\mathbb{F}]} 1 \, d\mathcal{R}_* \mu_{\psi_0}(\delta\psi) \quad (6)$$

for multipolynomial functionals $O : F \rightarrow \mathbb{R}$, where $\mathcal{R} : \mathbb{F} \rightarrow \mathbb{F}$ is some continuous linear operator, $\mathcal{R}[\mathbb{F}] \subset \mathbb{F}$ denotes the image of \mathbb{F} by \mathcal{R} , the symbol μ_{ψ_0} stands for the pushforward of the hypothetical finite measure $e^{i\frac{1}{\hbar}S(\cdot)} d\rho(\cdot)$ on F via the map $F \rightarrow \mathbb{F}$, $\psi \mapsto (\psi - \psi_0)$, and $\mathcal{R}_* \mu_{\psi_0}$ stands for the pushforward of the measure μ_{ψ_0} on \mathbb{F} to $\mathcal{R}[\mathbb{F}]$ by \mathcal{R} . The map \mathcal{R} can be called a *regulator*, it is chosen to be the convolution operator by some test function in case of theories over an affine spacetime (can be generalized for arbitrary spacetimes as well), and Eq.(6) means nothing but the natural pushforward Feynman integration on the subspace $\mathcal{R}[\mathbb{F}] \subset \mathbb{F}$, given that the original Feynman integration on F was meaningful. The map \mathcal{R} implements the high frequency damping. Using the fundamental formula of integral substitution, one infers that the Wilsonian regularized MDS equation on the field correlators reads

$$G^{(0)} = 1 \quad \text{and} \quad \mathcal{L}_{(\mathbf{E}_{\psi_0}|\delta\psi)} G = i \hbar L_{\mathcal{R} \delta\psi} G \quad (\forall \delta\psi \in \mathbb{F}) \quad (7)$$

in the analogy of Eq.(5), where again $L_{\mathcal{R}\delta\psi}$ is the left-multiplication in $\mathcal{T}(\mathbb{F})$ by the one-vector $\mathcal{R}\delta\psi \in \mathbb{F}$. As shall be expanded in Section 5, the pertinent objects can be made well-defined similarly to that of the unregularized MDS equation. The Wilsonian regularized MDS equation Eq.(7) does not seem to be described in the literature.

From this point on, we drop the heuristic arguments, and all the statements and formulas are intended to be mathematically rigorous. The aim of this paper is to show that the objects involved in Eq.(5) and Eq.(7) are mathematically well defined, and to establish the fundamental properties of the solution spaces of the pertinent equations.

3. Continuity properties of the Euler–Lagrange functional

Our presentation of the master Dyson–Schwinger (MDS) operator heavily relies on the precise definition of the Euler–Lagrange functional of a classical field theory. In order to pin down the topological properties of the involved spaces and precise continuity property of their operators, we need to briefly recall the standard variational formulation of a classical field theory. For the sake of simplified treatment, we will use neither the jet formalism, nor the theory of general connections over fiber bundles [22, 23]. We will rather concentrate on the TVS theory side (Appendix A), i.e. we keep the differential geometric treatment to a reasonable appropriate minimum. Let \mathcal{M} denote throughout the paper a finite dimensional real smooth orientable and oriented manifold, and let $m := \dim(\mathcal{M})$. It may be compact or noncompact, and may be with or without boundary (if with boundary, we assume the cone condition for it, so that locally the Sobolev and Maurin embedding theorems hold, see also Appendix A.9). The manifold \mathcal{M} is meant to model the spacetime manifold, or eventually, the compact manifold with boundary underlying the conformal compactification (Penrose diagram) of a spacetime. Let $V(\mathcal{M})$ be some real vector bundle over \mathcal{M} with finite dimensional fibers. Denote, as usual, by $\Gamma(\cdot)$ the space of smooth sections. In particular, the space $\Gamma(V(\mathcal{M}))$ denotes the real vector space of smooth sections of $V(\mathcal{M})$ (these are meant to model the matter fields). The covariant derivation operators over $\Gamma(V(\mathcal{M}))$ (which are meant to model the mediator fields) form an affine space with subordinate vector space $\Gamma(T^*(\mathcal{M}) \otimes V(\mathcal{M}) \otimes V^*(\mathcal{M}))$, as it is common knowledge. More particularly, covariant derivation operators can be considered as sections of an affine bundle over \mathcal{M} , which we will denote by $DV(\mathcal{M})$, and then a covariant derivation is an element of $\Gamma(DV(\mathcal{M}))$. The bundle $DV(\mathcal{M})$ is an affine bundle with subordinate vector bundle $T^*(\mathcal{M}) \otimes V(\mathcal{M}) \otimes V^*(\mathcal{M})$, the sections of which are the difference tensors (or gauge potentials, in field theory). As usual, the symbols \otimes shall denote pointwise tensor product of vector bundles over the same base, whereas \oplus will denote pointwise direct sum of vector bundles over the same base. The fibered product (pointwise cartesian product, or Whitney sum) of two generic fiber bundle over the same base shall be denoted by $V(\mathcal{M}) \times_w U(\mathcal{M})$, and their elements by $(v, u)_w$ in order to indicate that these are pairs of fields, over the same base points. The subscript $_w$ is used in order to distinguish the above from sections (v, u) of the cartesian product bundle $V(\mathcal{M}) \times U(\mathcal{M}')$, which would

On generally covariant mathematical formulation of Feynman integral in Lorentz signature 7

be a bundle over the product manifold $\mathcal{M} \times \mathcal{M}'$ (eventually $\mathcal{M}' = \mathcal{M}$).

Definition 1 Let \mathcal{M} , $V(\mathcal{M})$ and $DV(\mathcal{M})$ as above.

A **Lagrange form** is a base point preserving, smooth fiber bundle homomorphism

$$L : V(\mathcal{M}) \oplus T^*(\mathcal{M}) \otimes V(\mathcal{M}) \oplus T^*(\mathcal{M}) \wedge T^*(\mathcal{M}) \otimes V(\mathcal{M}) \otimes V^*(\mathcal{M}) \longrightarrow \wedge^m T^*(\mathcal{M}).$$

By construction, a Lagrange form takes some sections

$$v \in \Gamma(V(\mathcal{M})), \quad Dv \in \Gamma(T^*(\mathcal{M}) \otimes V(\mathcal{M})), \quad P \in \Gamma(T^*(\mathcal{M}) \wedge T^*(\mathcal{M}) \otimes V(\mathcal{M}) \otimes V^*(\mathcal{M}))$$

into a maximal form field $L(v, Dv, P) \in \Gamma(\wedge^m T^*(\mathcal{M}))$.

An element $(v, \nabla)_w \in \Gamma(V(\mathcal{M}) \times_w DV(\mathcal{M}))$ is called a **field configuration**. The field configurations form an affine space over the real vector space $\Gamma(V(\mathcal{M}) \oplus T^*(\mathcal{M}) \otimes V(\mathcal{M}) \otimes V^*(\mathcal{M}))$. An element $(\delta v, \delta C)_w$ from that space is called a **field variation**.

The map

$$\Gamma(V(\mathcal{M}) \times_w DV(\mathcal{M})) \longrightarrow \Gamma(\wedge^m T^*(\mathcal{M})), \quad (v, \nabla)_w \longmapsto L(v, \nabla v, P(\nabla))$$

is called the **Lagrangian expression**, where ∇v is the covariant derivative of the section v , and $P(\nabla)$ is the curvature tensor of ∇ . (Note that the expression $(v, \nabla v, P(\nabla))_w$ encodes the same information as the first jet of a field configuration $(v, \nabla)_w$, but we do not intend to use the jet formalism in the present paper.)

Given a Lagrange form L , its **action functional** is the real Radon measure valued map

$$S^L : \Gamma(V(\mathcal{M}) \times_w DV(\mathcal{M})) \longrightarrow \text{Rad}(\mathcal{M}, \mathbb{R}), \quad (v, \nabla)_w \longmapsto S^L(v, \nabla)_w$$

where on compact subsets $\mathcal{K} \subset \mathcal{M}$ the definition is $S^L_{\mathcal{K}}(v, \nabla)_w := \int_{\mathcal{K}} L(v, \nabla v, P(\nabla))$, i.e. the action functional is the Radon measure defined by local integrals of the Lagrangian expression, as usual.

We use the shorthand notation $F := \Gamma(V(\mathcal{M}) \times_w DV(\mathcal{M}))$ for the space of field configurations, moreover $\mathbb{F} := \Gamma(V(\mathcal{M}) \oplus T^*(\mathcal{M}) \otimes V(\mathcal{M}) \otimes V^*(\mathcal{M}))$ for the space of field variations. The space F is an affine space over the real vector space \mathbb{F} . The real vector space \mathbb{F} may be naturally endowed with the standard \mathcal{E} smooth function topology. (The \mathcal{E} topology is defined by the family of arbitrary order Sobolev norms of over compact patches of \mathcal{M} .) With this topology \mathbb{F} and thus F become Hausdorff locally convex topological vector and affine spaces, respectively. It is also common knowledge (Remark 58), that \mathbb{F} with the \mathcal{E} topology becomes a *nuclear Fréchet space*, which fact will be an important detail in the QFT construction.

The real vector space of real valued Radon measures $\text{Rad}(\mathcal{M}, \mathbb{R})$ can be also naturally endowed with a topology, defined by compact setwise total variations as family of seminorms, or equivalently, by the convergence of measure sequences over compact sets. With this, $\text{Rad}(\mathcal{M}, \mathbb{R})$ becomes a Hausdorff locally convex topological vector space.

Remark 2 It is not true in general that the continuity of a map between topological spaces is equivalent to its sequential continuity. It is common knowledge, however, that metrizable topological spaces are sequential (Remark 61), i.e. their topology is completely characterized by the convergence of sequences. Since \mathbb{F} is Fréchet space, by construction its topology is metrizable in a translationally invariant way, and therefore also is the topology of F . In particular, a map $S : F \rightarrow Y$ to any topological space Y is continuous if and only if S is sequentially continuous, i.e. it maps convergent sequences in F to convergent sequences in Y .

Remark 3 The following can be observed.

- (i) The action functional was defined to be a Radon measure valued map. That was motivated by the fact that no asymptotics was prescribed on the field configurations F , nor it was assumed that \mathcal{M} is compact. Because of that, one cannot guarantee that the smooth maximal form field $L(v, \nabla v, P(\nabla))$ is integrable throughout the full \mathcal{M} for sufficiently many field configurations $(v, \nabla)_w \in F$. It is, however, always locally integrable, hence the action functional as a Radon measure valued map is meaningful, and everywhere defined.
- (ii) Due to Lebesgue's theorem of dominated convergence, the action functional is sequentially continuous, and therefore by means of Remark 2, it is continuous.

The action functional is everywhere differentiable in the Fréchet–Hadamard sense (see also Appendix Appendix A.1), as it is common knowledge in Lagrangian field theory. In order to show its explicit form, we recall some differential geometric identities. We will use Penrose abstract indices for the tangent tensors throughout the section.

Remark 4 If ∇ is a covariant derivation over $T(\mathcal{M})$, then there is a unique covariant derivation $\tilde{\nabla}$ over $T(\mathcal{M})$ associated to it, having vanishing torsion tensor and having the same affine parametrized geodesics as ∇ . The covariant derivation $\tilde{\nabla}$ is called the torsion-free part of ∇ . In explicit formulae: whenever v^b is a smooth section of $T(\mathcal{M})$, then one has $\tilde{\nabla}_a v^b = \nabla_a v^b + \frac{1}{2} T(\nabla)_{ac}^b v^c$, where $T(\nabla)_{ac}^b$ denotes the torsion tensor of ∇ .

Theorem 5 The action functional S^L is everywhere differentiable, and its derivative at some fixed $(v, \nabla)_w \in F$ is a continuous linear map $DS^L(v, \nabla)_w : \mathbb{F} \rightarrow \text{Rad}(\mathcal{M}, \mathbb{R})$, given by the formula

$$\begin{aligned}
 (\delta v, \delta C)_w &\mapsto (DS^L_{\mathcal{K}}(v, \nabla)_w | (\delta v, \delta C)_w) = \\
 &\int_{\mathcal{K}} \left(D_1 L(v, \nabla v, P(\nabla)) \delta v + D_2^a L(v, \nabla v, P(\nabla)) (\nabla_a \delta v + \delta C_a v) + 2 D_3^{[ab]} L(v, \nabla v, P(\nabla)) \tilde{\nabla}_{[a} \delta C_{b]} \right),
 \end{aligned} \tag{8}$$

when evaluated on some compact subset $\mathcal{K} \subset \mathcal{M}$. Here, $D_1 L$, $D_2 L$, $D_3 L$ denote the spacetime pointwise partial derivative of L against its first, second and third field variable, respectively. It also follows that the derivative map $DS^L : F \times \mathbb{F} \rightarrow \text{Rad}(\mathcal{M}, \mathbb{R})$ is jointly continuous in its two variables. \square

Proof This is a simple consequence of the below elementary facts.

- The Lagrange form evaluation as a map $(v, Dv, P)_{\mathcal{W}} \mapsto L(v, Dv, P)$ acting on the space of sections is continuously differentiable in the \mathcal{E} topology, and the map $(v, \nabla)_{\mathcal{W}} \mapsto (v, \nabla v, P(\nabla))_{\mathcal{W}}$ is also continuously differentiable in the \mathcal{E} topology. Therefore, their composition, being the Lagrangian expression $(v, \nabla)_{\mathcal{W}} \mapsto L(v, \nabla v, P(\nabla))$, is also differentiable in the \mathcal{E} topology, and its derivative is given by the integrand of Eq.(8).
- The local integral evaluation of a smooth maximal form over a compact subset $\mathcal{K} \subset \mathcal{M}$ is sequentially continuous map in the \mathcal{E} topology due to Lebesgue theorem of dominated convergence, and therefore by means of Remark 2 it is continuous in the $\mathcal{E} \rightarrow \text{Rad}(\mathcal{M}, \mathbb{R})$ topologies. Due to its linearity then it is differentiable, and its derivative is itself.
- Chain rule for the differentiation of composite functions made out of the above two maps implies the first part of the theorem.
- Lebesgue's theorem of dominated convergence implies joint sequential continuity of DS^L . Therefore, by means of Remark 2, the derivative functional is jointly continuous as a $DS^L : F \times \mathbb{F} \rightarrow \text{Rad}(\mathcal{M}, \mathbb{R})$ map, since F, \mathbb{F} and thus their product $F \times \mathbb{F}$ is metrizable. This proves the second statement of the theorem. \blacksquare

Remark 6 Let us also recall the following differential geometric identities.

- Let $J_{[c_1 \dots c_m]}^a$ be a smooth section of $T(\mathcal{M}) \otimes \wedge^m T^*(\mathcal{M})$, i.e. a maximal form valued tangent vector field (the symbol $[\]$ denotes index antisymmetrization). Then, given any covariant derivation ∇ on $T(\mathcal{M})$, one has that the expression $\tilde{\nabla}_a J_{[c_1 \dots c_m]}^a$ is independent of the choice of the covariant derivation ∇ , where $\tilde{\nabla}$ denotes the torsion-free part of ∇ . That is, the divergence of a maximal form valued vector field is naturally defined without further assumptions. Similarly, for a smooth section $K_{[c_1 \dots c_m]}^{[ab]}$ of $(T(\mathcal{M}) \wedge T(\mathcal{M})) \otimes \wedge^m T^*(\mathcal{M})$ one has that $\tilde{\nabla}_a K_{[c_1 \dots c_m]}^{[ab]}$ is independent of the choice of the covariant derivation ∇ , and thus the divergence of such field is naturally defined without further assumptions.
- Let $J_{[c_1 \dots c_m]}^a$ be a smooth section of $T(\mathcal{M}) \otimes \wedge^m T^*(\mathcal{M})$, i.e. a maximal form valued tangent vector field. Then, given any covariant derivation ∇ on $T(\mathcal{M})$, one has that $\tilde{\nabla}_a J_{[c_1 \dots c_m]}^a = m d_{[c_1} (J_{a c_2 \dots c_m]}^a)$, where d denotes exterior differentiation (see [24]).

Theorem 7 The derivative $DS^L(v, \nabla)_{\mathcal{W}}$ of the action functional S^L at a fixed $(v, \nabla)_{\mathcal{W}} \in F$ can be re-expressed as

$$(\delta v, \delta C)_{\mathcal{W}} \mapsto (DS_{\mathcal{K}}^L(v, \nabla)_{\mathcal{W}} | (\delta v, \delta C)_{\mathcal{W}}) = \int_{\mathcal{K}} \left(D_1 L(v, \nabla v, P(\nabla))_{[c_1 \dots c_m]} \delta v - (\tilde{\nabla}_a D_2^a L(v, \nabla v, P(\nabla))_{[c_1 \dots c_m]} \delta v \right) +$$

$$\begin{aligned}
 & \left(D_2^a L(v, \nabla v, P(\nabla))_{[c_1 \dots c_m]} \delta C_a v - 2 \left(\tilde{\nabla}_a D_3^{[ab]} L(v, \nabla v, P(\nabla))_{[c_1 \dots c_m]} \right) \delta C_b \right) \\
 & + m \int_{\partial \mathcal{K}} \left(D_2^a L(v, \nabla v, P(\nabla))_{[ac_1 \dots c_{m-1}]} \delta v + 2 D_3^{[ab]} L(v, \nabla v, P(\nabla))_{[ac_1 \dots c_{m-1}]} \delta C_b \right), \quad (9)
 \end{aligned}$$

when evaluated over some compact subset $\mathcal{K} \subset \mathcal{M}$ with cone property boundary $\partial \mathcal{K}$. \square

Proof This can be proved as usual in Lagrangian field theory. Namely, we start out from the expression in Eq.(8), use Leibniz rule, apply the differential geometric identities of Remark 6, and then apply Stokes theorem for the boundary term. \blacksquare

Let us introduce $\mathbb{F}_T \subset \mathbb{F}$ to be either the vector space of compactly supported sections from \mathbb{F} , or if $\partial \mathcal{M} \neq \emptyset$ optionally they may be even required to vanish on $\partial \mathcal{M}$ together with all of their derivatives. Elements of \mathbb{F}_T will be called the **test field variations**. The space \mathbb{F}_T can be endowed with the standard \mathcal{D} test function topology, being stronger than the \mathcal{E} topology, defined by the restricted \mathcal{E} topology for sections with their supports within each fixed compact set of \mathcal{M} . It is common knowledge (Remark 58), that \mathbb{F}_T with its natural \mathcal{D} test function topology is a *strict inductive limit of a countable system of nuclear Fréchet spaces with closed adjacent images (LNF space)* whenever \mathcal{M} is noncompact, and it is *nuclear Fréchet (NF space)* if \mathcal{M} is compact. These are important details in the QFT construction. It is seen that due to Lebesgue's theorem of dominated convergence the integrand within the expression $(DS_{\mathcal{M}}^L(v, \nabla)_{\mathcal{W}} | (\delta v_T, \delta C_T)_{\mathcal{W}})$, see again Eq.(8) and Eq.(9), is absolutely integrable for all fields $(v, \nabla)_{\mathcal{W}} \in F$ and all test field variations $(\delta v_T, \delta C_T)_{\mathcal{W}} \in \mathbb{F}_T$. In other words: the measure $\mathcal{K} \mapsto (DS_{\mathcal{K}}^L(v, \nabla)_{\mathcal{W}} | (\delta v_T, \delta C_T)_{\mathcal{W}})$ has bounded total variation, and thus $(DS_{\mathcal{M}}^L(v, \nabla)_{\mathcal{W}} | (\delta v_T, \delta C_T)_{\mathcal{W}}) \in \mathbb{R}$ is finite. Consequently, the following definition is meaningful.

Definition 8 Let \mathcal{M} , $V(\mathcal{M})$, L , S^L as before. The map

$$E^L : F \times \mathbb{F}_T \rightarrow \mathbb{R}, \quad (\psi, \delta \psi_T) \mapsto (E^L(\psi) | \delta \psi_T) := (DS_{\mathcal{M}}^L(\psi) | \delta \psi_T) \quad (10)$$

is called the **Euler–Lagrange (EL) functional**. (Here, we used a shorthand notation $\psi := (v, \nabla)_{\mathcal{W}} \in F$ for a field, and $\delta \psi_T := (\delta v_T, \delta C_T)_{\mathcal{W}} \in \mathbb{F}_T$ for a test field variation.)

Note that it was possible to define the EL functional as real valued at the price of restricting its second argument to compactly supported field variations. This setting also explains why one can automatically discard the EL boundary terms in classical variational problems over noncompact manifolds without boundary. It is clear that for all $\psi \in F$ the map $(E^L(\psi) | \cdot) : \mathbb{F}_T \rightarrow \mathbb{R}$ is well defined. Moreover, it is linear, and continuous in the \mathcal{D} topology due to Lebesgue's theorem of dominated convergence. Therefore the EL functional may be viewed either as map $E^L : F \times \mathbb{F}_T \rightarrow \mathbb{R}$, or alternatively as a distribution valued map $E^L : F \rightarrow \mathbb{F}_T^*$, where $*$ denotes the strong dual. About their continuity properties, one can state the following.

Theorem 9 The EL functional $E^L : F \times \mathbb{F}_T \rightarrow \mathbb{R}$, with F and \mathbb{F}_T carrying the standard \mathcal{E} and \mathcal{D} topologies, respectively, has the following continuity properties.

- (i) It is jointly sequentially continuous.
- (ii) It is separately continuous in each variable.
- (iii) It is continuous as a $E^L : F \rightarrow \mathbb{F}_T^*$ map. □

Proof Property (i) is obviously seen via applying Lebesgue theorem of dominated convergence in the joint variables.

To see (ii), take first a fixed $\delta\psi_T \in \mathbb{F}_T$. Then, the map $E^L(\cdot, \delta\psi_T) : F \rightarrow \mathbb{R}$ is sequentially continuous by means of (i), and due to the metrizable of F , by means of Remark 2, then it is continuous. Take than a fixed $\psi \in F$. The linear map $E^L(\psi, \cdot) : \mathbb{F}_T \rightarrow \mathbb{R}$ is sequentially continuous by means of (i). Due to the facts in Remark 61, the space \mathbb{F}_T carries the bornological property, by means of which the sequentially continuous linear map $E^L(\psi, \cdot) : \mathbb{F}_T \rightarrow \mathbb{R}$ is continuous.

To see (iii), observe that due to (i) the map $E^L : F \rightarrow \mathbb{F}_T^*$ is sequentially continuous, whenever \mathbb{F}_T^* is endowed with the weak (pointwise) topology. Due to the facts in Remark 61, the space \mathbb{F}_T^* carries the Montel property, therefore weakly convergent sequences are also strongly convergent in \mathbb{F}_T^* . Thus, the pertinent map is also sequentially continuous when the target space \mathbb{F}_T^* is endowed with its standard strong dual topology (\mathcal{D}^* topology). Due to the metrizable of F , by means of Remark 2, then it is $\mathcal{E} \rightarrow \mathcal{D}^*$ continuous. ■

Definition 10 A tuple $(\mathcal{M}, V(\mathcal{M}), F, \mathbb{F}, \mathbb{F}_T, E, \mathcal{C})$ is called a **classical field theory**, where \mathcal{M} and $V(\mathcal{M})$ is as in Definition 1, F is the space of smooth sections of the affine bundle $V(\mathcal{M}) \times_w DV(\mathcal{M})$, the space \mathbb{F} consists of the smooth sections of the underlying vector bundle $V(\mathcal{M}) \oplus T^*(\mathcal{M}) \otimes V(\mathcal{M}) \otimes V^*(\mathcal{M})$, the space \mathbb{F}_T consists of compactly supported sections from \mathbb{F} (if $\partial\mathcal{M} \neq \emptyset$, optionally, elements of \mathbb{F}_T may be required to vanish on $\partial\mathcal{M}$ together with all of their derivatives — variation with boundary included/excluded). Furthermore, the object E is a map $F \times \mathbb{F}_T \rightarrow \mathbb{R}$, such that there exists a Lagrange form L as in Definition 1, such that $E = E^L$. Finally, $\mathcal{C} := \{\psi \in F \mid \forall \delta\psi_T \in \mathbb{F}_T : (E(\psi) | \delta\psi_T) = 0\}$. The set \mathcal{C} is called the **solution space** of the classical field theory.

Definition 11 Let $(\mathcal{M}', V'(\mathcal{M}'), F', \mathbb{F}', \mathbb{F}'_T, E', \mathcal{C}')$ and $(\mathcal{M}, V(\mathcal{M}), F, \mathbb{F}, \mathbb{F}_T, E, \mathcal{C})$ be two classical field theories. These are called **isomorphic**, if and only if there exists a vector bundle isomorphism $V'(\mathcal{M}') \rightarrow V(\mathcal{M})$ with an underlying diffeomorphism $\mathcal{M}' \rightarrow \mathcal{M}$ of the base manifold, such that L subordinate to E is pulled back to L' subordinate to E' . (Isomorphic classical field theories are postulated to describe the same physics.) Quite naturally, isomorphisms of a classical field theory with itself are called **automorphisms**, or **symmetries**.

A classical field theory $(\mathcal{M}, V(\mathcal{M}), F, \mathbb{F}, \mathbb{F}_T, E, \mathcal{C})$ is called **generally covariant**, if and only if all the vector bundle automorphisms $V(\mathcal{M}) \rightarrow V(\mathcal{M})$ are automorphisms of the classical field theory.

A classical field theory $(\mathcal{M}, V(\mathcal{M}), F, \mathbb{F}, \mathbb{F}_T, E, \mathcal{C})$ is called **diffeomorphism invariant**, if and only if for all the diffeomorphisms $\mathcal{M} \rightarrow \mathcal{M}$ of the base manifold there

exists a vector bundle automorphism $V(\mathcal{M}) \rightarrow V(\mathcal{M})$, such that it is an automorphism of the classical field theory.

Those automorphisms of a classical field theory $(\mathcal{M}, V(\mathcal{M}), F, \mathbb{F}, \mathbb{F}_T, E, C)$, for which the underlying $\mathcal{M} \rightarrow \mathcal{M}$ diffeomorphism is the identity of \mathcal{M} , are called **internal symmetries** or **gauge transformations**.

Definition 12 The **observables** of a classical field theory $(\mathcal{M}, V(\mathcal{M}), F, \mathbb{F}, \mathbb{F}_T, E, C)$ are the continuous maps $O : F \rightarrow \mathbb{R}$.

Remark 13 The presented formulation of a classical Lagrangian field theory formalizes the Palatini type variational principle, when applied to a setting eventually containing general relativity. That is: the spacetime metric field or its ingredients, if present in the theory, is treated just like any other of the fields. In particular, it is not assumed a priori that on $T(\mathcal{M})$ a Levi–Civita covariant derivation is present associated to some spacetime metric. If a metric and a covariant derivation on $T(\mathcal{M})$ is present, they are varied independently in the presented formulation. We also remark, that in this formulation, the Lagrange form of general relativity can be chosen to be polynomial in the field variables: one variable can be chosen to be the inverse spacetime metric densitised with the metric volume form, i.e. a field $\mathbf{g}_{[cdef]}^{ab}$ (this is in one-to-one correspondence with the ordinary spacetime metric field g_{ab}), and the other variable can simply be the $T(\mathcal{M})$ covariant derivation ∇_a . The Einstein–Hilbert Lagrangian expression is then $(\mathbf{g}_{[cdef]}^{ab}, \nabla_h) \mapsto \mathbf{g}_{[cdef]}^{ab} R(\nabla)_{abh}{}^h$ which is a third degree polynomial of its field variables, where $R(\nabla)_{abc}{}^d$ denotes the Riemann tensor of ∇ .

Remark 14 In QFT formalism, it will be necessary to also consider distributional sections of a vector bundle, not only smooth sections.

- (i) Let $U(\mathcal{M})$ be a vector bundle, as previously, and introduce the notation $U^\times(\mathcal{M}) := U^*(\mathcal{M}) \otimes \overset{m}{\wedge} T^*(\mathcal{M})$ for its **de Rham dual** or **densitised dual** bundle. By construction, one has $U^{\times\times}(\mathcal{M}) \equiv U(\mathcal{M})$. The space of **distributional sections** of $U(\mathcal{M})$ is defined to be the strong topological dual of the \mathcal{D} space, consisting of the compactly supported test sections from $\Gamma(U^\times(\mathcal{M}))$. Recall that we used the notation $\mathbb{F} := \Gamma(U(\mathcal{M}))$, with $U(\mathcal{M}) := V(\mathcal{M}) \oplus T^*(\mathcal{M}) \otimes V(\mathcal{M}) \otimes V^*(\mathcal{M})$, for the space of field variations, and \mathbb{F}_T for the space of compactly supported field variations. We will use the notation $\mathbb{F}^\times := \Gamma(U^\times(\mathcal{M}))$ and \mathbb{F}_T^\times correspondingly. The space of distributional field variations is defined to be $(\mathbb{F}_T^\times)^*$ with the \mathcal{D}^* topology, and the space of compactly supported distributional field variations is defined to be $(\mathbb{F}^\times)^*$ with the \mathcal{E}^* topology. There exist the natural nondegenerate bilinear forms $\mathbb{F} \times \mathbb{F}_T^\times \rightarrow \mathbb{R}$, $(\delta\psi, p_T) \mapsto \int_{\mathcal{M}} \delta\psi p_T$ and $\mathbb{F}_T \times \mathbb{F}^\times \rightarrow \mathbb{R}$, $(\delta\psi_T, p) \mapsto \int_{\mathcal{M}} \delta\psi_T p$, which are jointly sequentially continuous due to Lebesgue’s theorem of dominated convergence. Consequently, the natural embeddings of the dense subspaces $\mathbb{F} \subset (\mathbb{F}_T^\times)^*$ and $\mathbb{F}_T \subset (\mathbb{F}^\times)^*$ are sequentially continuous, and therefore are continuous in the $\mathcal{E} \rightarrow \mathcal{D}^*$ and $\mathcal{D} \rightarrow \mathcal{E}^*$ topology, respectively (the first statement is a consequence of Remark 2, the second is of Remark 61(viii)).

- (ii) A continuous linear operator $A : \mathbb{F} \rightarrow \mathbb{F}$ is said to possess a **formal transpose**, if there exists a continuous linear operator $A^t : \mathbb{F}_T^\times \rightarrow \mathbb{F}_T^\times$, such that for all $\delta\psi \in \mathbb{F}$ and $p_T \in \mathbb{F}_T^\times$ one has that $\int_{\mathcal{M}} (A \delta\psi) p_T = \int_{\mathcal{M}} \delta\psi (A^t p_T)$. The topological transpose $(A^t)^* : (\mathbb{F}_T^\times)^* \rightarrow (\mathbb{F}_T^\times)^*$ of the formal transpose operator A^t is called the **distributional extension** of the operator A . By construction, the restriction of the distributional extension of A to the dense subspace $\mathbb{F} \subset (\mathbb{F}_T^\times)^*$ coincides with the original operator A . It is important to note that an operator $A : \mathbb{F} \rightarrow \mathbb{F}$ is only extendible in the distributional sense, if it has a formal transpose, and not all operators have this property.
- (iii) Let $E : F \times \mathbb{F}_T \rightarrow \mathbb{R}$ be the EL functional of a classical field theory, and $J \in \mathbb{F}_T^*$, then we call an element $K_J \in F$ a **solution with a source J** whenever $\forall \delta\psi_T \in \mathbb{F}_T : (E(K_J) | \delta\psi_T) = (J | \delta\psi_T)$ holds. Specially, one may consider only $J \in \mathbb{F}_T^\times \subset \mathbb{F}^\times \subset \mathbb{F}_T^*$. If $K : \mathbb{F}_T^\times \rightarrow F$ is a continuous map, such that for all $J \in \mathbb{F}_T^\times$ the field $K(J) \in F$ is a solution with a source J , then K is called a **fundamental solution**. (In general, it may or may not exist, and if exists, it is usually not be unique.)

4. Mathematically rigorous definition of the unregularized MDS operator

In this section, if not otherwise mentioned, we will avoid using the knowledge that F , \mathbb{F} and \mathbb{F}_T are the concrete spaces of smooth sections of an affine bundle, its underlying vector bundle, and its corresponding compactly supported test sections, over some finite dimensional real smooth orientable and oriented manifold, as in Section 3. For the sake of genericity, they will be considered instead as abstract spaces, stated below. The symbol $*$ shall denote strong topological dual.

Definition 15 Let F be a real affine space, with an underlying real topological vector space \mathbb{F} . Let the topology on \mathbb{F} be nuclear Fréchet (NF space), see Remark 58. We call F the **space of classical field configurations** and the underlying vector space \mathbb{F} the **space of classical field variations**. Let $\mathbb{F}_T \subset \mathbb{F}$ be some subspace of \mathbb{F} , endowed with a topology not weaker in comparison to \mathbb{F} . Assume moreover \mathbb{F}_T to be either nuclear Fréchet or to be the strict inductive limit of a countable system of nuclear Fréchet spaces with closed adjacent images (LNF space), see Remark 58. Then, we call \mathbb{F}_T the **space of test field variations**.

See Appendix Appendix A and the Appendix of [11] for a condensed summary on the theory of topological vector spaces, especially on nuclear or Fréchet spaces and their duals.

Definition 16 Let F , \mathbb{F} , \mathbb{F}_T be as in Definition 15. Let $E : F \times \mathbb{F}_T \rightarrow \mathbb{R}$, $(\psi, \delta\psi_T) \mapsto E(\psi, \delta\psi_T)$ be a jointly sequentially continuous map which is linear in its second variable. Then, E will be called a **classical Euler–Lagrange (EL) functional**. (By means of Theorem 9(ii,iii), then E is separately continuous in its two variables, and when viewed

as a map $E : F \rightarrow \mathbb{F}_T^*$, $\psi \mapsto E(\psi)$, it is continuous.) Given a $\delta\psi_T \in \mathbb{F}_T$, when the second argument of E is evaluated, it will be denoted by $(E | \delta\psi_T) : F \rightarrow \mathbb{R}$, which is then a continuous map. When that map is evaluated at some $\psi \in F$, we denote it by $(E(\psi) | \delta\psi_T) \in \mathbb{R}$, or equivalently, by $(E | \delta\psi_T)(\psi) \in \mathbb{R}$. We call the equation

$$\text{we search for } \psi \in F \text{ such that : } \quad \forall \delta\psi_T \in \mathbb{F}_T : \quad (E(\psi) | \delta\psi_T) = 0 \quad (11)$$

the **classical Euler–Lagrange (EL) equation**. When E is viewed as a map $E : F \rightarrow \mathbb{F}_T^*$, given any fixed field $\psi_0 \in F$, we use the notation $E_{\psi_0} := E \circ (\mathbb{I}_F + \psi_0)$, which will then be a continuous map $E_{\psi_0} : \mathbb{F} \rightarrow \mathbb{F}_T^*$, and ψ_0 will be then called a **reference field**. (By construction, one has that for all $\psi \in F$, the identity $E(\psi) = E_{\psi_0}(\psi - \psi_0)$ holds.)

Remark 17 In the following, we will need to introduce the tensor algebra operations of \mathbb{F} . In the following remark block let \mathbb{U} denote a nuclear Fréchet (NF) or strong dual of a nuclear Fréchet (DNF) space. (See Remark 58.)

- (i) For all $n \in \mathbb{N}_0$, the completed topological tensor product $\overset{n}{\otimes} \mathbb{U}$ is meaningful (e.g. understood with the projective tensor product topology), and is NF or DNF, respectively. Moreover, in the analogy of finite dimensional vector spaces, the pertinent tensor product can be implemented via the multiplicative realization. That is, it is topologically isomorphic to the space of the jointly continuous multilinear forms on the strong dual space of \mathbb{U} . (See Remark 58)
- (ii) With the same assumptions, one has that for all $n \in \mathbb{N}_0$, the identity $(\overset{n}{\otimes} \mathbb{U})^* \equiv \overset{n}{\otimes} \mathbb{U}^*$ holds (Remark 58).
- (iii) Given a countable system of NF or a countable system of DNF spaces, their cartesian product can be equipped with a vector space structure and with the product (also called Tychonoff or initial or projective) topology (Remark 60). With this, it will become an NF or DNF space, respectively. (See Remark 60, 58) Therefore, the Tychonoff tensor algebra $\mathcal{T}(\mathbb{U}) := \bigoplus_{n=0}^{\infty} \overset{n}{\otimes} \mathbb{U}$ is meaningful and is NF or DNF, respectively. (The symbol $\bigoplus_{n=0}^{\infty} := \bigtimes_{n=0}^{\infty}$ as set operation, but we use rather \bigoplus for vector spaces.)
- (iv) Given a countable system of NF or a countable system of DNF spaces, in their cartesian product vector space there is the subspace of the elements with all zero entries except for finite entries, which subspace is called the algebraic direct sum space. This can be equipped with the locally convex direct sum (also called final or injective) topology. With this, it will become an NF or DNF space, respectively (see Remark 60, 58). Therefore, the algebraic tensor algebra $\mathcal{T}_a(\mathbb{U}) := \bigoplus_{n=0}^{\infty} \overset{n}{\otimes} \mathbb{U}$ with the locally convex direct sum topology is meaningful and is NF or DNF, respectively.
- (v) One has that $(\mathcal{T}(\mathbb{U}))^* \equiv \mathcal{T}_a(\mathbb{U}^*)$ and $(\mathcal{T}_a(\mathbb{U}))^* \equiv \mathcal{T}(\mathbb{U}^*)$. (See Remark 60.)

- (vi) The Tychonoff tensor algebra has a jointly continuous bilinear map $\otimes : \mathcal{T}(\mathbb{U}) \times \mathcal{T}(\mathbb{U}) \rightarrow \mathcal{T}(\mathbb{U})$, the tensor algebra multiplication, with a unit element $\mathbb{1} := (1, 0, 0, \dots) \in \mathcal{T}(\mathbb{U})$ (consequences of Remark 59). The subspaces of k -tensors provide a grading of $\mathcal{T}(\mathbb{U})$. Quite trivially, the left multiplication operator for all $u \in \mathcal{T}(\mathbb{U})$ is a continuous linear map $L_u : \mathcal{T}(\mathbb{U}) \rightarrow \mathcal{T}(\mathbb{U})$.
- (vii) Similarly, the algebraic tensor algebra has a jointly continuous bilinear map $\mathcal{T}_a(\mathbb{U}) \times \mathcal{T}_a(\mathbb{U}) \rightarrow \mathcal{T}_a(\mathbb{U})$, the tensor algebra multiplication, with a corresponding unit element (consequences of Remark 59). The subspaces of k -tensors provide a grading of $\mathcal{T}_a(\mathbb{U})$. Quite trivially, the left multiplication operator is a continuous linear map $\mathcal{T}_a(\mathbb{U}) \rightarrow \mathcal{T}_a(\mathbb{U})$.
- (viii) Since $\mathcal{T}(\mathbb{U})$ and $\mathcal{T}_a(\mathbb{U}^*)$ are strong duals to each-other, and both of these are graded unital associative algebras with jointly continuous multiplications, by transposing the algebra multiplication and unit from the duals, one infers that both $\mathcal{T}(\mathbb{U})$ and $\mathcal{T}_a(\mathbb{U}^*)$ are bialgebras, with corresponding coproduct and counit. The counit of $\mathcal{T}(\mathbb{U})$ is $b : \mathcal{T}(\mathbb{U}) \rightarrow \mathbb{R}$, $G := (G^{(0)}, G^{(1)}, \dots) \mapsto bG := G^{(0)}$, i.e. extraction of the scalar component.
- (ix) Due to the bialgebra nature of $\mathcal{T}(\mathbb{U})$, i.e. due to the existence of a continuous coproduct on $\mathcal{T}(\mathbb{U})$, for all $p \in \mathcal{T}_a(\mathbb{U}^*)$ the corresponding left insertion operator $\iota_p : \mathcal{T}(\mathbb{U}) \rightarrow \mathcal{T}(\mathbb{U})$ is meaningful, and is a continuous linear operator. More concretely, the left insertion operator $\iota_{p^{(n)}}$ with a $p^{(n)} \in \overset{n}{\otimes} \mathbb{U}^*$ ($n \in \mathbb{N}_0$) exists, because for all $m \in \mathbb{N}_0$ ($m \geq n$) the tensor product $\overset{m}{\otimes} \mathbb{U}$ can be identified with the space of $\overset{n}{\otimes} \mathbb{U}^* \times \overset{m-n}{\otimes} \mathbb{U}^* \rightarrow \mathbb{R}$ jointly continuous bilinear forms, as stated in (i). Similarly, the left insertion operators make sense in $\mathcal{T}_a(\mathbb{U})$, and is a continuous linear operator. (For the sake of distinction in terminology, we call merely the operators $\iota_{p^{(1)}}$ with $p^{(1)} \in \mathbb{U}^* \equiv \overset{1}{\otimes} \mathbb{U}^* \subset \mathcal{T}_a(\mathbb{U}^*)$ as insertion operators, whereas for generic $p^{(n)} \in \overset{n}{\otimes} \mathbb{U}^* \subset \mathcal{T}_a(\mathbb{U}^*)$ ($n \in \mathbb{N}_0$) or more generally for $p \in \mathcal{T}_a(\mathbb{U}^*)$, we call the corresponding $\iota_{p^{(n)}}$ or ι_p as multipolynomial insertion operator.) For all $p \in \mathcal{T}_a(\mathbb{U}^*)$, one has the identity $p = b \iota_p$. For the left insertion operator, we use the normalization convention such that for all $G^{(n)} \in \overset{n}{\otimes} \mathbb{U}$ and $u \in \mathbb{U}$ and $p \in \mathbb{U}^*$ one has $\iota_p L_u G^{(n)} = (n+1) (p|u) G^{(n)}$.
- (x) A historical note: over an affine (Minkowski) spacetime, one can define the space of rapidly decreasing (Schwartz) functions \mathcal{S} , which is an NF space. The tensor algebra $\mathcal{T}_a(\mathcal{S})$ is referred to as Borchers–Uhlmann (BU) algebra (original papers: [25, 26], and including a short review: [27]). The Wightman functionals in QFT are understood to be in the space $(\mathcal{T}_a(\mathcal{S}))^* \equiv \mathcal{T}(\mathcal{S}^*)$.
- (xi) By construction, the \mathcal{T}_a topology is strongest tensor algebra topology, whereas \mathcal{T} is the weakest. It is possible to define a natural topological tensor algebra which is in between the \mathcal{T} and \mathcal{T}_a , in terms of topology strength. It will be motivated and introduced later, in Section 6, and will be key to the presented construction, if one wishes to quantize analytic Lagrange forms, and not only polynomial ones.

Definition 18 Assume that the space of field variations as real nuclear Fréchet spaces has a direct sum splitting $\mathbb{F} = \mathbb{F}_r \oplus \mathbb{F}_c$, called **the real-complex splitting**, where both \mathbb{F}_r and \mathbb{F}_c are closed (and therefore nuclear Fréchet), and \mathbb{F}_c has a complex structure (i.e. it can be regarded as a complex nuclear Fréchet space). Denote by $\mathbb{F}_{r\mathbb{C}} := \mathbb{F}_r \otimes \mathbb{C}$ the complexification of \mathbb{F}_r . Then, we use the notation $\mathbb{F}_{(\mathbb{C})} := \mathbb{F}_{r\mathbb{C}} \oplus \mathbb{F}_c$, and call it the **space of field variations with complex structure**. (We assume that also $\mathbb{F}_T \subset \mathbb{F}$ respects this splitting.)

The above definition is necessary, because in field theory, certain fields (like Dirac fields) sit in an inherently complex vector space, whereas other fields, like gauge fields sit in an inherently real vector space, and QFT assumes that the sectors not being inherently complex are complexified. (In the most simple case, one has merely $\mathbb{F}_{(\mathbb{C})} = \mathbb{F}_{\mathbb{C}}$.)

Definition 19 Let the vector space of field variations admit a real-complex splitting $\mathbb{F} = \mathbb{F}_r \oplus \mathbb{F}_c$, as in Definition 18. Furthermore, assume a direct sum structure $\mathbb{F} = \bigoplus_{i=1}^f \mathbb{F}_i$, such that for each $i = 1, \dots, f$ the subspace \mathbb{F}_i is either entirely within \mathbb{F}_r or in \mathbb{F}_c and are closed (thus, also nuclear Fréchet), and let there be integers $s_i \in \{0, 1\}$ associated to each subspace \mathbb{F}_i ($i = 1, \dots, f$). Then, the subspaces $\mathbb{F}_1, \dots, \mathbb{F}_f$ are called the **flavor sectors**, and their associated integers s_1, \dots, s_f are called **bosonic or fermionic labels**. (We assume that also $\mathbb{F}_T \subset \mathbb{F}$ respects this splitting.)

In the most simple case, there is only one single flavor sector, globally endowed with a bosonic or fermionic label.

Remark 20 Whenever \mathbb{F} is split as $\mathbb{F} = \bigoplus_{i=1}^f \mathbb{F}_i$ into flavor sectors with bosonic / fermionic labels s_i ($i = 1, \dots, f$), as in Definition 19, then for all $n \in \mathbb{N}_0$ one may introduce a continuous linear representation U_π of a permutation group element $\pi \in \Pi_n$ on the space $\bigotimes^n \mathbb{F}_{(\mathbb{C})}$ as follows (see also [28] Chapter4). Take an element $x_1 \otimes \dots \otimes x_n \in \bigotimes^n \mathbb{F}_{(\mathbb{C})}$, where each factor x_i ($i = 1, \dots, n$) resides in some $\mathbb{F}_{(\mathbb{C})j}$ ($j = 1, \dots, f$). Then, set

$$U_\pi(x_1 \otimes \dots \otimes x_n) := (-1)^{s_1 \sigma_1(\pi) + \dots + s_f \sigma_f(\pi)} x_{\pi(1)} \otimes \dots \otimes x_{\pi(n)},$$

where $\sigma_i(\pi) \in \{0, 1\}$ ($i = 1, \dots, f$) is the parity of the permutation π within each index block. The map U_π can then be linearly extended in $\bigotimes^n \mathbb{F}_{(\mathbb{C})}$. Due to the NF property of the involved spaces, the topology defining seminorms on $\bigotimes^n \mathbb{F}_{(\mathbb{C})}$ may be taken to be such that U_π are continuous ([28] Chapter4), therefore can uniquely be extended as acting as a continuous linear map $U_\pi : \bigotimes^n \mathbb{F}_{(\mathbb{C})} \rightarrow \bigotimes^n \mathbb{F}_{(\mathbb{C})}$, thus defining the signed permutation operator on the entire space $\bigotimes^n \mathbb{F}_{(\mathbb{C})}$. Therefore, on each space $\bigotimes^n \mathbb{F}_{(\mathbb{C})}$ the continuous linear projection operator

$$P_n := \frac{1}{n!} \sum_{\pi \in \Pi_n} U_\pi$$

can be defined. The family of operators P_n ($n \in \mathbb{N}_0$) on the spaces $\bigotimes^n \mathbb{F}_{(\mathbb{C})}$ can be joined as a single grading preserving continuous linear projection operator $P : \mathcal{T}(\mathbb{F}_{(\mathbb{C})}) \rightarrow$

$\mathcal{T}(\mathbb{F}_{(\mathbb{C})})$. This signed symmetrizer projection operator P has the following properties against the tensor algebra multiplication:

$$P(xy) = P(P(x)P(y)) = P(P(x)y) = P(xP(y)) \quad (\forall x, y \in \mathcal{T}(\mathbb{F}_{(\mathbb{C})})).$$

Therefore, the closed subspace $\text{Ker}(P)$ is a two-sided ideal in $\mathcal{T}(\mathbb{F}_{(\mathbb{C})})$. (The presented approach was inspired by [28] Chapter 4.)

Using the fact that the closed subspace of an NF space is also NF and that the factor space of an NF space with a closed subspace is also NF (Remark 58), the following definition is meaningful.

Definition 21 *Let the space of field variations \mathbb{F} admit flavor sectors \mathbb{F}_i and bosonic / fermionic labels s_i ($i = 1, \dots, f$), as in Definition 19, and corresponding signed symmetrization projector P as in Remark 20. Then the factor algebra $A(\mathbb{F}_{(\mathbb{C})}) := \mathcal{T}(\mathbb{F}_{(\mathbb{C})}) / \text{Ker}(P)$ is called the **field algebra**. Clearly, it is a unital associative algebra, and a nuclear Fréchet (NF) topological vector space, with jointly continuous algebra multiplication $\bullet : A(\mathbb{F}_{(\mathbb{C})}) \times A(\mathbb{F}_{(\mathbb{C})}) \rightarrow A(\mathbb{F}_{(\mathbb{C})})$. The topological transpose P^* of P allows the analogous construction in the strong dual of $\mathcal{T}(\mathbb{F}_{(\mathbb{C})})$, which makes it also a unital associative algebra with jointly continuous algebra multiplication, and therefore $A(\mathbb{F}_{(\mathbb{C})})$ retains the bialgebra structure from $\mathcal{T}(\mathbb{F}_{(\mathbb{C})})$.*

Since the complementing projection operator $I-P$ to P is also continuous, as topological vector spaces one may naturally identify $A(\mathbb{F}_{(\mathbb{C})})$ with the closed subspace $\text{Ran}(P) = \text{Ker}(I-P) \subset \mathcal{T}(\mathbb{F}_{(\mathbb{C})})$. Using this linear topological identification, the algebraic product \bullet may be pushed forward from $A(\mathbb{F}_{(\mathbb{C})})$ to the subspace $\text{Ran}(P) \subset \mathcal{T}(\mathbb{F}_{(\mathbb{C})})$. That is, as usual, the algebra $A(\mathbb{F}_{(\mathbb{C})})$ may be regarded as a closed subspace of $\mathcal{T}(\mathbb{F}_{(\mathbb{C})})$. On that space the product \bullet can be traced back to the tensor algebra product \otimes , with the identity: for all $x \in \otimes^m \mathbb{F}_{(\mathbb{C})}$ and $y \in \otimes^n \mathbb{F}_{(\mathbb{C})}$, one has $x \bullet y = \frac{(m+n)!}{m!n!} P(x \otimes y)$. The unit element, the counit map, as well as the insertion operator by a one-form $p^{(1)} \in \mathbb{F}_{(\mathbb{C})}^*$ coincides to the one defined on $\mathcal{T}(\mathbb{F}_{(\mathbb{C})})$. The strong dual of $A(\mathbb{F}_{(\mathbb{C})})$ may be identified with the corresponding subspace of $\mathcal{T}_a(\mathbb{F}_{(\mathbb{C})}^*)$. Whenever not confusing, we will suppress the multiplication symbol \bullet .

The above definition was necessary, because in QFT certain sectors of fields are bosonic, others are fermionic. (In the most simple case one has that $A(\mathbb{F}_{(\mathbb{C})})$ is $\vee(\mathbb{F}_{(\mathbb{C})})$ or $\wedge(\mathbb{F}_{(\mathbb{C})})$ of $\mathcal{T}(\mathbb{F}_{(\mathbb{C})})$.) As seen from above, the left multiplication operator (given some $\delta\psi \in \mathbb{F}$) is the same as $L_{\delta\psi}$ in $\mathcal{T}(\mathbb{F}_{(\mathbb{C})})$, with a subsequent mixed symmetrization/antisymmetrization and normalization adjustment, respectively. It shall be denoted by the same symbol $L_{\delta\psi}$, when not confusing. Due to the symmetrization/antisymmetrization, one has that for all $G \in A(\mathbb{F}_{(\mathbb{C})})$, and for all $\delta\psi \in \mathbb{F}_{(\mathbb{C})}$, $\delta J \in \mathbb{F}_{(\mathbb{C})}^*$ from the same fermionic sector one has that $(\iota_{\delta J} L_{\delta\psi} + L_{\delta\psi} \iota_{\delta J}) G = (\delta J | \delta\psi) G$, whereas $(\iota_{\delta J} L_{\delta\psi} - L_{\delta\psi} \iota_{\delta J}) G = (\delta J | \delta\psi) G$ otherwise. Also recall that the counit of $\mathcal{T}(\mathbb{F})$ was the continuous linear map $b : \mathcal{T}(\mathbb{F}) \rightarrow \mathbb{R}$, $G = (G^{(0)}, G^{(1)}, \dots) \mapsto bG := G^{(0)}$, the letter “ b ” standing for “base” or “bottom form”. This can be also regarded as the counit of the field algebra $A(\mathbb{F}_{(\mathbb{C})})$. The expression bG simply picks out

the scalar component of G . The map $\mathbb{1} b$ thus defines a continuous projection operator onto the space of scalars, whereas $(\mathbb{I} - \mathbb{1} b)$ defines a continuous projection operator onto the algebra ideal of at-least-rank-1 multivectors.

Until Section 6, for the sake of simplicity we assume that the EL functional $E : F \rightarrow \mathbb{F}_T^*$ is multipolynomial, which we define as follows.

Definition 22 *Let $E : F \times \mathbb{F}_T \rightarrow \mathbb{R}$ be an EL functional as in Definition 16. We say that the EL functional E is **multipolynomial**, whenever there exists a reference field $\psi_0 \in F$, such that there exists an element $\mathbf{E}_{\psi_0} \in (A(\mathbb{F}_{(\mathbb{C})}))^* \otimes \mathbb{F}_T^* \subset \mathcal{T}_a(\mathbb{F}_{(\mathbb{C})}^*) \otimes \mathbb{F}_T^*$, for which*

$$\forall \psi \in F, \delta\psi_T \in \mathbb{F}_T : \quad (E_{\psi_0}(\psi - \psi_0) | \delta\psi_T) = \left(\mathbf{E}_{\psi_0} \left| \left(1, \overset{1}{\otimes}(\psi - \psi_0), \overset{2}{\otimes}(\psi - \psi_0), \dots \right) \otimes \delta\psi_T \right. \right)$$

holds. (Note that then for all $\psi_0 \in F$ there exists the corresponding element \mathbf{E}_{ψ_0} .) When an element $\delta\psi_T \in \mathbb{F}_T$ is contracted with \mathbf{E}_{ψ_0} in its last tensorial entry, we will use the notation $(\mathbf{E}_{\psi_0} | \delta\psi_T)$ to denote the corresponding element of $(A(\mathbb{F}_{(\mathbb{C})}))^ \subset \mathcal{T}_a(\mathbb{F}_{(\mathbb{C})}^*)$.*

Given the above definitions, an EL functional $E : F \times \mathbb{F}_T \rightarrow \mathbb{R}$, when a reference field $\psi_0 \in F$ is also fixed, may be viewed as a map $E_{\psi_0} : F \times \mathbb{F}_T \rightarrow \mathbb{R}$, and when a test field $\delta\psi_T \in \mathbb{F}_T$ is also fixed, the map $(E_{\psi_0} | \delta\psi_T) : F \rightarrow \mathbb{R}$ may be viewed as an element $(\mathbf{E}_{\psi_0} | \delta\psi_T) \in (A(\mathbb{F}_{(\mathbb{C})}))^* \subset \mathcal{T}_a(\mathbb{F}_{(\mathbb{C})}^*)$, if E was multipolynomial. Given $(\mathbf{E}_{\psi_0} | \delta\psi_T)$, it has a corresponding multipolynomial insertion operator over the tensor algebra $\mathcal{T}(\mathbb{F}_{(\mathbb{C})})$, as stated in Remark 17(ix). We shall denote that by the symbol $\mathcal{L}_{(\mathbf{E}_{\psi_0} | \delta\psi_T)}$.

Definition 23 *Let \hbar be a fixed real number. Let $F, \mathbb{F}, \mathbb{F}_T$ as in Definition 15. Let $E : F \times \mathbb{F}_T \rightarrow \mathbb{R}$ as in Definition 16, and assume that it is multipolynomial as in Definition 22, i.e. that for any reference field $\psi_0 \in F$ and test fields $\delta\psi_T \in \mathbb{F}_T$ there is an element $(\mathbf{E}_{\psi_0} | \delta\psi_T) \in \mathcal{T}_a(\mathbb{F}^*)$ reproducing the map $\delta\psi \mapsto (E_{\psi_0}(\delta\psi) | \delta\psi_T)$. Let $A(\mathbb{F}_{(\mathbb{C})})$ be the field algebra as in Definition 21. Then, for some fixed reference field $\psi_0 \in F$ and fixed test field $\delta\psi_T \in \mathbb{F}_T$ the operator*

$$\mathbf{M}_{\hbar, \psi_0, \delta\psi_T} : A(\mathbb{F}_{(\mathbb{C})}) \rightarrow A(\mathbb{F}_{(\mathbb{C})}), \quad G \mapsto \mathbf{M}_{\hbar, \psi_0, \delta\psi_T} G := \left(\mathcal{L}_{(\mathbf{E}_{\psi_0} | \delta\psi_T)} - i \hbar L_{\delta\psi_T} \right) G \quad (12)$$

*is called the **unregularized master Dyson–Schwinger (MDS) operator**. Moreover, we call the below equation the **unregularized master Dyson–Schwinger (MDS) equation**:*

we search for $(\psi_0, G_{\psi_0}) \in F \times A(\mathbb{F}_{(\mathbb{C})})$, such that :

$$b G_{\psi_0} = 1, \quad \text{and} \quad \forall \delta\psi_T \in \mathbb{F}_T : \quad \mathbf{M}_{\hbar, \psi_0, \delta\psi_T} G_{\psi_0} = 0. \quad (13)$$

(When not particularly relevant, the \hbar will be merged into the normalization of E .)

The MDS formulation of QFT can be thought of as a construction, where the objects of interest are elements of $F \times A(\mathbb{F}_{(\mathbb{C})})$, and the selection equation for the physically realized such elements is the MDS equation. In Section 5 it shall be shown that some finetuning (regularization) to this idea is needed, as is well known in the QFT literature.

Definition 24 Any continuous map $O : F \rightarrow \mathbb{R}$ is called an **observable**, similarly as in a classical field theory. Given a fixed $\psi_0 \in F$, we use the notation $O_{\psi_0} := O \circ (\mathbf{I}_{\mathbb{F}} + \psi_0)$, which is then a continuous map $O_{\psi_0} : \mathbb{F} \rightarrow \mathbb{R}$, and one has $O(\psi) = O_{\psi_0}(\psi - \psi_0)$ for all $\psi \in F$ and observable O . An observable $O : F \rightarrow \mathbb{R}$ is called **multipolynomial observable**, whenever for some reference fields $\psi_0 \in F$, there exists an element $\mathbf{O}_{\psi_0} \in \mathcal{T}_a(\mathbb{F}^*)$, such that for all $\psi \in F$, one has $O_{\psi_0}(\psi - \psi_0) = \left(\mathbf{O}_{\psi_0} \mid \left(1, \overset{1}{\otimes}(\psi - \psi_0), \overset{2}{\otimes}(\psi - \psi_0), \dots \right) \right)$. (If it holds, it then holds for any $\psi_0 \in F$.)

Definition 25 Given a solution $(\psi_0, G_{\psi_0}) \in F \times A(\mathbb{F}(\mathbb{C}))$ of the MDS equation, the **(Feynman type) quantum expectation value** of the multipolynomial observable $O : F \rightarrow \mathbb{R}$ at the solution (ψ_0, G_{ψ_0}) is $\mu_{(\psi_0, G_{\psi_0})}(O) := \text{Re} \left(\mathbf{O}_{\psi_0} \mid G_{\psi_0} \right)$.

We note in passing that the above definition of MDS operator and observables can eventually be well defined also for non-polynomial but analytic EL functionals and observables. For that, however, a stronger topology is needed on the tensor algebra of \mathbb{F} , which we will address later in Section 6.

5. The weak (distributional) and the Wilsonian regularized MDS operator

Throughout this section, we use the information that the EL functional $E : F \times \mathbb{F}_T \rightarrow \mathbb{R}$ is assumed to be that of a classical field theory as defined in Section 3. Due to that, the de Rham dual spaces \mathbb{F}^\times and \mathbb{F}_T^\times are naturally defined according to Remark 14. First, we define what we mean by a free or non-interacting theory.

Definition 26 Let $F, \mathbb{F}, \mathbb{F}_T, A(\mathbb{F}(\mathbb{C})), E$ be as in Definition 23. We call the EL functional $E : F \times \mathbb{F}_T \rightarrow \mathbb{R}$ to be **free** or **non-interacting**, whenever the corresponding continuous map $E : F \rightarrow \mathbb{F}_T^*$ is affine. We call the Euler–Lagrangian functional **interacting** otherwise. (Note that by construction, for a free EL functional, given any reference field $\psi_0 \in F$, the map $E_{\psi_0}(\cdot) - E_{\psi_0}(0) : \mathbb{F} \rightarrow \mathbb{F}_T^*$ is linear. If ψ_0 were an EL solution, then $E_{\psi_0}(\cdot) : \mathbb{F} \rightarrow \mathbb{F}_T^*$ is linear.)

Remark 27 It is quite evident to see that if $(\psi_0, G_{\psi_0}) \in F \times A(\mathbb{F}(\mathbb{C}))$ were a solution to the unregularized MDS equation Eq.(13), and the reference field $\psi_0 \in F$ is and EL solution, then $\iota_{(\mathbf{E}_{\psi_0} | \delta \psi_T)} G_{\psi_0}^{(2)} = i \hbar \delta \psi_T$ holds for all test fields $\delta \psi_T \in \mathbb{F}_T$.

Corollary 28 Let the EL functional E be the one of the free wave equation over Minkowski spacetime. In that case, the solution space of the unregularized MDS equation Eq.(13) is empty, whenever $\hbar \neq 0$. □

The above is rather evident by means of Remark 27: the second order correlator $G_{\psi_0}^{(2)}$ would need to be proportional to a fundamental solution (Green’s functional), recalled in Remark 14(iii). We know, however, from the theory of PDEs, that already e.g. for the case of the simple wave equation over Minkowski spacetime, there is no fundamental solution in the space of fields in $\mathbb{F} \otimes \mathbb{F}$. The fundamental solution exists, however, as an element of $\mathcal{L}(\mathbb{F}_T^\times, \mathbb{F})$, which can be considered as a distribution. It is tempting thus to

modify the definition of the MDS equation to a weak (distributional) sense, so that free theories do have MDS solutions.

Remark 29 In order to define the weak MDS equation, we will need to substitute $A(\mathbb{F}_{(\mathbb{C})})$ with its distributional version, which is expanded below.

- (i) Let us take the affine bundle of classical field configurations $V(\mathcal{M}) \times_w DV(\mathcal{M})$ over the base manifold \mathcal{M} . Subordinate to it there is the corresponding underlying vector bundle of field variations $U(\mathcal{M}) := V(\mathcal{M}) \oplus T^*(\mathcal{M}) \otimes V(\mathcal{M}) \otimes V^*(\mathcal{M})$. Take the densitised dual of that vector bundle, $U^\times(\mathcal{M})$, as in Remark 14(i). Take the n -fold external tensor product bundle $U^\times(\mathcal{M}) \boxtimes \cdots \boxtimes U^\times(\mathcal{M})$ of that, which will then be a vector bundle over the n -fold cartesian product $\mathcal{M} \times \cdots \times \mathcal{M}$ as base manifold. The smooth sections of this vector bundle shall be denoted by \mathbb{F}_n^\times , and has its natural \mathcal{E} topology which is nuclear Fréchet (NF), and is topologically isomorphic to $\bigotimes^n \mathbb{F}^\times$ by means of Schwartz kernel theorem. It has the subspace of compactly supported sections, denoted by \mathbb{F}_T^\times and is a dense subspace within \mathbb{F}_n^\times in the \mathcal{E} topology. The space \mathbb{F}_T^\times with its natural \mathcal{D} topology becomes a countable strict inductive limit of nuclear Fréchet spaces with closed adjacent images (LNF space) whenever the base manifold \mathcal{M} is noncompact, and is nuclear Fréchet (NF) if \mathcal{M} is compact (see Remark 58). The strong dual of the space \mathbb{F}_T^\times is denoted by $(\mathbb{F}_T^\times)^*$ with its natural \mathcal{D}^* topology. It is a DLNF space when \mathcal{M} is noncompact, and DNF when \mathcal{M} is compact. One has that $\bigotimes^n \mathbb{F} \subset (\mathbb{F}_T^\times)^*$, i.e. the latter space can be regarded as the space of distributional n -field correlators.
- (ii) In the above construction we avoided using completed topological tensor product $\bigotimes^n \mathbb{F}_T^\times$, as that space is topologically not isomorphic to \mathbb{F}_T^\times whenever we are in the realm of LNF spaces, i.e. when \mathcal{M} is noncompact (although they are isomorphic as linear spaces, the latter has a stronger topology). This slight complication is mentioned in more details in Remark 59(iv). The complication is absent, whenever \mathcal{M} is compact by means of Remark 59 ($\mathbb{F}_T^\times \equiv \bigotimes^n \mathbb{F}_T^\times$ topologically, in that case).
- (iii) One can form the algebraic tensor algebra $\mathcal{T}_a(\mathbb{F}_T^\times)$, defined as the algebraic direct sum $\bigoplus_{n=0}^{\infty} \mathbb{F}_T^\times$ equipped with the locally convex direct sum topology. Its topology will be LNF whenever \mathcal{M} is noncompact, and NF if \mathcal{M} is compact. As an algebra, $\mathcal{T}_a(\mathbb{F}_T^\times)$ forms a unital associative algebra, with (at least) separately continuous algebra multiplication.
- (iv) The distributional field correlators $\mathcal{T}((\mathbb{F}_T^\times)^*)$ is defined to be the space $(\mathcal{T}_a(\mathbb{F}_T^\times))^*$. It is topologically isomorphic to $\bigoplus_{n=0}^{\infty} (\mathbb{F}_T^\times)^*$, by means of Remark 60. It is a DLNF space when \mathcal{M} is noncompact, and DNF space if \mathcal{M} is compact. It is also a unital associative algebra, with an (at least) separately continuous algebra multiplication.
- (v) The mixedly symmetrized/antisymmetrized field algebra $A((\mathbb{F}_T^\times(\mathbb{C}))^*)$ can be defined in the analogy of Definition 21. Clearly, the field algebra $A(\mathbb{F}_{(\mathbb{C})})$, used in the previous sections, is dense in the distributional sense field algebra $A((\mathbb{F}_T^\times(\mathbb{C}))^*)$.

Remark 30 The MDS operator of a free EL functional of a classical field theory can be naturally extended in a distributional sense. This is seen as follows.

- (i) First of all, for all $\delta\psi_T \in \mathbb{F}_T$ and $G \in A(\mathbb{F}(\mathbb{C}))$ and $p \in \mathcal{T}_a(\mathbb{F}_T^\times)$ one has that $(p | L_{\delta\psi_T} G) = (\mathcal{L}_{\delta\psi_T} p | G)$. Moreover, the linear map $\mathcal{L}_{\delta\psi_T} : \mathcal{T}_a(\mathbb{F}_T^\times) \rightarrow \mathcal{T}_a(\mathbb{F}_T^\times)$ is continuous. Therefore, $\mathcal{L}_{\delta\psi_T}$ is the formal transpose of $L_{\delta\psi_T}$. Taking the topological transpose of $\mathcal{L}_{\delta\psi_T}$ one gets the distributional extension of $L_{\delta\psi_T}$ as a continuous linear map $A((\mathbb{F}_T^\times(\mathbb{C}))^*) \rightarrow A((\mathbb{F}_T^\times(\mathbb{C}))^*)$.
- (ii) Whenever E is the EL functional of a classical field theory, and a $\psi_0 \in F$ is fixed, then for each $\delta\psi_T \in \mathbb{F}_T$ there exists a unique element $\pi_T \in \mathbb{F}_T^\times$, such that $(\mathbf{E}_{\psi_0} | \delta\psi_T) \delta\psi - (E(\psi_0) | \delta\psi_T) = \int_{\mathcal{M}} \pi_T \delta\psi$ for all $\delta\psi \in \mathbb{F}$. Consequently, one has that $\mathcal{L}_{(\mathbf{E}_{\psi_0} | \delta\psi_T) - (E(\psi_0) | \delta\psi_T)} = \mathcal{L}_{\int_{\mathcal{M}} \pi_T}(\cdot)$. Because of that, for any $G \in A(\mathbb{F}(\mathbb{C}))$ and $p \in \mathcal{T}_a(\mathbb{F}_T^\times)$ one has the identity $(p | \mathcal{L}_{(\mathbf{E}_{\psi_0} | \delta\psi_T) - (E(\psi_0) | \delta\psi_T)} G) = (L_{\pi_T} p | G)$. Consequently, the formal transpose of $\mathcal{L}_{(\mathbf{E}_{\psi_0} | \delta\psi_T)}$ exists, being the continuous linear map $L_{\pi_T} + (E(\psi_0) | \delta\psi_T) I : \mathcal{T}_a(\mathbb{F}_T^\times) \rightarrow \mathcal{T}_a(\mathbb{F}_T^\times)$. The topological transpose of that operator will give rise to the distributional extension of the operator as a $\mathcal{L}_{(\mathbf{E}_{\psi_0} | \delta\psi_T)} : A((\mathbb{F}_T^\times(\mathbb{C}))^*) \rightarrow A((\mathbb{F}_T^\times(\mathbb{C}))^*)$ map.
- (iii) The above construction clearly fails for non-free classical field theories, since then the formal transpose of $\mathcal{L}_{(\mathbf{E}_{\psi_0} | \delta\psi_T)}$ as a continuous linear map $\mathcal{T}_a(\mathbb{F}_T^\times) \rightarrow \mathcal{T}_a(\mathbb{F}_T^\times)$ cannot be defined.

Definition 31 Let $F, \mathbb{F}, \mathbb{F}_T, A(\mathbb{F}(\mathbb{C})), E, \hbar$ be as in Definition 23, and let $E : F \times \mathbb{F}_T \rightarrow \mathbb{R}$ the EL functional of a classical field theory. Let it be non-interacting as in the Definition 26. Fix a reference field $\psi_0 \in F$. Then, by means of Remark 30, for all $\delta\psi_T \in \mathbb{F}_T$, the MDS operator can be extended as a continuous linear operator $\mathbf{M}_{\hbar, \psi_0, \delta\psi_T} : A((\mathbb{F}_T^\times(\mathbb{C}))^*) \rightarrow A((\mathbb{F}_T^\times(\mathbb{C}))^*)$, called to be the **weak or distributional master Dyson–Schwinger (MDS) operator**. We call the equation

$$\begin{aligned} & \text{we search for } G_{\psi_0} \in A((\mathbb{F}_T^\times(\mathbb{C}))^*), \text{ such that :} \\ & b G_{\psi_0} = 1, \quad \text{and} \quad \forall \delta\psi_T \in \mathbb{F}_T : \mathbf{M}_{\hbar, \psi_0, \delta\psi_T} G_{\psi_0} = 0. \end{aligned} \tag{14}$$

the **weak or distributional master Dyson–Schwinger (MDS) equation**.

Remark 32 Let $F, \mathbb{F}, \mathbb{F}_T, A((\mathbb{F}_T^\times(\mathbb{C}))^*), E, \hbar$ as in Definition 31, so that the weak MDS operator can be defined, and fix a reference field $\psi_0 \in F$. Assume that the EL equation admits a fundamental solution $\mathbf{K} : \mathbb{F}_T^\times \rightarrow F, J \mapsto \mathbf{K}(J)$ as in Remark 14(iii). In that case, one can construct a corresponding fundamental solution $\mathbf{K}_{\psi_0} : \mathbb{F}_T^\times \rightarrow \mathbb{F}, J \mapsto \mathbf{K}_{\psi_0}(J) := \mathbf{K}(J) - \psi_0$ of the EL functional $E_{\psi_0} : \mathbb{F} \rightarrow \mathbb{F}_T^*$. Assume moreover, that ψ_0 was chosen to be an EL solution, in which case $E_{\psi_0} : \mathbb{F} \rightarrow \mathbb{F}_T^*$ becomes linear, and assume that \mathbf{K}_{ψ_0} can be chosen to be linear. Such a linear fundamental solution $\mathbf{K}_{\psi_0} : \mathbb{F}_T^\times \rightarrow \mathbb{F}$ can be naturally considered as an element $\mathbf{K}_{\psi_0}^{(2)} \in \mathcal{L}(\mathbb{F}_T^\times, \mathbb{F}) \subset (\mathbb{F}_T^\times)_2^*$. Assume moreover, that $\mathbf{K}_{\psi_0}^{(2)}$ can be chosen to be invariant to the permutation symmetry of the field algebra.

- (i) Given these conditions, one may define the element $K_{\psi_0} := (0, 0, i\hbar K_{\psi_0}^{(2)}, 0, 0, \dots) \in A((\mathbb{F}_T^\times(\mathbb{C}))^*)$, called to be the connected correlator, and one can take the ansatz $G_{\psi_0} := \exp(K_{\psi_0}) \in A((\mathbb{F}_T^\times(\mathbb{C}))^*)$. Then, $(\psi_0, G_{\psi_0}) \in F \times A((\mathbb{F}_T^\times(\mathbb{C}))^*)$ solves the weak (distributional sense) MDS equation Eq.(14).
- (ii) For the bosonic case, the above statement is seen trivially, by the fact that for all $\delta\psi_T \in \mathbb{F}_T$ the insertion operator $\iota_{(\mathbf{E}_{\psi_0}|\delta\psi_T)}$ is an algebra derivation, and the field algebra is commutative, so one can use the formula for the derivative of exponential. If \mathbb{F} has fermionic flavor sectors as well, then one can still trace the problem back to derivations acting on exponential: whenever $\delta\psi_T \in \mathbb{F}_T$ resides in a single flavor sector, then for all $\delta\psi'_T$ from the same flavor sector, the linear map $L_{\delta\psi'_T} \iota_{(\mathbf{E}_{\psi_0}|\delta\psi_T)}$ is also an algebra derivation.
- (iii) Rather evidently, the above do not necessarily exhaust all the possible solutions. Typically, a fundamental solution K_{ψ_0} satisfying the above is not unique. Moreover, one may add any term δK_{ψ_0} to K_{ψ_0} satisfying $b\delta K_{\psi_0} = 0$ and $\forall \delta\psi_T \in \mathbb{F}_T : \iota_{(\mathbf{E}_{\psi_0}|\delta\psi_T)} \delta K_{\psi_0} = 0$, in which case $\exp(K_{\psi_0} + \delta K_{\psi_0})$ will still solve the weak MDS equation. In usual QFTs, these ambiguities are removed by further invariance requirements on G_{ψ_0} , which are not imposed in the present paper.
- (iv) The existence of the assumed type of fundamental solution is guaranteed for any EL functional over an affine base manifold, whenever E_{ψ_0} corresponds to a linear PDE with a multipolynomial differential operator, having constant coefficients. This is ensured by the celebrated Malgrange–Ehrenpreis theorem ([29] Chapitre I.1 Théorème 1 and [30] Chapter 6 Theorem 10).

Corollary 33 *Let the EL functional E be the one of a non-interacting classical field theory, which has a fundamental solution as in Remark 32. In that case, the solution space of the weak (distributional) MDS equation Eq.(14) is **not** empty.*

According to Remark 32, given a reference field $\psi_0 \in F$, the solutions of the weak MDS equation Eq.(14) are not unique for free EL functionals. In the usual QFT constructions, this ambiguity is removed by additional requirements, such as Poincaré invariance of the solutions. In the presented construction, however such auxiliary conditions are not imposed, since in a generally covariant setting, it is not evident that the vacuum state should be required to be unique or not.

Remark 34 *For interacting models, the followings can be stated.*

- (i) *On one hand, there is a negative result: for a generic non-interacting EL functional over Minkowski spacetime, the unregularized MDS equation Eq.(13) has no solutions.*
- (ii) *On the other hand, there is a positive result: for a generic non-interacting EL functional of a classical field theory having appropriate fundamental solution, the weak MDS equation Eq.(14) does have solutions, just as is the common wisdom in heuristic QFT.*

- (iii) For interacting models, the weak MDS operator cannot be defined as an everywhere defined continuous operator acting on the space of distributional correlators. For instance, in a fourth order type interaction Lagrangian over Minkowski spacetime \mathcal{M} , the interacting part of the EL functional would be $(\psi, \delta\psi_T) \mapsto \int_{\mathcal{M}} \delta\psi_T (\psi - \psi_0)^3 v$ (here v denoting the Lebesgue volume form). Its corresponding term in the MDS operator is $(G^{(n)}, \delta\psi_T) \mapsto \int_{x \in \mathcal{M}} \delta\psi_T(x) G^{(n)}(x, x, x, \dots) v(x)$. This has no distributional extension, i.e. the operator $\mathcal{L}_{(\mathbb{E}_{\psi_0} | \delta\psi_T)}$ fails to possess a formal transpose. Technically, this phenomenon occurs because the diagonal evaluation map of smooth functions $((x, y) \mapsto G(x, y)) \mapsto (z \mapsto G(z, z))$ $x, y, z \in \mathcal{M}$ cannot be extended to the distributions, in general.
- (iv) In order to remedy the above problem, one is tempted to view the everywhere defined continuous bilinear operator $\mathbf{M}_{h, \psi_0} : A(\mathbb{F}_{(\mathbb{C})}) \times \mathbb{F}_T \rightarrow A(\mathbb{F}_{(\mathbb{C})})$ as a densely defined bilinear operator $\hat{\mathbf{M}}_{h, \psi_0} : A((\mathbb{F}_T^{\times}(\mathbb{C}))^*) \times \mathbb{F}_T \rightarrow A((\mathbb{F}_T^{\times}(\mathbb{C}))^*)$, via the natural dense linear embedding $A(\mathbb{F}_{(\mathbb{C})}) \subset A((\mathbb{F}_T^{\times}(\mathbb{C}))^*)$ of the function sense correlators to the distributional sense correlators. Then, one is tempted to take its maximally extended operator, the extension understood e.g. by sequential closure. (That is, a distributional correlator would be in the domain of the extended $\hat{\mathbf{M}}_{h, \psi_0}$, whenever it admits a function sense approximating sequence converging to it in the distributional sense, such that the evaluated \mathbf{M}_{h, ψ_0} on the approximator sequence is convergent in the distributional sense. The closure would exist, whenever any two approximator sequence of the same domain element yielded the same result.) This strategy is made impossible by the fact that for all interacting EL functionals one can show that the MDS operator is not closable as a densely defined operator in the space of distributional sense correlators. (This occurs because the above diagonal evaluation map is so-called maximally non-closable, see Remark 62 for more details.)
- (v) The celebrated Hörmander's criterion [31] on the wave front set gives a sufficient condition for diagonal evaluation of multivariate distributions, but that condition is not applicable for the present problem. (E.g. already the wave front set of a solution to the distributional MDS equation generated from the Minkowski wave equation is known to fail Hörmander's sufficiency criterion.)
- (vi) One can prove that the solution space of the unregularized MDS equation (understood over the smooth correlators) is always empty, regardless of the structure of the underlying base manifold \mathcal{M} and the interactions in the EL functional (we plan to detail the proof in a different paper).

In summary, the problem is that for interacting models, only the function sense MDS operator is well defined, but its solution space is always empty. In the non-interacting case, the MDS operator can be extended in the distributional sense, and its solution space has the right properties. However, the distributional extension of the MDS operator cannot be achieved for interacting models. In order to overcome this difficulty, one needs the regularized MDS operator, introduced below.

Definition 35 Let $F, \mathbb{F}, \mathbb{F}_T, A(\mathbb{F}(\mathbb{C})), E, \hbar$ be as in Definition 23. Assume that E is the EL functional of a classical field theory, so that \mathbb{F}^\times and \mathbb{F}_T^\times is defined. Fix a continuous linear operator $\mathcal{R} : \mathbb{F} \rightarrow \mathbb{F}$. Given these, we call the operator

$$\mathbf{M}_{\hbar, \psi_0, \mathcal{R}, \delta\psi_T} : A(\mathbb{F}(\mathbb{C})) \rightarrow A(\mathbb{F}(\mathbb{C})), G \mapsto \mathbf{M}_{\hbar, \psi_0, \mathcal{R}, \delta\psi_T} G := \left(\mathcal{L}_{(\mathbf{E}_{\psi_0} | \delta\psi_T)} - i \hbar L_{\mathcal{R} \delta\psi_T} \right) G \quad (15)$$

the \mathcal{R} -regularized master Dyson–Schwinger (MDS) operator. Moreover, we call

we search for $(\psi_0, G_{\psi_0}) \in F \times A(\mathbb{F}(\mathbb{C}))$, such that :

$$b G_{\psi_0} = 1, \quad \text{and} \quad \forall \delta\psi_T \in \mathbb{F}_T : \mathbf{M}_{\hbar, \psi_0, \mathcal{R}, \delta\psi_T} G_{\psi_0} = 0 \quad (16)$$

the \mathcal{R} -regularized master Dyson–Schwinger (MDS) equation.

Clearly, the above definition is motivated by the Wilsonian regularization, heuristically stated in Eq.(7). If the base manifold \mathcal{M} were an affine space, in order to achieve a Wilsonian regularization (UV frequency damping), the regularizer operator \mathcal{R} should be taken to be the convolution operator by a test function on \mathcal{M} . It is not difficult to see that in such a setting, for the non-interacting case, the Wilsonian regularized MDS equation Eq.(16) does have solutions in the space of smooth field correlators. Thus, Definition 35 is expected to make sense also for interacting theories, since there is no problem with the diagonal evaluation map on the space of multivariate smooth functions. In order to adapt this construction to generic, non-affine manifolds \mathcal{M} , we invoke a notion of generalized convolution on smooth manifolds, see also [32, 33, 34].

Remark 36 Let $\mathcal{R} : \mathbb{F} \rightarrow \mathbb{F}$ be a continuous linear operator. Then, it can be uniquely extended as a continuous grading preserving algebra derivation $\mathcal{R} : \mathcal{T}(\mathbb{F}) \rightarrow \mathcal{T}(\mathbb{F})$ of the unital associative topological graded algebra $\mathcal{T}(\mathbb{F})$ via requiring the annihilation of unity ($\mathcal{R} \mathbb{1} = 0$), the preservation of the space of n -tensors ($n \in \mathbb{N}_0$), the Leibniz rule over tensor product, and coincidence with \mathcal{R} on the one-vectors. If \mathcal{R} is also preserving flavor sectors of \mathbb{F} , then it can be restricted to $A(\mathbb{F}(\mathbb{C})) \subset \mathcal{T}(\mathbb{F}(\mathbb{C}))$ as an algebra derivation. Similarly, the topological transpose operator $\mathcal{R}^* : \mathbb{F}^* \rightarrow \mathbb{F}^*$ extends as a continuous linear operator $\mathcal{R}^* : \mathcal{T}_a(\mathbb{F}^*) \rightarrow \mathcal{T}_a(\mathbb{F}^*)$. Assume moreover, that the pertinent operator \mathcal{R} on \mathbb{F} has a formal transpose $\mathcal{R}^t : \mathbb{F}_T^\times \rightarrow \mathbb{F}_T^\times$. Then, for the same reason it extends uniquely to $\mathcal{T}_a(\mathbb{F}_T^\times)$ in the above manner, and as $(\mathcal{R}^t)^*$ to $\mathcal{T}((\mathbb{F}_T^\times)^*)$, and thus also to $A((\mathbb{F}_T^\times(\mathbb{C}))^*) \subset \mathcal{T}((\mathbb{F}_T^\times(\mathbb{C}))^*)$, if \mathcal{R} was flavor sector preserving. The operator $(\mathcal{R}^t)^*$ will not be distinguished in notation from \mathcal{R} , since the former is the distributional extension of the latter. Similarly, \mathcal{R}^t will in general be denoted by \mathcal{R}^* , since the latter is the distributional extension of the former. Such an operator \mathcal{R} can also generate a corresponding continuous unital grading preserving algebra homomorphism $H_{\mathcal{R}}$ by acting as $\otimes^n \mathcal{R}$ on the subspaces of n -vectors. (If \mathcal{R} were injective with a continuous inverse, $H_{\mathcal{R}}$ would become an algebra automorphism.)

Remark 37 Following the terminologies are from the theory of pseudodifferential operators [32, 33, 34].

- (i) A continuous linear map $C : (\mathbb{F}^\times)^* \rightarrow \mathbb{F}$ is called a **smoothing operator**, their space is denoted by $\Psi^{-\infty}$ in the literature. By Schwartz kernel theorem, such an operator can be identified with an element $\kappa \in \mathbb{F} \otimes \mathbb{F}^\times$, i.e. κ is a smooth section of the vector bundle $U(\mathcal{M}) \boxtimes U^\times(\mathcal{M})$ over the base $\mathcal{M} \times \mathcal{M}$, with $U(\mathcal{M})$ being the vector bundle of field variations, i.e. $\mathbb{F} = \Gamma(U(\mathcal{M}))$. This is emphasized by writing C_κ instead, where $(C_\kappa \delta\psi_T)(x) = \int_{y \in \mathcal{M}} \kappa(x, y) \delta\psi_T(y)$ for all $\delta\psi_T \in \mathbb{F}_T \subset (\mathbb{F}^\times)^*$ and for all $x \in \mathcal{M}$.
- (ii) A smoothing operator C_κ is called **properly supported**, whenever the canonical projections from $\text{supp}(\kappa) \subset \mathcal{M} \times \mathcal{M}$ onto each factor \mathcal{M} is proper, i.e. the inverse images of compact sets are compact. In other words, for all compact subsets $\mathcal{K} \subset \mathcal{M}$ the closure of the sets $\{(x, y) \in \mathcal{M} \times \mathcal{M} \mid x \in \mathcal{K}, \kappa(x, y) \neq 0\}$ and $\{(x, y) \in \mathcal{M} \times \mathcal{M} \mid y \in \mathcal{K}, \kappa(x, y) \neq 0\}$ are compact. In that case, the map C_κ can act as continuous linear maps $\mathbb{F}_T \rightarrow \mathbb{F}_T$, $\mathbb{F} \rightarrow \mathbb{F}$, $(\mathbb{F}^\times)^* \rightarrow (\mathbb{F}^\times)^*$, $(\mathbb{F}_T^\times)^* \rightarrow (\mathbb{F}_T^\times)^*$.

Definition 38 Let C_κ be a properly supported smoothing operator as in Remark 37. If it preserves the flavor sectors, then κ is called a **mollifying kernel**.

Remark 39 A special example can shed some light on the role of κ in Definition 38. Let \mathcal{M} be a finite dimensional real affine space (“Minkowski spacetime”), the underlying finite dimensional real vector space denoted by T (“tangent space”). Let the vector bundle of fields be trivial, and trivialized compatibly with the affine structure. In that case, the fields, i.e. the elements of \mathbb{F} are simply smooth functions from \mathcal{M} to a finite dimensional real vector space, in which the classical fields take their values. Let us denote the identity operator of that finite dimensional real vector space by I . Due to the affineness of \mathcal{M} , up to a positive multiplier there exists a unique positive volume form field v which is parallel against the affine parallel transport (this corresponds to the Lebesgue measure). Take a compactly supported C^∞ real valued scalar field $\rho : T \rightarrow \mathbb{R}$. Then, the field $(x, y) \mapsto \kappa(x, y) := \rho(x-y) v(y) I$ is called a convolution kernel, and defines a mollifying kernel. For any element $\delta\psi \in \mathbb{F}$ one has that $C_\kappa \delta\psi = \rho \star \delta\psi$, i.e. C_κ is the convolution operator by ρ . Similarly, for any element $p \in \mathbb{F}^\times$ one has that $C_\kappa^t p = \rho^t \star p$, where ρ^t is the reflected ρ (for all $v \in T$ one has $\rho^t(v) := \rho(-v)$). Due to the compact support of ρ , the κ is indeed properly supported. Moreover, by construction, it is flavor sector preserving.

With the notion of mollifying kernel, one can define the Wilsonian regularization (UV frequency cutoff) also over generic manifolds. Namely, in the Definition 35, one sets $\mathcal{R} = C_\kappa$ for some mollifying kernel κ . In that case, we use the abbreviation $\mathbf{M}_{\hbar, \psi_0, \kappa, \delta\psi_T}$ for $\mathbf{M}_{\hbar, \psi_0, \mathcal{R}, \delta\psi_T} = \mathbf{M}_{\hbar, \psi_0, C_\kappa, \delta\psi_T}$. It is seen that the regularized MDS equation is the analogy of the unregularized MDS equation Eq.(13), but with a smoothing appearing in it.

Remark 40 Let $F, \mathbb{F}, \mathbb{F}_T, A(\mathbb{F}_{(\mathbb{C})}), E, \hbar$ as in Definition 35, and κ be a mollifying kernel, so that the κ -regularized MDS equation is defined, for fixed reference fields $\psi_0 \in F$. Let the EL functional $E : F \times \mathbb{F}_T \rightarrow \mathbb{R}$ be non-interacting as in the Definition 26, and choose $\psi_0 \in F$ to be EL solution, in which case $E_{\psi_0} : \mathbb{F} \rightarrow \mathbb{F}_T^*$ is linear. Let

$K_{\psi_0}^{(2)} \in \mathcal{L}(\mathbb{F}_T^\times, \mathbb{F}) \subset (\mathbb{F}_T^\times)_2^*$ be a fundamental solution (Green's functional) of the EL equation, with properties as in Remark 32. In that case, the κ -regularized fundamental solution $C_\kappa K_{\psi_0}^{(2)}$ resides in $\overset{2}{\otimes} \mathbb{F}$, and it is compatible with the permutation symmetry of the field algebra $A(\mathbb{F}_{(\mathbb{C})})$. Let the base manifold be affine, and let κ be specifically a convolution kernel by a symmetric test function. Then, $C_\kappa K_{\psi_0}^{(2)}$ satisfies $\forall \delta\psi_T \in \mathbb{F}_T : \mathcal{L}_{(\mathbf{E}_{\psi_0}|\delta\psi_T)} C_\kappa K_{\psi_0}^{(2)} = L_{C_\kappa \delta\psi_T} \mathbf{1}$. Define the element $K_{\psi_0, \kappa} := (0, 0, i\hbar C_\kappa K_{\psi_0}^{(2)}, 0, 0, \dots) \in A(\mathbb{F}_{(\mathbb{C})})$, called to be the connected correlator. Define the correlator with the ansatz $G_{\psi_0, \kappa} := \exp(K_{\psi_0, \kappa}) \in A(\mathbb{F}_{(\mathbb{C})})$. Then, $(\psi_0, G_{\psi_0, \kappa}) \in F \times A(\mathbb{F}_{(\mathbb{C})})$ solves the κ -regularized MDS equation Eq.(16). (In order to see this, one merely needs to repeat the proof of Remark 32.)

It is seen that in the above definition the trick is that although the fundamental solution $K_{\psi_0}^{(2)} \in \mathcal{L}(\mathbb{F}_T^\times, \mathbb{F}) \subset (\mathbb{F}_T^\times)_2^*$ is merely defined in the distributional sense, but its κ -regularized version $C_\kappa K_{\psi_0}^{(2)}$ sits in the space of fields $\mathbb{F} \otimes \mathbb{F}$. Therefore, free theories of such kind will have smooth solutions of the κ -regularized MDS equation, and therefore one does not need to go to the realm of distributional sense MDS equation, which is not applicable to interacting models. The replacement of the unregularized MDS operator with the κ -regularized MDS operator is called *regularization*. A further sanity check on the presented Wilsonian regularization scheme of the MDS equation via a mollifying kernel is the fact that such an equation would always have formal perturbative solutions if the field algebra were $\mathcal{T}(\mathbb{F}_{(\mathbb{C})})$, and the base manifold \mathcal{M} were affine. This can be seen to be an immediate consequence of the Malgrange–Ehrenpreis surjectivity theorem ([29] Chapitre I.1 Théorème 1 and [30] Chapter 6 Theorem 10), but will be expanded in a different paper.

Having a rigorous and generally covariant formulation of Wilsonian regularization at hand, one can naturally ask the question whether it is possible to formulate Wilsonian renormalization in a rigorous and generally covariant way. This we address below.

Remark 41 *It is rather straightforward to see that the space of mollifying kernels form a real vector space, naturally carrying a Hausdorff sequential convergence (CVS) structure which is Cauchy complete. A sequence $(\kappa_n)_{n \in \mathbb{N}}$ of mollifying kernels is said to converge to zero iff for all compact sets $\mathcal{K} \subset \mathcal{M}$ there exists some compact set $\mathcal{K}' \subset \mathcal{M}$, such that for all $n \in \mathbb{N}$ the closure of the sets $\{(x, y) \in \mathcal{M} \times \mathcal{M} \mid x \in \mathcal{K}, \kappa_n(x, y) \neq 0\}$ and $\{(x, y) \in \mathcal{M} \times \mathcal{M} \mid y \in \mathcal{K}, \kappa_n(x, y) \neq 0\}$ are contained in $\mathcal{K} \times \mathcal{K}'$ and $\mathcal{K}' \times \mathcal{K}$, respectively, moreover the sections $(\kappa_n)_{n \in \mathbb{N}}$ along with all their polynomial derivatives converge uniformly to zero over the compact sets $\mathcal{K} \times \mathcal{K}' \subset \mathcal{M} \times \mathcal{M}$ and $\mathcal{K}' \times \mathcal{K} \subset \mathcal{M} \times \mathcal{M}$. We do not address in the present note whether this convergence structure originates from a TVS structure or not, since we do not need it. It is also rather easy to see, that whenever the base manifold \mathcal{M} is affine, the convolution kernels form a sequentially closed vector subspace within the space of all mollifying kernels.*

Definition 42 *On the set of mollifying kernels, one may introduce a natural pre-ordering relation. Namely, for mollifying kernels κ'' and κ we say that $\kappa'' \lesssim \kappa$ (in words:*

κ'' is less ultraviolet than κ) iff either $C_{\kappa''} = C_{\kappa}$ or there exists some mollifying kernel κ' such that $C_{\kappa''} = C_{\kappa'} C_{\kappa}$ holds. It is evidently seen from the construction, that indeed this defines a pre-order, i.e. a relation which is transitive and reflexive. It is also seen that such relation may be also formulated on the set of convolution kernels, whenever convolution is meaningful, i.e. whenever the base manifold \mathcal{M} is affine.

Theorem 43 For a real valued smooth compactly supported test function φ_T over \mathcal{M} , denote by M_{φ_T} the multiplication operator by φ_T . The pre-order relation \lesssim , introduced in Definition 42, when restricted to the set of mollifying kernels κ which admit some φ_T such that $C_{\kappa} M_{\varphi_T}$ is not finite rank, becomes a partial order, i.e. it is antisymmetric. \square

Proof Let κ and κ' be any two mollifying kernels. We need to show that $\kappa' \lesssim \kappa$ and $\kappa \lesssim \kappa'$ implies $\kappa' = \kappa$ under the conditions of the theorem.

Writing out the condition $\kappa' \lesssim \kappa$ and $\kappa \lesssim \kappa'$ explicitly, there exist continuous linear operators $A, B : \mathbb{F} \rightarrow \mathbb{F}$, such that $C_{\kappa'} = A C_{\kappa}$ and $C_{\kappa} = B C_{\kappa'}$, where $A = I$ or $A = C_{\alpha}$ with some mollifying kernel α , and $B = I$ or $B = C_{\beta}$ with some mollifying kernel β . Putting these together, they imply $C_{\kappa'} = A B C_{\kappa'}$ and $C_{\kappa} = B A C_{\kappa}$. Taking any real valued compactly supported smooth test function φ_T over \mathcal{M} , these imply $C_{\kappa'} M_{\varphi_T} = A B C_{\kappa'} M_{\varphi_T}$ and $C_{\kappa} M_{\varphi_T} = B A C_{\kappa} M_{\varphi_T}$. Since κ and κ' was properly supported, then there exists some large enough compact region $\mathcal{K} \subset \mathcal{M}$ containing $\text{supp}(\varphi_T)$, such that the supports of the images of $C_{\kappa} M_{\varphi_T}$ and $C_{\kappa'} M_{\varphi_T}$ are also contained within \mathcal{K} . Let η_T be a real valued smooth compactly supported test function, which takes the value 1 within this set \mathcal{K} . Then, one has

$$C_{\kappa'} M_{\varphi_T} = A B M_{\eta_T} C_{\kappa'} M_{\varphi_T} \text{ and } C_{\kappa} M_{\varphi_T} = B A M_{\eta_T} C_{\kappa} M_{\varphi_T}. \quad (17)$$

One can choose an even larger compact region $\mathcal{K}' \subset \mathcal{M}$, which contains $\text{supp}(\eta_T)$ and also contains the supports of the images of $A B M_{\eta_T}$ and $B A M_{\eta_T}$. Under such conditions, the kernel function of $C_{\kappa'} M_{\varphi_T}$ and of $C_{\kappa} M_{\varphi_T}$ are square integrable, and therefore are Hilbert–Schmidt on the space of L^2 sections over \mathcal{K}' , so they are compact operators. If any of A or B is not the unit operator, then it is a mollifying operator by our assumptions, and then both $A B M_{\eta_T}$ and $B A M_{\eta_T}$ are also Hilbert–Schmidt on the above L^2 function space over \mathcal{K}' , for the same above reason, so they are also compact. Eq.(17) implies that $\text{Ran}(C_{\kappa'} M_{\varphi_T})$ is contained in the eigensubspace of $A B M_{\eta_T}$ with eigenvalue one, and $\text{Ran}(C_{\kappa} M_{\varphi_T})$ is contained in the eigensubspace of $B A M_{\eta_T}$ with eigenvalue one. But since nonzero eigenvalue eigensubspaces of compact operators are finite dimensional, $\text{Ran}(C_{\kappa'} M_{\varphi_T})$ and $\text{Ran}(C_{\kappa} M_{\varphi_T})$ must be finite dimensional if any of A or B are not the unity operator. But it was assumed that κ admitted some φ_T such that $\text{Ran}(C_{\kappa} M_{\varphi_T})$ is not finite dimensional (and for κ' the same was assumed with some φ'_T). Therefore, both A and B must be the unity operator, i.e. $\kappa' = \kappa$. \blacksquare

Remark 44 The pre-ordering \lesssim becomes a partial order in the following sense.

- (i) If for some test function φ_T the mollifying kernel κ is such that $C_{\kappa} M_{\varphi_T}$ is injective on an infinite dimensional linear subspace of the L^2 sections, then $C_{\kappa} M_{\varphi_T}$ is not

finite rank. That is because in the pertinent case, the L^2 adjoint of the continuous operator $C_\kappa M_{\varphi_T}$ is evidently non-finite rank, due to which the operator itself cannot be finite rank.

- (ii) If the mollifying kernel κ is such that the operator C_κ is injective over the space of test field variations \mathbb{F}_T , then it satisfies the above condition, and thus $C_\kappa M_{\varphi_T}$ is not finite rank, for any test function φ_T .
- (iii) If the base manifold \mathcal{M} is affine, then the convolutions are meaningful, and the convolution kernels by test functions are such that their C_κ operators are injective over the space of test field variations \mathbb{F}_T . That claim can be verified in Fourier space, using a consequence of the Paley–Wiener–Schwartz theorem ([31] Theorem 7.3.1), namely the fact that the Fourier transform of a compactly supported distribution (and hence, of a function) is an analytic function. (Alternatively, it also follows from [35] Theorem 4.4.) Therefore, if κ were a convolution kernel, by means of the above observation, $C_\kappa M_{\varphi_T}$ is not finite rank.

The above leads us to the following conclusion.

Corollary 45 *On the set of mollifying kernels which are injective on the space of test field variations, the pre-ordering \lesssim is antisymmetric, i.e. it is a partial order.*

In particular, when the base manifold is affine, over the set of nonvanishing convolution kernels the pre-ordering \lesssim is a partial order.

In such cases, we may use the symbol \preceq instead of \lesssim for clarity. □

Remark 46 In Section 2 it was argued that the Wilsonian regularization justifies our regularized MDS equation Eq.(16). Applying the heuristic integral substitution (measure pushforward) formula for composite maps in the Wilsonian Feynman integral Eq.(6), it would follow that if $(\psi_0, G_{\psi_0, \kappa}) \in F \times A(\mathbb{F}_{(\mathbb{C})})$ were a solution of the κ -regularized MDS equation, and $\kappa'' \lesssim \kappa$, then there should exist a solution $(\psi_0, G_{\psi_0, \kappa''}) \in F \times A(\mathbb{F}_{(\mathbb{C})})$ of the κ'' -regularized MDS equation, such that the identity

$$H_{C_{\kappa'}} G_{\psi_0, \kappa} = G_{\psi_0, \kappa''} \tag{18}$$

holds, where κ' is the corresponding mollifying kernel satisfying $C_{\kappa''} = C_{\kappa'} C_\kappa$ (because of $\kappa'' \lesssim \kappa$), and $H_{C_{\kappa'}}$ is the unital algebra homomorphism of $A(\mathbb{F}_{(\mathbb{C})})$ generated by the continuous linear operator $C_{\kappa'} : \mathbb{F} \rightarrow \mathbb{F}$. This equation is called the exact renormalization equation (ERGE) in the QFT literature, and $H_{C_{\kappa'}}$ is called a blocking transformation. It is seen that if Feynman integrals existed as a proper finite measure, the ERGE equation would be just the consequence of the fundamental formula for integral substitution, for the pushforward measures. In our rigorous formalism, defined on the field correlators, one needs to impose that by hand, as stated below.

Definition 47 *Let the index set \mathcal{I} be the set of mollifying kernels, and denote by $A(\mathbb{F}_{(\mathbb{C})})^{\mathcal{I}}$ the set of all maps $\mathcal{I} \rightarrow A(\mathbb{F}_{(\mathbb{C})})$. Then, the **solution space of the Wilsonian renormalized MDS equation** is*

$$Q_r := \left\{ (\psi_0, G_{\psi_0, \cdot}) \in F \times A(\mathbb{F}_{(\mathbb{C})})^{\mathcal{I}} \mid \forall \kappa, \kappa'' \in \mathcal{I} : \kappa'' \lesssim \kappa \text{ (with } \kappa') \Rightarrow H_{C_{\kappa'}} G_{\psi_0, \kappa} = G_{\psi_0, \kappa''} \right.$$

$$\text{and } \forall \kappa \in \mathcal{I} : \left. \forall \delta\psi_T \in \mathbb{F}_T : b G_{\psi_0, \kappa} = 1, \mathbf{M}_{\hbar, \psi_0, \kappa, \delta\psi_T} G_{\psi_0, \kappa} = 0 \right\}, \quad (19)$$

*i.e. they are the solution families of the regularized MDS equation, satisfying the ERGE relation. We say that a model is **Wilsonian renormalizable**, if Q_r is not empty. (One may use the notation Q_{r, ψ_0} for the subset when the reference field is fixed to $\psi_0 \in F$.)*

One may recognize that the solution families satisfying the ERGE relation are so-called projective families, and therefore, the solution space of the Wilsonian renormalized MDS equation is the corresponding projective limit ([36] Chapter 4.21). The theory is Wilsonian renormalizable whenever the corresponding projective limit exists as a nonempty set.

Remark 48 *It is not uncommon in QFT that running coupling factors need to be introduced. In that case, it is assumed that the EL function can be specified as a finite sum $E = g_1 E_1 + \dots + g_n E_n$, with each $E_i : F \times \mathbb{F}_T \rightarrow \mathbb{R}$ being jointly sequentially continuous, called the Euler–Lagrange terms, and g_i being nonzero real numbers, called to be the coupling factors ($i = 1, \dots, n$). Recall that the space of mollifying kernels \mathcal{I} was a Hausdorff complete sequential convergence vector space, due to which one can define (sequentially) continuous functions from \mathcal{I} to other convergence vector spaces. Given some (sequentially) continuous functionals $\gamma_i : \mathcal{I} \rightarrow \mathbb{R}$ ($i = 1, \dots, n$), one may define the running regularized MDS operator as*

$$\begin{aligned} \mathbf{M}_{\hbar, \psi_0, \kappa, (\gamma_1, \dots, \gamma_n), \delta\psi_T} : A(\mathbb{F}_{(\mathbb{C})}) &\rightarrow A(\mathbb{F}_{(\mathbb{C})}), \\ G \mapsto \mathbf{M}_{\hbar, \psi_0, \kappa, (\gamma_1, \dots, \gamma_n), \delta\psi_T} G &:= \left(\mathcal{L}_{\gamma_1(\kappa)(\mathbf{E}_{1, \psi_0} | \delta\psi_T) + \dots + \gamma_n(\kappa)(\mathbf{E}_{n, \psi_0} | \delta\psi_T)} - i \hbar L_{C_\kappa \delta\psi_T} \right) G, \end{aligned} \quad (20)$$

for fixed $\hbar \in \mathbb{R}$, reference field $\psi_0 \in F$, test field variation $\delta\psi_T \in \mathbb{F}_T$, mollifying kernel $\kappa \in \mathcal{I}$ and running couplings γ_i ($i = 1, \dots, n$). The solution space of the Wilsonian renormalized MDS equation with running couplings is then

$$\begin{aligned} &\left\{ (\psi_0, (\gamma_1, \dots, \gamma_n), G_{\psi_0, \cdot}) \in F \times C(\mathcal{I}, \mathbb{R})^n \times A(\mathbb{F}_{(\mathbb{C})})^{\mathcal{I}} \mid \right. \\ &\quad \left. \forall \kappa, \kappa'' \in \mathcal{I} : \kappa'' \preceq \kappa (\text{with } \kappa') \Rightarrow H_{C_{\kappa'}} G_{\psi_0, \kappa} = G_{\psi_0, \kappa''} \right. \\ &\quad \left. \text{and } \forall \kappa \in \mathcal{I} : \forall \delta\psi_T \in \mathbb{F}_T : b G_{\psi_0, \kappa} = 1, \mathbf{M}_{\hbar, \psi_0, \kappa, (\gamma_1, \dots, \gamma_n), \delta\psi_T} G_{\psi_0, \kappa} = 0 \right\}. \end{aligned} \quad (21)$$

6. An existence condition for regularized MDS solutions

In this section, we present an existence condition for the solutions of the Wilsonian regularized MDS equation.

Remark 49 *The followings spell out some facts about the topology of \mathbb{F} and \mathbb{F}_T .*

- (i) *Assume that the base manifold \mathcal{M} underlying a concrete classical field theory as in Definition 10 is compact (with or without boundary, and if with boundary, we assume the cone condition). This is a realistic assumption for conformally invariant models, as for those, the theory can be reformulated on the compact manifold with boundary, underlying the conformally compactified spacetime (which will admit cone condition).*

- (ii) With the assumption as above, rather obviously, the space \mathbb{F}_T shall become also an NF space, similarly to \mathbb{F} . That is, \mathbb{F}_T becomes metrizable with all of its benefits: its topology will be sequential, and separately continuous multilinear maps from it will become jointly continuous. (Recall that if \mathcal{M} is compact then, either $\mathbb{F}_T = \mathbb{F}$, or \mathbb{F}_T may be chosen to be the closed subspace of \mathbb{F} consisting of fields vanishing at $\partial\mathcal{M}$ together with all of their derivatives. Its natural topology will become an \mathcal{E} function topology instead of \mathcal{D} type.)
- (iii) It is also an elementary fact that over a compact base manifold \mathcal{M} , the space \mathbb{F} and \mathbb{F}_T will not only be nuclear Fréchet, but also will admit continuous norms instead of merely continuous seminorms. By means of Remark 63, then they become nuclear Fréchet spaces with a countable increasing system of topology defining Hilbertian norms. Since these norms are simply Sobolev norms, they are Gel'fand compatible (see Remark 63 and Remark 64), therefore by means of Remark 63 they become NF spaces with the countably Hilbert (CH) property. Recall that if \mathcal{H} is a CH type NF space, then there exists a countable family

$$H_0 \supset H_1 \supset \dots \supset H_n \supset \dots$$

of topological vector spaces, such that $\mathcal{H} = \bigcap_{n \in \mathbb{N}_0} H_n$ and is dense in all of the spaces H_n ($n \in \mathbb{N}_0$), moreover their topologies are gradually strictly strengthening

$$\tau_0|_{\mathcal{H}} \subset \tau_1|_{\mathcal{H}} \subset \dots \subset \tau_n|_{\mathcal{H}} \subset \dots,$$

and all of their topologies are complete and generated by a Hilbertian scalar product (that is, for each $n \in \mathbb{N}_0$, the space H_n can be taken to be a Hilbert space), and for all $n \in \mathbb{N}_0$ there exists an integer $m \geq 1$ such that the inclusion maps $i_{n+m,n} : H_{n+m} \rightarrow H_n$ are nuclear. (Specially, the spaces H_0, H_1, \dots may be chosen such that all adjacent inclusion maps are nuclear.) The respective topology generating Hilbertian norms thus form an increasing system

$$\|\cdot\|_0 \leq C_{0,1} \|\cdot\|_1 \leq \dots \leq C_{n-1,n} \|\cdot\|_n \leq \dots$$

(for some $C_{0,1}, \dots, C_{n-1,n}, \dots \in \mathbb{R}^+$).

The corresponding Hilbertian scalar products shall be denoted by

$$\langle \cdot, \cdot \rangle_0, \langle \cdot, \cdot \rangle_1, \dots, \langle \cdot, \cdot \rangle_n, \dots$$

The proposed existence theorem will hinge on the fact that on the field algebra $A(\mathbb{F}_{(\mathbb{C})})$ it is possible to naturally define a reasonable topology, somewhat stronger than the Tychonoff topology, such that it preserves the NF property coming from \mathbb{F} , and if present, the eventual CH property of \mathbb{F} as well.

Remark 50 We recall some findings on topologies of the tensor algebra of \mathbb{F} [37].

- (i) Dubin and Hennings in their work ([37] Chapter 3.1) introduces the notion of tensor algebra topology of the following kind. Let \mathbb{F} be a nuclear Fréchet space. Then, they define the vector space $\mathcal{T}(\mathbb{F}, \lambda, \mathfrak{p})$, where λ is some topological subspace of

the space of $\mathbb{N}_0 \rightarrow \mathbb{C}$ sequences (it is a so-called Köthe echelon space), and \mathfrak{p} is a family of Hilbertian seminorms on \mathbb{F} defining its NF topology (recall that multiple seminorm families \mathfrak{p} can define the same topology on \mathbb{F}). As a vector space, it is defined as follows:

$$\mathcal{T}(\mathbb{F}, \lambda, \mathfrak{p}) := \left\{ G \in \bigoplus_{n=0}^{\infty} \otimes^n \mathbb{F} \mid \forall \|\cdot\| \in \mathfrak{p} : (\|G^{(n)}\|^{\otimes n})_{n \in \mathbb{N}_0} \in \lambda \right\},$$

where for all topology defining Hilbertian seminorms $\|\cdot\| \in \mathfrak{p}$ on \mathbb{F} , the symbol $\|\cdot\|^{\otimes n}$ denotes the n -fold cross norm over $\otimes^n \mathbb{F}$ originating from $\|\cdot\|$, which is then also a Hilbertian seminorm ([37] Chapter3.1). The locally convex vector topology on $\mathcal{T}(\mathbb{F}, \lambda, \mathfrak{p})$ is defined by the system of seminorms $(G \mapsto \sum_{n=0}^{\infty} |u_n| \|G^{(n)}\|^{\otimes n})_{u \in \lambda^\times, \|\cdot\| \in \mathfrak{p}}$, where λ^\times denotes the so-called Köthe dual (which is, under, mild conditions, the strong topological dual) of the sequence space λ .

(ii) Notable Köthe echelon spaces include ([37] Chapter2.4):

$$\begin{aligned} \phi &:= \{(u_n)_{n \in \mathbb{N}_0} \in \mathbb{C}^{\mathbb{N}_0} \mid \exists m \in \mathbb{N}_0 : \forall n > m : u_n = 0\} \quad (\text{terminating sequences}), \\ h &:= \{(u_n)_{n \in \mathbb{N}_0} \in \mathbb{C}^{\mathbb{N}_0} \mid \forall m \in \mathbb{N}_0 : \exists C_m \in \mathbb{R}^+ : \forall n \in \mathbb{N}_0 : 2^{nm} |u_n| \leq C_m\}, \\ h' &:= \{(u_n)_{n \in \mathbb{N}_0} \in \mathbb{C}^{\mathbb{N}_0} \mid \exists c \in \mathbb{N} : \exists m_c \in \mathbb{N}_0 : \forall n \in \mathbb{N}_0 : |u_n| \leq c 2^{nm_c}\}, \\ \omega &:= \{(u_n)_{n \in \mathbb{N}_0} \in \mathbb{C}^{\mathbb{N}_0} \mid \text{any } u\} \quad (\text{all sequences}). \end{aligned}$$

These are understood with their so-called normal topologies ([37] Chapter2.1). The space ω is the space of all sequences with the natural Tychonoff topology, the space ϕ is the space of finitely terminating sequences with the natural locally convex direct sum topology, whereas the space h is known to be topologically isomorphic to the space $H(\mathbb{C})$ of entire complex functions ([37] p.978). All of them are Hausdorff locally convex topological vector spaces, and the pairs (ϕ, ω) , (h, h') , (h', h) , (ω, ϕ) are strong dual to each-other, and Köthe duals to each-other. The spaces ϕ , h , ω are metrizable, and thus Fréchet. The spaces ϕ , h , ω have the so-called “h” property, because of which they are nuclear ([37] Chapter2.5). Therefore, ϕ , h , ω are NF, and h' is DNF. Specially, the space h also has the countably Hilbert property.

- (iii) It is shown in [37] Chapter3.3 that specially for the sequence spaces $\lambda = \phi$ or $\lambda = h$ or $\lambda = \omega$, the tensor algebra $\mathcal{T}(\mathbb{F}, \lambda, \mathfrak{p})$ is independent of the choice of the representant \mathfrak{p} of the topology defining Hilbertian seminorms on \mathbb{F} , thus one may write merely $\mathcal{T}(\mathbb{F}, \lambda)$ instead. Moreover, they inherit the NF property of \mathbb{F} , making $\mathcal{T}(\mathbb{F}, \lambda)$ a unital associative algebra with jointly continuous multiplication, with NF topology.
- (iv) It is rather easy to see that $\mathcal{T}(\mathbb{F}, \phi)$ is simply the algebraic tensor algebra $\mathcal{T}_a(\mathbb{F})$ with its natural locally convex direct sum topology, $\mathcal{T}(\mathbb{F}, \omega)$ is simply the Tychonoff tensor algebra $\mathcal{T}(\mathbb{F})$ with its natural Tychonoff topology.
- (v) For λ of the above types, it is shown in [37] Chapter3.3 that the topology defining

seminorms on $\mathcal{T}(\mathbb{F}, \lambda)$ may be chosen to be Hilbertian seminorms

$$G \mapsto \sqrt{\sum_{n=0}^{\infty} |u_n|^2 (\|G^{(n)}\|^{\otimes n})^2} \quad (u \in \lambda^\times, \|\cdot\| \in \mathfrak{p}), \quad (22)$$

where \mathfrak{p} is a representant of a topology defining family of Hilbertian seminorms on \mathbb{F} . From this, it is explicitly seen that whenever \mathbb{F} is an NF space admitting a continuous Hilbertian norm, then the NF space $\mathcal{T}(\mathbb{F}, h)$ also admits a continuous Hilbertian norm.

- (vi) The explicit form of a representant of a topology defining countable family of increasing Hilbertian seminorms on $\mathcal{T}(\mathbb{F}, h)$, encoding its NF topology, can be given by:

$$G \mapsto \sqrt{\sum_{n=0}^{\infty} 2^{m n} (\|G^{(n)}\|_m^{\otimes n})^2} \quad (m \in \mathbb{N}_0), \quad (23)$$

where $\|\cdot\|_0, \|\cdot\|_1, \dots$ is a representant of a topology defining countable system of increasing Hilbertian seminorms on \mathbb{F} , defining its NF topology. (The formula is the consequence of [37] Proposition 3.7, but is also explicitly used in [38].) From the above formula it is seen that whenever \mathbb{F} is of CH type, then $\mathcal{T}(\mathbb{F}, h)$ is also of CH type (see also Remark 63). We will use the abbreviation $\mathcal{T}_h(\mathbb{F}) := \mathcal{T}(\mathbb{F}, h)$, and will call it the analytic tensor algebra of \mathbb{F} , since the topology defining sequence space h is isomorphic to the space of entire functions $H(\mathbb{C})$.

- (vii) From Eq.(23) it is trivially read off that the counit map $b : \mathcal{T}_h(\mathbb{F}) \rightarrow \mathbb{R}, G \mapsto bG := G^{(0)}$ is continuous. Therefore, the corresponding projection operator $\mathbb{1} b$ onto the scalar sector and its complement $I - \mathbb{1} b$ is also continuous. Moreover, the pertinent complementing projection operators $\mathbb{1} b$ and $I - \mathbb{1} b$ are orthogonal projections with respect to the representants of Hilbertian sesquilinear forms from Eq.(23), and $bG = \langle \mathbb{1}, G \rangle_m$ holds (for all $G \in \mathcal{T}_h(\mathbb{F})$ and all $m \in \mathbb{N}_0$).
- (viii) From the Eq.(23) form of the Hilbertian seminorms on $\mathcal{T}_h(\mathbb{F})$ it is seen that this representant family has the property that whenever \mathbb{F} is CH type NF space, then whenever its representant Hilbertian norm family is chosen to be such that the adjacent norms are nuclear against each-other, then the adjacent Hilbertian norms defined by Eq.(23) are also. Similarly, whenever the adjacent norms on \mathbb{F} are Hilbert-Schmidt against each-other, then the adjacent norms Eq.(23) are also Hilbert-Schmidt against each-other.

In order to state our existence condition for the solutions of the regularized MDS equation, we will need to reconsider the space of field correlators to be based on $\mathcal{T}_h(\mathbb{F})$ with the analytic topology, and not on $\mathcal{T}(\mathbb{F})$ with the Tychonoff direct sum topology. The reason is that for the construction to work, we need the eventual CH property of \mathbb{F} to be inherited by its tensor algebra. Therefore, from this point on, the field algebra $A(\mathbb{F}_{(\mathbb{C})})$ will be defined to be the appropriately symmetrized factor algebra (or subspace)

of $\mathcal{T}_h(\mathbb{F}_{(\mathbb{C})})$ instead of $\mathcal{T}(\mathbb{F}_{(\mathbb{C})})$ (see again Remark 20 and Definition 21 for the technical construction of the symmetrized algebra).

Remark 51 *It is worth to verify at this point whether the tensor algebra topology on the new field algebra $A(\mathbb{F}_{(\mathbb{C})})$, inherited from $\mathcal{T}_h(\mathbb{F}_{(\mathbb{C})})$, is not overly strict. For instance, one would like a typical solution of the regularized MDS equation for a non-interacting theory to be not excluded from our new, smaller field algebra $A(\mathbb{F}_{(\mathbb{C})})$. By recalling Remark 40, one sees from Eq.(23) that the pertinent existent solution of the regularized MDS equation for a non-interacting theory indeed resides in the new, stricter field algebra.*

In order to state an existence condition, we shall assume, like in Remark 49, that the base manifold \mathcal{M} under the concrete theory is compact (with or without boundary, and if with boundary, we assume the cone condition, so that Sobolev and Maurin theorems hold, see Remark 64). As stated before, this can be a realistic assumption in a conformally invariant theory, in which case the theory can be defined over a compact manifold with boundary (the conformal compactification of the would-be-spacetime).

Remark 52 *Assume that the base manifold \mathcal{M} of the model is compact and its boundary, if not empty, has the cone condition. Then, the followings hold.*

- (i) *With such assumption, \mathbb{F} and \mathbb{F}_T become countably Hilbert type NF spaces, which is then inherited by $\mathcal{T}_h(\mathbb{F}_{(\mathbb{C})})$, and thus by the field algebra $A(\mathbb{F}_{(\mathbb{C})})$. From now on, let us use the abbreviation $\mathcal{H} := A(\mathbb{F}_{(\mathbb{C})})$.*
- (ii) *In its original definition, the regularized MDS operator was a separately continuous bilinear map $\mathbf{M}_{\psi_0} : \mathcal{H} \times \mathbb{F}_T \rightarrow \mathcal{H}$, $(G, \delta\psi_T) \mapsto \mathbf{M}_{\psi_0, \delta\psi_T} G$ (see also the original definition Eq.(15), we suppress the fixed mollifying kernel in the notation in this chapter). Due to the compactness assumption on \mathcal{M} , the space \mathbb{F}_T becomes also metrizable, therefore the map \mathbf{M}_{ψ_0} becomes jointly continuous (Remark 59). Therefore, the regularized MDS operator, may be viewed as a continuous linear map $\mathbf{M}_{\psi_0} : \mathcal{H} \otimes \mathbb{F}_T \rightarrow \mathcal{H}$.*
- (iii) *Due to our compactness assumption on the base manifold \mathcal{M} , both \mathbb{F}_T and \mathcal{H} became countably Hilbert NF spaces, which technically means that in both spaces as well as on their tensor product, the properties Remark 49(iii) hold. Denote an associated chain of Hilbert spaces subordinate to \mathcal{H} by H_0, H_1, \dots , their Hilbertian norms by $\|\cdot\|_0, \|\cdot\|_1, \dots$ and their Hilbertian scalar products by $\langle \cdot, \cdot \rangle_0, \langle \cdot, \cdot \rangle_1, \dots$. Similarly, for \mathbb{F}_T denote by F_0, F_1, \dots an associated chain of Hilbert spaces, their corresponding Hilbertian norms by $\|\cdot\|_0^F, \|\cdot\|_1^F, \dots$, and their Hilbertian scalar products by $\langle \cdot, \cdot \rangle_0^F, \langle \cdot, \cdot \rangle_1^F, \dots$. The associated chain of Hilbert spaces subordinate to $\mathcal{H} \otimes \mathbb{F}_T$ can be taken to be the Hilbert–Schmidt tensor product of the spaces $H_m \otimes_{\text{HS}} F_n$ ($m, n \in \mathbb{N}_0$), with their canonical Hilbertian cross-norms and crossed Hilbertian scalar products. (Eventually, a subfamily of this, with strictly growing norms may also be considered instead.)*

(iv) Because of the nuclearity of the spaces \mathcal{H} and \mathbb{F}_T , each Hilbertian norm in the above chains will have a stronger norm in the chain for which the embedding map becomes Hilbert–Schmidt, and eventually becomes nuclear, for large enough norms in the chain. (This can also be seen less abstractly on our concrete spaces as a consequence of the Maurin embedding theorem Remark 64(ii).)

(v) The continuity of the linear map $\mathbf{M}_{\psi_0} : \mathcal{H} \otimes \mathbb{F}_T \rightarrow \mathcal{H}$ in terms of these Hilbert space chains means that

$$\forall k \in \mathbb{N}_0 : \exists m_k, n_k \in \mathbb{N}_0 : \exists C_{k,m_k,n_k} \in \mathbb{R}^+ : \\ \forall (G, \delta\psi_T) \in G \times \mathbb{F}_T : \quad \|\mathbf{M}_{\psi_0}(G \otimes \delta\psi_T)\|_k \leq C_{k,m_k,n_k} \|G\|_{m_k} \|\delta\psi_T\|_{n_k} \quad (24)$$

holds. Since the norms were ordered, the above identity implies that once it holds, it holds for all $m \geq m_k$ and $n \geq n_k$ as well with some constants $C_{k,m,n} \in \mathbb{R}^+$. That is, the map \mathbf{M}_{ψ_0} is a continuous linear map $(\mathcal{H} \otimes \mathbb{F}_T) \cap (H_m \otimes_{\text{HS}} F_n) \rightarrow \mathcal{H} \cap H_k$ for large enough indices $m, n \in \mathbb{N}_0$, given the index $k \in \mathbb{N}_0$. The continuous extension of the map \mathbf{M}_{ψ_0} will be denoted by the same symbol for brevity, and it is then a continuous linear map $\mathbf{M}_{\psi_0} : H_m \otimes_{\text{HS}} F_n \rightarrow H_k$, for such indices.

(vi) By means of (iv), between distant enough indices, the inclusion maps $H_m \supset \dots \supset H_{m'}$ ($m < m'$) and $F_n \supset \dots \supset F_{n'}$ ($n < n'$) become Hilbert–Schmidt, and eventually become nuclear. Therefore, given $k \in \mathbb{N}_0$, for large enough indices $m, n \in \mathbb{N}_0$ the map $\mathbf{M}_{\psi_0} : H_m \otimes_{\text{HS}} F_n \rightarrow H_k$ becomes Hilbert–Schmidt, and eventually becomes nuclear. (In concrete spaces, Maurin embedding theorem gives the concrete index bounds, see Remark 64(ii).)

(vii) As a particular case of the above statement, for all large enough indices $m, n \in \mathbb{N}_0$ one has that the linear map $\mathbf{M}_{\psi_0} : H_m \otimes_{\text{HS}} F_n \rightarrow H_0$ is Hilbert–Schmidt. The adjoint of this map $\mathbf{M}_{\psi_0}^\dagger : H_0 \rightarrow H_m \otimes_{\text{HS}} F_n$ is then also Hilbert–Schmidt. Therefore, the operator $\mathbf{M}_{\psi_0}^\dagger \mathbf{M}_{\psi_0} : H_m \otimes_{\text{HS}} F_n \rightarrow H_m \otimes_{\text{HS}} F_n$, becomes a positive nuclear (trace class) operator.

(viii) Fix a complete orthonormal basis $(e_i)_{i \in I}$ in F_n (since F_n is separable, one may set $I \equiv \mathbb{N}$). Then, for all $G \in H_m$ the estimate

$$B(G, G) := \sum_{i \in I} \left\langle (G \otimes_{\text{HS}} e_i), \mathbf{M}_{\psi_0}^\dagger \mathbf{M}_{\psi_0} (G \otimes_{\text{HS}} e_i) \right\rangle_{H_m \otimes_{\text{HS}} F_n} < \infty \quad (25)$$

is valid. That is because of the Hilbert–Schmidt property of the map $\mathbf{M}_{\psi_0} : H_m \otimes_{\text{HS}} F_n \rightarrow H_0$. Namely, for some (and therefore: for any) complete orthonormal basis $(g_j)_{j \in J}$ in H_m , one has that $\sum_{j \in J} \sum_{i \in I} \|\mathbf{M}_{\psi_0}(g_j \otimes_{\text{HS}} e_i)\|_{H_m \otimes_{\text{HS}} F_n}^2 < \infty$ holds (one may set $J \equiv \mathbb{N}$ as well, due to the separability of H_m). Taking specially an orthonormal basis $(g_j)_{j \in J}$ in H_m , such that one of its elements is $G/\|G\|_{H_m}$, one infers that indeed the estimate Eq.(25) holds.

(ix) Given $G \in H_m$, the corresponding expression Eq.(25) is independent of the chosen complete orthonormal basis $(e_i)_{i \in I}$ in F_n . That is because for a Hilbert–Schmidt operator A and an unitary operator U in a Hilbert space, one has that the Hilbert–Schmidt norm of A and $U^\dagger A U$ is the same.

- (x) Due to the Hilbert–Schmidt property of $\mathbf{M}_{\psi_0} : H_m \otimes_{\text{HS}} F_n \rightarrow H_0$, the quadratic form $H_m \rightarrow \mathbb{C}$, $G \mapsto B(G, G)$ is continuous, and therefore by the polarization formula it gives rise to a corresponding continuous sesquilinear form $H_m \times H_m \rightarrow \mathbb{C}$, $(G_1, G_2) \mapsto B(G_1, G_2)$. Therefore, by Riesz representation theorem, there is a corresponding unique continuous linear map $\hat{\mathbf{M}}_{\psi_0}^2 : H_m \rightarrow H_m$, such that for all $G_1, G_2 \in H_m$, the identity $B(G_1, G_2) = \langle G_1, \hat{\mathbf{M}}_{\psi_0}^2 G_2 \rangle_{H_m}$ holds.
- (xi) Due to the positive semidefiniteness of B , the map $\hat{\mathbf{M}}_{\psi_0}^2$ is a positive operator. Moreover, due to the Hilbert–Schmidt property of $\mathbf{M}_{\psi_0} : H_m \otimes_{\text{HS}} F_n \rightarrow H_0$, the map $\hat{\mathbf{M}}_{\psi_0}^2$ is nuclear (trace class). The nuclear (trace) norm of $\hat{\mathbf{M}}_{\psi_0}^2 : H_m \rightarrow H_m$, by construction, equals to the Hilbert–Schmidt norm of $\mathbf{M}_{\psi_0} : H_m \otimes_{\text{HS}} F_n \rightarrow H_0$. One can see that the operator $\hat{\mathbf{M}}_{\psi_0}^2$ is simply the absolute value squared version of the MDS operator, with its \mathbb{F}_T variable traced out.
- (xii) It is obvious from the construction that $\bigcap_{\delta\psi_T \in \mathbb{F}_T} \text{Ker}(\mathbf{M}_{\psi_0, \delta\psi_T}) = \text{Ker}(\hat{\mathbf{M}}_{\psi_0}^2)$.

If the reference field $\psi_0 \in F$ was chosen to be such that it satisfies the EL equation Eq.(11), then one has that $\mathbf{M}_{\psi_0, \delta\psi_T} \mathbf{1} = -i L_{\delta\psi_T} \mathbf{1} \ (\forall \delta\psi_T \in \mathbb{F}_T)$. Because of that, in this situation, $b \hat{\mathbf{M}}_{\psi_0}^2 \mathbf{1} = \langle \mathbf{1}, \hat{\mathbf{M}}_{\psi_0}^2 \mathbf{1} \rangle_m > 0$, and therefore $\mathbf{1} \notin \text{Ker}(\hat{\mathbf{M}}_{\psi_0}^2)$. Thus, generally, the trivial correlator $\mathbf{1}$ cannot be a solution of the strong or regularized MDS equation. One could still aim to find a projection of $\mathbf{1}$ which (up to normalization) satisfies the regularized MDS equation. Let us denote the orthogonal projection onto $\text{Ker}(\hat{\mathbf{M}}_{\psi_0}^2)$ in H_m by P . Then, $\text{Ker}(\hat{\mathbf{M}}_{\psi_0}^2) = \text{Ran}(P)$. One can state the following theorem on $P\mathbf{1}$.

Theorem 53 *Let P denote the orthoprojection in H_m onto $\text{Ker}(\hat{\mathbf{M}}_{\psi_0}^2)$. Then, the following statements are equivalent.*

- (i) *The solution space of the regularized MDS equation in H_m is not empty.*
- (ii) *One has that $P\mathbf{1} \neq 0$.*
- (iii) *One has that $bP\mathbf{1} \neq 0$.* □

Proof By construction, the MDS equation has solutions in H_m if and only if $b \text{Ker}(\hat{\mathbf{M}}_{\psi_0}^2) \neq \{0\}$, i.e. if and only if there exists some $G \in H_m$, such that $bPG \neq 0$. (That is because $\text{Ker}(\hat{\mathbf{M}}_{\psi_0}^2) = \text{Ran}(P)$ and because Remark 52(xii).) However, the identity $bPG = \langle \mathbf{1}, PG \rangle_m = \langle P\mathbf{1}, G \rangle_m$ holds, because of Remark 50(vii), and because P was orthoprojection in H_m . Therefore, (i) \Leftrightarrow (ii).

Moreover, one has that $\langle P\mathbf{1}, P\mathbf{1} \rangle_m = \langle \mathbf{1}, P\mathbf{1} \rangle_m = bP\mathbf{1}$, since P was an orthoprojection in H_m , and because of Remark 50(vii). Therefore, (ii) \Leftrightarrow (iii). ■

It is seen that the orthoprojection P in H_m onto $\text{Ker}(\hat{\mathbf{M}}_{\psi_0}^2)$ plays an important role in the problematics of existence of MDS solutions. One can approximate P as below.

Theorem 54 *For all $T > 0$ parameter, which is not smaller than the operator norm of $\hat{\mathbf{M}}_{\psi_0}^2$, one has that with the notation $\mathcal{P} := I - T^{-1} \hat{\mathbf{M}}_{\psi_0}^2$, the operator sequence $k \mapsto \mathcal{P}^k$ converges strongly (pointwise) to P in H_m .*

Proof The operator $\mathcal{P} = I - T^{-1} \hat{\mathbf{M}}_{\psi_0}^2$ is a positive continuous operator with spectrum in $[0, 1]$. Therefore, $k \mapsto \mathcal{P}^k$ is a monotonically decreasing sequence of such operators, bounded from below by the zero operator. Therefore the sequence $k \mapsto \mathcal{P}^k$ converges strongly (pointwise). Since it converges strongly, it converges also weakly (i.e. matrix element-wise), and its weak limit equals to the strong limit. We evaluate its strong limit via evaluating its weak limit, below.

Take any $f, g \in H_m$, then there exists a unique complex valued bounded variation Radon measure $\mu_{\mathcal{P}, f, g}$ over \mathbb{C} with $\text{supp}(\mu_{\mathcal{P}, f, g}) \subset \text{Sp}(\mathcal{P}) \subset [0, 1]$, and $\langle f, \mathcal{P}g \rangle_m = \int_{\lambda \in [0, 1]} \lambda d\mu_{\mathcal{P}, f, g}(\lambda)$ holds. Moreover, for any non-negative integer k , one has that $\langle f, \mathcal{P}^k g \rangle_m = \int_{\lambda \in [0, 1]} \lambda^k d\mu_{\mathcal{P}, f, g}(\lambda)$ holds. One has that $\int_{\lambda \in [0, 1]} \lambda^k d\mu_{\mathcal{P}, f, g}(\lambda) = \int_{\lambda \in [0, 1[} \lambda^k d\mu_{\mathcal{P}, f, g}(\lambda) + \int_{\lambda \in \{1\}} \lambda^k d\mu_{\mathcal{P}, f, g}(\lambda)$, where the second term equals to $\langle f, P g \rangle_m$ by construction. The function $\lambda \mapsto \lambda^k$ converges to zero pointwise on $[0, 1[$, and is bounded by the constant 1 function which is $\mu_{\mathcal{P}, f, g}$ absolute integrable on $[0, 1[$. Therefore, by Lebesgue's theorem of dominated convergence, the integral $\int_{\lambda \in [0, 1[} \lambda^k d\mu_{\mathcal{P}, f, g}(\lambda)$ tends to zero as a function of k . Therefore, $\langle f, \mathcal{P}^k g \rangle_m$ converges to $\langle f, P g \rangle_m$ in k , i.e. \mathcal{P}^k converges weakly to P . ■

By combining Theorem 53 and Theorem 54, one can draw the following conclusion.

Corollary 55 For all $T > 0$ parameter, which is not smaller than the operator norm of $\hat{\mathbf{M}}_{\psi_0}^2$, one has that the iteration

$$G_0 := \mathbb{1}, \quad G_{k+1} := G_k - T^{-1} \hat{\mathbf{M}}_{\psi_0}^2 G_k \quad (26)$$

converges in H_m . Therefore, there exists the finite real number $\nu := \lim_{k \rightarrow \infty} b G_k \in \mathbb{R}$.

The MDS equation has solutions in H_m if and only if $\nu \neq 0$.

Moreover, if $\nu \neq 0$, then $\frac{1}{\nu} \lim_{k \rightarrow \infty} G_k$ is a solution of the MDS equation in H_m . ■

Proof Clearly, by construction we have that $G_k = \mathcal{P}^k \mathbb{1}$ for all $k \in \mathbb{N}_0$, and we have just shown in Theorem 54, that $k \mapsto \mathcal{P}^k$ converges strongly to P . Therefore, $\lim_{k \rightarrow \infty} G_k = P \mathbb{1}$, and $\nu = b P \mathbb{1}$. Applying then Theorem 53, we get the stated result. ■

Remark 56 The following identities are useful for technical evaluation.

- (i) The minimal factor $T > 0$, which can be used in the above existence test is the operator norm of $\hat{\mathbf{M}}_{\psi_0}^2$. The operator norms are generally hard to estimate. However, it can be estimated from above by the trace norm of $\hat{\mathbf{M}}_{\psi_0}^2$, or equivalently, by the Hilbert–Schmidt norm of \mathbf{M}_{ψ_0} , which are technically easier to evaluate.
- (ii) It is a useful fact that the indicator sequence $k \mapsto \nu_k := b G_k \in \mathbb{R}$ consists of non-negative numbers, and is monotonically decreasing. That is because $\nu_k = b G_k = \langle \mathbb{1}, G_k \rangle_m = \langle \mathbb{1}, \mathcal{P}^k \mathbb{1} \rangle_m$. The operator sequence $k \mapsto \mathcal{P}^k$ consists of a sequence of positive operators, which are monotonically decreasing. Therefore

$k \mapsto \nu_k$ inherits this property. Thus, it is enough to test whether the indicator sequence $k \mapsto \nu_k$ is bounded away from zero. Moreover, the scalar component bG_k of the approximants G_k start from 1, they do stay real, and they do not flip sign from positive to negative, and they monotonically decrease.

- (iii) By means of Corollary 55, for concrete models, evaluating whether the indicator ν is bounded away from zero, is expected to involve elaborate Sobolev estimates. The corollary, however, pinpoints a well defined point where one has to invoke these estimates, and therefore this can be considered as a useful existence test condition.
- (iv) Since Corollary 55 is a necessary and sufficient condition, and not merely a sufficient condition, one may also use it in the reverse direction. Namely, if for a concrete model the regularized MDS equation had any solutions, then the iteration scheme of Corollary 55 is guaranteed to be good enough to be convergent, and to produce one particular MDS solution. This is a useful piece of information, even without actually performing the above Sobolev estimates.

7. Concluding remarks

In the QFT literature, the master Dyson–Schwinger (MDS) equation on the field correlators is known to be a differential reformulation of the Feynman integral formalism. In this paper it is shown that the MDS equation can be cast into a particular presentation, in which the involved function spaces and operators are perfectly well defined, regardless of a fixed background spacetime metric, or causal structure, or signature. Moreover, the Wilsonian regularized version of the construction is also shown to be well defined in such a generally covariant setting. A necessary and sufficient condition is proved for the solution space of the regularized MDS equation to be nonempty, for conformally invariant Lagrangians. The pertinent theorem is constructive in the sense that it provides an iterative algorithm to obtain an MDS solution. The algorithm is guaranteed to converge whenever the solution space is nonempty, and could be eventually used for a lattice QFT-like nonperturbative numerical solution scheme, capable of working in the original metric signature.

Acknowledgments

We would like to thank to the organizers of the Simplicity III workshop at the Perimeter Institute, and especially to Neil Turok and Job Feldbrugge for the inspiring exposition on the problematics of Feynman integral formulation in Lorentz signature, which inspired this work. We would also like to thank to Antal Jakovác for enlightening discussion on Feynman integral formulation and ERGE from the physical point of view. We would also like to thank Áron Szabó, Bence Racskó and Igor Khavkine for valuable feedback on the mathematical content of the manuscript. Special thanks to János Kristóf for the enlightening mathematical inputs concerning the theory of topological

vector spaces (TVS) and measure theory over TVS, moreover to Zsigmond Tarcsay for double-checking the mathematical content of the paper concerning the theory of TVS, especially regarding the questions of operator closability.

This work was supported in part by the Hungarian Scientific Research fund (NKFIH K-138152).

Appendix A. Some recalled facts on topological vector spaces

We intend to review here some notions and fundamental results on the theory of topological vector spaces. For a concise introduction see [39] Chapter 0,4,5,7 and also [40] as well as [41]. The main point is that nuclear Fréchet spaces and their strong duals behave well against tensor algebra operations.

Appendix A.1. Fréchet–Hadamard derivative

Let F and G be real topological affine spaces, both with some Hausdorff locally convex topology. Let their underlying vector spaces denoted by \mathbb{F} , \mathbb{G} , respectively. A mapping $S : F \rightarrow G$ is said to be **Fréchet–Hadamard differentiable at a point** $\psi \in F$ [42, 17], whenever there exists a continuous linear map $DS(\psi) : \mathbb{F} \rightarrow \mathbb{G}$, called the derivative of S at ψ , such that for any convergent sequence $n \mapsto h_n$ in \mathbb{F} and any nowhere zero sequence $n \mapsto t_n$ in \mathbb{R} which converges to zero, the continuous linear map $DS(\psi)$ satisfies

$$\lim_{n \rightarrow \infty} \left(\frac{S(\psi + t_n h_n) - S(\psi)}{t_n} - DS(\psi) h_n \right) = 0. \quad (\text{A.1})$$

It is called Fréchet–Hadamard differentiable, if it is so in every point of its domain. It is called continuously Fréchet–Hadamard differentiable, if it is differentiable, and the derivative map $F \rightarrow \mathcal{L}(\mathbb{F}, \mathbb{G})$, $\psi \mapsto DS(\psi)$ is continuous. Note that in general, in order to make sense of this latter definition, one needs to specify a topology on the space $\mathcal{L}(\mathbb{F}, \mathbb{G})$ of $\mathbb{F} \rightarrow \mathbb{G}$ continuous linear maps (there can be many natural topologies on $\mathcal{L}(\mathbb{F}, \mathbb{G})$, see also the followings).

Appendix A.2. Fundamentals on topological vector spaces

Remark 57 We recall some basic definitions and facts on topological vector spaces.

- (i) A Hausdorff topological vector space is called locally convex whenever it has a topological basis consisting of convex neighborhoods. One of the many equivalent definitions of a Hausdorff locally convex topological vector space (HLCTVS) is that it is a real or complex vector space, with a topology defined by a family of seminorms which separate points ([40] Chapter I.1 Theorem 1.36, 1.37 and [43] Chapter I.3, I.4, I.7 Proposition I.7.7 and [44] Chapter 2.1–4).
- (ii) Without loss of generality, the above HLCTVS topology can always be defined with an increasing family of seminorms ([44] Chapter 2.4 and [45] Remark 3.19).

Moreover, if the family of seminorms is countable, it can be recombined such that they are indexed by \mathbb{N}_0 in increasing order (using the above cited algorithm).

- (iii) Similarly to normed spaces, the notion of Cauchy completeness and (up to natural isomorphism) a unique Cauchy completion of a HLCTVS can always be defined ([41] Chapter1 and [43] ChapterI.5 and [44] Chapter2.9).
- (iv) A HLCTVS is metrizable if and only if its topology can be specified by an at most countable system of seminorms, and if complete it is called Fréchet. The TVS-sense Cauchy completion and the metric-sense Cauchy completion coincides. (See: [40] ChapterI.1 and [43] ChapterI.8,I.10 and [44] Chapter2.6.)
- (v) A set B of a HLCTVS is called von Neumann bounded or shortly as bounded, whenever for all open neighborhoods U of the origin there exists some $t_U \in \mathbb{R}^+$, such that for all numbers s with $|s| \geq t_U$ the relation $B \subset sU$ holds. If the space is metrizable, the von Neumann boundedness is usually not the same as the metric boundedness. On Fréchet spaces, the former is stronger or equal to the latter. The two notions coincide on normed spaces. By default, by boundedness we mean von Neumann boundedness ([40] ChapterI.1.29).
- (vi) A HLCTVS is normable if and only if its origin has a bounded neighborhood (Kolmogorov's normability criterion, [40] ChapterI.1 Theorem1.39 and [43] ChapterI.14 Proposition14.4 and [44] Chapter2.6 and [46] ChapterII.2 Theorem2.1 and [39] Theorem0.8.2).
- (vii) Let X be a HLCTVS, then X' denotes the vector space of the continuous linear functionals from X to the numbers, i.e. the continuous dual of X . The vector space X' may be endowed with many natural topologies. The most important two are the strong and weak dual topologies. The weak dual topology is generated by the open zero neighborhoods $G_{A,\varepsilon}^{X'}(0) := \{y \in X' \mid \sup_{x \in A} |(y|x)| < \varepsilon\}$ where $\varepsilon > 0$ and A runs through the finite sets of X . This topology is the weakest one such that the evaluation map $y \mapsto (y|x)$ is continuous for all $x \in X$. The convergence in this topology is equivalent to pointwise convergence. The strong dual topology is generated by the open zero neighborhoods $G_{A,\varepsilon}^{X'}(0) := \{y \in X' \mid \sup_{x \in A} |(y|x)| < \varepsilon\}$ where $\varepsilon > 0$ and A runs through the von Neumann bounded sets of X . The convergence in that topology is the uniform convergence on von Neumann bounded sets. We denote by X^* the space X' endowed with the strong dual topology. If X were normed, the strong dual topology on X' would coincide with the natural norm topology on the continuous linear functionals. (See: [41] Chapter4 Definition4.1 and [43] ChapterII.19 and [45] Chapter3.6.2-4 and [46] ChapterIV.1-5 and [39] Chapter0.6 and [47] Chapter2.)
- (viii) It is common knowledge that the space of smooth sections of a vector bundle over a smooth manifold with its natural \mathcal{E} topology forms a Fréchet space. So does its closed subspace $\mathcal{D}_{\mathcal{K}}$, the sections with support within a fixed compact subset \mathcal{K} of the manifold. (These facts can be also deduced by following arguments of

[41] Chapter 6.)

Appendix A.3. Tensor product and nuclear spaces

Remark 58 We recall some basic definitions and facts related to topological tensor products.

- (i) Let X and Y be HLCTVS. As usual in linear algebra, their algebraic tensor product is a space $X \otimes_a Y$ such that all bilinear maps $X \times Y \rightarrow Z$ to a third space Z factors through a unique linear map $X \otimes_a Y \rightarrow Z$. From the definition it follows that the algebraic tensor product is unique up to linear isomorphism, so one can speak about “the” algebraic tensor product space. The algebraic tensor product $X \otimes_a Y$ may be endowed with a natural HLCTVS topology by the projective topology. This is the finest locally convex topology such that the map $X \times Y \rightarrow X \otimes_a Y$ is jointly continuous. An other important topology, weaker than the projective, is the equicontinuous topology. That is based on viewing $X \otimes_a Y$ as a space of linear forms on $X' \otimes_a Y'$, and the equicontinuous topology is the one defined by uniform convergence on sets $S \otimes_a T \subset X' \otimes_a Y'$ where $S \subset X'$ and $T \subset Y'$ are equicontinuous sets. Since generally we will be dealing with Cauchy complete spaces, by default $X \otimes Y$ shall denote the Cauchy completion of $X \otimes_a Y$ in the projective tensor product topology. (See: [41] Chapter 3 and [43] Chapter III and [46] Chapter III.6.1,5 and [39] Chapter 7.1.)
- (ii) It is a well known phenomenon that in infinite dimensions the completed tensor product of Hilbert spaces leads out from the category of Hilbert spaces ([48] Appendix 19). It is possible, however, to introduce the Hilbert–Schmidt (HS) tensor product within the Hilbert space category. Care should be taken, however, that the HS tensor product fails to obey the universality property, i.e. it is not truly a tensor product operation ([48] Appendix 19). The HS tensor product is the operation used e.g. when constructing a Fock space.
- (iii) Let X and Y be Fréchet spaces and let $X \otimes Y$ denote their completed tensor product in the projective tensor product topology. Then, for all $u \in X$ there exists a sequence $(x_n)_{n \in \mathbb{N}_0}$ and $(y_n)_{n \in \mathbb{N}_0}$ both converging to zero in X and Y , respectively, and an absolute summable sequence of numbers $(\lambda_n)_{n \in \mathbb{N}_0}$, such that $u = \sum_{n=0}^{\infty} \lambda_n x_n \otimes y_n$ ([41] Chapter 3.9 Theorem 3.9 and [43] Chapter III.45 Theorem 45.1 [46] Chapter III.6 Theorem 6.4).
- (iv) One of the equivalent definitions of a HLCTVS to be nuclear is the following. A HLCTVS X is nuclear if and only if for all HLCTVS Y the completed projective and equicontinuous tensor product of X and Y are isomorphic. The completed tensor product of two nuclear spaces is nuclear. That is, nuclear spaces are distinguished by the fact that they behave well against the completed tensor product with any HLCTVS. (See: [43] Chapter III.50.3 Theorem 50.1 and [39] Chapter 5.)

- (v) An important equivalent defining property of nuclear spaces is the following. A complete nuclear HLCTVS space is the projective limit of a family of Hilbert spaces, with nuclear linking maps. A Fréchet space is nuclear if and only if it is the projective limit of an at most countable such family. (See e.g.: [41] Chapter5 Corollary5.14 and [46] ChapterIII.7.3 Corollary3 and [39] Chapter7.3.)
- (vi) In infinite dimensions, nuclear spaces cannot be normed: a normable HLCTVS is nuclear if and only if it is finite dimensional ([43] ChapterIII.50.11 Corollary2).
- (vii) Cauchy completion of a nuclear space is nuclear ([41] Chapter5 Corollary5.10 and [43] ChapterIII.50.5 Proposition50.1).
- (viii) A Fréchet space is nuclear if and only if its strong dual is nuclear ([43] ChapterIII.50.15 Proposition50.6). A nuclear Fréchet space is reflexive to the strong duality ([41] Chapter5 Proposition5.12).
- (ix) The projective limit of a family of nuclear spaces is nuclear, and the strict inductive limit of a countable family of nuclear spaces is nuclear ([41] Chapter5 Corollary5.17 and [43] ChapterIII.50.5 Proposition50.1).
- (x) Every linear subspace of a nuclear space is nuclear, and the quotient with a closed linear subspace is nuclear ([41] Chapter5 Proposition5.16 and [43] ChapterIII.50.5 Proposition50.1 and [39] Chapter5.1).
- (xi) The category of nuclear Fréchet (NF) spaces and the category of strong dual of nuclear Fréchet (DNF) spaces is closed under the completed tensor product, and they are dual to each-other in terms of the strong duality. Moreover, if X and Y are both NF or both DNF, then

$$\mathcal{L}(X, Y^*) \equiv X^* \otimes Y^* \equiv (X \otimes Y)^* \equiv \mathcal{B}(X, Y)$$

holds, where the space $\mathcal{L}(X, Y^*)$ of continuous linear maps is understood with the topology of uniform convergence on von Neumann bounded sets, and $\mathcal{B}(X, Y)$ is the space of jointly continuous bilinear functionals of $X \times Y$ into the numbers (Schwartz kernel theorem, see e.g.: [41] Chapter5 Theorem5.25). That is, for NF or for DNF spaces, tensor product can be implemented via the multiplicative realization, in the analogy to finite dimensional vector spaces. The closed linear subspace of an NF or DNF space is NF or DNF, respectively, and the quotient by closed linear subspace also preserves the NF or DNF property ([39] Chapter5.1.5,6,7,8). All NF or DNF spaces are separable ([39] Chapter4.4.10).

- (xii) It is common knowledge that the space of smooth sections of a vector bundle over a smooth manifold with its natural \mathcal{E} topology forms a nuclear Fréchet space. So does its closed subspace $\mathcal{D}_{\mathcal{K}}$, the sections with support within a fixed compact subset \mathcal{K} of the manifold. (These facts can be also deduced by following arguments of [41] Chapter6.)
- (xiii) A topological vector space is called LF space, if it is a countable strict inductive limit of Fréchet spaces ([43] ChapterI.13). An LF space is Cauchy complete

([43] Chapter I.13 Theorem 13.1). An LF space is metrizable iff the inductive system of subspaces are finite dimensional. An LF space is called LNF if it is countable strict inductive limit of nuclear Fréchet spaces, and as such, an LNF space is nuclear. Let us require from this point on that an LNF space is such a strict inductive limit of a countable system of NF spaces, that each member of the family has closed image within its adjacent member. It can be shown that such an LNF space is reflexive against strong duality and its dual, a DLNF space, is also nuclear ([41] Chapter 6 Proposition 6.8,9,10).

- (xiv) It is common knowledge that the space of compactly supported smooth sections of a vector bundle over a smooth non-compact manifold with its natural \mathcal{D} topology forms an LNF space, and its strong dual \mathcal{D}^* a corresponding DLNF space. (These facts can be also deduced by following arguments of [41] Chapter 6.)

Appendix A.4. Joint and separate continuity of bilinear maps

Remark 59 In a generic HLCTVS, certain continuity properties of bilinear forms are rather counterintuitive. E.g., separately continuous bilinear maps need not be jointly continuous.

- (i) The duality pairing form on a HLCTVS and its strong dual is jointly continuous if and only if the space is normable ([44] Chapter 4.7 p.359). Therefore, in an infinite dimensional nuclear space, the duality pairing cannot be jointly continuous. There are many examples in such spaces which are separately continuous, but not jointly.
- (ii) Let X, Y, Z be HLCTVS and consider a separately continuous bilinear map $X \times Y \rightarrow Z$. If X is Fréchet, then separate continuity implies joint sequential continuity. Moreover, if in addition Y is metrizable, then separate continuity implies joint continuity ([40] Chapter I.2 Theorem 2.17).
- (iii) Let X, Y, Z be HLCTVS and consider a separately continuous bilinear map $X \times Y \rightarrow Z$. Assume that X and Y are strong duals of nuclear Fréchet spaces. Then, the pertinent bilinear map is jointly continuous ([45] Chapter 3.9.1 Theorem 3.137(b)).
- (iv) For instance the bilinear map defined by $\mathcal{E}(\mathbb{R}^n) \times \mathcal{D}(\mathbb{R}^n) \rightarrow \mathcal{D}(\mathbb{R}^n)$, $(\delta\psi, \delta\psi_T) \mapsto \delta\psi\delta\psi_T$ (pointwise product) is separately continuous, but not jointly continuous. It is merely so-called hypocontinuous. (See: [43] Chapter III.41.4 p.423.) On the other hand, the pointwise product $\mathcal{D}(\mathbb{R}^n) \times \mathcal{D}(\mathbb{R}^n) \rightarrow \mathcal{D}(\mathbb{R}^n)$, $(\delta\psi_{T1}, \delta\psi_{T2}) \mapsto \delta\psi_{T1}\delta\psi_{T2}$ is jointly continuous ([49] Proposition 2.2). But $\mathcal{D}(\mathbb{R}^n) \times \mathcal{D}(\mathbb{R}^m) \rightarrow \mathcal{D}(\mathbb{R}^{n+m})$, $(\delta\psi_{T1}, \delta\psi_{T2}) \mapsto \delta\psi_{T1} \otimes \delta\psi_{T2}$ is not jointly continuous because although the completed tensor product $\mathcal{D}(\mathbb{R}^n) \otimes \mathcal{D}(\mathbb{R}^m) \equiv \mathcal{D}(\mathbb{R}^{n+m})$ as vector spaces, but the tensor product topology is weaker than the original (inductive limit) topology of $\mathcal{D}(\mathbb{R}^{n+m})$ ([49] Theorem 2.4). The above phenomena are explained by the fact that even forming a finite cartesian product topology cannot be exchanged with taking the (strict) inductive limit topology.

Appendix A.5. Cartesian products, locally convex direct sums

Remark 60 We collect some fundamental identities on the cartesian products and locally convex direct sums of HLCTVS spaces.

- (i) Let $(X_i)_{i \in I}$ a family of LCTVS. The cartesian product set $\prod_{i \in I} X_i$ endowed with the entrywise linear operations is denoted by $\bigoplus_{i \in I} X_i$, which can be naturally endowed with the product (Tychonoff or initial) topology, being the coarsest topology such that the canonical projection maps are continuous. The Tychonoff topology is, quite evidently, a LCTVS topology. Therefore, $\bigoplus_{i \in I} X_i$ is called the Tychonoff direct sum. The box topology on $\bigoplus_{i \in I} X_i$ is defined by the zero neighborhoods being the convex hulls of cartesian boxes composed of zero neighborhoods. The linear subspace of $\bigoplus_{i \in I} X_i$ which consists of tuples with only finitely many nonzero entries is denoted by $\bigoplus_{i \in I} X_i$, and is called the algebraic direct sum. It may be endowed with the locally convex direct sum topology (final topology), which is the finest locally convex topology on which the canonical injections are continuous. Clearly, the Tychonoff and the box topologies may be as well restricted to the algebraic direct sum space. These topologies are Hausdorff whenever each element of the family is Hausdorff. (See: [44] Chapter2.7 and [45] Chapter3.3.7.)
- (ii) For a general family of LCTVS, the relation: product (Tychonoff) topology \subset box topology \subset locally convex direct sum topology holds. For a countable family of LCTVS, the relation: product (Tychonoff) topology \subset box topology = locally convex direct sum topology holds. For a finite family of LCTVS, the relation: product (Tychonoff) topology = box topology = locally convex direct sum topology holds. (See e.g.: [50] Proposition11, Theorem21, Corollary22.)
- (iii) The Cauchy completion of a Tychonoff direct sum is the Tychonoff direct sum of Cauchy completed spaces, the Cauchy completion of a locally convex direct sum is the locally convex direct sum of Cauchy completed spaces ([44] Chapter2.7 and [46] ChapterII.6.1,2).
- (iv) The Tychonoff direct sum of countable family of metrizable HLCTVS is metrizable, the locally convex direct sum of a countable family of metrizable HLCTVS is metrizable ([44] Chapter2.7).
- (v) The Tychonoff direct sum of arbitrarily many nuclear spaces is nuclear, the locally convex direct sum of at most countable many nuclear spaces is nuclear ([41] Proposition5.15 and [39] Chapter5.2.1,2).
- (vi) Let $(X_i)_{i \in I}$ be a family of HLCTVS. Then, $(\bigoplus_{i \in I} X_i)^* = \bigoplus_{i \in I} X_i^*$ and $(\prod_{i \in I} X_i)^* = \prod_{i \in I} X_i^*$, where $(\cdot)^*$ denotes strong duality (see e.g. [45] Theorem3.105).

Appendix A.6. Continuity and sequential continuity

Remark 61 Since not all the HLCTVS have sequential topology, it is useful to recall some relations of continuity and sequential continuity of maps.

- (i) Since an NF space is metrizable, its topology is sequential ([40] Appendix A6).
- (ii) A DNF space is sequential ([51] Proposition 5.7).
- (iii) A Hausdorff topological vector space is sequential if and only if there exists no strictly finer topology with the same convergent sequences ([52] Theorem 7.4).
- (iv) A LNF or a DLNF space which are not metrizable or dual metrizable, respectively, is not sequential ([53] Theorem 1.1). An LNF is metrizable iff it is a strict inductive limit of a countable sequence of finite dimensional vector spaces. In particular, the test functions on non-compact manifolds and their strong duals are not sequential.
- (v) An NF or a DNF space is Montel ([41] Theorem 5.25).
An NF or a DNF space is bornological ([41] Theorem 5.25).
Strict inductive limit preserves the Montel property ([45] Theorem 3.68).
Strict inductive limit preserves the bornological property ([41] p.11).
Strong duality preserves the Montel property ([44] Chapter 3.9 Proposition 9, [45] Theorem 3.123, [46] Chapter IV.5.9 – assumes reflexivity, permanence properties automatic).
Strong duality does not preserve the bornological property.
But an DLNF space is still bornological, since LNF is the strict inductive limit of Montel and bornological Fréchet spaces and these are bornological ([44] Chapter 3.16 Theorem 2).
In summary: NF, DNF, LNF, DLNF spaces are Montel and bornological.
- (vi) On Montel spaces the weakly bounded sets are strongly bounded. On the strong dual of a Montel space the weak and the strong topology coincides over bounded sets ([43] Chapter II.34.7 Proposition 34.6).
Therefore, in Montel spaces sequences are weakly convergent if and only if they are strongly convergent ([43] Chapter II.34.7 Corollary 1).
- (vii) Arbitrary cartesian product of Montel spaces is Montel ([44] Chapter 3.9 Proposition 4 and Remark 1).
At most countable cartesian product of bornological spaces are bornological ([45] p.147).
Therefore at most countable cartesian product of NF, DNF, LNF, DLNF spaces are Montel and bornological.
- (viii) If a Hausdorff locally convex topological vector space X is bornological, and Y is locally convex topological vector space, then a linear map $F : X \rightarrow Y$ is continuous if and only if it is sequentially continuous ([46] Chapter II.8 Theorem 8.3). If X is Montel and bornological, then weak sequential continuity (in X) also implies continuity, for linear maps.

- (ix) Sequential topology is not preserved by cartesian product, but metrizability is preserved by at most countable cartesian product, which is then sequential. Therefore, at most countable cartesian product of an NF space is sequential. A DNF space is sequential, but since it is not metrizable, its cartesian products with itself or other spaces are generally not sequential.
- (x) LNF or DLNF spaces are not sequential. But they are still Montel and bornological, similarly to NF and DNF. Therefore, if X consists of at most countable cartesian product of NF, DNF, LNF, DLNF spaces, and Y is a locally convex topological vector space, then a linear map $F : X \rightarrow Y$ is continuous if and only if it is sequentially continuous (weak sequential continuity in X is enough).
- (xi) There is a theory of convergence vector spaces (CVS), capturing only the convergence aspects instead of topological aspects over some vector space [54]. In the analogy of TVS, is possible to define such a space to be Hausdorff as well as being locally convex (HLCVS), and to be complete. A CVS may or may not be originating from a corresponding topological vector space (TVS). In addition, a CVS may eventually be determined by convergence of countable sequences [55, 56, 52], called to be sequential CVS. An example for a CVS which does not originate from a TVS is the space of Lebesgue measurable functions over \mathbb{R}^N with the pointwise almost everywhere convergence. Most of the CVS, however admit some underlying TVS. The CVS category, however, harmonizes better with the notion of joint sequential continuity, see e.g. [56]. In this paper, as is most common, we will use an approach through TVS, and only refer to CVS properties when unavoidable.
- (xii) Let each of X , Y and Z be one of these spaces: NF, DNF, LNF or DLNF. Then, any separately sequentially continuous bilinear map $F : X \times Y \rightarrow Z$ is also jointly sequentially continuous ([57] Proposition 4.1 and Theorem 4.8). That is, in such topological spaces, when considered as CVS, separate continuity implies joint continuity in the sense of CVS continuity (but usually not as TVS continuity).

Appendix A.7. Closure and sequential closure

Remark 62 We recall some results from the literature on the closure of linear operators.

- (i) Let X and Y be TVS, and $F : X \rightarrow Y$ a linear map defined on a subspace $\text{Dom}(F) \subset X$, which may be smaller than X . The operator F is called closed iff its graph is closed in $X \times Y$. It is called sequentially closed iff its graph is sequentially closed in $X \times Y$. It is called closable or sequentially closable, iff it has a closed or sequentially closed extension as a linear map, and the smallest such extension is called its closure or its sequential closure, respectively. Clearly, closability and closure are stronger than sequential closability and sequential closure, even for maps between complete spaces. These are equivalent, however, if both X and Y are complete metrizable (e.g. for both being NF spaces). These are also equivalent whenever both X and Y are DNF spaces.

- (ii) Let X and Y be TVS, and $F : X \rightarrow Y$ a linear operator, defined on $\text{Dom}(F) \subset X$. Then, F is closable iff Y can be endowed with a Hausdorff vector topology which is not stronger than the original topology on Y , and with respect to which F becomes continuous ([58] Theorem1). Moreover, if X and Y was LCTVS, then F is closable iff Y can be endowed with a respective HLCTVS topology, not stronger than the original, making F continuous (see [58] p.108 or in proof of Theorem1 there).
- (iii) Let X and Y be TVS, and $F : X \rightarrow Y$ a densely defined closable linear operator, with its closure \tilde{F} . Then, the subspace $\text{Ker}(\tilde{F})$ is closed. (Consequence of [58] Theorem2.)
- (iv) It is elementary to check that a linear map $F : X \rightarrow Y$ is sequentially closable if and only if: for all $x \in X$, for any sequences $(x_n)_{n \in \mathbb{N}}$ and $(x'_n)_{n \in \mathbb{N}}$ with $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} x'_n = x$, one has that whenever both $(F x_n)_{n \in \mathbb{N}}$ and $(F x'_n)_{n \in \mathbb{N}}$ are convergent, then $\lim_{n \rightarrow \infty} F x'_n = \lim_{n \rightarrow \infty} F x_n$ holds. An other reformulation is that for all sequences $(x_n)_{n \in \mathbb{N}}$ with $\lim_{n \rightarrow \infty} x_n = 0$ one has that whenever $(F x_n)_{n \in \mathbb{N}}$ is convergent, then $\lim_{n \rightarrow \infty} F x_n = 0$ holds. A further equivalent reformulation of sequential closability is the following: for all $x \in X$, for any sequences $(x_n)_{n \in \mathbb{N}}$ and $(x'_n)_{n \in \mathbb{N}}$ with $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} x'_n = x$ and with $\lim_{n \rightarrow \infty} F x_n = 0$, one has that whenever $(F x'_n)_{n \in \mathbb{N}}$ is convergent, then $\lim_{n \rightarrow \infty} F x'_n = 0$ holds. The latter in words means that: the operator F is sequentially closable iff it is sequentially closable at its approximate kernel points (which will then be the actual kernel points of its sequential closure).
- (v) If X and Y are not both metrizable (e.g. NF) or DNF, then generally the sequential closability does not imply closability of a densely defined linear operator $F : X \rightarrow Y$, even if X and Y were complete Montel and bornological HLCTVS. That is because the dense subspace $\text{Dom}(F) \subset X$ may not inherit the bornological property of X , and therefore one cannot conclude from sequential behavior, despite of the linearity of F .
- (vi) For a densely defined linear operator $F : X \rightarrow Y$ between two complete HLCTVS, the notion of multivalued set can be introduced: $\text{Mul}(F) := \{y \in Y \mid \exists (x_n)_{n \in \mathbb{N}} \text{ in } \text{Dom}(F) : \lim_{n \rightarrow \infty} x_n = 0 \text{ and } \lim_{n \rightarrow \infty} F x_n = y\}$. By construction, $\text{Mul}(F)$ is a linear subspace (moreover, in Banach spaces it is also closed, see e.g. [59]). The operator F is sequentially closable iff $\text{Mul}(F) = \{0\}$. That is, $\text{Mul}(F)$ measures the non-closability of F . The operator F is called maximally non-closable or maximally singular, whenever $\text{Mul}(F)$ is dense in the closure of $\text{Ran}(F)$. In particular, whenever $\text{Mul}(F)$ is dense in Y , the operator F is necessarily maximally non-closable.
- (vii) Let \mathbb{F} denote the space of smooth sections of some vector bundle over the base manifold \mathcal{M} , as previously, with the standard \mathcal{E} smooth function topology. Recall that $\mathbb{F} \otimes \mathbb{F}$ may be identified with the corresponding space of smooth sections

over $\mathcal{M} \times \mathcal{M}$ with their \mathcal{E} topology. Introduce the diagonal evaluation map $\text{ev} : \mathbb{F} \otimes \mathbb{F} \longrightarrow \mathbb{F}$, $((x, y) \mapsto G(x, y)) \longmapsto (z \mapsto G(z, z))$. Such map appears e.g. in the interaction term of the MDS operator generated by an interacting Euler–Lagrange functional. The above map may be considered as well as a densely defined linear map $\widehat{\text{ev}} : (\mathbb{F}_T^\times)^* \otimes (\mathbb{F}_T^\times)^* \multimap (\mathbb{F}_T^\times)^*$ on the space of distributions, since there is the natural dense inclusion $\mathbb{F} \subset (\mathbb{F}_T^\times)^*$. If this operator were (sequentially) closable, it would provide a straightforward solution for the problematics of renormalization. It can be shown, however, that this map is maximally non-closable. In order to demonstrate that, take any function $g \in \mathbb{F}$ and a corresponding sequence $(G_n)_{n \in \mathbb{N}}$ in $\mathbb{F} \otimes \mathbb{F}$ such that for all $n \in \mathbb{N}$ the equality $G_n(z, z) = g(z)$ ($\forall z \in \mathcal{M}$) is satisfied, and that $(G_n)_{n \in \mathbb{N}}$ converges locally uniformly to zero around any point of $\mathcal{M} \times \mathcal{M} \setminus \{(x, y) \in \mathcal{M} \times \mathcal{M} | x = y\}$. Clearly, one can construct such a sequence for all $g \in \mathbb{F}$. (One can construct this explicitly on \mathbb{R}^N , and then one can bring this to a manifold \mathcal{M} via partition of unity arguments.) The above finding means, however, that all $g \in \mathbb{F} \subset (\mathbb{F}_T^\times)^*$ is in $\text{Mul}(\widehat{\text{ev}})$. Therefore, $\widehat{\text{ev}}$ is maximally non-closable.

Appendix A.8. Nuclear Fréchet spaces with countable Hilbertian structure

Remark 63 We recall some fundamental facts on a special type of nuclear Fréchet spaces, which will play an important role in QFT constructions.

- (i) Whenever the topology of a HLCTVS is defined by a countable family of seminorms, and there exists a continuous norm on that HLCTVS, then by means of Remark 57(ii): without loss of generality, the defining family of seminorms of the topology of the space can be taken to be an increasing family of norms instead of seminorms, indexed by \mathbb{N}_0 . Specially, the pertinent continuous norm may be chosen as one of the elements of the topology defining norm family. Moreover, it may be chosen also to be the weakest one. Let \mathcal{H} be such a HLCTVS, and let $\|\cdot\|_n$ ($n \in \mathbb{N}_0$) be a countable increasing family of topology defining norms on it. Denote the completion of \mathcal{H} in the $\|\cdot\|_n$ norm by H_n , then the natural inclusion $i_{\infty, n} : \mathcal{H} \rightarrow H_n$ is a continuous injective linear map. One may write $H_n \supset \mathcal{H}$ is dense (for all $n \in \mathbb{N}_0$).
- (ii) If the topology of a HLCTVS is defined by a countable system of Hilbertian seminorms, and there exists a continuous Hilbertian norm, then the analogy of the statements in (i) will hold, respectively, with Hilbertian norms and Hilbert spaces as completed spaces.
- (iii) Let \mathcal{H} be a HLCTVS as in (i), and let $\|\cdot\|_n$ ($n \in \mathbb{N}_0$) be a countable increasing family of topology defining norms on it, the completed Banach spaces denoted by H_n . The $\mathcal{H} \rightarrow \mathcal{H}$ identity map is continuous when the norm topology of $\|\cdot\|_{n+1}$ is taken on the starting space and the norm topology of $\|\cdot\|_n$ is taken on the image space. Therefore, its continuous extension in these norms is a continuous linear

map $i_{n+1,n} : H_{n+1} \rightarrow H_n$. The map $i_{n+1,n}$ is the identity over \mathcal{H} , and therefore it is injective on that subspace, but its continuous extension $i_{n+1,n} : H_{n+1} \rightarrow H_n$ may or may not be injective over the full completed space H_{n+1} . Whenever $i_{n+1,n}$ is injective, the adjacent norms are called Gel'fand compatible (see [60] Chapter 2.2 and [61] Chapter 2.2.8-2.2.11 and [62] and [63] Appendix B.10-13). An other way to formulate the Gel'fand compatibility of the adjacent norms is that any sequence in \mathcal{H} which is Cauchy in both of the adjacent norms, and which is convergent to zero in either of the norms, then it is convergent to zero also in the other norm. If Gel'fand compatibility of adjacent norms hold, then one may regard the completed spaces as nested in each-other: $H_0 \supset \dots \supset H_n \supset H_{n+1} \supset \dots \supset \mathcal{H}$.

- (iv) Combining the above observations and Remark 58(v), the topology of a nuclear Fréchet space \mathcal{H} admitting a continuous Hilbertian norm can be described by a countable system of increasing Hilbertian norms, indexed by \mathbb{N}_0 . Whenever these adjacent topology defining Hilbertian norms are Gel'fand compatible, then (iii) applies, and in addition $\bigcap_{n \in \mathbb{N}_0} H_n = \mathcal{H}$ holds. Moreover, for all $n \in \mathbb{N}_0$ there exists an integer $m \geq 1$, such that the inclusion map $i_{n+m,n} : H_{n+m} \rightarrow H_n$ is nuclear. (Specially, one may choose the system such, that all the adjacent inclusion maps $i_{n+1,n}$ are nuclear.) Nuclear Fréchet spaces of this kind are called countably Hilbert (CH) type NF spaces. (See [61] Chapter 2.2.8, and e.g. the review paper [47]. Note that the Gel'fand compatibility condition is often overlooked in the literature.) The CH type NF spaces give realization to a special form of a projective limit: the projection maps $i_{\infty,n}$ as well as the linking maps $i_{n+1,n}$ are all injective.
- (v) On a noncompact manifold, the space of smooth sections of a vector bundle with the standard \mathcal{E} topology is an NF space, but not of CH type. It is not difficult to see that on a compact manifold, they are of CH type NF spaces. Also, on a finite dimensional real affine space, one can define the space of rapidly decreasing (Schwartz) sections, which are well known to be NF spaces of CH type (see a review paper: [64]).
- (vi) The strong topological dual (or complex conjugate dual) space of CH type NF spaces have a corresponding inductive limit structure. Namely, let \mathcal{H} be a CH type NF space, with the notations as in (iv). Taking the strong topological dual (or complex conjugate dual) of the inclusion chain $H_0 \supset \dots \supset H_n \supset H_{n+1} \supset \dots \supset \mathcal{H}$, one infers that $\mathcal{H}^* \supset \dots \supset H_{n+1}^* \supset H_n^* \supset \dots \supset H_0^*$ and $\bar{\mathcal{H}}^* \supset \dots \supset \bar{H}_{n+1}^* \supset \bar{H}_n^* \supset \dots \supset \bar{H}_0^*$ holds. More precisely, for all $n \in \mathbb{N}_0$ one has the continuous linear injective maps $i_{\infty,n} : \mathcal{H} \rightarrow H_n$ and $i_{n+1,n} : H_{n+1} \rightarrow H_n$, whose topological transpose (or complex conjugate topological transpose) give rise to the continuous linear maps $i_{\infty,n}^* : H_n^* \rightarrow \mathcal{H}^*$ and $i_{n+1,n}^* : H_n^* \rightarrow H_{n+1}^*$, as well as $\bar{i}_{\infty,n}^* : \bar{H}_n^* \rightarrow \bar{\mathcal{H}}^*$ and $\bar{i}_{n+1,n}^* : \bar{H}_n^* \rightarrow \bar{H}_{n+1}^*$, respectively. Since for all $n \in \mathbb{N}_0$ the image of $i_{n+1,n}$ was dense (it contained $i_{\infty,n}[\mathcal{H}]$ whose completion was H_n), the transpose (or complex conjugate transpose) linking maps cannot have a kernel. Thus, $i_{\infty,n}^*$, $i_{n+1,n}^*$, $\bar{i}_{\infty,n}^*$, $\bar{i}_{n+1,n}^*$ are injective, which justifies the above inclusion chain of the dual (or com-

plex conjugate dual) spaces. Moreover, one has $\mathcal{H}^* \equiv \bigcup_{n \in \mathbb{N}_0} H_n^*$ and $\bar{\mathcal{H}}^* \equiv \bigcup_{n \in \mathbb{N}_0} \bar{H}_n^*$.

Since the inclusion $H_n \supset H_{n+m}$ was eventually Hilbert–Schmidt (and eventually nuclear), the above corresponding dual (or complex conjugate dual) inclusions also eventually become Hilbert–Schmidt (and eventually become nuclear).

- (vii) Whenever a preferred Hilbertian norm $\|\cdot\|_0$ is fixed on a countable Hilbert type NF space, it give rise to the well known rigged Hilbert spaces or Gel’fand triples (see [60, 61]). More concretely, using the notations of (vi), for all $n \in \mathbb{N}_0$ the Riesz representation theorem guarantees the natural linear unitary isomorphism $r_n : H_n \rightarrow \bar{H}_n^*$, and correspondingly, the natural antilinear antiunitary antiisomorphism $\bar{r}_n : H_n \rightarrow H_n^*$, via the relations $r_n(x) := \overline{\langle x, \cdot \rangle_{H_n}}$ and $\bar{r}_n(x) := \langle x, \cdot \rangle_{H_n}$ for all $x \in H_n$. Using this, in particular for H_0 , one has the continuous linear inclusions with dense image $\bar{\mathcal{H}}^* \supset \dots \supset H_{-n-1} \supset H_{-n} \supset \dots \supset H_0 \supset \dots \supset H_n \supset H_{n+1} \supset \dots \supset \mathcal{H}$ where the inclusions are eventually Hilbert–Schmidt (and eventually nuclear). Here, we used the notation $H_{-n} := \bar{H}_n^*$ for all $n \in \mathbb{N}_0$, and the identification $\bar{H}_0^* \equiv H_0$ via r_0 . One should note that the injective continuous linear maps (which are eventually Hilbert–Schmidt and eventually nuclear) $\bar{i}_{n,n+k}^* \circ r_n \circ i_{n+m,n} : H_{n+m} \rightarrow \bar{H}_{n+k}^*$ and $\bar{i}_{n',n'+k'}^* \circ r_{n'} \circ i_{n'+m',n'} : H_{n'+m'} \rightarrow \bar{H}_{n'+k'}^*$ ($n, m, k, n', m', k' \in \mathbb{N}_0$) are in general different maps for $n' \neq n$ even if $n' + m' = n + m$ and $n' + k' = n + k$ holds. That is because the Riesz identification maps r_n and $r_{n'}$, even if restricted to the common set \mathcal{H} , are different for $n' \neq n$. That is, the continuous linear injection $\mathcal{H} \rightarrow H_0 \rightarrow \bar{\mathcal{H}}^*$ is not natural, it explicitly depends on the choice of the Hilbertian norm $\|\cdot\|_0$ on \mathcal{H} . In Section Appendix A.9 we shall show an example when a similar embedding chain construction is natural.

Appendix A.9. Sobolev and Maurin embedding theorems

Remark 64 For some of the proofs, recalling the Sobolev and Maurin embedding theorems will be helpful. Let \mathcal{M} be \mathbb{R}^m , or an open set of \mathbb{R}^m with a boundary having the cone property, or a compact smooth m -dimensional real manifold with such boundary. Let $V(\mathcal{M})$ be a real vector bundle over \mathcal{M} . Fix some smooth Riemannian metric on $V(\mathcal{M})$. The symbol $C_b^k(V(\mathcal{M}))$ will denote the Banach space of k -times continuously differentiable sections of $V(\mathcal{M})$ with bounded derivatives, with the C^k supremum norm. The symbol $H_l(V(\mathcal{M}))$ will denote the Hilbert space of l -times weakly differentiable section Lebesgue equivalence classes of $V(\mathcal{M})$ which are square integrable together with all of their derivatives, equipped with the corresponding L^2 norm including all the derivatives (i.e. H_l is the l -th order L^2 type Sobolev space).

- (i) For \mathcal{M} as above, and $l > k + \frac{\dim(\mathcal{M})}{2}$ one has the inclusion $H_l(V(\mathcal{M})) \subset C_b^k(V(\mathcal{M}))$ and the embedding is continuous, i.e. the H_l norm is stronger than the C_b^k norm. (Sobolev embedding theorem, see e.g.: [65] Chapter4 Theorem4.12(I)(A) and [66] ChapterVI.2.II.3, [66] ChapterVI.16.3.)

- (ii) For \mathcal{M} being as above, but assumed to be bounded for the case $\mathcal{M} \subset \mathbb{R}^m$, then for $l > k + \frac{\dim(\mathcal{M})}{2}$ one has that the inclusion $H_l(V(\mathcal{M})) \subset H_k(V(\mathcal{M}))$ is Hilbert–Schmidt. (By construction of the H -type Sobolev spaces, one has that for all $l > k$ the inclusion $H_l(V(\mathcal{M})) \subset H_k(V(\mathcal{M}))$ is trivially valid, and is continuous, i.e. the H_l norm is stronger than the H_k norm.) The analogous statement holds for the H -type Sobolev spaces obtained from the completion of compactly supported sections. (Maurin embedding theorem, see e.g.: [65] Chapter 6 Theorem 6.61 or [66] Chapter VI.2.II.3, [66] Chapter VI.2.II.7, [66] Chapter VI.16.3 etc.)
- (iii) Knowing result (i) and Hölder’s inequality, it is easy to understand why on the \mathcal{E} or \mathcal{D} type spaces the family of supremum type C^k local norms and the L^2 type C^k local norms ($k \in \mathbb{N}_0$) are equivalent.
- (iv) Knowing result (ii), it is easy to understand the nuclearity of the \mathcal{E} or \mathcal{D} spaces. The theorem implies that the L^2 type C^k local norms over each fixed compact region are eventually getting gradually stronger with growing $k \in \mathbb{N}_0$, and eventually the embedding becomes Hilbert–Schmidt with large enough k . Consequently, the embedding eventually becomes nuclear (trace class) with large enough k (since composition of two Hilbert–Schmidt maps are nuclear).
- (v) Knowing that for all $l > k$ the inclusion $H_l(V(\mathcal{M})) \rightarrow H_k(V(\mathcal{M}))$ is injective, it is easy to understand that over compact manifolds the \mathcal{E} (or \mathcal{D}) spaces are not only NF spaces, but are also CH type NF spaces. It is obvious that they admit continuous Hilbertian norms, and because of the above injective inclusion, the topology defining countable family of increasing Hilbertian norms are also automatically Gel’fand compatible (see Remark 63).
- (vi) Denote by $\bar{V}^\times(\mathcal{M})$ the densitized complex conjugate dual vector bundle of $V(\mathcal{M})$, and their spaces of smooth sections by $\bar{\mathcal{E}}^\times$ and \mathcal{E} , respectively. Whenever the base manifold \mathcal{M} is compact, there is the natural jointly continuous sesquilinear form $\bar{\mathcal{E}}^\times \times \mathcal{E} \rightarrow \mathbb{C}$, $(p, \delta\psi) \mapsto \int_{\mathcal{M}} \bar{p} \delta\psi$. This gives rise to a natural continuous complex-linear injection $r : \mathcal{E} \rightarrow (\bar{\mathcal{E}}^\times)^*$. If a smooth complex (sesquilinear) Riemann metric on $V(\mathcal{M})$ is chosen which is densitized by a positive volume form on \mathcal{M} , it gives rise to corresponding Hilbertian norms $\|\cdot\|_0^\times$ and $\|\cdot\|_0$ on $\bar{\mathcal{E}}^\times$ and \mathcal{E} , respectively. The corresponding norm equivalence class, and therefore the norm topology does not depend on the particular choice of the densitized Riemann metric (this is elementary, but see also [67] Appendix A). The above sesquilinear form is continuous in these norms. On the other hand, both $\bar{\mathcal{E}}^\times$ and \mathcal{E} are CH type NF spaces, giving rise to the continuous complex-linear injections $\mathcal{E} \rightarrow E_0$ and $(\bar{E}_0^\times)^* \rightarrow (\bar{\mathcal{E}}^\times)^*$, referring to the notations of Remark 63(vii). With these choices, the above injection r induces a continuous complex-linear bijection $r : E_0 \rightarrow (\bar{E}_0^\times)^*$, which is isometric, and therefore is a unitary isomorphism. Consequently, one has the natural inclusions $(\bar{\mathcal{E}}^\times)^* \supset \dots \supset E_{-n-1} \supset E_{-n} \supset \dots \supset E_0 \supset \dots \supset E_n \supset E_{n+1} \supset \dots \supset \mathcal{E}$, where for $n \in \mathbb{N}_0$ one has $E_{-n} := (\bar{E}_n^\times)^*$, furthermore on E_0 and \bar{E}_0^\times the above particular

Hilbertian norms were used, and therefore one has the natural unitary isomorphism $E_0 \equiv \bar{E}_0^\times$ induced by the map r , regardless of the particular choice of the densitized Riemann metric.

- (vii) If $l > \frac{\dim(\mathcal{M})}{2}$, and the base manifold \mathcal{M} is compact, the Sobolev space $H_l(V(\mathcal{M}))$ becomes a reproducing kernel Hilbert space (see also [68] and e.g. [67] AppendixD), due to the Sobolev inequality (i). That is, for every $x \in \mathcal{M}$ the point evaluation $H_l(V(\mathcal{M})) \rightarrow \mathbb{C}$, $f \mapsto (p_x|f(x))$ becomes a well defined continuous complex-linear map, where p_x is any element of $V_x^*(\mathcal{M})$. The Riesz representation theorem ensures that for each $x \in \mathcal{M}$ and $p_x \in V_x^*(\mathcal{M})$, there exists a unique $K_x^{\bar{p}_x} \in H_l(V(\mathcal{M}))$ such that $\langle K_x^{\bar{p}_x}, f \rangle = (p_x|f(x))$ holds for all $f \in H_l(V(\mathcal{M}))$. If Penrose abstract indices $\mathcal{A}, \mathcal{B}, \dots$ are used on $V(\mathcal{M})$, this reads as $\langle \bar{p}_{x\mathcal{A}'} K_x^{\mathcal{A}'}, f \rangle = p_{x\mathcal{A}} f^{\mathcal{A}}(x)$. As $K_x^{\bar{p}_x}$ itself is a section of $V(\mathcal{M})$, it may also be evaluated at any point. The reproducing kernel function $K : \bar{V}^*(\mathcal{M}) \times V^*(\mathcal{M}) \rightarrow \mathbb{C}$ is defined as $K(x, y)^{\bar{p}_x, q_y} := (q_y|K_x^{\bar{p}_x}(y))$, or in Penrose abstract indices, $q_{y\mathcal{B}} \bar{p}_{x\mathcal{A}'} K(x, y)^{\mathcal{A}'\mathcal{B}} := q_{y\mathcal{B}} \bar{p}_{x\mathcal{A}'} K_x^{\mathcal{A}'}(y)^{\mathcal{B}}$, and thus it can be regarded as a section of the vector bundle $(\bar{V}^*(\mathcal{M}) \times V^*(\mathcal{M}))^*$ over the manifold $\mathcal{M} \times \mathcal{M}$. It may be verified that for any $x, y \in \mathcal{M}$:

$$(A) \langle K(x, \cdot)^{\mathcal{A}'}, K(y, \cdot)^{\mathcal{B}'}, \rangle = K(y, x)^{\mathcal{B}'\mathcal{A}},$$

$$(B) \overline{K(y, x)^{\mathcal{B}'\mathcal{A}}} = K(x, y)^{\mathcal{A}'\mathcal{B}},$$

- (C) if $(\Phi_i)_{i \in \mathcal{I}}$ comprises a complete orthonormal system in $H_l(V(\mathcal{M}))$, then the identity $K(x, \cdot)^{\mathcal{A}'\mathcal{B}} = \sum_{i \in \mathcal{I}} \bar{\Phi}_i^{\mathcal{A}'}(x) \Phi_i^{\mathcal{B}}(\cdot)$ holds, where the infinite summation is understood in the H_l norm topology.

References

- [1] M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems*. Princeton University Press, 1994.
- [2] N. N. Bogolubov, A. A. Logunov, A. I. Oksak and I. T. Todorov, *General Principles of Quantum Field Theory*. Kluwer Academic Publishers, 1990.
- [3] C. J. Fewster and K. Rejzner, *Algebraic quantum field theory – an introduction*, in *in Progress and Visions in Quantum Theory in View of Gravity – Bridging Foundations of Physics and Mathematics* (F. Finster, D. Giulini, J. Kleiner and J. Tolksdorf, eds.), 2020. 1904.04051.
- [4] E. Dable-Heath, C. J. Fewster, K. Rejzner and N. Woods, *Algebraic classical and quantum field theory on causal sets*, *Phys. Rev.* **D101** (2020) 065013, [1908.01973].
- [5] S. Hollands and R. M. Wald, *Existence of local covariant time ordered products of quantum fields in curved spacetime*, *Commun.Math.Phys.* **231** (2002) 309–345, [gr-qc/0111108].
- [6] M. Dütsch and K. Fredenhagen, *Algebraic quantum field theory, perturbation theory, and the loop expansion*, *Commun. Math. Phys.* **219** (2001) 5–30, [hep-th/0001129].
- [7] R. Brunetti, K. Fredenhagen and R. Verch, *The generally covariant locality principle – a new paradigm for local quantum physics*, *Commun. Math. Phys.* **237** (2003) 31–68, [math-ph/0112041].
- [8] Y. Dabrowski and C. Brouder, *Functional properties of Hörmander’s space of distributions having a specified wavefront set*, *Commun. Math. Phys.* **332** (2014) 1345–1380, [1308.1061].
- [9] M. Dütsch, *From Classical Field Theory to Perturbative Quantum Field Theory*. Springer, 2019.

- [10] S. Hollands, *Renormalized quantum Yang-Mills fields in curved spacetime*, *Rev. Math. Phys* **20** (2008) 1033–1172, [0705.3340].
- [11] K. Costello, *Renormalization and Effective Field Theory*. American Mathematical Society, 2011.
- [12] R. P. Feynman and A. R. Hibbs, *Quantum Mechanics and Path Integrals – Emended Edition by D. F. Styer*. Dover Publications, 2010.
- [13] J. Glimm and A. Jaffe, *Quantum Physics: a Functional Integral Point of View*. Springer, 1987.
- [14] J. Velhinho, *Topics of measure theory on infinite dimensional spaces*, *Mathematics* **5** (2017) 44.
- [15] S. A. Albeverio, R. J. Høegh-Krohn and S. Mazzucchi, *Mathematical Theory of Feynman Path Integrals*. Springer, 2008.
- [16] T. L. Gill and W. W. Zachary, *Banach spaces for the Feynman integral*, *Real Analysis Exchange* **34** (2008/2009) 267–310.
- [17] J. Montaldi and O. G. Smolyanov, *Feynman path integrals and lebesgue–feynman measures*, *Doklady Mathematics* **96** (2017) 368, [1612.06657].
- [18] J. Feldbrugge, J. Lehnert and N. Turok, *No smooth beginning for spacetime*, *Phys. Rev. Lett.* **119** (2017) 171301, [1705.00192].
- [19] J. Feldbrugge, J. Lehnert and N. Turok, *Lorentzian quantum cosmology*, *Phys. Rev.* **D95** (2017) 103508, [1703.02076].
- [20] A. Baldazzi, R. Percacci and V. Skrinjar, *Quantum fields without wick rotation*, *Symmetry* **11** (2019) 373, [1901.01891].
- [21] T. Weigand, “Quatum field theory II.” Heidelberg University Lecture Notes, 2014.
- [22] G. Sardanasvily, “Fibre bundles, jet manifolds and lagrangian theory. lectures for theoreticians.” Moscow State University Lecture Notes, 2009.
- [23] R. Cohen, *The Topology of Fiber Bundles*. AMS Open Math Notes Series, 2017.
- [24] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time*. Cambridge University Press, 1973.
- [25] H.-J. Borchers, *On structure of the algebra of field operators*, *Nuovo Cimento* **24** (1962) 214–236.
- [26] A. Uhlmann, *Über die Definitionen des Quantenfelder nach Wightman und Haag*, *Wiss. Z. KMU Leipzig* **11** (1962) 213–217.
- [27] J. Yngvason, *On the algebra of test functions for field operators*, *Commun. Math. Phys.* **34** (1973) 315–333.
- [28] D. A. Dubin and M. A. Hennings, *Symmetric tensor algebras and integral decompositions*, *Publ. RIMS. Kyoto Univ.* **25** (1989) 1001–1020.
- [29] B. Malgrange, *Existence et approximation des solutions des équations aux dérivées partielles et des équations de convolution*, *Annales de l’Institut Fourier* **6** (1956) 271–355.
- [30] L. Ehrenpreis, *Solution of some problems of division: Part I. Division by a polynomial of a derivation*, *Americal Journal of Mathematics* **76** (1954) 883–903.
- [31] L. Hörmander, *The Analysis of Linear Partial Differential Operators I*. Springer, 1990.
- [32] L. Hörmander, *The Analysis of Linear Partial Differential Operators III*. Springer, 2007.
- [33] M. A. Shubin, *Pseudodifferential Operators and Spectral Theory*. Springer, 2001.
- [34] M. J. Radzikowski, *Micro-local approach to the Hadamard condition in quantum field theory on curved space-time*, *Commun.Math.Phys.* **179** (1996) 529–553.
- [35] F. Andersson and M. Carlsson, *On general domain truncated correlation and convolution operators with finite rank*, *Integral Equations and Operator Theory* **82** (2015) 339–370.
- [36] J. de Jong et al, “Stacks project.” Columbia University lecture notes, 2021.
- [37] D. A. Dubin and M. A. Hennings, *Regular tensor algebras*, *Publ. RIMS. Kyoto Univ.* **25** (1989) 971–999.
- [38] D. Vogt, *The tensor algebra of power series spaces*, *Studia Mathematica* **193** (2009) 189–202.
- [39] A. Pietsch and W. H. Ruckle, *Nuclear Locally Convex Spaces*. Springer, 1972.
- [40] W. Rudin, *Functional Analysis*. McGraw-Hill, 1991.
- [41] J. L. Taylor, “Notes on locally convex topological vector spaces (supplementary material to Rudin: Functional Analysis).” University of Utah Lecture Notes, 1995.

- [42] M. Schechter, *Differentiation in abstract spaces*, *J. Diff. Eq.* **55** (1984) 330.
- [43] F. Trèves, *Topological Vector Spaces, Distributions and Kernels*. Academic Press New York, 1970.
- [44] J. Horváth, *Topological Vector Spaces and Distributions*. Addison–Wesley, 1966.
- [45] H. Bourlès, *Fundamentals of Advanced Mathematics 2*. Elsevier, 2018.
- [46] H. H. Schaefer and M. P. Wolff, *Topological Vector Spaces*. Springer, 1999.
- [47] J. Becnel and A. Sengupta, *Nuclear space facts, strange and plain*, *Mathematics* **4** (2016) 61.
- [48] P. Garrett, “Schwartz kernel theorems, tensor products, nuclearity.” University of Minnesota Lecture Notes, 2020.
- [49] T. Hirai, H. Shimomura, N. Tatsuuma and E. Hirai, *Inductive limits of topologies, their direct products, and problems related to algebraic structures*, *J. Math. Kyoto. Univ.* **41** (2001) 475–505.
- [50] M. J. Chasco and X. Domínguez, *Topologies on the direct sum of topological Abelian groups*, *Topology and its Applications* **133** (2003) 209–223.
- [51] J. H. Webb, *Sequential convergence in locally convex spaces*, *Proc. Camb. Phil. Soc.* **64** (1968) 341.
- [52] R. M. Dudley, *On sequential convergence*, *Transactions of the American Mathematical Society* **112** (1964) 483–507.
- [53] S. Gabrielyan, *Topological properties of strict (LF)-spaces and strong duals of Montel strict (LF)-spaces*, *Monatshefte für Mathematik* **189** (2019) 91–99, [1702.07867].
- [54] R. Beattie and H.-P. Butzmann, *Convergence Structures and Applications to Functional Analysis*. Springer, 2002.
- [55] R. Beattie and H.-P. Butzmann, *Sequentially determined convergence spaces*, *Czechoslovak Mathematical Journal* **37** (1987) 231–247.
- [56] R. Beattie, *Continuous convergence and functional analysis*, *Topology and its Applications* **70** (1996) 101–111.
- [57] R. Beattie and H.-P. Butzmann, *On the banach–steinhaus theorem and the continuity of bilinear mappings*, *Math. Nachr.* **153** (1991) 297–312.
- [58] N. Falkner, *Closable operators and semigroups*, *Proceedings of the AMS* **87** (1983) 107–110.
- [59] A. Antonevich, A. Buraczewski and Y. Radyno, *On closability of nonclosable operators*, *Panamerican Mathematical Journal* **7** (1997) 37–51.
- [60] I. M. Gel’fand and G. E. Shilov, *Generalized Functions (vol.2)*. Academic Press, 1968.
- [61] V. I. Bogachev and O. G. Smolyanov, *Topological Vector Spaces and Their Applications*. Springer, 2017.
- [62] M. Merkle, *Completion of countably seminormed spaces*, *Acta Math. Hungar.* **80** (1998) 1.
- [63] M. J. Merkle, “Multi-Hilbertian spaces and their duals.” Technical Report No.291 University of North Carolina, 1990.
- [64] J. Becnel and A. Sengupta, *The Schwartz space: tools for quantum mechanics and infinite dimensional analysis*, *Mathematics* **3** (2015) 527.
- [65] R. A. Adams and J. J. F. Fournier, *Sobolev spaces*. Elsevier, 2003.
- [66] Y. Choquet-Bruhat and C. DeWitt-Morette, *Analysis, Manifolds and Physics II*. Elsevier, 2000.
- [67] P. Csizmadia, A. László and I. Rácz, *On the use of multipole expansion in time evolution of nonlinear dynamical systems and some surprises related to superradiance*, *Class. Quant. Grav.* **30** (2013) 015010, [1207.5837].
- [68] N. Aronszajn, *Theory of reproducing kernels*, *Transactions of AMS* **68** (1950) 337–404.