

Strong dispersion property for the quantum walk on the hypercube

Martins Kokainis¹, Krišjānis Prūsis¹, Jevgēnijs Vihrovs¹, Vyacheslavs Kashcheyevs², and Andris Ambainis¹

¹Centre for Quantum Computer Science, Faculty of Computing, University of Latvia, Raiņa 19, Riga, Latvia, LV-1586

²Department of Physics, University of Latvia, Jelgavas 3, Riga, Latvia, LV-1004

We show that the discrete time quantum walk on the Boolean hypercube of dimension n has a strong dispersion property: if the walk is started in one vertex, then the probability of the walker being at any particular vertex after $O(n)$ steps is of an order $O(1.4818^{-n})$. This improves over the known mixing results for this quantum walk which show that the probability distribution after $O(n)$ steps is close to uniform but do not show that the probability is small for every vertex. A rigorous proof of this result involves an intricate argument about analytic properties of Bessel functions.

1 Introduction

Quantum walks are an important tool for designing quantum algorithms, from the exponential speedup for the “glued trees” problem by Childs et al. [9] and the element distinctness algorithm of [2] to general results about speeding up classes of Markov chains [4–6, 23]. Quantum walks also have applications to other areas (e.g. quantum state transfer [14] or Hamiltonian complexity [16]) and are interesting objects of study on their own.

Quantum walk on the hypercube (introduced by Moore and Russell [15]) is among the most popular quantum walks and has been studied by a number of researchers [1, 11, 12, 19]. The hypercube was the first graph for which a search by quantum walk algorithm was developed, by Shenvi et al. [22]. This was the starting point for a line of work which has now led to many interesting results, from [23] to [6].

The Boolean hypercube is also the simplest graph on which a quantum walk mixes rapidly. Namely, the n -dimensional hypercube has 2^n vertices but, as shown in [15], after $\lceil \frac{\pi}{4}n \rceil$ steps, the probability distribution over the vertices becomes close to uniform in variational distance (the variational distance is $o(n^{-7/6})$). The closeness in variational distance does not, however, imply that the probability of every vertex is small (because a substantial fraction of the difference can be concentrated on a small number of vertices).

In this paper, we show this is actually the case and the discrete quantum walk disperses so well that, if a walker is started in one vertex (with the coin register being in a uniform superposition of all possible directions), then, after $O(n)$ steps, the probability of the walker being in each vertex becomes exponentially small.

Our computer simulations show that, after $0.849\dots n$ steps, the probability of being at each vertex is at most $1.93\dots^{-n}$. Since the hypercube has 2^n vertices, the quantum walk is close to achieving

Martins Kokainis: martins.kokainis@lu.lv

Krišjānis Prūsis: krisjanis.prusis@lu.lv

Jevgēnijs Vihrovs: jevgenijs.vihrovs@lu.lv

Vyacheslavs Kashcheyevs: vjaceslavs.kascejevs@lu.lv

Andris Ambainis: andris.ambainis@lu.lv

the biggest possible dispersion. (In the uniform distribution, each vertex has the probability 2^{-n} .)

Proving this rigorously is surprisingly difficult and requires a complicated argument based on analytic properties of Bessel functions. With this argument, we can show a result of a similar asymptotic form with somewhat weaker constants: the probability of being at each vertex after $0.8663n$ steps is at most $O(1.4818^{-n})$.

Remark. We note that an even stronger dispersion property is known for the continuous-time walk on the hypercube. Namely [15], after $\frac{\pi}{4}n$ time, the probability distribution of the continuous-time walker over the vertices of the hypercube is exactly uniform and this is quite easy to prove, because the Hamiltonian of the continuous time walk can be expressed as a sum of Hamiltonians corresponding to each dimension of the hypercube. The matrix of the discrete time walk does not factorize into parts corresponding to each dimension and this makes the result for the discrete time walk much more challenging.

Remark. The original motivation for this work was a result by Brandao and Horodecki [8] that any unitary U , such that $|\langle a|U|b\rangle| \leq 2^{-k}$ for any states $|a\rangle, |b\rangle$ of the computational basis, can be used to construct a query problem that is solvable quantumly with $O(1)$ queries and $O(1)$ applications of U but requires $2^{\Omega(k)}$ queries classically. If a strong dispersion result similar to ours was true for the same quantum walk with a starting state $|x, i\rangle$ with the walker in one vertex x and the coin register in a basis state corresponding to one direction i , this would imply a superexponential quantum query separation with the quantum walk on the hypercube as the main non-query transformation in the quantum algorithm. However, due to the phenomenon of oscillatory localization explored in [3], the walker stays in one of the vertices close to x with a high probability if the walk is started in $|x, i\rangle$.

2 Results

For a positive integer n , let $[n]$ denote the set $\{1, \dots, n\}$.

The n -dimensional Boolean hypercube Q_n is a graph with 2^n vertices indexed by $x \in \{0, 1\}^n$ and edges (x, y) for x, y that differ in one coordinate. The notation $\{0\}^n$ is shortened to 0^n . Let $|x|$ denote the Hamming weight of a vertex x , defined as the number of $i \in [n]$ with $x_i = 1$. For $x \in \{0, 1\}^n$ and $i \in [n]$, $x^{(i)}$ denotes the vertex obtained from x by changing the i^{th} component:

$$x^{(i)} = (x_1, \dots, x_{i-1}, 1 - x_i, x_{i+1}, \dots, x_n). \quad (2.1)$$

We consider the standard discrete-time quantum walk on the hypercube [15], with the Grover diffusion operator as the coin flip. The state space of this walk has basis states $|x, i\rangle$ where $x \in \{0, 1\}^n$ is a vertex of Q_n and $i \in [n]$ is an index for one of the directions for the edges of the hypercube.

One step of the quantum walk consists of two parts:

1. We apply the transformation D_n defined by $D_n |x, i\rangle = -\frac{n-2}{n} |x, i\rangle + \frac{2}{n} \sum_{j \in [n] \setminus \{i\}} |x, j\rangle$ for all $x \in \{0, 1\}^n$ and $i \in [n]$.
2. We apply the shift transformation S defined by $S |x, i\rangle = |x^{(i)}, i\rangle$.

We denote the sequence of these two transformations by U . The walk is started in the state $|\psi_{start}\rangle = \sum_{i=1}^n \frac{1}{\sqrt{n}} |0^n, i\rangle$ where the walker is localized in one vertex and the direction register is in the uniform superposition of all possible directions.

Let

$$|\psi^{(t)}\rangle = \sum_{x \in \{0, 1\}^n, i \in [n]} \alpha_{x, i}^{(t)} |x, i\rangle \quad (2.2)$$

be the state of the quantum walk after t steps and $P(x, t) = \sum_i |\alpha_{x, i}^{(t)}|^2$ be the probability of the walker being at location x at this time.




Figure 1: Illustration of the relevant quantities for a quantum walk on the hypercube. **Upper left:** the maximum probability of a vertex during a walk on the 50-dimensional hypercube. **Upper right:** the number of steps t_{\min} to reach the minimum of $\max_x P(x, t)$ (the cross symbols). The solid line is the graph of the function $-0.754 + 0.849n$, which approximates t_{\min} . **Lower left:** the probability $\max_x P(x, t_{\min})$ over the vertices (at time $t_{\min} \approx 0.849n$) (the cross symbols). The solid line is the graph of the function $5 \cdot 1.93^{-n}$, which is an upper bound on the probability. **Lower right:** on the 50-dimensional hypercube, the maximum probability remains at the starting vertex on even steps until t_{\min} . The cross symbols denote the maximum probability amongst all the vertices. The square markers denote the probability at the 0^n vertex. Only even steps are shown as the probability at 0^n is 0 at odd steps. The data that support the graphs of this figure are available in the Zenodo repository <https://doi.org/10.5281/zenodo.5907185>.

After $t \approx 0.85n$ steps, the walker disperses over the vertices $x \in \{0, 1\}^n$ very well, with no particular vertex x having a substantial probability $P(x, t)$ of the walker being there.

In Fig. 1, the upper left panel shows the maximum probability of a single vertex $\max_x P(x, t)$ at every step of a quantum walk on a 50-dimensional hypercube. We can see that this probability reaches a minimum of about 10^{-14} (which is only slightly larger than the theoretical minimum of $2^{-50} = 0.88\dots \cdot 10^{-16}$) after a number of steps that is slightly less than $n = 50$.

We note that $\max_x P(x, t)$ fluctuates between odd and even numbered steps, due to the walker being at an odd distance from the starting vertex 0^n after an odd number of steps and at an even distance from 0^n after an even number of steps. This effect is particularly pronounced when $\max_x P(x, t)$ is large. Then, a large fraction of the probability is concentrated on 0^n after even steps but, after odd steps, this probability is equally divided among n vertices x with $|x| = 1$. As a result, the maximum $\max_x P(x, t)$ is visibly larger after an even number of steps. This effect becomes smaller when $\max_x P(x, t)$ is small.

The upper right panel of Fig. 1 shows that the number of steps t_{\min} required to minimize $\max_x P(x, t)$ grows linearly with n , and is achieved at approximately $-0.754 + 0.849n$.

The probability $\max_x P(x, t_{\min})$ achieved at this t_{\min} is approximately $5 \cdot 1.93^{-n}$ (see the lower left panel of Fig. 1).

Lastly, this maximum is achieved at $x = 0^n$ (at even numbered steps) or at vertices x with $|x| = 1$ (for odd numbered steps). This is shown in the lower right panel of Fig. 1 where we plot $\max_x P(x, t)$ and $P(0^n, t)$ for even numbered steps t . The panel shows that the maximum is achieved at $x = 0^n$ until the moment when the probability $\max_x P(x, t)$ starts increasing again.

Rigorously, we can prove a weaker bound:

Theorem 1. *For $t = \lfloor 0.8663n \rfloor$, we have $\max_x P(x, t) = O(1.4818^{-n})$.*

We first describe the strategy of the proof. Because of the symmetry of the walk, all vertices x with the same $|x| = k$ will have equal probabilities $P(x, t)$. Since there are $\binom{n}{k}$ vertices x , this immediately implies $P(x, t) \leq 1/\binom{n}{k}$. If k is such that $\binom{n}{k}$ is sufficiently large, we get the desired upper bound on $P(x, t)$.

It remains to handle the case when $\binom{n}{k}$ is small. This corresponds to k being either close to 0 ($0 \leq k < 0.13368n$) or close to n ($(1 - 0.13368)n < k \leq n$). The second case is trivial: the number of time steps $t = 0.8663n$ that we are considering is less than $(1 - 0.13368)n$, so, vertices x with $|x| > (1 - 0.13368)n$ cannot be reached in t steps.

For the first case, we show that if $P(x, t)$ is large, then $P(0^n, t')$ must also be non-negligible for some $t' \in \{t - |x|, \dots, t + |x|\}$ (Lemma 1). Therefore, one can show an upper bound on all $P(x, t)$ with $|x| < 0.13368n$ by upper bounding $P(0^n, t')$ for all $t' : t - 0.13368n \leq t' \leq t + 0.13368n$. This is done by Lemma 3 and Theorem 2, first expressing $P(0^n, t')$ in terms of Jacobi polynomials and then bounding their asymptotics.

The rest of the paper is devoted to the implementation of the above strategy.

3 Proof of the main result

3.1 Relating the probability to be at an arbitrary vertex with the probability to be at the initial vertex

We denote

$$|w, \rightarrow\rangle = \frac{1}{\sqrt{\binom{n}{w}(n-w)}} \sum_x \sum_{\substack{i \\ |x|=w \\ x_i=0}} |x, i\rangle, \quad (3.1)$$

$$|w, \leftarrow\rangle = \frac{1}{\sqrt{\binom{n}{w}w}} \sum_x \sum_{\substack{i \\ |x|=w \\ x_i=1}} |x, i\rangle. \quad (3.2)$$

By symmetry, the state of the quantum walk after any number of steps t is of the form

$$|\psi^{(t)}\rangle = \sum_{w=0}^{n-1} \alpha_{w,\rightarrow}^{(t)} |w, \rightarrow\rangle + \sum_{w=1}^n \alpha_{w,\leftarrow}^{(t)} |w, \leftarrow\rangle. \quad (3.3)$$

Let

$$P[w, t] = |\alpha_{w,\rightarrow}^{(t)}|^2 + |\alpha_{w,\leftarrow}^{(t)}|^2 \quad (3.4)$$

be the total probability of the walker being at one of vertices x with $|x| = w$ after t steps. By the symmetry of the quantum walk,

$$P(x, t) = P[w, t] / \binom{n}{w} \quad (3.5)$$

for any $x : |x| = w$. In particular, $P(0^n, t) = P[0, t]$.

Assume that the minimum of $P[0, t]$ is achieved at time t . We want to prove that at the same time $P[w, t]$ is also small for all w . For this, we have the following lemma:

Lemma 1. *Suppose that $n \geq 2$ and $P[0, t'] \leq p_0$ for all $t' \in [t - w', t + w']$, where $w' < n/2$. Then for all $w \in \{0, 1, \dots, w'\}$, we have*

$$P[w, t] \leq \frac{n^w}{w!} p_0. \quad (3.6)$$

Proof. We prove the contrapositive: suppose that $P[w, t] = p_w$ for some $0 < w < n/2$; then there exists a $t' \in [t - w, t + w]$ such that $P[0, t'] \geq \frac{w! p_w}{n^w}$.

We do this by showing two inequalities:

$$\max\left(|\alpha_{w,\leftarrow}^{(t)}|^2, |\alpha_{w-1,\rightarrow}^{(t+1)}|^2\right) \geq \frac{w}{n-w} |\alpha_{w,\rightarrow}^{(t)}|^2, \quad (3.7)$$

$$P[w-1, t-1] \geq |\alpha_{w,\leftarrow}^{(t)}|^2. \quad (3.8)$$

These two inequalities imply that one of $P[w-1, t-1], P[w-1, t+1]$ is at least

$$\min\left(\frac{w}{n-w} |\alpha_{w,\rightarrow}^{(t)}|^2, |\alpha_{w,\leftarrow}^{(t)}|^2\right). \quad (3.9)$$

Because of (3.4), we must either have $|\alpha_{w,\rightarrow}^{(t)}|^2 \geq \frac{n-w}{n} P[w, t]$ or $|\alpha_{w,\leftarrow}^{(t)}|^2 \geq \frac{w}{n} P[w, t]$. In both cases (3.9) is at least $\frac{w}{n} P[w, t]$.

By repeating this argument w times, we get that $P[0, t']$, for some $t' \in [t - w, t + w]$ is at least

$$p_w \cdot \frac{w}{n} \cdot \frac{w-1}{n} \cdot \dots \cdot \frac{1}{n} = \frac{w! p_w}{n^w}. \quad (3.10)$$

We now prove (3.7) and (3.8). To prove (3.7), consider vertices $x \in \{0, 1\}^n$ for which $|x| = w$. Before the coin flip, the amplitudes of $|x, i\rangle$ with $x_i = 0$ are equal to $\frac{\alpha_{w,\rightarrow}^{(t)}}{\sqrt{\binom{n-w}{w}}}$ and the amplitudes of $|x, i\rangle$ with $x_i = 1$ are equal to $\frac{\alpha_{w,\leftarrow}^{(t)}}{\sqrt{w \binom{n}{w}}}$. After applying the coin flip C , the amplitudes of $|x, i\rangle$ with $x_i = 1$ become equal to

$$\frac{2(n-w)}{n} \frac{\alpha_{w,\rightarrow}^{(t)}}{\sqrt{\binom{n-w}{w}}} - \frac{n-2w}{n} \frac{\alpha_{w,\leftarrow}^{(t)}}{\sqrt{w \binom{n}{w}}}. \quad (3.11)$$

After the shift operation, each of those becomes an amplitude of $|y, i\rangle$ with $|y| = w-1$ and $y_i = 0$. Since $|w-1, \rightarrow\rangle$ consists of $\binom{n}{w} - 1(n-w+1) = \binom{n}{w} w$ such $|y, i\rangle$, we have

$$\alpha_{w-1,\rightarrow}^{(t+1)} = \frac{2\sqrt{w(n-w)}}{n} \alpha_{w,\rightarrow}^{(t)} - \frac{n-2w}{n} \alpha_{w,\leftarrow}^{(t)}. \quad (3.12)$$

Assume that $|\alpha_{w,\leftarrow}^{(t)}| < \frac{\sqrt{w}}{\sqrt{n-w}} |\alpha_{w,\rightarrow}^{(t)}|$. (Otherwise, (3.7) is immediately true.) Then,

$$|\alpha_{w-1,\rightarrow}^{(t+1)}| \geq \left(\frac{2\sqrt{w(n-w)}}{n} - \frac{\sqrt{w}}{\sqrt{n-w}} \frac{n-2w}{n} \right) |\alpha_{w,\rightarrow}^{(t)}|. \quad (3.13)$$

The equation (3.7) now follows from

$$\frac{2\sqrt{w(n-w)}}{n} - \frac{\sqrt{w}}{\sqrt{n-w}} \frac{n-2w}{n} = \frac{\sqrt{w}}{\sqrt{n-w}}. \quad (3.14)$$

To prove (3.8), we simply note that $S|w-1,\rightarrow\rangle = |w,\leftarrow\rangle$. Since coin flip C moves the amplitudes between $|w-1,\leftarrow\rangle$ and $|w-1,\rightarrow\rangle$, this means that all the amplitude at $|w,\leftarrow\rangle$ after t steps must have been at $|w-1,\leftarrow\rangle$ or $|w-1,\rightarrow\rangle$ one step earlier. \square

3.2 Bounding the probability to be at the initial vertex

Next, we prove a bound on $P[0,t]$. For this, we utilize the following representation of Chebyshev polynomials of even order, valid for $|z| \leq 1$:

Lemma 2. *Let t be a positive even integer; then the following equality is valid for all $z \in [-1, 1]$:*

$$T_t(z) = (-1)^{t/2} t \int_0^\infty x^{-1} J_t(x) \cos(xz) dx, \quad (3.15)$$

where T_t is the t^{th} -degree Chebyshev polynomial of the first kind and J_t stands for the Bessel function of the first kind.

Proof. We start with a special case of the Sonine-Schafheitlin formula [24, p.405, §13.42, Eq. (3)]

$$\int_0^\infty \frac{J_\mu(ax) \cos(bx)}{x} dx = \mu^{-1} \cos(\mu \arcsin(b/a)), \quad 0 < b \leq a. \quad (3.16)$$

Taking $a = 1$, $b = z \in (0, 1]$ and $\mu = t$ in (3.16) gives

$$\int_0^\infty \frac{J_t(x) \cos(xz)}{x} dx = \frac{\cos(t \cdot \arcsin z)}{t}. \quad (3.17)$$

It remains to notice that

$$\cos(t \cdot \arcsin z) = \cos\left(t \left(\frac{\pi}{2} - \arccos z\right)\right) = (-1)^{t/2} \cos(t \arccos z) = (-1)^{t/2} T_t(z). \quad (3.18)$$

Continuity and evenness of \cos and T_t ensures the validity of (3.15) for $z \in [-1, 0]$. \square

This lemma allows us to characterize the probability $P[0,t]$ to obtain the vertex 0^n after t steps for even t as follows:

Lemma 3.

$$\sqrt{P[0,t]} = \frac{1}{2^n} \sum_{m=0}^n \binom{n}{m} T_t\left(1 - \frac{2m}{n}\right) = t \left| \int_0^\infty x^{-1} J_t(x) \cos^n \frac{x}{n} dx \right| \quad (3.19)$$

Proof. Due to the symmetry of the walk (w.r.t. permuting the coordinates of the hypercube), the amplitudes of $|0^n, j\rangle$ for all $j \in [n]$ remain equal after any number of steps. Therefore,

$$P[0,t] = \langle \psi_{start} | U^t | \psi_{start} \rangle^2. \quad (3.20)$$

Let $|v_j\rangle$ be the eigenvectors of U , with eigenvalues e^{iw_j} . We can decompose $|\psi_{start}\rangle$ as a sum of eigenvectors: $|\psi_{start}\rangle = \sum_j a_j |v_j\rangle$. Then,

$$\sqrt{P[0,t]} = \langle \psi_{start} | U^t | \psi_{start} \rangle = \sum_j |a_j|^2 e^{itw_j}. \quad (3.21)$$

As described in [15], U has 2^{n+1} eigenvectors that have non-zero overlap with the starting state. These eigenvectors can be indexed by $k \in \{0,1\}^n$. For each $k \in \{0,1\}^n$, we have a pair of eigenvectors $|v_k\rangle$ and $|v_{*k}\rangle$ with eigenvalues $e^{\pm iw_m}$ where $m = |k|$ is the Hamming weight of k and w_m is such that $\cos w_m = 1 - \frac{2m}{n}$. Furthermore, all of those eigenvectors have equal overlap with the starting state: $a_j = |\langle v_k | \psi_{start} \rangle|^2 = |\langle v_{*k} | \psi_{start} \rangle|^2 = \frac{1}{2^{n+1}}$. Therefore, we get

$$\sqrt{P[0,t]} = \frac{1}{2^n} \sum_{k \in \{0,1\}^n} \cos w_m t \quad (3.22)$$

$$= \frac{1}{2^n} \sum_{m=0}^n \binom{n}{m} \cos w_m t \quad (3.23)$$

$$= \frac{1}{2^n} \sum_{m=0}^n \binom{n}{m} \cos \left(t \cdot \arccos \left(1 - 2 \cdot \frac{m}{n} \right) \right) \quad (3.24)$$

$$= \frac{1}{2^n} \sum_{m=0}^n \binom{n}{m} T_t \left(1 - 2 \cdot \frac{m}{n} \right). \quad (3.25)$$

Now, using (3.15) with $z = (n - 2m)/n$ for even integer t ,

$$\frac{1}{2^n} \sum_{m=0}^n \binom{n}{m} T_t \left(1 - \frac{2m}{n} \right) = (-1)^{t/2} t \int_0^\infty x^{-1} J_t(x) \sum_{m=0}^n \frac{1}{2^n} \binom{n}{m} \cos \left(x - \frac{xm}{n} \right) dx \quad (3.26)$$

$$= (-1)^{t/2} t \int_0^\infty x^{-1} J_t(x) \cos^n \frac{x}{n} dx \quad (3.27)$$

with the last step following from the binomial expansion of $(e^{ix/n} + e^{-ix/n})^n$. \square

We point out that the RHS of Lemma 3 turns out more suitable (compared to the expression using only Chebyshev's polynomials) for estimating $\sqrt{P[0,t]}$ in part due to a clear separation between the oscillating part (J_t) and the exponential part ($\cos^n(x/n)$). This distinction helps to explain how the ratio t/n leads to the oscillating behavior of $P[0,t]$ observed in the lower right panel of Fig. 1 (with $n = 50$). When t/n is small (e.g., see the upper row in Fig. 2 with $n = 50, t = 0.12n$), the Bessel function's first maximum largely overlaps with the contribution of $\cos^n(\cdot)$ in the vicinity of $x = 0$. However, the subsequent maxima of $\cos^n(\cdot)$ occur where the oscillations of J_t cancel out, resulting in an insignificant contribution to the integral. Therefore, the value of the integral is dominated by the values of the integrand near $x = 0$ and the probability $P[0,t]$ is large.

In contrast, consider the case when t/n is "large" (see the middle row in Fig. 2 for the case $n = 50, t = 3.12n$). While the Bessel function's values near $x = 0$ are negligible, the following local maximum greatly overlaps with the next peak of $\cos^n(x/n)$ (in the vicinity of $x \approx \pi n$). This results in a large contribution to the integral, which is not canceled out by the following peaks of $\cos^n(x/n)$ (where the Bessel function's fluctuations make the contribution to the integral negligible).

Finally, when t/n is "just right" (see the lower row in Fig. 2 for the case $n = 50, t = 0.84n$), the first maxima of Bessel's functions occur when $\cos^n(x/n)$ is near zero. This ensures that the resulting integrand's oscillations near each peak of $\cos^n(x/n)$ largely cancel out. We proceed to quantify the remaining integral's value and show that it indeed is exponentially small in n .

We will upper-bound the integral appearing in Lemma 3, provided that t/n is suitably bounded. In what follows, we use $\nu \in \mathbb{R}$ instead of t (since the latter was assumed to be an even integer in our notation before).




Figure 2: Illustration of the integrand and its factors for $n = 50$ and $t = 0.12n$ (upper row), $t = 3.12n$ (middle row), $t = 0.84n$ (lower row). Panels on the left show $J_t(x)$, solid line, with units on the left axis and $\cos^n(x/n)$, dash-dotted line, with units on the right axis. Panels on the right depict the integrand $x^{-1} J_t(x) \cos^n(x/n)$. The dotted gridlines divide the real line into subintervals separating the peaks of $\cos^n(x/n)$.

Theorem 2. Denote $a_k = (k - 0.5)\pi$, $k = 1, 2, 3, \dots$; let a positive constant $\alpha \in (\frac{\pi}{6}, 1)$ be fixed.

For $n \geq 2$ and $\nu > 1$, satisfying $\nu \in (n\alpha, n) \subset (\frac{\pi n}{6}, n)$, the following estimates hold:

$$\text{The tail:} \quad \left| \int_{na_n}^{\infty} x^{-1} J_\nu(x) \cos^n\left(\frac{x}{n}\right) dx \right| < \frac{100\sqrt{n}}{2^n}; \quad (3.28)$$

$$\text{The middle part:} \quad \left| \int_{na_1}^{na_n} x^{-1} J_\nu(x) \cos^n\left(\frac{x}{n}\right) dx \right| < \frac{4000\sqrt{n}}{1.541^n}; \quad (3.29)$$

$$\text{The bulk:} \quad \left| \int_0^{na_1} x^{-1} J_\nu(x) \cos^n\left(\frac{x}{n}\right) dx \right| < \frac{3}{(1 + \alpha)^{0.5\alpha n}}. \quad (3.30)$$

The proof of the theorem is deferred to Appendix A; below we sketch a rough reasoning behind the proof.

At first glance, both expressions of $\sqrt{P[0, t]}$ (via the sum involving Chebyshev polynomials or via the integral) suffer the same drawback (or the advantage) of catastrophic cancellations. Indeed, while the integrand's extreme values near each peak of $\cos^n(x/n)$ are polynomially decaying in n , cancellations ensure that the overall contribution is exponentially small! This property prohibits from making simple estimates of $\sqrt{P[0, t]}$ in either case. However, here the main advantage of the integral representation manifests itself: instead of computing integrals over real intervals, we use the Cauchy integral theorem to transform them to contour integrals in the complex plane. Since the integrand turns out to be non-oscillatory in the complex plane in directions orthogonal to the real axis, this provides an efficient way to estimate the value of $P[0, t]$.

As Fig. 2 might suggest, it is advantageous to divide the whole integral into a sum of integrals over subintervals of the form $[(k - 0.5)\pi n, (k + 0.5)\pi n]$, separating the individual peaks of $\cos^n(x/n)$ and estimating the contribution of each integral separately. It turns out that there are three regimes to consider:

1. When $k \geq n$, the integral over each subinterval $[(k - 0.5)\pi n, (k + 0.5)\pi n]$ turns out to be of order $2^{-n}n^{1/2}k^{-5/2}$; the series (taking the sum over $k : k \geq n$) converges, leading to the tail estimate (3.28).
2. When $1 \leq k < n$, we estimate the integral over each subinterval $[(k - 0.5)\pi n, (k + 0.5)\pi n]$ as $1.541^{-n}n^{-1/2}$, with the constant 1.541 coming from an upper bound on a certain auxiliary function. Since there are $n - 1$ such subintervals, this leads to the estimate of the middle part (3.29).
3. In both previous cases, the integral is going to be estimated via an excursion into the complex plane. However, for the remaining interval $(0, 0.5\pi n)$ this is not a viable option anymore. Instead, here we combine direct bounds on the Bessel function J_ν and the $\cos^n(x/n)$ term when x is “small”. These estimates (in particular, the estimate of the integral over the interval $[0, \nu]$) lead to the dominating contribution (“the bulk”) (3.30).

To estimate the integrals of the form $\int_{na_k}^{na_{k+1}} x^{-1} J_\nu(x) \cos^n\left(\frac{x}{n}\right) dx$, we form a rectangle in the complex plane, consisting of the real interval $[na_k, na_{k+1}]$, a parallel interval $[na_k + iR, na_{k+1} + iR]$ for some $R > 0$, as well as two segments orthogonal to the real axis forming a closed contour. The Cauchy integral theorem then implies that the integral over the real integral coincides with the integral over the rest of the contour. While the integrand still is oscillatory over the set $[na_k + iR, na_{k+1} + iR]$, its absolute value vanishes as $R \rightarrow +\infty$. Thus, taking the limit $R \rightarrow +\infty$, we are left with integrals over the rays of the form $[na_k, na_k + i\infty)$.

For this strategy to work, we need the integrand to be holomorphic (in a domain containing the integration contour). We note that the integrand $x^{-1} J_\nu(x) \cos^n\left(\frac{x}{n}\right)$ is the real part of the function

$$f(z) = \frac{H_\nu^{(1)}(z) \cos^n(z/n)}{z}, \quad z \in \mathbb{C}, |z| > 1, \quad (3.31)$$

where $H_\nu^{(1)}$ is the Hankel function of the first kind¹ (and satisfies $\Re H_\nu^{(1)}(x) = J_\nu(x)$ for real x).

¹By slightly abusing terminology, we will refer to $H_\nu^{(1)}$ as *the* Hankel function.

Now f is holomorphic in the necessary domain, and we make use of the estimate

$$\left| \int_{na_k}^{na_{k+1}} x^{-1} J_\nu(x) \cos^n\left(\frac{x}{n}\right) dx \right| = \left| \Re \int_{na_k}^{na_{k+1}} f(x) dx \right| \leq \left| \int_{na_k}^{na_{k+1}} f(x) dx \right|. \quad (3.32)$$

Finally, to bound the values of f on the rays $[na_k, na_{k+1} + i\infty)$, we will employ asymptotic expansions of the Hankel function $H_\nu^{(1)}(z)$. The case of large z is straightforward [10, §10.17(iv)]. However, when the argument z and order ν are of similar magnitude, there is no suitable expansion of the Hankel function with error bounds (to the best of our knowledge). This leads to the most technical contribution of this paper, where we relate the Hankel function to the modified Bessel function of the second kind, consider its asymptotic expansion and bound the relevant error term, as detailed in Appendix A.8.

3.3 Concluding the proof

Finally, we are ready to show Theorem 1, restated here for convenience.

Theorem 1. *For $t = \lfloor 0.8663n \rfloor$, we have $\max_x P(x, t) = O(1.4818^{-n})$.*

Proof. Let² $t = \lfloor 0.8663n \rfloor$ and $w' = \lfloor 0.13368n \rfloor$; set $\alpha = 0.7326$ in Theorem 2. We will assume that n is sufficiently large, more precisely, that $n \geq 2$, $[t - w', t + w'] \subset (0.7326n, n)$, and $n > n_0$, where n_0 is described below.

Then $(1 + \alpha)^{0.5\alpha} > 1.22302$ and Theorem 2 gives the estimate (for $t' \in (\alpha n, n)$)

$$\left| \int_0^\infty x^{-1} J_{t'}(x) \cos^n\left(\frac{x}{n}\right) dx \right| < \frac{100\sqrt{n}}{2^n} + \frac{4000\sqrt{n}}{1.541^n} + \frac{3}{1.22302^n} = O(1.22302^{-n}). \quad (3.33)$$

This together with Lemma 3 implies that there are positive constants C_0, C_1 and a positive integer n_0 such that for all $n > n_0$ and all $t' \in (\alpha n, n) = (0.7326n, n)$ the probability $P[0, t']$ can be estimated as

$$P[0, t'] < \frac{C_0 n t'^2}{1.22302^{2n}} \leq \frac{C_0 n^3}{1.49578 \dots^n} < \frac{C_1}{1.49578^n}, \quad t' \in (0.7326n, n). \quad (3.34)$$

In particular,

$$P[0, t'] < \frac{C_1}{1.49578^n} \quad \text{for all } t' \in [t - w', t + w'] \subset (0.7326n, n). \quad (3.35)$$

Fix any $x \in \{0, 1\}^n$ and let $w = |x|$. If $w > n - w' > t$, then the discrete-time quantum walk cannot reach x in t steps, thus $P(x, t) = 0$. Therefore there are two possibilities to consider:

1. $w \leq w'$, then Eq. (3.5) and Lemma 1 with $p_0 = C_1 \cdot 1.49578^{-n}$ give

$$P(x, t) = \frac{P[w, t]}{\binom{n}{w}} \leq p_0 \frac{n^w (n - w)!}{n!} < \frac{C_1}{1.49578^n} \cdot n^{nc} \cdot \frac{(n - nc)!}{n!} \quad (3.36)$$

with $c := w'/n = 0.13368$. Lower and upper bounds [20] on the factorials yield

$$n! \geq \left(\frac{n}{e}\right)^n e^{\frac{1}{12n+1}} \sqrt{2\pi n} \quad \text{and} \quad (n - nc)! \leq \left(\frac{n(1-c)}{e}\right)^{n(1-c)} e^{\frac{1}{12n(1-c)}} \sqrt{2\pi n(1-c)}, \quad (3.37)$$

leading to

$$\frac{(n - nc)!}{n!} \leq \left(\frac{n^{(1-c)}(1-c)^{(1-c)} \cdot e}{e^{(1-c)} \cdot n}\right)^n e^{\frac{1}{12n(1-c)} - \frac{1}{12n+1}} \sqrt{(1-c)}. \quad (3.38)$$

²See the remark after the proof about the numerical values chosen here.

Now, since $e^{\frac{1}{12n(1-c)} - \frac{1}{12n+1}} < e^{\frac{1}{12(1-c)}} < 2$ and $\sqrt{(1-c)} < 1$, we have

$$\frac{(n-nc)!}{n!} < 2n^{-nc}e^{nc}(1-c)^{(1-c)n} < 2 \cdot 0.99068^{-n}n^{-nc}, \quad (3.39)$$

because $e^c(1-c)^{(1-c)} = 0.990681\dots^{-1} < 0.99068^{-1}$. Let $C_2 = 2C_1$, then we conclude

$$P(x, t) < \frac{C_2}{(1.49578 \cdot 0.99068)^n} < \frac{C_2}{1.4818^n} = O(1.4818^{-n}). \quad (3.40)$$

2. $w' < w \leq n - w'$, then (3.5) and a lower bound [13, Lemma 9.2] on the binomial coefficient in terms of the binary entropy function H gives

$$P(x, t) \leq \binom{n}{w'}^{-1} \leq (n+1)2^{-nH(w'/n)} = (n+1)\left(2^{H(0.13368)}\right)^{-n}. \quad (3.41)$$

Since $2^{H(0.13368)} = 1.48189\dots$, we arrive at $P(x, t) = (n+1) \cdot 1.48189\dots^{-n} = O(1.4818^{-n})$. \square

Remark. The constants appearing in the above theorem can be explained as follows: let $w' = cn$, $t = (1-c)n$ and $\alpha = (1-2c)n$. That corresponds to the case $[t-w', t+w'] = [\alpha n, n]$, and we need to balance the estimate (when $w \leq w'$)

$$\left(\frac{e(1-c)^{1-c}}{(2-2c)^{(1-2c)}}\right)^n = \left(\frac{e(1-c)^c}{2^{(1-2c)}}\right)^n \quad (3.42)$$

with the estimate (when $w' < w < n - w'$)

$$2^{-H(c)n} = \left(c^c(1-c)^{(1-c)}\right)^n, \quad (3.43)$$

leading to the equilibrium value $c = 0.133682\dots$. However, this reasoning ignores the fact that $[t-w', t+w']$ must be a proper subset of $(\alpha n, n)$, as well as the polynomial factor $(n+1)$ in the second estimate of $P(x, t)$. Therefore we choose slightly smaller values of α and t , as well as upper bound the true value of $2^{-H(c)}$, leading to a slightly worse overall estimate.

4 Acknowledgements

We thank Ashley Montanaro for suggesting the problem and the connection to [8] and Raqueline Santos for participating in the early work on this subject. This work has been supported by Latvian Council of Science (project no. lzp-2018/1-0173) and the ERDF project 1.1.1.5/18/A/020.

References

- [1] Gorjan Alagic and Alexander Russell. Decoherence in quantum walks on the hypercube. *Physical Review A*, 72(6):062304, 2005. DOI: [10.1103/PhysRevA.72.062304](https://doi.org/10.1103/PhysRevA.72.062304).
- [2] Andris Ambainis. Quantum walk algorithm for element distinctness. *SIAM J. Comput.*, 37(1):210–239, 2007. DOI: [10.1137/S0097539705447311](https://doi.org/10.1137/S0097539705447311).
- [3] Andris Ambainis, Krišjānis Prūsis, Jevgēnijs Vihrovs, and Thomas G. Wong. Oscillatory localization of quantum walks analyzed by classical electric circuits. *Physical Review A*, 94(6):062324, 2016. DOI: [10.1103/PhysRevA.94.062324](https://doi.org/10.1103/PhysRevA.94.062324).
- [4] Andris Ambainis, András Gilyén, Stacey Jeffery, and Martins Kokainis. Quadratic speedup for finding marked vertices by quantum walks. In Konstantin Makarychev, Yury Makarychev, Madhur Tulsiani, Gautam Kamath, and Julia Chuzhoy, editors, *Proceedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing, STOC 2020, Chicago, IL, USA, June 22–26, 2020*, pages 412–424. ACM, 2020. DOI: [10.1145/3357713.3384252](https://doi.org/10.1145/3357713.3384252).

- [5] Simon Apers and Alain Sarlette. Quantum fast-forwarding: Markov chains and graph property testing. *Quantum Inf. Comput.*, 19(3&4):181–213, 2019. DOI: [10.26421/QIC19.3-4-1](https://doi.org/10.26421/QIC19.3-4-1).
- [6] Simon Apers, András Gilyén, and Stacey Jeffery. A unified framework of quantum walk search. In Markus Bläser and Benjamin Monmege, editors, *38th International Symposium on Theoretical Aspects of Computer Science, STACS 2021, March 16-19, 2021, Saarbrücken, Germany (Virtual Conference)*, volume 187 of *LIPICs*, pages 6:1–6:13. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021. DOI: [10.4230/LIPICs.STACS.2021.6](https://doi.org/10.4230/LIPICs.STACS.2021.6).
- [7] Gang Bao and Haijun Wu. Convergence analysis of the perfectly matched layer problems for time-harmonic maxwell’s equations. *SIAM Journal on Numerical Analysis*, 43(5):2121–2143, 2006. ISSN 00361429. DOI: [10.1137/040604315](https://doi.org/10.1137/040604315).
- [8] Fernando G. S. L. Brandão and Michal Horodecki. Exponential quantum speed-ups are generic. *Quantum Information & Computation*, 13(11-12):901–924, 2013. DOI: [10.26421/QIC13.11-12-1](https://doi.org/10.26421/QIC13.11-12-1).
- [9] Andrew M. Childs, Richard Cleve, Enrico Deotto, Edward Farhi, Sam Gutmann, and Daniel A. Spielman. Exponential algorithmic speedup by a quantum walk. In Lawrence L. Larmore and Michel X. Goemans, editors, *Proceedings of the 35th Annual ACM Symposium on Theory of Computing, June 9-11, 2003, San Diego, CA, USA*, pages 59–68. ACM, 2003. DOI: [10.1145/780542.780552](https://doi.org/10.1145/780542.780552).
- [10] DLMF. *NIST Digital Library of Mathematical Functions*. <http://dlmf.nist.gov/>, Release 1.1.4 of 2022-01-15. URL <http://dlmf.nist.gov/>. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [11] Hari Krovi and Todd A. Brun. Hitting time for quantum walks on the hypercube. *Physical Review A*, 73(3):032341, 2006. DOI: [10.1103/PhysRevA.73.032341](https://doi.org/10.1103/PhysRevA.73.032341).
- [12] Franklin L Marquezino, Renato Portugal, Gonzalo Abal, and Raul Donangelo. Mixing times in quantum walks on the hypercube. *Physical Review A*, 77(4):042312, 2008. DOI: [10.1103/PhysRevA.77.042312](https://doi.org/10.1103/PhysRevA.77.042312).
- [13] Michael Mitzenmacher and Eli Upfal. *Probability and Computing: Randomized Algorithms and Probabilistic Analysis*. Cambridge University Press, 2005. DOI: [10.1017/CBO9780511813603](https://doi.org/10.1017/CBO9780511813603).
- [14] Masoud Mohseni, Patrick Rebentrost, Seth Lloyd, and Alan Aspuru-Guzik. Environment-assisted quantum walks in photosynthetic energy transfer. *The Journal of Chemical Physics*, 129(17):11B603, 2008. DOI: [10.1063/1.3002335](https://doi.org/10.1063/1.3002335).
- [15] Cristopher Moore and Alexander Russell. Quantum walks on the hypercube. In *International Workshop on Randomization and Approximation Techniques in Computer Science*, pages 164–178. Springer, 2002. DOI: [10.1007/3-540-45726-7_14](https://doi.org/10.1007/3-540-45726-7_14).
- [16] Daniel Nagaj, Pawel Wocjan, and Yong Zhang. Fast amplification of QMA. *Quantum Information & Computation*, 9(11):1053–1068, 2009. DOI: [10.26421/QIC9.11-12-8](https://doi.org/10.26421/QIC9.11-12-8).
- [17] Frank W. J. Olver. *Asymptotics and special functions*. A K Peters/CRC Press, 1997. ISBN 1-56881-069-5. DOI: [10.1201/9781439864548](https://doi.org/10.1201/9781439864548).
- [18] Richard B. Paris. An Inequality for the Bessel Function $J_\nu(\nu x)$. *SIAM Journal on Mathematical Analysis*, 15(1):203–205, 1984. DOI: [10.1137/0515016](https://doi.org/10.1137/0515016).
- [19] Václav Potoček, Aurél Gábris, Tamás Kiss, and Igor Jex. Optimized quantum random-walk search algorithms on the hypercube. *Physical Review A*, 79(1):012325, 2009. DOI: [10.1103/PhysRevA.79.012325](https://doi.org/10.1103/PhysRevA.79.012325).
- [20] Herbert Robbins. A remark on Stirling’s formula. *The American Mathematical Monthly*, 62(1):26–29, 1955. ISSN 00029890, 19300972. DOI: [10.2307/2308012](https://doi.org/10.2307/2308012).
- [21] Alberto G. Setti. L^p and operator norm estimates for the complex time heat operator on homogeneous trees. *Transactions of the American Mathematical Society*, 350(2):743–768, 1998. ISSN 00029947. DOI: [10.1090/S0002-9947-98-02042-X](https://doi.org/10.1090/S0002-9947-98-02042-X).
- [22] Neil Shenvi, Julia Kempe, and K. Birgitta Whaley. Quantum random-walk search algorithm. *Physical Review A*, 67(5):052307, 2003. DOI: [10.1103/PhysRevA.67.052307](https://doi.org/10.1103/PhysRevA.67.052307).
- [23] Mario Szegedy. Quantum speed-up of Markov Chain based algorithms. In *45th Symposium on Foundations of Computer Science (FOCS 2004), 17-19 October 2004, Rome, Italy, Proceedings*, pages 32–41. IEEE Computer Society, 2004. DOI: [10.1109/FOCS.2004.53](https://doi.org/10.1109/FOCS.2004.53).
- [24] George N. Watson. *A treatise on the theory of Bessel functions*. Cambridge University Press, 1922.

A Bounding the integral

A.1 Preliminaries

Throughout the proof, we will make use of standard notation of some special functions; Γ will stand for the gamma function; B denotes the beta function, satisfying $B(x, y) = \Gamma(x)\Gamma(y)/\Gamma(x, y)$. The Bessel functions of the first and second kind will be denoted as $J_\nu(x)$ and $Y_\nu(x)$, respectively; and the Hankel function of the first kind is denoted as $H_\nu^{(1)}(x) = J_\nu(x) + iY_\nu(x)$. As in the statement of the theorem, we denote $a_k = (k - 0.5)\pi$, $k = 1, 2, 3, \dots$, and let a positive constant $\alpha \in (\frac{\pi}{6}, 1)$ be fixed. Furthermore, let $\nu > 0$ and suppose an integer $n \geq 2$ satisfies

$$n \geq \frac{1}{\alpha} \quad \text{and} \quad \alpha < \frac{\nu}{n} < 1.$$

Also, for $a, b \in \mathbb{R}$, with $a < b$, we define $D_{a,b} = \{x + iy \mid x \in [a, b], 0 \leq y < \infty\}$ (i.e., $D_{a,b}$ stands for an ‘‘infinite rectangle’’ in the complex plane with one side being the real interval $[a; b]$).

A.2 Asymptotic expansions of the Hankel function

The case of large argument. When $z \in \mathbb{C}$, $|\arg z| \in [0, \pi - \delta)$ for an arbitrary small positive constant δ , we have [10, §10.17(iv)]

$$H_\nu^{(1)}(z) = \sqrt{\frac{2}{\pi z}} e^{i(z - \nu\pi/2 - \pi/4)} (1 + \rho(\nu, z)), \quad (\text{A.1})$$

where the principal branch of \sqrt{z} is used, and

$$|\rho(\nu, z)| \leq (\nu^2 - 1/4) |z|^{-1} \exp\left((\nu^2 - 1/4) |z|^{-1}\right). \quad (\text{A.2})$$

For future reference, we note that whenever z additionally satisfies $|z| \geq n^2$, we have (since $\nu < n$)

$$|\rho(\nu, z)| < \nu^2 |z|^{-1} \exp\left(\nu^2 |z|^{-1}\right) < e \quad (\text{A.3})$$

and

$$\left|H_\nu^{(1)}(z)\right| \leq \sqrt{\frac{2}{\pi}} (1 + e) |e^{iz}| |z|^{-1/2} < \frac{3e^{-\Im(z)}}{|z|^{1/2}}. \quad (\text{A.4})$$

The case when argument and order are of similar magnitude.

Lemma 4. *When $\arg w \in [0; \pi/2)$ and $\Re w > 1$, we have*

$$H_\nu^{(1)}(\nu w) = -i \sqrt{\frac{2}{\pi \nu}} \frac{e^{i\nu(\sqrt{w^2 - 1} - \arccos(1/w))}}{(1 - w^2)^{0.25}} (1 + \eta(\nu, -iw)), \quad (\text{A.5})$$

where the fractional powers and the logarithm take their principal values on the positive real axis and the term η can be bounded as

$$|\eta(\nu, -iw)| \leq \exp\left(\frac{2\mathcal{V}_{+\infty, -iw}}{\nu}\right) \frac{2\mathcal{V}_{+\infty, -iw}}{\nu},$$

and the quantity $\mathcal{V}_{+\infty, -iw}$ satisfies

$$\mathcal{V}_{+\infty, -iw} \leq \frac{1}{12} + \frac{1}{6\sqrt{5}} + \left(\frac{4}{27}\right)^{1/4} + \frac{c^2(c^2 + 2)}{\sqrt{8}(c^2 - 1)^{2.5}}, \quad c := \Re w > 1, \quad \Im w \geq 0.$$

Moreover, when $\nu \geq 1$ and $w \in \mathbb{C}$ satisfies $\Re(w) \geq \pi/2$ and $\Im(w) \geq 0$, we have

$$|1 + \eta(\nu, -iw)| \leq 430. \quad (\text{A.6})$$

The proof is postponed until Appendix A.8.

A.3 Auxiliary lemmata

Here we list a few somewhat disjoint auxiliary results that will be useful in the subsequent analysis. Proofs of those claims are postponed until Appendix A.6.

Lemma 5. *Suppose that $f : D \rightarrow \mathbb{C}$ is holomorphic in D , where D is a domain containing the region $D_{a,b} := \{x + iy \mid x \in [a, b], 0 \leq y < \infty\}$, for some reals $a < b$. Moreover, assume that*

1. $\lim_{R \rightarrow +\infty} \sup_{x \in [a,b]} |f(x + iR)| = 0$, and
2. integrals $\int_0^\infty f(a + iy) dy$, $\int_0^\infty f(b + iy) dy$ converge.

Then

$$\int_a^b f(x) dx = i \int_0^\infty (f(a + iy) - f(b + iy)) dy. \quad (\text{A.7})$$

Lemma 6. *For all $a > 0$ the following equalities hold:*

$$\int_0^\infty \frac{dy}{(a^2 + y^2)^{3/4}} = \frac{B(0.5, 0.25)}{2\sqrt{a}} \approx 2.62206 \dots \cdot a^{-0.5}, \quad (\text{A.8})$$

$$\int_0^\infty \frac{dy}{(a^2 + y^2)^{5/4}} = \frac{B(0.5, 0.75)}{2\sqrt{a^3}} \approx 1.1984 \dots \cdot a^{-1.5}. \quad (\text{A.9})$$

Lemma 7. *For all $t \in [0; \pi/2)$ the inequality $\cos t \leq e^{-t^2/2}$ holds.*

Lemma 8. *Let g be defined in the set $\{z \in \mathbb{C} \mid \Re z \geq 1, \Im z \geq 0\}$ via*

$$g(z) = z - \sqrt{z^2 - 1} + \arccos(1/z),$$

where the square root and the inverse cosine functions take their principal values on the positive real axis. Then $\Im(g(z)) < 0.2607$ for all z with $\Re z \geq 1$, $\Im z \geq 0$.

Remark. See Footnote 6 (on page 24) for an interpretation of Lemma 8 in the context of the modified Bessel function of the second kind.

A.4 Properties of the main function

It can be noticed that the integrand $x^{-1} J_\nu(x) \cos^n\left(\frac{x}{n}\right)$ is the real part of the function f defined by (3.31). Since $H_\nu^{(1)}$ is holomorphic throughout the complex plane cut along the negative real axis, f is holomorphic in the domain $\{z \in \mathbb{C} \mid \Re z \geq \epsilon\}$, for an arbitrary $\epsilon > 0$.

We intend to estimate the integral of f with the help of Lemma 5 by dividing the integration domain into subintervals with endpoints $0, na_1, na_2, \dots$; to that end, we will make use of the following properties of f . The proofs of the following claims are postponed until Appendix A.7.

First we characterize and bound the values of f :

Lemma 9. *If $x, y \geq 0$ are nonnegative reals such that f is defined at $x + iy$, then*

$$f(x + iy) = \left(\cos(x/n) \cosh(y/n) - i \sin(x/n) \sinh(y/n) \right)^n \frac{H_\nu^{(1)}(x + iy)}{x + iy}. \quad (\text{A.10})$$

Furthermore, if $\max\{x, y\} \geq n^2$, then the following inequality is satisfied

$$|f(x + iy)| \leq 3 |x + iy|^{-3/2}. \quad (\text{A.11})$$

We also characterize the values of f when its argument is of the form $na_k + iy$:

Lemma 10. *For all $y \geq 0$ and integer $k \geq 1$ it holds that*

$$f(na_k + iy) = 2^{-n} \frac{e^{ina_{k+1}} \cdot (e^{y/n} - e^{-y/n})^n H_\nu^{(1)}(na_k + iy)}{na_k + iy}, \quad (\text{A.12})$$

and, whenever $|na_k + iy| \geq n^2$,

$$|f(na_k + iy)| < 3 \cdot 2^{-n} |na_k + iy|^{-3/2}. \quad (\text{A.13})$$

Furthermore,

1. if $1 \leq k < n$, then for all $y \geq 0$ the following estimate holds:

$$|f(na_k + iy)| < 860 \cdot 1.541^{-n} |na_k + iy|^{-1.5}. \quad (\text{A.14})$$

2. if $2 \leq n \leq k$, then for all $y \geq 0$ the following estimate holds:

$$|f(na_k + iy) - f(na_{k+1} + iy)| < 15n^2 \cdot 2^{-n} |na_k + iy|^{-2.5} \quad (\text{A.15})$$

A.5 Proof of Theorem 2

Now we can prove Theorem 2, restated here for convenience:

Theorem 2. Denote $a_k = (k - 0.5)\pi$, $k = 1, 2, 3, \dots$; let a positive constant $\alpha \in (\frac{\pi}{6}, 1)$ be fixed.

For $n \geq 2$ and $\nu > 1$, satisfying $\nu \in (n\alpha, n) \subset (\frac{\pi n}{6}, n)$, the following estimates hold:

$$\text{The tail:} \quad \left| \int_{na_n}^{\infty} x^{-1} J_{\nu}(x) \cos^n \left(\frac{x}{n} \right) dx \right| < \frac{100\sqrt{n}}{2^n}; \quad (3.28)$$

$$\text{The middle part:} \quad \left| \int_{na_1}^{na_n} x^{-1} J_{\nu}(x) \cos^n \left(\frac{x}{n} \right) dx \right| < \frac{4000\sqrt{n}}{1.541^n}; \quad (3.29)$$

$$\text{The bulk:} \quad \left| \int_0^{na_1} x^{-1} J_{\nu}(x) \cos^n \left(\frac{x}{n} \right) dx \right| < \frac{3}{(1 + \alpha)^{0.5\alpha n}}. \quad (3.30)$$

Proof of Theorem 2. Denote

$$I_0(n, \nu) = \int_0^{na_1} x^{-1} J_{\nu}(x) \cos^n \left(\frac{x}{n} \right) dx;$$

$$I_k(n, \nu) = \int_{na_k}^{na_{k+1}} x^{-1} J_{\nu}(x) \cos^n \left(\frac{x}{n} \right) dx, \quad k \in \mathbb{N}.$$

The tail integral (proof of (3.28)). Let $k \geq n \geq 2$ and notice that $I_k(n, \nu)$ is the real part of the integral $\int_{na_k}^{na_{k+1}} f(x) dx$. The function f is holomorphic in a domain containing $D_{na_k, na_{k+1}}$; let us verify the other assumptions of Lemma 5.

Since $na_k \geq n(n - 0.5)\pi > n^2$, Eq. (A.11) in Lemma 9 implies

$$\sup_{x \in [na_k, na_{k+1}]} |f(x + iR)| \leq \frac{3}{|na_k + iR|^{1.5}} \xrightarrow{R \rightarrow +\infty} 0.$$

From (A.13) it follows that $|f(na_k + iy)| \leq 3 \cdot 2^{-n} |na_k + iy|^{-3/2}$, thus the integral $\int_0^{\infty} f(na_k + iy) dy$ converges and its absolute value, by Lemma 6, Eq. (A.8), is bounded by

$$\int_0^{\infty} |f(na_k + iy)| dy \leq 3 \cdot 2^{-n} \int_0^{\infty} \frac{dy}{(n^2 a_k^2 + y^2)^{3/4}} < \frac{9 \cdot 2^{-n}}{\sqrt{na_k}};$$

similarly, $\int_0^{\infty} f(na_{k+1} + iy) dy$ converges. Using Lemma 5 we bound

$$|I_k(n, \nu)| \leq \left| \int_{na_k}^{na_{k+1}} f(x) dx \right| \leq \int_0^{\infty} |f(na_k + iy) - f(na_{k+1} + iy)| dy. \quad (\text{A.16})$$

Now apply Lemma 10, Eq. (A.15) to upper-bound the difference of f values as

$$|f(na_k + iy) - f(na_{k+1} + iy)| \leq \frac{15n^2}{2^n} |na_k + iy|^{-5/2}. \quad (\text{A.17})$$

By Lemma 6, Eq. (A.9) we have

$$\int_0^\infty \frac{dy}{(n^2 a_k^2 + y^2)^{5/4}} < \frac{2}{\sqrt{n^3 a_k^3}} < \frac{30\sqrt{n}}{2^n k^{3/2}}, \quad (\text{A.18})$$

where the last inequality uses $k \leq a_k$.

Now, by combining (A.16)-(A.18), we arrive at

$$\left| \int_{na_n}^\infty x^{-1} J_\nu(x) \cos^n\left(\frac{x}{n}\right) dx \right| \leq \sum_{k=n}^\infty |I_k(n, \nu)| < \frac{30\sqrt{n}}{2^n} \sum_{k=n}^\infty k^{-3/2}.$$

Since $30 \sum_{k=n}^\infty k^{-3/2} < \sum_{k=1}^\infty k^{-3/2} \approx 30 \cdot 2.6124 \dots < 100$, we are done.

The middle part (proof of (3.29)). Let k satisfy $1 \leq k \leq n-1$ and notice that $I_k(n, \nu)$ is the real part of the integral $\int_{na_k}^{na_{k+1}} f(x) dx$. The function f is holomorphic in a domain containing $D_{na_k, na_{k+1}}$; let us verify the other assumptions of Lemma 5.

When $R > n^2$, Lemma 9 implies

$$\sup_{x \in [na_k, na_{k+1}]} |f(x + iR)| \leq \frac{3}{|na_k + iR|^{1.5}} \xrightarrow{R \rightarrow +\infty} 0.$$

Split

$$\int_0^\infty f(na_k + iy) dy = \int_0^{n^2} f(na_k + iy) dy + \int_{n^2}^\infty f(na_k + iy) dy,$$

then from (A.13) in Lemma 10 we conclude that $\int_{n^2}^\infty f(na_k + iy) dy$ converges and its absolute value is bounded by

$$3 \cdot 2^{-n} \int_0^\infty \frac{dy}{(n^2 a_k^2 + y^2)^{3/4}} < 9 \cdot 2^{-n} (na_k)^{-0.5}.$$

Thus $\int_0^\infty f(na_k + iy) dy$ converges as well (and so does $\int_0^\infty f(na_{k+1} + iy) dy$) and Lemma 5 applies.

Moreover, we also conclude

$$\left| \int_0^\infty f(na_k + iy) dy \right| < \left| \int_0^{n^2} f(na_k + iy) dy \right| + \frac{8}{2^n \sqrt{n}}. \quad (\text{A.19})$$

Here we have used $a_k \geq a_1 = \pi/2$ and

$$\left| \int_{n^2}^\infty f(na_k + iy) dy \right| < \frac{9}{2^n \sqrt{na_1}} = \frac{9\sqrt{2}/\sqrt{\pi}}{2^n \sqrt{n}} < \frac{8}{2^n \sqrt{n}}.$$

Due to Lemma 10, Eq. (A.14), we can estimate the remaining integral as

$$\left| \int_0^{n^2} f(na_k + iy) dy \right| < 860 \cdot 1.541^{-n} \int_0^\infty |na_k + iy|^{-1.5} dy < \frac{1800}{1.541^n \sqrt{n}}.$$

The last step applies (A.8) and inequalities

$$860 \cdot \frac{B(0.5, 0.25)}{2\sqrt{a_k}} \leq 860 \cdot \frac{B(0.5, 0.25)}{\sqrt{2\pi}} < 1800.$$

Now (A.19) gives

$$\left| \int_0^\infty f(na_k + iy) dy \right| < \frac{1800}{1.541^n \sqrt{n}} + \frac{8}{2^n \sqrt{n}} < \frac{2000}{1.541^n \sqrt{n}},$$

thus

$$|I_k(n, \nu)| \leq \left| \int_0^\infty f(na_k + iy) dy \right| + \left| \int_0^\infty f(na_{k+1} + iy) dy \right| < \frac{4000}{1.541^n \sqrt{n}}$$

and (3.29) is obtained as

$$\left| \int_{na_1}^{na_n} x^{-1} J_\nu(x) \cos^n\left(\frac{x}{n}\right) dx \right| \leq \sum_{k=1}^{n-1} |I_k(n, \nu)| < \frac{4000\sqrt{n}}{1.541^n}.$$

The bulk (proof of (3.30)). Denote $c = \nu/n \in (\alpha, 1)$. We start by splitting

$$I_0(n, \nu) = \int_0^\nu x^{-1} J_\nu(x) \cos^n\left(\frac{x}{n}\right) dx + \int_{nc}^{n\pi/2} x^{-1} J_\nu(x) \cos^n\left(\frac{x}{n}\right) dx. \quad (\text{A.20})$$

The second integral can be bounded by employing the fact that $\cos(x/n)^n/x$ is decreasing in $[nc, n\pi/2]$ and noting that [10, Eq. 10.14.1] the absolute value of the Bessel function J_ν is bounded by 1 for all real arguments, whence

$$\int_{nc}^{n\pi/2} x^{-1} J_\nu(x) \cos^n\left(\frac{x}{n}\right) dx \leq \frac{\cos^n(c)}{nc} \cdot (n\pi/2 - nc) < 2 \cos^n(\alpha). \quad (\text{A.21})$$

The last step relies on the assumption $\alpha > \pi/6$.

For the first integral in the RHS of (A.20), we apply Lemma 7, which gives $\cos^n\left(\frac{x}{n}\right) \leq \exp\left(-\frac{x^2}{2n}\right)$, as well as the estimate

$$|J_\nu(\nu t)| \leq J_\nu(\nu) \cdot t^\nu \exp\left(\frac{\nu^2(1-t^2)}{2\nu+4}\right),$$

valid [18, p.204] for all $t \in (0, 1]$ and $\nu > 0$. This inequality can be slightly weakened by bounding $\frac{\nu^2(1-t^2)}{2\nu+4} < \frac{\nu(1-t^2)}{2}$. Those imply that the integral can be bounded as

$$\begin{aligned} \left| \int_0^\nu x^{-1} J_\nu(x) \cos^n\left(\frac{x}{n}\right) dx \right| &\leq J_\nu(\nu) \nu^{-\nu} \cdot \int_0^\nu x^{\nu-1} \exp\left(\frac{\nu}{2} - \frac{x^2}{2\nu} - \frac{x^2}{2n}\right) dx \\ &= J_\nu(\nu) \left(\frac{e}{\nu^2}\right)^{\nu/2} \cdot \int_0^\nu x^{\nu-1} \exp\left(-\frac{(c+1)x^2}{2\nu}\right) dx \\ &= J_\nu(\nu) \left(\frac{e}{\nu^2}\right)^{\nu/2} \cdot \int_0^{(c+1)\nu/2} \left(\frac{2\nu y}{c+1}\right)^{\nu/2-1} \cdot \frac{\nu}{c+1} \exp(-y) dy \\ &= \frac{1}{2} J_\nu(\nu) \left(\frac{2e}{\nu(c+1)}\right)^{\nu/2} \cdot \int_0^{(c+1)\nu/2} y^{\nu/2-1} \exp(-y) dy \\ &< 0.5 J_\nu(\nu) \left(\frac{2e}{\nu(c+1)}\right)^{\nu/2} \cdot \underbrace{\int_0^\infty y^{\nu/2-1} \exp(-y) dy}_{\Gamma(\nu/2)}. \end{aligned}$$

Now we apply a bound on the gamma function, valid [10, Eq. 5.6.1] for positive arguments:

$$\Gamma(x) > \sqrt{\frac{2\pi}{x}} x^x \exp\left(-x + \frac{1}{12x}\right), \quad x > 0.$$

Taking $x = \nu/2$ this implies that

$$\Gamma(\nu/2) > 2\sqrt{\frac{\pi}{\nu}} \left(\frac{\nu}{2}\right)^{\nu/2} \exp\left(-\frac{\nu}{2} + \frac{1}{6\nu}\right)$$

and, since $\nu > 1$,

$$\left| \int_0^\nu x^{-1} J_\nu(x) \cos^n\left(\frac{x}{n}\right) dx \right| < J_\nu(\nu) \sqrt{\frac{\pi}{\nu}} e^{1/6} (c+1)^{-\nu/2}.$$

By [10, Eq. 10.14.2],

$$0 < J_\nu(\nu) < \frac{2^{1/3}}{3^{2/3}\Gamma(2/3)\nu^{1/3}} < \frac{0.45}{\nu^{1/3}}.$$

From $0.45\sqrt{\pi}e^{1/6} < 1$ and $\nu > 1$ we arrive at

$$\left| \int_0^\nu x^{-1} J_\nu(x) \cos^n\left(\frac{x}{n}\right) dx \right| < \frac{0.45\sqrt{\pi}e^{1/6}}{\nu^{5/6}} \exp\left(-\frac{nc \ln(c+1)}{2}\right) < (1+c)^{-cn/2}. \quad (\text{A.22})$$

We note that $c \mapsto (1+c)^{c/2}$ is increasing in c , therefore the bound on the RHS of (A.22) is decreasing in c and attains its maximal value at α .

Finally, an application of Lemma 7 gives $\cos^n(\alpha) < \exp(-0.5n\alpha^2)$; since $-\alpha^2 < -\alpha \ln(\alpha+1)$, we have

$$2 \cos^n(\alpha) < 2(1+\alpha)^{-\alpha n/2},$$

which yields the desired inequality (3.30). \square

A.6 Proofs of the auxiliary lemmata

Proof of Lemma 5. For every $R > 0$ consider the positively oriented rectifiable curve γ_R consisting of the lines

$$[a, b] \cup \{b + iy \mid y \in [0, R]\} \cup \{x + iR \mid x \in [a, b]\} \cup \{a + iy \mid y \in [0, R]\}.$$

By Cauchy's integral theorem we have $\oint_{\gamma_R} f(z) dz = 0$, i.e.,

$$\int_a^b f(x) dx + i \int_0^R f(b + iy) dy - \int_a^b f(x + iR) dx - i \int_0^R f(a + iy) dy = 0.$$

Rearrange this equality as

$$\int_a^b f(x) dx = i \left(\int_0^R f(a + iy) dy - \int_0^R f(b + iy) dy \right) + \int_a^b f(x + iR) dx$$

and take the limit $R \rightarrow \infty$. Since

$$\left| \int_a^b f(x + iR) dx \right| \leq (b-a) \sup_{x \in [a, b]} |f(x + iR)| \xrightarrow{R \rightarrow +\infty} 0,$$

we are done. \square

Proof of Lemma 6. The first equality can be proven as follows: substitute $t = a^2 y^2$, which gives $dy = 0.5at^{-1/2} dt$. Then

$$\int_0^\infty \frac{dy}{(a^2 + y^2)^{3/4}} = \frac{a}{2} \cdot \frac{1}{a^{3/2}} \int_0^\infty \frac{t^{-1/2} dt}{(1+t)^{3/4}} = \frac{1}{2a^{1/2}} \int_0^\infty \frac{t^{1/2-1} dt}{(1+t)^{1/2+1/4}}.$$

Now from the identity $B(x, y) = \int_0^\infty \frac{t^{x-1} dt}{(1+t)^{x+y}}$ we recognize the integral on the RHS as the beta function value $B(0.5, 0.25) = \Gamma(0.5)\Gamma(0.25)/\Gamma(1.25) \approx 5.24412\dots$. In a similar manner one shows the second equality. \square

Proof of Lemma 7. Let $h(t) = \ln(\cos(t)) + t^2/2$ and consider $h'(t) = t - \tan(t)$. Since $h'(0) = 0$ and $h'(t) < 0$ for $t \in (0; \pi/2)$, h attains its maximum at $t = 0$, i.e., $\ln(\cos t) \leq -t^2/2$ with equality at $t = 0$. Exponentiating gives the desired result. \square

Proof of Lemma 8. Let $R > 0$; we will assume that R is large enough, e.g., $R > 1$.

Since g is holomorphic on the domain $\{x + iy \mid x \in (1, R), y \in (0, R)\}$ and continuous on its closure, its imaginary part $\Im(g(z))$ is harmonic and attains its maximum on the boundary of this region. Moreover, as $g(z) \in \mathbb{R}$ when $\Im z = 0$ and

$$\left| z - \sqrt{z^2 - 1} \right| \xrightarrow{|z| \rightarrow \infty} 0 \quad \text{and} \quad \arccos(1/z) \xrightarrow{|z| \rightarrow \infty} \pi/2,$$

it follows that $\Im(g(z))$ must attain its maximum on the segment with $\Re z = 1$, $\Im z \in [0, R]$.

Let us separate the real and imaginary part of g , i.e., introduce real-valued bivariate functions u, v satisfying $g(x + iy) = u(x, y) + iv(x, y)$. By the arguments above, we need to show that

$v(1, y) < 0.2607$ for all $y \in [0, R]$, for arbitrarily large R . We will show that (for $R > 1$) the function $v(1, \cdot)$ has a single local maximum at $y_0 \approx 0.86883\dots$ where its value is less than 0.2607. To calculate the derivative of $y \mapsto v(1, y)$ and show that it is positive on $(0, y_0)$ and negative on (y_0, R) , we employ Cauchy-Riemann equations.

Consider the derivative of $g(z)$, denoted by $g^{(1)}(z)$:

$$g^{(1)}(z) := \frac{d}{dz}g(z) = 1 - \frac{z}{\sqrt{z^2-1}} + \frac{1}{z^2\sqrt{1-z^{-2}}} = 1 - \frac{\sqrt{z^2-1}}{z}.$$

Separate the real and imaginary parts of the derivative, i.e., $g^{(1)}(x+iy) = u^{(1)}(x, y) + iv^{(1)}(x, y)$, $x, y \in \mathbb{R}$, then Cauchy-Riemann equations imply

$$\begin{cases} u^{(1)}(x_0, y_0) = \frac{\partial u}{\partial x}(x_0, y_0) = \frac{\partial v}{\partial y}(x_0, y_0), \\ v^{(1)}(x_0, y_0) = -\frac{\partial u}{\partial y}(x_0, y_0) = \frac{\partial v}{\partial x}(x_0, y_0), \end{cases}$$

for any $(x_0, y_0) \in (1, R) \times (0, R)$. Moreover, these equations remain also true for $x_0 = 1, y_0 > 0$ (this follows from the fact that g is holomorphic in a neighborhood of any $z = 1 + iy_0$ with $y_0 > 0$). Since we are interested in $\frac{\partial v}{\partial y}(1, y)$ for $y \in (0, \infty)$, consider the real part of $g^{(1)}(z)$, where $z = 1 + iy$. A direct calculation gives

$$\frac{\partial v}{\partial y}(1, y) = u^{(1)}(1, y) = 1 - \frac{\sqrt{\frac{y^2-1}{(y^2+1)^2} + \frac{\sqrt{y^4+4y^2}}{y^2+1} + 1}}{\sqrt{2}}.$$

It is easy to verify that this expression is positive for $y \in (0, y_0)$ and negative for $y \in (y_0, \infty)$, where $y_0 > 0$ is the real solution of

$$\frac{y\sqrt{y^2+4}+1}{y^2+1} - \frac{2}{(y^2+1)^2} = 1.$$

Letting $t = y^2$ and simplifying gives the equation $t^3 + t^2 = 1$, whose only real solution is

$$t_0 = \frac{\left(\frac{25}{2} - \frac{3\sqrt{69}}{2}\right)^{1/3} + \left(\frac{3\sqrt{69}}{2} + \frac{25}{2}\right)^{1/3} - 1}{3} \approx 0.7548\dots,$$

thus $y_0 = \sqrt{t_0} \approx 0.86883\dots$ and $\Im(g(z))$ attains its maximum value at $z = 1 + iy_0$. Finally, numerical calculations yield $g(1 + iy_0) = 1.29797\dots + i0.26066\dots$, thus $\Im(g(z)) < 0.2607$ for the values of z under consideration. \square

A.7 Proving properties of the main function

Proof of Lemma 9. The equality in (A.10) follows from the definition of f and the identity

$$\cos(x/n + iy/n) = \cos(x/n) \cosh(y/n) - i \sin(x/n) \sinh(y/n).$$

To show (A.11), apply (A.4) to bound $H_\nu^{(1)}(x+iy)$ in (A.10) (notice that $\arg(x+iy) \in [0, \pi/2)$ for the x, y values under the consideration; moreover, $|x+iy| \geq \max\{x, y\} \geq n^2$), obtaining

$$\begin{aligned} |f(x+iy)| &= \left| \cos(x/n) \cosh(y/n) - i \sin(x/n) \sinh(y/n) \right|^n \cdot \left| \frac{3e^{-y}}{(x+iy)^{1.5}} \right| \\ &\leq \frac{3e^{-y} \left(|\cos(x/n) \cosh(y/n)| + |\sin(x/n) \sinh(y/n)| \right)^n}{|x+iy|^{1.5}} \\ &\leq 3e^{-y} \left(\cosh(y/n) + \sinh(y/n) \right)^n |x+iy|^{-1.5} = 3|x+iy|^{-1.5}. \end{aligned}$$

\square

Proof of Lemma 10. Set $x = na_k$ in (A.10); since $\cos(a_k) = 0$, $(-i \sin(a_k))^n = e^{-ina_k} = e^{ina_{k+1}}$ and

$$\sinh(y/n) = 0.5 \left(e^{y/n} - e^{-y/n} \right),$$

we obtain (A.12). Now, if $|na_k + iy| \geq n^2$, we can apply (A.4) to bound $H_\nu^{(1)}(x + iy)$ in (A.12), obtaining the desired estimate (A.13):

$$|f(na_k + iy)| \leq 2^{-n} \left(e^{y/n} - e^{-y/n} \right)^n \frac{3e^{-y}}{|na_k + iy|^{1.5}} < 3 \cdot 2^{-n} |na_k + iy|^{-1.5}.$$

Proof of (A.14). Suppose $1 \leq k \leq n-1$. Let $w = na_k/\nu + iy/\nu$ and apply Lemma 4 to rewrite (A.12) as

$$\begin{aligned} |f(na_k + iy)| &= \frac{|e^{y/n} - e^{-y/n}|^n |H_\nu^{(1)}(\nu w)|}{2^n |na_k + iy|} \\ &\leq \frac{e^y |e^{i\nu(\sqrt{w^2-1} - \arccos(1/w))}|}{2^n \sqrt{\nu\pi/2} |na_k + iy| |1 - w^2|^{0.25}} \cdot |1 + \eta(\nu, -iw)|. \\ &= \frac{e^{y - \nu \Im(\sqrt{w^2-1} - \arccos(1/w))}}{2^n \sqrt{\nu\pi/2} |na_k + iy| |1 - w^2|^{0.25}} \cdot |1 + \eta(\nu, -iw)|. \end{aligned}$$

Notice that $y - \nu \cdot \Im(\sqrt{w^2-1} - \arccos(1/w)) = \nu \Im(g(w))$, where g is defined as in Lemma 8. Moreover, $\Re w \geq na_1/\nu > \pi/2$, so (A.6) from Lemma 4 applies; hence we have

$$|f(na_k + iy)| \leq \frac{430e^{\nu \Im(g(w))}}{2^n \sqrt{\nu\pi/2} |na_k + iy| |1 - w^2|^{0.25}}. \quad (\text{A.23})$$

Since $|w| \geq na_1/\nu > \pi/2 > 1.5$, we have $4/9 |w|^2 > 1$ and by the triangle inequality,

$$|w^2 - 1| \geq |w|^2 - 1 \geq 5/9 |w|^2.$$

Thus

$$|w^2 - 1|^{0.25} \geq \left(5/9 |w|^2 \right)^{0.25} > |0.5w|^{0.5} = |na_k + iy|^{0.5} \cdot (2\nu)^{-0.5}.$$

This allows to estimate

$$\sqrt{\nu\pi/2} |na_k + iy| |w^2 - 1|^{0.25} > 0.5 |na_k + iy|^{1.5}$$

and simplify (A.23) to

$$|f(na_k + iy)| < \frac{860 \exp(\nu \Im(g(w)))}{2^n |na_k + iy|^{1.5}}. \quad (\text{A.24})$$

From Lemma 8 we have $\Im(g(w)) < 0.2607$, since $\Re w \geq \pi/2 > 1$, $\Im w \geq 0$. From $\nu < n$ and $e^{0.2607}/2 < 1.541^{-1}$ we obtain

$$\frac{e^{\nu \Im(g(w))}}{2^n} < (e^{0.2607}/2)^n < 1.541^{-n}$$

which together with (A.24) yields the desired bound:

$$|f(na_k + iy)| < 860 \cdot 1.541^{-n} |na_k + iy|^{-1.5}.$$

Proof of (A.15). Suppose, $k \geq n \geq 2$. We utilise (A.1) to rewrite (A.12) as

$$f(na_k + iy) = \frac{2^{-n} (1 - e^{-2y/n})^n e^{-i\pi(2\nu+1)/4}}{\sqrt{\pi/2}} \cdot \frac{1 + \rho(\nu, na_k + iy)}{(na_k + iy)^{3/2}}, \quad (\text{A.25})$$

since $i(na_k + iy) = ina_k - y$ and

$$e^{ina_{k+1}} \cdot e^{i(na_k - \nu\pi/2 - \pi/4)} = e^{-i\pi(2\nu+1)/4} \quad \text{and} \quad e^{-y} \left(e^{y/n} - e^{-y/n} \right)^n = \left(1 - e^{-2y/n} \right)^n.$$

Now, using (A.25) we can bound

$$|f(na_k + iy) - f(na_{k+1} + iy)| \leq 2^{-n} \left| \frac{1 + \rho(\nu, na_k + iy)}{(na_k + iy)^{3/2}} - \frac{1 + \rho(\nu, na_{k+1} + iy)}{(na_{k+1} + iy)^{3/2}} \right|. \quad (\text{A.26})$$

Since $na_k \geq n^2$ and $\nu < n$, from (A.3) we have

$$|\rho(\nu, na_k + iy)| \cdot |na_k + iy| < n^2 e;$$

similarly,

$$|\rho(\nu, na_{k+1} + iy)| \cdot |na_{k+1} + iy| < n^2 e.$$

Therefore we can estimate the RHS of (A.26) via

$$\begin{aligned} & \left| \frac{1 + \rho(\nu, na_k + iy)}{(na_k + iy)^{3/2}} - \frac{1 + \rho(\nu, na_{k+1} + iy)}{(na_{k+1} + iy)^{3/2}} \right| \\ & \leq \left| \frac{1}{(na_k + iy)^{3/2}} - \frac{1}{(na_{k+1} + iy)^{3/2}} \right| + \left| \frac{\rho(\nu, na_k + iy)}{(na_k + iy)^{3/2}} \right| + \left| \frac{\rho(\nu, na_{k+1} + iy)}{(na_{k+1} + iy)^{3/2}} \right| \\ & < \left| \frac{1}{(na_k + iy)^{3/2}} - \frac{1}{(na_{k+1} + iy)^{3/2}} \right| + \frac{n^2 e}{|na_k + iy|^{5/2}} + \frac{n^2 e}{|na_{k+1} + iy|^{5/2}} \end{aligned}$$

or (since $na_{k+1} > na_k$)

$$2^n |f(na_k + iy) - f(na_{k+1} + iy)| < \left| (na_k + iy)^{-3/2} - (na_{k+1} + iy)^{-3/2} \right| + \frac{2n^2 e}{|na_k + iy|^{5/2}}. \quad (\text{A.27})$$

By the mean value theorem (applied to the function $x \mapsto (x + iy)^{-3/2}$),

$$\frac{1}{(na_k + iy)^{3/2}} - \frac{1}{(na_{k+1} + iy)^{3/2}} = \frac{3\pi n}{2(\xi + iy)^{5/2}},$$

for some $\xi \in (na_k, na_{k+1})$, which gives us

$$\left| \frac{1}{(na_k + iy)^{3/2}} - \frac{1}{(na_{k+1} + iy)^{3/2}} \right| < \frac{1.5\pi n}{|na_k + iy|^{5/2}}.$$

Combining this with (A.27) yields (A.15):

$$|f(na_k + iy) - f(na_{k+1} + iy)| \leq 2^{-n} \cdot \frac{1.5\pi n + 2en^2}{|na_k + iy|^{5/2}} < \frac{15n^2}{2^n |na_k + iy|^{5/2}}.$$

□

A.8 The Hankel function's expansion

Proof of Lemma 4. We proceed to prove the asymptotic expansion of $H_\nu^{(1)}(z)$ when the argument and order are of similar magnitude. We relate the Hankel function to $K_\nu(z)$, the modified Bessel function of the second kind, via [10, Eq. 10.27.8]

$$K_\nu(z) = 0.5\pi e^{(\nu+1)\pi i/2} H_\nu^{(1)}(iz), \quad -\pi \leq \arg z \leq \pi/2. \quad (\text{A.28})$$

In Appendix A.8.1 we will investigate the expansion of $K_\nu(\nu z)$ due to Olver [17, Chapter 10] and sketch a brief overview of the techniques employed. These bounds (A.29)-(A.30) are not yet explicit in the sense that the bound (A.30) depends on an unspecified variational path and has to be estimated. In Appendix A.8.2 we make the bounds explicit. Finally, in Appendix A.8.3 we use (A.28) to derive the explicit bounds on the Hankel function.

A.8.1 Error bounds for the modified Bessel function

Denote

$$\xi(z) = (1 + z^2)^{1/2} + \ln \frac{z}{1 + \sqrt{1 + z^2}},$$

where all branches take their principal values on the positive real axis and are continuous elsewhere. Consider $K_\nu(\nu z)$, when $\nu > 0$ and $|\arg z| < \pi/2$. Then [17, Chapter 10, Eq. (7.17) & Eq. (7.15)]

$$K_\nu(\nu z) = \left(\frac{\pi}{2\nu}\right)^{0.5} \frac{e^{-\nu\xi(z)}}{(1 + z^2)^{0.25}} (1 + \eta(\nu, z)) \quad (\text{A.29})$$

$$|\eta(\nu, z)| \leq \exp\left(\frac{2\mathcal{V}_{+\infty, z}}{\nu}\right) \frac{2\mathcal{V}_{+\infty, z}}{\nu}, \quad (\text{A.30})$$

with $\mathcal{V}_{+\infty, z} \in \mathbb{R}$ explained later. These estimates are derived by approximating the solutions of the differential equation [17, Eq. (7.02), p. 374],

$$\frac{d^2 w}{dz^2} = \left(\nu^2 \frac{1 + z^2}{z^2} - \frac{1}{4z^2}\right) w,$$

which is satisfied by $z^{1/2} K_\nu(\nu z)$; in the differential equation z is confined to the half-plane $\Re z > 0$. In the subsequent analysis, change of variables $z \mapsto \xi(z)$ takes place, mapping the half-plane $\Re z > 0$ to a region in the ξ plane consisting of the half-plane $\Re \xi > 0$ and the half-strip $|\Im \xi| < \frac{1}{2}\pi, \Re \xi \leq 0$.




Figure 3: Sketch of the variational path

In Fig. 3 we illustrate both regions; note that Fig. 3 essentially reproduces Figs. 7.1-7.2 from [17, p.376] (except for the variational path). On the left, the z plane and the half-plane $\Re z > 0$ is shown; on the right, the ξ plane with the image of $\Re z > 0$ is illustrated. In the ξ plane, we depict a few contours and points on them; on the left, the preimages of these contours and points are shown. The shaded regions are the “shadow regions” in Olver’s terminology.

The quantity $\mathcal{V}_{+\infty, z}$ appearing in (A.30) is defined as the total variation of [17, Eq. (7.11), p.376] the function

$$U_1(p) = \frac{3p - 5p^3}{24}, \quad p := (1 + z^2)^{-1/2}, \quad (\text{A.31})$$

along any ξ -progressive path connecting $+\infty$ with $z : \Re z > 0$. A path γ is said to be ξ -progressive [17, p.222] if 1) γ is a piecewise C^2 -path and 2) $\Re \xi(\gamma)$ is non-increasing as γ passes from $+\infty$ to z .

The case $\Re z > 0$ is not sufficient for us, however, since (A.28) effectively rotates the argument of $H_\nu^{(1)}$ in the complex plane by $-\pi/2$. In order to prove (A.5) also for real w , we need the

expansion of $K_\nu(\nu z)$ also when $\arg z = -\pi/2$. Fortunately, the estimates (A.29)-(A.30) remain valid [17, Chapter 10, §8.2] also for $\arg z = -\pi/2$, provided that $|z|$ is bounded away from 1 and the variational path for $\mathcal{V}_{+\infty, z}$ is correctly constructed (i.e., the path is ξ -progressive).

Typically the variational path for $\mathcal{V}_{+\infty, z}$ is chosen so that in the ξ plane the image of the path travels³ from $\xi(z)$ parallel to the imaginary axis until the real axis is reached, then proceeding along the real axis to $+\infty$, see, e.g., [17, Chapter 10, §7.5], [21, p. 764] or [7, p. 2133].

However, when z might be of the form $z = -ci$ for some $c > 1$ (as in our setting), this approach is not suitable, since the path must avoid the point $z = -i$. Instead, for $z = -ci$ we form a path as follows: travel parallel to the real axis until $z_1 = c(1 - i)$ is reached (satisfying $\arg z_1 = -\pi/4$), then proceed from z_1 as described previously. The path is sketched in Fig. 3 (on the left), with its image on the right.

In Appendix A.8.2 we show that the described path is indeed ξ -progressive and estimate the total variation along this path.

A.8.2 Explicit error bounds for the modified Bessel function

Fix any $c > 1$ and define $\gamma_0(s) = c(s - i)$, $s \in [0, 1]$ and $z_0 = \gamma_0(0) = -ci$, $z_1 = \gamma_0(1) = c(1 - i)$. We shall show that

1. $\Re \xi(\gamma_0(s))$ is non-decreasing (i.e., the described path from $+\infty$ to z is valid);
2. $\mathcal{V}_{z_1, z_0} \leq \frac{c^2(c^2+2)}{\sqrt{8}(c^2-1)^{2.5}}$, where \mathcal{V}_{z_1, z_0} is the variation of U_1 along γ_0 .

Since the described path connects z_0 to z_1 and z_1 to $+\infty$ with $\Re \xi(z)$ is non-decreasing (and the path is clearly piecewise C^2), we conclude that the construction ensures a ξ -progressive path. Furthermore, the variation $\mathcal{V}_{+\infty, z_1}$ can [7, Eq. (5.13)] be bounded as $\frac{1}{12} + \frac{1}{6\sqrt{5}} + \left(\frac{4}{27}\right)^{1/4}$. Therefore, since $\mathcal{V}_{+\infty, z_0} = \mathcal{V}_{+\infty, z_1} + \mathcal{V}_{z_1, z_0}$, we can estimate $\mathcal{V}_{+\infty, z_0}$ as

$$\mathcal{V}_{+\infty, z_0} \leq \frac{1}{12} + \frac{1}{6\sqrt{5}} + \left(\frac{4}{27}\right)^{1/4} + \frac{c^2(c^2+2)}{\sqrt{8}(c^2-1)^{2.5}}, \quad c := -\Im z_0 > 1. \quad (\text{A.32})$$

Finally, the estimate (A.32) remains valid when $z_0 = -ci$ is replaced by $z = x - ci$ with any $x > 0$:

- If $x \in (0, c]$, then z lies on the described path from z_0 to $+\infty$, therefore $\mathcal{V}_{+\infty, z}$ is upper-bounded by $\mathcal{V}_{+\infty, z_0}$;
- if $x > c$, then $\arg z \in (-\pi/4, 0)$ and the bound [7, Eq. (5.13)] applies; then, $\mathcal{V}_{+\infty, z}$ is upper-bounded by $\frac{1}{12} + \frac{1}{6\sqrt{5}} + \left(\frac{4}{27}\right)^{1/4}$.

The path is ξ -progressive. Since ξ is symmetric around the real axis, i.e., $\xi(\bar{z}) = \overline{\xi(z)}$, we have $2\Re \xi(z) = \xi(z) + \xi(\bar{z})$. Define $h(s) = \xi(c(s - i)) + \xi(c(s + i))$, then $2\Re \xi(\gamma_0(s)) = h(s)$ and we need to show that $h : [0, 1] \rightarrow \mathbb{R}$ is non-decreasing.

Let $s \in [0, 1]$; denote $z = \gamma(s) = c(s - i)$ and $\omega = \sqrt{1 + z^2}$. Since⁴ $\frac{d\xi}{dz} = \frac{\sqrt{1+z^2}}{z}$, we find that

$$h'(s) = c \frac{d\xi}{dz} (c(s - i)) + c \frac{d\xi}{dz} (c(s + i)) = c \frac{\omega}{z} + c \frac{\bar{\omega}}{\bar{z}} = \frac{\omega \bar{z} + z \bar{\omega}}{c(s^2 + 1)} = \frac{2\Re(\omega \bar{z})}{c(s^2 + 1)}.$$

Since $z = c(s - i)$ satisfies $\arg(z) \in [-\pi/2, -\pi/4]$, it follows that $\arg(z^2) \in [-\pi, -\pi/2]$ and⁵ $\arg(z^2 + 1) \in [-\pi, 0)$. Consequently, $\arg(\omega) \in [-\pi/2, 0)$. Since $\arg(\bar{z}) \in [\pi/4, \pi/2]$, we obtain $\arg(\omega \bar{z}) = \arg(\omega) + \arg(\bar{z}) \in [-\pi/4, \pi/2)$, thus $\Re(\omega \bar{z}) > 0$. We see that $h'(s)$ is positive for $s \in [0, 1]$, and $h(s) = 2\Re \xi(\gamma_0(s))$ is non-decreasing as required.

³In fact, the construction describes the *reverse* path connecting z and $+\infty$; along the described path $\Re \xi(z)$ must be non-decreasing. This distinction is unimportant for the value of the variation and from now on we shall ignore it.

⁴In fact, in [17] this is the defining property of ξ .

⁵To exclude the possibility $\arg(z^2 + 1) = 0$, one must also take into account that $|z| \geq c > 1$.

Estimate of the variation \mathcal{V}_{z_1, z_0} . It is worth recalling that, for a holomorphic function f in a complex domain D containing a piecewise continuously differentiable path $\gamma(s)$, $s \in [s_0, s_1]$, the total variation of f along the path γ is defined as

$$\mathcal{V}_\gamma(f) = \int_{s_0}^{s_1} |f'(\gamma(s))\gamma'(s)| \, ds.$$

We have

$$\mathcal{V}_{z_1, z_0} = \int_0^1 \left| \frac{dU_1}{dz}(p(z)) \right|_{z=\gamma_0(s)} |\gamma'_0(s)| \, ds.$$

Since

$$\frac{dU_1(p)}{dp} = \frac{1-5p^2}{8}, \quad \frac{dp(z)}{dz} = \frac{-z}{(z^2+1)^{3/2}} \quad \text{and} \quad \gamma'_0(s) = c,$$

the integrand equals

$$\frac{1}{8} |1-5(1+z^2)^{-1}| \cdot \frac{|z|}{|z^2+1|^{-3/2}} \cdot c = \frac{c}{8} \cdot \frac{|z(z^2-4)|}{|z^2+1|^{5/2}}, \quad z := c(s-i).$$

From $|z \pm i| = |cs - (c \mp 1)i| \geq c \mp 1$ we have $|z^2+1|^{5/2} \geq (c^2-1)^{2.5}$. On the other hand, $|z| \leq c|1-i| = c\sqrt{2}$ and $|z^2-4| \leq |z|^2+4 \leq 2c^2+4$. Since the obtained bound is independent of z , the integral satisfies

$$\mathcal{V}_{z_1, z_0} \leq \frac{\sqrt{2}c^2(2c^2+4)}{8(c^2-1)^{2.5}} = \frac{c^2(c^2+2)}{\sqrt{8}(c^2-1)^{2.5}},$$

as claimed.

A.8.3 Explicit error bounds for the Hankel function

Let $w \in \mathbb{C}$ be such that $\arg w \in [0; \pi/2)$, then $z = -iw$ satisfies $\arg z \in [-\pi/2; 0)$ and (A.28) with (A.29) imply

$$H_\nu^{(1)}(\nu w) = \frac{2}{\pi} e^{-(\nu+1)\pi i/2} K_\nu(\nu z) = -i \sqrt{\frac{2}{\pi\nu}} \frac{e^{-\nu(\xi(-iw)+i\pi/2)}}{(1-w^2)^{0.25}} (1 + \eta(\nu, -iw)). \quad (\text{A.33})$$

To see that we can rewrite (A.33) as (A.5), simplify⁶ $\xi(-iw)$ as

$$\begin{aligned} \xi(-iw) &= (1-w^2)^{1/2} + \ln \frac{-iw}{1+\sqrt{1-w^2}} = (1-w^2)^{1/2} - \ln \frac{i(1+\sqrt{1-w^2})}{w} \\ &= (1-w^2)^{1/2} - \ln \left(w^{-1} - i\sqrt{1-w^2} \right) - \pi i/2 = (1-w^2)^{1/2} + i \arccos(1/w) - \pi i/2 \\ &= -i(w^2-1)^{1/2} + i \arccos(1/w) - \pi i/2 \end{aligned}$$

The show the last equality, notice that $\arg(w^2-1) \in [0; \pi)$, thus $\arg(i(w^2-1)^{1/2}) \in [\pi/2; \pi)$ and $\arg(\sqrt{1-w^2}) \in [-\pi/2, 0)$, hence $\sqrt{1-w^2} = -i\sqrt{w^2-1}$.

Finally, to show (A.6), notice that the function $\frac{1}{12} + \frac{1}{6\sqrt{5}} + \left(\frac{4}{27}\right)^{1/4} + \frac{c^2(c^2+2)}{\sqrt{8}(c^2-1)^{2.5}}$ is decreasing in c for $c > 1$ (seen by differentiating with respect to c). Since it takes value $2.272365\dots$ at $c = \pi/2$, for $\nu > 1$ and $c = \Re(w) \geq \pi/2$ we have

$$|1 + \eta(\nu, -iw)| \leq 1 + \exp\left(\frac{2\mathcal{V}_{+\infty, -iw}}{\nu}\right) \frac{2\mathcal{V}_{+\infty, -iw}}{\nu} \leq 1 + 2 \cdot 2.273e^{2 \cdot 2.273} < 430.$$

□

⁶In particular, this argument expresses $\xi(z) - z$ as $i(g(iz) - \pi/2)$ for $z : \arg z \in [-\pi/2, 0)$, with g defined as in Lemma 8. Thus, taking into account the symmetry of ξ around the real axis, Lemma 8 upper bounds the real part of $z - \xi(z)$ for $z : \Re z > 0, |\Im z| \geq 1$.