

Path-connectedness of the intersection of translates of $St(n, H)$

Nizar El Idrissi, Samir Kabbaj, and Brahim Moalige

February 1, 2022

Abstract

If H is a Hilbert space, the Stiefel manifold $St(n, H)$ is formed by all the independent n -tuples in H . In this article, we contribute to the topological study of Stiefel manifolds by proving a path-connectedness result. We prove that the intersection of translates of $St(n, H)$ is path-connected by polygonal paths under a condition on the codimension of the span of the components of the translating n -tuples. We rely on a lemma that we prove for the occasion.

Contents

1	Introduction	1
2	Preliminaries	2
2.1	Notation	2
2.2	Connectedness properties of $St(n, H)$	2
3	Path-connectedness	3

1 Introduction

If H is a Hilbert space, the Stiefel manifold $St(n, H)$ is the set of independent n -frames in H in the geometric sense. The term n -frame is used in geometry to simply denote an independent n -tuple. Stiefel manifolds are studied in differential

2020 *Mathematics Subject Classification.* 57N20; 54D05.

Key words and phrases. Stiefel manifold, path-connected space.

topology and are one of the fundamental examples in this area. The theory of finite dimensional Stiefel manifolds is summarized in [5] and also covered in ([4],[?]). The theory of infinite dimensional Stiefel manifolds is less studied and some recent results can be found in ([1],[3]).

In this article, we contribute to the topological study of Stiefel manifolds by proving a path-connectedness result. We prove that the intersection of translates of $St(n, H)$ is path-connected under a condition on the codimension of the span of the components of the translating n -tuples. We rely on a lemma that we prove for the occasion.

Plan of the article. This article is organized as follows. In section 2, we set some notations and present Stiefel manifolds and their connectedness properties. In the main section 3, we prove that the intersection of translates of $St(n, H)$ is path-connected by polygonal paths under a condition on the codimension of the span of the components of the translating n -tuples. We rely on a lemma that we prove for the occasion.

2 Preliminaries

2.1 Notation

\mathbb{N} denotes the set of natural numbers including 0 and $\mathbb{N}^* = \mathbb{N} \setminus \{0\}$. We denote by n an element of \mathbb{N}^* and by H a Hilbert space over \mathbb{R} or \mathbb{C} .

2.2 Connectedness properties of $St(n, H)$

In this subsection, we introduce $St(n, H)$ and $St_o(n, H)$ and recall their connectedness properties. We recall that n is a fixed element of \mathbb{N}^* and H is a Hilbert space. Observe that $St(n, H)$ is non-empty exactly when $\dim(H) \geq n$. In the following, we will always suppose this condition.

Definition 2.1. The *Stiefel manifold of independent n -frames in H* is defined by $St(n, H) := \{h = (h_1, \dots, h_n) \in H^n : \{h_1, \dots, h_n\} \text{ is free}\}$. The *Stiefel manifold of orthonormal n -frames in H* is defined by $St_o(n, H) := \{h = (h_1, \dots, h_n) \in H^n : \{h_1, \dots, h_n\} \text{ is an orthonormal system}\}$.

Definition 2.2. Let X be a topological space and $m \in \mathbb{N}$. Then X is said to be m -connected if its homotopy groups $\pi_i(X)$ are trivial for all $i \in [0, m]$.

Proposition 2.1. (see pp. 382-383 of [4] and [7]) We have

1. $St(n, \mathbb{R}^k)$ is $(k - n - 1)$ -connected.

2. $St(n, \mathbb{C}^k)$ is $(2k - 2n)$ -connected.
3. If H is infinite dimensional, then $St(n, H)$ is contractible.

3 Path-connectedness

Definition 3.1. Let E be a topological vector space and $\gamma : [0, 1] \rightarrow E$ be a continuous path. We say that γ is a *polygonal path* if there exists $q \in \mathbb{N}^*$, $(e_k)_{k \in [1, q]}$ and $(f_k)_{k \in [1, q]}$ two finite sequences with $e_k, f_k \in E$ for all $k \in [1, q]$, and $(\gamma_k)_{k \in [1, q]}$ a finite sequence of (continuous) straight paths with

$$\gamma_k = \begin{cases} [\frac{k-1}{q}, \frac{k}{q}] & \rightarrow E \\ t & \mapsto q(t - \frac{k-1}{q})f_k + q(\frac{k}{q} - t)e_k \end{cases} \text{ such that } \gamma = \gamma_1 * \cdots * \gamma_q, \text{ where } *$$

is the path composition operation. We say that a subset $S \subseteq E$ is *polygonally connected* if every two points of S are connected by a polygonal path.

In the following, when we say that $S \subseteq E$ is polygonally connected, we mean that each two points of S are connected by a polygonal path of the type $\gamma_1 * \gamma_2$ where γ_1 and γ_2 are two straight paths.

Before we prove the main proposition of this section, let's prove a useful lemma.

Lemma 3.1. 1. Suppose we have a family $(a(j))_{j \in J}$ indexed by J where each $a(j)$ belongs to $St(n, H)$. Then if $(a(j)_k)_{j \in J, k \in [1, n]}$ is free, we have

$$\text{span}(\{a(j)\}_{j \in J}) \setminus \{0\} \subseteq St(n, H)$$

2. Suppose we have a family indexed by J where each $a(j)$ belongs to $St_o(n, H)$ for all $j \in J$. Then if $(a(j)_k)_{j \in J, k \in [1, n]}$ is an orthonormal system, we have

$$\{x \in \text{span}(\{a(j)\}_{j \in J}) : \|x\| = \sqrt{n}\} \subseteq St_o(n, H)$$

Proof. 1. Let $h = (h_1, \dots, h_n) \in \text{span}(\{a(j)\}_{j \in J}) \setminus \{0\}$. We can write $h = \sum_{u=1}^r \lambda_u a(j_u)$ with $\lambda_u \in \mathbb{F}$ for all $u \in [1, r]$ and the λ_u 's are not all zeros. We need to show that (h_1, \dots, h_n) is an independent system. Suppose otherwise $\sum_{k=1}^n c_k h_k = 0$. This means that $\sum_{k=1}^n c_k (\sum_{u=1}^r \lambda_u a(j_u)_k) = 0$, and so $\sum_{k=1}^n \sum_{u=1}^r (\lambda_u c_k) a(j_u)_k = 0$. Since $\bigcup_{j \in J} \bigcup_{k \in [1, n]} \{a(j)_k\}$ is free, we deduce that $\lambda_u c_k = 0$ for all $u \in [1, r]$ and $k \in [1, n]$, which implies that $c_k = 0$ for all $k \in [1, n]$ since the λ_u 's are not all zeros. Therefore, $h \in St(n, H)$.

2. Let $h = (h_1, \dots, h_n) \in \text{span}(\{a(j)\}_{j \in J})$ such that $\|h\| = \sqrt{n}$. We can write $h = \sum_{u=1}^r \lambda_u a(j_u)$ with $\lambda_u \in \mathbb{F}$ for all $u \in [1, r]$. We need to show

that $\langle h_k, h_l \rangle = \delta_{k,l}$ for all $k, l \in [1, n]$. For $k \neq l$, we have : $\langle h_k, h_l \rangle = \langle \sum_{u=1}^r \lambda_u a(j_u)_k, \sum_{u=1}^r \lambda_u a(j_u)_l \rangle = \sum_{u=1}^r \sum_{v=1}^r \lambda_u \overline{\lambda_v} \langle a(j_u)_k, a(j_v)_l \rangle = 0$ since $\bigcup_{j \in J} \bigcup_{k \in [1, n]} \{a(j)_k\}$ is an orthogonal system. Moreover, $\|h_k\|^2 = \sum_{u=1}^r \sum_{v=1}^r \lambda_u \overline{\lambda_v} \langle a(j_u)_k, a(j_v)_k \rangle = \sum_{u=1}^r |\lambda_u|^2 \|a(j_u)_k\|^2 = \sum_{u=1}^r |\lambda_u|^2$ since $\bigcup_{j \in J} \bigcup_{k \in [1, n]} \{a(j)_k\}$ is an orthonormal system. By hypothesis, $n = \|h\|^2 = \sum_{k=1}^n \|h_k\|^2 = n(\sum_{u=1}^r |\lambda_u|^2)$, so $\|h_k\|^2 = \sum_{u=1}^r |\lambda_u|^2 = 1$ for all $k \in [1, n]$ as desired. \square

Proposition 3.1. *Let H be a Hilbert space with $\dim(H) \geq n$, J an index set and $(U(j))_{j \in [1, l]}$ a family with $U(j) \in H^n$ for all $j \in J$. If $\text{codim}_H(\text{span}(\{u(j)_k : j \in J, k \in [1, n]\})) \geq 3n$, then $\bigcap_{j \in J} (U(j) + \text{St}(n, H))$ is polygonally-connected.*

Proof. Let $X = (x_1, \dots, x_n)$ and $Y = (y_1, \dots, y_n)$ in $\bigcap_{j \in J} (U(j) + \text{St}(n, H))$. Let (z_1, \dots, z_n) be an independent family in H such that

$$\text{span}(\{z_k : k \in [1, n]\}) \cap \text{span}(\{x_k : k \in [1, n]\} \cup \{y_k : k \in [1, n]\} \cup \{u(j)_k : j \in J, k \in [1, n]\}) = \{0\}$$

This is possible since $\text{codim}_H(\text{span}(\{u(j)_k : j \in J, k \in [1, n]\})) \geq 3n$.

This ensures that we have for all $j \in J$

$$\text{span}(\{-u(j)_k + z_k : k \in [1, n]\}) \cap \text{span}(\{-u(j)_k + x_k : k \in [1, n]\}) = \{0\},$$

$$\text{span}(\{-u(j)_k + z_k : k \in [1, n]\}) \cap \text{span}(\{-u(j)_k + y_k : k \in [1, n]\}) = \{0\},$$

and

$$(-u(j)_k + z_k)_{k \in [1, n]} \text{ is independent.}$$

We define the straight paths

$$\begin{cases} \gamma_1 & : [0, 1] \rightarrow H^n \\ \gamma_2 & : [0, 1] \rightarrow H^n \end{cases}$$

by $\gamma_1(t) = tZ + (1-t)X$ and $\gamma_2(t) = tY + (1-t)Z$ respectively.

We have $\gamma_1(0) = X$, $\gamma_1(1) = \gamma_2(0) = Z$, and $\gamma_2(1) = Y$.

Since for all $j \in J$

$$\text{span}(\{-u(j)_k + z_k : k \in [1, n]\}) \cap \text{span}(\{-u(j)_k + x_k : k \in [1, n]\}) = \{0\},$$

we have for all $t \in [0, 1]$ and $j \in J$

$$-U(j) + tZ + (1-t)X = t(-U(j) + Z) + (1-t)(-U(j) + X) \in \text{St}(n, H)$$

by lemma 3.1, and so $\gamma_1(t) \in \bigcap_{j \in J} (U(j) + \text{St}(n, H))$.

Similarly, $\gamma_2(t) \in \bigcap_{j \in J} (U(j) + \text{St}(n, H))$ for all $t \in [0, 1]$.

Composing γ_1 with γ_2 , we see that $\bigcap_{j \in J} (U(j) + \text{St}(n, H))$ is polygonally connected. \square

Acknowledgement

The first author is financially supported by the *Centre National pour la Recherche Scientifique et Technique* of Morocco.

References

- [1] E. Bardelli and A. C. G. Mennucci. *Probability measures on infinite-dimensional Stiefel manifolds*. Journal of Geometric Mechanics, vol. 9, 2017, pp. 291-316.
- [2] J. Cahill, D. Mixon and N. Strawn. *Connectivity and irreducibility of algebraic varieties of finite unit norm tight frames*. SIAM Journal on Applied Algebra and Geometry, vol. 1, no. 1, 2017, pp. 38-72.
- [3] P. Harms and A. C. G. Mennucci. *Geodesics in infinite dimensional Stiefel and Grassmann manifolds*. Comptes Rendus Mathématique, vol. 350, no. 15–16, August 2012, pp. 773-776.
- [4] A. Hatcher. *Algebraic topology*. <https://pi.math.cornell.edu/~hatcher/AT/AT.pdf>, 2001.
- [5] I. M. James. *The topology of Stiefel manifolds*. London Mathematical Society Lecture Note Series, 1976.
- [6] T. Needham and C. Shonkwiler. *Symplectic geometry and connectivity of spaces of frames*. Preprint: arXiv:1804.05899v2, 2019.
- [7] math.stackexchange.com : Contractibility of the sphere and Stiefel manifolds of a separable H

Nizar El Idrissi.

Laboratoire : Equations aux dérivées partielles, Algèbre et Géométrie spectrales.
Département de mathématiques, faculté des sciences, université Ibn Tofail, 14000 Kénitra.
E-mail address : `nizar.elidrissi@uit.ac.ma`

Pr. Samir Kabbaj.

Laboratoire : Equations aux dérivées partielles, Algèbre et Géométrie spectrales.
Département de mathématiques, faculté des sciences, université Ibn Tofail, 14000 Kénitra.
E-mail address : `samir.kabbaj@uit.ac.ma`

Pr. Brahim Moalige.

Laboratoire : Equations aux dérivées partielles, Algèbre et Géométrie spectrales.
Département de mathématiques, faculté des sciences, université Ibn Tofail, 14000 Kénitra.
E-mail address : `brahim.moalige@uit.ac.ma`