

When do you Stop Supporting your Bankrupt Subsidiary? A Systemic Risk Perspective

Maxim Bichuch* Nils Detering†

May 10, 2022

Abstract

We consider a network of bank holdings, where every holding has two subsidiaries of different types. A subsidiary can trade with another holding's subsidiary of the same type. Holdings support their subsidiaries up to a certain level when they would otherwise fail to honor their financial obligations. We investigate the spread of contagion in this banking network when the number of bank holdings is large, and find the final number of defaulted subsidiaries under different rules for the holding support. We also consider resilience of this multilayered network to small shocks. Our work sheds light onto the role that holding structures can play in the amplification of financial stress. We find that depending on the capitalization of the network, a holding structure can be beneficial as compared to smaller separated entities. In other instances it can be harmful and actually increase contagion. We illustrate our results in a numerical case study and also determine the optimal level of holding support from a regulator perspective.

Keywords: systemic risk, financial contagion, holdings, multilayered networks

1 Introduction

At first, financial risk was managed at an individual firm level, and has been considered on a bilateral basis. However, after the financial crisis of 2007-2009, we have gained the understanding that this risk can spread akin to a virus through the entire banking system. And similar to a virus, the risk increases the more nodes succumb to the shock. This phenomenon is known as systemic risk – the risk that a (small) shock that hits the system spreads throughout the system to a degree that it endangers the entire system. This spread is also referred to as financial contagion. There are two distinct ways for this risk to spread – through local connections (for example when one bank cannot honor its obligations to another), and through global connection (for example when asset prices are impacted, as a result of liquidations). This paper will focus on the contagion spread through local connections.

One of the first papers on the spread of contagion through local connections is Eisenberg and Noe (2001); which finds a clearing equilibrium payment in a network of banks connected by liabilities. Numerous generalizations and follow up studies include e.g., Anand et al. (2015); Halaj and Kok (2015); Boss et al. (2004); Elsinger et al. (2013); Upper (2011); Gai et al. (2011);

*Department of Applied Mathematics & Statistics, Johns Hopkins University, Baltimore, MD 21218, USA. Work is partially supported by NSF grant DMS-1736414. Research is partially supported by the Acheson J. Duncan Fund for the Advancement of Research in Statistics. Email: mbichuch@jhu.edu.

†Department of Statistics and Applied Probability, University of California, Santa Barbara, CA 93106, USA. Email: detering@pstat.ucsb.edu.

Bardoscia et al. (2017). Generalizations such as fire sales and bankruptcy costs have been considered in Elsinger (2009); Rogers and Veraart (2013); Elliott et al. (2014); Glasserman and Young (2015); Weber and Weske (2017); Capponi et al. (2016); Elsinger (2009); Elliott et al. (2014); Weber and Weske (2017); Gouriéroux et al. (2012); Cifuentes et al. (2005); Nier et al. (2007); Arinaminpathy et al. (2012); Amini et al. (2013); Chen et al. (2016); Weber and Weske (2017); Amini et al. (2016b); Bichuch and Feinstein (2019, 2022)). We also refer to, e.g., Weber and Weske (2017); Staum (2013); Hüser (2015) for additional review of this literature.

Another stream of literature considers contagion effects in random networks (Gai and Kapadia (2010); Amini et al. (2016a); Detering et al. (2019, 2020), and Detering et al. (2021) for an extension incorporating fire sales). In these papers, asymptotic methods (as the number of banks increases to infinity) are facilitated to determine the final damage to the system after some initial shock hits some banks. From the perspective of random graph theory, the results on contagion in random networks link to a process called *bootstrap percolation*. This is a process which models the spread of some activity or infection within a graph via its edges. It has been analysed for Erdos-Rényi random graphs Janson et al. (2012) and random regular graphs Balogh and Pittel (2007), and these results have later been extended to inhomogeneous graph in Amini and Fountoulakis (2014); Amini et al. (2014); Detering et al. (2019).

Notably, in all these models the banks in the network have always been thought as an indivisible atom. In other words, the entire bank must be in one state only (e.g. bankrupt or solvent). In reality this is not the case. First, big international banks are usually big holding companies, with dozens if not hundreds of subsidiaries, that are divided by business lines and locations (e.g. JPMorgan Chase¹ and AIG²). Second, a financial loss—possibly amplified through contagion—does not necessarily affect the entire bank (at least not immediately), but often starts with one (or more) of its subsidiaries (see e.g. the classical example of AIG in Lewis (2010)). The contagion then can spread both between subsidiaries of different banks (e.g. through local connections), and between different subsidiaries of the same bank (e.g. through the holding).

For corporations, becoming a holding with subsidiaries has several advantages: It allows corporations to defer taxable business income and use these earnings for other business opportunities or, by channeling income to low tax countries, reduce taxes altogether. In other instances, it allows them to limit the spillover risk if one business line is in trouble. In this case, the holding acts as a pure shareholder and enjoys limited liability. Often bank holdings naturally divide into subsidiaries based on location (i.e. Europe, US) and/or business lines (i.e. broker-dealers, commercial bank, insurance, asset management). The precise judicial setup differs across different countries but two main types of subsidiaries are prevalent:

- *Type A*: The holding acts as an asset holder with full liability for the business activity of its subsidiary. This setup is usually accompanied by a profit transfer agreement ensuring fiscal unity. It is common in Europe; and is becoming more common in the US³⁴. For the holding this structure has the advantage that profits from the subsidiary are often collected on a pretax basis and can be netted with losses from other subsidiaries or the holding. For creditors of the subsidiary, an obvious advantage is that holdings support their subsidiaries, and cover some of their losses.
- *Type B*: The holding acts as a shareholder of its subsidiaries with limited liability. This is

¹<https://www.sec.gov/Archives/edgar/data/19617/000119312508043536/dex211.htm>

²<https://www.sec.gov/Archives/edgar/data/5272/000110465920023889/exhibit21.htm>

³<https://www.fdic.gov/regulations/reform/resplans/plans/boa-165-2107.pdf>,

⁴<https://www.federalreserve.gov/econres/notes/feds-notes/foreign-banks-asset-reallocation-intermediate-holding-company-rule-of-2016-20210512.htm>

the standard holding structure in the US⁵. From a risk perspective this holding structure is beneficial for the holdings, as the default of a subsidiary does not directly impact other business lines of the same holding. It might promote greater risk taking, because losses not covered by equity are immediately beared by other market participants, and not by the holding first.

While for the corporations themselves, the advantages are imminent, whether holdings are beneficial from a society perspective is less clear. In particular it is not well understood how shocks propagate in a network of bank holdings and what influence the holding structure has on the contagion process.

In this paper we address this important research question and try to understand the influence of the holding structures on the propagation of shocks through the financial system. We are particularly interested in the effect that support of the holding for its stressed subsidiary has on contagion. Holdings might support their subsidiaries beyond their financial liability for several reasons. For example they might fear the reputation cost of a bankrupt subsidiary, or the threat of a possible change in credit rating if a subsidiary defaults. In other cases, the holding might support its subsidiary because it still believes in its business model, and considers the solvency problems to be of temporary nature. We assume that all holdings will support their subsidiaries up to the same fixed capital $x \leq 0$, and we also allow for full support ($x = -\infty$). We consider two holding types which mimic the situation A and B described above. Depending on the holding type, the financial support of the holding for one subsidiary has different implications for the holdings other subsidiary.

We pursue our analysis in a random network setup which raises also some interesting new theoretical challenges. From a technical perspective our model setup provides a first instance of a multilayered random network in which contagion spreads through different channels and where interactions between layers is through the nodes. This complicates the analysis compared to a one layer network as the state of the nodes is not determined by a one dimensional quantity, but rather lies in a multidimensional domain. Our analysis includes a multi-layer analog to the classical *bootstrap percolation* process mentioned above.

We perform our analysis in a simple toy model. For this we consider a network of banks. Every bank has two subsidiaries, which we refer to as subsidiary 1 and 2. We assume that subsidiaries only trade with subsidiaries of the same kind, i.e. subsidiary 1 of one bank might trade with subsidiary 1 of another bank but not with its subsidiary 2. The capital of the holding is the sum of the capitals of the subsidiaries, because the holding itself is not involved in any business activity. We then consider a contagion process that is triggered by a financial shock and then propagates throughout the network according to the following rules:

1. If a subsidiary has exposure to a newly defaulted subsidiary of another holding, then its capital is reduced by this exposure.
2. A subsidiary defaults as soon as:
 - Its capital has reached the maximal holding support level $x < 0$ or,
 - Its capital is greater than x , but the holding has not enough capital to support it. For holding structure of type B this means that the subsidiary's capital is non-positive. Whereas, for holding structure of type A this means that the capital of the entire holding is non-positive, and both subsidiaries will default.

⁵<https://www.wsj.com/articles/paul-kupiec-and-peter-wallison-the-fdics-bank-holding-company-heist-1419292997>

3. For holding structure of type A any support for a stressed subsidiary negatively affects the holding's other subsidiary. For holdings of type B , the support does not affect its other subsidiary.

All holdings will support their subsidiaries up to the same fixed level x , when possible. For type A holdings, the support for a subsidiary in distress has to come directly from the other subsidiary of the same holding. It can be thought of the holding reallocating capital from the healthy subsidiary to the other distressed subsidiary. For holding structures of type B , the support for a troubled subsidiary does not change the capital of the other subsidiary. Instead, this support can be thought of as coming from a borrowing transaction of the holding against the capital of the healthy subsidiary. In any case, the holding supports its subsidiaries up to the level x if possible and gives up on them when faced with substantial losses larger than x . For holdings of type A , limiting the loss from a stressed subsidiary at x is only possible by either selling this subsidiary or by offloading the assets of this subsidiary into a bad bank construction. As mentioned already, our results also cover the case $x = -\infty$ of full liability.

We investigate the above described contagion mechanism in a banking network where the number of banks is large and, under both holding types A and B , we find the final number of defaulted type 1 and type 2 subsidiaries at the end of the cascade. We further investigate resilience of this multilayered network to small shocks. We find that depending on the capitalization of the network the holding structure can be beneficial as compared to smaller separated entities. This is because the holding is able to support a distressed subsidiary with capital from a subsidiary that is better off, therefore reducing the probability of default of the distressed subsidiary. However, if this support goes too far, it can actually amplify the spread of contagion because healthy parts of the network get weakened and trigger new rounds of feedback effects. In a numerical case study we determine the optimal support level x from a regulator perspective.

The paper is structured as follows: In Section 2 we state our model and our main result regarding the final number of defaulted subsidiaries of different types. In Section 3 we derive results that allow us to classify networks according to their resilience to small shocks. We give an example of a system in which the default of one subsidiary immediately triggers the default of the holding's second subsidiary. Still this system is more resilient than its fully separated counterpart due to the fact that the default of both subsidiaries can be slightly deferred with some holding support. Section 4 contains our numerical case study in which we also determine the optimal holding support. We summarize our results in Section 5. All proofs are in Section 6.

2 Model and main result

We consider a network of n banks and label each bank by an index $i \in [n] := \{1, \dots, n\}$. Each bank is structured as a holding with two subsidiaries of different type 1 and 2. We denote the type 1 subsidiary of bank i by i_1 and the type 2 subsidiary of bank i by i_2 . We denote the set of type 1 subsidiaries by $[n_1] := \{1_1, \dots, n_1\}$ and the set of type 2 subsidiaries by $[n_2] := \{1_2, \dots, n_2\}$. We assume that subsidiaries only trade with subsidiaries of the same type, i.e. subsidiary of type 1 of bank i might trade with subsidiary of type 1 of another bank j but not with subsidiary of type 2. The type can for example originate from a business line (e.g. fixed income or equities), or from a location (e.g. US and Europe). The assumption then implies that there is no cross trading between subsidiaries of different types, and linkage is only through the holdings. We consider a network that describes these trading activities. In this toy model we assume that every loan is of equal unit size 1. For this let $e_{i,j}^l \in \{0, 1\}$, $i, j \in [n]$, $i \neq j$, $l \in \{1, 2\}$ denote the exposure of subsidiary l of bank i towards subsidiary l of bank j . We then build

a network by drawing a directed link from subsidiary l of bank i to subsidiary l of bank j if $e_{i,j}^l = 1$. We do not allow for self loops and multiple edges.

We denote by $c_{i,1} \in \mathbb{Z}$ and $c_{i,2} \in \mathbb{Z}$ the capital of subsidiary 1 and 2 of bank i . The total capital of the bank is then equal to $c_i := c_{i,1} + c_{i,2}$. This is the initial capital structure of each bank holding. As mentioned in the introduction already, in this paper we consider two types of holding structure. Type A that mimics the situation where the holding acts as an asset holder of the assets of its subsidiary and type B where the holding acts as a shareholder of its subsidiary. We assume in the following that all holdings are of the same type, either all of type A or all of type B. Our results could however easily be extended to networks that consist of type A and type B holdings. In what follows, several quantities that are introduced depend on $J \in \{A, B\}$, but in order to lighten the notation we make this dependency only specific when necessary.

We assume that some subsidiaries are in default, i.e. the set $\mathcal{D}_0 = \mathcal{S}_1 \cup \mathcal{S}_2$ is not empty, where

$$\mathcal{S}_l = \{i_l \in [n_l] : i_l \text{ is in default}\}, \quad l = 1, 2.$$

Let $x \leq 0$ be a global support level. A subsidiary i_l of bank i is in default if either its capital $c_{i,l}$ is less than x or, despite being larger than x , the capital structure of the holding, expressed by $(c_{i,1}, c_{i,2})$, is such that subsidiary l defaults. For holding structure B this is the case if the subsidiary has defaulted ($c_{i,l} \leq 0$) and the holding has no capital to support it ($c_{i,1} + c_{i,2} \leq 0$). For holding structure A due to the joint liability this can even be the case if the subsidiary l would be solvent by itself ($c_{i,l} > 0$) but due to distress of the other subsidiary the total holding capital is negative ($c_{i,1} + c_{i,2} \leq 0$). However, if $x > -\infty$, then even for type A, subsidiary l can only default if $c_{i,l} \leq -x$, because we assume that a holding gives up on its subsidiary if the subsidiary reaches a capital of x which in turn limits the negative impact on the supporting subsidiary.

The initial defaults in \mathcal{D}_0 now trigger a cascade that then evolves in *generations* or *rounds*. In each round, the contagion expands, because a defaulted subsidiary causes a loss of 1 to every subsidiary it traded with and therefore reduces their capital by 1. We denote by $c_{i,1}^{(k)}$ and $c_{i,2}^{(k)}$ the capital of subsidiary 1 and 2 of bank i in round $k \geq 0$, and set $(c_{i,1}^{(0)}, c_{i,2}^{(0)}) := (c_{i,1}, c_{i,2})$. In round k , the total capital of the bank is then equal to $c_{i,1}^{(k)} + c_{i,2}^{(k)}$.

Following above considerations, for bank i with capital structure $(c_{i,1}^{(k)}, c_{i,2}^{(k)})$, subsidiary $l \in \{1, 2\}$ is in default in round k if one of the following conditions hold:

- $c_{i,l}^{(k)} \leq x$.
- For a holding structure of type A: $c_{i,1}^{(k)} + c_{i,2}^{(k)} \leq 0$ and $c_{i,l}^{(k)} \leq -x$.
For a holding structure of type B: $c_{i,1}^{(k)} + c_{i,2}^{(k)} \leq 0$ and $c_{i,l}^{(k)} \leq 0$.

If subsidiary i_l of bank i defaults in round k , then $c_{j,l}^{(k+1)} = c_{j,l}^{(k)} - 1$ for all $j \in \{j \in [n] : e_{i,j}^l \neq 0\}$ unless subsidiary j_l had already defaulted. Updating the capitals of all subsidiaries ends the round and the next round then starts.

In round k , the set of banks with defaulted type l subsidiary is denoted by

$$\mathcal{S}_{l,k} = \{i_l \in [n_l] : \text{subsidiary } i_l \text{ is in default in round } k\}, \quad (1)$$

and $\mathcal{D}_k = \mathcal{S}_{1,k} \cup \mathcal{S}_{2,k}$ denotes the set of all defaulted subsidiaries at step k . This then leads to the two cascades

$$\mathcal{S}_{l,0} \subset \mathcal{S}_{l,1} \subset \dots, \quad l = 1, 2,$$

where both cascade processes are strongly coupled through the holdings. The cascade of all subsidiaries is given by

$$\mathcal{D}_0 \subset \mathcal{D}_1 \subset \dots$$

This cascade stops after at most $2n$ iterations and we denote by $\mathcal{S}_{l,2n-1}$ the final sets of defaulted subsidiaries for $l \in \{1, 2\}$. Such a process is exemplified for a small network with $n = 3$ and support level $x = -1$ in Figure 1.

In the following we want to determine $n^{-1} |\mathcal{S}_{l,2n-1}|$ in a random network setting. For this let us first stress that when a holding supports a subsidiary that has a non-positive capital, we formalize this support not by actually increasing the capital of the supported subsidiary in order to lift it above 0, but by reducing the threshold at which it defaults by 1. Similarly in case of the holding type A , the support for one subsidiary does not reduce the other subsidiary's capital but instead increases the level at which the other subsidiary defaults to above zero. While both descriptions are equivalent and lead to the same defaulted subsidiaries, actually changing the capital to formalize the holding support is less convenient for the mathematical analysis. In fact, our approach of changing the default level has the advantage that in order to determine whether a subsidiary of a holding with capital structure $(c_{i,1}^{(k)}, c_{i,2}^{(k)})$ has defaulted, one simply needs to check whether $(c_{i,1}^{(k)}, c_{i,2}^{(k)})$ is in some subset of $\mathbb{Z} \times \mathbb{Z}$. This subset is the default region and it differs for types A and B . We shall specify the default regions now.

For simplicity we assume that all capitals are bounded by $R \in \mathbb{N}$, and because a subsidiary defaults always when the capital is less or equal than x , we do not need to consider any capitals less than x . We therefore define $\mathbf{D} := (\mathbb{Z} \cap [x, R]) \times (\mathbb{Z} \cap [x, R])$ as the domain for the holding capital. Above consideration then leads us to the default regions $\mathbf{D}_{A,1}, \mathbf{D}_{A,2} \subset \mathbf{D}$ ($\mathbf{D}_{B,1}, \mathbf{D}_{B,2} \subset \mathbf{D}$ respectively) for subsidiary 1 and subsidiary 2 and for holding of type A , respectively B . They are explicitly given by

$$\mathbf{D}_{A,1} = \{(j, k) \in \mathbf{D} | j \leq x \text{ or } j + k \leq 0, j \leq -x\}, \quad (2)$$

$$\mathbf{D}_{A,2} = \{(j, k) \in \mathbf{D} | k \leq x \text{ or } j + k \leq 0, k \leq -x\}, \quad (3)$$

$$\mathbf{D}_{B,1} = \{(j, k) \in \mathbf{D} | j \leq x \text{ or } j + k \leq 0, j \leq 0\}, \quad (4)$$

$$\mathbf{D}_{B,2} = \{(j, k) \in \mathbf{D} | k \leq x \text{ or } j + k \leq 0, k \leq 0\}. \quad (5)$$

With the help of these regions we may then rewrite

$$\mathcal{S}_{l,k} = \{i_l \in [n_l] : (c_{i,1}^{(k)}, c_{i,1}^{(k)}) \in \mathbf{D}_{J,l}\},$$

for $J \in \{A, B\}$. Recall that to simplify the notation the dependency on J of $\mathcal{S}_{l,k}$ is implicit. In Figure 2 we display the default regions for both holding structures of type A and B . By the definition of the contagion process for type B it follows trivially that smaller x leads to less infections. However, we will see in the following that the situation for type A is more complicated and interesting phenomena arise.

In order to determine $n^{-1} |\mathcal{S}_{l,2n-1}|$ we need to specify our random network setting. We fix $p_1 \in \mathbb{R}_+$ and $p_2 \in \mathbb{R}_+$ and assume that $e_{i,j}^l$ is present with probability p_l/n for $l \in \{1, 2\}$ and that all edges are mutually independent. Instead of a fixed network we analyse a sequence of networks of increasing size. For $n \in \mathbb{N}$, let as before $c_{i,1} = c_{i,1}(n)$ and $c_{i,2} = c_{i,2}(n)$ be the initial capital of subsidiary 1, respectively 2 of bank i . Let then $\mathbf{c}_1(n) = (c_{1,1}(n), \dots, c_{n,1}(n))$ and $\mathbf{c}_2(n) = (c_{1,2}(n), \dots, c_{n,2}(n))$ be the capitals of the subsidiaries for a network of size n . We will often drop the dependency on n in the notation when it does not lead to confusion. Let $X_{i,j}^l$ be the indicator that a directed edge from subsidiary l of bank i to subsidiary l of bank j is present. Let $D_i^{l-} := \sum_{j \in [n] \setminus \{i\}} X_{j,i}^l$ and $D_i^{l+} := \sum_{j \in [n] \setminus \{i\}} X_{i,j}^l$ the in- respectively out-degree




Figure 1: Illustration of an exemplary contagion process for a network with three holdings of type A with a support level $x = -1$. The pair of two circles arranged vertically form a holding with the top circle denoting the type 1 subsidiary and the bottom circle the type 2 subsidiary. The edges between type 1 subsidiaries are drawn in black and the edges between type 2 subsidiaries are drawn in green. The cascade start (round 0) is shown in the center upper graph. The type 1 subsidiary of bank 1 has defaulted because its capital has reached the maximal support level and therefore defaults although the holding capital is still positive. This leads to $\mathcal{S}_{1,0} = \{1_1\}$ and $\mathcal{S}_{2,0} = \emptyset$ which triggers a short cascade evolving in two rounds. In round 1, shown in the lower left graph, subsidiary 1 of bank 3 loses 1 unit in capital, so its capital then becomes $c_{3,1}^{(1)} = -1$, which causes the entire bank holding 3 to default and the round ends with $(\mathcal{S}_{1,1} = \{1_1, 3_1\}$ and $\mathcal{S}_{2,1} = \{3_2\})$. The result of round 2 which is pictured in the lower right graph is that subsidiary 2 of bank 2 loses one unit of capital, and its terminal capital is $c_{2,1}^{(2)} = 2$ which does not lead to any further defaults ($\mathcal{S}_{1,2} = \{1_1, 3_1\}$ and $\mathcal{S}_{2,2} = \{3_2\}$).

of subsidiary l of bank i . Clearly then $\mathbb{E}[D_i^{l\pm}] = p_l + o(1)$ and $D_i^{l\pm} \xrightarrow{p} \text{Poi}(p_l)$ as $n \rightarrow \infty$, where \xrightarrow{p} denotes convergence in probability.

In order to ensure that our asymptotic statements are relevant also for networks of finite size we shall need some regularity. In particular we require that the proportion of banks with a certain capital structure stabilizes as the network size increases. This is ensured by the following standing assumption for the rest of this paper:

Assumption 2.1. For each $n \in \mathbb{N}$, denote the joint empirical distribution function of $\mathbf{c}_1(n)$ and $\mathbf{c}_2(n)$ by

$$F_n(x, y) = n^{-1} \sum_{i \in [n]} \mathbf{1}_{\{c_{i,1}(n) \leq x, c_{i,2}(n) \leq y\}}, \quad (x, y) \in \mathbf{D}.$$

We assume that there exists a distribution F on \mathbf{D} such that it holds $\lim_{n \rightarrow \infty} F_n(x, y) = F(x, y)$ for all $(x, y) \in \mathbf{D}$. Denote by (C_1, C_2) a random vector distributed according to F .

Considering a network of holding types $J \in \{A, B\}$, we note that for fixed $n \in \mathbb{N}$ the contagion process only starts if there exists at least one bank i such that $(c_{i,1}, c_{i,2}) \in \mathbf{D}_{J,1} \cup \mathbf{D}_{J,2}$. Under Assumption 2.1, this will be the case for n large if $\mathbb{P}((C_1, C_2) \in \mathbf{D}_{J,1} \cup \mathbf{D}_{J,2}) > 0$.




Figure 2: (a) Illustration for type *A* holding. (b) illustration for type *B* holding. Blue filled area is the default region of subsidiary 1, red filled area is the default region of subsidiary 2. The blue and red dotted lines correspond to the boundaries $\partial\mathbf{D}_{A,1}$ and $\partial\mathbf{D}_{A,2}$ for type *A* (respectively $\partial\mathbf{D}_{B,1}$ and $\partial\mathbf{D}_{B,2}$ for type *B*.)

We further define boundaries of the default regions by:

$$\partial\mathbf{D}_{A,1} = \{(j, k) \in \mathbf{D}_{A,1}^c | j = x + 1 \text{ or } j + k = 1, x < j \leq -x \text{ or } j = -x + 1, k \leq x\}, \quad (6)$$

$$\partial\mathbf{D}_{A,2} = \{(j, k) \in \mathbf{D}_{A,2}^c | k = x + 1 \text{ or } j + k = 1, x < k \leq -x \text{ or } k = -x + 1, j \leq x\}, \quad (7)$$

$$\partial\mathbf{D}_{B,1} = \{(j, k) \in \mathbf{D}_{B,1}^c | j = x + 1 \text{ or } j + k = 1, j \leq 0 \text{ or } j = 1, k \leq 0\}, \quad (8)$$

$$\partial\mathbf{D}_{B,2} = \{(j, k) \in \mathbf{D}_{B,2}^c | k = x + 1 \text{ or } j + k = 1, k \leq 0 \text{ or } k = 1, j \leq 0\}. \quad (9)$$

For example for a holding to be in $\partial\mathbf{D}_{A,1}$ means that its subsidiary 1 is at risk of default. This is the case either because the subsidiary defaults as soon as it has a new defaulted neighbor ($j = x + 1$ or $j + k = 1$, $x < j \leq -x$) itself or because it defaults as soon as the other subsidiary has a new defaulted neighbor ($j = x + 1$). We refer again to Figure 2 for an illustration of these sets.

Moreover, we define the following subsets $\partial\mathbf{D}_{Jlm}$ to be the sets of all points such that a link to subsidiary m results in a default of subsidiary l , for $J \in \{A, B\}, l, m \in \{1, 2\}$. The points on the boundary from which transition to the default region is possible through links from subsidiary 1 are given by:

$$\partial\mathbf{D}_{A11} = \{(j, k) \in \mathbf{D}_{A,1}^c | j = x + 1 \text{ or } j + k = 1, x < j \leq -x \text{ or } j = -x + 1, k \leq x\}, \quad (10)$$

$$\partial\mathbf{D}_{A21} = \{(j, k) \in \mathbf{D}_{A,2}^c | j + k = 1, k \leq -x\}, \quad (11)$$

$$\partial\mathbf{D}_{B11} = \{(j, k) \in \mathbf{D}_{B,1}^c | j = x + 1 \text{ or } j = 1, k \leq 0 \text{ or } j + k = 1, j \leq 0\}, \quad (12)$$

$$\partial\mathbf{D}_{B21} = \{(j, k) \in \mathbf{D}_{B,2}^c | j + k = 1, k \leq 0\}, \quad (13)$$

and for those through links from subsidiary 2:

$$\partial\mathbf{D}_{A^{12}} = \{(j, k) \in \mathbf{D}_{A,1}^c | j + k = 1, j \leq -x\}, \quad (14)$$

$$\partial\mathbf{D}_{A^{22}} = \{(j, k) \in \mathbf{D}_{A,2}^c | k = x + 1, x < k \leq -x \text{ or } j + k = 1 \text{ or } k = -x + 1, j \leq x\}, \quad (15)$$

$$\partial\mathbf{D}_{B^{12}} = \{(j, k) \in \mathbf{D}_{B,1}^c | j + k = 1, j \leq 0\}, \quad (16)$$

$$\partial\mathbf{D}_{B^{22}} = \{(j, k) \in \mathbf{D}_{B,2}^c | k = x + 1 \text{ or } k = 1, j \leq 0 \text{ or } j + k = 1, k \leq 0\}. \quad (17)$$

It clearly holds that $\partial\mathbf{D}_{A,l} = \partial\mathbf{D}_{A^{l1}} \cup \partial\mathbf{D}_{A^{l2}}$ and $\partial\mathbf{D}_{B,l} = \partial\mathbf{D}_{B^{l1}} \cup \partial\mathbf{D}_{B^{l2}}$ for $l \in \{1, 2\}$. These boundary regions turn out to be important as they allow us to derive criteria for the contagion process to stop. Heuristically, bank holdings with capital on the boundary have very vulnerable subsidiaries and if there are too many of them, the contagion process can quickly regain momentum even after it had slowed down.

Our first result, Theorem 2.3, describes the size of the final fraction of defaulted type 1 and type 2 subsidiaries. Its proof is based on a sequential reformulation of the process. This formulation leads to the same final outcome, but instead of exploring the effect of all defaulted subsidiaries at once, in each step only the effect of one defaulted subsidiary is considered. It turns out that in this reformulated process, the entire state of the system can be described in the limit ($n \rightarrow \infty$) by a system of continuous functions. In the following we introduce these functions and provide some heuristic explanation for the results whose rigorous proof is given in Section 6.

Let $\psi_i(\lambda) := e^{-\lambda} \frac{\lambda^i}{i!}$ for $\lambda \geq 0$ be the probability that a Poisson distributed random variable with parameter λ takes value i . Let further $\Psi_k(\lambda) = \sum_{j=k}^{\infty} e^{-\lambda} \frac{\lambda^j}{j!}$ for $\lambda \geq 0$, the probability that a Poisson distributed random variable with parameter λ is greater or equal k . We also define $\psi_{i,j}(\lambda_1, \lambda_2) := e^{-\lambda_1} \frac{\lambda_1^i}{i!} e^{-\lambda_2} \frac{\lambda_2^j}{j!}$ for $\lambda_1, \lambda_2 \geq 0$ the probability that two independent Poisson distributed random variables X and Y with parameter λ_1 and λ_2 respectively, take the values i and j , i.e. $\mathbb{P}(X = i, Y = j) = \psi_{i,j}(\lambda_1, \lambda_2)$.

Now define the functions $a_{j,k}(z_1, z_2)$ for $(j, k) \in \mathbf{D}$ by

$$a_{j,k}(z_1, z_2) = \mathbb{E}[\mathbf{1}_{\{(C_1, C_2) \geq (j, k)\}} \psi_{C_1-j, C_2-k}(p_1 z_1, p_2 z_2)], \quad (18)$$

where the inequality $(C_1, C_2) \geq (j, k)$ is meant component-wise. In the continuous approximation mentioned above, the two arguments z_1 and z_2 describe the number of defaulted type 1 respectively type 2 subsidiaries whose effect on the system has already been taken into account, divided by n . For a holding with original subsidiary capitals $(C_1, C_2) \geq (j, k)$, the probability that due to links from these defaulted institutions, the capital has been reduced to (j, k) , is approximately given by $\psi_{C_1-j, C_2-k}(p_1 z_1, p_2 z_2)$ and thus heuristically $a_{j,k}(z_1, z_2)$ describes the fraction of holdings that have the capital (j, k) , after a fraction z_1 (respectively z_2) of type 1 (respectively type 2) subsidiaries have defaulted.

With the functions $\{a_{j,k}\}_{(j,k) \in \mathbf{D}}$ defined, we may sum over all those (j, k) that are in the default region of subsidiary l in order to get the total fraction of defaulted type l subsidiaries $\sum_{(j,k) \in \mathbf{D}_{J,l}} a_{j,k}(z_1, z_2)$. The effect of the fraction z_l on the system has already been considered and it remains to explore the impact of the rest, i.e.

$$f_l(z_1, z_2) = \sum_{(j,k) \in \mathbf{D}_{J,l}} a_{j,k}(z_1, z_2) - z_l \quad (19)$$

for $J \in \{A, B\}, l \in \{1, 2\}$. Recall that to simplify the notation the dependency on J of $f_l(z_1, z_2)$ is implicit.

A calculation based on Figure 2 concludes that

$$f_1^A(z_1, z_2) = -z_1 + \mathbb{P}((C_1, C_2) \in \mathbf{D}_{A,1}) + \mathbb{E}[\mathbf{1}_{\{(C_1, C_2) \in \mathbf{D}_{A,1}^c\}} \Psi_{C_1-x}(p_1 z_1)] \\ + \mathbb{E} \left[\sum_{j=(C_1 \vee -x)}^{-C_2 \wedge x+1} \mathbf{1}_{\{(C_1, C_2) \in \mathbf{D}_{A,1}^c\}} \psi_{C_1-j}(p_1 z_1) \Psi_{C_2+j}(p_2 z_2) \right],$$

and

$$f_1^B(z_1, z_2) = -z_1 + \mathbb{P}((C_1, C_2) \in \mathbf{D}_{B,1}) + \mathbb{E}[\mathbf{1}_{\{(C_1, C_2) \in \mathbf{D}_{B,1}^c\}} \Psi_{C_2-x}(p_2 z_2)] \\ + \mathbb{E} \left[\sum_{j=(C_1 \vee 0)}^{-C_2 \wedge x+1} \mathbf{1}_{\{(C_1, C_2) \in \mathbf{D}_{A,1}^c\}} \psi_{C_1-j}(p_1 z_1) \Psi_{C_2+j}(p_2 z_2) \right].$$

The corresponding expressions f_2^A and f_2^B for subsidiary 2 can be derived similarly.

It will turn out that the first joint zero of the functions f_1 and f_2 allows us to determine the final number of defaulted type 1 and type 2 subsidiaries. Note that for $x = 0$ we get that $f_1^A(z_1, z_2) = f_1^B(z_1, z_2) = \mathbb{P}((C_1, C_2) \in \mathbf{D}_1) - z_1 + \mathbb{E}[\mathbf{1}_{\{(C_1, C_2) \in \mathbf{D}_1^c\}} \Psi_{C_1}(p_1 z_1)]$ and similarly for $f_2^A(z_1, z_2) = f_2^B(z_1, z_2)$ and thus the systems decouple.

Let us also calculate the partial derivatives of f_l as they appear in the statement of the following theorem. With $\mathbf{z} = (z_1, z_2)$, for $J \in \{A, B\}, l, m \in \{1, 2\}$ one obtains that

$$\frac{\partial f_l}{\partial z_m}(\mathbf{z}) = -\delta_{l,m} + p_m \sum_{(j,k) \in \partial \mathbf{D}_{Jlm}} a_{j,k}(\mathbf{z}), \quad (20)$$

where $\delta_{i,j} = \mathbf{1}_{\{i=j\}}$. For a vector $\mathbf{v} = (v_1, v_2) \in \mathbb{R}^2$ we obtain the directional derivatives

$$D_{\mathbf{v}} f_l(\mathbf{z}) = -v_l + v_1 p_1 \sum_{(j,k) \in \partial \mathbf{D}_{Jl1}} a_{j,k}(\mathbf{z}) + v_2 p_2 \sum_{(j,k) \in \partial \mathbf{D}_{Jl2}} a_{j,k}(\mathbf{z}). \quad (21)$$

In the following proposition we collect some more properties of the function f_l that will be important for the statement and proof of our first main Theorem 2.3.

Proposition 2.2. *Let $J \in \{A, B\}$. The functions $f_l, l \in \{1, 2\}$ are continuous and:*

1. $f_l(0, 0) = \mathbb{P}((C_1, C_2) \in \mathbf{D}_{J,l})$ and thus $(0, 0)$ is a joint zero of the functions f_1, f_2 if and only if $\mathbb{P}((C_1, C_2) \in \mathbf{D}_{J,1} \cup \mathbf{D}_{J,2}) = 0$.
2. f_1 is increasing in its second argument and f_2 is increasing in its first argument.

We are now ready to state the first result that allows us to determine the final number of defaulted type 1 and type 2 subsidiaries. We provide several applications of this result in Section 4.

Theorem 2.3. *Let $J \in \{A, B\}$. Consider a sequence of financial systems satisfying Assumption 2.1 and let $\mathbb{P}((C_1, C_2) \in \mathbf{D}_{J,1} \cup \mathbf{D}_{J,2}) > 0$. Then there exists a unique smallest (component-wise) positive joint root $\hat{\mathbf{z}} = (\hat{z}_1, \hat{z}_2)$ of the functions $f_i(z_1, z_2), i \in \{1, 2\}$. Moreover, the following holds for $i \in \{1, 2\}$:*

1. For all $\epsilon > 0$, with high probability $n^{-1} \mathcal{S}_{i,2n-1} \geq \hat{z}_i - \epsilon$.

2. If in addition there exists a vector $\mathbf{v} \in \mathbb{R}_+^2$ such that $D_{\mathbf{v}}f_1(\hat{z}), D_{\mathbf{v}}f_2(\hat{z}) < 0$, then

$$n^{-1}\mathcal{S}_{l,2n-1} \xrightarrow{p} \hat{z}_i, \quad \text{as } n \rightarrow \infty.$$

Here \xrightarrow{p} denotes convergence in probability.

3 Resilience

In this section we consider how small infections can propagate through a financial network and amplify to lead to significant damage. For this we start with an a-priori uninfected network, parametrized by (C_1, C_2) with $\mathbb{P}((C_1, C_2) \in \mathbf{D}_{J,1} \cup \mathbf{D}_{J,2}) = 0$ and then apply ex-post infections. The ex-post infections will lead to $(\tilde{C}_1, \tilde{C}_2)$, a network with lower capitalized banks, and we determine the spread of contagion in this new network based on the characteristics of (C_1, C_2) . This provides a measure of resilience of the system (C_1, C_2) to shocks.

Since our results hold for both, type A and type B holdings, and the proofs are generic, we drop the holding type $J \in \{A, B\}$ in the notation, and since we will work with different financial systems we now include the random variable (C_1, C_2) in the notation of the functionals f_l defined in (19) and write $f_l(C_1, C_2; z_1, z_2)$. Before we move on let us first collect some additional properties of these functions which we need in what follows.

Proposition 3.1. *Let (C_1, C_2) a configuration for the financial system and let $f_l(C_1, C_2; \cdot), l \in \{1, 2\}$ its functional. Let further $\Lambda_{n,h}^l(z_1 p_1, z_2 p_2) := \mathbb{P}((n, h) - (X_1(p_1 z_1), X_2(p_2 z_2)) \in \mathbf{D}_{J,l})$ with X_1 and X_2 mutually independent Poisson distributed random variables with parameter $p_1 z_1$ and $p_2 z_2$ respectively. Then,*

$$f_l(C_1, C_2; z_1, z_2) = \mathbb{E}[\Lambda_{C_1, C_2}^l(z_1 p_1, z_2 p_2)] - z_l. \quad (22)$$

Moreover, by the law of total probability

$$f_l(C_1, C_2; z_1, z_2) = \mathbb{P}((C_1, C_2) - (X_1(p_1 z_1), X_2(p_2 z_2)) \in \mathbf{D}_{J,l}) - z_l, \quad (23)$$

with $X_1(p_1 z_1)$ and $X_2(p_2 z_2)$ mutually independent and independent of (C_1, C_2) Poisson distributed random variables with parameter $p_1 z_1$ and $p_2 z_2$.

With the help of the previous proposition it becomes clear from the representation of f_l that a larger default region leads to a larger number of defaulted subsidiaries. In particular in the case of type B subsidiaries, a decrease of x leads to fewer infections. Of course this observation can also be made directly from the specification of the process in Section 2, but in order to derive our results in this section the representations provided in Proposition 3.1 will turn out useful.

As a direct consequence of the last proposition we obtain the following two corollaries:

Corollary 3.2. *Let $\{(C_1^n, C_2^n)\}$ be a sequence of random variables on \mathbf{D} that converges weakly to (C_1, C_2) . Then $f_l(C_1^n, C_2^n; \cdot, \cdot)$ and $D_{\mathbf{v}}f_l(C_1^n, C_2^n; \mathbf{z})$ converge to $f_l(C_1, C_2; \cdot, \cdot)$ and $D_{\mathbf{v}}f_l(C_1, C_2; \mathbf{z})$ for all $\mathbf{v} \in \mathbb{R}^2$, uniformly on compacts.*

Corollary 3.3. *Let (C_1, C_2) and $(\tilde{C}_1, \tilde{C}_2)$ be configurations for the financial system and let $f_l(C_1, C_2; \cdot)$ respective $f_l(\tilde{C}_1, \tilde{C}_2; z_1, z_2), l \in \{1, 2\}$ be their functional. Then:*

1. *If $\mathbb{P}((\tilde{C}_1, \tilde{C}_2) \leq (j, k)) \geq \mathbb{P}((C_1, C_2) \leq (j, k))$ for all $(j, k) \in \mathbf{D}$, that is (C_1, C_2) stochastically dominates $(\tilde{C}_1, \tilde{C}_2)$, then*

$$f_l(C_1, C_2; z_1, z_2) \leq f_l(\tilde{C}_1, \tilde{C}_2; z_1, z_2)$$

for all $z_1, z_2 \geq 0$.

2. Moreover, denote by $\mathbf{B}_{J,l}$ the set of entrance points to the default region $\mathbf{D}_{J,l}$, that is

$$\mathbf{B}_{A,1} = \{(j, k) \in \mathbf{D}_A | j = x, k \geq -x \text{ or } j + k = 0, x \leq k \leq -x \text{ or } j = -x, k \leq x\}, \quad (24)$$

$$\mathbf{B}_{A,2} = \{(j, k) \in \mathbf{D}_A | k = x, j \geq -x \text{ or } j + k = 0, x \leq j \leq -x \text{ or } j = -x, j \leq x\}, \quad (25)$$

$$\mathbf{B}_{B,1} = \{(j, k) \in \mathbf{D}_B | j = x, k \geq -x \text{ or } j + k = 0, 0 \leq k \leq -x \text{ or } j = 0, k \leq 0\}, \quad (26)$$

$$\mathbf{B}_{B,2} = \{(j, k) \in \mathbf{D}_B | k = x, k \geq -x \text{ or } j + k = 0, 0 \leq j \leq -x \text{ or } j = 0, j \leq 0\}. \quad (27)$$

If in addition there exists $(j, k) \in \mathbf{D}_{J,l}^c \cup \mathbf{B}_{J,l}$ such that $\mathbb{P}((\tilde{C}_1, \tilde{C}_2) \leq (j, k)) > \mathbb{P}((C_1, C_2) \leq (j, k))$, then

$$f_l(C_1, C_2; z_1, z_2) < f_l(\tilde{C}_1, \tilde{C}_2; z_1, z_2)$$

for all $z_1, z_2 \geq 0$.

We now move on to the statement of our main result in this section Theorem 3.4 and first specify what is exactly meant by a shock to the system.

Specification of financial shock: Following previous literature on contagion in financial networks Amini et al. (2016a); Detering et al. (2019, 2021, 2022) we now consider a network that has a-priori no defaults, meaning $\mathbb{P}((C_1, C_2) \in \mathbf{D}_{J,1} \cup \mathbf{D}_{J,2}) = 0$. We then apply a small shock to the system. This shock has the effect that the capital of some subsidiaries (and therefore their holdings) is reduced and we assume this reduction to be such that some subsidiaries actually default.

For a shock that affects a fraction ε this leads to a new financial system $(\tilde{C}_1, \tilde{C}_2)$ which is such that:

1. $\mathbb{P}((\tilde{C}_1, \tilde{C}_2) \leq (C_1, C_2)) = 1$ (no one is better off after the shock),
2. $\mathbb{P}((\tilde{C}_1, \tilde{C}_2) \neq (C_1, C_2)) = \varepsilon$ (an ε fraction of holdings is affected by the shock),
3. $\mathbb{P}(\tilde{C}_1, \tilde{C}_2) \in \mathbf{D}_{J,1} \cup \mathbf{D}_{J,2}) > 0$ (some defaults occur due to the shock).

Resilience: We now call a financial system described by (C_1, C_2) *resilient* if for every given $\tilde{\varepsilon} > 0$, there exists ε such that for all shocked systems $(\tilde{C}_1, \tilde{C}_2)$ of (C_1, C_2) with $\mathbb{P}((\tilde{C}_1, \tilde{C}_2) \neq (C_1, C_2)) \leq \varepsilon$, the number of defaulted subsidiaries $n^{-1}(\mathcal{S}_{1,2n-1} + \mathcal{S}_{2,2n-1})$ at the end of the default processes is less than $\tilde{\varepsilon}$ w.h.p..

If on the contrary, there exists a lower bound Δ , such that for every shocked system $(\tilde{C}_1, \tilde{C}_2)$ with $\mathbb{P}((\tilde{C}_1, \tilde{C}_2) \in \mathbf{D}_{J,1} \cup \mathbf{D}_{J,2}) > 0$ and $\mathbb{P}((\tilde{C}_1, \tilde{C}_2) \leq (C_1, C_2)) = 1$, it holds that $n^{-1}(\mathcal{S}_{1,2n-1} + \mathcal{S}_{2,2n-1}) \geq \Delta$ w.h.p., then we call the system *non-resilient*.

The following theorem gives criteria for resilience and non-resilience based on the derivative in $\mathbf{0}$ of the functional of the uninfected network, parametrized by (C_1, C_2) .

Theorem 3.4. *We have the following statement regarding resilience:*

1. If there exists a vector $\mathbf{v} \in \mathbb{R}_+^2$ such that $D_{\mathbf{v}}f_1(C_1, C_2; \mathbf{0}), D_{\mathbf{v}}f_2(C_1, C_2; \mathbf{0}) < 0$, then the network is resilient.
2. If there exists a vector $\mathbf{v} \in \mathbb{R}_+^2$ such that $D_{\mathbf{v}}f_1(C_1, C_2; \mathbf{0}), D_{\mathbf{v}}f_2(C_1, C_2; \mathbf{0}) > 0$, then the network is non-resilient.

In particular the conditions 1. and 2. are mutually exclusive.

Let us provide some rough sketch of the proof of Theorem 3.4, which is fully worked out in the Appendix: We know by Proposition 2.2 that for the network without defaults it holds that $f_1(C_1, C_2; \mathbf{0}) = f_2(C_1, C_2; \mathbf{0}) = 0$. If condition 1. holds and there exists a vector \mathbf{v} such that the directional derivative of both functions $f_1(C_1, C_2, \cdot)$ and $f_2(C_1, C_2, \cdot)$ is negative, then this implies that in the direction \mathbf{v} both functions become negative. As long as the shock is small, i.e. $\mathbb{P}((\tilde{C}_1, \tilde{C}_2) \neq (C_1, C_2)) = \varepsilon$ small, we obtain by Corollary 3.2 that $f_l(C_1, C_2; z_1, z_2)$ is close to $f_l(\tilde{C}_1, \tilde{C}_2; z_1, z_2)$ and $D_{\mathbf{v}}f_l(C_1, C_2; z_1, z_2)$ is close to $D_{\mathbf{v}}f_l(\tilde{C}_1, \tilde{C}_2; z_1, z_2)$ for $l \in \{1, 2\}$ and (z_1, z_2) in some compact set. Despite $\max\{f_1(\tilde{C}_1, \tilde{C}_2; \mathbf{0}), f_2(\tilde{C}_1, \tilde{C}_2; \mathbf{0})\} > 0$ this will imply that for small ε there exists a small t_0 such that $f_l(C_1, C_2; t_0\mathbf{v}) \leq \mathbf{0}$ for $l \in \{1, 2\}$. The following Lemma 3.5 then shows that $t_0\mathbf{v}$ serves as an upper bound for the first joint zero of $f_1(\tilde{C}_1, \tilde{C}_2; \cdot)$ and $f_2(\tilde{C}_1, \tilde{C}_2; \cdot)$ which then leads to the conclusion of resilience.

Lemma 3.5. *Let $\tilde{\mathbf{z}} = (\tilde{z}_1, \tilde{z}_2)$ be such that $(f_1(\tilde{\mathbf{z}}), f_2(\tilde{\mathbf{z}})) \leq \mathbf{0}$, then it holds that $\hat{\mathbf{z}} \leq \tilde{\mathbf{z}}$.*

In order to show non-resilience we first observe that condition 2. implies that in the direction \mathbf{v} the functions $f_1(C_1, C_2; \cdot)$ and $f_2(C_1, C_2; \cdot)$ are both becoming positive. By continuity of the derivative this in particular implies that $\min\{f_1(C_1, C_2; t\mathbf{v}), f_2(C_1, C_2; t\mathbf{v})\} > 0$ for all $t \in (0, t_0]$ for some $t_0 > 0$. We can then use Corollary 3.3 to show that also $\min\{f_1(\tilde{C}_1, \tilde{C}_2; t\mathbf{v}), f_2(\tilde{C}_1, \tilde{C}_2; t\mathbf{v})\} > 0$ for all $t \in (0, t_0]$. Then one can use the properties of f derived in Proposition 2.2 to conclude that the smallest fixed-point of $f_1(\tilde{C}_1, \tilde{C}_2; \cdot)$ and $f_2(\tilde{C}_1, \tilde{C}_2; \cdot)$ is bounded by $t_0\mathbf{v}$. Because the choice of t_0 only depends on the specification of (C_1, C_2) and not on the precise specification of $(\tilde{C}_1, \tilde{C}_2)$, this allows us to conclude non-resilience.

In the following we provide some examples that demonstrate the effect that the holding support can have on the resilience/non-resilience of the network.

Example 3.6. *Consider a financial network with $\mathbb{P}((C_1, C_2) = (1, 1)) = 1$ and $p_i > 1$ for $i \in \{1, 2\}$. We assume that the holdings are of type A. We first consider the decoupled system $x = 0$. For this we obtain*

$$\begin{aligned} D_{\mathbf{v}}f_1(\mathbf{z}) &= -v_1 + v_1p_1 \sum_{k \leq 1} a_{1,k}(\mathbf{z}) = -v_1 + v_1p_1 \sum_{k \geq 0} \psi_{0,k}(p_1z_1, p_2z_2) = -v_1 + v_1p_1e^{-z_1p_1} \\ D_{\mathbf{v}}f_2(\mathbf{z}) &= -v_2 + v_2p_2 \sum_{k \leq 1} a_{k,1}(\mathbf{z}) = -v_2 + v_2p_2 \sum_{k \geq 0} \psi_{k,0}(p_1z_1, p_2z_2) = -v_2 + v_2p_2e^{-z_2p_2} \end{aligned}$$

and therefore $D_{\mathbf{v}}f_l(\mathbf{0}) = -v_l + v_l p_l$ for $l \in \{1, 2\}$, which is positive for any $(v_1, v_2) > \mathbf{0}$ and the system is therefore non-resilient. This is as expected as for $x = 0$ because the systems are decoupled and the propagation of distress in one system is not influenced by the other system. It is well known then (see for example Hofstad (2016)) that each system $l \in \{1, 2\}$ separately forms a giant component for $p_l > 1$. We have $C_l = 1$, the capital of each subsidiary equals 1 and it defaults as soon as one of their debtors default. This implies that as soon as one subsidiary in the giant component defaults, then the entire giant component defaults.

We now look at the coupled case $x \leq -1$ and note that because of the capital structure of the uninfected network, the area of the default region that can actually be reached does not depend on x for $x \leq -1$. It is therefore sufficient to consider $x = -1$. We observe that

$$D_{\mathbf{v}}f_l(\mathbf{z}) = -v_l + v_l p_1(a_{0,1}(\mathbf{z}) + a_{1,0}(\mathbf{z})) + v_l p_2(a_{0,1}(\mathbf{z}) + a_{1,0}(\mathbf{z}))$$

with

$$a_{0,1}(z_1, z_2) = \psi_{1,0}(p_1 z_1, p_2 z_2) = e^{-p_1 z_1} \frac{(p_1 z_1)^1}{1!} e^{-p_2 z_2} \quad (28)$$

$$a_{1,0}(z_1, z_2) = \psi_{0,1}(p_1 z_1, p_2 z_2) = e^{-p_1 z_1} e^{-p_2 z_2} \frac{(p_2 z_2)^1}{1!} \quad (29)$$

and thus $a_{0,1}(\mathbf{0}) = a_{1,0}(\mathbf{0}) = 0$. It follows that $D_{\mathbf{v}} f_l(\mathbf{0}) < (0, 0)$ for any $\mathbf{v} = (v_1, v_2) > (0, 0)$. The system is therefore resilient. The fact that the system is resilient for $x = -1$ might be surprising at first for the following reason. If a subsidiary defaults, then the default of the holding's other subsidiary is triggered immediately as well. This is because when the default happens the holdings capital is 0. From this perspective the coupling can lead to many additional defaults as a result of a defaulted subsidiary. On the other hand however, it allows to contain second order effects as healthy holdings with both subsidiaries still having capital equal to 1 require a total of 2 links to defaulted subsidiaries in order to trigger the default of one and then automatically both subsidiaries. This example shows also that it can be beneficial from a regulator perspective if entities form little support groups. Although the distress of one member of the group can then pull down all the others in that group, the formation of support groups can in some instances still lead to a more resilient system.

Example 3.7. We consider now a network with $\mathbb{P}((C_1, C_2) = (1, 0)) = 1$, which means that for all banks their type 1 subsidiary would be in default if it was not supported by the holding through the other subsidiary. We assume that the holdings are of type A. Again, we only need to consider the case $x = -1$. Then

$$D_{\mathbf{v}} f_l(\mathbf{z}) = -v_l + v_1 p_1 a_{0,1}(\mathbf{z}) + v_2 p_2 a_{0,1}(\mathbf{z})$$

with

$$a_{0,1}(z_1, z_2) = \psi_{0,0}(p_1 z_1, p_2 z_2) = e^{-p_1 z_1} e^{-p_2 z_2}.$$

Then, we obtain that

$$D_{(1/2, 1/2)} f_l(\mathbf{0}) = 1/2(p_1 - 1) + 1/2p_2$$

which is (> 0) for $p_1 + p_2 > 1$ and (< 0) for $p_1 + p_2 < 1$. It follows that the system is resilient for $p_1 + p_2 < 1$ and non-resilient for $p_1 + p_2 > 1$. Note that in this example it is clear from the discussion above about the giant component that for $p_1, p_2 > 1$ the system must be non-resilient because one link from a defaulted subsidiary leads automatically to the default of both subsidiaries of the same holding. Because for $p_1, p_2 > 1$ each subsidiary network has its own giant component, it is clear that this network is more prone to large cascades than two separated networks with the same connection probability. For $2 > p_1 + p_2 > 1$ and $p_1 < 1$ the coupling then has actually a negative effect for the type 1 subsidiary network. The effect of connecting to the under-capitalized type 2 network is the same as adding additional second edges between the type 1 subsidiaries with probability p_2 . Subsidiary 1 inherits the exposures of subsidiary 2. Due to these additional edges, the network gets non-resilient. It is not surprising that the additional edges have the same effect as increasing the connection probability, because for two nodes i and j , the probability of having neither an edge between the type 1 subsidiaries nor between the type 2 subsidiaries is given by $1 - (p_1 + p_2)/n + (p_1 p_2)/n^2$ where the term $(p_1 p_2)/n^2$ becomes negligible for large n .

4 Case Study

We now study the contagion process for a number of different system configurations and illustrate a few intuitive findings. For different system configurations we determine the optimal support level. We wish to highlight that the optimal support level is viewed from the point of a (benevolent) regulator, who wishes to stabilize the financial system (by some measure), by imposing rules on the financial entities to support their own subsidiaries. These support actions may not be optimal from the point of view of an individual agent, and may not be a Nash equilibrium.

We assume that all subsidiaries of type 1 have capital r_1 except for an ε_1 share which has capital x and thus defaults. Similarly, all type 2 subsidiaries have capital equal to r_2 except for an ε_2 share that has capital x and thus defaults. Assuming that the defaulted subsidiaries of type 1 are chosen independently of those of type 2, this of course implies that (C_1, C_2) is such that $\mathbb{P}(C_l = x) = \varepsilon_l = 1 - \mathbb{P}(C_l = r_l)$ for $l \in \{1, 2\}$ and C_1 and C_2 independent. Then we obtain that

$$\begin{aligned} f_1^A(z_1, z_2) &= \varepsilon_1 - z_1 + (1 - \varepsilon_1)\Psi_{r_1-x}(p_1 z_1) \\ &+ (1 - \varepsilon_1) \left((1 - \varepsilon_2) \sum_{j=x+1}^{-x-1} \psi_{r_1+j}(p_1 z_1) \Psi_{\max\{r_2-j, 0\}}(p_2 z_2) \right. \\ &\left. + \varepsilon_2 \sum_{j=x}^{-x-1} \psi_{r_1+j}(p_1 z_1) \right), \end{aligned}$$

and

$$\begin{aligned} f_1^B(z_1, z_2) &= \varepsilon_1 - z_1 + (1 - \varepsilon_1)\Psi_{r_1-x}(p_1 z_1) \\ &+ (1 - \varepsilon_1) \left((1 - \varepsilon_2) \sum_{j=1}^{-x-1} \psi_{r_1+j}(p_1 z_1) \Psi_{\max\{r_2-j, 0\}}(p_2 z_2) \right. \\ &\left. + \varepsilon_2 \sum_{j=0}^{-x-1} \psi_{r_1+j}(p_1 z_1) \right). \end{aligned}$$

The functions f_2^A and f_2^B are simply derived by exchanging the role of p_1, z_1, r_1 and p_2, z_2, r_2 . Observe that for $-x \geq r_1$ respectively $-x \geq r_2$ our specification implies that the default of type 2, respectively type 1, subsidiaries triggers immediately the default of the type 1, respectively type 2, subsidiary of the same holding.

In our first example we choose the parameters $r_1 = 5$ and $r_2 = 9$ and $p_1 = p_2 = 14$, so that both layers in the networks are equally connected. We assume that in both subnetworks 15% of the subsidiaries are defaulted, i.e. $\varepsilon_1 = \varepsilon_2 = 0.15$. We consider a holding type A. The only difference between the type 1 network and the type 2 network is then the capitalization. The type 2 network is better capitalized. Also note that for $x \leq -5$, each default of a type 2 subsidiary immediately triggers the default of its corresponding type 1 subsidiary, which implies that in this case the actual fraction of initially defaulted type 1 subsidiaries is equal to $0.15 + 0.85 \cdot 0.15 = 0.2775$. The same holds for the type 2 subsidiaries if $-x \leq 9$. For $-x$ ranging from 0 to 9 we determine the first joint zero (\hat{z}_1, \hat{z}_2) of the functions f_1^A and f_2^A which gives us the final number of defaulted type 1 and type 2 subsidiaries according to Theorem 2.3. The results are displayed in Figure 3. This is a typical example where the number of defaults




Figure 3: Graph with defaults of: subsidiary of type 1 (orange squares), subsidiary of type 2 (green rhombuses), total defaults (blue circles). The network parameters are: Holding A , $r_1 = 5$, $r_2 = 9$, $p_1 = 14$, $p_2 = 14$, $\epsilon_1 = \epsilon_2 = 15\%$.

in both subsidiaries first decreases, reaches a minimum and then increases as $(-x)$ increases. While at first, this may seem counterintuitive, as more support, should mean less defaults, we remind that if a holding supports its distressed subsidiary (in case of type A) then the holdings other subsidiary takes a hit of up to $-x$. With this in mind it is not surprising that large $-x$ may cause the second subsidiary to default.

For $x = 0$ the systems are decoupled. We observe that in this case, as a result of strong contagion, almost all type 1 subsidiaries default. This is due to the relatively low capital in this network and the relatively strong connectivity. The contagion is very much driven by higher order effects. This can be seen by calculating the probability that a subsidiary that is not initially defaulted defaults as a result of edges to initially defaulted subsidiaries. This probability is equal to the probability that a Poisson distributed random variable with parameter $14 \cdot 0.15$ is at least 5, and calculates as $\approx 6.21\%$ which thus accounts only for a very low share of the final defaults. Subsidiary network 2 is different. Due to the larger capital $r_2 = 9$ almost no contagion takes place. This is not surprising because the probability that an initially not defaulted subsidiary defaults as the result of edges from initially defaulted subsidiaries is very low. It is equal to the probability that a Poisson distributed random variable with parameter $14 \cdot 0.15$ is larger than 9. This probability is less than 0.04%.

With support for the subsidiaries $x < 0$, the contagion in the type 1 network can be significantly contained. In fact we see that with a support level of $x = -3$, the fraction of defaulted type 1 subsidiaries is reduced to $\approx 33.5\%$ without significantly increasing the number of defaulted type 2 subsidiaries. As a result also the total number of defaulted subsidiaries (type 1 together with type 2) is reduced. In fact the lowest fraction of all the defaults is obtained for $x = -5$. With larger support, however, the subsidiaries in the type 2 network get weakened substantially up to the point where more of them default. These defaults then channel back into the type 1 network and so forth. For $x = -8$ this leads to almost all subsidiaries defaulting.

In comparison we now consider a network in which we change the connection probability p_2 to 4 while keeping all other parameters unchanged. The results are shown in Figure 4. Due to the lower overall connectivity in the type 2 network, the weakening of the type 2 subsidiaries, as a result of their support for the type 1 counterparts, has much less effect because defaults do not amplify. In particular we do not see the feedback effects in between the two layers that we saw in the previous example. While for strong support ($x = -8$ and $x = -9$), the fraction of defaulted type 2 subsidiaries has increased from 0.150 to 0.209 and 0.278, this increase does not

trigger new default rounds in the type 1 network. Observe that for both examples for $x = -9$, the number of defaulted type 1 and type 2 subsidiaries equal. In fact, because a subsidiary supports the holdings other subsidiary up to the point where it defaults itself, for $x = -9$, the subsidiaries always default together.




Figure 4: Graph with defaults of: subsidiary of type 1 (orange squares), subsidiary of type 2 (green rhombuses), total defaults (blue circles). The network parameters are: Holding A, $r_1 = 5$, $r_2 = 9$, $p_1 = 14$, $p_2 = 4$, $\epsilon_1 = \epsilon_2 = 15\%$.

Clearly a decrease in the initial shock ϵ_1 decreases the number of subsidiaries that default (Figure 5). It also should be noted that at a low level of support, and with a low degree of connectivity there is very little influence on the defaults in the more capitalized subnetwork as the holding support increases.




Figure 5: Graph with defaults of: subsidiary of type 1 (orange squares), subsidiary of type 2 (green rhombuses), total defaults (blue circles). The network parameters are: Holding A, $r_1 = 5$, $r_2 = 9$, $p_1 = 14$, $p_2 = 14$, $\epsilon_1 = 10\%$, $\epsilon_2 = 15\%$.

We also illustrate that this holding structure can be a stabilizing force ensuring resilience in the sense of Section 3. We consider the case from Example 3.7, with a holding with two (identical) subsidiaries $C_1 = 1 = C_2$, $p_1 = 1.5 = p_2$, and with a small initial shock: $\epsilon_1 = 1\% = \epsilon_2$. Without the holding with the two networks of subsidiaries separated, there is a cluster of defaults (approximately 0.5936), as a result of the propagation of the initial (small) shock. Whereas, once the holding is added, and the support is non-zero ($x = -1$), the defaults decrease to 0.01051.

As a last example we consider again our original setting with $r_1 = 5$, $r_2 = 9$, $p_1 = p_2 = 14$ and $\epsilon_1 = \epsilon_2 = 15\%$, but now with a holding structure B . For this holding structure the support for one subsidiary has no negative effect on the other subsidiary of the same holding. As we mentioned previously, the support can be thought of as a loan that is taken against the positive capital of the holdings solvent subsidiary. The external lender is thus faced with the risk of both subsidiaries defaulting at the end. This external lender is not part of the network. As expected, we therefore see a significant reduction of the number of defaulted subsidiaries with stronger holding support and the maximal holding support $x = -9$ leads to the fewest defaults. This is illustrated in Figure 6.




Figure 6: Graph with defaults of: subsidiary of type 1 (orange squares), subsidiary of type 2 (green rhombuses), total defaults (blue circles). The network parameters are: Holding B , $r_1 = 5$, $r_2 = 9$, $p_1 = 14$, $p_2 = 14$, $\epsilon_1 = \epsilon_2 = 15\%$.

5 Conclusion

We proposed a simple model for a financial network of bank holdings with subsidiaries. This model allows us to study the role that the holding structure and the holding's support for troubled subsidiaries has on the propagation of the contagion in a financial system. We observe that when (financial) firms are grouped into holdings this can have both, positive and negative effects. The support of a holding for a subsidiary in distress can mitigate or defer defaults and thus increase financial stability. However, in low capitalized systems, it can also amplify contagion as distress can reach previously healthy parts of the system. We show this phenomena in two ways: with a numerical case study that uses our theoretical results about the final default fraction, and analytically, based on a notion of resilience which allows us to investigate the spread of small initial shocks. We also determine the optimal support level from a regulator perspective. We stress that this is a simple model, which allows for many interesting generalizations, including more subsidiaries, more heterogeneous degrees, or holding that have additional capital.

6 Proofs

All the proofs in this section are generic for $J \in \{A, B\}$ therefore we continue to omit J from the notation.

Proof of Proposition 2.2. Continuity follows by continuity of the functions $\psi_k(\lambda)$ and $\Psi_k(\lambda)$ and dominated convergence.

For part 1. recall that $f_l(0, 0) = \sum_{(j,k) \in \mathbf{D}_{J,l}} a_{j,k}(0, 0)$ with $a_{j,k}(0, 0) \geq 0$. Since $\psi_{i,j}(0, 0) = 0$ if i or j is larger than 0, it follows that $a_{j,k}(0, 0) = \mathbb{E}[\mathbf{1}_{\{(C_1, C_2) = (j, k)\}}] = \mathbb{P}((C_1, C_2) = (j, k))$ and thus $f_l(0, 0) = \sum_{(j,k) \in \mathbf{D}_{J,l}} \mathbb{P}((C_1, C_2) = (j, k)) = \mathbb{P}((C_1, C_2) \in \mathbf{D}_{J,l})$.

Part 2. follows from (20), and the fact that $a_{j,k} \geq 0$. \square

Proof of Theorem 2.3. First observe that by Proposition 2.2 the function f_1 is increasing in its second argument while f_2 is increasing in its first argument. Moreover, $f_1(z_1, z_2) \leq 0$ for $z_1 \geq 1$ and $f_2(z_1, z_2) \leq 0$ for $z_2 \geq 1$ since $\sum_{(j,k) \in \mathbf{D}_{J,l}} a_{j,k}(z_1, z_2) \leq 1$. These properties allow to apply the proof of Lemma 3.2. in Detering et al. (2020) without any changes and we can conclude that a component-wise smallest joint root $\hat{z} = (\hat{z}_1, \hat{z}_2)$ exists.

As mentioned above, instead of directly considering the contagion process based on rounds as described in (1), we explore the set $\mathcal{S}_{i,2n-1}$ by sequentially quantifying the effect of a defaulted subsidiary on the rest of the system. So, in each step $t \in \mathbb{N}$ we only expose the effect of one (yet unexposed) defaulted subsidiary. Such a sequential reformulation is standard in the literature and was first used for exploration of the giant component (see Hofstad (2016) for instance) and later used for bootstrap percolation and financial contagion (see Janson et al. (2012); Amini et al. (2016a); Detering et al. (2019)). The situation here, however, is conceptually more complicated because of the complexity of the contagion process and we therefore describe it in some detail. First note that whenever a subsidiary reaches the capital x we know for sure that it has defaulted, independent of the second subsidiary's financial situation. Therefore, there is no need to reduce the capital of such a subsidiary further in case it receives additional links from defaulted subsidiaries. This implies that each holding only takes on a finite number of *relevant* states.

During the exploration process we keep track of several quantities:

- The sets $\tilde{C}_{j,k}(t)$ for $(j, k) \in \mathbf{D}$ consisting of those holdings $i \in [n]$ with capital $c_{i,1}$ of subsidiary 1 being equal to j and capital $c_{i,2}$ of subsidiary 2 being equal to k at step $t \in \mathbb{N}$. At the beginning of the process ($t = 0$) we set $\tilde{C}_{j,k}(0) := \{i \in [n] \mid (c_{i,1}, c_{i,2}) = (j, k)\}$ for all $j, k \in \mathbf{D}$.
- The sets $U_l(t), l \in \{1, 2\}$ of unexplored, defaulted type l subsidiaries. At the beginning of the process we have $U_l(0) \subset [n_l]$ defined by: $i_l \in U_l(0)$ if and only if $(c_{i,1}, c_{i,2}) \in \mathbf{D}_{J,l}$. So, $U_l(0)$ contains those subsidiaries of type $l \in \{1, 2\}$ which are in default at the beginning of the process.
- The number $n_1(t)$ and $n_2(t)$ of already explored type 1 and type 2 subsidiaries. We set $n_1(0) = n_2(0) = 0$.

We now describe one step $t \in \mathbb{N}$ of this exploration process: At the very beginning of the step we choose uniformly a defaulted subsidiary from $U_1(t) \cup U_2(t)$. We denote this subsidiary by S . We then draw the links from S to all the holdings in the sets $\tilde{C}_{j,k}(t)$. We then update these sets as follows. If in step t we selected a type $l \in \{1, 2\}$ subsidiary and the subsidiary l of holding $i \in \tilde{C}_{j,k}(t)$ receives a link from S , then i will be moved into $\tilde{C}_{j-1,k}(t+1)$ unless $j = x$ in which case it will just remain in its current set ($i \in \tilde{C}_{j,k}(t+1)$). During this exploration, some subsidiaries might move into their default region. If this happens and a type 1 or type 2 subsidiary goes bankrupt, we generate a *copy* of it and place it into the set $U_1(t)$, respectively $U_2(t)$. At the end of the step, if the selected subsidiary was in $U_1(t)$ we set $n_1(t) = n_1(t-1)+1$ and $n_2(t) = n_2(t-1)$, and we remove the subsidiary from the set of defaulted unexplored subsidiaries (i.e. we set $U_1(t+1) = U_1(t) \setminus \{S\}$). If the selected subsidiary was from $U_2(t)$, then we set $n_2(t) = n_2(t-1)+1$ and $n_1(t) = n_1(t-1)$ and remove the subsidiary ($U_2(t+1) = U_2(t) \setminus \{S\}$). Note that elements in $U_1(t)$ and $U_2(t)$ are distinguishable. The process ends at $\hat{t} = \min\{t \in \mathbb{N} \mid U_1(t) \cup U_2(t) = \emptyset\}$

and \hat{t} corresponds to the total number of defaulted subsidiaries. Moreover, $\hat{t} = n_1(\hat{t}) + n_2(\hat{t})$, because in each step we have exactly explored one subsidiary. Note that the procedure is such that the capital of some subsidiaries is still being updated although they have already defaulted (and possibly have been explored) but this updating has no effect. This happens if a subsidiary defaults with more capital than x .

We start with proving the first statement of the theorem, the lower bound for $n^{-1}\mathcal{S}_{i,2n-1}^J$ based on this sequential exploration. Let $u_1(t) = |U_1(t)|$ and $u_2(t) = |U_2(t)|$ and $\tilde{c}_{j,k}(t) = |\tilde{C}_{j,k}(t)|$ and let $h(t)$ describe the entire system, i.e.

$$h(t) = (u_1(t), u_2(t), n_1(t), n_2(t), \{\tilde{c}_{j,k}\}_{(j,k) \in \mathbf{D}}).$$

We would like to stress that the the vector $h(t) = (u_1(t), u_2(t), n_1(t), n_2(t), \{\tilde{c}_{j,k}\}_{(j,k) \in \mathbf{D}^x})$ is a random vector and that we have chosen the lower case letters in order to distinguish from the random sets $U_1(t), U_2(t)$, and the $\tilde{C}_{j,k}(t)$ for $(j, k) \in \mathbf{D}$.

With probability $u_1(t)/(u_1(t) + u_2(t))$ in step t a type 1 subsidiary is selected that leads to a reduction of the number of vertices in $U_1(t)$ by 1. On the other hand, each subsidiary in $\partial \mathbf{D}_{J11}$ receives an edge from the selected subsidiary with probability $\frac{p_1}{n}$ and thus defaults and is added to $U_1(t+1)$. Similar, if a subsidiary of type 2 is selected, which happens with probability $u_2(t)/(u_1(t) + u_2(t))$, then each subsidiary in $\partial \mathbf{D}_{J12}$ receives an edge from this subsidiary with probability $\frac{p_2}{n}$, defaults and is added to $U_1(t+1)$. These considerations lead to

$$\mathbb{E}[u_1(t+1) - u_1(t)|h(t)] \tag{30}$$

$$= \frac{u_1(t)}{u_1(t) + u_2(t)} \left(-1 + \sum_{(j,k) \in \partial \mathbf{D}_{J11}} \frac{p_1}{n} \tilde{c}_{j,k}(t) \right) + \frac{u_2(t)}{u_1(t) + u_2(t)} \left(\sum_{(j,k) \in \partial \mathbf{D}_{J12}} \frac{p_2}{n} \tilde{c}_{j,k}(t) \right) \tag{31}$$

$$= g_1(u_1(t)/n, u_2(t)/n, \{\tilde{c}_{j,k}(t)/n\}_{(j,k) \in \mathbf{D}}) \tag{32}$$

for $t \leq \hat{t}$ with

$$g_1(\nu_1, \nu_2, \{\gamma_{l,m}\}_{m,n}) := \frac{\nu_1}{\nu_1 + \nu_2} \left(-1 + \sum_{(j,k) \in \partial \mathbf{D}_{J11}} p_1 \gamma_{j,k} \right) + \frac{\nu_2}{\nu_1 + \nu_2} \left(\sum_{(j,k) \in \partial \mathbf{D}_{J12}} p_2 \gamma_{j,k} \right) \tag{33}$$

and similarly for $u_2(t)$. Moreover, a similar combinatorial argument yields that

$$\begin{aligned} \mathbb{E}[\tilde{c}_{k,j}(t+1) - \tilde{c}_{k,j}(t)|h(t)] &= \frac{u_1(t)}{u_1(t) + u_2(t)} \frac{p_1}{n} (\tilde{c}_{k+1,j}(t) \mathbf{1}_{\{k+1 \leq R\}} - \tilde{c}_{k,j}(t) \mathbf{1}_{\{k-1 \geq x\}}) \\ &+ \frac{u_2(t)}{u_1(t) + u_2(t)} \frac{p_2}{n} (\tilde{c}_{k,j+1}(t) \mathbf{1}_{\{j+1 \leq R\}} - \tilde{c}_{k,j}(t) \mathbf{1}_{\{j-1 \geq x\}}) \\ &= \frac{u_1(t)/n}{u_1(t)/n + u_2(t)/n} p_1 ((\tilde{c}_{k+1,j}(t)/n) \mathbf{1}_{\{k+1 \leq R\}} - (\tilde{c}_{k,j}(t)/n) \mathbf{1}_{\{k-1 \geq x\}}) \\ &+ \frac{u_2(t)/n}{u_1(t)/n + u_2(t)/n} p_2 ((\tilde{c}_{k,j+1}(t)/n) \mathbf{1}_{\{j+1 \leq R\}} - (\tilde{c}_{k,j}(t)/n) \mathbf{1}_{\{j-1 \geq x\}}) \\ &= g_{k,j}(u_1(t)/n, u_2(t)/n, \{\tilde{c}_{l,m}(t)/n\}_{m,n}) \end{aligned}$$

with

$$g_{k,j}(\nu_1, \nu_2, \{\gamma_{l,m}\}_{m,n}) := \frac{\nu_1}{\nu_1 + \nu_2} p_1 (\gamma_{k+1,j} \mathbf{1}_{\{k+1 \leq R\}} - \gamma_{k,j} \mathbf{1}_{\{k-1 \geq x\}}) \\ + \frac{\nu_2}{\nu_1 + \nu_2} p_2 (\gamma_{k,j+1} \mathbf{1}_{\{j+1 \leq R\}} - \gamma_{k,j} \mathbf{1}_{\{j-1 \geq x\}})$$

for $(j, k) \in \mathbf{D}$. And finally

$$\mathbb{E}[n_l(t+1) - n_l(t) | h(t)] = \frac{u_l(t)}{u_1(t) + u_2(t)} = q_l(u_1(t)/n, u_2(t)/n) \quad (34)$$

with $q_l(\nu_1, \nu_2) = \frac{\nu_l}{\nu_1 + \nu_2}$ for $l \in \{1, 2\}$.

While the above calculated differences are random and very much depend on the degree of the chosen subsidiary in each step, it turns out that the sum of a few successive steps becomes concentrated at the mean and can be described by a vector valued continuous function. For this consider therefore the vector valued function $(\nu_1, \nu_2, \{\gamma_{k,j}\}_{(j,k) \in \mathbf{D}}, \beta_1, \beta_2)$ defined by the system of differential equations:

$$\frac{d\nu_l(\tau)}{d\tau} = g_l(\nu_1(\tau), \nu_2(\tau), \{\gamma_{j,k}(\tau)\}_{(j,k) \in \mathbf{D}}), \quad \nu_l(0) = \mathbb{P}((C_1, C_2) \in \mathbf{D}_{J,l}), \quad l \in \{1, 2\}, \quad (35)$$

$$\frac{d\gamma_{k,j}(\tau)}{d\tau} = g_{k,j}(\nu_1(\tau), \nu_2(\tau), \{\gamma_{j,k}(\tau)\}_{(j,k) \in \mathbf{D}}), \quad \gamma_{k,j}(0) = \mathbb{P}(C_1 = k, C_2 = j), \quad (k, j) \in \mathbf{D}, \quad (36)$$

$$\frac{d\beta_l(\tau)}{d\tau} = q_l(\nu_1(\tau), \nu_2(\tau)), \quad \beta_l(0) = 0, \quad l \in \{1, 2\}. \quad (37)$$

Now note that the functions $g_1, g_2, \{g_{k,j}\}_{(j,k) \in \mathbf{D}^x}, q_1, q_2$ fulfill a Lipchitz condition on the domain

$$D_{\delta_0, \delta_1} = \left\{ (\tau, \nu_1, \nu_2, \{\gamma_{j,k}\}_{(j,k) \in \mathbf{D}}) \in \mathbb{R}^{5+|\mathbf{D}|} \mid \begin{array}{l} -\delta_0 < \tau < 2, \\ -\delta_0 < \gamma_{j,k} < 2, \\ -\delta_0 < \nu_1, \nu_2 < 2, \\ \delta_1 < \nu_1 + \nu_2 \end{array} \right\}, \quad (38)$$

for any $\delta_0, \delta_1 > 0$ as can easily be seen by calculating the partial derivatives and observing that they are bounded. This implies existence of a solution to the system (35) and (36). Since the solution exists on D_{δ_0, δ_1} for any positive δ_1 , we can of course extend the solution to $D_{\delta_0, 0}$.

The next step of the proof is to solve the system described by (35) and (36). In order to do so, define the following implicitly given functions

$$h_1(\tau) = \frac{\nu_1(\tau)}{\nu_1(\tau) + \nu_2(\tau)}, \quad h_2(\tau) = \frac{\nu_2(\tau)}{\nu_1(\tau) + \nu_2(\tau)} \quad (39)$$

$$\beta_1(\tau) = \int_0^\tau h_1(s) ds, \quad \beta_2(\tau) = \int_0^\tau h_2(s) ds, \quad (40)$$

defined on $D_{\delta_0, 0}$. Note that $h_1(\tau) + h_2(\tau) = 1$ and thus $\beta_1(\tau) + \beta_2(\tau) = \tau$. The equation for $\gamma_{R,R}(\tau)$ is a simple linear equation solved by

$$\gamma_{R,R}(\tau) = \gamma_{R,R}(0) e^{-p_1 H_1(\tau)} e^{-p_2 H_2(\tau)} = \gamma_{R,R}(0) \psi_0(p_1 H_1(\tau)) \psi_0(p_2 H_2(\tau)). \quad (41)$$

Note that $\gamma_{R,R}(\tau)$ is a constant term multiplied by the probability that two independent Poisson distributed random variables with parameters $p_1 H_1(\tau)$ and $p_2 H_2(\tau)$ are zero. This allows to guess that for $(j, k) \in \mathbf{D}$,

$$\begin{aligned} \gamma_{j,k}(\tau) &= \sum_{\substack{j \leq l_1 \leq R \\ k \leq l_2 \leq R}} \gamma_{l_1, l_2}(0) (\psi_{l_1-j}(p_1 H_1(\tau)) \mathbf{1}_{\{j > x\}} + \Psi_{l_1-j}(p_1 H_1(\tau)) \mathbf{1}_{\{j=x\}}) \cdot \\ &\quad \cdot (\psi_{l_2-k}(p_2 H_2(\tau)) \mathbf{1}_{\{k > x\}} + \Psi_{l_2-k}(p_2 H_2(\tau)) \mathbf{1}_{\{k=x\}}). \end{aligned} \quad (42)$$

To see that above expression does in fact satisfy (36), first note that

$$\frac{\partial}{\partial \lambda} \psi_k(\lambda) = -e^{-\lambda} \frac{\lambda^k}{k!} + e^{-\lambda} \frac{\lambda^{k-1}}{(k-1)!} \mathbf{1}_{\{k \geq 1\}} = \psi_{k-1}(\lambda) \mathbf{1}_{\{k \geq 1\}} - \psi_k(\lambda)$$

and

$$\frac{\partial}{\partial \lambda} \Psi_k(\lambda) = e^{-\lambda} \frac{\lambda^{k-1}}{k-1!} = \psi_{k-1}(\lambda) \mathbf{1}_{\{k \geq 1\}}.$$

Then for $j, k \notin \{x, R\}$ we get from (42) that

$$\begin{aligned} &\frac{\partial}{\partial \tau} \gamma_{j,k}(\tau) \quad (43) \\ &= \sum_{\substack{j \leq l_1 \leq R \\ k \leq l_2 \leq R}} \gamma_{l_1, l_2}(0) p_1 h_1(\tau) (\psi_{l_1-j-1}(p_1 H_1(\tau)) \mathbf{1}_{\{l_1-j \geq 1\}} - \psi_{l_1-j}(p_1 H_1(\tau))) \psi_{l_2-k}(p_2 H_2(\tau)) \\ &+ \sum_{\substack{j \leq l_1 \leq R \\ k \leq l_2 \leq R}} \gamma_{l_1, l_2}(0) p_2 h_2(\tau) \psi_{l_1-j}(p_1 H_1(\tau)) (\psi_{l_2-k-1}(p_2 H_2(\tau)) \mathbf{1}_{\{l_2-k \geq 1\}} - \psi_{l_2-k}(p_2 H_2(\tau))) \\ &= p_1 h_1(\tau) \sum_{\substack{j+1 \leq l_1 \leq R \\ k \leq l_2 \leq R}} \gamma_{l_1, l_2}(0) (\psi_{l_1-(j+1)}(p_1 H_1(\tau)) \psi_{l_2-k}(p_2 H_2(\tau)) \\ &- p_1 h_1(\tau) \sum_{\substack{j \leq l_1 \leq R \\ k \leq l_2 \leq R}} \gamma_{l_1, l_2}(0) \psi_{l_1-j}(p_1 H_1(\tau)) \psi_{l_2-k}(p_2 H_2(\tau)) \\ &+ p_2 h_2(\tau) \sum_{\substack{j \leq l_1 \leq R \\ k+1 \leq l_2 \leq R}} \gamma_{l_1, l_2}(0) (\psi_{l_1-j}(p_1 H_1(\tau)) \psi_{l_2-(k+1)}(p_2 H_2(\tau)) \\ &- p_2 h_2(\tau) \sum_{\substack{j \leq l_1 \leq R \\ k \leq l_2 \leq R}} \gamma_{l_1, l_2}(0) \psi_{l_1-j}(p_1 H_1(\tau)) \psi_{l_2-k}(p_2 H_2(\tau)) \\ &= p_1 h_1(\tau) \gamma_{j+1, k}(\tau) - p_1 h_1(\tau) \gamma_{j, k}(\tau) + p_2 h_2(\tau) \gamma_{j, k+1}(\tau) - p_2 h_2(\tau) \gamma_{j, k}(\tau), \end{aligned} \quad (44)$$

which verifies (36). The case where j or k is in $\{x, R\}$ can be verified similarly and is left to the reader.

In order to derive $\nu_1(\tau)$ and $\nu_2(\tau)$, recall that

$$\nu_l(0) = \mathbb{P}((C_1, C_2) \in \mathbf{D}_{J,l}) = \sum_{(r_1, r_2) \in \mathbf{D}_{J,l}^x} \gamma_{r_1, r_2}(0), \quad l \in \{1, 2\}.$$

We then make the following Ansatz for the solution:

$$\nu_l(\tau) = \sum_{(j,k) \in \mathbf{D}_{J,l}} \gamma_{j,k}(\tau) - H_1(\tau). \quad (45)$$

Note that the ν_l are well defined on $D_{\delta_0,0}$ as the sum is only over finitely many terms. A tedious, but straightforward calculation of taking derivatives of (45), verifies that ν_l^J satisfies (35) as most of the terms in the sum cancel out and only the terms on the boundaries $\partial \mathbf{D}_{Jml}, l, m \in \{1, 2\}$ remain.

For $(k, j) \in \mathbf{D}$ we now want to approximate the quantities $\tilde{c}_{j,k}(t)$ with $\gamma_{j,k}(t/n)$, and for $i \in \{1, 2\}$ the quantities $u_i(t)$ with $\nu_i(t/n)$ respectively. We use Wormald (1995)[Theorem 2] to show that in fact

$$u_l(t) = \nu_l(t/n)n + o(n), l \in \{1, 2\}, \quad (46)$$

$$\tilde{c}_{j,k}(t) = \gamma_{j,k}(t/n)n + o(n), (k, j) \in \mathbf{D}, \quad (47)$$

$$n_l(t) = \beta_l(t/n)n + o(n), l \in \{1, 2\}, \quad (48)$$

with high probability. By above considerations, assumptions (ii) and (iii) of Wormald (1995)[Theorem 2] are satisfied. To see that also assumption (i) is satisfied, note that in each step of the exploration, the number of vertices receiving an edge from the currently explored defaulted subsidiary is bounded by the degree of this subsidiary. Let D^+ be the largest out-degree in the graph. A simple probabilistic bound for D^+ is given by

$$\mathbb{P}(D^+ \geq \log(n)) \leq n \binom{n-1}{\log(n)} (\max\{p_1, p_2\}/n)^{\log(n)} \leq n (\max\{p_1, p_2\})^{\log(n)} / (\log(n)!).$$

By Stirling's formula, it follows that $\log(n)! \geq n^m (\max\{p_1, p_2\})^{\log(n)}$ for any m and thus

$$\mathbb{P}(D^+ \geq \log(n)) \leq n^{-(m-1)}.$$

We may choose $\lambda(n) = n^{1/10}$ and $\omega(n) = n^{5/10}$ for condition (i) in Wormald's Theorem. This implies that the approximation (46)-(48) holds uniformly for

$$(t/n, \nu_1(t/n), \nu_2(t/n), \{\gamma_{j,k}(t/n)\}_{(j,k) \in \mathbf{D}}, \beta_1(t/n), \beta_2(t/n)) \in D_{\delta_0, \delta_1}$$

with high probability.

Let now $\tau_{\delta_1} := \sup\{\tau \in \mathbb{R}_+ : (\tau, \nu_1, \nu_2, \{\gamma_{j,k}\}_{(j,k) \in \mathbf{D}}) \in D_{\delta_0, \delta_1}\}$. By the definition of the system $(u_1(t), u_2(t), \{\tilde{c}_{k,j}(t)\}_{(j,k) \in \mathbf{D}})$ it is clear that all quantities are in between 0 and $2n$ and thus by (46)-(48) and the definition of D_{δ_0, δ_1} it follows that $\nu_1(\tau_{\delta_1}) + \nu_2(\tau_{\delta_1}) = \delta_1$.

Let us now set $\hat{\tau} := \lim_{\delta_1 \rightarrow 0} \tau_{\delta_1}$, where the limit clearly exists as τ_{δ_1} is increasing in δ_1 and bounded by 2. Moreover, $\lim_{\delta_1 \rightarrow 0} \nu_1(\tau_{\delta_1}) = \lim_{\delta_1 \rightarrow 0} \nu_2(\tau_{\delta_1}) = 0$ and we shall thus define $\nu_1(\hat{\tau}) = \nu_2(\hat{\tau}) = 0$.

Note that $f_l(\beta_1(\tau), \beta_2(\tau)) = \nu_l(\tau)$ for $\tau \leq \hat{\tau}$. Moreover, since (z_1, z_2) is the smallest (component-wise) joint zero of f_1 and f_2 , and due to the monotonicity and continuity of β_1 and β_2 , and of f_l as derived in Proposition 2.2, it follows that $(\beta_1(\hat{\tau}), \beta_2(\hat{\tau})) = (\hat{z}_1, \hat{z}_2)$. Let $\hat{t} := \min\{t \mid u_1(t) + u_2(t) = 0\}$. By the previous considerations it follows that $\hat{t}/n \geq \tau_{\delta_1}$ w.h.p. for every δ_1 and thus by (46) $n^{-1} \mathcal{S}_{i, 2n-1} \geq \beta_i(\tau_{\delta_1}) - \varepsilon$ w.h.p. for any $\varepsilon > 0$ and by $\lim_{\delta_1 \rightarrow 0} \beta_i(\tau_{\delta_1}) = \hat{z}_i$ it follows that

$$n^{-1} \mathcal{S}_{i, 2n-1} \geq \hat{z}_i - \varepsilon$$

with high probability. This proves the first part of the theorem.

Now we will show that under the additional assumption that there exists $\mathbf{v} = (\mathbf{v}_1, \mathbf{v}_2) \in \mathbb{R}_+^2$ such that $D_{\mathbf{v}}f_1(\hat{z}), D_{\mathbf{v}}f_2(\hat{z}) < 0$, it holds in fact that

$$n^{-1}\mathcal{S}_{l,2n-1} \xrightarrow{p} \hat{z}_i, \quad \text{as } n \rightarrow \infty.$$

We know that $D_{\mathbf{v}}f_1(\hat{z}) = D_{\mathbf{v}}f_1(\beta_1(\hat{\tau}), \beta_2(\hat{\tau})), D_{\mathbf{v}}f_2(\hat{z}) = D_{\mathbf{v}}f_2(\beta_1(\hat{\tau}), \beta_2(\hat{\tau})) < 0$ and by continuity of the derivative and because of

$$\lim_{\tau \rightarrow \hat{\tau}} (\beta_1(\tau), \beta_2(\tau)) = (\beta_1(\hat{\tau}), \beta_2(\hat{\tau})) = (\hat{z}_1, \hat{z}_2)$$

it follows that also $D_{\mathbf{v}}f_1(\beta_1(\tau_{\delta_1}), \beta_2(\tau_{\delta_1})), D_{\mathbf{v}}f_2(\beta_1(\tau_{\delta_1}), \beta_2(\tau_{\delta_1})) < 0$ for δ_1 small enough. Since $a_{j,k}(\beta_1(\tau), \beta_2(\tau)) = \gamma_{j,k}(\tau)$ it follows by (20) that

$$v_1 p_1 \sum_{(j,k) \in \partial \mathbf{D}_{j11}} \gamma_{j,k}(\tau_{\delta_1}) + v_2 p_2 \sum_{(j,k) \in \partial \mathbf{D}_{j12}} \gamma_{j,k}(\tau_{\delta_1}) < c_1 v_1 \quad (49)$$

$$v_1 p_1 \sum_{(j,k) \in \partial \mathbf{D}_{j21}} \gamma_{j,k}(\tau_{\delta_0}) + v_2 p_2 \sum_{(j,k) \in \partial \mathbf{D}_{j22}} \gamma_{j,k}(\tau_{\delta_1}) < c_2 v_2 \quad (50)$$

for some $c_1, c_2 < 1$ sufficiently close to 1. We shall track the process until the step $\lfloor \tau_{\delta_1} n \rfloor$ by making use of (46) - (48), and then we explore the remaining subsidiaries in the sets $U_1(\lfloor \tau_{\delta_1} n \rfloor) \cup U_2(\lfloor \tau_{\delta_1} n \rfloor)$ in rounds. We place the holdings with capital $(j, k) \in \mathbf{D}$ in four sets which we define now and where a holding might be placed in more than one set. For this, let us call a subsidiary *weak* if it can default by either receiving one edge from a defaulted subsidiary itself or because the other subsidiary of the same holding receives a link from a defaulted subsidiary. We call a subsidiary *strong* if it can only default if this subsidiary and the other subsidiary of the same holding, receive in total at least two links from defaulted subsidiaries. For $l \in \{1, 2\}$ let \mathbb{W}_l be the set that includes those holdings with weak type l subsidiary and \mathbb{S}_l those with strong type l subsidiary. Note that $\mathbb{W}_l \cap \mathbb{S}_l = \emptyset$ for $l \in \{1, 2\}$ but the other four possible intersections of these sets are not necessary empty. Further for each $l \in \{1, 2\}$ we define the subsets \mathbb{W}_l^i of holdings whose weak type l subsidiary defaults in round i of the exploration process and the subsets \mathbb{S}_l^i of holdings whose strong type l subsidiary defaults in round i of the exploration process.

Note that from (49) and (50) we get together with the approximation (47) that for n large enough

$$v_1 p_1 \sum_{(j,k) \in \partial \mathbf{D}_{j11}} \frac{\tilde{c}_{j,k}(\lfloor \tau_{\delta_1} n \rfloor)}{n} + v_2 p_2 \sum_{(j,k) \in \partial \mathbf{D}_{j12}} \frac{\tilde{c}_{j,k}(\lfloor \tau_{\delta_1} n \rfloor)}{n} < c_1 v_1 \quad (51)$$

$$v_1 p_1 \sum_{(j,k) \in \partial \mathbf{D}_{j21}} \frac{\tilde{c}_{j,k}(\lfloor \tau_{\delta_1} n \rfloor)}{n} + v_2 p_2 \sum_{(j,k) \in \partial \mathbf{D}_{j22}} \frac{\tilde{c}_{j,k}(\lfloor \tau_{\delta_1} n \rfloor)}{n} < c_2 v_2 \quad (52)$$

w.h.p.. This allows us now to bound the expected number of weak subsidiaries of type $l \in \{1, 2\}$

that default in the first round through a combinatorial counting argument by

$$\begin{aligned}
& n^{-1} \mathbb{E} \left[|\mathbb{W}_l^1| \mid h(\lfloor \tau_{\delta_1} n \rfloor) \right] \\
& \leq \left(\frac{u_1(\lfloor \tau_{\delta_1} n \rfloor)}{n} p_1 \sum_{(j,k) \in \partial \mathbf{D}_{j1}} \frac{\tilde{c}_{k,j}(\lfloor \tau_{\delta_1} n \rfloor)}{n} + \frac{u_2(\lfloor \tau_{\delta_1} n \rfloor)}{n} p_2 \sum_{(j,k) \in \partial \mathbf{D}_{j2}} \frac{\tilde{c}_{k,j}(\lfloor \tau_{\delta_1} n \rfloor)}{n} \right) \\
& < c_l v_l \varepsilon
\end{aligned}$$

and in particular

$$n^{-1} \sum_{l \in \{1,2\}} \mathbb{E} \left[|\mathbb{W}_l^1| \mid h(\lfloor \tau_{\delta_1} n \rfloor) \right] < \sum_{l \in \{1,2\}} c_l v_l \varepsilon \quad (53)$$

for any ε provided that $\frac{u_1(\lfloor \tau_{\delta_1} n \rfloor)}{n} < v_1 \varepsilon$ and $\frac{u_2(\lfloor \tau_{\delta_1} n \rfloor)}{n} < v_2 \varepsilon$, which we can ensure by possibly decreasing δ_1 further. We further choose $c \in (\max\{c_1, c_2\}, 1)$ and obtain that

$$\sum_{l \in \{1,2\}} c_l v_l \varepsilon < c \varepsilon (v_1 + v_2).$$

Similarly we find for $l \in \{1,2\}$ the very rough bound

$$\begin{aligned}
& n^{-1} \mathbb{E} \left[|\mathbb{S}_l^1| \mid h(\lfloor \tau_{\delta_1} n \rfloor) \right] \\
& = \left(\frac{u_1(\lfloor \tau_{\delta_1} n \rfloor) + u_2(\lfloor \tau_{\delta_1} n \rfloor)}{n} \right)^2 \max\{p_1, p_2\} \sum_{(j,k) \in \mathbf{D}_{j,l}^c} \frac{\tilde{c}_{k,j}(\lfloor \tau_{\delta_1} n \rfloor)}{n} \\
& < ((v_1 + v_2)\varepsilon)^2 \max\{p_1, p_2\} < (c'/4)\varepsilon v_l
\end{aligned} \quad (54)$$

with $c' > 0$ such that $2c' < 1 - c$, provided that ε is chosen small enough. In particular it then follows that

$$n^{-1} \sum_{l \in \{1,2\}} \mathbb{E} [\mathbb{W}_l^1 + \mathbb{S}_l^1] < (c + c'/2)\varepsilon(v_1 + v_2). \quad (55)$$

Our plan is now to show that for $l \in \{1,2\}$ it holds for every $m > 0$ that

$$\mathbb{E} [n^{-1} \mathbb{W}_l^m] < (c_l + c'/4)(c + c')^{(m-1)} \varepsilon v_l \quad (56)$$

$$\mathbb{E} [n^{-1} \mathbb{S}_l^m] < (c'/4)(c + c')^{(m-1)} \varepsilon v_l. \quad (57)$$

This implies

$$n^{-1} \sum_{l \in \{1,2\}} \mathbb{E} [\mathbb{W}_l^m + \mathbb{S}_l^m] < (c + c')^m \varepsilon (v_1 + v_2), \quad (58)$$

and from this it follows that

$$n^{-1} \sum_{m=1}^{2n} \sum_{l \in \{1,2\}} \mathbb{E} [\mathbb{W}_l^m + \mathbb{S}_l^m] < \frac{1}{1 - (c + c')} \varepsilon (v_1 + v_2). \quad (59)$$

and therefore

$$\mathbb{P} \left(n^{-1} \sum_{m=1}^{2n} \sum_{l \in \{1,2\}} (\mathbb{W}_l^m + \mathbb{S}_l^m) \geq \sqrt{\frac{1}{1 - (c + c')} \varepsilon (v_1 + v_2)} \right) \leq \sqrt{\frac{1}{1 - (c + c')} \varepsilon (v_1 + v_2)}. \quad (60)$$

The quantity on the right hand side of (60) can be made arbitrarily small by decreasing ε . So in order to conclude, it remains to show (56) and (57).

For $m = 1$ this is done above. Suppose that (56) and (57) holds for $m > 0$. Then it follows for $l \in \{1, 2\}$ by (51) and (52) that

$$n^{-1}\mathbb{E}\left[|\mathbb{W}_l^{m+1}|\left|h(\lfloor\tau_{\delta_1}n\rfloor)\right.\right] \quad (61)$$

$$\leq \frac{\sum_{x \in \mathbb{W}_1 \cup \mathbb{S}_1} \mathbb{P}(x \in \mathbb{W}_1^m \cup \mathbb{S}_1^m)}{n} p_1 \sum_{(j,k) \in \partial \mathbf{D}_{j^{l1}}} \frac{\tilde{c}_{k,j}(\lfloor\tau_{\delta_1}n\rfloor)}{n} \quad (62)$$

$$+ \frac{\sum_{x \in \mathbb{W}_2 \cup \mathbb{S}_2} \mathbb{P}(x \in \mathbb{W}_2^m \cup \mathbb{S}_2^m)}{n} p_2 \sum_{(j,k) \in \partial \mathbf{D}_{j^{l2}}} \frac{\tilde{c}_{k,j}(\lfloor\tau_{\delta_1}n\rfloor)}{n}$$

$$\leq (\mathbb{E}[n^{-1}\mathbb{W}_1^m] + \mathbb{E}[n^{-1}\mathbb{S}_1^m]) p_1 \sum_{(j,k) \in \partial \mathbf{D}_{j^{l1}}} \frac{\tilde{c}_{k,j}(\lfloor\tau_{\delta_1}n\rfloor)}{n} \quad (63)$$

$$+ (\mathbb{E}[n^{-1}\mathbb{W}_2^m] + \mathbb{E}[n^{-1}\mathbb{S}_2^m]) p_2 \sum_{(j,k) \in \partial \mathbf{D}_{j^{l2}}} \frac{\tilde{c}_{k,j}(\lfloor\tau_{\delta_1}n\rfloor)}{n}$$

$$< (c_l + c'/2)(c + c')^m \varepsilon v_l \quad (64)$$

and in particular

$$n^{-1}\mathbb{E}\left[\sum_{l \in \{1,2\}} |\mathbb{W}_l^{m+1}| |h(\lfloor\tau_{\delta_1}n\rfloor)|\right] < (c + c'/2)(c + c')^m \varepsilon (v_1 + v_2)$$

For a subsidiary to be in \mathbb{S}_l^{m+1} it must receive a link from the set $\mathbb{W}_l^m \cup \mathbb{S}_l^m$ and one from $\bigcup_{k=1}^m \mathbb{W}_l^k \cup \mathbb{S}_l^k$. It follows from a very rough bound that

$$n^{-1}\mathbb{E}[|\mathbb{S}_l^{m+1}| |h(\lfloor\tau_{\delta_1}n\rfloor)|]$$

$$\leq \frac{\sum_{x \in \mathbb{W}_1 \cup \mathbb{S}_1 \cup \mathbb{W}_2 \cup \mathbb{S}_2} \mathbb{P}(x \in \mathbb{W}_1^m \cup \mathbb{S}_1^m \cup \mathbb{W}_2^m \cup \mathbb{S}_2^m)}{n} \max\{p_1, p_2\} \cdot$$

$$\cdot \frac{\sum_{x \in \mathbb{W}_1 \cup \mathbb{S}_1 \cup \mathbb{W}_2 \cup \mathbb{S}_2} \mathbb{P}(x \in \bigcup_{k=1}^m \mathbb{W}_1^k \cup \mathbb{S}_1^k \cup \mathbb{W}_2^k \cup \mathbb{S}_2^k)}{n} \max\{p_1, p_2\}$$

$$< (c + c'/2)^2 (c + c')^m 1/(1 - c + c') \varepsilon^2 (v_1 + v_2)^2$$

$$< c'/4 (c + c')^m \varepsilon (v_1 + v_2) \quad (65)$$

for ε sufficiently small. This finishes the proof. \square

Proof of Proposition 3.1. First note that the graph that describes the bound of the default region for subsidiary 1 is a non-decreasing function of the capital of subsidiary 2. Similarly the graph that describes the bound of the default region of subsidiary 2 is a non-decreasing function

of the capital of subsidiary 1 (see Figure 2). We rewrite (19):

$$\begin{aligned}
f_l(C_1, C_2; z_1, z_2) &= \sum_{(j,k) \in \mathbf{D}_{J,l}} a_{j,k}(z_1, z_2) - z_l \\
&= \sum_{(j,k) \in \mathbf{D}_{J,l}} \mathbb{E}[\mathbf{1}_{\{(C_1, C_2) \geq (j,k)\}} \psi_{C_1-j, C_2-k}(p_1 z_1, p_2 z_2)] - z_l \\
&= \sum_{(j,k) \in \mathbf{D}_{J,l}} \sum_{(n,h) \geq (j,k)} \mathbb{E}[\mathbf{1}_{\{(C_1, C_2) = (n,h)\}} \psi_{n-j, h-k}(p_1 z_1, p_2 z_2)] - z_l \\
&= \sum_{(n,h)} \sum_{\substack{(j,k) \in \mathbf{D}_{J,l} \\ (j,k) \leq (n,h)}} \mathbb{E}[\mathbf{1}_{\{(C_1, C_2) = (n,h)\}} \psi_{n-j, h-k}(p_1 z_1, p_2 z_2)] - z_l. \tag{66}
\end{aligned}$$

For fixed (n, h) because of $\sum_{k,l \geq 0} \psi_{k,l}(p_1 z_1, p_2 z_2) = 1$, it follows that

$$\sum_{(j,k) \leq (n,h)} \mathbb{E}[\mathbf{1}_{\{(C_1, C_2) = (n,h)\}} \psi_{n-j, h-k}(p_1 z_1, p_2 z_2)] = \mathbb{P}((C_1, C_2) = (n, h)).$$

The shape of the default region is such that if $(n, h) \in \mathbf{D}_{J,l}$ and $(j, k) \leq (n, h)$, then also $(j, k) \in \mathbf{D}_{J,l}$. This implies that in the inner sum in (66) for $(n, h) \in \mathbf{D}_{J,l}$ we can actually drop the condition $(j, k) \in \mathbf{D}_{J,l}$. Thus we can rewrite the double sum in (66) as

$$\mathbb{P}((C_1, C_2) \in \mathbf{D}_{J,l}) + \sum_{(n,h) \in \mathbf{D}_{J,l}^c} \sum_{\substack{(j,k) \in \mathbf{D}_{J,l} \\ (j,k) \leq (n,h)}} \mathbb{E}[\mathbf{1}_{\{(C_1, C_2) = (n,h)\}} \psi_{n-j, h-k}(p_1 z_1, p_2 z_2)]. \tag{67}$$

Now observe that $\mathbb{E}[\mathbf{1}_{\{(C_1, C_2) = (n,h)\}} \psi_{n-j, h-k}(p_1 z_1, p_2 z_2)] / \mathbb{P}((C_1, C_2) = (n, h))$ is the conditional expectation of the random variable $\psi_{C_1-j, C_2-k}(p_1 z_1, p_2 z_2)$, given that $(C_1, C_2) = (n, h)$ and

$$(1/\mathbb{P}((C_1, C_2) = (n, h))) \cdot \sum_{\substack{(j,k) \in \mathbf{D}_{J,l} \\ (j,k) \leq (n,h)}} \mathbb{E}[\mathbf{1}_{\{(C_1, C_2) = (n,h)\}} \psi_{n-j, h-k}(p_1 z_1, p_2 z_2)],$$

the conditional expectation of the random variable

$$\Lambda_{C_1, C_2}^l(p_1 z_1, p_2 z_2) = \sum_{\substack{(j,k) \in \mathbf{D}_{J,l} \\ (j,k) \leq (n,h)}} \psi_{C_1-j, C_2-k}(p_1 z_1, p_2 z_2), \tag{68}$$

given that $(C_1, C_2) = (n, h)$. By the definition of ψ , this is just the probability that $(n, h) + (X_1(p_1 z_1), X_2(p_2 z_2))$ is in the default region where X_1 and X_2 being independent Poisson distributed random variables with parameter $p_1 z_1$ and $p_2 z_2$ respectively. Summing over (n, h) , as we do in (67), now gives us by the law of total expectation the unconditional probability of $\Lambda_{C_1, C_2}(p_1 z_1, p_2 z_2)$ which shows (22). By the law of total probability it follows that

$$f_l(C_1, C_2; z_1, z_2) = \mathbb{P}((C_1, C_2) + (X_1(p_1 z_1), X_2(p_2 z_2)) \in \mathbf{D}_{J,l}),$$

with $X_1(p_1 z_1)$ and $X_2(p_2 z_2)$ being mutually independent Poisson distributed random variables with parameter $p_1 z_1$ and $p_2 z_2$ independent of (C_1, C_2) . \square

Proof of Corollary 3.2. We prove the result for f_l , the proof for $D_v f_l$ is very similar. For fixed $(z_1, z_2) \in \mathbb{R}_+ \times \mathbb{R}_+$, point-wise convergence follows directly from Proposition 3.1 since for every

(z_1, z_2) , the integrand $\Lambda_{C_1, C_2}(z_1 p_1, z_2 p_2)$ in (22) is bounded by 1. By weak convergence of (C_1^n, C_2^n) it then follows that $f_l(C_1^n, C_2^n; z_1, z_2)$ converges to $f_l(C_1, C_2; z_1, z_2)$. Let now $A \subset \mathbb{R}_+ \times \mathbb{R}_+$ compact. For simplicity we assume that A is an interval of the form $A = [a, b]$ with $a, b \in \mathbb{R}_+ \times \mathbb{R}_+$ but the proof for general compact sets does not pose any further complications. For a given δ , choose now $K \in \mathbb{N}$ and points $x^1 = (x_1^1, x_2^1), \dots, x^K = (x_1^K, x_2^K) \in A$ such that for every $x = (x_1, x_2) \in A$ there exists $0 \leq k \leq K$, such that $|x - x_k| \leq \delta/2$. Now choose n large enough such that $|f_l(C_1^n, C_2^n; x_1^k, x_2^k) - f_l(C_1, C_2; x_1^k, x_2^k)| \leq \delta/2(1 + \max\{p_1, p_2\})$ for all $0 \leq k \leq K$. By (20), the partial derivatives of f_l are bounded by $1 + \max\{p_1, p_2\}$. Therefore, for arbitrary x , choose x_k such that $|x - x_k| \leq \delta/2$. By the triangular inequality

$$\begin{aligned} & |f_l(C_1^n, C_2^n; x_1, x_2) - f_l(C_1, C_2; x_1, x_2)| \\ &= \left| f_l(C_1^n, C_2^n; x_1, x_2) - f_l(C_1^n, C_2^n; x_1^k, x_2^k) + f_l(C_1^n, C_2^n; x_1^k, x_2^k) - f_l(C_1, C_2; x_1, x_2) \right| \\ &\leq \left| f_l(C_1^n, C_2^n; x_1, x_2) - f_l(C_1^n, C_2^n; x_1^k, x_2^k) \right| + \left| f_l(C_1^n, C_2^n; x_1^k, x_2^k) - f_l(C_1, C_2; x_1, x_2) \right| \\ &\leq \delta, \end{aligned}$$

which shows uniform convergence. \square

Proof of Corollary 3.3. The first statement follows directly from the representation (23) since by the shape of the default region for $(j, k) < (n, h)$, it holds that

$$\Lambda_{j,k}^l(p_1 z_1, p_2 z_2) \geq \Lambda_{n,h}^l(p_1 z_1, p_2 z_2).$$

To prove 2. observe that for $(j, k) < (n, h)$, it actually holds that

$$\Lambda_{j,k}^l(p_1 z_1, p_2 z_2) > \Lambda_{n,h}^l(p_1 z_1, p_2 z_2)$$

for $(z_1, z_2) \neq (0, 0)$ unless $(j, k), (n, h) \in \mathbf{D}_{J,l}$. The condition $\mathbb{P}((\tilde{C}_1, \tilde{C}_2) \leq (j, k)) < \mathbb{P}((C_1, C_2) \leq (j, k))$ for some $(j, k) \in \mathbf{D}_{J,l}^c \cup \mathbf{B}_{J,l}$ ensures then exactly that $\mathbb{P}(\Lambda_{\tilde{C}_1, \tilde{C}_2}(p_1 z_1, p_2 z_2) \leq (j, k)) < \mathbb{P}(\Lambda_{C_1, C_2}(p_1 z_1, p_2 z_2) \leq (j, k))$ and by (22) the claim follows. \square

Proof Theorem 3.4. We start with the proof for 1. Let $\tilde{\varepsilon} > 0$ be given. We know by Corollary 3.2 that if $(\tilde{C}_1, \tilde{C}_2)$ and (C_1, C_2) are close (in distribution), then the functionals and their derivatives are close. Now, if there exists a direction $\mathbf{v} \in \mathbb{R}^2$ such that $D_{\mathbf{v}} f_l(C_1, C_2; \mathbf{0}) < 0$, then for ε small enough, we know that also $D_{\mathbf{v}} f_l(\tilde{C}_1, \tilde{C}_2; \mathbf{0}) =: \kappa < 0$ for $\mathbb{P}((\tilde{C}_1, \tilde{C}_2) \neq (C_1, C_2)) \leq \varepsilon$. In fact, it follows by continuity that $D_{\mathbf{v}} f_l(\tilde{C}_1, \tilde{C}_2; \mathbf{z}) < \kappa/2$ for \mathbf{z} in some neighborhood $B_\delta(\mathbf{0})$ of $\mathbf{0}$. By possibly decreasing ε even further to the point that $f_1(\tilde{C}_1, \tilde{C}_2; \mathbf{0}), f_2(\tilde{C}_1, \tilde{C}_2; \mathbf{0}) \leq -\tilde{\varepsilon}\kappa/4$ and ensuring that $\varepsilon/2 \leq \delta$, we get that $f_l(\tilde{C}_1, \tilde{C}_2; (\varepsilon/2)\mathbf{v}) < 0$ for $l \in \{1, 2\}$. By Lemma 3.5 below, it follows that the first joint zero $\hat{\mathbf{z}}$ of $f_1(\tilde{C}_1, \tilde{C}_2; \cdot)$ and $f_2(\tilde{C}_1, \tilde{C}_2; \cdot)$ is bounded by $(\varepsilon/2)\mathbf{v}$. By (21) we may assume that \mathbf{v} is such that $v_1 + v_2 = 1$ where $\mathbf{v} = (v_1, v_2)$. It then follows by Theorem 2.3 that $n^{-1}(\mathcal{S}_{1,2n-1} + \mathcal{S}_{2,2n-1}) \leq \varepsilon$.

To show part 2. about non-resilience: Again we start with an uninfected network (C_1, C_2) , which implies that $f_1(C_1, C_2; \mathbf{0}) = f_2(C_1, C_2; \mathbf{0}) = 0$. By assumption,

$$\min\{D_{\mathbf{v}} f_1(C_1, C_2; \mathbf{0}), D_{\mathbf{v}} f_2(C_1, C_2; \mathbf{0})\} =: \kappa > 0,$$

and by continuity on the directional derivative in the direction \mathbf{v} , there exists some neighborhood $B_\delta(\mathbf{0})$ of $\mathbf{0}$ such that

$$\min\{D_{\mathbf{v}} f_1(C_1, C_2; \mathbf{z}), D_{\mathbf{v}} f_2(C_1, C_2; \mathbf{z})\} =: \kappa/2 > 0,$$

for $\mathbf{z} \in \mathbf{B}_\delta(\mathbf{0})$. This implies that $f_1(C_1, C_2; t\mathbf{v}), f_2(C_1, C_2; t\mathbf{v}) > 0$ for $t \leq \delta$.

We show that this in fact implies that $\max\{f_1(C_1, C_2; \mathbf{z}), f_2(C_1, C_2; \mathbf{z})\} > 0$ for any $\mathbf{z} \in (\mathbf{0}, \delta\mathbf{v}]$. Let now $\tilde{\mathbf{z}} \in (\mathbf{0}, \delta\mathbf{v}]$ and assume without loss of generality that $\tilde{\mathbf{z}}$ is to the right of the line $\delta\mathbf{v}$ in \mathbb{R}^2 . Let $t_{\tilde{\mathbf{z}}}\mathbf{v}$ the unique point on the line with the same first coordinate as the point $\tilde{\mathbf{z}}$, i.e. $t_{\tilde{\mathbf{z}}}v_1 = z_1$ where $\mathbf{v} = (v_1, v_2)$. We then know that $f_1(C_1, C_2; t_{\tilde{\mathbf{z}}}\mathbf{v}), f_2(C_1, C_2; t_{\tilde{\mathbf{z}}}\mathbf{v}) > 0$. However, by monotonicity of $f_2(C_1, C_2; \mathbf{z})$ with respect to z_1 , it follows that $f_2(C_1, C_2; \tilde{\mathbf{z}}) \geq f_2(C_1, C_2; t_{\tilde{\mathbf{z}}}\mathbf{v}) > 0$. If instead $\tilde{\mathbf{z}}$ was on the left of the line $\delta\mathbf{v}$ the same argument applied to the second coordinate and the function f_1 would show that $f_1(C_1, C_2; \tilde{\mathbf{z}}) > 0$. Because $\tilde{\mathbf{z}} \in (\mathbf{0}, \delta\mathbf{v}]$ was chosen arbitrary, we conclude that $\max\{f_1(C_1, C_2; \mathbf{z}), f_2(C_1, C_2; \mathbf{z})\} > 0$ for any $\mathbf{z} \in (\mathbf{0}, \delta\mathbf{v}]$.

Now consider the infected network parameterized by $(\tilde{C}_1, \tilde{C}_2)$ with $\mathbb{P}((\tilde{C}_1, \tilde{C}_2) \leq (C_1, C_2)) = 1$ and $\mathbb{P}(\tilde{C}_1, \tilde{C}_2) \in \mathbf{D}_{J,1} \cup \mathbf{D}_{J,2}) > 0$. It follows by Corollary 3.3 item 2. that

$$f_l(C_1, C_2; z_1, z_2) < f_l(\tilde{C}_1, \tilde{C}_2; z_1, z_2)$$

for all $z_1, z_2 \geq 0$ and $l \in \{1, 2\}$. By above consideration we know that no point $\mathbf{z} \in (\mathbf{0}, \delta\mathbf{v}]$ exists such that both $f_1(C_1, C_2; \cdot)$ and $f_2(C_1, C_2; \cdot)$ are ≤ 0 . This implies that there exists also no point $\mathbf{z} \in (\mathbf{0}, \delta\mathbf{v}]$ such that both $f_1(\tilde{C}_1, \tilde{C}_2; \cdot)$ and $f_2(\tilde{C}_1, \tilde{C}_2; \cdot)$ are ≤ 0 . In particular because of $\max\{f_1(\tilde{C}_1, \tilde{C}_2; \mathbf{0}), f_2(\tilde{C}_1, \tilde{C}_2; \mathbf{0})\} > 0$, there can not be any joint zero of $f_1(\tilde{C}_1, \tilde{C}_2; \cdot)$ and $f_2(\tilde{C}_1, \tilde{C}_2; \cdot)$ in $[\mathbf{0}, \delta\mathbf{v}]$. Again, by Theorem 2.3 part 1. it follows that $n^{-1}(\mathcal{S}_{1,2n-1} + \mathcal{S}_{2,2n-1}) \geq \delta v_1 + \delta v_2$, independent of the specification of $(\tilde{C}_1, \tilde{C}_2)$. \square

Proof Lemma 3.5. First note that $f_1(0, z_2) \geq 0$ for any $z_2 \geq 0$ and $f_2(0, z_2) \geq 0$ for any $z_1 \geq 0$. We now construct a decreasing sequence of points $\{\mathbf{z}_n = (z_{1,n}, z_{2,n})\}_{n \in \mathbb{N}}$ with $\mathbf{z}_0 = \tilde{\mathbf{z}}$ and such that $\lim_{n \rightarrow \infty} \mathbf{z}_n$ exists and is a joint zero of f_1 and f_2 .

For this, set $\mathbf{z}_0 = \tilde{\mathbf{z}}$, and then for every $n > 0$ odd we define $\mathbf{z}_n = (z_{1,n}, z_{2,n})$ by $z_{2,n} = z_{2,n-1}$ and

$$z_{1,n} = \sup\{t \leq z_{1,n-1} \mid f_1(t, z_{2,n-1}) = 0\} \quad (69)$$

and for even n we define $\mathbf{z}_n = (z_{1,n}, z_{2,n})$ by $z_{1,n} = z_{1,n-1}$ and

$$z_{2,n} = \sup\{t \leq z_{2,n-1} \mid f_2(z_{1,n-1}, t) = 0\}. \quad (70)$$

We first need to show that the above sets are not empty and that $(z_{1,n}, z_{2,n})$ is actually a well defined point in $[0, \infty) \times [0, \infty)$ for every n . By assumption, we have that $f_1(\tilde{\mathbf{z}}), f_2(\tilde{\mathbf{z}}) \leq 0$. For the set defined in (69) as long as $f_1(\mathbf{z}_{n-1}) \leq 0$, it follows by continuity of f_1 and by $f_1(0, z_2) \geq 0$ that the set is not empty and $z_{1,n} \in [0, \infty)$. Moreover, by definition of $z_{1,n}$, it holds that $f_1(\tilde{\mathbf{z}}_n) = 0$ and by monotonicity of f_2 in its first argument it holds that $f_2(\tilde{\mathbf{z}}_n) \leq 0$ as long as $f_2(\tilde{\mathbf{z}}_{n-1}) \leq 0$. The same argument applies to (70).

Clearly, the sequence \mathbf{z}_n is decreasing and bounded from below by $\mathbf{0}$. This implies that the sequence converges to some point $\bar{\mathbf{z}} \leq \tilde{\mathbf{z}}$. Moreover, because $f_1(\mathbf{z}_n) = 0$ whenever n is odd and $f_2(\mathbf{z}_n) = 0$ whenever n is even, it follows by continuity of f that $f_1(\bar{\mathbf{z}}) = f_2(\bar{\mathbf{z}}) = 0$ and thus $\bar{\mathbf{z}}$ is a joint zero. Note that the sequence $\tilde{\mathbf{z}}_n$ is defined to approximate the joint zero $\bar{\mathbf{z}}$ from a region where both function values are ≤ 0 . \square

References

- Amini, H., Cont, R., and Minca, A. (2016a). Resilience to Contagion in Financial Networks. *Mathematical Finance*, 26(2):329–365.
- Amini, H., Filipović, D., and Minca, A. (2013). Systemic risk with central counterparty clearing. Swiss Finance Institute Research Paper No. 13-34, Swiss Finance Institute.

- Amini, H., Filipović, D., and Minca, A. (2016b). Uniqueness of equilibrium in a payment system with liquidation costs. *Operations Research Letters*, 44(1):1–5.
- Amini, H. and Fountoulakis, N. (2014). Bootstrap Percolation in Power-Law Random Graphs. *Journal of Statistical Physics*, 155(1):72–92.
- Amini, H., Fountoulakis, N., and Panagiotou, K. (2014). Bootstrap percolation in Inhomogeneous random graphs. *arXiv:1402.2815*.
- Anand, K., Craig, B., and Von Peter, G. (2015). Filling in the blanks: Network structure and interbank contagion. *Quantitative Finance*, 15(4):625–636.
- Arinaminpathy, N., Kapadia, S., and May, R. M. (2012). Size and complexity in model financial systems. *Proceedings of the National Academy of Sciences*, 109(45):18338–18343.
- Balogh, J. and Pittel, B. G. (2007). Bootstrap percolation on the random regular graph. *Random Structures Algorithms*, 30(1-2):257–286.
- Bardoscia, M., Barucca, P., Brinley Codd, A., and Hill, J. (2017). The decline of solvency contagion risk. *Bank of England Staff Working Paper*, 662.
- Bichuch, M. and Feinstein, Z. (2019). Optimization of fire sales and borrowing in systemic risk. *SIAM Journal on Financial Mathematics*, 10(1):68–88.
- Bichuch, M. and Feinstein, Z. (2022). A repo model of fire sales with vwap and lob pricing mechanisms. *European Journal of Operational Research*, 296(1):353–367.
- Boss, M., Elsinger, H., Summer, M., and Thurner, S. (2004). Network topology of the interbank market. *Quantitative Finance*, 4(6):677–684.
- Capponi, A., Chen, P.-C., and Yao, D. D. (2016). Liability concentration and systemic losses in financial networks. *Operations Research*, 64(5):1121–1134.
- Chen, N., Liu, X., and Yao, D. D. (2016). An optimization view of financial systemic risk modeling: The network effect and the market liquidity effect. *Operations Research*, 64(5).
- Cifuentes, R., Shin, H. S., and Ferrucci, G. (2005). Liquidity risk and contagion. *Journal of the European Economic Association*, 3(2-3):556–566.
- Detering, N., Meyer-Brandis, T., and Panagiotou, K. (2019). Bootstrap Percolation in Directed Inhomogeneous Random Graphs. *The Electronic Journal of Combinatorics*, 26(3).
- Detering, N., Meyer-Brandis, T., Panagiotou, K., and Ritter, D. (2019). Managing default contagion in inhomogeneous financial networks. *SIAM Journal on Financial Mathematics*, 10(2):578–614.
- Detering, N., Meyer-Brandis, T., Panagiotou, K., and Ritter, D. (2021). An integrated model for fire sales and default contagion. *Mathematics and Financial Economics*, 15(1):59–101.
- Detering, N., Meyer-Brandis, T., Panagiotou, K., and Ritter, D. (2022). Suffocating fire sales. *SIAM Journal on Financial Mathematics*, 13(1):70–108.
- Detering, N., Meyer-Brandis, T., Panagiotou, K., Ritter, D., et al. (2020). Financial contagion in a stochastic block model. *International Journal of Theoretical and Applied Finance*, 23(08):1–53.
- Eisenberg, L. and Noe, T. H. (2001). Systemic risk in financial systems. *Management Science*, 47(2):236–249.
- Elliott, M., Golub, B., and Jackson, M. O. (2014). Financial networks and contagion. *American Economic Review*, 104(10):3115–3153.

- Elsinger, H. (2009). Financial networks, cross holdings, and limited liability. *Österreichische Nationalbank (Austrian Central Bank)*, 156.
- Elsinger, H., Lehar, A., and Summer, M. (2013). Network models and systemic risk assessment. In *Handbook on Systemic Risk*, pages 287–305. Cambridge University Press.
- Gai, P., Haldane, A., and Kapadia, S. (2011). Complexity, concentration and contagion. *Journal of Monetary Economics*, 58(5):453–470.
- Gai, P. and Kapadia, S. (2010). Contagion in Financial Networks. *Proc. R. Soc. A*, page 2401–2423.
- Glasserman, P. and Young, H. P. (2015). How likely is contagion in financial networks? *Journal of Banking and Finance*, 50:383–399.
- Gouriéroux, C., Héam, J.-C., and Monfort, A. (2012). Bilateral exposures and systemic solvency risk. *Canadian Journal of Economics*, 45(4):1273–1309.
- Halaç, G. and Kok, C. (2015). Modelling the emergence of the interbank networks. *Quantitative Finance*, 15(4):653–671.
- Hofstad, R. v. d. (2016). *Random Graphs and Complex Networks*, volume 1 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press.
- Hüser, A.-C. (2015). Too interconnected to fail: A survey of the interbank networks literature. *Journal of Network Theory in Finance*, 1(3):1–50.
- Janson, S., Luczak, T., Turova, T., and Vallier, T. (2012). Bootstrap percolation on the random graph $G_{n,p}$. *Ann. Appl. Probab.*, 22(5):1989–2047.
- Lewis, M. (2010). *The Big Short : Inside the Doomsday Machine*. New York :Simon & Schuster.
- Nier, E., Yang, J., Yorulmazer, T., and Alentorn, A. (2007). Network models and financial stability. *Journal of Economic Dynamics and Control*, 31(6):2033–2060.
- Rogers, L. C. and Veraart, L. A. (2013). Failure and rescue in an interbank network. *Management Science*, 59(4):882–898.
- Staum, J. (2013). Counterparty contagion in context: Contributions to systemic risk. In *Handbook on Systemic Risk*, pages 512–548. Cambridge University Press.
- Upper, C. (2011). Simulation methods to assess the danger of contagion in interbank markets. *Journal of Financial Stability*, 7(3):111–125.
- Weber, S. and Weske, K. (2017). The joint impact of bankruptcy costs, fire sales and cross-holdings on systemic risk in financial networks. *Probability, Uncertainty and Quantitative Risk*, 2(1):9.
- Wormald, N. C. (1995). Differential Equations for Random Processes and Random Graphs. *The Annals of Applied Probability*, 5(4):1217 – 1235.