

Convex analysis on Hadamard spaces and scaling problems

Hiroshi HIRAI

Graduate School of Mathematics,
Nagoya University, Nagoya, 464-8602, Japan.

hirai.hiroshi@math.nagoya-u.ac.jp

June 1, 2023

Abstract

In this paper, we address the bounded/unbounded determination of geodesically convex optimization on Hadamard spaces. In Euclidean convex optimization, the recession function is a basic tool to study the unboundedness, and provides the domain of the Legendre-Fenchel conjugate of the objective function. In a Hadamard space, the asymptotic slope function (Kapovich, Leeb, and Millson 2009), which is a function on the boundary at infinity, plays a role of the recession function. We extend this notion by means of convex analysis and optimization, and develop a convex analysis foundation for the unbounded determination of geodesically convex optimization on Hadamard spaces, particularly on symmetric spaces of nonpositive curvature. We explain how our developed theory is applied to operator scaling and related optimization on group orbits, which are our motivation.

Keywords: Convex analysis, Hadamard space, CAT(0) space, recession function, Legendre-Fenchel conjugate, symmetric space, Euclidean building, matrix and operator scaling, null-cone membership, moment polytope, submodular function, Busemann function

1 Introduction

Hadamard spaces are complete geodesic metric spaces having nonpositive curvature. In such a space, a geodesic connecting any two points is uniquely determined. A function on a Hadamard space is called (*geodesically*) *convex* if it is convex along any geodesics. The theory of convex optimization on Hadamard spaces is a promising direction of research, though it has just started and is still undeveloped; see e.g., [5]. Since the influential paper [22] by Garg, Gurvits, Oliveira, and Wigderson on *operator scaling* [25], apparently unrelated problems in diverse fields of mathematical sciences have been formulated and partially/completely solved via geodesically convex optimization on Riemannian manifolds; see [2, 13, 15, 19, 23, 28] and references therein. These manifolds are, in fact, *Hadamard manifolds* (Riemannian manifolds that are Hadamard spaces), more specifically, *symmetric spaces of nonpositive curvature*.

In these problems, as well as finding near-optimal solutions, deciding boundedness of the optimization problem,

$$\inf_{x \in X} f(x) > -\infty \text{ or } = -\infty \quad (1.1)$$

becomes an important issue. Examples are the approximate scalability in operator scaling and its invariant theoretic generalizations (null-cone membership, moment polytope membership); see the above references.

The present paper addresses this bounded/unbounded determination by means of convex analysis. For explaining our approach, let us recall the Euclidean situation. In Euclidean convex optimization, *recession functions* (also called *asymptotic functions*) are a basic tool to study the boundedness property; see [29, Section 3.2] and [40, Section 8]. For a convex function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, the recession function $f^\infty : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$ is defined by

$$f^\infty(u) := \lim_{t \rightarrow \infty} f(x + tu)/t \quad (u \in \mathbb{R}^n), \quad (1.2)$$

where $f^\infty(u)$ is independent of $x \in \mathbb{R}^n$. The recession function f^∞ is a positively homogeneous convex function, and links with *Legendre-Fenchel duality* as follows. Recall the *Legendre-Fenchel conjugate* $f^* : \mathbb{R}^n \rightarrow \mathbb{R}^n \cup \{\infty\}$ of f , which is defined by $f^*(p) := \sup_{x \in \mathbb{R}^n} \langle p, x \rangle - f(x)$. Its domain $\text{dom } f^* := \{p \in \mathbb{R}^n \mid f^*(p) < \infty\}$ is precisely the set of vectors $p \in \mathbb{R}^n$ for which $\inf_{x \in \mathbb{R}^n} f(x) - \langle p, x \rangle$ is bounded below. Then, the recession function f^∞ equals the *support function* of the domain $\text{dom } f^*$. Namely, $f^\infty(u) = \sup_{p \in \text{dom } f^*} \langle u, p \rangle$ holds, and the closure $\overline{\text{dom } f^*}$ of $\text{dom } f^*$ equals

$$B(f^\infty) := \{p \in \mathbb{R}^n \mid \langle u, p \rangle \leq f^\infty(u) \ (u \in \mathbb{R}^n)\}. \quad (1.3)$$

See [29, Example 2.4.6] and [40, Theorem 13.3]. In particular, f^∞ provides an inequality description of $\text{dom } f^*$.

Suppose further that f is smooth. The image $\nabla f(\mathbb{R}^n)$ of gradient map $x \mapsto \nabla f(x)$ is precisely the set of vectors p for which the infimum of $\inf_{x \in \mathbb{R}^n} f(x) - \langle p, x \rangle$ is attained. Then, the gradient space $\nabla f(\mathbb{R}^n)$ contains the relative interior $\text{ri } B(f^\infty)$ of $B(f^\infty) = \overline{\text{dom } f^*}$ [40, Corollary 26.4.1]. Thus, the following relation holds:

$$\text{ri } B(f^\infty) \subseteq \nabla f(\mathbb{R}^n) \subseteq \text{dom } f^* \subseteq B(f^\infty). \quad (1.4)$$

Particularly, $\overline{\nabla f(\mathbb{R}^n)} = \overline{\text{dom } f^*} = B(f^\infty)$ holds. If $\text{dom } f^*$ is closed, then the following conditions are equivalent for $p \in \mathbb{R}^n$:

- (a) $\inf_{x \in \mathbb{R}^n} \|\nabla f(x) - p\| = 0$.
- (b) $-f^*(p) = \inf_{x \in \mathbb{R}^n} f(x) - \langle p, x \rangle > -\infty$.
- (c) $p \in B(f^\infty)$.

This equivalence plays fundamental roles in *matrix scaling* [44]—the origin of operator scaling and related group orbits optimization. Given an $n \times n$ nonnegative matrix $A = (a_{ij})$ and positive vectors $r, c \in \mathbb{R}^n$ with the same sum $\sum r_i = \sum_i c_i = l$, the matrix scaling problem is to ask positive diagonal matrices R, C such that $((RAC)^\top \mathbf{1}, RAC\mathbf{1}) = (r, c)$, where $\mathbf{1}$ denotes the all-ones vector. The matrix A is said to be *scalable* if there exist such R, C , and *approximately scalable* if there exist R, C

such that $((RAC)^\top \mathbf{1}, RAC\mathbf{1})$ is arbitrary close to (r, c) . In fact, the approximate scalability is written as the condition (a) for convex function $f_A(s, t) := l \log \sum_{i,j} e^{s_i} a_{ij} e^{t_j}$ and vector $p = (r, c)$. The recession function f_A^∞ is given by $f_A^\infty(u, v) = l \max\{u_i + v_j \mid i, j : a_{ij} \neq 0\}$. The condition (c) is written as a network flow LP and efficiently verified. The combinatorial characterization of the approximate scalability by Rothblum and Schneider [41] was proved via f^∞ in this way. From (b) and the fact that $\text{dom } f_A^*$ is closed, approximate scaling RAC is obtained by solving convex optimization $\inf_{s,t} f_A(s, t) - \langle r, s \rangle - \langle c, t \rangle$. *Sinkhorn algorithm* [44] is viewed as an alternating minimization algorithm for this problem.

The goal of this paper is to establish analogues of the equivalence of (a), (b), and (c) for geodesically convex optimization on Hadamard spaces, and to provide a convex analysis foundation to the above mentioned problems. We particularly focus on an analogous notion of the recession function on a Hadamard space. In fact, such a notion was already introduced by Kapovich, Leeb, and Millson [31] (see also [46, Section 5.4]), who defined the *asymptotic slope* of a convex function via a geodesic analogue of (1.2). By utilizing the asymptotic slope, they studied the boundedness (semistability) for a class of convex functions related to a generalization of Horn’s problem. Our presented theory reformulates and extends some of their arguments from viewpoints of convex analysis and optimization.

The structure and results of this paper are outlined as follows. In Section 2, we provide necessary backgrounds on a Hadamard space X , particularly, the *boundary* X^∞ at infinity and its *Euclidean cone* CX^∞ . The cone CX^∞ of the boundary is also a Hadamard space, and plays a role of the “dual space” of X . With a point $p \in CX^\infty$, we associate the *Busemann function* b_p as a correspondent of a linear function, and regard the boundary CX^∞ as the space of Busemann functions. This viewpoint leads to the notion of the *asymptotic Legendre-Fenchel conjugate* f^* of f , which is a function on CX^∞ defined by $p \mapsto f^*(p) := \sup_{x \in X} -b_p(x) - f(x)$. The asymptotic slope is a function on X^∞ . We extend it, in a homogeneous way, to introduce the recession function f^∞ on CX^∞ . We also define the associated subset $B(f^\infty)$ via an analogue of (1.3). We show that f^∞ is positively homogeneous convex on CX^∞ and that $p \in B(f^\infty)$ is a necessary condition for $f^*(p) < \infty$, i.e., $\text{dom } f^* \subseteq B(f^\infty)$.

We then move to smooth convex optimization on a Hadamard manifold M . The boundary cone CM^∞ is identified with the tangent space T_x at any point x , though CM^∞ has a different topology from the usual one $T_x \simeq \mathbb{R}^n$, and is not a manifold in general. We define the *asymptotic gradient* $\nabla^\infty f(x)$ as the image of the gradient $\nabla f(x)$ in CM^∞ . This notion fits into our purpose: We verify a weaker analogue of (1.4) in which ∇ is replaced by ∇^∞ and ri is placed by the interior. We also verify an analogue of the equivalence of (a) and (c).

We further restrict our study to *symmetric spaces of nonpositive curvature*—a representative class of Hadamard manifolds having rich group symmetry. The boundary CM^∞ has a polyhedral cone complex structure, known as a *Euclidean building*. We utilize the symmetry property and building structure to show that the conjugate f^* is convex on each cone (*Weyl chamber*). We also provide some fundamental results on $\text{dom } f^*$ and $B(f^\infty)$. Then we present detailed calculations and specializations of several concepts for the symmetric space $P_n = GL(n, \mathbb{C})/U(n)$ of positive definite matrices. The boundary CP_n^∞ is viewed as the order complex of flags of vector subspaces, which naturally links *submodular functions* on the lattice of vector subspaces and their

Lovász extension [26, 27]. Via the Lovász extension, submodular functions correspond to convex functions on CP_n^∞ that are linear on each cone. For a convex function f on M inducing a submodular function at infinity, called an *asymptotically submodular* function, $B(f^\infty)$ is an analogue of the *base polyhedron* and the membership of $B(f^\infty)$ becomes discrete convex optimization (i.e., *submodular function minimization*) over the modular lattice of all vector subspaces of \mathbb{C}^n .

In Section 3, we explain how the developed theory is applied to operator scaling and related optimization on group orbits. These problems are viewed as convex optimization on symmetric spaces of nonpositive curvature. We take up the *operator scaling with marginals* by Franks [19]. We compute the recession function, from which Franks' characterization of the approximate triangular scalability is naturally deduced.

We finally consider the null-cone and moment polytope membership for a linear action of a reductive group G , formulated by Bürgisser, Franks, Garg, Oliveira, Walter, and Wigderson [15]. These are convex optimization of the *Kempf-Ness function* f_v over symmetric space $M = G/K$. In this setting, asymptotic gradients give rise to the *moment map* and *moment polytope*, for which f_v^∞ gives an inequality description. We establish the link with Hilbert-Mumford criterion and Kempf-Ness theorem, and verify from our approach shifting trick and convexity theorem for the moment polytope.

The main implication of this paper is that finding a one-parameter subgroup in the Hilbert-Mumford criterion and membership of the moment polytope (after randomization) reduce to minimization of recession functions f_v^∞ over Euclidean building CM^∞ . This is a far-reaching generalization of the approach by Hamada and Hirai [26] for the *noncommutative-rank* computation (= the null-cone problem of the left-right action), in which their algorithm is now viewed as minimizing f_v^∞ . The current approach (e.g., [15, 28]) for these problems is mainly based on smooth convex optimization (b) on G/K . We hope that our results will motivate to develop nonsmooth convex optimization techniques on non-manifold Hadamard spaces, for verifying (c).

2 Convex analysis on Hadamard spaces

2.1 Hadamard spaces

Here we introduce Hadamard spaces; see [5, 6, 11] for details. Let X be a metric space with distance function d . A *path* in X is a continuous map from interval $[0, l] \subseteq \mathbb{R}$ to X , where $l \geq 0$. We say that a path $c : [0, l] \rightarrow X$ connects $c(0)$ and $c(l)$. A path $c : [0, l] \rightarrow X$ is said to be *geodesic* if $d(c(s), c(t)) = |s - t|$ for every $s, t \in [0, l]$. A *geodesic metric space* is a metric space X in which any pair of two points is connected by a geodesic path.

Suppose that X is a geodesic metric space. Let $x, y, z \in X$. A *geodesic triangle* of x, y, z is the union of geodesic paths connecting x, y , x, z , and y, z . The *comparison triangle* is the triangle in Euclidean plane \mathbb{R}^2 with vertices $\bar{x}, \bar{y}, \bar{z}$ such that $d(x, y) = \|\bar{x} - \bar{y}\|_2$, $d(y, z) = \|\bar{y} - \bar{z}\|_2$ and $d(z, x) = \|\bar{z} - \bar{x}\|_2$. In the geodesic triangle, suppose that points x, y and x, z are connected by geodesic paths c and c' , respectively, with $c(0) = c'(0) = x$. For $t, t' \in [0, 1]$, let $p := c(td(x, y))$ and $q := c'(t'd(x, z))$. Let $\bar{p} := (1 - t)\bar{x} + t\bar{y}$ and $\bar{q} := (1 - t')\bar{x} + t'\bar{z}$ be the corresponding points in \mathbb{R}^2 . Then the *CAT(0)-inequality* is given by

$$d(p, q) \leq \|\bar{p} - \bar{q}\|_2. \quad (2.1)$$

A geodesic metric space X is said to be $CAT(0)$ if the $CAT(0)$ inequality (2.1) holds for every choice of a geodesic triangle and $t \in [0, 1]$. There are several ways of defining $CAT(0)$ spaces. A useful one is the following: X is $CAT(0)$ if for every triple of points $x, y, z \in X$, geodesic path c with $c(0) = x$ and $c(d(x, y)) = y$, and $t \in [0, 1]$, it holds

$$d(p_t, z)^2 \leq (1-t)d(x, z)^2 + td(y, z)^2 - t(1-t)d(x, y)^2, \quad (2.2)$$

where $p_t = c(td(x, y))$. Notice that the RHS equals the squared comparison distance between $\bar{p}_t = (1-t)\bar{x} + t\bar{y}$ and \bar{z} .

It is known [11, II.1.4] that $CAT(0)$ spaces are *uniquely geodesic*, i.e., a geodesic path connecting any two points is unique. In this case, for points $x, y \in X$, let $[x, y]$ denote the image of the unique geodesic path connecting x, y . For $t \in [0, 1]$, let $(1-t)x + ty$ denote the point $z \in [x, y]$ with $d(x, z)/d(x, y) = t$.

A *Hadamard space* is a $CAT(0)$ space that is complete as a metric space. Let X be a Hadamard space. A subset $S \subseteq X$ is called *convex* if $x, y \in X$ implies $[x, y] \subseteq S$. The smallest convex set including a given $S \subseteq X$ is called the *convex hull* of S . A function $f : X \rightarrow \mathbb{R} \cup \{\infty\}$ is said to be *convex* if for all $x, y \in X, t \in [0, 1]$ it satisfies

$$(1-t)f(x) + tf(y) \geq f((1-t)x + ty). \quad (2.3)$$

f is called *strictly convex* if $<$ holds in (2.3) for any $t \in (0, 1)$. If $-f$ is convex, then f is said to be *concave*. An *affine function* is a convex and concave function. A function $f : X \rightarrow \mathbb{R}$ is said to be *L-Lipschitz* with parameter $L \geq 0$ if it satisfies $|f(x) - f(y)| \leq Ld(x, y)$ for all $x, y \in X$.

2.1.1 Boundary at infinity and its Euclidean cone

To study the asymptotic behavior of convex functions, we consider the boundary of a Hadamard space X ; see [6, Chapter II] and [11, Chapter II.8] for the boundary.

A *geodesic ray* is a continuous map $c : [0, \infty) \rightarrow X$ such that $d(c(s), c(t)) = |s-t|$ for $s, t \in [0, \infty)$. With a geodesic ray c and $r \geq 0$, the map written as $[0, \infty) \ni t \mapsto c(rt)$ is called a *constant-speed ray*, or simply, a *ray*. If $c(0) = x$, we say that ray c *issues from* x . Two geodesic rays c, c' are said to be *asymptotic* if there is a positive constant $K > 0$ such that $d(c(t), c'(t)) \leq K$ for all $t \geq 0$. The asymptotic relation is an equivalence relation on the set of all geodesic rays. Let X^∞ denote the set of all equivalence classes, which is called the *boundary of X at infinity*. The equivalence class of c is called the *asymptotic class* of c , and is denoted by $c(\infty)$. For any point $x \in X$, and each $\xi \in X^\infty$ there is a unique geodesic ray c issuing from x such that $c(\infty) = \xi$. Therefore, X^∞ can be identified with the set of all geodesic rays issuing from any fixed x .

We next introduce a metric on X^∞ . For points $x, y, z \in X$ (with $y \neq x \neq z$), the *comparison angle* $\bar{\angle}_x(y, z)$ between y and z at x is the inner angle of the comparison triangle of x, y, z at x . For two geodesic rays c, c' issuing from x , let the angle $\angle_x(c, c')$ of c, c' at x be defined by

$$\angle_x(c, c') := \lim_{t \rightarrow 0} \bar{\angle}_x(c(t), c'(t)), \quad (2.4)$$

where the limit indeed exists [11, II.3.1]. For $\xi, \xi' \in X^\infty$, the *angle* $\angle(\xi, \xi') \in [0, \pi]$ is defined by

$$\angle(\xi, \xi') := \sup_{x \in X} \angle_x(c, c') = \lim_{t, t' \rightarrow \infty} \bar{\angle}_x(c(t), c'(t')) = \sup_{t, t' > 0} \bar{\angle}_x(c(t), c'(t')), \quad (2.5)$$

where c, c' are geodesic rays issuing from x with $c(\infty) = \xi, c'(\infty) = \xi'$. The equalities in (2.5) follow from [11, II. 9.8 (1)] (where x is arbitrary). Here $(\xi, \xi') \mapsto \angle(\xi, \xi')$ defines a metric on X^∞ , which is called the *angular metric*. Then X^∞ becomes a metric space, and a topological space accordingly.

In addition to X^∞ , we consider its *Euclidean cone* CX^∞ ; see [11, Chapter I.5] for generalities of the Euclidean cone construction. Let \mathbb{R}_+ denote the set of nonnegative numbers. The set CX^∞ is the quotient of $\mathbb{R}_+ \times X^\infty$ by the equivalence relation $(r, \xi) \simeq (r', \xi')$ if $r = r' = 0$ or $(r, \xi) = (r', \xi')$. The equivalence class of (r, ξ) is denoted by $r\xi$. By identifying ξ with 1ξ , we regard X^∞ as a subset of CX^∞ . Let 0 denote the class of $(0, \xi)$. The angle $\angle(p, p')$ of (nonzero) points $p = r\xi, p' = r'\xi' \in CX^\infty$ is defined as $\angle(\xi, \xi')$.

The space CX^∞ is viewed as the space of all constant-speed rays issuing from any fixed point. Indeed, $r\xi \in CX^\infty$ is associated with a constant-speed ray $t \mapsto c^*(t) = c(rt)$, where c is a geodesic ray with $c(\infty) = \xi$. In this case, we let $c^*(\infty) := r\xi$. The space CX^∞ is metrized by the following distance d^∞ :

$$d^\infty(r\xi, r'\xi')^2 := r^2 + r'^2 - 2rr' \cos \angle(\xi, \xi') \quad (r\xi, r'\xi' \in CX^\infty). \quad (2.6)$$

The topology of CX^∞ is given accordingly, which is called the *d^∞ -topology*.

Lemma 2.1 (see [6, II.4.8]). *CX^∞ is a Hadamard space.*

A *flat triangle* in a Hadamard space is a geodesic triangle whose convex hull is isometric to the convex hull of their comparison triangle in \mathbb{R}^2 . It is known (see [11, II.2.9]) that if one of the vertex angles of the triangle is equal to the corresponding angle of the comparison triangle, then it is flat. By construction of CX^∞ , the angle at 0 is always equal to the corresponding angle of the comparison triangle. Hence we have:

Lemma 2.2. *In CX^∞ , any three points containing 0 form a flat triangle.*

Therefore, the convex hull of $\{0, tp, t'p'\}_{t, t' \in \mathbb{R}_+}$ is isometric to the convex cone C in \mathbb{R}^2 . Then, via the isometry to $C \subseteq \mathbb{R}^2$, we can consider nonnegative combinations in two nonzero points in CX^∞ . For a point $p = t\xi \in CX^\infty$ and $a \in \mathbb{R}_+$, define $ap := (at)\xi \in CX^\infty$. It is clear that $p \mapsto ap$ is continuous and $a(bp) = (ab)p$. For two points $p, q \in CX^\infty$, the sum $p + q$ of p, q is defined by

$$p + q := 2((1/2)p + (1/2)q).$$

Recall that $(1/2)p + (1/2)q$ is the midpoint of the geodesic path between p and q . Map $(p, q) \mapsto p + q$ is continuous; this follows from the fact [11, II.1.4] that the geodesic segment in a Hadamard space varies continuously with its endpoints. Notice that $+$ is not associative in general; so there may be many “inverses” q of p with $p + q = 0$. From (2.6), it holds $d^\infty(\alpha p, \alpha q) = \alpha d^\infty(p, q)$. This means that by $p \mapsto \alpha p$, geodesic segment $[p, q]$ is mapped to geodesic segment $[\alpha p, \alpha q]$. Then we have a linearity relation

$$\alpha(p + q) = \alpha p + \alpha q \quad (\alpha \in \mathbb{R}_+). \quad (2.7)$$

Define the inner product $\langle p, q \rangle$ of $p, q \in CX^\infty$ by

$$2\langle p, q \rangle := d^\infty(0, p)^2 + d^\infty(0, q)^2 - d^\infty(p, q)^2. \quad (2.8)$$

For $p = t\xi \in CX^\infty$, define the norm $\|p\|$ by $\|p\| := t$. Observe from definitions (2.6) (2.8) that $\|p\| = d^\infty(0, p) = \sqrt{\langle p, p \rangle}$. Then we have

$$\langle p, q \rangle = \|p\| \|q\| \cos \angle(p, q). \quad (2.9)$$

A function $f : CX^\infty \rightarrow \mathbb{R} \cup \{\infty\}$ is called *positively homogeneous* if it holds

$$f(\alpha p) = \alpha f(p) \quad (a \in \mathbb{R}_+).$$

Lemma 2.3. *For any $q \in CX^\infty$, the function $p \mapsto \langle p, q \rangle$ is continuous and positively homogeneous concave.*

Proof. The continuity follows from (2.8) that $\langle \cdot, \cdot \rangle$ is written by continuous function d^∞ . The positive homogeneity follows from (2.9). For concavity, it suffices to show

$$\langle p + p', q \rangle \geq \langle p, q \rangle + \langle p', q \rangle. \quad (2.10)$$

Twice the RHS is equal to

$$-d^\infty(p, q)^2 - d^\infty(p', q)^2 + \|p\|^2 + \|p'\|^2 + 2\|q\|^2. \quad (2.11)$$

On the other hand, twice the LHS is equal to

$$4\langle (p + p')/2, q \rangle = -2d^\infty((p + p')/2, q)^2 + 2\|(p + p')/2\|^2 + 2\|q\|^2. \quad (2.12)$$

Since $0, p, p'$ form a flat triangle (Lemma 2.2), the CAT(0) inequality (2.2) with $(x, y, z) = (p, p', 0)$ and $t = 1/2$ holds in equality:

$$2\|(p + p')/2\|^2 = \|p\|^2 + \|p'\|^2 - d^\infty(p, p')^2/2. \quad (2.13)$$

From (2.11), (2.12), and (2.13), we see that (2.10) is equivalent to

$$2d^\infty((p + p')/2, q)^2 \leq d^\infty(p, q)^2 + d^\infty(p', q)^2 - d^\infty(p, p')^2/2. \quad (2.14)$$

This is the CAT(0)-inequality (2.2) with $(x, y, z) = (p, p', q)$ and $t = 1/2$, which holds by Lemma 2.1. \square

Example 2.4. Consider the case of $X = \mathbb{R}^n$. Two geodesic rays $t \rightarrow \xi t + b$ and $t \rightarrow \xi' t + b'$ are asymptotic if and only if $\xi = \xi'$. The angle of the corresponding asymptotic classes is given by $\cos^{-1}\langle \xi, \xi' \rangle$. Therefore, X^∞ is isometric to the sphere S^{n-1} . The Euclidean cone CX^∞ is isometric to the Euclidean space $\mathbb{R}^n = \mathbb{R}_+ \times S^{n-1} / \simeq$. Also the distance d^∞ and product $\langle \cdot, \cdot \rangle$ coincides with the Euclidean ones.

2.1.2 Busemann functions and asymptotic Legendre-Fenchel conjugate

For a geodesic ray c , the *Busemann function* $b_c : X \rightarrow \mathbb{R}$ is defined by

$$b_c(x) := \lim_{t \rightarrow \infty} d(x, c(t)) - t \quad (x \in X). \quad (2.15)$$

Lemma 2.5 (see [11, II.8.22]). *Busemann functions for geodesic rays are 1-Lipschitz convex functions.*

Two asymptotic geodesic rays c, c' yields the same Busemann functions up to additive constant:

Lemma 2.6 (see [11, II.8.20]). *c and c' are asymptotic if and only if $b_c - b_{c'}$ is a constant function.*

The Busemann function b_c is extended for a constant-speed ray $c : t \mapsto c^*(rt)$ by $b_c := rb_{c^*}$, where c^* is a geodesic ray. Fix $x_0 \in X$. For $p \in CX^\infty$, there is a unique ray c issuing x_0 with $c(\infty) = p$, and hence b_c is also written as $b_{x_0, p} = b_p$. In this way, we can identify CX^∞ with the space of Busemann functions.

Example 2.7. In the case of $X = \mathbb{R}^n$, the Busemann function b_c for a geodesic ray $c(t) = \xi t + x_0$ is given by

$$b_c(x) = -\langle \xi, x - x_0 \rangle.$$

This follows from $\|x - \xi t - x_0\|_2 - t = t - \langle \xi, x - x_0 \rangle + O(1/t) - t$. Thus, Busemann functions (for constant-speed rays) are precisely affine functions. If the origin 0 is chosen as x_0 , the space $CX^\infty (\simeq \mathbb{R}^n)$ is identified with the space of linear functions $x \mapsto -\langle p, x \rangle$, i.e., the dual space of \mathbb{R}^n .

For a (convex) function $f : X \rightarrow \mathbb{R}$, define the *asymptotic Legendre-Fenchel conjugate* $f^* : CX^\infty \rightarrow \mathbb{R} \cup \{\infty\}$ of f by

$$f^*(p) := \sup_{x \in X} -b_p(x) - f(x) \quad (p \in CX^\infty). \quad (2.16)$$

Note that this notion of Legendre-Fenchel conjugate is rather different from those introduced by [4, 36]. If $X = \mathbb{R}^n$, then this matches the usual Legendre-Fenchel conjugate (see Examples 2.4 and 2.7). Notice that f^* is not necessarily convex in CX^∞ . We give a simple lemma providing explicit examples of asymptotic Legendre-Fenchel conjugates.

Lemma 2.8. *For a function $h : \mathbb{R}_+ \rightarrow \mathbb{R}$, define $f : X \rightarrow \mathbb{R}$ by*

$$f(x) := h(d(x, x_0)) \quad (x \in X).$$

Then the asymptotic Legendre-Fenchel conjugate f^ is given by*

$$f^*(p) := h^*(\|p\|) \quad (p \in CX^\infty),$$

where h^ is the Legendre-Fenchel conjugate of h , i.e., $h^*(\xi) := \sup_{r \in \mathbb{R}} \xi r - h(r)$ with $h(r) := \infty$ for $r < 0$.*

Proof. By definitions of f^* and b_p , we have

$$f^*(p) = \sup_{x \in X} -\|p\| \lim_{t \rightarrow \infty} (d(x, c(t)) - t) - h(d(x, x_0)),$$

where c is a geodesic ray with $c(0) = x_0$ and $c(\infty) = p/\|p\|$. For $x \in X$, letting $r := d(x, x_0)$, we have $h(d(x, x_0)) = h(r)$, and $d(x, c(t)) \geq d(c(t), x_0) - d(x, x_0) = t - r = d(c(t), c(0))$ for every large t . This implies that sup can be restricted to c . Thus $f^*(p) = \sup_{r \in \mathbb{R}_+} \|p\| r - h(r) = h^*(\|p\|)$. \square

Example 2.9. Suppose that $f(x) = \frac{1}{2}d(x, x_0)^2$. By the CAT(0)-inequality (2.2), f is (strongly) convex. By the above lemma, we have $f^*(p) = \frac{1}{2}\|p\|^2$. In this case, f^* is also convex. The convexity is seen by considering the flat triangle of vertices $0, p, q$.

Further investigation of f^* is left for future research. As mentioned in the introduction, our central interest is how to describe the domain $\text{dom } f^* = \{p \in CX^\infty \mid f^*(p) < \infty\}$ of the conjugate f^* .

2.1.3 Recession functions (asymptotic slope functions)

Let $f : X \rightarrow \mathbb{R}$ be a continuous convex function. The *recession function* $f^\infty : CX^\infty \rightarrow \mathbb{R} \cup \{\infty\}$ of f is defined by

$$f^\infty(p) := \lim_{t \rightarrow \infty} \{f(c(t)) - f(c(0))\}/t = \lim_{t \rightarrow \infty} f(c(t))/t \quad (p \in CX^\infty) \quad (2.17)$$

where c is a constant-speed ray with $c(\infty) = p$. The recession function is just a homogeneous extension of the *asymptotic slope function* by Kapovich, Leeb, and Millson [31], which is defined on X^∞ . By convexity, $(f(c(t)) - f(c(0)))/t$ is monotone nondecreasing, and it converges to a finite value or ∞ . The recession function is indeed independent of the choice of a ray c .

Lemma 2.10 ([33, Lemma 2.10]). $\lim_{t \rightarrow \infty} f(c(t))/t = \lim_{t \rightarrow \infty} f(c'(t))/t$ holds for two asymptotic geodesic rays c, c' .

It is easy to see the positive homogeneity of f^∞ (by the change of variable in (2.17)). In particular, $f^\infty(r\xi) = rf^\infty(\xi)$ for $\xi \in X^\infty$. A partial convexity property of asymptotic slope functions is obtained in [31, Lemma 3.2 (ii)]. In the setting of recession functions the following general convexity holds.

Theorem 2.11. *The recession function f^∞ is positively homogeneous convex.*

Proof. We have already seen the positive homogeneity. Hence it suffices to show

$$f^\infty(p) + f^\infty(p') \geq f^\infty(p + p') \quad (p, p' \in CX^\infty).$$

Let $p, p' \in CX^\infty$. We can assume that both p and p' are nonzero and both $f^\infty(p)$ and $f^\infty(p')$ are finite. Suppose that $p = a\xi$ and $p' = a'\xi'$ for $a \geq a' > 0$ and $\xi, \xi' \in X^\infty$. Let $x \in X$ and let σ, σ' be geodesic rays issuing from x with $\sigma(\infty) = \xi$ and $\sigma'(\infty) = \xi'$.

Suppose first that $p + q \neq 0$. Then $\angle(\xi, \xi') < \pi$, or $\angle(\xi, \xi') = \pi$ and $a > a'$. Let $m(t)$ be the midpoint of $\sigma(at)$ and $\sigma'(a't)$. We will consider geodesic segment $[x, m(t)]$ with $t \rightarrow \infty$. We show $e := \lim_{t \rightarrow \infty} d(x, m(t))/t > 0$. Indeed, by triangle inequality $d(x, m(t)) \geq d(x, \sigma(at)) - d(\sigma(at), m(t))$ with $d(x, \sigma(at)) = at$ and $d(\sigma(at), m(t)) = d(\sigma(at), \sigma'(a't))/2$ we have

$$d(x, m(t))/t \geq a - d(\sigma(at), \sigma'(a't))/2t = a - \frac{1}{2} \sqrt{a^2 + a'^2 - 2aa' \cos \bar{Z}_x(\sigma(at), \sigma'(a't))},$$

where the equality follows from the law of cosine in the comparison triangle of $x, \sigma(at), \sigma(a't)$. Letting $t \rightarrow \infty$, we have $\bar{Z}_x(\sigma(at), \sigma'(a't)) \rightarrow \angle(\xi, \xi')$ and

$$e \geq a - \frac{1}{2} \sqrt{a^2 + a'^2 - 2aa' \cos \angle(\xi, \xi')} \geq (a - a')/2.$$

By $a > a'$ or $\angle(\xi, \xi') < \pi$, we have $e > 0$. By [6, II.4.4], it holds $p + p' = 2e\eta$ for some $\eta \in X^\infty$. Let ρ denote the geodesic ray with $\rho(0) = x$ and $\rho(\infty) = \eta$.

We can assume $f(x) = 0$ by replacing f with $f - f(x)$. Consider the constant-speed path $\rho_t : [0, et] \rightarrow X$ with $\rho(0) = x$ and $\rho(et) = m(t)$. Then, by [6, II.4.4], the path ρ_t converges to geodesic ray ρ for $t \rightarrow \infty$. For every $\epsilon > 0$ and $s \geq 0$, there is t_0 such that for every $t \geq t_0$ it holds

$$f(\rho(s)) \leq \epsilon + f(\rho_t(s)) \leq \epsilon + f(m(t))d(x, \rho_t(s))/d(x, m(t)),$$

where the second inequality follows from $f(x) = 0$ and the convexity of f along $[x, m(t)]$. For all large s, t we have

$$\frac{f(\sigma(at))}{t} + \frac{f(\sigma'(a't))}{t} \geq 2\frac{f(m(t))}{t} \geq 2\left(\frac{f(\rho(s))}{s} - \frac{\epsilon}{s}\right) \frac{s}{d(x, \rho_t(s))} \frac{d(x, m(t))}{t}$$

By $t \rightarrow \infty$, we have $d(x, \rho_t(s)) \rightarrow s$ and $d(x, m(t))/t \rightarrow e$. By $s \rightarrow \infty$, we have

$$f^\infty(p) + f^\infty(p') \geq \lim_{s \rightarrow \infty} f(\rho(s))2e/s = 2ef^\infty(\eta) = f^\infty(p + p').$$

Finally, consider the case $p + p' = 0$, i.e., $\angle(\xi, \xi') = \pi$ and $a = a'$. For $\epsilon > 0$, it holds $p + (1 + \epsilon)p' \neq 0$. Then, from the above case, we have

$$f^\infty(p) + (1 + \epsilon)f^\infty(p') = f^\infty(p) + f^\infty((1 + \epsilon)p') \geq f^\infty(p + (1 + \epsilon)p') = \epsilon f^\infty(p').$$

For the last equality, observe from (2.6) that the midpoint between $p, (1 + \epsilon)p'$ is $(\epsilon/2)p'$. Hence $p + (1 + \epsilon)p' = \epsilon p'$ and $f^\infty(\epsilon p') = \epsilon f^\infty(p')$. By $\epsilon \rightarrow 0$, we obtain the desired inequality. \square

It is known [31, p. 318] that the asymptotic slope of a Busemann function is given by $b_\xi(\eta)^\infty = -\cos \angle(\xi, \eta)$ for $\xi, \eta \in X^\infty$. Hence we have:

Lemma 2.12. *It holds $(b_p)^\infty(q) = -\langle p, q \rangle$ for $p, q \in CX^\infty$.*

Motivated by (1.3), for any positively homogeneous function $h : CX^\infty \rightarrow \mathbb{R} \cup \{\infty\}$ we define a subset $B(h) \subseteq CX^\infty$ by

$$B(h) := \{p \in CX^\infty \mid \langle u, p \rangle \leq h(u) \ (u \in X^\infty \subseteq CX^\infty)\}. \quad (2.18)$$

Since $p \mapsto \langle u, p \rangle$ is continuous (Lemma 2.3), we have:

Lemma 2.13. *$B(h)$ is a closed subset in CX^∞ .*

We mainly consider $B(f^\infty)$ for recession function f^∞ . Belonging to $B(f^\infty)$ is a necessary condition for $p \in CX^\infty$ for which $f + b_p$ is bounded below.

Lemma 2.14. $\text{dom } f^* \subseteq B(f^\infty)$.

Proof. Suppose that $\langle u, p \rangle > f^\infty(u)$ for some $u \in X^\infty$. Then $f^\infty(u) - \langle u, p \rangle = (f + b_p)^\infty(u) = \lim_{t \rightarrow \infty} (f(c(t)) + b_p(c(t)))/t$ is negative, where c is a ray with $c(\infty) = u$. Therefore, for some $\alpha > 0$, it holds $f(c(t)) + b_p(c(t)) < -\alpha t$ for every large $t > 0$. This means that $\inf_{x \in X} f(x) + b_p(x) = -\infty$, and $p \notin \text{dom } f^*$. \square

For a subset $R \subseteq CX^\infty$, let \overline{R} denote the closure of R with respect to the d^∞ -topology. By the above lemma, it holds $\overline{\text{dom } f^*} \subseteq B(f^\infty)$. We do not know whether the equality holds in general.




Figure 1: The boundary of hyperbolic space

2.2 Hadamard manifolds

From here, we restrict our study to smooth convex optimization on a *Hadamard manifold*, i.e., a simply-connected complete Riemannian manifold having nonpositive sectional curvature. We utilize elementary concepts in Riemannian geometry; see e.g., [42]. A recent book [9] for optimization perspectives is also useful. For Hadamard manifolds, we consult [7, 17].

Let M be an n -dimensional Hadamard manifold. For $x \in M$, let $\langle \cdot, \cdot \rangle_x$ denote the inner product of the tangent space T_x . Let d denote the distance function on M obtained from the Riemannian connection. The metric space M is known to be a Hadamard space. A geodesic ray issuing from x is given by the *exponential map* $\exp_x : T_x \rightarrow M$. Namely, for $v \in T_x$, the map $t \mapsto \exp_x(vt)$ is a constant-speed ray with speed $\|v\|_x := \sqrt{\langle v, v \rangle_x}$. The map \exp_x is a diffeomorphism from T_x to M .

Consider the boundary M^∞ and its Euclidean cone CM^∞ of M . Via the exponential map, the boundary M^∞ is identified with the unit sphere at T_x , and the cone CM^∞ is identified with $T_x \simeq \mathbb{R}^n$. This identification gives another topology to M^∞ and to CM^∞ , which is called the *standard topology*, and is independent of the choice of x . In the standard topology, M^∞ and CM^∞ are homeomorphic to sphere S^{n-1} and Euclidean space \mathbb{R}^n , respectively. It is known [11, II.9.7(1)] that the identity map on CM^∞ from the d^∞ -topology to the standard topology is continuous (and is not homeomorphic in general).

Example 2.15. Suppose that M is a hyperbolic space. For distinct $\xi, \eta \in M^\infty$ there is a geodesic line $c : \mathbb{R} \rightarrow M$ such that $c(\infty) = \xi$ and $c(-\infty) = \eta$. This means that $\angle(\xi, \eta) = \pi$ for every distinct $\xi, \eta \in M^\infty$. Therefore, the boundary M^∞ is a discrete topological space. See [11, II.9.6 (ii)]. The cone CM^∞ is a *star* obtained from infinitely many half-lines \mathbb{R}_+ , each associated with $\xi \in M^\infty$, by identifying the origin of all \mathbb{R}_+ . See Figure 1. This can be seen from the distance formula (2.6) as $d^\infty(r\xi, s\eta) = r + s$ if $\xi \neq \eta$ and $|r - s|$ if $\xi = \eta$.

Let $f : M \rightarrow \mathbb{R}$ be a smooth convex function.

Lemma 2.16. (1) f^∞ is lower semicontinuous in the standard topology [33, Lemma 2.11].

(2) $(p, q) \mapsto \langle p, q \rangle$ is upper semicontinuous in the standard topology.

Proof. (1). Take an arbitrary $x \in M$, and identify CM^∞ with T_x (by $v \mapsto \exp_x v \infty$). Let $f_t(p) := (f(\exp_x(pt)) - f(x))/t$ for $p \in T_x$, which is monotone nondecreasing in t and continuous on T_x (in the standard topology). Therefore $f^\infty(p) = \sup_{t \geq 0} f_t(p)$. It is well-known that the supremum of continuous functions f_t is lower semicontinuous.

(2). It is known [11, II.9.5] that $(\xi, \eta) \mapsto \angle(\xi, \eta)$ is lower semicontinuous. Then $(p, q) \mapsto \langle p, q \rangle$ is upper semicontinuous, since $p \mapsto d^\infty(0, p) = \|p\|_x$ is continuous in the standard topology and cosine is a decreasing function. \square

As in the Euclidean case, a minimizer of f is characterized by the gradient. The *gradient* $\nabla f(x) \in T_x$ of f at x is defined via

$$\langle \nabla f(x), v \rangle_x = df(x)(v) = \lim_{t \rightarrow 0} \{f(\exp_x tv) - f(x)\}/t \quad (v \in T_x),$$

where $df(x) : T_x \rightarrow \mathbb{R}$ is the differential of f at x . It is easy to see:

Lemma 2.17 (see [9, Corollary 11.22]). *$x \in M$ is a minimizer of f if and only if $\nabla f(x) = 0$.*

In Euclidean case $M = \mathbb{R}^n$, x is a minimizer of $x \mapsto f(x) - \langle p, x \rangle$ if and only if $\nabla f(x) = p$. To extend it, we consider the gradient of Busemann functions. Note that Busemann functions are continuously differentiable; see [17, 1.10.2 (1)]. As in the previous subsection, we fix $x_0 \in M$ and identify CM^∞ with the space of Busemann functions for rays issuing from x_0 .

Lemma 2.18 (see [17, 1.10.2 (2)]). *For $x \in M$ and $p \in CM^\infty$, it holds $\nabla b_p(x) = -u$ for $u \in T_x$ with $p = \exp_x \infty u$.*

The *asymptotic gradient* $\nabla^\infty f(x)$ for $x \in M$ is defined as the asymptotic class of the ray $t \mapsto \exp_x t \nabla f(x)$, that is,

$$\nabla^\infty f(x) := \exp_x \infty \nabla f(x).$$

Notice that $\|\nabla^\infty f(x)\| = d^\infty(0, \nabla^\infty f(x))$ is the speed of $t \mapsto \exp_x t \nabla f(x)$. Therefore we have

$$\|\nabla^\infty f(x)\| = \|\nabla f(x)\|_x. \quad (2.19)$$

Then we have the following analogue of the one in Euclidean convex analysis.

Lemma 2.19. *For $p \in CM^\infty$, the following conditions are equivalent:*

- (i) x is a minimizer of $f + b_p$ over M .
- (ii) $\nabla^\infty f(x) = p$.
- (iii) $f(x) + f^*(p) = -b_p(x)$

Proof. (i) $\Leftrightarrow \nabla(f + b_p)(x) = 0 \Leftrightarrow \nabla f(x) = u$ for $p = \exp_x \infty u \Leftrightarrow$ (ii). (i) \Leftrightarrow (iii) is obvious from the definition (2.16) of conjugate f^* . \square

As in the Euclidean case, the conjugate of a (smooth) convex function recovers the original function via the inverse transformation.

Lemma 2.20. $f(x) = \sup_{p \in CM^\infty} -b_p(x) - f^*(p) \quad (x \in M)$.

Proof. By definition, $f^*(p) \geq -b_p(x) - f(x)$ for every $x \in M$ and $p \in CM^\infty$. Therefore $f(x) \geq \sup_{p \in CM^\infty} -b_p(x) - f^*(p)$. For $p = \nabla^\infty f(x)$, the equality is attained by Lemma 2.19. \square

This gives rise to an interesting question of characterizing the class of functions g on CM^∞ for which $g^*(x) := \sup_{p \in CM^\infty} -b_p(x) - g(p)$ is convex on M . However this is beyond the theme of this paper, and we leave it for future research.

For a set $R \subseteq CM^\infty$, let $\text{int } R$ denote the interior of R in the d^∞ -topology.

Lemma 2.21. *Let $h : CM^\infty \rightarrow \mathbb{R} \cup \{\infty\}$ be a positively homogeneous function. Suppose that h is lower semicontinuous in the standard topology. Then it holds*

$$\text{int } B(h) = \{p \in CM^\infty \mid \langle \xi, p \rangle < h(\xi) \ (\xi \in M^\infty \subseteq CM^\infty)\}. \quad (2.20)$$

Proof. Let $p \in B(h)$. Suppose that $\langle \xi, p \rangle = h(\xi)$ for some $\xi \in M^\infty$. Then, for arbitrary $\epsilon > 0$, $\langle \xi, p + \epsilon \xi \rangle \geq \langle \xi, p \rangle + \epsilon \|\xi\|^2 > h(\xi)$ (Lemma 2.3). This means $p + \epsilon \xi \notin B(h)$. Since $\langle \cdot, \cdot \rangle$ is continuous (Lemma 2.3), p is never an interior point of $B(h)$.

Suppose that $\langle \xi, p \rangle < h(\xi)$ for all $\xi \in M^\infty$. For $q \in CM^\infty$, let $\Delta(q) := \inf_{\xi \in M^\infty} h(\xi) - \langle \xi, q \rangle$. Since M^∞ is compact and $\xi \mapsto h(\xi) - \langle \xi, q \rangle$ is lower semicontinuous in the standard topology (Lemma 2.16 (2)), the infimum is always attained. Let $\epsilon := \Delta(p) > 0$. Since $(q, \xi) \mapsto h(\xi) - \langle \xi, q \rangle$ is lower semicontinuous, for each $\xi \in M^\infty$ there is an open neighborhood $U_\xi \times V_\xi$ of (p, ξ) (in the standard topology) such that for each $(q, \eta) \in U_\xi \times V_\xi$ we have $h(\eta) - \langle \eta, q \rangle \geq h(\xi) - \langle \xi, p \rangle - \epsilon \geq \Delta(p) - \epsilon = 0$. Since M^∞ is compact, there are $\xi_1, \xi_2, \dots, \xi_m$ with $M^\infty = \bigcup_{i=1}^m V_{\xi_i}$. Let $U := \bigcap_{i=1}^m U_{\xi_i}$, which is an open neighborhood of p (in the standard topology). For any $q \in U$, $\Delta(q) = h(\xi') - \langle \xi', q \rangle$ for some ξ' . Since ξ' belongs to V_{ξ_i} for some i , it holds $\Delta(q) = h(\xi') - \langle \xi', q \rangle \geq 0$. Hence, we have $p \in U \subseteq B(h)$. Since the identify map on CM^∞ from the d^∞ -topology to the standard topology is continuous, U is an open neighborhood of p in d^∞ -topology. \square

As the proof shows, the RHS of (2.20) is the interior of $B(h)$ also in the standard topology, although $B(h)$ may not be closed in this topology.

We apply this lemma to the recession function f^∞ and the associated subset $B(f^\infty)$.

Proposition 2.22. *For $p \in \text{int } B(f^\infty)$, there is a minimizer of $f + b_p$. In particular, any point in $B(f^\infty) \setminus \text{dom } f^*$ belongs to the boundary of $B(f^\infty)$.*

See [31, Lemma 3.2 (iv)] for a related argument.

Proof. Since $\xi \mapsto f^\infty(\xi) - \langle \xi, p \rangle$ is lower semicontinuous on compact set M^∞ in the standard topology, the minimum value $\alpha > 0$ exists. Let $\alpha' \in (0, \alpha)$ and let $g := f + b_p$. Fix an arbitrary $x \in M$. Then, for every $\xi \in M^\infty$ there is $t_\xi \in \mathbb{R}_+$ such that $g(\exp_x t_\xi) - g(x) > \alpha' t_\xi$ for all $t \geq t_\xi$. Define $h : M^\infty \rightarrow \mathbb{R}_+$ by

$$h(\xi) := \inf\{t \geq 0 \mid g(\exp_x t\xi) - g(x) > \alpha't\} \quad (\xi \in M^\infty).$$

Then h is upper semicontinuous, since the epigraph $\{(\xi, t) \in M^\infty \times \mathbb{R} \mid t \leq h(\xi)\}$ is the closed set $\{(\xi, t) \in M^\infty \times \mathbb{R}_+ \mid g(\exp_x t\xi) - g(x) \leq \alpha't\} \cup \{(\xi, t) \in M^\infty \times \mathbb{R} \mid t \leq 0\}$. Since M^∞ is compact, the maximum t^* of h over M^∞ exists. Then, for every $\xi \in M^\infty$, we have $g(\exp_x t\xi) - g(x) \geq \alpha't$ for all $t \geq t^*$. This means that the level set $\{y \in M \mid g(y) \leq g(x) + \alpha't^*\}$ belongs to the metric ball at center x with radius t^* , which is compact by Hopf-Rinow theorem (see [42, III.1]). A minimizer of g exists in this set. \square

Thus we have

$$\text{int } B(f^\infty) \subseteq \nabla^\infty f(M) \subseteq \text{dom } f^* \subseteq B(f^\infty). \quad (2.21)$$

We next provide a characterization of $B(f^\infty)$, which sharpens the following important result by Kapovich, Leeb, and Millson [31].

Theorem 2.23 ([31, Lemma 3.4]). *If $f^\infty(u) \geq 0$ for all $u \in M^\infty$, i.e., $0 \in B(f^\infty)$, then $\inf_{x \in M} \|\nabla f(x)\|_x = 0$.*

An outline of the proof is as follows: If $\inf_{x \in M} \|\nabla f(x)\|_x > 0$, then a trajectory of the normalized gradient flow of f goes to $u \in M^\infty$ with $f^\infty(u) < 0$; See also [46, section 5.4].

We now obtain an analogue of the equivalence between (a) and (c) in the introduction.

Theorem 2.24. *For $p \in CM^\infty$, the following are equivalent:*

$$(a) \inf_{x \in M} \|\nabla(f + b_p)(x)\|_x = 0.$$

$$(c) p \in B(f^\infty).$$

Proof. (c) \Rightarrow (a) follows from applying the above theorem to $f + b_p$. We verify (a) \Rightarrow (c). Let $g := f + b_p$. For arbitrary $\epsilon > 0$, there is $x \in M$ such that $\|\nabla g(x)\|_x < \epsilon$. Consider $u \in M^\infty$ and the geodesic ray $t \mapsto \exp_x vt$ with $\exp_x v\infty = u$. Then $\lim_{t \rightarrow 0} (g(\exp_x vt) - g(x))/t = \langle \nabla g(x), v \rangle_x = \|\nabla g(x)\|_x \cos \theta > -\epsilon$, where θ is the angle between $\nabla g(x)$ and v in T_x . Since $(g(\exp_x vt) - g(x))/t$ is monotone nondecreasing, by $t \rightarrow \infty$ we have $g^\infty(u) = f^\infty(u) - \langle u, p \rangle > -\epsilon$. Thus, for every $\epsilon > 0$ and $u \in M^\infty$, it holds $\langle u, p \rangle < f^\infty(u) + \epsilon$. This implies $p \in B(f^\infty)$. \square

The condition (a) $\inf_{x \in M} \|\nabla(f + b_p)(x)\|_x = 0$ may be viewed as a correspondent of $\inf_{x \in \mathbb{R}^n} \|\nabla f(x) - p\| = 0$ of the Euclidean case. If $\nabla^\infty f(x_i)$ ($i = 1, 2, \dots$) converges to $p \in CM^\infty$ in the d^∞ -topology, then (a) holds. Indeed, by Lemma 2.18 and (2.19) it holds

$$\begin{aligned} \|\nabla(f + b_p)(x)\|_x^2 &= \|\nabla f(x)\|_x^2 + \|u\|_x^2 - 2\|\nabla f(x)\|_x \|u\|_x \cos \theta_x \\ &\leq \|\nabla^\infty f(x)\|^2 + \|p\|^2 - 2\|\nabla^\infty f(x)\| \|p\| \cos \theta \\ &= d^\infty(\nabla^\infty f(x), p)^2, \end{aligned}$$

where $u := -\nabla b_p(x)$, θ_x is the angle between $\nabla f(x)$ and u in T_x , and $\theta := \angle(\nabla^\infty f(x), p)$ ($\geq \theta_x$ by definition (2.5)). However, the converse is not true. Also (a) does not mean the convergence in the standard topology.

2.3 Symmetric spaces of nonpositive curvature

Here we consider symmetric spaces of nonpositive curvature, which constitute a fundamental class of Hadamard manifolds. Our argument basically consults [17, Chapters 2 and 3]. In a Hadamard manifold M and a point x , the *geodesic symmetry* $\sigma_x : M \rightarrow M$ at x is defined by $\sigma_x(\exp_x(v)) = \exp_x(-v)$ for $v \in T_x$. A *symmetric space of nonpositive curvature* is a Hadamard manifold M such that for every $x \in M$ the geodesic symmetry σ_x is an isometry on M . Via the de Rham decomposition theorem (see [42, III.6]), M is (uniquely) decomposed as Riemannian product $M = M_0 \times N$, where M_0 is isometric

to Euclidean space \mathbb{R}^k ($k \geq 0$) (called the Euclidean de Rham factor of M) and a symmetric space N of noncompact type, i.e., it is given by $N = G/K$ for a (real) semisimple Lie group G and its maximal compact subgroup K . Then N has a trivial Euclidean de Rham factor, and its Riemannian structure is given by a G -invariant metric. We will see more concrete constructions in Sections 2.4 and 3.2.

Let M be a symmetric space of nonpositive curvature, and let $x_0 \in M$. A k -dimensional *flat* is a submanifold of M isometric to \mathbb{R}^k . A *maximal flat* is a flat that is not contained in another flat of a larger dimension. It is a basic fact that all maximal flats have the same dimension d , which is called the *rank* of M . By a *geodesic line* we mean a map $l : \mathbb{R} \rightarrow M$ with $d(l(s), l(t)) = |s - t|$ for $s, t \in \mathbb{R}$, which is just a 1-dimensional flat. A geodesic line is called *regular* if it is contained by a unique maximal flat. Let $x \in M$ and let F be a maximal flat containing x . A *Weyl chamber* at tip x is the closure of a connected component of the set of points $y \in F \setminus \{x\}$ such that the unique geodesic line containing x, y is regular. When F is viewed as \mathbb{R}^d with origin x , Weyl chambers are polyhedral cones. They have the same shape, since the group G acts transitively on them.

We next explain the building structure of the boundary M^∞ and its cone CM^∞ ; [7, Appendix 5] is a useful reference. It is clear (from Example 2.4) that the boundary F^∞ of a maximal flat F is isometric to sphere S^{d-1} . The boundary C^∞ of a Weyl chamber C (at some tip) is called an (asymptotic) *Weyl chamber*. For two Weyl chambers C, D (at possibly different tips), the asymptotic ones C^∞, D^∞ are the same or have disjoint interiors. Then, the set of all Weyl chambers and their faces give rise to a cell-complex structure on M^∞ , where each cell is isometric to a polyhedral cell in a sphere (the intersection of a sphere and a polyhedral cone). This structure is an M_1 -*polyhedral complex* in the sense of [11, Chapter I.7], and forms a *spherical building*, where an *apartment* is precisely the subcomplex formed by the boundary of a maximal flat. Although Weyl chambers here may not be simplices, one can subdivide them to obtain a simplicial complex \mathcal{C} so that each apartment is a spherical Coxeter complex. Then the apartments are glued nicely. That is, they satisfy, as an abstract simplicial complex, the axiom of building:

- Any two simplices in \mathcal{C} are contained in a common apartment.
- For two apartments A, A' including simplices C, C' , there is an isomorphism $A \rightarrow A'$ fixing C, C' pointwise.

See e.g., [1] and [11, II.10. Appendix] for (formal) theory of building. By considering the Euclidean cone, CM^∞ has the structure of a *Euclidean building*. A maximal cone is also called a Weyl chamber. Apartments are the subcomplexes induced by CF^∞ for all maximal flats F . We simply call CF^∞ an apartment. Each apartment is a convex subspace of CM^∞ isometric to \mathbb{R}^d . We can identify an apartment $E \subseteq CM^\infty$ with \mathbb{R}^d so that the origins coincide. In this identification, \langle, \rangle in E is precisely the Euclidean inner product of \mathbb{R}^d .

Suppose that $M = \mathbb{R}^k \times N$, where N is a symmetric space G/K of noncompact type. Then $CM^\infty = \mathbb{R}^k \times CN^\infty$. The structure of building CM^∞ is determined by CN^∞ . We can suppose (as in [17]) that G is the identity component of the group of isometries of N , where G acts isometrically on $M = \mathbb{R}^k \times N$ by $\mathbb{R}^k \times N \ni (x', x) \mapsto (x', gx)$. Since any isometry on M induces an isometry on M^∞ , the group G acts isometrically on M^∞ and on CM^∞ by $gp = \|p\|g(p/\|p\|)$. Also G acts on the set of Weyl chambers

and their faces. The facial incidence structure of the building M^∞ is described by the inclusion relation of all parabolic subgroups of G . Here a *parabolic subgroup* is the subgroup consisting of $g \in G$ with $g\xi = \xi$ for some $\xi \in N^\infty$, which is denoted by G_ξ . If ξ, ξ' belong to the relative interior of the same face of a Weyl chamber, then $G_\xi = G_{\xi'}$. Therefore, minimal parabolic subgroups correspond to Weyl chambers. Fix a Weyl chamber C_0 , and regard it as a polyhedral cone in \mathbb{R}^d . Let P be the minimal parabolic subgroup for C_0 . The set of Weyl chambers is identified with the flag variety G/P . If $p \in CM^\infty$ belongs to a Weyl chamber corresponding to $\mathcal{F} \in G/P$ and the orbit of $p \in CM^\infty$ by G -action meets a (unique) point λ in $C_0 \subseteq \mathbb{R}^d$, then we denote p by

$$p = \lambda \cdot \mathcal{F}. \quad (2.22)$$

This notation designates the coordinate λ of p in the Weyl chamber indexed by \mathcal{F} .

Two geodesic lines $c, c' : \mathbb{R} \rightarrow M$ are said to be *parallel* if $d(c(t), c'(t))$ ($t \in \mathbb{R}$) is bounded above. For a geodesic line c , let $F(c)$ denote the union of all geodesic lines parallel to c . It is known that $F(c)$ is a totally geodesic submanifold of M . If c is regular, $F(c)$ is a maximal flat.

For $\xi \in N^\infty$, the *horospherical subgroup* N_ξ of G_ξ consists of $n \in G$ such that $\lim_{t \rightarrow \infty} d(n(c(t)), c(t)) = 0$, where c is the geodesic ray with $c(\infty) = \xi$ and $c(0) = x_0$ (independent of x_0). Then N_ξ keeps the Busemann function b_ξ as $b_\xi(nx) = b_\xi(x)$, since $|b_\xi(nx) - b_\xi(x)| = \lim_{t \rightarrow \infty} |d(x, n^{-1}c(t)) - d(x, c(t))| \leq \lim_{t \rightarrow \infty} d(c(t), n^{-1}c(t)) = 0$. The *generalized Iwasawa decomposition* [17, 2.17.5 (5)] implies that M is diffeomorphic to $N_\xi \times F(c)$ by $(n, y) \mapsto n(y)$.

In this setting, let us start our convex analysis on M . Let $f : M \rightarrow \mathbb{R}$ be a smooth convex function. For a maximal flat F , let $f_F : F \rightarrow \mathbb{R}$ denote the restriction of f to F . The asymptotic Legendre-Fenchel conjugate $f_F^* : CF^\infty \rightarrow \mathbb{R} \cup \{\infty\}$ is defined by $f_F^*(p) := \sup_{x \in F} -b_p(x) - f(x)$, where x_0 may be outside of F . By $p \in CF^\infty$ and Example 2.7, b_p is an affine function on F . Therefore, f_F^* is viewed as the ordinary Legendre-Fenchel conjugate (up to an additive constant) under identification $CF^\infty = \mathbb{R}^d$. Hence we have:

Lemma 2.25. *For any maximal flat F , f_F^* is a convex function on CF^∞ .*

Still, f^* may be nonconvex but “partially” convex in the following sense:

Proposition 2.26. *For a Weyl chamber $C \subseteq CM^\infty$, it holds*

$$f^*(p) = \sup_F f_F^*(p) \quad (p \in C), \quad (2.23)$$

where \sup is taken over all maximal flats F with $CF^\infty \supseteq C$. In particular, f^* is a convex function on C , and $C \cap \text{dom } f^*$ is a convex set contained by a convex set $\bigcap_F C \cap \text{dom } f_F^*$:

$$C \cap \text{dom } f^* \subseteq \bigcap_F C \cap \text{dom } f_F^*. \quad (2.24)$$

Note that \subseteq may be strict since $f^*(p) = \sup_F f_F^*(p)$ may be ∞ even if $f_F^*(p) < \infty$ for every F .

Proof. Consider the Iwasawa decomposition $M = N \times F_0$, where F_0 is a maximum flat containing C at infinity and N is the holospherical subgroup for C . Then NF_0 ranges

over all maximal flats containing C at infinity. Thus M is the (disjoint) union of all maximal flats F such that $F^\infty \supseteq C \ni p$. Then we have $f^*(p) = \sup_F \sup_{x \in F} -b_p(x) - f(x) = \sup_F f_F^*(p)$. \square

Next, we consider the recession function f^∞ and the associated subset $B(f^\infty)$. For each apartment E , we also consider

$$B_E(f^\infty) := \{p \in E \mid \langle u, p \rangle \leq f^\infty(u) \ (u \in E)\}. \quad (2.25)$$

Proposition 2.27. *For a Weyl chamber $C \subseteq E$, it holds*

$$C \cap B(f^\infty) = \bigcap_F C \cap B_{CF^\infty}(f^\infty) = \bigcap_F C \cap \overline{\text{dom } f_F^*}. \quad (2.26)$$

where \bigcap is taken over all maximal flats F with $CF^\infty \supseteq C$. In particular, $C \cap B(f^\infty)$ is a convex set in C .

Proof. Since CM^∞ is the union of all apartments E containing any fixed Weyl chamber C , we have the first equality. When E is viewed as Euclidean space \mathbb{R}^n , then C is a convex cone, $\langle u, p \rangle \leq f^\infty(u)$ is a linear inequality, and therefore $C \cap B_E(f^\infty)$ is convex. Necessarily the intersection $\bigcap_E C \cap B_E(f^\infty)$ is convex. By Lemma 2.16, the restriction of f^∞ to Euclidean space E is lower semicontinuous and positively homogeneous convex, and must be the support function of $\text{dom } f_F^*$. Hence we have $\overline{\text{dom } f_F^*} = B_E(f^\infty)$, and the second equality. \square

In the view of (2.24) and (2.26), the closure $\overline{\text{dom } f^*}$ seems very close to $B(f^\infty)$, although we do not know whether they really differ. Under non-degeneracy assumptions, three spaces $\nabla^\infty f(M)$, $\overline{\text{dom } f^*}$, and $B(f^\infty)$ are equal, as in Euclidean case.

Proposition 2.28. *For a Weyl chamber C , if $C \cap B(f^\infty)$ has nonempty interior, then $C \cap \nabla^\infty f(M) = C \cap \overline{\text{dom } f^*} = C \cap B(f^\infty)$.*

Proof. By (2.21), it holds

$$C \cap \text{int } B(f^\infty) \subseteq C \cap \overline{\nabla^\infty f(M)} \subseteq C \cap \overline{\text{dom } f^*} \subseteq C \cap B(f^\infty). \quad (2.27)$$

In Euclidean space, for any closed convex set D having nonempty interior, it holds $\overline{\text{int } D} = D$. This can be applied to $C \cap B(f^\infty)$, which is viewed as a Euclidean convex set. From $C \cap B(f^\infty) = \overline{\text{int}(C \cap B(f^\infty))} \subseteq \overline{C \cap \text{int } B(f^\infty)} \subseteq C \cap B(f^\infty)$, by taking the closure in (2.27), we have the claim. \square

Proposition 2.29. *Suppose that f is strictly convex. Then the gradient map $x \mapsto \nabla^\infty f(x)$ is a bijection from M to $\text{int } B(f^\infty)$. For a Weyl chamber C , if $C \cap \nabla^\infty f(M) \neq \emptyset$, then $C \cap \overline{\nabla^\infty f(M)} = C \cap \overline{\text{dom } f^*} = C \cap B(f^\infty)$.*

Proof. We first verify $\nabla^\infty f : M \rightarrow CM^\infty$ is injective. Indeed, if $\nabla^\infty f(x) = \nabla^\infty f(y) = p$, then x and y are minimizers of $f + b_p$, and it must hold $x = y$ by strict convexity of $f + b_p$. For surjectivity, from Proposition 2.22 it suffices to show that $\nabla^\infty f(x)$ for any $x \in M$ belongs to $\text{int } B(f^\infty)$. We utilize the following property, where C is a Weyl chamber.

$$b_p + b_q = b_{p+q} \quad (p, q \in C). \quad (2.28)$$

Indeed, consider a maximal flat F containing x_0 and C at infinity, and consider horospherical subgroup N of C and Iwasawa decomposition $M = N \times F$. If $x = n(y)$ for $n \in N$ and $y \in F$, then $b_p(x) + b_q(x) = b_p(y) + b_q(y) = -\langle p, y - x_0 \rangle - \langle q, y - x_0 \rangle = -\langle p + q, y - x_0 \rangle = b_{p+q}(y) = b_{p+q}(x)$, where $p + q \in C$ and F is viewed as \mathbb{R}^d (see Example 2.7).

Let $p := \nabla^\infty f(x)$ for $x \in M$. Suppose that $0 \neq p = \|p\|\xi$ for $\xi \in M^\infty$. Consider sufficiently small $\epsilon > 0$. Let B_ϵ be the set of all points q represented as $q = (\|p\| - t)\xi + 2t\eta$, where $t \leq \epsilon$ and $\xi, \eta \in CM^\infty$ belong to the same Weyl chamber. Then B_ϵ contains p in its interior.

We show that $B_\epsilon \subseteq \nabla^\infty f(M)$ for small $\epsilon > 0$. By strict convexity of f , x is a unique minimizer of $f + b_p$. Let $\alpha > 0$ be the minimum of $(f + b_p)(\exp_x v) - (f + b_p)(x)$ over all $v \in T_x$ with $\|v\|_x = 1$, which exists by compactness. For $q = (\|p\| - t)\xi + 2t\eta \in B_\epsilon$, it holds $f + b_q = f + b_p - tb_\xi + 2tb_\eta$ by (2.28). Here the additional term $-tb_\xi + 2tb_\eta$ is 3ϵ -Lipschitz (Lemma 2.5). For $3\epsilon \leq \alpha$, a minimizer of $f + b_q$ exists in the unit ball around x . Thus $q \in \nabla^\infty f(M)$, as required. The proof of the case $p = 0$ is similar; omit $-t\xi$ above.

If $\nabla^\infty f(M) \cap C \neq \emptyset$, then $\nabla^\infty f(M) \cap C$ has nonempty interior, and Proposition 2.28 is applicable to obtain the latter statement. \square

Note that there is a possibility that $C \cap \nabla^\infty f(M)$ is empty but $C \cap B(f^\infty)$ is nonempty and has no interior. Further refined study on such a degenerate situation is left for future research.

Via the inverse of $x \mapsto \nabla^\infty f(x)$, the symmetric space M is coordinated by the asymptotic gradient space $\nabla^\infty f(M)$. This can be viewed as a generalization of the *dual coordinate* of dually-flat manifolds in information geometry [3]. It is an interesting research direction to develop an information-geometrical theory based on this idea.

In the case of $\text{dom } f^* = B(f)$ which we will face in Section 3, the boundedness of $\inf_{x \in M} (f + b_p)(x)$ is verified by convex optimization on Euclidean building CM^∞ :

$$\inf. \quad f^\infty(u) - \langle u, p \rangle \quad \text{s.t.} \quad u \in U, \quad (2.29)$$

where $U \subseteq CM^\infty$ is any convex neighborhood of the origin, such as a ball.

2.4 Symmetric space $P_n = GL(n, \mathbb{C})/U(n)$

Here we consider the symmetric space $P_n = GL(n, \mathbb{C})/U(n)$ of positive definite Hermitian $n \times n$ matrices, and present concrete descriptions and specializations of several concepts introduced above. Arguments regarding the PSD-cone as a symmetric space are found in [11, Chapter II.10] and [17, 2.2.13], where they consider $GL(n, \mathbb{R})/O(n)$ or $SL(n, \mathbb{R})/SO(n)$ but the arguments are analogous.

When regarding P_n as a Riemannian manifold, the tangent space T_x at $x \in P_n$ is identified with the space S_n of $n \times n$ Hermitian matrices and the inner product is given by $\langle H, H' \rangle_x = \text{tr } x^{-1} H x^{-1} H'$. The cotangent space T_x^* is also identified with S_n by $H(H') := \text{tr } H H'$ for $H, H' \in S_n$. Let d denote the corresponding distance function on M . The exponential map $\exp_x : T_x \rightarrow P_n$ at x is given by $H \mapsto x^{1/2} e^{x^{-1/2} H x^{-1/2}} x^{1/2}$. In particular, any constant-speed ray is written as $t \mapsto g e^{t \text{diag } \lambda} g^\dagger$ for $\lambda \in \mathbb{R}^n$ and $g \in GL(n, \mathbb{C})$, where $\text{diag } \lambda$ denotes the diagonal matrix with diagonal entries $\lambda_1, \lambda_2, \dots, \lambda_n$ in order, and $(\cdot)^\dagger$ denotes the complex conjugate. By $(g, x) \mapsto g x g^\dagger$, $GL(n, \mathbb{C})$ acts

isometrically and transitively on P_n . The isotropy group at I is the group $U(n)$ of unitary matrices. So P_n is a symmetric space $GL(n, \mathbb{C})/U(n) = \mathbb{R} \times (SL(n, \mathbb{C})/SU(n))$, where the geodesic symmetry at $x \in P_n$ is given by $y \mapsto xy^{-1}x$. The identity matrix I is naturally chosen as the base point x_0 of P_n .

Any maximal flat is the set of matrices of form

$$F(g) := \{ge^{\text{diag } \lambda}g^\dagger \mid \lambda \in \mathbb{R}^n\} \quad (2.30)$$

for $g \in GL(n, \mathbb{C})$, where $\lambda \mapsto ge^{\text{diag } \lambda}g^\dagger$ is an isometry from \mathbb{R}^n to $F(g)$. From this, we see that a geodesic ray $t \mapsto ge^{t \text{diag } \lambda}g^\dagger$ is regular if and only if all values λ_i are different. A vector $\lambda \in \mathbb{R}^n$ is said to be *arranged* if $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$. The next lemma characterizes asymptotic classes of geodesic rays.

Lemma 2.30 (See [11, II.10.64]). *For arranged vectors $\lambda, \mu \in \mathbb{R}^n$ with $\|\lambda\|_2 = \|\mu\|_2 = 1$ and $g, h \in GL(n, \mathbb{C})$, two geodesic rays $t \mapsto ge^{t \text{diag } \lambda}g^\dagger$ and $t \mapsto he^{t \text{diag } \mu}h^\dagger$ are asymptotic if and only if $\lambda = \mu$ and $(h^{-1}g)_{ij} = 0$ for all $i \geq j$ with $\lambda_i < \lambda_j$.*

Proof. Let $b(t) := e^{-t \text{diag } \mu/2}h^{-1}ge^{t \text{diag } \lambda/2}$, where $b(t)_{ij} = e^{-t(\mu_i - \lambda_j)/2}(h^{-1}g)_{ij}$. Then we have $d(ge^{t \text{diag } \lambda}g^\dagger, he^{t \text{diag } \mu}h^\dagger) = d(b(t)b(t)^\dagger, I)$. Consider the smallest i with $\lambda_i \neq \mu_i$ if it exists. Say $\lambda_i < \mu_i$. Then $\lim_{t \rightarrow \infty} b(t)$ has a zero block of i rows and $n - i + 1$ columns, and is singular (and/or has a ∞ entry). Necessarily $d(b(t)b(t)^\dagger, I) \rightarrow \infty$. If $\lambda = \mu$ and $(h^{-1}g)_{ij} \neq 0$ for some $i \geq j$ with $\lambda_i < \lambda_j$, then $(b(t)b(t)^\dagger)_{ii} \rightarrow \infty$, implying $d(b(t)b(t)^\dagger, I) \rightarrow \infty$. Conversely, suppose that the condition is satisfied. Then $b(t)$ converges to a constant matrix. This means that $d(b(t)b(t)^\dagger, I)$ is bounded, and the two rays are asymptotic. \square

Thus, the minimal parabolic subgroup for Weyl chamber $C_0 := \{e^{\infty \text{diag } \lambda} \mid \lambda : \text{arranged}\}$ is the group B of upper triangular matrices. Then $GL(n, \mathbb{C})/B$ is viewed as the space of complete flags $U_1 \subset U_2 \subset \dots \subset U_n = \mathbb{C}^n$ of vector subspaces U_i of \mathbb{C}^n . As an abstract simplicial complex, the building CP_n^∞ is the order complex of the lattice of all (nonzero) vector subspaces of \mathbb{C}^n . As consistent with (2.22), any point p in CP_n^∞ is represented as $p = \lambda \cdot \mathcal{U}$ for an arranged vector λ and complete flag \mathcal{U} . Specifically, if p is written as $p = ge^{\infty \lambda}g^\dagger$ for an arranged vector λ and $g \in GL(n, \mathbb{C})$, then \mathcal{U} is the complete flag consisting of vector subspaces spanned by the first i columns of g for $i = 1, 2, \dots, n$. The i -th vector subspaces of complete flags \mathcal{U} and \mathcal{V} are denoted by U_i and V_i , respectively. Lemma 2.30 rephrases as: $\lambda \cdot \mathcal{U} = \mu \cdot \mathcal{V}$ if and only if $\lambda = \mu$ and $U_i = V_i$ for each $i \in [n - 1]$ with $\lambda_i > \lambda_{i+1}$. By associating $\lambda \cdot \mathcal{U}$ with formal sum $\sum_{i=1}^n (\lambda_i - \lambda_{i+1})U_i$ with $\lambda_{n+1} := 0$, the boundary CP_n^∞ is also identified with the set of all formal sums

$$p = \sum_i \alpha_i U_i, \quad (2.31)$$

where vector subspaces U_i form a (partial) flag and nonzero coefficients $\alpha_i \in \mathbb{R}$ are positive if $U_i \neq \mathbb{C}^n$. Note that this expression (2.31) is unique. In this way, any single vector subspace $X (= 1X)$ is viewed as a point of CP_n^∞ .

A Weyl chamber $C \subseteq CP_n^\infty$ consists of $\lambda \cdot \mathcal{U}$ over all arranged vectors λ for a (fixed) flag \mathcal{U} . The distance $d^\infty(p, q)$ of two points $\lambda \cdot \mathcal{U}$ and $\mu \cdot \mathcal{U}$ in the same chamber is given by $\|\lambda - \mu\|_2$. Accordingly, the distance of any two points in CP_n^∞ is given by the length metric of the two points (the infimum of the length of a path connecting them). The whole space CP_n^∞ is a polyhedral cone complex obtained by gluing these




Figure 2: The boundary CP_2^∞ of P_2

Euclidean polyhedral cones. An apartment $E(g) := F(g)^\infty = \{ge^{\infty \text{diag } \lambda} g^\dagger \mid \lambda \in \mathbb{R}^n\}$ for $g \in GL(n, \mathbb{C})$ consists of all points of form $\lambda \cdot \mathcal{U}$ such that each subspace U_i in flag \mathcal{U} is spanned by column vectors of g . $GL(n, \mathbb{C})$ acts isometrically on CP_n^∞ by $(g, \lambda \cdot \mathcal{U}) \mapsto \lambda \cdot g\mathcal{U}$.

Example 2.31. Consider the case of $n = 2$. Any complete flag \mathcal{U} is uniquely determined by its 1-dimensional subspace $U = U_1$. The corresponding Weyl chamber is a form of $\{\lambda \cdot \mathcal{U} \mid \lambda \in \mathbb{R}^2, \lambda_1 \geq \lambda_2\}$, and is isometric to half-plane $\{x \in \mathbb{R}^2 \mid x_1 \geq x_2\} := C_U \subseteq \mathbb{R}^2$. Then CP_2^∞ is obtained by gluing C_U for all 1-dimensional subspaces U , along the line of $x_1 = x_2$. Specifically, it is the disjoint union $\coprod_U C_U$ over all 1-dimensional subspaces U modulo the equivalence relation: $(U, x) \sim (U', x')$ if and only if $x_1 = x_2 = x'_1 = x'_2$. Subspaces U and \mathbb{C}^2 are the points of CP_2^∞ that are the images of $(U, (1, 0))$ and $(U, (1, 1))$, respectively. See Figure 2. This shape of CP_2^∞ can be directly seen from the expression $CP_2^\infty = \mathbb{R} \times C(SL(2, \mathbb{C})/SU(2))^\infty$. Here $SL(2, \mathbb{C})/SU(2)$ is a 3-dimensional hyperbolic space, and $C(SL(2, \mathbb{C})/SU(2))^\infty$ is an infinite star (Example 2.15).

In this setting, we present explicit descriptions of Busemann functions, asymptotic gradients, and inner product \langle, \rangle on CP_n^∞ . For $g \in GL(n, \mathbb{C})$, let $[g]$ denote the image in G/B , which is the complete flag of vector subspaces spanned by the first i columns of g for $i = 1, 2, \dots, n$. For a Hermitian matrix H and subset $I \subseteq [n] := \{1, 2, \dots, n\}$, let $H[I]$ denote the principal matrix of H consisting of row/column indices in I , where $H[\emptyset] := 1$.

Lemma 2.32 (see [11, II.10.69]). *For $p = \lambda \cdot [u] \in CP_n^\infty$ with $u \in U(n)$, it holds*

$$b_p(x) = -\sum_{i=1}^n \lambda_i \log \frac{\det(u^\dagger x u)[\{i, \dots, n\}]}{\det(u^\dagger x u)[\{i+1, \dots, n\}]}, \quad (2.32)$$

$$\nabla b_p(x) = -u b \text{diag } \lambda b^\dagger u^\dagger, \quad (2.33)$$

where $u^\dagger x^{1/2} = bk$ for upper triangular matrix b and unitary matrix k (Gram-Schmidt orthonormalization).

Proof. It suffices to consider $p \in P_n^\infty$. The geodesic ray c issuing from I with $c(\infty) = p$ is written as $c(t) = ue^{t \text{diag } \lambda} u^\dagger$. Therefore, we have $b_p(x) = \lim_{t \rightarrow \infty} d(ue^{t \text{diag } \lambda} u^\dagger, x) - t =$

$\lim_{t \rightarrow \infty} d(e^{t \text{diag } \lambda}, u^\dagger x u) - t$. Suppose that all λ_i are different. Decompose $u^\dagger x u$ as $u^\dagger x u = n \text{diag } r n^\dagger$, where n is an upper triangular matrix having 1 on each diagonal and $r \in \mathbb{R}^n$ is a positive vector with r_i written as

$$r_i = \frac{\det(u^\dagger x u)[\{i, \dots, n\}]}{\det(u^\dagger x u)[\{i+1, \dots, n\}]}.$$
 (2.34)

As in the proof of Lemma 2.30, it holds $\lim_{t \rightarrow \infty} d(n^{-1} e^{t \text{diag } \lambda} n^{-\dagger}, e^{t \text{diag } \lambda}) = 0$, and we have $b_p(x) = \lim_{t \rightarrow \infty} d(e^{t \text{diag } \lambda}, \text{diag } r) - t = \lim_{t \rightarrow \infty} \|t \text{diag } \lambda - \log r\|_2 - t = -\sum_{i=1}^n \lambda_i \log r_i$ (see Example 2.7), where $\log r \in \mathbb{R}^n$ is defined by $(\log r)_i := \log r_i$. Then we obtain (2.32) from (2.34). If some of λ_i are equal, we decompose $u^\dagger x u$ to $n y n^\dagger$, where n is an upper triangular matrix satisfying $n_{ii} = 1$ and $n_{ij} = n_{ji} = 0$ if $\lambda_i = \lambda_j$ and $i \neq j$, and y is a block diagonal matrix with $y_{ij} = y_{ji} = 0$ if $\lambda_i \neq \lambda_j$. As above, it holds $b_p(x) = \lim_{t \rightarrow \infty} d(e^{t \text{diag } \lambda}, y) - t$. Diagonalizing y in each block by unitary matrices, we obtain the same formula.

Next we verify the second equation. For $H := u b \text{diag } \lambda b^\dagger u^\dagger$, consider the geodesic $t \mapsto x^{1/2} e^{t x^{-1/2} H x^{-1/2}} x^{1/2}$ (issuing from x). By $x^{1/2} = u b k = k^\dagger b^\dagger u^\dagger$ and $x^{-1/2} = k^\dagger b^{-1} u^\dagger = u b^{-\dagger} k$, we have $x^{1/2} e^{t x^{-1/2} H x^{-1/2}} x^{1/2} = u b e^{t \text{diag } \lambda} b^\dagger u^\dagger$. Hence, by Lemma 2.30, this geodesic is asymptotic to c , i.e., $t \mapsto u e^{t \text{diag } \lambda} u^\dagger$. From Lemma 2.18, we have $\nabla b_p(x) = -H$. \square

Let $\|\cdot\|_F$ denote the Frobenius norm; then $\|H\|_x = \|x^{-1/2} H x^{-1/2}\|_F$ for $H \in T_x$.

Proposition 2.33. *Let $f : P_n \rightarrow \mathbb{R}$ be a smooth convex function.*

(1) *For $p = \lambda \cdot [u]$ with unitary matrix u , it holds*

$$\|\nabla(f + b_p)(x)\|_x = \|b^\dagger u^\dagger df(x) u b - \text{diag } \lambda\|_F,$$
 (2.35)

where $u^\dagger x^{1/2} = b k$ for upper triangular matrix b and unitary matrix k . In particular, if $\nabla^\infty f(x) = p$, then $x^{1/2} df(x) x^{1/2} = k^\dagger \text{diag } \lambda k$.

(2) *Let s denote the projection $\lambda \cdot \mathcal{U} \mapsto \lambda$. Then it holds*

$$\overline{s \nabla^\infty f(P_n)} = \overline{s \text{dom } f^*} = s B(f^\infty) = \bigcup_C s(C \cap B(f^\infty)),$$
 (2.36)

where C ranges over all Weyl chambers.

We will see in Section 3.2 that $\overline{s \nabla^\infty f(P_n)}$ is viewed as an analogue of the moment polytope.

Proof. (1). From $\langle \nabla f(x), H \rangle_x = \text{tr } df(x) H$, we have $\nabla f(x) = x df(x) x$. Then $\|\nabla(f + b_p)(x)\|_x = \|x^{-1/2} (x df(x) x - u b \text{diag } \lambda b^\dagger u^\dagger) x^{-1/2}\|_F = \|k^\dagger (b^\dagger u^\dagger df(x) u b - \text{diag } \lambda) k\|_F = \|b^\dagger u^\dagger df(x) u b - \text{diag } \lambda\|_F$.

(2). By (2.21), the inclusion $\overline{s \nabla^\infty f(P_n)} \subseteq \overline{s \text{dom } f^*} \subseteq s B(f^\infty)$ is clear. Take $p = \lambda \cdot [u] \in B(f^\infty)$. By Theorem 2.24, there is a sequence (x_i) in P_n such that $\lim_{i \rightarrow \infty} \|\nabla(f + b_p)(x_i)\|_{x_i} = 0$. Suppose that $\nabla^\infty f(x_i) = \lambda_i \cdot [u_i]$. Via the decomposition $u^\dagger x_i^{1/2} = b_k k_i$ and $u_i^\dagger x_i^{1/2} = b'_i h_i$, it holds $x_i^{1/2} df(x) x_i^{1/2} = h_i \text{diag } \lambda_i h_i^\dagger$, and $s \nabla^\infty f(x_i) = \lambda_i$. By the above calculation, we have $\|\nabla(f + b_p)(x_i)\|_{x_i} = \|h_i^\dagger \text{diag } \lambda_i h_i - k_i^\dagger \text{diag } \lambda k_i\|_F \rightarrow 0$ ($i \rightarrow \infty$). This implies that $s \nabla^\infty f(x_i) = \lambda_i \rightarrow \lambda$, and $\lambda \in \overline{s \nabla^\infty f(P_n)}$. \square

Lemma 2.34. For two points $p = \lambda \cdot \mathcal{U}$, $q = \mu \cdot \mathcal{V}$ in CP_n^∞ , it holds

$$\langle p, q \rangle = \sum_{1 \leq i, j \leq n} (\lambda_i - \lambda_{i+1})(\mu_j - \mu_{j+1}) \dim U_i \cap V_j. \quad (2.37)$$

Proof. It is well-known (see [11, II. 10.80]) that there are $g \in GL(n, \mathbb{C})$ and a permutation matrix σ such that $\mathcal{U} = [g]$ and $\mathcal{V} = [g\sigma]$; this is nothing but an axiom of building. In particular, both $p = ge^{\infty \text{diag } \lambda} g^\dagger$ and $q = ge^{\infty \sigma \text{diag } \mu \sigma^\top} g^\dagger$ belong to the apartment $F(g)^\infty$. They are regarded as points in \mathbb{R}^n : $p = \lambda = \sum_i (\lambda_i - \lambda_{i+1}) \mathbf{1}_{[i]}$ and $q = \sum_i (\mu_i - \mu_{i+1}) \mathbf{1}_{\sigma[i]}$, where $\mathbf{1}_J$ denotes the n -dimensional 0,1-vector taking 1 only on indices in $J \subseteq [n]$. Then $\langle p, q \rangle = \sum_{i,j} (\lambda_i - \lambda_{i+1})(\mu_j - \mu_{j+1}) |[i] \cap \sigma[j]|$. Here $|[i] \cap \sigma[j]| = \dim U_i \cap V_j$. \square

Kapovich, Leeb, and Millson [31, Lemma 6.1] gives the corresponding formula of the angle of two vector subspaces regarded as points in $(SL(n, \mathbb{C})/SU(n))^\infty$.

Connections to submodular functions. Let $\mathcal{S}(\mathbb{C}^n)$ denote the family of all vector subspaces of \mathbb{C}^n . A function $\rho : \mathcal{S}(\mathbb{C}^n) \rightarrow \mathbb{R} \cup \{\infty\}$ is called *submodular* if it satisfies

$$\rho(X) + \rho(Y) \geq \rho(X \cap Y) + \rho(X + Y) \quad (X, Y \in \mathcal{S}(\mathbb{C}^n)). \quad (2.38)$$

This extends the classical submodular functions, which are functions κ on $2^{[n]}$ satisfying $\kappa(X) + \kappa(Y) \geq \kappa(X \cap Y) + \kappa(X \cup Y)$; see e.g., [21, 38, 43]. A submodular function ρ with $\rho(\{0\}) = 0$ gives rise to a positively homogeneous function $\bar{\rho} : CP_n^\infty \rightarrow \mathbb{R} \cup \{\infty\}$ via piecewise linear extension

$$\bar{\rho}(p) := \sum_{i=1}^n (\lambda_i - \lambda_{i+1}) \rho(U_i) \quad (p = \lambda \cdot \mathcal{U} \in CP_n^\infty). \quad (2.39)$$

This is an analogy of the *Lovász extension* [37] in the classical setting. So we call $\bar{\rho}$ the Lovász extension of ρ . This is equivalent to the one considered in [26, 27], where λ is restricted to $\lambda \in [0, 1]^n$.

Proposition 2.35 ([27, Theorem 3.9]). For a positively homogeneous function $h : CP_n^\infty \rightarrow \mathbb{R} \cup \{\infty\}$, the following are equivalent:

- (i) h is a convex function that is affine on each Weyl chamber.
- (ii) h is the Lovász extension of a submodular function on $\mathcal{S}(\mathbb{C}^n)$.

The proof reduces to the classical convexity characterization [37] by restricting h to each apartment.

Lemma 2.36. Let $q = \mu \cdot \mathcal{V} \in CP_n^\infty$. Then $p \mapsto -\langle q, p \rangle$ is the Lovász extension of submodular function

$$X \mapsto -\langle q, X \rangle = - \sum_{i=1}^n (\mu_i - \mu_{i+1}) \dim V_i \cap X. \quad (2.40)$$

Proof. By Lemma 2.12, $p \mapsto -\langle q, p \rangle$ is convex. Also it is an affine function on any apartment containing q . Necessarily, it is affine on every Weyl chamber. Therefore, it suffices to show that (2.40) is submodular. We first show that $X \mapsto -\dim V \cap X$ is submodular. This follows from $\dim V \cap X + \dim V \cap Y = \dim V \cap X \cap Y + \dim((V \cap X) + (V \cap Y))$ and $V \cap X + V \cap Y \subseteq V \cap (X + Y)$, we have submodularity of each summand in (2.40) with $i \neq n$. For $i = n$ ($V_n = \mathbb{C}$), the equality $\dim X \cap V_n + \dim X \cap V_n = \dim(X \cap Y) \cap V_n + \dim(X \cup Y) \cap V_n$ holds, and hence (2.40) is submodular (with taking zero on $\{0\}$). \square

For a submodular function $\rho : \mathcal{S}(\mathbb{C}^n) \rightarrow \mathbb{R} \cup \{\infty\}$ with $\rho(\{0\}) = 0$, the subset $B(\bar{\rho})$ is described by fewer inequalities indexed by vector subspaces: It equals the set of points $p \in CP_n^\infty$ satisfying

$$\langle X, p \rangle \leq \rho(X) \quad (X \in \mathcal{S}(\mathbb{C}^n)), \quad \langle \mathbb{C}^n, p \rangle = \rho(\mathbb{C}^n). \quad (2.41)$$

Indeed, if p satisfies (2.41), then for $q = \mu \cdot \mathcal{V} \in CP_n^\infty$, it holds $\langle q, p \rangle = \sum_i (\mu_i - \mu_{i+1}) \langle V_i, p \rangle \leq \sum_i (\mu_i - \mu_{i+1}) h(V_i) = h(q)$.

Then $B(\bar{\rho})$ is called the *base polyhedron* of ρ , which is clearly an analogue of the classical one. A convex function f on P_n is said to be *asymptotically submodular* if f^∞ is the Lovász extension of a submodular function. In this case, the condition (c) in Theorem 3.15 can be replaced by (2.41) with $\rho = f^\infty$. Accordingly, the convex optimization problem (2.29) becomes “discrete” convex optimization (submodular function minimization) over the lattice of vector subspaces:

$$\inf. \quad f^\infty(X) - \langle p, X \rangle \quad \text{s.t.} \quad X \in \mathcal{S}(\mathbb{C}^n). \quad (2.42)$$

3 Scaling problems

In this section, we explain how the results in the previous sections are applied to operator scaling and its generalizations. In our argument, the following convex function on \mathbb{R}^n plays important roles. This function appears in proving (semi)stability results (Kempf-Ness theorem, Hilbert-Mumford criterion) in invariant theory; see [32, 45].

Lemma 3.1. *For $a_i > 0$ and $w_i \in \mathbb{R}^n$ ($i = 1, 2, \dots, m$), define $f : \mathbb{R}^n \rightarrow \mathbb{R}$ by*

$$f(x) := \log \sum_{i=1}^m a_i e^{\langle w_i, x \rangle} \quad (x \in \mathbb{R}^n). \quad (3.1)$$

Then f is convex, where:

$$f^\infty(p) = \max_{i=1,2,\dots,m} \langle w_i, p \rangle \quad (p \in \mathbb{R}^n),$$

$$\text{dom } f^* = B(f^\infty) = \text{the convex hull of } w_1, w_2, \dots, w_m.$$

Proof. The convexity of f is well-known. Computing f^∞ is also a little exercise [40, p. 68]: Letting $v^* := \max_i \langle w_i, p \rangle$, we have $f^\infty(p) = \lim_{t \rightarrow \infty} \frac{1}{t} \log \sum_{i=1}^m a_i e^{\langle w_i, pt \rangle} = v^* + \lim_{t \rightarrow \infty} \frac{1}{t} \log \sum_{i=1}^m a_i e^{t(\langle w_i, p \rangle - v^*)} = v^*$.

Thus $B(f^\infty)$ equals the convex hull of w_i . We show that it equals $\text{dom } f^*$. Suppose $p = \sum_i \lambda_i w_i \in B(f^\infty)$ with $\lambda_i \geq 0$ and $\sum_i \lambda_i = 1$. Then, for small $c > 0$, it holds $c\lambda_i \leq a_i$ for all i . From $\log \sum_{i=1}^m a_i e^{\langle w_i, x \rangle} \geq \log \sum_{i=1}^m c\lambda_i e^{\langle w_i, x \rangle} \geq \log c + \log e^{\sum_i \lambda_i \langle w_i, x \rangle} = \log c + \langle p, x \rangle$, we have $f(x) - \langle p, x \rangle \geq \log c$ for all $x \in \mathbb{R}^n$. Hence $B(f^\infty) \subseteq \text{dom } f^*$; the reverse inclusion is generally true. \square

3.1 Operator scaling with specified marginals

Let $A = (A_1, A_2, \dots, A_m)$ be an m tuple of nonzero complex $n \times n$ matrices. Let $\lambda, \mu \in \mathbb{R}^n$ be nonnegative arranged vectors with the same sum $\sum_i \lambda_i = \sum_i \mu_i = n$ (say). The *operator scaling problem with marginal* λ, μ , introduced by Franks [19], is to find a pair of nonsingular matrices $g, h \in GL(n, \mathbb{C})$ such that

$$\sum_{k=1}^m g^\dagger A_k h h^\dagger A_k^\dagger g = \text{diag } \lambda, \quad \sum_{k=1}^m h^\dagger A_k^\dagger g g^\dagger A_k h = \text{diag } \mu. \quad (3.2)$$

If such g, h exist, then A is said to be (λ, μ) -scalable. If for every $\epsilon > 0$ there are g, h such that

$$\left\| \sum_{k=1}^m g^\dagger A_k h h^\dagger A_k^\dagger g - \text{diag } \lambda \right\|_{\mathbb{F}} < \epsilon, \quad \left\| \sum_{k=1}^m h^\dagger A_k^\dagger g g^\dagger A_k h - \text{diag } \mu \right\|_{\mathbb{F}} < \epsilon, \quad (3.3)$$

then A is said to be *approximately* (λ, μ) -scalable. If $\lambda = \mu = \mathbf{1}$, it is the original operator scaling problem by Gurvits [25]. In this case, the approximate scalability is equivalent to the *noncommutative nonsingularity* of symbolic matrix $\sum_k A_k x_k$ [18, 30]. See [19, 20, 22] for further applications of operator scaling.

For simplicity, we assume that at least one of $\bigcap_{k=1}^m \ker A_k$ and $\bigcap_{k=1}^m \ker A_k^\dagger$ is trivial $\{0\}$. Otherwise, by coordinate change, we can make A satisfy $(A_k)_{in} = (A_k)_{nj} = 0$ for i, j, k . Then the problem reduces to the upper left $(n-1) \times (n-1)$ submatrices.

The operator scaling problem is viewed as the problem of finding a point (x, y) in $P_n \times P_n$ at which the following convex function $f_A : P_n \times P_n \rightarrow \mathbb{R}$ has a specified asymptotic gradient:

$$f_A(x, y) := n \log \sum_{k=1}^m \text{tr } x A_k y A_k^\dagger. \quad (3.4)$$

This function is known to be (geodesically) convex.

Lemma 3.2 (See e.g., [2]). *f_A is convex.*

Indeed, on a maximal flat $F = F(g) \times F(h) = \{g e^{\text{diag } \alpha} g^\dagger\}_{\alpha \in \mathbb{R}^n} \times \{h e^{\text{diag } \beta} h^\dagger\}_{\beta \in \mathbb{R}^n}$, f_A is written as

$$f_A(g e^{\text{diag } \alpha} g^\dagger, h e^{\text{diag } \beta} h^\dagger) = n \log \sum_{1 \leq i, j \leq n} a_{ij}(g, h) e^{\alpha_i + \beta_j} \quad (\alpha, \beta \in \mathbb{R}^n), \quad (3.5)$$

where $a_{ij}(g, h) := \sum_{k=1}^m |(g^\dagger A_k h)_{ij}|^2$. By Lemma 3.1, f_A is convex in every flat.

In addition to the (λ, μ) -scalability, we consider a sharper scalability concept. Let \mathcal{U}, \mathcal{V} be complete flags, and consider points $(\lambda \cdot \mathcal{U}, \lambda \cdot \mathcal{V})$ in the boundary $C(P_n \times P_n)^\infty = CP_n^\infty \times CP_n^\infty$. We say that A is $(\lambda \cdot \mathcal{U}, \mu \cdot \mathcal{V})$ -scalable if there are $g, h \in GL(n, \mathbb{C})$ such that $([g], [h]) = (\mathcal{U}, \mathcal{V})$ and (3.2) hold. Accordingly, we say that A is *approximately* $(\lambda \cdot \mathcal{U}, \mu \cdot \mathcal{V})$ -scalable if for every $\epsilon > 0$ there are $g, h \in GL(n, \mathbb{C})$ such that $([g], [h]) = (\mathcal{U}, \mathcal{V})$ and (3.3) hold. By definition, A is (approximately) (λ, μ) -scalable if and only if A is (approximately) $(\lambda \cdot \mathcal{U}, \mu \cdot \mathcal{V})$ -scalable for some flags \mathcal{U}, \mathcal{V} .

When \mathcal{U} and \mathcal{V} are standard flag $\mathcal{E} := [I]$, scaling matrices g, h are upper triangular, and hence the (approximate) $(\lambda \cdot \mathcal{E}, \mu \cdot \mathcal{E})$ -scalability is equivalent to (approximate) (λ, μ) -scalability by triangular matrices in the sense of Franks [19]. Note that the

$(\lambda \cdot \mathcal{U}, \mu \cdot \mathcal{V})$ -scalability reduces to the triangular scalability, since A is $(\lambda \cdot \mathcal{U}, \mu \cdot \mathcal{V})$ -scalable if and only if $g^\dagger A h$ is $(\lambda \cdot \mathcal{E}, \mu \cdot \mathcal{E})$ -scalable for $([g], [h]) = (\mathcal{U}, \mathcal{V})$.

The $(\lambda \cdot \mathcal{U}, \mu \cdot \mathcal{V})$ -scalability is rephrased by using asymptotic gradient ∇^∞ and Busemann functions.

Proposition 3.3. (1) A is $(\lambda \cdot \mathcal{U}, \mu \cdot \mathcal{V})$ -scalable if and only if there are points x, y in P_n such that $\nabla^\infty f_A(x, y) = (\lambda \cdot \mathcal{U}, \mu \cdot \mathcal{V})$.

(2) A is approximately $(\lambda \cdot \mathcal{U}, \mu \cdot \mathcal{V})$ -scalable if and only if $\inf_{x, y \in P_n} \|\nabla(f_A + b_{\lambda \cdot \mathcal{U}, \mu \cdot \mathcal{V}})(x, y)\|_{x, y} = 0$.

Proof. From $df_A(x, y)(H, G) = \frac{d}{dt} \Big|_{t=0} n \log \sum_{k=1}^m \text{tr}(x + tH) A_k (y + tG) A_k^\dagger$, we have

$$df_A(x, y) = C_{x, y} \left(\sum_k A_k y A_k^\dagger, \sum_k A_k^\dagger x A_k \right), \quad (3.6)$$

where $C_{x, y} := n / \sum_{k=1}^m \text{tr} x A_k y A_k^\dagger$. Let $\mathcal{U} = [u]$ and $\mathcal{V} = [v]$ for $u, v \in U(n)$. By Proposition 2.33, we have

$$\begin{aligned} \|\nabla(f_A + b_{\lambda \cdot \mathcal{U}, \mu \cdot \mathcal{V}})(x, y)\|_{x, y}^2 &= \left\| C_{x, y} \sum_k b^\dagger u^\dagger A_k v c c^\dagger v^\dagger A_k^\dagger u b - \text{diag } \lambda \right\|_{\mathbb{F}}^2 \\ &\quad + \left\| C_{x, y} \sum_k c^\dagger v^\dagger A_k^\dagger u b b^\dagger u^\dagger A_k v c - \text{diag } \mu \right\|_{\mathbb{F}}^2, \end{aligned} \quad (3.7)$$

where $u^\dagger x^{\frac{1}{2}} = bk$ and $v^\dagger y^{\frac{1}{2}} = ck'$ for $k, k' \in U(n)$ and upper-triangular matrices b, c . From this, we have the claims, where required scaling matrices g, h are given as $g = C_{x, y}^{1/4} u b$ and $h = C_{x, y}^{1/4} v c$ with $[g] = [u] = \mathcal{U}$ and $[h] = [v] = \mathcal{V}$. \square

For the function f_A , the inclusion $\text{dom } f_A^* \subseteq \overline{\text{dom } f_A^*} \subseteq B(f_A^\infty)$ becomes equality.

Proposition 3.4. $\text{dom } f_A^* = B(f_A^\infty)$.

We will prove a general version (Proposition 3.14) in Section 3.2. Thus, the (p, q) -scalability with $p = \lambda \cdot \mathcal{U}$ and $q = \mu \cdot \mathcal{V}$ can be decided by the boundeness of convex optimization:

$$\inf. (f_A + b_{p, q})(x, y) = f_A(x, y) + b_p(x) + b_q(y) \text{ s.t. } (x, y) \in P_n \times P_n,$$

where Busemann functions b_p and b_q are explicitly given by Lemma 2.32. By optimizing y under a fixed $x \in P_n$, we may minimize function $g_{A, q} : P_n \rightarrow \mathbb{R}$:

$$g_{A, q}(x) := \inf_{y \in P_n} f_A(x, y) + b_q(y) \quad (x \in P_n). \quad (3.8)$$

One can see from (3.6) and (3.7) that optimal y is obtained by $y = h h^\dagger$ for $h \in GL(n, \mathbb{C})$ with $\mathcal{V} = [h]$ and $h^\dagger (\sum_k A_k x A_k^\dagger) h = \text{diag } \mu$. When $\mathcal{U} = \mathcal{V} = \mathcal{E}$, the infimum of $f_A + b_p + b_q$ (or $g_{A, q} + b_p$) equals (up to constant) the logarithm of the *capacity of specified marginal* in Franks [19].¹

We compute explicit descriptions of the recession functions of f_A and the associated subset $B(f_A^\infty)$. See Remark 3.8 (1) for $g_{A, q}$ and $B(g_{A, q}^\infty)$. Let \mathcal{S}_A be the family of all pairs (X, Y) of vector subspaces in \mathbb{C}^n such that $u^\dagger A_k v = 0$ for all $u \in X, v \in Y, k \in [m]$.

¹To see the consistency with his formulation, use the relation $b_{\lambda, \mathcal{E}}(g g^\dagger) = -\log \det(\text{diag } \lambda, g^\dagger g)$ for any upper-triangular matrix g , where $\det(\text{diag } \lambda, g^\dagger g)$ is the *relative determinant* in the sense of [19].

Proposition 3.5. (1) The recession function f_A^∞ is given by

$$f_A^\infty(p, q) = n \max\{\alpha_i + \beta_j \mid i, j \in [n] : (g^\dagger A_k h)_{ij} \neq 0 \ (\exists k \in [m])\}, \quad (3.9)$$

where $p = \alpha \cdot [g], q = \beta \cdot [h] \in CP_n^\infty$.

(2) $B(f_A^\infty)$ is the set of $(p, q) \in CP_n^\infty \times CP_n^\infty$ satisfying

$$\langle X, p \rangle + \langle Y, q \rangle \leq n \quad ((X, Y) \in \mathcal{S}_A). \quad (3.10)$$

Proof. (1) follows from Lemma 3.1 and the expression (3.5). (2). Let $(p, q) \in CP_n^\infty \times CP_n^\infty$. For $(X, Y) \in \mathcal{S}_A$, choose $g, h \in GL(n, \mathbb{C})$ such that g and h span X and Y in the first k and l column subsets, respectively. Then, each $g^\dagger A_k h$ has a $k \times l$ zero block in the upper left corner. If X, Y are viewed as points in CP_n^∞ by (2.31), then $X = \mathbf{1}_{[k]} \cdot [g]$ and $Y = \mathbf{1}_{[l]} \cdot [h]$. By the assumption that $\bigcap_{k=1}^m \ker A_k = \{0\}$ or $\bigcap_{k=1}^m \ker A_k^\dagger = \{0\}$, the maximum in (3.9) is attained by $i \in [k], j \in [n] \setminus [l]$ or $i \in [n] \setminus [k], j \in [l]$, and we have $f_A^\infty(X, Y) = n$. Thus (3.10) is a necessary condition for $(p, q) \in B(f_A^\infty)$. Consider an apartment $E(g) \times E(h) = F(g)^\infty \times F(h)^\infty$ containing (p, q) and identify it with $\mathbb{R}^n \times \mathbb{R}^n$ by $(ge^{\infty\alpha} g^\dagger, he^{\infty\beta} h^\dagger) \mapsto (\alpha, \beta)$. From Lemma 3.1 and (3.5), we have

$$B_{E(g) \times E(h)}(f^\infty) = n \text{ the convex hull of } e_i + f_j \text{ for all } i, j \text{ with } a_{ij}(g, h) \neq 0, \quad (3.11)$$

where e_i and f_i denote the i -th unit vectors of $E(g)$ and $E(h)$, respectively. This is n times the *clique polytope* of the bipartite graph G with vertex set $[n] \sqcup [n]$ and edge set $\{ij \mid a_{ij}(g, h) \neq 0\}$; see [43, Section 65.4]. By a standard network flow argument, we obtain the inequality description of $B_{E(g) \times E(h)}$ as

$$\sum_{i \in S} \alpha_i + \sum_{j \in T} \beta_j \leq n \quad ((S, T) \subseteq [n] \times [n] : a_{ij}(g, h) = 0 \ (i \in S, j \in T)), \quad (3.12)$$

where $S \sqcup T$ is nothing but a stable set of the graph G . Notice that (3.12) is the subsystem for (3.10) such that vector subspace X and Y are spanned by columns vectors of g and h . By Proposition 2.27, satisfying all such inequalities is also sufficient for $(p, q) \in B(f_A^\infty)$. \square

Thus, by Theorem 2.24, Propositions 3.3, 3.4, 3.5, and Lemma 2.34, we have:

Theorem 3.6 ([19]). *The following conditions are equivalent:*

(a) A is approximately $(\lambda \cdot \mathcal{U}, \mu \cdot \mathcal{V})$ -scalable.

(b) $\inf_{x, y \in P_n} f_A(x, y) + b_{\lambda \cdot \mathcal{U}}(x) + b_{\mu \cdot \mathcal{V}}(y) > -\infty$.

(c) For all $(X, Y) \in \mathcal{S}_A$, it holds

$$\sum_{i=1}^n (\lambda_i - \lambda_{i+1}) \dim U_i \cap X + \sum_{i=1}^n (\mu_i - \mu_{i+1}) \dim V_i \cap Y \leq n. \quad (3.13)$$

Franks [19] showed that the approximate scalability reduces to the triangular scalability in the generic case.

Theorem 3.7 ([19]). *A is approximately (λ, μ) -scalable if and only if $g^\dagger Ah$ is approximately $(\lambda \cdot \mathcal{E}, \mu \cdot \mathcal{E})$ -scalable for generic $g, h \in GL(n, \mathbb{C})$.*

Here “generic” means that there is an affine variety $V \subseteq GL(n, \mathbb{C})^2$ such that the latter property holds for all $(g, h) \in GL(n, \mathbb{C})^2 \setminus V$. We will verify this theorem for a general setting of the moment polytope membership in the next section.

Remark 3.8. (1) One can show that the recession function $g_{A,q}^\infty$ of $g_{A,q}$ is the Lovász extension of submodular function

$$X \mapsto n - \langle X^{\perp A}, q \rangle, \quad (3.14)$$

where $X^{\perp A}$ denotes the maximum subspace Y with $(X, Y) \in \mathcal{S}_A$. In particular, $g_{A,q}$ is asymptotically submodular, and $B(g_{A,q}^\infty)$ coincides with the base polyhedron of $g_{A,q}^\infty$.

- (2) Computation of the constant (*BL-constant*) of the *Brascamp-Lieb inequality* [10, 35] is formulated as the same type of convex optimization over the product of PSD-cones (over \mathbb{R}) [23]. The objective function is also asymptotically submodular. A finiteness characterization of the BL-constant by [8] can be deduced by the same way as for (3.13) above.
- (3) Since $(X, Y), (X', Y') \in \mathcal{S}_A$ implies $(X \cap X', Y + Y'), (X + X', Y \cap Y') \in \mathcal{S}_A$, the function $(X, Y) \mapsto -\langle X, p \rangle - \langle Y, q \rangle$ also admits a submodular function structure on the lattice \mathcal{S}_A with $\wedge = (\cap, +), \vee = (+, \cap)$. Its Lovász extension (in the sense of [26, 27]) coincides with a part of $f_{A,p,q}^\infty$. The nc-rank computation algorithm in [26] is interpreted as minimizing $f_{A,p,q}^\infty$ (with $\mu = \lambda = \mathbf{1}$) over a convex neighborhood of 0.

3.2 Optimization on group orbits

The operator scaling and its generalizations (e.g., tensor scaling [13, 14]) can be formulated as optimization over an orbit of a group action. We finally consider the generalized scaling problems formulated by Bürgisser, Franks, Garg, Oliveira, Walter, and Wigderson [15]. Let $G \subseteq GL(n, \mathbb{C})$ be a reductive algebraic group over \mathbb{C} , i.e., G is defined by the zero set of a finite number of polynomials with complex coefficients, and $g \in G$ implies $g^\dagger \in G$. We assume that G is connected. Since G is a closed subgroup of Lie group $GL(n, \mathbb{C})$, it is also a Lie group. Let $K := G \cap U(n)$ be a maximal compact subgroup of G . Let \mathfrak{g} and \mathfrak{u} denote the Lie algebras of G and K , respectively, where $\mathfrak{g} = \mathfrak{u} + i\mathfrak{u}$ is the complexification of \mathfrak{u} (or Cartan decomposition of involution $X \mapsto -(X)^\dagger$). This is a situation of [34, VII. 2. Example (2)].

Let $\pi : G \rightarrow GL(N, \mathbb{C})$ be a rational representation, i.e., each entry of matrix $\pi(g)$ is a polynomial of g_{ij} and $(\det g)^{-1}$. Let $\langle \cdot, \cdot \rangle$ denote a K -invariant inner product on \mathbb{C}^N , i.e., it satisfies $\langle \pi(k)u, \pi(k)v \rangle = \langle u, v \rangle$ for all $k \in K$. Let $\Pi := d\pi(I)$ be the Lie algebra representation of π . Then $\pi(e^H) = e^{\Pi(H)}$ holds for $H \in \mathfrak{g}$. The conjugate of $g \in GL(N, \mathbb{C})$ with respect to $\langle \cdot, \cdot \rangle$ is denoted by g^\dagger (the matrix satisfying $\langle g^\dagger u, v \rangle = \langle u, gv \rangle$ for all $u, v \in \mathbb{C}^N$). Then $\Pi(H^\dagger) = \Pi(H)^\dagger$ holds for $H \in \mathfrak{g}$.² Also

²From $\frac{d}{dt} \langle \pi(e^{tH_0})u, \pi(e^{tH_0})v \rangle = 0$ for $H_0 \in \mathfrak{u}$, we have $\langle \Pi(H_0)u, v \rangle + \langle u, \Pi(H_0)v \rangle = 0$. Thus $\Pi(H_0)^\dagger = -\Pi(H_0)$. For $H = H_0 + iH_1$ with $H_0, H_1 \in \mathfrak{u}$, we have $\Pi(H)^\dagger = \Pi(H_0)^\dagger - i\Pi(H_1)^\dagger = -\Pi(H_0) + i\Pi(H_1) = \Pi(-H_0 + iH_1) = \Pi(H^\dagger)$.

$\pi(g)^\dagger = \pi(g^\dagger)$ holds for $g \in G$.³

Given a vector $v \in \mathbb{C}^N$, consider minimization of log-norm $\log \|\pi(g)v\|^2$ (twice of the *Kempf-Ness function* in [15]) over the G -orbit of v :

$$\inf. \log \|\pi(g)v\|^2 \quad \text{s.t. } g \in G. \quad (3.15)$$

This optimization can decide whether $0 \in \overline{\pi(G)v}$ (the closure of orbit $\pi(G)v$), via the unboundedness. This is equivalent to the membership of v in the *null-cone* of the invariant ring of π . The operator scaling in the previous section corresponds to the *left-right action* $(g, h) \mapsto g^\dagger Ah$, where f_A is a constant multiple of the Kempf-Ness function.

Since the norm is K -invariant, the optimization problem (3.15) is viewed as that the quotient space G/K , which turns out to be a symmetric space of nonpositive curvature. We formulate the optimization problem (3.15) more explicitly, as in [15, Remark 3.4]. Note that $\|\pi(g)v\|^2 = \langle v, \pi(g^\dagger g)v \rangle$, where $g^\dagger g \in G \cap P_n$. Since G is algebraic, $x \in G \cap P_n$ implies $x^{1/2} \in G \cap P_n$; see [11, II.10.59]. Therefore (3.15) is also written as

$$\inf. f_v(x) := \log \langle v, \pi(x)v \rangle \quad \text{s.t. } x \in M := G \cap P_n. \quad (3.16)$$

Here $M = G \cap P_n$ is a totally geodesic subspace of P_n , and hence is a symmetric space of nonpositive curvature (see [11, II. 10. 50]). By polar decomposition $G = Ke^{iu}$, we have $M = e^{iu} \simeq G/K$, where iu is viewed as the tangent space at I with inner product $(\lambda, \nu) \mapsto \text{tr } \lambda \nu$.

Then, M is decomposed as a Euclidean space and symmetric space of noncompact type as follows. Since G is reductive, the Lie algebra \mathfrak{g} is the direct sum of the center \mathfrak{z} and semisimple Lie algebra $\mathfrak{g}_1 := [\mathfrak{g}, \mathfrak{g}]$, where \mathfrak{z} and \mathfrak{g}_1 are orthogonal in the inner product $X, Y \mapsto \text{Re tr}(XY^\dagger)$; see [34, Proposition 1.59]. Now G is commuting product ZG_1 of the center Z for \mathfrak{z} and semisimple Lie group G_1 for \mathfrak{g}_1 , where $iu = \mathfrak{z} \cap iu + \mathfrak{g}_1 \cap iu$; see [34, Proposition 7.19 (e)]. Thus $M = G \cap P_n$ is Riemannian product of Euclidean space $\mathbb{R}^k \simeq e^{\mathfrak{z} \cap iu}$ and symmetric space $M_1 := G_1 \cap P_n = e^{\mathfrak{g}_1 \cap iu}$ of noncompact type. If $g = zg_1$ for $z \in Z$ and $g_1 \in G_1$, then the action of g on $M = \mathbb{R}^k \times M_1$ is given so that g_1 acts on M_1 as $x \mapsto g_1 x g_1^\dagger$ (as before) and z acts on \mathbb{R}^k as translation. Particularly, z acts trivially on the boundary $CM^\infty = \mathbb{R}^k \times CM_1^\infty$.

Maximal flats of M are the intersection of maximal flats of P_n with G , and are given by e^{it} for maximal commutative subspaces (maximal tori) \mathfrak{t} of \mathfrak{u} . Fix a maximal torus \mathfrak{t} of \mathfrak{u} . Then, any maximal flat is written as $F(g) := \{ge^\lambda g^\dagger \mid \lambda \in \mathfrak{t}\}$ for $g \in G$, where $\lambda \mapsto ge^\lambda g^\dagger$ is an isometry from Euclidean space \mathfrak{t} to $F(g)$. The dimension d of \mathfrak{t} is equal to the rank of M . Via $\lambda \mapsto e^{\infty \lambda}$, we regard \mathfrak{t} as a subset, particularly, an apartment of CM^∞ . Let \mathfrak{t}^+ be any fixed (asymptotic) Weyl chamber in \mathfrak{t} , and let B denote the minimal parabolic subgroup (Borel subgroup) for \mathfrak{t}^+ . Any point p in CM^∞ is written as $\lambda \cdot \mathcal{F}$ for $\lambda \in \mathfrak{t}^+$ and $\mathcal{F} \in G/B$.

Since $M = G \cap P_n$, any Weyl chamber of M is written as the intersection of a Weyl chamber of P_n and a maximal flat of M . Consequently, M^∞ is an isometric subspace of P_n^∞ . By using the notation in Section 2.4, for some $k_0 \in U(n)$, vectors $\lambda \in \mathfrak{t}^+$ are written as $\bar{\lambda} \cdot [k_0]$, where $\bar{\lambda}$ ranges over a subspace of arranged vectors.

It is known [15, 46] that the Kempf-Ness function f_v is convex on M . Indeed, consider the expression of f_v in the maximal flat $F(g)$. Since $\{e^\lambda\}_{\lambda \in \mathfrak{t}}$ is a commutative

³By $\pi(k^\dagger) = \pi(k^{-1}) = \pi(k)^{-1} = \pi(k)^\dagger$ for $k \in K$ and polar decomposition $g = ke^{iH}$ for $k \in K$ and $H \in \mathfrak{u}$, we have $\pi(g^\dagger) = e^{-i\Pi(H^\dagger)} \pi(k^\dagger) = e^{-i\Pi(H)} \pi(k)^\dagger = (\pi(k)e^{i\Pi(H)})^\dagger = \pi(g)^\dagger$.

subgroup, there is a finite set $\Omega(\pi)$ of vectors, called *weights*, in \mathfrak{it} such that matrices $\pi(e^\lambda) = e^{\Pi(\lambda)}$ ($\lambda \in \mathfrak{it}$) are simultaneously diagonalized to a diagonal matrix of diagonals $e^{\text{tr}\omega\lambda}$ for $\omega \in \Omega(\pi)$. Therefore, we have

$$f_v(ge^\lambda g^\dagger) = \log \sum_{\omega \in \Omega(\pi)} \|(\pi(g^\dagger)v)_\omega\|^2 e^{\text{tr}\omega\lambda} \quad (\lambda \in \mathfrak{it}), \quad (3.17)$$

where $(\pi(g^\dagger)v)_\omega$ denotes the orthogonal projection of $\pi(g^\dagger)v$ to the eigenspace of ω . By Lemma 3.1 and the expression (3.17), we have:

Lemma 3.9. *f_v is convex, where:*

(1) *The recession function f_v^∞ is given by*

$$f_v^\infty(p) = \max\{\text{tr}\omega\lambda \mid \omega \in \Omega(\pi) : (\pi(g^\dagger)v)_\omega \neq 0\}, \quad (3.18)$$

where $p = ge^{\lambda^\infty}g^\dagger$ for $g \in G$ and $\lambda \in \mathfrak{it}$.

(2) *$B_{CF(g)^\infty}(f_v^\infty)$ is the convex hull of ω over all $\omega \in \Omega(\pi)$ with $(\pi(g^\dagger)v)_\omega \neq 0$, where ω are viewed as points in $CF(g)^\infty$ by $\omega \mapsto ge^{\omega^\infty}g^\dagger$.*

To study the boundedness of f_v , the following criterion is fundamental:

Theorem 3.10 (Hilbert-Mumford criterion; see [45, Section 3.4.2]). *If $\inf_{g \in G} \|\pi(g)v\| = 0$, then there is $u \in \mathfrak{iu}$ such that $\lim_{t \rightarrow \infty} \|\pi(e^{tu})v\| = 0$.*

The reference [45, Theorem 3.23] also includes an elementary proof. As noticed in [31, 46], the nonnegativity of the asymptotic slope function of f_v is equivalent to the Hilbert-Mumford criterion:

Theorem 3.11 (see [31, 46]). *The following conditions are equivalent:*

(a) $\inf_{x \in M} \|\nabla f_v(x)\|_x = 0$.

(b) $\inf_{x \in M} f_v(x) > -\infty$.

(c) $0 \in B(f_v^\infty)$.

The equivalence (a) \Leftrightarrow (b) is known as the Kempf-Ness theorem [32], and is called the *noncommutative duality* in [15].

Proof. We have already seen (a) \Leftrightarrow (c) and (b) \Rightarrow (c) in general situation; see Lemma 2.14 and Theorem 2.24. We verify (c) \Rightarrow (b). Suppose that $\inf_{x \in M} f_v(x) = -\infty$. By the Hilbert-Mumford criterion, there is $u \in \mathfrak{iu}$ such that $\lim_{t \rightarrow \infty} f_v(e^{tu}) = -\infty$. Consider a maximal flat F containing geodesic $t \mapsto e^{tu}$. Then f_v is unbounded on F . By Lemma 3.1, we have $0 \notin B((f_v)_F^\infty) = B_{CF^\infty}(f_v^\infty)$. By Proposition 2.27, we have $0 \notin B(f_v^\infty)$. \square

In particular, a one-parameter subgroup $t \rightarrow e^{tu}$ in the Hilbert-Mumford criterion can be found by convex optimization of f_v^∞ on Euclidean building CM^∞ :

$$\inf. \quad f_v^\infty(u) \quad \text{s.t.} \quad u \in U, \quad (3.19)$$

where U is any convex neighborhood of the origin.

We are going to extend Theorem 3.11 for $f_v + b_p$ with giving a whole description of $B(f_v^\infty)$ and $\text{dom } f_v^*$. For this, we need a representation theoretic interpretation of Busemann functions. By a *weight* we mean a point in \mathfrak{it} that arises as a weight of some representation. It is known that the set of weights is a discrete subgroup (*weight lattice*) in \mathfrak{it} , and is generated by weights in \mathfrak{it}^+ . Any weight λ in \mathfrak{it}^+ determines an irreducible representation π_λ of G such that λ is a highest weight. The eigenspace for λ is one dimensional, and the unit eigenvector is denoted by v_λ .

Lemma 3.12. *For a weight λ , it holds $b_{e^{-\lambda\infty}}(g^\dagger g) = \log \|\pi_\lambda(g)v_\lambda\|^2$ ($g \in G$).*

Proof. Consider Iwasawa decomposition $g = ke^\nu n$ for $k \in K, \nu \in \mathfrak{it}, n \in N$, where N is interpreted as the horospherical subgroup for \mathfrak{it}^+ (see [17, Section 2.17]). From $\pi_\lambda(n)v_\lambda = v_\lambda$ (see [34, Theorem 5.5]), the RHS equals $2 \text{tr } \nu \lambda$. On the other hand, n^\dagger is an element of the horospherical subgroup of the opposite Weyl chamber $-\mathfrak{it}^+$, since $\lim_{t \rightarrow \infty} d(n^{-\dagger} e^{-\lambda t} n^{-1}, e^{-\lambda t}) = \lim_{t \rightarrow \infty} d(ne^{\lambda t} n^\dagger, e^{\lambda t}) = 0$ for $\lambda \in \mathfrak{it}^+$. Then the LHS equals $b_{e^{-\lambda\infty}}(n^\dagger e^{2\nu} n) = b_{e^{-\lambda\infty}}(e^{2\nu}) = -2 \text{tr}(-\lambda \nu)$ (by Example 2.7). \square

Therefore, the minimization of $f_v + b_p$ is essentially the *p-scaling problem* in [15]. A point ν in \mathfrak{it} is said to be *rational* if $\alpha\nu$ is a weight for some positive integer α . Let $s : CM^\infty \rightarrow \mathfrak{it}^+$ denote the projection $\lambda \cdot \mathcal{F} \mapsto \lambda$.

Proposition 3.13. *There is a finite set $\Lambda \subseteq \mathfrak{it}^+$ (independent of v) such that*

$$B(f_v^\infty) = \{p \in CM^\infty \mid \langle \nu \cdot \mathcal{F}, p \rangle \leq f_v^\infty(\nu \cdot \mathcal{F}) \ (\nu \in \Lambda, \mathcal{F} \in G/B)\}. \quad (3.20)$$

For any Weyl chamber C , the projection $s(C \cap B(f_v^\infty))$ is a rational convex polytope.

Proof. Take a Weyl chamber C of CP_n^∞ , and consider all apartments E containing C . When all E are regarded as \mathbb{R}^d with a common convex cone C , by Proposition 2.27, $C \cap B(f_v^\infty)$ is the intersection of C and finitely many (integral) polytopes $B_E(f_v^\infty)$, which are convex hulls of finite subsets of weights in $\Omega(\pi) \subseteq \mathbb{R}^n$. Consequently, $s(C \cap B(f_v^\infty))$ is a rational convex polytope.

In particular, the affine span of a facet of $B_E(f_v^\infty)$ is spanned by a subset of $\Omega(\pi)$, and its normal vector is chosen from \mathfrak{it} . The corresponding inequality is written as $\langle \nu \cdot \mathcal{F}, p \rangle \leq f_v^\infty(\nu \cdot \mathcal{F})$ for some $\mathcal{F} \in G/B$ and $\nu \in \mathfrak{it}^+$ (with $\nu \cdot \mathcal{F} \in F$). Thus, Λ can be chosen as a (finite) set of vectors arising as normal vectors of $d - 1$ -spaces spanned by subsets of $\Omega(\pi)$. \square

Proposition 3.14. $\text{dom } f_v^* = B(f_v^\infty)$.

Proof. By rationality and convexity (Proposition 3.13), it suffices to show that for $p = \lambda \cdot \mathcal{F} \in CM^\infty$ with rational $\lambda \in \mathfrak{it}^+$ it holds $\inf_{x \in M} (f_v + b_p)(x) > -\infty$ if and only if $p \in B(f_v^\infty)$. Suppose that $\lambda = \nu/\alpha$ where ν is a weight and α is a positive integer. By Lemma 3.12, $\alpha b_p = b_{\nu \cdot \mathcal{F}}$ is the Kempf-Ness function for some representation π_ν and vector v' . Then $\alpha f_v + b_{\nu \cdot \mathcal{F}}$ is the Kempf-Ness function for representation $\overbrace{\pi \otimes \pi \otimes \cdots \otimes \pi}^\alpha \otimes \pi_\nu$ and vector $\overbrace{v \otimes v \otimes \cdots \otimes v}^\alpha \otimes v'$; see [15, Section 3.6]. Therefore, by Theorem 3.11, we have $\inf_{x \in M} (f_v + b_p)(x) = \inf_{x \in M} (1/\alpha)(\alpha f_v + b_{\nu \cdot \mathcal{F}})(x) > -\infty \Leftrightarrow 0 \in B((\alpha f_v + b_{\nu \cdot \mathcal{F}})^\infty) \Leftrightarrow 0 \in B(f_v^\infty + b_p^\infty) \Leftrightarrow p \in B(f_v^\infty)$. \square

Summarizing, we obtain a convex analysis formulation of *p-scalability*.

Theorem 3.15. For $p \in CM^\infty$, the following conditions are equivalent:

- (a) $\inf_{x \in M} \|\nabla(f_v + b_p)(x)\|_x = 0$.
- (b) $-f^*(p) = \inf_{x \in M} (f_v + b_p)(x) > -\infty$.
- (c) $p \in B(f_v^\infty)$.

We finally consider the moment polytope membership. Here, $\overline{s\nabla^\infty f_v(M)}$ is nothing but the *moment polytope* for π, v in the sense of [15].⁴

Lemma 3.16. $\overline{s\nabla^\infty f_v(M)} = \bigcup_{g \in G} it^+ \cap B(f_{\pi(g)v}^\infty)$.

Proof. Proposition 2.33 holds in this setting by replacing $\text{diag } \lambda$ with $\lambda \in it^+$ and bk with Iwasawa decomposition bk ($b \in B, k \in K$). By (2.36), it suffices to show that $it^+ \cap B(f_{\pi(g)v}^\infty) = s(C \cap B(f_v^\infty))$ for $g \in G$ and Weyl chamber $C = g^\dagger(it^+)$. Indeed, from $f_{\pi(g)v}^\infty(u) = f_v^\infty(g^\dagger u)$ (by (3.18)), we have $\lambda \in it^+ \cap B(f_{\pi(g)v}^\infty) \Leftrightarrow \langle u, \lambda \rangle \leq f_{\pi(g)v}^\infty(u)$ ($\forall u \in M^\infty$) $\Leftrightarrow \langle g^\dagger u, g^\dagger \lambda \rangle \leq f_v^\infty(g^\dagger u)$ ($\forall u \in M^\infty$) $\Leftrightarrow \langle u', g^\dagger \lambda \rangle \leq f_v^\infty(u')$ ($\forall u' \in M^\infty$) $\Leftrightarrow g^\dagger \lambda \in B(f_v^\infty) \Leftrightarrow \lambda \in s(C \cap B(f_v^\infty))$, where $g^\dagger \lambda$ is written as $\lambda \cdot \mathcal{F}$ for $\mathcal{F} \in G/B$ corresponding to C . \square

The polytope $it^+ \cap B(f_{\pi(g)v}^\infty) = s(C \cap B(f_v^\infty))$ is what should be called the *Borel polytope*; see [14]. Notice that the approximate (λ, μ) -scalability of the previous section is nothing but the moment polytope membership $(\lambda, \mu) \in \overline{s\nabla^\infty f_A(P_n \times P_n)}$.

The convexity theorem of the moment polytope says:

Theorem 3.17 (Convexity theorem [24]). *The moment polytope $\overline{s\nabla^\infty f_v(M)}$ is a rational convex polytope.*

The *shifting trick* [12, 39] reduces the membership of the moment polytope to a single optimization problem.

Theorem 3.18 (Shifting trick [12, 39]). *A rational vector $\lambda \in it^+$ belongs to $\overline{s\nabla^\infty f_v(M)}$ if and only if $\inf_{x \in M} (f_{\pi(g)v} + b_{e\lambda})(x) > -\infty$ for generic $g \in G$.*

We prove a slightly stronger statement from our formulation, which implies Theorems 3.17 and 3.18.

Theorem 3.19. $\overline{s\nabla^\infty f_v(M)} = it^+ \cap B(f_{\pi(g)v}^\infty)$ for generic $g \in G$.

Our proof is a direct adaptation of [19, Lemma 54 and Proposition 55].

Proof. Let $\Lambda \subseteq it^+$ be a finite set in Proposition 3.13. Let $g \in G$. Then $it^+ \cap B(f_{\pi(g)v}^\infty)$ is the set of $\lambda \in it^+$ satisfying

$$\langle \nu \cdot \mathcal{F}, \lambda \rangle \leq f_{\pi(g)v}^\infty(\nu \cdot \mathcal{F}) \quad (\nu \in \Lambda, \mathcal{F} \in G/B). \quad (3.21)$$

We use the notation $\lambda = \bar{\lambda} \cdot [k_0]$ to deduce an explicit inequality description. Represent $\nu \cdot \mathcal{F}$ as $\nu \cdot \mathcal{F} = \bar{\nu} \cdot [g^{-\dagger} h^\dagger k_0]$ for $h \in G$. Then $\langle \nu \cdot \mathcal{F}, \lambda \rangle = \langle \bar{\nu} \cdot [g^{-\dagger} h^\dagger k_0], \bar{\lambda} \cdot [k_0] \rangle = \langle \bar{\nu} \cdot [k_0^\dagger g^{-\dagger} h^\dagger k_0], \bar{\lambda} \cdot [I] \rangle$, and $f_{\pi(g)v}^\infty(\nu \cdot \mathcal{F}) = f_{\pi(g)v}^\infty(\bar{\nu} \cdot [g^{-\dagger} h^\dagger k_0]) = f_{\pi(h)v}^\infty(\bar{\nu})$, where

⁴This fact can be seen from Proposition 2.33 and the fact that the *moment map* $\mu : \pi(G)v \setminus \{0\} \rightarrow i\mathfrak{u}(= T_I)$ in [15] is written as $\mu(\pi(g)v) = g^\dagger df_v(gg^\dagger)g$.

$[I] = \{E_i\}$ is the standard flag. Let $\mathcal{U}^{g,h} = \{U_j^{g,h}\}$ be the flag generated by $k_0^\dagger g^{-\dagger} h^\dagger k_0$. By Lemma 2.34, (3.21) is written as

$$\sum_{i,j=1}^n (\bar{\lambda}_i - \bar{\lambda}_{i+1})(\bar{\nu}_j - \bar{\nu}_{j+1}) \dim E_i \cap U_j^{g,h} \leq f_{\pi(h)v}^\infty(\nu) \quad (\nu \in \Lambda, h \in G). \quad (3.22)$$

We next consider the quantities $f_{\pi(h)v}^\infty(\nu)$ and $\dim E_i \cap U_j^{g,h}$ involving h . By (3.18), the former quantity $f_{\pi(h)v}^\infty(\nu)$ takes a value from finite set $A_\nu := \{\text{tr } \nu \omega \mid \omega \in \Omega(\pi)\}$. For $\alpha \in A_\nu$, let $G_{\nu,\alpha} \subseteq G$ be the affine subvariety of consisting of h with $f_{\pi(h)v}^\infty(\nu) \leq \alpha$, which is defined by algebraic conditions $(\pi(h)v)_\omega = 0$ for all $\omega \in \Omega(\pi)$ with $\text{tr } \nu \omega > \alpha$. The latter quantity $\dim E_i \cap U_j^{g,h}$ takes a value in $\{0, 1, 2, \dots, n\}$. Let D denote the set of all $n \times n$ matrices $d = (d_{ij})$ such that each entry d_{ij} is one of $0, 1, 2, \dots, n$, where the partial order \leq on D is defined by $d \leq d' \Leftrightarrow d_{ij} \leq d'_{ij} \ (\forall i, j)$. For $\nu \in \Lambda, \alpha \in A_\nu$, let $D_{\nu,\alpha}(g) \subseteq D$ be the set of all $d = (d_{ij})$ such that there is $h \in G_{\nu,\alpha}$ such that $d_{ij} = \dim E_i \cap U_j^{g,h}$ for $1 \leq i, j \leq n$, where $d_{nj} = d_{jn} = j$ holds for all $j \in [n]$. Let $\bar{D}_{\nu,\alpha}(g) \subseteq D_{\nu,\alpha}(g)$ denote the set of maximal members with respect to \leq . Then (3.22) is written as

$$\sum_{i,j=1}^n (\bar{\lambda}_i - \bar{\lambda}_{i+1})(\bar{\nu}_j - \bar{\nu}_{j+1}) d_{ij} \leq \alpha \quad (\nu \in \Lambda, \alpha \in A_\nu, d \in \bar{D}_{\nu,\alpha}(g)). \quad (3.23)$$

Let $S_{\nu,\alpha,d}$ be the subvariety of $G \times G_{\nu,\alpha}$ consisting of g, h with $\dim E_i \cap U_j^{g,h} \geq d_{ij}$ for i, j , which is defined by vanishing of subdeterminants of $k_0^\dagger g^{-\dagger} h^\dagger k_0$. Indeed, $\dim E_i \cap U_j^{g,h}$ is j minus the rank of lower left $(n-i) \times j$ submatrix $k_0^\dagger g^{-\dagger} h^\dagger k_0$. Let π be the projection $(g, h) \mapsto g$. We claim:

(*) $\pi(S_{\nu,\alpha,d})$ is an affine subvariety of G .

The proof is given in the end. Let $D_{\nu,\alpha}^* \subseteq D$ be the set of all maximal d with $\pi(S_{\nu,\alpha,d}) = G$. Consider the set Q^* of $\lambda \in \mathfrak{it}^+$ satisfying

$$\sum_{i,j=1}^n (\bar{\lambda}_i - \bar{\lambda}_{i+1})(\bar{\nu}_j - \bar{\nu}_{j+1}) d_{ij} \leq \alpha \quad (\nu \in \Lambda, \alpha \in A_\nu, d \in D_{\nu,\alpha}^*). \quad (3.24)$$

For $d \in D_{\nu,\alpha}^*$, there is $d' \in D_{\nu,\alpha}(g)$ with $d \leq d'$. That is, (3.24) is looser than (3.23). Thus, Q^* contains $\mathfrak{it}^+ \cap B(f_{\pi(g)v}^\infty)$ for every $g \in G$. Consider the finite union $H := \bigcup_{\nu \in \Lambda, \alpha \in A_\nu, d \in D \setminus D_{\nu,\alpha}^*} \pi(S_{\nu,\alpha,d})$, which is a proper subvariety of G . Then we can choose a generic $g^* \in G \setminus H$. For such g^* , it must hold $\bar{D}_{\nu,\alpha}(g^*) = D_{\nu,\alpha}^*$ for all $\nu \in \Lambda, \alpha \in A_\nu$. This means $Q^* = \mathfrak{it}^+ \cap B(f_{\pi(g^*)v}^\infty) = \bigcup_{g \in G} \mathfrak{it}^+ \cap B(f_{\pi(g)v}^\infty) = \overline{s\nabla^\infty f_v(M)}$.

Finally we verify (*). By the closure theorem (see [16, Section 4.7, Theorem 7]), $\pi(G_{\nu,\alpha,d})$ is a constructible set, i.e., it is an affine variety Z_0 minus an affine variety Z' . We show that $\pi(G_{\nu,\alpha,d})$ is a closed set in the Euclidean topology, which implies that $\pi(G_{\nu,\alpha,d}) = Z_0$ is an affine variety. Consider a sequence g_1, g_2, \dots in $\pi(G_{\nu,\alpha,d})$ converging to $g \in G$. For each i , there is $h_k \in G$ with such that $f_{\pi(h_k)v}^\infty(\nu) \leq \alpha$ and $\dim E_i \cap U_j^{g,h_k} \geq d_{ij}$ ($i, j \in [n]$). These quantities are determined by $\bar{\nu} \cdot [g^{-\dagger} h_k^\dagger k_0] = g^{-\dagger} h_k^\dagger \nu$. By Iwasawa decomposition, h_k can be chosen from the compact group K . By taking a subsequence, we may assume that h_k converges to $h \in K$. From $h_k \in G_{\nu,\alpha}$ for each k , it is clear that

$h \in G_{\nu,\alpha}$. As mentioned, the condition $\dim E_i \cap U_j^{g_k, h_k} \geq d_{ij}$ is written as vanishing of subdeterminants of $k_0^\dagger g_k^{-\dagger} h_k^\dagger k_0$. These subdeterminants vanish in the limit $k_0^\dagger g^{-\dagger} h^\dagger k_0$ as well. Then $\dim E_i \cap U_j^{g, h} \geq d_{ij}$. Thus $(g, h) \in S_{\nu,\alpha,d}$, and $\pi(S_{\nu,\alpha,d})$ is closed. \square

In particular, $B(f_v^\infty)$ contains the moment polytope in a “generic” chamber. Thus, the moment polytope membership for a given vector $\lambda \in \mathfrak{it}^+$ also reduces, after taking generic $g \in G$, to the convex optimization problem on Euclidean building CM^∞ :

$$\inf. \quad f_{\pi(g)v}^\infty(u) - \langle p, u \rangle \quad \text{s.t.} \quad u \in U, \quad (3.25)$$

where $p := e^{\infty\lambda} \in CM^\infty$ and U is any convex neighborhood of 0.

This gives rise to a challenging research problem to develop algorithms solving convex optimization problems (3.19), (3.25). On a single Weyl chamber C (or an apartment), it is a usual Euclidean convex optimization. However, at a boundary point q of C , one have to search a descent direction from infinitely many Weyl chambers containing q . This seems impossible in principle. So one have to exploit and utilize special properties of the objective function, particularly, the recession function of the Kempf-Ness function, as in [26]. Moreover, to keep variable $q = \lambda \cdot \mathcal{F}$, one should keep basis vectors of flag \mathcal{F} with bounded bit-length. This is also a highly nontrivial problem. A recent work [20] for finding a violating vector subspace in (3.13) may give hints toward this direction.

Acknowledgments

The author thanks Hiroyuki Ochiai for helpful discussion, and thanks for Zhiyuan Zhan for corrections. The author also thanks Harold Nieuwboer and Michael Walter for discussion on the Legendre-Fenchel duality. The work was partially supported by JST PRESTO Grant Number JPMJPR192A, Japan.

References

- [1] P. Abramenko and K. S. Brown, *Buildings—Theory and Applications*. Springer, New York, 2008.
- [2] Z. Allen-Zhu, A. Garg, Y. Li, R. Oliveira, and A. Wigderson, Operator scaling via geodesically convex optimization, invariant theory and polynomial identity testing. [arXiv:1804.01076](https://arxiv.org/abs/1804.01076), 2018, the conference version in STOC 2018.
- [3] S. Amari and K. Nagaoka, *Methods of Information Geometry*. American Mathematical Society, Providence, RI, 2000.
- [4] R. Bergmann, R. Herzog, M. S. Louzeiro, D. Tenbrinck, and J. Vidal-Núñez, Fenchel duality theory and a primal-dual algorithm on Riemannian Manifolds. *Foundations of Computational Mathematics* **2** (2021), 1465–1504, 2021.
- [5] M. Bačák, *Convex Analysis and Optimization in Hadamard Spaces*. De Gruyter, Berlin, 2014.
- [6] W. Ballmann, *Lectures on Spaces of Nonpositive Curvature*. Birkhäuser Verlag, Basel, 1995.

- [7] W. Ballmann, M. Gromov, and V. Schroeder *Manifolds of Nonpositive Curvature*, Birkhäuser, Boston MA, 1985.
- [8] J. Bennett, A. Carbery, M. Christ, and T. Tao, The Brascamp-Lieb inequalities: finiteness, structure and extremals. *Geometric and Functional Analysis* **17** (2007) 1343–1415.
- [9] N. Boumal, *An Introduction to Optimization on Smooth Manifolds*. Cambridge University Press, Cambridge, 2023.
- [10] H. Brascamp and E. Lieb, Best constants in Young’s inequality, its converse, and its generalization to more than three functions. *Advances in Mathematics***20** (1976), 151–173.
- [11] M. R. Bridson and A. Haefliger, *Metric Spaces of Non-positive Curvature*. Springer-Verlag, Berlin, 1999.
- [12] M. Brion, Sur l’image de l’application moment. In: M. -P. Malliavin (eds): *Séminaire d’algèbre Paul Dubreil et Marie-Paule Malliavin (Paris, 1986)*, *Lecture Notes in Mathematics* **1296**, Springer, Berlin (1987), pp 177–192.
- [13] P. Bürgisser, A. Garg, R. Oliveira, M. Walter, and A. Wigderson, Alternating minimization, scaling algorithms, and the null-cone problem from invariant theory, [arXiv:1711.08039](#), 2017, the conference version in ITCS 2018.
- [14] P. Bürgisser, C. Franks, A. Garg, R. Oliveira, M. Walter, and A. Wigderson, Efficient algorithms for tensor scaling, quantum marginals and moment polytopes. [arXiv:1804.04739](#), 2018, the conference version in FOCS 2018.
- [15] P. Bürgisser, C. Franks, A. Garg, R. Oliveira, M. Walter, and A. Wigderson, Towards a theory of non-commutative optimization: geodesic first and second order methods for moment maps and polytopes. [arXiv:1910.12375](#), 2019, the conference version in FOCS 2019.
- [16] D. A. Cox, J. Little, D. O’Shea, *Ideals, Varieties, and Algorithms, 4th edition*, Springer, Cham, 2015.
- [17] P. B. Eberlein, *Geometry of Nonpositively Curved Manifolds*. University of Chicago Press, Chicago, IL, 1996.
- [18] M. Fortin and C. Reutenauer, Commutative/non-commutative rank of linear matrices and subspaces of matrices of low rank. *Séminaire Lotharingien de Combinatoire* **52** (2004), B52f.
- [19] C. Franks, Operator scaling with specified marginals. [arXiv:1801.01412](#), 2018, the conference version in STOC 2018.
- [20] C. Franks, T. Soma, and M. X. Goemans, Shrunk subspaces via operator Sinkhorn iteration. [arXiv:2207.08311](#), 2022, the conference version in SODA 2023.
- [21] S. Fujishige, *Submodular Functions and Optimization, 2nd Edition*. Elsevier, Amsterdam, 2005.
- [22] A. Garg, L. Gurvits, R. Oliveira, and A. Wigderson, Operator scaling: theory and applications. *Foundations of Computational Mathematics* **20** (2020), 223–290.

- [23] A. Garg, L. Gurvits, R. Oliveira, and A. Wigderson, Algorithmic and optimization aspects of Brascamp-Lieb inequalities, via Operator Scaling. *Geometric and Functional Analysis* **28** (2018) 100–145.
- [24] V. Guillemin and S. Sternberg, Convexity properties of the moment mapping, *Inventiones Mathematicae* **67** (1982) 491–513.
- [25] L. Gurvits, Classical complexity and quantum entanglement, *Journal of Computer and System Sciences* **69** (2004), 448–484.
- [26] M. Hamada and H. Hirai, Computing the nc-rank via discrete convex optimization on CAT(0) spaces, *SIAM Journal on Applied Geometry and Algebra* **5** (2021), 455–478.
- [27] H. Hirai, L-convexity on graph structures. *Journal of the Operations Research Society of Japan* **61** (2018), 71–109.
- [28] H. Hirai, H. Nieuwboer, and M. Walter, Interior-point methods on manifolds: theory and applications, [arXiv:2303.04771](https://arxiv.org/abs/2303.04771), 2023.
- [29] J.-B. Hiriart-Urruty and C. Lemaréchal, *Fundamentals of Convex Analysis*. Springer-Verlag, Berlin, 2001.
- [30] G. Ivanyos, Y. Qiao, and K. V. Subrahmanyam, Non-commutative Edmonds’ problem and matrix semi-invariants. *Computational Complexity* **26** (2017), 717–763.
- [31] M. Kapovich, B. Leeb, and J. Millson, Convex functions on symmetric spaces, side lengths of polygons and the stability inequalities for weighted configurations at infinity. *Journal of Differential Geometry* **81** (2009), 297–354.
- [32] G. Kempf and L. Ness, The length of vectors in representation spaces, In K. Lønsted (ed.) *Algebraic Geometry (Summer Meeting, Copenhagen, August 7–12, 1978)*, *Lecture Notes in Mathematics* **732**, Springer, Berlin, 1979, pp. 233–243.
- [33] B. Kleiner and B. Leeb, Rigidity of invariant convex sets in symmetric spaces. *Inventiones Mathematicae* **163** (2006), 657–676.
- [34] A. W. Knap, *Lie Groups Beyond an Introduction, Second Edition*, Birkhäuser, Boston, 2002.
- [35] E. Lieb, Gaussian kernels have only Gaussian maximizers. *Inventiones Mathematicae* **102** (1990), 179–208.
- [36] M. S. Louzeiro, R. Bergmann, and R. Herzog, Fenchel duality and a separation theorem on Hadamard manifolds. *SIAM Journal on Optimization* **32** (2022), 854–873.
- [37] L. Lovász, Submodular functions and convexity. In A. Bachem, M. Grötschel, and B. Korte (eds.): *Mathematical Programming—The State of the Art* (Springer-Verlag, Berlin, 1983), 235–257.
- [38] K. Murota, *Discrete Convex Analysis*. SIAM, Philadelphia, 2004.
- [39] L. Ness and D. Mumford, A stratification of the null cone via the moment map. *American Journal of Mathematics* **106** (1984), 1281–1329.
- [40] R. T. Rockafellar, *Convex Analysis*, Princeton University Press, NJ, 1970.

- [41] U. G. Rothblum and H. Schneider, Scalings of matrices which have prespecified row sums and column sums via optimization. *Linear Algebra and Its Applications* **114/115** (1989), 737–764.
- [42] T. Sakai, *Riemannian Geometry*, American Mathematical Society, Providence RI, 1996.
- [43] A. Schrijver, *Combinatorial Optimization—Polyhedra and Efficiency*. Springer, Berlin, 2003.
- [44] R. Sinkhorn, A relationship between arbitrary positive matrices and doubly stochastic matrices. *Annals of Mathematics Statistics* **35** (1964), 876–879.
- [45] N. R. Wallach, *Geometric Invariant Theory*. Springer, Cham, 2017.
- [46] C. Woodward, Moment maps and geometric invariant theory, [arXiv:0912.1132](https://arxiv.org/abs/0912.1132), 2009.