

G-STABLE RANK OF SYMMETRIC TENSORS AND LOG CANONICAL THRESHOLD

ZHI JIANG

ABSTRACT. Shitov recently gave a counterexample to Comon's conjecture that the symmetric tensor rank and tensor rank of a symmetric tensor are the same. In this paper we show that an analog of Comon's conjecture for the G -stable rank introduced by Derksen is true: the symmetric G -stable rank and G -stable rank of a symmetric tensor are the same. We also show that the log-canonical threshold of a singularity is bounded by the G -stable rank of the defining ideal.

1. INTRODUCTION

An order d tensor is a vector in a tensor product of d vector spaces. There are several generalizations of the rank of a matrix to tensors of order ≥ 3 , for example the tensor rank, border rank, sub-rank, slice rank and G -stable rank. A simple tensor in $V_1 \otimes V_2 \otimes \cdots \otimes V_d$ is a tensor of the form $v_1 \otimes v_2 \otimes \cdots \otimes v_d$, where $v_i \in V_i$. The tensor rank of $T \in V_1 \otimes V_2 \otimes \cdots \otimes V_d$ is the smallest number of simple tensors that sum up to T . The G -stable rank of a tensor was introduced by Derksen in [2]. The slice rank and G -stable rank have been used to find bounds for the cap set problem (see [2], [3], [4], [10]).

If $V_1 = V_2 = \cdots = V_d = V$ then there is a natural action of the symmetric group S_d on $V^{\otimes d} = V \otimes V \otimes \cdots \otimes V$. A tensor invariant under this action is called a symmetric tensor of order d . The Waring rank or symmetric rank of a symmetric tensor $T \in V^{\otimes d}$ is the smallest number d such that T can be written as a sum of d tensors of the form $v^{\otimes d} = v \otimes v \otimes \cdots \otimes v$. It is clear that the tensor rank is less than or equal to the symmetric rank. It was conjectured by Comon [1] that the symmetric rank and tensor rank of a symmetric tensor are equal. Recently, Shitov gave a counterexample [9]. In this paper, we study the notion of G -stable rank of a tensor. The G -stable rank of a tensor is defined in terms of geometric invariant theory and the notion of stability for algebraic group actions on tensors. It is also natural to define a symmetric G -stable rank for a symmetric tensor. One main result of this paper is that the symmetric G -stable rank and G -stable rank of a symmetric tensor are the same.

In algebraic geometry and singularity theory, the log canonical threshold is an important invariant of singularities. We will show that the symmetric G -stable rank and the log canonical threshold are closely related. We extend the notion of G -stable rank to ideals in a coordinate ring of a smooth irreducible affine variety. In this context, we show that the log canonical threshold is less than or equal to the G -stable rank. In the case of monomial ideals in the polynomial ring we show equality.

1.1. Stability of tensors. Let K be a perfect field, and V a finite dimensional vector space over K , we consider the action of the group of product of special linear groups $\mathrm{SL}(V)^d = \mathrm{SL}(V) \times \mathrm{SL}(V) \times \cdots \times \mathrm{SL}(V)$ on the tensor product space $V^{\otimes d} = V \otimes V \otimes \cdots \otimes V$. A 1-parameter subgroup of an algebraic group G is a homomorphism of algebraic groups $\lambda : \mathbb{G}_m \rightarrow G$, where \mathbb{G}_m is the multiplicative group. For any integer m , we define the multiple of

λ by m , denoted by $m \cdot \lambda$, which is also a 1-parameter subgroup with $(m \cdot \lambda)(t) = (\lambda(t))^m$. A 1-parameter subgroup is *indivisible* if it is not a multiple of any other 1-parameter subgroup with factor $m \geq 2$. We say a tensor $v \in V^{\otimes d}$ is $\mathrm{SL}(V)^d$ -*unstable* if there is a 1-parameter subgroup $\lambda : \mathbb{G}_m \rightarrow \mathrm{SL}(V)^d$, such that

$$\lim_{t \rightarrow 0} \lambda(t) \cdot v = 0.$$

If no such 1-parameter subgroup exists, then v is called $\mathrm{SL}(V)^d$ -*semistable*. Let G be a reductive algebraic group over K . By a G -scheme X we mean a separated, finite type scheme X over K as well as a morphism $G \times X \rightarrow X$ mapping (g, x) to $g \cdot x$, such that $g \cdot (h \cdot x) = (gh) \cdot x$, for all $g, h \in G$ and for all $x \in X$. A morphism $f : X \rightarrow Y$ between two G -schemes X and Y is G -equivariant if for all $g \in G$ and $x \in X$, we have $f(g \cdot x) = g \cdot f(x)$. A subscheme S of a G -scheme X is called a G -subscheme if S is a G -scheme and the immersion $S \hookrightarrow X$ is G -equivariant.

Throughout this paper, we will work over a perfect field K . In [5], Kempf proved a K -rational version of the *Hilbert-Mumford* criterion:

Theorem 1 ([5], Corollary 4.3). *Let G be a reductive algebraic group. Suppose that X is a G -scheme and $x \in X$ is a K -point. Assume S is a closed G -subscheme of X which does not contain x and S meets the closure of the orbit $G \cdot x$. Then there exists a K -rational 1-parameter subgroup $\lambda : \mathbb{G}_m \rightarrow G$, such that*

$$\lim_{t \rightarrow 0} \lambda(t) \cdot x \in S.$$

If $G = \mathrm{SL}(V)^d$, $X = V^{\otimes d}$, then by Theorem 1, v is unstable if and only if 0 is in the closure of the orbit of v , i.e. $0 \in \overline{\mathrm{SL}(V)^d \cdot v}$.

A tensor $T \in V^{\otimes d}$ is called symmetric if it is invariant under the action of symmetric group S_d . Let $D^d V \subseteq V^{\otimes d}$ be the space of symmetric tensors. As a representation of $\mathrm{GL}(V)$, this is the space of divided powers, which is isomorphic to the d -th symmetric power $S^d V$ if the characteristic of K is 0 or $> d$.

It is interesting to look at the diagonal action of $\mathrm{SL}(V)$ on $D^d V$ via the diagonal embedding:

$$(1) \quad \Delta : \mathrm{SL}(V) \hookrightarrow \mathrm{SL}(V)^d.$$

Definition 2. A symmetric tensor $v \in D^d V$ is $\mathrm{SL}(V)$ -*unstable* if there is a 1-parameter subgroup $\lambda : \mathbb{G}_m \rightarrow \mathrm{SL}(V)$, such that

$$\lim_{t \rightarrow 0} \lambda(t) \cdot v = 0.$$

Otherwise we say v is $\mathrm{SL}(V)$ -*semistable*.

1.2. G-stable rank for tensors. In [2], Derksen introduced G -stable rank for tensors. Suppose the base field K is perfect. If $\lambda : \mathbb{G}_m \rightarrow \mathrm{GL}_n$ is a 1-parameter subgroup, then we can view $\lambda(t)$ as an invertible $n \times n$ matrix whose entries lie in the ring $K[t, t^{-1}]$ of Laurent polynomials. We say that $\lambda(t)$ is a polynomial 1-parameter subgroup of GL_n if all these entries lie in the polynomial ring $K[t]$. Consider the action of the group $G = \mathrm{GL}(V_1) \times \mathrm{GL}(V_2) \times \cdots \times \mathrm{GL}(V_d)$ on the tensor product space $W = V_1 \otimes V_2 \otimes \cdots \otimes V_d$. A 1-parameter subgroup $\lambda : \mathbb{G}_m \rightarrow G$ can be written as

$$\lambda(t) = (\lambda_1(t), \cdots, \lambda_d(t)),$$

where $\lambda_i(t)$ is a 1-parameter subgroup of $\mathrm{GL}(V_i)$ for all i . We say that $\lambda(t)$ is polynomial if and only if $\lambda_i(t)$ is a polynomial 1-parameter subgroup for all i .

The t -valuation $\mathrm{val}(a(t))$ of a polynomial $a(t) \in K[t]$ is the biggest integer n such that $a(t) = t^n b(t)$ for some $b(t) \in K[t]$. For $a(t), b(t) \in K[t]$, the t -valuation $\mathrm{val}\left(\frac{a(t)}{b(t)}\right)$ of the rational function $\frac{a(t)}{b(t)} \in K(t)$ is $\mathrm{val}\left(\frac{a(t)}{b(t)}\right) = \mathrm{val}(a(t)) - \mathrm{val}(b(t))$. For a tuple $u(t) = (u_1(t), u_2(t), \dots, u_d(t)) \in K(t)^d$, we define the t -valuation of $u(t)$ as

$$(2) \quad \mathrm{val}(u(t)) = \min_i \{\mathrm{val}(u_i(t)) \mid 1 \leq i \leq d\}.$$

If λ is a 1-parameter subgroup of G and $v \in W$ is a tensor, then we have $\lambda(t) \cdot v \in K(t) \otimes W$. We view $K(t) \otimes W$ as a vector space over $K(t)$ and define the t -valuation $\mathrm{val}(\lambda(t) \cdot v)$ as in (2). Assume $\mathrm{val}(\lambda(t) \cdot v) > 0$, then for any $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_d) \in \mathbb{R}_{>0}^d$, we define the slope

$$(3) \quad \mu_\alpha(\lambda(t), v) = \frac{\sum_{i=1}^d \alpha_i \mathrm{val}(\det(\lambda_i(t)))}{\mathrm{val}(\lambda(t) \cdot v)}.$$

The G -stable rank for $v \in W$ is the infimum of the slope with respect to all such 1-parameter subgroups. More precisely:

Definition 3 ([2], Theorem 2.4). If $\alpha \in \mathbb{R}_{>0}^d$, then the G -stable rank $\mathrm{rk}_\alpha^G(v)$ is the infimum of $\mu_\alpha(\lambda(t), v)$ where $\lambda(t)$ is a polynomial 1-parameter subgroup of $G = \mathrm{GL}(V_1) \times \dots \times \mathrm{GL}(V_n)$ and $\mathrm{val}(\lambda(t) \cdot v) > 0$. If $\alpha = (1, 1, \dots, 1)$, we simply write $\mathrm{rk}^G(v)$.

Let V be a finitely dimensional vector space over K . Let $D^d V \subset V^{\otimes d}$ be the space of all symmetric tensors. Assume the group $\mathrm{GL}(V)$ acts on $D^d V$ via the diagonal embedding: $\mathrm{GL}(V) \hookrightarrow \mathrm{GL}(V)^d$.

Definition 4. Let $v \in D^d V$ be a symmetric tensor, the symmetric G -stable rank $\mathrm{symmrk}^G(v)$ of v is the infimum of $\mu(\lambda(t), v) = d \frac{\mathrm{val}(\det(\lambda(t)))}{\mathrm{val}(\lambda(t) \cdot v)}$, where $\lambda(t)$ is a polynomial 1-parameter subgroup of $\mathrm{GL}(V)$ and $\mathrm{val}(\lambda(t) \cdot v) > 0$.

Since any 1-parameter subgroup of $\mathrm{GL}(V)$ is also a 1-parameter subgroup of $\mathrm{GL}(V)^d$ via the diagonal embedding, we have $\mathrm{symmrk}^G(v) \geq \mathrm{rk}^G(v)$ for any $v \in D^d V$. It turns out that the other inequality is also true,

Theorem 5. *Let $v \in D^d V$ be a symmetric tensor, then we have*

$$\mathrm{symmrk}^G(v) = \mathrm{rk}^G(v).$$

Example 6. Suppose that $V = K^2$, and $v = e_2 \otimes e_1 \otimes e_1 + e_1 \otimes e_2 \otimes e_1 + e_1 \otimes e_1 \otimes e_2 \in V^{\otimes 3}$, where $\{e_1, e_2\}$ is the standard basis of $V = K^2$. Let $\lambda(t) = \begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix}$ be a polynomial 1-parameter subgroup of $\mathrm{GL}(K^2)$. Then $\lambda(t) \cdot v = t^2 v$, $\det(\lambda(t)) = t$, the slope is

$$\mu(\lambda(t), v) = 3 \frac{\mathrm{val}(\det(\lambda(t)))}{\mathrm{val}(\lambda(t) \cdot v)} = \frac{3}{2}.$$

Therefore we have $\mathrm{symmrk}^G(v) \leq \frac{3}{2}$. It was proved in [2] that $\mathrm{rk}^G(v) = \frac{3}{2}$. Hence by the fact $\mathrm{symmrk}^G(v) \geq \mathrm{rk}^G(v)$ we have $\mathrm{symmrk}^G(v) = \frac{3}{2}$.

A 1-parameter subgroup of $\mathrm{SL}(V)$ is also a 1-parameter subgroup of $\mathrm{SL}(V)^d$ via the diagonal embedding. It follows that if a symmetric tensor $v \in D^d V$ is $\mathrm{SL}(V)$ -unstable, then v is also $\mathrm{SL}(V)^d$ -unstable. Equivalently, if v is $\mathrm{SL}(V)^d$ -semistable, then v is also $\mathrm{SL}(V)$ -semistable. It follows from Theorem 5 that the converse direction is also true:

Corollary 7. *Let $v \in D^d V$ be a symmetric tensor, then v is $\mathrm{SL}(V)^d$ -semistable if and only if it is $\mathrm{SL}(V)$ -semistable.*

1.3. G -stable rank for ideals and log canonical threshold. Let V be an n -dimensional vector space over a perfect field K . By choosing a basis of V and a dual basis $\{x_1, x_2, \dots, x_n\}$ of V^* , we have an isomorphism of algebras $SV^* \cong K[x_1, \dots, x_n]$, where SV^* is the symmetric algebra on the vector space V^* . We have defined the symmetric G -stable rank for symmetric tensors, it is natural to extend this idea to polynomials and more generally to ideals in the polynomial ring $K[x_1, \dots, x_n]$. Furthermore, let X be a smooth irreducible affine variety with coordinate ring $K[X]$, and let $\mathfrak{a} \subset K[X]$ be an ideal. We can define the G -stable rank $\mathrm{rk}^G(P, \mathfrak{a})$ for the ideal \mathfrak{a} at a point $P \in V(\mathfrak{a})$. We postpone the precise definition of G -stable rank for ideals to Section 5. It turns out that the G -stable rank $\mathrm{rk}^G(P, \mathfrak{a})$ of an ideal \mathfrak{a} at P is closely related to the *log canonical threshold* $\mathrm{lct}_P(\mathfrak{a})$ of the ideal \mathfrak{a} at the point $P \in V(\mathfrak{a})$.

Log canonical threshold is an invariant of singularities in algebraic geometry, [6] gives a comprehensive introduction to this subject. Let $K = \mathbb{C}$ be the complex field. Let $H \subset \mathbb{C}^n$ be a hypersurface defined by a polynomial $f \in \mathbb{C}[x_1, \dots, x_n]$, and let $P \in H$ be a closed point. The log canonical threshold $\mathrm{lct}_P(f)$ of f at the point P tells us how singular f is at the point P . More precisely, $\mathrm{lct}_P(f)$ is a rational number bounded above by 1, and equal to 1 if P is a smooth point of H .

There are several equivalent ways to define the log canonical threshold, here we give an analytic definition, which we will use later.

Definition 8. Let X be a smooth irreducible affine variety. Let $\mathfrak{a} = (f_1, \dots, f_r) \subset \mathbb{C}[X]$ be an ideal, and $P \in V(\mathfrak{a})$ is a closed point. The *log canonical threshold* $\mathrm{lct}_P(\mathfrak{a})$ of the ideal \mathfrak{a} at P is

$$(4) \quad \mathrm{lct}_P(\mathfrak{a}) = \sup \left\{ s > 0 \mid \frac{1}{(\sum_{i=1}^r |f_i|^2)^s} \text{ is integrable around } P \right\}.$$

The log canonical threshold $\mathrm{lct}_P(f)$ of a polynomial $f \in \mathbb{C}[X]$ is the log canonical threshold of the principle ideal $\mathfrak{a} = (f)$. We have the following relation between the log canonical threshold and the G -stable rank:

Theorem 9. *In the situation of Definition 8, the log canonical threshold of \mathfrak{a} is less than or equal to the G -stable rank of \mathfrak{a} at P :*

$$(5) \quad \mathrm{lct}_P(\mathfrak{a}) \leq \mathrm{rk}^G(P, \mathfrak{a}).$$

When \mathfrak{a} is a monomial ideal, i.e. \mathfrak{a} is generated by monomials, the equality holds.

Theorem 10. *Suppose $\mathfrak{a} \subset \mathbb{C}[x_1, \dots, x_n]$ is a proper nonzero ideal generated by monomials and $P = (0, \dots, 0)$ is the origin. Then we have*

$$(6) \quad \mathrm{lct}_P(\mathfrak{a}) = \mathrm{rk}^G(P, \mathfrak{a}).$$

2. KEMPF'S THEORY OF OPTIMAL SUBGROUPS

Let G be a reductive algebraic group over a perfect field K . In our case, G is one of $\mathrm{SL}(V)^d$, $\mathrm{SL}(V)$, $\mathrm{GL}(V)^d$ or $\mathrm{GL}(V)$, depend on the situation. Let $\Gamma(G)$ denote the set of all 1-parameter subgroups of G . In [5], Kempf provided a way to approach the boundary of an orbit. Following [5], we have the definition:

Definition 11. Let X be a G -scheme over a perfect field K and let $x \in X$ be a K -point. We define $|X, x|$ to be the set of all 1-parameter subgroups of G such that $\lim_{t \rightarrow 0} \lambda(t) \cdot x$ exists in X . Assume S is a G -invariant closed sub-scheme of X not containing x , we define a subset $|X, x|_S \subset |X, x|$ by

$$(7) \quad |X, x|_S = \{\lambda \in |X, x| \mid \lim_{t \rightarrow 0} \lambda(t) \cdot x \in S\}.$$

Remark 12. If $S \cap \overline{G \cdot x} \neq \emptyset$, then by Theorem 1, there exists a 1-parameter subgroup $\lambda(t) \in \Gamma(G)$, such that $\lim_{t \rightarrow 0} \lambda(t) \cdot x \in S$. Hence $|X, x|_S \neq \emptyset$. If $X = V^{\otimes d}$, $S = 0$ and $v \in V^{\otimes d}$ is $\mathrm{SL}(V)^d$ -unstable, then $|V^{\otimes d}, v|_{\{0\}} \neq \emptyset$.

Let $\lambda \in |X, x|$ be a 1-parameter subgroup of G , we get a morphism $\phi_\lambda : \mathbb{A}^1 \rightarrow X$ by $\phi_\lambda(t) = \lambda(t) \cdot x$ if $t \neq 0$ and $\phi_\lambda(0) = \lim_{t \rightarrow 0} \lambda(t) \cdot x$. Assume S is a G -invariant closed sub-scheme of X not containing x , the inverse image $\phi_\lambda^{-1}(S)$ is an effective divisor supported inside $t = 0$. Let $a_{S,x}(\lambda)$ denote the degree of the divisor $\phi_\lambda^{-1}(S)$ for $\lambda \in |X, x|$. Note that we have a natural conjugate action of G on the set of 1-parameter subgroups $\Gamma(G)$ by $(g \cdot \lambda)(t) = g\lambda(t)g^{-1}$, where $g \in G$, $\lambda \in \Gamma(G)$.

Definition 13. A *length* function $\|\cdot\|$ is a non-negative real-valued function on $\Gamma(G)$ such that

- (1) $\|g \cdot \lambda\| = \|\lambda\|$ for any $\lambda \in \Gamma(G)$ and $g \in G$.
- (2) For any maximal torus $T \subseteq G$, we have $\Gamma(T) \subseteq \Gamma(G)$, the restriction of $\|\cdot\|$ on $\Gamma(T)$ is integral valued and extends to a *norm* on the vector space $\Gamma(T) \otimes_{\mathbb{Z}} \mathbb{R}$.

Remark 14. Such a length function exists. Let T be a maximal torus of G . Let N be the normalizer of T . Then the Weyl group with respect to T is defined by $W = N/T$. By the fact that $\Gamma(G)/G \cong \Gamma(T)/W$, it suffices to define a W -invariant norm on $\Gamma(T) \otimes_{\mathbb{Z}} \mathbb{R}$. Since W is a finite group, any norm on $\Gamma(T) \otimes_{\mathbb{Z}} \mathbb{R}$ and then average over W will work.

Remark 15. In the original paper [5], Kempf defined a length function $\|\cdot\|$ that satisfies a different condition (2): for any maximal torus T of G , there is a positive definite integral-valued bilinear form (\cdot, \cdot) on $\Gamma(T)$, such that $(\lambda, \lambda) = \|\lambda\|^2$ for any λ in $\Gamma(T)$. But the proof in [5] of the theorem below is also valid for our slightly weaker definition of length function.

Theorem 16 (Kempf [5]). *Let X be an affine G -scheme over a perfect field K . Let $x \in X$ be a K -point. Assume S is an G -invariant closed sub-scheme not containing x such that $S \cap \overline{G \cdot x} \neq \emptyset$. Fix a length function $\|\cdot\|$ on $\Gamma(G)$, then we have*

- (1) The function $\frac{a_{S,x}(\lambda)}{\|\lambda\|}$ has a maximum positive value $B_{S,x}$ on the set of non-trivial 1-parameter subgroups in $|X, x|$.
- (2) Let $\Lambda_{S,x}$ be the set of indivisible 1-parameter subgroups $\lambda \in |X, x|$ such that $a_{S,x}(\lambda) = B_{S,x} \cdot \|\lambda\|$, then we have
 - (a) $\Lambda_{S,x} \neq \emptyset$.
 - (b) For $\lambda \in \Lambda_{S,x}$, Let $P(\lambda) = \{g \in G \mid \lim_{t \rightarrow 0} \lambda(t) \cdot g \cdot \lambda(t)^{-1} \text{ exists}\}$, then $P(\lambda)$ is a parabolic subgroup and independent of λ . We denote it by $P_{S,x}$.

(c) Any maximal torus of $P_{S,x}$ contains a unique member of $\Lambda_{S,x}$.

3. G -STABLE RANK AND SYMMETRIC G -STABLE RANK

Let V be an n dimensional vector space over K , fix a maximal torus T of $\mathrm{GL}(V)$, we have an isomorphism $\Gamma(T) \cong \mathbb{Z}^n$. Any 1-parameter subgroup λ of the maximal torus T is given by a tuple of n integers (ν_1, \dots, ν_n) , we define a function on $\Gamma(T) \cong \mathbb{Z}^n$ by

$$(8) \quad \|\lambda\| = \sum_{i=1}^n |\nu_i|.$$

This function extends linearly to a norm on the vector space $\Gamma(T) \otimes_{\mathbb{Z}} \mathbb{R}$. The Weyl group of $\mathrm{GL}(V)$ with respect to T is the symmetric group S_n . It is clear that the function is invariant under the action of S_n by permutation, therefore by Remark 14, it defines a length function on $\Gamma(\mathrm{GL}(V))$. Let $G = \mathrm{GL}(V)^d$, fix a maximal torus $T_i \subset \mathrm{GL}(V)$ for each component of $\mathrm{GL}(V)^d$, then $T = T_1 \times \dots \times T_d$ is a maximal torus of G . We have $\Gamma(T) \cong (\mathbb{Z}^n)^d$, Let $\lambda = (\lambda_1, \dots, \lambda_d)$ be a 1-parameter subgroup of T , where

$$\lambda_i = (\lambda_{i,1}, \dots, \lambda_{i,n}), \quad \lambda_{i,j} \in \mathbb{Z} \text{ for all } j$$

is a tuple of n integers. We define a function on $\Gamma(T)$ by

$$(9) \quad \|\lambda\| = \sum_{i=1}^d \|\lambda_i\|,$$

where $\|\lambda_i\| = \sum_{j=1}^n |\lambda_{i,j}|$. This extends to a length function on $\Gamma(G) = \Gamma(\mathrm{GL}(V)^d)$.

Let $G = \mathrm{GL}(V)^d$, $X = V^{\otimes d}$ and $S = \{0\}$. Recall the definition of t -valuation in equation (2).

Lemma 17. *Let $v \in D^d V \subset V^{\otimes d}$ be a symmetric tensor, and $G = \mathrm{GL}(V)^d$ acts on $V^{\otimes d}$ in the usual way. If $\lambda(t)$ is a 1-parameter subgroup of G , then*

- (1) $|X, v| = \{\lambda \in \Gamma(G) \mid \mathrm{val}(\lambda(t) \cdot v) \geq 0\}$.
- (2) $|X, v|_{\{0\}} = \{\lambda \in \Gamma(G) \mid \mathrm{val}(\lambda(t) \cdot v) > 0\}$.
- (3) $a_{\{0\}, v}(\lambda) = \mathrm{val}(\lambda(t) \cdot v)$ for $\lambda \in |X, v|$.

Proof. This follows immediately from the definition. □

Lemma 18. *Let $v \in D^d V \subset V^{\otimes d}$ be a symmetric tensor, then the function $\frac{\mathrm{val}(\lambda(t) \cdot v)}{\|\lambda\|} : \Gamma(G) \rightarrow \mathbb{R}$ attains its maximal value at some 1-parameter subgroup $\lambda \in \Gamma(\mathrm{GL}(V)^d)$. There exists a maximal torus $T \subset \mathrm{GL}(V)^d$, such that $\lambda \in \Gamma(T)$ and under the isomorphism $\Gamma(T^d) \cong (\mathbb{Z}^n)^d$, we can write $\lambda = (\lambda_1, \dots, \lambda_d)$, where $\lambda_i = (\lambda_{i,1}, \dots, \lambda_{i,n})$ such that $\lambda_{i,j} \in \mathbb{Z}$ and $\lambda_{i,j} \geq 0$ for all $1 \leq i \leq d, 1 \leq j \leq n$, in other words, λ is a polynomial 1-parameter subgroup.*

Proof. By Theorem 16, the maximal value of $\frac{\mathrm{val}(\lambda(t) \cdot v)}{\|\lambda\|}$ exists. Let $T \subset G$ be a maximal torus and $\lambda \in \Gamma(T)$ such that the function attains its maximum at λ . Assume λ is of the form in the lemma and $\lambda_{i,j} < 0$ for some i and j . If we replace $\lambda_{i,j}$ by $-\lambda_{i,j}$, $\mathrm{val}(\lambda(t) \cdot v)$ never decreases and $\|\lambda\|$ does not change. Therefore by the maximality of $\frac{\mathrm{val}(\lambda(t) \cdot v)}{\|\lambda\|}$, the value $\frac{\mathrm{val}(\lambda(t) \cdot v)}{\|\lambda\|}$ does not change after the replacement. Hence without loss of generality we can assume all $\lambda_{i,j} \geq 0$. □

Recall that for a tensor $v \in V^{\otimes d}$ and a polynomial 1-parameter subgroup λ of $G = \mathrm{GL}(V)^d$ such that $\mathrm{val}(\lambda(t) \cdot v) > 0$, we have the slope function

$$(10) \quad \mu(\lambda(t), v) = \frac{\sum_{i=1}^d \mathrm{val}(\det(\lambda_i(t)))}{\mathrm{val}(\lambda(t) \cdot v)}.$$

Let $\lambda = (\lambda_1, \dots, \lambda_d) \in \Gamma(G)$ be a polynomial 1-parameter subgroup of $G = \mathrm{GL}(V)^d$, then by Lemma 18, $\sum_{i=1}^d \mathrm{val}(\det(\lambda_i(t)))$ is the restriction of the length function defined by equation (9). Let S_d be the symmetric group acting on $G = \mathrm{GL}(V)^d$ by permuting the d components. Then the length function defined by equation (9) is invariant under the action of S_d . From now on, fix this length function on $\Gamma(G)$. We have a corollary following from Theorem 16:

Corollary 19. *Let $v \in D^d V \subset V^{\otimes d}$ be a symmetric tensor. Let $\Lambda_{\{0\},v}$ be the set of indivisible 1-parameter subgroups $\lambda \in |V^{\otimes d}, v|$ such that $\frac{\mathrm{val}(\lambda(t) \cdot v)}{\|\lambda\|}$ attains the maximum value. Then we have*

- (1) $\Lambda_{\{0\},v}$ is invariant under S_d .
- (2) $P_{\{0\},v}$ is S_d invariant. In other words, $P_{\{0\},v} = P^d \subset \mathrm{GL}(V)^d$ for some parabolic subgroup $P \subset \mathrm{GL}(V)$.

Proof.

- (1) It is clear that $\frac{\mathrm{val}(\lambda(t) \cdot v)}{\|\lambda\|}$ is S_d invariant. Indeed, Let $\sigma \in S_d$, since $v \in D^d V$ is a symmetric tensor and $\|\cdot\|$ is S_d invariant, we have

$$\frac{\mathrm{val}((\sigma\lambda)(t) \cdot v)}{\|\sigma\lambda\|} = \frac{\mathrm{val}((\sigma\lambda)(t) \cdot (\sigma v))}{\|\sigma\lambda\|} = \frac{\mathrm{val}(\sigma(\lambda(t) \cdot v))}{\|\sigma\lambda\|} = \frac{\mathrm{val}(\lambda(t) \cdot v)}{\|\lambda\|}.$$

Therefore if $\lambda \in \Lambda_{\{0\},v}$, so is $\sigma(\lambda)$.

- (2) Since $G = \mathrm{GL}(V)^d$, the parabolic subgroup $P_{\{0\},v}$ is a product of parabolic subgroups of $\mathrm{GL}(V)$, the symmetric group S_d acts on $P_{\{0\},v}$ by permuting the components. Let $\lambda \in \Lambda_{\{0\},v}$, for any $\sigma \in S_d$, we have

$$\sigma(P_{\{0\},v}) = P(\sigma(\lambda)) = P_{\{0\},v}.$$

We used the fact that $P_{\{0\},v} = P(\lambda)$ is independent of $\lambda \in \Lambda_{\{0\},v}$ and $\sigma(\lambda) \in \Lambda_{\{0\},v}$. So $P_{\{0\},v}$ is S_d invariant and we can find a parabolic subgroup $P \subset \mathrm{GL}(V)$ such that $P_{\{0\},v} = P^d$. □

Let $T \subset P \subset \mathrm{GL}(V)$ be a maximal torus, then T^d is a maximal torus of $P^d = P_{\{0\},v}$. By (2.c) in Theorem 16 and Lemma 18, there is a polynomial 1-parameter subgroup $\lambda = (\lambda_1, \dots, \lambda_n)$ of $T^d \subset \mathrm{GL}(V)^d$, such that the slope function

$$\mu(\lambda(t), v) = \frac{\sum_{i=1}^d \mathrm{val}(\det(\lambda_i(t)))}{\mathrm{val}(\lambda(t) \cdot v)} = \frac{\|\lambda\|}{\mathrm{val}(\lambda(t) \cdot v)}$$

has a minimum value at λ . The minimal value of $\mu(\lambda(t), v)$ is by definition the G -stable rank $\mathrm{rk}^G(v)$ of v . In rest of the section, we fix such a maximal torus $T \subset \mathrm{GL}(V)$.

Let $\lambda = (\lambda_1, \dots, \lambda_d)$ be a polynomial 1-parameter subgroup of $T^d \subset \mathrm{GL}(V)^d$, then $\gamma = \prod_{i=1}^d \lambda_i$ is a polynomial 1-parameter subgroup of $\mathrm{GL}(V)$. Furthermore, γ acts on $v \in D^d V$ via the diagonal embedding $\mathrm{GL}(V) \hookrightarrow \mathrm{GL}(V)^d$. We have the following lemma:

Lemma 20. *For any symmetric tensor $v \in D^dV$, we have $\text{val}(\gamma(t) \cdot v) \geq d \cdot \text{val}(\lambda(t) \cdot v)$.*

Proof. Let $C = \text{val}(\lambda(t) \cdot v)$, we define a subspace W of $V^{\otimes d}$ as following

$$W = \{w \in V^{\otimes d} \mid \text{val}(\sigma(\lambda(t)) \cdot w) \geq C, \forall \sigma \in S_d\}.$$

Since $\text{val}(\sigma(\lambda(t)) \cdot v) = \text{val}(\sigma(\lambda(t) \cdot v)) = \text{val}(\lambda(t) \cdot v) = C$, we have $v \in W$. For any $\sigma \in S_d$, it is clear that $\sigma(\lambda(t)) \cdot W \subset t^C K[t] \cdot W$. We can write

$$\begin{aligned} \gamma(t) \cdot v &= (\prod_{i=1}^d \lambda_i, \dots, \prod_{i=1}^d \lambda_i) \cdot v \\ &= (\lambda_1, \lambda_2, \dots, \lambda_d)(\lambda_2, \lambda_3, \dots, \lambda_d, \lambda_1) \cdots (\lambda_d, \lambda_1, \dots, \lambda_{d-1}) \cdot v \\ &= (\lambda_1, \lambda_2, \dots, \lambda_d) \sigma(\lambda_1, \lambda_2, \dots, \dots, \lambda_d) \cdots \sigma^{d-1}(\lambda_1, \lambda_2, \dots, \lambda_d) \cdot v \\ &= \lambda \sigma(\lambda) \cdots \sigma^{d-1}(\lambda) \cdot v, \end{aligned}$$

where $\sigma \in S_d$ satisfies $\sigma(1) = 2, \sigma(2) = 3, \dots, \sigma(d) = 1$. Therefore $\gamma(t) \cdot v \in t^{dC} K[t] \cdot W$, hence we get $\text{val}(\gamma(t) \cdot v) \geq dC = d \cdot \text{val}(\lambda(t) \cdot v)$. \square

Next we prove that the symmetric G -stable rank is the same as the G -stable rank for symmetric tensors.

Proof of Theorem 5. Let T be the chosen maximal torus of $\text{GL}(V)$ as above. Let $\lambda = (\lambda_1, \dots, \lambda_d)$ be a polynomial 1-parameter subgroup of $T^d \subset \text{GL}(V)^d$ such that the slope function $\mu(\lambda(t), v)$ attains its minimum value. In other words, λ computes the G -stable rank $\text{rk}^G(v)$ of v :

$$\text{rk}^G(v) = \frac{\sum_{i=1}^d \text{val}(\det(\lambda_i(t)))}{\text{val}(\lambda(t) \cdot v)}.$$

Let $\gamma = \prod_{i=1}^d \lambda_i$ as above, then we have

$$\text{symmrk}^G(v) \leq \frac{\sum_{i=1}^d \text{val}(\det(\gamma(t)))}{\text{val}(\gamma(t) \cdot v)} \leq \frac{d \sum_{i=1}^d \text{val}(\det(\lambda_i(t)))}{d \cdot \text{val}(\lambda(t) \cdot v)} = \text{rk}^G(v).$$

On the other hand, it is clear that $\text{symmrk}^G(v) \geq \text{rk}^G(v)$. Therefore $\text{symmrk}^G(v) = \text{rk}^G(v)$, this completes the proof. \square

4. STABILITY OF SYMMETRIC TENSORS

As a result of Theorem 5, we prove Corollary 7, which says that for a symmetric tensor $v \in D^dV$, v is $\text{SL}(V)^d$ -semistable if and only if v is $\text{SL}(V)$ -semistable. It is clear that $\text{SL}(V)^d$ -semistability implies $\text{SL}(V)$ -semistability. To prove the other direction, we will use a result which relates semistability with G -stable rank.

Proposition 21 ([2], Proposition 2.6). *Suppose that $\alpha = (\frac{1}{n_1}, \dots, \frac{1}{n_d})$ where $n_i = \dim V_i$. For $v \in V_1 \otimes V_2 \otimes \cdots \otimes V_d$ we have $\text{rk}_\alpha^G(v) \leq 1$. Moreover, $\text{rk}_\alpha^G(v) = 1$ if and only if v is semistable with respect to the group $H = \text{SL}(V_1) \times \text{SL}(V_2) \times \cdots \times \text{SL}(V_d)$.*

If $v \in D^dV$ is a symmetric tensor, $\alpha = (1, 1, \dots, 1)$ and $n = \dim V$, then by the above proposition, $\text{rk}^G(v) = n$ if and only if v is $\text{SL}(V)^d$ -semistable. We have a similar result for symmetric G -stable rank.

Proposition 22. *For a symmetric tensor $v \in D^dV$, we have $\text{symmrk}^G(v) \leq n$, where $n = \dim V$. Moreover, $\text{symmrk}^G(v) = n$ if and only if v is $\text{SL}(V)$ -semistable.*

Proof. The first statement is clear from Theorem 5. If $\text{symmrk}^G(v) = n$, by Proposition 21, we have $\text{rk}^G(v) = n$ and v is $\text{SL}(V)^d$ -semistable, hence v is $\text{SL}(V)$ -semistable. On the other hand, assume v is $\text{SL}(V)$ -semistable. Let λ be a polynomial 1-parameter subgroup of $\text{GL}(V)$ such that $\lim_{t \rightarrow 0} \lambda(t) \cdot v = 0$. Then we can define another 1-parameter subgroup $\lambda'(t) = \lambda(t)^n t^{-e}$, where $\det(\lambda(t)) = t^e$, such that $\det(\lambda') = 1$ and $\lambda' \in \text{SL}(V)$. Since v is $\text{SL}(V)$ -semistable, we have $\text{val}(\lambda'(t) \cdot v) \leq 0$. It follows that

$$\text{val}(\lambda'(t) \cdot v) = \text{val}(\lambda(t)^d t^{-e} \cdot v) = n \text{val}(\lambda(t) \cdot v) - ed \leq 0.$$

The slope function

$$\mu(\lambda(t), v) = \frac{d \text{val}(\det(\lambda(t)))}{\text{val}(\lambda(t) \cdot v)} = \frac{de}{\text{val}(\lambda(t) \cdot v)} \geq n.$$

We get $\text{symmrk}^G(v) = n$. □

Proof of Corollary 7. It suffices to prove that if v is $\text{SL}(V)$ -semistable, then v is $\text{SL}(V)^d$ -semistable. Let us assume v is $\text{SL}(V)$ -semistable, then by Proposition 22 and Theorem 5,

$$\text{rk}^G(v) = \text{symmrk}^G(v) = n.$$

It follows from Proposition 21 that v is $\text{SL}(V)^d$ -semistable. □

5. G-STABLE RANK FOR IDEALS AND LOG CANONICAL THRESHOLD

5.1. G-stable rank for ideals. Let $X = \text{Spec}(R)$ be a nonsingular irreducible complex affine algebraic variety of dimension n , and $\mathfrak{a} \subset R$ be a nonzero ideal, and let $P \in V(\mathfrak{a})$ be a closed point, \mathcal{O}_P be the local ring at P and \mathfrak{m}_P be the maximal ideal corresponding to P .

Definition 23 ([8]). Functions $x_1, \dots, x_n \in \mathcal{O}_P$ are a *system of local parameters* at P if each $x_i \in \mathfrak{m}_P$, and the images of x_1, \dots, x_n form a basis of the vector space $\mathfrak{m}_P/\mathfrak{m}_P^2$.

Let $T = \{x_1, x_2, \dots, x_n\}$ be a system of local parameters at P . Let $\mathbb{C}\{x_1, x_2, \dots, x_n\}$ be the ring of convergent power series in x_1, x_2, \dots, x_n . The ring \mathcal{O}_P is contained in $\mathbb{C}\{x_1, x_2, \dots, x_n\}$. If y_1, y_2, \dots, y_n is any system of local parameters, then $\mathbb{C}\{x_1, x_2, \dots, x_n\} = \mathbb{C}\{y_1, y_2, \dots, y_n\}$. For any $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n) \in \mathbb{Z}_{\geq 0}^n$, we have a natural action of \mathbb{C}^* on $\mathbb{C}\{x_1, x_2, \dots, x_n\}$ by $t \cdot x_i = t^{\lambda_i} x_i$ for any $t \in \mathbb{C}^*$.

Definition 24. Let $T = \{x_1, x_2, \dots, x_n\}$ be a system of local parameters at P and $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n) \in \mathbb{Z}_{\geq 0}^n$ a tuple of non-negative integers. Let $f \in \mathbb{C}\{x_1, x_2, \dots, x_n\}$ be a convergent power series. We define the *valuation* of f with respect to T by

$$(11) \quad \text{val}_\lambda^T(f) = \max\{k \mid f(t^{\lambda_1} x_1, t^{\lambda_2} x_2, \dots, t^{\lambda_n} x_n) = t^k g(x_1, \dots, x_n, t), \text{ for some } g \in \mathbb{C}\{x_1, \dots, x_n, t\}\}.$$

Let $\mathfrak{a} \subset R$ be a nonzero ideal as before, the *order* of \mathfrak{a} with respect to this system of local parameters T and $\lambda \in \mathbb{Z}_{\geq 0}^n$ is defined as

$$(12) \quad \text{ord}_\lambda^T(\mathfrak{a}) = \min\{\text{val}_\lambda^T(f) \mid f \in \mathfrak{a}\}.$$

Remark 25. If \mathfrak{a} is generated by f_1, \dots, f_r , then

$$\text{ord}_\lambda^T(\mathfrak{a}) = \min\{\text{val}_\lambda^T(f_i) \mid i = 1, \dots, r\}.$$

Indeed, it is clear that $\min\{\text{val}_\lambda^T(f) \mid f \in \mathfrak{a}\} \leq \min\{\text{val}_\lambda^T(f_i) \mid i = 1, \dots, r\}$. On the other hand, if $f \in \mathfrak{a}$ computes $\text{ord}_\lambda^T(\mathfrak{a})$, then we can write $f = \sum_i a_i f_i$, for some $a_i \in R$, we have $\text{val}_\lambda^T(f) = \text{val}_\lambda^T(\sum_i a_i f_i) \geq \min\{\text{val}_\lambda^T(a_i f_i) \mid i = 1, \dots, r\} \geq \min\{\text{val}_\lambda^T(f_i) \mid i = 1, \dots, r\}$.

Definition 26. Assume $T = \{x_1, \dots, x_n\}$ is a system of local parameters at P and $\lambda = \{\lambda_1, \dots, \lambda_n\} \in \mathbb{Z}_{\geq 0}^n$, we define the *slope* function $\mu_P(\lambda, \mathfrak{a})$ at P as

$$(13) \quad \mu_P(\lambda, \mathfrak{a}) = \frac{\sum_{i=1}^n \lambda_i}{\text{ord}_{\lambda}^T(\mathfrak{a})}.$$

The *T-stable rank* of \mathfrak{a} at P is the infimum of the slope function $\mu_P(\lambda, \mathfrak{a})$ with respect to the tuple $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n) \in \mathbb{Z}_{\geq 0}^n$,

$$(14) \quad \text{rk}^T(P, \mathfrak{a}) = \inf_{\lambda} \mu_P(\lambda, \mathfrak{a}) = \inf_{\lambda} \frac{\sum_{i=1}^n \lambda_i}{\text{ord}_{\lambda}^T(\mathfrak{a})}.$$

The *G-stable rank* of \mathfrak{a} is defined by taking the infimum of *T-stable rank* with respect to all system of local parameters T at P ,

$$(15) \quad \text{rk}^G(P, \mathfrak{a}) = \inf_T (\text{rk}^T(P, \mathfrak{a})).$$

If $P \notin V(\mathfrak{a})$, we define $\text{rk}^G(P, \mathfrak{a}) = \infty$, we write $\text{rk}^G(\mathfrak{a})$ and $\text{rk}^T(\mathfrak{a})$ if P is known in the context. In the following example, we see that an ideal \mathfrak{a} can have different *T-stable rank* with respect to different system of local parameters T at a point P .

Example 27. Let $R = \mathbb{C}[x, y]$, $T = \{x, y\}$ and assume $\mathfrak{a} = (x^2 + 2xy + y^2)$ is a principle ideal generated by a polynomial $f(x, y) = x^2 + 2xy + y^2$, $P = (0, 0)$ is the origin. Let $\lambda = (\lambda_1, \lambda_2) \in \mathbb{Z}_{\geq 0}^2$, then we have

$$\text{rk}^T(f) = \inf_{\lambda} \frac{\lambda_1 + \lambda_2}{\min(2\lambda_1, \lambda_1 + \lambda_2, 2\lambda_2)} = 1$$

Let us choose a different system of local parameters $T' = \{u = x + y, v = x - y\}$, then $\mathfrak{a} = (u^2)$, and $f(u, v) = u^2$, then

$$\text{rk}^{T'}(f) = \inf_{\lambda} \frac{\lambda_1 + \lambda_2}{2\lambda_1} = \frac{1}{2}$$

In fact, $\text{lct}_P(u^2) = \frac{1}{2}$, and by Theorem 9, we have $\text{lct}_P(\mathfrak{a}) \leq \text{rk}^G(\mathfrak{a})$, therefore we get $\text{rk}^G(f) = \frac{1}{2}$.

Example 28. Let $\mathfrak{a} = (x^2y, y^2z, z^2x) \subset \mathbb{C}[x, y, z]$, $T = \{x, y, x\}$, $P = (0, 0, 0)$, then we get

$$\text{rk}^T(\mathfrak{a}) = \inf_{\lambda} \frac{\lambda_1 + \lambda_2 + \lambda_3}{\min(2\lambda_1 + \lambda_2, 2\lambda_2 + \lambda_3, 2\lambda_3 + \lambda_1)} = 1$$

The ideal $\mathfrak{a} = (x^2y, y^2z, z^2x)$ is a monomial ideal and we will see later that for a monomial ideal \mathfrak{a} , we have $\text{rk}^G(\mathfrak{a}) = \text{lct}_P(\mathfrak{a})$. Using the fact that $\text{lct}_P(\mathfrak{a}) = 1$, we obtain $\text{rk}^G(\mathfrak{a}) = 1$.

Remark 29. We have a short exact sequence

$$(16) \quad 1 \rightarrow K \rightarrow \text{Aut}(\mathbb{C}\{x_1, \dots, x_n\}) \rightarrow \text{GL}(n) \rightarrow 1,$$

where K is a normal subgroup of the group $\text{Aut}(\mathbb{C}\{x_1, \dots, x_n\})$ of local holomorphic automorphisms. The morphism $\text{Aut}(\mathbb{C}\{x_1, \dots, x_n\}) \rightarrow \text{GL}(n)$ is given by computing the Jacobian matrix at $(0, \dots, 0)$. Furthermore, this sequence splits, we have $\text{Aut}(\mathbb{C}\{x_1, \dots, x_n\}) = K \rtimes \text{GL}(n)$.

By Remark 29, there is an action of $\mathrm{GL}(n)$ on the set of system of local parameters. Assume $T = \{x_1, \dots, x_n\}$ is a system of local parameters at P . For $g \in \mathrm{GL}(n)$, $g \cdot T$ is another system of local parameters. We say a system of local parameters $T = \{x_1, \dots, x_n\}$ is *good* for \mathfrak{a} if $\mathrm{rk}^G(P, \mathfrak{a}) = \mathrm{rk}^{g \cdot T}(P, \mathfrak{a})$ for some $g \in \mathrm{GL}(n)$. In other words, to compute the G -stable rank of \mathfrak{a} , it is enough to consider all systems of local parameters obtained from T by actions of $\mathrm{GL}(n)$.

Example 30. Let $f(x, y) = x + y^2 \in \mathbb{C}[x, y]$, $P = (0, 0)$, we take $T = \{x, y\}$. It can be shown that $\mathrm{rk}^T(f) = \frac{3}{2}$. However, if we choose another system of local parameters $T' = \{u = x + y^2, v = y\}$, then $f(u, v) = u$, and $\mathrm{rk}^{T'}(f) = 1$. Indeed, this system of local parameters is optimal, in other words, we can compute the G -stable rank in this system of local parameters, and we have $\mathrm{rk}^G(f) = 1$.

Proposition 31. *If \mathfrak{a} is homogeneous in local parameters $T = \{x_1, \dots, x_n\}$, then T is good for \mathfrak{a} .*

Proof. Since \mathfrak{a} is a homogeneous ideal, we can find a set of generators which are homogeneous polynomials. By Remark 25, it is enough to assume that \mathfrak{a} is generated by a single homogeneous polynomial f . Let $g \in K \subseteq \mathrm{Aut}(\mathbb{C}\{x_1, \dots, x_n\})$, we can write the action of g on T as following

$$g(x_i) = x_i + p_i(x_1, \dots, x_n),$$

where $p_i \in \mathbb{C}\{x_1, \dots, x_n\}$ with no constant and degree 1 terms. Since f is a homogeneous polynomial, we have

$$\mathrm{val}_\lambda^T(f(g(x_1), \dots, g(x_n))) \leq \mathrm{val}_\lambda^T(f(x_1, \dots, x_n)).$$

Let T' be the system of local parameters obtained from T by the action of g , then we have

$$(17) \quad \frac{\sum_{i=1}^n \lambda_i}{\mathrm{ord}_\lambda^{T'}(f)} \geq \frac{\sum_{i=1}^n \lambda_i}{\mathrm{ord}_\lambda^T(f)}.$$

By Lemma 29, given any $h \in \mathrm{Aut}(\mathbb{C}\{x_1, \dots, x_n\})$, we can decompose the action of h into an action of K following by an action of $\mathrm{GL}(n)$. By inequality (17), in the system of local parameters obtained by the action of K from T , we have larger slope than the slope computed in T , hence to compute the G -stable rank of f , it suffices to consider the action of $\mathrm{GL}(n)$. This shows that T is good for \mathfrak{a} . \square

Corollary 32. *If $f \in \mathbb{C}[x_1, \dots, x_n]$ is a homogeneous polynomial of degree $d \geq 2$ and f has isolated singularity at $P = (0, \dots, 0)$. Then $\mathrm{rk}^G(f) = \frac{n}{d}$.*

Proof. By Corollary 31, the system of local parameters $T = \{x_1, \dots, x_n\}$ is good for f , therefore we only need to consider the group action of $\mathrm{GL}(n)$.

We claim that f is $\mathrm{SL}(n)$ -semistable in the sense of Definition 2. Indeed, since f has an isolated singularity at origin, $\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n}$ only have a common zero at origin. By [7], chapter 13, their resultant $\mathrm{Res}(\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n})$ is nonzero and invariant under the action of $\mathrm{SL}(n)$. Now assume there is a one parameter subgroup $\lambda : \mathbb{G}_m \rightarrow \mathrm{SL}(n)$, such that

$$\lim_{t \rightarrow 0} \lambda(t) \cdot v = 0.$$

Then the resultant $\mathrm{Res}(\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n})$ is 0, which is impossible. This proves the claim.

By the claim that f is $\mathrm{SL}(n)$ -semistable, the corollary follows immediately from Proposition 2.6 in [2] \square

5.2. Relation to log canonical threshold. Let $X = \text{Spec}(R)$ be a nonsingular irreducible complex affine variety, $\mathfrak{a} \subset R$ is an ideal, $P \in V(\mathfrak{a})$. If $\mathfrak{a} = (f_1, \dots, f_r) \subseteq R$ is a nonzero ideal, recall the Definition 8, the log canonical threshold $\text{lct}_P(\mathfrak{a})$ of the ideal \mathfrak{a} at point P is

$$(18) \quad \text{lct}_P(\mathfrak{a}) = \sup\{s > 0 \mid \frac{1}{(\sum_{i=0}^r |f_i|^2)^s} \text{ is integrable around } P\}.$$

Theorem 9 says that the log canonical threshold of an ideal is less than or equal to the G -stable rank of that ideal

$$(19) \quad \text{lct}_P(\mathfrak{a}) \leq \text{rk}^G(P, \mathfrak{a}).$$

Proof of Theorem 9. Let $s > 0$ be such that $\frac{1}{(\sum_{i=0}^r |f_i|^2)^s}$ is integrable around P , then there is a neighborhood U_P of P , such that

$$\int_{U_P} \frac{dV}{(\sum_{i=0}^r |f_i|^2)^s} < C < \infty,$$

for some constant C . Choose a system of local parameters $T = \{x_1, \dots, x_n\}$ at P , let $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{Z}_{\geq 0}^n$, then let $t \in \mathbb{C}^*$ act on the coordinates by $x_i \rightarrow t^{\lambda_i} x_i$. We denote $t \cdot U_P$ for the image of U_P under this action. If $|t| < 1$, we have $t \cdot U_P \subset U_P$, therefore

$$(20) \quad \int_{t \cdot U_P} \frac{dV}{(\sum_{i=0}^r |f_i|^2)^s} < \int_{U_P} \frac{dV}{(\sum_{i=0}^r |f_i|^2)^s} < C$$

Let $y_i = t^{-\lambda_i} x_i$ for $i = 1, \dots, n$, then we have

$$(21) \quad \int_{t \cdot U_P} \frac{dx_1 d\bar{x}_1 \cdots dx_n d\bar{x}_n}{(\sum_{i=0}^r |f_i(x_1, \dots, x_n)|^2)^s} = \int_{U_P} \frac{|t|^{2\sum_{i=1}^n \lambda_i} dy_1 d\bar{y}_1 \cdots dy_n d\bar{y}_n}{(\sum_{i=1}^r |f_i(t^{\lambda_1} y_1, \dots, t^{\lambda_n} y_n)|^2)^s} < C$$

Recall the Definition 24 for the valuation (11) and order of an ideal (12), we can write

$$\begin{aligned} \sum_{i=0}^r |f_i(t^{\lambda_1} y_1, \dots, t^{\lambda_n} y_n)|^2 &= |t|^{2\min_i(\text{val}_\lambda^T(f_i))} \left(\sum_{i=1}^r |\tilde{f}_i(y_1, \dots, y_n, t)|^2 \right) \\ &= |t|^{2\text{ord}_\lambda^T(\mathfrak{a})} \left(\sum_{i=1}^r |\tilde{f}_i(y_1, \dots, y_n, t)|^2 \right), \end{aligned}$$

for some $\tilde{f}_i(y_1, \dots, y_n, t) \in \mathbb{C}\{y_1, y_2, \dots, y_n, t\}$. In particular, we know that $\sum_{i=1}^r |\tilde{f}_i(y_1, \dots, y_n, 0)|^2$ is not constantly zero in U_P . So we can find a point $Q \in U_P$ such that $0 < \sum_{i=1}^r |\tilde{f}_i(Q, 0)|^2 < B$ for some constant $B > 0$. By the continuity, there is a neighborhood U_Q such that $Q \in U_Q \subset U_P$ and some $\epsilon > 0$, such that $0 < \sum_{i=1}^r |\tilde{f}_i(y_1, \dots, y_n, t)|^2 < B$ for any $(y_1, \dots, y_n) \in U_Q$ and $0 \leq t < \epsilon$. We can write integral (21) as

$$(22) \quad \int_{U_P} \frac{|t|^{2\sum_{i=1}^n \lambda_i} dy_1 d\bar{y}_1 \cdots dy_n d\bar{y}_n}{|t|^{2\text{ord}_\lambda^T(\mathfrak{a})} (\sum_{i=1}^r |\tilde{f}_i(y_1, \dots, y_n, t)|^2)^s} = \int_{U_P} \frac{|t|^{2(\sum_{i=1}^n \lambda_i - s \text{ord}_\lambda^T(\mathfrak{a}))} dV}{(\sum_{i=1}^r |\tilde{f}_i(y_1, \dots, y_n, t)|^2)^s} < C.$$

Therefore we get

$$|t|^{2(\sum_{i=1}^n \lambda_i - s \text{ord}_\lambda^T(\mathfrak{a}))} \int_{U_Q} \frac{dV}{B^s} = \int_{U_Q} \frac{|t|^{2(\sum_{i=1}^n \lambda_i - s \text{ord}_\lambda^T(\mathfrak{a}))} dV}{B^s} < \int_{U_P} \frac{|t|^{2(\sum_{i=1}^n \lambda_i - s \text{ord}_\lambda^T(\mathfrak{a}))} dV}{(\sum_{i=1}^r |\tilde{f}_i(y_1, \dots, y_n, t)|^2)^s} < C$$

for all $t \in [0, \epsilon)$, where $\epsilon > 0$. Since $\int_{U_Q} \frac{dV}{B^s} > 0$, we must have

$$\sum_{i=1}^n \lambda_i - s \operatorname{ord}_\lambda^T(\mathbf{a}) \geq 0.$$

Hence we get $s \leq \frac{\sum_{i=1}^n \lambda_i}{\operatorname{ord}_\lambda^T(\mathbf{a})}$. This holds for any system of local coordinates and $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{Z}_{\geq 0}^n$, therefore

$$\operatorname{lct}_P(\mathbf{a}) \leq \operatorname{rk}^G(P, \mathbf{a}).$$

This completes the proof of Theorem 9. \square

Example 33. Let $f = x_1^{u_1} + x_2^{u_2} + \dots + x_n^{u_n} \in \mathbb{C}[x_1, \dots, x_n]$, $P = (0, \dots, 0)$ be the origin. It was shown in [6] that $\operatorname{lct}_P(f) = \min(1, \sum_{i=1}^n \frac{1}{u_i})$. However, we will show that $\operatorname{rk}^G(f) = \sum_{i=1}^n \frac{1}{u_i}$. So we get $\operatorname{lct}_P(f) \leq \operatorname{rk}^G(f)$. If $n = 3$, $u_1 = u_2 = u_3 = 2$, then we have $f = x_1^2 + x_2^2 + x_3^2$ and $\operatorname{lct}_P(f) = 1 < \operatorname{rk}^G(f) = \frac{3}{2}$.

Suppose $\mathbf{a} = (m_1, \dots, m_r) \subset \mathbb{C}[x_1, \dots, x_n]$ is a proper nonzero ideal generated by monomials $\{m_1, \dots, m_r\}$ and let $P = (0, \dots, 0)$ be the origin. Given $u = (u_1, \dots, u_n) \in \mathbb{Z}_{\geq 0}^n$, we write $x^u = x_1^{u_1} \dots x_n^{u_n}$. The *Newton Polyhedron* of \mathbf{a} is

$$(23) \quad P(\mathbf{a}) = \operatorname{convex hull}(\{u \in \mathbb{Z}_{\geq 0}^n \mid x^u \in \mathbf{a}\}).$$

It was shown in [6] that

$$(24) \quad \operatorname{lct}_P(\mathbf{a}) = \max\{\nu \in \mathbb{R}_{\geq 0} \mid (1, 1, \dots, 1) \in \nu \cdot P(\mathbf{a})\}.$$

In other words, $\operatorname{lct}_P(\mathbf{a})$ is equal to the largest ν such that $\sum_{i=1}^n \lambda_i \geq \nu \cdot \min_{u \in P(\mathbf{a})} \langle u, \lambda \rangle$ for any $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{Z}_{\geq 0}^n$, where we use the standard inner product $\langle u, \lambda \rangle = \sum_{i=1}^n u_i \lambda_i$.

Theorem 10 says that the log canonical threshold is equal to the G -stable rank for monomial ideals. More precisely, suppose $\mathbf{a} \subset \mathbb{C}[x_1, \dots, x_n]$ is a proper nonzero ideal generated by monomials and $P = (0, \dots, 0)$ is the origin. Then

$$(25) \quad \operatorname{lct}_P(\mathbf{a}) = \operatorname{rk}^G(\mathbf{a}).$$

Proof of Theorem 10. Let $\mathbf{a} = (m_1, \dots, m_r)$ and $\{m_i = x_1^{l_{i1}} x_2^{l_{i2}} \dots x_n^{l_{in}}\}_{i=1, \dots, r}$ be a set of generators, where $l_i = (l_{i1}, \dots, l_{in}) \in \mathbb{Z}_{\geq 0}^n$, we have

$$\begin{aligned} \operatorname{lct}_P(\mathbf{a}) &= \max\{\nu \in \mathbb{R}_{\geq 0} \mid (1, \dots, 1) \in \nu \cdot P(\mathbf{a})\} \\ &= \max\{\nu \in \mathbb{R}_{\geq 0} \mid \sum_j \lambda_j \geq \nu \cdot \min_{u \in P(\mathbf{a})} \langle u, \lambda \rangle, \forall \lambda \in \mathbb{Z}_{\geq 0}^n\} \\ &= \max\{\nu \in \mathbb{R}_{\geq 0} \mid \sum_j \lambda_j \geq \nu \cdot \min_i \langle l_i, \lambda \rangle, \forall \lambda \in \mathbb{Z}_{\geq 0}^n\} \\ &= \max\{\nu \in \mathbb{R}_{\geq 0} \mid \nu \leq \frac{\sum_j \lambda_j}{\min_i (\sum_j l_{ij} \lambda_j)}, \forall \lambda \in \mathbb{Z}_{\geq 0}^n\}. \end{aligned}$$

In this system of local parameters $T = \{x_1, \dots, x_n\}$, we have $\operatorname{ord}_\lambda^T(\mathbf{a}) = \min_i (\sum_j l_{ij} \lambda_j)$, therefore we get

$$\begin{aligned} \operatorname{lct}_P(\mathbf{a}) &= \max\{\nu \in \mathbb{R}_{\geq 0} \mid \nu \leq \frac{\sum_i \lambda_i}{\operatorname{ord}_\lambda^T(\mathbf{a})}, \forall \lambda \in \mathbb{Z}_{\geq 0}^n\} \\ &= \max\{\nu \in \mathbb{R}_{\geq 0} \mid \nu \leq \operatorname{rk}^T(\mathbf{a})\} = \operatorname{rk}^T(\mathbf{a}). \end{aligned}$$

Since log canonical threshold does not depend on the local coordinates, hence we have

$$\text{lct}_P(\mathfrak{a}) = \text{rk}^G(\mathfrak{a}).$$

This completes the proof of Theorem 10. \square

Example 34. Suppose $\mathfrak{a} = (x_1^{u_1}, \dots, x_n^{u_n})$ and $P = (0, \dots, 0)$, then $\text{lct}_P(\mathfrak{a}) = \sum_{i=1}^n \frac{1}{u_i}$. Therefore we also have $\text{rk}^G(\mathfrak{a}) = \sum_{i=1}^n \frac{1}{u_i}$.

5.3. Some properties of G -stable rank for ideals. Some results for log canonical threshold can be found in [6]. Here, we also prove similar results for the G -stable rank of ideals.

Proposition 35. *If $\mathfrak{a} \subseteq \mathfrak{b}$ are nonzero ideals on X , then we have $\text{lct}_P(\mathfrak{a}) \leq \text{lct}_P(\mathfrak{b})$ and $\text{rk}^G(P, \mathfrak{a}) \leq \text{rk}^G(P, \mathfrak{b})$.*

Proof. The first inequality was shown in [6]. If $P \notin V(\mathfrak{b})$, it is trivial. Assume $P \in V(\mathfrak{b})$, let T be a system of local parameters at P and $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{Z}_{\geq 0}^n$. Since $\mathfrak{a} \subseteq \mathfrak{b}$, we have $\text{ord}_\lambda^T(\mathfrak{b}) \leq \text{ord}_\lambda^T(\mathfrak{a})$, therefore $\mu_P(\lambda, \mathfrak{a}) \leq \mu_P(\lambda, \mathfrak{b})$, it follows immediately that $\text{rk}^G(P, \mathfrak{a}) \leq \text{rk}^G(P, \mathfrak{b})$. \square

Proposition 36. *We have $\text{lct}_P(\mathfrak{a}^r) = \frac{\text{lct}_P(\mathfrak{a})}{r}$ and $\text{rk}^G(P, \mathfrak{a}^r) = \frac{\text{rk}^G(P, \mathfrak{a})}{r}$ for every $r \geq 1$.*

Proof. The first claim was shown in [6] and the second claim follows from the fact that $\text{ord}_\lambda^T(\mathfrak{a}^r) = r \cdot \text{ord}_\lambda^T(\mathfrak{a})$ and Definition 26. \square

Proposition 37. *If \mathfrak{a} and \mathfrak{b} are ideals of X , then*

$$\frac{1}{\text{lct}_P(\mathfrak{a} \cdot \mathfrak{b})} \leq \frac{1}{\text{lct}_P(\mathfrak{a})} + \frac{1}{\text{lct}_P(\mathfrak{b})}, \quad \frac{1}{\text{rk}^G(P, \mathfrak{a} \cdot \mathfrak{b})} \leq \frac{1}{\text{rk}^G(P, \mathfrak{a})} + \frac{1}{\text{rk}^G(P, \mathfrak{b})}.$$

Proof. The first inequality was shown in [6], we show the second inequality. Let T be a system of local parameters at P and $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{Z}_{\geq 0}^n$, then

$$\text{ord}_\lambda^T(\mathfrak{a} \cdot \mathfrak{b}) = \text{ord}_\lambda^T(\mathfrak{a}) + \text{ord}_\lambda^T(\mathfrak{b}).$$

Therefore, we have

$$\begin{aligned} \sup_{T, \lambda} \frac{\text{ord}_\lambda^T(\mathfrak{a} \cdot \mathfrak{b})}{\sum \lambda_i} &= \sup_{T, \lambda} \left(\frac{\text{ord}_\lambda^T(\mathfrak{a})}{\sum \lambda_i} + \frac{\text{ord}_\lambda^T(\mathfrak{b})}{\sum \lambda_i} \right) \\ &\leq \sup_{T, \lambda} \frac{\text{ord}_\lambda^T(\mathfrak{a})}{\sum \lambda_i} + \sup_{T, \lambda} \frac{\text{ord}_\lambda^T(\mathfrak{b})}{\sum \lambda_i}. \end{aligned}$$

\square

The following two propositions are from [6]. A lot of evidence suggests the same results for G -stable rank of ideals, we give them as conjectures.

Proposition 38. *If $H \subset X$ is a nonsingular hypersurface such that $\mathfrak{a} \cdot \mathcal{O}_H$ is nonzero, then $\text{lct}_P(\mathfrak{a} \cdot \mathcal{O}_H) \leq \text{lct}_P(\mathfrak{a})$.*

Proposition 39. *If \mathfrak{a} and \mathfrak{b} are ideals on X , then*

$$\text{lct}_P(\mathfrak{a} + \mathfrak{b}) \leq \text{lct}_P(\mathfrak{a}) + \text{lct}_P(\mathfrak{b})$$

for every $P \in X$.

Conjecture 40. *If $H \subset X$ is a nonsingular hypersurface such that $\mathfrak{a} \cdot \mathcal{O}_H$ is nonzero, then $\text{rk}^G(P, \mathfrak{a} \cdot \mathcal{O}_H) \leq \text{rk}^G(P, \mathfrak{a})$ for any $P \in H$.*

Conjecture 41. *Let \mathfrak{a} and \mathfrak{b} be two nonzero proper ideals of X , then for any point $P \in X$, we have*

$$\mathrm{rk}^G(P, \mathfrak{a} + \mathfrak{b}) \leq \mathrm{rk}^G(P, \mathfrak{a}) + \mathrm{rk}^G(P, \mathfrak{b}).$$

6. ACKNOWLEDGEMENTS

I would like to thank Harm Derksen for direction and discussion, and Visu Makam for comments on an earlier draft of this paper.

REFERENCES

- [1] Pierre Comon, Gene Golub, Lek-Heng Lim, Bernard Mourrain, *Symmetric tensors and symmetric tensor rank*, SIAM J. Matrix Anal. Appl. 2008, 30, 1254–1279.
- [2] Harm Derksen, *The G -stable rank for tensors*, arXiv:2002.08435.
- [3] Jordan S. Ellenberg, Dion Gijswijt, *On large subsets of \mathbb{F}_q^n with no three-term arithmetic progression*, Ann. of Math. (2) 185 (2017), no. 1, 339–343.
- [4] Zhi Jiang, *Improved explicit upper bounds for the Cap Set Problem*, arXiv:2103.06481.
- [5] George R. Kempf, *Instability in invariant theory*, Ann. of Math. (2) **108** (1978), no. 2, 299–316.
- [6] Mircea Mustața, *Impanga Lectures Notes on Log Canonical Thresholds, Notes by Tomasz Szemberg* in Contributions to Algebraic Geometry, Editor: Piotr Pragacz, EMS Series of Congress Reports.
- [7] Israel M. Gelfand, Mikhail M. Kapranov, Andrei V. Zelevinsky, *Discriminants, Resultants, and Multidimensional Determinants*, Boston, Birkhauser, 1994.
- [8] Igor R. Shafarevich, *Basic Algebraic Geometry 1*, Springer-Verlag, Berlin, Heidelberg, 2013.
- [9] Yaroslav Shitov, *A counterexample to Comon’s conjecture*, SIAM Journal on Applied Algebra and Geometry, Vol.2, Iss.3.
- [10] Terence Tao, *A symmetric formulation of the Croot–Lev–Pach–Ellenberg–Gijswijt capset bound (2016)*, available at <https://terrytao.wordpress.com/2016/05/18/>. blog post.