

MODULAR SHEAVES ON HYPERKÄHLER FOURFOLDS OF KUMMER TYPE

ABSTRACT. We partially extend to hyperkähler fourfolds of Kummer type the results on hyperkähler (HK) varieties of type $K3^{[2]}$ proved in [O’G22]. We show that if (M, h) is a general polarized HK fourfold of Kummer type such that $q_M(h) \equiv -6 \pmod{16}$ and $\text{div}(h) = 2$, then there exists a slope stable rank 4 vector bundle \mathcal{F} on M such that $c_1(\mathcal{F}) = h$ and $\Delta(\mathcal{F}) = c_2(M)$. Moreover if \mathcal{F} is such a vector bundle and φ is an automorphism of M acting trivially on $H^2(M)$, then $\varphi^*(\mathcal{F})$ is isomorphic to \mathcal{F} . Lastly, the general such vector bundle is rigid. (There is a similar result for polarizations h such that $q_M(h) \equiv -6 \pmod{144}$ and $\text{div}(h) = 6$.) We are motivated by the desire to describe explicitly a locally complete family of polarized HK fourfolds of Kummer type.

1. INTRODUCTION

1.1. **Motivation.** Several locally complete families of polarized $K3$ surfaces (S, L) have been explicitly described. The classical families correspond to very low degrees $c_1(L)^2 \in \{2, 4, 6, 8\}$. If (S, L) is a general polarized surface of degree in $\{4, 6, 8\}$ then the image of the map $S \rightarrow |L|^\vee$ is isomorphic to S and is a complete intersection, and hence readily described (if $c_1(L)^2 = 2$, the map $S \rightarrow |L|^\vee$ is the double cover branched over a sextic curve). There are also the non classical families with degrees $c_1(L)^2 \in \{10, \dots, 18\}$ described by Mukai, see [Muk88]. If (S, L) is such a general polarized surface, then one does not describe explicitly its ideal sheaf in $|L|^\vee$ (S is not a complete intersection). Instead Mukai describes explicitly S as the zero-locus of a (homogeneous) vector bundle on a Grassmannian (or an orthogonal Grassmannian, or similar homogeneous varieties). Such a description is possible because over S there is a stable spherical vector bundle of determinant isomorphic to L (given by the restriction of the tautological quotient vector bundle on the Grassmannian).

What happens if we consider higher dimensional analogues of $K3$ surfaces, i.e. HK manifolds of dimension greater than 2? Some locally complete families of polarized HK varieties of dimension greater than 2 have been explicitly described, for example the varieties of lines on a smooth cubic hypersurface in \mathbb{P}^5 [BD85], double EPW sextics [O’G06], Debarre-Voisin varieties [DV10], the LLSvS varieties attached to smooth cubic hypersurface in \mathbb{P}^5 [LLSvS17], the infinite family of moduli spaces of stable objects in the Kuznetsov category attached to smooth cubic hypersurface in \mathbb{P}^5 [BLM⁺21]. All of the locally complete polarized families of HK’s which have been explicitly described parametrize varieties of type $K3^{[n]}$. The present work is motivated by the desire to describe explicitly a locally complete family of polarized HK fourfolds of Kummer type by mimicking Mukai. In order to explain our strategy, we note that the explicit families given by the varieties of lines on a smooth cubic hypersurface in \mathbb{P}^5 and the Debarre-Voisin varieties are very similar to the Mukai families. In fact the restrictions of the tautological quotient vector bundles are stable (for a general variety of that kind) rank 4 vector bundles which extend to all deformations of the polarized (X, L) , and actually the

Date: 9th March 2022.

Partially supported by PRIN 2017.

projectivization of those vector bundles extends to all small deformations of the unpolarized X (this last property differentiates the tautological quotient from the tautological subbundle). In a nutshell our plan is to mimic such a construction for polarized HK fourfolds of Kummer type. In order to explain in broad terms what we do here, we recall that a torsion free sheaf \mathcal{F} on a HK manifold M of dimension $2n$ is modular if there exists $d(\mathcal{F}) \in \mathbb{Q}$ such that

$$\int_M (2rc_2(\mathcal{F}) - (r-1)c_1(\mathcal{F})^2) \smile \alpha^{2n-2} = d(\mathcal{F}) \cdot (2n-3)!! \cdot q_M(\alpha, \alpha)^{n-1} \quad (1.1.1)$$

for all $\alpha \in H^2(M)$, where q_M is the Beauville-Bogomolov-Fujiki (BBF for short) quadratic form of M . The rank 4 tautological quotient vector bundles on the varieties of lines on a smooth cubic hypersurface in \mathbb{P}^5 and the Debarre-Voisin varieties are examples of modular vector bundles. Variation of stability for modular sheaves is independent of the dimension, in other words even if $n > 1$ it behaves as if the base were a surface, see [O'G22]. In [O'G22] we took advantage of this fact to prove results on stable modular sheaves on polarized HK varieties of type $K3^{[2]}$.

In the present paper we partially extend those results to HK fourfolds of Kummer type.

1.2. Main result. Before stating our main result we recall a few results on hyperkähler manifolds of Kummer type. First, if M is such a manifold of dimension $2n \geq 4$, then the subgroup $\text{Aut}^0(M) < \text{Aut}(M)$ of automorphisms acting trivially on $H^2(M)$ is isomorphic to a semidirect product of $\mathbb{Z}/(n+1)^4$ and $\mathbb{Z}/(2)$.

Next we discuss 4 dimensional polarized HK's of Kummer type. Let (M, h) be such a variety (the ample class $h \in H_{\mathbb{Z}}^{1,1}(M)$ is primitive). Let $e := q_M(h)$, and let $q_M(h, H^2(X; \mathbb{Z})) = (i)$, i.e. e is the *square of h* and i is the *divisibility of h* , denoted $\text{div}(h)$. Then one of the following holds:

$$i = 1, \quad e \equiv 0 \pmod{2}, \quad (1.2.1)$$

$$i = 2, \quad e \equiv -6 \pmod{8}, \quad (1.2.2)$$

$$i = 3, \quad e \equiv -6 \pmod{18}, \quad (1.2.3)$$

$$i = 6, \quad e \equiv -6 \pmod{72}. \quad (1.2.4)$$

Conversely, if e is a positive integer, $i \in \{1, 2, 3, 6\}$ and one of (1.2.1), (1.2.2) (1.2.3) or (1.2.4) holds, then there exists a 4 dimensional polarized HK (M, h) of Kummer type such that $q_M(h) = e$ and $\text{div}(h) = i$. Let Kum_e^i be the corresponding moduli stack of 4 dimensional polarized HK's (M, h) of Kummer type (note that Kum_e^i is 4 dimensional). The number of irreducible components of Kum_e^i has been computed by Onorati in [Ono16]. In particular Kum_e^2 is irreducible (we assume that (1.2.2) holds), and Kum_e^6 has two irreducible components (here we assume that (1.2.4) holds). Let $\text{M}_e^i \rightarrow \text{Kum}_e^i$ be the tautological family of HK varieties, and let $\text{VB}_e^i \rightarrow \text{Kum}_e^i$ be the relative moduli stack of slope semistable (of course stability is with respect to the polarization) vector bundles \mathcal{F} on fibers M of $\text{M}_e^i \rightarrow \text{Kum}_e^i$ such that

$$r(\mathcal{F}) = 4, \quad c_1(\mathcal{F}) = h, \quad \Delta(\mathcal{F}) := 8c_2(\mathcal{F}) - 3c_1(\mathcal{F})^2 = c_2(M). \quad (1.2.5)$$

(Chern classes are in cohomology. Recall that $\Delta(\mathcal{F})$ is the discriminant of \mathcal{F} .)

Theorem 1.1. *Let e be a positive integer such that $e \equiv -6 \pmod{16}$. Then there exists at least one irreducible component of VB_e^2 , say $\text{VB}_e^2(0)$, such that the map*

$$\text{VB}_e^2(0) \rightarrow \text{Kum}_e^2$$

is dominant of finite degree. Similarly, let e be a positive integer such that $e \equiv -6 \pmod{144}$. Then there exists at least one irreducible component of VB_e^6 , say $\mathrm{VB}_e^6(0)$, such that the map

$$\mathrm{VB}_e^6(0) \rightarrow \mathrm{Kum}_e^6$$

is dominant of finite degree over one of the two irreducible components of Kum_e^6 . If \mathcal{F} is a vector bundle parametrized by $\mathrm{VB}_e^6(0)$ or $\mathrm{VB}_e^6(0)$ and $\varphi \in \mathrm{Aut}^0(M)$, then $\varphi^*(\mathcal{F})$ is isomorphic to \mathcal{F} . Lastly, if \mathcal{F} is a general such vector bundle then $H^1(M, \mathrm{End}^0(\mathcal{F})) = 0$.

1.3. Outline of the paper. In Section 2 we collect a few preliminary results on modular sheaves and HK manifolds of Kummer type.

In Section 3 we consider a particular case of the following construction. Let $f: B \rightarrow A$ be an isogeny of abelian surfaces. Let $\rho: K_n(B) \dashrightarrow K_n(A)$ be the map between generalized Kummars which associates to $[Z] \in K_n(B)$ the point $[f(Z)] \in K_n(A)$, and let $\tilde{\rho}: X \rightarrow K_n(A)$ be a resolution of indeterminacies of ρ . (As soon as $\deg f \geq 2$ and $n \geq 2$ the map ρ is not regular.) If \mathcal{L} is a line bundle on X , let $\mathcal{E}(\mathcal{L}) := \tilde{\rho}_*(\mathcal{L})$. Thus $\mathcal{E}(\mathcal{L})$ is a torsion free sheaf on $K_n(A)$, of rank $(\deg f)^n$. Section 3 is devoted to the study of $\mathcal{E}(\mathcal{L})$ for $n = 2$ and $\deg f = 2$. We determine exactly under which hypotheses the rank 4 sheaf $\mathcal{E}(\mathcal{L})$ is modular. In particular we show that if \mathcal{L} is the pull-back of a line bundle on $K_2(B)$ then $\mathcal{E}(\mathcal{L})$ is modular, it is locally free and $\Delta(\mathcal{E}(\mathcal{L})) = c_2(K_2(A))$. These are the vector bundles which are studied in the remaining part of the paper.

In Section 4 we prove that, under additional assumptions, the vector bundle $\mathcal{E}(\mathcal{L})$ is slope stable with respect to suitable polarizations and moreover it deforms if we deform $K_2(A)$ maintaining $\det \mathcal{E}(\mathcal{L})$ of type $(1, 1)$. For the proof of the stability result the modularity of $\mathcal{E}(\mathcal{L})$ is a crucial ingredient. In fact we prove stability for A isogenous to a product, so that there is a Lagrangian fibration $\pi_A: K_2(A) \rightarrow \mathbb{P}^2$. Since $\mathcal{E}(\mathcal{L})$ is modular, it is slope stable with respect to polarizations very close to $\pi_A^*(\mathcal{O}_{\mathbb{P}^2}(1))$ if it restricts to a slope stable vector bundle on a general Lagrangian fiber. This holds (under additional assumptions on $c_1(\mathcal{L})$) for $\mathcal{E}(\mathcal{L})$ precisely because of the definition as push-forward of a line bundle, actually this fact prompted us to study the sheaves $\mathcal{E}(\mathcal{L})$. Once we know that $\mathcal{E}(\mathcal{L})$ is slope stable with respect to some polarization, it follows from Verbitsky's results (see [Ver96b]) that $\mathcal{E}(\mathcal{L})$ is projectively hyperholomorphic, and hence it deforms along deformations of $K_2(A)$ maintaining $\det \mathcal{E}(\mathcal{L})$ of type $(1, 1)$.

In Section 5 we prove that, under assumptions slightly stronger than those of Section 4, the vector bundle $\mathcal{E}(\mathcal{L})$ is rigid, i.e. $H^1(K_2(A), \mathrm{End}^0(\mathcal{E}(\mathcal{L}))) = 0$. This allows us to show that if M is a small deformation of $K_2(A)$ maintaining $\det \mathcal{E}(\mathcal{L})$ of type $(1, 1)$, then the corresponding deformation of $\mathcal{E}(\mathcal{L})$ is left invariant by the action of $\mathrm{Aut}^0(M)$. In fact by rigidity it suffices to show that $\mathrm{Aut}^0(K_2(A))$ acts trivially on $\mathcal{E}(\mathcal{L})$, and this follows from the definition of $\mathcal{E}(\mathcal{L})$.

The last section, namely Section 6, wraps everything up to give the proof of Theorem 1.1. The idea is to show that there exists a vector bundle $\mathcal{E}(\mathcal{L})$ such that $\det \mathcal{E}(\mathcal{L})$ is a primitive ample line bundle of the given square e and divisibility i . One finishes the proof by dealing with slope stability with respect to $\det \mathcal{E}(\mathcal{L})$. We do not prove that this holds, but rather that it holds for certain “small” deformations of the couple $(\mathcal{E}(\mathcal{L}), \det \mathcal{E}(\mathcal{L}))$.

1.4. Comments. Following are a few questions that arise naturally regarding the results of the present paper.

Let r_0 be a positive number coprime to 3. Our method for constructing stable modular vector bundles of rank 4 should provide moduli stacks of stable torsion free sheaves of rank r_0^2 dominating components of Kum_e^i for suitable e, i . In fact,

mimicing our approach for rank 4, one would start from an isogeny $f: B \rightarrow A$ of abelian surfaces of degree r_0 , then one would consider the induced rational map $\rho: K_2(B) \dashrightarrow K_2(A)$. The main problem is to resolve the indeterminacies of ρ and reproduce the computations of Section 3. Of course, there might be an approach which does not rely on explicit computations, see Markman's recent preprint [Mar21].

In [O'G22] we constructed a stable rank r_0^2 vector bundle \mathcal{F} on the general polarized HK variety (M, h) of type $K3^{[2]}$ (under certain numerical assumptions on the polarization) with $c_1(\mathcal{F}) = r_0/i$, where i is the divisibility of h , and $\Delta(\mathcal{F}) = r(\mathcal{F})(r(\mathcal{F}) - 1)c_2(M)/12$. We proved also that, up to isomorphism, there is a unique stable vector bundle with the given rank and Chern classes. Note that the vector bundles appearing in Theorem 1.1 for $e \equiv -6 \pmod{16}$ and $i = 2$ are numerically analogous to the vector bundles on polarized HK varieties of type $K3^{[2]}$ that one gets setting $r_0 = 2$ and $i = 2$. We expect that unicity up to isomorphism holds also for the vector bundles that we consider in the present paper. The proof of unicity given in [O'G22] consists in a series of steps, which can be carried out also in the case of our vector bundles on fourfolds of Kummer type, except possibly for the following one: we need that on a general deformation of our vector bundles on a Lagrangian fourfold, the restriction to the Lagrangian fibers is slope stable in codimension 1. We know that the restriction to smooth Lagrangian fibers is slope stable (under certain numerical hypotheses), and that the restriction to a general singular fiber is a simple vector bundle. It looks as if we are almost there, but something is still missing because the Lagrangians that we analyzed explicitly are generalized Kummers of abelian surfaces isogenous to products, so that the singular fibers are reducible and non reduced, and the restriction to a general such singular fiber is definitely not slope stable.

Let \mathcal{F} be a general vector bundle parametrized by $\text{VB}_e^2(0)$ or by $\text{VB}_e^6(0)$ (notation as in Theorem 1.1); we expect that $\dim \text{Ext}^p(\text{End}^0 \mathcal{F}) = 0$ for all p , as in the analogous case of modular stable vector bundles on HK varieties of type $K3^{[2]}$ considered in [O'G22]. What is missing in the present case is the vanishing of $\text{Ext}^2(\text{End}^0 \mathcal{F})$. In particular we expect that \mathcal{F} is a \mathbb{P}^2 object in $D^b(M)$, and hence that it is an atomic sheaf according to Beckmann's definition, see [Bec22].

1.5. Conventions.

- Algebraic variety is synonymous of complex quasi projective scheme, and sheaf is synonymous of coherent sheaf on an algebraic variety. Occasionally we do not differentiate between divisor classes, line bundles and invertible sheaves on a smooth variety.
- Chern classes of a sheaf on a smooth complex quasi projective variety X are elements of the Betti cohomology ring $H^*(X(\mathbb{C}); \mathbb{Z})$ unless we state the contrary.
- Let \mathcal{F} be a sheaf of positive rank on a smooth projective variety X . If L is an ample line bundle then $\mu_L(\mathcal{F})$ is the slope of \mathcal{F} with respect to L , and similarly $\mu_h(\mathcal{F})$ is the slope of \mathcal{F} with respect to the numerical equivalence class of an ample divisor on X . Hopefully there will be no confusion with Donaldson's map μ discussed in Subsection 2.2.
- A *polarized variety* is a couple (X, h) where X is a projective scheme and h is the numerical equivalence class of an ample divisor on X . If X is smooth of positive rank, slope (semi)stability of \mathcal{F} refers to the slope function defined by h .
- "Abelian variety" often means a variety isomorphic to a bona fide abelian variety A , in other words a torsor over a bona fide abelian variety.

2. PRELIMINARIES

2.1. Modular sheaves. We start by presenting an equivalent characterization of modular sheaves on a HK manifold M . Let $V \subset H(M)$ be the image of the map $\text{Sym } H^2(M) \rightarrow H(M)$ defined by cup product, and let $U := V^\perp \subset H(M)$ be the orthogonal (with respect to the intersection form) of V . We let $V^d := V \cap H^d(M)$ and $U^d := U \cap H^d(M)$. By Verbitsky [Ver96a] we have a direct sum decomposition $H(M) = V \oplus U$, in particular

$$H^4(M) = V^4 \oplus U^4.$$

The projection of $c_2(M)$ onto V^4 is non zero (because the tangent bundle of M is stable by Yau's Theorem but not flat), let us denote it (following Markman) by $\bar{c}_2(M)$. Then \mathcal{F} is modular if and only if the projection of $\Delta(\mathcal{F})$ onto $\text{Sym}^2 H^2(M)$ is a multiple of $\bar{c}_2(M)$.

2.2. Generalized Kummers. Let A be an abelian surface. The generalized Kummer $K_n(A)$ is the fiber over 0 of the map $A^{[n+1]} \rightarrow A$ given by the composition

$$A^{[n+1]} \xrightarrow{\mathfrak{h}} A^{(n+1)} \xrightarrow{\sigma} A$$

where $\mathfrak{h}[Z] := \sum_{a \in A} \ell(\mathcal{O}_{Z,a})(a)$ and $\sigma((a_1) + \dots + (a_{n+1})) := a_1 + \dots + a_{n+1}$ is the summation map in the group A . Here and in the rest of the paper we denote by (a) the generator of the group of 0 cycles on A that corresponds to the point $a \in A$. Hence if k_1, \dots, k_{n+1} are integers and $a_1, \dots, a_{n+1} \in A$ then $k_1(a_1) + \dots + k_{n+1}(a_{n+1})$ is a 0 cycle while $k_1 a_1 + \dots + k_{n+1} a_{n+1}$ is an element of A .

The cohomology group $H^2(K_n(A); \mathbb{Z})$ is described as follows. There is a homomorphism $\mu_{n,A}: H^2(A) \rightarrow H^2(K_n(A))$ given by the composition

$$H^2(A) \xrightarrow{s_{n+1}} H^2(A^{(n+1)}) \xrightarrow{\mathfrak{h}^*} H^2(K_n(A))$$

where s_{n+1} is the natural symmetrization map. The map $\mu_{n,A}$ is injective but not surjective because \mathfrak{h} contracts the prime divisor (here we assume that $n \geq 2$)

$$\Delta_n(A) := \{[Z] \in A^{[n+1]} \mid Z \text{ is not reduced}\}.$$

The cohomology class of $\Delta_n(A)$ is (uniquely) divisible by 2 in integral cohomology. We let $\delta_n(A) \in H^2(A^{(n+1)}; \mathbb{Z})$ be the class such that

$$2\delta_n(A) = \text{cl}(\Delta_n(A)). \quad (2.2.1)$$

(Beware of the potential misunderstanding: $\delta_n(A)$ is *not* the class of $\Delta_n(A)$.) One has

$$H^2(K_n(A); \mathbb{Z}) = \mu_{n,A}(H^2(A; \mathbb{Z})) \oplus \mathbb{Z}\delta_n(A), \quad (2.2.2)$$

where orthogonality is with respect to the BBF quadratic form. Moreover the BBF quadratic form is given by

$$q(\mu_{n,A}(\alpha) + x\delta_n(A)) = (\alpha, \alpha)_A - 2(n+1)x^2. \quad (2.2.3)$$

(Here $(\alpha, \alpha)_A$ is the self intersection of $\alpha \in H^2(A)$.)

Let $\beta_1, \dots, \beta_{2n} \in H^2(K_n(A))$. Then

$$\int_{K_n(A)} \beta_1 \smile \dots \smile \beta_{2n} = (n+1) \cdot \widetilde{\sum} q(\beta_{i_1}, \alpha_{i_2}) \cdot \dots \cdot q(\beta_{i_{2n-1}}, \beta_{i_{2n}}), \quad (2.2.4)$$

where $\widetilde{\sum}$ means that in the summation we avoid repeating addends which are formally equal (i.e. are equal modulo reordering of the factors $q_X(\cdot, \cdot)$'s and switching

the entries in $q_X(\cdot, \cdot)$). We also recall that if \mathcal{L} is a line bundle on a generalized Kummer X of dimension $2n$ then

$$\chi(X; \mathcal{L}) = (n+1) \binom{\frac{1}{2}q_X(c_1(\mathcal{L})) + n}{n}. \quad (2.2.5)$$

From now on we deal only with 4 dimensional generalized Kummers $K_2(A)$. We replace $\mu_{2,A}, \Delta_2(A), \delta_2(A)$ by $\mu_A, \Delta(A), \delta(A)$ respectively.

Lastly let M be a HK fourfold of Kummer type, i.e. a deformation of $K_2(A)$; if $\zeta \in H^2(M)$ then (see for example Equation (6) in [HT13])

$$\int_M c_2(M) \smile \zeta^2 = 54q_M(\zeta). \quad (2.2.6)$$

3. MODULAR BASIC SHEAVES ON 4 DIMENSIONAL GENERALIZED KUMMERS

3.1. Statement of the main result. Let $f: B \rightarrow A$ be a homomorphism of (bona fide) abelian surfaces of degree 2. Then f defines a rational map

$$\begin{array}{ccc} K_2(B) & \xrightarrow{\rho} & K_2(A) \\ [Z] & \mapsto & [f(Z)] \end{array} \quad (3.1.1)$$

The map ρ is regular away from

$$V(f) := \{[Z] \in K_2(B) \mid \ell(f(Z)) < \ell(Z) = 2\}. \quad (3.1.2)$$

The result below is proved in Subsection 3.3.

Proposition 3.1. *Keep notation and hypotheses as above. Then $V(f)$ (see (3.1.2)) is irreducible of codimension 2 and smooth. Moreover by blowing up $V(f)$ one resolves the indeterminacies of ρ .*

Let $\nu: X \rightarrow K_2(B)$ be the blow up of $V(f)$, and let $\tilde{\rho}: X \rightarrow K_2(A)$ be the regular map lifting the rational map ρ . Thus we have the commutative diagram

$$\begin{array}{ccc} & X & \\ \nu \swarrow & & \searrow \tilde{\rho} \\ K_2(B) & \overset{\rho}{\dashrightarrow} & K_2(A) \end{array} \quad (3.1.3)$$

For a line bundle \mathcal{L} on X , we let $\mathcal{E}(\mathcal{L}) := \tilde{\rho}_*(\mathcal{L})$ and, if there is no ambiguity regarding \mathcal{L} , we denote it by \mathcal{E} . Since the map $\tilde{\rho}$ has degree 4 and \mathcal{L} is torsion free, $\mathcal{E}(\mathcal{L})$ is a rank 4 torsion free sheaf on $K_2(A)$. We prove that for suitable choices of \mathcal{L} the sheaf $\mathcal{E}(\mathcal{L})$ is modular. In order to state our result we introduce some notation. Let $D \subset X$ be the exceptional divisor of the blow up map ν (notice that D is irreducible by Proposition 3.1). There exist $\omega_B \in H_{\mathbb{Z}}^{1,1}(B)$ and integers x, y such that

$$c_1(\mathcal{L}) = \nu^*(\mu_B(\omega_B) + x\delta(B)) + y \text{cl}(D). \quad (3.1.4)$$

Below is the main result of the present section.

Theorem 3.2. *The sheaf $\mathcal{E}(\mathcal{L})$ is modular if and only if $y = x$ or $y = x + 1$. If this is the case then*

$$\Delta(\mathcal{E}(\mathcal{L})) = c_2(K_2(A)). \quad (3.1.5)$$

Moreover if $y = x$ then $\mathcal{E}(\mathcal{L})$ is locally free.

Theorem 3.2 is proved in Subsections 3.7, 3.9 and 3.10.

Remark 3.3. Let ξ be the square root of the line bundle $\mathcal{O}_{K_2(A)}(\Delta)$. Thus $c_1(\xi) = \delta(A)$. For $t \in \mathbb{Z}$ we have

$$\tilde{\rho}_*(\mathcal{L} \otimes \tilde{\rho}^* \xi^{\otimes t}) = \mathcal{E}(\mathcal{L}) \otimes \xi^{\otimes t}, \quad (3.1.6)$$

$$c_1(\mathcal{L} \otimes \tilde{\rho}^* \xi^{\otimes t}) = \nu^*(\mu_B(\omega_B) + (x+t)\delta(B)) + (y+t)\text{cl}(D). \quad (3.1.7)$$

This explains why the hypotheses on (x, y) that ensure that $\mathcal{E}(\mathcal{L})$ is modular are invariant under translation by multiples of $(1, 1)$.

Remark 3.4. Let (X, h) be a general polarized HK of type $K3^{[2]}$. In [O'G22] we constructed stable modular vector bundles \mathcal{F} on X of rank an arbitrary square (but there are conditions on h) and

$$\Delta(\mathcal{F}) = \frac{r(\mathcal{F})(r(\mathcal{F}) - 1)}{12} c_2(X). \quad (3.1.8)$$

Notice that if $r(\mathcal{F}) = 4$ then the equation in (3.1.8) is the exact analogue of the equation in (3.1.5). In both cases the stable vector bundle is the unique stable vector bundle with given rank, first and second Chern class. In order to gain some perspective we notice that every vector bundle \mathcal{G} on a $K3$ surface is modular and that if it is stable with zero dimensional moduli space then

$$\Delta(\mathcal{G}) = \frac{r(\mathcal{G})^2 - 1}{12} c_2(S). \quad (3.1.9)$$

3.2. From $S^{[2]}$ to $S^{[3]}$ according to Ellingrud and Strømme. Let S be a smooth surface, and let

$$S^{[2,3]} := \{[W], [Z]\} \in S^{[2]} \times S^{[3]} \mid W \subset Z\}.$$

Let $\alpha: S^{[2,3]} \rightarrow S^{[2]} \times S$ be the product of the projection $S^{[2,3]} \rightarrow S^{[2]}$ and the map $([W], [Z]) \mapsto \text{supp}(\mathcal{I}_W/\mathcal{I}_Z)$. Let $\beta: S^{[2,3]} \rightarrow S^{[3]}$ be the projection. We have a commutative diagram

$$\begin{array}{ccc} & S^{[2,3]} & \\ \alpha \swarrow & & \searrow \beta \\ S^{[2]} \times S & \xrightarrow{\xi} & S^{[3]}, \end{array}$$

where $\xi([W], p) := [W \sqcup \{p\}]$ if $p \notin \text{supp } W$. Let $\mathcal{Z}_n(S) \subset S^{[n]} \times S$ be the universal subscheme.

Proposition 3.5 (Ellingrud-Strømme, Prop. 2.1 and 2.2 of [ES98]). *The map α is the blow up of $\mathcal{Z}_2(S)$. The Stein factorization of β is*

$$S^{[2,3]} \xrightarrow{\gamma} \mathcal{Z}_3(S) \longrightarrow S^{[3]},$$

where $\mathcal{Z}_3(S) \longrightarrow S^{[3]}$ is the projection.

The map γ is an isomorphism away from the subset $\{(V(\mathfrak{m}_p^2), p) \mid p \in S\}$. Moreover

$$\gamma^{-1}((V(\mathfrak{m}_p^2), p)) = \beta^{-1}([V(\mathfrak{m}_p^2)]) = \{([W], [V(\mathfrak{m}_p^2)]) \mid W \subset V(\mathfrak{m}_p^2)\} \cong \mathbb{P}(\Theta_p(S)).$$

3.3. The maps ρ and $\tilde{\rho}$.

Proof of Proposition 3.1. Let $\rho: B^{[3]} \dashrightarrow A^{[3]}$ be defined as ρ , i.e. $\rho([Z]) = [f(Z)]$, and let $\mathbf{V}^{[3]}(f) \subset B^{[3]}$ be the set of $[Z]$ such that $\ell(f(Z)) < \ell(Z)$. It suffices to prove that $\mathbf{V}^{[3]}(f)$ is irreducible of codimension 2 and smooth, and that blowing it up we resolve the indeterminacies of ρ . Let $\varepsilon \in B$ be the non zero element of $\ker f$:

$$\ker f = \{0, \varepsilon\}. \quad (3.3.1)$$

A simple but useful observation is contained in the following equality:

$$\mathbf{V}^{[3]}(f) = \{[Z] \in B^{[3]} \mid Z = W \sqcup \{b\}, \quad [W] \in B^{[2]}, \quad (b + \varepsilon) \in \text{supp } W\}. \quad (3.3.2)$$

Let $\sigma: B^{[2]} \times B \rightarrow B^{[2]} \times B$ be defined by $\sigma([W], b) := ([W], b + \varepsilon)$. Consider the maps

$$B^{[2]} \times B \xrightarrow{\sigma} B^{[2]} \times B \xrightarrow{\xi} B^{[3]} \xrightarrow{\rho} A^{[3]}.$$

Let $[Z] \in \mathbf{V}^{[3]}(f)$. Then we have $Z = W \sqcup \{b\}$ as in (3.3.2). Since $b \notin \text{supp } W$, the map ξ is regular at $([W], b)$ and defines an isomorphism between a small neighborhood (in the analytic topology) \hat{U} of $([W], b)$ and a small neighborhood (in the analytic topology) U of $[Z]$. Moreover σ maps isomorphically $\hat{U} \cap \mathcal{Z}_2(B)$ to $U \cap \xi^{-1}(\mathbf{V}^{[3]}(f))$. Since $\mathcal{Z}_2(B)$ is isomorphic to the blow up of $B \times B$ along the diagonal it is irreducible of codimension 2 and smooth. It follows that the same holds for $\mathbf{V}^{[3]}(f)$.

Moreover σ identifies (locally around $([W], b)$ and $[Z]$) the blow ups of $B^{[2]} \times B$ with centers $\mathcal{Z}_2(B)$ and $\mathbf{V}^{[3]}(f)$. Since $\rho \circ \xi \circ \sigma = \rho \circ \xi$ the blow up of $\mathbf{V}^{[3]}(f)$ resolves the indeterminacies of ρ by Ellingsrud-Strømme's Proposition 3.5. \square

Given $b \in B[3]$ let

$$R_b := \{[\{b\} \sqcup W] \mid W \in B^{[2]}, \text{supp } W = \{b + \varepsilon\}\} \subset K_2(B). \quad (3.3.3)$$

Notice that R_b is isomorphic to \mathbb{P}^1 .

Proposition 3.6. *Let $X \xrightarrow{\tilde{\rho}_2} \overline{X} \xrightarrow{\tilde{\rho}_1} K_2(A)$ be the Stein factorization of $\tilde{\rho}$.*

- (a) $\tilde{\rho}_1$ is a finite map of degree 4.
- (b) There exists an embedding $F: B[3] \hookrightarrow \overline{X}$ with the following properties. Let $b \in B[3]$, and let $a := f(b) \in A[3]$. There is a curve $\tilde{R}_b \subset X$ mapped isomorphically to R_b by ν and contracted by $\tilde{\rho}_2$ to a point $F(b) \in \tilde{\rho}_1^{-1}(a)$. Moreover $\tilde{R}_b = \tilde{\rho}_2^{-1}(F(b))$.
- (c) Away from the union of the \tilde{R}_b 's (for $b \in B[3]$) the map $\tilde{\rho}_2$ is an isomorphism onto its image.
- (d) \overline{X} has rational singularities.

Proof. Items (a)-(c) follow easily from the proof of Proposition 3.1 together with Ellingsrud and Strømme's Proposition 3.5. Let us prove Item (d). First \overline{X} is smooth away from $F(B[3])$. On the other hand we claim that \overline{X} in a neighborhood of $F(b)$, for $b \in B[3]$, is isomorphic to the fiber over 0 of the composition

$$\mathcal{Z}_3(A) \longrightarrow A^{[3]} \longrightarrow A^{(3)} \longrightarrow A,$$

where the last map is the summation map. In fact our assertion follows from the proof of Proposition 3.1 and from Proposition 3.5. Since $\mathcal{Z}_3(A) \longrightarrow A$ is a locally (in the classical or étale topology) trivial fibration, \overline{X} has a rational singularity at $F(b)$ if $\mathcal{Z}_3(A)$ has rational singularities. The last assertion holds by the main result of [Son16]. \square

3.4. Chern classes, I. We follow the notation introduced in Theorem 3.2. We will set $\mathcal{E} = \mathcal{E}(\mathcal{L})$.

Lemma 3.7. *For $\mathcal{E} = \mathcal{E}(\mathcal{L})$ the following hold:*

$$\begin{aligned} \text{ch}_1(\mathcal{E}) &= \tilde{\rho}_* \left[c_1(\mathcal{L}) + \frac{c_1(X)}{2} \right], \\ \text{ch}_2(\mathcal{E}) &= \tilde{\rho}_* \left[\frac{c_1(\mathcal{L})^2}{2} + \frac{c_1(\mathcal{L}) \smile c_1(X)}{2} + \frac{c_1(X)^2 + c_2(X)}{12} \right] - \frac{c_2(K_2(A))}{3}. \end{aligned}$$

Proof. By Proposition 3.6 the higher direct images sheaves $R^i \tilde{\rho}_* \mathcal{L}$ (for $i > 0$) vanish over $(K_2(A) \setminus A[3])$. Hence the Chern character of \mathcal{E} over $(K_2(A) \setminus A[3])$ is computed by GRR. Since $A[3]$ has codimension 4 in $K_2(A)$ it follows that the same formula gives $\text{ch}_p(\mathcal{E})$ on $K_2(A)$ for $p \leq 3$. Writing out GRR we get the formulae of the corollary. \square

Proposition 3.8. *For $\mathcal{E} = \mathcal{E}(\mathcal{L})$ and $\alpha \in H^2(K_2(A))$ we have*

$$\int_{K_2(A)} \text{ch}_1(\mathcal{E})^2 \smile \alpha^2 = 6 \left(2 \int_A (f_* \omega_B)^2 - 3(2x + 2y - 1)^2 \right) q(\alpha) + 6q(2\mu_A(f_* \omega_B) + (2x + 2y - 1)\delta(A), \alpha)^2. \quad (3.4.1)$$

Proof. Since X is the blow up of $K_2(B)$ with center $V(f)$, we have

$$c_1(X) = -\text{cl}(D). \quad (3.4.2)$$

Thus Lemma 3.7 gives

$$\text{ch}_1(\mathcal{E}) = \tilde{\rho}_* c_1(\mathcal{L}) - \frac{1}{2} \tilde{\rho}_* \text{cl}(D). \quad (3.4.3)$$

We claim that

$$\tilde{\rho}_* \nu^* \mu_B(\omega_B) = 2\mu_A(f_* \omega_B), \quad \tilde{\rho}_* \nu^* \Delta(B) = 2\Delta(A), \quad \tilde{\rho}_* D = \Delta(A). \quad (3.4.4)$$

One gets the first equality by representing the Poincaré dual of ω_B by a C^∞ immersed submanifold Σ , and the Poincaré dual of $\mu_B(\omega_B)$ by the immersed submanifold I_Σ of $[Z] \in K_2(B)$ meeting Σ . It follows that the Poincaré dual of $\tilde{\rho}_* \nu^* \mu_B(\omega_B)$ is represented by the immersed submanifold $I_{f(\Sigma)}$ of $[Z] \in K_2(A)$ meeting $f(\Sigma)$ (the point being that I_Σ meets properly $V(f)$ for a generic choice of Σ), counted with multiplicity 2 because the map $I_\Sigma \rightarrow I_{f(\Sigma)}$ has degree 2.

In order to prove the second equality in (3.4.4) notice that $\Delta(B)$ intersects properly $V(f)$, and hence $\tilde{\rho}_* \nu^* \Delta(B)$ is represented by the closure of $\tilde{\rho}(\Delta(B) \setminus V(f))$, which is $\Delta(A)$, with multiplicity the degree of the map $(\Delta(B) \setminus V(f)) \rightarrow \Delta(A)$, which is 2.

The proof of Proposition 3.1 gives the third equality in (3.4.4).

Plugging into (3.4.3) the equations in (3.4.4), we get that

$$\text{ch}_1(\mathcal{E}) = 2\mu_A(f_* \omega_B) + (2x + 2y - 1)\delta(A). \quad (3.4.5)$$

The proposition follows from (3.4.5) and the formula in (2.2.4). \square

3.5. Analysis of $V(f)$. In order to compute $\int_{K_2(A)} \text{ch}_2(\mathcal{E}) \smile \alpha^2$ for $\alpha \in H^2(K_2(A))$ we must examine more closely $V(f)$. Let $\mathfrak{h}: K_2(B) \rightarrow B^{(3)}$ be the Hilbert-Chow map. The image of \mathfrak{h} is equal to

$$\mathfrak{S} := \{(b_1) + (b_2) + (b_3) \mid b_1 + b_2 + b_3 = 0\}.$$

Let $\mathfrak{S}_\epsilon \subset \mathfrak{S}$ be the subset of $(b_1) + (b_2) + (b_3)$ such that $b_i - b_j = \epsilon$ for some $i, j \in \{1, 2, 3\}$. Then $V(f) = \mathfrak{h}^{-1}(\mathfrak{S}_\epsilon)$, and the restriction of \mathfrak{h} to $V(f)$ is a birational map $\mathfrak{h}_{V(f)}: V(f) \rightarrow \mathfrak{S}_\epsilon$. Let

$$\begin{array}{ccc} B & \xrightarrow{g} & \mathfrak{S}_\epsilon \\ b & \mapsto & (b) + (b + \epsilon) + (-2b + \epsilon) \end{array} \quad (3.5.1)$$

Then g identifies \mathfrak{S}_ϵ with $B/\ker f = A$.

Proposition 3.9. *Identifying S_ϵ with A via the map g , the map $\mathfrak{h}_{V(f)}: V(f) \rightarrow S_\epsilon$ is identified with the blow up of $A[3]$. Let R_1, \dots, R_{81} be the exceptional divisors of $\mathfrak{h}_{V(f)}$ (notice that they are equal to $R_{b_1}, \dots, R_{b_{81}}$ where $b_j \in B[3]$, notation as in (3.3.3)). Then*

$$\Delta(B)|_{V(f)} = 2 \sum_{i=1}^{81} R_i. \quad (3.5.2)$$

Proof. By Proposition 3.1 we know that $V(f)$ is smooth. Moreover $\mathfrak{h}_{V(f)}$ is an isomorphism over $(A \setminus A[3])$ and it has fiber \mathbb{P}^1 over each point of $A[3]$. It follows that $\mathfrak{h}_{V(f)}$ is the blow up of $A[3]$. The remaining part of the proposition is straightforward. \square

Corollary 3.10. *Let $V(f) \subset K_2(B)$ be as in (3.1.2), and let $\zeta \in H^2(B)$. Then*

$$\int_{V(f)} (\mu_B(\zeta) + t\delta(B))^2 = 18 \left(\int_B \zeta^2 \right) - 81t^2.$$

Proof. We have $\mu_B(\zeta) = \mathfrak{h}^*(\zeta^{(3)})$, where $\mathfrak{h}: K_2(B) \rightarrow B^{(3)}$ be the Hilbert-Chow map and $\zeta^{(3)} \in H^2(B^{(3)})$ is the symmetrization of the class ζ . Since $g^*(\zeta^{(3)}) = 6\zeta$ and g is surjective of degree 2, we get that

$$\int_{V(f)} \mu_B(\zeta)^2 = \int_W \zeta^{(3)} \smile \zeta^{(3)} = \frac{1}{2} \int_B (6\zeta)^2 = 18 \int_B \zeta^2.$$

Next we notice that

$$\int_{V(f)} \mu_B(\zeta) \smile \delta(B) = \int_W \zeta^{(3)} \smile \mathfrak{h}_{V(f),*}(\delta(B)) = 0. \quad (3.5.3)$$

(The last equality holds because $\mathfrak{h}_{V(f),*}(\Delta(B)) = 0$.)

Lastly, by (3.5.2) we have

$$\int_{V(f)} \delta(B)^2 = -81 \quad (3.5.4)$$

\square

Lemma 3.11. *The restriction of $\tilde{\rho}$ defines a map $\nu^*\Delta(B) \rightarrow \Delta(A)$ of degree 2, and a birational map $D \rightarrow \Delta(A)$. We have*

$$\tilde{\rho}^*\delta(A) = \nu^*\delta(B) + \text{cl}(D). \quad (3.5.5)$$

Proof. The first sentence is immediate. We also see that ρ is unramified at the generic point of $\Delta(B)$, and hence

$$\tilde{\rho}^*\Delta(A) = \nu^*\Delta(B) + m \text{cl}(D). \quad (3.5.6)$$

Since $K_X \equiv D$, we get $m = 2$ by the adjunction formula for the map $\tilde{\rho}$. Dividing by 2 the equality in (3.5.6) we get (3.5.5). \square

Proposition 3.12. *We have*

$$c_1(\mathcal{N}_{V(f)/K_2(B)}) = \sum_{i=1}^{81} \text{cl}(R_i) = \delta(B)|_{V(f)}, \quad (3.5.7)$$

$$\int_{V(f)} c_2(\mathcal{N}_{V(f)/K_2(B)}) = 3^4, \quad (3.5.8)$$

$$\int_{V(f)} c_2(K_2(B)) = 3^5. \quad (3.5.9)$$

Proof. By Proposition 3.9 we have $c_1(V(f)) = -\sum_{i=1}^{81} \text{cl}(R_i)$, and hence (3.5.7) holds because $c_1(K_2(B)) = 0$ (remember Proposition 3.9).

In order to prove (3.5.8) we notice that

$$\begin{aligned} 4 \cdot 81 &= \int_{K_2(A)} \delta(A)^4 = \frac{1}{4} \int_X (\nu^* \delta(B) + \text{cl}(D))^4 = \\ &= \frac{1}{4} \int_{K_2(B)} \delta(B)^4 + \int_X \nu^* \delta(B)^3 \smile \text{cl}(D) + \frac{3}{2} \int_X \nu^* \delta(B)^2 \smile \text{cl}(D)^2 + \\ &+ \int_X \nu^* \delta(B) \smile \text{cl}(D)^3 + \frac{1}{4} \int_X \text{cl}(D)^4 = 81 + \frac{3}{2} \cdot 81 + 81 + \frac{1}{4} \int_X \text{cl}(D)^4. \end{aligned} \quad (3.5.10)$$

(To compute the third and fourth integrals in the second line use Corollary 3.10 and (3.5.7).) It follows that

$$2 \cdot 81 = \int_X \text{cl}(D)^4 = \int_{V(f)} c_2(\mathcal{N}_{V(f)/K_2(B)}) - c_1(\mathcal{N}_{V(f)/K_2(B)})^2. \quad (3.5.11)$$

By (3.5.7) we get that (3.5.8) holds.

Lastly (3.5.9) follows from (3.5.7), (3.5.8) and the normal exact sequence for the restriction of Θ_X to $V(f)$. \square

3.6. Chern classes, II. By definition D is $\mathbb{P}(\mathcal{N}_{V(f)/K_2(B)})$, where $\mathcal{N}_{V(f)/K_2(B)}$ is the normal bundle of $V(f)$ in $K_2(B)$. Let $\nu_D: D \rightarrow V(f)$ be the structure map (i.e. the restriction of ν), and let

$$0 \longrightarrow \lambda \longrightarrow \nu_D^* \mathcal{N}_{V(f)/K_2(B)} \longrightarrow \xi \longrightarrow 0 \quad (3.6.1)$$

be the tautological exact sequence, where $\lambda = \mathcal{O}_D(-1)$ is the normal bundle of D in X . Let $i: D \hookrightarrow X$ be the inclusion map.

Lemma 3.13. *We have*

$$c_2(X) = \nu^* c_2(K_2(B)) + i_* [c_1(\xi)]. \quad (3.6.2)$$

Proof. The exact sequence

$$0 \longrightarrow \Theta_X \xrightarrow{d\nu} \nu^* \Theta_{K_2(B)} \longrightarrow i_* \xi \longrightarrow 0$$

gives

$$c(X) = \nu^* c(K_2(B)) \smile c(i_* \xi)^{-1}. \quad (3.6.3)$$

The Chern classes of $i_* \xi$ are expressed via GRR. One gets $c_1(i_* \xi) = \text{cl}(D)$ and $c_2(i_* \xi) = i_* [c_1(\lambda) - c_1(\xi)]$. Plugging this into (3.6.3) we get the lemma. \square

Lemma 3.14. *For $\mathcal{E} = \mathcal{E}(\mathcal{L})$ we have*

$$\begin{aligned} \mathrm{ch}_2(\mathcal{E}) &= \frac{1}{2} \tilde{\rho}_* \{ \nu^*(\mu_B(\omega_B) + x\delta(B))^2 + (2y-1)\nu^*(\mu_B(\omega_B) + x\delta(B)) \smile \mathrm{cl}(D) \} + \\ &\quad + \frac{1}{12} \tilde{\rho}_* \{ i_* [c_1(\xi) + (6y^2 - 6y + 1)c_1(\lambda)] \} \\ &\quad + \frac{1}{12} \tilde{\rho}_* (\nu^* c_2(K_2(B))) - \frac{1}{3} c_2(K_2(A)). \end{aligned} \quad (3.6.4)$$

Proof. Plug (3.4.2) and (3.6.2) in the second equation of Lemma 3.7. \square

We are now ready to compute $\int_{K_2(A)} \Delta(\mathcal{E}) \smile \alpha^2$ for $\alpha \in H^2(K_2(A))$.

Proposition 3.15. *For $\mathcal{E} = \mathcal{E}(\mathcal{L})$ and $\gamma \in H^2(A)$ we have*

$$\int_{K_2(A)} \Delta(\mathcal{E}) \smile \mu_A(\gamma)^2 = 18(4(x-y)^2 + 4(x-y) + 3) q(\mu_A(\gamma)). \quad (3.6.5)$$

Proof. First we notice that

$$\int_B f^* \gamma^2 = 2 \int_A \gamma^2, \quad \int_A (f_* \omega_B)^2 = 2 \int_B \omega_B^2, \quad \tilde{\rho}^* \mu_A(\gamma) = \nu^* \mu_B(f^* \gamma). \quad (3.6.6)$$

The first equality holds because $\deg f = 2$. The second equality holds by the following series of equalities

$$\int_A (f_* \omega_B)^2 = \int_A f_*(\omega_B \smile f^*(f_* \omega_B)) = \int_B \omega_B \smile (\omega_B + T_\epsilon^* \omega_B) = 2 \int_B \omega_B^2. \quad (3.6.7)$$

(We let $T_\epsilon: B \rightarrow B$ be the translation by ϵ .) The third equality in (3.6.6) is proved by representing the Poincaré dual of $\mu_A(\gamma)$ and $\mu_B(f^* \gamma)$ as in the proof of the first equality in (3.4.4).

By Proposition 3.8 we have

$$\int_{K_2(A)} \mathrm{ch}_1(\mathcal{E})^2 \smile \alpha^2 = \left(24 \int_B \omega_B^2 - 18(2x+2y-1)^2 \right) \cdot \left(\int_A \gamma^2 \right) + 24 \left(\int_A f_* \omega_B \smile \gamma \right)^2. \quad (3.6.8)$$

On the other hand Lemma 3.14 and the third equality in (3.6.6) give that

$$\begin{aligned} \int_{K_2(A)} \mathrm{ch}_2(\mathcal{E}) \smile \mu_A(\gamma)^2 &= \frac{1}{2} \int_{K_2(B)} (\mu_B(\omega_B) + x\delta(B))^2 \smile \mu_B(f^* \gamma)^2 \\ &\quad - \frac{1}{2} (y^2 - y) \int_{V(f)} \mu_B(f^* \gamma)^2 + \\ &\quad + \frac{1}{12} \int_{K_2(B)} c_2(K_2(B)) \smile \mu_B(f^* \gamma)^2 - \frac{1}{3} \int_{K_2(A)} c_2(K_2(A)) \smile \mu_A(\gamma)^2. \end{aligned} \quad (3.6.9)$$

Recalling the results of Subsection 2.2 and Corollary 3.10 we get that

$$\begin{aligned} \int_{K_2(A)} \mathrm{ch}_2(\mathcal{E}) \smile \mu_A(\gamma)^2 &= \\ &= 3 \left(\left(\int_B \omega_B^2 \right) - 6x^2 - 6y^2 + 6y - 3 \right) \cdot \left(\int_A \gamma^2 \right) + 3 \left(\int_B \omega_B \smile f^* \gamma \right)^2 \end{aligned} \quad (3.6.10)$$

Since \mathcal{E} has rank 4 we have $\Delta(\mathcal{E}) = \text{ch}_1(\mathcal{E})^2 - 8 \text{ch}_2(\mathcal{E})$. The proposition follows from (3.6.8) and (3.6.10) (recall that $\mu: H^2(A) \rightarrow H^2(K_2(A))$ is an isometry). \square

Proposition 3.16. *For $\mathcal{E} = \mathcal{E}(\mathcal{L})$ and $\gamma \in H^2(A)$ we have*

$$\int_{K_2(A)} \Delta(\mathcal{E}) \smile \mu_A(\gamma) \smile \delta(A) = 0. \quad (3.6.11)$$

Proof. By Proposition 3.8 we have

$$\int_{K_2(A)} \text{ch}_1^2(\mathcal{E}) \smile \mu_A(\gamma) \smile \delta(A) = -72(2x + 2y - 1) \int_A \gamma \smile f_*\omega_B.$$

Lemmas 3.14, 3.11, Corollary 3.10 and (3.5.7) give that

$$\begin{aligned} \int_{K_2(A)} \text{ch}_2(\mathcal{E}) \smile \mu_A(\gamma) \smile \delta(A) &= \frac{1}{2} \int_{K_2(B)} (\mu_B(\omega_B) + x\delta(B))^2 \smile \mu_B(f^*\gamma) \smile \delta(B) + \\ &+ \frac{1}{2}(2y - 1)\nu^*(\mu_B(\omega_B) + x\delta(B)) \smile \mu_B(f^*\gamma) \smile \text{cl}(D)^2 = \\ &= -9(2x + 2y - 1) \int_B \omega_B \smile f^*\gamma. \end{aligned}$$

Since $\Delta(\mathcal{E}) = \text{ch}_1(\mathcal{E})^2 - 8 \text{ch}_2(\mathcal{E})$ the proposition follows from the above equalities. \square

Proposition 3.17. *Keeping notation as above, we have*

$$\int_{K_2(A)} \Delta(\mathcal{E}(\mathcal{L})) \smile \delta(A)^2 = 54((x - y)^2 + (x - y) + 1)q(\delta(A)). \quad (3.6.12)$$

Proof. Let $\mathcal{E} = \mathcal{E}(\mathcal{L})$. By Proposition 3.8 we have

$$\int_{K_2(A)} \text{ch}_1(\mathcal{E})^2 \smile \delta(A)^2 = 324(2x + 2y - 1)^2 - 72 \int_A (f_*\omega_B)^2. \quad (3.6.13)$$

By Lemmas 3.14 and 3.11 we have

$$\begin{aligned}
\int_{K_2(A)} \text{ch}_2(\mathcal{E}) \smile \delta(A)^2 &= -\frac{1}{3} \int_{K_2(A)} c_2(K_2(A)) \smile \delta(A)^2 + \\
+ \frac{1}{2} \int_{K_2(B)} (\mu_B(\omega_B) + x\delta(B))^2 \smile \delta(B)^2 &+ \frac{1}{12} \int_D [c_1(\xi) + (6y^2 - 6y + 1)c_1(\lambda)] \smile \delta(B)^2 + \\
+ \frac{1}{12} \int_{K_2(B)} c_2(K_2(B)) \smile \delta(B)^2 &+ (2y-1) \int_X \nu^*((\mu_B(\omega_B) + x\delta(B)) \smile \delta(B)) \smile \text{cl}(D)^2 + \\
&+ \frac{1}{6} \int_D [c_1(\xi) \cup c_1(\lambda) + (6y^2 - 6y + 1)c_1(\lambda)^2] \smile \delta(B) + \\
+ \frac{1}{2} \int_X \nu^*(\mu_B(\omega_B) + x\delta(B))^2 \smile \text{cl}(D)^2 &+ \frac{1}{2}(2y-1) \int_X \nu^*(\mu_B(\omega_B) + x\delta(B)) \smile \text{cl}(D)^3 + \\
&+ \frac{1}{12} \int_D [c_1(\xi) \smile c_1(\lambda)^2 + (6y^2 - 6y + 1)c_1(\lambda)^3] + \\
&+ \frac{1}{12} \int_X \nu^* c_2(K_2(B)) \smile \text{cl}(D)^2. \quad (3.6.14)
\end{aligned}$$

We have proved all the results needed to compute each integral above. In particular notice that (3.6.1) and Proposition 3.12 give the following relations in the cohomology ring of D :

$$c_1(\xi) = \sum_{i=1}^{81} \text{cl}(R_i) - c_1(\lambda), \quad c_1(\lambda)^2 = \nu_D^* \left(\sum_{i=1}^{81} \text{cl}(R_i) \right) \smile c_1(\lambda) - 81\nu_D^*(\eta_{V(f)}),$$

where $\nu_D: D \rightarrow V(f)$ is the natural map (the restriction of ν to D) and $\eta_{V(f)}$ is the fundamental class of $V(f)$. It follows that

$$\int_{K_2(A)} \text{ch}_2(\mathcal{E}) \smile \delta(A)^2 = \frac{81}{2} (5x^2 + 6xy + 5y^2 - 3x - 5y + 2) - 18 \int_B \omega_B^2. \quad (3.6.15)$$

The proposition follows from (3.6.13) and (3.6.15) because $\Delta(\mathcal{E}) = -8\text{ch}_2(\mathcal{E}) + \text{ch}_1(\mathcal{E})^2$. \square

Notice that the right hand sides of (3.6.5), (3.6.11) and (3.6.12) are polynomials in $(x - y)$; this is explained by Remark 3.3.

3.7. Modularity. By Propositions 3.15, 3.16 and 3.17 the sheaf $\mathcal{E}(\mathcal{L})$ is modular if and only if

$$18(4(x - y)^2 + 4(x - y) + 3) = 18(3(x - y)^2 + 3(x - y) + 3). \quad (3.7.1)$$

The solutions are given by $x - y = 0$ and $x - y = -1$. For modular $\mathcal{E}(\mathcal{L})$ one gets that

$$d(\mathcal{E}(\mathcal{L})) = 54 \quad (3.7.2)$$

by plugging the values $x - y = 0$ and $x - y = -1$ into the polynomial on the right hand side of (3.6.5) or (3.6.12) (we recall that $d(\mathcal{E}(\mathcal{L}))$ is defined by the equation in (1.1.1)).

3.8. Action of $\text{Aut}^0(K_2(A))$ on $\mathcal{E}(\mathcal{L})$. We recall that if M is a HK manifold, then $\text{Aut}^0(M)$ is the group of automorphisms of M acting trivially on $H^2(M)$. The isomorphism class of $\text{Aut}^0(M)$ depends only on the deformation class of M . If M is a $2n$ dimensional HK manifold of Kummer type ($n \geq 2$) then $\text{Aut}^0(M)$ is a semidirect product of $\mathbb{Z}/(n+1)^4$ and $\mathbb{Z}/(2)$. More precisely, $\text{Aut}^0(K_2(A))$ is generated by the following automorphisms. For $\tau \in A[3]$ let $g_\tau: K_2(A) \rightarrow K_2(A)$ be the automorphism mapping a subscheme Z to $\tau(Z)$. Then $g_\tau \in \text{Aut}^0(K_2(A))$, and the set of such automorphism is a normal subgroup isomorphic to $\mathbb{Z}/(n+1)^4$. Let $\iota: K_2(A) \rightarrow K_2(A)$ be the automorphism mapping a subscheme Z to $-1(Z)$ (here $-1: A \rightarrow A$ is multiplication by -1). Then ι belongs to $\text{Aut}^0(K_2(A))$ and moreover $\text{Aut}^0(K_2(A))$ is generated by the collection of the g_τ 's and ι .

Proposition 3.18. *Let \mathcal{L} be a line bundle on X as in Subsection 3.1 (no further assumptions on \mathcal{L}). If $\varphi \in \text{Aut}^0(K_2(A))$ then*

$$\varphi^*(\mathcal{E}) \cong \mathcal{E}. \quad (3.8.1)$$

Proof. We must prove that $g_\tau^*(\mathcal{E}) \cong \mathcal{E}$ for all $\tau \in A[3]$, and that $\iota^*(\mathcal{E}) \cong \mathcal{E}$. We start by noting that the restriction of f to $B[3]$ defines an isomorphism $B[3] \rightarrow A[3]$. Let $\lambda \in B[3]$ be the element such that $f(\lambda) = \tau$. The corresponding automorphism $g_\lambda: K_2(B) \rightarrow K_2(B)$ acts trivially on $H^2(K_2(B))$ and hence on $\text{Pic}(K_2(B))$. Thus $g_\lambda^*(\mathcal{L}) \cong \mathcal{L}$. Now g_λ lifts to an automorphism $\tilde{g}_\lambda: X \rightarrow X$ because g_λ maps $V(f)$ to itself, and of course $\tilde{g}_\lambda^*(\mathcal{L}) \cong \mathcal{L}$. Since $\tilde{\rho} \circ \tilde{g}_\lambda = g_\tau \circ \tilde{\rho}$, we get $g_\tau^*(\mathcal{E}) \cong \mathcal{E}$. The proof that $\iota^*(\mathcal{E}) \cong \mathcal{E}$ is similar. \square

3.9. The discriminant. Next we prove that the equation in (3.1.5) holds. We assume that $\mathcal{E}(\mathcal{L})$ is modular, i.e. either $y = x$ or $y = x + 1$. As mentioned in Subsection 2.1, this means that

$$\Delta(\mathcal{E}) \in \mathbb{Q}\bar{c}_2(K_2(A)) \oplus \text{Sym}^2 H^2(M)_{\mathbb{Q}}^{\perp}, \quad (3.9.1)$$

where $\bar{c}_2(K_2(A))$ is the orthogonal projection of $c_2(K_2(A))$ in the image of the map $\text{Sym}^2 H^2(K_2(A)) \rightarrow H^4(K_2(A))$. The right hand side of (3.9.1) is studied in [HT13]. In Theorem 4.4 of loc.cit. one finds the definition of submanifolds $Z_\tau \subset K_2(A)$ for $\tau \in A[3]$ with the following properties: the cohomology classes $\{\text{cl}(Z_\tau)\}_{\tau \in A[3]}$ give a basis the right hand side of (3.9.1) and the action of $A[3]$ on the Z_τ 's is by translation of the index. It follows that the subspace of invariants for this action has dimension 1, and since $c_2(K_2(A))$ is a non zero class it generates the space of invariants (by Proposition 5.1 in [HT13] we have $c_2(K_2(A)) = \frac{1}{3} \sum_{\tau} \text{cl}(Z_\tau)$). Since (3.8.1) holds for all $\varphi = g_\tau$, where $\tau \in A[3]$, $\Delta(\mathcal{E})$ is invariant and hence it is a multiple of $c_2(K_2(A))$. By (6.1.6) and (2.2.6) it follows that $\Delta(\mathcal{E}) = c_2(K_2(A))$.

3.10. Local structure of modular \mathcal{E} .

Proposition 3.19. *Let \mathcal{L} be a line bundle on X such that*

$$c_1(\mathcal{L}) = \nu^*(\mu_B(\omega_B) + x\delta(B)) + x \text{cl}(D),$$

i.e. $x = y$ in (3.1.4). Then $R^i \tilde{\rho}_{2,} \mathcal{L} = 0$ for $i > 0$, and $\mathcal{E}(\mathcal{L})$ is locally free.*

Proof. Let $X \xrightarrow{\tilde{\rho}_2} \bar{X} \xrightarrow{\tilde{\rho}_1} K_2(A)$ be the Stein factorization of $\tilde{\rho}$ - see Proposition 3.6. It suffices to prove that $\tilde{\rho}_{2,*} \mathcal{L}$ is an invertible sheaf and that

$$R^i \tilde{\rho}_{2,*} \mathcal{L} = 0, \quad i > 0. \quad (3.10.1)$$

By (3.1.6) and (3.1.7) we may assume that $x = y = 0$. Let $F: B[3] \hookrightarrow \bar{X}$ be the embedding of Item (b) of Proposition 3.6. The map $\tilde{\rho}_2$ is an isomorphism over $(\bar{X} \setminus F(B[3]))$. Hence away from $F(B[3])$ the sheaf $\tilde{\rho}_{2,*} \mathcal{L}$ is invertible and (3.10.1) holds.

Let $b \in B[3]$. The fiber $\tilde{R}_b = \tilde{\rho}_2^{-1}(F(b))$ is mapped isomorphically to $R_b \subset K_2(B)$, where R_b is defined in (3.3.3). Since $x = y = 0$ the line bundle \mathcal{L} is the pull back of a line bundle on $B^{(3)}$ via the Hilbert-Chow map $K_2(B) \rightarrow B^{(3)}$. The curve R_b (a \mathbb{P}^1) is contracted by the Hilbert-Chow map. It follows that \mathcal{L} is trivial on an open neighborhood \mathcal{U} of \tilde{R}_b . Shrinking \mathcal{U} we may assume that $\tilde{\rho}_2(\mathcal{U})$ is open (because $\tilde{R}_b = \tilde{\rho}_2^{-1}(F(a))$, where $a = f(b)$, and $\tilde{\rho}_2$ is projective), and hence $\tilde{\rho}_{2,*}\mathcal{L}$ is invertible in a neighborhood of $F(b)$.

By Proposition 3.6 we know that \overline{X} has rational singularities. Hence (3.10.1) holds because in a neighborhood of \tilde{R}_b we have $\mathcal{L} \cong \mathcal{O}_X$. \square

4. MODULAR BASIC VECTOR BUNDLES WHICH EXTEND SIDEWAYS

4.1. Main result. Let $\nu: X \rightarrow K_2(B)$ be the blow up of $V(f)$, as in Subsection 3.1, and let \mathcal{L} be a line bundle on X as in (3.1.4) such that $x = y$. Then $\mathcal{E}(\mathcal{L})$ is a modular vector bundle on $K_2(B)$ by Theorem 3.2. In the present section we prove that the natural map $\text{Def}(K_2(A), \mathcal{E}(\mathcal{L})) \rightarrow \text{Def}(K_2(A), \det \mathcal{E}(\mathcal{L}))$ is surjective under suitable additional hypotheses.

In order to state our result we go through some preliminaries.

Hypothesis 4.1. (1) B is an abelian surface containing a (bona fide) elliptic curve $C_B \subset B$ with Poincarè dual denoted by γ_B .
(2) $\omega_B \in H_{\mathbb{Z}}^{1,1}(B)$ is a class such that the subgroup $(\mathbb{Z}\omega_B + \mathbb{Z}\gamma_B) < H_{\mathbb{Z}}^{1,1}(B)$ is saturated, and

$$\omega_B \cdot \omega_B = 2a, \quad \omega_B \cdot \gamma_B = d. \quad (4.1.1)$$

(When no confusion is possible \cdot denotes the intersection pairing.)

Assume that Hypothesis 4.1 holds, and choose a non zero $\epsilon \in C_B[2]$. Let $A := B/\langle \epsilon \rangle$ and let $f: B \rightarrow A$ be the quotient map. Thus we are in the set up of Subsection 3.1. Let \mathcal{L} be the line bundle on X such that

$$c_1(\mathcal{L}) = \nu^*(\mu_B(m\omega_B)). \quad (4.1.2)$$

In other words we set $x = y = 0$ in (3.1.4).

Proposition 4.2. *Assume that Hypothesis 4.1 and Equation (4.1.2) hold, and that $m \cdot d$ is odd, where d is as in (4.1.1) and m is as in (4.1.2). Then the natural map $\text{Def}(K_2(A), \mathcal{E}(\mathcal{L})) \rightarrow \text{Def}(K_2(A), \det \mathcal{E}(\mathcal{L}))$ is surjective.*

The ingredients in the proof of Proposition 4.2 are the following. First one proves that $\mathcal{E}(\mathcal{L})$ is slope stable with respect to certain polarizations. By Hypothesis 4.1 we have a Lagrangian fibration $\pi_A: K_2(A) \rightarrow \mathbb{P}^2$ (see Subsection 4.4) and hence, by the results proved in [O'G22], in order to prove slope stability of $\mathcal{E}(\mathcal{L})$ for polarizations close to $\pi_A^* \mathcal{O}_{\mathbb{P}^2}(1)$ it suffices to show the restriction to a general fiber of π_A is slope stable: this is done in Subsection 4.4. Secondly, by applying Verbitsky's theory of (projectively) hyperholomorphic vector bundles one proves that the projectivization $\mathbb{P}(\mathcal{E}(\mathcal{L}))$ extends to all small deformations of $K_2(A)$ as a fibration locally trivial in the analytic topology. For this step to work one needs to know that $\mathcal{E}(\mathcal{L})$ is slope stable with respect to Kähler classes belonging to an open (non empty) subset of the Kähler cone. Lastly a simple Brauer class computation gives Proposition 4.2. The last two steps are carried out in Subsection 4.5. In Subsection 4.2 we discuss the wall and chamber decomposition of the Kähler cone associated to a modular sheaf, and in Subsection 4.3 we recall results on stability of modular sheaves on lagrangian fibrations.

4.2. Variation of stability with respect to Kähler classes. Let X be a HK and let \mathcal{F} be a modular sheaf on X . There is a decomposition of the Kähler cone $\mathcal{K}(X) \subset H_{\mathbb{R}}^{1,1}(X)$ into open chambers with the property that if Kähler classes ω_0, ω_1 belong to the same chamber then \mathcal{F} is ω_0 slope-stable if and only if it is ω_1 slope-stable. Moreover the decomposition depends only on the rank and discriminant of \mathcal{F} . This is the analogue of the decomposition of the ample cone of a projective HK described in [O’G22]. In order to describe the relevant walls and chambers we need a couple of definitions.

Definition 4.3. Let a be a positive real number. An a -wall of $\mathcal{K}(X)$ is the intersection $\lambda^\perp \cap \mathcal{K}(X)$, where $\lambda \in H_{\mathbb{Z}}^{1,1}(X)$, $-a \leq q_X(\lambda) < 0$, and orthogonality is with respect to the BBF quadratic form q_X .

As is well-known, the set of a -walls is locally finite, in particular the union of all the a -walls is closed in $\mathcal{K}(X)$.

Definition 4.4. An *open a -chamber* of $\mathcal{K}(X)$ is a connected component of the complement of the union of all the a -walls of $\mathcal{K}(X)$.

Definition 4.5. Let X be a HK manifold, and let \mathcal{F} be a modular torsion free sheaf on X . Then

$$a(\mathcal{F}) := \frac{r(\mathcal{F})^2 \cdot d(\mathcal{F})}{4c_X}, \quad (4.2.1)$$

where $d(\mathcal{F})$ is as in Definition 1.1.1.

Example 4.6. Let \mathcal{E} be a modular sheaf as in Theorem 3.2 (i.e. corresponding to $y = x$ or $y = x + 1$). Then $a(\mathcal{E}) = 72$. In fact $r(\mathcal{E}) = 4$, the Fujiki constant $c_{K_2(A)}$ equals 3 by (2.2.4), and $d(\mathcal{E}) = 54$ by (6.1.6).

Proposition 4.7. *Let X be a HK variety of dimension $2n$ and let \mathcal{F} be a torsion free modular sheaf on X . Then the following hold:*

- (1) *Suppose that ω is a Kähler class on X which belongs to an open $a(\mathcal{F})$ -chamber. If \mathcal{F} is strictly ω slope-semistable there exists an exact sequence of torsion free non zero sheaves*

$$0 \longrightarrow \mathcal{E} \longrightarrow \mathcal{F} \longrightarrow \mathcal{G} \longrightarrow 0 \quad (4.2.2)$$

such that $r(\mathcal{F})c_1(\mathcal{E}) - r(\mathcal{E})c_1(\mathcal{F}) = 0$.

- (2) *Suppose that ω_0, ω_1 are Kähler classes on X belonging to the same open $a(\mathcal{F})$ -chamber. Then \mathcal{F} is ω_0 slope-stable if and only if it is ω_1 slope-stable.*

Proof. Proposition 3.4 in [O’G22] is the analogous result that one gets upon replacing the Kähler cone by the ample cone. We show how to adapt that proof in the present context. The proof is based on two ingredients. The first ingredient is Bogomolov’s inequality, which holds for slope semistable sheaves with respect to a Kähler class by the Donaldson-Uhlenbeck-Yau Theorem. The other ingredient one needs is the following statement: “if a torsion free sheaf \mathcal{F} is ω_0 slope-stable and not ω_1 slope-stable then there exists $t \in [0, 1]$ such that \mathcal{F} is strictly $t\omega_0 + (1-t)\omega_1$ slope-semistable, i.e. slope semistable but not slope stable”. This is Lemma 6.2 in [GT17] (the proof is in [GT]). \square

4.3. Modular sheaves on Lagrangian fibrations. Let $\pi: M \rightarrow \mathbb{P}^n$ be a Lagrangian fibration of a HK manifold of dimension $2n$. If $M_t = \pi^{-1}(t)$ is a smooth fiber the image of the restriction map $H^2(M; \mathbb{R}) \rightarrow H^2(M_t; \mathbb{R})$ is of dimension 1 and is generated by the class of an ample divisor class θ_t on Y_t , see [Wie16]. *Stability of a sheaf on M_t means stability with respect to θ_t .*

Remark 4.8. In the case relevant to us, i.e. M a 4 dimensional HK of Kummer type, the ample divisor class θ_t has elementary divisors $(1, 3)$, see [Wie18].

Let h be an ample divisor on M and let \mathcal{F} be a modular torsion free sheaf on M . There is a strong link between h stability of \mathcal{F} and stability of the restriction of \mathcal{F} to a general Lagrangian fiber if h satisfies a suitable numerical condition. In order to give a precise result, we need a definition.

Definition 4.9. Let $\pi: M \rightarrow \mathbb{P}^n$ be a Lagrangian fibration of a HK manifold of dimension $2n$. If $a > 0$ we say that a polarization h of M is *a-suitable with respect to π* if the following holds: letting

$$f := \pi^* c_1(\mathcal{O}_{\mathbb{P}^n}(1)), \quad (4.3.1)$$

then, given any $\lambda \in H_{\mathbb{Z}}^{1,1}(M)$ such that $-a \leq q_M(\lambda) < 0$ the values of $q_M(\lambda, h)$ and $q_M(\lambda, f)$ have the same sign or are both zero. (Whenever there is no ambiguity regarding the fibration we simply write that h is *a-suitable*.)

Proposition 4.10 (Prop. 3.6 in [O'G22]). *Let $\pi: M \rightarrow \mathbb{P}^n$ be a Lagrangian fibration of a HK variety of dimension $2n$. Let \mathcal{F} be a torsion free modular sheaf on M such that $\text{sing } \mathcal{F}$ does not dominate \mathbb{P}^n . Let h be an ample divisor class on M which is $a(\mathcal{F})$ -suitable. Then the following hold:*

- (i) *If the restriction of \mathcal{F} to a generic fiber of π is slope-stable, then \mathcal{F} is h slope-stable.*
- (ii) *If \mathcal{F} is h slope-stable then the restriction of \mathcal{F} to the generic fiber of π is slope-semistable.*

The next result guarantees the existence of a -suitable polarizations.

Lemma 4.11 (R. Friedman). *Let $\pi: M \rightarrow \mathbb{P}^n$ be a Lagrangian fibration of a HK manifold of dimension $2n$. Let h_0 be a polarization of M and let $a > 0$. If*

$$t > \frac{1}{2}q(h_0, f)a$$

then $h_0 + tf$ is a -suitable.

Proof. One gets a proof upon replacing, in the proof of Lemma (2.3) in [Fri89], the intersection form on the elliptic surface with the BBF form on the Lagrangian fibration (note: use also that $q(h_0, f) > 0$). \square

Lastly we recall from [O'G22] the link between modular sheaves on Lagrangian HK's and semi homogeneous vector bundles on abelian varieties.

Proposition 4.12 (Lemma 2.5+Prop. 2.7 in [O'G22]). *Let $\pi: X \rightarrow \mathbb{P}^n$ be a Lagrangian fibration of a HK manifold of dimension $2n$, and let \mathcal{F} be a modular torsion free sheaf on X . Suppose that $t \in \mathbb{P}^n$ is a regular value of π and that \mathcal{F} is locally-free in a neighborhood of X_t . Then*

$$\int_{X_t} \Delta(\mathcal{F}_t) \smile \theta_t^{n-2} = 0. \quad (4.3.2)$$

If moreover \mathcal{F}_t is slope-stable then it is a (simple) semi homogeneous vector bundle.

4.4. Stability of modular basic vector bundles. In the present subsection we show that, under the assumptions of Proposition 4.2, the modular vector bundles $\mathcal{E}(\mathcal{L})$ are slope stable with respect to certain polarizations. In order to state precisely hypotheses on the polarization which guarantee stability, we go through a few preliminaries.

Assume that Hypothesis 4.1 holds, and that we have chosen a non zero $\epsilon \in C_B[2]$. We recall that $A := B/\langle \epsilon \rangle$ and that $f: B \rightarrow A$ is the quotient map (see

Subsection 4.1). Let $E := B/C_B$ and let $B \xrightarrow{\varphi_B} E$ be the quotient map. Then φ_B descends to a homomorphism

$$A \xrightarrow{\varphi_A} E \quad (4.4.1)$$

with kernel the quotient $C_A := C_B/\langle \epsilon \rangle$. We let $\gamma_A \in H_{\mathbb{Z}}^{1,1}(A)$ be the Poincaré dual of C_A . Since the restriction of f to C_B defines the degree-2 quotient map $C_B \rightarrow C_A$, we have the equalities

$$f_*(\gamma_B) = 2\gamma_A, \quad f^*(\gamma_A) = \gamma_B. \quad (4.4.2)$$

The homomorphisms φ_B and φ_A induce Lagrangian fibrations

$$\begin{array}{ccc} K_2(B) & \xrightarrow{\pi_B} & |\mathcal{O}_E(3(0_E))| \\ [Z] & \mapsto & \varphi_B(|Z|) \end{array} \quad \begin{array}{ccc} K_2(A) & \xrightarrow{\pi_A} & |\mathcal{O}_E(3(0_E))| \\ [Z] & \mapsto & \varphi_A(|Z|) \end{array}$$

where $|Z|$ is the cycle associated to the scheme Z . The commutative diagram in (3.1.3) extends to the following commutative diagram:

$$\begin{array}{ccc} & X & \\ \nu \swarrow & & \searrow \tilde{\rho} \\ K_2(B) & \overset{\rho}{\dashrightarrow} & K_2(A) \\ \pi_B \searrow & & \swarrow \pi_A \\ & |\mathcal{O}_E(3(0_E))| & \end{array} \quad (4.4.3)$$

Let \mathcal{L} be the line bundle on X such that the equality in (4.1.2) holds. Below is our stability result.

Proposition 4.13. *Assume that Hypothesis 4.1 and Equation (4.1.2) hold, and that $m \cdot d$ is odd, where d is as in (4.1.1) and m is as in (4.1.2). Then $\mathcal{E}(\mathcal{L})$ is slope stable for any polarization of $K_2(A)$ which is 72-suitable with respect to π_A .*

The key step in the proof of Proposition 4.13 is provided by the following result.

Proposition 4.14. *Keep hypotheses as in Proposition 4.13. The restriction of $\mathcal{E}(\mathcal{L})$ to any smooth fiber of the Lagrangian fibration π_A is slope stable.*

Proof of Proposition 4.13 assuming Proposition 4.14. We have $a(\mathcal{E}(\mathcal{L})) = 72$ by Example 4.6. Hence Proposition 4.13 follows from Propositions 4.14 and 4.10. \square

Before proving Proposition 4.14 we go through a series of preliminary results.

Proposition 4.15. *Suppose that Hypothesis 4.1 holds and that d is odd, where d is as in (4.1.1). Let $\omega_A := f_*(\omega_B)$. Then $\mathbb{Z}\omega_A + \mathbb{Z}\gamma_A$ is a saturated subgroup of $H_{\mathbb{Z}}^{1,1}(A)$, and*

$$\omega_A \cdot \gamma_A = d, \quad \omega_A \cdot \omega_A = 4a. \quad (4.4.4)$$

In particular ω_A has elementary divisors $(1, 2a)$.

Proof. Suppose that $\mathbb{Z}\omega_A + \mathbb{Z}\gamma_A$ is not a saturated subgroup of $H_{\mathbb{Z}}^{1,1}(A)$. By hypothesis $\mathbb{Z}\omega_B + \mathbb{Z}\gamma_B$ is a saturated subgroup of $H_{\mathbb{Z}}^{1,1}(B)$; since $f^*(\omega_A) = 2\omega_B$ and $f^*(\gamma_A) = \gamma_B$, it follows that $\mathbb{Z}\omega_A + \mathbb{Z}\gamma_A$ is of index 2 in its saturation. This is a contradiction because the discriminant of the intersection form on $\mathbb{Z}\omega_A + \mathbb{Z}\gamma_A$ is equal to $-d^2$ which is odd. The first equality in (4.4.4) follows from

$$2 \int_B \omega_B \cup \gamma_B = \int_B f^*(\omega_A) \cup f^*(\gamma_A) = 2 \int_A \omega_A \cup \gamma_A,$$

and the second one follows from a similar computation. Lastly, ω_A has elementary divisors $(1, 2a)$ because it is primitive and $\omega_A \cdot \omega_A = 4a$. \square

We recall the following fact.

Proposition 4.16. *Let $D_0 \in |\mathcal{O}_E(3(0_E))|$. Then $\pi_A^{-1}(D_0)$ is smooth if and only if D_0 is reduced.*

In order to describe the smooth fibers of π_A (or π_B) we introduce the following notation: if Y is a (bona fide) elliptic curve we let

$$Z_n(Y) := \{(y_1, \dots, y_{n+1}) \in Y^{n+1} \mid y_1 + \dots + y_{n+1} = 0\}. \quad (4.4.5)$$

Let $D_0 \in |\mathcal{O}_E(3(0_E))|$ be reduced, given by $D_0 = (x_1) + (x_2) + (x_3)$. Let $C_{B,i} := \varphi_B^{-1}(x_i)$, $C_{A,i} := \varphi_A^{-1}(x_i)$, where φ_B and φ_A are as in Hypothesis 4.1 and in (4.4.1) respectively. For $i \in \{1, 2\}$ choose $b_1, b_2 \in B$ such that $T_{b_i}(C_B) = C_{B,i}$, where $T_{b_i}: B \rightarrow B$ is the translation by b_i . Let $a_i := f(b_i)$, so that $T_{a_i}(C_A) = C_{A,i}$. We have isomorphisms

$$\begin{aligned} Z_2(C_B) & \xrightarrow{\sim} \pi_B^{-1}((x_1) + (x_2) + (x_3)) \\ (y_1) + (y_2) + (y_3) & \mapsto (y_1 + b_1) + (y_2 + b_2) + (y_3 - b_1 - b_2) \end{aligned} \quad (4.4.6)$$

and

$$\begin{aligned} Z_2(C_A) & \xrightarrow{\sim} \pi_A^{-1}((x_1) + (x_2) + (x_3)) \\ (y_1) + (y_2) + (y_3) & \mapsto (y_1 + a_1) + (y_2 + a_2) + (y_3 - a_1 - a_2) \end{aligned} \quad (4.4.7)$$

(Note: since elements of $\pi_A^{-1}((x_1) + (x_2) + (x_3))$ are reduced subschemes, we identify them with the corresponding 0-cycles.)

Proposition 4.17. *Keep hypotheses as in Proposition 4.13. Then the restriction of $\mathcal{E}(\mathcal{L})$ to a smooth fiber of π is a simple semi-homogeneous vector bundle.*

Proof. Let $\pi_A^{-1}(D_0)$ be a smooth fiber of π_A . Then $\pi_A^{-1}(D_0)$ does not meet $\Delta(A)$, and hence the last equation in (3.4.4) gives that $\tilde{\rho}^{-1}(\pi_A^{-1}(D_0))$ does not meet the exceptional divisor of $\nu: X \rightarrow K_2(B)$ (watch out for the notational similarity between the exceptional divisor $D \subset X$ and $D_0 \in |\mathcal{O}_E(3(0_E))|$). It follows that $\tilde{\rho}^{-1}(\pi_A^{-1}(D_0))$ is equal to $\pi_B^{-1}(D_0)$. Moreover the étale map

$$\pi_B^{-1}(D_0) = \tilde{\rho}^{-1}(\pi_A^{-1}(D_0)) \longrightarrow \pi_A^{-1}(D_0)$$

is identified, given the isomorphisms in (4.4.6) and (4.4.7), with the étale map

$$\begin{aligned} Z_2(C_B) & \xrightarrow{\psi^2} Z_2(C_A) \\ (z_1, \dots, z_3) & \mapsto (f(z_1), \dots, f(z_3)) \end{aligned} \quad (4.4.8)$$

We claim that the thesis of the proposition follows from Proposition A.2; in fact $\deg \psi^2 = 4$ and the degree d_0 of Proposition A.2 is equal to md , which is odd by hypothesis, and hence $\deg \psi^2$ is coprime to $3 \cdot d_0$. \square

Proof of Proposition 4.14. Let $\mathcal{E} := \mathcal{E}(\mathcal{L})$. Let S be a smooth fiber of π_A , i.e. $S = \pi^{-1}(D_0)$ where $D_0 \in |\mathcal{O}_E(3(0_E))|$ is reduced. Let θ be the polarization of S induced by π_A (see the discussion at the beginning of Subsection 4.3). We claim that $\mathcal{E}|_S$ is θ slope stable because the hypotheses of Corollary B.2 are satisfied by (S, θ) and $\mathcal{E}|_S$. In fact by Proposition 4.17 the restriction $\mathcal{E}|_S$ is a simple semi-homogeneous vector bundle, and hence it is Gieseker stable with respect to an arbitrary polarization by Proposition 6.16 in [Muk78]. In particular $\mathcal{E}|_S$ is θ slope semistable. Moreover, since \mathcal{E} is a modular vector bundle we have $\Delta(\mathcal{E}|_S) = 0$ by Proposition 4.12. Lastly, since $r(\mathcal{E}) = 4$, it remains to show that $c_1(\mathcal{E}|_S) = 2b_0\theta$ with b_0 odd. By (3.4.5) we have $c_1(\mathcal{E}|_S) = (2\mu_A(m\omega_A) - \delta_A)|_S$. Since S does not intersect $\Delta(A)$ (see Proposition 4.16), $c_1(\mathcal{E}|_S) = 2\mu_A(m\omega_A)|_S$. We identify S with $Z_2(C_A)$ via the isomorphism in (4.4.8), and we let $p_i: Z_2(C_A) \rightarrow C_A$ be the i -th projection for $i \in \{1, 2, 3\}$. By the first equation in (4.4.4) we get that

$$c_1(\mathcal{E}|_S) = 2\mu_A(m\omega_A)|_S = 2md(p_1^*(\eta_{C_A}) + p_2^*(\eta_{C_A}) + p_3^*(\eta_{C_A})), \quad (4.4.9)$$

where $\eta_{C_A} \in H^2(C_A; \mathbb{Z})$ is the fundamental class. Since θ is a polarization of type $(1, 3)$ we get that $c_1(\mathcal{E}|_S) = 2md\theta$. Since md is odd, this finishes the proof that the hypotheses of Corollary B.2 are satisfied by (S, θ) and $\mathcal{E}|_S$. \square

4.5. Sideways deformations of modular basic vector bundles. In this subsection we prove Proposition 4.2.

Given the stability result of Proposition 4.13, the key input is Verbitsky's theory of hyperholomorphic bundles. Let $\text{Def}(K_2(A), \mathbb{P}(\mathcal{E}(\mathcal{L})))$ be the deformation space parametrizing couple (Y, \mathbf{P}) , where Y is a deformation of $K_2(A)$ and \mathbf{P} is a \mathbb{P}^3 bundle locally trivial in the classical topology (but not necessarily in the Zariski topology) which is a deformation of $\mathbb{P}(\mathcal{E}(\mathcal{L}))$.

Proposition 4.18. *Let hypotheses and notation be as in Proposition 4.2, and let $\mathbb{P}(\mathcal{E}(\mathcal{L}))$ be the projectivization of the vector bundle $\mathcal{E}(\mathcal{L})$ on $K_2(A)$. Then the natural map*

$$\text{Def}(K_2(A), \mathbb{P}(\mathcal{E}(\mathcal{L}))) \longrightarrow \text{Def}(K_2(A)) \quad (4.5.1)$$

is surjective.

Proof. Let $\mathcal{E} = \mathcal{E}(\mathcal{L})$. By Proposition 4.14 there exists a polarization h of $K_2(A)$ such that \mathcal{E} is h slope stable. By Proposition 4.7 there exists an open subcone $\mathcal{U} \subset \mathcal{K}_{K_2(A)}$ of the Kähler cone containing h with the property that \mathcal{E} is ω slope stable for all $\omega \in \mathcal{U}$.

Let $\omega \in \mathcal{U}$. We claim that $\mathbb{P}(\mathcal{E})$ extends to a holomorphic projective bundle on every deformation of $K_2(A)$ parametrized by the twistor line $T(K_2(A), \omega)$ associated to ω . In fact, by Donaldson-Uhlenbeck-Yau there is an ω Hermite-Einstein connection ∇ on the C^∞ vector bundle \mathbf{E} associated to \mathcal{E} because \mathcal{E} is ω slope stable. The connection on $\det \mathbf{E}$ defined by ∇ is also Hermite-Einstein (HE), call it $\det \nabla$. Let $\{V_\alpha\}$ be an open covering of $K_2(A)$ such that the restriction of $\det \mathbf{E}$ to each V_α has a 4-th root L_α . Let D_α be the connection on L_α^{-1} defined by the restriction of $\det \nabla$ to V_α (it is well defined up to a choice of 4-th root of 1 on connected components of V_α). On $\mathbf{E}|_{V_\alpha} \otimes L_\alpha^{-1}$ the connection $\nabla|_{V_\alpha} \otimes D_\alpha$ is HE (for the restriction of ω to V_α); let F_α be its curvature.

By Theorem 11.1 in [Ver96b] (together with the third equality in (3.1.5)), the connection ∇ is projectively hyperholomorphic, i.e. F_α is of type $(1, 1)$ for all complex structures J parametrized by the twistor line $T(K_2(A), \omega)$. Hence we have local holomorphic frames of $\mathbf{E}|_{V_\alpha} \otimes L_\alpha^{-1}$ varying holomorphically with J , giving holomorphic transition functions matching up to a multiple of the identity on triple intersections (there is no global 4-th root of $\det \mathbf{E}$), i.e. a holomorphic family of holomorphic projective bundles, with the bundle over $K_2(A)$ (with the "initial" nice projective complex structure) equal to $\mathbb{P}(\mathcal{E})$.

The image of the map in (4.5.1) is a complex analytic subset of $H^{1,1}(K_2(A)) = H_{\mathbb{R}}^{1,1}(K_2(A)) \otimes_{\mathbb{R}} \mathbb{C}$, and by what we have just proved it contains the intersection of the open cone $\mathcal{U} \subset H_{\mathbb{R}}^{1,1}(K_2(A))$ and a neighborhood of 0 in $H_{\mathbb{R}}^{1,1}(K_2(A))$; it follows that it contains a neighborhood of 0 in $H^{1,1}(K_2(A))$. \square

Proof of Proposition 4.2. We set $\mathcal{E}_0 := \mathcal{E}(\mathcal{L})$. Let $\mathcal{Y} \rightarrow U$ be a (sufficiently small) representative of the deformation space $\text{Def}(K_2(A), \det \mathcal{E}_0)$. Thus $\mathcal{Y} \rightarrow U$ is a proper map of complex manifolds, there is a point $0 \in U$ such that $Y_0 \cong K_2(A)$ (for $t \in U$ we let Y_t be the fiber of $\mathcal{Y} \rightarrow U$ over t), and there is a line bundle \mathcal{D} on \mathcal{Y} such that $\mathcal{D}_0 \cong \det \mathcal{E}_0$. By Proposition 4.18 the \mathbb{P}^3 bundle $\mathbb{P}(\mathcal{E}_0)$ deforms to a projective bundle \mathbf{P}_t on Y_t for every $t \in U$. We must prove that \mathbf{P}_t is the projectivization of a vector bundle. Let $g: \mathbb{P}(\mathcal{E}_0) \rightarrow Y_0$ be the projection, and let $L_0 := \mathcal{O}_{\mathbb{P}(\mathcal{E}_0)}(-1)$ be the tautological sub line-bundle of $g^*(\mathcal{E}_0)$. A straightforward

computation gives that

$$4c_1(L_0) = c_1(K_{\mathbb{P}_0/Y_0} + g^*(\mathcal{D}_0)). \quad (4.5.2)$$

It follows that for every $t \in U$ there exists $\alpha_t \in H^2(\mathbb{P}_t; \mathbb{Z})$ such that $4\alpha_t = c_1(K_{\mathbb{P}_t/Y_t} + g^*(\mathcal{D}_t))$. Since α_t is of type $(1, 1)$ it is the first Chern class of a holomorphic line-bundle L_t . Since the restriction of L_t to the \mathbb{P}^3 -fibers of $\mathbb{P}_t \rightarrow Y_t$ is $\mathcal{O}(-1)$, it follows that \mathbb{P}_t is the projectivization of a vector bundle. \square

5. RIGIDITY AND ACTION OF THE AUTOMORPHISM GROUP

5.1. Main results. The first main result of the present section is the following.

Proposition 5.1. *Assume that Hypothesis 4.1 and Equation (4.1.2) hold, and that $m \cdot d$ is odd, where d is as in (4.1.1) and m is as in (4.1.2). Then*

$$H^1(K_2(A), \text{End}^0(\mathcal{E}(\mathcal{L}))) = 0.$$

Notice that the hypotheses of the above proposition are the same hypotheses of Proposition 4.2. Hence, if we assume that the above hypotheses hold, the natural map $\text{Def}(K_2(A), \mathcal{E}(\mathcal{L})) \rightarrow \text{Def}(K_2(A), \det \mathcal{E}(\mathcal{L}))$ is an isomorphism, and as a consequence the following holds. Let $\mathcal{M} \rightarrow U$ be a sufficiently small representative of $\text{Def}(K_2(A), \det \mathcal{E}(\mathcal{L}))$, with M_t being the fiber over $t \in U$ and $M_0 = K_2(A)$ (thus $0 \in U$ and the germ $(U, 0)$ is identified with $\text{Def}(K_2(A), \det \mathcal{E}(\mathcal{L}))$); then, letting $\mathcal{E}_0 := \mathcal{E}(\mathcal{L})$, for every $t \in U$ there exists one and only one (up to isomorphism) vector bundle on M_t which is a deformation of \mathcal{E}_0 (we denote it \mathcal{E}_t). The second main result of the present section is about the action of $\text{Aut}^0(M_t)$ on \mathcal{E}_t .

Proposition 5.2. *Keep notation as above. Assume that Hypothesis 4.1 and Equation (4.1.2) hold, and that $m \cdot d$ is odd, where d is as in (4.1.1) and m is as in (4.1.2). If $t_0 \in U$ and $\varphi_{t_0} \in \text{Aut}^0(M_{t_0})$ then the vector bundle $\varphi_{t_0}^*(\mathcal{E}_{t_0})$ is isomorphic to \mathcal{E}_{t_0} .*

The proofs of Proposition 5.1 and Proposition 5.2 are in Subsection 5.5. Actually Proposition 5.2 follows from Proposition 5.1 and Proposition 3.18, so the key result is Proposition 5.1. The latter proposition follows from the following result (i.e. Proposition 5.13): if $D_0 \in |\mathcal{O}_E(3(0))|$ is not a flex divisor, then the restriction of $\mathcal{E}(\mathcal{L})$ to the scheme theoretic fiber $\pi_A^{-1}(D_0)$ is simple. Since we already know that this holds if $\pi_A^{-1}(D_0)$ is smooth, i.e. D_0 is reduced (Proposition 4.17), the greater part of this section is devoted to proving that the analogue holds if $D_0 = 2x_0 + y_0$ where $x_0, y_0 \in E$ are distinct.

5.2. The general singular Lagrangian fiber. Let $E^\vee \subset |\mathcal{O}_E(3(0_E))|$ be the sextic curve parametrizing non reduced divisors, i.e. the dual of the smooth cubic $E \subset |\mathcal{O}_E(3(0_E))|^\vee$. We have $\pi_A(\Delta(A)) = E^\vee$, but there is another irreducible component of $\pi_A^{-1}(\Delta(A))$.

Definition 5.3. Let $V(B) \subset K_2(B)$ be the closure of the set parametrizing reduced subschemes Z such that $\pi_B(Z)$ has length smaller than 3, and define similarly $V(A) \subset K_2(A)$ (by definition $V(B), V(A)$ are reduced schemes). Let $\mathbf{V}(B) \subset K_2(B)$ be the subscheme with locally principal ideal such that the associated cycle is $2V(B)$, and define similarly $\mathbf{V}(A) \subset K_2(A)$.

Note that we have the equality of closed sets

$$\pi_A^{-1}(E^\vee)_{\text{red}} = V(A) \cup \Delta(A). \quad (5.2.1)$$

Proposition 5.4. *Keep notation and assumptions as above. Then $\pi_A^{-1}(E^\vee) = \mathbf{V}(A) \cup \Delta(A)$.*

Proof. The proof is similar to the proof of Proposition 6.8 in [O'G22]. It suffices to prove that we have the equality of Cartier divisors

$$\pi_A^*(E^\vee) = 2V(A) + \Delta(A). \quad (5.2.2)$$

By (5.2.1) there exist positive integers a, b such that $\pi_A^*(E^\vee) = aV(A) + b\Delta(A)$. Let $X_3(A)$ be the isospectral Hilbert scheme obtained by blowing up the big diagonal in A^3 . By Proposition 3.4.2 in [Hai01] there exists a regular finite map $X_3(A) \rightarrow A^{[3]}$, invariant under the natural action of the permutation group \mathcal{S}_3 , which is the obvious map on the open dense subset of $X_3(A)$ parameterizing 3-tuples of distinct points. Let $X_3^0(A) \subset X_3(A)$ be the preimage of $K_2(A) \subset A^{[3]}$, and let $\alpha: X_3^0(A) \rightarrow K_2(A)$ be the restriction of the map $X_3(A) \rightarrow A^{[3]}$. Restricting to $X_3^0(A)$ the natural map $X_3(A) \rightarrow A^3$ we get a map $X_3^0(A) \rightarrow Z_2(A)$, where $Z_2(A) \subset A^3$ is the kernel of the summation map $A^3 \rightarrow A$. Let $\lambda_A: X_3^0(A) \rightarrow Z_2(E)$ be the composition of $X_3^0(A) \rightarrow Z_2(A)$ and the map $Z_2(A) \rightarrow Z_2(E)$ defined by $\varphi_A: A \rightarrow E$. Lastly, let $\beta: Z_2(E) \rightarrow |\mathcal{O}_E(3(0_E))|$ be the map sending (x_1, x_2, x_3) to $(x_1) + (x_2) + (x_3)$. We have a commutative square

$$\begin{array}{ccc} X_3^0(A) & \xrightarrow{\alpha} & K_2(A) \\ \lambda_A \downarrow & & \downarrow \pi_A \\ Z_2(E) & \xrightarrow{\beta} & |\mathcal{O}_E(3(0_E))|. \end{array} \quad (5.2.3)$$

As is easily checked we have $\lambda_A^*(\beta^*E^\vee) = 2V(A) + 2\Delta(A)$. On the other hand, since $\pi_A^*(E^\vee) = aV(A) + b\Delta(A)$, we have $\alpha^*(\pi_A^*E^\vee) = aV(A) + 2b\Delta(A)$. Since $\lambda_A^* \circ \beta^* = \alpha^* \circ \pi_A^*$, we get that $a = 2$ and $b = 1$. \square

Definition 5.5. If $D_0 \in E^\vee$ we let $V(B)_{D_0} \subset V(B)$ be the fiber over D_0 of the map $V(B) \rightarrow E^\vee$ given by the restriction of π_B , and we define similarly $\mathbf{V}(B)_{D_0} \subset \mathbf{V}(B)$. Ditto for $V(A)_{D_0} \subset V(A)$ and $\mathbf{V}(A)_{D_0} \subset \mathbf{V}(A)$. We let $\Delta(B)_{D_0} \subset \Delta(B)$ be the fiber over D_0 of the map $\Delta(B) \rightarrow E^\vee$ given by the restriction of π_B , and we define similarly $\Delta(A)_{D_0} \subset \Delta(A)$.

Let $D_0 \in E^\vee$. By Proposition 5.4 we have

$$\pi_A^{-1}(D_0) = \mathbf{V}(A)_{D_0} \cup \Delta(A)_{D_0}. \quad (5.2.4)$$

In the present section we are concerned with fibers of π_A over a $D_0 \in E^\vee$ which is not an inflection divisor, i.e. such that

$$D_0 = 2(x_0) + (y_0), \quad 2x_0 + y_0 = 0, \quad x_0 \neq y_0 \in E. \quad (5.2.5)$$

Let $\xi(A)$ be the line bundle on $K_2(A)$ such that

$$c_1(\xi(A)) = \delta(A). \quad (5.2.6)$$

Proposition 5.6. *Suppose that (5.2.5) holds. Then we have an exact sequence*

$$0 \longrightarrow \mathcal{O}_{V(A)_{D_0}} \otimes \left(\xi(A)|_{V(A)_{D_0}} \right) \longrightarrow \mathcal{O}_{\mathbf{V}(A)_{D_0}} \longrightarrow \mathcal{O}_{V(A)_{D_0}} \longrightarrow 0, \quad (5.2.7)$$

where the right hand map is given by restriction.

Proof. By the equality in (5.2.2) we have $-2V(A) + \pi_A^*(\mathcal{O}(6)) \equiv \Delta(A)$ (the degree of E^\vee is 6). Since $K_2(A)$ is simply connected, we get that

$$-V(A) + \pi_A^*(\mathcal{O}(3)) \equiv \xi(A). \quad (5.2.8)$$

Restricting to $\mathbf{V}(A)_{D_0}$ we get that the ideal sheaf of $V(A)_{D_0} \subset \mathbf{V}(A)_{D_0}$ is isomorphic to the restriction of the invertible sheaf $\xi(A)$. \square

5.3. Geometry of $V(B)_{D_0}$, $V(A)_{D_0}$. Assume that $D_0 \in E^\vee$ is not an inflection divisor, as in (5.2.5). Let $C_{B,x_0} := \varphi_B^{-1}(x_0)$ and $C_{B,y_0} := \varphi_B^{-1}(y_0)$. Choose $\tilde{x}_0 \in C_{B,x_0}$ and let $\tilde{y}_0 := -2\tilde{x}_0$. Notice that $\tilde{y}_0 \in C_{B,y_0}$. We have an isomorphism

$$\begin{array}{ccc} C_B^{(2)} & \xrightarrow{\sim} & V(B)_{D_0} \\ (z_1) + (z_2) & \mapsto & (z_1 + \tilde{x}_0) + (z_2 + \tilde{x}_0) + (-z_1 - z_2 + \tilde{y}_0) \end{array} \quad (5.3.1)$$

A word about notation: a subscheme parametrized by a point of $V(B)_{D_0}$ is the disjoint union of a length 2 subscheme of C_{B,\tilde{x}_0} and a reduced point, and for this reason we identify it with the corresponding cycle. In other words, if $z_1 = z_2$, then $(z_1) + (z_2)$ is to be understood as the unique length 2 subscheme of C_{B,\tilde{x}_0} supported at $z_1 = z_2$. Of course we have an analogous identification between $C_A^{(2)}$ and $V(A)_{D_0}$.

Recall that the degree 4 rational map $\rho: K_2(B) \dashrightarrow K_2(A)$ (see (3.1.1)) has indeterminacy locus $V(f)$ (see (3.1.2)). The intersection of $V(f)$ and $V(B)_{D_0}$ is the reduced curve identified by the isomorphism in (5.3.1) with the curve

$$\{(z) + (z + \epsilon) \mid z \in C_B\}.$$

(Recall that $\epsilon \in B[2]$ is the generator of $\ker f$.) Hence the inclusion $V(B)_{D_0} \hookrightarrow K_2(B)$ lifts to an inclusion $V(B)_{D_0} \hookrightarrow X$ (recall that $\nu: X \rightarrow K_2(B)$ is the blow up of $V(f)$). With this understood we have the schematic equality

$$\tilde{\rho}^{-1}(V(A)_{D_0}) = V(B)_{D_0}. \quad (5.3.2)$$

Moreover, once we make the identification in (5.3.1) and the analogous identification between $C_A^{(2)}$ and $V(A)_{D_0}$, the restriction of $\tilde{\rho}$ to $V(B)_{D_0}$ is identified with the degree 4 regular map

$$\begin{array}{ccc} V(B)_{D_0} = C_B^{(2)} & \xrightarrow{(f|_{C_B})^{(2)}} & C_A^{(2)} = V(A)_{D_0} \\ (z_1) + (z_2) & \mapsto & (f(z_1)) + (f(z_2)) \end{array} \quad (5.3.3)$$

Restricting the line bundle \mathcal{L} to $V(B)_{D_0}$ and pulling it back to $C_B^{(2)}$ via the isomorphism in (5.3.1) we get a line bundle \mathbf{L} on $C_B^{(2)}$. By the equality in (5.3.2) we have the isomorphism (we make the identification in (5.3.1))

$$\mathcal{E}(\mathcal{L})|_{V(A)_{D_0}} \cong (f|_{C_B})_*^{(2)}(\mathbf{L}). \quad (5.3.4)$$

In order to analyze the vector bundle on the right hand side in (5.3.4), we recall that we have \mathbb{P}^1 fibrations

$$\begin{array}{ccc} \mathbb{P}^1 & \longrightarrow & C_B^{(2)} = V(B)_{D_0}, & \mathbb{P}^1 & \longrightarrow & C_A^{(2)} = V(A)_{D_0}, \\ & & \downarrow g_B & & & \downarrow g_A \\ & & C_B & & & C_A \end{array} \quad (5.3.5)$$

where $g_B((z_1) + (z_2)) = z_1 + z_2$ and g_A is defined similarly. For use later on, we notice that if $x \in C_B$ we have

$$\deg(\mathbf{L}|_{g_B^{-1}(x)}) = md. \quad (5.3.6)$$

The fibrations in (5.3.5) are the projectivizations of rank 2 vector bundles:

$$V(B)_{D_0} = C_B^{(2)} = \mathbb{P}(\mathcal{F}_B), \quad V(A)_{D_0} = C_A^{(2)} = \mathbb{P}(\mathcal{F}_A). \quad (5.3.7)$$

(of course \mathcal{F}_B is determined only up to tensorization by the pull-back of a line bundle on C_B , and similarly for \mathcal{F}_A). Let $\mathcal{O}_{C_B^{(2)}}(1) = \mathcal{O}_{V(B)_{D_0}}(1)$ be the corresponding line bundle on $C_B^{(2)} = V(B)_{D_0}$ and similarly for $V(A)_{D_0} = C_A^{(2)}$. (We follow the classical definition of projectivization of a vector bundle: the space of global

sections of $\mathcal{O}_{C_B^{(2)}}(1)$ is $H^0(C_B, \mathcal{F}_B^\vee)$.) The result below is well-known (see [Ati57], p.451).

Proposition 5.7. *Keeping notation as above, the rank 2 vector bundles \mathcal{F}_B and \mathcal{F}_A are stable (in particular they have odd degrees).*

Let $\Gamma \subset V(A)_{D_0}$ be (the class of) a fiber of g_A , and let $\Sigma := V(A)_{D_0} \cap \Delta(A)_{D_0}$ (note that the intersection $V(A) \cap \Delta(A)$ is generically transverse, and hence Σ is reduced). For the sake of simplicity we omit to record A and D_0 in our notation for Γ and Σ . Then $\{\Gamma, \Sigma\}$ is a basis of $H_{\mathbb{Q}}^{1,1}(V(A)_{D_0})$. The following result is used later on - the proof is left to the reader.

Proposition 5.8. *The ample cone of $V(A)_{D_0}$ is the interior of the convex cone generated by Γ and Σ .*

5.4. Restriction of $\mathcal{E}(\mathcal{L})$ to a general singular Lagrangian fiber.

Proposition 5.9. *Assume that Hypothesis 4.1 and Equation (4.1.2) hold, and that $m \cdot d$ is odd, where d is as in (4.1.1) and m is as in (4.1.2). Suppose that $D_0 \in E^\vee$ is not an inflection divisor. Let $g_A: V(A)_{D_0} \rightarrow C_A$ be the fibration in (5.3.5). Then the restriction of $\mathcal{E}(\mathcal{L})$ to $V(A)_{D_0}$ is isomorphic to $g_A^*(\mathcal{V}) \otimes \mathcal{O}_{V(A)_{D_0}}((md-1)/2)$ where \mathcal{V} is a stable rank 4 vector bundle on C_A .*

Proof. Throughout the proof we identify $V(A)_{D_0}$ with $C_A^{(2)}$. In order to simplify notation we let $\psi := (f|_{C_B})^{(2)}$. Let

$$L' := L \otimes \psi^* \mathcal{O}_{C_A^{(2)}}(-(md-1)/2). \quad (5.4.1)$$

By the push pull formula, it suffices to prove that there exists a stable rank 4 vector bundle \mathcal{V} on C_A such that

$$\psi_*(L') \cong g_A^*(\mathcal{V}). \quad (5.4.2)$$

We factor ψ as the composition of two maps. Let $\epsilon \in C_B[2]$ be the generator of the kernel of $f: B \rightarrow A$. Then $\psi: C_B^{(2)} \rightarrow C_A^{(2)}$ is the quotient map for the action of $\mathbb{Z}/(2)^2$ on $C_B^{(2)}$ defined by

$$(z_1) + (z_2) \mapsto (z_1 + k_1\epsilon) + (z_2 + k_2\epsilon),$$

where $(k_1, k_2) \in \mathbb{Z}/(2)^2$. Let i be the involution of $C_B^{(2)}$ defined by $i((z_1) + (z_2)) = (z_1 + \epsilon) + (z_2 + \epsilon)$, and let $Y := C_B^{(2)} / \langle i \rangle$. We let $p: C_B^{(2)} \rightarrow Y$ be the quotient map. Since i commutes with g_B , the \mathbb{P}^1 fibration g_B induces a \mathbb{P}^1 fibration $g_Y: Y \rightarrow C_B$. The restriction of p to a fiber $g_B^{-1}(x)$ is a map

$$\mathbb{P}^1 \cong g_B^{-1}(x) \xrightarrow{p|_{g_B^{-1}(x)}} g_Y^{-1}(x) \cong \mathbb{P}^1. \quad (5.4.3)$$

of degree 2.

Next, by mapping $[(z_1) + (z_2)]$ to $[(z_1 + \epsilon) + (z_2)]$ we get a well defined involution $j: Y \rightarrow Y$ such that $Y / \langle j \rangle$ is identified with $C_A^{(2)}$. Let $q: Y \rightarrow Y / \langle j \rangle \cong C_A^{(2)}$ be the quotient map. Notice that the fibration $g_Y: Y \rightarrow C_B$ is identified with the pull-back of the fibration g_A via the double cover $C_B \rightarrow C_A$. In other words we have an identification $Y = C_B \times_{C_A} C_A^{(2)}$. Summing up, we have a factorization $\psi = q \circ p$. Hence the isomorphism in (5.4.2) holds if and only if

$$q_*(p_*(L')) \cong g_A^*(\mathcal{V}) \quad (5.4.4)$$

for a stable rank 4 vector bundle \mathcal{V} on C_A . Let us prove that there exists a line bundle λ on C_B such that

$$p_*(L') \cong g_Y^*(\mathcal{F}_B^\vee \otimes \lambda), \quad (5.4.5)$$

where \mathcal{F}_B is a vector bundle on C_B such that the second equality in (5.3.7) holds. Since the map in (5.4.3) (for $x \in C_B$) has degree 2, and since $g_Y: Y \rightarrow C_B$ is identified with the pull-back of g_A via the double cover $C_B \rightarrow C_A$, we have $\psi^* \left(\mathcal{O}_{C_A^{(2)}}(1) \right) \cong \mathcal{O}_{C_B^{(2)}}(2) \otimes g_B^*(\xi)$ for a suitable line bundle ξ on C_B . By the equality in (5.3.6) it follows that the restriction of L' to any fiber $g_B^{-1}(x)$ has degree 1. As is easily checked, if $\alpha: \mathbb{P}^1 \rightarrow \mathbb{P}^1$ is a degree 2 map then $\alpha_*(\mathcal{O}_{\mathbb{P}^1}(1))$ is the trivial rank 2 vector bundle $\mathcal{O}_{\mathbb{P}^1}^2$. Notice that we have an identification $H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(1)) \xrightarrow{\sim} H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}^2)$. From this we get that there exists a vector bundle \mathcal{W} on C_B such that $p_*(L') \cong g_Y^*(\mathcal{W})$, and moreover we get an isomorphism $g_{B,*}(\mathcal{O}_{C_B^{(2)}}(1)) \otimes \lambda \cong \mathcal{W}$ for a suitable line λ on C_B . Since $g_{B,*}(\mathcal{O}_{C_B^{(2)}}(1)) \cong \mathcal{F}_B^\vee$, this proves (5.4.5).

Let $\tau: C_B \rightarrow C_A$ be the double cover map. From the isomorphism in (5.4.5) we get that

$$q_*(p_*(L')) \cong g_A^*(\tau_*(\mathcal{F}_B^\vee \otimes \lambda)). \quad (5.4.6)$$

This proves that (5.4.2) holds with $\mathcal{V} := \tau_*(\mathcal{F}_B^\vee \otimes \lambda)$. It remains to prove that \mathcal{V} is stable. The Grothendieck-Riemann-Roch Theorem and a straightforward computation give that $\deg(\mathcal{V}) = \deg(\mathcal{F}_B^\vee \otimes \lambda)$. Since $\mathcal{F}_B^\vee \otimes \lambda$ has odd degree (see Proposition 5.7), the rank (i.e. 4) and the degree of \mathcal{V} are coprime. Hence in order to prove that \mathcal{V} is stable it suffices to show that \mathcal{V} is semistable. The vector bundle $\mathcal{F}_B^\vee \otimes \lambda$ is stable by Proposition 5.7, hence it is semi-homogeneous. Since the map $\text{Pic}^0(C_A) \rightarrow \text{Pic}^0(C_B)$ defined by pull-back is surjective, it follows that \mathcal{V} is semi-homogeneous as well. By Proposition 6.13 in [Muk78] it follows that \mathcal{V} is semistable, and we are done. \square

Corollary 5.10. *Suppose that Hypothesis 4.1 and Equation (4.1.2) both hold, and that the intersection number d appearing in (4.1.1) is odd. Suppose also $D_0 \in E^\vee$ is not an inflection divisor. Then the restriction of $\mathcal{E}(\mathcal{L})$ to $V(A)_{D_0}$ is slope stable with respect to any polarization.*

Proof. Let \mathcal{V} be the rank 4 stable vector bundle on C_A of Proposition 5.9. It suffices to prove that $g_A^*(\mathcal{V})$ is slope stable with respect to any polarization. First we notice that $g_A^*(\mathcal{V})$ is slope stable with respect to a polarization represented by a section Σ of g_A , because the restriction of $g_A^*(\mathcal{V})$ to Σ is identified with \mathcal{V} , and hence is slope stable. Because of this (existence of polarizations for which $g_A^*(\mathcal{V})$ is slope stable), it suffices to prove that there does not exist a polarization for which $g_A^*(\mathcal{V})$ is strictly slope semistable, i.e. slope semistable but not semistable. We argue by contradiction. Suppose that

$$0 \longrightarrow \mathcal{A} \longrightarrow g_A^*(\mathcal{V}) \longrightarrow \mathcal{B} \longrightarrow 0 \quad (5.4.7)$$

is an h slope destabilizing exact sequence. Since $g_A^*(\mathcal{V})$ is h slope semistable, \mathcal{B} is torsion free, \mathcal{A} is locally free, and both \mathcal{A} and \mathcal{B} are h slope semistable. A Chern class computation gives the equality

$$4(r(\mathcal{B})\Delta(\mathcal{A}) + r(\mathcal{A})\Delta(\mathcal{B})) = r(\mathcal{A}) \cdot r(\mathcal{B})\Delta(g_A^*(\mathcal{V})) + (4c_1(\mathcal{A}) - r(\mathcal{A})c_1(g_A^*(\mathcal{V})))^2. \quad (5.4.8)$$

Since the exact sequence in (5.4.7) is h slope destabilizing (but $g_A^*(\mathcal{V})$ is h slope semistable), we have $(4c_1(\mathcal{A}) - r(\mathcal{A})c_1(g_A^*(\mathcal{V}))) \cdot h = 0$. By the Hodge index Theorem it follows that $(4c_1(\mathcal{A}) - r(\mathcal{A})c_1(g_A^*(\mathcal{V})))^2 \leq 0$, and if equality holds then $(4c_1(\mathcal{A}) - r(\mathcal{A})c_1(g_A^*(\mathcal{V}))) = 0$ (in cohomology). On the other hand $\Delta(g_A^*(\mathcal{V})) = 0$, and $\Delta(\mathcal{A}) \geq 0$, $\Delta(\mathcal{B}) \geq 0$ by Bogomolov's inequality. By the equality in (5.4.8) we get that $(4c_1(\mathcal{A}) - r(\mathcal{A})c_1(g_A^*(\mathcal{V})))^2 = 0$, and hence

$$4c_1(\mathcal{A}) = r(\mathcal{A})c_1(g_A^*(\mathcal{V})).$$

This is a contradiction because $1 \leq r(\mathcal{A}) \leq 3$ and $\deg \mathcal{V}$ is coprime to $r(\mathcal{V}) = 4$ (by stability of \mathcal{V}). \square

Proposition 5.11. *Assume that Hypothesis 4.1 and Equation (4.1.2) hold, and that $m \cdot d$ is odd, where d is as in (4.1.1) and m is as in (4.1.2). Suppose also that $D_0 \in E^\vee$ is not an inflection divisor. Then we have an exact sequence*

$$0 \longrightarrow \mathcal{E}(\mathcal{L}) \otimes \xi(A)|_{V(A)_{D_0}} \xrightarrow{\alpha} \mathcal{E}(\mathcal{L})|_{V(A)_{D_0}} \xrightarrow{\beta} \mathcal{E}(\mathcal{L})|_{V(A)_{D_0}} \longrightarrow 0, \quad (5.4.9)$$

where β is given by restriction. Moreover the restriction $\mathcal{E}(\mathcal{L})|_{V(A)_{D_0}}$ is slope stable for any polarization of $V(A)_{D_0}$.

Proof. This is a straightforward consequence of Proposition 5.6 and Corollary 5.10. \square

Next, we analyze the restriction of $\mathcal{E}(\mathcal{L})$ to $\Delta(A)_{D_0}$ where $D_0 \in E^\vee$ is not an inflection divisor. We recall that $\Delta(A)_{D_0}$ is isomorphic to $C_A \times \mathbb{P}^1$.

Proposition 5.12. *Assume that Hypothesis 4.1 and Equation (4.1.2) hold, and that $m \cdot d$ is odd, where d is as in (4.1.1) and m is as in (4.1.2). Suppose also that $D_0 \in E^\vee$ is not an inflection divisor. Then we have the Harder-Narasimhan filtration (with respect to any polarization)*

$$0 \longrightarrow \mathcal{O}_{C_A}(P) \boxtimes \mathcal{O}_{\mathbb{P}^1}(1) \longrightarrow \mathcal{E}(\mathcal{L})|_{\Delta(A)_{D_0}} \longrightarrow \bigoplus_{k=1}^3 \mathcal{O}_{C_A}(P_k) \boxtimes \mathcal{O}_{\mathbb{P}^1} \longrightarrow 0, \quad (5.4.10)$$

where P_k and P are divisors on C_A of degree $3d$.

Proof. In the proof of Lemma 3.11 we showed that the equation in (3.5.6) holds with $m = 2$, i.e. that we have

$$\tilde{\rho}^* \Delta(A) = \nu^*(\Delta(B)) + 2D. \quad (5.4.11)$$

Let

$$\Xi(A) := \tilde{\rho}(\nu^{-1}(\Delta(B)) \cap D) = \{[Z] \in K_2(A) \mid \mathfrak{h}(Z) = 3(p), p \in A[3]\},$$

(recall that $\mathfrak{h}: K_2(A) \rightarrow A^{(3)}$ is the Hilbert-Chow map), and let $\Delta(A)^0 := \Delta(A) \setminus \Xi(A)$. It follows from the equality in (5.4.11) that we have

$$\mathcal{E}(\mathcal{L})|_{\Delta(A)^0} \cong \mu_{1,*}(\mathcal{L}|_{\nu^*(\Delta(B))})|_{\Delta(A)^0} \oplus \mu_{2,*}(\mathcal{L}|_{2D})|_{\Delta(A)^0}, \quad (5.4.12)$$

where μ_1 and μ_2 are the restrictions of $\tilde{\rho}$ to $\nu^*(\Delta(B))$ and $2D$ respectively. Since $\Delta(A)_{D_0} \subset \Delta(A)^0$, we get that

$$\mathcal{E}(\mathcal{L})|_{\Delta(A)_{D_0}} \cong \mu_{1,*}(\mathcal{L}|_{\nu^*(\Delta(B))})|_{\Delta(A)_{D_0}} \oplus \mu_{2,*}(\mathcal{L}|_{2D})|_{\Delta(A)_{D_0}}. \quad (5.4.13)$$

We proceed to describe the direct summands in the right hand side of (5.4.13). First, we have $\tilde{\rho}^{-1}(\Delta(A)_{D_0}) = \nu^{-1}(\Delta(B)_{D_0})$. Second, since $\Delta(B)_{D_0}$ is disjoint from the center of the blow up $\nu: X \rightarrow K_2(B)$, the map $\nu^{-1}(\Delta(B)_{D_0}) \rightarrow \Delta(B)_{D_0}$ is an isomorphism; for this reason we identify $\nu^{-1}(\Delta(B)_{D_0})$ and $\Delta(B)_{D_0}$. Third, the restriction of μ_1 to $\Delta(B)_{D_0}$ is the étale double cover $C_B \times \mathbb{P}^1 \rightarrow C_A \times \mathbb{P}^1$ defined by the quotient map $\tau: C_B \rightarrow C_A$. It follows that, letting $\sigma_B: C_B \hookrightarrow \Delta(B)_{D_0} = C_B \times \mathbb{P}^1$ be a section, we have

$$\mu_{1,*}(\mathcal{L}|_{\nu^*(\Delta(B))})|_{\Delta(A)_{D_0}} \cong \tau_*(\sigma_B^*(\mathcal{L})) \boxtimes \mathcal{O}_{\mathbb{P}^1}. \quad (5.4.14)$$

A computation gives that $\deg \sigma_B^*(\mathcal{L}) = 6d$. Hence there exists a divisor P_1 on C_A of degree $3d$ such that $\tau^* \mathcal{O}_{C_A}(P_1) \cong \sigma_B^*(\mathcal{L})$. It follows that

$$\tau_*(\sigma_B^*(\mathcal{L})) \cong \mathcal{O}_{C_A}(P_1) \oplus \mathcal{O}_{C_A}(P_2), \quad (5.4.15)$$

where $\deg P_2 = 3d$. (In fact $P_2 \equiv P_1 + \eta$ where η is the divisor class of order 2 determined by the double cover $\tau: C_B \rightarrow C_A$.)

Next we notice that the map $D \rightarrow \Delta(A)$ given by the restriction of $\tilde{\rho}$ is an isomorphism away from $\Xi(A)$ (see Subsection 3.5) hence we have an open inclusion $\iota: \Delta(A)^0 \hookrightarrow D$ that composed with $\tilde{\rho}$ gives the identity. We have an exact sequence

$$0 \longrightarrow \iota^*(\mathcal{L}) \otimes \iota^*(\mathcal{O}_D(-D)) \longrightarrow \mu_{2,*}(\mathcal{L}|_{2D})|_{\Delta(A)^0} \longrightarrow \iota^*(\mathcal{L}) \longrightarrow 0.$$

Restricting the above exact sequence to $\Delta(A)_{D_0}$ we get an exact sequence

$$0 \longrightarrow \mathcal{O}_{C_A}(P') \boxtimes \mathcal{O}_{\mathbb{P}^1}(1) \longrightarrow \mu_{2,*}(\mathcal{L}|_{2D})|_{\Delta(A)_{D_0}} \longrightarrow \mathcal{O}_{C_A}(P') \boxtimes \mathcal{O}_{\mathbb{P}^1} \longrightarrow 0 \quad (5.4.16)$$

where P' is a divisor of degree $3d$. Hence we get that $\mathcal{E}(\mathcal{L})|_{\Delta(A)_{D_0}}$ fits into an exact sequence as in (5.4.10), where $P_3 = P = P'$. It is clear that the exact sequence is the Harder-Narsimhan filtration. \square

5.5. Simplicity of the restriction to a general singular fiber. The key result is the following.

Proposition 5.13. *Assume that Hypothesis 4.1 and Equation (4.1.2) hold, and that $m \cdot d$ is odd, where d is as in (4.1.1) and m is as in (4.1.2). Suppose also that $D_0 \in E^\vee$ is not an inflection divisor. Then the restriction of $\mathcal{E}(\mathcal{L})$ to $\pi_A^{-1}(D_0)$ is a simple sheaf.*

Proof of Proposition 5.13. Let $\mathcal{E} = \mathcal{E}(\mathcal{L})$, and let $\varphi: \mathcal{E} \rightarrow \mathcal{E}$ be an endomorphism. Then φ maps the kernel of α in (5.4.9) to itself (because, as subsheaf of $\mathcal{E}|_{\mathbf{V}(A)_{D_0}}$, it is the kernel of multiplication by the ideal of $V(A)_{D_0}$ in $\mathbf{V}(A)_{D_0}$). Since $\mathcal{E}|_{V(A)_{D_0}}$ is slope stable it is simple, and hence the restriction of φ to $\ker(\alpha)$ is multiplication by a certain $\lambda \in \mathbb{C}$. Thus $\varphi = \lambda \text{Id}_{\mathcal{E}} + \bar{\varphi}$, where $\bar{\varphi}$ annihilates $\ker(\alpha)$. Let us prove that $\bar{\varphi}$ vanishes. By definition the restriction of $\bar{\varphi}$ to $\mathbf{V}(A)_{D_0}$ factors as follows:

$$\mathcal{E}|_{\mathbf{V}(A)_{D_0}} \longrightarrow \mathcal{E}|_{V(A)_{D_0}} \xrightarrow{\psi} \mathcal{E} \otimes \xi(A)|_{V(A)_{D_0}}. \quad (5.5.1)$$

We start by proving that the determinant of ψ vanishes. In fact, by the factorization in (5.5.1), the restriction of $\bar{\varphi}$ to $\Delta(A)_{D_0}$ is a homomorphism

$$\mathcal{E}|_{\Delta(A)_{D_0}} \rightarrow \mathcal{E}|_{\Delta(A)_{D_0}} \otimes \mathcal{O}|_{\Delta(A)_{D_0}}(-\Sigma).$$

The exact sequence in (5.4.10) gives that the restriction of $\bar{\varphi}$ to $\Delta(A)_{D_0}$ has image contained in the sub line bundle $\mathcal{O}_{C_A}(P) \boxtimes \mathcal{O}_{\mathbb{P}^1}(1) \otimes \mathcal{O}|_{\Delta(A)_{D_0}}(-\Sigma)$. In particular the restriction to Σ of such a homomorphism has rank at most 1. It follows that $\det \psi$ vanishes on Σ with order at least 3. Since $\det \psi$ is a section of $\xi(A)|_{V(A)_{D_0}}^{\otimes 4} \cong \mathcal{O}_{V(A)_{D_0}}(2\Sigma)$ we get that the determinant of ψ is identically zero.

By Proposition 5.9, the restriction of \mathcal{E} to $V(A)_{D_0}$ is the tensor product of $g_A^*(\mathcal{V})$ (notation as in Proposition 5.9) and a line bundle. Thus ψ defines a map

$$\psi': g_A^*(\mathcal{V}) \longrightarrow g_A^*(\mathcal{V}) \otimes \xi(A)|_{V(A)_{D_0}} \quad (5.5.2)$$

with (generic) rank 1, 2 or 3. We reach a contradiction in each of the three cases. We have $c_1(\text{Im } \psi') = a\Sigma + b\Gamma$ with a, b rational numbers (see Subsection 5.3). Let P be a polarization of $V(A)_{D_0}$; thus $c_1(P) = s\Sigma + t\Gamma$ where s, t are positive rational numbers.

Assume that the (generic) rank of ψ' is 1. By Corollary 5.10, the vector bundle $g_A^*(\mathcal{V})$ is P slope stable. Hence the surjection $g_A^*(\mathcal{V}) \twoheadrightarrow \text{Im } \psi'$ gives the inequality

$$s \deg \mathcal{V} = \mu_P(g_A^*(\mathcal{V})) < \mu_P(\text{Im } \psi') = 4at + 4bs \quad (5.5.3)$$

and the injection $\text{Im } \psi' \hookrightarrow g_A^*(\mathcal{V}) \otimes \xi(A)|_{V(A)_{D_0}}$ gives the inequality

$$4at + 4bs = \mu_P(\text{Im } \psi') < \mu_P(g_A^*(\mathcal{V}) \otimes \xi(A)|_{V(A)_{D_0}}) = s \deg \mathcal{V} + 2t. \quad (5.5.4)$$

By Proposition 5.8 the coefficient t may be arbitrarily small (and s is positive); it follows that $\deg \mathcal{V} \leq 4b \leq \deg \mathcal{V}$. Hence $\deg \mathcal{V} = 4b$. On the other hand $2b$ is an

integer because $c_1(\xi(A)|_{V(A)_{D_0}}) \cdot c_1(\text{Im } \psi') = \frac{1}{2}\Sigma \cdot (a\Sigma + b\Gamma) = 2b$, and hence we get that \mathcal{V} has even degree. This contradicts the stability of \mathcal{V} .

Next we deal with the remaining two cases, i.e. the (generic) rank of ψ' is $r \in \{2, 3\}$. Then we have inclusions of sheaves on $V(A)_{D_0}$

$$\text{Im}(\psi') \subset \mathcal{F} \subset g_A^*(\mathcal{V}) \otimes \xi(A)|_{V(A)_{D_0}},$$

where \mathcal{F} is the saturation of $\text{Im}(\psi')$, i.e. a sheaf of rank r and with torsion free cokernel. The key observation is that, since the rank of $\text{Im}(\psi)$ on the divisor Σ is at most 1 (this was proved above, when we showed that ψ has vanishing determinant), we have $c_1(\mathcal{F}) = c_1(\text{Im}(\psi')) + m\Sigma + \text{Eff}$ where $m \geq (r-1)$ and Eff is an effective divisor. By stability of $g_A^*(\mathcal{V})$ we get the inequality

$$s \deg \mathcal{V} = \mu_P(g_A^*(\mathcal{V})) < \mu_P(\text{Im } \psi'),$$

and the inequality

$$\mu_P(\text{Im } \psi') + \frac{4(r-1)t}{r} \leq \mu_P(\mathcal{F}) < \mu_P(g_A^*(\mathcal{V}) \otimes \xi(A)|_{V(A)_{D_0}}) = s \deg \mathcal{V} + 2t.$$

Since $r \in \{2, 3\}$, the second equality gives that $\mu_P(\text{Im } \psi') < s \deg \mathcal{V}$, and this contradicts the first inequality. This proves that $\psi = 0$, i.e. that the restriction of $\bar{\varphi}$ to $V(A)_{D_0}$ vanishes.

It follows that the restriction of $\bar{\varphi}$ to $\Delta(A)_{D_0}$ is a homomorphism

$$\mathcal{E}|_{\Delta(A)_{D_0}} \rightarrow \mathcal{E}|_{\Delta(A)_{D_0}} \otimes \mathcal{O}_{\Delta(A)_{D_0}}(-2\Sigma).$$

By Proposition 5.12 such a homomorphism vanishes. This proves that the restriction of $\bar{\varphi}$ to $\Delta(A)_{D_0}$ is zero, and hence $\bar{\varphi} = 0$. \square

5.6. Proof of Propositions 5.1 and 5.2.

Proof of Proposition 5.1. The proof is analogous to the proof of Proposition 6.7 in [O'G22]. Suppose that $D_0 \in |\mathcal{O}_C(3(0_E))|$ is not a flex divisor, and let \mathcal{W}_{D_0} be the restriction of $\text{End}^0 \mathcal{E}(\mathcal{L})$ to $\pi_A^{-1}(D_0)$. By Proposition 4.14 and Proposition 5.13 there are no non zero global sections of \mathcal{W}_{D_0} . Since $\pi_A^{-1}(D_0)$ is a local complete intersection surface with trivial dualizing sheaf, it follows that $H^2(\pi_A^{-1}(D_0), \mathcal{W}_{D_0})$ vanishes as well. On the other hand $\chi(\pi_A^{-1}(D_0), \mathcal{W}_{D_0}) = 0$ because \mathcal{W}_{D_0} is a semi-homogeneous vector bundle if D_0 is reduced (see Proposition 4.17), and hence all the cohomology of \mathcal{W}_{D_0} vanishes. Since the set of flex divisors has codimension 2 in $|\mathcal{O}_C(3(0_E))|$, it follows that $R^q \pi_{A,*} \text{End}^0 \mathcal{E}(\mathcal{L}) = 0$ for $q \in \{0, 1\}$ (see Proposition 2.26 in [Muk87]). By the Leray spectral sequence for π_A we get that $H^1(K_2(A), \text{End}^0(\mathcal{E}(\mathcal{L}))) = 0$. \square

Proof of Proposition 5.2. By Propositions 4.2 and 5.1 we have the isomorphism

$$\text{Def}(K_2(A), \mathcal{E}(\mathcal{L})) \xrightarrow{\sim} \text{Def}(K_2(A), \det \mathcal{E}(\mathcal{L}))$$

given by the natural morphism. Let $\mathcal{M} \rightarrow U$ be a sufficiently small representative of $\text{Def}(K_2(A), \det \mathcal{E}(\mathcal{L}))$, with M_t being the fiber over $t \in U$ and $M_0 = K_2(A)$. Then, letting $\mathcal{E}_0 := \mathcal{E}(\mathcal{L})$, for every $t \in U$ there exists one and only one (up to isomorphism) vector bundle on M_t which is a deformation of \mathcal{E}_0 ; we denote it \mathcal{E}_t . We let \mathbf{E} be a vector bundle on \mathcal{M} such that $\mathbf{E}|_{M_t} \cong \mathcal{E}_t$. Now let $\varphi_{t_0} \in \text{Aut}^0(M_{t_0})$. The family of groups over U with fiber $\text{Aut}^0(M_t)$ over $t \in U$ is trivial (since U is small), hence there exists a unique $\Phi \in \text{Aut}(\mathcal{M})$ which restricts to φ_{t_0} over t_0 . Let φ_0 be the restriction of Φ to $M_0 = K_2(A)$. Since $\mathcal{E}_0 := \mathcal{E}(\mathcal{L})$, we have $\varphi_0^*(\mathcal{E}_0) \cong \mathcal{E}_0$ by Proposition 3.18. Hence the vector bundle $\Phi^*(\mathbf{E})$ over \mathcal{M} restricts to \mathcal{E}_0 over M_0 , and therefore it is isomorphic to \mathbf{E} because of the properties of $\text{Def}(K_2(A), \mathcal{E}(\mathcal{L}))$ discussed above (note that $\text{Pic}(U)$ is trivial because U is small). This proves that $\varphi_{t_0}^*(\mathcal{E}_{t_0}) \cong \mathcal{E}_{t_0}$.

□

6. PROOF OF THEOREM 1.1

6.1. Ampleness of $\det \mathcal{E}(\mathcal{L})$. In the setting of Proposition 4.2, we have $c_1(\mathcal{E}(\mathcal{L})) = 2m\mu_A(\omega_A) - \delta(A)$ by (3.4.5). Below is the main result of the present subsection. (The hypotheses of the proposition below are motivated by Hypothesis 4.1 and Proposition 4.15.)

Proposition 6.1. *Let a be a positive integer. If $d \gg 0$ (the lower bound depends on a) the following holds. Let A be an abelian surface such that $H_{\mathbb{Z}}^{1,1}(A) = \mathbb{Z}\omega_A \oplus \mathbb{Z}\gamma_A$, where ω_A is ample, γ_A is the Poincarè dual of a (bona fide) elliptic curve C_A , and*

$$\omega_a \cdot \omega_a = 4a, \quad \omega_A \cdot \gamma_A = d. \quad (6.1.1)$$

Then for any positive integer m the class $2m\mu_A(\omega_A) - \delta(A)$ is ample on $K_2(A)$.

In order to prove Proposition 6.1, we make explicit the general results of [Yos16] and [KLCM19] in the case of $\text{Kum}_2(A)$.

If M is a HK manifold, let $\mathcal{C}(M) \subset H_{\mathbb{R}}^{1,1}(M)$ be the cone of $x \in H_{\mathbb{R}}^{1,1}(M)$ such that $q_M(x) > 0$, and let $\mathcal{C}^+(M)$ the connected component of $\mathcal{C}(M)$ containing Kähler classes. We recall that the Kähler cone $\mathcal{K}(M) \subset \mathcal{C}^+(M)$ is a connected component (open chamber) of the complement of the wall divisors, given by $w^\perp \cap \mathcal{C}^+(M)$ for certain classes $w \in H_{\mathbb{Z}}^{1,1}(M)$ of negative square, see [?].

Proposition 6.2. *Suppose that $w \in H_{\mathbb{Z}}^{1,1}(K_2(A))$ is primitive and $w^\perp \cap \mathcal{C}^+(K_2(A))$ is a wall divisor. Then $q(w) = -6$ and $\text{div}(w) \in \{2, 3, 6\}$.*

Proof. First we recall that the Mukai pairing $\langle \cdot, \cdot \rangle$ on $H^{2*}(A; \mathbb{Z})$ is defined by

$$\langle (r, \ell, s), (r', \ell', s') \rangle := -rs' - r's + \int_S \ell \cup \ell'.$$

If $v^2 \geq 6$, then the cohomology of the Albanese fiber of a moduli space of stable sheaves on A with assigned primitive Mukai vector $v \in H^{2*}(A; \mathbb{Z})$ is identified with the orthogonal $v^\perp \subset H^{2*}(A; \mathbb{Z})$, see [?] for details. In our case, i.e. $K_2(A)$, the Mukai vector is $v := (1, 0, -3)$.

According to Theorem 2.9 in [KLCM19], a class $w \in v^\perp$ defines a wall divisor (i.e. $w^\perp \cap \mathcal{C}^+(K_2(A))$ is a wall divisor) if $\langle w, w \rangle$ is negative and the saturated sublattice of $H^{2*}(A; \mathbb{Z})$ generated by v and w contains an $s \in H^{2*}(A; \mathbb{Z})$ such that

$$0 \leq \langle s, s \rangle < \langle s, v \rangle \leq \frac{1}{2}(\langle v, v \rangle + \langle s, s \rangle) = 3 + \frac{1}{2}\langle s, s \rangle. \quad (6.1.2)$$

Note that we may assume that s is primitive. Clearly

$$w = \frac{1}{m}(\langle s, v \rangle v - 6s), \quad (6.1.3)$$

for a certain integer m which is a multiple of $\text{gcd}(\langle v, s \rangle, 6)$. Thus we get that

$$\langle w, w \rangle = -\frac{6}{m^2}(\langle s, v \rangle^2 - 6\langle s, s \rangle), \quad \frac{6}{m} \text{ divides } \text{div}(w). \quad (6.1.4)$$

The inequalities in (6.1.2) give that $\langle s, s \rangle < 6$, and hence $\langle s, s \rangle \in \{0, 2, 4\}$ because the Mukai pairing is even, and also that $\langle s, v \rangle$ belongs to a finite list. Explicitly, either $\langle s, s \rangle = 0$ and $\langle s, v \rangle \in \{1, 2, 3\}$, or $\langle s, s \rangle = 2$ and $\langle s, v \rangle \in \{3, 4\}$, or $\langle s, s \rangle = 4$ and $\langle s, v \rangle = 5$. A case by case analysis gives that $m = \text{gcd}(\langle v, s \rangle, 6)$ in all cases, and the proposition follows (recall that $\text{div}(x)$ is a divisor of 6 for every $x \in H^2(K_2(A); \mathbb{Z})$). □

We will use the following elementary result.

Lemma 6.3 (Lemma 4.3 in [O'G22]). *Let (Λ, q) be a non degenerate rank 2 lattice which represents 0, and hence $\text{disc}(\Lambda) = -d_0^2$ where d_0 is a strictly positive integer. Let $\alpha \in \Lambda$ be primitive isotropic, and complete it to a basis $\{\alpha, \beta\}$ such that $q(\beta) \geq 0$. If $\gamma \in \Lambda$ has strictly negative square (i.e. $q(\gamma) < 0$) then*

$$q(\gamma) \leq -\frac{2d_0}{1 + q(\beta)}. \quad (6.1.5)$$

Proof of Proposition 6.2. Since ω_A is ample, the class $\mu_A(\omega_A)$ is in the closure of the ample cone. Suppose that $2m\mu_A(\omega_A) - \delta(A)$ is not ample. By Proposition 6.2 there exist $\beta \in H_{\mathbb{Z}}^{1,1}(A)$, $x \in \mathbb{Z}$ such that

$$q(\beta^2 - x\delta(A)) = -6 \leq \beta^2 - 6x^2 < 0, \quad (6.1.6)$$

and either

$$q(\mu_A(\beta) - x\delta(A), \mu_A(\omega_A)) = q(\mu_A(\beta) - x\delta(A), 2m\mu_A(\omega_A) - \delta(A)) = 0, \quad (6.1.7)$$

or

$$q(\mu_A(\beta) - x\delta(A), \mu_A(\omega_A)) > 0, \quad q(\mu_A(\beta) - x\delta(A), 2m\mu_A(\omega_A) - \delta(A)) \leq 0. \quad (6.1.8)$$

In other words, either both $2m\mu_A(\omega_A)$ and $2m\mu_A(\omega_A) - \delta(A)$ lie on a wall, or there exists a wall separating $2m\mu_A(\omega_A)$ from $2m\mu_A(\omega_A) - \delta(A)$.

Assume that (6.1.7) holds. Then $(\beta, \omega_A) = 0$, and hence $\beta^2 \leq 0$ by Hodge index. The last inequality is strict because if it were an equality then we would have $\beta = 0$, and hence also $x = 0$. By (6.1.6) we get that $-6 \leq \beta^2 < 0$. By Lemma 6.3 this is impossible if $d > 12a + 3$.

Next assume that (6.1.8) holds. We rewrite (6.1.6) as

$$\beta^2 < 6x^2 \leq \beta^2 + 6. \quad (6.1.9)$$

By (6.1.8) we have

$$0 < m(\beta, \omega_A) \leq 3x. \quad (6.1.10)$$

In particular x is positive. Since by Hodge index we have $(\omega_A^2) \cdot (\beta^2) \leq (\beta, \omega_A)^2$, it follows that

$$m^2 \cdot 4a\beta^2 \leq m^2 \cdot (\beta, \omega_A)^2 \leq 9x^2 \leq \frac{3}{2}\beta^2 + 9. \quad (6.1.11)$$

It follows that $\beta^2 \leq 18/5$. Feeding this into the second inequality in (6.1.9) we get that $x = 1$. This together with (6.1.10) gives that $(\beta, \omega_A) \in \{1, 2, 3\}$. (Notice that if $m > 3$ we are done.) The class $(4a\beta - (\beta, \omega_A)\omega_A)$ is orthogonal to ω_A , hence

$$-12a \leq (4a\beta - (\beta, \omega_A)\omega_A)^2 \leq 0, \quad (6.1.12)$$

where the first inequality follows from $0 \leq \beta^2$ (see (6.1.9)) and $(\beta, \omega_A) \in \{1, 2, 3\}$. By Lemma 6.3, if $d > 24a^2 + 6a$ this forces the square to vanish, and hence β to be a multiple of ω_A . This is absurd by (6.1.10). \square

Remark 6.4. Arguing geometrically one can give a lower bound on d in Proposition 6.1 which is much stronger than the bound one gets by following the proof, at least if $a \neq 2$. In fact one can show that if $a = 1$ then $d \geq 2$ suffices for ampleness of $2m\mu_A(\omega_A) - \delta(A)$, and that if $a \geq 3$ then $d \geq 4$ suffices. We sketch the proof. Let L be a line bundle on A such that $c_1(L) = 2m\omega_A$. Then L is ample on A and it is globally generated. We claim that the map $A \rightarrow |L|^\vee \cong \mathbb{P}^{8am^2-1}$ is an embedding and that the image is cut out by quadrics. If $a = 1$ this follows from the results in [Bar87], and if $a \geq 3$ the statement holds because \mathcal{L} satisfies property (N_1) by the main result in [KL19] or [Ito18]. Thus we have $A \subset |L|^\vee \cong \mathbb{P}^{8am^2-1}$. Since A

is cut out by quadrics and it contains no lines every length 3 subscheme of A spans a plane in \mathbb{P}^{8am^2-1} and hence we have a regular map

$$\begin{array}{ccc} A^{[3]} & \xrightarrow{\varphi} & \mathrm{Gr}(3, \mathbb{C}^{8am^2}) \\ [Z] & \mapsto & \langle Z \rangle \end{array} \quad (6.1.13)$$

Next, one proves that φ has finite fibers, and hence the pullback $\varphi^*(\mathcal{P})$ of the Plücker (ample) line-bundle on $\mathrm{Gr}(3, \mathbb{C}^{8am^2})$ is ample on $A^{[3]}$. At this point we are done because the restriction of $\varphi^*(\mathcal{P})$ to $K_2(A)$ is $2m\mu_A(\omega_A) - \delta(A)$.

6.2. Proof of Theorem 1.1. In the first case, i.e. $e \equiv -6 \pmod{16}$, we write $e = 16a - 6$ where a is a positive integer, in the second case, i.e. $e \equiv -6 \pmod{144}$, we write $e = 144a - 6$ where a is a positive integer. Let B be an abelian surface such that Hypothesis 4.1 holds. We adopt the notation introduced in Section 4. In particular $A = B/\langle \epsilon \rangle$, and we have the ample class ω_A on A with $\omega_A^2 = 4a$. We assume throughout that the positive integer d (see (4.1.1)) is odd and large (the lower bound on d depends only on a). Now we choose the line bundle \mathcal{L} on X :

- (1) If $e \equiv -6 \pmod{16}$ we let $c_1(\mathcal{L}) = \nu^*(\mu_B(\omega_B))$,
- (2) if $e \equiv -6 \pmod{144}$ we let $c_1(\mathcal{L}) = \nu^*(3\mu_B(\omega_B))$.

Set $\mathcal{E}_0 := \mathcal{E}(\mathcal{L})$ and $h_0 := c_1(\mathcal{E}_0)$. By (3.4.5) we have

- (1) $h_0 = 2\mu_A(\omega_A) - \delta(A)$ if $e \equiv -6 \pmod{16}$, and
- (2) $h_0 = 6\mu_A(\omega_A) - \delta(A)$ if $e \equiv -6 \pmod{144}$.

Hence $q(h_0) = e$ in both cases, and $\mathrm{div}(h_0) = 2$ in the first case, while $\mathrm{div}(h_0) = 6$ in the second case. Moreover h_0 is ample by Proposition 6.1 (recall that $d \gg 0$, this is where we need d to be large). This proves that h_0 is a polarization of the type appearing in Theorem 1.1.

Let $\mathcal{M} \rightarrow T$ be a (sufficiently small) representative of $\mathrm{Def}(K_2(A), h_0)$. By Proposition 4.2 the vector bundle \mathcal{E}_0 has an extension to a vector bundle \mathcal{E}_t on M_t for every $t \in T$ (the extension is unique by Proposition 5.1).

Proposition 6.5. *Keeping notation as above the vector bundle \mathcal{E}_t is h_t slope stable for general $t \in T$, where h_t is the polarization (deformation of h_0) on M_t .*

Proof. Let $N \subset T$ be the Noether-Lefschetz locus parametrizing M_t such that $\mu_A(\gamma_A)$ remains of type (1,1). If $t \in N$ then we have a Lagrangian fibration $\pi_t: M_t \rightarrow \mathbb{P}^2$ extending the Lagrangian fibration $\pi_A: K_2(A) \rightarrow |\mathcal{O}_E(3(0))|$. By Proposition 4.14 and openness of stability the restriction of \mathcal{E}_t to a general fiber of π_t is slope stable. Now assume in addition that t is very general in N , so that the Neron-Severi group of M_t has rank 2, and hence is obtained via parallel transport from the subgroup of $\mathrm{NS}(K_2(A))$ generated by $(2\mu_A(\omega_A) - \delta(A)), \mu_A(\gamma_A)$ in Case (1), respectively $6\mu_A(\omega_A) - \delta(A), \mu_A(\gamma_A)$ in Case (2). Applying Lemma 6.3 we get that there is a single 72-chamber in $\mathrm{NS}(M_t)$, and hence h_t is 72-suitable. By Proposition 4.10 it follows that \mathcal{E}_t is h_t slope stable (recall that $a(\mathcal{E}_t) = 72$ by Example 4.6). \square

Let $d \gg 0$. By Proposition 6.5 (and its proof) we have an irreducible component $\mathrm{VB}_e^i(d)$ of the moduli stack VB_e^i , where $i = 2$ in the Case (1) and $i = 6$ in Case (2), such that the map $\mathrm{VB}_e^i(d) \rightarrow \mathrm{Kum}_e^i$ is dominant of finite degree. Since Kum_e^2 is irreducible (see Example 2.10 in [Ono16]), this finishes the proof for $i = 2$. On the other hand there are two irreducible components of Kum_e^6 and, as is easily checked using the results in [Ono16], the image of $\mathrm{VB}_e^6(d) \rightarrow \mathrm{Kum}_e^6$ lands into the same irreducible component of Kum_e^6 for all $d \gg 0$.

APPENDIX A. EXAMPLES OF SEMI-HOMOGENEOUS VECTOR BUNDLES

A.1. Basics. In the present section “abelian variety means abelian variety”, and similarly for “elliptic curve”. Let A be an abelian variety. For $a \in A$, we let $T_a: A \rightarrow A$ be the translation by a . If \mathcal{L} is a line bundle on A , we let

$$\begin{array}{ccc} A & \xrightarrow{\varphi_{\mathcal{L}}} & A^\vee \\ a & \mapsto & T_a^*(\mathcal{L}) \otimes \mathcal{L}^{-1}. \end{array} \quad (\text{A.1.1})$$

We recall that a vector bundle \mathcal{F} on A is *semi-homogeneous* if, for every $a \in A$, there exists a line bundle \mathcal{L}_a such that $T_a^* \mathcal{F} \cong \mathcal{F} \otimes \mathcal{L}_a$. We recall the following result of Oda (Theorem 1.2 in [Oda71]) and Mukai (Theorem 5.8 in [Muk78]).

Theorem A.1 (Oda, Mukai). *A vector bundle \mathcal{E} on an abelian variety A is simple and semi-homogeneous if and only if it is isomorphic to $f_* \mathcal{L}$ where $f: \tilde{A} \rightarrow A$ is an isogeny and \mathcal{L} is a line bundle on \tilde{A} such that $\ker(f) \cap \ker(\varphi_{\mathcal{L}}) = \{0\}$.*

A.2. Semi-homogeneous vector bundles on powers of elliptic curves. Let C be an elliptic curve, let \tilde{C} be an elliptic curve and let $f: \tilde{C} \rightarrow C$ be an isogeny. For $i \in \{1, \dots, n+1\}$ let L_i be a line bundle on \tilde{C} of degree d_0 (independent of i). Let $Z_n(C) \subset C^{n+1}$, $Z_n(\tilde{C}) \subset \tilde{C}^{n+1}$ be as in (4.4.5) and for $i \in \{1, \dots, n+1\}$ let $\tilde{p}_i: Z_n(\tilde{C}) \rightarrow \tilde{C}$ be the i -th projection. Lastly let

$$\begin{array}{ccc} Z_n(\tilde{C}) & \xrightarrow{\psi^n} & Z_n(C) \\ (z_1, \dots, z_{n+1}) & \mapsto & (f(z_1), \dots, f(z_{n+1})) \end{array}$$

Proposition A.2. *Keeping notation as above, suppose that $\deg f$ is coprime to $(n+1) \cdot d_0$. Then $\mathcal{G} := \psi_*^n \left(\bigotimes_{i=1}^{n+1} \tilde{p}_i^* L_i \right)$ is a simple semi-homogeneous vector bundle on $Z_n(C)$ of rank $(\deg f)^n$.*

Proof. The sheaf \mathcal{G} is locally free of rank equal to $\deg(\psi^n)$ because ψ^n is finite. Since $\ker(\psi^n) \cong (\ker f)^n$, it follows that \mathcal{G} has rank $(\deg f)^n$. In order to simplify the notation we let

$$\xi := \bigotimes_{i=1}^{n+1} \tilde{p}_i^* L_i.$$

Let us prove that

$$\int_{Z_n(\tilde{C})} c_1(\xi)^n = (n+1) \cdot d_0^n \quad (\text{A.2.2})$$

In fact, let $\{\alpha, \beta\}$ be a \mathbb{Z} -basis of $H^1(\tilde{C}; \mathbb{Z})$, and for $i \in \{1, \dots, n\}$ (we stop at n) let

$$x_i := \tilde{p}_i^*(\alpha)|_{Z_n(\tilde{C})}, \quad y_i := \tilde{p}_i^*(\beta)|_{Z_n(\tilde{C})}.$$

Then $\{x_1, \dots, x_n, y_1, \dots, y_n\}$ is a basis of $H^1(Z_n(\tilde{C}); \mathbb{Z})$, and we have

$$c_1(\xi) = d_0 \cdot \sum_{i=1}^n x_i \wedge (y_1 + \dots + y_{i-1} + 2y_i + y_{i+1} + \dots + y_n). \quad (\text{A.2.3})$$

Equation (A.2.2) follows from the above formula and a straightforward computation. By (A.2.2), the order of $\ker(\varphi_\xi)$ is equal to $(n+1)^2 \cdot d_0^{2n}$, and hence is coprime to $(\deg f)^n$ by our hypothesis. Since the order of $\ker(\psi^n)$ is $(\deg f)^n$, it follows that

$$\ker(\psi^n) \cap \ker(\varphi_\xi) = \{0\}. \quad (\text{A.2.4})$$

(Here φ_ξ is as in (A.1.1).) Hence \mathcal{G} is simple semi-homogeneous by Theorem A.1. \square

APPENDIX B. PROPERLY SEMISTABLE VECTOR BUNDLES WITH VANISHING
DISCRIMINANT ON ABELIAN SURFACES

In the present subsection we extend Proposition 4.4 and Corollary 4.5 of [O'G22] to arbitrary polarized abelian surfaces.

Proposition B.1. *Let (A, θ) be a $(1, e)$ polarized abelian surface. Let \mathcal{F} be a slope semistable vector bundle on A such that $\Delta(\mathcal{F}) = 0$ and $c_1(\mathcal{F}) = a\theta$. Then there exist coprime integers r_0, b_0 , with r_0 positive, and a positive integer m such that*

$$r(\mathcal{F}) = m \frac{r_0^2}{g}, \quad a = m \frac{r_0}{g} b_0, \quad (\text{B.0.1})$$

where $g := \gcd\{r_0, e\}$. If \mathcal{F} is slope strictly semistable, i.e. slope semistable but not slope stable, then $m \geq 2$.

Proof. Suppose first that \mathcal{F} is slope stable. Then \mathcal{F} is semi-homogeneous because $\Delta(\mathcal{F}) = 0$, see Proposition A.2 in [O'G22]. Since it is also simple, the thesis holds by Proposition A.3 in [O'G22].

Now suppose that \mathcal{F} is strictly slope semistable. Then there exist an integer $m \geq 2$ and a (slope stability) Jordan-Holder filtration of \mathcal{F}

$$0 = \mathcal{G}_0 \subsetneq \mathcal{G}_1 \subsetneq \dots \subsetneq \mathcal{G}_m = \mathcal{F} \quad (\text{B.0.2})$$

such that for each $i \in \{1, \dots, m\}$ the sheaf $\mathcal{G}_i/\mathcal{G}_{i-1}$ is locally free, $c_1(\mathcal{G}_i/\mathcal{G}_{i-1}) = a_i\theta$, and $\Delta(\mathcal{G}_i/\mathcal{G}_{i-1}) = 0$, see the proof of Proposition 4.4 in [O'G22]. Arguing as above we get that $\mathcal{G}_i/\mathcal{G}_{i-1}$ is a simple semi-homogeneous vector bundle. Let $i \in \{1, \dots, m\}$. By Proposition A.3 in [O'G22] there exist coprime integers r_i, b_i , with r_i positive, such that

$$r(\mathcal{G}_i/\mathcal{G}_{i-1}) = \frac{r_i^2}{g_i}, \quad a_i = \frac{r_i}{g_i} b_i,$$

where $g_i := \gcd\{r_i, e\}$. Hence the slope of $\mathcal{G}_i/\mathcal{G}_{i-1}$ is equal to $(b_i/r_i)\theta^2$. Since the filtration in (B.0.2) is the (slope stability) Jordan-Holder filtration of \mathcal{F} , it follows that $(b_1/r_1) = \dots = (b_m/r_m)$. Since r_i, b_i are coprime we get that $r_1 = \dots = r_m$ and $b_1 = \dots = b_m$. Hence there exist coprime integers r_0, b_0 , with r_0 positive, such that

$$r(\mathcal{G}_i/\mathcal{G}_{i-1}) = \frac{r_0^2}{g_0}, \quad c_1(\mathcal{G}_i/\mathcal{G}_{i-1}) = \frac{r_0}{g_0} b_0.$$

where $g_0 := \gcd\{r_0, e\}$. This proves the thesis because $m \geq 2$. \square

Corollary B.2. *Let (A, θ) be a $(1, e)$ polarized abelian surface. Let \mathcal{F} be a slope semistable vector bundle on A such that $\Delta(\mathcal{F}) = 0$. Suppose that there exist coprime integers s_0, c_0 , with s_0 positive, such that $\gcd\{s_0, e\} = 1$ and*

$$r(\mathcal{F}) = s_0^2, \quad c_1(\mathcal{F}) = s_0 c_0 \theta. \quad (\text{B.0.3})$$

Then \mathcal{F} is slope stable.

Proof. By Proposition B.1 there exist coprime integers r_0, b_0 , with r_0 positive, and a positive integer m such that

$$s_0^2 = m \frac{r_0^2}{g}, \quad s_0 c_0 = m \frac{r_0}{g} b_0, \quad (\text{B.0.4})$$

where $g := \gcd\{r_0, e\}$. It follows that $r_0 c_0 = s_0 b_0$. Since r_0, b_0 are coprime and also s_0, c_0 , it follows that $r_0 = s_0$ and $b_0 = c_0$. This gives that $m = g$. Since $r_0 = s_0$ and by hypothesis s_0 is coprime to e , we have $g = 1$, and thus $m = 1$. By Proposition B.1 it follows that \mathcal{F} is slope stable. \square

REFERENCES

- [Ati57] Michael F. Atiyah, *Vector bundles over an elliptic curve*, Proc. London Math. Soc. (3) **7** (1957), 414–452. (1957), no. 7, 414–452.
- [Bar87] Wolf Barth, *Abelian surfaces with (1, 2)-polarization*, Algebraic geometry, Sendai, 1985, Adv. Stud. Pure Math., vol. 10, North-Holland, Amsterdam, 1987, pp. 41–84. MR 946234
- [BD85] Arnaud Beauville and Ron Donagi, *La variété des droites d’une hypersurface cubique de dimension 4*, C. R. Acad. Sci. Paris Sér. I Math. **301** (1985), no. 14, 703–706. MR 818549
- [Bec22] Thorsten Beckmann, *Atomic objects on hyper-kähler manifolds*, arXiv:2202.01184 (2022).
- [BLM⁺21] Arend Bayer, Martí Lahoz, Emanuele Macrì, Howard Nuer, Alexander Perry, and Paolo Stellari, *Stability conditions in families*, Publ. Math. Inst. Hautes Études Sci. **133** (2021), 157–325. MR 4292740
- [DV10] Olivier Debarre and Claire Voisin, *Hyper-Kähler fourfolds and Grassmann geometry*, J. Reine Angew. Math. **649** (2010), 63–87. MR 2746467
- [ES98] Geir Ellingsrud and Stein Arild Stromme, *An intersection number for the punctual Hilbert scheme of a surface*, Trans. Amer. Math. Soc. **350** (1998), no. 6, 2547–2552. MR 1432198
- [Fri89] Robert Friedman, *Rank two vector bundles over regular elliptic surfaces*, Invent. Math. **96** (1989), no. 2, 283–332. MR 989699
- [GT] Daniel Greb and Matei Toma, *Compact moduli spaces for slope-semistable sheaves*, arXiv:1303.2480 [math.AG].
- [GT17] Daniel Greb and Matei Toma, *Compact moduli spaces for slope-semistable sheaves*, Algebr. Geom. **4** (2017), no. 1, 40–78. MR 3592465
- [Hai01] Mark Haiman, *Hilbert schemes, polygraphs and the Macdonald positivity conjecture*, J. Amer. Math. Soc. **14** (2001), no. 4, 941–1006. MR 1839919
- [HT13] Brendan Hassett and Yuri Tschinkel, *Hodge theory and Lagrangian planes on generalized Kummer fourfolds*, Mosc. Math. J. **13** (2013), no. 1, 33–56, 189. MR 3112215
- [Ito18] Atsushi Ito, *A remark on higher syzygies on abelian surfaces*, Comm. Algebra **46** (2018), no. 12, 5342–5347. MR 3923760
- [KL19] Alex Küronya and Victor Lozovanu, *A Reider-type theorem for higher syzygies on abelian surfaces*, Algebr. Geom. **6** (2019), no. 5, 548–570. MR 4009173
- [KLCM19] Andreas Leopold Knutsen, Margherita Lelli-Chiesa, and Giovanni Mongardi, *Wall divisors and algebraically coisotropic subvarieties of irreducible holomorphic symplectic manifolds*, Trans. Amer. Math. Soc. **371** (2019), no. 2, 1403–1438. MR 3885184
- [LLSvS17] Christian Lehn, Manfred Lehn, Christoph Sorger, and Duco van Straten, *Twisted cubics on cubic fourfolds*, J. Reine Angew. Math. **731** (2017), 87–128. MR 3709061
- [Mar21] Eyal Markman, *Stable vector bundles on a hyper-kähler manifold with a rank 1 obstruction map are modular*, arXiv:2107.13991 (2021).
- [Muk78] Shigeru Mukai, *Semi-homogeneous vector bundles on an Abelian variety*, J. Math. Kyoto Univ. **18** (1978), no. 2, 239–272. MR 0498572
- [Muk87] S. Mukai, *On the moduli space of bundles on K3 surfaces. I*, Vector bundles on algebraic varieties (Bombay, 1984), Tata Inst. Fund. Res. Stud. Math., vol. 11, Tata Inst. Fund. Res., Bombay, 1987, pp. 341–413. MR 893604
- [Muk88] Shigeru Mukai, *Curves, K3 surfaces and fano 3-folds of genus 10*, Algebraic geometry and commutative algebra, vol. I, Kinokuniya, Tokyo, 1988, pp. 357–377.
- [Oda71] Tadao Oda, *Vector bundles on an elliptic curve*, Nagoya Math. J. **43** (1971), 41–72. MR 0318151
- [O’G06] Kieran G. O’Grady, *Irreducible symplectic 4-folds and Eisenbud-Popescu-Walter sextics*, Duke Math. J. **134** (2006), no. 1, 99–137.
- [O’G22] ———, *Modular sheaves on hyperkähler varieties*, Algebraic Geometry **9** (2022), no. 1, 1–38.
- [Ono16] Claudio Onorati, *Connected components of moduli spaces of generalised kummer varieties*, arXiv:1608.06465 (2016).
- [Son16] Lei Song, *On the universal family of Hilbert schemes of points on a surface*, J. Algebra **456** (2016), 348–354. MR 3484148
- [Ver96a] Mikhail Verbitsky, *Cohomology of compact hyper-Kähler manifolds and its applications*, Geom. Funct. Anal. **6** (1996), no. 4, 601–611. MR 1406664
- [Ver96b] ———, *Hyperholomorphic bundles over a hyper-Kähler manifold*, J. Algebraic Geom. **5** (1996), no. 4, 633–669. MR 1486984

- [Wie16] Benjamin Wieneck, *On polarization types of Lagrangian fibrations*, Manuscripta Math. **151** (2016), no. 3-4, 305–327. MR 3556822
- [Wie18] ———, *Monodromy invariants and polarization types of generalized Kummer fibrations*, Math. Z. **290** (2018), no. 1-2, 347–378. MR 3848435
- [Yos16] Kota Yoshioka, *Bridgeland’s stability and the positive cone of the moduli spaces of stable objects on an abelian surface*, Development of moduli theory—Kyoto 2013, Adv. Stud. Pure Math., vol. 69, Math. Soc. Japan, [Tokyo], 2016, pp. 473–537. MR 3616985