

CONICS ON GUSHEL-MUKAI FOURFOLDS, EPW SEXTICS AND BRIDGELAND MODULI SPACES

HANFEI GUO, ZHIYU LIU, SHIZHUO ZHANG

ABSTRACT. We identify the double dual EPW sextic \tilde{Y}_{A^\perp} and the double EPW sextic \tilde{Y}_A , associated with a very general Gushel-Mukai fourfold X , with the Bridgeland moduli spaces of stable objects of character Λ_1 and Λ_2 in the Kuznetsov component $\mathcal{K}u(X)$. This provides an affirmative answer to a question of Perry-Pertusi-Zhao. As an application, we prove a conjecture of Kuznetsov-Perry for very general Gushel-Mukai fourfolds.

1. INTRODUCTION

1.1. Hyperkähler varieties as Bridgeland moduli spaces for Kuznetsov components of Fano fourfolds. Compact hyperkähler varieties are higher-dimensional analogues of K3 surfaces, which are important building blocks of algebraic geometry. However, constructing a compact hyperkähler variety is involved so that only a few of examples are known. The major examples are moduli spaces of stable sheaves on a K3 surface, by the work [Bea83], [Muk84], [O’G95], [Yos99], [Yos01] and many others.

To produce more examples of hyperkähler varieties, one could consider the moduli spaces of stable objects on a non-commutative K3 surface, which we now briefly explain. In [Kuz04], Kuznetsov constructs a semi-orthogonal decomposition of the derived category of a cubic fourfold X

$$D^b(X) = \langle \mathcal{K}u(X), \mathcal{O}_X, \mathcal{O}_X(H), \mathcal{O}_X(2H) \rangle,$$

where H is the ample line bundle $\mathcal{O}_{\mathbb{P}^5}(1)|_X$. Kuznetsov observes that the non-trivial semi-orthogonal component $\mathcal{K}u(X)$ is a K3 category in the sense that it has the same Serre functor and Hochschild cohomology as the derived category of a K3 surface. On the other hand, it has now been expected that the Kuznetsov component of a smooth Fano variety encodes essential birational geometric information. In [BLMS17], the authors construct stability conditions on the Kuznetsov component of a series of Fano varieties, including $\mathcal{K}u(X)$. In particular, one could construct Bridgeland moduli spaces of stable objects in $\mathcal{K}u(X)$ with respect to the stability conditions. Under certain circumstances, these moduli spaces provide new examples of hyperkähler varieties. In the present article, we focus on the case of a Gushel-Mukai(GM) fourfold X , which is a degree 10 and index two Fano variety. A general GM fourfold is defined by a smooth transverse intersection of $\mathrm{Gr}(2, V_5)$ with a linear section \mathbb{P}^8 and a quadric section Q in \mathbb{P}^9 after the Plücker embedding

$$X := \mathrm{Gr}(2, V_5) \cap \mathbb{P}^8 \cap Q.$$

By the work [DIM15], [DK18], [KP18], [DK19] and many others, it is shown that GM fourfolds share many similarities with cubic fourfolds. For example, a GM fourfold X also admits a semi-orthogonal decomposition

$$D^b(X) = \langle \mathcal{K}u(X), \mathcal{O}_X, \mathcal{U}^\vee, \mathcal{O}_X(H), \mathcal{U}^\vee(H) \rangle.$$

Here \mathcal{U} is the pull back of the tautological bundle on $\mathrm{Gr}(2, V_5)$ and $\mathcal{O}_X(H)$ is the restriction of the Plücker line bundle $\mathcal{O}_{\mathbb{P}^9}(1)$. Furthermore, $\mathcal{K}u(X)$ is also a K3 category. In particular,

2010 *Mathematics Subject Classification.* Primary 14F05; secondary 14J45, 14D20, 14D23.

Key words and phrases. Bridgeland moduli spaces, Kuznetsov components, Gushel-Mukai fourfolds, hyperkähler variety.

there is a rank two lattice inside the numerical Grothendieck group $\mathcal{N}(\mathcal{K}u(X))$ generated by

$$\Lambda_1 = -2 + (H^2 - \Sigma') - \frac{1}{2}P, \quad \Lambda_2 = -4 + 2H - \frac{5}{3}L,$$

where Σ' is the class of a degree 6 surface. In [PPZ19], the authors construct stability conditions on $\mathcal{K}u(X)$ and they show that for a non-zero primitive Mukai vector v and a generic stability condition σ , if the moduli space $\mathcal{M}_\sigma(v)$ is non-empty, it is a smooth projective hyperkähler variety of dimension $(v, v) + 2$. In particular, they prove that if X is very general, $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1)$ is either isomorphic to the double dual EPW sextic \tilde{Y}_{A^\perp} or the double EPW sextic \tilde{Y}_A , where A is the Lagrangian data associated with X . Furthermore, they expect that there exist two isomorphisms, $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1) \cong \tilde{Y}_{A^\perp}$ and $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_2) \cong \tilde{Y}_A$.

1.2. Main Results. The first main result of our article answers the question of Perry-Pertusi-Zhao.

Theorem 1.1. *Let X be a very general GM fourfold. Then, for a generic stability condition σ on $\mathcal{K}u(X)$, we have*

- (1) $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1) \cong \tilde{Y}_{A^\perp}$.
- (2) $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_2) \cong \tilde{Y}_A$.
- (3) *There is an involutive auto-equivalence on $\mathcal{K}u(X)$ and the induced involution on $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1)$ coincides with the natural involution on \tilde{Y}_{A^\perp} .*

Remark 1.2. Here *very general* means *general non-Hodge-special*, see Definition 2.1. We only use this assumption in Theorem 7.1 to prove the stability of projection objects. Once Theorem 7.1 is known for general GM fourfolds, Theorem 1.1 can be generalized to the case of general GM fourfolds. This is a work in progress.

Our approach to Theorem 1.1 is projecting objects related to conics to the Kuznetsov component. The Hilbert scheme of conics $F_g(X)$ of a general GM fourfold X is a smooth projective variety of dimension 5. By [IM11], there exists a morphism

$$f : F_g(X) \longrightarrow \tilde{Y}_{A^\perp}$$

and this morphism is an essential \mathbb{P}^1 -fibration in that f contracts a \mathbb{P}^1 -family of generic conics, while taking two special types of conics to two different points. Starting with a twisted structure sheaf $\mathcal{O}_C(H)$ of a conic $C \subset X$, we show that the projection functor to $\mathcal{K}u(X)$ produces an essential \mathbb{P}^1 -fibration over $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1)$. More precisely, if X is a very general GM fourfold, we prove that the morphism $p : F_g(X) \rightarrow \mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1)$ induced by the projection functor coincides with the morphism f constructed in [IM11]. As a result, we prove that $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1) \cong \tilde{Y}_{A^\perp}$.

In [KP19], the authors study GM varieties of arbitrary dimension and propose the following conjecture.

Conjecture 1.3 ([KP19, Conjecture 1.7]). *If X and X' are GM varieties of the same dimension such that there exists an equivalence $\mathcal{K}u(X) \simeq \mathcal{K}u(X')$, then X and X' are birationally equivalent.*

In [JLLZ21], we show the conjecture is true for general GM threefolds. In this article, we prove the following theorem.

Theorem 1.4. *Let X and X' be very general GM fourfolds. If there is an equivalence $\mathcal{K}u(X) \simeq \mathcal{K}u(X')$, then X and X' are period partners or period duals. In particular, X is birational to X' .*

We will review the definitions of period partners and period duals in Section 2. Roughly speaking, X and X' have the same period point up to an involution of the period domain.

The idea is very similar to the proof for GM threefolds as in [JLLZ21, Theorem 10.1]. The equivalence $\Phi : \mathcal{K}u(X) \simeq \mathcal{K}u(X')$ would induce an isomorphism from the moduli space $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_2)$ to either $\mathcal{M}_\sigma(\mathcal{K}u(X'), \Lambda'_2)$ or $\mathcal{M}_\sigma(\mathcal{K}u(X'), \Lambda'_1)$. By Theorem 1.1, the former case shows that X and X' are period partners while the latter shows that they are period duals. In both cases, X is birationally equivalent to X' by [DK18, Corollary 4.16, Theorem 4.20] and [DK19, Remark 5.28].

1.3. Related work.

1.3.1. *Hyperkähler varieties as Bridgeland moduli spaces for Kuznetsov components.* In [LPZ18], the authors reconstruct the Fano variety of lines for any cubic fourfold and the LLSvS eightfold for cubic fourfolds not containing a plane as the moduli spaces of stable objects on the Kuznetsov component with primitive Mukai vector $\lambda_1 + \lambda_2$ and $2\lambda_1 + \lambda_2$ respectively. In [LPZ20], the authors show that a symplectic resolution of the moduli space $\mathcal{M}_\sigma(2\lambda_1 + 2\lambda_2)$ is a hyperkähler variety, deformation equivalent to O'Grady 10.

1.3.2. *Birational categorical Torelli for GM varieties.* In [JLLZ21], we show that the Kuznetsov component determines the birational isomorphism class for general GM threefolds while in the present article, we prove a similar statement for very general GM fourfolds. Since GM fivefolds and sixfolds are all rational, the analogous statements automatically hold in these cases.

1.4. Further questions.

1.4.1. *Structure of the moduli space $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_2)$.* It would be interesting to know if the Bridgeland moduli space $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_2)$ can be realized as a Gieseker moduli space on X . Inspired by our previous work [JLLZ21] for GM threefolds, we wonder if the moduli space $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_2)$ is isomorphic to the moduli space $M_X(4 - 2H + \frac{1}{6}H^3)$ of semistable sheaves on X .

1.4.2. *Refined categorical Torelli for GM fourfolds.* The duality conjecture [KP19, Theorem 1.6] tells us that the Kuznetsov component of GM varieties cannot determine the isomorphism class. In [JLLZ21], we prove what we called *Refined categorical Torelli theorem* for GM threefolds, meaning that an extra assumption can be made on the equivalence $\Phi : \mathcal{K}u(X) \simeq \mathcal{K}u(X')$ of GM threefolds X and X' to deduce that $X \cong X'$. It is natural to see if similar statements can be proved for GM fourfolds, as asked in [PS22, Question 6.5].

1.5. **Organization of the paper.** In Section 2, we review the basic terminologies of GM fourfolds and the associated hyperkähler varieties, double EPW sextics and double dual EPW sextics. In Section 3, we introduce the semi-orthogonal decomposition of GM fourfolds and construct an involutive auto-equivalence on the Kuznetsov component. In Section 4, we briefly review the concepts of weak stability conditions on a general triangulated category and stability conditions on the Kuznetsov component $\mathcal{K}u(X)$ for a GM fourfold X . Then we prove some properties of stability conditions on $\mathcal{K}u(X)$ which will be used later. In Section 5, we introduce three types of conics on GM fourfolds. In Section 6, we compute the projection objects for conics of each type. In Section 7, we prove the stability of the projection objects of conics. In Section 8, we show that the morphism induced by the projection functor coincides with the classical map defined by Iliev-Manivel in [IM11], and as a consequence we prove Theorem 1.1. In Section 9, using results in Section 8, we prove Theorem 1.4.

1.6. Notation and conventions.

- We work over $k = \mathbb{C}$.
- The term K3 surface means a smooth projective K3 surface.
- We denote the bounded derived category of coherent sheaves on a smooth variety X by $D^b(X)$. The derived dual functor $R\mathcal{H}om_X(-, \mathcal{O}_X)$ on $D^b(X)$ is denoted by $\mathbb{D}(-)$.
- If $X \rightarrow Y$ is a morphism between varieties and $F \in D^b(Y)$, then we often write F_X for the pullback of F to X . By abuse of notation, if D is a divisor on Y , we often still denote its pullback by D .
- We will use V_i to denote a complex vector space of dimension i .
- We use hom and ext^i to represent the dimension of the vector spaces Hom and Ext^i for some $i > 0$.
- We denote the Hilbert scheme of conics on a variety X by $F_g(X)$.
- The symbol \simeq denotes an equivalence of categories and a birational equivalence of varieties. The symbol \cong denotes an isomorphism between varieties, complexes or functors.

Acknowledgements. Firstly, it is our pleasure to thank Arend Bayer and Qizheng Yin for very useful discussions on the topics of this project. We would like to thank Sasha Kuznetsov, Kieran G. O’Grady, Alexander Perry, Laura Pertusi and Xiaolei Zhao for helpful comments. The first author would like to thank her best friend Guolei Zhong for his suggestions and companion. The second author would like to thank Huizhi Liu for encouragement and support. The third author thanks Tingyu Sun for constant support and encouragement. The third author is supported by the ERC Consolidator Grant WallCrossAG, no. 819864.

2. GEOMETRY OF GUSHEL-MUKAI FOURFOLDS AND THE ASSOCIATED EPW SEXTICS

Let X be an ordinary GM fourfold, which is defined by a smooth transverse intersection of $\text{Gr}(2, V_5)$ with a linear section \mathbb{P}^8 and a quadric section Q in \mathbb{P}^9 after the Plücker embedding

$$X := \text{Gr}(2, V_5) \cap \mathbb{P}^8 \cap Q.$$

There is a natural embedding $\gamma_X : X \rightarrow \text{Gr}(2, V_5)$, which is usually called the Gushel map. We define $\mathcal{O}_X(H) := \mathcal{O}_{\mathbb{P}^9}(1)|_X$ and $\mathcal{U} := \gamma_X^* \mathcal{U}_{\text{Gr}(2, V_5)}$, where $\mathcal{U}_{\text{Gr}(2, V_5)}$ is the tautological rank two sub-bundle of $\text{Gr}(2, V_5)$.

We denote $\sigma_{i,j} \in H^{2(i+j)}(\text{Gr}(2, V_5), \mathbb{Z})$ the Schubert cycles of $\text{Gr}(2, V_5)$ for any integers $3 \geq i \geq j \geq 0$. By [DK19, Proposition 3.4], the cohomology group $H^4(X, \mathbb{Z})$ is torsion free. The image of the embedding $\gamma_X^* : H^4(\text{Gr}(2, V_5), \mathbb{Z}) \rightarrow H^4(X, \mathbb{Z})$ is a rank two sub-lattice generated by $\gamma_X^*(\sigma_1)^2$ and $\gamma_X^*(\sigma_2)$.

Definition 2.1. An ordinary GM fourfold X is called *non-Hodge-special* if

$$H^{2,2}(X) \cap H^4(X, \mathbb{Z}) = \gamma_X^* H^4(\text{Gr}(2, V_5), \mathbb{Z}).$$

It means $H^{2,2}(X) \cap H^4(X, \mathbb{Z})$ is a rank two integral lattice. X is called *Hodge-special* if the lattice $H^{2,2}(X) \cap H^4(X, \mathbb{Z})$ is of rank at least three.

According to [DIM15], [DK19] and [Deb20, Section 4.5], there is a period map from the moduli stack of GM fourfolds to the period domain

$$\wp_4 : \mathbf{M}_4^{\text{GM}} \longrightarrow \mathcal{D}.$$

In particular, the locus of periods of the Hodge-special GM fourfolds constitute a countably union of hypersurfaces.

In the current paper, we will always assume X to be *very general* in the sense that it is general among the locus of non-Hodge-special GM fourfolds. For a comment on the general case, see Remark 1.2.

2.1. EPW sextics and conics on Gushel-Mukai fourfolds. As a cubic fourfold admits an associated hyperkähler variety, which is called Fano variety of lines, a general GM fourfold also admits its associated hyperkähler variety.

Here we briefly review the definition and some properties of EPW sextics introduced by Eisenbud, Popescu, and Walter, in particular their relations with GM varieties.

Let X be an ordinary GM fourfold, following [DK19], one can naturally associate a triple $(A(X), V_5(X), V_6(X))$ with X , which is called a *Lagrangian data* of X . Here $V_5(X)$ is a hyperplane in $V_6(X)$ and $A(X) \subset \bigwedge^3 V_6(X)$ is Lagrangian with respect to the natural symplectic structure on $\bigwedge^3 V_6(X)$ given by wedge product. When X is clear, we will use the notation (A, V_5, V_6) .

Starting from a Lagrangian data, one can construct various varieties which are important to the geometry of GM fourfolds. For any integer $l \geq 0$, we define closed subschemes

$$Y_A^{\geq l} := \{[U_1] \in \mathbf{P}(V_6) \mid \dim(A \cap (U_1 \wedge \bigwedge^2 V_6)) \geq l\} \subset \mathbf{P}(V_6),$$

$$Y_{A^\perp}^{\geq l} := \{[U_5] \in \mathbf{P}(V_6^\vee) \mid \dim(A \cap \bigwedge^3 U_5) \geq l\} \subset \mathbf{P}(V_6^\vee).$$

At the same time, we set $Y_A^\ell := Y_A^{\geq \ell} \setminus Y_A^{\geq \ell+1}$ and $Y_{A^\perp}^\ell := Y_{A^\perp}^{\geq \ell} \setminus Y_{A^\perp}^{\geq \ell+1}$.

If X is general, we can assume that A is also general. Then $Y_A := Y_A^{\geq 1} \subset \mathbf{P}(V_6)$ is a normal integral sextic hypersurface, which is called an EPW sextic. The fourfold Y_A is singular at the integral surface $Y_A^{\geq 2}$. In [O'G10, Section 1.2], the author constructs a canonical double cover

$$\tilde{Y}_A \rightarrow Y_A,$$

branched over the integral surface $Y_A^{\geq 2}$, which is called the double EPW sextic. Since A is general, $Y_A^{\geq 3} = \emptyset$, \tilde{Y}_A is a smooth hyperkähler fourfold. The analogue statements also hold for $Y_{A^\perp}^{\geq l}$.

Many properties of a GM fourfold X depend on $A(X)$, its even part of the corresponding Lagrangian subspace. Here we introduce two important notions called period partner and period dual.

Definition 2.2 ([DK18, Definition 3.20]). Two GM fourfolds X_1 and X_2 are called *period partners* if there exists an isomorphism $\phi : V_6(X_1) \cong V_6(X_2)$ such that we have $(\bigwedge^3 \phi)(A(X_1)) = A(X_2)$. They are called *period duals* if there exists an isomorphism $\phi : V_6(X_1) \cong V_6(X_2)^\vee$ such that $(\bigwedge^3 \phi)(A(X_1)) = A(X_2)^\perp$.

By definition, period partners are constructed by a same Lagrangian subspace A , but possibly with different hyperplanes of V_6 . For GM fourfolds, period partners share the same period point as it is shown in the following theorem.

Theorem 2.3 ([DK19, Theorem 5.1]). *Let X be a GM fourfold, with associated Lagrangian data (A, V_5, V_6) . Assume that A is general so that \tilde{Y}_A is smooth, then there is an isomorphism between two polarized Hodge structure*

$$H^4(X, \mathbf{Z})_{00} \simeq H^2(\tilde{Y}_A, \mathbf{Z})_0(-1).$$

According to [O'G06], for any $A \in \mathbb{L}\mathbb{G}(\bigwedge^3 V_6)^{00}$ (i.e., the associated double cover of the EPW sextic and its dual \tilde{Y}_A and \tilde{Y}_{A^\perp} are both smooth), the period points of \tilde{Y}_A and \tilde{Y}_{A^\perp} differ by an involution \bar{r} .

Theorem 2.4 ([O'G06, Theorem 1.1]). *Let \mathcal{P} be the period map of double EPW sextics and \bar{r} be the involution on the period domain,*

$$\mathcal{P} : \tilde{\mathbf{M}}_{\text{EPW}} \rightarrow \mathcal{Q}_2.$$

For $A \in \mathbb{L}\mathbb{G}(\bigwedge^3 V_6)^{00}$, we have $\mathcal{P}(A^\perp) = \bar{r} \circ \mathcal{P}(A)$,

Here the involution $\bar{r} : \mathcal{Q}_2 \rightarrow \mathcal{Q}_2$ is non-trivial, which implies $\mathcal{P}(A) \neq \mathcal{P}(A^\perp)$, for a generic $A \in \mathbb{L}\mathbb{G}(\wedge^3 V_6)^{00}$. As a corollary, \tilde{Y}_A is not isomorphic to \tilde{Y}_{A^\perp} for general A .

In [IM11], the authors show that the double EPW sextic can be constructed from the Hilbert scheme of conics on a general GM fourfold. Here we give a short review of their construction.

Denote by $F_g(X)$ the Hilbert scheme of conics lying on X . When X is general, it is a smooth projective variety of dimension five. $F_g(X)$ admits a natural map to a sextic hypersurface $Y_X^\vee \subset \mathbb{P}^5$, over which $F_g(X)$ is essentially a fibration in projective lines. By Stein factorization, we get

$$F_g(X) \xrightarrow{f} \tilde{Y}_X^\vee \rightarrow Y_X^\vee.$$

It turns out \tilde{Y}_X^\vee is also a smooth fourfold and the morphism f is birational to a \mathbb{P}^1 -bundle.

By [IM11, Theorem 3.2], there are three types of conics on X which are τ -conics, ρ -conics and σ -conics. The locus of the last two types $F_g^\rho(X)$ and $F_g^\sigma(X)$ are isomorphic to a three dimensional quadric Q^3 and a blow-up $\text{Bl}_o(\mathbb{P}^4)$ respectively.

Proposition 2.5 ([IM11]). *The morphism $f : F_g(X) \rightarrow \tilde{Y}_X^\vee$ is a birational \mathbb{P}^1 -bundle in that*

- (1) $f(F_g^\rho(X)) = p_1$ and $f(F_g^\sigma(X)) = p_2$, where $p_1, p_2 \in \tilde{Y}_X^\vee$ are preimages of the Plücker point under the double cover $\tilde{Y}_X^\vee \rightarrow Y_X^\vee$.
- (2) $f(F_g^\tau(X)) = \tilde{Y}_X^\vee - \{p_1, p_2\}$ and the restriction $f|_{F_g^\tau(X)} : F_g^\tau(X) \rightarrow \tilde{Y}_X^\vee - \{p_1, p_2\}$ is a \mathbb{P}^1 -bundle.

Thus the natural holomorphic two form on $F_g(X)$, which is induced by the generator of $H^{3,1}(X)$, descends to \tilde{Y}_X^\vee . This makes \tilde{Y}_X^\vee a hyperkähler fourfold. Indeed, $\tilde{Y}_X^\vee \rightarrow Y_X^\vee$ is a double cover and the natural involution is anti-symplectic as in [IM11, Proposition 4.17]. This implies Y_X^\vee is an EPW sextic and \tilde{Y}_X^\vee coincides with the double cover constructed by O'grady in Section 2.1. In the followings, we will use the notations Y_A and \tilde{Y}_A uniformly to refer to an EPW sextic and its double cover.

3. SEMI-ORTHOGONAL DECOMPOSITION AND KUZNETSOV COMPONENTS OF GUSHEL-MUKAI FOURFOLDS

Let X be a GM fourfold, the derived category $D^b(X)$ admits a semi-orthogonal decomposition, given by [KP18, Prop 4.2]

$$D^b(X) = \langle \mathcal{K}u(X), \mathcal{O}_X, \mathcal{U}^\vee, \mathcal{O}_X(H), \mathcal{U}^\vee(H) \rangle.$$

In this case, we define the projection functor as $\text{pr}_1 := \mathbf{L}\mathcal{O}_X \mathbf{L}\mathcal{U}^\vee \mathbf{L}\mathcal{O}_X(H) \mathbf{L}\mathcal{U}^\vee(H)$.

Using Serre duality, there is an alternative semi-orthogonal decomposition

$$D^b(X) = \langle \mathcal{O}_X(-H), \mathcal{U}, \mathcal{K}u(X), \mathcal{O}_X, \mathcal{U}^\vee \rangle.$$

Under this decomposition, we denote the projection functor by $\text{pr}_2 := \mathbf{R}\mathcal{U} \mathbf{R}\mathcal{O}_X(-H) \mathbf{L}\mathcal{O}_X \mathbf{L}\mathcal{U}^\vee$.

We denote the Grothendieck group of $\mathcal{K}u(X)$ by $K_0(\mathcal{K}u(X))$ and χ is the Euler form over it. Its numerical Grothendieck group is defined as $\mathcal{N}(\mathcal{K}u(X)) := K_0(\mathcal{K}u(X)) / \ker(\chi)$.

Lemma 3.1 ([KP18, Proposition 2.25]). *Let X be a very general GM fourfold, then $\mathcal{N}(\mathcal{K}u(X)) \cong \mathbb{Z}^2$. Furthermore, it is generated by two Mukai vectors Λ_1 and Λ_2 , where $\Lambda_1 = -2 + (H^2 - \Sigma') - \frac{1}{2}P$ and $\Lambda_2 = -4 + 2H - \frac{5}{3}L$. The Euler form $\chi(-, -)$ on $\langle \Lambda_1, \Lambda_2 \rangle$ is in the form*

$$\begin{bmatrix} -2 & 0 \\ 0 & -2 \end{bmatrix}. \quad (1)$$

Here $H := \gamma_X^* \sigma_1$ and $\Sigma' := \gamma_X^* \sigma_2$. By standard computation, $H \cdot \Sigma' = 6L$ and

$$H^2 \Sigma' = H^2 \cdot \gamma_X^* \sigma_2 = (\gamma_X^* \sigma_1)^2 \cdot \gamma_X^* \sigma_2 = \sigma_1^2 \cdot \sigma_2 = (\sigma_2)^2 + \sigma_{1,1} \cdot \sigma_2 = 6.$$

In [Per19], the Todd class of X is calculated

$$\mathrm{td}(X) = 1 + H + \left(\frac{2}{3} H^2 - \frac{1}{12} \Sigma' \right) + \frac{17}{60} H^3 + \frac{1}{10} H^4.$$

Then for any $\kappa = a + bH + (cH^2 + d\Sigma') + eL + fP \in \mathcal{N}(\mathcal{K}u(X))$, the Euler characteristic is given by

$$\chi(X, \kappa) = (\mathrm{ch}(\kappa) \cdot \mathrm{td}(X))_4 = a + \frac{17}{6}b + \frac{37}{6}c + \frac{11}{3}d + e + f.$$

Now we are going to introduce a functor T on $\mathcal{K}u(X)$, which is defined by $T := \mathbf{L}_{\mathcal{O}_X} \circ \mathbb{D}$.

Proposition 3.2. *The functor T is an involutive auto-equivalence on $\mathcal{K}u(X)$.*

Proof. First we prove that for any object $E \in \mathcal{K}u(X)$, we have $T(E) \in \mathcal{K}u(X)$. Indeed, by the definition of $\mathbf{L}_{\mathcal{O}_X}$, we have a triangle

$$\mathrm{RHom}(\mathcal{O}_X, \mathbb{D}(E)) \otimes \mathcal{O}_X \rightarrow \mathbb{D}(E) \rightarrow \mathbf{L}_{\mathcal{O}_X}(\mathbb{D}(E)) = T(E). \quad (2)$$

By Serre duality, we know that $\mathbb{D}(E) \in \langle \mathcal{K}u(X), \mathcal{O}_X \rangle$. As $\mathcal{O}_X \in \langle \mathcal{K}u(X), \mathcal{O}_X \rangle$, we have $\mathbf{L}_{\mathcal{O}_X}(\mathbb{D}(E)) \in \langle \mathcal{K}u(X), \mathcal{O}_X \rangle$. Thus if we apply $\mathrm{Hom}(\mathcal{O}_X, -)$ to the triangle (2), we have $\mathrm{RHom}(\mathcal{O}_X, \mathbf{L}_{\mathcal{O}_X}(\mathbb{D}(E))) = 0$, which implies $T(E) = \mathbf{L}_{\mathcal{O}_X}(\mathbb{D}(E)) \in \mathcal{K}u(X)$.

Then for any object $F \in \mathcal{K}u(X)$, if we apply $\mathrm{Hom}(\mathbb{D}(F), -)$ to the triangle (2), since $\mathrm{RHom}(\mathbb{D}(F), \mathcal{O}_X) = \mathrm{RHom}(\mathcal{O}_X, F) = 0$, we obtain natural isomorphisms

$$\mathrm{RHom}(\mathbb{D}(F), \mathbb{D}(E)) \cong \mathrm{RHom}(\mathbb{D}(F), T(E)) \cong \mathrm{RHom}(T(F), T(E)),$$

thus T is fully faithful. Note that the last isomorphism follows from $T(E) \in \mathcal{K}u(X)$ and the adjunction of $\mathbf{L}_{\mathcal{O}_X}$.

Now applying the functor \mathbb{D} to the triangle (2), we obtain a triangle

$$\mathbb{D}(T(E)) \rightarrow E \rightarrow \mathrm{RHom}(\mathcal{O}_X, \mathbb{D}(E))^\vee \otimes \mathcal{O}_X. \quad (3)$$

Taking the functor $\mathbf{L}_{\mathcal{O}_X}$ again, by $E \in \mathcal{K}u(X)$ and exceptionality of \mathcal{O}_X , we have $T(T(E)) \cong \mathbf{L}_{\mathcal{O}_X} E = E$, which shows that $T \circ T \cong \mathrm{id}_{\mathcal{K}u(X)}$, i.e., T is an involution on $\mathcal{K}u(X)$. \square

Remark 3.3. It is easy to see that the induced action of T on $\mathcal{N}(\mathcal{K}u(X))$ will fix Λ_1 and map Λ_2 to $-\Lambda_2$.

4. STABILITY CONDITIONS ON KUZNETSOV COMPONENTS OF GUSHEL-MUKAI FOURFOLDS

In this section, we recall (weak) Bridgeland stability conditions on a triangulated category and the notion of stability conditions on the Kuznetsov component of an ordinary GM fourfold. We follow from [BLMS17, Section 2].

4.1. Weak stability conditions. Let \mathcal{D} be a triangulated category and $K_0(\mathcal{D})$ be its Grothendieck group. Fix a surjective morphism to a finite rank lattice $v : K_0(\mathcal{D}) \rightarrow \Lambda$.

Definition 4.1. The *heart of a bounded t -structure* on \mathcal{D} is an abelian subcategory $\mathcal{A} \subset \mathcal{D}$ such that the following conditions are satisfied

- (1) for any $E, F \in \mathcal{A}$ and $n < 0$, we have $\mathrm{Hom}(E, F[n]) = 0$,
- (2) for any object $E \in \mathcal{D}$, there exists a sequence of morphisms

$$0 = E_0 \xrightarrow{\phi_1} E_1 \xrightarrow{\phi_2} \dots \xrightarrow{\phi_m} E_m = E$$

such that $\mathrm{cone}(\phi_i)$ is in the form $A_i[k_i]$, for some sequence $k_1 > k_2 > \dots > k_m$ of integers and $A_i \in \mathcal{A}$.

Definition 4.2. Let \mathcal{A} be an abelian category and $Z : K_0(\mathcal{A}) \rightarrow \mathbb{C}$ be a group homomorphism such that for any $E \in \mathcal{A}$ we have $\text{Im } Z(E) \geq 0$ and if $\text{Im } Z(E) = 0$ then $\text{Re } Z(E) \leq 0$. Then we call Z a *weak stability function* on \mathcal{A} . Furthermore, for any $0 \neq E \in \mathcal{A}$, if we have $\text{Im } Z(E) \geq 0$ and $\text{Im } Z(E) = 0$ implies that $\text{Re } Z(E) < 0$, then we call Z a *stability function* on \mathcal{A} .

Definition 4.3. A *weak stability condition* on \mathcal{D} is a pair $\sigma = (\mathcal{A}, Z)$, where \mathcal{A} is the heart of a bounded t-structure on \mathcal{D} and $Z : \Lambda \rightarrow \mathbb{C}$ is a group homomorphism such that

- (1) the composition $Z \circ v : K_0(\mathcal{A}) \cong K_0(\mathcal{D}) \rightarrow \mathbb{C}$ is a weak stability function on \mathcal{A} .
From now on, we write $Z(E)$ rather than $Z(v(E))$.

Much like the slope from classical μ -stability, we can define a *slope* μ_σ for σ using Z . For any $E \in \mathcal{A}$, set

$$\mu_\sigma(E) := \begin{cases} -\frac{\text{Re } Z(E)}{\text{Im } Z(E)}, & \text{Im } Z(E) > 0 \\ +\infty, & \text{else.} \end{cases}$$

We say an object $0 \neq E \in \mathcal{A}$ is σ -(semi)stable if $\mu_\sigma(F) < \mu_\sigma(E)$ (respectively $\mu_\sigma(F) \leq \mu_\sigma(E)$) for any proper subobject $F \subset E$.

- (2) Any object $E \in \mathcal{A}$ has a Harder-Narasimhan filtration in terms of σ -semistability defined above.
- (3) There exists a quadratic form Q on $\Lambda \otimes \mathbb{R}$ such that $Q|_{\ker Z}$ is negative definite and $Q(E) \geq 0$ for all σ -semistable objects $E \in \mathcal{A}$. This is known as the *support property*.

If the composition $Z \circ v$ is a stability function, then σ is a *stability condition* on \mathcal{D} .

4.2. Stability conditions on Kuznetsov components of Gushel-Mukai fourfolds.

In [PPZ19], inspired by the idea of constructing stability conditions on Kuznetsov components of cubic fourfolds in [BLMS17], the authors embed the Kuznetsov component $\mathcal{Ku}(X)$ of a general ordinary GM fourfold X into a twisted derived category of modules over a quadric threefold, associated to a conic fibration of X . In particular, they prove the following theorem.

Theorem 4.4. *Let X be a GM fourfold over an algebraically closed field k with $\text{char}(k) = 0$ or $\text{char}(k) > 5$. Then the category $\mathcal{Ku}(X)$ has a stability condition.*

We prove some simple properties for stability conditions on $\mathcal{Ku}(X)$.

Proposition 4.5. *Let X be a non-Hodge-special GM fourfold and σ be a stability condition on $\mathcal{Ku}(X)$ and \mathcal{A} be its heart. Then*

- (1) the homological dimension of \mathcal{A} is 2.
- (2) If X is a very general GM fourfold, then we have $\text{ext}^1(A, A) \geq 4$ for any non-trivial object $A \in \mathcal{A}$.

Proof. Let $A, B \in \mathcal{A}$, then $\text{Hom}(A, B[i]) \cong \text{Hom}(B[i], A[2]) \cong \text{Hom}(B, A[2-i]) = 0$ for $i \geq 3$. Thus (1) holds. Let A be a non-trivial object in \mathcal{A} , then by (1),

$$\chi(A, A) = \text{hom}(A, A) - \text{ext}^1(A, A) + \text{ext}^2(A, A).$$

Note that $\chi(A, A) \leq -2$. Then $\text{ext}^1(A, A) = 2\text{hom}(A, A) - \chi(A, A) \geq 4$, which proves (2). \square

Since the Serre functor $S_{\mathcal{Ku}(X)} \cong [2]$, by the same argument as in [BB17, Lemma 2.4], we have

Lemma 4.6. (*Weak Mukai Lemma*) *Let $A \rightarrow E \rightarrow B$ be a triangle in $\mathcal{Ku}(X)$ with $\text{Hom}(A, B) = 0$, then we have*

$$\text{ext}^1(A, A) + \text{ext}^1(B, B) \leq \text{ext}^1(E, E).$$

Proposition 4.7. *Let X be a non-Hodge-special GM fourfold. If $E \in \mathcal{K}u(X)$ is an object with $\chi(E, E) = -2$ and $\text{ext}^1(E, E) \leq 9$, then E is stable with respect to every stability condition σ on $\mathcal{K}u(X)$.*

Proof. The proof is very similar to the argument in [Zha20, Section 4.2]. Firstly, we show that up to a shift, E is in the heart \mathcal{A} of the stability condition σ . To this end, we apply the spectral sequence for objects in $\mathcal{K}u(X)$, whose second page

$$E_2^{p,q} = \bigoplus_i \text{Hom}^p(\mathcal{H}^i(E), \mathcal{H}^{i+q}(E)) \implies \text{Hom}^{p+q}(E, E),$$

where the cohomology is taken with respect to the heart \mathcal{A} . By Proposition 4.5, the homological dimension of \mathcal{A} is 2, it follows that $E_\infty^{1,q} = E_2^{1,q}$, so if we take $q = 0$,

$$9 \geq \text{ext}^1(E, E) = \dim\left(\bigoplus_{p+q=1} E_\infty^{p,q}\right) \geq \dim(E_\infty^{1,0}) = \sum_i \text{ext}^1(\mathcal{H}^i(E), \mathcal{H}^i(E)) \geq 4r,$$

where $r > 0$ is the number of non-zero cohomology objects of E . Then $r = 1$ or $r = 2$. If $r = 1$, we are done and $E \in \mathcal{A}$ up to a shift. If $r = 2$, let M and N be two non-zero cohomology objects of E in \mathcal{A} . Then we have $\text{ext}^1(M, M) + \text{ext}^1(N, N) = 8$. Indeed, $\text{ext}^1(M, M) \geq 4$ and $\text{ext}^1(N, N) \geq 4$ by Proposition 4.5, on the other hand, $\text{ext}^1(E, E) \leq 9$ implies that $\text{ext}^1(M, M) = 4$ and $\text{ext}^1(N, N) = 4$ since both of them must be even numbers. Then

$$\text{hom}(M, M) = \text{hom}(N, N) = 1, \quad \chi(M, M) = \chi(N, N) = -2,$$

which implies $[M], [N] \in \{\pm\Lambda_1, \pm\Lambda_2\}$. But as a class in $\mathcal{N}(\mathcal{K}u(X))$, $[E]$ is in the form $[E] = (-1)^i \cdot [M] + (-1)^j \cdot [N]$ for some $i, j \neq 0$, which is impossible since $[E] \in \{\pm\Lambda_1, \pm\Lambda_2\}$ as well. Then $r = 1$ and E is in the heart \mathcal{A} up to a shift. Now we show E is σ -stable. The argument is almost the same as [PY20, Lemma 5.13] and [Zha20, Lemma 4.16]. We assume that E is unstable, then there is a destabilizing triangle in the heart \mathcal{A}

$$A \rightarrow E \rightarrow B$$

with both A and B are σ -semistable objects and the phase $\phi(A)$ is greater than $\phi(B)$. Then $\text{Hom}(A, B) = 0$. Thus Mukai Lemma 4.6 applies and

$$\text{ext}^1(A, A) + \text{ext}^1(B, B) \leq \text{ext}^1(E, E) \leq 9.$$

But now $E \in \mathcal{A}$ and $\text{ext}^1(E, E)$ can only be even. On the other hand,

$$\text{ext}^1(A, A) + \text{ext}^1(B, B) \geq 8,$$

so $\text{ext}^1(E, E) = 8$ and $\text{ext}^1(A, A) = \text{ext}^1(B, B) = 4$. By the same reason as above, $[A] + [B]$ can never equal to $[E]$ in $\mathcal{N}(\mathcal{K}u(X))$. Thus E is σ -semistable. It remains to show that E is not σ -strictly semistable while the argument is almost the same as in [PY20, Lemma 5.13]. \square

5. CONICS ON GUSHEL-MUKAI FOURFOLDS

In [IM11], the moduli space of conics $F_g(X)$ on a general ordinary GM fourfold X plays an important role in the construction of the hyperkähler variety \tilde{Y}_{A^\perp} . In this section, we present some basic properties of conics in $F_g(X)$ and list some calculation results which heavily rely on the geometry of X .

According to [DIM15], there are two types of planes in $\text{Gr}(2, V_5)$ which are called σ -planes and ρ -planes. A σ -plane is in the form $\mathbb{P}(V_1 \wedge V_4)$ and a ρ -plane is in the form $\mathbb{P}(\wedge^2 V_3)$ for some subspaces V_1, V_3 and V_4 in V_5 .

Definition 5.1 ([IM11, Section 3.1]). Let X be an ordinary GM fourfold, there are three types of conics

- (1) τ -conics are conics spanning planes which are not contained in $\text{Gr}(2, V_5)$.

- (2) σ -conics are conics parametrizing lines passing through a common point, i.e., they span σ -planes.
- (3) ρ -conics are conics parametrizing lines contained in a common plane, i.e., they span ρ -planes.

Lemma 5.2 ([IM11, Section 3.2]). *Let X be a general ordinary GM fourfold and \mathcal{U}, \mathcal{Q} be the tautological sub and the quotient bundle on X . For a smooth conic C on X , we have*

- (1) $\mathcal{U}^\vee|_C \cong \mathcal{O}_C(1) \oplus \mathcal{O}_C(1)$ and $\mathcal{Q}|_C \cong \mathcal{O}_C(1) \oplus \mathcal{O}_C(1) \oplus \mathcal{O}_C$ if C is a τ -conic;
- (2) $\mathcal{U}^\vee|_C \cong \mathcal{O}_C(2) \oplus \mathcal{O}_C$ and $\mathcal{Q}|_C \cong \mathcal{O}_C(1) \oplus \mathcal{O}_C(1) \oplus \mathcal{O}_C$ if C is a σ -conic;
- (3) $\mathcal{U}^\vee|_C \cong \mathcal{O}_C(1) \oplus \mathcal{O}_C(1)$ and $\mathcal{Q}|_C \cong \mathcal{O}_C(2) \oplus \mathcal{O}_C \oplus \mathcal{O}_C$ if C is a ρ -conic.

In fact, the type of a conic on X is detected by the numbers $\text{hom}(\mathcal{U}, I_C)$ and $\text{hom}(\mathcal{Q}^\vee, I_C)$.

Lemma 5.3. *Let X be a general ordinary GM fourfold and C be a conic on X .*

- (1) *If C is a τ -conic, then $\text{Hom}(\mathcal{U}, I_C) = k$ and $\text{Hom}(\mathcal{Q}^\vee, I_C) = 0$.*
- (2) *If C is a σ -conic, then $\text{Hom}(\mathcal{U}, I_C) = k$ and $\text{Hom}(\mathcal{Q}^\vee, I_C) = k$.*
- (3) *If C is a ρ -conic, then $\text{Hom}(\mathcal{U}, I_C) = k^2$ and $\text{Hom}(\mathcal{Q}^\vee, I_C) = 0$.*

Proof. Note that for a conic C , if $\text{Hom}(\mathcal{U}, I_C) = k^a$, for some integer $a \geq 0$, then C is contained in $\text{Gr}(2, V_{5-a}) \cap X$. Since for any conic C , there is some V_4 such that C lies in $\text{Gr}(2, V_4)$, then we have $\text{hom}(\mathcal{U}, I_C) \geq 1$. Now if $\text{hom}(\mathcal{U}, I_C) \geq 2$, we know that C is contained in a ρ -plane $\text{Gr}(2, V_3)$. For a τ -conic C , $\langle C \rangle$ is not contained in $\text{Gr}(2, V_4)$ for any $V_4 \subset V_5$ and a σ -conic C generates a σ -plane $\mathbb{P}(V_1 \wedge V_4)$. Thus for such two types of conics, we have $\text{Hom}(\mathcal{U}, I_C) = k$. For a ρ -conic C , since $\langle C \rangle$ is in the form $\text{Gr}(2, V_3)$, we have $\text{hom}(\mathcal{U}, I_C) \geq 2$. But if $\text{hom}(\mathcal{U}, I_C) \geq 3$, we know that $C \subset \text{Gr}(2, V_2)$, which is impossible. Hence for a ρ -conic C , we have $\text{Hom}(\mathcal{U}, I_C) = k^2$.

On the other hand, if $\text{Hom}(\mathcal{Q}^\vee, I_C) = k^b$ for an integer $b \geq 0$, C is contained in $\text{Gr}(2-b, V_{5-b}) \cap X$. Thus we have $\text{hom}(\mathcal{Q}^\vee, I_C) \leq 1$ for any conic C . It is easy to see $\text{hom}(\mathcal{Q}^\vee, I_C) = 1$ if and only if C is contained in the zero locus of a global section of \mathcal{Q} , which is a σ -3-plane $\mathbb{P}(V_1 \wedge V_5)$ of $\text{Gr}(2, V_5)$. This implies that $\text{Hom}(\mathcal{Q}^\vee, I_C) = 0$ for conics of type τ or ρ and $\text{Hom}(\mathcal{Q}^\vee, I_C) = k$ for conics of type σ . \square

Now we state two lemmas, which are useful in the following sections.

Lemma 5.4. *Let X be an ordinary GM fourfold, then we have*

- (1) $\text{RHom}(\mathcal{U}, \mathcal{Q}^\vee) = k$.
- (2) $\text{RHom}(\mathcal{U}^\vee, \mathcal{Q}) = 0$.
- (3) $\text{RHom}(\mathcal{U}^\vee, \mathcal{U}) = 0$.
- (4) $\text{RHom}(\mathcal{Q}^\vee, \mathcal{Q}(-H)) = k[-2]$.
- (5) $\text{RHom}(\mathcal{U}^\vee, \mathcal{Q}^\vee) = k[-1]$.

Proof. The Koszul resolution of an ordinary GM fourfold is in the form

$$0 \rightarrow \mathcal{O}_{\text{Gr}(2, V_5)}(-3) \rightarrow \mathcal{O}_{\text{Gr}(2, V_5)}(-2) \oplus \mathcal{O}_{\text{Gr}(2, V_5)}(-1) \rightarrow \mathcal{O}_{\text{Gr}(2, V_5)} \rightarrow \mathcal{O}_X \rightarrow 0.$$

Then the result follows from this resolution and applying the Borel-Weil-Bott theorem. \square

Lemma 5.5. *Let X be a general ordinary GM fourfold and $C \subset X$ be a conic. Then we have*

- (1) $\text{RHom}(\mathcal{O}_X(H), \mathcal{O}_C) = k[-1]$.
- (2) $\text{RHom}(\mathcal{U}^\vee, \mathcal{O}_C) = 0$ when C is of type τ or ρ .
- (3) $\text{RHom}(\mathcal{U}^\vee, \mathcal{O}_C) = k \oplus k[-1]$ when C is a σ -conic.
- (4) $\text{RHom}(\mathcal{U}^\vee(H), \mathcal{O}_C) = k^4[-1]$.
- (5) $\text{RHom}(\mathcal{Q}^\vee(H), \mathcal{O}_C) = k[-1]$ when C is of type τ or σ .
- (6) $\text{RHom}(\mathcal{Q}^\vee(H), \mathcal{O}_C) = k \oplus k^2[-1]$ when C is of type ρ .

Proof. When C is smooth, the result follows from $C \cong \mathbb{P}^1$ and Lemma 5.2.

When C is not smooth but reduced, we have an exact sequence

$$0 \rightarrow \mathcal{O}_C \rightarrow \mathcal{O}_{l_1} \oplus \mathcal{O}_{l_2} \rightarrow \mathcal{O}_x \rightarrow 0,$$

where l_i are lines such that $l_1 \cup l_2 = C$, $l_1 \cap l_2 = x$. When C is non-reduced, we have

$$0 \rightarrow \mathcal{O}_l(-H) \rightarrow \mathcal{O}_C \rightarrow \mathcal{O}_l \rightarrow 0,$$

where $l = C_{red}$ is a line. Then the results follow from applying the Hom-functor to two exact sequences above, taking long exact sequences and applying Lemma 5.3. \square

Finally, we introduce a notion called a σ -quadric surface, denoted by q . As in [DIM15, Section 3], q is defined by a σ -3-plane $\mathbb{P}(V_1^\omega \wedge V_5)$ in $\text{Gr}(2, V_5)$ intersecting with X . Indeed, $\langle q \rangle = \mathbb{P}(V_1^\omega \wedge V_5)$, i.e., q is the zero locus of a section of \mathcal{Q} . Moreover, when X is general, q is smooth and is the only quadric surface contained in X due to [PPZ19, Remark 2.2] and [DIM15].

6. PROJECTION OBJECTS OF CONICS

In this section, we find out the objects $\text{pr}_1(\mathcal{O}_C(H))$ and $\text{pr}_2(I_C)$ for any conic C on a general ordinary GM fourfold, then we relate them via the involution T defined in Section 3.

Recall that $\text{pr}_1 := \mathbf{L}_{\mathcal{O}_X} \mathbf{L}_{\mathcal{U}^\vee} \mathbf{L}_{\mathcal{O}_X(H)} \mathbf{L}_{\mathcal{U}^\vee(H)}$ and $\text{pr}_2 := \mathbf{R}_{\mathcal{U}} \mathbf{R}_{\mathcal{O}_X(-H)} \mathbf{L}_{\mathcal{O}_X} \mathbf{L}_{\mathcal{U}^\vee}$. The characters $\text{ch}(\text{pr}_1(\mathcal{O}_C(H)))$ and $\text{ch}(\text{pr}_2(I_C))$ of a conic C are Λ_1 . We start with two lemmas.

Lemma 6.1. *Let X be an ordinary GM fourfold and C be a conic on X . Let $j : Y \hookrightarrow X$ be any hyperplane section containing C . Then we have*

$$\text{pr}_2(I_C) \cong \text{pr}_2(I_{C/Y}).$$

Proof. Note that we have an exact sequence

$$0 \rightarrow \mathcal{O}_X(-H) \rightarrow I_C \rightarrow I_{C/Y} \rightarrow 0.$$

Since $\mathbf{L}_{\mathcal{O}_X} \mathbf{L}_{\mathcal{U}^\vee}(\mathcal{O}_X(-H)) = \mathcal{O}_X(-H)$, thus we have

$$\text{pr}_2(\mathcal{O}_X(-H)) = \mathbf{R}_{\mathcal{U}} \mathbf{R}_{\mathcal{O}_X(-H)} \mathbf{L}_{\mathcal{O}_X} \mathbf{L}_{\mathcal{U}^\vee}(\mathcal{O}_X(-H)) = 0.$$

Therefore we obtain $\text{pr}_2(I_C) \cong \text{pr}_2(I_{C/Y})$. \square

Lemma 6.2. *Let X be a general ordinary GM fourfold and C be a conic on X . Then we have*

$$T(\text{pr}_2(I_C)) \cong \text{pr}_1(\mathcal{O}_C(H))[-2].$$

Proof. Note that we have

$$T(\text{pr}_2(I_C)) = \mathbf{L}_{\mathcal{O}_X} \mathbf{L}_{\mathcal{U}^\vee} \mathbf{L}_{\mathcal{O}_X(H)} \mathbf{R}_{\mathcal{O}_X} \mathbf{R}_{\mathcal{U}}(\mathbb{D}(I_C)).$$

Applying $\mathbf{R}_{\mathcal{O}_X} \mathbf{R}_{\mathcal{U}}(\mathbb{D}(-))$ to the standard exact sequence of C , we obtain

$$\mathbf{R}_{\mathcal{O}_X} \mathbf{R}_{\mathcal{U}}(\mathbb{D}(I_C)) \cong \mathbf{R}_{\mathcal{O}_X} \mathbf{R}_{\mathcal{U}}(\mathbb{D}(\mathcal{O}_C))[1],$$

which means

$$T(\text{pr}_2(I_C)) \cong \mathbf{L}_{\mathcal{O}_X} \mathbf{L}_{\mathcal{U}^\vee} \mathbf{L}_{\mathcal{O}_X(H)} \mathbf{R}_{\mathcal{O}_X} \mathbf{R}_{\mathcal{U}}(\mathbb{D}(\mathcal{O}_C))[1].$$

Firstly, we deal with τ -conics and ρ -conics. According to Lemma 5.5 we have

$$\begin{aligned} \mathbf{L}_{\mathcal{O}_X(H)} \mathbf{L}_{\mathcal{U}^\vee(H)}(\mathcal{O}_C(H)) &= I_C(H)[1], \\ \mathbf{R}_{\mathcal{O}_X} \mathbf{R}_{\mathcal{U}}(\mathcal{O}_C(H)) &= \mathbf{R}_{\mathcal{O}_X}(\mathcal{O}_C(H)) \end{aligned}$$

and a triangle

$$\mathbf{R}_{\mathcal{O}_X} \mathbf{R}_{\mathcal{U}}(\mathcal{O}_C(H)) \rightarrow \mathcal{O}_C(H) \rightarrow \mathcal{O}_X[3].$$

Applying $\mathbf{L}_{\mathcal{O}_X} \mathbf{L}_{\mathcal{U}^\vee} \mathbf{L}_{\mathcal{O}_X(H)}$ to this triangle, we obtain that

$$\mathbf{L}_{\mathcal{O}_X} \mathbf{L}_{\mathcal{U}^\vee} \mathbf{L}_{\mathcal{O}_X(H)} \mathbf{R}_{\mathcal{O}_X} \mathbf{R}_{\mathcal{U}}(\mathcal{O}_C(H)) \cong \mathbf{L}_{\mathcal{O}_X} \mathbf{L}_{\mathcal{U}^\vee} \mathbf{L}_{\mathcal{O}_X(H)}(\mathcal{O}_C(H)) \cong \mathbf{L}_{\mathcal{O}_X} \mathbf{L}_{\mathcal{U}^\vee}(I_C(H)[1]).$$

The last complex is just $\mathrm{pr}_1(\mathcal{O}_C(H))$. In order to establish $T(\mathrm{pr}_2(I_C)) \cong \mathrm{pr}_1(\mathcal{O}_C(H))[-2]$, we only need to show that $\mathbb{D}(\mathcal{O}_C) \cong \mathcal{O}_C(H)[-3]$.

- Let C be a smooth conic of type τ or ρ . By Grothendieck-Verdier duality, we have $\mathbb{D}(\mathcal{O}_C) \cong \mathcal{O}_C(H)[-3]$.
- Let C be a non-smooth reduced conic of type τ or ρ . We have the exact sequence

$$0 \rightarrow \mathcal{O}_C \rightarrow \mathcal{O}_{l_1} \oplus \mathcal{O}_{l_2} \rightarrow \mathcal{O}_x \rightarrow 0$$

where l_i are lines such that $l_1 \cup l_2 = C$, $l_1 \cap l_2 = x$. By Grothendieck-Verdier duality, we have $\mathbb{D}(\mathcal{O}_{l_i}) \cong \mathcal{O}_{l_i}[-3]$ and $\mathbb{D}(\mathcal{O}_x) \cong \mathcal{O}_x[-4]$. Thus applying \mathbb{D} to the exact sequence above, we obtain a triangle

$$\mathcal{O}_x[-4] \rightarrow (\mathcal{O}_{l_1} \oplus \mathcal{O}_{l_2})[-3] \rightarrow \mathbb{D}(\mathcal{O}_C).$$

It means we have an exact sequence

$$0 \rightarrow \mathcal{O}_{l_1} \oplus \mathcal{O}_{l_2} \rightarrow \mathbb{D}(\mathcal{O}_C)[3] \rightarrow \mathcal{O}_x \rightarrow 0,$$

then we obtain $\mathbb{D}(\mathcal{O}_C) \cong \mathcal{O}_C(H)[-3]$.

- Let C be a double line of type τ or ρ . We have

$$0 \rightarrow \mathcal{O}_l(-H) \rightarrow \mathcal{O}_C \rightarrow \mathcal{O}_l \rightarrow 0,$$

where $l = C_{red}$ is a line. Applying \mathbb{D} to this exact sequence, we have a triangle

$$\mathcal{O}_l[-3] \rightarrow \mathbb{D}(\mathcal{O}_C) \rightarrow \mathcal{O}_l(H)[-3],$$

then we obtain $\mathbb{D}(\mathcal{O}_C) \cong \mathcal{O}_C(H)[-3]$.

When C is a σ -conic, the computation is similar. We omit details here. \square

Next, we compute projection objects of all three types of conics to the Kuznetsov component $\mathcal{K}u(X)$.

6.1. τ -conic. In this subsection, we compute the projection objects of τ -conics.

Proposition 6.3. *Let X be a general ordinary GM fourfold and C be a τ -conic on X . Then we have*

$$\mathrm{pr}_1(\mathcal{O}_C(H)) \cong \mathbf{L}_{\mathcal{O}_X}(I_{C/\Sigma}(H))[1],$$

where Σ is the zero locus of a section of \mathcal{U}^\vee containing C . Moreover, there is an exact sequence

$$0 \rightarrow \mathcal{U}^{\oplus 4} \rightarrow K_1 \rightarrow \mathrm{pr}_2(I_C) \rightarrow I_C \rightarrow 0,$$

where $K_1 := \mathrm{cok}(\mathcal{O}_X(-H) \hookrightarrow \mathcal{U}^{\oplus 5}) = \mathbf{R}_{\mathcal{U}}\mathcal{O}_X(-H)[1]$.

Proof. By definition of pr_1 , we have

$$\begin{aligned} \mathrm{pr}_1(\mathcal{O}_C(H)) &= \mathbf{L}_{\mathcal{O}_X} \mathbf{L}_{\mathcal{U}^\vee} \mathbf{L}_{\mathcal{O}_X(H)} \mathbf{L}_{\mathcal{U}^\vee(H)}(\mathcal{O}_C(H)) \\ &\cong \mathbf{L}_{\mathcal{O}_X} \mathbf{L}_{\mathcal{U}^\vee} \mathbf{L}_{\mathcal{O}_X(H)}(\mathcal{O}_C(H)) \\ &\cong \mathbf{L}_{\mathcal{O}_X} \mathbf{L}_{\mathcal{U}^\vee}(I_C(H))[1]. \end{aligned}$$

The first isomorphism follows from Lemma 5.5. The second isomorphism follows from the standard exact sequence associated with C . Next, we have an exact triangle

$$\mathrm{RHom}(\mathcal{U}^\vee, I_C(H)) \otimes \mathcal{U}^\vee \rightarrow I_C(H) \rightarrow \mathbf{L}_{\mathcal{U}^\vee}(I_C(H)).$$

Note that $\mathrm{RHom}(\mathcal{U}^\vee, I_C(H)) \cong \mathrm{RHom}(\mathcal{U}, I_C)$. By Lemma 5.3, $\mathrm{RHom}(\mathcal{U}, I_C) = k[0]$. Then the object $\mathbf{L}_{\mathcal{U}^\vee}(I_C(H))$ fits into the triangle

$$\mathcal{U}^\vee \rightarrow I_C(H) \rightarrow \mathbf{L}_{\mathcal{U}^\vee}(I_C(H)).$$

The image of $\pi : \mathcal{U} \rightarrow I_C$ is the ideal sheaf I_Σ , where Σ is the zero locus of a section of \mathcal{U}^\vee containing C , which is a surface $\Sigma = \mathrm{Gr}(2, V_4) \cap X$ for some V_4 . Then we have two short exact sequences

$$0 \rightarrow \ker \pi \rightarrow \mathcal{U} \rightarrow I_\Sigma \rightarrow 0$$

and

$$0 \rightarrow I_\Sigma \rightarrow I_C \rightarrow I_{C/\Sigma} \rightarrow 0.$$

Note that $\ker \pi$ is a rank one reflexive sheaf on X , hence is a line bundle. Thus we have $\ker \pi \cong \mathcal{O}_X(-H)$ and there is an exact triangle

$$\mathcal{O}_X(-H)[2] \rightarrow \mathbf{L}\mathcal{U}(I_C)[1] \rightarrow I_{C/\Sigma}[1].$$

Tensoring with $\mathcal{O}_X(H)$, we get

$$\mathcal{O}_X[2] \rightarrow \mathbf{L}\mathcal{U}^\vee(I_C(H))[1] \rightarrow I_{C/\Sigma}(H)[1].$$

Finally, applying $\mathbf{L}\mathcal{O}_X$ to this triangle, we get $\mathrm{pr}_1(\mathcal{O}_C(H)) \cong \mathbf{L}\mathcal{O}_X(I_{C/\Sigma}(H))[1]$.

Now we compute $\mathrm{pr}_2(I_C)$. Since $\mathrm{RHom}(\mathcal{O}_X, I_C) = \mathrm{RHom}(\mathcal{U}^\vee, I_C) = 0$ by Lemma 5.5, we have

$$\mathrm{pr}_2(I_C) = \mathbf{R}\mathcal{U}\mathbf{R}_{\mathcal{O}_X(-H)}(I_C).$$

Since we have $\mathrm{RHom}(I_C, \mathcal{O}_X(-H)) = k[-2]$ and $\mathrm{RHom}(I_C, \mathcal{U}) = k^4[-2]$, then we obtain triangles

$$\mathbf{R}_{\mathcal{O}_X(-H)}(I_C) \rightarrow I_C \rightarrow \mathcal{O}_X(-H)[2],$$

$$\mathbf{R}\mathcal{U}(I_C) \rightarrow I_C \rightarrow \mathcal{U}^{\oplus 4}[2]$$

and

$$\mathbf{R}\mathcal{U}\mathbf{R}_{\mathcal{O}_X(-H)}I_C \rightarrow \mathbf{R}\mathcal{U}I_C \rightarrow K_1[1],$$

where $K_1 := \mathrm{cok}(\mathcal{O}_X(-H) \hookrightarrow \mathcal{U}^{\oplus 5}) = \mathbf{R}\mathcal{U}\mathcal{O}_X(-H)[1]$. Therefore, taking the long exact sequence of cohomology, we get

$$0 \rightarrow \mathcal{H}^{-1}(\mathrm{pr}_2(I_C)) \rightarrow \mathcal{U}^{\oplus 4} \xrightarrow{\alpha} K_1 \rightarrow \mathcal{H}^0(\mathrm{pr}_2(I_C)) \rightarrow I_C \rightarrow 0. \quad (4)$$

Now from the definition of α , we have a commutative diagram

$$\begin{array}{ccccc} \mathrm{Hom}(I_C[-2], \mathcal{U})^\vee \otimes \mathcal{U}[1] & \xrightarrow{f_2} & \mathbf{R}\mathcal{U}I_C & \longrightarrow & I_C \\ \downarrow f_1 & & \downarrow v' & & \downarrow v \\ \mathrm{Hom}(\mathcal{O}_X(-H), \mathcal{U})^\vee \otimes \mathcal{U}[1] & \longrightarrow & \mathbf{R}\mathcal{U}\mathcal{O}_X(-H)[2] & \longrightarrow & \mathcal{O}_X(-H)[2] \end{array}.$$

Here $v' \circ f_2 = \alpha[1]$ and rows are induced by the definition of functor $\mathbf{R}\mathcal{U}$. Let v be a non-zero element in $\mathrm{Hom}(I_C, \mathcal{O}_X(-H)[2]) = k$, v' and f_1 be the morphisms induced by v and the right mutation functor. To determine f_1 , we only need to determine the natural map $f_3 : \mathrm{Hom}(I_C[-2], \mathcal{U})^\vee \rightarrow \mathrm{Hom}(\mathcal{O}_X(-H), \mathcal{U})^\vee$ induced by v due to the fact $f_3 \otimes \mathrm{id}_{\mathcal{U}[1]} = f_1$. To this end, using Serre duality and the adjunction of mutations, f_3 is naturally isomorphic to $f'_3 : \mathrm{Hom}(\mathcal{U}^\vee(H), I_C[2]) \rightarrow \mathrm{Hom}(\mathcal{O}_X(H)^{\oplus 5}, I_C[2])$, which is induced by the triangle

$$\mathrm{RHom}(\mathcal{O}_X(H), \mathcal{U}^\vee(H)) \otimes \mathcal{O}_X(H) \rightarrow \mathcal{U}^\vee(H) \rightarrow \mathbf{L}_{\mathcal{O}_X(H)}\mathcal{U}^\vee(H).$$

Hence f'_3 fits into an exact sequence

$$0 \rightarrow \mathrm{Hom}(\mathbf{L}_{\mathcal{O}_X(H)}\mathcal{U}^\vee(H), I_C[2]) \rightarrow \mathrm{Hom}(\mathcal{U}^\vee(H), I_C[2]) \xrightarrow{f'_3} \mathrm{Hom}(\mathcal{O}_X(H)^{\oplus 5}, I_C[2]).$$

Since $\mathbf{L}_{\mathcal{O}_X(H)}\mathcal{U}^\vee(H) \cong \mathcal{Q}^\vee(H)[1]$, we know that $\ker(f'_3) \cong \mathrm{Hom}(\mathcal{Q}^\vee(H), \mathcal{O}_C) = 0$ by Lemma 5.5. Thus $\ker(\alpha) = 0$, $\mathcal{H}^{-1}(\mathrm{pr}_2(I_C)) = 0$ and the sequence of (4) becomes

$$0 \rightarrow \mathcal{U}^{\oplus 4} \xrightarrow{\alpha} K_1 \rightarrow \mathrm{pr}_2(I_C) \rightarrow I_C \rightarrow 0.$$

□

6.2. Projection of \mathcal{U} . In this subsection, we find out the projection object of \mathcal{U} .

Proposition 6.4. *Let X be a general ordinary GM fourfold. Then we have a triangle*

$$\mathcal{U} \rightarrow \mathrm{pr}_1(\mathcal{U}) \rightarrow K_2[-1],$$

where $K_2 := \mathbf{L}_{\mathcal{O}_X}(I_q(H))[-1]$ is a μ -stable reflexive sheaf, q is the unique σ -quadric in X .

Proof. We apply the first mutation $\mathbf{L}_{\mathcal{U}^\vee(H)}$ to \mathcal{U} ,

$$\mathrm{RHom}(\mathcal{U}^\vee(H), \mathcal{U}) \otimes \mathcal{U}^\vee(H) \rightarrow \mathcal{U} \rightarrow \mathbf{L}_{\mathcal{U}^\vee(H)}\mathcal{U}.$$

By Serre duality and exceptionality of \mathcal{U} , we have $\mathrm{RHom}(\mathcal{U}^\vee(H), \mathcal{U}) = k[-4]$. Then the triangle becomes

$$\mathcal{U}^\vee(H)[-4] \rightarrow \mathcal{U} \rightarrow \mathbf{L}_{\mathcal{U}^\vee(H)}\mathcal{U}.$$

Applying $\mathbf{L}_{\mathcal{O}_X(H)}$ to this triangle, we get

$$(\mathbf{L}_{\mathcal{O}_X}\mathcal{U}^\vee) \otimes \mathcal{O}_X(H)[-4] \rightarrow \mathbf{L}_{\mathcal{O}_X(H)}\mathcal{U} \rightarrow \mathbf{L}_{\mathcal{O}_X(H)}\mathbf{L}_{\mathcal{U}^\vee(H)}\mathcal{U}.$$

By Serre duality, $\mathrm{RHom}(\mathcal{O}_X(H), \mathcal{U}) = 0$, so that $\mathbf{L}_{\mathcal{O}_X(H)}\mathcal{U} \cong \mathcal{U}$. Then we have the triangle

$$\mathcal{Q}^\vee(H)[-3] \rightarrow \mathcal{U} \rightarrow \mathbf{L}_{\mathcal{O}_X(H)}\mathbf{L}_{\mathcal{U}^\vee(H)}\mathcal{U}.$$

Applying $\mathbf{L}_{\mathcal{U}^\vee}$, we obtain

$$(\mathbf{L}_{\mathcal{U}}\mathcal{Q}^\vee)(H)[-3] \rightarrow \mathbf{L}_{\mathcal{U}^\vee}\mathcal{U} \rightarrow \mathbf{L}_{\mathcal{U}^\vee}\mathbf{L}_{\mathcal{O}_X(H)}\mathbf{L}_{\mathcal{U}^\vee(H)}\mathcal{U}.$$

Since $\mathrm{RHom}(\mathcal{U}^\vee, \mathcal{U}) = 0$ by Lemma 5.4, we have the following triangle

$$(\mathbf{L}_{\mathcal{U}}\mathcal{Q}^\vee)(H)[-3] \rightarrow \mathcal{U} \rightarrow \mathbf{L}_{\mathcal{U}^\vee}\mathbf{L}_{\mathcal{O}_X(H)}\mathbf{L}_{\mathcal{U}^\vee(H)}\mathcal{U}.$$

Now from the fact $\mathrm{RHom}(\mathcal{U}, \mathcal{Q}^\vee) = k[0]$, there is a triangle

$$\mathcal{U} \rightarrow \mathcal{Q}^\vee \rightarrow \mathbf{L}_{\mathcal{U}}\mathcal{Q}^\vee.$$

Since \mathcal{U} and \mathcal{Q}^\vee are both μ -stable with slopes $\mu(\mathcal{U}) = -\frac{1}{2}$ and $\mu(\mathcal{Q}^\vee) = -\frac{1}{3}$, the map $\mathcal{U} \xrightarrow{s} \mathcal{Q}^\vee$ is injective and $\mathbf{L}_{\mathcal{U}}\mathcal{Q}^\vee \cong \mathrm{cok}(s) \cong I_q$, where q is the unique σ -quadric surface contained in X (cf. Section 5.1). Then the triangle becomes

$$I_q \otimes \mathcal{O}_X(H)[-3] \rightarrow \mathcal{U} \rightarrow \mathbf{L}_{\mathcal{U}^\vee}\mathbf{L}_{\mathcal{O}_X(H)}\mathbf{L}_{\mathcal{U}^\vee(H)}\mathcal{U}.$$

Applying $\mathbf{L}_{\mathcal{O}_X}$, we get

$$\mathbf{L}_{\mathcal{O}_X}(I_q \otimes \mathcal{O}_X(H))[-3] \rightarrow \mathcal{U} \rightarrow \mathrm{pr}_1(\mathcal{U}).$$

As in Section 5.1, $q = \langle q \rangle \cap X = \langle q \rangle \cap Q$, which means q is cut out by five hyperplane sections of \mathbb{P}^8 . This implies the morphism $t : \mathcal{O}_X^{\oplus 5} \rightarrow I_q(H)$ is surjective and $\mathbf{L}_{\mathcal{O}_X}(I_q(H))[-1] \cong \ker(t)$, which is denoted by K_2 . By [Har80, Proposition 1.1], K_2 is a reflexive sheaf. Finally, the stability of K_2 follows from the poly-stability of $\mathcal{O}_X^{\oplus 5}$ and the fact $\mathrm{RHom}(\mathcal{O}_X, K_2) = 0$. \square

6.3. ρ -conic. In this subsection, we compute the projection objects of ρ -conics.

At first, we offer two lemmas which will be very useful in the proof of Proposition 6.7.

Lemma 6.5. *Let X be an ordinary GM fourfold and F be a μ -semistable sheaf on X with $\mathrm{rk}(F) = 3$, $\mathrm{ch}_1(F) = -H$ and $H \cdot \mathrm{ch}_2(F) = eL$. Then we have $e \leq -1$.*

Proof. By Mayamura's restriction theorem, we can take a general smooth hyperplane section Y such that $F|_Y$ remains μ -semistable. Then $\mathrm{ch}_{\leq 2}(F|_Y) = (3, -H, eL)$. The result follows from [Li15, Proposition 3.2]. \square

Lemma 6.6. *Use the notations as in Proposition 6.4, we have*

- (1) $\mathrm{RHom}(I_q(H), \mathcal{U}) = k[-3]$.
- (2) $\mathrm{RHom}(K_2, \mathcal{U}) = k[-2]$.
- (3) $\mathrm{Ext}^1(I_q(H), I_C) = k$.
- (4) $\mathrm{Hom}(K_2, I_C) = k$.

Proof. For (1), by Serre duality we only need to compute $\mathrm{RHom}(\mathcal{U}^\vee, I_q)$. Then we apply $\mathrm{Hom}(\mathcal{U}^\vee, -)$ to the exact sequence

$$0 \rightarrow \mathcal{U} \rightarrow \mathcal{Q}^\vee \rightarrow I_q \rightarrow 0,$$

the result follows from Lemma 5.4.

For (3), we apply $\mathrm{Hom}(-, I_C)$ to the exact sequence

$$0 \rightarrow \mathcal{U}(H) \rightarrow \mathcal{Q}^\vee(H) \rightarrow I_q(H) \rightarrow 0,$$

since C is a ρ -conic, the result follows from $\mathrm{RHom}(\mathcal{U}^\vee, I_C) = 0$ in Lemma 5.3 and Lemma 5.5.

Now if we apply $\mathrm{Hom}(-, \mathcal{U})$ to the exact sequence

$$0 \rightarrow K_2 \rightarrow \mathcal{O}_X^{\oplus 5} \rightarrow I_q(H) \rightarrow 0,$$

then (2) follows from (1). If we apply $\mathrm{Hom}(-, I_C)$ to the exact sequence above, we have $\mathrm{Hom}(K_2, I_C) \cong \mathrm{Ext}^1(I_q(H), I_C)$, then (4) follows from (3). \square

Proposition 6.7. *Let X be a general ordinary GM fourfold and C be a ρ -conic on X . Then we have*

$$\mathrm{pr}_2(I_C) \cong \mathrm{pr}_1(\mathcal{U})[1].$$

Proof. As in the proof of Proposition 6.3, we have a long exact sequence

$$0 \rightarrow \mathcal{H}^{-1}(\mathrm{pr}_2(I_C)) \rightarrow \mathcal{U}^{\oplus 4} \xrightarrow{\alpha} K_1 \rightarrow \mathcal{H}^0(\mathrm{pr}_2(I_C)) \rightarrow I_C \rightarrow 0. \quad (5)$$

Note that in this case $\mathrm{Hom}(\mathcal{Q}^\vee(H), \mathcal{O}_C) = k$ by Lemma 5.5. Thus

$$\mathcal{H}^{-1}(\mathrm{pr}_2(I_C)) \cong \ker(\alpha) \cong \mathcal{U}.$$

And we have an exact sequence

$$0 \rightarrow K_3 \rightarrow \mathcal{H}^0(\mathrm{pr}_2(I_C)) \rightarrow I_C \rightarrow 0, \quad (6)$$

where $0 \rightarrow \mathcal{U}^{\oplus 3} \rightarrow K_1 \rightarrow K_3 \rightarrow 0$. Note that we have a commutative diagram

$$\begin{array}{ccccccc} & & & 0 & & & \\ & & & \downarrow & & & \\ & & & \mathcal{O}_X(-H) & & & \\ & & & \downarrow & & & \\ 0 & \longrightarrow & \mathcal{U}^{\oplus 3} & \longrightarrow & \mathcal{U}^{\oplus 5} & \longrightarrow & \mathcal{U}^{\oplus 2} \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{U}^{\oplus 3} & \longrightarrow & K_1 & \longrightarrow & K_3 \longrightarrow 0 \\ & & & & \downarrow & & \\ & & & & 0 & & \end{array}$$

Hence K_3 also fits into exact sequences

$$0 \rightarrow \mathcal{O}_X(-H) \rightarrow \mathcal{U}^{\oplus 2} \rightarrow K_3 \rightarrow 0 \quad (7)$$

and

$$0 \rightarrow \mathcal{U} \rightarrow K_3 \rightarrow I_{\Sigma_1} \rightarrow 0, \quad (8)$$

where Σ_1 is the zero locus of a regular section of \mathcal{U}^\vee . In particular, K_3 is torsion free, hence $\mathcal{H}^0(\mathrm{pr}_2(I_C))$ is also torsion free.

First we claim that K_3 is μ -(semi)stable. Indeed, by the stability of \mathcal{U} and I_{Σ_1} , the only possible case is that the maximal destabilizing subsheaf of K_3 is in the form I_W , where W is a closed subscheme containing the surface Σ_1 . It is easy to see $\mathrm{rk}(K_3/I_W) = 2$ and $\mathrm{ch}_1(K_3/I_W) = -H$. Thus $(K_3/I_W)^{\vee\vee}$ is also μ -semistable with rank two and $\mathrm{ch}_1 = -H$.

If we apply $\mathrm{Hom}(-, (K_3/I_W)^{\vee\vee})$ to the sequence (8), we have $\mathrm{Hom}(\mathcal{U}, (K_3/I_W)^{\vee\vee}) \neq 0$. By the stability of \mathcal{U} and $(K_3/I_W)^{\vee\vee}$, we know that $\mathcal{U} \subset (K_3/I_W)^{\vee\vee}$, but this is impossible. This is because \mathcal{U} is locally free and $(K_3/I_W)^{\vee\vee}$ is reflexive, the support of the quotient is of codimension ≤ 1 , which contradicts with the fact $(K_3/I_W)^{\vee\vee}/\mathcal{U}$ is supported in codimension ≥ 2 .

Next we claim that K_3 is reflexive. Indeed, we have a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{U} & \longrightarrow & K_3 & \longrightarrow & I_{\Sigma_1} \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{U} & \longrightarrow & K_3^{\vee\vee} & \xrightarrow{\theta} & \mathcal{O}_X \end{array}$$

and $\mathrm{Im}(\theta) = I_{Z_1}$, where Z_1 is a closed subscheme contained in Σ_1 . If $Z_1 \neq \Sigma_1$, we can assume that $\mathrm{ch}(I_{Z_1}) = 1 - eL + fP$, where $e \geq 0$. In this case $K_3^{\vee\vee}$ is also μ -semistable. But $H \cdot \mathrm{ch}_2(K_3^{\vee\vee}) = L$ and this contradicts Lemma 6.5. Thus $Z_1 = \Sigma_1$ and we know that $K_3 \cong K_3^{\vee\vee}$.

Then we claim that $\mathcal{H}^0(\mathrm{pr}_2(I_C))$ is μ -(semi)stable. Indeed, if $\mathcal{H}^0(\mathrm{pr}_2(I_C))$ is not μ -semistable, let K_4 be its minimal destabilizing quotient sheaf. Then by (6) and the stability of K_3 and I_C , it is not hard to see that the only possible case is $\mathrm{rk}(K_4) = 3$ and $\mathrm{ch}_1(K_4) = -H$. Then if we apply $\mathrm{Hom}(-, K_4^{\vee\vee})$ to the triangle (6), we obtain $\mathrm{Hom}(K_3, K_4^{\vee\vee}) \neq 0$. Since they have the same rank and ch_1 , by stability we have $K_3 \subset K_4^{\vee\vee}$, which is impossible since they are both reflexive but $K_4^{\vee\vee}/K_3$ is supported in codimension ≥ 2 .

Finally, we show that $\mathrm{Hom}(K_2, \mathcal{H}^0(\mathrm{pr}_2(I_C))) = k$, then using the μ -stability of K_2 and $\mathcal{H}^0(\mathrm{pr}_2(I_C))$, we obtain $K_2 \cong \mathcal{H}^0(\mathrm{pr}_2(I_C))$. From the definition of K_2 , it is not hard to see that $\mathrm{RHom}(K_2, \mathcal{O}_X(-H)) = 0$. Now applying $\mathrm{Hom}(K_2, -)$ to the exact sequence (7), we obtain $\mathrm{RHom}(K_2, \mathcal{U}^{\oplus 2}) = \mathrm{RHom}(K_2, K_3)$. By Lemma 6.6, we know that $\mathrm{RHom}(K_2, K_3) = k^2[-2]$. Therefore, if we apply $\mathrm{Hom}(K_2, -)$ to the exact sequence (6), we obtain

$$\mathrm{Hom}(K_2, \mathcal{H}^0(\mathrm{pr}_2(I_C))) = \mathrm{Hom}(K_2, I_C),$$

which equals to k by Lemma 6.6.

Recall that $\mathrm{pr}_1(\mathcal{U}[1])$ sits in the triangle

$$\mathcal{U}[1] \rightarrow \mathrm{pr}_1(\mathcal{U}[1]) \rightarrow K_2.$$

Now we have establish $\mathrm{pr}_2(I_C)$ and $\mathrm{pr}_1(\mathcal{U}[1])$ share the same cohomology objects. Then the result $\mathrm{pr}_2(I_C) \cong \mathrm{pr}_1(\mathcal{U}[1])$ follows from the fact $\mathrm{Ext}^1(K_2, \mathcal{U}[1]) = k$. \square

6.4. σ -conic. In this subsection, we compute the projection objects of σ -conics.

Proposition 6.8. *Let X be a general ordinary GM fourfold and C be a σ -conic on X . Then we have a triangle*

$$\mathbb{D}(I_q(H))[1] \rightarrow \mathrm{pr}_2(I_C) \rightarrow \mathcal{Q}^\vee,$$

where q is the unique σ -quadric on X .

Proof. Let Y be a hyperplane section of X containing C , it is easy to see Y is integral because X is of Picard number one. Since X is general, it does not contain any plane, then $\langle C \rangle \cap Y = C$. At the same time, $\langle C \rangle$ is a σ -plane in Y , which means C is the zero locus of a section of \mathcal{Q}_Y . Thus we have an exact sequence on Y

$$0 \rightarrow \mathcal{U}_Y \rightarrow \mathcal{Q}_Y^\vee \rightarrow I_{C/Y} \rightarrow 0.$$

Note that we have exact sequences on X

$$0 \rightarrow \mathcal{U}(-H) \rightarrow \mathcal{U} \rightarrow \mathcal{U}_Y \rightarrow 0 \tag{9}$$

and

$$0 \rightarrow \mathcal{Q}^\vee(-H) \rightarrow \mathcal{Q}^\vee \rightarrow \mathcal{Q}_Y^\vee \rightarrow 0. \tag{10}$$

Hence $\mathrm{RHom}(\mathcal{U}^\vee, \mathcal{U}_Y) = k[-3]$ and $\mathrm{RHom}(\mathcal{U}^\vee, \mathcal{Q}_Y^\vee) = k[-1]$. Applying the mutation $\mathbf{L}_{\mathcal{O}_X}$ to the defining complexes of $\mathbf{L}_{\mathcal{U}^\vee}\mathcal{U}_Y$ and $\mathbf{L}_{\mathcal{U}^\vee}\mathcal{Q}_Y^\vee$ respectively, we get triangles

$$\mathcal{U}_Y \rightarrow \mathbf{L}_{\mathcal{O}_X}\mathbf{L}_{\mathcal{U}^\vee}\mathcal{U}_Y \rightarrow \mathcal{Q}^\vee[-1]$$

and

$$\mathcal{Q}^\vee \rightarrow \mathcal{Q}_Y^\vee \rightarrow \mathbf{L}_{\mathcal{O}_X}\mathbf{L}_{\mathcal{U}^\vee}\mathcal{Q}_Y^\vee.$$

Now applying the mutation $\mathbf{R}_{\mathcal{O}_X(-H)}$ on the sequences (9) and (10), since $\mathrm{Ext}^1(\mathcal{U}, \mathcal{Q}^\vee) = 0$ by Lemma 5.4, we have $\mathbf{R}_{\mathcal{O}_X(-H)}\mathcal{U}_Y \cong \mathcal{U} \oplus \mathcal{Q}(-H)$ and $\mathbf{R}_{\mathcal{O}_X(-H)}\mathcal{Q}_Y^\vee \cong \mathcal{U} \oplus \mathcal{Q}^\vee$.

Applying $\mathbf{R}_{\mathcal{O}_X(-H)}$ to the above triangles respectively, we get triangles

$$\mathcal{U} \oplus \mathcal{Q}(-H) \rightarrow \mathbf{R}_{\mathcal{O}_X(-H)}\mathbf{L}_{\mathcal{O}_X}\mathbf{L}_{\mathcal{U}^\vee}\mathcal{U}_Y \rightarrow \mathcal{Q}^\vee[-1]$$

and

$$\mathcal{Q}^\vee \rightarrow \mathcal{U} \oplus \mathcal{Q}^\vee \rightarrow \mathbf{R}_{\mathcal{O}_X(-H)}\mathbf{L}_{\mathcal{O}_X}\mathbf{L}_{\mathcal{U}^\vee}\mathcal{Q}_Y^\vee.$$

After taking the mutation $\mathbf{R}_{\mathcal{U}}$, we have

$$\mathbf{R}_{\mathcal{U}}\mathcal{Q}(-H) \rightarrow \mathrm{pr}_2(\mathcal{U}_Y) \rightarrow \mathcal{Q}^\vee[-1]$$

and

$$\mathrm{pr}_2(\mathcal{Q}_Y^\vee) \cong 0.$$

Therefore, combined with the sequence $0 \rightarrow \mathcal{U}_Y \rightarrow \mathcal{Q}_Y^\vee \rightarrow I_{C/Y} \rightarrow 0$ and $\mathrm{pr}_2(\mathcal{Q}_Y^\vee) \cong 0$, we obtain

$$\mathrm{pr}_2(I_C) \cong \mathrm{pr}_2(\mathcal{U}_Y)[1].$$

Under this case, $\mathrm{pr}_2(I_C)$ sits in the triangle

$$\mathbf{R}_{\mathcal{U}}\mathcal{Q}(-H)[1] \rightarrow \mathrm{pr}_2(I_C) \rightarrow \mathcal{Q}^\vee.$$

Now the result follows from $\mathbf{R}_{\mathcal{U}}\mathcal{Q}(-H) = \mathbb{D}(I_q(H))$. To this end, we only need to prove that $I_q(H) \cong \mathbb{D}(\mathbf{R}_{\mathcal{U}}\mathcal{Q}(-H)) \cong \mathbf{L}_{\mathcal{U}^\vee}\mathcal{Q}^\vee(H)$, which is implied by the fact $\mathbf{L}_{\mathcal{U}}\mathcal{Q}^\vee \cong I_q$ in Proposition 6.4. \square

7. STABILITY OF PROJECTION OBJECTS OF CONICS

In this section, we apply Proposition 4.7 to show $\mathrm{pr}_2(I_C)$ is stable with respect to every stability condition on $\mathcal{K}u(X)$ for a *very general* ordinary GM fourfold X . It is worth mentioning that this is the only section we use the very general (general non-Hodge-special) assumption.

Theorem 7.1. *Let X be a very general ordinary GM fourfold and C be a conic on X . Then the objects $\mathrm{pr}_2(I_C)$ and $\mathrm{pr}_1(\mathcal{O}_C(H))$ are stable with respect to every stability condition on $\mathcal{K}u(X)$.*

Proof. By the identification in Lemma 6.2, for a conic C , we only need to show that one of the objects $\mathrm{pr}_1(\mathcal{O}_C(H))$ and $\mathrm{pr}_2(I_C)$ is stable. In the followings, we prove the theorem in Proposition 7.4, Proposition 7.6 and Proposition 7.7. \square

7.1. Stability of projection objects of τ -conics. At first, we list two lemmas which are useful when we compute the spectral sequences in Proposition 7.4.

Lemma 7.2. *Let X be a general ordinary GM fourfold and C be a τ -conic on X . Let Σ be the zero locus of a section of \mathcal{U}^\vee containing C . Then we have*

- (1) $\mathrm{RHom}(I_C, I_C) = k[0] \oplus k^5[-1] \oplus k^7[-2]$.
- (2) $\mathrm{RHom}(I_\Sigma, I_\Sigma) = k[0] \oplus k^4[-1]$
- (3) $\mathrm{RHom}(I_C, I_\Sigma) = k[-1] \oplus k^4[-2]$.
- (4) $\mathrm{RHom}(I_\Sigma, I_C) = k[0] \oplus k^6[-1]$.

Proof. (1): It is clear that $\text{hom}(I_C, I_C) = 1$, by Serre duality we have

$$\text{ext}^4(I_C, I_C) = \text{hom}(I_C, I_C(-2H)) = 0.$$

Using [IM11, Theorem 3.2], we know $\text{ext}^1(I_C, I_C) = 5$. Since $\chi(I_C, I_C) = 3$, we only need to show that $\text{ext}^3(I_C, I_C) = 0$. To this end, we apply $\text{Hom}(I_C, -)$ to the exact sequence

$$0 \rightarrow I_C(-2H) \rightarrow \mathcal{O}_X(-2H) \rightarrow \mathcal{O}_C(-2H) \rightarrow 0.$$

Since $\text{Hom}(I_C, \mathcal{O}_X(-2H)) \cong \text{Ext}^1(I_C, \mathcal{O}_X(-2H)) = 0$, we have

$$\text{Ext}^1(I_C, I_C(-2H)) \cong \text{Hom}(I_C, \mathcal{O}_C(-2H)) = 0.$$

By Serre duality, we obtain $\text{Ext}^3(I_C, I_C) \cong \text{Ext}^1(I_C, I_C(-2H)) = 0$.

(2): Note that $\chi(I_\Sigma, I_\Sigma) = -3$. Recall that Σ is the zero locus of a section of \mathcal{U}^\vee , hence we have the Koszul resolution

$$0 \rightarrow \mathcal{O}_X(-H) \rightarrow \mathcal{U} \rightarrow I_\Sigma \rightarrow 0.$$

Then the result follows from applying [Pir20, Lemma 2.27] to this exact sequence.

(3): It is clear that $\text{Hom}(I_C, I_\Sigma) = \text{Ext}^4(I_C, I_\Sigma) = 0$. Now the result follows from applying $\text{Hom}(I_C, -)$ to the Koszul resolution of I_Σ .

(4): Applying $\text{Hom}(-, I_C)$ to the resolution of I_Σ , we obtain $\text{Ext}^i(I_\Sigma, I_C) = 0$ for $i \neq 0, 1$ and an exact sequence

$$0 \rightarrow \text{Hom}(I_\Sigma, I_C) \rightarrow k \rightarrow k^6 \rightarrow \text{Ext}^1(I_\Sigma, I_C) \rightarrow 0.$$

Since $\text{hom}(I_\Sigma, I_C) = 1$, we obtain $\text{ext}^1(I_\Sigma, I_C) = 6$. □

Lemma 7.3. *Let X be a general ordinary GM fourfold and C be a τ -conic on X . Let Σ be the zero locus of a section of \mathcal{U}^\vee containing C . Then we have:*

- (1) $\text{RHom}(I_{C/\Sigma}, I_{C/\Sigma}) = k[0] \oplus k^n[-1] \oplus k^{n+1}[-2]$ for some $n \geq 3$.
- (2) $\text{RHom}(I_{C/\Sigma}(H), \mathcal{O}_X) = k^2[-2]$.
- (3) $\text{RHom}(\mathcal{O}_X, I_{C/\Sigma}(H)) = k^2[0]$.

Proof. Note that $\chi(I_{C/\Sigma}, I_{C/\Sigma}) = 2$. Then (1) follows from Lemma 7.2 and applying [Pir20, Lemma 2.27] to the exact sequence $0 \rightarrow I_\Sigma \rightarrow I_C \rightarrow I_{C/\Sigma} \rightarrow 0$.

(2) and (3) follow from applying $\text{Hom}(-, \mathcal{O}_X)$ and $\text{Hom}(\mathcal{O}_X, -)$ to the exact sequence $0 \rightarrow I_\Sigma(H) \rightarrow I_C(H) \rightarrow I_{C/\Sigma}(H) \rightarrow 0$. □

Proposition 7.4. *Let X be a very general ordinary GM fourfold and C be a τ -conic on X . Then $\text{pr}_1(\mathcal{O}_C(H))$ is stable with respect to every stability condition on $\mathcal{K}u(X)$.*

Proof. In Proposition 6.3, we have $\text{pr}_1(\mathcal{O}_C(H)) \cong \mathbf{L}_{\mathcal{O}_X}(I_{C/\Sigma}(H))[1]$. Then we apply [Pir20, Lemma 2.27] to the triangle

$$\mathcal{O}_X^{\oplus 2} \rightarrow I_{C/\Sigma}(H) \rightarrow \text{pr}_1(\mathcal{O}_C(H))[-1].$$

From Lemma 7.3, for $i \notin \{0, 1, 2\}$,

$$\text{Hom}(\text{pr}_1(\mathcal{O}_C(H)), \text{pr}_1(\mathcal{O}_C(H))) = k, \quad \text{Ext}^i(\text{pr}_1(\mathcal{O}_C(H)), \text{pr}_1(\mathcal{O}_C(H))) = 0.$$

Then by Serre duality in $\mathcal{K}u(X)$, we have

$$\text{Ext}^2(\text{pr}_1(\mathcal{O}_C(H)), \text{pr}_1(\mathcal{O}_C(H))) = \text{Hom}(\text{pr}_1(\mathcal{O}_C(H)), \text{pr}_1(\mathcal{O}_C(H))) = k.$$

Since $\chi(\text{pr}_1(\mathcal{O}_C(H)), \text{pr}_1(\mathcal{O}_C(H))) = -2$, we obtain $\text{Ext}^1(\text{pr}_1(\mathcal{O}_C(H)), \text{pr}_1(\mathcal{O}_C(H))) = k^4$. Then by Proposition 4.7, $\text{pr}_1(\mathcal{O}_C(H))$ is stable with respect to every stability condition on $\mathcal{K}u(X)$. □

7.2. Stability of projection objects of ρ -conics. Let C be a ρ -conic on X , by Proposition 6.7, $\mathrm{pr}_2(I_C) \cong \mathrm{pr}_1(\mathcal{U})[1]$, where $\mathrm{pr}_1(\mathcal{U})$ fits into the triangle as in Proposition 6.4

$$\mathcal{U} \rightarrow \mathrm{pr}_1(\mathcal{U}) \rightarrow \mathbf{L}_{\mathcal{O}_X}(I_q(H))[-2].$$

Now we only need to prove $\mathrm{pr}_1(\mathcal{U})$ is stable.

Lemma 7.5.

- (1) $\mathrm{RHom}(\mathcal{U}, \mathbf{L}_{\mathcal{O}_X}(I_q(H))[-3]) = k^m[-2] \oplus k^{m-3}[-3]$, for some integer $3 \leq m \leq 25$.
(2) $\mathrm{RHom}(\mathcal{U}, \mathrm{pr}_1(\mathcal{U})) \cong \mathrm{RHom}(\mathrm{pr}_1(\mathcal{U}), \mathrm{pr}_1(\mathcal{U})) = k[0] \oplus k^4[-1] \oplus k[-2]$.

Proof. (1): Applying $\mathrm{Hom}(\mathcal{U}, -)$ to the short exact sequence

$$0 \rightarrow \mathcal{U}^\vee \rightarrow \mathcal{Q}^\vee(H) \rightarrow I_q(H) \rightarrow 0,$$

we get $\mathrm{RHom}(\mathcal{U}, I_q(H)) = k^{22}[0]$. Then we apply $\mathrm{Hom}(\mathcal{U}, -)$ to the triangle

$$\mathcal{O}_X^{\oplus 5} \rightarrow I_q(H) \rightarrow \mathbf{L}_{\mathcal{O}_X}(I_q(H)).$$

Since $\mathrm{RHom}(\mathcal{U}, \mathcal{O}_X^{\oplus 5}) = k^{25}[0]$, by the long exact sequence we have

$$\mathrm{RHom}(\mathcal{U}, \mathbf{L}_{\mathcal{O}_X}(I_q(H))) = k^m[1] \oplus k^{m-3}[0],$$

for some integer $3 \leq m \leq 25$. Then we obtain

$$\mathrm{RHom}(\mathcal{U}, \mathbf{L}_{\mathcal{O}_X}(I_q(H))[-3]) = k^m[-2] \oplus k^{m-3}[-3].$$

(2): Since $\mathrm{pr}_1(\mathcal{U})$ fits into the triangle

$$\mathbf{L}_{\mathcal{O}_X}(I_q \otimes \mathcal{O}_X(H))[-3] \rightarrow \mathcal{U} \rightarrow \mathrm{pr}_1(\mathcal{U}).$$

Using (1) and $\mathrm{RHom}(\mathcal{U}, \mathcal{U}) = k[0]$, we have $\mathrm{RHom}(\mathcal{U}, \mathrm{pr}_1(\mathcal{U})) = k[0] \oplus k^m[-1] \oplus k^{m-3}[-2]$. By Serre duality in $\mathcal{K}u(X)$ and adjunction, we have

$$\mathrm{Hom}(\mathcal{U}, \mathrm{pr}_1(\mathcal{U})) = \mathrm{Hom}(\mathrm{pr}_1(\mathcal{U}), \mathrm{pr}_1(\mathcal{U})) = \mathrm{Hom}(\mathrm{pr}_1(\mathcal{U}), \mathrm{pr}_1(\mathcal{U})[2]) = \mathrm{Hom}(\mathcal{U}, \mathrm{pr}_1(\mathcal{U})[2]).$$

Therefore, we obtain $m-3 = 1$, which means $\mathrm{RHom}(\mathcal{U}, \mathrm{pr}_1(\mathcal{U})) = k[0] \oplus k^4[-1] \oplus k[-2]$. \square

Proposition 7.6. *Let X be a very general ordinary GM fourfold and C be a ρ -conic on X , then $\mathrm{pr}_2(I_C)$ is stable with respect to every stability condition on $\mathcal{K}u(X)$.*

Proof. By Lemma 7.5, $\mathrm{Ext}^1(\mathrm{pr}_1(\mathcal{U}), \mathrm{pr}_1(\mathcal{U})) = k^4$. Then using Proposition 4.7, $\mathrm{pr}_1(\mathcal{U})$ is stable, which implies $\mathrm{pr}_2(I_C)$ is stable with respect to every stability condition on $\mathcal{K}u(X)$. \square

7.3. Stability of projection objects of σ -conics.

Proposition 7.7. *Let X be a very general GM fourfold and C be a σ -conic on X , then $\mathrm{pr}_2(I_C)$ is stable with respect to every stability condition on $\mathcal{K}u(X)$.*

Proof. By Proposition 6.8, the object $\mathrm{pr}_2(I_C)$ fits into the triangle

$$\mathbb{D}(I_q(H))[1] \rightarrow \mathrm{pr}_2(I_C) \rightarrow \mathcal{Q}^\vee.$$

It is easy to check that $T(\mathrm{pr}_2(I_C)) \cong \mathrm{pr}_1(\mathcal{U})[1]$. Then the result follows from Proposition 7.6. \square

Remark 7.8. For any ρ -conic C , $\mathrm{pr}_2(I_C) \cong \mathrm{pr}_1(\mathcal{U})[1]$, for any σ -conic C , we have $T(\mathrm{pr}_2(I_C)) \cong \mathrm{pr}_1(\mathcal{U})[1]$. Indeed, it is easy to check $T(\mathrm{pr}_1(\mathcal{U})[1]) \not\cong \mathrm{pr}_1(\mathcal{U})[1]$. This means that the morphism p induced by pr_2 contracts the locus σ -conics and ρ -conics in $F_g(X)$ to two different points in the moduli space $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1)$. Moreover, the induced action of T on $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1)$ will take one point to another.

8. BRIDGELAND MODULI SPACES AND THE DOUBLE EPW SEXTICS

8.1. **Moduli space of stable objects** $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1)$. For a very general GM fourfold X , let $F_g(X)$ be the Hilbert scheme of conics on X . In this section, we show that the projection functor $\mathrm{pr}_1 : D^b(X) \rightarrow \mathcal{K}u(X)$ (or equivalently, pr_2) induces a dominant proper morphism p from $F_g(X)$ to the Bridgeland moduli space $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1)$. Moreover, this morphism p is compatible with the morphism $f : F_g(X) \rightarrow \tilde{Y}_{A^\perp}$ defined in [IM11, Section 4.4]. In particular, we show that $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1) \cong \tilde{Y}_{A^\perp}$ and $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_2) \cong \tilde{Y}_A$.

By Theorem 7.1, the object $\mathrm{pr}_1(\mathcal{O}_C(H))$ is σ -stable for any conic $C \in F_g(X)$. On the other hand, $F_g(X)$ admits an universal family and the functor pr_1 is of Fourier-Mukai type. By the standard argument as in [Zha20, Lemma 6.1, Proposition 6.2] and [LPZ18, Theorem 3.9], the projection functor pr_1 induces a morphism $p : F_g(X) \rightarrow \mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1)$. According to Remark 7.8, the morphism p contracts ρ -conics and σ -conics to two different points in $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1)$. Next, when restricting on the locus of τ -conics, we show that p is a \mathbb{P}^1 -bundle.

Proposition 8.1. *Let C and C' be two τ -conics on X . Then $f(C) = f(C')$ if and only if $\mathrm{Hom}(\mathrm{pr}_1(\mathcal{O}_C(H)), \mathrm{pr}_1(\mathcal{O}_{C'}(H))) \neq 0$.*

Proof. First we assume that $C \subset \Sigma$ and $C' \subset \Sigma'$, where the two surfaces $\Sigma := \mathrm{Gr}(2, V_4) \cap X$ and $\Sigma' := \mathrm{Gr}(2, V'_4) \cap X$ are zero locus of two sections of \mathcal{U}^\vee .

From the definition of left mutation, we have two triangles

$$\mathcal{O}_X^{\oplus 2} \rightarrow I_{C/\Sigma}(H) \rightarrow \mathrm{pr}_1(\mathcal{O}_C(H))[-1]$$

and

$$\mathcal{O}_X^{\oplus 2} \rightarrow I_{C'/\Sigma'}(H) \rightarrow \mathrm{pr}_1(\mathcal{O}_{C'}(H))[-1].$$

If we apply [Pir20, Lemma 2.27] to these two triangles, we obtain a spectral sequence with the first page $E_1^{p,q}$ in the form

$$\begin{array}{ccc|ccc} \vdots & & & \vdots & & \vdots \\ \mathrm{Ext}^1(I_{C/\Sigma}(H), \mathcal{O}_X^{\oplus 2}) & & \mathrm{Ext}^1(\mathcal{O}_X^{\oplus 2}, \mathcal{O}_X^{\oplus 2}) \oplus \mathrm{Ext}^1(I_{C/\Sigma}(H), I_{C'/\Sigma'}(H)) & & \mathrm{Ext}^1(\mathcal{O}_X^{\oplus 2}, I_{C'/\Sigma'}(H)) & \\ \mathrm{Hom}(I_{C/\Sigma}(H), \mathcal{O}_X^{\oplus 2}) & & \mathrm{Hom}(\mathcal{O}_X^{\oplus 2}, \mathcal{O}_X^{\oplus 2}) \oplus \mathrm{Hom}(I_{C/\Sigma}(H), I_{C'/\Sigma'}(H)) & & \mathrm{Hom}(\mathcal{O}_X^{\oplus 2}, I_{C'/\Sigma'}(H)) & \\ \hline 0 & & 0 & & 0 & \end{array}$$

Since $\mathrm{Hom}(I_{C/\Sigma}(H), \mathcal{O}_X^{\oplus 2}) = \mathrm{Ext}^1(I_{C/\Sigma}(H), \mathcal{O}_X^{\oplus 2}) = 0$ and $\mathrm{pr}_1(\mathcal{O}_{C'}(H)) \in \mathcal{K}u(X)$,

$$\mathrm{Hom}(\mathrm{pr}_1(\mathcal{O}_C(H)), \mathrm{pr}_1(\mathcal{O}_{C'}(H))) = E_\infty^{0,0} = \ker(E_1^{0,0} \rightarrow E_1^{1,0}) = \mathrm{Hom}(I_{C/\Sigma}, I_{C'/\Sigma'}).$$

Then we need to prove that $\mathrm{Hom}(I_{C/\Sigma}, I_{C'/\Sigma'}) \neq 0$ if and only if $f(C) = f(C')$.

To this end, we first review the construction of $f : F_g(X) \rightarrow \tilde{Y}_{A^\perp}$. In [IM11, Section 4.4], $f(C) = f(C')$ if and only if there exists some $V_4 \subset V_5$ such that $C, C' \subset \mathrm{Gr}(2, V_4) \cap X$, where we denote the surface $\mathrm{Gr}(2, V_4) \cap X$ by $\tilde{\Sigma}$. This means $\Sigma = \Sigma' = \tilde{\Sigma}$. At the same time, the planes $\langle C \rangle$ and $\langle C' \rangle$ in $\mathbb{P}(\wedge^2 V_4) \cap H \cong \mathbb{P}^4$ should be contained in a same quadric Q' , where Q' is in the pencil $|P_{V_4}, Q_{V_4}|$. Here $Q_{V_4} := Q \cap \mathbb{P}(\wedge^2 V_4) \cap H$ and $P_{V_4} := \mathrm{Gr}(2, V_4) \cap H$.

First we claim that if $\mathrm{Hom}(I_{C/\Sigma}, I_{C'/\Sigma'}) \neq 0$, then $\Sigma = \Sigma'$, i.e., $\mathrm{Hom}(I_{\Sigma/\mathbb{P}^4}, I_{\Sigma'/\mathbb{P}^4}) \neq 0$. Indeed, this follows from applying [Pir20, Lemma 2.27] to the exact sequences

$$0 \rightarrow I_{\Sigma/\mathbb{P}^4} \rightarrow I_{C/\mathbb{P}^4} \rightarrow I_{C/\Sigma} \rightarrow 0$$

and

$$0 \rightarrow I_{\Sigma'/\mathbb{P}^4} \rightarrow I_{C'/\mathbb{P}^4} \rightarrow I_{C'/\Sigma'} \rightarrow 0.$$

Therefore in the following we can assume that C and C' are both contained in $\mathrm{Gr}(2, V_4)$ for a same V_4 .

As in [IM11, Proposition 4.9], Q' is either a cone over a smooth quadric surface and planes in Q' are parametrized by two projective lines, or a double cone over a smooth conic and planes in Q' are parametrized by that smooth conic. From the definition, there is a one-to-one correspondence between a τ -conic C with $\langle C \rangle \subset Q'$ and a plane contained in Q' . If $f(C) = f(C')$, from the construction of f , we know that C and C' come from a same family of planes in Q' . This means that they are linearly equivalent as Weil divisors in Σ , which implies $I_{C/\Sigma} \cong I_{C'/\Sigma}$. Hence we have $\text{Hom}(I_{C/\Sigma}, I_{C'/\Sigma}) \neq 0$.

Conversely, if $\text{Hom}(I_{C/\Sigma}, I_{C'/\Sigma}) \neq 0$, then we know that C and C' are linearly equivalent as Weil divisors in Σ . From the construction of f , we obtain that $f(C) = f(C')$. \square

Now, we are ready to prove the first main result of our paper.

Theorem 8.2. *Let X be a very general GM fourfold with a Lagrangian data (A, V_5, V_6) . For any generic stability condition σ on $\mathcal{K}u(X)$, the projection functor pr_1 will induce an isomorphism*

$$i : \tilde{Y}_{A^\perp} \cong \mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1).$$

Proof. Now we have the morphism $p : F_g(X) \rightarrow \mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1)$ induced by pr_1 . In Proposition 8.1, on the locus of τ -conics, p is a \mathbb{P}^1 -bundle and coincides with f . Furthermore, it follows from Proposition 6.7 and Proposition 6.8 that the morphism p coincides with f on the whole $F_g(X)$. By [PPZ19, Proposition 1.5], we know that $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1)$ is a smooth projective variety of dimension four. On the other hand, since the dimension of $F_g(X)$ is five and the general fiber of p is one dimensional, p is a proper dominant morphism onto $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1)$. Then according to [Vak17, Exercise 29.5.C], we have an isomorphism $i : \tilde{Y}_{A^\perp} \cong \mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1)$ such that $i \circ f = p$. \square

Remark 8.3. Let Y be a smooth hyperplane section of X . Then by [Log12] and [DK22], there is a contraction $g : F_g(Y) \rightarrow \tilde{Y}_{A^\perp}^{\geq 2}$ from the Fano surface of conics $F_g(Y)$ to the associated double dual EPW surface $\tilde{Y}_{A^\perp}^{\geq 2}$. Moreover, we have a commutative diagram as in [IM11]

$$\begin{array}{ccc} F_g(Y) & \xrightarrow{g} & \tilde{Y}_{A^\perp}^{\geq 2} \\ \downarrow & & \downarrow h \\ F_g(X) & \xrightarrow{f} & \tilde{Y}_{A^\perp} \end{array}$$

By [JLLZ21, Theorem 7.13], $\tilde{Y}_{A^\perp}^{\geq 2}$ is isomorphic to the moduli space $\mathcal{M}_\tau(\mathcal{K}u(Y), x)$ of stable objects of a (-1) -class x in the Kuznetsov component $\mathcal{K}u(Y)$, with respect to some stability conditions τ . Thus the diagram above becomes

$$\begin{array}{ccc} F_g(Y) & \xrightarrow{p'} & \mathcal{M}_\tau(\mathcal{K}u(Y), x) \\ \downarrow & & \downarrow h \\ F_g(X) & \xrightarrow{p} & \mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1) \end{array}$$

such that the morphisms p and p' are induced by the projection functors of Kuznetsov components. In our upcoming preprint [GLZ22], we show that the embedding h is induced by the functor $\text{pr}_2 \circ \iota_*$, where $\iota : Y \hookrightarrow X$ is the closed embedding.

8.2. Moduli space of stable objects $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_2)$. In Theorem 8.2, we have shown that for a very general GM fourfold X , the moduli space $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1)$ is isomorphic to the double dual EPW sextic \tilde{Y}_{A^\perp} . Now we prove the moduli space $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_2)$ is isomorphic to another hyperkähler fourfold, the double EPW sextic \tilde{Y}_A .

Theorem 8.4. *Let X be a very general GM fourfold with a Lagrangian data (A, V_5, V_6) . For any generic stability condition σ on $\mathcal{K}u(X)$, we have an isomorphism*

$$\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_2) \cong \tilde{Y}_A.$$

Proof. Let X' be the period dual of X , we can always choose X' to be very general as well. Then by [KP19, Theorem 1.6], there is an equivalence of Fourier-Mukai type $\Phi : \mathcal{K}u(X) \simeq \mathcal{K}u(X')$. We claim that Φ induces an isomorphism between Bridgeland moduli spaces $\phi : \mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1) \rightarrow \mathcal{M}_{\sigma'}(\mathcal{K}u(X'), \Lambda'_2)$. Indeed, as a corollary of [BP22, Theorem 5.12], the induced isomorphism

$$[\Phi] : \mathcal{N}(\mathcal{K}u(X)) \cong \mathcal{N}(\mathcal{K}u(X'))$$

will identify the canonical rank 2 lattices $\langle \Lambda_1, \Lambda_2 \rangle$ and $\langle \Lambda'_1, \Lambda'_2 \rangle$ on each side. Then up to sign, $[\Phi](\Lambda_1) = \Lambda'_1$ or Λ'_2 . If $[\Phi](\Lambda_1) = \Lambda'_1$, then we have

$$\phi : \tilde{Y}_{A^\perp} \cong \mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1) \cong \mathcal{M}_{\sigma'}(\mathcal{K}u(X'), \Lambda'_1) \cong \tilde{Y}_{A'^\perp}.$$

But X' is the period dual of X , $A' = A^\perp$, then we get an isomorphism $\tilde{Y}_{A'^\perp} \cong \tilde{Y}_A$, which is impossible by Theorem 2.4. Thus $[\Phi](\Lambda_1) = \Lambda'_2$ and $[\Phi](\Lambda_2) = \Lambda'_1$. Then the moduli space $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_2) \cong \mathcal{M}_\sigma(\mathcal{K}u(X'), \Lambda'_1) \cong \tilde{Y}_{A'^\perp}$, again by $A' = A^\perp$, we have $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_2) \cong \tilde{Y}_A$. \square

Remark 8.5. In fact, [BP22, Theorem 5.12] holds for any GM fourfold. As a result, once we extend Theorem 1.1(1) to a general GM fourfold, Theorem 1.1(2) automatically holds. The point is that even though the rank of the numerical Grothendieck group is bigger than two, the induced equivalence of period duals still fixes the canonical $A_1^{\oplus 2}$ lattice. Then our method in Theorem 8.4 still works.

8.3. An involution on $\mathcal{K}u(X)$ and its induced action on the double dual EPW sextic. Now we are going to discuss two involutions acting on \tilde{Y}_{A^\perp} and $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1)$ respectively. One is naturally induced by the structure of double cover and the other is induced by the involutive functor T of $\mathcal{K}u(X)$ defined in Lemma 3.2.

By the result of Theorem 8.2, we have the following diagram

$$\begin{array}{ccccc} & & \tilde{Y}_{A^\perp} & \xrightarrow{\eta} & \tilde{Y}_{A^\perp} \\ & \nearrow f & \downarrow i & & \\ F_g(X) & \xrightarrow{p} & \mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1) & \xrightarrow{\eta'} & \mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1), \end{array}$$

where i is the isomorphism such that $i \circ f = p$ in Theorem 8.2 and the two involutions are denoted by η and η' respectively. To prove η and η' coincide, it suffices to show that for a general conic $C \in F_g(X)$, there exists another conic C' such that $\eta \circ f(C) = f(C')$ and $\eta' \circ p(C) = p(C')$.

Firstly, we briefly review the involution η described in [IM11, Lemma 4.19]. For a general conic C , if there exists another conic C' such that $\eta \circ f(C) = f(C')$, then the spanning planes $\langle C \rangle$ and $\langle C' \rangle$ lie in a same quadric threefold $Q_{C, V_4} = Q_{C', V_4}$. Here Q_{C, V_4} is the unique singular quadric of the pencil $\langle \text{Gr}(2, V_4) \cap H, H \cap Q \rangle$ contained in $\mathbb{P}(\wedge^2 V_4) \cap H$. However, $\langle C \rangle$ and $\langle C' \rangle$ do not belong to the same ruling, as a result, $\langle C \rangle$ meets $\langle C' \rangle$ along a line. Furthermore, the generating 3-plane $\langle C, C' \rangle$ cuts X at a degenerate elliptic curve

$$e := C \cup C' = \langle C, C' \rangle \cap \text{Gr}(2, V_4) \cap Q.$$

Now we only need to show such C' satisfies $\eta' \circ p(C) = p(C')$ as well, which is equivalent to prove $T(\text{pr}_1(\mathcal{O}_C(H))) \cong \text{pr}_1(\mathcal{O}_{C'}(H))$.

Lemma 8.6. *Let X be a general ordinary GM fourfold. Let C be a general τ -conic such that it is contained in a smooth surface Σ , defined by the zero locus of a section of \mathcal{U}^\vee . Then we have a triangle*

$$0 \rightarrow I_\Sigma \rightarrow \mathrm{pr}_2(I_C) \rightarrow I_C \rightarrow 0.$$

Moreover, $T(\mathrm{pr}_1(\mathcal{O}_C(H))) \cong \mathrm{pr}_1(\mathcal{O}_{C'}(H))$ for another τ -conic $C' \subset \Sigma$ such that as divisors of Σ , $C' = -C - K_\Sigma$.

Proof. The first statement follows from Proposition 6.3, note that $I_\Sigma \cong \mathrm{cok}(\mathcal{U}^{\oplus 4} \hookrightarrow K_1)$. Since I_Σ and I_C are both subsheaves of \mathcal{O}_X , $\mathrm{pr}_2(I_C)$ is also a subsheaf of $\mathcal{O}_X^{\oplus 2}$. The cokernel of $\mathrm{pr}_2(I_C) \xrightarrow{i} \mathcal{O}_X^{\oplus 2}$ is the sheaf obtained by an extension of \mathcal{O}_Σ by \mathcal{O}_C . Consider the standard exact sequence of $C \subset \Sigma$,

$$0 \rightarrow \mathcal{O}_\Sigma(-C) \rightarrow \mathcal{O}_\Sigma \rightarrow \mathcal{O}_C \rightarrow 0.$$

Tensoring with $\mathcal{O}_\Sigma(C)$, since $C \cdot C = 0$, we get

$$0 \rightarrow \mathcal{O}_\Sigma \rightarrow \mathcal{O}_\Sigma(C) \rightarrow \mathcal{O}_C \rightarrow 0.$$

Then $\mathrm{pr}_2(I_C) \cong \ker(\mathcal{O}_X^{\oplus 2} \rightarrow \mathcal{O}_\Sigma(C))$. Note that $K_\Sigma \cong \mathcal{O}_\Sigma(-H)$ and $-C - K_\Sigma$ is a divisor of conic as well.

We take $C' \in |-C - K_\Sigma|$, then $C \cup C' \in |C' + C| = |-K_\Sigma|$, which is a degenerate degree 4 elliptic curve on Σ . Thus we have

$$\mathrm{pr}_2(I_C) \cong \ker(\mathcal{O}_X^{\oplus 2} \rightarrow \mathcal{O}_\Sigma(-C' + H)) \cong \ker(\mathcal{O}_X^{\oplus 2} \rightarrow I_{C'/\Sigma}(H)) \cong \mathrm{pr}_1(\mathcal{O}_{C'}(H))[-2].$$

By Lemma 6.2, $\mathrm{pr}_2(I_C) \cong T(\mathrm{pr}_1(\mathcal{O}_C(H)))[-2]$, we obtain $T(\mathrm{pr}_1(\mathcal{O}_C(H))) \cong \mathrm{pr}_1(\mathcal{O}_{C'}(H))$.

Now it remains to show that C' is also a τ -conic. If C' is of σ -type or ρ -type, then $\mathrm{pr}_1(\mathcal{O}_{C'}(H)) \cong \mathrm{pr}_1(\mathcal{U})[3]$ or $T(\mathrm{pr}_1(\mathcal{U}))[3]$, neither of them is a rank two sheaf up to a shift. Then the result follows. \square

Proposition 8.7. *Let X be a very general ordinary GM fourfold, then via the isomorphism i , the two involutions η and η' coincide.*

Proof. By Lemma 8.6, for a general conic C , if $\eta' \circ p(C) = p(C')$ for another conic C' , C and C' lie in a same surface Σ of degree four. In particular, $C \cup C'$ is an elliptic curve of degree 4, which spans a 3-plane in $\mathbb{P}(\wedge^2 V_4) \cap H$, cutting along X by $C \cup C'$. This coincides with the choice of C' in [IM11], which means for a general conic C , we can always find another C' such that $\eta \circ f(C) = f(C')$ and $\eta' \circ p(C) = p(C')$. In conclusion, the two involutions essentially are the same. \square

Remark 8.8. There is an easier proof of Proposition 8.7 using the fact that for a very general GM fourfold X , the automorphism group of \tilde{Y}_{A^\perp} is isomorphic to \mathbb{Z}_2 , generated by the natural involution η . Thus to show η coincides with η' , it suffices to prove η' is non-trivial, which is obvious. However, our method in Proposition 8.7 is independent with this fact. Thus Proposition 8.7 will still be true in general once we identify \tilde{Y}_{A^\perp} with $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1)$ for a general ordinary GM fourfold X .

9. BIRATIONAL CATEGORICAL TORELLI FOR GUSHEL-MUKAI FOURFOLDS

Consider the period map of GM fourfolds,

$$\wp_4 : \mathbf{M}_4^{\mathrm{GM}} \longrightarrow \mathcal{D}.$$

It is known that the fibers of \wp_4 are of dimension four. On the other hand, it is shown in [KP19] that GM fourfolds share the equivalent Kuznetsov components if they are in the same fiber of the period map. It is natural to ask if $\mathcal{K}u(X)$ determines the birational isomorphism class of X . In fact, Kuznetsov-Perry propose the following conjecture in [KP19, Conjecture 1.9].

Conjecture 9.1. *Let X and X' be GM varieties of the same dimension such that there is an equivalence $\mathcal{K}u(X) \simeq \mathcal{K}u(X')$, then X is birational to X' .*

In [JLLZ21, Theorem 1.5], we prove the conjecture for general GM threefolds. In the current article, we prove this conjecture for very general GM fourfolds.

Theorem 9.2. *Let X and X' be very general ordinary GM fourfolds with equivalent Kuznetsov components $\mathcal{K}u(X) \simeq \mathcal{K}u(X')$. Then X is either the period partner or the period dual of X' . In particular, X is birational to X' .*

Proof. Let Φ denote the equivalence $\mathcal{K}u(X) \simeq \mathcal{K}u(X')$, then Φ induces an isometry between numerical Grothendieck groups. Since X and X' are non-Hodge-special, the isometry will map the canonical rank two lattice $\langle \Lambda_1, \Lambda_2 \rangle$ to $\langle \Lambda'_1, \Lambda'_2 \rangle$. Thus Φ induces a bijection ϕ or ϕ' between Bridgeland moduli spaces

$$\begin{array}{ccc} \mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1) & \xrightarrow{\phi} & \mathcal{M}_{\sigma'}(\mathcal{K}u(X'), \Lambda'_2) \\ & \searrow \phi' & \\ & & \mathcal{M}_{\sigma'}(\mathcal{K}u(X'), \Lambda'_1) \end{array}$$

By Theorem 8.2, the moduli space $\mathcal{M}_\sigma(\mathcal{K}u(X), \Lambda_1)$ is constructed from the Hilbert scheme of conics on the GM fourfold X by the projection functor. Then according to the standard argument in [BMMS12, Section 5], ϕ or ϕ' is actually an isomorphism.

In either case, after taking the primitive cohomology on both sides, we have a Hodge isometry

$$\langle \Lambda_1, \Lambda_2 \rangle^\perp \cong \langle \Lambda'_1, \Lambda'_2 \rangle^\perp.$$

Using the result in [PPZ19, Proposition 4.14], this is equivalent to an isomorphism between weight two Hodge structure of X and X'

$$H^4(X, \mathbb{Z})_0(1) \cong H^4(X', \mathbb{Z})_0(1).$$

Thus we deduce that X and X' are period partners or period duals (cf. [DK19, Remark 5.28]). In particular, X is birational to X' by [DK18, Corollary 4.16, Theorem 4.20]. \square

Remark 9.3. As a corollary of Theorem 9.2, for very general ordinary GM fourfolds X and X' , they are period partners or duals if and only if there exists an equivalence $\Phi : \mathcal{K}u(X) \simeq \mathcal{K}u(X')$. In a forthcoming preprint [LPZ22], the authors prove that any equivalence Φ between the Kuznetsov components of GM fourfolds is of Fourier-Mukai type. Thus Φ induces a Hodge isometry between the numerical Grothendieck groups. Using this property, combined with the result in [BP22], there is a more general version of Theorem 9.2. More precisely, using [BP22, Theorem 5.12] and the same argument in [PS22, Remark 6.16], one can prove that any two GM fourfolds X and X' are period partners or duals if and only if there exists an equivalence $\Phi : \mathcal{K}u(X) \simeq \mathcal{K}u(X')$ such that the induced Hodge isometry $[\Phi] : \mathcal{N}(\mathcal{K}u(X)) \rightarrow \mathcal{N}(\mathcal{K}u(X'))$ maps $\langle \Lambda_1, \Lambda_2 \rangle$ to $\langle \Lambda'_1, \Lambda'_2 \rangle$.

REFERENCES

- [BB17] Arend Bayer and Tom Bridgeland. Derived automorphism groups of K3 surfaces of Picard rank 1. *Duke Math. J.*, 166(1), Jan 2017.
- [Bea83] Arnaud Beauville. Variétés Kählériennes dont la première classe de Chern est nulle. *J. Diff. Geom.*, 18(4):755–782, 1983.
- [BLMS17] Arend Bayer, Martí Lahoz, Emanuele Macrì, and Paolo Stellari. Stability conditions on Kuznetsov components. (*Appendix joint with Xiaolei Zhao*) To appear in *Ann. Sci. Éc. Norm. Supér.*, arXiv:1703.10839, 2017.
- [BMMS12] Marcello Bernardara, Emanuele Macrì, Sukhendu Mehrotra, and Paolo Stellari. A categorical invariant for cubic threefolds. *Adv. Math.*, 229(2):770–803, 2012.

- [BP22] Arend Bayer and Alexander Perry. Kuznetsov’s Fano threefold conjecture via K3 categories and enhanced group actions. *arXiv preprint arXiv:2202.04195*, 2022.
- [Deb20] Olivier Debarre. Gushel-Mukai varieties. *arXiv preprint arXiv:2001.03485*, 2020.
- [DIM15] Olivier Debarre, Atanas Iliev, and Laurent Manivel. Special prime Fano fourfolds of degree 10 and index 2. *Recent advances in algebraic geometry*, 417:123, 2015.
- [DK18] Olivier Debarre and Alexander Kuznetsov. Gushel-Mukai varieties: classification and birationalities. *Algebr. Geom.*, 5:15–76, 01 2018.
- [DK19] Olivier Debarre and Alexander Kuznetsov. Gushel-Mukai varieties: linear spaces and periods. *Kyoto J. Math.*, 59(4):897–953, 2019.
- [DK22] Oliver Debarre and Alexander Kuznetsov. Gushel-Mukai varieties: quadrics. *In preparation*, 2022.
- [GLZ22] Hanfei Guo, Zhiyu Liu, and Shizhuo Zhang. Moduli space theoretical approach to Lagrangian subvariety of hyperkähler variety: Examples. *preprint*, 2022.
- [Har80] Robin Hartshorne. Stable reflexive sheaves. *Math. Ann.*, 254:121–176, 1980.
- [IM11] Atanas Iliev and Laurent Manivel. Fano manifolds of degree ten and EPW sextics. *Ann. Sci. Éc. Norm. Supér.*, 44(3):393–426, 2011.
- [JLLZ21] Augustinas Jacovskis, Xun Lin, Zhiyu Liu, and Shizhuo Zhang. Categorical Torelli theorems for Gushel-Mukai threefolds. *arXiv preprint, arXiv:2108.02946*, 2021.
- [KP18] Alexander Kuznetsov and Alexander Perry. Derived categories of Gushel-Mukai varieties. *Compos. Math.*, 154(7):1362–1406, 2018.
- [KP19] Alexander Kuznetsov and Alexander Perry. Categorical cones and quadratic homological projective duality. *To appear in Ann. Sci. Éc. Norm. Supér.*, *arXiv:1902.09824*, 2019.
- [Kuz04] Alexander Kuznetsov. Derived categories of cubic and V14 threefolds. *Tr. Mat. Inst. Steklova*, 246:171–194, 2004.
- [Li15] Chunyi Li. Stability conditions on Fano threefolds of Picard number 1. *J. Eur. Math. Soc.*, 21, 10 2015.
- [Log12] Dmitry Logachev. Fano threefolds of genus 6. *Asian Journal of Mathematics*, 16(3):515–560, 2012.
- [LPZ18] Chunyi Li, Laura Pertusi, and Xiaolei Zhao. Twisted cubics on cubic fourfolds and stability conditions. *arXiv preprint arXiv:1802.01134*, 2018.
- [LPZ20] Chunyi Li, Laura Pertusi, and Xiaolei Zhao. Elliptic quintics on cubic fourfolds, O’Grady 10, and Lagrangian fibrations. *arXiv preprint arXiv:2007.14108*, 2020.
- [LPZ22] Chunyi Li, Laura Pertusi, and Xiaolei Zhao. Derived categories of hearts on Kuznetsov components. *In preparation*, 2022.
- [Muk84] Shigeru Mukai. On the moduli space of bundles on K3 surfaces. I. *Vector bundles on algebraic varieties (Bombay, 1984)*, 11:341–413, 1984.
- [O’G95] Kieran O’Grady. The weight-two Hodge structure of moduli spaces of sheaves on a K3 surface. *J. Algebraic Geom.*, 6, 11 1995.
- [O’G06] Kieran O’Grady. Dual Double EPW-sextics and Their Periods. *Pure Appl. Math. Q.*, 4, 06 2006.
- [O’G10] Kieran O’Grady. Double covers of EPW-sextics. *Michigan Math. J.*, 62, 07 2010.
- [Per19] Laura Pertusi. On the double EPW sextic associated to a Gushel-Mukai fourfold. *J. London Math. Soc.*, 100(1):83–106, 2019.

- [Pir20] Dmitrii Pirozhkov. Admissible subcategories of del Pezzo surfaces. *arXiv preprint, arXiv: 2006.07643*, 2020.
- [PPZ19] Alexander Perry, Laura Pertusi, and Xiaolei Zhao. Stability conditions and moduli spaces for Kuznetsov components of Gushel-Mukai varieties. *To appear in Geom. Topol., arXiv: 1912.06935*, 2019.
- [PS22] Laura Pertusi and Paolo Stellari. Categorical Torelli theorems: results and open problems. *arXiv preprint arXiv:2201.03899*, 2022.
- [PY20] Laura Pertusi and Song Yang. Some remarks on Fano threefolds of index two and stability conditions. *To appear in Int. Math. Res. Not., arXiv:2004.02798*, 2020.
- [Vak17] Ravi Vakil. The rising sea: Foundations of Algebraic Geometry. <http://math.stanford.edu/~vakil/216blog/FOAGnov1817public.pdf>, 2017.
- [Yos99] Kota Yoshioka. Irreducibility of moduli spaces of vector bundles on K3 surfaces. *arXiv preprint math/9907001*, 1999.
- [Yos01] Kōta Yoshioka. Moduli spaces of stable sheaves on abelian surfaces. *Math. Ann.*, 321(4):817–884, 2001.
- [Zha20] Shizhuo Zhang. Bridgeland moduli spaces and Kuznetsov’s Fano threefold conjecture. *arXiv preprint arXiv:2012.12193*, 2020.

BEIJING INTERNATIONAL CENTER FOR MATHEMATICAL RESEARCH, PEKING UNIVERSITY, BEIJING, CHINA

Email address: hfguo@pku.edu.cn

COLLEGE OF MATHEMATICS, SICHUAN UNIVERSITY, CHENGDU, SICHUAN PROVINCE 610064 P. R. CHINA

Email address: zhiyuliu@stu.scu.edu.cn

SCHOOL OF MATHEMATICS, THE UNIVERSITY OF EDINBURGH, JAMES CLERK MAXWELL BUILDING, KINGS BUILDINGS, EDINBURGH, UNITED KINGDOM, EH9 3FD

Email address: Shizhuo.Zhang@ed.ac.uk