
RELATIVE REGULAR RIEMANN-HILBERT CORRESPONDENCE II

by

Luisa Fiorot, Teresa Monteiro Fernandes & Claude Sabbah

Abstract. We develop the theory of relative regular holonomic \mathcal{D} -modules with a smooth complex manifold S of arbitrary dimension as parameter space, together with their main functorial properties. In particular, we establish in this general setting the relative Riemann-Hilbert correspondence proved in a previous work in the one-dimensional case.

Contents

1. Introduction.....	1
2. A review on relative coherent and holonomic \mathcal{D} -modules.....	5
3. Regular holonomic $\mathcal{D}_{X \times S/S}$ -modules.....	9
4. Holonomic $\mathcal{D}_{X \times S/S}$ -modules of D-type and applications.....	20
5. Construction of the relative Riemann-Hilbert functor RH_X^S	35
6. Proof of Theorem 2.....	42
References.....	49

1. Introduction

In this article, we are concerned with holomorphic families of coherent \mathcal{D} -modules on a complex manifold X , parametrized by a complex manifold S , that is, coherent modules over the ring $\mathcal{D}_{X \times S/S}$ of linear relative differential operators with respect to the projection $p_X : X \times S \rightarrow S$ (simply denoted by p when no confusion is possible). More specifically, we consider families for which the characteristic variety in the relative cotangent space $(T^*X) \times S$ is contained in the product by S of a fixed closed conic Lagrangian analytic subset $\Lambda \subset T^*X$. Following [8, 29, 31], we call these systems *relative holonomic $\mathcal{D}_{X \times S/S}$ -modules*. Here are some examples.

2020 Mathematics Subject Classification. 14F10, 32C38, 32S40, 32S60, 35Nxx, 58J10.

Key words and phrases. Holonomic relative D -module, regularity, relative constructible sheaf, relative perverse sheaf.

The research of T. Monteiro Fernandes was supported by Fundação para a Ciência e Tecnologia, under the project: UIDB/04561/2020.

(1) Deligne considered in [6] the case of vector bundles E on $X \times S$ with a flat relative connection ∇ , and established an equivalence with the category of locally constant sheaves of coherent $p^{-1}\mathcal{O}_S$ -modules on $X \times S$. In this case, the relative characteristic variety is the product of the zero section T_X^*X by S .

(2) For any holonomic \mathcal{D}_X -module \mathcal{M} on X with characteristic variety Λ , the pullback $q^*\mathcal{M} = \mathcal{O}_{X \times S} \otimes_{q^{-1}\mathcal{O}_X} q^{-1}\mathcal{M}$ by the projection $q : X \times S \rightarrow X$ is naturally endowed with a $\mathcal{D}_{X \times S/S}$ -module structure, and the relative characteristic variety of \mathcal{M} is equal to $\Lambda \times S$. For (E, ∇) as in (1), the characteristic variety of $q^*\mathcal{M} \otimes_{\mathcal{O}_{X \times S}} E$ (equipped with its natural $\mathcal{D}_{X \times S/S}$ -module structure) is contained in $\Lambda \times S$.

(3) Some integral transformations from objects on X to objects on S have kernels which are such flat bundles (E, ∇) . One of them is the Fourier-Mukai transformation FM introduced by Laumon [19] and Rothstein [35], which attaches to any bounded complex of \mathcal{D} -modules with coherent cohomology on an abelian variety A a bounded complex of \mathcal{O} -modules with coherent cohomology on the moduli space A^\sharp of line bundles with integrable connection on A (cf. [39] and the references therein). It is obtained as the integral transform with kernel \mathcal{P} on $A \times A^\sharp$ associated to the Poincaré bundle on the product $A \times \text{Pic}^0(A)$. By construction, \mathcal{P} is equipped with a flat relative connection, i.e., is a $\mathcal{D}_{A \times A^\sharp/A^\sharp}$ -module. Then $q^*\mathcal{M} \otimes_{\mathcal{O}_{A \times A^\sharp}} \mathcal{P}$ is an instance of (2), and $\text{FM}(\mathcal{M})$ is the pushforward $\text{Rp}_*(q^*\mathcal{M} \otimes_{\mathcal{O}_{A \times A^\sharp}} \mathcal{P})$. It is an object of $\text{D}_{\text{coh}}^b(\mathcal{O}_{A^\sharp})$.

(4) Given a holonomic \mathcal{D}_X -module \mathcal{M} and holomorphic functions f_1, \dots, f_p on X defining a divisor $Y = \{\prod_i f_i = 0\}$, the S -analytic counterpart of [21, Prop. 13] asserts that the $\mathcal{D}_{X \times S/S}$ -submodule generated by $q^*\mathcal{M} \cdot (\prod_i f_i^{s_i})$ in the twisted coherent $\mathcal{D}_{X \times S/S}(*)(Y \times S)$ -module $q^*\mathcal{M}(*)(Y \times S) \cdot (\prod_i f_i^{s_i})$ is relative holonomic.

(5) In his construction of moduli spaces for regular holonomic \mathcal{D} -modules, Nitsure [32] fixes a divisor with normal crossings in X and deforms pre- \mathcal{D} -modules (extending the notion of vector bundle with flat logarithmic connection) relative to this divisor and its canonical stratification. The corresponding holomorphic family of regular holonomic \mathcal{D} -modules has its characteristic variety adapted to this stratification, hence of the form $\Lambda \times S$.

(6) Mixed twistor \mathcal{D} -modules (cf. [26]) are compound objects defined on the product of X by the complex line \mathbb{C} , whose module components are holomorphic families of holonomic \mathcal{D} -modules parametrized by $S = \mathbb{C}^*$ degenerating at $0 \in \mathbb{C}$ to a coherent module on the cotangent space T^*X . On S , the characteristic variety of each holonomic \mathcal{D} -module is, by definition, contained in a fixed Lagrangian variety Λ . Of particular interest are the regular mixed twistor \mathcal{D} -modules, which have furnished the first example of families we are dealing with (cf. [29]).

Our definition of relative holonomicity imposes the following: the only possible changes in the characteristic variety of the restricted \mathcal{D}_X -module to a fixed parameter, when the parameter varies, is a change of multiplicities on each irreducible component of Λ . This condition is reasonable, as shown by the previous examples.

Besides, not any relative holonomic $\mathcal{D}_{X \times S/S}$ -module \mathcal{K} can serve as the kernel of an integral transformation as in (3), because it cannot be ensured that, for a holonomic \mathcal{D}_X -module \mathcal{M} , the tensor product $q^*\mathcal{M} \otimes_{\mathcal{O}_{X \times S}} \mathcal{K}$ is $\mathcal{D}_{X \times S/S}$ -coherent, in contrast with the classical result when S is reduced to a point. Indeed, in general, Bernstein-Sato theory only applies on some open subset of S depending on \mathcal{M} , as exemplified in

[30, Ex. 2.4]. Fortunately, adding the regularity property—as defined in [31]—of the kernel \mathcal{K} and of \mathcal{M} overcomes this difficulty, as already shown in [8] when $\dim S = 1$.

In order to replace Bernstein-Sato theory, the main tool is the stability of regular holonomicity by \mathcal{D} -module pullback [8, Th. 2], that we generalize to the case where $\dim S$ can be arbitrary.

Theorem 1 (Stability by pullback). *Let $\mathcal{M} \in \mathbf{D}_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$ and let $f : Y \rightarrow X$ be a morphism of complex manifolds. Then ${}_{\mathcal{D}}f^*\mathcal{M} \in \mathbf{D}_{\text{rhol}}^b(\mathcal{D}_{Y \times S/S})$.*

A possible application of our approach is that, for an integral transformation as in (3) with regular holonomic kernel \mathcal{K} , the analysis of local properties of $q^*\mathcal{M} \otimes_{\mathcal{O}_{X \times S}}^L \mathcal{K}$ leads to a better understanding of the \mathcal{K} -transform ${}_{\mathcal{D}}p_*(q^*\mathcal{M} \otimes_{\mathcal{O}_{X \times S}}^L \mathcal{K})$ of \mathcal{M} .

Here are natural examples of regular holonomic $\mathcal{D}_{X \times S/S}$ -modules:

(a) Coherent $\mathcal{D}_{X \times S/S}$ -submodules of regular holonomic $\mathcal{D}_{X \times S}$ -modules, provided that the irreducible components of the characteristic variety of the latter decompose as products with respect to the product decomposition $T^*X \times T^*S$ (cf. Section 3.c); they can be called *integrable* regular holonomic $\mathcal{D}_{X \times S/S}$ -modules.

(b) In the setting of (4) above, let us assume that \mathcal{M} is regular holonomic. Then the coherent $\mathcal{D}_{X \times S/S}(*Y \times S)$ -module $q^*\mathcal{M}(*Y) \cdot \prod_i f_i^{s_i}$ is $\mathcal{D}_{X \times S/S}$ -regular holonomic, hence $\mathcal{D}_{X \times S/S}$ -coherent (cf. Example 4.36) and this enables us to recover the S -analytic version of the result of Maisonobe in (4) when \mathcal{M} is regular holonomic.

We have developed in [29] the notion of relative \mathbb{C} -constructible complex and relative perverse complex on $X \times S$, generalizing the notion of relative local system of coherent $p^{-1}\mathcal{O}_S$ -modules considered by Deligne. For each stratum X_α of a suitable complex stratification (X_α) of X , each cohomology of such a complex is locally isomorphic to $p_{X_\alpha}^{-1}G_\alpha$ for some *coherent* \mathcal{O}_S -module G_α . We prove here a Riemann-Hilbert correspondence for any $\dim S$ (the $\dim S = 1$ case was proved in [8]).

Theorem 2 (Relative Riemann-Hilbert correspondence). *The functors*

$$\begin{aligned} \mathbf{P}\text{Sol}_X &: \mathbf{D}_{\text{rhol}}^b(\mathcal{D}_{X \times S/S}) \longrightarrow \mathbf{D}_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S) \\ \text{RH}_X^S &: \mathbf{D}_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S) \longrightarrow \mathbf{D}_{\text{rhol}}^b(\mathcal{D}_{X \times S/S}) \end{aligned}$$

are quasi-inverse equivalences of categories.

The proof of the correspondence is now made possible owing to [28], as we construct the relative Riemann-Hilbert functor RH^S by means of the site $X_{\text{sa}} \times S$ introduced in [28] instead of the subanalytic site $X_{\text{sa}} \times S_{\text{sa}}$ considered in [27] and used in [31, 8]. The main results are the exact analogues of Theorems 2 and 1 in [8] for S of arbitrary dimension, with the same notation for relative regular holonomic complexes and S - \mathbb{C} -constructible complexes, and the same meaning for the perverse solution functor $\mathbf{P}\text{Sol}_X$. We change the order of the theorems with respect to loc. cit. since their proof is done the other way round in the present paper.

The strategy for the proof of these theorems is similar to that of [8]: it is made precise in Sections 4.e and 6. However, when $\dim S \geq 2$, we have to distinguish between S -torsion-freeness and S -flatness (also called strictness in loc. cit.). While this is not much trouble for some of the results, owing to the analysis of t-structures made in [7], we are led to using Rossi's flattening theorem of [34] at some point.

Along the way, we complement the results of [31, 8] by giving further characterizations and properties of regular holonomic $\mathcal{D}_{X \times S/S}$ -modules or objects of the bounded derived category $\mathcal{D}_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$, which are needed for the proof.

Comparison with other works on regular holonomic \mathcal{D} -modules with parameters

(1) In [5], the authors introduce a formal parameter \hbar and the corresponding rings $\mathbb{C}[[\hbar]]$, $\mathcal{O}_X[[\hbar]]$, $\mathcal{D}_X[[\hbar]]$ denoted by \mathcal{D}_X^\hbar . They define the notion of regular holonomic \mathcal{D}_X^\hbar -module by asking that the restriction to the closed point $\hbar = 0$ is a regular holonomic \mathcal{D}_X -module which is different from our point view since we ask regular holonomicity at any value of the parameter. They prove a Riemann-Hilbert correspondence with the category of \mathbb{C} -constructible complexes of $\mathbb{C}_X[[\hbar]]$ -modules by means of a functor they denote by TH_\hbar , which would correspond to the notation RH^S in the present paper. The main difference with the notions introduced in the present paper (or more precisely with those introduced in [8], which only considered the case where $\dim S = 1$) is that the sheaves considered in [5] are sheaves of $\mathbb{C}[[\hbar]]$ -modules on X , while we consider sheaves of $p^{-1}\mathcal{O}_S$ -modules ($p : X \times S \rightarrow S$ the projection) on the product space $X \times S$. The authors also consider the restriction at the generic point of $\mathbb{C}[[\hbar]]$, that is, modules over the ring $\mathcal{D}_X((\hbar))$ which has no counterpart in our setting.

(2) If instead of considering \mathcal{D}_X -modules one considers flat meromorphic bundles on X (i.e., meromorphic connections) with fixed pole divisor, there are many works in the algebraic or formal setting, with base fields or rings that can be different from \mathbb{C} . Since the literature is vast, let us only mention [1] and the more recent preprint [10]. In the latter preprint, the authors extend Deligne's equivalence result [6, Th. II.5.9] to an equivalence with a formal parameter. The corresponding equivalence in the present article would be that for $\mathcal{D}_{X \times S/S}$ -modules of D-type, as defined in Section 4.

(3) Wu [41] has established a Riemann-Hilbert correspondence similar to that of Theorem 2 in the case of Alexander complexes that occur in Example (4) above.

Organization of the paper. In Section 2, we review and complete various results on coherent $\mathcal{D}_{X \times S/S}$ -modules obtained in our previous works [29, 31, 7, 8]. We emphasize the behavior of holonomicity with respect to pullback and proper pushforward both with respect to X and to the parameter space S . Note that the parameter space S is always assumed to be a complex manifold, while one should be able to generalize various statements to any complex analytic space. For example, the sheaf $\mathcal{D}_{X \times T/T}$ is well-defined if T is a possibly singular and non reduced complex analytic space with sheaf of functions \mathcal{O}_T . In this work, we restrict the setting to those complex spaces T that are embeddable in a smooth complex manifold S , with ideal $\mathcal{I}_T \subset \mathcal{O}_S$, and we regard $\mathcal{D}_{X \times T/T}$ -modules as $\mathcal{D}_{X \times S/S}$ -modules annihilated by \mathcal{I}_T .

In Section 3, we complement various results of [8] on the regularity property. In particular, we give details on [8, Rem. 1.11]. Furthermore, the relation with the usual notion of regularity of holonomic $\mathcal{D}_{X \times S}$ -modules is made precise in Section 3.c. Stability under base pullback and proper base pushforward is established in Section 3.b (as usual, under a goodness assumption for proper pushforward). The case of pushforward with respect to a proper morphism $f : X \rightarrow Y$ (with a goodness assumption) has already been treated in [31], and stability by pullback, which is the content of Theorem 1 needs first a detailed analysis of holonomic $\mathcal{D}_{X \times S/S}$ -modules of D-type.

This analysis is performed in Section 4. The reasoning made in [31] when $\dim S = 1$, relying on the property that the torsion-free quotient of a coherent \mathcal{O}_S -module is locally free, has to be adapted by using base changes with respect to S , so that the base functoriality properties considered in Sections 2 and 3 are most useful. Theorem 1 is proved in Section 4.e in a way similar to that done in [8]. In Section 4.f, we give a characterization of relative regular holonomicity in terms of formal solutions.

Section 6 gives details on the main steps of the proof of Theorem 2. The strategy is similar to the one in [31], although we need various new technical details contained in Section 5. On the one hand, the construction of the Riemann-Hilbert functor RH^S is now performed by using the partial subanalytic site $X_{\mathrm{sa}} \times S$ introduced in [28]. On the other hand, the comparison between two definitions of Deligne's extension, one using the standard notion of moderate growth and the other one obtained via the complex of tempered holomorphic functions on the partial subanalytic site $X_{\mathrm{sa}} \times S$, is also done in Section 5, relying once more on results of [28]. We emphasize that the proof given here is simpler, when $\dim S = 1$, than that given in [31].

Acknowledgements. We thank Luca Prelli and Pierre Schapira for useful discussions and suggestions. We thank H el ene Esnault for pointing out to us the preprint [10], and its authors for interesting discussions.

2. A review on relative coherent and holonomic \mathcal{D} -modules

For complex analytic manifolds X and S , we denote by

$$p_X : X \times S \longrightarrow S \quad \text{and} \quad q_S : X \times S \longrightarrow X$$

the projections, and we use the notation p, q when there is no confusion possible. The sheaf of relative differential operators $\mathcal{D}_{X \times S/S}$ is naturally defined as

$$(2.1) \quad \mathcal{D}_{X \times S/S} = q^{-1} \mathcal{D}_X \otimes_{q^{-1} \mathcal{O}_X} \mathcal{O}_{X \times S} = \mathcal{O}_{X \times S} \otimes_{q^{-1} \mathcal{O}_X} q^{-1} \mathcal{D}_X.$$

2.a. Coherence, goodness and holonomicity. We adapt the definitions of [13, §4.7], which we refer to for properties and proofs.

Definition 2.2. We say that an $\mathcal{O}_{X \times S}$ -module \mathcal{F} is

(1) *\mathcal{O} -good on $X \times S$* if, on any relatively compact open set U of $X \times S$, $\mathcal{F}|_U$ is the direct limit of an increasing sequence of \mathcal{O}_U -coherent submodules (equivalently, the direct limit of an inductive system of \mathcal{O}_U -coherent modules).

(2) *\mathcal{O} -quasi-coherent* if each point of $X \times S$ has an open neighbourhood U on which \mathcal{F} is \mathcal{O} -good.

We say that a $\mathcal{D}_{X \times S/S}$ -module \mathcal{M} is

(3) *good on $X \times S$* if it is \mathcal{O} -good on $X \times S$ and $\mathcal{D}_{X \times S/S}$ -coherent.

The category $\mathrm{Mod}_{\mathcal{O}\text{-q-coh}}(\mathcal{O}_{X \times S})$ of \mathcal{O} -quasi-coherent $\mathcal{O}_{X \times S}$ -modules is an abelian full subcategory of $\mathrm{Mod}(\mathcal{O}_{X \times S})$ closed under extensions. Let Y be a hypersurface of X . We denote by $\bullet(*Y)$, instead of $\bullet(*(Y \times S))$ the localization functor along the hypersurface $Y \times S$ of $X \times S$. We will use the notation $X^* = X \setminus Y$.

Lemma 2.3.

- (1) Let \mathcal{L} be an \mathcal{O} -quasi-coherent module, then $\mathcal{L}(*Y)$ is \mathcal{O} -quasi-coherent.
- (2) Let \mathcal{L} be an \mathcal{O} -quasi-coherent module supported on $Y \times S$ and localized along $Y \times S$ (i.e., $\mathcal{L} \simeq \mathcal{L}(*Y)$). Then $\mathcal{L} = 0$.
- (3) Let $\mathcal{M} \xrightarrow{\phi} \mathcal{N}$ be a morphism of \mathcal{O} -quasi-coherent modules which are localized along $Y \times S$. If the restriction of ϕ at $X^* \times S$ is an isomorphism (resp. zero) then ϕ is an isomorphism (resp. zero).

Proof.

(1) By definition any point of $X \times S$ has an open neighbourhood U , which we can suppose to be a relatively compact open set, on which $\mathcal{L}|_U = \bigcup_i \mathcal{L}_i$ is the direct limit of an increasing sequence of \mathcal{O} -coherent submodules. Thus $\mathcal{L}|_U(*Y) = \bigcup_i \mathcal{L}_i(*Y)$ and any $\mathcal{L}_i(*Y) = \bigcup_n \mathcal{J}_{Y \times S}^{-n} \mathcal{L}_i$ which proves that $\mathcal{L}(*Y)$ is \mathcal{O} -quasi-coherent.

(2) The question is local. Since \mathcal{L} is an \mathcal{O} -quasi-coherent module supported on $Y \times S$, we can suppose (up to shrinking the neighborhood) that it is a direct limit $\mathcal{L} = \bigcup_i \mathcal{L}_i$ of an increasing sequence of \mathcal{O} -coherent submodules which are supported on $Y \times S$. Thus $\mathcal{L}(*Y) = \bigcup_i \mathcal{L}_i(*Y) = 0$ since $\mathcal{L}_i(*Y) = 0$ because \mathcal{L}_i are \mathcal{O} -coherent modules supported on $Y \times S$.

(3) Let us denote by \mathcal{L} and \mathcal{L}' respectively the kernel and the cokernel of ϕ . They are \mathcal{O} -quasi-coherent modules. If the restriction of ϕ to $X^* \times S$ is an isomorphism, they are supported on $Y \times S$. By applying the localization functor along Y , we get the following exact sequence of \mathcal{O} -quasi-coherent modules (by the first point)

$$0 \longrightarrow \mathcal{L}(*Y) \longrightarrow \mathcal{M}(*Y) \longrightarrow \mathcal{N}(*Y) \longrightarrow \mathcal{L}'(*Y) \longrightarrow 0.$$

By hypothesis $\mathcal{M}(*Y) \simeq \mathcal{M}$ and $\mathcal{N}(*Y) \simeq \mathcal{N}$, thus $\mathcal{L}(*Y) \simeq \mathcal{L}$, $\mathcal{L}'(*Y) \simeq \mathcal{L}'$ and by the previous point \mathcal{L} and \mathcal{L}' are zero. It $\phi|_{X^* \times S}$ is zero, $\mathcal{L} \rightarrow \mathcal{M}$ is an isomorphism on $X^* \times S$, hence it is an isomorphism and so ϕ is zero. q.e.d.

Let $\mathcal{D}_{X \times S/S}(*Y)$ be the coherent sheaf of rings $\mathcal{O}_{X \times S}(*Y) \otimes_{\mathcal{O}_{X \times S}} \mathcal{D}_{X \times S/S}$. Then any coherent $\mathcal{D}_{X \times S/S}$ - or $\mathcal{D}_{X \times S/S}(*Y)$ -module is \mathcal{O} -quasi-coherent. On the other hand, any \mathcal{O} -quasi-coherent $\mathcal{D}_{X \times S/S}$ -submodule \mathcal{N} of a coherent $\mathcal{D}_{X \times S/S}$ -module \mathcal{M} is $\mathcal{D}_{X \times S/S}$ -coherent. The next lemma follows by an easy adaptation of [13, Prop. 4.23].

Lemma 2.4.

- (1) The category of coherent, resp. \mathcal{O} -good, $\mathcal{D}_{X \times S/S}$ -modules, is abelian and stable by extensions in $\text{Mod}(\mathcal{D}_{X \times S/S})$, and $\mathcal{D}_{\text{coh}}^b(\mathcal{D}_{X \times S/S})$, resp. $\mathcal{D}_{\text{good}}^b(\mathcal{D}_{X \times S/S})$, is a triangulated full subcategory of $\mathcal{D}^b(\mathcal{D}_{X \times S/S})$.
- (2) Let \mathcal{M}^\bullet be a bounded complex of \mathcal{O} -good $\mathcal{D}_{X \times S/S}$ -modules. Then each cohomology $\mathcal{H}^\ell(\mathcal{M}^\bullet)$ is \mathcal{O} -good on $X \times S$.
- (3) On any relatively compact open subset $U \subseteq X \times S$, each good $\mathcal{D}_{X \times S/S}$ -module \mathcal{N} comes in an exact sequence

$$0 \longrightarrow \mathcal{N}' \longrightarrow \mathcal{D}_{X \times S/S} \otimes_{\mathcal{O}_{X \times S}} \mathcal{L} \longrightarrow \mathcal{N}|_U \longrightarrow 0$$

with \mathcal{L} being \mathcal{O}_U -coherent and \mathcal{N}' being good on U . In particular, $\mathcal{N}|_U$ has a coherent $F_\bullet \mathcal{D}_{X \times S/S}$ -filtration.

Remark 2.5.

- (1) With a slight adaptation of the proof of Lemma 2.3(1) we conclude that if \mathcal{M} is good (resp. $\mathcal{D}_{X \times S/S}$ -coherent) then $\mathcal{M}(*Y)$ is \mathcal{O} -good (resp. \mathcal{O} -quasi-coherent).

(2) If a local section m of an \mathcal{O} -quasi-coherent module satisfying $\mathcal{M} = \mathcal{M}(*Y)$ is zero when restricted to $X^* \times S$, then it is zero. Indeed if m is defined on $U \times U_S$, m generates an \mathcal{O} -coherent submodule of $\mathcal{M}(*Y)|_{U \times U_S}$ which is supported on $Y \times S$; given a local defining equation $f = 0$ for Y , we have $f^k m = 0$ for some k , thus $m = 0$.

Characteristic varieties. To any object \mathcal{M} of $\text{Mod}_{\text{coh}}(\mathcal{D}_{X \times S/S})$ is associated, by means of local coherent $\mathcal{O}_{X \times S}$ -filtrations, its characteristic variety $\text{Char}(\mathcal{M})$, which is contained in $T^*X \times S$. A coherent $\mathcal{D}_{X \times S/S}$ -module is *holonomic* if its characteristic variety is contained in $\Lambda \times S$, with Λ closed complex analytic Lagrangian \mathbb{C}^* -homogeneous in T^*X . Correspondingly are defined the derived categories $\text{D}_{\text{coh}}^{\text{b}}(\mathcal{D}_{X \times S/S})$ and $\text{D}_{\text{hol}}^{\text{b}}(\mathcal{D}_{X \times S/S})$ and the characteristic variety $\text{Char}(\mathcal{M})$ for such objects \mathcal{M} .

The structure of the characteristic variety of a holonomic $\mathcal{D}_{X \times S/S}$ -module \mathcal{M} is described in [7, Lem. 2.10]: for each irreducible component Λ_i of Λ ($i \in I$) there exists a locally finite family $(T_{ij})_{j \in J_i}$ of closed analytic subsets of S such that

$$(2.6) \quad \text{Char}(\mathcal{M}) = \bigcup_{i \in I} \bigcup_{j \in J_i} \Lambda_i \times T_{ij} = \bigcup_{i \in I} \Lambda_i \times T_i \quad \text{with } T_i = \bigcup_j T_{ij}.$$

The projection to X of Λ_i is an irreducible closed analytic subset of X that we denote by Z_i . These subsets form a locally finite family of closed analytic subsets of X . We have

$$(2.7) \quad \text{Supp}(\mathcal{M}) = \bigcup_{i \in I} Z_i \times T_i,$$

and we set

$$\text{Supp}_X(\mathcal{M}) = \bigcup_{i \in I} Z_i, \quad \text{Supp}_S(\mathcal{M}) = \bigcup_{i \in I} T_i,$$

that we call respectively the X -support (which is a closed analytic subset of X) and the S -support of \mathcal{M} (which may be not closed analytic if I is infinite). Anyway, we set $\dim \text{Supp}_S(\mathcal{M}) = \max_{i,j} \dim T_{ij} \leq \dim S$.

Any $\mathcal{D}_{X \times S/S}$ -coherent submodule or quotient module of \mathcal{M} is holonomic and its characteristic variety is the union of some irreducible components of $\text{Char}(\mathcal{M})$.

Lemma 2.8. *The category $\text{Mod}_{\text{hol}}(\mathcal{D}_{X \times S/S})$ of holonomic $\mathcal{D}_{X \times S/S}$ -modules is closed under taking extensions in the category $\text{Mod}(\mathcal{D}_{X \times S/S})$, and under taking sub-quotients in the category $\text{Mod}_{\text{coh}}(\mathcal{D}_{X \times S/S})$.*

We say that a local section m of a $\mathcal{D}_{X \times S/S}$ -module \mathcal{M} is an *S -torsion section* if it is annihilated by some holomorphic function on S . The *S -torsion submodule* \mathcal{M}_t of \mathcal{M} is the submodule consisting of S -torsion local sections. Note that if \mathcal{M} is a holonomic $\mathcal{D}_{X \times S/S}$ -module then the $\mathcal{D}_{X \times S/S}$ -submodule \mathcal{M}_t is holonomic since it is an \mathcal{O} -quasi coherent submodule of \mathcal{M} . We say that \mathcal{M} is *S -torsion-free* if $\mathcal{M}_t = 0$. We denote by $\mathcal{M}_{\text{tf}} : \mathcal{M}/\mathcal{M}_t$ the torsion-free quotient.

We recall that the duality functor \mathbf{D} for $\mathcal{D}_{X \times S/S}$ -modules was introduced in [29, Def. 3.4] and that $\text{D}_{\text{hol}}^{\text{b}}(\mathcal{D}_{X \times S/S})$ is stable under duality which is an involution.

2.b. Behaviour with respect to pullback, pushforward and external product

Notation 2.9.

(1) For a holomorphic map $f : X \rightarrow X'$, we also denote by f the morphism of complex manifolds $f \times \text{Id} : X \times S \rightarrow X' \times S$ and by ${}_{\text{D}}f^*$ and ${}_{\text{D}}f_*$ the pullback and pushforward functors in the derived category of relative \mathcal{D} -modules.

(2) For a morphism $\pi : S \rightarrow S'$ between analytic spaces, we denote by π^* and $R\pi_*$ the natural extension to the category of relative \mathcal{D} -modules of the similar functors defined on the categories of \mathcal{O} -modules

Definition 2.10. We say that a $\mathcal{D}_{X \times S/S}$ -module \mathcal{M} is *f-(\mathcal{O} -)good* (resp. *π -(\mathcal{O} -)good*) if it is (\mathcal{O} -)good in some neighbourhood of each fiber of f (resp. π).

We recall results concerning the behaviour with respect to a morphism of complex manifolds.

Proposition 2.11. *Let \mathcal{M} be a coherent $\mathcal{D}_{X \times S/S}$ -module.*

(1) *If $f : X' \rightarrow X$ is a holomorphic map of complex manifolds then, for each $\ell \in \mathbb{Z}$, $\mathcal{H}_D^\ell f^* \mathcal{M}$ is an f - \mathcal{O} -good $\mathcal{D}_{X' \times S/S}$ -module.*

(2) *Let $f : X \rightarrow X'$ be a proper holomorphic map. If \mathcal{M} is f -good (resp. f -good and holonomic), then for each ℓ , $\mathcal{H}_D^\ell f_* \mathcal{M}$ is $\mathcal{D}_{X' \times S/S}$ -coherent (resp. holonomic).*

Proof. We refer to [38, Th. 4.2 & Cor. 4.3] for the proof of (2) (or one can adapt the proof of [13, Th. 4.25 & 4.27]), while for the proof of (1) we can easily adapt the argument of [20, Prop. 2.1] which gives the absolute case. q.e.d.

Base pullback. Let $\pi : S' \rightarrow S$ be a morphism of complex analytic manifolds. We also denote by π the induced map $\text{Id} \times \pi : X \times S' \rightarrow X \times S$. We have

$$\mathcal{D}_{X \times S'/S'} = \mathcal{O}_{X \times S'} \otimes_{\pi^{-1} \mathcal{O}_{X \times S}} \pi^{-1} \mathcal{D}_{X \times S/S} =: \pi^* \mathcal{D}_{X \times S/S},$$

hence $\mathcal{D}_{X \times S'/S'}$ is a $\pi^{-1} \mathcal{D}_{X \times S/S}$ -module. One then defines in a natural way the functor $L\pi^* : \mathcal{D}(\mathcal{D}_{X \times S/S}) \rightarrow \mathcal{D}(\mathcal{D}_{X \times S'/S'})$ as $L\pi^*(\bullet) = \mathcal{D}_{X \times S'/S'} \otimes_{\pi^{-1} \mathcal{D}_{X \times S/S}}^L (\bullet)$.

The following is straightforward.

Lemma 2.12. *Let \mathcal{L} be an \mathcal{O} -quasi-coherent $\mathcal{O}_{X \times S}$ -module. Then each cohomology sheaf $L^j \pi^* \mathcal{L}$ is π - \mathcal{O} -good.*

Lemma 2.13. *The functor $L\pi^*$ induces a functor $\mathcal{D}_{\text{hol}}^b(\mathcal{D}_{X \times S/S}) \rightarrow \mathcal{D}_{\text{hol}}^b(\mathcal{D}_{X \times S'/S'})$. For a holonomic $\mathcal{D}_{X \times S/S}$ -module \mathcal{M} and for each j , the characteristic variety $\text{Char } L^j \pi^* \mathcal{M}$ is contained in the pullback $\pi^{-1} \text{Char } \mathcal{M}$ (here, π denotes the map $T^*X \times S' \rightarrow T^*X \times S$).*

Proof. By using that, locally, a coherent $\mathcal{D}_{X \times S/S}$ -module has a resolution of length $2 \dim X$ by free $\mathcal{D}_{X \times S/S}$ -modules of finite rank, one gets the first point for $\mathcal{D}_{\text{coh}}^b(\mathcal{D}_{X \times S/S})$. It suffices thus to prove the second point, which is a local question. One can find such a resolution which is strictly compatible with coherent filtrations relative to $F_\bullet \mathcal{D}_{X \times S/S}$. In such a way, one finds that for each j , $\text{Char } L^j \pi^* \mathcal{M}$ is contained in the support of $L\pi^* \text{gr}^F \mathcal{M}$, hence the assertion. q.e.d.

Base pushforward. We now consider pushforward by a proper morphism $\pi : S \rightarrow S'$. There is a natural morphism $\mathcal{D}_{X \times S'/S'} \rightarrow \pi_* \mathcal{D}_{X \times S/S}$ obtained as follows, since $\mathcal{D}_{X \times S'/S'}$ is $\mathcal{O}_{X \times S'}$ -locally free. We have, owing to (2.1),

$$\begin{aligned} R\pi_* \mathcal{D}_{X \times S/S} &\simeq R\pi_*(\mathcal{O}_{X \times S} \otimes_{\pi^{-1} \mathcal{O}_{X \times S'}}^L \pi^{-1} \mathcal{D}_{X \times S'/S'}) \\ &\simeq R\pi_* \mathcal{O}_{X \times S} \otimes_{\mathcal{O}_{X \times S'}}^L \mathcal{D}_{X \times S'/S'} \simeq R\pi_* \mathcal{O}_{X \times S} \otimes_{\mathcal{O}_{X \times S'}} \mathcal{D}_{X \times S'/S'}, \end{aligned}$$

and thus, by taking cohomology in degree zero,

$$\pi_* \mathcal{D}_{X \times S/S} \simeq \pi_* \mathcal{O}_{X \times S} \otimes_{\mathcal{O}_{X \times S'}} \mathcal{D}_{X \times S'/S'}.$$

The desired morphism is obtained by using the natural morphism $\mathcal{O}_{X \times S'} \rightarrow \pi_* \mathcal{O}_{X \times S}$. We recall [31, Prop. 1.6]:

Proposition 2.14. *Assume that π is proper and that \mathcal{M} is $\mathcal{D}_{X \times S/S}$ -holonomic and π -good. Then, for each j , $R^j \pi_* \mathcal{M}$ is $\mathcal{D}_{X \times S'/S'}$ -holonomic with characteristic variety contained in $\pi(\text{Char } \mathcal{M})$.*

An adjunction formula. Let $\pi : S' \rightarrow S$ be a morphism of complex manifolds. We will make use of the following adjunction formula.

Lemma 2.15. *Let \mathcal{M}, \mathcal{N} be objects of $\mathbf{D}_{\text{coh}}^b(\mathcal{D}_{X \times S/S})$. Then there is a bi-functorial isomorphism in $\mathbf{D}^b(\mathcal{D}_{X \times S/S})$:*

$$(2.15^*) \quad R\pi_* R\mathcal{H}om_{\mathcal{D}_{X \times S'/S'}}(L\pi^* \mathcal{M}, L\pi^* \mathcal{N}) \simeq R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, R\pi_* L\pi^* \mathcal{N}).$$

If \mathcal{N} is a coherent $\mathcal{D}_{X \times S/S}$ -module, there are functorial morphisms in $\mathbf{D}^b(\mathcal{D}_{X \times S/S})$:

$$(2.15^{**}) \quad \mathcal{N} \longrightarrow R\pi_* L\pi^* \mathcal{N} \longrightarrow R\pi_* \pi^* \mathcal{N},$$

and if π is proper, $R\pi_ L\pi^* \mathcal{N} \simeq (R\pi_* \mathcal{D}_{X \times S'/S'}) \otimes_{\mathcal{D}_{X \times S/S}}^L \mathcal{N}$ belongs to $\mathbf{D}_{\text{coh}}^b(\mathcal{D}_{X \times S/S})$.*

Proof. We have (cf. [13, p. 241])

$$R\mathcal{H}om_{\mathcal{D}_{X \times S'/S'}}(L\pi^* \mathcal{M}, L\pi^* \mathcal{N}) \simeq R\mathcal{H}om_{\pi^{-1} \mathcal{D}_{X \times S/S}}(\pi^{-1} \mathcal{M}, L\pi^* \mathcal{N}),$$

hence (2.15*) is obtained by adjunction (cf. [15, (2.6.15)]). By setting $\mathcal{M} = \mathcal{N}$ in (2.15*) we get $\text{Hom}_{\mathcal{D}_{X \times S'/S'}}(L\pi^* \mathcal{N}, L\pi^* \mathcal{N}) \simeq \text{Hom}_{\mathcal{D}_{X \times S/S}}(\mathcal{N}, R\pi_* L\pi^* \mathcal{N})$. The image of Id by this isomorphism is the first morphism in (2.15**) while the natural morphism $L\pi^* \mathcal{N} \rightarrow \pi^* \mathcal{N}$ in $\mathbf{D}^b(\mathcal{D}_{X \times S'/S'})$ provides the desired morphism $R\pi_* L\pi^* \mathcal{N} \rightarrow R\pi_* \pi^* \mathcal{N}$. The last isomorphism is obtained by the projection formula (cf. [15, Prop. 2.6.6]).
q.e.d.

Remark 2.16 ((External) tensor product). For $\mathcal{M}, \mathcal{N} \in \mathbf{D}^-(\mathcal{D}_{X \times S/S})$, the tensor product $\mathcal{M} \otimes_{\mathcal{O}_{X \times S}}^L \mathcal{N} \in \mathbf{D}^-(\mathcal{D}_{X \times S/S})$ is isomorphic to the pullback ${}_{\mathbf{D}}\delta^*(\mathcal{M} \boxtimes_{\mathcal{O}_S}^L \mathcal{N})$ of the S -external tensor product $\mathcal{M} \boxtimes_{\mathcal{O}_S}^L \mathcal{N}$ by $\delta : X \times S \hookrightarrow (X \times X) \times S$ (the diagonal embedding over S). If $\mathcal{M}, \mathcal{N} \in \mathbf{D}_{\text{coh}}^b(\mathcal{D}_{X \times S/S})$, $\mathcal{M} \boxtimes_{\mathcal{O}_S}^L \mathcal{N}$ belongs to $\mathbf{D}_{\text{coh}}^b(\mathcal{D}_{X \times X \times S/S})$ and we have $C := \text{Char}(\mathcal{M} \boxtimes_{\mathcal{O}_S}^L \mathcal{N}) \subset \text{Char}(\mathcal{M}) \times_S \text{Char}(\mathcal{N})$. This is seen by considering local resolutions of \mathcal{M} resp. \mathcal{N} by free $\mathcal{D}_{X \times S/S}$ -modules, showing both inclusions $C \subset T^*X \times \text{Char}(\mathcal{N})$ and $C \subset \text{Char}(\mathcal{M}) \times T^*X$. Therefore, if \mathcal{M}, \mathcal{N} are holonomic, so is $\mathcal{M} \boxtimes_{\mathcal{O}_S}^L \mathcal{N}$.

3. Regular holonomic $\mathcal{D}_{X \times S/S}$ -modules

In the special case of Lemma 2.13 where π is the inclusion $i_{s_o} : \{s_o\} \hookrightarrow S$ of a point s_o in S , we will often refer to [29, Prop. 3.1] for the compatibility between the functor $L i_{s_o}^*$ and the bi-functors $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\cdot, \cdot)$ and $R\mathcal{H}om_{p^{-1}\mathcal{O}_S}(\cdot, \cdot)$.

3.a. Characterization of relative regular holonomicity. The category of *regular holonomic* $\mathcal{D}_{X \times S/S}$ -modules was introduced in [31] as well as the full subcategory $\mathbf{D}_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$ of $\mathbf{D}_{\text{hol}}^b(\mathcal{D}_{X \times S/S})$ of bounded complexes of $\mathcal{D}_{X \times S/S}$ -modules having regular holonomic cohomology. According to [31], we say that an object $\mathcal{M} \in \mathbf{D}_{\text{hol}}^b(\mathcal{D}_{X \times S/S})$ is *regular* if it satisfies

(Reg 1) For each $s_o \in S$ and any $j \in \mathbb{Z}$, $Li_{s_o}^* \mathcal{H}^j(\mathcal{M}) \in \mathbf{D}_{\text{rhol}}^b(\mathcal{D}_X)$.

An alternative and natural property of regularity would be the following:

(Reg 2) For each $s_o \in S$, $Li_{s_o}^* \mathcal{M} \in \mathbf{D}_{\text{rhol}}^b(\mathcal{D}_X)$.

Regularity in either sense is the same property for objects of $\text{Mod}(\mathcal{D}_{X \times S/S})$. Property (Reg 1) is by definition compatible with the truncation functors while Property (Reg 2) is compatible with base change on S , meaning that, for any morphism $\pi : S' \rightarrow S$ of complex manifolds and any object $\mathcal{M} \in \mathbf{D}_{\text{hol}}^b(\mathcal{D}_{X \times S/S})$ which satisfies (Reg 2), the pullback $L\pi^*(\mathcal{M}) \in \mathbf{D}_{\text{hol}}^b(\mathcal{D}_{X' \times S/S})$ satisfies (Reg 2) too. We enlarge the setting for further use since both conditions make sense for any complex in $\mathbf{D}^b(\mathcal{D}_{X \times S/S})$.

Proposition 3.1. *Let Y be a hypersurface of X . On any complex manifold S ,*

(i) *for a complex in $\mathbf{D}_{\text{coh}}^b(\mathcal{D}_{X \times S/S})$ (resp. in $\mathbf{D}_{\text{coh}}^b(\mathcal{D}_{X \times S/S}(*Y))$), the condition (Reg 1) is equivalent to (Reg 2), and we denote both by (Reg),*

(ii) *the category of coherent $\mathcal{D}_{X \times S/S}$ (resp. coherent $\mathcal{D}_{X \times S/S}(*Y)$)-modules satisfying (Reg) is closed under taking extensions in the category $\text{Mod}(\mathcal{D}_{X \times S/S})$ (resp. in $\text{Mod}(\mathcal{D}_{X \times S/S}(*Y))$) and sub-quotients in the category $\text{Mod}_{\text{coh}}(\mathcal{D}_{X \times S/S})$ (resp. the category $\text{Mod}_{\text{coh}}(\mathcal{D}_{X \times S/S}(*Y))$),*

(iii) *the category of regular holonomic $\mathcal{D}_{X \times S/S}$ -modules is closed under taking extensions in $\text{Mod}(\mathcal{D}_{X \times S/S})$ and sub-quotients in $\text{Mod}_{\text{coh}}(\mathcal{D}_{X \times S/S})$.*

We note that (iii) follows from (i) and (ii) together with Lemma 2.8, so that we will focus on the latter properties. The proof in both cases is the same because it is based on the coherence of the rings involved. We provide it in the localized case.

Lemma 3.2. *Condition (Reg 1) implies (Reg 2).*

Proof. We argue by induction on the amplitude of the complex \mathcal{M} . We may assume that $\mathcal{M} \in \mathbf{D}_{\text{coh}}^{\geq 0}(\mathcal{D}_{X \times S/S}(*Y))$ and we consider the following distinguished triangle

$$(3.3) \quad \mathcal{H}^0 \mathcal{M} \longrightarrow \mathcal{M} \longrightarrow \tau^{\geq 1} \mathcal{M} \xrightarrow{+1}$$

(where $\tau^{\geq 1}$ is the truncation functor with respect to the natural t -structure on $\mathbf{D}_{\text{coh}}^b(\mathcal{D}_{X \times S/S}(*Y))$). Let us assume that \mathcal{M} satisfies (Reg 1), hence by definition both $\mathcal{H}^0 \mathcal{M}$ and $\tau^{\geq 1} \mathcal{M}$ satisfy (Reg 1). As remarked, $\mathcal{H}^0 \mathcal{M}$ satisfies (Reg 2) too and by induction on the amplitude of \mathcal{M} , $\tau^{\geq 1} \mathcal{M}$ satisfies (Reg 2). q.e.d.

Proof of Proposition 3.1. For $d \geq 0$, we denote by (Reg 1) $_d$, resp. (Reg 2) $_d$, the corresponding condition for $\dim S \leq d$. If $\dim S = 0$, (i) $_0$ holds true, and (ii) $_0$ is proved e.g. in [4, Th. 5.3.4]. We thus assume from now on that $d \geq 1$ and we proceed by induction on $d := \dim S$, denoting by (i) $_d$ and (ii) $_d$ the statements of the proposition restricted to $\dim S \leq d$. We will prove the following implications for $d \geq 1$:

- (a) (i) $_{d-1} \wedge$ (ii) $_{d-1} \implies$ (i) $_d$;
- (b) (i) $_d \wedge$ (ii) $_{d-1} \implies$ (ii) $_d$.

Let us start with (a). Assuming that both (i) $_{d-1}$ and (ii) $_{d-1}$ hold, we have to prove that (Reg 2) $_d \Rightarrow$ (Reg 1) $_d$. Due to the induction hypothesis (i) $_{d-1}$ we simply write (Reg) $_{d-1}$ for either (Reg 1) $_{d-1}$ or (Reg 2) $_{d-1}$.

We note that $\mathcal{M} \in \mathbf{D}_{\text{coh}}^b(\mathcal{D}_{X \times S/S}(*Y))$ satisfies (Reg 2) $_d$ if and only if for each smooth codimension-one germ (H, s_o) , $Li_H^* \mathcal{M}$ satisfies (Reg) $_{d-1}$. It is then enough to prove that, for such an \mathcal{M} , $Li_H^* \mathcal{H}^j \mathcal{M}$ satisfies (Reg) $_{d-1}$ for any j and H . We shall argue by induction on the amplitude of \mathcal{M} . We may assume that $\mathcal{M} \in \mathbf{D}_{\text{coh}}^{\geq 0}(\mathcal{D}_{X \times S/S}(*Y))$ and we consider the distinguished triangle (3.3). We deduce an isomorphism

$$(3.4) \quad \mathcal{H}^{-1} Li_H^* \mathcal{H}^0 \mathcal{M} \simeq \mathcal{H}^{-1} Li_H^* \mathcal{M}$$

and an exact sequence

$$0 \longrightarrow \mathcal{H}^0 Li_H^* \mathcal{H}^0 \mathcal{M} \longrightarrow \mathcal{H}^0 Li_H^* \mathcal{M} \longrightarrow \mathcal{H}^0 Li_H^* \tau^{\geq 1} \mathcal{M} \longrightarrow 0.$$

(Note that $\mathcal{H}^k Li_H^* \mathcal{H}^0 \mathcal{M} = 0$ for $k \neq 0, -1$.) Since $Li_H^* \mathcal{M}$ satisfies (Reg) $_{d-1}$, so does $\mathcal{H}^k Li_H^* \mathcal{M}$ ($k = -1, 0$), and so does $\mathcal{H}^{-1} Li_H^* \mathcal{H}^0 \mathcal{M}$ by (3.4). Since (ii) $_{d-1}$ is assumed to hold, any coherent sub-quotient of $\mathcal{H}^0 Li_H^* \mathcal{M}$ satisfies (Reg) $_{d-1}$, hence so does $\mathcal{H}^0 Li_H^* \mathcal{H}^0 \mathcal{M}$, which proves, following Lemma 3.2, that $Li_H^* \mathcal{H}^0 \mathcal{M}$ satisfies (Reg) $_{d-1}$. Thus $\mathcal{H}^0 \mathcal{M}$ satisfies (Reg 2) $_d$ and, by the distinguished triangle (3.3), $\tau^{\geq 1} \mathcal{M}$ satisfies (Reg 2) $_d$. Induction on the cohomological length applied to $\tau^{\geq 1} \mathcal{M}$ implies that $Li_H^* \mathcal{H}^j \mathcal{M}$ satisfies (Reg) $_{d-1}$ for any $j \geq 1$, which concludes the proof of (a) and we now simply write (Reg) $_d$.

Let us now prove (b). The extension property in (ii) $_d$ is clear. Let us consider stability by sub-quotients in $\text{Mod}_{\text{coh}}(\mathcal{D}_{X \times S/S}(*Y))$. Let $\mathcal{M} \in \text{Mod}_{\text{coh}}(\mathcal{D}_{X \times S/S}(*Y))$ satisfy (Reg) $_d$. Given any short exact sequence

$$(3.5) \quad 0 \longrightarrow \mathcal{M}_1 \longrightarrow \mathcal{M} \longrightarrow \mathcal{M}_2 \longrightarrow 0$$

of coherent $\mathcal{D}_{X \times S/S}(*Y)$ -modules, we wish to prove that \mathcal{M}_1 and \mathcal{M}_2 satisfy (Reg) $_d$. Owing to our assumption on \mathcal{M} , $Li_H^* \mathcal{M}$ satisfies (Reg) $_{d-1}$ for any smooth codimension-one germ $(H, s_o) \subset (S, s_o)$, and it is enough to prove that either $Li_H^* \mathcal{M}_1$ or $Li_H^* \mathcal{M}_2$ satisfies (Reg) $_{d-1}$. From (3.5) we obtain the long exact sequence:

$$(3.6) \quad 0 \longrightarrow \mathcal{H}^{-1} Li_H^* \mathcal{M}_1 \longrightarrow \mathcal{H}^{-1} Li_H^* \mathcal{M} \longrightarrow \mathcal{H}^{-1} Li_H^* \mathcal{M}_2 \\ \longrightarrow \mathcal{H}^0 Li_H^* \mathcal{M}_1 \longrightarrow \mathcal{H}^0 Li_H^* \mathcal{M} \longrightarrow \mathcal{H}^0 Li_H^* \mathcal{M}_2 \longrightarrow 0.$$

Since $Li_H^* \mathcal{M}$ satisfies (Reg) $_{d-1}$ then so do $\mathcal{H}^{-1} Li_H^* \mathcal{M}_1$ and $\mathcal{H}^0 Li_H^* \mathcal{M}_2$ owing to (ii) $_{d-1}$, and it remains to be proved that either $\mathcal{H}^{-1} Li_H^* \mathcal{M}_2$, or $\mathcal{H}^0 Li_H^* \mathcal{M}_1$, satisfies (Reg) $_{d-1}$. Let s be a local coordinate on S vanishing on H .

Let us denote by \mathcal{M}' the pullback of $\mathcal{M}'_2 := \text{tors}_H(\mathcal{M}_2)$ in \mathcal{M} and by $\mathcal{M}_{2,\text{tf}}$ the quotient $\mathcal{M}_2/\mathcal{M}'_2$. The commutative diagram below is Cartesian and its columns and rows are short exact sequences:

$$(3.7) \quad \begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{M}_1 & \longrightarrow & \mathcal{M}' & \longrightarrow & \mathcal{M}'_2 & \longrightarrow & 0 \\ & & \parallel & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathcal{M}_1 & \longrightarrow & \mathcal{M} & \longrightarrow & \mathcal{M}_2 & \longrightarrow & 0 \\ & & & & \downarrow & & \downarrow & & \\ & & & & \mathcal{M}_{2,\text{tf}} & \xlongequal{\quad} & \mathcal{M}_{2,\text{tf}} & & \end{array}$$

Since $\mathcal{M}_{2,\text{tf}}$ is i_H^* -acyclic, the exact sequence (3.6) for the middle column splits as

$$\begin{aligned} & \mathcal{H}^{-1}Li_H^*\mathcal{M}' \xrightarrow{\sim} \mathcal{H}^{-1}Li_H^*\mathcal{M}, \\ 0 \longrightarrow & \mathcal{H}^0Li_H^*\mathcal{M}' \longrightarrow \mathcal{H}^0Li_H^*\mathcal{M} \longrightarrow \mathcal{H}^0Li_H^*\mathcal{M}_{2,\text{tf}} \longrightarrow 0, \end{aligned}$$

which implies that $Li_H^*\mathcal{M}'$ and $Li_H^*\mathcal{M}_{2,\text{tf}}$ satisfy $(\text{Reg})_{d-1}$ by $(ii)_{d-1}$, hence \mathcal{M}' and $\mathcal{M}_{2,\text{tf}}$ satisfy $(\text{Reg})_d$. We now prove that $Li_H^*\mathcal{M}'_2$ satisfies $(\text{Reg})_{d-1}$, that will conclude the proof. Since \mathcal{M}'_2 is $\mathcal{D}_{X \times S/S}(*Y)$ -coherent, there exists locally an integer $k \geq 1$ such that $s^k\mathcal{M}'_2 = 0$. We will prove by induction on k that any $\mathcal{D}_{X \times S/S}(*Y)$ -coherent torsion quotient of a $\mathcal{D}_{X \times S/S}(*Y)$ -coherent module satisfying $(\text{Reg})_d$, satisfies $(\text{Reg})_d$ too.

If $k = 1$, we have $\mathcal{H}^{-1}Li_H^*\mathcal{M}'_2 \simeq \mathcal{H}^0Li_H^*\mathcal{M}'_2$, and, by $(ii)_{d-1}$ the latter satisfies $(\text{Reg})_{d-1}$, being a quotient of $\mathcal{H}^0Li_H^*\mathcal{M}'$.

If $k > 1$, we argue with the following Cartesian commutative diagram, analogous to (3.7):

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{M}_1 & \longrightarrow & \mathcal{M}'' & \longrightarrow & \mathcal{M}''_2 & \longrightarrow & 0 \\ & & \parallel & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathcal{M}_1 & \longrightarrow & \mathcal{M}' & \longrightarrow & \mathcal{M}'_2 & \longrightarrow & 0 \\ & & & & \downarrow & & \downarrow s & & \\ & & & & s \cdot \mathcal{M}'_2 & \xlongequal{\quad} & s \cdot \mathcal{M}'_2 & & \end{array}$$

By the induction hypothesis on k , $Li_H^*(s \cdot \mathcal{M}'_2)$ satisfies $(\text{Reg})_{d-1}$ since $s \cdot \mathcal{M}'_2$ is a $\mathcal{D}_{X \times S/S}(*Y)$ -coherent quotient of \mathcal{M}' which is annihilated by s^{k-1} . It follows that, by considering the middle vertical sequence, so does $Li_H^*\mathcal{M}''$. Since \mathcal{M}''_2 is a $\mathcal{D}_{X \times S/S}(*Y)$ -coherent quotient of \mathcal{M}'' annihilated by s , $Li_H^*(\mathcal{M}''_2)$ satisfies $(\text{Reg})_{d-1}$. It follows that, by considering the first horizontal sequence, so does $Li_H^*\mathcal{M}_1$, hence finally, by considering the middle horizontal sequence, so does $Li_H^*\mathcal{M}'_2$, as wanted. q.e.d.

Corollary 3.8. *The category $D_{\text{rhohol}}^b(\mathcal{D}_{X \times S/S})$ is a full triangulated subcategory of $D_{\text{coh}}^b(\mathcal{D}_{X \times S/S})$ stable by duality.*

Proof. We note that $(\text{Reg} 2)$ implies that $D_{\text{rhohol}}^b(\mathcal{D}_{X \times S/S})$ is a full triangulated subcategory of $D_{\text{hol}}^b(\mathcal{D}_{X \times S/S})$. Since the latter is a full triangulated subcategory of $D_{\text{coh}}^b(\mathcal{D}_{X \times S/S})$, the first assertion follows. Stability by duality follows from the same property in the absolute case (cf. e.g. [4, Th. 5.4.15 (4)], together with the isomorphism $Li_{s_o}^*DM \simeq DLi_{s_o}^*M$, which follows from [13, (A.10)] in a way similar to that used in [29, Prop. 2.1 & 3.1]. q.e.d.

3.b. Stability of regular holonomicity under base pullback and base push-forward

For a proper morphism $f : X \rightarrow Y$, it has been shown in [31, Cor. 2.4] that if \mathcal{M} is an object of $D_{\text{rhohol}}^b(\mathcal{D}_{X \times S/S})$ with f -good cohomology, then ${}_Df_*\mathcal{M}$ belongs to $D_{\text{rhohol}}^b(\mathcal{D}_{Y \times S/S})$. On the other hand, stability of regular holonomicity (and hence its coherence) by pullback ${}_Df^*$ has been shown in [8] only if $\dim S = 1$ as a consequence of the Riemann-Hilbert correspondence proved there. This will be obtained in general by the proof of Theorem 1 in Section 4.e. In this section, we consider on the other hand the behaviour with respect to base pullback and pushforward.

Proposition 3.9 (Stability under base pullback). *Let $\pi : S' \rightarrow S$ be a morphism of complex manifolds and let \mathcal{M} be an object of $D_{\text{rhoh}}^b(\mathcal{D}_{X \times S/S})$. Then $L\pi^*\mathcal{M}$ belongs to $D_{\text{rhoh}}^b(\mathcal{D}_{X \times S'/S'})$.*

Proof. We already know that $L\pi^*\mathcal{M}$ belongs to $D_{\text{rhoh}}^b(\mathcal{D}_{X \times S'/S'})$ by Lemma 2.13. Regularity follows from the isomorphism of functors $Li_{s'_o}^*L\pi^* \simeq Li_{\pi(s'_o)}^*$ for any $s'_o \in S'$.
q.e.d.

Let t_o be a *fat point* of S , that is, a complex subspace of S supported on a reduced point $|t_o| \in S$. In other words, the ideal $\mathcal{J}_{t_o} \subset \mathcal{O}_S$, which satisfies $\mathcal{J}_{t_o} \subset \mathfrak{m}_{|t_o|}$ (which is the ideal associated to $|t_o|$ such that its fiber in $|t_o|$ is the maximal ideal of $\mathcal{O}_{S,|t_o|}$), contains some power $\mathfrak{m}_{|t_o|}^k$. By abuse of notation we will still denote by \mathcal{J}_{t_o} (resp. $\mathfrak{m}_{|t_o|}$) the ideal $\mathcal{J}_{t_o,|t_o|}$ (resp. $\mathfrak{m}_{|t_o|,|t_o|}$) and also the sheaf $p^{-1}\mathcal{J}_{t_o}$ (resp. $p^{-1}\mathfrak{m}_{|t_o|}$). Let $i_{t_o} : t_o \rightarrow S$ denote the natural morphism of complex spaces defined by the surjective morphism $\mathcal{O}_S \rightarrow \mathcal{O}_S/\mathcal{J}_{t_o}$. For an $\mathcal{O}_{X \times S}$ -module, resp. a $\mathcal{D}_{X \times S/S}$ -module \mathcal{M} , we set

$$i_{t_o}^*\mathcal{M} := (p^{-1}\mathcal{O}_S/p^{-1}\mathcal{J}_{t_o}) \otimes_{p^{-1}\mathcal{O}_S} \mathcal{M},$$

that we regard in a natural way as a \mathcal{D}_X -module since $\mathcal{O}_{S,|t_o|}/\mathcal{J}_{t_o}$ is a finite-dimensional vector space. We define thereby the pullback functor

$$Li_{t_o}^* : D^b(\mathcal{D}_{X \times S/S}) \longrightarrow D^b(\mathcal{D}_X).$$

We note that, endowed with its natural structure of \mathcal{D}_X -module, $i_{t_o}^*\mathcal{D}_{X \times S/S}$ is coherent. As a consequence, if \mathcal{M} is $\mathcal{D}_{X \times S/S}$ -coherent, then $Li_{t_o}^*\mathcal{M}$ has \mathcal{D}_X -coherent cohomology. In other words, $Li_{t_o}^*$ induces a functor $D_{\text{coh}}^b(\mathcal{D}_{X \times S/S}) \rightarrow D_{\text{coh}}^b(\mathcal{D}_X)$.

Corollary 3.10. *Let \mathcal{M} be an object of $D_{\text{coh}}^b(\mathcal{D}_{X \times S/S})$. Then, for any fat point t_o of S , $Li_{t_o}^*\mathcal{M}$ belongs to $D_{\text{rhoh}}^b(\mathcal{D}_X)$ if and only if $Li_{|t_o|}^*\mathcal{M}$ does so.*

Proof. For $k \geq 0$ we set $\mathcal{J}_k = \mathcal{J}_{t_o} \cap \mathfrak{m}_{|t_o|}^k$ (with $\mathfrak{m}_{|t_o|}^0 := \mathcal{O}_S$), so that $\mathcal{J}_1 = \mathcal{J}_{t_o}$ and $\mathcal{J}_k = \mathfrak{m}_{|t_o|}^k$ for k large enough. It is enough to prove that $p^{-1}(\mathcal{J}_k/\mathcal{J}_{k+1}) \otimes_{p^{-1}\mathcal{O}_S}^L \mathcal{M}$ belongs to $D_{\text{rhoh}}^b(\mathcal{D}_X)$ for any k if and only if $Li_{|t_o|}^*\mathcal{M}$ does so. Since the sheaf $\mathcal{J}_k/\mathcal{J}_{k+1}$ is an $\mathcal{O}_S/\mathfrak{m}_{|t_o|}$ -module supported on $|t_o|$ whose fiber is a finite dimensional vector space, we have

$$\begin{aligned} p^{-1}(\mathcal{J}_k/\mathcal{J}_{k+1}) \otimes_{p^{-1}\mathcal{O}_S}^L \mathcal{M} &\simeq p^{-1}(\mathcal{J}_k/\mathcal{J}_{k+1}) \otimes_{p^{-1}(\mathcal{O}_S/\mathfrak{m}_{|t_o|})}^L (p^{-1}(\mathcal{O}_S/\mathfrak{m}_{|t_o|}) \otimes_{p^{-1}\mathcal{O}_S}^L \mathcal{M}) \\ &= (\mathcal{J}_k/\mathcal{J}_{k+1}) \otimes Li_{|t_o|}^*\mathcal{M}, \end{aligned}$$

and the conclusion follows.

q.e.d.

Setting 3.11. We consider a local setting where S is a polydisc Δ^d written as $\Delta^{d-1} \times \Delta = S' \times \Delta$, and we denote by $q : S \rightarrow S'$ the projection $(s', t) \mapsto s'$, and we keep the same notation after taking the product with X . Recall (cf. e.g. [13, Prop. A.14]) that the sheaves of rings $q^{-1}\mathcal{O}_{S'}$, $q^{-1}\mathcal{O}_{X \times S'}$ and $q^{-1}\mathcal{D}_{X \times S'/S'}$ are Noetherian.

Let $h(s', t) = t^k + \sum_{i=0}^{k-1} h_i(s')t^i$ be a Weierstrass polynomial, with h_i holomorphic on S' and let $T = h^{-1}(0)$. The equivalence between the categories of

- coherent \mathcal{O}_S -modules supported on T ,
- h -nilpotent coherent $q^{-1}\mathcal{O}_{S'}[t]$ -modules,
- h -nilpotent $q^{-1}\mathcal{O}_{S'}[t]$ -modules which are $q^{-1}\mathcal{O}_{S'}$ -coherent,

extends in a natural way to $\mathcal{O}_{X \times S}$, to $\mathcal{D}_{X \times S/S}$ and to $\mathcal{D}_{X \times S/S}(*Y)$ for a given hypersurface Y of X . For example, in one direction, if \mathcal{M} is $\mathcal{D}_{X \times S/S}$ -coherent and supported on $X \times T$, then \mathcal{M} is $q^{-1}\mathcal{D}_{X \times S'/S'}$ -coherent and each local section is annihilated by some power of h . By taking local $\mathcal{D}_{X \times S/S}$ -generators of \mathcal{M} , we conclude that there exists locally an integer $\ell \geq 1$ such that $h^\ell \mathcal{M} = 0$. Conversely, if \mathcal{M} satisfies the latter property, we can regard it as an h -nilpotent coherent $q^{-1}\mathcal{D}_{X \times S'/S'}[t]$ -module and the associated $\mathcal{D}_{X \times S/S}$ -module is $\mathcal{D}_{X \times S/S} \otimes_{q^{-1}\mathcal{D}_{X \times S'/S'}[t]} \mathcal{M}$.

By definition, the characteristic variety of a coherent $q^{-1}\mathcal{D}_{X \times S'/S'}$ -module is (locally) the support in $(T^*X) \times S$ of the graded coherent $q^{-1}\text{gr}^F \mathcal{D}_{X \times S'/S'}$ -module with respect to any local coherent $q^{-1}F_\bullet \mathcal{D}_{X \times S'/S'}$ -filtration. Such a module is said to be holonomic if this support is contained in $\Lambda \times S$ for some Lagrangian variety $\Lambda \subset T^*X$.

Remark 3.12 (Holonomic and regular holonomic $q^{-1}\mathcal{D}_{X \times S'/S'}$ -modules)

Given a coherent $\mathcal{D}_{X \times S/S}$ -module \mathcal{M} supported on T as above, one checks that a coherent $F_\bullet \mathcal{D}_{X \times S/S}$ -filtration is also a coherent $q^{-1}F_\bullet \mathcal{D}_{X \times S'/S'}$ -filtration by the above correspondence. As a consequence, such a module is $\mathcal{D}_{X \times S/S}$ -holonomic if and only if it is $q^{-1}\mathcal{D}_{X \times S'/S'}$ -holonomic.

On the other hand, we say that a holonomic $q^{-1}\mathcal{D}_{X \times S'/S'}$ -module \mathcal{M} is regular if, for any $s'_o \in S'$ and any $s_o \in q^{-1}(s'_o)$, the holonomic \mathcal{D}_X -module $i_{s'_o}^{-1}(Li_{q^{-1}(s'_o)}^* \mathcal{M})$ is regular.

Corollary 3.13. *A coherent $\mathcal{D}_{X \times S/S}$ -module \mathcal{M} supported on $X \times T$ is regular holonomic if and only if, when regarded as a $q^{-1}\mathcal{D}_{X \times S'/S'}$ -module, it is regular holonomic.*

Proof. By Remark 3.12, we only need to check regularity, and we can suppose from the start that \mathcal{M} is $\mathcal{D}_{X \times S/S}$ -holonomic supported on $X \times T$. We assume first that \mathcal{M} is $q^{-1}\mathcal{D}_{X \times S'/S'}$ -regular. We wish to prove that $Li_{s'_o}^* \mathcal{M}$ is \mathcal{D}_X -regular holonomic for any $s_o \in S$. It is enough to prove this for $s_o \in T$. Let us choose coordinates $(s'_1, \dots, s'_{d-1}, t)$ of $S' \times \Delta$ centered at s_o . There exists $k \geq 1$ such that $\mathcal{O}_S/(h^k) \otimes_{\mathcal{O}_S}^L \mathcal{M} \simeq \mathcal{M}[1] \oplus \mathcal{M}$, since $h^k \mathcal{M} = 0$ for some $k \geq 1$, so, by the assumption,

$$\mathcal{O}_S/(s'_1, \dots, s'_{d-1}) \otimes_{\mathcal{O}_S}^L [\mathcal{O}_S/(h^k) \otimes_{\mathcal{O}_S}^L \mathcal{M}] = \mathcal{O}_S/(s'_1, \dots, s'_{d-1}, h^k) \otimes_{\mathcal{O}_S}^L \mathcal{M}$$

is a regular holonomic \mathcal{D}_X -module. Since the support of $\mathcal{O}_S/(s'_1, \dots, s'_{d-1}, h^k)$ is a finite union of fat points in S (defined by the ideal generated by $h^k(0, t)$), one of which is supported at s_o , we conclude that $Li_{s'_o}^* \mathcal{M}$ is a regular holonomic \mathcal{D}_X -module by applying Corollary 3.10. The converse is proved similarly. q.e.d.

Theorem 3.14 (Stability under projective base pushforward)

Let $\pi : S \rightarrow S'$ be a projective morphism of complex manifolds and let \mathcal{M} be an object of $\mathbb{D}_{\text{hol}}^b(\mathcal{D}_{X \times S/S})$. Assume that the cohomology of \mathcal{M} is π -good. Then $R\pi_ \mathcal{M}$ belongs to $\mathbb{D}_{\text{hol}}^b(\mathcal{D}_{X \times S'/S'})$.*

Proof. By a standard argument, one can reduce to the case where \mathcal{M} is a single regular holonomic $\mathcal{D}_{X \times S/S}$ -module which is π -good. By Proposition 2.14, $R\pi_* \mathcal{M}$ belongs to $\mathbb{D}_{\text{hol}}^b(\mathcal{D}_{X \times S'/S'})$. Furthermore, we note that the question is local with respect to X and to S' .

Step 1: Reduction to the case where S' is a point. Since π is projective, we can regard π (locally with respect to S') as the composition of the inclusion $S \hookrightarrow \mathbb{P}^m \times S'$ and the projection $\mathbb{P}^m \times S' \rightarrow S'$ for a suitable m . Moreover, we note that the result is easy if π is a closed embedding. We can thus assume that π is a projection $S = \mathbb{P}^m \times S' \rightarrow S'$. For a regular holonomic $\mathcal{D}_{X \times S/S}$ -module, proving the $\mathcal{D}_{X \times S'/S'}$ -regularity of $R\pi_*\mathcal{M}$ amounts to proving the \mathcal{D}_X -regularity of $Li_{s'_o}^* R\pi_*\mathcal{M}$ for any $s'_o \in S'$. Let $\mathfrak{m}_{s'_o}$ be the maximal ideal sheaf of $\mathcal{O}_{S'}$ at s'_o . Let us set $\mathbb{P}_{s'_o}^m = \mathbb{P}^m \times \{s'_o\}$ and consider the following Cartesian square

$$\begin{array}{ccc} X \times \mathbb{P}_{s'_o}^m & \xrightarrow{\pi'} & X \times \{s'_o\} \\ i' \downarrow & & \downarrow i_{s'_o} \\ X \times S & \xrightarrow{\pi} & X \times S' \end{array}$$

Then we have

$$\begin{aligned} Li_{s'_o}^* R\pi_*\mathcal{M} &= i_{s'_o}^{-1}((\mathcal{O}_{X \times S'/p^{-1}(\mathfrak{m}_{s'_o})} \mathcal{O}_{X \times S'}) \otimes_{\mathcal{O}_{X \times S'}}^L R\pi_*\mathcal{M}) \\ &\simeq i_{s'_o}^{-1} R\pi_* (\pi^{-1}(\mathcal{O}_{X \times S'/p^{-1}(\mathfrak{m}_{s'_o})} \mathcal{O}_{X \times S'}) \otimes_{\pi^{-1}\mathcal{O}_{X \times S'}}^L \mathcal{M}) \\ &\simeq R\pi'_* i'^{-1} (\pi^{-1}(\mathcal{O}_{X \times S'/p^{-1}(\mathfrak{m}_{s'_o})} \mathcal{O}_{X \times S'}) \otimes_{\pi^{-1}\mathcal{O}_{X \times S'}}^L \mathcal{M}) \\ &\stackrel{(*)}{\simeq} R\pi'_* (\mathcal{O}_{X \times \mathbb{P}_{s'_o}^m} \otimes_{i'^{-1}\mathcal{O}_{X \times S}}^L i'^{-1}\mathcal{M}) \\ &= R\pi'_*(Li'^*\mathcal{M}), \end{aligned}$$

where the isomorphism $(*)$ follows by extension of scalars since \mathcal{M} is an $\mathcal{O}_{X \times S}$ -module. By Proposition 3.9, $Li'^*\mathcal{M}$ is regular holonomic, so that if we know the theorem for π' , we deduce it for π .

From now on, we assume that S' is a point. In such a case, S is a projective space \mathbb{P}^m . Since the question is local with respect to X and since S is compact, we can assume that the sets I and J_i occurring in (2.6) are finite, so the S -support of \mathcal{M} is a closed analytic subset of S . We argue by induction on the dimension of the S -support of \mathcal{M} .

Step 2: Case $\dim \text{Supp}_S \mathcal{M} = 0$. If the S -support of \mathcal{M} has dimension zero, it consists of a finite number of points, and it is enough to consider the case where the support consists of one point $s_o \in S$ and locally we can assume that there exists $k \geq 1$ such that, denoting by \mathfrak{m}_{s_o} the maximal ideal sheaf of s_o in S , we have $\mathfrak{m}_{s_o}^k \mathcal{M} = 0$. If we denote by t_o the fat point supported by s_o with ring $\mathcal{O}_S/\mathfrak{m}_{s_o}^k \mathcal{O}_S$, we conclude that $\mathcal{M} = i_{s_o*} i_{t_o}^* \mathcal{M}$. Therefore, $R\pi_*\mathcal{M} = R\pi_* i_{s_o*} i_{t_o}^* \mathcal{M} = i_{t_o}^* \mathcal{M}$. In this case, the theorem follows from Corollary 3.10.

Step 3: Case $\dim \text{Supp}_S \mathcal{M} \geq 1$. We recall that, in this step, π is the constant map on $S = \mathbb{P}^m$.

(i) Proof of the regularity of $\pi_*\mathcal{M}$. We now assume that the S -support T of \mathcal{M} has dimension $d \geq 1$ and we argue by induction on the dimension of the S -support of \mathcal{M} . We first prove that $\pi_*\mathcal{M}$ is regular holonomic (instead of all modules $R^k\pi_*\mathcal{M}$).

One checks that the $\mathcal{D}_{X \times S/S}$ -submodule \mathcal{M}' of \mathcal{M} consisting of local sections m such that the S -support of $\mathcal{D}_{X \times S/S} \cdot m$ has dimension $< d$ is holonomic (denoting by $T_{<d}$ the union of irreducible components of T of dimension $< d$, \mathcal{M}' is locally defined as

the sheaf of local sections annihilated by some power of the ideal $\mathcal{J}_{T_{<d}}$ of $T_{<d}$ in \mathcal{O}_S). By Proposition 3.1(iii), \mathcal{M}' is regular holonomic and, by the induction hypothesis, $\pi_*\mathcal{M}'$ is regular holonomic. It is thus enough to prove regularity of $\pi_*(\mathcal{M}/\mathcal{M}')$, and we can likewise assume that

(3.15) \mathcal{M} has no nonzero coherent submodule with S -support of dimension $< d$.

Let us denote by Λ a Lagrangian subvariety of T^*X such that the characteristic variety of \mathcal{M} is contained in $\Lambda \times S$. Since T is compact, it has a finite number of irreducible components and we index by I_d those which are of dimension d . We choose a point s_i on the smooth part of each T_i ($i \in I_d$). For the sake of simplicity, we denote by \mathbf{s}_o the finite set $\{s_i \mid i \in I_d\}$.

There is a natural morphism of \mathcal{D}_X -modules

$$(3.16) \quad \pi_*\mathcal{M} \longrightarrow i_{\mathbf{s}_o}^{-1}\mathcal{M}$$

which associates to a section $m \in \Gamma(U; \pi_*\mathcal{M}) = \Gamma(U \times S; \mathcal{M})$ (U open in X) its germ along $U \times \mathbf{s}_o$. Since \mathcal{M} has no S -torsion supported in dimension $< d$, this morphism is *injective*. Indeed, let $x_o \in U \subseteq X$ and let $m \in \Gamma(U \times S; \mathcal{M})$ belong to the kernel, that is, such that its germ at (x_o, \mathbf{s}_o) vanishes, i.e., m is zero in some neighbourhood of (x_o, \mathbf{s}_o) that we write $U \times V$, up to shrinking U . Let us suppose by contradiction that $m \neq 0$. Let us consider the coherent $\mathcal{D}_{U \times S/S}$ -submodule $\mathcal{D}_{U \times S/S} \cdot m \subset \mathcal{M}|_{U \times S}$ which is holonomic with S -support contained in T . By (3.15) $\mathcal{D}_{U \times S/S} \cdot m$ has S -support of dimension d . Hence any irreducible d -dimensional component of its S -support is a T_i for a suitable $i \in I_d$, but since m is zero on $U \times V$ this is not possible.

Let $\mathfrak{m}_{\mathbf{s}_o}$ denote the ideal sheaf of \mathbf{s}_o in \mathcal{O}_S . For each $k \geq 1$, we have the map

$$\pi_*\mathcal{M} \longrightarrow i_{\mathbf{s}_o}^{-1}\mathcal{M}/i_{\mathbf{s}_o}^{-1}\mathfrak{m}_{\mathbf{s}_o}^k\mathcal{M}.$$

The proposition will be proved if we prove that this morphism is injective for k large enough, since the right-hand side is regular holonomic by Corollary 3.10.

Its kernel \mathcal{N}_k is a coherent, hence holonomic, \mathcal{D}_X -submodule of $\pi_*\mathcal{M}$ and the sequence (\mathcal{N}_k) is decreasing with characteristic variety contained in Λ . It is thus stationary. Let $\mathcal{N} \subset \pi_*\mathcal{M}$ denote this constant value. We conclude that the map

$$\mathcal{N} \longrightarrow i_{\mathbf{s}_o}^{-1}\mathcal{M}/i_{\mathbf{s}_o}^{-1}\mathfrak{m}_{\mathbf{s}_o}^k\mathcal{M}$$

is zero for any k . Since $\mathcal{N} = \mathcal{N}_k$ for k large enough, we aim at proving that $\mathcal{N} = 0$.

The image of \mathcal{N} by (3.16) is contained in $\bigcap_k i_{\mathbf{s}_o}^{-1}\mathfrak{m}_{\mathbf{s}_o}^k\mathcal{M}$, and since (3.16) is injective, it suffices to prove that $\mathcal{M}' := \bigcap_k i_{\mathbf{s}_o}^{-1}\mathfrak{m}_{\mathbf{s}_o}^k\mathcal{M} = 0$. For $x_o \in X$, let us denote by $\mathcal{D}_{(x_o, \mathbf{s}_o)}$ the germ at (x_o, \mathbf{s}_o) of $\mathcal{D}_{X \times S/S}$ and similarly by $\mathcal{M}_{(x_o, \mathbf{s}_o)}$ that of \mathcal{M} . It suffices thus to prove that, for all $x_o \in X$, the germ $\mathcal{M}'_{(x_o, \mathbf{s}_o)} = \bigcap_k \mathfrak{m}_{\mathbf{s}_o}^k \mathcal{M}_{(x_o, \mathbf{s}_o)}$ is zero. We note that $\mathcal{M}'_{(x_o, \mathbf{s}_o)}$ is of finite type over $\mathcal{D}_{(x_o, \mathbf{s}_o)}$ since the latter ring is Noetherian, hence $\mathcal{M}'_{(x_o, \mathbf{s}_o)}$ is holonomic with characteristic variety contained in $\Lambda_{x_o} \times (S, \mathbf{s}_o)$, where Λ_{x_o} is the germ of Λ along $T_{x_o}^*X$. Furthermore, it satisfies $\mathcal{M}'_{(x_o, \mathbf{s}_o)} = \mathfrak{m}_{\mathbf{s}_o} \mathcal{M}'_{(x_o, \mathbf{s}_o)}$. The Nakayama-type argument given in the proof of [31, Prop. 1.9(1)] shows that $\mathcal{M}'_{(x_o, \mathbf{s}_o)} = 0$ for each x_o . This concludes the proof of Step 3(i).

(ii) Regularity of $R^k\pi_*\mathcal{M}$. We use the result of (i) in order to prove holonomicity and regularity of $R^k\pi_*\mathcal{M}$ for all k .

For any $\ell \in \mathbb{Z}$, we consider the line bundle $\mathcal{O}_S(\ell)$ that we can realize as $\mathcal{O}_S(\ell H)$ for any hyperplane H of S . For a $\mathcal{D}_{X \times S/S}$ -module or an $\mathcal{O}_{X \times S}$ -module \mathcal{N} , we set $\mathcal{N}(\ell) = p^{-1}\mathcal{O}_S(\ell) \otimes_{p^{-1}\mathcal{O}_S} \mathcal{N}$.

We apply the technical lemma 3.17 below to \mathcal{M} . We note that $\mathcal{M}(\ell)$ is $\mathcal{D}_{X \times S/S}$ regular holonomic: indeed, this is a local property on S , and locally $\mathcal{O}_S(\ell) \simeq \mathcal{O}_S$. Also, $\mathcal{M}(\ell)$ is π -good. Furthermore, we choose a hyperplane $H \subset S$ such that $\dim H \cap T < \dim T$, and we realize $\mathcal{O}_S(\ell)$ as $\mathcal{O}_S(\ell H)$. We also consider $\mathcal{M}(\ell H)$ with such a choice of H . From Step 3(i), we know that $\pi_*\mathcal{M}(\ell H)$ is \mathcal{D}_X -regular. Let us consider $\mathcal{O}_{\ell H}$ given by the exact sequence

$$0 \longrightarrow \mathcal{O}_S \longrightarrow \mathcal{O}_S(\ell H) \longrightarrow \mathcal{O}_{\ell H} \longrightarrow 0$$

giving rise to the distinguished triangle

$$\mathcal{M} \longrightarrow \mathcal{M}(\ell H) \longrightarrow \mathcal{N} \xrightarrow{+1}$$

where \mathcal{N} is a complex in $\mathbf{D}_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$ with S -support contained in $T \cap H$. We deduce the long exact sequence

$$0 \longrightarrow H^{-1}R\pi_*\mathcal{N} \longrightarrow \pi_*\mathcal{M} \longrightarrow \pi_*\mathcal{M}(\ell H) \longrightarrow H^0R\pi_*\mathcal{N} \longrightarrow R^1\pi_*\mathcal{M} \longrightarrow 0$$

and the isomorphisms $R^{k+1}\pi_*\mathcal{M} \simeq H^kR\pi_*\mathcal{N}$ for $k \geq 1$. The induction hypothesis implies the regularity of $R^{k+1}\pi_*\mathcal{M}$ for $k \geq 1$ and, together with Corollary 3.8, it implies that of $\pi_*\mathcal{M}$ and $R^1\pi_*\mathcal{M}$. q.e.d.

Lemma 3.17. *Let \mathcal{N} be a coherent π -good $\mathcal{D}_{X \times S/S}$ -module. Then, for any $x_o \in X$, there exists a neighbourhood $\text{nb}(x_o) \subset X$ and an integer $\ell \geq 0$ such that, for any $k \geq 1$, $R^k\pi_*\mathcal{N}(\ell)|_{\text{nb}(x_o)} = 0$.*

A similar result is well-known to hold for a coherent $\mathcal{O}_{X \times S}$ -module, as a consequence of Grauert-Remmert's Theorems A and B (cf. [9] and [3, Th. IV.2.1]).

Proof. For any $N \geq 1$, we consider a resolution of \mathcal{N} as in Lemma 2.4(3) on $\text{nb}(x_o) \times S$. We thus obtain a distinguished triangle on this neighbourhood:

$$\mathcal{N}'[N] \longrightarrow \mathcal{D}_{X \times S/S} \otimes_{\mathcal{O}_{X \times S}} \mathcal{L}_\bullet \longrightarrow \mathcal{N} \xrightarrow{+1}.$$

Let us choose $N = 2m + 1$ and let ℓ be an integer given by the Grauert-Remmert theorems for \mathcal{L}_i ($i = -N, \dots, 0$) in $\text{nb}(x_o)$. We consider the same triangle obtained after tensoring with $p^{-1}\mathcal{O}_S(\ell)$. Since π has cohomological dimension $2m$, we find that $R^k\pi_*\mathcal{N}'(\ell)[N] = 0$ for any $k \geq 0$. On the other hand, on $\text{nb}(x_o)$ (cf. (2.1)),

$$\begin{aligned} R\pi_*(\mathcal{D}_{X \times S/S} \otimes_{\mathcal{O}_{X \times S}} \mathcal{L}_i(\ell)) &\simeq R\pi_*(\pi^{-1}\mathcal{D}_X \otimes_{\pi^{-1}\mathcal{O}_X} \mathcal{L}_i(\ell)) \\ &\simeq \mathcal{D}_X \otimes_{\mathcal{O}_X} R\pi_*\mathcal{L}_i(\ell) = \mathcal{D}_X \otimes_{\mathcal{O}_X} \pi_*\mathcal{L}_i(\ell). \end{aligned}$$

Since $R\pi_*\mathcal{N}(\ell)|_{\text{nb}(x_o)}$ is in non negative degrees it follows that $R\pi_*\mathcal{N}(\ell)|_{\text{nb}(x_o)}$ is isomorphic to the complex $\tau^{\geq 0}(\mathcal{D}_X \otimes_{\mathcal{O}_X} \pi_*\mathcal{L}_\bullet(\ell))$ and since the latter is in non-positive degrees, we conclude that $R\pi_*\mathcal{N}(\ell)|_{\text{nb}(x_o)} \simeq \pi_*\mathcal{N}(\ell)|_{\text{nb}(x_o)}$. q.e.d.

3.c. Integrable regular holonomic $\mathcal{D}_{X \times S/S}$ -modules

The following proposition, answering a question of Lei Wu, shows that, under a suitable condition on the characteristic variety, a coherent $\mathcal{D}_{X \times S/S}$ -submodule of a regular holonomic $\mathcal{D}_{X \times S}$ -module is relatively regular holonomic. We call such $\mathcal{D}_{X \times S/S}$ -modules *integrable* since their relative connection can be lifted as an integrable connection, i.e., a $\mathcal{D}_{X \times S}$ -module structure. The condition on the characteristic variety is due to the restrictive definition of a holonomic $\mathcal{D}_{X \times S/S}$ -module (cf. after Lemma 2.4).

Proposition 3.18. *Let X and S be complex manifolds and let \mathcal{N} be a regular holonomic $\mathcal{D}_{X \times S}$ -module. Assume that the characteristic variety of \mathcal{N} is contained in $\Lambda \times T^*S$ for some conic Lagrangian closed analytic subset Λ in T^*X . Let \mathcal{M} be a coherent $\mathcal{D}_{X \times S/S}$ -submodule of \mathcal{N} . Then \mathcal{M} is a regular holonomic $\mathcal{D}_{X \times S/S}$ -module.*

Let us first recall two properties ((3.19) and (3.20) below) of regular holonomic $\mathcal{D}_{X \times S}$ -modules that will be useful for the proof and do not depend on the assumption on $\text{Char } \mathcal{N}$ made in the proposition.

Let \mathcal{N} be any regular holonomic $\mathcal{D}_{X \times S}$ -module and let \mathcal{M} be a coherent $\mathcal{D}_{X \times S/S}$ -submodule of \mathcal{N} . Any irreducible component of $\text{Char } \mathcal{N}$ projects to $X \times S$ as an irreducible closed analytic subset Z and, denoting by Z° the smooth part of Z , this irreducible component is the closure of the conormal bundle $T_{Z^\circ}^*(X \times S)$, that we denote by $T_Z^*(X \times S)$. We also denote by $Z^{\circ\circ}$ the open set of the smooth locus Z° on which $p_{Z^\circ} : Z^\circ \rightarrow S$ has maximal rank. The closure in the relative cotangent space $T^*(X \times S/S)$ of the relative conormal bundle $T_{p|Z^{\circ\circ}}^*(X \times S/S)$ is denoted by $T_{p|Z}^*(X \times S/S)$, and is a conic analytic subset of $T^*(X \times S/S)$.

(3.19) *According to [36, Th. 3.2] any irreducible component of $\text{Char } \mathcal{M}$ is equal to $T_{p|Z}^*(X \times S/S)$ for some Z such that $T_Z^*(X \times S)$ is an irreducible component of $\text{Char } \mathcal{N}$.*

Assume furthermore that $\dim S = 1$. For each $s_o \in S$, we denote by s a local coordinate on S vanishing at s_o . We postpone after the proof of Proposition 3.18 that of the following classical result.

(3.20) *Under the above assumptions, the kernel and cokernel of $s : \mathcal{M} \rightarrow \mathcal{M}$ are regular holonomic \mathcal{D}_X -modules.*

Proof of Proposition 3.18, Step 1. We now add the assumption on $\text{Char } \mathcal{N}$. We first show that \mathcal{M} is relative holonomic with characteristic variety contained in $\Lambda \times S$. We note that $T_Z^*(X \times S)$ is contained in $\Lambda \times T^*S$ if and only if Z decomposes as the product $Y \times T$ for some irreducible closed analytic subsets $Y \subset X$ and $T \subset S$ and T_Y^*X is an irreducible component of Λ (this seen by considering first the smooth part Z°). It is then easily seen that $T_{p|Z}^*(X \times S/S) = (T_Y^*X) \times T$, hence is contained in $\Lambda \times S$. The conclusion follows from (3.19). q.e.d.

Proof of Proposition 3.18, Step 2. It remains to show the relative regularity of \mathcal{M} . We argue by induction on $d = \dim S$. If $d = 1$, relative regularity is provided by (3.20), since we already have relative holonomicity by Step 1. We thus assume that the statement of the proposition holds whenever $\dim S \leq d - 1$ and we assume $\dim S = d \geq 2$. The question being local, we fix a local coordinate s in S , defining a smooth

hypersurface $H := \{s = 0\}$. Locally, we can assume that $S = H \times \mathbb{C}$. According to Proposition 3.1, we are reduced to proving that

(3.21) *the kernel and cokernel of $s: \mathcal{M} \rightarrow \mathcal{M}$ are regular holonomic $\mathcal{D}_{X \times H/H}$ -modules.*

For the sake of clarity we set

$$X' = X \times H, \quad S' = \mathbb{C}, \quad p' : X' \times S' \longrightarrow S' \quad \text{so that} \quad X = X' \times S', \quad S = H \times S',$$

We denote by \mathcal{M}_1 the $\mathcal{D}_{X' \times S'/S'}$ -submodule of the regular holonomic $\mathcal{D}_{X' \times S'}$ -module \mathcal{N} generated by \mathcal{M} . It is $\mathcal{D}_{X' \times S'/S'}$ -coherent, so by (3.20), the kernel \mathcal{N}_1 and the cokernel \mathcal{N}'_1 of $s : \mathcal{M}_1 \rightarrow \mathcal{M}_1$ are $\mathcal{D}_{X'}$ -regular holonomic.

Claim. *The characteristic varieties $\text{Char } \mathcal{N}_1$ and $\text{Char } \mathcal{N}'_1$ are contained in $\Lambda \times T^*H$.*

Proof. We apply (3.19) to $\mathcal{M}_1 \subset \mathcal{N}$ and to the map $X' \times S' \rightarrow S'$. Since any irreducible component of $\text{Char } \mathcal{N}$ takes the form $T_Y^*X \times T_T^*S = T_Z^*(X' \times S')$ with $Z = Y \times T$, any irreducible component of $\text{Char } \mathcal{M}_1$ takes the form $T_{p'_Z}^*(X' \times S'/S')$ for some such Z . Denoting by $q : S \rightarrow S'$ the projection $H \times S' \rightarrow S'$, the latter space reads

$$(T_Y^*X) \times T_{q|_T}^*(H \times S'/S').$$

The fiber of the composition $T_{q|_T}^*(H \times S'/S') \rightarrow T \rightarrow S'$ above $s = 0$ (s is the coordinate on S') is contained in T^*H and is Lagrangian: indeed, it has dimension $\dim H$ since $T_{s|_T}^*(H \times S'/S')$ has dimension $\dim H + 1$, and it is easily seen to be isotropic. We denote it by Λ_T . The characteristic varieties of \mathcal{N}_1 and \mathcal{N}'_1 are contained in the fiber above $s = 0$ of $\text{Char } \mathcal{M}_1$, hence in the union, over Y, T occurring in $\text{Char } \mathcal{N}$, of the Lagrangian subsets $T_Y^*X \times \Lambda_T$, as claimed, since Λ is nothing but $\bigcup_Y T_Y^*X$.
q.e.d.

As a consequence, \mathcal{N}_1 is a regular holonomic $\mathcal{D}_{X \times H}$ -module satisfying the assumption of the proposition, and $\ker s : \mathcal{M} \rightarrow \mathcal{M}$ is a coherent $\mathcal{D}_{X \times H/H}$ -submodule of it. We can thus apply the induction hypothesis to conclude that (3.21) holds for $\ker s$.

On the other hand, for each $k \geq 0$, by applying (3.20) to $s^k \mathcal{M}_1$, we obtain, according to the claim, that $s^k \mathcal{M}_1 / s^{k+1} \mathcal{M}_1$ is a regular holonomic $\mathcal{D}_{X \times H}$ -module with characteristic variety contained in $\Lambda \times T^*H$. Then

$$\mathcal{M}^{(k)} := (\mathcal{M} \cap s^k \mathcal{M}_1) / (\mathcal{M} \cap s^{k+1} \mathcal{M}_1)$$

is a coherent $\mathcal{D}_{X \times H/H}$ -submodule of $s^k \mathcal{M}_1 / s^{k+1} \mathcal{M}_1$. The induction hypothesis implies that $\mathcal{M}^{(k)}$ is $\mathcal{D}_{X \times H/H}$ -regular holonomic. It follows by Corollary 3.8 that $\mathcal{M} / (\mathcal{M} \cap s^{k+1} \mathcal{M}_1)$ is also a regular holonomic $\mathcal{D}_{X \times H/H}$ -module. It remains to notice that coherence implies that, locally on X , there exists k such that $\mathcal{M} \cap s^{k+1} \mathcal{M}_1 \subset s\mathcal{M}$, so that $\mathcal{M}/s\mathcal{M}$ is a coherent quotient of a regular holonomic $\mathcal{D}_{X \times H/H}$ -module. Corollary 3.8 implies that it is also a regular holonomic $\mathcal{D}_{X \times H/H}$ -module, concluding the proof of (3.21).
q.e.d.

Proof of (3.20). We consider the Kashiwara-Malgrange V -filtration $V_\bullet \mathcal{N}$ of \mathcal{N} relative to the function s (cf. e.g. [24, 22]). This is a filtration indexed by \mathbb{Z} which satisfies, owing to the regularity of \mathcal{N} , the following properties:

- $V_k \mathcal{N}$ is $\mathcal{D}_{X \times S/S}$ -coherent for any $k \in \mathbb{Z}$ ([24, Th. (4.12.1)]),
- for $k \leq -1$, the multiplication by s on $V_k \mathcal{N}$ is injective and has image $V_{k-1} \mathcal{N}$ (cf. e.g. [24, Prop. (4.5.2)]),

- each $\mathrm{gr}_k^V \mathcal{N}$ is a regular holonomic \mathcal{D}_X -module (cf. e.g. [22, Cor. 4.7-5]).

From the second point we deduce that, for $k \leq -1$, $Li_{s_o}^* V_k \mathcal{N} = i_{s_o}^* V_k \mathcal{N} = \mathrm{gr}_k^V \mathcal{N}$, and by the third point the latter is \mathcal{D}_X -regular holonomic. For $k \geq 0$, $V_k \mathcal{N}$ is a successive extension of $V_{-1} \mathcal{N}$ by regular holonomic \mathcal{D}_X -modules $\mathrm{gr}_\ell^V \mathcal{N}$. Therefore, for each k , the kernel and cokernel of $s : V_k \mathcal{N} \rightarrow V_k \mathcal{N}$ are \mathcal{D}_X -regular holonomic, as one can prove by induction on k since it is clear for $k < 0$, and the inductive step follows by considering the snake lemma applied to the following commutative diagram of exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & V_{k-1} \mathcal{N} & \longrightarrow & V_k \mathcal{N} & \longrightarrow & \mathrm{gr}_k^V \mathcal{N} \longrightarrow 0 \\ & & \downarrow s & & \downarrow s & & \downarrow 0 \\ 0 & \longrightarrow & V_{k-1} \mathcal{N} & \longrightarrow & V_k \mathcal{N} & \longrightarrow & \mathrm{gr}_k^V \mathcal{N} \longrightarrow 0 \end{array}$$

We deduce that $sV_k \mathcal{N}/s^\ell V_k \mathcal{N}$ is regular holonomic for any $\ell \geq 1$. As any $\mathcal{D}_{X \times S/S}$ -coherent submodule of \mathcal{N} , locally on $X \times S$, \mathcal{M} is contained in $V_k \mathcal{N}$ for some $k \gg 0$ and, by the Artin-Rees lemma, $s\mathcal{M}$ contains $\mathcal{M} \cap s^\ell V_k \mathcal{N}$ for some $\ell \geq 1$. Since $\ker s : \mathcal{M} \rightarrow \mathcal{M}$ is contained in $\ker s : V_k \mathcal{N} \rightarrow V_k \mathcal{N}$, the former is regular holonomic since the latter is so. On the other hand, by considering the inclusion and the quotient maps

$$sV_k \mathcal{N}/s^\ell V_k \mathcal{N} \longleftarrow (\mathcal{M} \cap sV_k \mathcal{N})/(\mathcal{M} \cap s^\ell V_k \mathcal{N}) \longrightarrow (\mathcal{M} \cap sV_k \mathcal{N})/s\mathcal{M}$$

we conclude that $(\mathcal{M} \cap sV_k \mathcal{N})/s\mathcal{M}$ is also \mathcal{D}_X -regular holonomic. Furthermore, by considering the exact sequence

$$0 \longrightarrow (\mathcal{M} \cap sV_k \mathcal{N})/s\mathcal{M} \longrightarrow \mathcal{M}/s\mathcal{M} \longrightarrow V_k \mathcal{N}/sV_k \mathcal{N},$$

we conclude that $\mathcal{M}/s\mathcal{M}$ is \mathcal{D}_X -regular holonomic.

q.e.d.

4. Holonomic $\mathcal{D}_{X \times S/S}$ -modules of D-type and applications

4.a. S -locally constant sheaves and their associated relative connections

Let X be a connected complex manifold and let L be an S -locally constant sheaf of $p^{-1}\mathcal{O}_S$ -modules on $X \times S$ (cf. [31, App.]). For any $x_o \in X$, L is uniquely determined from a monodromy representation $\pi_1(X, x_o) \rightarrow \mathrm{Aut}_{\mathcal{O}_S}(G)$ with $G = i_{x_o}^{-1}L$. As a consequence, there exists an \mathcal{O}_S -module G such that, for any 1-connected open subset U of $X \times S$, there exists an isomorphism $L|_{U \times S} \simeq p_U^{-1}G$. Two choices of G are isomorphic, but non-canonically in general. Furthermore, L is $p^{-1}\mathcal{O}_S$ -coherent if and only if G is \mathcal{O}_S -coherent.

To any S -locally constant sheaf L one can associate an exact sequence of sheaves of $p^{-1}\mathcal{O}_S$ -modules

$$(4.1) \quad 0 \longrightarrow L_t \longrightarrow L \longrightarrow L_{\mathrm{tf}} \longrightarrow 0$$

where L_t denotes the subsheaf of $p^{-1}\mathcal{O}_S$ -torsion and L_{tf} the maximal $p^{-1}\mathcal{O}_S$ -torsion-free quotient. Then L_t and L_{tf} are S -locally constant and the previous exact sequence gives rise (for any choice of G) to the exact sequence of \mathcal{O}_S -modules

$$0 \longrightarrow G_t \longrightarrow G \longrightarrow G_{\mathrm{tf}} \longrightarrow 0,$$

with G_t and G_{tf} defined similarly. The following is straightforward.

Lemma 4.2. *Let $\pi : S \rightarrow S'$ be a holomorphic map between complex manifolds.*

(1) *If L is an S -locally constant sheaf on $X \times S$ with associated \mathcal{O}_S -module G , then $\pi_! L$ is S' -locally constant on $X \times S'$ and $\pi_! G$ is an associated $\mathcal{O}_{S'}$ -module to $\pi_! L$.*

(2) *If L' is an S' -locally constant sheaf on $X \times S'$ with associated \mathcal{O}'_S -module G' , then $\pi^* L'$ is S -locally constant on $X \times S$ and $\pi^* G'$ is an associated \mathcal{O}_S -module to $\pi^* L'$.*

Let us recall the Riemann-Hilbert correspondence for coherent S -local systems proved in [6, Th.2.23 p.14], in the particular case of a projection $X \times S \rightarrow S$, where X and S are complex manifolds.

The functor $L \mapsto (E_L, \nabla) = (\mathcal{O}_{X \times S} \otimes_{p^{-1}\mathcal{O}_S} L, d_{X \times S/S} \otimes \text{Id})$ induces an equivalence between the category of coherent S -locally constant sheaves of $p^{-1}\mathcal{O}_S$ -modules and the category of coherent $\mathcal{O}_{X \times S}$ -modules E equipped with an integrable relative connection $\nabla : E \rightarrow \Omega^1_{X \times S/S} \otimes_{\mathcal{O}_{X \times S}} E$. A quasi-inverse is given by $(E, \nabla) \mapsto E^\nabla = \ker \nabla$. The monodromy representation attached to the local system E^∇ is also called the monodromy representation of ∇ on E . Let us emphasize a direct consequence:

Corollary 4.3. *Let (E, ∇) be a coherent $\mathcal{O}_{X \times S}$ -module equipped with an integrable relative connection. Then the natural $\mathcal{O}_{X \times S}$ -linear morphism*

$$\mathcal{O}_{X \times S} \otimes_{p^{-1}\mathcal{O}_S} E^\nabla \longrightarrow E$$

is an isomorphism compatible with the integrable connections $d_{X \times S/S} \otimes \text{Id}$ and ∇ .

Proposition 4.4. *Notation as in Lemma 4.2.*

(1) *Let L be a coherent S -locally constant sheaf. If π is proper, then $\pi_* L$ is S' -coherent and there exists a natural morphism $E_{\pi_* L} \rightarrow \pi_* E_L$.*

(2) *Let L' be a coherent S' -locally constant sheaf. Then $E_{\pi^* L'} \simeq \pi^*(E_{L'})$.*

Proof.

(1) Since π is proper, $\pi_* L$ is a coherent S' -locally constant sheaf on $X \times S'$ (Lemma 4.2(1) and Grauert's theorem). The natural morphism $\mathcal{O}_{X \times S'} \rightarrow \pi_* \mathcal{O}_{X \times S}$ induces a composed morphism

$$\mathcal{O}_{X \times S'} \otimes_{p'^{-1}\mathcal{O}_{S'}} \pi_* L \longrightarrow \pi_* \mathcal{O}_{X \times S} \otimes_{p^{-1}\mathcal{O}_S} \pi_* L \longrightarrow \pi_*(\mathcal{O}_{X \times S} \otimes_{p^{-1}\mathcal{O}_S} L),$$

which is clearly compatible with the relative differential $d \otimes \text{Id}$. We note that if π is surjective with connected fibers, the first morphism is an isomorphism since $\mathcal{O}_{X \times S'} \rightarrow \pi_* \mathcal{O}_{X \times S}$ is then an isomorphism. This is the case if for example S' is a complex manifold and π is a proper modification of S' .

(2) The second point is straightforward. q.e.d.

4.b. The Deligne extension of an S -locally constant sheaf. We recall Theorem 2.6 and extend Corollary 2.8 of [31] to the case where $\dim S > 1$.

Notation 4.5. Let Y be a hypersurface in X (assumed to be connected) and let $j : X^* := X \setminus Y \hookrightarrow X$ denote the inclusion. Let L be a coherent S -locally constant sheaf on $X^* \times S$. Let $(E_L, \nabla) = (\mathcal{O}_{X^* \times S} \otimes_{p^{-1}\mathcal{O}_S} L, d_{X^* \times S/S} \otimes \text{Id})$ be the associated coherent $\mathcal{O}_{X^* \times S}$ -module with flat relative connection $\nabla = d_{X^* \times S/S} \otimes \text{Id}$, so that $L = E_L^\nabla$. We simply set $E = E_L$ when the context is clear, that we consider as a left $\mathcal{D}_{X^* \times S/S}$ -module.

Lemma 4.6. *The functor $j_* E_\bullet : L \mapsto j_* E_L$ with values in $\text{Mod}(\mathcal{D}_{X \times S/S})$ is exact.*

Proof. Any $(x_o, s_o) \in Y \times S$ has a fundamental system of open neighborhood $U \times U(s_o)$ such that $(U \setminus D) \times U(s_o)$ is Stein. Since E_L is $\mathcal{O}_{X^* \times S}$ -coherent, the result follows. q.e.d.

We assume from now on that $Y = D$ is a divisor with normal crossings in X . Let $\varpi : \tilde{X} \rightarrow X$ denote the real blowing up of X along the components of D . We denote by $\tilde{j} : X^* \hookrightarrow \tilde{X}$ the inclusion, so that $j = \varpi \circ \tilde{j}$. Let $x_o \in D$, $\tilde{x}_o \in \varpi^{-1}(x_o)$ and let $s_o \in S$. Choose local coordinates (x_1, \dots, x_n) at x_o such that $D = \{x_1 \cdots x_\ell = 0\}$ and consider the associated polar coordinates $(\boldsymbol{\rho}, \boldsymbol{\theta}, \mathbf{x}') := (\rho_1, \theta_1, \dots, \rho_\ell, \theta_\ell, x_{\ell+1}, \dots, x_n)$ so that \tilde{x}_o has coordinates $\boldsymbol{\rho}_o = 0$, $\boldsymbol{\theta}_o$, $\mathbf{x}'_o = 0$. We also denote by ϖ the induced map $\tilde{X} \times S \rightarrow X \times S$ and by $\tilde{p} : \tilde{X} \times S \rightarrow S$ the projection.

Definition 4.7. The subsheaf $\mathcal{A}_{\tilde{X} \times S}^{\text{mod}}$ of $\tilde{j}_* \mathcal{O}_{X^* \times S}$ of holomorphic functions having *moderate growth* along $\varpi^{-1}(D)$ is defined by the following two requirements:

- $\tilde{j}^{-1} \mathcal{A}_{\tilde{X} \times S}^{\text{mod}} = \mathcal{O}_{X^* \times S}$.
- For any $x_o \in D$, $\tilde{x}_o \in \varpi^{-1}(x_o)$ and $s_o \in S$, a germ $\tilde{h} \in (\tilde{j}_* \mathcal{O}_{X^* \times S})_{(\tilde{x}_o, s_o)}$ is said to have *moderate growth*, i.e., to belong to $\mathcal{A}_{\tilde{X} \times S, (\tilde{x}_o, s_o)}^{\text{mod}}$, if there exist open sets

$$(4.8) \quad \tilde{U}_\varepsilon := \{\|\boldsymbol{\rho}\| < \varepsilon, \|\mathbf{x}'\| < \varepsilon, \|\boldsymbol{\theta} - \boldsymbol{\theta}_o\| < \varepsilon\} \quad \text{and} \quad U(s_o) \ni s_o$$

(ε small enough) such that, setting $U_\varepsilon^* := \tilde{U}_\varepsilon \setminus \{\rho_1 \cdots \rho_\ell = 0\}$, \tilde{h} is defined on $U_\varepsilon^* \times U(s_o)$ and $|\tilde{h}|$ is bounded on this open set by $C\|\boldsymbol{\rho}\|^{-N}$, for some $C, N > 0$.

We recall the following properties of the sheaf $\mathcal{A}_{\tilde{X} \times S}^{\text{mod}}$:

- (1) $\mathcal{A}_{\tilde{X} \times S}^{\text{mod}}$ is a $\varpi^{-1} \mathcal{D}_{X \times S/S}(*D)$ -module (cf. [37, §1.1.6]) which is $\varpi^{-1} \mathcal{O}_{X \times S}$ -flat (cf. [25, Th. 4.6.1]),
- (2) for any coherent $\mathcal{O}_{X \times S}$ -module M , the natural morphism

$$\mathcal{A}_{\tilde{X} \times S}^{\text{mod}} \otimes_{\varpi^{-1} \mathcal{O}_{X \times S}} \varpi^{-1} M \longrightarrow \tilde{j}_* j^{-1} M$$

is injective (cf. loc. cit.),

- (3) $R\varpi_* \mathcal{A}_{\tilde{X} \times S}^{\text{mod}} = \varpi_* \mathcal{A}_{\tilde{X} \times S}^{\text{mod}} = \mathcal{O}_{X \times S}(*D)$ (cf. [37, Cor. II.1.1.18]).

Let us already notice for later use that these properties imply, for any \mathcal{O}_S -module G ,

$$(4.9) \quad \varpi_* (\mathcal{A}_{\tilde{X} \times S}^{\text{mod}} \otimes_{\tilde{p}^{-1} \mathcal{O}_S} \tilde{p}^{-1} G) \simeq \mathcal{O}_{X \times S}(*D) \otimes_{p^{-1} \mathcal{O}_S} p^{-1} G.$$

Indeed, we have

$$\begin{aligned} R\varpi_* (\mathcal{A}_{\tilde{X} \times S}^{\text{mod}} \otimes_{\tilde{p}^{-1} \mathcal{O}_S} \tilde{p}^{-1} G) &\simeq R\varpi_* (\mathcal{A}_{\tilde{X} \times S}^{\text{mod}} \otimes_{\tilde{p}^{-1} \mathcal{O}_S}^L \tilde{p}^{-1} G) \quad \text{by (1)} \\ &\simeq R\varpi_* (\mathcal{A}_{\tilde{X} \times S}^{\text{mod}}) \otimes_{p^{-1} \mathcal{O}_S}^L p^{-1} G \\ &\simeq \mathcal{O}_{X \times S}(*D) \otimes_{p^{-1} \mathcal{O}_S}^L p^{-1} G \quad \text{by (3)} \\ &\simeq \mathcal{O}_{X \times S}(*D) \otimes_{p^{-1} \mathcal{O}_S} p^{-1} G, \end{aligned}$$

where the latter isomorphism follows from flatness of $\mathcal{O}_{X \times S}(*D)$ over $p^{-1} \mathcal{O}_S$. Furthermore, (2) implies that, for any coherent \mathcal{O}_S -module G , the natural morphism

$$\mathcal{A}_{\tilde{X} \times S}^{\text{mod}} \otimes_{\tilde{p}^{-1} \mathcal{O}_S} \tilde{p}^{-1} G \longrightarrow \tilde{j}_* (\mathcal{O}_{X^* \times S} \otimes_{p^{-1} \mathcal{O}_S} p^{-1} G)$$

is injective.

Note that, U_ε^* being contractible, we have $L|_{U_\varepsilon^* \times U(s_o)} \simeq p_{U_\varepsilon^*}^{-1} G|_{U(s_o)}$ (cf. [31, Prop. A.12]). We thus have an identification

$$(4.10) \quad (\tilde{j}_* E)|_{\tilde{U}_\varepsilon \times U(s_o)} \simeq \tilde{j}_* \left(\mathcal{O}_{U_\varepsilon^* \times U(s_o)} \otimes_{p^{-1}\mathcal{O}_{U(s_o)}} p_{U_\varepsilon^*}^{-1} G|_{U(s_o)}, d_{U_\varepsilon^* \times U(s_o)/U(s_o)} \otimes \text{Id} \right) \\ \xleftarrow{\sim} \left((\tilde{j}_* \mathcal{O}_{U_\varepsilon^* \times U(s_o)}) \otimes_{\tilde{p}^{-1}\mathcal{O}_{U(s_o)}} \tilde{p}_{U_\varepsilon}^{-1} G|_{U(s_o)}, d_{\tilde{U}_\varepsilon \times U(s_o)/U(s_o)} \otimes \text{Id} \right).$$

Indeed, for a polysector $\tilde{V}_\varepsilon \subset \tilde{U}_\varepsilon$ and $V \subset U(s_o)$, the natural morphism

$$\Gamma(V_\varepsilon^* \times V, \mathcal{O}_{X \times S}) \otimes_{\mathcal{O}(V)} \Gamma(\tilde{V}_\varepsilon \times V, \tilde{p}^{-1} G) \longrightarrow \Gamma(V_\varepsilon^* \times V, \mathcal{O}_{X \times S}) \otimes_{\mathcal{O}(V)} \Gamma(V_\varepsilon^* \times V, p^{-1} G)$$

is an isomorphism, since the adjunction morphism $p_{W^*} p_W^{-1} G \rightarrow G$ is an isomorphism for both $W = V_\varepsilon^*$ and $W = \tilde{V}_\varepsilon$ (both being connected, cf. e.g. [31, Prop. A.1(2)]). The assertion is obtained by passing from the pre-sheaf isomorphism to a sheaf isomorphism.

Definition 4.11 (The Deligne extension). Let $x_o \in X$, $\tilde{x}_o \in \varpi^{-1}(x_o)$ and $s_o \in S$.

(1) A germ \tilde{v} of $(\tilde{j}_* E_L)_{(\tilde{x}_o, s_o)}$ is said to have *moderate growth* if for some open set $\tilde{U}_\varepsilon \times U(s_o)$ as above, and for some (equivalently, any) identification $L|_{U_\varepsilon^* \times U(s_o)} \simeq p_{U_\varepsilon^*}^{-1} G|_{U(s_o)}$, \tilde{v} belongs to the image of the *injective* morphism

$$\Gamma(\tilde{U}_\varepsilon \times U(s_o); \mathcal{A}_{X \times S}^{\text{mod}} \otimes_{\tilde{p}^{-1}\mathcal{O}_S} \tilde{p}^{-1} G) \longrightarrow \Gamma(\tilde{U}_\varepsilon \times U(s_o); \tilde{j}_* E_L).$$

(2) A germ v of $(j_* E_L)_{(x_o, s_o)}$ is said to have *moderate growth* if for each \tilde{x}_o in $\varpi^{-1}(x_o)$, the corresponding germ in $(\tilde{j}_* E_L)_{(\tilde{x}_o, s_o)}$ has moderate growth. In particular, this holds for any local section of E_L at (x_o, s_o) if $x_o \notin D$.

(3) The subsheaf of $j_* E_L$ consisting of local sections whose germs have moderate growth is denoted by \tilde{E}_L . It satisfies $j^* \tilde{E}_L = E_L$. It is called *the Deligne extension* of E_L .

Remark 4.12. By definition, with the previous notations, v has moderate growth if and only if, on any such polysector $U_\varepsilon \times U(s_o)$, for any isomorphism $L|_{U_\varepsilon^* \times U(s_o)} \simeq p_{U_\varepsilon^*}^{-1} G|_{U(s_o)}$, for any family of local generators (g_j) of G on $U(s_o)$, v can be written as $\sum_j v_j \otimes g_j$ with v_j being holomorphic functions on the corresponding polysector in $X^* \times S$ with moderate growth with respect to D .

Since $\mathcal{A}_{X \times S}^{\text{mod}}$ is stable by derivations with respect to X , \tilde{E}_L is a $\mathcal{D}_{X \times S/S}(*D)$ -submodule of $j_* E_L$.

Proposition 4.13 (First properties of the Deligne extension \tilde{E}_L)

Let L be a coherent S -locally constant sheaf on $X^* \times S$ and let $(E_L, \nabla) = (\mathcal{O}_{X^* \times S} \otimes_{p^{-1}\mathcal{O}_S} L, d_{X^* \times S/S} \otimes \text{Id})$ be the associated $\mathcal{O}_{X^* \times S}$ -module with flat relative connection.

(1) The assignment $L \mapsto (\tilde{E}_L, \nabla)$ with values in $\text{Mod}(\mathcal{D}_{X \times S/S})$ is functorial.

(2) Let $\pi : S' \rightarrow S$ be a proper holomorphic map. Then the natural morphism $j_*(E_{\pi^* L}, \nabla) \rightarrow \pi_* j_*(E_L, \nabla)$ sends the subsheaf $\tilde{E}_{\pi^* L}$ to $\pi_* \tilde{E}_L$.

(3) Let $\pi : S' \rightarrow S$ be a holomorphic map. Then the natural morphism $\pi^* j_*(E_L, \nabla) \rightarrow j_*(E_{\pi^* L}, \nabla)$ sends isomorphically $\pi^*(\tilde{E}_L, \nabla)$ to $(\tilde{E}_{\pi^* L}, \nabla)$.

Proof. We only prove (1) and (2), and (3) will be a consequence of Theorem 4.15 below.

(1) A morphism of S -locally constant sheaves $\varphi : L \rightarrow L'$ defines a morphism $\varphi : E_L \rightarrow E_{L'}$ compatible with ∇ , hence $j_*E_L \rightarrow j_*E_{L'}$ compatible with ∇ , and we only need to check that it sends \tilde{E}_L to $\tilde{E}_{L'}$. This is straightforward from the definition.

(2) From the commutative diagram

$$(4.14) \quad \begin{array}{ccc} X^* \times S & \xrightarrow{j} & X \times S \\ \pi \downarrow & & \downarrow \pi \\ X^* \times S' & \xrightarrow{j} & X \times S' \end{array}$$

together with the natural morphism of Proposition 4.2(1), we obtain the morphism $j_*(E_{\pi_*L}, \nabla) \rightarrow \pi_*j_*(E_L, \nabla)$ and similarly $\tilde{j}_*(E_{\pi_*L}, \nabla) \rightarrow \pi_*\tilde{j}_*(E_L, \nabla)$. Let us consider an open subset $\tilde{U}_\varepsilon \subset \tilde{X}$ as in (4.8). Then U_ε^* is contractible and for any $s'_o \in S'$, $L|_{U_\varepsilon^* \times \pi^{-1}(U(s'_o))} \simeq p_{U_\varepsilon^*}^{-1}G|_{\pi^{-1}(U(s'_o))}$. In order to simplify the notation, we set $S' = U(s'_o)$ and $S = \pi^{-1}(U(s'_o))$. We also set $p = p_{\tilde{U}_\varepsilon}$, so that $pU_\varepsilon^* = p \circ \tilde{j}$. According to (4.10), we are led to proving that the natural morphism

$$(\tilde{j}_*\mathcal{O}_{U_\varepsilon^* \times S'}) \otimes_{\tilde{p}^{-1}\mathcal{O}_S} \tilde{p}'^{-1}\pi_*G \longrightarrow \pi_*((\tilde{j}_*\mathcal{O}_{U_\varepsilon^* \times S}) \otimes_{p^{-1}\mathcal{O}_S} p^{-1}G)$$

sends $\mathcal{A}_{\tilde{U}_\varepsilon \times S'}^{\text{mod}} \otimes_{\tilde{p}'^{-1}\mathcal{O}_S} \tilde{p}'^{-1}\pi_*G$ to $\pi_*(\mathcal{A}_{\tilde{U}_\varepsilon \times S}^{\text{mod}} \otimes_{p^{-1}\mathcal{O}_S} p^{-1}G)$. Following the proof of Proposition 4.4(1), we are led to proving that the natural morphism $\tilde{j}_*\mathcal{O}_{U_\varepsilon^* \times S'} \rightarrow \pi_*(\tilde{j}_*\mathcal{O}_{U_\varepsilon^* \times S})$ sends $\mathcal{A}_{\tilde{U}_\varepsilon \times S'}^{\text{mod}}$ to $\pi_*\mathcal{A}_{\tilde{U}_\varepsilon \times S}^{\text{mod}}$. This follows from the definition of moderate growth, owing to the properness of π . q.e.d.

4.c. Regular holonomicity of the Deligne extension of an S -locally constant sheaf. We continue to refer to Notation 4.5 and assume that $Y = D$ has normal crossings.

Theorem 4.15. *Assume that L is a coherent S -locally constant sheaf on $X^* \times S$. Then*

- (1) *the subsheaf \tilde{E}_L of j_*E_L is $\mathcal{O}_{X \times S}(*D)$ -coherent;*
- (2) *the functor $L \mapsto \tilde{E}_L$ from the category of coherent S -locally constant sheaves on $X^* \times S$ to that of $\mathcal{D}_{X \times S/S}$ -modules is fully faithful;*
- (3) *as a $\mathcal{D}_{X \times S/S}$ -module, (\tilde{E}_L, ∇) is regular holonomic.*

We will make use of the following flattening result for a coherent \mathcal{O}_S -module (here, we only need a local version, but the corresponding global one also holds true).

Proposition 4.16. *Near each $s_o \in S$, there exists a projective modification $\pi : S' \rightarrow S$ with S' smooth such that the torsion-free quotient of π^*G is $\mathcal{O}_{S'}$ -locally free.*

Proof. We first apply the flattening theorem [34, Th.3.5] to the coherent sheaf G . There exists thus a projective modification $\pi'' : S'' \rightarrow S$ such that π''^*G , when quotiented by its $\mathcal{O}_{S''}$ -torsion, is $\mathcal{O}_{S''}$ -locally free. We then apply resolution of singularities $\pi' : S' \rightarrow S''$ of S'' in the neighbourhood of the projective subset $\pi''^{-1}(s_o)$ (cf. [11, §7, Main Th. I']) and denote by π the morphism $\pi'' \circ \pi'$, which answers the question. q.e.d.

Proof of Theorem 4.15(1). We recall the proof of [31, Th.2.6] for the $\mathcal{O}_{X \times S}(*D)$ -coherence. The problem is local on $X \times S$. We thus assume that $X \times S$ is a small neighbourhood of (x_o, s_o) as above. In such a neighbourhood, giving the local system

is equivalent to giving $T_1, \dots, T_\ell \in \text{Aut}(G)$ which pairwise commute. Let $U(s_o)$ be an open neighbourhood of s_o isomorphic to an open polydisc. The formula [40, (2.11)] defining a logarithm of T_i can be used to show that there exist $A_1(s), \dots, A_\ell(s) \in \text{End}_{\mathcal{O}_S}(G|_{U(s_o)})$ which pairwise commute, such that $T_i = \exp(-2\pi i A_i(s))$ for each i on $U(s_o)$ (cf. [40, Cor. 2.3.10 & Chap. 3]). Set $\widehat{E}_G := \mathcal{O}_{X \times S}(*D) \otimes_{p^{-1}\mathcal{O}_S} p^{-1}G$, equipped with the connection $\widehat{\nabla}$ such that $\widehat{\nabla}_{x_i \partial_{x_i}}$ acts on $1 \otimes_{p^{-1}\mathcal{O}_S} p^{-1}G$ by $\text{Id} \otimes A_i(s)$ if $i = 1, \dots, \ell$, and zero otherwise. Then the monodromy representation of $\widehat{\nabla}$ on $\widehat{E}_G|_{X^* \times S}$ is given by T_1, \dots, T_ℓ , hence an isomorphism $L \simeq (\widehat{E}_G|_{X^* \times S})^{\widehat{\nabla}}$, from which one deduces, according to Corollary 4.3, an isomorphism

$$(4.17) \quad \begin{aligned} (E_L, \nabla) &= (\mathcal{O}_{X^* \times S} \otimes_{p^{-1}\mathcal{O}_S} L, d_{X^* \times S/S} \otimes \text{Id}) \\ &\simeq (\mathcal{O}_{X^* \times S} \otimes_{p^{-1}\mathcal{O}_S} (\widehat{E}_G|_{X^* \times S})^{\widehat{\nabla}}, d_{X^* \times S/S} \otimes \text{Id}) \simeq (\widehat{E}_G|_{X^* \times S}, \widehat{\nabla}). \end{aligned}$$

It follows that $\widetilde{E}_L \simeq \widetilde{\widehat{E}_G|_{X^* \times S}}$ and we are thus reduced to proving that $\widetilde{\widehat{E}_G|_{X^* \times S}} = \widetilde{E}_G$ (we have trivialized the locally constant sheaf but the connection is not trivial anymore).

Remark 4.18. Let $p_{X^*}^{-1}G$ be the constant S -local system on $X^* \times S$. Thus, locally on $X \times S$, there exist $\mathcal{O}_{X^* \times S}$ -linear isomorphisms $E_L \simeq \widehat{E}_G|_{X^* \times S} \simeq E_{p_{X^*}^{-1}G}$, but in general we do not have $(E_L, \nabla) \simeq (E_{p_{X^*}^{-1}G}, \widehat{\nabla})$, i.e., this isomorphism is not $\mathcal{D}_{X^* \times S/S}$ -linear.

Let us prove the inclusion $\widetilde{\widehat{E}_G|_{X^* \times S}} \subset \widetilde{E}_G$. Let $v \in (j_* \widehat{E}_G)_{(x_o, s_o)}$ be locally defined on $U \times U(s_o)$. On any polysector $U_\varepsilon^* \times U(s_o)$, we can choose logarithms of x_i ($i = 1, \dots, \ell$) and the automorphism $\prod_{i=1}^\ell x_i^{A_i(s)}$ of $\widehat{E}_G|_{X^* \times S} = \mathcal{O}_{U_\varepsilon^* \times U(s_o)} \otimes_{p^{-1}\mathcal{O}_{U(s_o)}} p^{-1}G$ is well-defined by setting $x_i^{A_i(s)} = \sum_k [(\log x_i)^k / k!] \otimes A_i(s)^k$. By definition, v has moderate growth, i.e., is a local section of $\widetilde{\widehat{E}_G|_{X^* \times S}}$, if and only if, on any such polysector $\widetilde{U}_\varepsilon \times U(s_o)$, the section $w := (\prod_{i=1}^\ell x_i^{-A_i(s)}) \cdot v$ is a section of $\mathcal{A}_{\widetilde{X} \times S}^{\text{mod}} \otimes_{\widetilde{p}^{-1}\mathcal{O}_S} \widetilde{p}^{-1}G$.

Let (g_j) be a family of local generators of G on $U(s_o)$. The entries of the matrices of $\prod_{i=1}^\ell x_i^{A_i(s)}$ and of $\prod_{i=1}^\ell x_i^{-A_i(s)}$ with respect to this family have moderate growth. If, on such a polysector, w writes $\sum_j w_j \otimes g_j$, where w_j are sections of $\mathcal{A}_{\widetilde{X} \times S}^{\text{mod}}$, then $v = \prod_{i=1}^\ell x_i^{A_i(s)} \cdot w$ writes similarly $\sum_j v_j \otimes g_j$, where v_j are sections of $\mathcal{A}_{\widetilde{X} \times S}^{\text{mod}}$. Conversely, if v has the previous expression, then so does $w = \prod_{i=1}^\ell x_i^{-A_i(s)} \cdot v$. We conclude that, for any such polysector, the moderate growth condition on v is equivalent to $v \in \Gamma(\widetilde{U}_\varepsilon \times U(s_o); \mathcal{A}_{\widetilde{X} \times S}^{\text{mod}} \otimes_{\widetilde{p}^{-1}\mathcal{O}_S} \widetilde{p}^{-1}G)$. In other words, if v has moderate growth we have

$$\begin{aligned} v \in \Gamma(U \times U(s_o); \varpi_* \mathcal{A}_{\widetilde{X} \times S}^{\text{mod}} \otimes_{\widetilde{p}^{-1}\mathcal{O}_S} \widetilde{p}^{-1}G) &= \Gamma(U \times U(s_o); \mathcal{O}_{X \times S}(*D) \otimes_{p^{-1}\mathcal{O}_S} p^{-1}G) \\ &= \Gamma(U \times U(s_o); \widehat{E}_G), \end{aligned}$$

hence $\widetilde{\widehat{E}_G|_{X^* \times S}} \subset \widetilde{E}_G$. The reverse inclusion also follows from the moderate growth of the entries of the matrices of $\prod_{i=1}^\ell x_i^{A_i(s)}$ and of $\prod_{i=1}^\ell x_i^{-A_i(s)}$. q.e.d.

Proof of Theorem 4.15(2). We already know, by the Riemann-Hilbert correspondence of [6, Th. 2.23], that the functor $L \mapsto E_L$ is fully faithful, and it is clearly exact, so it remains to prove that $E_L \mapsto \widetilde{E}_L$ is so. Faithfulness is clear. Let us check fullness. Let

$\varphi : \widetilde{E}_L \rightarrow \widetilde{E}_{L'}$ be a morphism and set $\widetilde{\varphi} = \widetilde{j^* \varphi} : \widetilde{E}_L \rightarrow \widetilde{E}_{L'}$. The kernel and cokernel of $\varphi - \widetilde{\varphi}$ are coherent $\mathcal{O}_{X \times S}(*D)$ -modules by 4.15(1), and are zero when restricted to $X^* \times S$. Therefore they are zero, and $\varphi = \widetilde{\varphi}$, proving the assertion. q.e.d.

Proof of Theorem 4.15(3). The question is local at a point $(x_o, s_o) \in X \times S$. By the proof of 4.15(1), we can work with $(\widehat{E}_G, \widehat{\nabla}) = (\mathcal{O}_{X \times S}(*D) \otimes_{p^{-1}\mathcal{O}_S} p^{-1}G, \widehat{\nabla})$ (with $\widehat{\nabla} = d \otimes \text{Id} + \sum_i dx_i/x_i \otimes A_i$ and $A_i \in \text{End}_{\mathcal{O}_S}(G)$). Our method is to reduce to the case where G is locally free of rank one and then prove (3) for it.

We argue by induction on the lexicographically ordered pair $(\dim S, \text{rk } G)$, where $\text{rk } G$ is the rank of G at a general point of S . The case $\dim S = 0$ and $\text{rk } G$ arbitrary is well-known. We thus assume that $d = \dim S \geq 1$ and $r = \text{rk } G \geq 0$, and that (3) holds for $(\widehat{E}_G, \widehat{\nabla})$ for any $p'^{-1}\mathcal{O}_{S'}$ -module G' with $(\dim S', \text{rk } G') < (d, r)$.

Step 1: The case $(d, 0)$. If $\text{rk } G = 0$, G is an \mathcal{O}_S -torsion module. Since the question is local, we can assume that the support of G is contained in a hypersurface $T \subset S$ locally presented as in the local setting 3.11. Then G is $q^{-1}\mathcal{O}_{S'}$ -coherent and is endowed with an endomorphism t (i.e., multiplication by t), so that the natural \mathcal{O}_S -linear morphism $\mathcal{O}_S \otimes_{q^{-1}\mathcal{O}_{S'}[t]} G \rightarrow G$ is an isomorphism. Furthermore, the endomorphisms A_i are $q^{-1}\mathcal{O}_{S'}$ -linear (via the natural inclusion $q^{-1}\mathcal{O}_{S'} \hookrightarrow \mathcal{O}_S$). Let us set $(\widehat{E}'_G, \widehat{\nabla}) = (\widehat{E}_G, \widehat{\nabla})$ when regarded as a $q^{-1}\mathcal{D}_{X \times S'/S'}$ -module. By the induction hypothesis, it is regular holonomic. By Corollary 3.13 we conclude that $(\widehat{E}_G, \widehat{\nabla})$ is regular holonomic, so that (3) holds for $(\widehat{E}_G, \widehat{\nabla})$.

Step 2: Reduction after proper surjective generically finite base change

Let $\pi : S' \rightarrow S$ be a proper surjective generically finite morphism of complex manifolds and let $p' : X \times S' \rightarrow S'$ denote the projection. Let us assume that (3) holds for $(\mathcal{O}_{X \times S'}(*D) \otimes_{p'^{-1}\mathcal{O}_{S'}} p'^{-1}\pi^*G, \pi^*\widehat{\nabla})$. The purpose of Step 2 is to show that, under such an assumption, (3) holds for $(\widehat{E}_G, \widehat{\nabla})$.

From now on, we assume $\dim S = d$ and $\text{rk } G \geq 1$. Let G' be the \mathcal{O}_S -torsion subsheaf of G . We note that the endomorphisms A_i of G preserve G' , so the exact sequence $0 \rightarrow G' \rightarrow G \rightarrow G'' \rightarrow 0$ with G'' being torsion-free gives rise to an exact sequence $0 \rightarrow (\widehat{E}_{G'}, \widehat{\nabla}) \rightarrow (\widehat{E}_G, \widehat{\nabla}) \rightarrow (\widehat{E}_{G''}, \widehat{\nabla}) \rightarrow 0$. Since the (3) holds for $(\widehat{E}_{G'}, \widehat{\nabla})$ by Step 1, it is enough to prove (3) for $(\widehat{E}_{G''}, \widehat{\nabla})$. In other words, we may assume that G is torsion-free.

By assumption, there exists a closed analytic subset T of codimension ≥ 1 in S such that, setting $T' = \pi^{-1}(T)$, the morphism $\pi : S' \setminus T' \rightarrow S \setminus T$ is finite étale. We have $\dim S' = \dim S$ and $\dim T' < \dim S$. Recall that we still denote by π the map $\text{Id} \times \pi : X \times S' \rightarrow X \times S$.

Remark that we have, for any holomorphic map $\pi : S' \rightarrow S$, and denoting for clarity by $\widetilde{\pi}$ the map $\text{Id} \times \pi : X \times S' \rightarrow X \times S$, the following list of canonical isomorphisms:

$$\begin{aligned}
& L\widetilde{\pi}^*(\mathcal{O}_{X \times S}(*D) \otimes_{p^{-1}\mathcal{O}_S} p^{-1}G) \\
&= \mathcal{O}_{X \times S'}(*D) \otimes_{\widetilde{\pi}^{-1}\mathcal{O}_{X \times S}(*D)} \widetilde{\pi}^{-1}(\mathcal{O}_{X \times S}(*D) \otimes_{p^{-1}\mathcal{O}_S} p^{-1}G) \\
(4.19) \quad & \stackrel{(*)}{=} \mathcal{O}_{X \times S'}(*D) \otimes_{\widetilde{\pi}^{-1}p^{-1}\mathcal{O}_S} \widetilde{\pi}^{-1}p^{-1}G \\
&= \mathcal{O}_{X \times S'}(*D) \otimes_{p'^{-1}\mathcal{O}_{S'}}^{L} (p'^{-1}\mathcal{O}_{S'} \otimes_{\widetilde{\pi}^{-1}p^{-1}\mathcal{O}_S} \widetilde{\pi}^{-1}p^{-1}G) \\
&= \mathcal{O}_{X \times S'}(*D) \otimes_{p'^{-1}\mathcal{O}_{S'}}^{L} p'^{-1}L\pi^*G = \mathcal{O}_{X \times S'}(*D) \otimes_{p'^{-1}\mathcal{O}_{S'}} p'^{-1}L\pi^*G,
\end{aligned}$$

which reads

$$(4.20) \quad L\pi^*\widehat{E}_G \simeq \mathcal{O}_{X \times S'}(*D) \otimes_{p'^{-1}\mathcal{O}_{S'}} p'^{-1}L\pi^*G.$$

We conclude that, for each j , we have $L^j\pi^*\widehat{\nabla}_{x_k\partial_{x_k}} = d \otimes \text{Id} + \text{Id} \otimes L^j\pi^*A_k$.

Furthermore, under the assumption on π for this step, according to the projection formula for $R\tilde{\pi}_*$ applied to $(*)$, we have

$$R\pi_*(L\pi^*\widehat{E}_G) \simeq R\pi_*\mathcal{O}_{X \times S'}(*D) \otimes_{p^{-1}\mathcal{O}_S}^L p^{-1}G.$$

By induction, if $j \neq 0$, (3) holds for $(\mathcal{O}_{X \times S'}(*D) \otimes_{p'^{-1}\mathcal{O}_{S'}} p'^{-1}L^j\pi^*G, L^j\pi^*\widehat{\nabla})$, according to the argument given in Step 1, since $L^j\pi^*G$ is supported on T' , hence is a torsion module. The assumption thus yields that (3) holds for $L\pi^*\widehat{E}_G$ (i.e., for each of its cohomology modules) since for $j = 0$ it follows from Proposition 4.13(3). Since \widehat{E}_G is \mathcal{O} -quasi-coherent (cf. Remark 2.5(1)), one deduces from Lemma 2.12 that each $L^j\pi^*\widehat{E}_G$ is π -good. Then, by Theorem 3.14, (3) holds for $R\pi_*(L\pi^*\widehat{E}_G)$, hence for $\mathcal{H}^0(R\pi_*(L\pi^*\widehat{E}_G))$.

The natural morphism $\mathcal{O}_{X \times S}(*D) \rightarrow R\pi_*\mathcal{O}_{X \times S'}(*D)$ yields a morphism $\widehat{E}_G \rightarrow \mathcal{H}^0(R\pi_*(L\pi^*\widehat{E}_G))$. Both modules are $\mathcal{O}_{X \times S}(*D)$ -coherent by (1), hence so is the kernel of this morphism.

On $S \setminus T$, we claim that this morphism is injective: indeed, since $\pi : S' \setminus T' \rightarrow S \setminus T$ is finite étale, the trace morphism $\text{tr}_\pi : \pi_*\mathcal{O}_{S' \setminus T'} \rightarrow \mathcal{O}_{S \setminus T}$ defined by $\text{tr}_\pi(\varphi)(s) = (1/\deg \pi) \sum_{s' \in \pi^{-1}(s)} \varphi(s')$ satisfies $\text{tr}_\pi \circ \iota = \text{Id}$, if ι denotes the natural morphism $\mathcal{O}_{S \setminus T} \rightarrow \pi_*\mathcal{O}_{S' \setminus T'}$; hence $\widehat{E}_G|_{S \setminus T}$ is locally a direct summand of the right-hand side. As G , hence \widehat{E}_G , is assumed to be \mathcal{O}_S -torsion-free, the kernel is \mathcal{O}_S -torsion-free, $\mathcal{O}_{X \times S}(*D)$ -coherent and S -supported on T . It is thus zero and this morphism is injective.

Furthermore, \widehat{E}_G is $\mathcal{D}_{X \times S/S}$ -coherent: indeed, since (3) holds for $\mathcal{H}^0(R\pi_*(L\pi^*\widehat{E}_G))$ and \widehat{E}_G is \mathcal{O} -good, it follows that \widehat{E}_G is $\mathcal{D}_{X \times S/S}$ -coherent, and then regular holonomic by Proposition 3.1(iii), so that (3) holds for \widehat{E}_G .

Step 3: Reduction to the case where G is \mathcal{O}_S -locally free. We choose π as in Proposition 4.16 and by Step 2 we can assume from the start that the torsion-free quotient G'' of G is \mathcal{O}_S -locally free. By Step 1, it is enough to prove (3) for $(\widehat{E}_{G''}, \widehat{\nabla})$, i.e., we can assume (and we do assume from now on) that G is \mathcal{O}_S -locally free.

Step 4: The case where G is \mathcal{O}_S -locally free. We still work locally on S and we assume that G is \mathcal{O}_S -free of rank $r \geq 1$. By the induction hypothesis, (3) holds for any $(\widehat{E}_{G'}, \widehat{\nabla})$ with $\text{rk } G' < r$. Locally let us fix an \mathcal{O}_S -basis of G and let $A_1(s)$ denote the matrix of A_1 in this basis. Let $\Sigma \subset S \times \mathbb{C}$ be the zero locus of $\det(\alpha_1 \text{Id} - A_1)$. Since this function is a Weierstrass polynomial of degree $\deg \sigma$ with respect to the variable α_1 (considered as a coordinate on \mathbb{C}) with coefficients in \mathcal{O}_S , Σ is defined by the corresponding reduced Weierstrass polynomial. We note the following two properties related to σ and similarly to $\text{Id}_X \times \sigma$:

$$(4.21) \quad \sigma \text{ is finite and } \sigma_*\mathcal{O}_\Sigma \text{ is } \mathcal{O}_S\text{-free of degree } \deg \sigma \text{ (with basis } 1, \alpha_1, \dots, \alpha_1^{\deg \sigma - 1}\text{)}.$$

(4.22) There exist dense analytic open subsets $\Sigma^\circ \subset \Sigma$ and $S^\circ \subset S$ such that $\sigma : \Sigma^\circ \rightarrow S^\circ$ is a finite covering of degree $\deg \sigma$. The corresponding trace morphism $\text{tr}_\sigma : \sigma_*\mathcal{O}_{\Sigma^\circ} \rightarrow \mathcal{O}_{S^\circ}$ has been defined above. If φ is a section of $\sigma_*\mathcal{O}_\Sigma$, its trace on Σ° is bounded on S° , hence extends holomorphically to S , so that tr_σ extends as a

morphism $\sigma_*\mathcal{O}_\Sigma \rightarrow \mathcal{O}_S$. If ι denotes the natural morphism $\mathcal{O}_S \rightarrow \sigma_*\mathcal{O}_\Sigma$, we clearly have $\text{tr}_\sigma \circ \iota = \text{Id}$, making \mathcal{O}_S a direct summand of $\sigma_*\mathcal{O}_\Sigma$.

Let $\pi : S' \rightarrow \Sigma$ be a resolution of singularities of Σ , so that the natural composed map $(\sigma \circ \pi) : S' \rightarrow S$ is projective and generically finite. Let us set $G' = \ker(\alpha_1 \text{Id} - A_1 \circ \sigma \circ \pi) \subset (\sigma \circ \pi)^*G$, where we regard α_1 as a function $S' \rightarrow \mathbb{C}$. Noting that $\sigma \circ \pi$ is generically a local isomorphism, it follows by construction that $\text{rk } G' \geq 1$. We consider the exact sequence $0 \rightarrow G' \rightarrow (\sigma \circ \pi)^*G \rightarrow G'' \rightarrow 0$ which satisfies $0 < \text{rk } G' < \text{rk } G'' < \text{rk } G$, and which is preserved by the endomorphisms $(\sigma \circ \pi)^*A_i$, so that it induces an exact sequence of $\mathcal{D}_{X \times S/S}$ -modules

$$0 \longrightarrow (\widehat{E}_{G'}, \widehat{\nabla}) \longrightarrow (\sigma \circ \pi)^*(\widehat{E}_G, \widehat{\nabla}) \longrightarrow (\widehat{E}_{G''}, \widehat{\nabla}) \longrightarrow 0.$$

If $\text{rk } G' < \text{rk } G$, we can apply induction to $(\widehat{E}_{G'}, \widehat{\nabla})$, $(\widehat{E}_{G''}, \widehat{\nabla})$ and conclude that (3) holds for $(\sigma \circ \pi)^*(\widehat{E}_G, \widehat{\nabla})$, hence for $(\widehat{E}_G, \widehat{\nabla})$ according to Step 2.

If $\text{rk } G' = \text{rk } G$, then $\text{rk } G'' = 0$, so Step 1 applies to $(\widehat{E}_{G''}, \widehat{\nabla})$, and we are reduced to proving (3) for $(\widehat{E}_{G'}, \widehat{\nabla})$, i.e., we can assume that $A_1 = \alpha_1 \text{Id}$. Iterating the argument, we are reduced to the case where $A_i = \alpha_i \text{Id}$ for $i = 1, \dots, \ell$, where $\alpha_1, \dots, \alpha_\ell$ are holomorphic functions on S , and G is \mathcal{O}_S -locally free. By considering a local basis of G , it suffices to consider the case where $\text{rk } G = 1$.

Step 5: The case where G is \mathcal{O}_S -locally free of rank one. We now consider $(\widehat{E}_G, \widehat{\nabla}) = (\mathcal{O}_{X \times S}(*D), d_{X \times S/S} + \sum_{i=1}^{\ell} \alpha_i(s) dx_i/x_i)$. The argument for obtaining (3) is then similar to that used in the proof of [31, Cor.2.8]. One can assume that, in the neighbourhood of s_o and for any i , $\alpha_i(s) \in \mathbb{Z} \Rightarrow \alpha_i(s) = 0$. Then there exists a surjective morphism

$$\mathcal{D}_{X \times S/S} / ((x_i \partial_{x_i} - \alpha_i(s) + 1)_{i=1, \dots, \ell}, (\partial_{x_j})_{j=\ell+1, \dots, n}) \longrightarrow (\widehat{E}_G, \widehat{\nabla})$$

sending 1 to $1/x_1 \cdots x_\ell$, which is easily seen to be an isomorphism by the condition on $(\alpha_i)_{i=1, \dots, \ell}$. q.e.d.

Proof of Proposition 4.13(3). The statement is local, so, as a consequence of (4.17) in the proof of Theorem 4.15(1) and keeping the same notation, we are reduced to proving $\pi^*(\widehat{E}_G, \widehat{\nabla}) \simeq (\widehat{E}_{\pi^*G}, \widehat{\nabla})$, that is,

$$\pi^*(\mathcal{O}_{X \times S}(*D) \otimes_{p^{-1}\mathcal{O}_S} p^{-1}G, \widehat{\nabla}) \simeq (\mathcal{O}_{X \times S'}(*D) \otimes_{p'^{-1}\mathcal{O}_{S'}} p'^{-1}\pi^*G, \pi^*\widehat{\nabla}),$$

which follows by taking the 0-cohomology in (4.19). q.e.d.

4.d. $\mathcal{D}_{X \times S/S}$ -modules of D-type. Recall Notation 4.5. In this section, we exhibit a family of regular holonomic $\mathcal{D}_{X \times S/S}$ -modules, that we call *of D-type* and we prove in Proposition 4.26 a special case of the analogue in the relative setting of [14, Th. 2.3.2] asserting that the restriction functor to the complement of the divisor is an equivalence of categories. The general case will be obtained in Theorem 6.17 below.

Let Y be a closed hypersurface of X .

Definition 4.23. We say that a coherent $\mathcal{D}_{X \times S/S}(*Y)$ -module \mathcal{L} is of *D-type* if

(a) there exists a coherent S -locally constant sheaf L on $X^* \times S$ such that $\mathcal{L}|_{X^* \times S} \simeq E_L = (\mathcal{O}_{X^* \times S} \otimes_{p^{-1}\mathcal{O}_S} L, d_{X^* \times S/S} \otimes \text{Id})$, equivalently, $\mathcal{L}|_{X^* \times S}$ is $\mathcal{D}_{X^* \times S/S}$ -holonomic with characteristic variety contained in the zero section,

(b) for each $s \in S$, the cohomology of $Li_s^* \mathcal{L}$ is \mathcal{D}_X -regular holonomic (in particular, \mathcal{D}_X -coherent). In other words \mathcal{L} satisfies condition (Reg 2) and thus (Reg 1) (cf. Proposition 3.1).

We say that \mathcal{L} is *strict* if L is $p^{-1}\mathcal{O}_S$ -locally free.

We denote by $\text{Mod}_Y(\mathcal{D}_{X \times S/S}(*Y))$ the full subcategory of $\text{Mod}(\mathcal{D}_{X \times S/S}(*Y))$ whose objects are coherent $\mathcal{D}_{X \times S/S}(*Y)$ -modules of D-type.

Lemma 4.24. *For $\pi : S' \rightarrow S$ and $\mathcal{L} \in \text{Mod}_Y(\mathcal{D}_{X \times S/S}(*Y))$, $L\pi^* \mathcal{L}$ has cohomology in $\text{Mod}_Y(\mathcal{D}_{X \times S'/S'}(*Y))$.*

Proof. The good behaviour of 4.23(a) by base pullback is clear. Let us check that of 4.23(b). Arguing as in Lemma 2.13, we see that $L\pi^* \mathcal{L}$ is an object of $\text{D}_{\text{coh}}^b(\mathcal{D}_{X \times S'/S'}(*D))$. For any $s' \in S'$ we have $Li_{s'}^* L\pi^* \mathcal{L} \simeq Li_{\pi(s')}^* \mathcal{L}$, so the complex $Li_{s'}^* L\pi^* \mathcal{L}$ has \mathcal{D}_X -regular holonomic cohomology. By Proposition 3.1(i), each $Li_{s'}^* L^j \pi^* \mathcal{L}$ also has \mathcal{D}_X -regular holonomic cohomology. q.e.d.

Proposition 4.25. *Assume that $Y = D$ is a divisor with normal crossings in X .*

- (1) *If \mathcal{L} is $\mathcal{D}_{X \times S/S}(*D)$ -coherent of D-type and strict, then the natural morphism $\psi : \mathcal{L} \rightarrow j_* E_L$ sends \mathcal{L} isomorphically to \widetilde{E}_L .*
- (2) *If \mathcal{L} is $\mathcal{D}_{X \times S/S}(*D)$ -coherent of D-type, then \mathcal{L} is $\mathcal{D}_{X \times S/S}$ -regular holonomic.*

Proof. We note that the question is local on $X \times S$, so we may replace E_L with $\widehat{E}_{G|X^* \times S}$ as in (4.17). Namely, we have

$$\mathcal{L}|_{X^* \times S} \simeq (\mathcal{O}_{X^* \times S} \otimes_{p^{-1}\mathcal{O}_S} p^{-1}G, \widehat{\nabla}).$$

We note that:

- If \mathcal{L} is strict, the second assertion follows from the first one, because \widetilde{E}_L is $\mathcal{D}_{X \times S/S}$ -regular holonomic (Theorem 4.15(3)).
- For any coherent $\mathcal{D}_{X \times S/S}(*D)$ -module of D-type, the morphism ψ is injective. Indeed, according to the first point of Definition 4.23, the restriction of ψ to $X^* \times S$ is an isomorphism. The assertion follows from the \mathcal{O} -quasi-coherence of \mathcal{L} (cf. Remark 2.5(2)). We thus identify \mathcal{L} with a $\mathcal{D}_{X \times S/S}(*D)$ -submodule of $j_* E_L$.
- Due to 4.23(b), 4.25(2) amounts to holonomicity of \mathcal{L} (in particular, $\mathcal{D}_{X \times S/S}$ -coherence).

Proof of Proposition 4.25(1). We assume that G is \mathcal{O}_S -locally free of finite rank. We can mimic the end of the proof of [31, Prop. 2.11] to directly show that ψ is an isomorphism $\mathcal{L} \xrightarrow{\sim} \widetilde{E}_L$ because, although in loc. cit. we assumed the $\mathcal{D}_{X \times S/S}$ -coherence of \mathcal{L} , that proof works under the weaker assumption of its $\mathcal{D}_{X \times S/S}(*Y)$ -coherence. This shows 4.25(1). In particular this implies the $\mathcal{D}_{X \times S/S}$ -regular holonomicity of \mathcal{L} .

Proof of Proposition 4.25(2). We now prove the holonomicity of \mathcal{L} by assuming only that G is \mathcal{O}_S -coherent. As in the proof of Theorem 4.15(3), we argue by induction on the dimension of S . The case where S is a point is well-known ([14, Th. 2.3.2]). We assume that the result holds if $\dim S \leq d - 1$ ($d \geq 1$) and that $\dim S = d$.

Step 1: The case where G is a torsion \mathcal{O}_S -module. Since we work locally, we can assume that the support of G is contained in a hypersurface T of S , having equation $h = 0$, endowed with a finite morphism q to S' of dimension $d - 1$, and that $h^k G = 0$ for some $k \geq 1$. We claim that any local section m of \mathcal{L} is annihilated by h^k . Indeed, for any such m , $h^k m$ is zero on $X^* \times S$ and we can apply the result of Remark 2.5(2).

Therefore, $\text{Supp}_S \mathcal{L}$ is contained in T . By the induction hypothesis and the equivalence recalled in Setting 3.11, we deduce that \mathcal{L} is $q^{-1}\mathcal{D}_{X \times S'/S'}$ -holonomic, hence, arguing as in Remark 3.12, we conclude that \mathcal{L} is $\mathcal{D}_{X \times S/S}$ -holonomic, as desired.

Step 2: The case where G_{tf} is \mathcal{O}_S -locally free. We consider the exact sequence (4.1) and we assume that G_{tf} is \mathcal{O}_S -locally free. We also consider the similar exact sequence $0 \rightarrow \mathcal{L}_t \rightarrow \mathcal{L} \rightarrow \mathcal{L}_{\text{tf}} \rightarrow 0$. Let us prove that \mathcal{L}_{tf} is of D-type. First, \mathcal{L}_t is easily seen to be $\mathcal{D}_{X \times S/S}(*D)$ -coherent, hence so is \mathcal{L}_{tf} . Next, \mathcal{L}_{tf} satisfies the first point of Definition 4.23, with local system $p^{-1}G_{\text{tf}}$.

For the second point, we note that the image of $i_s^* \mathcal{L}_t \rightarrow i_s^* \mathcal{L}$ is \mathcal{D}_X -coherent since $i_s^* \mathcal{L}$ is so and $i_s^* \mathcal{L}_t$ is \mathcal{O} -quasi-coherent. As a consequence, it is \mathcal{D}_X -regular holonomic since $i_s^* \mathcal{L}$ is so, and thus $i_s^* \mathcal{L}_{\text{tf}}$ is regular holonomic. On the other hand, for $j < 0$, $L^j i_s^* \mathcal{L}_{\text{tf}}$ is \mathcal{O} -quasi-coherent and supported on D by our assumption on G_{tf} , so $L^j i_s^* \mathcal{L}_{\text{tf}} = 0$ by Lemma 2.3(2).

In conclusion, \mathcal{L}_{tf} is of D-type, and we also deduce that \mathcal{L}_t is of D-type. By Step 1 and 4.25(1), \mathcal{L} is holonomic if G_{tf} is \mathcal{O}_S -locally free.

Step 3: The general case. Let $\pi : S' \rightarrow S$ be a projective modification as in Proposition 4.16 such that the S' -torsion-free quotient of $\pi^* G_{\text{tf}}$ is $\mathcal{O}_{S'}$ -locally free. The S' -torsion-free quotient of $\pi^* G$, being equal to it, is then $\mathcal{O}_{S'}$ -locally free.

By Lemma 4.24, $L\pi^* \mathcal{L}$ has cohomology in $\text{Mod}_D(\mathcal{D}_{X \times S'/S'}(*D))$. We can therefore apply Steps 1 and 2 to deduce that $\pi^* \mathcal{L}$ is $\mathcal{D}_{X \times S'/S'}$ -holonomic, and it is regular by Definition 4.23(b). It is moreover π -good (cf. Lemma 2.12). Hence, $\pi_* \pi^* \mathcal{L}$ is $\mathcal{D}_{X \times S/S}$ -regular holonomic (Theorem 3.14).

Since \mathcal{L} is $\mathcal{D}_{X \times S/S}(*D)$ -coherent, the image \mathcal{L}' of the adjunction morphism $\mathcal{L} \rightarrow \pi_* \pi^* \mathcal{L}$ is $\mathcal{D}_{X \times S/S}$ -coherent, hence regular holonomic (Proposition 3.1(iii)) and its kernel is $\mathcal{D}_{X \times S/S}(*D)$ -coherent. It follows that the latter, which clearly satisfies the first point of Definition 4.23, also satisfies the second point since \mathcal{L} and \mathcal{L}' do so (cf. Proposition 3.1). Since π is biholomorphic above $S \setminus T$ for some closed analytic subset T of codimension ≥ 2 , this kernel satisfies the assumption of Step 1. It is thus $\mathcal{D}_{X \times S/S}$ -holonomic, hence \mathcal{L} is $\mathcal{D}_{X \times S/S}$ -holonomic, as desired. q.e.d.

Note that the assignment $\mathcal{L} \mapsto L = \mathcal{H}^0 \text{DR}(\mathcal{L}|_{X^* \times S})$ is a functor which takes values in the category of coherent S -locally constant sheaves, since the characteristic variety of $\mathcal{L}|_{X^* \times S}$ is contained in the zero section.

Proposition 4.26. *The category of strict regular holonomic $\mathcal{D}_{X \times S/S}$ -modules of D-type with singularities along D is equivalent to the category of S -locally constant sheaves on $X^* \times S$ which are $p^{-1}\mathcal{O}_S$ -locally free of finite rank, under the correspondences $\mathcal{L} \mapsto L = \mathcal{H}^0 \text{DR}(\mathcal{L}|_{X^* \times S})$ and $L \mapsto \mathcal{L} = \tilde{E}_L$.*

Proof. Owing to the Riemann-Hilbert correspondence on $X^* \times S$ (cf. [31, Rem. A.10]), we have $\mathcal{L}|_{X^* \times S} \simeq (\mathcal{O}_{X^* \times S} \otimes_{p^{-1}\mathcal{O}_S} L, d_{X \times S/S} \otimes \text{Id}) = E_L$. Since the natural morphism $\mathcal{L} \rightarrow j_* E_L$ sends isomorphically \mathcal{L} to \tilde{E}_L (Proposition 4.25(1)), the functor $L \mapsto \tilde{E}_L$

from the category of S -locally constant sheaves on $X^* \times S$ which are $p^{-1}\mathcal{O}_S$ -locally free of finite rank to that of regular holonomic $\mathcal{D}_{X \times S/S}$ -modules of D-type is essentially surjective. That it is fully faithful has been proved in Theorem 4.15(2). q.e.d.

4.e. Proof of Theorem 1. Although the next proposition is not general enough to prove Theorem 1, it will be one of the main tools for its proof.

Let $f : X \rightarrow Y$ be a morphism of real or complex analytic manifolds, we denote by $\mathcal{D}_{Y \leftarrow X/S}$ and by $\mathcal{D}_{X \rightarrow Y/S}$ the relative transfer bi-modules.

Proposition 4.27. *Let \mathcal{M} be a regular holonomic $\mathcal{D}_{X \times S/S}$ -module with X -support $Z = \bigcup_i Z_i$ (see (2.7)). Let $Y \subset X$ be a hypersurface such that $Y \supset Z_i$ if $\dim Z_i < \dim Z$, and $Z_o := Z \setminus (Y \cap Z)$ is smooth of pure dimension $\dim Z$. Then the localized $\mathcal{D}_{X \times S/S}$ -module $\mathcal{M}(*Y)$ is regular holonomic and locally isomorphic to the projective pushforward of a relative \mathcal{D} -module of D-type.*

Proof. The question is local. The assumption on Y implies that the characteristic variety of $\mathcal{M}|_{(X \setminus Y) \times S}$ is contained in $(T_{Z_o}^* X) \times S$. By Kashiwara's equivalence, $\mathcal{M}|_{(X \setminus Y) \times S}$ is the pushforward by the inclusion map of a coherent $\mathcal{O}_{Z_o \times S}$ -module with flat relative connection, which is thus of the form $(\mathcal{O}_{Z_o \times S} \otimes_{p^{-1}\mathcal{O}_S} L, d_{Z_o \times S/S})$ for some coherent locally constant $p_{Z_o}^{-1}\mathcal{O}_S$ -module L .

One can find a complex manifold X' together with a divisor with normal crossings $Y' \subset X'$ and a projective morphism $f : X' \rightarrow X$ which induces a biholomorphism $X' \setminus Y' \xrightarrow{\sim} Z_o$ (cf. e.g. [2, Prologue, Th. 4]). We set $\delta = \dim Z - \dim X = \dim X' - \dim X \leq 0$. For each ℓ , we consider the $\mathcal{D}_{X' \times S/S}$ -module $\mathcal{M}^\ell := \mathcal{H}_D^\ell f_* \mathcal{M}$. Although it is not yet known to be coherent, it is f - \mathcal{O} -good in the sense of Definition 2.2 (Proposition 2.11(1)). By considering the filtration by the order of the pole along Y' , one checks that $\mathcal{M}^\ell(*Y')$ is also f - \mathcal{O} -good (cf. Remark 2.5(1)).

If $\ell \neq \delta$, the sheaf-theoretic restriction of \mathcal{M}^ℓ to $(X' \setminus Y') \times S$ is zero, therefore $\mathcal{M}^\ell(*Y') = 0$ owing to \mathcal{O} -quasi-coherence (cf. Lemma 2.3(2)). Since $\mathcal{O}_{X' \times S}(*Y')$ is flat over $\mathcal{O}_{X' \times S}$, we conclude that

$$(4.28) \quad {}_D f^*(\mathcal{M}(*Y))[\delta] \simeq ({}_D f_* \mathcal{M})(*Y')[\delta] \simeq \mathcal{M}'^\delta(*Y').$$

We can interpret ${}_D f^*(\mathcal{M}(*Y))$ as the pullback of $\mathcal{M}(*Y)$ as a $\mathcal{D}_{X \times S/S}(*Y)$ -module. Since f is a modification and $Y' = f^{-1}(Y)$, we have

$$(4.29) \quad \begin{aligned} \mathcal{D}_{X' \times S/S}(*Y') &\simeq \mathcal{O}_{X' \times S}(*Y') \otimes_{f^{-1}(\mathcal{O}_{X \times S}(*Y))} f^{-1} \mathcal{D}_{X \times S/S}(*Y) \\ &= \mathcal{D}_{X' \rightarrow X/S}(*Y') \end{aligned}$$

(in local coordinates x' in X' and x in X , the determinant of the matrix expressing $\partial_{x'_i}$ in terms of ∂_{x_j} is invertible in $\mathcal{O}_{X' \times S}(*Y')$). It follows that $\mathcal{M}'^\delta(*Y') = \mathcal{D}_{X' \times S/S}(*Y') \otimes_{f^{-1} \mathcal{D}_{X \times S/S}(*Y)} f^{-1} \mathcal{M}(*Y)$ is $\mathcal{D}_{X' \times S/S}(*Y')$ -coherent.

Since Li_s^* commutes with ${}_D f^*$, $\mathcal{M}'^\delta(*Y')$ is $\mathcal{D}_{X' \times S/S}(*Y')$ -coherent of D-type (Definition 4.23) so, by Proposition 4.25(2), it is $\mathcal{D}_{X' \times S/S}$ -regular holonomic. Since $\mathcal{M}'^\delta(*Y')$ is f - \mathcal{O} -good and $\mathcal{D}_{X' \times S/S}$ -coherent, it is f -good.

According to [31, Cor. 2.4], ${}_D f_*(\mathcal{M}'^\delta(*Y'))$ has regular holonomic cohomology. Furthermore, since $\mathcal{H}^j({}_D f_* \mathcal{M}'^\delta(*Y'))$ is supported on $Y \times S$ for $j \neq 0$, and since ${}_D f_*(\mathcal{M}'^\delta(*Y')) \simeq ({}_D f_* \mathcal{M}'^\delta)(*Y)$, we have

$${}_D f_*(\mathcal{M}'^\delta(*Y')) \simeq \mathcal{H}^0 {}_D f_*(\mathcal{M}'^\delta(*Y')) \simeq \mathcal{H}^0({}_D f_* \mathcal{M}'^\delta)(*Y).$$

On the other hand, there is a natural adjunction morphism (cf. [13, Lem. 4.28 & Prop. 4.34])

$${}_{\mathcal{D}}f_* {}_{\mathcal{D}}f^* \mathcal{M}[\delta] \longrightarrow \mathcal{M},$$

which induces a morphism of coherent $\mathcal{D}_{X \times S/S}(*Y)$ -modules

$$(4.30) \quad \mathcal{H}^0({}_{\mathcal{D}}f_* \mathcal{M}'^\delta(*Y')) \simeq (\mathcal{H}^0 {}_{\mathcal{D}}f_* \mathcal{M}'^\delta)(*Y) \longrightarrow \mathcal{M}(*Y),$$

where the left-hand side is $\mathcal{D}_{X \times S/S}$ -coherent and regular holonomic. Its cokernel is zero on $(X \setminus Y) \times S$ and $\mathcal{D}_{X \times S/S}(*Y)$ -coherent, hence it is zero according to Lemma 2.3(2), so that this morphism is an isomorphism. In conclusion, $\mathcal{M}(*Y)$ is regular holonomic. q.e.d.

Corollary 4.31. *Let \mathcal{M} be a regular holonomic $\mathcal{D}_{X \times S/S}$ -module and let Y be any hypersurface in X . Then the localized $\mathcal{D}_{X \times S/S}$ -module $\mathcal{M}(*Y)$ is regular holonomic (hence $\mathcal{D}_{X \times S/S}$ -coherent).*

Proof. The question is local on $X \times S$. Let Z denote the X -support of \mathcal{M} . We argue by induction on the dimension of Z . The case where $\dim Z = 0$ is clear because either $Z \subset Y$ and then $\mathcal{M}(*Y) = 0$, or $Z \not\subset Y$ and then $\mathcal{M}(*Y) = \mathcal{M}$.

Let Y' be a hypersurface satisfying the properties as in Proposition 4.27. Then $Y \cup Y'$ satisfies the same properties. We consider the following commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \Gamma_{[Y']} \mathcal{M} & \longrightarrow & \mathcal{M} & \longrightarrow & \mathcal{M}(*Y') & \longrightarrow & R^1 \Gamma_{[Y']} \mathcal{M} & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & (\Gamma_{[Y']} \mathcal{M})(*Y) & \longrightarrow & \mathcal{M}(*Y) & \longrightarrow & \mathcal{M}(*(Y' \cup Y)) & \longrightarrow & (R^1 \Gamma_{[Y']} \mathcal{M})(*Y) & \longrightarrow & 0 \end{array}$$

By Proposition 4.27, the terms of the top horizontal line, together with $\mathcal{M}(*(Y' \cup Y))$, are regular holonomic. On the other hand, the support of the regular holonomic modules $\Gamma_{[Y']} \mathcal{M}$ and $R^1 \Gamma_{[Y']} \mathcal{M}$ is of dimension $< \dim Z$, hence the assertion holds for these modules by the induction hypothesis, and the extreme terms of the lower horizontal line are regular holonomic. It follows that the remaining term $\mathcal{M}(*Y)$ is regular holonomic. q.e.d.

Corollary 4.32. *Let \mathcal{M} be a regular holonomic $\mathcal{D}_{X \times S/S}$ -module and let Y be any closed analytic subset in X . Then $R\Gamma_{[Y]} \mathcal{M}$ belongs to $\mathcal{D}_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$.*

Proof. The question is local. The case of a hypersurface follows from Corollary 4.31. The general case follows by writing locally Y as the intersection of hypersurfaces. q.e.d.

Corollary 4.33. *Let Y be a closed hypersurface of X and let \mathcal{M} be a coherent $\mathcal{D}_{X \times S/S}(*Y)$ -module \mathcal{L} of D -type. Then \mathcal{M} belongs to $\text{Mod}_{\text{rhol}}(\mathcal{D}_{X \times S/S})$.*

Proof. This is obtained by the arguments used in the proof of Proposition 4.27. q.e.d.

Proof of Theorem 1. We regard the morphism f as the composition of a closed inclusion and a projection. The latter case is clear, so we only consider the case where $f : Y \hookrightarrow X$ is the inclusion of a closed submanifold. Then Corollary 4.32 implies that $R\Gamma_{[Y]} \mathcal{M}$ is regular holonomic. By Kashiwara's equivalence, we conclude that ${}_{\mathcal{D}}f^* \mathcal{M}$ belongs to $\mathcal{D}_{\text{rhol}}^b(\mathcal{D}_{Y \times S/S})$. q.e.d.

Proposition 4.34. *If \mathcal{M}, \mathcal{N} are objects of $D_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$, then so is $\mathcal{M} \otimes_{\mathcal{O}_{X \times S}}^L \mathcal{N}$.*

Proof. Recall that the tensor product has been defined in Remark 2.16. According to Theorem 1 applied to the diagonal embedding δ , it is enough to prove that the S -external tensor product $\mathcal{M} \boxtimes_{\mathcal{O}_S}^L \mathcal{N}$ is an object of $D_{\text{rhol}}^b(\mathcal{D}_{(X \times X) \times S/S})$. Holonomicity has been observed in Remark 2.16. Regularity follows from the isomorphism $Li_s^*(\mathcal{M} \boxtimes_{\mathcal{O}_S}^L \mathcal{N}) \simeq (Li_s^* \mathcal{M}) \boxtimes_{\mathbb{C}} (Li_s^* \mathcal{N})$ and from the regular holonomicity of the latter as a complex in $D^b(\mathcal{D}_{X \times X})$. q.e.d.

Let Y_i ($i = 1, \dots, p$) be hypersurfaces of X defined as the zero set of holomorphic functions $h_i : X \rightarrow \mathbb{C}$, set $Y = \bigcup_i Y_i$ and let \mathcal{N} be a $\mathcal{D}_{X \times S/S}(*Y)$ -module. We regard \mathcal{N} as an $\mathcal{O}_{X \times S}(*Y)$ -module with flat relative connection ∇ , and for a tuple $\alpha = (\alpha_1, \dots, \alpha_p)$ of holomorphic functions $\alpha_i : S \rightarrow \mathbb{C}$, we denote by $\mathcal{N}h^\alpha$ the $\mathcal{O}_{X \times S}$ -module \mathcal{N} , endowed with the flat relative connection $\nabla + \sum_i \alpha_i \text{Id} \otimes dh_i/h_i$. The functor $\mathcal{N} \rightarrow \mathcal{N}h^\alpha$ is an auto-equivalence of the category $\text{Mod}(\mathcal{D}_{X \times S/S}(*Y))$, as well as of $\text{Mod}_{\text{coh}}(\mathcal{D}_{X \times S/S}(*Y))$. We have a functorial isomorphism $\mathcal{N}h^\alpha \simeq \mathcal{O}_{X \times S}(*Y)h^\alpha \otimes_{\mathcal{O}_{X \times S}} \mathcal{N} \simeq \mathcal{O}_{X \times S}(*Y)h^\alpha \otimes_{\mathcal{O}_{X \times S}}^L \mathcal{N}$.

Corollary 4.35. *Assume that \mathcal{M} is a regular holonomic $\mathcal{D}_{X \times S/S}$ -module. Then so is $\mathcal{M}(*Y)h^\alpha$.*

Sketch of proof. One checks that the coherent $\mathcal{D}_{X \times S/S}(*Y)$ -module $\mathcal{O}_{X \times S}(*Y)h^\alpha$ is of D-type along Y by using that $\mathcal{O}_X(*Y)h^{\alpha(s)}$ is regular holonomic for each $s \in S$. Hence $\mathcal{O}_{X \times S}(*Y)h^\alpha$ is regular holonomic by Corollary 4.33. Then, on noting that $\mathcal{M}(*Y)h^\alpha \simeq \mathcal{O}_{X \times S}(*Y)h^\alpha \otimes_{\mathcal{O}_{X \times S}} \mathcal{M}$, we conclude by applying Proposition 4.34. q.e.d.

Example 4.36 (Generalized Mellin transform). Let f_1, \dots, f_p be meromorphic functions on X , i.e., locally each f_i is the quotient of two holomorphic functions h_i, g_i without common factor. Let Y be the union of the divisors of zeros and poles of the functions f_i , i.e., locally the divisors of zeros of h_i, g_i . We also set $S = \mathbb{C}^p$ with its analytic topology. It is usual to denote by $\mathcal{O}_{X \times S}(*Y)f^s$ the $\mathcal{O}_{X \times S}$ -module $\mathcal{O}_{X \times S}(*Y)$ equipped with the twisted connection $d + \sum_i s_i df_i/f_i$. The same argument as in Corollary 4.35 shows that $\mathcal{O}_{X \times S}(*Y)f^s$ is regular holonomic and that, if \mathcal{M} is a regular holonomic \mathcal{D}_X -module, then the $\mathcal{D}_{X \times S/S}$ -module $q^*(\mathcal{M}(*Y))f^s := q^* \mathcal{M} \otimes_{\mathcal{O}_{X \times S}} \mathcal{O}_{X \times S}(*Y)f^s$ (where $q : X \times S \rightarrow X$ denotes the projection) is also regular holonomic.

Furthermore, the $\mathcal{D}_{X \times S/S}$ -submodule of $q^*(\mathcal{M}(*Y))f^s$ generated by the image of $q^* \mathcal{M} \otimes 1 \cdot f^s$ is also regular holonomic, according to Proposition 3.1(iii), as it is clearly coherent (being locally of finite type in a coherent $\mathcal{D}_{X \times S/S}$ -module). This property is the S -analytic variant of [21, Prop. 13], with the regularity assumption however.

Since \mathcal{M} is regular holonomic, it is good, and so is $q^*(\mathcal{M}(*Y))f^s$. Therefore, if X is compact, the \mathcal{D} -module pushforward ${}_{\text{D}p_*} [q^*(\mathcal{M}(*Y))f^s]$ is an object of $D_{\text{coh}}^b(\mathcal{O}_S)$. This is the generalized Mellin transform of \mathcal{M} with respect to (f_1, \dots, f_p) .

4.f. Another characterization of regular holonomicity. For a closed analytic subset Y of X , we denote by $\mathcal{O}_{\widehat{Y \times S}} = \varprojlim_{k \in \mathbb{Z}} \mathcal{O}_{X \times S}/\mathcal{I}^k$ the formal completion of $\mathcal{O}_{X \times S}$

along $Y \times S$, where \mathcal{J} denotes the defining ideal of $Y \times S$ in $X \times S$. Let $i : Y \hookrightarrow X$ denote the inclusion. We consider the exact sequence of sheaves supported on $Y \times S$:

$$0 \longrightarrow Ri_*i^{-1}\mathcal{O}_{X \times S} \longrightarrow \mathcal{O}_{\widehat{Y \times S}} \longrightarrow \mathcal{Q}_{Y \times S} \longrightarrow 0.$$

Corollary 4.37. *If $\mathcal{M} \in \mathbf{D}_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$, then the complexes*

$$R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{O}_{\widehat{Y \times S}}) \quad \text{and} \quad R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{Q}_{Y \times S})$$

belong to $\mathbf{D}_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$.

Proof. It follows from Corollary 4.32 that $R\Gamma_{[Y]}\mathcal{M} \in \mathbf{D}_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$, so the complex $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(R\Gamma_{[Y]}\mathcal{M}, \mathcal{O}_{X \times S})$ belongs to $\mathbf{D}_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$, according to [29, Th. 3.7]. On the one hand, by mimicking the proof when S is reduced to a point (cf. e.g. [23, Cor. 2.7-2]), one finds a natural isomorphism

$$R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(R\Gamma_{[Y]}\mathcal{M}; \mathcal{O}_{X \times S}) \xrightarrow{\sim} R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{O}_{\widehat{Y \times S}}),$$

hence the S - \mathbb{C} -constructibility of the latter complex. On the other hand, we have natural isomorphisms

$$\begin{aligned} Ri_*i^{-1}R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{O}_{X \times S}) &\xrightarrow{\sim} Ri_*R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(i^{-1}\mathcal{M}, i^{-1}\mathcal{O}_{X \times S}) \\ &\simeq R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, Ri_*i^{-1}\mathcal{O}_{X \times S}), \end{aligned}$$

showing S - \mathbb{C} -constructibility of the latter complex, and therefore that of the complex $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{Q}_{Y \times S})$. q.e.d.

Theorem 4.38. *Let \mathcal{M} belong to $\mathbf{D}_{\text{hol}}^b(\mathcal{D}_{X \times S/S})$. Then \mathcal{M} is regular holonomic if and only if for any germ of closed analytic subset $Y \subset X$, $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{Q}_{Y \times S})$ is isomorphic to zero.*

Lemma 4.39. *For any closed analytic subset $Y \subset X$ we have*

$$Li_s^*\mathcal{O}_{\widehat{Y \times S}} \simeq \mathcal{O}_{\widehat{Y}} \quad \text{and} \quad Li_s^*\mathcal{Q}_{Y \times S} \simeq \mathcal{Q}_Y.$$

Proof. Thanks to the properties of Li_s^* (cf. [29, Prop. 3.1]) the result follows from Mittag-Leffler's condition since the morphisms $\mathcal{O}_{X \times S}/\mathcal{J}^{k+1} \rightarrow \mathcal{O}_{X \times S}/\mathcal{J}^k$ are surjective. q.e.d.

Proof of Theorem 4.38. We first remark that the theorem holds if S is reduced to a point, according to [14, (6.4.6) & (6.4.7)].

If $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{Q}_{Y \times S}) = 0$ for all germ $Y \subset X$, then for any $s \in S$

$$(4.40) \quad \begin{aligned} R\mathcal{H}om_{\mathcal{D}_X}(Li_s^*\mathcal{M}, \mathcal{Q}_Y) &\simeq R\mathcal{H}om_{\mathcal{D}_X}(Li_s^*\mathcal{M}, Li_s^*\mathcal{Q}_{Y \times S}) \\ &\simeq Li_s^*R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{Q}_{Y \times S}) = 0. \end{aligned}$$

From the preliminary remark we conclude that $Li_s^*\mathcal{M} \in \mathbf{D}_{\text{rhol}}^b(\mathcal{D}_X)$ for any $s \in S$, hence $\mathcal{M} \in \mathbf{D}_{\text{rhol}}^b(\mathcal{D}_{X \times S/S})$.

Conversely, if \mathcal{M} is regular holonomic, then $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{Q}_{Y \times S})$ is S - \mathbb{C} -constructible, and by the variant of Nakayama's lemma (cf. [29, Prop. 2.2]), it is zero if (and only if) $Li_s^*R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{Q}_{Y \times S}) = 0$ for any $s \in S$. Reading (4.40) backwards and according to the preliminary remark, we find that the latter property is satisfied, hence $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{Q}_{Y \times S}) = 0$. q.e.d.

5. Construction of the relative Riemann-Hilbert functor RH_X^S

In this section, we extend the definition of the functor RH_X^S introduced in [31] when $\dim S = 1$ to the case $\dim S \geq 2$. We will check that it satisfies properties similar to those explained in [31, 8].

5.a. Reminder on the subanalytic site and complements. We first recall the main results in [28]. We consider the site in the real analytic manifold $S_{\mathbb{R}}$ given by the usual topology, that is, where the family $\mathrm{Op}(S)$ consisting of all open sets. On the other hand, we have the subanalytic site X_{sa} underlying the real analytic manifold $X_{\mathbb{R}}$ for which the family of open subsets $\mathrm{Op}(X_{\mathrm{sa}})$ consists of subanalytic open subsets in $X_{\mathbb{R}}$. Lastly, we let $X_{\mathrm{sa}} \times S$ be the subanalytic site underlying the real analytic manifold $X_{\mathbb{R}} \times S_{\mathbb{R}}$, for which the family of open sets $\mathrm{Op}(X_{\mathrm{sa}} \times S)$ consists of those which are finite unions of products $U \times V$ with $U \in \mathrm{Op}(X_{\mathrm{sa}})$ and V is open in S . A subset $T \subset \mathrm{Op}(X_{\mathrm{sa}} \times S)$ is a covering of $W = U \times V \in \mathrm{Op}(X_{\mathrm{sa}} \times S)$ if and only if it admits a refinement $\{U_i \times V_j\}_{i \in I, j \in J}$ such that $\{U_i\}_{i \in I}$ is a covering of U (in X_{sa}) and $\{V_j\}_{j \in J}$ is a covering of V (in S). In particular, when U is relatively compact I is finite but J needs not to be so even if V is relatively compact.

We have the following commutative diagram, where the arrows are natural morphisms of sites induced by the inclusion of families of open subsets.

$$(5.1) \quad \begin{array}{ccc} & X_{\mathrm{sa}} \times S & \\ \rho_S \nearrow & & \searrow a \\ X \times S & \xrightarrow{\rho'} & X_{\mathrm{sa}} \times S_{\mathrm{sa}} \\ \rho \searrow & & \nearrow \eta \\ & (X \times S)_{\mathrm{sa}} & \end{array}$$

We recall that

(5.2) ρ_S^{-1} commutes with tensor products (cf. [18, Lem. 18.3.1(ii)(c)]) and $\rho_S^{-1} \rho_{S^*} \simeq \mathrm{Id}$. Furthermore, ρ_S^{-1} admits a left adjoint $\rho_S!$ which is exact and commutes with tensor products (cf. [28, §3]).

The following proposition generalizes [31, Props. 3.3 & 3.5]:

Proposition 5.3. *The category $\mathrm{Mod}_{\mathbb{R}\text{-c}}(p_X^{-1} \mathcal{O}_S)$ is acyclic for ρ_S :*

$$\forall F \in \mathrm{Mod}_{\mathbb{R}\text{-c}}(p_X^{-1} \mathcal{O}_S), \quad \forall k \geq 1, \quad \mathcal{H}^k \rho_{S^*} F = 0.$$

In particular ρ_S is exact on $\mathrm{Mod}_{\mathbb{R}\text{-c}}(p_X^{-1} \mathcal{O}_S)$.

Proof. We have to show that $R^k \Gamma(\Omega; F) = 0$ for $k \neq 0$, where $\Omega = U \times V$ runs on a basis of the topology of $X_{\mathrm{sa}} \times S$; hence we may assume that U is relatively compact subanalytic in X and V a relatively compact open Stein subset in S . Hence it is sufficient to prove that, for each $k \neq 0$, there exists a covering $\{U_i \times V_j\}_{i \in I, j \in J}$ of $U \times V$ (thus I is finite) such that $H^k(U_i \times V_j, F) = 0$.

Let $X = \bigcup_{\alpha} X_{\alpha}$ be a Whitney stratification adapted to F . By [15, Prop. 8.2.5], there exist a simplicial complex $\mathbf{K} = (K, \Delta)$ and a homeomorphism $\iota : |\mathbf{K}| \simeq X$ such that, for any simplex $\sigma \in \Delta$, there exists α such that $\iota(|\sigma|) \subset X_{\alpha}$ and $\iota(|\sigma|)$ is a subanalytic manifold of X . Moreover, we may assume that U is a finite union of

the images by ι of open subsets $U(\sigma)$ of $|\mathbf{K}|$, with $U(\sigma) = \bigcup_{\tau \in \Delta, \tau \supset \sigma} |\tau|$. We shall see that we may take for $U_i \times V_j$ the open sets $i(U(\sigma)) \times V$. Still denoting by ι the homeomorphism $|\mathbf{K}| \times S \rightarrow X \times S$, it is enough to prove that for any $\sigma \in \Delta$ and any $x \in |\sigma|$, we have:

- (i) $H^0(U(\sigma) \times V, \iota^* F) \simeq H^0(V, F_{\{x\} \times S})$,
- (ii) $H^k(U(\sigma) \times V, \iota^* F) = H^k(V, F_{\{x\} \times S}) = 0$, for $k \neq 0$.

The proof of (i) and (ii) proceeds mimicking the proof of [15, Prop. 8.1.4], by using [31, Prop. A.7], the fact that the $F|_{\{x\} \times S}$ is \mathcal{O}_S -coherent and that V is Stein. \square e.d.

The following proposition can be proved in a way similar to [31, Prop. 3.5], where it is assumed that $\dim S = 1$.

Proposition 5.4 (cf. [31, Prop. 3.5]). *Let $F \in D_{\mathbb{R}\text{-c}}^b(p^{-1}\mathcal{O}_S)$. Then F is isomorphic to a complex*

$$0 \longrightarrow \bigoplus_{i_\alpha \in I_\alpha} p^{-1}\mathcal{O}_S \otimes \mathbb{C}_{U_\alpha, i_\alpha \times V_\alpha, i_\alpha} \longrightarrow \cdots \longrightarrow \bigoplus_{i_\beta \in I_\beta} p^{-1}\mathcal{O}_S \otimes \mathbb{C}_{U_\beta, i_\beta \times V_\beta, i_\beta} \longrightarrow 0,$$

where $\{U_\gamma, i_\gamma\}_{\gamma, i_\gamma}$ are locally finite families of relatively compact open subanalytic subsets of X and $\{V_\gamma, i_\gamma\}_{\gamma, i_\gamma}$ is a (locally finite) covering of S .

Remark 5.5. We remark that, for any S , the site $X_{\text{sa}} \times S$ is a ringed site both relatively to the sheaf $\rho_{S*}(p^{-1}\mathcal{O}_S)$ and to the sheaf $\rho_{S!}\mathcal{O}_{X \times S}$ (cf. [18, p. 449]), and ρ_S is a morphism of ringed sites in both cases. Thus, according to [18, Lem. 18.3.1, Th. 18.6.9], ρ_S^{-1} commutes with \otimes , $\otimes_{\rho_{S*}p_X^{-1}\mathcal{O}_S}$, $\otimes_{\rho_{S!}\mathcal{O}_{X \times S}}$, $\otimes_{\rho_{S*}p_X^{-1}\mathcal{O}_S}^L$ and $\otimes_{\rho_{S!}\mathcal{O}_{X \times S}}^L$.

Let $\pi : S' \rightarrow S$ be a morphism of complex manifolds. Since $\mathcal{O}_{S'}$ is a $\pi^{-1}\mathcal{O}_S$ -module, $\rho_{S'*}(p^{-1}\mathcal{O}'_S)$ is a $\rho_{S'*}(p^{-1}\pi^{-1}\mathcal{O}_S)$ -module hence a $\pi^{-1}\rho_{S*}(p^{-1}\mathcal{O}_S)$ -module. Similarly, $\rho_{S'!}\mathcal{O}_{X \times S'}$ is a $\pi^{-1}\rho_{S!}\mathcal{O}_{X \times S}$ -module. In other words, π induces a morphism of ringed sites with respect to both sheaves of rings. Consequently, according to [18, Th. 18.6.9(i)], the derived functors $L\pi^* : D(\rho_{S*}p^{-1}\mathcal{O}_S) \rightarrow D(\rho_{S'*}p'^{-1}\mathcal{O}'_S)$ resp. (keeping the same notation π for the morphism $\text{Id} \times \pi$), $L\pi^* : D(\rho_{S!}\mathcal{O}_{X \times S}) \rightarrow D(\rho_{S'!}\mathcal{O}_{X \times S'})$ are well-defined.

We recall that ρ_{S*} commutes with $R\pi_*$ and ρ_S^{-1} commutes with π^{-1} according to [18, Prop. 17.5.3]. Furthermore (π^{-1}, π_*) is a pair of adjoint functors according to [18, Th. 17.5.2(i)] and ρ_S^{-1} commutes with $R\pi_*$ which follows by copying the proof of [33, Prop. 2.2.1(ii)] since ρ_S^{-1} admits a left adjoint ($\rho_{S!}$) hence takes injective sheaves to injective sheaves.

Note that, in general, we do not have $\pi^{-1}\rho_{S*} = \rho_{S'*}\pi^{-1}$ while we have $\pi^{-1}\rho_{S!} = \rho_{S'!}\pi^{-1}$.

We thus have isomorphisms of functors

$$\rho_{S'}^{-1}\pi^{-1}\rho_{S*} \simeq \pi^{-1}, \quad \rho_{S'*}\rho_{S'}^{-1}\pi^{-1}\rho_{S*} \simeq \rho_{S'*}\pi^{-1}$$

and, composing with the natural morphism $\text{Id} \rightarrow \rho_{S'*}\rho_{S'}^{-1}$, we deduce a natural morphism

$$\pi^{-1}\rho_{S*} \longrightarrow \rho_{S'*}\pi^{-1}.$$

Remark 5.6. From the fact that a_* is fully faithful and that $a^{-1}a_* = \text{Id}$, thus $\rho_{S*} = a^{-1}\rho'_*$ as explained in [28] before Proposition 3.1, we deduce that, for any

open subset $U \times V \in \text{Op}^c(X_{\text{sa}} \times S)$, the constant sheaf $\mathbb{C}_{U \times V}$ on the site $X_{\text{sa}} \times S$ coincides with $\rho_{S*} \mathbb{C}_{U \times V}$. Similarly

$$\rho_{S*}(\mathbb{C}_{U \times V} \otimes p^{-1} \mathcal{O}_S) \simeq \rho_{S*}(\mathbb{C}_{U \times V}) \otimes \rho_{S*}(p^{-1} \mathcal{O}_S)$$

since the isomorphism holds true with ρ' instead of ρ_S , and that

$$\pi^{-1} \rho_{S*}(\mathbb{C}_{U \times V}) = \pi^{-1}(\mathbb{C}_{U \times V}) = \mathbb{C}_{U \times \pi^{-1}V} = \rho_{S'*}(\mathbb{C}_{U \times \pi^{-1}V}) = \rho_{S'*} \pi^{-1}(\mathbb{C}_{U \times V}).$$

Lemma 5.7. *Let $\pi : S' \rightarrow S$ be a morphism. Then we have an isomorphism of functors on $\mathbb{D}_{\mathbb{R}\text{-c}}^b(p^{-1} \mathcal{O}_S)$ with values in $\mathbb{D}_{\mathbb{R}\text{-c}}(\rho_{S'*} p'^{-1} \mathcal{O}_{S'})$:*

$$L\pi^* \rho_{S*}(F) \simeq \rho_{S'*}(L\pi^* F).$$

Proof. By Proposition 5.4 and [8, Prop.1.6 & Rem.1.8], we may assume that $F = \mathbb{C}_{U \times V} \otimes p^{-1} \mathcal{O}_S$, for some relatively compact open subsets respectively of X and S , with U subanalytic. We have

$$\begin{aligned} L\pi^* \rho_{S*}(F) &= \rho_{S'*}(p'^{-1} \mathcal{O}_{S'}) \otimes_{\pi^{-1} \rho_{S*}(p^{-1} \mathcal{O}_S)}^L \pi^{-1} \rho_{S*}(\mathbb{C}_{U \times V} \otimes p^{-1} \mathcal{O}_S) \\ &\simeq \rho_{S'*}(p'^{-1} \mathcal{O}_{S'}) \otimes_{\pi^{-1} \rho_{S*}(p^{-1} \mathcal{O}_S)}^L \pi^{-1} \rho_{S*}(\mathbb{C}_{U \times V}) \otimes \pi^{-1} \rho_{S*}(p^{-1} \mathcal{O}_S) \\ &\simeq \rho_{S'*}(p'^{-1} \mathcal{O}_{S'}) \otimes \pi^{-1} \rho_{S*}(\mathbb{C}_{U \times V}) \simeq \rho_{S'*}(p'^{-1} \mathcal{O}_{S'}) \otimes \rho_{S'*} \pi^{-1}(\mathbb{C}_{U \times V}) \\ &\simeq \rho_{S'*}(p'^{-1} \mathcal{O}_{S'} \otimes \mathbb{C}_{U \times \pi^{-1}V}) \simeq \rho_{S'*}(L\pi^* F). \end{aligned} \quad \text{q.e.d.}$$

5.b. The construction of RH_X^S and behaviour under pushforward. For any object $F \in \mathbb{D}_{\mathbb{R}\text{-c}}^b(p^{-1} \mathcal{O}_S)$ we define:

$$\begin{aligned} \text{TH}_X^S(F) &:= \rho_S^{-1} R\mathcal{H}om_{\rho_{S*} p^{-1} \mathcal{O}_S}(\rho_{S*} F, \mathfrak{D}\mathfrak{b}_{X \times S}^{t,S}), \\ \text{RH}_X^S(F) &:= \rho_S^{-1} R\mathcal{H}om_{\rho_{S*} p^{-1} \mathcal{O}_S}(\rho_{S*} F, \mathcal{O}_{X \times S}^{t,S}[d_X]). \end{aligned}$$

Each expression gives a triangulated functor on $F \in \mathbb{D}_{\mathbb{R}\text{-c}}^b(p^{-1} \mathcal{O}_S)^{\text{op}}$ with values on $\mathbb{D}^b(\mathcal{D}_{X \times S/S})$. Moreover, if $\dim S = 1$ we recover the definitions of [31]:

$$\begin{aligned} \text{TH}_X^S(F) &:= \rho_S^{-1} R\mathcal{H}om_{\rho_{S*} p^{-1} \mathcal{O}_S}(\rho_{S*} F, \mathfrak{D}\mathfrak{b}_{X \times S}^{t,S,\sharp}), \\ \text{RH}_X^S(F) &:= \rho_S^{-1} R\mathcal{H}om_{\rho_{S*} p^{-1} \mathcal{O}_S}(\rho_{S*} F, \mathcal{O}_{X \times S}^{t,S,\sharp}[d_X]). \end{aligned}$$

This follows by adjunction upon noting that a (cf. (5.1)) is proper, $a^1 = a^{-1}$, $a^{-1} a_* = \text{Id}$ and $a^{-1} \mathcal{O}_{X \times S}^{t,S,\sharp} = \mathcal{O}_{X \times S}^{t,S}$.

Let U be a subanalytic open subset in X and let us denote by $j : U \times S \rightarrow X \times S$ the inclusion.

Lemma 5.8 (Extension in the case of an open subanalytic set)

Let $F \in \mathbb{D}_{\mathbb{R}\text{-c}}^b(p_U^{-1} \mathcal{O}_S)$. Then there are natural isomorphisms in $\mathbb{D}^b(\mathcal{D}_{X \times S/S})$

$$(5.8^*) \quad \text{TH}_X^S(j_! F) \simeq \rho_S^{-1} Rj_* R\mathcal{H}om_{\rho_{S*} p_U^{-1} \mathcal{O}_S}(\rho_{S*} F, j^{-1} \mathfrak{D}\mathfrak{b}_{X \times S}^{t,S}),$$

$$(5.8^{**}) \quad \text{RH}_X^S(j_! F)[-d_X] \simeq \rho_S^{-1} Rj_* R\mathcal{H}om_{\rho_{S*} p_U^{-1} \mathcal{O}_S}(\rho_{S*} F, j^{-1} \mathcal{O}_{X \times S}^{t,S}).$$

Proof. This lemma is a variant of [31, Lem. 3.25] and its proof is similar to that of loc. cit. q.e.d.

We have the relative version of [12, Th. 4.1]:

Theorem 5.9. *Let $f : X \rightarrow Y$ be a proper morphism of real analytic manifolds, let $F \in \mathbf{D}_{\mathbb{R}\text{-c}}^{\mathbf{b}}(p_Y^{-1}\mathcal{O}_S)$, Then we have a canonical morphism in $\mathbf{D}(\mathcal{D}_{Y \times S/S})$*

$${}_{\mathbf{D}}f_* \mathrm{TH}_X^S(f^{-1}F) \longrightarrow \mathrm{TH}_Y^S(F).$$

The proof is stepwise similar to that with $\dim S = 1$ in [8, Lem. 2.5] so we omit it. Proposition 7.1 of [16] (see also [16, Th. 4.4]) has a relative version already used in [31], whose proof is given below:

Theorem 5.10. *Let $f : X \rightarrow Y$ be a morphism of complex analytic manifolds, let $F \in \mathbf{D}_{\mathbb{R}\text{-c}}^{\mathbf{b}}(p_X^{-1}\mathcal{O}_S)$, and assume that f is proper on $\mathrm{Supp} F$. Then there is a canonical isomorphism in $\mathbf{D}^{\mathbf{b}}(\mathcal{D}_{X \times S/S})$*

$$(5.10*) \quad {}_{\mathbf{D}}f_* \mathrm{RH}_X^S(F)[d_X] \longrightarrow \mathrm{RH}_Y^S(Rf_*F)[d_Y].$$

Proof. We recall that

$$(5.11) \quad \mathrm{RH}_X^S(F) \simeq R\mathcal{H}om_{\mathcal{D}_{\overline{X} \times \overline{S}/\overline{S}}}(\mathcal{O}_{\overline{X} \times \overline{S}}, \mathrm{TH}^S(F))[d_X].$$

In view of Theorem 5.9, where we replace F by Rf_*F , and by adjunction, we derive a natural morphism

$$(5.12) \quad \mu_F : {}_{\mathbf{D}}f_* \mathrm{TH}_X^S(F) \longrightarrow \mathrm{TH}_Y^S(Rf_*F).$$

Lemma 5.13. *The morphism μ_F is an isomorphism.*

Proof. We will follow the proof of [16, Th. 4.4] which contains the statement in the absolute case. We decompose f by the graph embedding so that we first assume that $f : X \rightarrow Y$ is a closed embedding and next we treat the case of a smooth morphism.

Step 1. Let us assume that $f : X \rightarrow Y$ is the embedding of a closed manifold. The morphism being defined, to check it is an isomorphism we may consider a local situation. In view of [8, Prop. 1.6] (with the trivial adaptation of the category \mathcal{S} to the new site $Y_{\mathrm{sa}} \times S$) and Theorem 5.9, together with Proposition 5.4, we are thus conducted to assume that $F \simeq \mathbb{C}_{\Omega \times V} \otimes p^{-1}\mathcal{O}_S$, for some open subanalytic subset Ω of X and V open in S . Up to restricting to V , we may write S instead of V .

As in the proof of [8, Prop. 2.4], let us start by proving the statement for F of the form $\mathbb{C}_{Z \times S} \otimes p^{-1}\mathcal{O}_S$, for any closed subanalytic subset Z of X . The conclusion for $\Omega := X \setminus Z$ will follow easily by functoriality, considering the exact sequence

$$0 \longrightarrow \mathbb{C}_{\Omega \times S} \longrightarrow \mathbb{C}_{X \times S} \longrightarrow \mathbb{C}_{Z \times S} \longrightarrow 0.$$

We have $\mathrm{TH}_X^S(F) \simeq \Gamma_{Z \times S}(\mathfrak{D}\mathbf{b}_{X \times S})$ regarded as a $\mathcal{D}_{Y_{\mathbb{R}} \times S_{\mathbb{R}}/S}$ -module. We note that the local structure of distributions supported by a submanifold entail that

$$\Gamma_{X \times S}(\mathcal{D}_{Y \times S}) \simeq f_*(\mathcal{D}_{Y \leftarrow X/S} \otimes_{\mathcal{D}_{X \times S/S}} \mathfrak{D}\mathbf{b}_{X \times S}).$$

Then

$$\begin{aligned} {}_{\mathbf{D}}f_* \Gamma_{Z \times S}(\mathfrak{D}\mathbf{b}_{X \times S}) &\simeq f_*(\mathcal{D}_{Y \leftarrow X/S} \otimes_{\mathcal{D}_{X \times S/S}} \Gamma_{Z \times S}(\mathfrak{D}\mathbf{b}_{X \times S})) \\ &\simeq \Gamma_{Z \times S}(\mathfrak{D}\mathbf{b}_{Y \times S}) \simeq T\mathcal{H}om(f_*\mathbb{C}_{Z \times X}, \mathfrak{D}\mathbf{b}_{Y \times S}) \simeq \mathrm{TH}_Y^S(f_*(\mathbb{C}_{Z \times S} \otimes p^{-1}\mathcal{O}_S)). \end{aligned}$$

Step 2. Let us now assume that f is smooth and $f \times \text{Id}_S$ is proper on the support of F . Let $W = \text{Supp}_X F$. Then W is closed subanalytic in X and the assumption entails that $f|_W$ is proper. The question being local we may assume as in the proof of [16, Th. 3.5(ii)] that $f : X = Y \times \mathbb{R} \rightarrow \mathbb{R}$ is the projection and, in view of Proposition 5.4, that $F \simeq \mathbb{C}_{\Omega \times S} \otimes p^{-1}\mathcal{O}_S$ for some subanalytic subset Ω contained in W .

As in Step 1, we prove first the statement for $F \simeq \mathbb{C}_{Z \times S} \otimes p^{-1}\mathcal{O}_S$ where Z is closed subanalytic contained in W . Furthermore, (3.9) of Lemma 3.6 in loc. cit. entails that we may assume that for any $x \in Y$, $Z \cap f^{-1}(x)$ is a closed interval in \mathbb{R} containing 0. With this assumption we have

$$T\mathcal{H}om(Rf_*\mathbb{C}_{Z \times S}, \mathfrak{D}\mathfrak{b}_{Y \times S}) \simeq T\mathcal{H}om(\mathbb{C}_{f(Z) \times S}, \mathfrak{D}\mathfrak{b}_{Y \times S}) \simeq \Gamma_{f(Z) \times S}(\mathfrak{D}\mathfrak{b}_{Y \times S}).$$

Noting that $\mathcal{D}_{Y \leftarrow X/S}$ is $\mathcal{D}_{Y \times S/S}$ -isomorphic to $\mathcal{D}_{Y \times S/S}/\mathcal{D}_{Y \times S/S}\partial_t$, where t is the coordinate in \mathbb{R} , the result follows by the exact sequence

$$0 \longrightarrow f_*\Gamma_{Z \times S}(\mathfrak{D}\mathfrak{b}_{X \times S}) \xrightarrow{\partial_t} f_*\Gamma_{Z \times S}(\mathfrak{D}\mathfrak{b}_{X \times S}) \xrightarrow{\int \cdot dt} \Gamma_{f(Z) \times S}(\mathfrak{D}\mathfrak{b}_{Y \times S}) \longrightarrow 0$$

proved in [12, Lem. 4.5]. The isomorphism for Ω follows by functoriality considering the exact sequence

$$0 \longrightarrow \mathbb{C}_{\Omega \times S} \longrightarrow \mathbb{C}_{\overline{\Omega} \times S} \longrightarrow \mathbb{C}_{\delta(\Omega) \times S} \longrightarrow 0. \quad \text{q.e.d.}$$

Let us now finish the proof of Theorem 5.10. Applying $R\mathcal{H}om_{\mathcal{D}_{\overline{Y \times S}/\overline{S}}}(\mathcal{O}_{\overline{Y \times S}}, \bullet)[d_Y]$ to both terms of (5.12) the right term becomes $\text{RH}_Y^S(Rf_*F)$. For the left side of (5.12) we obtain

$$\begin{aligned} & R\mathcal{H}om_{\mathcal{D}_{\overline{Y \times S}/\overline{S}}}(\mathcal{O}_{\overline{Y \times S}}, {}_{\mathcal{D}}f_! \text{TH}_X^S(F))[d_Y] \\ & \simeq Rf_*(R\mathcal{H}om_{\mathcal{D}_{\overline{Y \times S}/\overline{S}}}(\mathcal{O}_{\overline{Y \times S}}, \mathcal{D}_{Y_{\mathbb{R}} \leftarrow X_{\mathbb{R}}/S_{\mathbb{R}}} \otimes_{\mathcal{D}_{Y_{\mathbb{R}} \times S_{\mathbb{R}}/S_{\mathbb{R}}}}^L \text{TH}_X^S(F)))[d_Y] \\ & \simeq Rf_*(R\mathcal{H}om_{\mathcal{D}_{\overline{Y \times S}/\overline{S}}}(\mathcal{O}_{\overline{Y \times S}}, \mathcal{D}_{Y_{\mathbb{R}} \leftarrow X_{\mathbb{R}}/S_{\mathbb{R}}}) \otimes_{\mathcal{D}_{Y_{\mathbb{R}} \times S_{\mathbb{R}}/S_{\mathbb{R}}}}^L \text{TH}_X^S(F))[d_Y]. \end{aligned}$$

According to the relative version of [12, Lem. 7.2] we have

$$\begin{aligned} & R\mathcal{H}om_{\mathcal{D}_{\overline{Y \times S}/\overline{S}}}(\mathcal{O}_{\overline{Y \times S}}, \mathcal{D}_{Y_{\mathbb{R}} \leftarrow X_{\mathbb{R}}/S_{\mathbb{R}}}) \\ & \simeq \mathcal{D}_{Y \leftarrow X/S} \otimes_{\mathcal{D}_{X \times S/S}}^L R\mathcal{H}om_{\mathcal{D}_{\overline{X \times S}/\overline{S}}}(\mathcal{O}_{\overline{Y \times S}}, \mathcal{D}_{Y_{\mathbb{R}} \times S_{\mathbb{R}}/S_{\mathbb{R}}})[d_X - d_Y]. \end{aligned}$$

Therefore, by applying $R\mathcal{H}om_{\mathcal{D}_{\overline{Y \times S}/\overline{S}}}(\mathcal{O}_{\overline{Y \times S}}, \bullet)[d_Y]$ to the left-hand side of (5.12) we obtain an isomorphism with ${}_{\mathcal{D}}f_! \text{RH}_X^S(F)$ which concludes the construction of the morphism (5.10*). Lemma 5.13 shows that it is an isomorphism. q.e.d.

5.c. Riemann-Hilbert correspondence for Deligne's extensions. We consider the setting of Notation 4.5 and assume that $Y = D$ has normal crossings in X .

Proposition 5.14. *Let L be a coherent S -local system on $X^* \times S$ and let \tilde{E}_L be the associated Deligne extension. Then*

- the complex of $\mathcal{D}_{X \times S/S}$ -modules $\text{RH}_X^S(j_! D' L)[-d_X]$ is isomorphic to \tilde{E}_L and thus it is regular holonomic,
- ${}^{\text{p}}\text{Sol } \tilde{E}_L \simeq j_! D' L$.

Proof. We adapt the idea of proof of [31, Lem. 4.2]. Let us prove the second statement assuming the first one holds true. Firstly, the following lemma is similar to [31, Lem. 3.19]:

Lemma 5.15. *There exists a functorial isomorphism in $D_{\mathbb{R}\text{-c}}^b(p^{-1}\mathcal{O}_S)$:*

$${}^p\text{DR}_X(\text{RH}_X^S(F)) \xrightarrow{\sim} \mathbf{D}F.$$

Since $\text{RH}_X^S(j_! \mathbf{D}'L)[-d_X]$ is holonomic by the first point, we have

$${}^p\text{Sol}_X \text{RH}_X^S(j_! \mathbf{D}'L)[-d_X] \simeq \mathbf{D} {}^p\text{DR} \text{RH}_X^S(j_! \mathbf{D}'L)[-d_X] \stackrel{(*)}{\simeq} \mathbf{D}\mathbf{D}(j_! \mathbf{D}'L) = j_! \mathbf{D}'L,$$

where $(*)$ follows from Lemma 5.15.

Let us now prove the first statement. We will set

$$\tilde{E}_L^S = \text{RH}_X^S(j_! \mathbf{D}'L)[-d_X].$$

Then, according to (5.8**) and to [31, Lem. 3.22], we have

$$(5.16) \quad \begin{aligned} \tilde{E}_L^S &= \rho_S^{-1} Rj_* R\mathcal{H}om_{\rho_{S*} p_{X^*}^{-1} \mathcal{O}_S}(\rho_{S*} \mathbf{D}'L, j^{-1} \mathcal{O}_{X \times S}^{t,S}) \\ &\simeq \rho_S^{-1} Rj_*(\rho_{S*} L \otimes_{\rho_{S*} p_{X^*}^{-1} \mathcal{O}_S}^L j^{-1} \mathcal{O}_{X \times S}^{t,S}). \end{aligned}$$

We make use of the following result of [28], the proof of which we recall with details as it is used in an essential way below.

Lemma 5.17 ([28, Prop. 3.32]). *The complex \tilde{E}_L^S is concentrated in degree zero.*

Proof. Assume first that L is the constant local system $p_{X^*}^{-1}G$ with G being \mathcal{O}_S -coherent. Since the question is local on $X \times S$, we can assume that G admits a finite resolution $\mathcal{O}_S^\bullet \rightarrow G$ by free \mathcal{O}_S -modules of finite rank. Then we have

$$(5.18) \quad \begin{aligned} \rho_S^{-1} Rj_*(\rho_{S*} p_{X^*}^{-1} G \otimes_{p_{X^*}^{-1} \mathcal{O}_S}^L j^{-1} \mathcal{O}_{X \times S}^{t,S}) \\ \xleftarrow{\sim} \rho_S^{-1} Rj_*(\rho_{S*} p_{X^*}^{-1} \mathcal{O}_S^\bullet \otimes_{p_{X^*}^{-1} \mathcal{O}_S} j^{-1} \mathcal{O}_{X \times S}^{t,S}) \\ \simeq p_X^{-1} \mathcal{O}_S^\bullet \otimes_{p_X^{-1} \mathcal{O}_S} \rho_S^{-1} Rj_* j^{-1} \mathcal{O}_{X \times S}^{t,S}. \end{aligned}$$

We recall that, according to [17, Prop. 2.4.4 (2.4.4)], we have isomorphisms of functors on $D^b(\mathbb{C}_{X_{\text{sa}} \times S})$

$$R\mathcal{H}om(\rho_{S*} j_! \mathbb{C}_{X^* \times S}, (\bullet)) \xrightarrow{\sim} Rj_* R\mathcal{H}om(\rho_{S*} \mathbb{C}_{X^* \times S}, (\bullet)) \simeq Rj_* j^{-1}(\bullet).$$

As a consequence, we obtain an isomorphism

$$\rho_S^{-1} Rj_* j^{-1} \mathcal{O}_{X \times S}^{t,S} \simeq \rho_S^{-1} R\mathcal{H}om(\mathbb{C}_{X^* \times S}, \mathcal{O}_{X \times S}^{t,S}).$$

It follows then from [28, Prop. 3.24(1)] applied to $\mathbb{C}_{X^* \times S} = \mathbb{C}_{X^*} \boxtimes \mathbb{C}_S$, that the latter complex is isomorphic to $T\mathcal{H}om(\mathbb{C}_{X^* \times S}, \mathcal{O}_{X \times S})$, so that, according to [12, Lem. 7.5]

$$(5.19) \quad \rho_S^{-1} Rj_* j^{-1} \mathcal{O}_{X \times S}^{t,S} \simeq \mathcal{O}_{X \times S}(*D).$$

We thus deduce from (5.18) and (5.19):

$$(5.20) \quad \begin{aligned} \rho_S^{-1} Rj_*(\rho_{S*} p_{X^*}^{-1} G \otimes_{p_{X^*}^{-1} \mathcal{O}_S}^L j^{-1} \mathcal{O}_{X \times S}^{t,S}) \\ \simeq \rho_S^{-1} Rj_*(\rho_{S*} p_{X^*}^{-1} \mathcal{O}_S^\bullet \otimes_{p_{X^*}^{-1} \mathcal{O}_S} j^{-1} \mathcal{O}_{X \times S}^{t,S}) \\ \simeq p_X^{-1} \mathcal{O}_S^\bullet \otimes_{p_X^{-1} \mathcal{O}_S} \rho_S^{-1} Rj_* j^{-1} \mathcal{O}_{X \times S}^{t,S} \\ \simeq p_X^{-1} \mathcal{O}_S^\bullet \otimes_{p_X^{-1} \mathcal{O}_S} \mathcal{O}_{X \times S}(*D) \\ \simeq p_X^{-1} G \otimes_{p_X^{-1} \mathcal{O}_S} \mathcal{O}_{X \times S}(*D), \end{aligned}$$

where the latter isomorphism follows from the $p_X^{-1} \mathcal{O}_S$ -flatness of $\mathcal{O}_{X \times S}(*D)$. We conclude that $\tilde{E}_{p_X^{-1} G}^S$ is concentrated in degree zero.

Let us now consider the case of a possibly non constant S -local system L locally isomorphic to $p^{-1}G$, with G being \mathcal{O}_S -coherent. According to Remark 4.18, locally on $X \times S$ we have an isomorphism of $\mathcal{O}_{X^* \times S}$ -modules

$$(5.21) \quad L \otimes_{p_X^{-1} \mathcal{O}_S} \mathcal{O}_{X^* \times S} \simeq p_{X^*}^{-1} G \otimes_{p_{X^*}^{-1} \mathcal{O}_S} \mathcal{O}_{X^* \times S},$$

and by flatness of $\mathcal{O}_{X^* \times S}$ over $p_{X^*}^{-1} \mathcal{O}_S$, the same equality holds with the derived tensor product \otimes^L . According to [28, Lem. 3.15], the natural morphism $\rho_{S^*} L \rightarrow R\rho_{S^*} L$ is an isomorphism and [28, Prop. 3.16] implies that $\rho_{S^!} L \simeq R\rho_{S^*} L$. Recall also that, by [28, Prop. 3.13], the functor $\rho_{S^!}$ is exact and commutes with \otimes . From (5.21) we then derive $\mathcal{O}_{X^* \times S}$ -linear isomorphisms (they are a priori not $\mathcal{D}_{X^* \times S/S}$ -linear since we use a tensor product over $\mathcal{O}_{X^* \times S}$):

$$\begin{aligned} \rho_{S^*} L \otimes_{\rho_{S^*} p_{X^*}^{-1} \mathcal{O}_S}^L j^{-1} \mathcal{O}_{X^* \times S}^{t,S} &\simeq (\rho_{S^!} L \otimes_{\rho_{S^!} p_{X^*}^{-1} \mathcal{O}_S}^L \rho_{S^!} \mathcal{O}_{X^* \times S}) \otimes_{\rho_{S^!} \mathcal{O}_{X^* \times S}}^L j^{-1} \mathcal{O}_{X^* \times S}^{t,S} \\ &\simeq (\rho_{S^!} (L \otimes_{p_{X^*}^{-1} \mathcal{O}_S}^L \mathcal{O}_{X^* \times S})) \otimes_{\rho_{S^!} \mathcal{O}_{X^* \times S}}^L j^{-1} \mathcal{O}_{X^* \times S}^{t,S} \\ &\simeq (\rho_{S^!} (p_{X^*}^{-1} G \otimes_{p_{X^*}^{-1} \mathcal{O}_S}^L \mathcal{O}_{X^* \times S})) \otimes_{\rho_{S^!} \mathcal{O}_{X^* \times S}}^L j^{-1} \mathcal{O}_{X^* \times S}^{t,S} \\ &\simeq (\rho_{S^!} p_{X^*}^{-1} G \otimes_{\rho_{S^!} p_{X^*}^{-1} \mathcal{O}_S}^L \rho_{S^!} \mathcal{O}_{X^* \times S}) \otimes_{\rho_{S^!} \mathcal{O}_{X^* \times S}}^L j^{-1} \mathcal{O}_{X^* \times S}^{t,S} \\ &\simeq \rho_{S^*} p_X^{-1} G \otimes_{\rho_{S^*} p_{X^*}^{-1} \mathcal{O}_S}^L j^{-1} \mathcal{O}_{X^* \times S}^{t,S}. \end{aligned}$$

Then, due to (5.16), we deduce an $\mathcal{O}_{X \times S}$ -linear isomorphism

$$(5.22) \quad \tilde{E}_L^S \simeq \tilde{E}_{p_{X^*}^{-1} G}^S \simeq p_X^{-1} G \otimes_{p_X^{-1} \mathcal{O}_S} \mathcal{O}_{X \times S}(*D).$$

By the first part of the proof, we deduce that \tilde{E}_L^S is concentrated in degree zero. This concludes the proof of Proposition 5.17. q.e.d.

Lemma 5.23. *Both \tilde{E}_L and \tilde{E}_L^S are naturally $\mathcal{D}_{X \times S/S}$ -submodules of $j_* E_L$.*

Proof. Firstly, applying the commutation of ρ_S^{-1} with j^{-1} together with the analogue of [31, Cor. 3.24] entails that \tilde{E}_L^S and \tilde{E}_L coincide with E_L when restricted to $X^* \times S$.

On the one hand, by construction, \tilde{E}_L is naturally a $\mathcal{D}_{X \times S/S}$ -submodule of $j_* E_L$. Let us check on the other hand that \tilde{E}_L^S is also naturally a $\mathcal{D}_{X \times S/S}$ -submodule of $j_* E_L$. From the natural morphism of functors $\text{Id} \rightarrow R\rho_{S^*} \rho_S^{-1}$ on $\text{Mod}(\mathbb{C}_{X_{\text{sa}} \times S})$ we derive a natural morphism $\rho_S^{-1} Rj_* \rightarrow Rj_* \rho_S^{-1}$: denoting for a moment by j_*^S the morphism in the subanalytic site, we have an isomorphism of functors $Rj_*^S \circ R\rho_{S^*} \xrightarrow{\sim} R\rho_{S^*} \circ Rj_*$ (cf. [33, Prop. 2.2.1(i)], by replacing there ρ by ρ_S and using the same argument), hence a morphism

$$Rj_*^S \longrightarrow Rj_*^S \circ R\rho_{S^*} \rho_S^{-1} \xrightarrow{\sim} R\rho_{S^*} Rj_*^S \rho_S^{-1},$$

and, by applying ρ_S^{-1} on the left, we obtain the desired morphism. Recall also, as already used, that $\rho_{S*}L \xrightarrow{\sim} R\rho_{S*}L$. We then deduce a $\mathcal{D}_{X \times S/S}$ -linear morphism

$$\begin{aligned} \Psi : \widetilde{E}_L^S &\simeq \rho_S^{-1} Rj_*(\rho_{S*}L \otimes_{\rho_{S*}p_{X^*}^{-1}\mathcal{O}_S}^L j^{-1}\mathcal{O}_{X \times S}^{t,S}) \\ &\longrightarrow Rj_*\rho_S^{-1}(\rho_{S*}L \otimes_{\rho_{S*}p_{X^*}^{-1}\mathcal{O}_S}^L j^{-1}\mathcal{O}_{X \times S}^{t,S}) \\ &\stackrel{(a)}{\simeq} Rj_*(L \otimes_{p_{X^*}^{-1}\mathcal{O}_S}^L \rho_S^{-1}j^{-1}\mathcal{O}_{X \times S}^{t,S}) \\ &\simeq Rj_*(L \otimes_{p_{X^*}^{-1}\mathcal{O}_S}^L \mathcal{O}_{X^* \times S}) \\ &\stackrel{(b)}{\simeq} Rj_*(L \otimes_{p_{X^*}^{-1}\mathcal{O}_S} \mathcal{O}_{X^* \times S}) = Rj_*E_L \stackrel{(c)}{\simeq} j_*E_L, \end{aligned}$$

where (a) follows from (5.2), (b) follows from the $p_{X^*}^{-1}\mathcal{O}_S$ -flatness of $\mathcal{O}_{X^* \times S}$ and (c) from the Steinness of j .

In order to check that Ψ is injective, it is enough to consider it as an $\mathcal{O}_{X \times S}$ -linear morphism, and by (5.22), it is enough to check injectivity when $L = p_{X^*}^{-1}G$. Furthermore, the question is local on S .

In such a case, we consider a resolution $\mathcal{O}_S^\bullet \rightarrow G$ as in the proof of Proposition 5.17. Then, locally on S , (5.20) identifies $\widetilde{E}_{p_{X^*}^{-1}G}^S$ with $p_X^{-1}G \otimes_{p_X^{-1}\mathcal{O}_S} \mathcal{O}_{X \times S}(*D)$ and the morphism Ψ with the natural morphism

$$p_X^{-1}G \otimes_{p_X^{-1}\mathcal{O}_S} \mathcal{O}_{X \times S}(*D) \longrightarrow j_*(p_{X^*}^{-1}G \otimes_{p_{X^*}^{-1}\mathcal{O}_S} \mathcal{O}_{X^* \times S}),$$

which is injective since G is \mathcal{O}_S -coherent, as wanted.

q.e.d.

End of the proof of Proposition 5.14. By Lemma 5.23, we are reduced to showing that both $\mathcal{D}_{X \times S/S}$ -submodules \widetilde{E}_L^S and \widetilde{E}_L of j_*E_L coincide, and since the $\mathcal{D}_{X \times S/S}$ -structure is induced by that j_*E_L for both, it is enough to check that they coincide as $\mathcal{O}_{X \times S}$ -submodules. The question is local on $X \times S$. By Remark 4.18, there exists a local $\mathcal{O}_{X \times S}$ -linear isomorphism $j_*E_L \simeq j_*\widehat{E}_{p_{X^*}^{-1}G}$ under which \widetilde{E}_L is identified with $\widehat{E}_G = p_X^{-1}G \otimes_{p_X^{-1}\mathcal{O}_S} \mathcal{O}_{X \times S}(*D)$. Besides, (5.22) also identifies \widetilde{E}_L^S with $p_X^{-1}G \otimes_{p_X^{-1}\mathcal{O}_S} \mathcal{O}_{X \times S}(*D)$ as an $\mathcal{O}_{X \times S}$ -module under the same local isomorphism, so the $\mathcal{O}_{X \times S}$ -submodules \widetilde{E}_L and \widetilde{E}_L^S of j_*E_L locally coincide, as desired.

q.e.d.

6. Proof of Theorem 2

The proof of Theorem 2 is similar to that of [8, Th. 1] where $\dim S = 1$. However, various improvements are necessary in order to handle the case $\dim S \geq 2$.

6.a. First part of the proof of Theorem 2. We prove that RH_X^S is a right quasi-inverse of the functor ${}^p\mathrm{Sol}_X : \mathrm{D}_{\mathrm{rhol}}^b(\mathcal{D}_{X \times S/S}) \rightarrow \mathrm{D}_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$ by exhibiting an isomorphism of functors $\alpha : \mathrm{Id}_{\mathrm{D}_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)} \xrightarrow{\sim} {}^p\mathrm{Sol}_X \circ \mathrm{RH}_X^S$. This is the analogue in possibly higher dimension for S of [31, Th. 3]:

(6.1) For F in $\mathrm{D}_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$, $\mathrm{RH}_X^S(F)$ is an object of $\mathrm{D}_{\mathrm{rhol}}^b(\mathcal{D}_{X \times S/S})$ and there exists a functorial isomorphism $\alpha_F : F \simeq {}^p\mathrm{Sol}_X(\mathrm{RH}_X^S(F))$ in $\mathrm{D}_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$.

Proof of (6.1). We recall the notation for the duality functor (cf. [29, Prop. 2.23]): for F in $\mathrm{D}_{\mathbb{R}\text{-c}}^b(p^{-1}\mathcal{O}_S)$, we set $\mathbf{D}'F = R\mathcal{H}om_{p^{-1}\mathcal{O}_S}(F, p^{-1}\mathcal{O}_S) \in \mathrm{D}_{\mathbb{R}\text{-c}}^b(p^{-1}\mathcal{O}_S)$ and $\mathbf{D}F = \mathbf{D}'F[2d_X]$. Functoriality in (6.1) is obtained by means of Lemma 5.15: once

we know that, for F in $D_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$, $\mathrm{RH}_X^S(F)$ belongs to $D_{\mathrm{rhol}}^b(\mathcal{D}_{X \times S/S})$, we can apply [29, Cor. 3.9] together with bi-duality in $D_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$ (cf. [29, Prop. 2.23]) to obtain a functorial isomorphism

$$\alpha_F : F \xrightarrow{\sim} {}^p\mathrm{Sol}_X(\mathrm{RH}_X^S(F)).$$

For the first part of (6.1), the main step is provided by Proposition 5.14, where we proved the case $F = j_! \mathbf{D}' L$ for an S -coherent local system L in the setting of Section 4.b (from which we keep the notation), that is, D is a normal crossing divisor in X and L is an S -local system on $X^* \times S := (X \setminus D) \times S$.

We now conclude the proof of (6.1) by a standard induction on the dimension of the X -support of F , based on Theorem 5.10, analogous to that of [12, §7.3] (cf. [31, Th. 3] for the case $\dim S = 1$).

We thus assume that (6.1) holds if $\dim \mathrm{Supp}_X F < k$ ($k \geq 1$) and we prove that it holds for any F with $\dim \mathrm{Supp}_X F \leq k$. By functoriality, it is enough to prove the first part of (6.1), so that the question is local. By induction on the amplitude of the complex F , we can also assume that F is an S - \mathbb{C} -constructible sheaf. We can find a projective morphism $f : X' \rightarrow X$ with $\dim X' = k$, which is biholomorphic from the complement X'^* of a normal crossing divisor D' in X' to the smooth locus of dimension k of $\mathrm{Supp}_X F$, so that $f^{-1}F$ is a coherent S -local system on X'^* . Let $j' : X'^* \hookrightarrow X'$ denote the inclusion. Lemma 5.14 implies the regular holonomicity of $\mathrm{RH}_X^S(j'_! \mathbf{D}' j'^{-1} f^{-1} F)$ and Theorem 5.10, together with [31, Cor. 2.4], implies regular holonomicity of

$$\mathrm{RH}_X^S(Rf_* j'_! \mathbf{D}' j'^{-1} f^{-1} F) \simeq \mathrm{RH}_X^S(j_! \mathbf{D}' j^{-1} F),$$

where j denotes the inclusion of $f(X'^*)$ in X . Hence, if F is an S - \mathbb{C} -constructible sheaf with X -support of dimension $\leq k$, $\mathrm{RH}_X^S(\mathbf{D}' F)$ is regular holonomic since it fits in a distinguished triangle whose third term is regular holonomic by the induction hypothesis. The same holds for $\mathbf{D}' F$ for any $F \in D_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$ with $\dim \mathrm{Supp}_X F \leq k$, and replacing F with $\mathbf{D}' F$, which has the same X -support, we conclude that $\mathrm{RH}_X^S(F)$ is regular holonomic. q.e.d.

6.b. Second part of the proof of Theorem 2. We prove that RH_X^S is a left quasi-inverse of the functor ${}^p\mathrm{Sol}_X : D_{\mathrm{rhol}}^b(\mathcal{D}_{X \times S/S}) \rightarrow D_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$ by exhibiting an isomorphism of functors $\beta : \mathrm{Id}_{D_{\mathrm{rhol}}^b(\mathcal{D}_{X \times S/S})} \xrightarrow{\sim} \mathrm{RH}_X^S \circ {}^p\mathrm{Sol}_X$.

(6.2) For each object \mathcal{M} of $D_{\mathrm{rhol}}^b(\mathcal{D}_{X \times S/S})$, there exists an isomorphism, functorial in \mathcal{M} , $\beta_{\mathcal{M}} : \mathcal{M} \xrightarrow{\sim} \mathrm{RH}_X^S({}^p\mathrm{Sol}_X(\mathcal{M}))$.

Proof of (6.2). The argument is similar to that given in [8, §3.b]. It is enough to construct a bi-functorial isomorphism (with respect to $\mathcal{M}, \mathcal{N} \in D_{\mathrm{rhol}}^b(\mathcal{D}_{X \times S/S})$)

$$(6.3) \quad \mathrm{Hom}_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathrm{RH}_X^S({}^p\mathrm{Sol}_X(\mathcal{N}))) \xrightarrow{\sim} \mathrm{Hom}_{p^{-1}\mathcal{O}_S}({}^p\mathrm{Sol}_X(\mathcal{N}), {}^p\mathrm{Sol}_X(\mathcal{M})),$$

that we also denote as $(6.3)_{\mathcal{M}, \mathcal{N}}$. We then define the morphism

$$\beta_{\mathcal{M}} : \mathcal{M} \longrightarrow \mathrm{RH}_X^S({}^p\mathrm{Sol}_X(\mathcal{M}))$$

as the unique morphism such that $(6.3)_{\mathcal{M}, \mathcal{M}}(\beta_{\mathcal{M}}) = \mathrm{Id}_{{}^p\mathrm{Sol}_X(\mathcal{M})}$. It is then clearly an isomorphism. In order to check that $\beta_{\mathcal{M}}$ is functorial with respect to \mathcal{M} , that is, it

defines an isomorphism of functors $\mathrm{Id}_{\mathcal{D}_{\mathrm{rhol}}^b(\mathcal{D}_{X \times S/S})} \xrightarrow{\sim} \mathrm{RH}_X^S \circ \mathrm{PSol}_X$, we consider as in [31, p. 668], for a morphism $\varphi : \mathcal{M} \rightarrow \mathcal{N}$, the commutative diagram

$$\begin{array}{ccc}
\mathrm{Hom}_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathrm{RH}_X^S(\mathrm{PSol}_X(\mathcal{M}))) & \xrightarrow{(6.3)_{\mathcal{M}, \mathcal{M}}} & \mathrm{Hom}_{p^{-1}\mathcal{O}_S}(\mathrm{PSol}_X(\mathcal{M}), \mathrm{PSol}(\mathcal{M})) \\
\mathrm{RH}^S(\mathrm{PSol}_X(\varphi)) \circ \bullet \downarrow & & \downarrow \bullet \circ \mathrm{PSol}_X(\varphi) \\
\mathrm{Hom}_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathrm{RH}_X^S(\mathrm{PSol}_X(\mathcal{N}))) & \xrightarrow{(6.3)_{\mathcal{M}, \mathcal{N}}} & \mathrm{Hom}_{p^{-1}\mathcal{O}_S}(\mathrm{PSol}_X(\mathcal{N}), \mathrm{PSol}_X(\mathcal{M})) \\
\bullet \circ \varphi \uparrow & & \mathrm{PSol}_X(\varphi) \circ \bullet \uparrow \\
\mathrm{Hom}_{\mathcal{D}_{X \times S/S}}(\mathcal{N}, \mathrm{RH}_X^S(\mathrm{PSol}_X(\mathcal{N}))) & \xrightarrow{(6.3)_{\mathcal{N}, \mathcal{N}}} & \mathrm{Hom}_{p^{-1}\mathcal{O}_S}(\mathrm{PSol}_X(\mathcal{N}), \mathrm{PSol}_X(\mathcal{N}))
\end{array}$$

Since $\mathrm{Id}_{\mathrm{PSol}_X(\mathcal{M})} \circ \mathrm{PSol}_X(\varphi) = \mathrm{PSol}_X(\varphi) \circ \mathrm{Id}_{\mathrm{PSol}_X(\mathcal{N})}$, we obtain from the commutativity of the diagram the

$$\mathrm{RH}_X^S(\mathrm{PSol}_X(\varphi)) \circ \beta_{\mathcal{M}} = \beta_{\mathcal{N}} \circ \varphi,$$

which is the desired functoriality.

In order to obtain (6.3), it is enough to construct a bi-functorial morphism in $\mathcal{D}^b(p^{-1}\mathcal{O}_S)$

$$(6.4) \quad R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathrm{RH}_X^S(F)) \longrightarrow R\mathcal{H}om_{p^{-1}\mathcal{O}_S}(F, \mathrm{PSol}_X(\mathcal{M}))$$

for $\mathcal{M} \in \mathcal{D}_{\mathrm{rhol}}^b(\mathcal{D}_{X \times S/S})$ and $F \in \mathcal{D}_{\mathrm{C-c}}^b(p^{-1}\mathcal{O}_S)$, and to show that it is an isomorphism. Then (6.3) $_{\mathcal{M}, \mathcal{N}}$ is obtained by taking global sections of $\mathcal{H}^0(6.4)$ applied to $F = \mathrm{PSol}(\mathcal{N})$.

Such a morphism (6.4) is given by [8, (14)]. In loc. cit., it is also shown that, for each $s_o \in S$, $Li_{s_o}^*(6.4)$ can be identified with the morphism constructed in the absolute case by Kashiwara ([12, Cor. 8.6]), hence it is an isomorphism. To conclude that (6.4) is an isomorphism, we apply a Nakayama-type argument [29, Prop. 2.2]. For that we need to check a finiteness property. The proof of (6.2) will thus be concluded with the proof of the following assertion.

(6.5) *For any $\mathcal{M} \in \mathcal{D}_{\mathrm{rhol}}^b(\mathcal{D}_{X \times S/S})$ and for any $F \in \mathcal{D}_{\mathrm{C-c}}^b(p^{-1}\mathcal{O}_S)$, the complex $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathrm{RH}_X^S(F))$ belongs to $\mathcal{D}_{\mathrm{C-c}}^b(p^{-1}\mathcal{O}_S)$.*

We also consider the statement:

(6.6) *For any $\mathcal{M}, \mathcal{N} \in \mathcal{D}_{\mathrm{rhol}}^b(\mathcal{D}_{X \times S/S})$, the complex $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{N})$ belongs to $\mathcal{D}_{\mathrm{C-c}}^b(p^{-1}\mathcal{O}_S)$.*

We shall argue by induction on pair $(\dim S, \dim \mathrm{Supp}_X \mathcal{M})$ ordered lexicographically. For that purpose we introduce the following notations:

- For $d \geq 0$, we denote by (6.5) $_d$, resp. (6.6) $_d$ the corresponding statement concerning any S satisfying $\dim S \leq d$.
- For $k \geq 0$ we introduce the assertion (6.6) $_{d,k}$ by requiring that the property holds for $\dim S \leq d-1$ or $\dim S = d$ and $\dim \mathrm{Supp}_X \mathcal{M} \leq k$.

Because of the first part of (6.1) already proved, we have

$$(6.6)_d \implies (6.5)_d.$$

Conversely, the reverse implication also holds:

$$(6.5)_d \implies (6.6)_d.$$

Indeed we deduce from (6.5)_d an isomorphism $\beta_{\mathcal{N}} : \mathcal{N} \xrightarrow{\sim} \mathrm{RH}_X^S(\mathrm{pSol}_X(\mathcal{N}))$, so applying (6.5)_d with $F = \mathrm{pSol}_X(\mathcal{N})$ gives (6.6)_d. In the following, we will focus on (6.6)_d. Since belonging to $\mathrm{D}_{\mathrm{C-c}}^b(p^{-1}\mathcal{O}_S)$ is a local property (cf. [29, §2.5]), we will allow restriction to an open neighbourhood of a point of $X \times S$ when needed.

Proof of (6.6)_d. The proof of (6.6)_d is done by induction on d , in order to reduce the assertion to the particular case of Lemma 6.15 below. Let us recall that (6.6)₀ holds, according to [12]. So we assume that (6.6)_{d-1} holds ($d \geq 1$) and we consider S with $\dim S = d$, together with any $\mathcal{M}, \mathcal{N} \in \mathrm{D}_{\mathrm{rhol}}^b(\mathcal{D}_{X \times S/S})$.

Step 1: The S -torsion case

Lemma 6.7. *Let $\dim S = d$ and let us assume that (6.6)_{d-1} holds. Let $\mathcal{M}, \mathcal{N} \in \mathrm{D}_{\mathrm{rhol}}^b(\mathcal{D}_{X \times S/S})$ and assume that the cohomology of \mathcal{M} or that of \mathcal{N} is of S -torsion. Then (6.6)_d holds for \mathcal{M}, \mathcal{N} , that is,*

$$R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{N}) \in \mathrm{D}_{\mathrm{C-c}}^b(p^{-1}\mathcal{O}_S).$$

Proof. By the biduality isomorphism of [29, (3)], we have

$$(6.8) \quad R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{N}) \simeq R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathbf{D}\mathcal{N}, \mathbf{D}\mathcal{M}),$$

and we recall that both $\mathbf{D}\mathcal{N}$ and $\mathbf{D}\mathcal{M}$ are regular holonomic according to Corollary 3.8. It is thus enough to prove Lemma 6.7 when the cohomology of \mathcal{N} is of S -torsion. On the other hand, owing to the definition of $\mathrm{D}_{\mathrm{C-c}}^b(p^{-1}\mathcal{O}_S)$ in terms of micro-support (cf. [29, §2.5]), if two terms of a distinguished triangle in $\mathrm{D}^b(p^{-1}\mathcal{O}_S)$ are objects of $\mathrm{D}_{\mathrm{C-c}}^b(p^{-1}\mathcal{O}_S)$, then so does the third one. This property allows us to assume that \mathcal{M}, \mathcal{N} are regular holonomic $\mathcal{D}_{X \times S/S}$ -modules instead of complexes. Lastly, the same argument implies that belonging to $\mathrm{D}_{\mathrm{C-c}}^b(p^{-1}\mathcal{O}_S)$ is a local question on $X \times S$.

The assumption on \mathcal{N} entails that $\mathrm{Supp}_S \mathcal{N}$ is contained in a closed analytic subset T of S such that $\dim T < \dim S$. We will argue by induction on $\dim T$.

If $\dim T = 0$, as we consider a local situation, we may assume $T = \{s_o\}$ with maximal ideal sheaf \mathfrak{m}_{s_o} . By considering the (locally) finite filtration of \mathcal{N} by the $\mathcal{D}_{X \times S/S}$ -regular holonomic submodules $\mathfrak{m}_{s_o}^k \mathcal{N}$ ($k \in \mathbb{N}$), we may reduce to assume $\mathfrak{m}_{s_o} \mathcal{N} = 0$. We then have $\mathcal{N} \simeq i_{s_o*} i_{s_o}^* \mathcal{N}$ and by adjunction

$$R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, i_{s_o*} i_{s_o}^* \mathcal{N}) \simeq Ri_{s_o*} R\mathcal{H}om_{\mathcal{D}_X}(Li_{s_o}^* \mathcal{M}, i_{s_o}^* \mathcal{N}),$$

so that the proof is reduced to applying (6.6)₀ or, equivalently, the absolute case proved by Kashiwara in [12].

Assume now that $\dim T \geq 1$ and let $T_0 \subset T$ be a closed analytic subset of dimension $< \dim T$ such that $T \setminus T_0$ is non singular of dimension $\dim T$. Let $\pi : S' \rightarrow S$ be a projective morphism of complex manifolds satisfying the two conditions: $\pi(S') = T$ and, setting $T'_0 := \pi^{-1}(T_0)$, $\pi' := \pi|_{S' \setminus T'_0} : S' \setminus T'_0 \rightarrow T \setminus T_0$ is a biholomorphism. In particular $\dim S' = \dim T < d$.

Note that, according to Proposition 3.9, $L\pi^* \mathcal{M}$, $L\pi^* \mathcal{N}$ are regular holonomic on $X \times S'$ and, by Theorem 3.14, $R\pi_* L\pi^* \mathcal{N}$ belongs to $\mathrm{D}_{\mathrm{rhol}}^b(\mathcal{D}_{X \times S/S})$. By (6.6)_{d-1}, $R\mathcal{H}om_{\mathcal{D}_{X \times S'/S'}}(L\pi^* \mathcal{M}, L\pi^* \mathcal{N}) \in \mathrm{D}_{\mathrm{C-c}}^b(p'^{-1}\mathcal{O}_{S'})$ thus, by the properness of π and Lemma 2.15,

$$R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, R\pi_* L\pi^* \mathcal{N}) \in \mathrm{D}_{\mathrm{C-c}}^b(p^{-1}\mathcal{O}_S).$$

The cones of the natural morphisms $R\pi_*L\pi^*\mathcal{N} \rightarrow R\pi_*\pi^*\mathcal{N}$ and $\mathcal{N} \rightarrow R\pi_*\pi^*\mathcal{N}$ are supported in T_0 , so that, by the induction hypothesis we deduce that $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, R\pi_*\pi^*\mathcal{N}) \in \mathbb{D}_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$ and thus $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{N}) \in \mathbb{D}_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$. q.e.d.

Step 2: Induction on $\dim \text{Supp}_X \mathcal{M}$

Lemma 6.9. *Let $\dim S = d \geq 1$ and let us assume that $(6.6)_{d-1}$ holds. Then $(6.6)_{d,0}$ holds.*

Proof. This is a consequence of Theorem 1 and adjunction. Indeed, we can assume that $\text{Supp}_X \mathcal{M} = \{x\}$. Denoting by $i : \{x\} \times S \hookrightarrow X \times S$ the inclusion, there exists a coherent \mathcal{O}_S -module \mathcal{M}_0 such that $\mathcal{M} = {}_D i_* \mathcal{M}_0$ by Kashiwara's equivalence, and we have (cf. [13, Th. 4.33], which can be proved in a simple way in the present setting)

$$Ri_* R\mathcal{H}om_{\mathcal{O}_S}(\mathcal{M}_0, {}_D i^* \mathcal{N}) \xrightarrow{\sim} R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{N})[\dim X].$$

By Theorem 1, ${}_D i^* \mathcal{N}$ has \mathcal{O}_S -coherent cohomology, and the assertion follows. q.e.d.

We are thus reduced to proving:

(6.10) *Let $\dim S = d \geq 1$. Assume that $(6.6)_{d,k-1}$ holds (with $k \geq 1$). Then $(6.6)_{d,k}$ holds.*

We can reduce to the case \mathcal{M}, \mathcal{N} are regular holonomic $\mathcal{D}_{X \times S/S}$ -modules. Let Z denote the X -support of \mathcal{M} . Since the assertion is local, we can assume that there exists a hypersurface Y of X such that Y contains the singular locus of Z and $\dim Z \cap Y < k$. Recall that localization along $Y \times S$ preserves regular holonomicity (cf. Corollary 4.31).

Lemma 6.11. *It is enough to prove the assertion (6.10) for \mathcal{M}, \mathcal{N} such that $\mathcal{M} = \mathcal{M}(*Y)$ and $\mathcal{N} = \mathcal{N}(*Y)$.*

Proof. The assertion for \mathcal{M} follows from the property that $R\Gamma_{[Y]}\mathcal{M}$ belongs to $\mathbb{D}_{\text{hol}}^b(\mathcal{D}_{X \times S/S})$ (Corollary 4.32) and has X -support of dimension $< k$.

For the assertion concerning \mathcal{N} , we recall the argument given at the end of the proof of [8, Th. 3]. It is enough to prove that (6.10) holds if $\mathcal{N} = R\Gamma_{[Y]}\mathcal{N}$. We have, by [29, (3)],

$$R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{N}) \simeq R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathbf{D}\mathcal{N}, \mathbf{D}\mathcal{M}).$$

Since \mathcal{N} has $\mathcal{D}_{X \times S/S}$ -coherent cohomology and is supported on $Y \times S$, we have

$$R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathbf{D}\mathcal{N}, (\mathbf{D}\mathcal{M})(*Y)) = 0.$$

Furthermore, $\mathbf{D}\mathcal{M}$ being regular holonomic, so is $R\Gamma_{[Y]}(\mathbf{D}\mathcal{M})$ by Corollary 4.32, has well as $\mathcal{M}' := \mathbf{D}R\Gamma_{[Y]}(\mathbf{D}\mathcal{M})$. Finally, applying once more [29, (3) & (1)], we obtain

$$R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{N}) \simeq R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}', \mathcal{N}),$$

with $\dim \text{Supp}_X \mathcal{H}^j \mathcal{M}' < k$ for any j , so the latter complex is S - \mathbb{C} -constructible by (6.6) _{$d,k-1$} . q.e.d.

Step 3: Reduction to the case where \mathcal{M} is of D -type. We take up the notation of the proof of Proposition 4.27, so that $f : X' \rightarrow X$ is a projective morphism inducing a biholomorphism $X' \setminus D \xrightarrow{\sim} Z \setminus Z \cap Y$, and we set $\delta = \dim X' - \dim X$.

Lemma 6.12. *Let \mathcal{M}, \mathcal{N} be regular holonomic $\mathcal{D}_{X \times S/S}$ -modules such that $\mathcal{M} = \mathcal{M}(*Y)$. Then*

$$Rf_* R\mathcal{H}om_{\mathcal{D}_{X' \times S/S}}({}_{\mathbb{D}}f^*\mathcal{M}, {}_{\mathbb{D}}f^*\mathcal{N}) \simeq R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{N}).$$

Proof. There is a natural morphism of adjunction (cf. [13, Th. 4.33(1)])

$$Rf_* R\mathcal{H}om_{\mathcal{D}_{X' \times S/S}}({}_{\mathbb{D}}f^*\mathcal{M}, {}_{\mathbb{D}}f^*\mathcal{N}) \longrightarrow R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}({}_{\mathbb{D}}f_* {}_{\mathbb{D}}f^*\mathcal{M}[\delta], \mathcal{N}).$$

We note that ${}_{\mathbb{D}}f^*\mathcal{M}, {}_{\mathbb{D}}f^*\mathcal{N}$ are f -good. Therefore, this morphism is a quasi-isomorphism. Furthermore, the natural adjunction morphism ${}_{\mathbb{D}}f_* {}_{\mathbb{D}}f^*\mathcal{M}[\delta] \rightarrow \mathcal{M}$ already used in the proof of Proposition 4.27 is an isomorphism if $\mathcal{M} = \mathcal{M}(*Y)$.
q.e.d.

Step 4: The case where \mathcal{M} is of D -type along a normal crossing divisor. Recall Definition 4.23. From Step 3, we are reduced to proving (6.10) in the following setting:

(6.13) $\dim S = d, \dim X = k, \mathcal{M}$ is of D -type along a normal crossing divisor $D \subset X$, and $\mathcal{N} = \mathcal{N}(*D)$.

Since the assertion is local, we can assume that $(E_L, \nabla) = (\widehat{E}_{G|X^* \times S}, \widehat{\nabla})$ as in (4.17) for some \mathcal{O}_S -coherent module G . We denote by $r \geq 0$ the generic rank of G as an \mathcal{O}_S -module. The case $r = 0$ means that G is a torsion \mathcal{O}_S -module.

Let us consider X, S, D with $\dim S = d \geq 1, \dim X = k \geq 1$ and D is a divisor with normal crossings in X . We shall argue by induction on r . For that purpose we consider, for $r \geq 0$, the assertion:

(6.6) _{d, k, r} For any such X, S, D , the property (6.6) _{$d, k-1$} holds, as well as (6.6) _{d, k} if \mathcal{M} is of D -type with $E_L = \widehat{E}_{G|X^* \times S}$ and G being \mathcal{O}_S -coherent of generic rank $\leq r$, and \mathcal{N} being regular holonomic satisfying $\mathcal{N} = \mathcal{N}(*D)$.

Then, proving (6.13) amounts to proving that (6.6) _{d, k, r} for any $r \geq 0$, according to Lemma 6.11.

Lemma 6.14. *If (6.6) _{$d-1$} holds, then*

- (1) (6.6) _{$d, k, 0$} holds;
- (2) if $r \geq 1$ and if (6.6) _{d, k, r} holds for any G which is \mathcal{O}_S -locally free and any $\mathcal{N} = \mathcal{N}(*D)$, then (6.6) _{d, k, r} holds.

Proof. The first point follows from Lemma 6.7. In order to prove the second point, we choose (locally on S) a projective modification $\pi : S' \rightarrow S$ as in Proposition 4.16. Since the cones of the natural morphisms $\mathcal{N} \rightarrow R\pi_* \pi^* \mathcal{N}$ and $R\pi_* L\pi^* \mathcal{N} \rightarrow R\pi_* \pi^* \mathcal{N}$ are S -supported in dimension $< d$, Lemmas 6.7 and 2.15 imply that (6.6) holds for \mathcal{M}, \mathcal{N} on $X \times S$ if it holds for $L\pi^* \mathcal{M}, L\pi^* \mathcal{N}$ on $X \times S'$. Furthermore, $L^j \pi^* \mathcal{M}, L^j \pi^* \mathcal{N}$ are also S' -supported in dimension $< d$ if $j \neq 0$, so, by the same argument, (6.6) holds for \mathcal{M}, \mathcal{N} if it holds $\pi^* \mathcal{M}$ and $\pi^* \mathcal{N}$. Lastly, by Lemma 6.7, we can replace $\pi^* \mathcal{M}$ and $\pi^* \mathcal{N}$ with their quotient $\mathcal{M}', \mathcal{N}'$ by their S' -torsion subsheaves. Note that we still have $\mathcal{N}' = \mathcal{N}'(*D)$ and \mathcal{M}' is of D -type with corresponding $E_{L'}$ isomorphic to $\widehat{E}_{G'|X^* \times S'}$, where G' is $p'^{-1} \mathcal{O}_{S'}$ -locally free by the choice of π . The assumption in (2) therefore implies that (6.6) holds for any G of generic rank r , as wanted.
q.e.d.

Step 5: Induction on the generic rank of \mathcal{M} . We use the same notation as in Step 4 of the proof of Theorem 4.15.

Lemma 6.15. *Assume that (6.6)_{d,k-1} holds. Then (6.6)_{d,k,1} holds.*

Proof. Assume \mathcal{M} is of D-type with $E_L = \widehat{E}_G|_{X^* \times S}$ and G being \mathcal{O}_S -coherent and of generic rank one, and let \mathcal{N} be regular holonomic with $\mathcal{N} = \mathcal{N}(*D)$. By a projective modification we can assume that G is \mathcal{O}_S -locally free of rank one. Then, by Proposition 4.25(1), we have $\mathcal{M} = \widehat{E}_G$.

The matrix of the relative connection $\widehat{\nabla}$ on \widehat{E}_G in some local \mathcal{O}_S -basis of G writes $d_{X \times S/S} \otimes \text{Id} + \sum_{i=1}^{\ell} \alpha_i(s) dx_i/x_i \otimes \text{Id}$. Let us denote by $\mathcal{N}x^{-\alpha}$ the $\mathcal{O}_{X \times S}$ -module $\mathcal{N}(*D)$ for which we add to the connection the relative 1-form $-\sum_{i=1}^{\ell} \alpha_i(s) dx_i/x_i \otimes \text{Id}$. Then $\mathcal{N}x^{-\alpha}$ is also regular holonomic, according to Corollary 4.35. Since $\mathcal{M} = \mathcal{M}(*D)$ and $\mathcal{N} = \mathcal{N}(*D)$, they also are $\mathcal{D}_{X \times S/S}(*D)$ -modules and, as recalled before Corollary 4.35, the functor $(\bullet)x^{\alpha}$ is an auto-equivalence of $\text{Mod}(\mathcal{D}_{X \times S/S}(*D))$. Therefore, we have

$$\begin{aligned} R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{N}) &= R\mathcal{H}om_{\mathcal{D}_{X \times S/S}(*D)}(\mathcal{M}, \mathcal{N}) \\ &\simeq R\mathcal{H}om_{\mathcal{D}_{X \times S/S}(*D)}(\mathcal{M}x^{-\alpha}, \mathcal{N}x^{-\alpha}) \\ &= R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}x^{-\alpha}, \mathcal{N}x^{-\alpha}). \end{aligned}$$

Furthermore, $\mathcal{M}x^{-\alpha} \simeq (\mathcal{O}_{X \times S}(*D), d_{X \times S/S})$. Since $\text{DR}(\mathcal{N}x^{-\alpha})$ is S - \mathbb{C} -constructible (cf. [29, Th. 3.7]) and $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}([\mathcal{O}_{X \times S}(*D)/\mathcal{O}_{X \times S}], \mathcal{N}x^{-\alpha})$ also by (6.6)_{d,k-1}, we conclude that $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{N})$ belongs to $D_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$. q.e.d.

Lemma 6.16. *Let us assume that (6.6)_{d,k,r-1} holds ($r \geq 2$). Then (6.6)_{d,k,r} holds.*

Proof. Let us assume that $r \geq 2$. Together with the assumptions of (6.13) in Step 4, we can now assume that G is \mathcal{O}_S -locally free of finite rank r (Lemma 6.14) and thus, by Proposition 4.25(1), $\mathcal{M} \simeq (\widehat{E}_G, \widehat{\nabla})$. We argue in a way similar to that in Step 4 in the proof of Theorem 4.15(3).

If each endomorphism $A_i(s)$ occurring in the connection matrix of ∇ is scalar, i.e., of the form $\alpha_i(s) \text{Id}$, then Lemma 6.15 allows us to conclude, by obvious reduction to rank one, the proof of (6.6) for $\mathcal{M} = (\widehat{E}_G, \widehat{\nabla})$ and \mathcal{N} as above.

We can thus assume that some endomorphism $A_i(s)$, say $A_1(s)$, is not scalar. We use the notation of Step 4 in the proof of Theorem 4.15(3), in particular (4.21) and (4.22). Recall that we consider the composition $S' \xrightarrow{\pi} \Sigma \xrightarrow{\sigma} S$, with $\Sigma := \{\det(\alpha_1 \text{Id} - A_1(s)) = 0\}$ and $\pi : S' \rightarrow \Sigma$ is a resolution of singularities.

Recall also that we set $G' = \ker(\alpha_1 \text{Id} - A_1 \circ \sigma \circ \pi) \subset (\sigma \circ \pi)^*G$ and we consider the exact sequence $0 \rightarrow G' \rightarrow (\sigma \circ \pi)^*G \rightarrow G'' \rightarrow 0$. By construction and assumption, we have $1 \leq \text{rk } G' < \text{rk } G$ and thus $\text{rk } G'' < \text{rk } G$.

By applying the induction hypothesis (6.6)_{d,k,r-1} to G', G'' , we deduce that

$$R\mathcal{H}om_{\mathcal{D}_{X \times S'/S'}}(L(\sigma \circ \pi)^*\mathcal{M}, L(\sigma \circ \pi)^*\mathcal{N})$$

belongs to $D_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$, hence so does its pushforward by $\sigma \circ \pi$. Thus

$$R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, R(\sigma \circ \pi)_*L(\sigma \circ \pi)^*\mathcal{N}),$$

belongs to $D_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$ by Lemma 2.15.

The cones of the natural morphisms $R(\sigma \circ \pi)_* L(\sigma \circ \pi)^* \mathcal{N} \rightarrow R(\sigma \circ \pi)_*(\sigma \circ \pi)^* \mathcal{N}$, $(\sigma \circ \pi)_*(\sigma \circ \pi)^* \mathcal{N} \rightarrow R(\sigma \circ \pi)_*(\sigma \circ \pi)^* \mathcal{N}$ and $\sigma_* \sigma^* \mathcal{N} \rightarrow (\sigma \circ \pi)_*(\sigma \circ \pi)^* \mathcal{N}$ are supported on $X \times \sigma(\text{Sing}(\Sigma))$. Therefore, $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, (\sigma \circ \pi)_*(\sigma \circ \pi)^* \mathcal{N}) \in \mathbb{D}_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$. Recalling (cf. (4.21)) that locally $\sigma_* \mathcal{O}_{X \times \Sigma}$ is $\mathcal{O}_{X \times S}$ -free of rank $\deg \sigma$, $\sigma_* \sigma^* \mathcal{N}$, which is isomorphic to $\sigma_* \mathcal{O}_{X \times \Sigma} \otimes_{\mathcal{O}_{X \times S}} \mathcal{N}$, is regular holonomic. Besides, according to (4.22), \mathcal{N} is a direct summand of $\sigma_* \sigma^* \mathcal{N}$. Therefore $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{N})$ is a direct summand of $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \sigma_* \sigma^* \mathcal{N})$. Furthermore, the cone of the natural morphism $\sigma_* \sigma^* \mathcal{N} \rightarrow (\sigma \circ \pi)_*(\sigma \circ \pi)^* \mathcal{N}$ is supported on $X \times \sigma(\text{Sing}(\Sigma))$, thus, again by (6.6)_{d-1}, $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \sigma_* \sigma^* \mathcal{N})$ is an object of $\mathbb{D}_{\mathbb{C}\text{-c}}^b(p^{-1}\mathcal{O}_S)$, hence the same holds true for $R\mathcal{H}om_{\mathcal{D}_{X \times S/S}}(\mathcal{M}, \mathcal{N})$. q.e.d.

This lemma concludes Step 5, therefore the proof of (6.6)_d, and thereby that of Theorem 2. q.e.d.

As a consequence of Theorem 2 we can now generalize Proposition 4.26:

Theorem 6.17. *Let Y be a hypersurface in X . Then the category of $\mathcal{D}_{X \times S/S}$ -modules of D -type along Y is equivalent to the category of S -locally constant sheaves on $X^* \times S$ with finite rank, under the functor $\mathcal{L} \mapsto L = \mathcal{H}^0 \text{DR}(\mathcal{L}|_{X^* \times S})$ with quasi-inverse $L \rightarrow \text{RH}_X^S(j_! \mathbf{D}'L)[-d_X]$.*

Proof. Let us prove that the second functor takes values in the category of modules of D -type along Y . Let L be an S -locally constant sheaf on $X^* \times S$ of finite rank. Since, by [8, Prop. 2.6] (valid without any restriction on $\dim S$), $\text{RH}_X^S(j_! \mathbf{D}'L)$ is localized along Y , and regular holonomic by Theorem 2, hence $\mathcal{D}_{X \times S/S}(*Y)$ -coherent, it remains to prove that $\text{RH}_X^S(j_! \mathbf{D}'L)[-d_X]$ is in degree zero and that $\text{RH}_X^S(j_! \mathbf{D}'L)[-d_X]|_{X^* \times S} \simeq \mathcal{O}_{X^* \times S} \otimes_{p_{X^*}^{-1} \mathcal{O}_S} L$. The first assertion follows by the usual Nakayama-type argument. The second assertion follows from [31, Cor. 3.24] (valid without any restriction on $\dim S$), which shows that $\text{RH}_{X^*}^S(\mathbf{D}'L)[-d_X] \simeq \mathcal{O}_{X^* \times S} \otimes_{p_{X^*}^{-1} \mathcal{O}_S} L$.

Let \mathcal{L} be a $\mathcal{D}_{X \times S/S}$ -module of D -type along Y with associated S -local system L , that is $\text{DR}(\mathcal{L})|_{X^* \times S} \simeq L$; we have a natural morphism $\text{DR}(\mathcal{L}) \rightarrow j_* L$. By Corollary 4.33, \mathcal{L} is regular holonomic. According to Lemma 5.15, $\text{DR}(\text{RH}_X^S(j_! \mathbf{D}'L)[-d_X]) \simeq j_* L$ thus, by Theorem 2, there exists a natural morphism of localized regular holonomic $\mathcal{D}_{X \times S/S}$ -modules $\mathcal{L} \rightarrow \text{RH}_X^S(j_! \mathbf{D}'L)[-d_X]$, which has to be an isomorphism since that is so on $X^* \times S$. q.e.d.

References

- [1] Y. André, F. Baldassarri, and M. Cailotto, *De Rham cohomology of differential modules on algebraic varieties*, 2nd revised ed., Progress in Math., vol. 189, Birkhäuser, Basel, 2020.
- [2] J. M. Aroca, H. Hironaka, and J. L. Vicente, *Complex analytic desingularization*, Springer, Tokyo, 2018.
- [3] C. Bănică and O. Stănăşilă, *Algebraic methods in the global theory of complex spaces*, Editura Academiei, Bucharest, 1976.
- [4] J.-E. Björk, *Analytic \mathcal{D} -modules and applications*, Kluwer Academic Publisher, Dordrecht, 1993.
- [5] A. D'Agnolo, S. Guillermou, and P. Schapira, *Regular holonomic $\mathcal{D}[[\hbar]]$ -modules*, Publ. RIMS, Kyoto Univ. **47** (2011), no. 1, 221–255.

- [6] P. Deligne, *Équations différentielles à points singuliers réguliers*, Lect. Notes in Math., vol. 163, Springer-Verlag, 1970.
- [7] L. Fiorot and T. Monteiro Fernandes, *t-structures for relative \mathcal{D} -modules and t-exactness of the de Rham functor*, J. Algebra **509** (2018), 419–444.
- [8] L. Fiorot, T. Monteiro Fernandes, and C. Sabbah, *Relative regular Riemann-Hilbert correspondence*, Proc. London Math. Soc. (3) **122** (2021), no. 3, 434–457, Erratum: *Ibid.*, **123** (2021) no. 6, p. 649–654.
- [9] H. Grauert and R. Remmert, *Bilder und Urbilder analytischer Garben*, Ann. of Math. (2) **68** (1958), 393–443.
- [10] P. H. Hai, J. P. dos Santos, and P. T. Tãm, *Algebraic theory of formal regular-singular connections with parameters*, [arXiv:2107.06474](https://arxiv.org/abs/2107.06474), 2021.
- [11] H. Hironaka, *Resolution of singularities of an algebraic variety over a field of characteristic zero. I*, Ann. of Math. (2) **79** (1964), 109–203.
- [12] M. Kashiwara, *The Riemann-Hilbert problem for holonomic systems*, Publ. RIMS, Kyoto Univ. **20** (1984), 319–365.
- [13] ———, *D-modules and microlocal calculus*, Translations of Mathematical Monographs, vol. 217, American Mathematical Society, Providence, R.I., 2003.
- [14] M. Kashiwara and T. Kawai, *On the holonomic systems of differential equations (systems with regular singularities) III*, Publ. RIMS, Kyoto Univ. **17** (1981), 813–979.
- [15] M. Kashiwara and P. Schapira, *Sheaves on manifolds*, Grundlehren Math. Wiss., vol. 292, Springer-Verlag, Berlin, Heidelberg, 1990.
- [16] ———, *Moderate and formal cohomology associated with constructible sheaves*, Mém. Soc. Math. France (N.S.), vol. 64, Société Mathématique de France, Paris, 1996.
- [17] ———, *Ind-sheaves*, Astérisque, vol. 271, Société Mathématique de France, Paris, 2001.
- [18] ———, *Categories and sheaves*, Grundlehren Math. Wiss., vol. 332, Springer-Verlag, Berlin, Heidelberg, 2006.
- [19] G. Laumon, *Transformation de Fourier généralisée*, [arXiv:9603004](https://arxiv.org/abs/9603004), 1996.
- [20] Y. Laurent and P. Schapira, *Images inverses des modules différentiels*, Compositio Math. **61** (1987), no. 2, 229–251.
- [21] Ph. Maisonobe, *Filtration relative, l'idéal de Bernstein et ses pentes*, 2016, [hal-01285562v2](https://arxiv.org/abs/1601.01285).
- [22] Ph. Maisonobe and Z. Mebkhout, *Le théorème de comparaison pour les cycles évanescents*, Éléments de la théorie des systèmes différentiels géométriques, Séminaires & Congrès, vol. 8, Société Mathématique de France, Paris, 2004, pp. 311–389.
- [23] Z. Mebkhout, *Le théorème de positivité, le théorème de comparaison et le théorème d'existence de Riemann*, Éléments de la théorie des systèmes différentiels géométriques, Séminaires & Congrès, vol. 8, Société Mathématique de France, Paris, 2004, pp. 165–310.
- [24] Z. Mebkhout and C. Sabbah, *§III.4 \mathcal{D} -modules et cycles évanescents*, Le formalisme des six opérations de Grothendieck pour les \mathcal{D} -modules cohérents, Travaux en cours, vol. 35, Hermann, Paris, 1989, pp. 201–239.
- [25] T. Mochizuki, *Holonomic \mathcal{D} -modules with Betti structure*, Mém. Soc. Math. France (N.S.), vol. 138–139, Société Mathématique de France, Paris, 2014.
- [26] ———, *Mixed twistor D -modules*, Lect. Notes in Math., vol. 2125, Springer, Heidelberg, New York, 2015.
- [27] T. Monteiro Fernandes and L. Prelli, *Relative subanalytic sheaves*, Fund. Math. **226** (2014), no. 1, 79–100.
- [28] ———, *Relative subanalytic sheaves II*, Milan J. Math. **89** (2021), 387–411.
- [29] T. Monteiro Fernandes and C. Sabbah, *On the de Rham complex of mixed twistor \mathcal{D} -modules*, Internat. Math. Res. Notices (2013), no. 21, 4961–4984.
- [30] ———, *Relative Riemann-Hilbert correspondence in dimension one*, Portugal. Math. **74** (2017), no. 2, 149–159.

- [31] ———, *Riemann-Hilbert correspondence for mixed twistor \mathcal{D} -modules*, J. Inst. Math. Jussieu **18** (2019), no. 3, 629–672.
- [32] N. Nitsure, *Moduli of regular holonomic \mathcal{D} -modules with normal crossing singularities*, Duke Math. J. **99** (1999), 1–39.
- [33] L. Prelli, *Sheaves on subanalytic sites*, Rend. Sem. Mat. Univ. Padova **120** (2008), 167–216.
- [34] H. Rossi, *Picard variety of an isolated singular point*, Rice Univ. Studies **54** (1968), no. 4, 63–73.
- [35] M. Rothstein, *Sheaves with connection on abelian varieties*, Duke Math. J. **84** (1996), 565–598.
- [36] C. Sabbah, *Proximité évanescence, II. Équations fonctionnelles pour plusieurs fonctions analytiques*, Compositio Math. **64** (1987), 213–241.
- [37] ———, *Équations différentielles à points singuliers irréguliers et phénomène de Stokes en dimension 2*, Astérisque, vol. 263, Société Mathématique de France, Paris, 2000.
- [38] P. Schapira and J.-P. Schneiders, *Index theorem for elliptic pairs*, Astérisque, vol. 224, Société Mathématique de France, Paris, 1994.
- [39] C. Schnell, *Holonomic D -modules on abelian varieties*, Publ. Math. Inst. Hautes Études Sci. **121** (2015), 1–55, Erratum: *Ibid.* **123** (2016), 361–362.
- [40] L. Wang, *The constructibility theorem for differential modules*, Ph.D. thesis, University of Illinois at Chicago, 2008, <https://hdl.handle.net/10027/13547>.
- [41] L. Wu, *Riemann-Hilbert correspondence for Alexander complexes*, [arXiv:2104.06941](https://arxiv.org/abs/2104.06941), 2021.

L. FIOROT, Dipartimento di Matematica “Tullio Levi-Civita” Università degli Studi di Padova, Via Trieste, 63, 35121 Padova Italy • *E-mail* : luisa.fiorot@unipd.it

T. MONTEIRO FERNANDES, Centro de Matemática e Aplicações Fundamentais – Centro de investigação Operacional e Departamento de Matemática da FCUL, Edifício C 6, Piso 2, Campo Grande, 1700, Lisboa, Portugal • *E-mail* : mtfernandes@fc.ul.pt

C. SABBAB, CMLS, CNRS, École polytechnique, Institut Polytechnique de Paris, 91128 Palaiseau cedex, France • *E-mail* : Claude.Sabbah@polytechnique.edu
Url : <http://www.math.polytechnique.fr/perso/sabbah>