

ON PSEUDOEFFECTIVE CONES OF PROJECTIVE BUNDLES AND VOLUME FUNCTION

SNEHAJIT MISRA AND ANOOP SINGH

ABSTRACT. In this article, we compute the pseudoeffective cones of various projective bundles $\mathbb{P}_X(E)$ over higher dimensional varieties X under some assumptions on X as well as on the vector bundle E . We also compute the volume function on fibre product of two projective bundles $\mathbb{P}(F) \times_C \mathbb{P}(E)$ over a smooth irreducible complex projective curve C .

1. INTRODUCTION

In the last few decades, a number of notions of positivity have been used to understand the geometry of the higher dimensional projective varieties. A fundamental invariant of a smooth irreducible projective variety X is the closure of the cone generated by effective \mathbb{R} -divisors on X , called as pseudo-effective cone of X and denoted by $\overline{\text{Eff}}^1(X)$, which play a very crucial role in this understanding. The interior of $\overline{\text{Eff}}^1(X)$ is called Big cone of X , and its element are called as big divisors. Most of the developments in this direction has been successfully summarized in [L07], [L11].

In his paper [Miy87], Yoichi Miyaoka initiated the study of nef and effective divisors on a projective bundle $\mathbb{P}(E)$ over a smooth irreducible curve C in characteristics 0, where E is any rank r vector bundle on C , and gives a numerical criterion for semi-stability of E in terms of nefness of the normalized hyperplane class λ_E on $\mathbb{P}(E)$. More generally, in [F11], the cone of effective k -cycles in $\mathbb{P}(E)$ over an irreducible curve C is described in terms of the numerical data appearing in Harder-Narasimhan filtration of the bundle E , which generalizes Miyaoka's result in the semistable case. Later the pseudo-effective cone of the Grassmannian bundle $\text{Gr}_s(E)$ parametrizing all the s dimensional quotients of the fibres of E where $1 \leq s \leq r - 1$, has been studied in [BHP14]. However, in most of these cases, the Picard number of the space X is 2, and hence the pseudo-effective cones of divisors are finite polyhedra (see [F11]) generated by two extremal rays in a two dimensional space. When the Picard number is at least 3, there are very few examples where the pseudo-effective cones are computed. Recently, the pseudo-effective cones of the fiber product $\mathbb{P}(F)_C \times_C \mathbb{P}_C(E)$ of two projective bundles over a smooth irreducible complex projective curve C is computed in a different context in [M21], which has Picard number 3.

2020 *Mathematics Subject Classification*. Primary 14C20, 14E30, 14E99 ; Secondary 14D20, 14D06 .

Key words and phrases. Pseudo-effective cone, Big Cone, Semistability, Projective bundle, Volume Function.

In this article, our first main result is the following.

Theorem 1.1. *Let E be a semistable vector bundle of rank r on a smooth irreducible projective surface X such that $c_2(\text{End}(E)) = 2rc_2(E) - (r-1)c_1^2(E) = 0 \in H^4(X, \mathbb{Q})$. Then*

$$\overline{\text{Eff}}^1(\mathbb{P}_X(E)) = \left\{ x\lambda_E + \pi^*\gamma \mid x \in \mathbb{R}_{\geq 0}, \gamma \in \overline{\text{Eff}}^1(X) \right\},$$

where $\pi : \mathbb{P}_X(E) \rightarrow X$ is the projection and $\lambda_E := c_1(\mathcal{O}_{\mathbb{P}_X(E)}(1)) - \frac{1}{r}\pi^*c_1(E)$.

Further assume that $\overline{\text{Eff}}^1(X)$ is a finite polyhedra generated by $\{E_1, E_2, \dots, E_n\}$. Then

$$\overline{\text{Eff}}^1(\mathbb{P}_X(E)) = \left\{ x_0\lambda_E + x_1\pi^*E_1 + x_2\pi^*E_2 + \dots + x_n\pi^*E_n \mid x_i \in \mathbb{R}_{\geq 0} \right\}.$$

We also give several Corollaries and examples to illustrate Theorem 1.1 (see Corollary 4.2, Corollary 4.5, Example 4.6, Example 4.7 etc).

Next we restrict our attention in finding pseudo-effective cone of $\mathbb{P}_X(E)$ over higher dimensional varieties X where $c_2(\text{End}(E)) \neq 0$. In this context, we prove the following Theorems.

Theorem 1.2. *Let X be a smooth irreducible complex projective surface of Picard number 1, and L_X be an ample generator of $N^1(X)_{\mathbb{R}}$. Let $E = \mathcal{O}_X \oplus \mathcal{L}$ be rank 2 bundle over X such that $\mathcal{L} \equiv nL_X$ for some positive integer $n > 0$. Then*

$$\overline{\text{Eff}}^1(\mathbb{P}_X(E)) = \left\{ a(\xi - n\pi^*L_X) + b\pi^*L_X \mid a, b \in \mathbb{R}_{\geq 0} \right\},$$

where $\pi : \mathbb{P}_X(E) \rightarrow X$ is the projection and ξ denote the numerical class of $\mathcal{O}_{\mathbb{P}_X(E)}(1)$.

In particular, if $E = \mathcal{O}_{\mathbb{P}^2} \oplus \mathcal{O}_{\mathbb{P}^2}(n)$ is a rank 2 bundle on \mathbb{P}^2 for some positive integer n , then

$$\overline{\text{Eff}}^1(\mathbb{P}_{\mathbb{P}^2}(E)) = \left\{ a(\xi - n\pi^*H) + b\pi^*H \mid a, b \in \mathbb{R}_{\geq 0} \right\}.$$

where H is the numerical class of $\mathcal{O}_{\mathbb{P}^2}(1)$.

Theorem 1.3. *Let X be a smooth irreducible complex projective variety of Picard number 1 with an ample generator L_X of $N^1(X)_{\mathbb{R}}$. We consider the rank 3 bundle $E = \mathcal{O}_X \oplus L \oplus M$ such that $L \equiv mL_X$ and $M \equiv nL_X$ with $m \geq n > 0$. Consider the Harder Narasimhan filtration of E with respect to L_X as follow:*

$$\mathcal{O}_X \subsetneq L \oplus M \subsetneq E$$

with $E_1 = L \oplus M$ and $Q_1 = E/E_1 = X$. The projection map $p : \mathbb{P}(E) \setminus \mathbb{P}(Q_1) \rightarrow \mathbb{P}(E_1)$ can be seen as a rational map. The indeterminacies of this rational map are resolved by blowing up $\mathbb{P}(Q_1)$. Let $\beta : \text{Bl}_{\mathbb{P}(Q_1)}\mathbb{P}(E) \rightarrow \mathbb{P}(E)$ be the blow up map. Consider the following commutative diagram.

$$\begin{array}{ccc} \mathbb{P}_Y(\mathcal{F}) = \text{Bl}_{\mathbb{P}_X(Q_1)}\mathbb{P}_X(E) & \xrightarrow{\eta} & \mathbb{P}_X(E_1) = \mathbb{P}(L \oplus M) = Y \\ \downarrow \beta & & \downarrow \rho \\ \mathbb{P}_X(E) & \xrightarrow{\pi} & X \end{array}$$

Then the cone map

$$C^{(1)} : \overline{\text{Eff}}^1(\mathbb{P}(L \oplus M)) \longrightarrow \overline{\text{Eff}}^1(\mathbb{P}(E))$$

defined as follows :

$$C^{(1)}(\alpha) := \beta_* \eta^*(\alpha),$$

induces an isomorphism of $\overline{\text{Eff}}^1(\mathbb{P}(L \oplus M))$ onto $\overline{\text{Eff}}^1(\mathbb{P}(E))$.

Theorem 1.4. *Let C be a smooth irreducible complex projective curve and $\rho : X = \mathbb{P}_C(W) \longrightarrow C$ is a ruled surface defined by a normalized vector bundle (in the sense of Hartshorne) of rank 2 with $\mu_{\min}(W) = \deg(W)$. We denote the numerical classes of the tautological bundle $\mathcal{O}_X(1)$ and a fiber of the map ρ by η and f respectively. Let $E = \mathcal{O}_X \oplus \mathcal{L}$ be a rank 2 vector bundle on X such that \mathcal{L} is nef with $c_1(\mathcal{L}) \equiv a\eta + bf \in N^1(X)_{\mathbb{R}}$. Then*

$$\overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ x(\xi - a\pi^*\eta - b\pi^*f) + y(\pi^*\eta) + z(\pi^*f) \mid x, y, z \in \mathbb{R}_{\geq 0} \right\}.$$

Let L be a line bundle on a smooth irreducible complex projective variety X of dimension n . The volume of L , denoted by $\text{vol}(L)$ is defined as follows:

$$\text{vol}_X(L) := \limsup_{n \rightarrow \infty} \frac{\dim_k H^0(X, L^{\otimes m})}{m^n/n!}.$$

It is true, but not entirely obvious that the limsup is actually a limit (see [F94],[T07]). The volume function is invariant under numerical equivalence. Hence we have a well-defined mapping from the Néron-Severi group $N^1(X)_{\mathbb{Q}}$ to \mathbb{Q}_+ which we again denote by vol_X . Infact, the function

$$\xi \mapsto \text{vol}_X(\xi)$$

on $N^1(X)_{\mathbb{Q}}$ extends uniquely to a continuous function $\text{vol}_X : N^1(X)_{\mathbb{R}} \longrightarrow \mathbb{R}_+$ on the real Néron-Severi group $N^1(X)_{\mathbb{R}}$. If L is an ample line bundle, then by asymptotic Riemann-Roch one has $\text{vol}_X(L) = L^d$. In particular, the volume function is polynomial type on the ample cone of X . By definition of big divisors, a line bundle L is big if and only if $\text{vol}_X(L) > 0$. However, on the larger cone of big divisors, the behaviour of the volume function is not completely understood in general, and in such cases the volume functions are not so easy to calculate. More details about volume functions can be found in [ELMNP05] and [L11].

When X is a surface, then the volume function is piecewise polynomial on the pseudoeffective cone of the surface X , and the volume function can be computed using the Zariski decomposition of pseudoeffective divisors on X (see Chapter 2, [L11]).

In [C11], the volume function has been calculated on projective bundle $\mathbb{P}_C(E)$ over a smooth irreducible curve C defined over a field of characteristic zero . It is proved in [C11] that there is a

subdivision of the real Néron-Severi group $N^1(\mathbb{P}_C(E))_{\mathbb{R}}$ into sectors such that the volume function is polynomial on each sector.

In his thesis [W05], Wolfe consider the case of split \mathbb{P}^1 -bundle on a smooth projective variety X . More precisely, he computed the volume function on $\mathbb{P}_X(L_1 \oplus L_2)$ for some line bundles L_1 and L_2 . It is shown in [W05] that when X is a surface S , then there exists a dense, open subset $U \subseteq \text{Big}(\mathbb{P}_S(L_1 \oplus L_2))$ inside the big cone so that $\text{vol}|_U$ is real analytic.

In [ELMNP05], the following question is suggested about volume function on projective bundles over higher dimensional varieties (see Problem 2.13, Page 389, [ELMNP05]).

Question : What is the behaviour of volume function of line bundles on projective bundles $\mathbb{P}_X(E)$ over a higher dimensional varieties X ? It is also interesting to see to what extent the geometry of E is related to the volume function on $\mathbb{P}_X(E)$.

Motivated by the above question, we calculate the volume function on fibre product $\mathbb{P}_C(E) \times_C \mathbb{P}_C(F)$ of two projective bundles over a smooth curve C defined over a field of characteristic zero. Our proofs closely follow the ideas in [C11]. Note that the fibre product $X = \mathbb{P}_C(E) \times_C \mathbb{P}_C(F)$ has Picard number 3, and the behaviours of different cones inside $N^1(X)_{\mathbb{R}}$ are complicated in general.

Let C be a smooth projective irreducible curve defined over the field \mathbb{C} , and $K := k(C)$ be its function field. If E is a non-zero vector bundle on C , then there exists a unique filtration, called as Harder-Narasimhan filtration of E , as follows

$$0 = E_d \subsetneq E_{d-1} \subsetneq \cdots \subsetneq E_1 \subsetneq E_0 = E$$

such that each sub-quotient E_i/E_{i+1} is semistable for each $i \in \{0, 1, 2, \dots, d-1\}$, and

$$\mu_0 < \mu_1 < \cdots < \mu_{d-1}$$

where $\mu_i = \mu(E_i/E_{i+1}) = \frac{\deg(E_i/E_{i+1})}{\text{rk}(E_i/E_{i+1})}$ for each i . Then one can define a probability measure ν_E on \mathbb{R} as follows:

$$\nu_E := \frac{1}{\text{rk}(E)} \sum_{i=0}^{d-1} \{\text{rk}(E_{i+1}) - \text{rk}(E_i)\} \delta_{\mu_i},$$

where δ_x is the Dirac measure concentrated on $\{x\}$. For any real number $\varepsilon > 0$, let T_ε be the operator on the set of all Borel probability measure on \mathbb{R} which is defined by the following property:

if ν is Borel probability measure and if f is a continuous function of compact support on \mathbb{R} , then

$$\int f(x) T_\varepsilon \nu(dx) = \int f(\varepsilon x) \nu(dx).$$

In this article, we fix two vector bundles E and F over a smooth irreducible projective curve C over a field of characteristic 0 having ranks e and f respectively. We consider the fibre product $\mathbb{P}(F) \times_C \mathbb{P}(E)$ of two projective bundle over C .

Let

$$(1) \quad 0 = E_d \subsetneq E_{d-1} \subsetneq \cdots \subsetneq E_1 \subsetneq E_0 = E$$

be the Harder-Narasimhan filtration of E such that each sub-quotient E_i/E_{i+1} is semistable for each $i \in \{0, 1, 2, \dots, d-1\}$, and

$$\mu_0 < \mu_1 < \cdots < \mu_{d-1}$$

where $\mu_i = \mu(E_i/E_{i+1})$ for each i .

Similarly, let

$$(2) \quad 0 = F_h \subsetneq F_{h-1} \subsetneq F_{h-2} \subsetneq \cdots \subsetneq F_1 \subsetneq F_0 = F$$

be the Harder-Narasimhan filtration of F , and let $\mu'_j = \mu(F_j/F_{j+1})$ for each j so that we have

$$\mu'_{h-1} > \mu'_{h-2} > \cdots > \mu'_1 > \mu'_0.$$

Consider the following commutative fibre product diagram:

$$\begin{array}{ccc} X = \mathbb{P}(\mathcal{E}) = \mathbb{P}(F) \times_C \mathbb{P}(E) = \mathbb{P}(\psi^*E) & \xrightarrow{\pi_2} & \mathbb{P}(E) \\ \downarrow \pi_1 & & \downarrow \\ \mathbb{P}(F) & \xrightarrow{\psi} & C \end{array}$$

We call $\pi = \psi \circ \pi_1$ and $\mathcal{E} = \psi^*E$. Consider a line bundle $L = \mathcal{O}_{\mathbb{P}(\mathcal{E})}(m) \otimes \pi_1^* \mathcal{O}_{\mathbb{P}(F)}(l) \otimes \pi_1^* \psi^* \mathcal{M}$ on X , for some line bundle \mathcal{M} of degree a on C such that L_K is big.

We consider the vector $\underline{s} := (s_1, s_2, \dots, s_e, s'_1, s'_2, \dots, s'_f) \in \mathbb{R}^{e+f}$ such that the value μ_i appear exactly $\text{rk}(E_i/E_{i+1})$ times in the first e coordinates and the value μ'_j appears exactly $\text{rk}(F_j/F_{j+1})$ times in the last f coordinates. Let $\Delta \times \Delta' \subset \mathbb{R}^{e+f}$ be the set, where simplexes Δ and Δ' are defined as follows:

$$\Delta = \left\{ (x_1, x_2, \dots, x_e) \in \mathbb{R}^e \mid 0 \leq x_i \leq 1, \sum_{i=1}^e x_i = 1 \right\},$$

$$\Delta' = \left\{ (x'_1, x'_2, \dots, x'_f) \in \mathbb{R}^f \mid 0 \leq x'_j \leq 1, \sum_{j=1}^f x'_j = 1 \right\}.$$

Our main results in this article are as follows:

Theorem 1.5. *Let η be the Lebesgue measure on $\Delta \times \Delta'$ normalized such that $\eta(\Delta \times \Delta') = 1$. Then $T_{\frac{1}{n}} \nu_{\text{Sym}^{mn}(E) \otimes \text{Sym}^{ln}(F)}$ converges vaguely to $\phi_{\underline{s}} \eta$, where*

$$\phi_{\underline{s}} : \Delta \times \Delta' \longrightarrow \mathbb{R}$$

such that

$$(x_1, x_2, \dots, x_e, x'_1, \dots, x'_f) \mapsto \sum_{i=1}^e s_i x_i + \sum_{j=1}^f s'_j x'_j.$$

Theorem 1.6. *Let $L = \mathcal{O}_{\mathbb{P}(E)}(m) \otimes \pi_1^* \mathcal{O}_{\mathbb{P}(F)}(l) \otimes \pi_1^* \psi^* \mathcal{M}$ be a line bundle on $X = \mathbb{P}(F) \times_C \mathbb{P}(E)$ for some line bundle \mathcal{M} of degree a on C such that L_K is big. Then*

$$\mathrm{vol}_X(L) = \dim(X) \mathrm{vol}_{X_K}(L_K) \int (x+a)_+ \phi_{\underline{s}^*} \eta dx,$$

where $x_+ = \max\{x, 0\}$.

2. NOTATION AND CONVENTION

Throughout this article, all the algebraic varieties are assumed to be irreducible and defined over the field of complex numbers \mathbb{C} .

The projective bundle $\mathbb{P}_X(E)$ associated to a vector bundle E over a projective variety X is defined as $\mathbb{P}_X(E) := \mathrm{Proj}\left(\bigoplus_{m \geq 0} \mathrm{Sym}^m(E)\right)$ together with the projection map $\pi : \mathbb{P}_X(E) \rightarrow X$. We will simply write $\mathbb{P}(E)$ whenever the base space X is clear from the context. The numerical class of the tautological line bundle $\mathcal{O}_{\mathbb{P}(E)}(1)$ will be denoted by ξ unless otherwise specified.

The rank of a vector bundle E will be denoted by $\mathrm{rk}(E)$. The numerical class of a divisor D will be denoted by $[D]$.

The set of non-negative (resp. positive) real numbers will be denoted by $\mathbb{R}_{\geq 0}$ (resp. \mathbb{R}_+).

3. PRELIMINARIES

In this section we recall basic definition of pseudo-effective cones and its various properties. We also recall the characterization of semistable bundles with vanishing discriminant over higher dimensional projective varieties which we will use to prove one of our main results.

3.1. Nef cone and Pseudo-effective cone. Let X be a projective variety of dimension n and $\mathcal{Z}_k(X)$ (respectively $\mathcal{Z}^k(X)$) denotes the free abelian group generated by k -dimensional (respectively k -codimensional) subvarieties on X . The Chow groups $A_k(X)$ are defined as the quotient of $\mathcal{Z}_k(X)$ modulo rational equivalence. When X is a smooth irreducible projective variety, we denote $A^k(X) := A_{n-k}(X)$.

Two cycles $Z_1, Z_2 \in \mathcal{Z}^k(X)$ are said to be numerically equivalent, denoted by $Z_1 \equiv Z_2$ if $Z_1 \cdot \gamma = Z_2 \cdot \gamma$ for all $\gamma \in \mathcal{Z}_k(X)$. The *numerical groups* $N^k(X)_{\mathbb{R}}$ are defined as the quotient of $\mathcal{Z}^k(X) \otimes \mathbb{R}$ modulo numerical equivalence. When X is smooth, we define $N_k(X)_{\mathbb{R}} := N^{n-k}(X)_{\mathbb{R}}$ for all k .

Let

$$\mathrm{Div}^0(X) := \{D \in \mathrm{Div}(X) \mid D \cdot C = 0 \text{ for all curves } C \text{ in } X\} \subseteq \mathrm{Div}(X).$$

be the subgroup of $\mathrm{Div}(X)$ consisting of numerically trivial divisors. The quotient $\mathrm{Div}(X)/\mathrm{Div}^0(X)$ is called the Néron Severi group of X , and is denoted by $N^1(X)_{\mathbb{Z}}$. The Néron Severi group $N^1(X)_{\mathbb{Z}}$ is a free abelian group of finite rank. Its rank, denoted by $\rho(X)$ is called the Picard number of

X . In particular, $N^1(X)_{\mathbb{R}}$ is called the real Néron Severi group and $N^1(X)_{\mathbb{R}} := N^1(X)_{\mathbb{Z}} \otimes \mathbb{R} := (\text{Div}(X)/\text{Div}^0(X)) \otimes \mathbb{R}$. For X smooth, the intersection product induces a perfect pairing

$$N^k(X)_{\mathbb{R}} \times N_k(X)_{\mathbb{R}} \longrightarrow \mathbb{R}$$

which implies $N^k(X)_{\mathbb{R}} \simeq (N_k(X)_{\mathbb{R}})^{\vee}$ for every k satisfying $0 \leq k \leq n$. The direct sum $N(X) := \bigoplus_{k=0}^n N^k(X)_{\mathbb{R}}$ is a graded \mathbb{R} -algebra with multiplication induced by the intersection form.

The convex cone generated by the set of all effective k -cycles in $N_k(X)_{\mathbb{R}}$ is denoted by $\text{Eff}_k(X)$ and its closure $\overline{\text{Eff}}_k(X)$ is called the *pseudo-effective cone* of k -cycles in X . For any $0 \leq k \leq n$, $\overline{\text{Eff}}^k(X) := \overline{\text{Eff}}_{n-k}(X)$. The *nef cone* of divisors are defined as follows :

$$\text{Nef}^1(X) := \{ \alpha \in N^1(X) \mid \alpha \cdot \beta \geq 0 \ \forall \beta \in \overline{\text{Eff}}_1(X) \}.$$

The interior $\text{int}(\overline{\text{Eff}}^1(X))$ is called Big cone of big divisors in $N^1(X)_{\mathbb{R}}$, and denoted by $\text{Big}(X)$.

An irreducible curve C in X is called *movable* if there exists an algebraic family of irreducible curves $\{C_t\}_{t \in T}$ such that $C = C_{t_0}$ for some $t_0 \in T$ and $\bigcup_{t \in T} C_t \subset X$ is dense in X .

A class $\gamma \in N_1(X)_{\mathbb{R}}$ is called movable if there exists a movable curve C such that $\gamma = [C]$ in $N_1(X)_{\mathbb{R}}$. The closure of the cone generated by movable classes in $N_1(X)_{\mathbb{R}}$, denoted by $\overline{\text{Mov}}(X)$ is called the *movable cone*. By [BDPP13] $\overline{\text{Mov}}(X)$ is the dual cone to $\overline{\text{Eff}}^1(X)$. Also, $\text{Nef}^1(X) \subseteq \overline{\text{Eff}}^1(X)$ always (see Ch 2 [L07]). We refer the reader to [L07],[L11] for more details about these cones.

3.2. Semistability of Vector bundles. Let X be a smooth complex projective variety of dimension n with a fixed ample line bundle H on it. For a torsion-free coherent sheaf \mathcal{G} of rank r on X , the H -slope of \mathcal{G} is defined as

$$\mu_H(\mathcal{G}) := \frac{c_1(\mathcal{G}) \cdot H^{n-1}}{r} \in \mathbb{Q}.$$

A torsion-free coherent sheaf \mathcal{G} on X is said to be H -semistable if $\mu_H(\mathcal{F}) \leq \mu_H(\mathcal{G})$ for all subsheaves \mathcal{F} of \mathcal{G} . A vector bundle E on X is called H -unstable if it is not H -semistable. For every vector bundle E on X , there is a unique filtration

$$0 = E_d \subsetneq E_{d-1} \subsetneq E_{d-2} \subsetneq \cdots \subsetneq E_1 \subsetneq E_0 = E$$

of subbundles of E , called the Harder-Narasimhan filtration of E , such that E_i/E_{i+1} is H -semistable torsion-free sheaf for each $i \in \{0, 1, 2, \dots, d-1\}$ and $\mu_H(E_{d-1}/E_d) > \mu_H(E_{d-2}/E_{d-1}) > \cdots > \mu_H(E_0/E_1)$. We define $Q_1 := E_0/E_1$ and $\mu_{\min}(E) := \mu_H(Q_1) = \mu_H(E_0/E_1)$ and $\mu_{\max}(E) := \mu_H(E_{d-1}/E_d)$.

For a vector bundle E of rank r over X , the element $c_2(\text{End}(E)) \in H^4(X, \mathbb{Q})$ is called the discriminant of E . We recall the following result from [N99] or [Theorem 1.2,[BB08]] about semistable vector bundles with vanishing discriminant.

Theorem 3.1. *Let E be a vector bundle of rank r on a smooth complex projective variety X . Then the following are equivalent*

- (1) E is semistable and $c_2(\text{End}(E)) = 0$.
- (2) $\lambda_E := c_1(\mathcal{O}_{\mathbb{P}(E)}(1)) - \frac{1}{r}\pi^*c_1(E) \in \text{Nef}^1(\mathbb{P}(E))$.
- (3) For every pair of the form (ϕ, C) , where C is a smooth projective curve and $\phi : C \rightarrow X$ is a non-constant morphism, $\phi^*(E)$ is semistable.

Since nefness of a line bundle does not depend on the fixed polarization (i.e. fixed ample line bundle H) on X , the Theorem 3.1 implies that the semistability of a vector bundle E with $c_2(\text{End}(E)) = 0$ is independent of the fixed polarization H . We will not mention about the polarization H from now on whenever we speak of semistability of such bundles E with $c_2(\text{End}(E)) = 0$. We have the following lemmata as easy applications of Theorem 3.1.

Lemma 3.2. [Lemma 2.2, [M21]] *Let $\psi : X \rightarrow Y$ be a morphism between two smooth complex projective varieties and E is a semistable bundle on Y with $c_2(\text{End}(E)) = 0$. Then the pullback bundle $\psi^*(E)$ is also semistable with $c_2(\text{End}(\psi^*E)) = 0$.*

Lemma 3.3. [Lemma 2.3, [M21]] *Let E be a semistable vector bundle of rank r on a smooth complex projective variety X with $c_2(\text{End}(E)) = 0$. Then for any positive integer m and any line bundle \mathcal{L} on X , $\text{Sym}^m(E) \otimes \mathcal{L}$ is also semistable bundle on X with $c_2(\text{End}(\text{Sym}^m(E) \otimes \mathcal{L})) = 0$.*

4. PSEUDOEFFECTIVE CONE OF PROJECTIVE BUNDLE

Theorem 4.1. *Let E be a semistable vector bundle of rank r on a smooth irreducible projective surface X such that $c_2(\text{End}(E)) = 2rc_2(E) - (r-1)c_1^2(E) = 0 \in H^4(X, \mathbb{Q})$. Then*

$$\overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ x\lambda_E + \pi^*\gamma \mid x \in \mathbb{R}_{\geq 0}, \gamma \in \overline{\text{Eff}}^1(X) \right\}.$$

Further assume that $\overline{\text{Eff}}^1(X)$ is a finite polyhedra generated by $\{E_1, E_2, \dots, E_n\}$. Then

$$\overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ x_0\lambda_E + x_1\pi^*E_1 + x_2\pi^*E_2 + \dots + x_n\pi^*E_n \mid x_i \in \mathbb{R}_{\geq 0} \right\}.$$

Proof. Let D be an effective divisor on $\mathbb{P}(E)$ such that $\mathcal{O}_{\mathbb{P}(E)}(D) \simeq \mathcal{O}_{\mathbb{P}(E)}(m) \otimes \pi^*(\mathcal{L})$ for some integer m and a line bundle $\mathcal{L} \in \text{Pic}(X)$. Then

$$H^0(\mathbb{P}(E), \mathcal{O}_{\mathbb{P}(E)}(m) \otimes \pi^*(\mathcal{L})) = H^0(X, \text{Sym}^m(E) \otimes \mathcal{L}) \neq 0,$$

which implies $m \geq 0$.

Let $\gamma = [C]$ be a movable class in $N_1(X)_{\mathbb{R}}$. Then C belongs to an algebraic family of curves $\{C_t\}_{t \in T}$ such that $\bigcup_{t \in T} C_t$ covers a dense subset of X . So we can find a curve C_{t_1} in this family

such that

$$H^0(C_{t_1}, \text{Sym}^m(E)|_{C_{t_1}} \otimes \mathcal{L}|_{C_{t_1}}) \neq 0.$$

Let $\eta_{t_1} : \tilde{C}_{t_1} \rightarrow C_{t_1}$ be the normalization of the curve C_{t_1} and we call $\phi_{t_1} := i \circ \eta_{t_1}$ where $i : C_{t_1} \hookrightarrow X$ is the inclusion.

As E is a semistable bundle on X with $c_2(\text{End}(E)) = 0$, $\text{Sym}^m(E) \otimes \mathcal{L}$ is also semistable on X and $c_2(\text{Sym}^m(E) \otimes \mathcal{L}) = 0$. Therefore we have $\phi_{t_1}^*(\text{Sym}^m(E) \otimes \mathcal{L})$ is also semistable on \tilde{C}_{t_1} .

Since η_{t_1} is a surjective map, we have

$$H^0(\eta_{t_1}^*(\text{Sym}^m(E)|_{C_{t_1}} \otimes \mathcal{L}|_{C_{t_1}})) = H^0(\phi_{t_1}^*(\text{Sym}^m(E) \otimes \mathcal{L})) \neq 0.$$

This implies that $c_1(\phi_{t_1}^*(\text{Sym}^m(E) \otimes \mathcal{L})) \geq 0$, and hence

$$c_1(\text{Sym}^m(E) \otimes \mathcal{L}) \cdot C_{t_1} = \{c_1(\text{Sym}^m(E) \otimes \mathcal{L}) \cdot \gamma\} \geq 0 \text{ for a movable class } \gamma \in N_1(X)_{\mathbb{R}}.$$

Using the duality property of movable cone $\overline{\text{Mov}}(X)$ we conclude that

$$c_1(\text{Sym}^m(E) \otimes \mathcal{L}) = c_1(\pi_* \mathcal{O}_{\mathbb{P}(E)}(D)) \in \overline{\text{Eff}}^1(X).$$

Now $c_1(\pi_* \mathcal{O}_{\mathbb{P}(E)}(D)) = c_1(\text{Sym}^m(E) \otimes \mathcal{L}) = \text{rk}(\text{Sym}^m(E)) \left\{ \frac{m}{r} c_1(E) + c_1(\mathcal{L}) \right\} \in \overline{\text{Eff}}^1(X)$, so that

$$\frac{m}{r} c_1(E) + c_1(\mathcal{L}) \in \overline{\text{Eff}}^1(X).$$

Since E is a semistable bundle on X with $c_2(\text{End}(E)) = 0$, we also have

$$c_1(\mathcal{O}_{\mathbb{P}(E)}(1)) - \frac{1}{r} \pi^* c_1(E) \in \text{Nef}^1(\mathbb{P}(E)) \subseteq \overline{\text{Eff}}^1(\mathbb{P}(E)).$$

$$\begin{aligned} & \text{Hence, } \mathcal{O}_{\mathbb{P}(E)}(m) \otimes \pi^*(\mathcal{L}) \equiv m(c_1(\mathcal{O}_{\mathbb{P}(E)}(1))) + \pi^* c_1(\mathcal{L}) \\ & \equiv m \left\{ c_1(\mathcal{O}_{\mathbb{P}(E)}(1)) - \frac{1}{r} \pi^* c_1(E) \right\} + \frac{m}{r} (\pi^* c_1(E) + \pi^* c_1(\mathcal{L})) \\ & \equiv m \left\{ c_1(\mathcal{O}_{\mathbb{P}(E)}(1)) - \frac{1}{r} \pi^* c_1(E) \right\} + \pi^* \left(\frac{m}{r} c_1(E) + c_1(\mathcal{L}) \right) \in \overline{\text{Eff}}^1(\mathbb{P}(E)). \end{aligned}$$

This shows that

$$\overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ x \lambda_E + \pi^* \gamma \mid x \in \mathbb{R}_{\geq 0}, \gamma \in \overline{\text{Eff}}^1(X) \right\}.$$

Moreover, if $\overline{\text{Eff}}^1(X)$ is a finite polyhedra generated by E_1, E_2, \dots, E_n , then

$$\frac{m}{r} c_1(E) + c_1(\mathcal{L}) = y_1 E_1 + y_2 E_2 + \dots + y_n E_n$$

for some $y_1, y_2, \dots, y_n \in \mathbb{R}_{\geq 0}$, say.

$$\begin{aligned} & \text{Thus } \mathcal{O}_{\mathbb{P}(E)}(m) \otimes \pi^*(\mathcal{L}) \equiv m(c_1(\mathcal{O}_{\mathbb{P}(E)}(1))) + \pi^* c_1(\mathcal{L}) \\ & \equiv m \left\{ c_1(\mathcal{O}_{\mathbb{P}(E)}(1)) - \frac{1}{r} \pi^* c_1(E) \right\} + y_1 \pi^* E_1 + y_2 \pi^* E_2 + \dots + y_n \pi^* E_n \end{aligned}$$

$$\text{Therefore, } \text{Eff}^1(\mathbb{P}(E)) \subseteq \left\{ x_0 \lambda_E + x_1 \pi^* E_1 + x_2 \pi^* E_2 + \dots + x_n \pi^* E_n \mid x_i \in \mathbb{R}_{\geq 0} \right\} \subseteq \overline{\text{Eff}}^1(\mathbb{P}(E)).$$

$$\text{Taking closure, we get } \overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ x_0 \lambda_E + x_1 \pi^* E_1 + x_2 \pi^* E_2 + \dots + x_n \pi^* E_n \mid x_i \in \mathbb{R}_{\geq 0} \right\}. \quad \square$$

Next we illustrate Theorem 4.1 through various corollaries and corresponding examples.

Corollary 4.2. *Let X be a smooth complex projective variety of dimension n with Picard number $\rho(X) = 1$ and E be a semistable vector bundle of rank r on X with $c_2(\text{End}(E)) = 0$. Then*

$$\overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ x_0 \lambda_E + x_1 (\pi^* L_X) \mid x_0, x_1 \in \mathbb{R}_{\geq 0} \right\},$$

where L_X is the numerical class of an ample generator of the Néron-Severi group $N^1(X)_{\mathbb{Z}} \simeq \mathbb{Z}$.

Proof. Let D be an effective divisor on X and $D \equiv mL_X$ for some $m \in \mathbb{Z}$. Then $L_X^{n-1} \cdot D = mL_X^n \geq 0$. As L_X is ample, by Nakai criterion for ampleness we have $L_X^n > 0$. This shows that $m \geq 0$ and $D \equiv mL_X \in \overline{\text{Eff}}^1(X)$. So $\overline{\text{Eff}}^1(X)$ is generated by L_X , and hence the result follows from Theorem 4.1. \square

Example 4.3. Let X be a smooth projective variety with Picard number $\rho(X) = 1$, and $E = \mathcal{L}_1 \oplus \mathcal{L}_2 \oplus \cdots \oplus \mathcal{L}_n$ be a rank n bundle on X such that $\mathcal{L}_1 \cdot L_X = \mathcal{L}_2 \cdot L_X = \cdots = \mathcal{L}_n \cdot L_X$ (Here L_X is the ample generator for $N^1(X)_{\mathbb{Z}}$). Then E is semistable with $c_2(\text{End}(E)) = 4c_2(E) - c_1^2(E) = 0$. Therefore,

$$\overline{\text{Eff}}^1(\mathbb{P}(\mathcal{L}_1 \oplus \mathcal{L}_2 \oplus \cdots \oplus \mathcal{L}_n)) = \left\{ x_0 \lambda_E + x_1 (\pi^* L_X) \mid x_0, x_1 \in \mathbb{R}_{\geq 0} \right\}.$$

Example 4.4. Let X be a smooth complex projective variety with Picard number 1 and $q = H^1(X, \mathcal{O}_X) \neq 0$. Then for any line bundle L on X , there is a nonsplit extension

$$0 \longrightarrow L \longrightarrow E \longrightarrow L \longrightarrow 0.$$

In this case, E is a semistable bundle of rank 2 with $c_2(\text{End}(E)) = 0$. Moreover, for any positive integer r , the vector bundles of the forms $E^{\oplus r} \oplus L$ and $E^{\oplus r}$ are examples of semistable bundles of ranks $2r + 1$ and $2r$ respectively with $c_2(\text{End}(E^{\oplus r} \oplus L)) = c_2(\text{End}(E^{\oplus r})) = 0$. In these cases, $\overline{\text{Eff}}^1(\mathbb{P}(E))$ can be computed using Corollary 4.2.

Corollary 4.5. *Let X be a smooth irreducible complex projective variety such that $\overline{\text{Eff}}^1(X)$ is a finite polyhedra with generators C_1, C_2, \dots, C_n . Let $\rho : X \longrightarrow Y$ be a nonconstant morphism onto another smooth irreducible projective variety Y . Then for any semistable bundle V on Y with $c_2(\text{End}(V)) = 0$, we have*

$$\overline{\text{Eff}}^1(\mathbb{P}_X(\rho^*V)) = \left\{ x_0 \lambda_E + x_1 \pi^* C_1 + \cdots + x_n \pi^* C_n \mid x_i \in \mathbb{R}_{\geq 0} \right\}.$$

Proof. Note that ρ^*V is semistable with $c_2(\text{End}(\rho^*V)) = \rho^*c_2(\text{End}(V)) = 0$. Therefore the result follows from Theorem 4.1. \square

We now exhibit Corollary 4.5 through some concrete examples.

Example 4.6. Let C be a general elliptic curve and $X = C \times C$ be the self product. Then X is an abelian surface and $\overline{\text{Eff}}^1(X)$ is a non-polyhedral cone (see Lemma 1.5.4 in [L07]). Let $p_i : X \longrightarrow C$

be the projection maps. For any semistable vector bundle F of rank r on C , the pullback bundle $E := p_i^*(F)$ is a semistable bundle with $c_2(\text{End}(E)) = 0$. Therefore by Theorem 4.1,

$$\overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ a\xi + b\pi^*\gamma \mid \gamma \in \overline{\text{Eff}}^1(X) \right\}.$$

In this case $\overline{\text{Eff}}^1(\mathbb{P}(E))$ is a non-polyhedral cone.

Example 4.7. Let $\rho : X = \mathbb{P}(W) \rightarrow C$ be a ruled surface on a smooth complex projective curve C defined by a normalized bundle (in the sense of Hartshorne [see [H77], Ch 5]) W . Then

$$\overline{\text{Eff}}^1(X) = \left\{ a(\zeta - \mu_{\max}(W)f) + bf \mid a, b \in \mathbb{R}_{\geq 0} \right\}$$

is a finite polyhedron generated by classes $\zeta - \mu_{\max}(W)$ and f , where $\zeta = [\mathcal{O}_X(\sigma)]$ (here σ is the normalized section such that $\mathcal{O}_{\mathbb{P}(W)}(1) \cong \mathcal{O}_X(\sigma)$, and f is the numerical equivalence class of a fibre of ρ). Let V be a semistable (resp. stable) vector bundle of rank r and degree d on C . Then the pullback bundle $E := \rho^*(V)$ is also semistable (resp. stable) on X with

$$c_2(\text{End}(E)) = c_2(\text{End}(\rho^*V)) = 2rc_2(\rho^*V) - (r-1)c_1^2(\rho^*V) = 0 - (r-1)df \cdot df = 0.$$

Consider the following commutative diagram:

$$\begin{array}{ccc} \mathbb{P}(E) = \mathbb{P}(W) \times_C \mathbb{P}(V) & \xrightarrow{\eta} & \mathbb{P}(V) \\ \downarrow \pi & & \downarrow \psi \\ X = \mathbb{P}(W) & \xrightarrow{\rho} & C \end{array}$$

One can compute $\overline{\text{Eff}}^1(\mathbb{P}(E))$ using Corollary 4.5. Infact

$$\overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ a\lambda_E + b\pi^*(\zeta - \mu_{\max}(W)f) + c\pi^*f \mid a, b, c \in \mathbb{R}_{\geq 0} \right\}.$$

For example, any indecomposable normalized rank 2 bundle W over an elliptic curve C is one of the following type (see Proposition 2.15, Chapter V, [H77]):

- Either (i) $0 \rightarrow \mathcal{O}_C \rightarrow W \rightarrow \mathcal{O}_C \rightarrow 0$,
- or (ii) $0 \rightarrow \mathcal{O}_C \rightarrow W \rightarrow \mathcal{O}_C(p) \rightarrow 0$ for some closed point $p \in C$.

In both the cases, $\deg(W) \geq 0$, and hence W is semistable.

We consider the ruled surface $\rho : X = \mathbb{P}(W) \rightarrow C$ over the elliptic curve C defined by the nonsplit extension $0 \rightarrow \mathcal{O}_C \rightarrow W \rightarrow \mathcal{O}_C \rightarrow 0$. Let us define $[C_1] := [\mathcal{O}_X(\sigma)] = [\mathcal{O}_{\mathbb{P}(W)}(1)]$, and $[C_2] := f =$ numerical class of a fibre of the map ρ . Then

$$\overline{\text{Eff}}^1(X) = \left\{ a[C_1] + b[C_2] \mid a, b \in \mathbb{R}_{\geq 0} \right\}.$$

Let V be the bundle on C which sits in the nonsplit extension

$$0 \rightarrow \mathcal{O}_C \rightarrow V \rightarrow \mathcal{O}_C(p) \rightarrow 0$$

for some closed point p in C . Then, $E := \rho^*(V)$ is a stable bundle on X with $c_2(\text{End}(E)) = 0$. Note that, in this example, the Picard number of $\mathbb{P}(E)$ is 3.

Therefore

$$\overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ x_0 \lambda_E + x_1 \pi^*([C_1]) + y_2 \pi^*([C_2]) \mid x_0, x_1, x_2 \in \mathbb{R}_{\geq 0} \right\}$$

Example 4.8. Let $\rho : X = \mathbb{P}(W) \rightarrow C$ be a ruled surface over a smooth elliptic curve C defined by the rank two bundle $W = \mathcal{O}_C \oplus \mathcal{O}_C$. Then

$$\overline{\text{Eff}}^1(X) = \left\{ a\zeta + bf \mid a, b \in \mathbb{R}_{\geq 0} \right\},$$

where $\zeta = [\mathcal{O}_{\mathbb{P}(W)}(1)] \in N^1(X)$ and f is the numerical equivalence class of a fibre of ρ .

Let

$$\rho_x : \tilde{X}_x = \text{Bl}_x(X) \rightarrow X$$

be the blow up of X at a closed point x in a section σ such that $\mathcal{O}_X(\sigma) \cong \mathcal{O}_{\mathbb{P}(W)}(1)$, and E_x be the exceptional divisor. Then we claim that

$$\overline{\text{Eff}}^1(\tilde{X}_x) = \left\{ a[C_1] + b[C_2] + c[C_3] \mid a, b, c \in \mathbb{R}_{\geq 0} \right\},$$

where $[C_1] = \rho_x^* f - E_x$, $[C_2] = \rho_x^* \zeta - E_x$ and $[C_3] = E_x$.

To prove our claim enough to show that if V is an irreducible curve in X which is not a fibre i.e. $[V] = a\zeta + bf$; $a > 0$ and $b \geq 0$, then $\text{mult}_x V = r \leq a$.

Let F be a fibre passing through x . Then $\text{mult}_x F = 1$. Hence

$$a = V \cdot F = \sum_{P \in V \cap F} (V \cdot F)_P \geq (V \cdot F)_x \geq r \text{ (see [H77], Chapter V, Proposition 1.4)}$$

Let $E = \rho_x^*(\rho^*(V))$, where V is the semistable bundle on C given by the nonsplit extension

$$0 \rightarrow \mathcal{O}_C \rightarrow V \rightarrow \mathcal{O}_C(p) \rightarrow 0$$

for some closed point $p \in C$. Then E is also semistable bundle with $c_2(\text{End}(E)) = 0$.

Note that, in this example, the Picard number of $\mathbb{P}(E)$ is 4.

Therefore, the pseudo-effective cone of divisors in the projective bundle $\pi : \mathbb{P}(E) \rightarrow \tilde{X}_x$ is given by

$$\overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ x_0 \lambda_E + \sum_{i=1}^3 x_i \pi^*([C_i]) \mid x_i \in \mathbb{R}_{\geq 0} \right\}.$$

Next we restrict our attention in finding pseudo-effective cone of $\mathbb{P}_X(E)$ over higher dimensional varieties X where $c_2(\text{End}(E)) \neq 0$. Our first result in this direction is the following.

Theorem 4.9. *Let X be a smooth irreducible complex projective surface of Picard number 1, and L_X be an ample generator of $N^1(X)_{\mathbb{R}}$. Let $E = \mathcal{O}_X \oplus \mathcal{L}$ be rank 2 bundle over X such that $\mathcal{L} \equiv nL_X$ for some positive integer $n > 0$. Then*

$$\overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ a(\xi - n\pi^*L_X) + b\pi^*L_X \mid a, b \in \mathbb{R}_{\geq 0} \right\}.$$

In particular, if $E = \mathcal{O}_{\mathbb{P}^2} \oplus \mathcal{O}_{\mathbb{P}^2}(n)$ is a rank 2 bundle on \mathbb{P}^2 for some positive integer n , then

$$\overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ a(\xi - n\pi^*H) + b\pi^*H \mid a, b \in \mathbb{R}_{\geq 0} \right\}.$$

where H is the numerical class of $\mathcal{O}_{\mathbb{P}^2}(1)$.

Proof. The Harder Narasimhan filtration of E with respect to ample divisor L_X is as follows :

$$0 \subsetneq \mathcal{L} \subsetneq E$$

with $Q_1 = E/\mathcal{L} = \mathcal{O}_X$.

Our claim is that

$$\overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ a[\mathbb{P}(Q_1)] + b\pi^*L_X \mid a, b \in \mathbb{R}_{\geq 0} \right\}.$$

Note that $[\mathbb{P}(Q_1)] = \xi - c_1(\pi^*\mathcal{L}) = \xi - n\pi^*L_X$.

Clearly,

$$\left\{ a[\mathbb{P}(Q_1)] + b\pi^*L_X \mid a, b \in \mathbb{R}_{\geq 0} \right\} \subseteq \overline{\text{Eff}}^1(\mathbb{P}(E)).$$

Note that $\mathbb{P}(E)$ is a 3-fold. Hence $L_1 \cdot L_2 \in \overline{\text{Mov}}(X)$ for any two $L_1, L_2 \in \text{Nef}^1(\mathbb{P}(E))$.

We have using [Corollary 3.5, [MR20](#)]

$$\text{Nef}^1(\mathbb{P}(E)) = \left\{ a\xi + b\pi^*L_X \mid a, b \in \mathbb{R}_{\geq 0} \right\}.$$

Also $\xi^3 - c_1(\pi^*E)\xi^2 + c_2(\pi^*E)\xi - c_3(\pi^*E) = 0$

As $c_2(E) = c_3(E) = 0$, we conclude $\xi^3 - c_1(\pi^*E)\xi^2 = \xi^2(\xi - n\pi^*L_X) = 0$.

Let $a\xi + b\pi^*L_X \in \overline{\text{Eff}}^1(\mathbb{P}(E))$. Then by the above observation, we have

$$(a\xi + b\pi^*L_X) \cdot (\pi^*L_X)^2 = a\xi(\pi^*L_X)^2 = a \geq 0,$$

and $(a\xi + b\pi^*L_X) \cdot \xi^2 = a\xi^3 + b(\pi^*L_X)\xi^2 = (an + b)(\pi^*L_X) \cdot \xi^2 \geq 0$

Since π^*L_X is nef, $(\pi^*L_X) \cdot \xi^2 \geq 0$. Thus

$$(an + b) \geq 0.$$

Now $a\xi + b\pi^*L_X = a(\xi - n\pi^*L_X) + (an + b)(\pi^*L_X)$.

This shows that

$$\overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ a(\xi - n\pi^*L_X) + b\pi^*L_X \mid a, b \in \mathbb{R}_{\geq 0} \right\}.$$

□

Theorem 4.10. *Let X be a smooth irreducible complex projective surface of Picard rank 1 with an ample generator L_X of $N^1(X)_{\mathbb{R}}$. We consider the rank 3 bundle $E = \mathcal{O}_X \oplus L \oplus M$ such that $L \equiv mL_X$ and $M \equiv nL_X$ with $m \geq n > 0$. Consider the Harder Narasimhan filtration of E with respect to L_X as follow:*

$$\mathcal{O}_X \subsetneq L \oplus M \subsetneq E$$

with $E_1 = L \oplus M$ and $Q_1 = E/E_1 = X$. The projection map $p : \mathbb{P}(E) \setminus \mathbb{P}(Q_1) \rightarrow \mathbb{P}(E_1)$ can be seen as a rational map. The indeterminacies of this rational map are resolved by blowing up $\mathbb{P}(Q_1)$. Let $\beta : \text{Bl}_{\mathbb{P}(Q_1)} \mathbb{P}(E) \rightarrow \mathbb{P}(E)$ be the blow up map. Consider the following commutative diagram.

$$\begin{array}{ccc} \mathbb{P}_Y(\mathcal{F}) = \text{Bl}_{\mathbb{P}(Q_1)} \mathbb{P}(E) & \xrightarrow{\eta} & \mathbb{P}(E_1) = \mathbb{P}(L \oplus M) = Y \\ \downarrow \beta & & \downarrow \rho \\ \mathbb{P}(E) & \xrightarrow{\pi} & X \end{array}$$

Then the cone map

$$C^{(1)} : \overline{\text{Eff}}^1(\mathbb{P}(L \oplus M)) \rightarrow \overline{\text{Eff}}^1(\mathbb{P}(E))$$

defined as follows :

$$C^{(1)}(\alpha) := \beta_* \eta^*(\alpha),$$

induces an isomorphism of $\overline{\text{Eff}}^1(\mathbb{P}(L \oplus M))$ onto $\overline{\text{Eff}}^1(\mathbb{P}(E))$.

Proof. By following the ideas of Proposition 2.4 in [F11] (See Remark 2.5 [F11]), one can find a locally free sheaf \mathcal{F} on Y such that $(\text{Bl}_{\mathbb{P}(Q_1)} \mathbb{P}(E), Y, \eta)$ is the projective bundle $\mathbb{P}_Y(\mathcal{F})$ over Y with η as the projection map. Thus we have the following commutative diagram :

$$\begin{array}{ccc} \mathbb{P}_Y(\mathcal{F}) = \text{Bl}_{\mathbb{P}(Q_1)} \mathbb{P}(E) & \xrightarrow{\eta} & \mathbb{P}(E_1) = \mathbb{P}(\mathcal{O}_{\mathbb{P}^2}(n) \oplus \mathcal{O}_{\mathbb{P}^2}(m)) = Y \\ \downarrow \beta & & \downarrow \rho \\ \mathbb{P}(E) & \xrightarrow{\pi} & X \end{array}$$

Moreover, if $\gamma := \mathcal{O}_{\mathbb{P}_Y(\mathcal{F})}(1)$, $\xi := \mathcal{O}_{\mathbb{P}(E)}(1)$ and $\xi_1 := \mathcal{O}_{\mathbb{P}(E_1)}(1)$, then

$$\gamma = \beta^* \xi, \quad \eta^* \xi_1 = \beta^* \xi - E$$

where E is numerical class of the exceptional divisor of the map β .

Note that \mathcal{F} sits in the following exact sequence:

$$0 \rightarrow \mathcal{O}_{\mathbb{P}(E_1)}(1) \rightarrow \mathcal{F} \rightarrow \rho^*(Q_1) \rightarrow 0.$$

Therefore the total Chern class

$$c(\mathcal{F}) = c(\mathcal{O}_{\mathbb{P}(E_1)}(1)) \cdot c(\rho^*Q_1) = (1 + \xi_1)\rho^*(1 + c_1(Q_1) + c_2(Q_1))$$

Therefore

$$c_1(\mathcal{F}) = \xi_1, \quad c_2(\mathcal{F}) = 0$$

Now $\gamma^2 - c_1(\eta^*\mathcal{F})\gamma + c_2(\eta^*\mathcal{F}) = 0$.

$$\implies \gamma^2 - \eta^*(\xi_1)\gamma = 0$$

$$\implies \gamma^2 - \{\gamma - E\}\gamma = 0.$$

$$\implies E \cdot \gamma = E \cdot \beta^*\xi = 0$$

Now we have $N^1(Y)_{\mathbb{R}} = \{a\xi_1 + b\rho^*L_X \mid a, b \in \mathbb{R}\}$ and $N^1(\mathbb{P}(E))_{\mathbb{R}} = \{a\xi + b\pi^*L_X \mid a, b \in \mathbb{R}\}$.

We define

$$\phi_1 : N^1(\mathbb{P}(E))_{\mathbb{R}} \longrightarrow N^1(Y)_{\mathbb{R}}$$

by

$$\phi_1(a\xi + b\pi^*L_X) = a\xi_1 + b\rho^*L_X.$$

This gives an isomorphism between real vector spaces $N^1(\mathbb{P}(E))_{\mathbb{R}}$ and $N^1(Y)_{\mathbb{R}}$.

Also we define $U_1 : N^1(Y)_{\mathbb{R}} \longrightarrow N^1(\mathbb{P}(E))_{\mathbb{R}}$ as follows :

$$U_1(\alpha) = \beta_*\eta^*(\alpha).$$

In particular, $U_1(a\xi_1 + b\rho^*L_X) = a\xi + b\pi^*L_X$. As η is flat and β is bi-rational, the above map U_1 is well-defined. We construct an inverse for U_1 .

Define

$$D_1 : N^1(\mathbb{P}(E))_{\mathbb{R}} \longrightarrow N^1(Y)_{\mathbb{R}}$$

as follows :

$$D_1(\alpha) = \eta_*(\delta \cdot \beta^*\alpha), \quad \text{where } \delta := \beta^*(\xi).$$

Note that $D_1(\xi) = \eta_*(\delta \cdot \beta^*\xi) = \eta_*(\delta \cdot (\eta^*\xi_1 + E)) = \eta_*(\delta \cdot \eta^*\xi_1 + \delta \cdot E) = \eta_*\delta \cdot \xi_1 = [Y] \cdot \xi_1$. Similarly, $D_1(\pi^*L_X) = \rho^*L_X$. This shows that the maps D_1 and ϕ_1 are the same maps.

Next we will show that the map D_1 sends effective divisor in $\mathbb{P}(E)$ to $\overline{\text{Eff}}^1(Y)$. For any effective divisor α , we can write $\beta^*(\alpha) = \alpha' + j_*\tilde{\alpha}$ for some $\tilde{\alpha}$, where α' is the strict transform under the map β and $j : E \longrightarrow \mathbb{P}_Y(\mathcal{F})$ is the canonical inclusion. (Here by abuse of notation E is also the support of E). Since δ is intersection of nef divisors and η_* maps pseudo-effective divisors to pseudo-effective divisors, it is enough to prove that $\delta \cdot j_*\tilde{\alpha} = 0$. This is true because $E \cdot \delta = 0$ and $E \cdot N(\mathbb{P}_Y(\mathcal{F})) = j_*N(E)$.

Note that $\overline{\text{Eff}}^1(L \oplus M)$ can be computed after a suitable twist by M^{-1} using Theorem 4.9 or Example 4.3 depending on $m > n$ or $m = n$ respectively. \square

Remark 1. Let E and F be two vector bundles of ranks e and f respectively on a smooth complex projective curve C . Consider the following fibre product diagram:

$$\begin{array}{ccc} \mathbb{P}(\psi^*E) = \mathbb{P}(F) \times_C \mathbb{P}(E) & \xrightarrow{\pi_2} & \mathbb{P}(E) \\ \downarrow \pi_1 & & \downarrow \\ X = \mathbb{P}(F) & \xrightarrow{\psi} & C \end{array}$$

Note that $c_2(\text{End}(\psi^*E)) = 0$ and ψ^*E is a semistable bundle if and only if E is semistable. Also $\overline{\text{Eff}}^1(\mathbb{P}(F))$ is a finitely polyhedra (see [F11]). Thus $\overline{\text{Eff}}^1(\mathbb{P}(\psi^*E))$ can be computed using Theorem 4.1 if E is semistable.

Now consider the case that E is unstable, and let

$$0 = E_d \subsetneq E_{d-1} \subsetneq \cdots \subsetneq E_1 \subset E_0 = E$$

be the Harder-Narasimhan filtrations of E . Since ψ is a smooth map, in particular it is flat and hence ψ^* is an exact functor. Also we observe that for any semistable vector bundle V on C and for any ample line bundle H on $\mathbb{P}(F)$, $\psi^*(V)$ is H -semistable with $\mu_H(\psi^*V) = \mu(V)(f_1 \cdot H^{n-1})$ and $f_1 \cdot H^{n-1} > 0$, where f_1 denotes a fiber of the map ψ . These observations immediately imply that

$$0 = \psi^*E_d \subsetneq \psi^*E_{d-1} \subsetneq \cdots \subsetneq \psi^*E_1 \subsetneq \psi^*E_0 = \psi^*E$$

is the unique Harder-Narasimhan filtration of ψ^*E with respect to any ample line bundle H on $\mathbb{P}(F)$ with $Q_1 := \psi^*E/\psi^*E_1 = \psi^*Q_1$. In this situation $\overline{\text{Eff}}^1\mathbb{P}(\psi^*E)$ is calculated in [M21] by considering the following three cases :

CASE 1 : [Lemma 4.1, [M21]] We assume that F is semistable and E is unstable with $\text{rk}(Q_1) = e - 1$.

CASE 2 : [Lemma 4.2, [M21]] We assume that both E and F are unstable with $\text{rk}(Q_1) = e - 1$ and $\text{rk}(Q'_1) = f - 1$, where

$$0 = F_h \subsetneq F_{h-1} \subsetneq F_{h-2} \subsetneq \cdots \subsetneq F_1 \subsetneq F_0 = F$$

is the Harder Narasimhan filtration for F with $Q' = F/F_1$.

CASE 3 : [Lemma 4.3, [M21]] In this case, we assume that E is unstable with $\text{rk}(Q_1) < e - 1$. We define a cone map

$$C^{(1)} : \overline{\text{Eff}}^1(\mathbb{P}(\psi^*E_1)) \longrightarrow \overline{\text{Eff}}^1(\mathbb{P}(\psi^*E))$$

which is an isomorphism. Inductively, the Harder-Narasimhan filtration of E gives the isomorphisms

$$\overline{\text{Eff}}^1(\mathbb{P}(\psi^*E_{d-1})) \simeq \cdots \simeq \overline{\text{Eff}}^1(\mathbb{P}(\psi^*E_1)) \simeq \overline{\text{Eff}}^1(\mathbb{P}(\psi^*E))$$

of cone maps. Now $\overline{\text{Eff}}^1(\mathbb{P}(\psi^*E_{d-1}))$ is computed using CASE 1.

Theorem 4.11. Let C be a smooth irreducible complex projective curve and $\rho : X = \mathbb{P}_C(W) \rightarrow C$ is a ruled surface defined by a normalized vector bundle (in the sense of Hartshorne) of rank 2 with $\mu_{\min}(W) = \deg(W)$. We denote the numerical classes of the tautological bundle $\mathcal{O}_X(1)$ and a fiber of the map ρ by η and f respectively. Let $E = \mathcal{O}_X \oplus \mathcal{L}$ be a rank 2 vector bundle on X such that \mathcal{L} is nef with $c_1(\mathcal{L}) \equiv a\eta + bf \in N^1(X)_{\mathbb{R}}$. Then

$$\overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ x(\xi - a\pi^*\eta - b\pi^*f) + y(\pi^*\eta) + z(\pi^*f) \mid x, y, z \in \mathbb{R}_{\geq 0} \right\}.$$

Proof. Firstly, we observe that

$$\text{Nef}^1(X) = \left\{ a(\eta - \mu_{\min}(W)f) + bf \mid a, b \in \mathbb{R}_{\geq 0} \right\},$$

and

$$\overline{\text{Eff}}^1(X) = \left\{ a\eta + bf \mid a, b \in \mathbb{R}_{\geq 0} \right\}.$$

Let us fix the quotient $Q_1 := E/\mathcal{L} = \mathcal{O}_X$. Our claim is that

$$\overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ x[\mathbb{P}(Q_1)] + y(\pi^*\eta) + z(\pi^*f) \mid x, y, z \in \mathbb{R}_{\geq 0} \right\}.$$

Clearly,

$$\left\{ x[\mathbb{P}(Q_1)] + y(\pi^*\eta) + z(\pi^*f) \mid x, y, z \in \mathbb{R}_{\geq 0} \right\} \subseteq \overline{\text{Eff}}^1(\mathbb{P}(E)).$$

Note that $[\mathbb{P}(Q_1)] = \xi - c_1(\pi^*\mathcal{L}) = \xi - a\pi^*\eta - b\pi^*f$.

Note that, as $\mathbb{P}(E)$ is a 3-fold, if $L_1, L_2 \in \text{Nef}^1(\mathbb{P}(E))$, then $L_1 \cdot L_2 \in \overline{\text{Mov}}(X)$.

We have using [Corollary 3.4, [MR20](#)]

$$\text{Nef}^1(\mathbb{P}(E)) = \left\{ a\xi + b\pi^*\eta + c\pi^*f \mid a, b, c \in \mathbb{R}_{\geq 0} \right\}.$$

Consider the following intersection products:

$$\xi^2(\pi^*\eta) = \mathcal{L} \cdot \eta = (a\eta + bf) \cdot \eta = a \deg(W) + b, \quad \xi^2(\pi^*f) = \mathcal{L} \cdot f = a,$$

$$\xi \cdot (\pi^*\eta) \cdot (\pi^*f) = 1, \quad \xi\pi^*(\eta^2) = \deg(W), \quad \xi(\pi^*f)^2 = 0.$$

Let $x_1\xi + x_2\pi^*\eta + x_3\pi^*f \in \overline{\text{Eff}}^1(\mathbb{P}(E))$. Then by the above observations, we have

$$(x_1\xi + x_2\pi^*\eta + x_3\pi^*f) \cdot (\pi^*\eta - \mu_{\min}(W)\pi^*f) \cdot (\pi^*f) = x_1 \geq 0,$$

and

$$(x_1\xi + x_2\pi^*\eta + x_3\pi^*f) \cdot \pi^*\eta \cdot (\pi^*\eta - \mu_{\min}(W)\pi^*f)\xi = bx_1 + x_3 \geq 0,$$

and

$$(x_1\xi + x_2\pi^*\eta + x_3\pi^*f) \cdot (\pi^*f) \cdot \xi = ax_1 + x_2 \geq 0,$$

Note that $x_1\xi + x_2\pi^*\eta + x_3\pi^*f = x_1(\xi - a\pi^*\eta - b\pi^*f) + (ax_1 + x_2)(\pi^*\eta) + (bx_1 + x_3)(\pi^*f)$.

This shows that

$$\overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ x(\xi - a\pi^*\eta - b\pi^*f) + y\pi^*\eta + z\pi^*f \mid x, y, z \in \mathbb{R}_{\geq 0} \right\}.$$

□

Example 4.12. Let W be the semistable bundle of rank 2 on an elliptic curve C which sits in the following non-split extension:

$$0 \longrightarrow \mathcal{O}_C \longrightarrow W \longrightarrow \mathcal{O}_C.$$

Let $\rho : X = \mathbb{P}_C(W) \longrightarrow C$ be the ruled surface defined by this normalized bundle with $\deg(W) = 0$. We denote the numerical classes of the tautological bundle $\mathcal{O}_X(1)$ and a fiber of the map ρ by η and f respectively. Let $E = \mathcal{O}_{\mathbb{P}(W)} \oplus \mathcal{L}$ be a rank 2 bundle, where \mathcal{L} is nef line bundle on $\mathbb{P}_C(W)$ such that $c_1(\mathcal{L}) \equiv a\eta + bf$. Then by Theorem 4.11

$$\overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ x(\xi - a\pi^*\eta - b\pi^*f) + y\pi^*\eta + z\pi^*f \mid x, y, z \in \mathbb{R}_{\geq 0} \right\}.$$

Note that $c_2(\text{End}(E)) = 4c_2(E) - c_1^2(E) = 2ab \neq 0$ for suitable choices of a, b . This shows that E is not a pullback of some rank 2 bundle on C . Thus $\overline{\text{Eff}}^1(\mathbb{P}(E))$ can not be computed using strategy mentioned in Remark 1.

Example 4.13. Let $\rho : \mathbb{F}_n = \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-n)) \longrightarrow \mathbb{P}^1$ be an Hirzebruch surface for some positive integer $n > 0$. We denote the numerical class of the tautological bundle $\mathcal{O}_{\mathbb{F}_n}(1)$ and a fiber of the map ρ by τ and f respectively. Let $E = \mathcal{O}_{\mathbb{F}_n} \oplus \mathcal{L}$ be a rank 2 bundle on \mathbb{F}_n such that $c_1(\mathcal{L}) \equiv \tau + nf \in N^1(\mathbb{F}_n)_{\mathbb{R}}$. Thus \mathcal{L} is nef. Hence by Theorem 4.11

$$\overline{\text{Eff}}^1(\mathbb{P}(E)) = \left\{ a(\xi - \pi^*\tau - n\pi^*f) + b\pi^*\tau + c\pi^*f \mid a, b, c \in \mathbb{R}_{\geq 0} \right\}.$$

Remark 2. The nefness condition of the line bundle \mathcal{L} in Theorem 4.11 is a technical one so that $\text{Nef}^1(\mathbb{P}(E))$ can be computed using Corollary 3.4 in [MR20] easily. The pseudo-effective cone $\overline{\text{Eff}}^1(\mathbb{P}(E))$ can still be computed without the assumption of nefness of \mathcal{L} . However, the computations in those cases will be complicated in general.

5. VOLUME FUNCTION OF FIBER PRODUCT OF PROJECTIVE BUNDLES

A sequence of Borel probability measure $\{\nu_n\}_{n \geq 1}$ is said to converge vaguely to a Borel measure ν if for any continuous function f on \mathbb{R} of compact support, one has

$$\int f(x)\nu(dx) = \lim_{n \rightarrow \infty} \int f(x)\nu_n(dx).$$

We recall the following theorem from [C11]. We denote the function field of a smooth irreducible complex projective variety C by $K := k(C)$.

Theorem 5.1. *Let $\pi : X \rightarrow C$ be a projective and flat morphism, and L be an arbitrary line bundle on X such that L_K is big. Then*

(1) *The sequence of Borel probability measures $(T_{\frac{1}{n}} \nu_{\pi_* (L^{\otimes n})})_{n \geq 1}$ converges vaguely to a Borel measure ν_L^π .*

(2) *the following equality holds*

$$\text{vol}_X(L) = \dim(X) \text{vol}_{X_K}(L_K) \int x_+ \nu_L^\pi(dx),$$

where $x_+ = \max\{x, 0\}$.

Let G be a vector bundle on a smooth irreducible complex projective curve C having Harder-Narasimhan filtration

$$0 = G_r \subsetneq G_{r-1} \subsetneq \cdots \subsetneq G_0 = G$$

with successive slopes μ_i . For a line bundle M of degree a on C , we then have

$$0 = G_r \otimes M \subsetneq G_{r-1} \otimes M \subsetneq \cdots \subsetneq G_0 \otimes M = G \otimes M$$

is the Harder-Narasimhan filtration of $G \otimes M$ having successive slopes $\mu_i + a$. This shows that

$$\nu_{G \otimes M} = \tau_a \nu_E.$$

Here τ_a is defined as

$$\int f(x) \tau_a \nu dx = \int f(x+a) \nu dx.$$

Now recall the following commutative fibre product diagram:

$$\begin{array}{ccc} X = \mathbb{P}(\mathcal{E}) = \mathbb{P}(F) \times_C \mathbb{P}(E) = \mathbb{P}(\psi^* E) & \xrightarrow{\pi_2} & \mathbb{P}(E) \\ \downarrow \pi_1 & & \downarrow \\ \mathbb{P}(F) & \xrightarrow{\psi} & C \end{array}$$

Recall that $\pi = \psi \circ \pi_1$ and $\mathcal{E} = \psi^* E$. Let $L = \mathcal{O}_{\mathbb{P}(\mathcal{E})}(m) \otimes \pi_1^* \mathcal{O}_{\mathbb{P}(F)}(l) \otimes \pi_1^* \psi^* \mathcal{M}$ be a line bundle on X for some line bundle \mathcal{M} of degree a on C such that L_K is big. In particular, if L is big, then L_K is also big.

$$\begin{aligned} & \text{Note that } \pi_* (\mathcal{O}_{\mathbb{P}(\mathcal{E})}(m) \otimes \pi_1^* \mathcal{O}_{\mathbb{P}(F)}(l) \otimes \pi_1^* \psi^* \mathcal{M})^{\otimes n} \\ &= \psi_* (\pi_1)_* ((\mathcal{O}_{\mathbb{P}(\mathcal{E})}(m) \otimes \pi_1^* \mathcal{O}_{\mathbb{P}(F)}(l) \otimes \pi_1^* \psi^* \mathcal{M})^{\otimes n}) \\ &= \psi_* (\pi_1)_* (\mathcal{O}_{\mathbb{P}(\mathcal{E})}(mn) \otimes \pi_1^* \mathcal{O}_{\mathbb{P}(F)}(ln) \otimes \pi_1^* \psi^* (\mathcal{M}^{\otimes n})). \\ &= \psi_* (\text{Sym}^{mn}(\psi^* E) \otimes \mathcal{O}_{\mathbb{P}(F)}(ln) \otimes \psi^* (\mathcal{M}^{\otimes n})), \\ &= \text{Sym}^{mn}(E) \otimes \text{Sym}^{ln}(F) \otimes \mathcal{M}^{\otimes n}. \end{aligned}$$

Thus by the previous observations, we have

$$\begin{aligned} & \mathcal{V}_{\pi_*}(\mathcal{O}_{\mathbb{P}(\mathcal{E})}(m) \otimes \pi_1^* \mathcal{O}_{\mathbb{P}(F)}(l) \otimes \pi_1^* \psi^* \mathcal{M})^{\otimes n} \\ &= \mathcal{V}_{\text{Sym}^{mn}(E) \otimes \text{Sym}^{ln}(F) \otimes \mathcal{M}^{\otimes n}} \\ &= \tau_{na} \mathcal{V}_{\text{Sym}^{mn}(E) \otimes \text{Sym}^{ln}(F)} \end{aligned}$$

One also has

$$T_\varepsilon \tau_a \mathcal{V} = \tau_{a\varepsilon} T_\varepsilon \mathcal{V}.$$

$$\text{Thus } T_{\frac{1}{n}} \mathcal{V}_{\pi_* L^{\otimes n}} = T_{\frac{1}{n}} \tau_{an} \mathcal{V}_{\text{Sym}^{mn}(E) \otimes \text{Sym}^{ln}(F)} = \tau_a T_{\frac{1}{n}} \mathcal{V}_{\text{Sym}^{mn}(E) \otimes \text{Sym}^{ln}(F)}.$$

Note that by definition $\mathcal{V}_{\mathcal{O}_{\mathbb{P}(\mathcal{E})}(m) \otimes \pi_1^* \mathcal{O}_{\mathbb{P}(F)}(l)}^\pi$ is the vague limit of

$$T_{\frac{1}{n}} \mathcal{V}_{\text{Sym}^{mn}(E) \otimes \text{Sym}^{ln}(F)} = T_{\frac{1}{n}} \mathcal{V}_{\pi_* (\mathcal{O}_{\mathbb{P}(\mathcal{E})}(m) \otimes \pi_1^* \mathcal{O}_{\mathbb{P}(F)}(l))^{\otimes n}}.$$

Therefore

$$\mathcal{V}_L^\pi = \tau_a \mathcal{V}_{\mathcal{O}_{\mathbb{P}(\mathcal{E})}(m) \otimes \pi_1^* \mathcal{O}_{\mathbb{P}(F)}(l)}^\pi.$$

Thus by Theorem 5.1 we have

$$\text{vol}_X(L) = \dim(X) \text{vol}_{X_K}(L_K) \int x_+ \mathcal{V}_L^\pi(dx) = \dim(X) \text{vol}_{X_K}(L_K) \int (x+a)_+ \mathcal{V}_{\mathcal{O}_{\mathbb{P}(\mathcal{E})}(m) \otimes \pi_1^* \mathcal{O}_{\mathbb{P}(F)}(l)}^\pi dx.$$

6. HARDER-NARASIMHAN FILTRATION OF TENSOR PRODUCT

In this section we determine the Harder-Narasimhan filtration of $E \otimes F$ in terms of the Harder-Narasimhan filtrations of E and F . Recall that

$$(3) \quad 0 = E_d \subsetneq E_{d-1} \subsetneq \cdots \subsetneq E_1 \subsetneq E_0 = E$$

and

$$(4) \quad 0 = F_h \subsetneq F_{h-1} \subsetneq F_{h-2} \subsetneq \cdots \subsetneq F_1 \subsetneq F_0 = F$$

are the Harder-Narasimhan filtrations for E and F respectively. We fix the notations $Q_i = E_i/E_{i+1}$ for each $i \in \{0, 1, 2, \dots, d-1\}$ and $Q'_j = F_j/F_{j+1}$ for each $j \in \{0, 1, 2, \dots, h-1\}$.

Let $J_1 = \{0, 1, 2, \dots, d-1\}$ and $J_2 = \{0, 1, 2, \dots, h-1\}$. Consider the set $\Theta = J_1 \times J_2$. We define a partial order on Θ as follows:

$$(a, b) \leq (c, d) \iff a \leq c, b \leq d.$$

We say a subset $A \subseteq \Theta$ is saturated if $\alpha \in A, \beta \in \Theta$ and $\beta \geq \alpha \implies \beta \in A$. For a subset $A \subseteq \Theta$ we define the saturation of A , denoted by \overline{A} as follows:

$$\overline{A} = \{\beta \in \Theta \mid \exists \alpha \in A \text{ such that } \beta \geq \alpha\}.$$

For a nonempty subset $A \subseteq \Theta$ and $\alpha = (a, b) \in A$, we define

$$(5) \quad \mathfrak{S}_\alpha := E_a \otimes F_b \text{ and } \mathfrak{S}_A := \bigoplus_{\alpha \in A} \mathfrak{S}_\alpha.$$

Then $\mathfrak{S}_A = \mathfrak{S}_{\overline{A}}$. Also for any two subsets $A_1 \subseteq A_2 \subseteq \Theta$, we have $\mathfrak{S}_{A_1} \subseteq \mathfrak{S}_{A_2}$.

For any $\alpha = (a, b) \in \Theta$, we define

$$(6) \quad \mathcal{Q}_\alpha = Q_a \otimes Q'_b.$$

Since the ground field \mathbb{C} has characteristic 0, we have \mathcal{Q}_α is a semistable bundle for each $\alpha \in \Theta$.

Proposition 6.1. *Assume that $A \subseteq \Theta$ is non-empty and saturated, and that A'' is a subset of A consisting of maximal elements. Then $A' := A \setminus A''$ is also saturated. Furthermore, one has an isomorphism $\mathfrak{S}_A/\mathfrak{S}_{A'} \cong \bigoplus_{\alpha \in A''} \mathcal{Q}_\alpha$.*

Proof. Locally in Zariski topology, the Harder-Narasimhan filtrations split. Therefore locally we have the required isomorphism. These natural isomorphisms do not depend on the choice of the splitting, and therefore we have the isomorphism $\mathfrak{S}_A/\mathfrak{S}_{A'} \cong \bigoplus_{\alpha \in A''} \mathcal{Q}_\alpha$. \square

For any $\alpha = (a, b) \in \Theta$, let $\mu_\alpha = \mu(Q_a) + \mu(Q'_b)$. With this notation, μ_α is the slope of \mathcal{Q}_α , where \mathcal{Q}_α is defined as in (6). Denote Σ is the set of all real numbers of the form μ_α , where α takes through vectors in Θ . Note that, from the two Harder-Narasimhan filtrations (3) and (4), we have $\alpha \leq \beta$ implies $\mu_\alpha \leq \mu_\beta$, and $\alpha \not\leq \beta$ implies $\mu_\alpha \not\leq \mu_\beta$.

Theorem 6.2. *The set Σ identifies with that of successive slopes of $E \otimes F$. Furthermore, suppose that the elements in Σ are ordered as*

$$v_0 < v_1 < v_2 < \cdots < v_{m-1},$$

then

$$(7) \quad 0 = \mathfrak{S}_{A_m} \subsetneq \mathfrak{S}_{A_{m-1}} \subsetneq \cdots \subsetneq \mathfrak{S}_{A_1} \subsetneq \mathfrak{S}_{A_0}$$

is the Harder-Narasimhan filtration of $E \otimes F$, where

$$A_j = \left\{ \alpha \in \Theta \mid \mu_\alpha \geq v_j \right\} \subseteq \Theta$$

for each $j \in \{0, 1, \dots, m-1\}$ and $A_m = \emptyset$.

Proof. We have $A_j \setminus A_{j+1} = \{\alpha \mid \mu_\alpha = v_j\}$. Moreover if $\mu_\alpha = v_j$, then α is a maximal element of A_j . Therefore by the previous Proposition 6.1 we get

$$\mathfrak{S}_{A_j} \setminus \mathfrak{S}_{A_{j+1}} \cong \bigoplus_{\alpha \in A_j \setminus A_{j+1}} \mathcal{Q}_\alpha$$

is semistable of slope v_j . \square

7. COMPUTATION OF THE VOLUME FUNCTION

7.1. **Harder Narasimhan Filtration for $\text{Sym}^{mn}(E)$** : Recall that

$$(8) \quad 0 = E_d \subsetneq E_{d-1} \subsetneq E_{d-2} \subsetneq \cdots \subsetneq E_1 \subsetneq E_0 = E$$

is the Harder-Narasimhan filtration of E , and $Q_i = E_i/E_{i+1}$ for each i . Let $\mu_i = \mu(Q_i)$ so that we have

$$\mu_{d-1} > \mu_{d-2} > \cdots > \mu_1 > \mu_0.$$

Let $\Phi = \{0, 1, \dots, d-1\}^{mn}$. Then S^{mn} acts on Φ . Let $\tilde{\Phi} := \Phi/S^{mn}$. Note that the elements in $\tilde{\Phi}$ are in bijective correspondence with the set of partitions of mn into sum of d non-negative integers $(a_0, a_1, \dots, a_{d-1})$, i.e. $\sum_{i=0}^{d-1} a_i = mn$. Any element $[\alpha]$ in $\tilde{\Phi}$ will be denoted by $(a_0, a_1, \dots, a_{d-1})$ under this correspondence. For such a class $[\alpha]$, we use the symbol $\tilde{Q}_{[\alpha]}(E)$ to denote the tensor product $\bigotimes_{i=0}^{d-1} \text{Sym}^{a_i} Q_i$.

Let $A \subseteq \Phi$ be such that A is invariant under the action of S^{mn} . Then we have a map

$$\theta : E_A \longrightarrow \text{Sym}^{mn} E.$$

Define $S_A E := \text{Im}(\theta)$. This gives $E_A \twoheadrightarrow S_A E$ and hence $E_A/S^{mn} \cong S_A E$.

Let Σ_{mn} be the set of all μ_α for $\alpha \in \Phi$. We order the elements in Σ_{mn} as follows:

$$v_0 < v_1 < \cdots < v_{k-1}.$$

Let $A_j = \{\alpha \mid \mu_\alpha \geq v_j\}$ for $j \in \{0, 1, \dots, k-1\}$. We further assume that $A_k = \emptyset$.

Then

$$(9) \quad 0 = S_{A_k} E \subsetneq S_{A_{k-1}} E \subsetneq \cdots \subsetneq S_{A_1} E \subsetneq S_{A_0} E = S_A E$$

is the Harder-Narasimhan filtration of $\text{Sym}^{mn}(E)$ such that

$$S_{A_j} E / S_{A_{j+1}} E \cong \bigoplus_{[\alpha] \in (A_j \setminus A_{j+1}) / S^{mn}} \tilde{Q}_{[\alpha]}(E)$$

are semistable bundles of slope v_j .

We also have the expression for the rank R_j of the successive quotients. Take an arbitrary class $[\alpha]$ in $\tilde{\Phi}$ which correspondence to the partition $\underline{a} = (a_i)_{i=0}^{d-1}$ of mn . The value $\mu_\alpha = \sum_{i=0}^{d-1} a_i \mu_i$. For each $i \in \{0, 1, \dots, d-1\}$, let r_i be the rank of Q_i . Then the rank of $\tilde{Q}_{[\alpha]}(E)$ is just

$$r_{\underline{a}} := \prod_{i=0}^{d-1} \binom{a_i + r_i - 1}{r_i - 1}.$$

Thus we obtain

$$R_j = \text{rk}(S_{A_j}E/S_{A_{j+1}}E) = \sum_{\underline{a}=(a_i)_{i=0}^{d-1} \in \mathbb{N}^d, \sum_{i=0}^{d-1} a_i = mn, \sum_{i=0}^{d-1} a_i \mu_i = v_j} r_{\underline{a}}.$$

7.2. Harder Narasimhan Filtration for $\text{Sym}^{ln}(F)$: Recall that

$$(10) \quad 0 = F_h \subsetneq F_{h-1} \subsetneq F_{h-2} \subsetneq \cdots \subsetneq F_1 \subsetneq F_0 = F$$

is the Harder-Narasimhan filtration of F , and $Q'_j = F_j/F_{j+1}$ for each j with rank $\text{rk}(Q'_j) = r'_j$. Let $\mu'_j = \mu(Q'_j)$ so that we have

$$\mu'_{h-1} > \mu'_{h-2} > \cdots > \mu'_1 > \mu'_0.$$

Similar steps can be taken to get the Harder-Narasimhan filtration for $\text{Sym}^{ln}(F)$, and the rank R'_j for the succesive slopes of the Harder-Narasimhan filtration. Let

$$(11) \quad 0 = S_{A'_p}F \subsetneq S_{A'_{p-1}}F \subsetneq \cdots \subsetneq S_{A'_1}F \subsetneq S_{A'_0}F = S'_A F$$

be the Harder-Narasimhan filtration of $\text{Sym}^{ln}(F)$ with slopes $v'_0 < v'_1 < \cdots < v'_{p-1}$.

Then

$$R'_k = \text{rk}(S_{A'_k}F/S_{A'_{k+1}}F) = \sum_{\underline{a}'=(a'_i)_{i=0}^{h-1} \in \mathbb{N}^h, \sum_{i=0}^{h-1} a'_i = ln, \sum_{i=0}^{h-1} a'_i \mu'_i = v'_k} r_{\underline{a}'}$$

where

$$r_{\underline{a}'} = \prod_{j=0}^{h-1} \binom{a'_j + r'_j - 1}{r'_j - 1}.$$

7.3. Harder Narasimhan Filtration for $\text{Sym}^{mn}(E) \otimes \text{Sym}^{ln}(F)$: We will use the notations and conventions used in Section 6 from now on to get the Harder-Narasimhan filtration of the tensor product $\text{Sym}^{mn}(E) \otimes \text{Sym}^{ln}(F)$.

Recall that

$$0 = S_{A_k}E \subsetneq S_{A_{k-1}}E \subsetneq \cdots \subsetneq S_{A_1}E \subsetneq S_{A_0}E = S_A E$$

and

$$0 = S_{A'_p}F \subsetneq S_{A'_{p-1}}F \subsetneq \cdots \subsetneq S_{A'_1}F \subsetneq S_{A'_0}F = S'_A F$$

are the Harder Narasimhan filtrations of $\text{Sym}^{mn}(E)$ and $\text{Sym}^{ln}(F)$ respectively as in (9) and (11).

Let $J_1 = \{0, 1, 2, \dots, k-1\}$ and $J_2 = \{0, 1, 2, \dots, p-1\}$. Consider the product $\Theta = J_1 \times J_2$.

Then for $\alpha = (a, b) \in \Theta$, we have

$$\mathfrak{S}_\alpha = S_{A_a}E \otimes S_{A'_b}F.$$

We denote $\mathcal{Q}_\alpha = (S_{A_a}E/S_{A_{a+1}}E) \otimes (S_{A'_b}F/S_{A'_{b+1}}F)$. Then \mathcal{Q}_α is semistable.

Let

$$\Sigma = \left\{ \mu_\alpha \mid \alpha \in \Theta \right\},$$

where $\mu_\alpha := \mu(\mathcal{Q}_\alpha) = \mu(S_{A_a}E/S_{A_{a+1}}E) + \mu(S_{A'_b}F/S_{A'_{b+1}}F) = v_a + v'_b$ for $\alpha = (a, b) \in \Theta$.

We order the elements of Σ as follows:

$$w_0 < w_1 < \cdots < w_{q-1}.$$

Let $B_j = \{\alpha \in \Theta \mid \mu_\alpha \geq w_j\}$ for $j \in \{0, 1, \dots, q-1\}$, with $B_q = \emptyset$. Then by Theorem 6.2 we have

$$(12) \quad \mathfrak{S}_{B_{q-1}} \subsetneq \cdots \subsetneq \mathfrak{S}_{B_1} \subsetneq \mathfrak{S}_{B_0}$$

is the Harder-Narasimhan filtration of $\text{Sym}^{mn}(E) \otimes \text{Sym}^{ln}(F)$ such that $\mathfrak{S}_{B_j}/\mathfrak{S}_{B_{j+1}} \cong \bigoplus_{\alpha \in B_j \setminus B_{j+1}} \mathcal{Q}_\alpha$.

Note that $\text{rk}(\mathcal{Q}_\alpha) = R_a \cdot R'_b$, where $\alpha = (a, b) \in \Theta$.

$$\text{Therefore } R_j'' := \text{rk}(\mathfrak{S}_{B_j}/\mathfrak{S}_{B_{j+1}}) = \bigoplus_{\alpha=(a,b) \in B_j \setminus B_{j+1}} R_a \cdot R'_b = \bigoplus_{\mu_\alpha=w_j} R_a \cdot R'_b = \bigoplus_{v_a+v'_b=w_j} R_a \cdot R'_b.$$

Thus

$$\nu_{\text{Sym}^{mn} E \otimes \text{Sym}^{ln} F} = \frac{1}{\text{rk}(\text{Sym}^{mn} E \otimes \text{Sym}^{ln} F)} \sum_{j=0}^{q-1} R_j'' \delta_{w_j}.$$

Next we compute the term $\sum_{j=0}^{q-1} R_j'' \delta_{w_j}$. Recall $\text{rk}(E) = e$ and $\text{rk}(F) = f$.

Consider the vector $\underline{s} := (s_1, s_2, \dots, s_e, s'_1, s'_2, \dots, s'_f) \in \mathbb{R}^{e+f}$ such that the value μ_i appear exactly $\text{rk}(E_i/E_{i+1})$ times in the first e coordinates and the value μ'_j appears exactly $\text{rk}(F_j/F_{j+1})$ times in the last f coordinates.

Recall that

$$R_a = \text{rk}(S_{A_a}E/S_{A_{a+1}}E) = \sum_{\underline{a}=(a_i)_{i=0}^{d-1} \in \mathbb{N}^d, \sum_{i=0}^{d-1} a_i=mn, \sum_{i=0}^{d-1} a_i \mu_i=v_a} r_{\underline{a}} = \sum_{\underline{b}=(b_i)_{i=1}^e \in \mathbb{N}^e, \sum_{i=1}^e b_i=mn, \sum_{i=1}^e b_i s_i=v_a} 1.$$

Similarly

$$R'_b = \text{rk}(S_{A'_b}F/S_{A'_{b+1}}F) = \sum_{\underline{a}'=(a'_i)_{i=0}^{h-1} \in \mathbb{N}^h, \sum_{i=0}^{h-1} a'_i=ln, \sum_{i=0}^{h-1} a'_i \mu'_i=v'_b} r_{\underline{a}'} = \sum_{\underline{b}'=(b'_i)_{i=1}^f \in \mathbb{N}^f, \sum_{i=1}^f b'_i=ln, \sum_{i=1}^f b'_i s'_i=v'_b} 1.$$

Let $\Delta \times \Delta' \subset \mathbb{R}^{e+f}$ be the set, where simplexes Δ and Δ' are defined as follows:

$$\Delta = \left\{ (x_1, x_2, \dots, x_e) \in \mathbb{R}^e \mid 0 \leq x_i \leq 1, \sum_{i=1}^e x_i = 1 \right\},$$

$$\Delta' = \left\{ (x'_1, x'_2, \dots, x'_f) \in \mathbb{R}^f \mid 0 \leq x'_j \leq 1, \sum_{i=1}^f x'_j = 1 \right\}.$$

Thus $\nu_{\text{Sym}^{mn}(E) \otimes \text{Sym}^{ln}(F)}$

$$= \frac{1}{\text{rk}(\text{Sym}^{mn} E) \cdot \text{rk}(\text{Sym}^{ln} F)} \sum_{(\underline{b}, \underline{b}') \in \mathbb{N}^{e+f}, \sum_{i=1}^e b_i = mn, \sum_{i=1}^f b'_i = ln} \delta_{b_1 s_1 + b_2 s_2 + \dots + b_e s_e + b'_1 s'_1 + b'_2 s'_2 + \dots + b'_f s'_f}$$

Theorem 7.1. *Let η be the Lebesgue measure on $\Delta \times \Delta'$ normalized such that $\eta(\Delta \times \Delta') = 1$.*

Then $T_{\frac{1}{n}} \nu_{\text{Sym}^{mn}(E) \otimes \text{Sym}^{ln}(F)}$ converges vaguely to $\phi_{\underline{s}_} \eta$, where*

$$\phi_{\underline{s}} : \Delta \times \Delta' \longrightarrow \mathbb{R}$$

such that

$$(x_1, x_2, \dots, x_e, x'_1, \dots, x'_f) \mapsto \sum_{i=1}^e s_i x_i + \sum_{j=1}^f s'_j x'_j.$$

Proof. Note that $T_{\frac{1}{n}} \nu_{\text{Sym}^{mn}(E) \otimes \text{Sym}^{ln}(F)}$

$$\begin{aligned} &= T_{\frac{1}{n}} \left[\frac{1}{\text{rk}(\text{Sym}^{mn} E) \cdot \text{rk}(\text{Sym}^{ln} F)} \sum_{(\underline{b}, \underline{b}') \in \mathbb{N}^{e+f}, \sum_{i=1}^e b_i = mn, \sum_{i=1}^f b'_i = ln} \delta_{b_1 s_1 + b_2 s_2 + \dots + b_e s_e + b'_1 s'_1 + b'_2 s'_2 + \dots + b'_f s'_f} \right] \\ &= T_{\frac{1}{n}} \left[\frac{1}{\text{rk}(\text{Sym}^{mn} E) \cdot \text{rk}(\text{Sym}^{ln} F)} \sum_{(\underline{b}, \underline{b}') \in (\frac{1}{mn} \mathbb{N}^e) \times (\frac{1}{ln} \mathbb{N}^f), \sum_{i=1}^e b_i = 1, \sum_{i=1}^f b'_i = 1} \delta_{b_1 s_1 + b_2 s_2 + \dots + b_e s_e + b'_1 s'_1 + b'_2 s'_2 + \dots + b'_f s'_f} \right] \\ &= T_{\frac{1}{n}} \left[\frac{1}{\text{rk}(\text{Sym}^{mn} E) \cdot \text{rk}(\text{Sym}^{ln} F)} \sum_{(\underline{b}, \underline{b}') \in (\frac{1}{mn} \mathbb{N}^e \cap \Delta) \times (\frac{1}{ln} \mathbb{N}^f \cap \Delta')} \delta_{(\underline{b}, \underline{b}')} \right]. \end{aligned}$$

Thus $T_{\frac{1}{n}} \nu_{\text{Sym}^{mn}(E) \otimes \text{Sym}^{ln}(F)}$ converges vaguely to $\phi_{\underline{s}_*} \eta$. □

Theorem 7.2. *Let $L = \mathcal{O}_{\mathbb{P}(E)}(m) \otimes \pi_1^* \mathcal{O}_{\mathbb{P}(F)}(l) \otimes \pi_1^* \psi^* \mathcal{M}$ be a line bundle on $X = \mathbb{P}(F) \times_C \mathbb{P}(E)$ for some line bundle \mathcal{M} of degree a on C such that L_K is big. Then*

$$\text{vol}_X(L) = \dim(X) \text{vol}_{X_K}(L_K) \int (x + a)_+ \phi_{\underline{s}_*} \eta dx,$$

where $x_+ = \max\{x, 0\}$.

Proof. By Theorem 5.1 one has

$$\begin{aligned} \text{vol}_X(L) &= \text{vol}_X(\mathcal{O}_{\mathbb{P}(E)}(m) \otimes \pi_1^* \mathcal{O}_{\mathbb{P}(F)}(l) \otimes \pi_1^* \psi^* \mathcal{M}) \\ &= \dim(X) \text{vol}_{X_K}(L_K) \int x_+ \nu_L^\pi(dx) \\ &= \dim(X) \text{vol}_{X_K}(L_K) \int (x + a)_+ \nu_{\mathcal{O}_{\mathbb{P}(E)}(m) \otimes \pi_1^* \mathcal{O}_{\mathbb{P}(F)}(l)}^\pi dx \\ &= \dim(X) \text{vol}_{X_K}(L_K) \int (x + a)_+ \phi_{\underline{s}_*} \eta dx. \end{aligned}$$

□

Remark 3. One can compute the volume function on $\mathbb{P}_C(E_1) \times_C \mathbb{P}_C(E_2) \times_C \cdots \times_C \mathbb{P}_C(E_n)$ for finitely many vector bundles E_1, E_2, \dots, E_n over a smooth irreducible projective curve C defined over a field of characteristic zero by a similar approach.

8. ACKNOWLEDGEMENT

The authors would like to thank Krishna Hanumanthu and Nabanita Ray for many fruitful discussions. The first author is supported financially by SERB-NPDF fellowship (File no : PDF/2021/00028). The first author also like to thank IIT Bombay for its hospitality where the work is initiated.

REFERENCES

- [BB08] Indranil Biswas and Ugo Bruzzo, *On Semistable Principal Bundles over a Complex Projective Manifold*, International Mathematics Research Notices, Article ID rnn035, (2008).
- [BDPP13] S. Boucksom, J.P. Demailly, M. Păun and T Peternell, *The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension*, Journal of Algebraic Geometry. 22(2), (2013) 201-248.
- [BHP14] Indranil Biswas, Amit Hogadi and A.J.Parameswaran, *Pseudo-effective cone of Grassmann Bundles over a curve*, Geom Dedicata, 172, (2014) 69-77.
- [C11] Huayi Chen, *Computing volume function on projective bundle over a curve*, RIMS Kôkyûroku **1745** (2011), 169-182.
- [ELMNP05] Lawrence Ein, Robert Lazarsfeld, Mircea Mustatã, Michael Nakamaye and Mihnea Popa, *Asymptotic invariants of line bundles*, Pure Appl. Math. Q.1 (2005), no. 2, 379-403.
- [F94] T. Fujita, *Approximating Zariski decomposition of big line bundles*, Kodai Mathematical Journal **17** (1994), no. 1, p. 1-3.
- [F11] Mihai Fulger, *The cones of effective cycles on projective bundles over curves*, Math.Z., **269**, (2011) 449-459.
- [F98] William Fulton. *Intersection Theory*, Second Edition, Springer, (1998).
- [H77] Robin Hartshorne, *Algebraic Geometry*, Graduate Text in Mathematics, Springer, (1977).
- [L07] Robert Lazarsfeld, *Positivity in Algebraic Geometry, Volume I*, Springer, (2007).
- [L11] R. K. Lazarsfeld, *Positivity in Algebraic Geometry I*, A Series of Modern Surveys in Mathematics **48** (Springer-Verlag Berlin Heidelberg).

- [MR20] Snehajit Misra and Nabanita Ray, *Nef cones of projective bundles over surfaces and Seshadri constants*, to appear in Osaka Journal of Mathematics.
- [M21] Snehajit Misra, *Pseudo-effective cones of projective bundles and weak Zariski decomposition*, European Journal of Mathematics, **7**, 1438-1457 (2021).
- [Miy87] Yoichi Miyaoka, *The Chern classes and Kodaira dimension of a minimal variety*, Algebraic geometry, Sendai, 1985, Adv. Stud. Pure Math., vol. 10, North-Holland, Amsterdam, 1987, pp. 449-476.
- [N99] Noboru Nakayama, *Normalized Tautological divisors of semi-stable vector bundles*, (Japanese) Free resolutions of coordinate rings of projective varieties and related topics (Japanese) (Kyoto, 1998). Sūrikaisekikenkyūsho Kōkyūroku No. **1078**(1999), 167-173.
- [T07] S. Takagi, *Fujita's approximation theorem in positive characteristics*, Journal of Mathematics of Kyoto University **47** (2007), no. 1, p. 179-202.
- [W05] Wolfe Alexandre, *Asymptotic invariants of graded systems of ideals and linear systems on projective bundles*, Thesis (Ph.D.)-University of Michigan. 2005. 102 pp. ISBN: 978-0542-30309-8.

CHENNAI MATHEMATICAL INSTITUTE, H1 SIPCOT IT PARK, SIRUSERI, KELAMBAKKAM 603103, INDIA.
Email address, Snehajit Misra: snehajitm@cmi.ac.in

TATA INSTITUTE OF FUNDAMENTAL RESEARCH (TIFR), HOMI BHABHA ROAD, MUMBAI 400005, INDIA.
Email address, Anoop Singh: anoops@math.tifr.res.in