

LOGARITHMIC  $A$ -HYPERGEOMETRIC SERIES II

GO OKUYAMA AND MUTSUMI SAITO

ABSTRACT. In this paper, following [6], we continue to develop the perturbing method of constructing logarithmic series solutions to a regular  $A$ -hypergeometric system.

Fixing a fake exponent of an  $A$ -hypergeometric system, we consider some spaces of linear partial differential operators with constant coefficients. Comparing these spaces, we construct a fundamental system of series solutions with the given exponent by the perturbing method. In addition, we give a sufficient condition for a given fake exponent to be an exponent. As important examples of the main results, we give fundamental systems of series solutions to Aomoto-Gel'fand systems and to Lauricella's  $F_C$  systems with special parameter vectors, respectively.

## 1. INTRODUCTION

Let  $A = (\mathbf{a}_1, \dots, \mathbf{a}_n) = (a_{ij})$  be a  $d \times n$ -matrix of rank  $d$  with coefficients in  $\mathbb{Z}$ . Throughout this paper, we assume the homogeneity of  $A$ , i.e., we assume that all  $\mathbf{a}_j$  belong to one hyperplane off the origin in  $\mathbb{Q}^d$ . Let  $\mathbb{N}$  be the set of nonnegative integers. Let  $I_A$  denote the toric ideal in the polynomial ring  $\mathbb{C}[\partial_{\mathbf{x}}] = \mathbb{C}[\partial_1, \dots, \partial_n]$ , i.e.,

$$I_A = \langle \partial_{\mathbf{x}}^{\mathbf{u}} - \partial_{\mathbf{x}}^{\mathbf{v}} \mid A\mathbf{u} = A\mathbf{v}, \mathbf{u}, \mathbf{v} \in \mathbb{N}^n \rangle \subseteq \mathbb{C}[\partial_{\mathbf{x}}].$$

Here and hereafter we use the multi-index notation; for example,  $\partial_{\mathbf{x}}^{\mathbf{u}}$  means  $\partial_1^{u_1} \cdots \partial_n^{u_n}$  for  $\mathbf{u} = (u_1, \dots, u_n)^T$ . Given a column vector  $\boldsymbol{\beta} = (\beta_1, \dots, \beta_d)^T \in \mathbb{C}^d$ , let  $H_A(\boldsymbol{\beta})$  denote the left ideal of the Weyl algebra

$$D = \mathbb{C}\langle \mathbf{x}, \partial_{\mathbf{x}} \rangle = \mathbb{C}\langle x_1, \dots, x_n, \partial_1, \dots, \partial_n \rangle$$

generated by  $I_A$  and

$$\sum_{j=1}^n a_{ij} \theta_j - \beta_i \quad (i = 1, \dots, d),$$

where  $\theta_j = x_j \partial_j$ . The quotient  $M_A(\boldsymbol{\beta}) = D/H_A(\boldsymbol{\beta})$  is called the  $A$ -hypergeometric system with parameter  $\boldsymbol{\beta}$ , and a formal series annihilated by  $H_A(\boldsymbol{\beta})$  an  $A$ -hypergeometric series with parameter  $\boldsymbol{\beta}$ . The

---

2020 *Mathematics Subject Classification.* Primary: 33C70.

*Key words and phrases.*  $A$ -hypergeometric systems; the method of Frobenius.

homogeneity of  $A$  is known to be equivalent to the regularity of  $M_A(\boldsymbol{\beta})$  by Hotta [4] and Schulze, Walther [9].

Logarithm-free series solutions to  $M_A(\boldsymbol{\beta})$  were constructed by Gel'fand et al. [2, 3] for a generic parameter  $\boldsymbol{\beta}$ , and more generally in [8].

Note that the logarithmic coefficients of  $A$ -hypergeometric series solutions are polynomials of  $\log x^{\mathbf{b}}$  ( $\mathbf{b} \in L$ ) [5, Proposition 5.2], where

$$L := \text{Ker}_{\mathbb{Z}}(A) = \{\mathbf{u} \in \mathbb{Z}^n \mid A\mathbf{u} = \mathbf{0}\}.$$

To construct logarithmic series solutions, the second author [6] introduced a method of perturbation by a finite subset  $B = \{\mathbf{b}^{(1)}, \dots, \mathbf{b}^{(h)}\} \subset L$ , and explicitly described logarithmic series solutions for a fake exponent and a set  $B$  that satisfy certain conditions [6, Theorems 5.4, 6.2 and Remarks 5.6, 6.3].

In this paper, following [6], we continue to develop the perturbing method of constructing logarithmic series solutions to a regular  $A$ -hypergeometric system.

Fixing a fake exponent of an  $A$ -hypergeometric system, we consider some spaces of linear partial differential operators with constant coefficients. Comparing these spaces, we construct a fundamental system of series solutions with the given exponent by the perturbing method. In addition, we give a sufficient condition for a given fake exponent to be an exponent. As important examples of the main results, we give fundamental systems of series solutions to Aomoto-Gel'fand systems and to Lauricella's  $F_C$  systems with special parameter vectors, respectively.

This paper is organized as follows. In Section 2, we first recall a power series to perturb from [6], associated with a fake exponent  $\mathbf{v}$  and a linearly independent subset  $B$  of  $L$ . In particular, we discuss properties of each term  $a_{\mathbf{u}}(\mathbf{s})$  appearing in the series (for the definition of  $a_{\mathbf{u}}(\mathbf{s})$ , see (1)), and modify the series by changing the range of the sum from  $\text{NS}_{\mathbf{w}}(\mathbf{v})$  in [6] to  $\mathcal{N}$  which incorporates  $B$ . We give a refinement of [6, Theorem 6.2] as Theorem 2.7.

In Section 3, for a fake exponent  $\mathbf{v}$  of the  $A$ -hypergeometric ideal  $H_A(\boldsymbol{\beta})$  with respect to a generic weight vector  $\mathbf{w}$ , we recall the structure of the ideal  $Q_{\mathbf{v}}$  associated with the fake indicial ideal  $\text{find}_{\mathbf{w}}(H_A(\boldsymbol{\beta}))$  and that of its orthogonal complement  $Q_{\mathbf{v}}^{\perp}$  defined in [8, Sections 2.3 and 3.6]. We introduce ideals  $P_{\mathcal{N}}$  and  $P_B$  of  $\mathbb{C}[\mathbf{s}]$ , and their orthogonal complements  $P_{\mathcal{N}}^{\perp}$  and  $P_B^{\perp}$ . Then we discuss relations among these ideals. Under a certain condition, we can derive  $Q_{\mathbf{v}}^{\perp}$  as the image of a linear map from  $P_{\mathcal{N}}^{\perp}$  (Proposition 3.11, Theorem 3.14).

In Section 4, we give a sufficient condition for a fake exponent  $\mathbf{v}$  to be an exponent (Proposition 4.1). Then we construct a fundamental system of solutions with the exponent  $\mathbf{v}$  (Theorem 4.4) by applying

Theorem 2.7 and the results of Section 3 under the condition that  $B$  is a basis of  $L$ , which is the main theorem of this paper.

In Sections 5 and 6, we deal with the Aomoto-Gel'fand systems and Lauricella's  $F_C$  systems, which are important examples of  $H_A(\boldsymbol{\beta})$ . We discuss a fundamental system of solutions to  $H_A(\mathbf{0})$  in each system. In each case, we have a unique fake exponent  $\mathbf{v} = \mathbf{0}$ . Taking a basis  $B$  of  $L$ , we can obtain a fundamental system of series solutions for  $\boldsymbol{\beta} = \mathbf{0}$ .

## 2. REFINEMENT OF [6, THEOREM 6.2]

In this section, we refine [6, Theorem 6.2].

Recall that for  $\mathbf{v} = (v_1, \dots, v_n)^T \in \mathbb{C}^n$  its support  $\text{supp}(\mathbf{v})$  and its *negative support*  $\text{nsupp}(\mathbf{v})$  are defined as

$$\begin{aligned}\text{supp}(\mathbf{v}) &:= \{j \in \{1, \dots, n\} \mid v_j \neq 0\}, \\ \text{nsupp}(\mathbf{v}) &:= \{j \in \{1, \dots, n\} \mid v_j \in \mathbb{Z}_{<0}\},\end{aligned}$$

respectively.

For  $\mathbf{v} \in \mathbb{C}^n$  and  $\mathbf{u} \in \mathbb{N}^n$ , set

$$[\mathbf{v}]_{\mathbf{u}} := \prod_{j=1}^n v_j (v_j - 1) \cdots (v_j - u_j + 1).$$

Here recall that  $\mathbb{N} = \{0, 1, 2, \dots\}$ .

Note that we can uniquely write  $\mathbf{u} \in \mathbb{Z}^n$  as the sum  $\mathbf{u} = \mathbf{u}_+ - \mathbf{u}_-$  with  $\mathbf{u}_+, \mathbf{u}_- \in \mathbb{N}^n$  and  $\text{supp}(\mathbf{u}_+) \cap \text{supp}(\mathbf{u}_-) = \emptyset$ .

Let  $B = \{\mathbf{b}^{(1)}, \dots, \mathbf{b}^{(h)}\} \subset L$ . We write the same symbol  $B$  for the  $n \times h$  matrix  $(\mathbf{b}^{(1)}, \dots, \mathbf{b}^{(h)})$ .

Set

$$\text{supp}(B) := \bigcup_{k=1}^h \text{supp}(\mathbf{b}^{(k)}) \subset \{1, \dots, n\},$$

which means the set of all labels for nonzero rows in  $B$ .

Let  $\mathbf{s} = (s_1, \dots, s_h)^T$  be indeterminates, and let

$$(B\mathbf{s})_j := \sum_{k=1}^h b_j^{(k)} s_k \in \mathbb{C}[\mathbf{s}] := \mathbb{C}[s_1, \dots, s_h]$$

for  $j = 1, \dots, n$ . Set

$$(B\mathbf{s})^J := \prod_{j \in J} (B\mathbf{s})_j \in \mathbb{C}[\mathbf{s}]$$

for  $J \subset \{1, \dots, n\}$ . Note that  $(B\mathbf{s})_j = 0$  if  $j \notin \text{supp}(B)$ , hence we have  $(B\mathbf{s})^J = 0$  if  $J \not\subset \text{supp}(B)$ .

**Lemma 2.1.** *Let  $B = \{\mathbf{b}^{(1)}, \dots, \mathbf{b}^{(h)}\} \subset L$ ,  $\mathbf{u}, \mathbf{u}' \in L$  and  $\mathbf{v} \in \mathbb{C}^n$ . Let  $\mathbf{s} = (s_1, \dots, s_h)^T$  be indeterminates. Then  $[\mathbf{v} + B\mathbf{s} + \mathbf{u}]_{\mathbf{u}'_+} \neq 0$  if and only if  $\text{nsupp}(\mathbf{v} + \mathbf{u} - \mathbf{u}') \subset \text{supp}(B) \cup \text{nsupp}(\mathbf{v} + \mathbf{u})$ . In particular,  $[\mathbf{v} + B\mathbf{s} + \mathbf{u}]_{\mathbf{u}_+} \neq 0$  if and only if  $\text{nsupp}(\mathbf{v}) \subset \text{supp}(B) \cup \text{nsupp}(\mathbf{v} + \mathbf{u})$ .*

*Proof.* Note that

$$\begin{aligned} & [\mathbf{v} + B\mathbf{s} + \mathbf{u}]_{\mathbf{u}'_+} \\ &= \prod_{j; u'_j > 0} (v_j + (B\mathbf{s})_j + u_j) \cdots (v_j + (B\mathbf{s})_j + u_j - u'_j + 1). \end{aligned}$$

Hence,  $[\mathbf{v} + B\mathbf{s} + \mathbf{u}]_{\mathbf{u}'_+} = 0$  if and only if there exists  $j$  such that  $v_j + u_j - u'_j \in \mathbb{Z}_{<0}$ ,  $v_j + u_j \in \mathbb{N}$ , and  $b_j^{(k)} = 0$  for all  $k$ .

Hence  $[\mathbf{v} + B\mathbf{s} + \mathbf{u}]_{\mathbf{u}'_+} = 0$  if and only if  $\text{nsupp}(\mathbf{v} + \mathbf{u} - \mathbf{u}') \not\subset \text{supp}(B) \cup \text{nsupp}(\mathbf{v} + \mathbf{u})$ .  $\square$

Let  $\mathbf{w}$  be a generic weight. Recall that  $\mathbf{v}$  is called a *fake exponent* of  $H_A(\beta)$  with respect to  $\mathbf{w}$  if  $A\mathbf{v} = \beta$  and  $[\mathbf{v}]_{\mathbf{u}_+} = 0$  for all  $\mathbf{u} \in L$  with  $\mathbf{u}_+ \cdot \mathbf{w} > \mathbf{u}_- \cdot \mathbf{w}$ , where  $\mathbf{u} \cdot \mathbf{w} = \sum_{j=1}^n u_j w_j$ .

Throughout this paper, fix a generic weight  $\mathbf{w}$ , a fake exponent  $\mathbf{v}$  of  $H_A(\beta)$  with respect to  $\mathbf{w}$ . In addition, we assume the following for a subset  $B = \{\mathbf{b}^{(1)}, \dots, \mathbf{b}^{(h)}\} \subset L$ .

**Assumption 2.2.** A subset  $B = \{\mathbf{b}^{(1)}, \dots, \mathbf{b}^{(h)}\} \subset L$  is linearly independent, hence  $\text{rank}(B) = h$ , and satisfies

$$\text{nsupp}(\mathbf{v}) \subset \text{supp}(B) \cup \text{nsupp}(\mathbf{v} + \mathbf{u})$$

for any  $\mathbf{u} \in L$ .

*Remark 2.3.* (1) If  $B$  satisfies  $\text{nsupp}(\mathbf{v}) \subset \text{supp}(B)$ , then Assumption 2.2 holds. For example, this condition holds for each of the following cases:

- (i)  $\text{supp}(B) = \{1, \dots, n\}$ ,
- (ii)  $\text{nsupp}(\mathbf{v}) = \emptyset$ .

(2) If  $B$  is a basis of  $L$ , then  $B$  satisfies

$$\text{supp}(B) \cup \text{nsupp}(\mathbf{v} + \mathbf{u}) = \text{supp}(B) \cup \text{nsupp}(\mathbf{v})$$

for all  $\mathbf{u} \in L$ . Indeed, since  $B$  is a basis of  $L$ , we see that if  $j \notin \text{supp}(B)$  then  $\mathbf{u}_j = 0$  for all  $\mathbf{u} \in L$ . This implies that  $\text{nsupp}(\mathbf{v} + \mathbf{u}) \setminus \text{supp}(B) = \text{nsupp}(\mathbf{v}) \setminus \text{supp}(B)$  for all  $\mathbf{u} \in L$ .

We abbreviate  $\text{nsupp}(\mathbf{v} + \mathbf{u})$  to  $I_{\mathbf{u}}$  for  $\mathbf{u} \in L$ . In particular,  $I_{\mathbf{0}} = \text{nsupp}(\mathbf{v})$ . Assumption 2.2 can be rewritten as

$$I_{\mathbf{0}} \subset \text{supp}(B) \cup I_{\mathbf{u}}$$

for all  $\mathbf{u} \in L$ .

For  $\mathbf{u} \in L$ , let

$$(1) \quad a_{\mathbf{u}}(\mathbf{s}) := \frac{[\mathbf{v} + B\mathbf{s}]_{\mathbf{u}_-}}{[\mathbf{v} + B\mathbf{s} + \mathbf{u}]_{\mathbf{u}_+}}.$$

**Lemma 2.4.** *Let  $\mathbf{u}, \mathbf{u}' \in L$ . Under Assumption 2.2, the following hold.*

- (i)  $a_{\mathbf{u}}(\mathbf{s}) \neq 0$  if and only if  $I_{\mathbf{u}} \subset \text{supp}(B) \cup I_0$ , if and only if  $\text{supp}(B) \cup I_{\mathbf{u}} = \text{supp}(B) \cup I_0$ .
- (ii) If  $I_{\mathbf{u}} \cup I_{\mathbf{u}-\mathbf{u}'} \not\subset \text{supp}(B) \cup I_0$ , then  $\partial^{\mathbf{u}'_+}(a_{\mathbf{u}}(\mathbf{s})x^{\mathbf{v}+B\mathbf{s}+\mathbf{u}}) = 0$ .

*Proof.* First, the denominator of  $a_{\mathbf{u}}(\mathbf{s})$  is not zero by Lemma 2.1.

(i) We have  $[\mathbf{v} + B\mathbf{s}]_{\mathbf{u}_-} = 0$  if and only if there exists  $j$  such that  $[v_j + \sum_{k=1}^h s_k b_j^{(k)}]_{-u_j} = [v_j]_{-u_j} = 0$ , namely  $v_j \in \mathbb{N}$ ,  $v_j + u_j \in \mathbb{Z}_{<0}$ , and  $b_j^{(k)} = 0$  for all  $k$ . Hence it is equivalent to saying that there exists  $j$  such that  $j \in I_{\mathbf{u}} \setminus (\text{supp}(B) \cup I_0)$ , or  $I_{\mathbf{u}} \not\subset \text{supp}(B) \cup I_0$ .

By the assumption, the inclusion  $I_{\mathbf{u}} \subset \text{supp}(B) \cup I_0$  is equivalent to the equality  $\text{supp}(B) \cup I_{\mathbf{u}} = \text{supp}(B) \cup I_0$ .

(ii) Note that

$$\partial^{\mathbf{u}'_+}(a_{\mathbf{u}}(\mathbf{s})x^{\mathbf{v}+B\mathbf{s}+\mathbf{u}}) = a_{\mathbf{u}}(\mathbf{s})[\mathbf{v} + B\mathbf{s} + \mathbf{u}]_{\mathbf{u}'_+} x^{\mathbf{v}+B\mathbf{s}+\mathbf{u}-\mathbf{u}'_+}.$$

Hence, if  $I_{\mathbf{u}} \not\subset \text{supp}(B) \cup I_0$ , then  $\partial^{\mathbf{u}'_+}(a_{\mathbf{u}}(\mathbf{s})x^{\mathbf{v}+B\mathbf{s}+\mathbf{u}}) = 0$  by (i).

Suppose that  $I_{\mathbf{u}-\mathbf{u}'} \setminus I_{\mathbf{u}} \not\subset \text{supp}(B) \cup I_0$ . Then, by Assumption 2.2,  $I_{\mathbf{u}-\mathbf{u}'} \setminus (\text{supp}(B) \cup I_{\mathbf{u}}) \neq \emptyset$ . Hence, by Lemma 2.1,  $[\mathbf{v} + B\mathbf{s} + \mathbf{u}]_{\mathbf{u}'_+} = 0$  and we have  $\partial^{\mathbf{u}'_+}(a_{\mathbf{u}}(\mathbf{s})x^{\mathbf{v}+B\mathbf{s}+\mathbf{u}}) = 0$ .  $\square$

We recall the definitions related to  $\text{NS}_{\mathbf{w}}(\mathbf{v})$  for a fake exponent  $\mathbf{v}$  from [6] and modify them.

Let

$$\mathcal{G} := \left\{ \partial^{\mathbf{g}^{(i)}_+} - \partial^{\mathbf{g}^{(i)}_-} \mid i = 1, \dots, m \right\}$$

denote the reduced Gröbner basis of  $I_A$  with respect to  $\mathbf{w}$  with  $\partial^{\mathbf{g}^{(i)}_+} \in \text{in}_{\mathbf{w}}(I_A)$  for all  $i$ . Note that the  $\mathcal{G}$  in [6, Section 4] should be *the reduced* Gröbner basis. Set

$$C(\mathbf{w}) := \sum_{i=1}^m \mathbb{N}\mathbf{g}^{(i)}.$$

A collection  $\text{NS}_{\mathbf{w}}(\mathbf{v})$  of negative supports  $I_{\mathbf{u}}$  ( $\mathbf{u} \in L$ ) is defined by

$$\text{NS}_{\mathbf{w}}(\mathbf{v}) := \{I_{\mathbf{u}} \mid \mathbf{u} \in L. \text{ If } I_{\mathbf{u}} = I_{\mathbf{u}'} \text{ for } \mathbf{u}' \in L, \text{ then } \mathbf{u}' \in C(\mathbf{w}).\}.$$

In addition, define

$$\text{NS}_{\mathbf{w}}(\mathbf{v})^c := \{I_{\mathbf{u}} \mid \mathbf{u} \in L\} \setminus \text{NS}_{\mathbf{w}}(\mathbf{v}).$$

We modify the definition of  $\text{NS}_{\mathbf{w}}(\mathbf{v})$  under Assumption 2.2. Define

$$(2) \quad \mathcal{N} := \{I \in \text{NS}_{\mathbf{w}}(\mathbf{v}) \mid \text{supp}(B) \cup I = \text{supp}(B) \cup I_0\},$$

and set

$$\mathcal{N}^c := \{I_{\mathbf{u}} \mid \mathbf{u} \in L, \text{supp}(B) \cup I_{\mathbf{u}} = \text{supp}(B) \cup I_0\} \setminus \mathcal{N}.$$

Consider the subset  $L'$  of  $L$  defined by

$$(3) \quad L' := \{\mathbf{u} \in L \mid I_{\mathbf{u}} \in \mathcal{N}\}.$$

By definition, we see that  $L' \subset C(\mathbf{w})$ .

Let

$$K_{\mathcal{N}} := \bigcap_{I \in \mathcal{N}} I,$$

and define the homogeneous ideal  $P_{\mathcal{N}}$  of  $\mathbb{C}[\mathbf{s}]$  for  $\mathcal{N}$  as

$$(4) \quad P_{\mathcal{N}} := \left\langle (B\mathbf{s})^{I \cup J \setminus K_{\mathcal{N}}} \mid I \in \mathcal{N}, J \in \mathcal{N}^c \right\rangle.$$

In addition, we define the orthogonal complement  $P^{\perp}$  for a homogeneous ideal  $P \subset \mathbb{C}[\mathbf{s}]$  as

$$(5) \quad P^{\perp} := \{q(\partial_{\mathbf{s}}) \in \mathbb{C}[\partial_{\mathbf{s}}] \mid (q(\partial_{\mathbf{s}}) \bullet h(\mathbf{s}))|_{\mathbf{s}=\mathbf{0}} = 0 \text{ for all } h(\mathbf{s}) \in P\} \\ = \{q(\partial_{\mathbf{s}}) \in \mathbb{C}[\partial_{\mathbf{s}}] \mid q(\partial_{\mathbf{s}}) \bullet P \subset \langle s_1, \dots, s_h \rangle\},$$

where  $\mathbb{C}[\partial_{\mathbf{s}}] := \mathbb{C}[\partial_{s_1}, \dots, \partial_{s_h}]$ . Since  $P$  and  $\langle s_1, \dots, s_h \rangle$  are both homogeneous,  $P^{\perp}$  is homogeneous with respect to the usual total ordering.

**Example 2.5.** (cf. [6, Examples 3.3, 4.8, 6.4]) Let  $n = 5$ ,  $d = 3$ , and

$$A = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ -1 & 1 & 1 & -1 & 0 \\ -1 & -1 & 1 & 1 & 0 \end{bmatrix}.$$

Let  $\boldsymbol{\beta} = (1, 0, 0)^T$  and  $\mathbf{w} = (1, 1, 1, 1, 0)$ . Then  $\mathbf{v} = (0, 0, 0, 0, 1)^T$  is a unique exponent, and

$$\mathcal{G} = \{\underline{\partial_{x_1} \partial_{x_3}} - \partial_{x_5}^2, \underline{\partial_{x_2} \partial_{x_4}} - \partial_{x_5}^2\},$$

where the underlined terms are the leading ones. Put  $\mathbf{g}^{(1)} := (1, 0, 1, 0, -2)^T$  and  $\mathbf{g}^{(2)} := (0, 1, 0, 1, -2)^T$ . Recall that

$$\text{NS}_{\mathbf{w}}(\mathbf{v}) = \{\emptyset = I_0, \{5\}\}, \\ \text{NS}_{\mathbf{w}}(\mathbf{v})^c = \{\{1, 3\}, \{2, 4\}, \{1, 3, 5\}, \{2, 4, 5\}, \{1, 2, 3, 4\}\}.$$

Let  $B := \{\mathbf{g}^{(1)}, \mathbf{g}^{(2)}\}$ . Then we have  $\text{supp}(B) = \{1, 2, 3, 4, 5\}$ , and

$$\begin{aligned}\mathcal{N} &= \text{NS}_{\mathbf{w}}(\mathbf{v}), \\ \mathcal{N}^c &= \text{NS}_{\mathbf{w}}(\mathbf{v})^c, \\ K_{\mathcal{N}} &= \emptyset.\end{aligned}$$

The homogeneous ideal  $P_{\mathcal{N}} \subset \mathbb{C}[\mathbf{s}] = \mathbb{C}[s_1, s_2]$  and the vector space  $P_{\mathcal{N}}^{\perp} \subset \mathbb{C}[\partial_{\mathbf{s}}] = \mathbb{C}[\partial_{s_1}, \partial_{s_2}]$  are given as

$$\begin{aligned}P_{\mathcal{N}} &= \langle (B\mathbf{s})^{\{1,3\}}, (B\mathbf{s})^{\{2,4\}} \rangle = \langle s_1^2, s_2^2 \rangle, \\ P_{\mathcal{N}}^{\perp} &= \{q(\partial_{s_1}, \partial_{s_2}) \in \mathbb{C}[\partial_{s_1}, \partial_{s_2}] \mid q(\partial_{s_1}, \partial_{s_2}) \bullet \langle s_1^2, s_2^2 \rangle \subset \langle s_1, s_2 \rangle\} \\ &= \mathbb{C}1 + \mathbb{C}\partial_{s_1} + \mathbb{C}\partial_{s_2} + \mathbb{C}\partial_{s_1}\partial_{s_2}.\end{aligned}$$

We consider another case. Let  $B_1 = \{\mathbf{g}^{(1)}\}$ . Then we have  $\text{supp}(B_1) = \{1, 3, 5\}$  and

$$\begin{aligned}\mathcal{N}_1 &= \text{NS}_{\mathbf{w}}(\mathbf{v}) = \{\emptyset = I_0, \{5\}\}, \\ \mathcal{N}_1^c &= \{\{1, 3\}, \{1, 3, 5\}\}, \\ K_{\mathcal{N}_1} &= \emptyset.\end{aligned}$$

The homogeneous ideal  $P_{\mathcal{N}_1} \subset \mathbb{C}[s]$  and the vector space  $P_{\mathcal{N}_1}^{\perp} \subset \mathbb{C}[\partial_s]$  are given as

$$\begin{aligned}P_{\mathcal{N}_1} &= \langle (B\mathbf{s})^{\{1,3\}} \rangle = \langle s^2 \rangle, \\ P_{\mathcal{N}_1}^{\perp} &= \{q(\partial_s) \in \mathbb{C}[\partial_s] \mid q(\partial_s) \bullet \langle s^2 \rangle \subset \langle s \rangle\} = \mathbb{C}1 + \mathbb{C}\partial_s.\end{aligned}$$

Throughout this paper, put

$$m(\mathbf{s}) := (B\mathbf{s})^{I_0 \setminus K_{\mathcal{N}}}.$$

The following lemma guarantees that we may plug  $\mathbf{s} = \mathbf{0}$  into the series appearing in Theorem 2.7.

**Lemma 2.6.** *Let  $\mathcal{N}$  be the one defined by (2), and let  $\mathbf{u}, \mathbf{u}' \in L$ . Then, under Assumption 2.2, each term of the power series for  $m(\mathbf{s}) \cdot a_{\mathbf{u}}(\mathbf{s}) \cdot [\mathbf{v} + B\mathbf{s} + \mathbf{u}]_{\mathbf{u}'_+}$  in the indeterminates  $\mathbf{s}$  is divided by  $(B\mathbf{s})^{I_{\mathbf{u} \cup I_{\mathbf{u}'}} \setminus K_{\mathcal{N}}}$ .*

*Proof.* By [6, Lemma 6.1], there exists a formal power series  $g(\mathbf{y})$  in the indeterminates  $\mathbf{y} = (y_1, \dots, y_n)$  such that

$$\begin{aligned} & a_{\mathbf{u}}(\mathbf{s}) \cdot [\mathbf{v} + B\mathbf{s} + \mathbf{u}]_{\mathbf{u}'_+} \\ &= \frac{[\mathbf{v} + B\mathbf{s}]_{\mathbf{u}_-}}{[\mathbf{v} + B\mathbf{s} + \mathbf{u}]_{\mathbf{u}_+}} \cdot [\mathbf{v} + B\mathbf{s} + \mathbf{u}]_{\mathbf{u}'_+} \\ &= \left( \frac{(B\mathbf{s})^{I_{\mathbf{u}} \setminus I_0}}{(B\mathbf{s})^{I_0 \setminus I_{\mathbf{u}}}} \cdot (B\mathbf{s})^{I_{\mathbf{u}-\mathbf{u}'} \setminus I_{\mathbf{u}}} \right) \cdot g((B\mathbf{s})_1, \dots, (B\mathbf{s})_n) \\ &= \frac{(B\mathbf{s})^{(I_{\mathbf{u}} \cup I_{\mathbf{u}-\mathbf{u}'}) \setminus I_0}}{(B\mathbf{s})^{I_0 \setminus (I_{\mathbf{u}} \cup I_{\mathbf{u}-\mathbf{u}'})}} \cdot g((B\mathbf{s})_1, \dots, (B\mathbf{s})_n). \end{aligned}$$

Hence we have

$$\begin{aligned} & m(\mathbf{s}) \cdot a_{\mathbf{u}}(\mathbf{s}) \cdot [\mathbf{v} + B\mathbf{s} + \mathbf{u}]_{\mathbf{u}'_+} \\ &= (B\mathbf{s})^{I_0 \setminus K_{\mathcal{N}}} \cdot \frac{(B\mathbf{s})^{(I_{\mathbf{u}} \cup I_{\mathbf{u}-\mathbf{u}'}) \setminus I_0}}{(B\mathbf{s})^{I_0 \setminus (I_{\mathbf{u}} \cup I_{\mathbf{u}-\mathbf{u}'})}} \cdot g((B\mathbf{s})_1, \dots, (B\mathbf{s})_n) \\ &= (B\mathbf{s})^{I_{\mathbf{u}} \cup I_{\mathbf{u}-\mathbf{u}'} \setminus K_{\mathcal{N}}} \cdot g((B\mathbf{s})_1, \dots, (B\mathbf{s})_n), \end{aligned}$$

and the assertion holds.  $\square$

We can refine the main results [6, Theorem 5.4, Theorem 6.2] as follows.

**Theorem 2.7.** *Let  $\mathcal{N}$  be the one defined by (2). Set*

$$F_{\mathcal{N}}(\mathbf{x}, \mathbf{s}) := \sum_{\mathbf{u} \in L'} a_{\mathbf{u}}(\mathbf{s}) x^{v+B\mathbf{s}+\mathbf{u}},$$

and

$$\tilde{F}_{\mathcal{N}}(\mathbf{x}, \mathbf{s}) := m(\mathbf{s}) F_{\mathcal{N}}(\mathbf{x}, \mathbf{s}),$$

where  $L'$  is defined by (3).

Then  $(q(\partial_{\mathbf{s}}) \bullet \tilde{F}_{\mathcal{N}}(\mathbf{x}, \mathbf{s}))|_{\mathbf{s}=\mathbf{0}}$  are solutions to  $M_A(\boldsymbol{\beta})$  for any  $q(\partial_{\mathbf{s}}) \in P_{\mathcal{N}}^{\perp}$ .

*Proof.* Let  $\mathbf{u}' \in L$  and  $\mathbf{u} \in L'$ . If  $\partial^{\mathbf{u}'_+}(a_{\mathbf{u}}(\mathbf{s})x^{v+B\mathbf{s}+\mathbf{u}}) \neq 0$ , then we have  $I_{\mathbf{u}} \cup I_{\mathbf{u}-\mathbf{u}'} \subset \text{supp}(B) \cup I_0$  by Lemma 2.4, and hence  $\text{supp}(B) \cup I_{\mathbf{u}-\mathbf{u}'} = \text{supp}(B) \cup I_0$  by Assumption 2.2. Thus  $I_{\mathbf{u}-\mathbf{u}'} \notin \mathcal{N}$  implies  $I_{\mathbf{u}-\mathbf{u}'} \in \mathcal{N}^c$ .



Similar to the arguments in the proofs of [6, Theorem 5.4, Theorem 6.2], we see that

$$\begin{aligned} & (\partial^{\mathbf{u}'_+} - \partial^{\mathbf{u}'_-}) \bullet \tilde{F}_{\mathcal{N}}(\mathbf{x}, \mathbf{s}) \\ &= \sum_{\mathbf{u} \in L', I_{\mathbf{u}-\mathbf{u}'_-} \in \mathcal{N}^c} m(\mathbf{s}) \partial^{\mathbf{u}'_+} (a_{\mathbf{u}}(\mathbf{s}) x^{\mathbf{v}+B\mathbf{s}+\mathbf{u}}) \\ & \quad - \sum_{\mathbf{u} \in L', I_{\mathbf{u}+\mathbf{u}'_+} \in \mathcal{N}^c} m(\mathbf{s}) \partial^{(-\mathbf{u})'_+} (a_{\mathbf{u}}(\mathbf{s}) x^{\mathbf{v}+B\mathbf{s}+\mathbf{u}}). \end{aligned}$$

Let  $q(\partial_{\mathbf{s}}) \in P_{\mathcal{N}}^{\perp}$ . Then the series  $(q(\partial_{\mathbf{s}}) \bullet \tilde{F}_{\mathcal{N}}(\mathbf{x}, \mathbf{s}))|_{\mathbf{s}=\mathbf{0}}$  is a solution to  $M_A(\boldsymbol{\beta})$  if

$$\left( q(\partial_{\mathbf{s}}) \bullet \left( m(\mathbf{s}) \partial^{\mathbf{u}'_+} a_{\mathbf{u}}(\mathbf{s}) x^{\mathbf{v}+B\mathbf{s}+\mathbf{u}} \right) \right) \Big|_{\mathbf{s}=\mathbf{0}} = 0$$

for any  $\mathbf{u} \in L'$  and  $\mathbf{u}' \in L$  with  $I_{\mathbf{u}-\mathbf{u}'_-} \in \mathcal{N}^c$ .

By Lemma 2.6, each coefficient of

$$\begin{aligned} & m(\mathbf{s}) (\partial^{\mathbf{u}'_+} a_{\mathbf{u}}(\mathbf{s}) x^{\mathbf{v}+B\mathbf{s}+\mathbf{u}}) \\ &= m(\mathbf{s}) a_{\mathbf{u}}(\mathbf{s}) [\mathbf{v} + B\mathbf{s} + \mathbf{u}]_{\mathbf{u}'_+} x^{\mathbf{v}+B\mathbf{s}+\mathbf{u}-\mathbf{u}'_+} \end{aligned}$$

in the indeterminates  $\mathbf{s}$  is divided by  $(B\mathbf{s})^{I_{\mathbf{u} \cup I_{\mathbf{u}-\mathbf{u}'_-} \setminus K_{\mathcal{N}}}}$ , hence belongs to  $P_{\mathcal{N}}$ . By the definition of  $P_{\mathcal{N}}^{\perp}$ , the assertion holds.  $\square$

### 3. RELATIONS BETWEEN $P_{\mathcal{N}}^{\perp}$ AND $Q_{\mathbf{v}}^{\perp}$

In this section, we recall  $Q_{\mathbf{v}}$  and its orthogonal complement  $Q_{\mathbf{v}}^{\perp}$  defined in [8, Section 2.3], and discuss relations between  $P_{\mathcal{N}}^{\perp}$  and  $Q_{\mathbf{v}}^{\perp}$ . For the definitions of  $P_{\mathcal{N}}$  and  $P_{\mathcal{N}}^{\perp}$ , see (4) and (5).

Consider the fake indicial ideal  $\text{find}_{\mathbf{w}}(H_A(\boldsymbol{\beta}))$  of  $H_A(\boldsymbol{\beta})$  with respect to  $\mathbf{w}$ :

$$\text{find}_{\mathbf{w}}(H_A(\boldsymbol{\beta})) := \langle A\theta_{\mathbf{x}} - \boldsymbol{\beta} \rangle + \tilde{\text{in}}_{\mathbf{w}}(I_A) \subset \mathbb{C}[\theta_{\mathbf{x}}] := \mathbb{C}[\theta_1, \dots, \theta_n].$$

Here  $\tilde{\text{in}}_{\mathbf{w}}(I_A)$  is the distraction of the initial ideal  $\text{in}_{\mathbf{w}}(I_A)$  with respect to  $\mathbf{w}$  (cf. [8, Section 3.1]). Related to the reduced Gröbner basis  $\mathcal{G} = \{\mathbf{g}^{(i)} \mid i = 1, \dots, m\}$  of  $I_A$  with respect to  $\mathbf{w}$ , define

$$G^{(i)} := I_{-\mathbf{g}^{(i)}} \setminus I_{\mathbf{0}} = \{j \in \{1, \dots, n\} \mid v_j \in \mathbb{N}, g_j^{(i)} - v_j > 0\}$$

for  $i = 1, \dots, m$ . Since

$$\tilde{\text{in}}_{\mathbf{w}}(I_A) = \left\langle [\theta_{\mathbf{x}}]_{\mathbf{g}_+^{(i)}} := \prod_{j; g_j^{(i)} > 0} \prod_{\nu=0}^{g_j^{(i)}-1} (\theta_j - \nu) \mid i = 1, \dots, m \right\rangle$$

by [8, Theorem 3.2.2], we see that its primary component at a fake exponent  $\mathbf{v}$  is

$$(6) \quad \tilde{\text{in}}_{\mathbf{w}}(I_A)_{\mathbf{v}} = \left\langle ([\theta]_{\mathbf{g}_+^{(i)}})_{\mathbf{v}} := \prod_{j \in G^{(i)}} (\theta_j - v_j) \mid i = 1, \dots, m \right\rangle.$$

We obtain the homogeneous ideal  $Q_{\mathbf{v}}$  of  $\mathbb{C}[\theta_{\mathbf{x}}]$  from  $\text{find}_{\mathbf{w}}(H_A(\boldsymbol{\beta}))_{\mathbf{v}}$  by replacing  $\theta_j \mapsto \theta_j + v_j$  for  $j = 1, \dots, n$  (cf. [8, Section 2.3]). Namely,

$$(7) \quad Q_{\mathbf{v}} = \langle A\theta_{\mathbf{x}} \rangle + \left\langle \prod_{j \in G^{(i)}} \theta_j \mid i = 1, \dots, m \right\rangle.$$

The orthogonal complement  $Q_{\mathbf{v}}^{\perp}$  of  $Q_{\mathbf{v}}$  is defined by

$$Q_{\mathbf{v}}^{\perp} := \{f \in \mathbb{C}[\mathbf{x}] \mid \phi(\partial_{\mathbf{x}})(f) = 0 \text{ for all } \phi = \phi(\theta_{\mathbf{x}}) \in Q_{\mathbf{v}}\}.$$

Note that  $Q_{\mathbf{v}}^{\perp}$  is a graded  $\mathbb{C}$ -vector space with the usual grading.

**Proposition 3.1.** *Let  $f(\mathbf{x})$  be a polynomial. Then  $x^{\mathbf{v}}f(\log \mathbf{x})$  is a solution to  $\text{find}_{\mathbf{w}}(H_A(\boldsymbol{\beta}))$  if and only if  $f(\mathbf{x})$  satisfies the following conditions:*

- (i)  $f(\mathbf{x}) \in \mathbb{C}[\mathbf{x}G] := \mathbb{C}[\mathbf{x}\mathbf{g}^{(1)}, \dots, \mathbf{x}\mathbf{g}^{(m)}]$ .
- (ii)  $\partial_{\mathbf{x}}^{G^{(i)}} \bullet f(\mathbf{x}) = 0$  for all  $i = 1, \dots, m$ .

Here

$$\mathbf{x}G := (\mathbf{x}\mathbf{g}^{(1)}, \dots, \mathbf{x}\mathbf{g}^{(m)}) = \left( \sum_{j=1}^n g_j^{(1)} x_j, \dots, \sum_{j=1}^n g_j^{(m)} x_j \right)$$

for  $\mathbf{x} = (x_1, \dots, x_n)$ .

*Proof.* By [8, Theorem 2.3.11], the function  $x^{\mathbf{v}}f(\log \mathbf{x})$  is a solution to  $\text{find}_{\mathbf{w}}(H_A(\boldsymbol{\beta}))$  if and only if  $f(\mathbf{x}) \in Q_{\mathbf{v}}^{\perp}$ . From  $f(\mathbf{x}) \in \langle A\partial_{\mathbf{x}} \rangle^{\perp}$ , we see (i) [5, Lemma 5.1]. (ii) follows from Equation (6).  $\square$

**Example 3.2** (Continuation of Example 2.5). Note that  $\mathbf{v} - \mathbf{g}^{(1)} = (-1, 0, -1, 0, 3)^T$  and  $\mathbf{v} - \mathbf{g}^{(2)} = (0, -1, 0, -1, 3)^T$ . Thus we see that

$$\begin{aligned} G^{(1)} &= I_{-\mathbf{g}^{(1)}} \setminus I_{\mathbf{0}} = \text{nsupp}(\mathbf{v} - \mathbf{g}^{(1)}) \setminus \text{nsupp}(\mathbf{v}) = \{1, 3\}, \\ G^{(2)} &= I_{-\mathbf{g}^{(2)}} \setminus I_{\mathbf{0}} = \text{nsupp}(\mathbf{v} - \mathbf{g}^{(2)}) \setminus \text{nsupp}(\mathbf{v}) = \{2, 4\}. \end{aligned}$$

The ideal  $Q_{\mathbf{v}} \subset \mathbb{C}[\theta_{\mathbf{x}}] = \mathbb{C}[\theta_1, \theta_2, \theta_3, \theta_4, \theta_5]$  is given as

$$Q_{\mathbf{v}} = \langle \theta_1 + \theta_2 + \theta_3 + \theta_4 + \theta_5, -\theta_1 + \theta_2 + \theta_3 - \theta_4, -\theta_1 - \theta_2 + \theta_3 + \theta_4, \theta_1\theta_3, \theta_2\theta_4 \rangle.$$

In addition, we see that

$$Q_{\mathbf{v}}^{\perp} = \mathbb{C} \cdot 1 + \mathbb{C} \cdot \mathbf{x}\mathbf{g}^{(1)} + \mathbb{C} \cdot \mathbf{x}\mathbf{g}^{(2)} + \mathbb{C} \cdot (\mathbf{x}\mathbf{g}^{(1)}) \cdot (\mathbf{x}\mathbf{g}^{(2)}).$$

To compare  $Q_{\mathbf{v}}$  with  $P_{\mathcal{N}}$ , we consider the graded ring homomorphism  $\Phi_B : \mathbb{C}[\theta_{\mathbf{x}}] \rightarrow \mathbb{C}[\mathbf{s}]$  defined by  $\theta_j \mapsto (B\mathbf{s})_j$  for  $j = 1, \dots, n$ . By the linear independence of  $B$ , we see that  $\Phi_B$  is surjective. Define  $P_B := \Phi_B(Q_{\mathbf{v}})$ . By the ring isomorphism theorem,  $\Phi_B$  induces the ring isomorphism

$$\tilde{\Phi}_B : \mathbb{C}[\theta_{\mathbf{x}}]/\Phi_B^{-1}(P_B) \simeq \mathbb{C}[\mathbf{s}]/P_B.$$

Since  $\langle A\theta_{\mathbf{x}} \rangle$  is vanished by  $\Phi_B$ , we have

$$(8) \quad P_B = \left\langle (B\mathbf{s})^{G^{(i)}} \mid i = 1, \dots, m \right\rangle.$$

**Proposition 3.3.** *Let  $J \in \text{NS}_{\mathbf{w}}(\mathbf{v})^c$ . Then  $G^{(i)} \subset J \setminus I_0$  for some  $i$ .*

*Proof.* By definition and [6, Lemma 4.2], we see that there exists  $\mathbf{u} \in L \setminus C(\mathbf{w})$  such that  $J = I_{\mathbf{u}}$  and  $\partial^{\mathbf{u}+} \notin \text{in}_{\mathbf{w}}(I_A)$ . Hence  $\partial^{\mathbf{u}-} = \text{in}_{\mathbf{w}}(\partial^{\mathbf{u}-} - \partial^{\mathbf{u}+})$  is divided by some  $\partial^{\mathbf{g}^{(i)}}$ . Let  $j \in G^{(i)} = I_{-\mathbf{g}^{(i)}} \setminus I_0$ . Then  $v_j \in \mathbb{N}$  and  $v_j - g_j^{(i)} \in \mathbb{Z}_{<0}$ . Since  $g_j^{(i)} \in \mathbb{Z}_{>0}$ , we see that  $g_j^{(i)} \leq -u_j$  and  $v_j + u_j \leq v_j - g_j^{(i)} < 0$ . Thus we have  $j \in I_{\mathbf{u}} \setminus I_0 = J \setminus I_0$ .  $\square$

Three ideals  $Q_{\mathbf{v}}$ ,  $P_{\mathcal{N}}$ , and  $P_B$  are related as follows.

**Proposition 3.4.** *Let  $Q_{\mathbf{v}}$ ,  $P_{\mathcal{N}}$ , and  $P_B$  be the ones in (7), (4), and (8), respectively. Then, the following hold.*

- (i)  $m(\mathbf{s}) \cdot P_B \subset P_{\mathcal{N}} \subset P_B$ . In particular, if  $K_{\mathcal{N}} = I_0$ , then  $P_{\mathcal{N}} = P_B$ .
- (ii) If  $B$  is a basis of  $L$ , then  $\Phi_B^{-1}(P_B) = Q_{\mathbf{v}}$ .

*Proof.* (i) Let  $I \in \mathcal{N}$  and  $J \in \mathcal{N}^c$ . Since  $J \in \text{NS}_{\mathbf{w}}(\mathbf{v})^c$  and  $K_{\mathcal{N}} \subset I_0$ ,  $I \cup J \setminus K_{\mathcal{N}}$  contains some  $G^{(i)}$  by Proposition 3.3. Hence the inclusion  $P_{\mathcal{N}} \subset P_B$  holds.

For any  $i = 1, \dots, m$ , since  $-\mathbf{g}^{(i)} \notin C(\mathbf{w})$  we see that  $I_{-\mathbf{g}^{(i)}} \in \text{NS}_{\mathbf{w}}(\mathbf{v})^c$ . If  $I_{-\mathbf{g}^{(i)}} \notin \mathcal{N}^c$ , then

$$(9) \quad \text{supp}(B) \cup I_{-\mathbf{g}^{(i)}} \neq \text{supp}(B) \cup I_0.$$

By Assumption 2.2, (9) implies that  $G^{(i)} = I_{-\mathbf{g}^{(i)}} \setminus I_0 \not\subset \text{supp}(B)$ . Hence we have  $(B\mathbf{s})^{G^{(i)}} = 0$ . If  $I_{-\mathbf{g}^{(i)}} \in \mathcal{N}^c$ , then

$$m(\mathbf{s})(B\mathbf{s})^{G^{(i)}} = (B\mathbf{s})^{I_0 \cup I_{-\mathbf{g}^{(i)}} \setminus K_{\mathcal{N}}} \in P_{\mathcal{N}}.$$

Hence we have  $m(\mathbf{s}) \cdot P_B \subset P_{\mathcal{N}}$ .

(ii) Since  $B$  is a basis of  $L$ , we have  $\text{Ker}(\Phi_B) = \langle A\theta_{\mathbf{x}} \rangle$ . Thus the assertion  $\Phi_B^{-1}(P_B) = Q_{\mathbf{v}}$  holds from (7).  $\square$

**Example 3.5** (Continuation of Example 2.5 and 3.2). Consider the case where  $B = \{\mathbf{g}^{(1)}, \mathbf{g}^{(2)}\}$ . Then we have  $m(\mathbf{s}) = (B\mathbf{s})^\emptyset = 1$ , and

$$P_B = \langle (B\mathbf{s})^{G^{(1)}}, (B\mathbf{s})^{G^{(2)}} \rangle = \langle s_1^2, s_2^2 \rangle = P_{\mathcal{N}}.$$

Furthermore, since  $B$  is a basis of  $L$ , we see that

$$\Phi_B^{-1}(P_B) = \Phi_B^{-1}(\langle (B\mathbf{s})^{G^{(1)}}, (B\mathbf{s})^{G^{(2)}} \rangle) = \langle \theta_1\theta_3, \theta_2\theta_4 \rangle + \langle A\theta_x \rangle = Q_{\mathbf{v}}.$$

Consider the other case where  $B_1 = \{\mathbf{g}^{(1)}\}$ . Then we have  $m(s) = (B_1s)^\emptyset = 1$ , and

$$P_{B_1} = \langle (B_1s)^{G^{(1)}} \rangle = \langle s^2 \rangle = P_{\mathcal{N}_1}.$$

We see that  $B_1$  does not span  $L$  and that

$$\Phi_{B_1}^{-1}(P_{B_1}) = \Phi_{B_1}^{-1}(\langle (B_1s)^{G^{(1)}} \rangle) = \langle \theta_1\theta_3 \rangle + \langle A\theta_x \rangle \subsetneq Q_{\mathbf{v}}.$$

We consider relations between  $P_{\mathcal{N}}^\perp$  and  $P_B^\perp$ , and between  $P_B^\perp$  and  $Q_{\mathbf{v}}^\perp$ . Recall the construction of a basis of orthogonal complements in [8, Section 2.3].

Let  $P$  be a homogeneous ideal of  $\mathbb{C}[\mathbf{s}]$ . Fix any term order  $\prec$  on  $\mathbb{C}[\mathbf{s}]$ , and let  $\mathcal{H} \subset \mathbb{C}[\mathbf{s}]$  be the reduced Gröbner basis of  $P$  with respect to  $\prec$ . For any  $\boldsymbol{\mu} \in \mathbb{N}^h$  with  $\mathbf{s}^\boldsymbol{\mu} \in \text{in}_\prec(P)$ , there exist unique  $c_{\boldsymbol{\mu}, \boldsymbol{\nu}} \in \mathbb{C}$  for  $\boldsymbol{\nu} \in \mathbb{N}^h$  with  $|\boldsymbol{\nu}| = |\boldsymbol{\mu}|$  and  $\mathbf{s}^\boldsymbol{\nu} \notin \text{in}_\prec(P)$  such that

$$p_\boldsymbol{\mu}(\mathbf{s}) := \mathbf{s}^\boldsymbol{\mu} - \sum_{\substack{\boldsymbol{\nu} \in \mathbb{N}^h; |\boldsymbol{\nu}|=|\boldsymbol{\mu}|, \\ \mathbf{s}^\boldsymbol{\nu} \notin \text{in}_\prec(P)}} c_{\boldsymbol{\mu}, \boldsymbol{\nu}} \mathbf{s}^\boldsymbol{\nu} \in P.$$

We obtain  $p_\boldsymbol{\mu}(\mathbf{s})$  by taking the normal form modulo  $\mathcal{H}$  for the monomial  $\mathbf{s}^\boldsymbol{\mu}$ . For  $\boldsymbol{\nu} \in \mathbb{N}^h$  with  $\mathbf{s}^\boldsymbol{\nu} \notin \text{in}_\prec(P)$ , define the homogeneous polynomial  $q_\boldsymbol{\nu}(\partial_s)$  of degree  $|\boldsymbol{\nu}|$  by

$$q_\boldsymbol{\nu}(\partial_s) := \frac{1}{\boldsymbol{\nu}!} \partial_s^\boldsymbol{\nu} + \sum_{\substack{\boldsymbol{\mu} \in \mathbb{N}^h; |\boldsymbol{\mu}|=|\boldsymbol{\nu}|, \\ \mathbf{s}^\boldsymbol{\mu} \in \text{in}_\prec(P)}} \frac{c_{\boldsymbol{\mu}, \boldsymbol{\nu}}}{\boldsymbol{\mu}!} \partial_s^\boldsymbol{\mu} \in \mathbb{C}[\partial_s].$$

**Lemma 3.6.** *Let  $P$  be a homogeneous ideal of  $\mathbb{C}[\mathbf{s}]$ . Fix any term order  $\prec$  on  $\mathbb{C}[\mathbf{s}]$ , and let  $\mathcal{H} \subset \mathbb{C}[\mathbf{s}]$  be the reduced Gröbner basis of  $P$  with respect to  $\prec$ . Then*

$$\{p_\boldsymbol{\mu}(\mathbf{s}) \mid \mathbf{s}^\boldsymbol{\mu} \in \text{in}_\prec(P)\}$$

and

$$\{q_\boldsymbol{\nu}(\partial_s) \mid \boldsymbol{\nu} \in \mathbb{N}^h \text{ with } \mathbf{s}^\boldsymbol{\nu} \notin \text{in}_\prec(P)\}$$

form  $\mathbb{C}$ -bases of  $P$  and  $P^\perp$ , respectively.

*Proof.* This is similar to [8, Proposition 2.3.13].  $\square$

**Lemma 3.7.** *Let  $P$  and  $\tilde{P}$  be homogeneous ideals of  $\mathbb{C}[\mathbf{s}]$ . Then  $P \subset \tilde{P}$  if and only if  $P^\perp \supset \tilde{P}^\perp$ .*

*Proof.* Assume that  $P \subset \tilde{P}$ . By definition,  $P^\perp \supset \tilde{P}^\perp$  is clear.

Conversely, assume that  $P^\perp \supset \tilde{P}^\perp$ . Fix any term order  $\prec$  on  $\mathbb{C}[\mathbf{s}]$ , and let  $\tilde{\mathcal{H}}$  be the reduced Gröbner basis of  $\tilde{P}$ .

Let  $\{\tilde{q}_\nu(\partial_s) \mid \nu \in \mathbb{N}^h \text{ with } \mathbf{s}^\nu \notin \text{in}_\prec(\tilde{P})\}$  be the  $\mathbb{C}$ -basis of  $\tilde{P}^\perp$  as in Lemma 3.6. Let  $p \in P$ . Applying the division algorithm with respect to  $\tilde{\mathcal{H}}$  to  $p$ , we can express  $p$  as

$$p = \tilde{p} + \sum_{\lambda: \mathbf{s}^\lambda \notin \text{in}_\prec(\tilde{P})} d_\lambda \mathbf{s}^\lambda$$

with some  $\tilde{p} \in \tilde{P}$  and  $d_\lambda \in \mathbb{C}$ . For each  $\nu \in \mathbb{N}^h$  with  $\mathbf{s}^\nu \notin \text{in}_\prec(\tilde{P})$ , since  $[\partial_s^\mu \bullet \mathbf{s}^\lambda]_{|s=0} = \boldsymbol{\mu}! \delta_{\mu, \lambda}$  for any  $\boldsymbol{\mu}$  and  $\boldsymbol{\lambda}$ , we have

$$\begin{aligned} & [\tilde{q}_\nu(\partial_s) \bullet p]_{|s=0} \\ &= \left[ \tilde{q}_\nu(\partial_s) \bullet \left( \tilde{p} + \sum_{\lambda: \mathbf{s}^\lambda \notin \text{in}_\prec(\tilde{P})} d_\lambda \mathbf{s}^\lambda \right) \right]_{|s=0} \\ &= \sum_{\lambda: \mathbf{s}^\lambda \notin \text{in}_\prec(\tilde{P})} d_\lambda [\tilde{q}_\nu(\partial_s) \bullet \mathbf{s}^\lambda]_{|s=0} \\ &= \sum_{\lambda: \mathbf{s}^\lambda \notin \text{in}_\prec(\tilde{P})} d_\lambda \left[ \left( \frac{1}{\boldsymbol{\nu}!} \partial_s^\nu + \sum_{\substack{\boldsymbol{\mu} \in \mathbb{N}^h; |\boldsymbol{\mu}| = |\boldsymbol{\nu}|, \\ \mathbf{s}^\mu \in \text{in}_\prec(\tilde{P})}} \frac{c_{\boldsymbol{\mu}, \boldsymbol{\nu}}}{\boldsymbol{\mu}!} \partial_s^\mu \right) \bullet \mathbf{s}^\lambda \right]_{|s=0} \\ &= d_\nu. \end{aligned}$$

It follows from the assumption  $P^\perp \supset \tilde{P}^\perp$  that  $d_\nu = 0$  for all  $\nu \in \mathbb{N}^h$  with  $\mathbf{s}^\nu \notin \text{in}_\prec(\tilde{P})$ . Hence, we have  $p \in \tilde{P}$ .  $\square$

Let  $\mathbb{C}[\partial_z] := \mathbb{C}[\partial_{z_1}, \dots, \partial_{z_h}]$  be the ring of partial differential operators with constant coefficients in indeterminates  $\mathbf{z} = (z_1, \dots, z_h)$ . To describe relations between  $P_N^\perp$  and  $P_B^\perp$ , and between  $P_B^\perp$  and  $Q_v^\perp$ , we define an action of  $\mathbb{C}[\partial_z]$  on  $\mathbb{C}[\partial_s]$  and a ring homomorphism  $\Psi_B$  from  $\mathbb{C}[\partial_s]$  to  $\mathbb{C}[\mathbf{x}]$ .

For  $U(\partial_z) \in \mathbb{C}[\partial_z]$  and  $q(\partial_s) \in \mathbb{C}[\partial_s]$ , we define a  $\mathbb{C}$ -linear operation  $U(\partial_{z_1}, \dots, \partial_{z_h}) \star q(\partial_s)$  by

$$U(\partial_{z_1}, \dots, \partial_{z_h}) \star q(\partial_s) := (U(\partial_z) \bullet q(\mathbf{z}))|_{\mathbf{z}=\partial_s} \in \mathbb{C}[\partial_s].$$

**Lemma 3.8.** *The following hold for the  $\star$ -operation.*

(i) *Let  $k = 1, \dots, h$  and  $q(\partial_s) \in \mathbb{C}[\partial_s]$ . Then*

$$\partial_{z_k} \star q(\partial_s) = q(\partial_s)s_k - s_k q(\partial_s) \in \mathbb{C}\langle \mathbf{s}, \partial_s \rangle.$$

(ii) *Let  $U(\partial_z), U'(\partial_z) \in \mathbb{C}[\partial_z]$ , and  $q(\partial_s) \in \mathbb{C}[\partial_s]$ . Then*

$$U(\partial_z) \star (U'(\partial_z) \star q(\partial_s)) = (U(\partial_z)U'(\partial_z)) \star q(\partial_s).$$

(iii) *Let  $U(\partial_z) = \prod_{\nu=1}^N l_\nu(\partial_z) \in \mathbb{C}[\partial_z]$  be the product of non-zero linear homogeneous polynomials  $l_\nu(\partial_z)$ , and let  $q(\partial_s) \in \mathbb{C}[\partial_s]$ . Then there exists  $r(\partial_s) \in \mathbb{C}[\partial_s]$  such that  $U(\partial_z) \star r(\partial_s) = q(\partial_s)$ .*

*Proof.* (i) For any  $k = 1, \dots, h$ , and  $\boldsymbol{\mu} \in \mathbb{N}^h$ , we have

$$\partial_{z_k} \star \partial_s^\boldsymbol{\mu} = \mu_k \partial_s^{\boldsymbol{\mu} - \mathbf{e}_k} = \partial_s^\boldsymbol{\mu} s_k - s_k \partial_s^\boldsymbol{\mu}.$$

(ii) It suffices to show that the equality holds for  $U(\partial_z) = \partial_z^\boldsymbol{\lambda}$ ,  $U'(\partial_z) = \partial_z^\boldsymbol{\mu}$ , and  $q(\partial_s) = \partial_s^\boldsymbol{\nu}$  with  $\boldsymbol{\lambda}, \boldsymbol{\mu}, \boldsymbol{\nu} \in \mathbb{N}^h$ . We see that

$$\begin{aligned} \partial_z^\boldsymbol{\lambda} \star (\partial_z^\boldsymbol{\mu} \star \partial_s^\boldsymbol{\nu}) &= \partial_z^\boldsymbol{\lambda} \star ((\partial_z^\boldsymbol{\mu} \bullet \mathbf{z}^\boldsymbol{\nu})|_{z=\partial_s}) \\ &= [\boldsymbol{\nu}]_\mu \partial_z^\boldsymbol{\lambda} \star \partial_s^{\boldsymbol{\nu} - \boldsymbol{\mu}} \\ &= [\boldsymbol{\nu}]_\mu (\partial_z^\boldsymbol{\lambda} \bullet \mathbf{z}^{\boldsymbol{\nu} - \boldsymbol{\mu}})|_{z=\partial_s} \\ &= [\boldsymbol{\nu}]_\mu [\boldsymbol{\nu} - \boldsymbol{\mu}]_\lambda \partial_s^{\boldsymbol{\nu} - \boldsymbol{\mu} - \boldsymbol{\lambda}} \\ &= [\boldsymbol{\nu}]_{\lambda + \boldsymbol{\mu}} \partial_s^{\boldsymbol{\nu} - (\lambda + \boldsymbol{\mu})} \\ &= (\partial_z^{\lambda + \boldsymbol{\mu}} \bullet \partial_s^\boldsymbol{\nu})|_{z=\partial_s} \\ &= \partial_z^{\lambda + \boldsymbol{\mu}} \star \partial_s^\boldsymbol{\nu}, \end{aligned}$$

and hence the assertion holds.

(iii) We show the statement by induction on  $N$ . First, let  $U(\partial_z)$  be a non-zero linear homogeneous polynomial, and let  $q(\partial_s) \in \mathbb{C}[\partial_s]$ . By changing coordinates, we may assume that  $U(\partial_z) = \partial_{z_1}$ . Put

$$q(\partial_s) = \sum_{\boldsymbol{\nu} \in \mathbb{N}^h} d_\nu \partial_s^\boldsymbol{\nu}.$$

Then

$$r(\partial_s) = \sum_{\boldsymbol{\nu} \in \mathbb{N}^h} \frac{d_\nu}{\nu_1 + 1} \partial_s^{\boldsymbol{\nu} + \mathbf{e}_1}$$

satisfies  $U(\partial_z) \star r(\partial_s) = q(\partial_s)$ .

Next, fix  $N > 1$ , and let  $U(\partial_z) = \prod_{\nu=1}^N l_\nu(\partial_z)$  such that  $l_\nu(\partial_z)$  are non-zero linear homogeneous polynomials. Assume that the assertion holds for any product of non-zero linear homogeneous polynomial of degree less than  $N$ . By the induction hypothesis, there exist

$r(\partial_s), \tilde{r}(\partial_s) \in \mathbb{C}[\partial_s]$  such that

$$l_1(\partial_z) \star \tilde{r}(\partial_s) = q(\partial_s), \quad \left( \prod_{\nu=2}^N l_\nu(\partial_z) \right) \star r(\partial_s) = \tilde{r}(\partial_s).$$

By (ii), we have

$$\begin{aligned} U(\partial_z) \star r(\partial_s) &= \left( l_1(\partial_z) \left( \prod_{\nu=2}^N l_\nu(\partial_z) \right) \right) \star r(\partial_s) \\ &= l_1(\partial_z) \star \left( \left( \prod_{\nu=2}^N l_\nu(\partial_z) \right) \star r(\partial_s) \right) \\ &= l_1(\partial_z) \star \tilde{r}(\partial_s) \\ &= q(\partial_s), \end{aligned}$$

and hence the assertion holds.  $\square$

**Lemma 3.9.** *Let  $U(\partial_z) \in \mathbb{C}[\partial_z]$ ,  $q(\partial_s) \in \mathbb{C}[\partial_s]$ , and  $f(\mathbf{s}) \in \mathbb{C}[[\mathbf{s}]]$ . Then*

$$[q(\partial_s) \bullet (U(\mathbf{s})f(\mathbf{s}))]_{|\mathbf{s}=\mathbf{0}} = [(U(\partial_z) \star q(\partial_s)) \bullet f(\mathbf{s})]_{|\mathbf{s}=\mathbf{0}}.$$

*Proof.* We show that the statement holds for any monomial operator  $U(\partial_z) = \partial_z^\mu$  by induction on  $|\mu|$ . Firstly, the assertion is clear for  $\mu = \mathbf{0}$ .

Secondly, assume that  $\mu = e_k$  for  $k = 1, \dots, h$ , hence  $U(\partial_z) = \partial_z^\mu = \partial_{z_k}$ . Note that  $[s_k q(\partial_s) \bullet f(\mathbf{s})]_{|\mathbf{s}=\mathbf{0}} = 0$  for any  $k = 1, \dots, h$ ,  $q(\partial_s) \in \mathbb{C}[\partial_s]$ , and  $f(\mathbf{s}) \in \mathbb{C}[[\mathbf{s}]]$ . Thus, by Lemma 3.8, we see that

$$\begin{aligned} &[q(\partial_s) \bullet (U(\mathbf{s})f(\mathbf{s}))]_{|\mathbf{s}=\mathbf{0}} \\ &= [(q(\partial_s)s_k) \bullet f(\mathbf{s})]_{|\mathbf{s}=\mathbf{0}} \\ &= [(s_k q(\partial_s) + \partial_{z_k} \star q(\partial_s)) \bullet f(\mathbf{s})]_{|\mathbf{s}=\mathbf{0}} \\ &= [s_k(q(\partial_s) \bullet f(\mathbf{s}))]_{|\mathbf{s}=\mathbf{0}} + [(\partial_{z_k} \star q(\partial_s)) \bullet f(\mathbf{s})]_{|\mathbf{s}=\mathbf{0}} \\ &= [(\partial_{z_k} \star q(\partial_s)) \bullet f(\mathbf{s})]_{|\mathbf{s}=\mathbf{0}}. \end{aligned}$$

Hence the assertion holds for  $|\mu| = 1$ .

Finally, fix  $\mu \in \mathbb{N}^h$  with  $|\mu| > 1$ . Let  $U(\partial_z) = \partial_z^\mu$ ,  $q(\partial_s) \in \mathbb{C}[\partial_s]$ , and  $f(\mathbf{s}) \in \mathbb{C}[[\mathbf{s}]]$ . Assume that the assertion holds for any  $\tilde{U}(\partial_z) = \partial_z^{\tilde{\mu}}$  with  $|\tilde{\mu}| < |\mu|$ . Then there exists  $k$  such that  $\mu_k > 0$ . Applying the induction hypothesis to the operators  $\partial_{z_k}$  and  $\partial_z^{\mu - e_k}$ , respectively, we

see from Lemma 3.8 (ii) that

$$\begin{aligned}
& [q(\partial_s) \bullet (U(\mathbf{s})f(\mathbf{s}))]_{|\mathbf{s}=\mathbf{0}} \\
&= [q(\partial_s) \bullet (s_k \cdot \mathbf{s}^{\mu-e_k} f(\mathbf{s}))]_{|\mathbf{s}=\mathbf{0}} \\
&= [(\partial_{z_k} \star q(\partial_s)) \bullet (\mathbf{s}^{\mu-e_k} f(\mathbf{s}))]_{|\mathbf{s}=\mathbf{0}} \\
&= [(\partial_z^{\mu-e_k} \star (\partial_{z_k} \star q(\partial_s))) \bullet f(\mathbf{s})]_{|\mathbf{s}=\mathbf{0}} \\
&= [(U(\partial_z) \star q(\partial_s)) \bullet f(\mathbf{s})]_{|\mathbf{s}=\mathbf{0}}.
\end{aligned}$$

Hence the assertion holds.  $\square$

We define a ring homomorphism  $\Psi_B : \mathbb{C}[\partial_s] \rightarrow \mathbb{C}[\mathbf{x}]$  as

$$(10) \quad \Psi_B(q(\partial_s))(\mathbf{x}) := q(\mathbf{x}B) = q\left(\sum_{j=1}^n b_j^{(1)} x_j, \dots, \sum_{j=1}^n b_j^{(h)} x_j\right)$$

for  $q(\partial_s) \in \mathbb{C}[\partial_s]$ . Note that  $\Psi_B$  is injective by the linear independence of  $B$ .

**Proposition 3.10.** *Let  $q(\partial_s) \in \mathbb{C}[\partial_s]$ . Then*

$$[q(\partial_s) \bullet (m(\mathbf{s})x^{v+B\mathbf{s}})]_{|\mathbf{s}=\mathbf{0}} = x^v \Psi_B(m(\partial_z) \star q(\partial_s))(\log \mathbf{x}),$$

where  $\log \mathbf{x} := (\log x_1, \dots, \log x_n)$ .

*Proof.* Note that we can regard  $x^{v+B\mathbf{s}}$  as the formal series  $x^v e^{(\log \mathbf{x})B\mathbf{s}}$  in  $\mathbf{s}$ , where

$$(\log \mathbf{x})B\mathbf{s} := \sum_{j=1}^n \sum_{k=1}^h (\log x_j) b_j^{(k)} s_k.$$

Put  $r(\partial_s) := m(\partial_z) \star q(\partial_s)$ . Then, by Lemma 3.9,

$$\begin{aligned}
[r(\partial_s) \bullet (m(\mathbf{s})x^{v+B\mathbf{s}})]_{|\mathbf{s}=\mathbf{0}} &= [(m(\partial_z) \star q(\partial_s)) \bullet x^{v+B\mathbf{s}}]_{|\mathbf{s}=\mathbf{0}} \\
&= [r(\partial_s) \bullet x^{v+B\mathbf{s}}]_{|\mathbf{s}=\mathbf{0}} \\
&= [r((\log \mathbf{x})B)x^{v+B\mathbf{s}}]_{|\mathbf{s}=\mathbf{0}} \\
&= x^v \Psi_B(r(\partial_s))(\log \mathbf{x}).
\end{aligned}$$

$\square$

**Proposition 3.11.** *The following hold.*

- (i)  $m(\partial_z) \star P_N^\perp \subset P_B^\perp \subset P_N^\perp$ . In particular, if  $K_N = I_0$ , then  $P_N^\perp = P_B^\perp$ .
- (ii)  $m(\mathbf{s}) \in P_N$  if and only if  $m(\partial_z) \star P_N^\perp = \{0\}$ .
- (iii) If  $P_N = m(\mathbf{s}) \cdot P_B$ , then  $m(\partial_z) \star P_N^\perp = P_B^\perp$ .



*Proof.* (i)  $P_B^\perp \subset P_N^\perp$  is clear by Lemma 3.4 (i) and Lemma 3.7. Let  $q(\partial_s) \in P_N^\perp$ . Then, for any  $f(\mathbf{s}) \in P_B$ , Lemma 3.9 shows that

$$[(m(\partial_z) \star q(\partial_s)) \bullet f(\mathbf{s})]_{|\mathbf{s}=\mathbf{0}} = [q(\partial_s) \bullet (m(\mathbf{s})f(\mathbf{s}))]_{|\mathbf{s}=\mathbf{0}}.$$

It follows from Lemma 3.4 (i) that the right hand side is 0. Hence we have  $m(\partial_z) \star P_N^\perp \subset P_B^\perp$ .

(ii) Assume that  $m(\mathbf{s}) \in P_N$ . Let  $q(\partial_s) \in P_N^\perp$ . Put  $m(\partial_z) \star q(\partial_s) = \sum_{\nu} a_{\nu} \partial_s^{\nu}$ , where  $a_{\nu} \in \mathbb{C}$ . Then, by Lemma 3.9, we have

$$\begin{aligned} \nu! a_{\nu} &= [(m(\partial_z) \star q(\partial_s)) \bullet \mathbf{s}^{\nu}]_{|\mathbf{s}=\mathbf{0}} \\ &= [q(\partial_s) \bullet (m(\mathbf{s})\mathbf{s}^{\nu})]_{|\mathbf{s}=\mathbf{0}} = 0 \end{aligned}$$

for any  $\nu \in \mathbb{N}^h$ . Hence, we have  $m(\partial_z) \star q(\partial_s) = 0$ .

Conversely, assume that  $m(\partial_z) \star P_N^\perp = \{0\}$ . Let  $q(\partial_s) \in P_N^\perp$ . Then, Lemma 3.9 shows that

$$[q(\partial_s) \bullet (m(\mathbf{s})f(\mathbf{s}))]_{|\mathbf{s}=\mathbf{0}} = [(m(\partial_z) \star q(\partial_s)) \bullet f(\mathbf{s})]_{|\mathbf{s}=\mathbf{0}} = 0$$

for any  $f(\mathbf{s}) \in \mathbb{C}[\mathbf{s}]$ . Thus we have  $q(\partial_s) \in \langle m(\mathbf{s}) \rangle^\perp$ , that is,  $P_N^\perp \subset \langle m(\mathbf{s}) \rangle^\perp$ . By Lemma 3.7,  $m(\mathbf{s}) \in P_N$ .

(iii) In (i), we have seen  $m(\partial_z) \star P_N^\perp \subset P_B^\perp$ . We show its reverse inclusion. Let  $q(\partial_s) \in P_B^\perp$ . By Lemma 3.8 (iii), there exists  $r(\partial_s) \in \mathbb{C}[\partial_s]$  such that  $q(\partial_s) = m(\partial_z) \star r(\partial_s)$ . It suffices to show that  $r(\partial_s) \in P_N^\perp$ . Let  $f(\mathbf{s}) \in P_N$ . By the assumption, we have  $f(\mathbf{s}) = m(\mathbf{s})g(\mathbf{s})$  for some  $g(\mathbf{s}) \in P_B$ . By Lemma 3.9, we see that

$$\begin{aligned} [r(\partial_s) \bullet f(\mathbf{s})]_{|\mathbf{s}=\mathbf{0}} &= [r(\partial_s) \bullet (m(\mathbf{s})g(\mathbf{s}))]_{|\mathbf{s}=\mathbf{0}} \\ &= [(m(\partial_z) \star r(\partial_s)) \bullet g(\mathbf{s})]_{|\mathbf{s}=\mathbf{0}} \\ &= [q(\partial_s) \bullet g(\mathbf{s})]_{|\mathbf{s}=\mathbf{0}} = 0. \end{aligned}$$

Hence we have the assertion.  $\square$

**Example 3.12.** (cf. [6, Examples 3.2 and 4.7]) Let  $A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 3 & 4 \end{bmatrix}$  and let  $\mathbf{w} = (3, 1, 0, 0)$ . Then the reduced Gröbner basis of  $I_A$  is

$$\mathcal{G} = \{ \underline{\partial_{x_1} \partial_{x_3}^2} - \partial_{x_2}^2 \partial_{x_4}, \underline{\partial_{x_2} \partial_{x_4}^2} - \partial_{x_3}^3, \underline{\partial_{x_1}^2 \partial_{x_3}} - \partial_{x_2}^3, \underline{\partial_{x_1} \partial_{x_4}} - \partial_{x_2} \partial_{x_3} \}.$$

Here underlined terms are the leading ones. Thus we have

$$\text{in}_{\mathbf{w}}(I_A) = \langle \partial_{x_1} \partial_{x_3}^2, \partial_{x_2} \partial_{x_4}^2, \partial_{x_1}^2 \partial_{x_3}, \partial_{x_1} \partial_{x_4} \rangle.$$

Put

$$\begin{aligned} \mathbf{g}^{(1)} &= (1, -2, 2, -1)^T, & \mathbf{g}^{(2)} &= (0, 1, -3, 2)^T, \\ \mathbf{g}^{(3)} &= (2, -3, 1, 0)^T, & \mathbf{g}^{(4)} &= (1, -1, -1, 1)^T. \end{aligned}$$

Let  $\boldsymbol{\beta} = (-2, -1)^T$ , and let

$$B = (\mathbf{g}^{(1)}, \mathbf{g}^{(2)}) = \begin{bmatrix} 1 & 0 \\ -2 & 1 \\ 2 & -3 \\ -1 & 2 \end{bmatrix}.$$

Note that  $\text{supp}(B) = \{1, 2, 3, 4\}$ . Take  $\mathbf{v} = (0, -2, -1, 1)^T$  as a fake exponent. Then we have

$$\begin{aligned} \mathcal{N} &= \{\{2\}, \{3\}, \{2, 3\} = I_0\}, \\ \mathcal{N}^c &= \{\{1, 2\}, \{1, 3\}, \{1, 4\}, \{2, 4\}, \{1, 2, 4\}, \{1, 3, 4\}\}, \\ K_{\mathcal{N}} &= \emptyset. \end{aligned}$$

Furthermore, we have

$$\begin{aligned} G^{(1)} &= I_{-\mathbf{g}^{(1)}} \setminus I_0 = \{1\}, & G^{(2)} &= I_{-\mathbf{g}^{(2)}} \setminus I_0 = \{4\}, \\ G^{(3)} &= I_{-\mathbf{g}^{(3)}} \setminus I_0 = \{1\}, & G^{(4)} &= I_{-\mathbf{g}^{(4)}} \setminus I_0 = \{1\}. \end{aligned}$$

Thus the ideals  $P_{\mathcal{N}}$  and  $P_B$  are

$$\begin{aligned} P_{\mathcal{N}} &= \langle (B\mathbf{s})^{\{1,2\}}, (B\mathbf{s})^{\{1,3\}}, (B\mathbf{s})^{\{2,4\}} \rangle \\ &= \langle s_1(-2s_1 + s_2), s_1(2s_1 - 3s_2), (-2s_1 + s_2)(-s_1 + 2s_2) \rangle \\ &= \langle s_1^2, s_1s_2, s_2^2 \rangle \end{aligned}$$

and

$$P_B = \langle (B\mathbf{s})^{\{1\}}, (B\mathbf{s})^{\{4\}} \rangle = \langle s_1, -s_1 + 2s_2 \rangle = \langle s_1, s_2 \rangle,$$

respectively. The orthogonal complements  $P_{\mathcal{N}}^\perp$  and  $P_B^\perp$  are

$$P_{\mathcal{N}}^\perp = \mathbb{C}1 + \mathbb{C}\partial_{s_1} + \mathbb{C}\partial_{s_2}$$

and

$$P_B^\perp = \mathbb{C}1,$$

respectively. In this case, note that

$$m(\mathbf{s}) = (B\mathbf{s})^{I_0 \setminus K_{\mathcal{N}}} = (B\mathbf{s})^{\{2,3\}} = (-2s_1 + s_2)(2s_1 - 3s_2) \in P_{\mathcal{N}}.$$

Hence, by Proposition 3.11, we have

$$m(\partial_{\mathbf{z}}) \star P_{\mathcal{N}}^\perp = \{0\}.$$

**Lemma 3.13.** *Let  $q(\mathbf{z}) \in \mathbb{C}[\mathbf{z}] := \mathbb{C}[z_1, \dots, z_h]$  be a homogeneous polynomial in indeterminates  $\mathbf{z}$  of degree  $r$ . Then*

$$q(\partial_{\mathbf{s}}) \bullet \left( \frac{1}{r!} (\mathbf{x}B\mathbf{s})^r \right) = q(\mathbf{x}B) = \Psi_B(q(\partial_{\mathbf{s}}))(\mathbf{x}).$$

Here,

$$\mathbf{x}B\mathbf{s} := \sum_{j=1}^n \sum_{k=1}^h x_j b_j^{(k)} s_k$$

denotes the quadratic form associated with  $B$ .

*Proof.* We show the assertion by induction on  $r$ . In the case of  $r = 1$ , the assertion is clear. Fix  $r > 1$  and assume that the assertion holds for any homogeneous polynomial of degree less than  $r$ . Let  $\boldsymbol{\mu} \in \mathbb{N}^h$  with  $|\boldsymbol{\mu}| = r$  and  $\mu_k > 0$ . Then, by the chain rule and the induction hypothesis, we see that

$$\begin{aligned} \partial_s^{\boldsymbol{\mu}} \bullet \left( \frac{1}{r!} (\mathbf{x}B\mathbf{s})^r \right) &= \partial_s^{\boldsymbol{\mu}-\mathbf{e}_k} \bullet \left( \partial_{s_k} \bullet \left( \frac{1}{r!} (\mathbf{x}B\mathbf{s})^r \right) \right) \\ &= (\mathbf{x}B)_k \cdot \partial_s^{\boldsymbol{\mu}-\mathbf{e}_k} \bullet \left( \frac{1}{(r-1)!} (\mathbf{x}B\mathbf{s})^{r-1} \right) \\ &= (\mathbf{x}B)_k (\mathbf{x}B)^{\boldsymbol{\mu}-\mathbf{e}_k} = (\mathbf{x}B)^{\boldsymbol{\mu}} = \Psi_B(\partial_s^{\boldsymbol{\mu}})(\mathbf{x}). \end{aligned}$$

□

Two vector spaces  $Q_v^\perp$  and  $P_B^\perp$  are related as follows.

**Theorem 3.14.** *Let  $\Psi_B$  be the homomorphism in (10). Then,  $P_B^\perp = \Psi_B^{-1}(Q_v^\perp)$  and  $\dim_{\mathbb{C}}(P_B^\perp) \leq \dim_{\mathbb{C}}(Q_v^\perp)$ . Furthermore, if  $B$  is a basis of  $L$ , then  $\Psi_B(P_B^\perp) = Q_v^\perp$  and  $\dim_{\mathbb{C}}(Q_v^\perp) = \dim_{\mathbb{C}}(P_B^\perp)$ .*

*Proof.* Let  $\deg(q(\mathbf{z})) = r$ . Then, it follows from Lemma 3.13 that

$$\begin{aligned}
\partial_{\mathbf{x}}^{G^{(i)}} \bullet q(\mathbf{x}B) &= \partial_{\mathbf{x}}^{G^{(i)}} \bullet \left\{ q(\partial_{\mathbf{s}}) \bullet \left( \frac{1}{r!} (\mathbf{x}B\mathbf{s})^r \right) \right\} \\
&= q(\partial_{\mathbf{s}}) \bullet \left\{ \partial_{\mathbf{x}}^{G^{(i)}} \bullet \left( \frac{1}{r!} (\mathbf{x}B\mathbf{s})^r \right) \right\} \\
&= q(\partial_{\mathbf{s}}) \bullet \left\{ \partial_{\mathbf{x}}^{G^{(i)}} \bullet \left( \frac{1}{r!} \sum_{\mu \in \mathbb{N}^n; |\mu|=r} \frac{r!}{\mu!} \mathbf{x}^\mu (B\mathbf{s})^\mu \right) \right\} \\
&= q(\partial_{\mathbf{s}}) \bullet \left\{ \sum_{\mu \in \mathbb{N}^n; |\mu|=r} \frac{\partial_{\mathbf{x}}^{G^{(i)}} \bullet \mathbf{x}^\mu}{\mu!} (B\mathbf{s})^\mu \right\} \\
&= q(\partial_{\mathbf{s}}) \bullet \left\{ \sum_{\mu \in \mathbb{N}^n; |\mu|=r, \text{supp}(\mu) \supset G^{(i)}} \frac{\mathbf{x}^{\mu - \mathbf{e}_{G^{(i)}}}}{(\mu - \mathbf{e}_{G^{(i)}})!} (B\mathbf{s})^\mu \right\} \\
&= \sum_{\mu \in \mathbb{N}^n; |\mu|=r, \text{supp}(\mu) \supset G^{(i)}} \frac{\mathbf{x}^{\mu - \mathbf{e}_{G^{(i)}}}}{(\mu - \mathbf{e}_{G^{(i)}})!} \{q(\partial_{\mathbf{s}}) \bullet (B\mathbf{s})^\mu\}
\end{aligned}$$

for any  $i$ . Here  $\mathbf{e}_{G^{(i)}} := \sum_{j \in G^{(i)}} \mathbf{e}_j$  denotes the indicator vector of  $G^{(i)}$ . Since  $\partial_{\mathbf{s}}^{\mathbf{p}} \bullet \mathbf{s}^{\mathbf{q}} = \mathbf{p}! \delta_{\mathbf{p}, \mathbf{q}}$  for any  $\mathbf{p}, \mathbf{q} \in \mathbb{N}^h$ , by Proposition 3.1 we have

$$\begin{aligned}
q(\mathbf{x}B) \in Q_{\mathbf{v}}^\perp &\iff \partial_{\mathbf{x}}^{G^{(i)}} \bullet q(\mathbf{x}B) = 0 \text{ for all } i = 1, \dots, m \\
&\iff q(\partial_{\mathbf{s}}) \bullet (B\mathbf{s})^\mu = 0 \\
&\quad \text{for all } i \text{ and all } \mu \in \mathbb{N}^n \text{ with } |\mu| = r, \text{supp}(\mu) \supset G^{(i)} \\
&\iff q(\partial_{\mathbf{s}}) \in P_B^\perp.
\end{aligned}$$

Here, the second equivalence follows from the linear independence of the monomials  $\mathbf{x}^\mu$ . The third equivalence follows from the linear independence of  $B$ , because it yields that the ideal  $P_B$  is spanned as a vector space by the polynomials whose terms are of the form  $(B\mathbf{s})^\mu$  with  $\text{supp}(\mu) \supset G^{(i)}$  for some  $i$ . Thus we have  $P_B^\perp = \Psi_B^{-1}(Q_{\mathbf{v}}^\perp)$ . Moreover, we have the inequality  $\dim_{\mathbb{C}}(P_B^\perp) \leq \dim_{\mathbb{C}}(Q_{\mathbf{v}}^\perp)$  because  $\Psi_B(P_B^\perp) = \Psi_B(\Psi_B^{-1}(Q_{\mathbf{v}}^\perp)) \subset Q_{\mathbf{v}}^\perp$  and  $\Psi_B$  is injective.

Assume that  $B$  is a basis of  $L$ . Note that each  $\mathbf{x}\mathbf{g}^{(k)} = \sum_{j=1}^n g_j^{(k)} x_j$  can be represented by a linear combination of  $(\mathbf{x}B)_1, \dots, (\mathbf{x}B)_h$ . Let  $f(\mathbf{x}) \in Q_{\mathbf{v}}^\perp$ . Then, by Proposition 3.1 and the above result, there exists  $q(\partial_{\mathbf{s}}) \in P_B^\perp$  such that  $f(\mathbf{x}) = q(\mathbf{x}B)$ . Thus we have  $f(\mathbf{x}) = \Psi_B(q(\partial_{\mathbf{s}}))(\mathbf{x})$ .  $\square$

## 4. FUNDAMENTAL SYSTEMS OF SOLUTIONS

In this section, we construct a fundamental system of series solutions with a given exponent to  $M_A(\boldsymbol{\beta})$ . We recall that the homogeneity of  $A$  yields the regular holonomicity of  $M_A(\boldsymbol{\beta})$ . This means that, for a fixed generic weight  $\mathbf{w}$ , the solution space to  $M_A(\boldsymbol{\beta})$  has a basis consisting of canonical series with starting monomial  $x^{\mathbf{v}}(\log \mathbf{x})^{\mathbf{b}}$  for some exponent  $\mathbf{v}$  and  $\mathbf{b} \in \mathbb{N}^n$ . Note that each  $x^{\mathbf{v}}(\log \mathbf{x})^{\mathbf{b}}$  is derived as the initial monomial of a solution to the indicial ideal  $\text{ind}_{\mathbf{w}}(H_A(\boldsymbol{\beta}))_{\mathbf{v}}$ , or of an element of  $Q_{\mathbf{v}}^{\perp}$ . For the detail, see [8, Sections 2.3, 2.4, and 2.5] and Proposition 3.1.

Throughout this section, we assume that  $B$  is a basis of  $L$ . Since  $B$  satisfies Assumption 2.2 (see Remark 2.3), we have the following homomorphisms by Propositions 3.1, 3.10, and Theorem 3.14:

$$(11) \quad \begin{array}{ccc} P_{\mathcal{N}}^{\perp} & \rightarrow & P_B^{\perp} \\ q(\partial_{\mathbf{s}}) & \mapsto & m(\partial_{\mathbf{z}}) \star q(\partial_{\mathbf{s}}) \end{array} \simeq \text{Sol}(\text{find}_{\mathbf{w}}(H_A(\boldsymbol{\beta}))_{\mathbf{v}}),$$

$$\leftrightarrow x^{\mathbf{v}}\Psi_B(m(\partial_{\mathbf{z}}) \star q(\partial_{\mathbf{s}}))(\log \mathbf{x}).$$

Here,  $\text{Sol}(\text{find}_{\mathbf{w}}(H_A(\boldsymbol{\beta}))_{\mathbf{v}})$  denotes the solution space of the fake indicial ideal  $\text{find}_{\mathbf{w}}(H_A(\boldsymbol{\beta}))_{\mathbf{v}}$ .

**Proposition 4.1.** *If  $m(\mathbf{s}) \notin P_{\mathcal{N}}$ , then  $\mathbf{v}$  is an exponent.*

*Proof.* Assume that  $m(\mathbf{s}) \notin P_{\mathcal{N}}$ . Then, by Proposition 3.11 (ii), there exists  $q(\partial_{\mathbf{s}}) \in P_{\mathcal{N}}^{\perp}$  such that  $m(\partial_{\mathbf{z}}) \star q(\partial_{\mathbf{s}}) \neq 0$ . We see from Theorem 2.7 that

$$\begin{aligned} & (q(\partial_{\mathbf{s}}) \bullet \tilde{F}_{\mathcal{N}}(\mathbf{x}, \mathbf{s}))|_{\mathbf{s}=\mathbf{0}} \\ &= \sum_{\mathbf{u} \in L'} (q(\partial_{\mathbf{s}}) \bullet (m(\mathbf{s})a_{\mathbf{u}}(\mathbf{s})x^{\mathbf{v}+B\mathbf{s}+\mathbf{u}}))|_{\mathbf{s}=\mathbf{0}} \\ &= (q(\partial_{\mathbf{s}}) \bullet (m(\mathbf{s})x^{\mathbf{v}+B\mathbf{s}}))|_{\mathbf{s}=\mathbf{0}} \\ & \quad + \sum_{\mathbf{u} \in L' \setminus \{\mathbf{0}\}} (q(\partial_{\mathbf{s}}) \bullet (m(\mathbf{s})a_{\mathbf{u}}(\mathbf{s})x^{\mathbf{v}+B\mathbf{s}+\mathbf{u}}))|_{\mathbf{s}=\mathbf{0}} \end{aligned}$$

is a solution to  $M_A(\boldsymbol{\beta})$ . By Proposition 3.10, we have

$$(q(\partial_{\mathbf{s}}) \bullet (m(\mathbf{s})x^{\mathbf{v}+B\mathbf{s}}))|_{\mathbf{s}=\mathbf{0}} = x^{\mathbf{v}}\Psi_B(m(\partial_{\mathbf{z}}) \star q(\partial_{\mathbf{s}}))(\log \mathbf{x}),$$

hence this solution has a *non-zero* starting term. Hence  $\mathbf{v}$  is an exponent.  $\square$

**Example 4.2** (Continuation of Example 3.12). Let  $A$  and  $\mathbf{v}$  be the ones in Example 3.12. Recall that  $m(\mathbf{s}) \in P_{\mathcal{N}}$ , which is a necessary condition for the fake exponent  $\mathbf{v}$  not to be an exponent. We see that  $\mathbf{v}$  is not an exponent from the following calculation. Note that

$$\theta_1 - 2\theta_3 - 3\theta_4 + 1 \in \langle A\theta_{\mathbf{x}} - \boldsymbol{\beta} \rangle.$$

Then we have

$$\begin{aligned}
0 &\equiv \theta_1\theta_3(\theta_1 - 2\theta_3 - 3\theta_4 + 1) \\
&= \theta_1^2\theta_3 - 2\theta_1\theta_3^2 - 3\theta_1\theta_3\theta_4 + \theta_1\theta_3 \\
&= \theta_1(\theta_1 - 1)\theta_3 - 2\theta_1\theta_3(\theta_3 - 1) - 3\theta_1\theta_3\theta_4 \\
&= x_1^2x_3\partial_{x_1}^2\partial_{x_3} - 2x_1x_3^2\partial_{x_1}\partial_{x_3}^2 - 3x_1x_3x_4\partial_{x_1}\partial_{x_3}\partial_{x_4} \\
&\equiv x_1^2x_3\partial_{x_2}^3 - 2x_1x_3^2\partial_{x_2}^2\partial_{x_4} - 3x_1x_3x_4\partial_{x_2}\partial_{x_3}^2
\end{aligned}$$

modulo  $H_A(\boldsymbol{\beta})$ . Hence

$$x_1^2x_3\partial_{x_2}^3 - 2x_1x_3^2\partial_{x_2}^2\partial_{x_4} - 3x_1x_3x_4\partial_{x_2}\partial_{x_3}^2 \in H_A(\boldsymbol{\beta}),$$

and

$$x_1x_3^2\partial_{x_2}^2\partial_{x_4} \in \text{in}_{(-\mathbf{w}, \mathbf{w})}(H_A(\boldsymbol{\beta})).$$

Since

$$x_1x_3^2\partial_{x_2}^2\partial_{x_4} \bullet x^{\mathbf{v}} = x_1x_3^2\partial_{x_2}^2\partial_{x_4} \bullet x_2^{-2}x_3^{-1}x_4 \neq 0,$$

$\mathbf{v}$  is not an exponent.

**Corollary 4.3.** *Assume that  $B$  is a basis of  $L$ . If  $|I \cup J| > |I_0|$  for any  $I \in \mathcal{N}$  and  $J \in \mathcal{N}^c$ , then  $\mathbf{v}$  is an exponent.*

*Proof.* For any  $I \in \mathcal{N}$  and  $J \in \mathcal{N}^c$ , we see that

$$|I \cup J \setminus K_{\mathcal{N}}| = |I \cup J| - |K_{\mathcal{N}}| > |I_0| - |K_{\mathcal{N}}| = |I_0 \setminus K_{\mathcal{N}}|$$

because both of  $I$  and  $I_0$  contain  $K_{\mathcal{N}}$ . Since the degree of  $m(\mathbf{s}) = (B\mathbf{s})^{I_0 \setminus K_{\mathcal{N}}}$  is less than that of any  $(B\mathbf{s})^{I \cup J \setminus \mathcal{N}}$ ,  $m(\mathbf{s})$  cannot belong to  $P_{\mathcal{N}}$ . By Proposition 4.1,  $\mathbf{v}$  is an exponent.  $\square$

**Theorem 4.4.** *Assume that  $B$  is a basis of  $L$ , and that  $P_{\mathcal{N}} = m(\mathbf{s}) \cdot P_B$ . Then  $\mathbf{v}$  is an exponent, and the set*

$$\{(q(\partial_{\mathbf{s}}) \bullet \widetilde{F}_{\mathcal{N}}(x, \mathbf{s}))|_{\mathbf{s}=\mathbf{0}} \mid q(\partial_{\mathbf{s}}) \in P_{\mathcal{N}}^{\perp}\}.$$

*spans the space of series solutions in the direction of  $\mathbf{w}$  to  $M_A(\boldsymbol{\beta})$  with exponent  $\mathbf{v}$ . In particular, for  $q(\partial_{\mathbf{s}}) \in P_{\mathcal{N}}^{\perp}$ , the solution  $(q(\partial_{\mathbf{s}}) \bullet \widetilde{F}_{\mathcal{N}}(x, \mathbf{s}))|_{\mathbf{s}=\mathbf{0}}$  has the starting term  $x^{\mathbf{v}}\Psi_B(m(\partial_{\mathbf{z}}) \star q(\partial_{\mathbf{s}}))(\log \mathbf{x})$ .*

*Proof.* By definition,  $P_B \neq \mathbb{C}[\mathbf{s}]$ . It follows from the assumption that  $m(\mathbf{s}) \notin P_{\mathcal{N}}$ . Hence, by Proposition 4.1,  $\mathbf{v}$  is an exponent. Moreover, by Proposition 3.11, the homomorphism (11) is surjective. Hence we have

$$\begin{aligned}
\dim_{\mathbb{C}}(P_{\mathcal{N}}^{\perp}) &\geq \dim_{\mathbb{C}}(P_B^{\perp}) = \dim_{\mathbb{C}}(Q_{\mathbf{v}}^{\perp}) = \dim_{\mathbb{C}}(\text{Sol}(\text{find}_{\mathbf{w}}(H_A(\boldsymbol{\beta}))_{\mathbf{v}})) \\
&\geq \dim_{\mathbb{C}}(\text{Sol}(\text{ind}_{\mathbf{w}}(H_A(\boldsymbol{\beta}))_{\mathbf{v}})).
\end{aligned}$$

By the regularity of  $M_A(\boldsymbol{\beta})$ , the dimension of the space of series solutions with the exponent  $\mathbf{v}$  coincides with  $\dim_{\mathbb{C}}(\text{Sol}(\text{ind}_{\mathbf{w}}(H_A(\boldsymbol{\beta}))_{\mathbf{v}}))$ . Hence, by Theorem 2.7, we have the former half of the assertion.

The latter half of the assertion follows from Proposition 3.10.  $\square$

**Example 4.5** (Continuation of Examples 2.5, 3.2, and 3.5). Consider the case where  $B = \{\mathbf{g}^{(1)}, \mathbf{g}^{(2)}\}$ . Then,  $B$  satisfies the assumption in Theorem 4.4. Recall that  $m(\mathbf{s}) = (B\mathbf{s})^{\emptyset} = 1$ , and

$$P_B = \langle (B\mathbf{s})^{G^{(1)}}, (B\mathbf{s})^{G^{(2)}} \rangle = \langle s_1^2, s_2^2 \rangle = P_{\mathcal{N}}.$$

Hence, we see by Proposition 4.1 that  $\mathbf{v}$  is an exponent, and that

$$\{1, \partial_{s_1}, \partial_{s_2}, \partial_{s_1} \partial_{s_2}\}$$

is a basis of  $P_B^{\perp}$ . Hence  $x^{\mathbf{v}} f(\log \mathbf{x})$  is a solution to  $\text{find}_{\mathbf{w}}(H_A(\boldsymbol{\beta}))_{\mathbf{v}}$  if and only if

$$f \in \langle 1, \mathbf{x}\mathbf{g}^{(1)}, \mathbf{x}\mathbf{g}^{(2)}, (\mathbf{x}\mathbf{g}^{(1)}) \cdot (\mathbf{x}\mathbf{g}^{(2)}) \rangle_{\mathbb{C}}.$$

By the uniqueness of an exponent, the above space coincides with the space of solutions to  $M_A(\boldsymbol{\beta})$ . Note that the holonomic rank of  $M_A(\boldsymbol{\beta})$  is four (cf. [8, Example 3.5.2]).

**Example 4.6.** [8, Examples 3.6.3, 3.6.11, 3.6.16] Let  $d = 3$ ,  $n = 9$  and

$$A = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 0 & 1 & 2 & 0 & 1 & 2 \\ 0 & 0 & 0 & 1 & 1 & 1 & 2 & 2 & 2 \end{bmatrix}.$$

Let  $\mathbf{w} = (2, 0, 0, 0, -1, 0, 0, 0, 2)$ ,  $\boldsymbol{\beta} = (1, 1, 1)^T = \mathbf{a}_5$ . Consider an exponent  $\mathbf{v} = (0, 0, 0, 0, 1, 0, 0, 0, 0)^T$ . Then  $K_{\mathcal{N}_B} = I_{\mathbf{0}} = \emptyset$ . The reduced Gröbner basis consists of the following twenty vectors

$$\{\mathbf{g}^{(1)} := (0, 1, -1, 0, -1, 1, 0, 0, 0)^T, \mathbf{g}^{(2)} := (0, 0, 0, 1, -1, 0, -1, 1, 0)^T,$$

$$\mathbf{g}^{(3)} := (0, 0, 0, 1, -2, 1, 0, 0, 0)^T, \mathbf{g}^{(4)} := (0, 1, 0, 0, -2, 0, 0, 1, 0)^T,$$

$$\mathbf{g}^{(5)} := (1, -1, 0, -1, 1, 0, 0, 0, 0), \mathbf{g}^{(6)} := (0, 0, 0, 0, 1, -1, 0, -1, 1),$$

$$\mathbf{g}^{(7)}, \dots, \mathbf{g}^{(20)}\}.$$

Hence we have

$$\begin{aligned} & \{G^{(i)} \mid i = 1, \dots, 20\} \\ & = \{G^{(1)} = \{2, 6\}, G^{(2)} = \{4, 8\}, G^{(3)} = \{4, 6\}, G^{(4)} = \{2, 8\}, \\ & \quad G^{(5)} = \{1\}, G^{(6)} = \{9\}, \{1, 8\}, \{2, 7\}, \{1, 3\}, \{1, 6\}, \{3, 4\}, \\ & \quad \{1, 9\}, \{3, 7\}, \{4, 9\}, \{6, 7\}, \{7, 9\}, \{2, 9\}, \{3, 8\}, \{3, 9\}, \{1, 7\}\}. \end{aligned}$$

Let

$$B = [\mathbf{g}^{(1)}, \mathbf{g}^{(2)}, \mathbf{g}^{(3)}, \mathbf{g}^{(4)}, \mathbf{g}^{(5)}, \mathbf{g}^{(6)}] = \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 & -1 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & -1 & 0 \\ -1 & -1 & -2 & -2 & 1 & 1 \\ 1 & 0 & 1 & 0 & 0 & -1 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Then

$$\begin{aligned} P_{\mathcal{N}} &= P_B \\ &= \langle (s_1 + s_4 - s_5)(s_1 + s_3 - s_6), (s_2 + s_3 - s_5)(s_2 + s_4 - s_6), \\ & \quad (s_2 + s_3 - s_5)(s_1 + s_3 - s_6), (s_1 + s_4 - s_5)(s_2 + s_4 - s_6), \\ & \quad s_5, s_6, s_2(s_1 + s_4 - s_5), s_1(s_2 + s_3 - s_5), \\ & \quad s_1 s_2, s_2(s_1 + s_3 - s_6), s_1(s_2 + s_4 - s_6) \rangle \\ &= \langle s_1 s_2, s_1 s_3, s_1 s_4, s_2 s_3, s_2 s_4, s_3^2, s_4^2, s_1^2 + s_3 s_4, s_2^2 + s_3 s_4, s_5, s_6 \rangle, \end{aligned}$$

where the last generator set gives the reduced Gröbner basis with respect to the lexicographic order  $<$  with  $s_1 > s_2 > s_3 > s_4 > s_5 > s_6$ .

We see that

$$\{\boldsymbol{\nu} \in \mathbb{N}^6 \mid \mathbf{s}^{\boldsymbol{\nu}} \notin \text{in}_{<}(P_B)\} = \{\mathbf{0}, \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4, \mathbf{e}_3 + \mathbf{e}_4\},$$

and hence we immediately have

$$q_{\mathbf{0}} = 1, q_{\mathbf{e}_1} = \partial_{s_1}, q_{\mathbf{e}_2} = \partial_{s_2}, q_{\mathbf{e}_3} = \partial_{s_3}, q_{\mathbf{e}_4} = \partial_{s_4}.$$

As to the last generator  $q_{\mathbf{e}_3 + \mathbf{e}_4}$ , since

$$c_{\boldsymbol{\mu}, \mathbf{e}_3 + \mathbf{e}_4} = \begin{cases} -1 & (\text{if } \boldsymbol{\mu} = 2\mathbf{e}_1, 2\mathbf{e}_2) \\ 0 & (\text{otherwise}) \end{cases},$$

we have

$$q_{\mathbf{e}_3 + \mathbf{e}_4} = \partial_{s_3} \partial_{s_4} + \sum_{\boldsymbol{\mu}; |\boldsymbol{\mu}|=2} c_{\boldsymbol{\mu}, \mathbf{e}_3 + \mathbf{e}_4} \frac{1}{\boldsymbol{\nu}!} \partial_{\mathbf{s}}^{\boldsymbol{\mu}} = \partial_{s_3} \partial_{s_4} - \frac{1}{2} \partial_{s_1}^2 - \frac{1}{2} \partial_{s_2}^2.$$



Hence a  $\mathbb{C}$ -basis of  $Q_v^\perp$  is given by

$$\{1, \mathbf{xg}^{(1)}, \mathbf{xg}^{(2)}, \mathbf{xg}^{(3)}, \mathbf{xg}^{(4)}, (\mathbf{xg}^{(3)}) \cdot (\mathbf{xg}^{(4)}) - \frac{1}{2}(\mathbf{xg}^{(1)})^2 - \frac{1}{2}(\mathbf{xg}^{(2)})^2\}.$$

## 5. AOMOTO-GEL'FAND SYSTEMS

In this section, let

$$A = \{\mathbf{a}_{i,j} \mid 1 \leq i \leq m, m+1 \leq j \leq m+l\},$$

where  $\mathbf{a}_{i,j} = \mathbf{e}_i + \mathbf{e}_j$ , and  $\{\mathbf{e}_1, \dots, \mathbf{e}_{l+m}\}$  is the standard basis of  $\mathbb{Z}^{l+m}$ .

Then  $\mathbb{Z}A = \{\mathbf{a} \in \mathbb{Z}^{l+m} \mid \sum_{i=1}^m a_i = \sum_{j=m+1}^{m+l} a_j\}$ ,  $\text{rank}(A) = m+l-1$ , and  $\text{rank}(L) = ml - (m+l-1) = (m-1)(l-1)$ , where

$$L = \{[c_{ij}]_{1 \leq i \leq m, m+1 \leq j \leq m+l} \in M_{m \times l}(\mathbb{Z}) \mid \sum_{i,j} c_{ij} \mathbf{a}_{ij} = \mathbf{0}\}.$$

Since  $A$  is normal, (that is,  $\mathbb{N}A = \mathbb{Z}A \cap \mathbb{R}_{\geq 0}A$ ),  $I_A$  is a Cohen-Macaulay ideal and hence  $\text{rank}(M_A(\boldsymbol{\beta})) = \text{vol}(A)$  for any  $\boldsymbol{\beta}$  (see [3]).

Take a weight vector  $\mathbf{w}$  satisfying  $w_{i,j} > w_{p,q}$  whenever  $(i,j) \neq (p,q)$ ,  $i \leq p$ , and  $j \leq q$ .

Then the reduced Gröbner basis of  $I_A$  with respect to  $\mathbf{w}$  equals

$$G := \{\underline{\partial^{(\mathbf{g}_{(p,q)}^{(i,j)})^+}} - \partial^{(\mathbf{g}_{(p,q)}^{(i,j)})^-} \mid i < p, j < q\},$$

and

$$\text{in}_{\mathbf{w}}(I_A) = \langle \partial_{i,j} \partial_{p,q} \mid i < p, j < q \rangle,$$

where  $\mathbf{g}_{(p,q)}^{(i,j)} := E_{i,j} + E_{p,q} - E_{i,q} - E_{p,j} \in L$  and  $E_{i,j}$  are matrix units.

The weight  $\mathbf{w}$  induces a staircase regular triangulation, which is unimodular (cf. [10, Example 8.12]); for example, let  $m = 2, l = 4$ , the standard pairs are

$$\begin{bmatrix} * & * & * & * \\ * & 0 & 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & * & * & * \\ * & * & 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 & * & * \\ * & * & * & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 & 0 & * \\ * & * & * & * \end{bmatrix},$$

where let the row-numbers be  $1, \dots, m$  and the column-numbers  $m+1, \dots, m+l$ .

For general  $l, m$ , the standard pairs correspond to the paths from the southwest corner to the northeast corner going only northward or eastward. In this way, we see

$$\text{vol}(A) = \binom{m+l-2}{m-1}.$$

Let

$$B := \{\mathbf{b}^{(i,j)} := \mathbf{g}_{(i+1,j+1)}^{(i,j)} \mid 1 \leq i < m, m+1 \leq j < m+l\}.$$

Then  $B$  is a basis of  $L$  and  $\text{supp}(B) = \{1, \dots, m+l\}$ , hence  $B$  satisfies Assumption 2.2. Let  $\mathbf{s} = (s_{(i,j)})_{1 \leq i < m, m+1 \leq j < m+l}$  be indeterminates such that  $s_{(i,j)}$  corresponds to  $\mathbf{b}^{(i,j)}$ . For convenience, set  $s_{(i,j)} := 0$  unless  $(i,j) \in \{1, \dots, m-1\} \times \{m+1, \dots, m+l-1\}$ . Then

$$\begin{aligned} (B\mathbf{s})_{(\mu,\nu)} &= \sum_{\substack{1 \leq i < m \\ m+1 \leq j < m+l}} s_{(i,j)} (\mathbf{g}_{(i+1,j+1)}^{(i,j)})_{(\mu,\nu)} \\ &= s_{(\mu,\nu)} - s_{(\mu,\nu-1)} - s_{(\mu-1,\nu)} + s_{(\mu-1,\nu-1)}. \end{aligned}$$

**Lemma 5.1.** *Let  $\boldsymbol{\beta} = \mathbf{0}$ . Then  $\mathbf{v} = \mathbf{0}$  is a unique exponent.*

*Proof.* Since  $\text{in}_{\mathbf{w}}(I_A)$  is square-free, every fake exponent is an exponent by Theorem 3.6.6 in [8].

Let  $\mathbf{v}$  be an exponent. Then there exists a standard pair  $(\mathbf{a}, \sigma) = (\mathbf{0}, \sigma)$  corresponding to  $\mathbf{v}$  such that

$$v_j = 0 \quad (j \notin \sigma), \quad A\mathbf{v} = \boldsymbol{\beta} = \mathbf{0}.$$

Since the submatrix  $A_\sigma = (\mathbf{a}_j)_{j \in \sigma}$  is invertible and satisfies  $A_\sigma \mathbf{v}_\sigma = \mathbf{0}$ , we have  $\mathbf{v} = \mathbf{0}$ .  $\square$

From now on, let  $\boldsymbol{\beta} = \mathbf{0}$  and  $\mathbf{v} = \mathbf{0}$ . Hence  $I_{\mathbf{0}} = \emptyset = K_{\mathcal{N}_B}$ .

**Lemma 5.2.** (i)  $\{(ij), (pq)\} \in \mathcal{N}^c = \text{NS}_{\mathbf{w}}(\mathbf{v})^c$  for  $i < p, j < q$ .  
(ii) Let  $J \in \mathcal{N}^c = \text{NS}_{\mathbf{w}}(\mathbf{v})^c$ . Then there exist  $i < p, j < q$  such that  $J \supset \{(ij), (pq)\}$ .

*Proof.* (i) This follows from  $G_{(pq)}^{(ij)} = \text{nsupp}(\mathbf{v} - \mathbf{g}_{(pq)}^{(ij)}) = \text{nsupp}(-\mathbf{g}_{(pq)}^{(ij)}) = \{(ij), (pq)\}$ .

(ii) This is immediate from Proposition 3.3.  $\square$

**Proposition 5.3.**

$$P_B = \langle (B\mathbf{s})^{\{(i,j),(p,q)\}} \mid i < p, j < q \rangle = \langle s_{(i,j)} s_{(p,q)} \mid i \leq p, j \leq q \rangle.$$

*Proof.* The first equality follows from Lemma 5.2. We show the second equality. Since, for any  $i, j, p, q$  with  $i < p$  and  $j < q$ ,

$$\begin{aligned} (B\mathbf{s})^{\{(i,j),(p,q)\}} &= (s_{(i,j)} - s_{(i,j-1)} - s_{(i-1,j)} + s_{(i-1,j-1)}) \\ &\quad \times (s_{(p,q)} - s_{(p,q-1)} - s_{(p-1,q)} + s_{(p-1,q-1)}) \in \text{RHS}, \end{aligned}$$

we only need to show the reverse inclusion. We introduce the total order  $<$  on  $\mathcal{X} := \{((i,j), (p,q)) \mid 1 \leq i \leq p < m, m+1 \leq j \leq q < m+l\}$

defined by

$$\begin{aligned} ((i, j), (p, q)) &< ((i', j'), (p', q')) \\ \iff &\begin{cases} i < i', \\ \text{or } [i = i' \text{ and } j < j'], \\ \text{or } [(i, j) = (i', j') \text{ and } p > p'], \\ \text{or } [(i, j, p) = (i', j', p') \text{ and } q > q']. \end{cases} \end{aligned}$$

We prove by induction on the totally ordered set  $(\mathcal{X}, <)$ .

First, for the minimum element  $((1, m+1), (m-1, m+l-1))$ , we have

$$s_{(1, m+1)} s_{(m-1, m+l-1)} = (B\mathbf{s})^{\{(1, m+1), (m, m+l)\}} \in \text{LHS}.$$

Next, fix  $((i, j), (p, q)) \neq ((1, m+1), (m-1, m+l-1))$  with  $1 \leq i \leq p < m$  and  $m+1 \leq j \leq q < m+l$ . Suppose that  $s_{(i', j')} s_{(p', q')} \in \text{LHS}$  for all  $((i', j'), (p', q')) < ((i, j), (p, q))$ . Then the leading monomial in

$$\begin{aligned} (\text{LHS} \ni) (B\mathbf{s})^{\{(i, j), (p+1, q+1)\}} &= (s_{(i, j)} - s_{(i, j-1)} - s_{(i-1, j)} + s_{(i-1, j-1)}) \\ &\quad \times (s_{(p+1, q+1)} - s_{(p+1, q)} - s_{(p, q+1)} + s_{(p, q)}) \end{aligned}$$

is  $s_{(i, j)} s_{(p, q)}$ . It follows from the induction hypothesis that  $s_{(i, j)} s_{(p, q)} \in \text{LHS}$ . Hence we have the assertion.  $\square$

#### Corollary 5.4.

$$\begin{aligned} P_B^\perp &= \langle \partial_{\mathbf{s}}^{\mathbf{p}} \mid \mathbf{s}^{\mathbf{p}} \notin \langle s_{(i, j)} s_{(p, q)} \mid i \leq p, j \leq q \rangle_{\mathbb{C}} \rangle_{\mathbb{C}} \\ &= \langle \partial_{s_{(i_1, j_1)}} \cdots \partial_{s_{(i_r, j_r)}} \mid i_1 < \cdots < i_r, j_1 > \cdots > j_r \rangle_{\mathbb{C}}. \end{aligned}$$

Furthermore,

$$\dim_{\mathbb{C}}(P_B^\perp) = \sum_{r=0}^{\min\{l-1, m-1\}} \binom{l-1}{r} \binom{m-1}{r} = \binom{l+m-2}{m-1}.$$

*Proof.* The first part follows from Proposition 5.3. For the second part, compare the coefficients of  $z^{m-1}$  in the following:

$$\begin{aligned} \sum_{k=0}^{l+m-2} \binom{l+m-2}{k} z^k &= (1+z)^{l+m-2} = (1+z)^{l-1} (1+z)^{m-1} \\ &= \left( \sum_{p=0}^{l-1} \binom{l-1}{p} z^p \right) \left( \sum_{q=0}^{m-1} \binom{m-1}{q} z^{m-1-q} \right) \\ &= \sum_{p, q} \binom{l-1}{p} \binom{m-1}{q} z^{m-1-q+p}. \end{aligned}$$

□

**Corollary 5.5.**

$\{((\partial_{s_{(i_1, j_1)}} \cdots \partial_{s_{(i_r, j_r)}}) \bullet F_{\mathcal{N}}(\mathbf{x}, \mathbf{s}))|_{\mathbf{s}=\mathbf{0}} \mid i_1 < \cdots < i_r, j_1 > \cdots > j_r\}$   
forms a fundamental system of solutions to  $M_A(\mathbf{0})$ .

**Proposition 5.6.**

$$Q_{\mathbf{0}}^{\perp} = \left\langle \prod_{k=1}^r (\mathbf{x} \mathbf{g}_{(i_k+1, j_k+1)}^{(i_k, j_k)}) \mid \begin{array}{l} i_1 < i_2 < \cdots < i_r \\ j_1 > j_2 > \cdots > j_r \end{array} \right\rangle_{\mathbb{C}}.$$

*Proof.* This is immediate from Theorem 3.14 and Corollary 5.4. □

**Example 5.7.** Let  $m = 2$ . This case corresponds to the Lauricella's  $F_D$  (e.g. see [1, §3.1.3]). Then  $\text{vol}(A) = \binom{l+m-2}{m-1} = l$ , and

$$P_B^{\perp} = \langle 1, \partial_{s_{(1,3)}}, \partial_{s_{(1,4)}}, \dots, \partial_{s_{(1, l+1)}} \rangle_{\mathbb{C}}.$$

**Example 5.8.** Let  $l = m = 3$ . Then  $\text{vol}(A) = \binom{l+m-2}{m-1} = \binom{4}{2} = 6$ , and

$$P_B^{\perp} = \langle 1, \partial_{s_{(1,4)}}, \partial_{s_{(1,5)}}, \partial_{s_{(2,4)}}, \partial_{s_{(2,5)}}, \partial_{s_{(1,5)}} \partial_{s_{(2,4)}} \rangle_{\mathbb{C}}.$$

6. LAURICELLA'S  $F_C$ 

Let

$$A := \{\mathbf{a}_i := \mathbf{e}_0 + \mathbf{e}_i, \mathbf{a}_{-i} := \mathbf{e}_0 - \mathbf{e}_i \mid i = 1, 2, \dots, m\}.$$

In this case, the  $A$ -hypergeometric systems correspond to Lauricella's  $F_C$  [7]. Then

$$\mathbb{Z}A = \{\mathbf{a} \in \mathbb{Z}^{m+1} \mid \sum_{i=0}^m a_i \in 2\mathbb{Z}\},$$

and

$$L = \{\mathbf{l} \in \mathbb{Z}^{2m} \mid \sum_{i=\pm 1, \dots, \pm m} l_i = 0, l_i - l_{-i} = 0 \ (1 \leq i \leq m)\}.$$

We have  $\text{rank}(L) = 2m - (m + 1) = m - 1$ .

Take a weight  $\mathbf{w}$  so that

$$w_1 + w_{-1} > w_2 + w_{-2} > \cdots > w_m + w_{-m}.$$

Then

$$\text{in}_{\mathbf{w}}(I_A) = \langle \partial_1 \partial_{-1}, \partial_2 \partial_{-2}, \dots, \partial_{m-1} \partial_{-(m-1)} \rangle,$$

and the reduced Gröbner basis  $G$  is given by

$$G = \{\mathbf{g}^{(1)}, \mathbf{g}^{(2)}, \dots, \mathbf{g}^{(m-1)}\},$$

where  $\mathbf{g}^{(i)} = \mathbf{e}_i + \mathbf{e}_{-i} - \mathbf{e}_m - \mathbf{e}_{-m}$ . We see that  $G$  is a basis of the free  $\mathbb{Z}$ -module  $L$ . Set  $B := G$ , then  $B$  satisfies Assumption 2.2. Note that  $\text{supp}(B) = \{\pm 1, \dots, \pm m\}$ . Let  $\mathbf{s} = (s_i)_{1 \leq i \leq m-1}$  be indeterminates such that  $s_i$  corresponds to  $\mathbf{b}^{(i)} = \mathbf{g}^{(i)}$ . The standard pairs are pairs of  $*$ -place  $\{\epsilon(i)i \mid i \in [1, m-1]\} \cup \{\pm m\}$  ( $\epsilon : [1, m-1] \rightarrow \{\pm 1\}$ ) and 0-place its complement.

Hence  $\text{vol}(A) = 2^{m-1}$ .

**Lemma 6.1.** *Let  $\beta = \mathbf{0}$ . Then  $\mathbf{v} = \mathbf{0}$  is a unique exponent.*

*Proof.* The proof is similar to Lemma 5.1. □

**Proposition 6.2.**

$$P_B = \langle (B\mathbf{s})^{\{\pm i\}} \mid i = 1, \dots, m-1 \rangle = \langle s_i^2 \mid 1 \leq i \leq m-1 \rangle.$$

*Proof.* We have  $G^{(i)} = \{\pm i\}$  and

$$(B\mathbf{s})^{\{\pm i\}} = \left( \sum_{\nu=1}^{m-1} s_\nu \mathbf{g}_{+i}^{(\nu)} \right) \left( \sum_{\nu=1}^{m-1} s_\nu \mathbf{g}_{-i}^{(\nu)} \right) = s_i^2.$$

□

**Proposition 6.3.**

$$\begin{aligned} P_B^\perp &= \langle \partial_{\mathbf{s}}^{\mathbf{p}} \mid \mathbf{s}^{\mathbf{p}} \notin \langle s_i^2 \mid 1 \leq i \leq m-1 \rangle_{\mathbb{C}} \rangle_{\mathbb{C}} \\ &= \langle \partial_{\mathbf{s}}^I = \prod_{i \in I} \partial_{s_i} \mid I \subset \{1, \dots, m-1\} \rangle_{\mathbb{C}}, \end{aligned}$$

and

$$Q_{\mathbf{0}}^\perp = \langle \prod_{i \in I} (\mathbf{x} \mathbf{g}^{(i)}) \mid I \subset \{1, \dots, m-1\} \rangle_{\mathbb{C}}.$$

Furthermore,

$$\dim_{\mathbb{C}}(Q_{\mathbf{0}}^\perp) = 2^{m-1}.$$

*Proof.* This is immediate from Lemma 3.6, Theorem 3.14, and Proposition 6.2. □

**Corollary 6.4.**  $\{((\prod_{i \in I} \partial_{s_i}) \bullet F_{\mathcal{N}}(\mathbf{x}, \mathbf{s}))|_{\mathbf{s}=\mathbf{0}} \mid I \subset [1, m-1]\}$  forms a basis of solutions of  $M_A(\mathbf{0})$ .

## REFERENCES

- [1] Aomoto, K., Kita, M.: *Theory of hypergeometric functions*, Springer, New York, 2011.
- [2] Gel'fand, I. M., Graev, M. I., Zelevinskii, A. V.: Holonomic systems of equations and series of hypergeometric type, *Soviet Mathematics Doklady* 36 (1988) 5–10.

- [3] Gel'fand, I. M., Zelevinskii, A. V., Kapranov, M. M.: Hypergeometric functions and toral manifolds, *Funktsional. Anal. i Prilozhen* 23 (1989) 12–26; *Funct. Anal. Appl.* (English translation), 23 (1989) 94–106.
- [4] Hotta, R.: Equivariant  $D$ -modules, *Proceeding of ICPAM Spring School in Wuhan, 1991*, available at [arXiv:math/9805021](https://arxiv.org/abs/math/9805021).
- [5] Saito, M.: Logarithm-free  $A$ -hypergeometric series. *Duke Math. J.* **115** (2002) 53–73.
- [6] Saito, M.: Logarithmic  $A$ -hypergeometric series. *International Journal of Mathematics* No. 31, Issue No. 13, Article No. 2050110, Year 2020.
- [7] Saito, M.: Contiguity Relations for the Lauricella Functions, *Funkcialaj Ekvacioj*, **38** (1995), 37–58.
- [8] Saito, M., Sturmfels, B. and Takayama, N.: *Gröbner deformations of hypergeometric differential equations*. *Algorithms and Computation in Mathematics*, vol. 6, Springer, New York, 2000.
- [9] Schulze, M. and Walther, U.: Irregularity of hypergeometric systems via slopes along coordinate subspaces. *Duke Math. J.* **142** (2008) 465–509.
- [10] Sturmfels, B.: *Gröbner bases and convex polytopes* *University Lecture Notes*, vol. 8, American Mathematical Society, Providence, 1995.

(GO OKUYAMA) HIGHER EDUCATION SUPPORT CENTER

HOKKAIDO UNIVERSITY OF SCIENCE

SAPPORO, 006-8585, JAPAN,

(MUTSUMI SAITO) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE

HOKKAIDO UNIVERSITY

SAPPORO, 060-0810, JAPAN

*Email address:* [gokuyama@hus.ac.jp](mailto:gokuyama@hus.ac.jp) (G.Okuyama), [saito@math.sci.hokudai.ac.jp](mailto:saito@math.sci.hokudai.ac.jp) (M.Saito)