

TROPICAL MEDIANS BY TRANSPORTATION

ANDREI COMĂNECI AND MICHAEL JOSWIG

ABSTRACT. Fermat–Weber points with respect to an asymmetric tropical distance function are studied. It turns out that they correspond to the optimal solutions of a transportation problem. The results are applied to obtain a new method for computing consensus trees in phylogenetics. This method has several desirable properties; e.g., it is Pareto and co-Pareto on rooted triplets.

1. INTRODUCTION

The following optimization problem can be studied in any metric space. Given a finite number of points, sometimes called *sites*, find a point which minimizes the sum of the distances to the sites. Such a point is called a *Fermat–Weber* point, and this is some version of a geometric median of the sites, which is known to be robust in a certain sense [7, §21]. Computing Fermat–Weber points is a rich topic with a remarkable history; see [7, Chapter II]. Here we consider a specific distance function, d_Δ , which occurs in tropical geometry [2, 23]. This function is asymmetric, i.e., $d_\Delta(a, b)$ may differ from $d_\Delta(b, a)$. So we call d_Δ the *asymmetric tropical distance*, to differentiate from the *symmetric tropical distance*, which is more common [16, §5.3]. The symmetric tropical Fermat–Weber problem was investigated by Lin, Sturmfels, Tang and Yoshida [20] and Lin and Yoshida [21].

As our main result we prove that one (asymmetric tropical) Fermat–Weber point can be computed by solving a transportation problem. The transportation problem is an optimization classic, with numerous applications, both in theory and practice. For an overview we refer to Schrijver’s monograph [27, §21.6]; see also the survey by De Loera and Kim for the polyhedral geometry point of view [12]. Efficient methods for solving transportation problems include algorithms by Tokuyama and Nakano [30], Kleinschmidt and Schannath [18], and Brenner [8]. In general, Fermat–Weber points are not unique, so one part of the present work is devoted to understanding the entire (asymmetric tropical) *Fermat–Weber set* for a given set of sites. This is tightly related to the study of tropical hyperplane arrangements and tropical convex hulls, which are at the core of tropical combinatorics [16]. The latter subject is concerned with the rich interplay between tropical geometry and optimization.

One motivation for studying the Fermat–Weber problem in the setting of tropical geometry comes from phylogenetics [20, 21]. In that field, a part of computational biology, the goal is to associate trees to input data. A typical example are trees encoding ancestral relations among species, where the data originates from strands of DNA. In tropical geometry spaces of metric trees with n labeled leaves occur naturally as the tropical Grassmannians $T\text{Gr}(2, n)$; see [22, §4.3] and [16, §10.6]. In phylogenetics many different

2020 *Mathematics Subject Classification.* 90C08 (14T15, 46B20, 92B10).

Key words and phrases. Fermat–Weber points; tropical convexity; polyhedral norms; consensus trees.

Support by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy – The Berlin Mathematics Research Center MATH⁺ (EXC-2046/1, project ID 390685689) gratefully acknowledged. M. Joswig has further been supported by “Symbolic Tools in Mathematics and their Application” (TRR 195, project-ID 286237555). A. Comăneci was supported by “Facets of Complexity” (GRK 2434, project-ID 385256563).

methods are known to construct trees from a fixed data set. Since those methods usually do not come up with the same tree, there is a need to find the common ground. This gives rise to some *consensus tree*, which describes where the several methods agree. Finding consensus trees is a topic of its own [9], and the authors of [20] argue that “tropical convexity and tropical linear algebra . . . behave better” than other methods. Here we show that passing from the symmetric tropical distance function to its asymmetric sibling leads to a new method for computing metric consensus trees which is even better behaved. This is because the asymmetric tropical Fermat–Weber sets are nicer geometrically.

Our paper is organized as follows. We start out with a brief summary of facts from tropical combinatorics which are relevant for the Fermat–Weber problem. Then we prove that the Fermat–Weber set arises as a cell in the covector decomposition of the tropical torus induced by the sites. That covector cell is then characterized in several ways. The first approach employs regular subdivisions of products of simplices; see [13, §6.2]. That leads to a linear programming formulation of the Fermat–Weber problem, and the dual linear program is a transportation problem. The latter then provides efficient algorithms. The final chapter is devoted to computing Fermat–Weber sets in spaces of equidistant trees.

Related work. As an early contribution of tropical geometry to data science Gärtner and Jaggi [15] developed a concept for “tropical support vector machines”, with applications to classification in mind. A different train of thought was developed by Pachter and Sturmfels [24, §2.4] who related tropical geometry to phylogenetic trees. Later, Lin et al. [20] connected these ideas to the geometry of tree spaces studied by Billera, Holmes and Vogtman [5]. Yoshida, Zhang and Zhang [31] proposed a method to analyze data, which they call “tropical principal component analysis”. In a way, the latter may be viewed as a synthesis of the above. A key contribution here is work of Ardila and Klivans, who saw that spaces of equidistant trees form the Bergman fans of the graphic matroids of complete graphs [3]. The term “tropical convexity” was coined by Develin and Sturmfels [14] to connect tropical geometry with the older topic of $(\max, +)$ -linear algebra [10].

2. TROPICAL CONVEXITY

The purpose of this section is to set our notation and to collect a few facts which are key to our methods; for the details we refer to [16]. We consider the *tropical semiring* $\mathbb{T}^{\max} = (\mathbb{R} \cup \{-\infty\}, \oplus, \odot)$ with $\oplus = \max$ as the tropical addition and $\odot = +$ as the tropical multiplication. The additive neutral element is $-\infty$, and 0 is neutral with respect to the multiplication. Usually, we abbreviate $\mathbb{T} = \mathbb{T}^{\max}$. The set \mathbb{T}^n inherits the structure of a semimodule by componentwise tropical addition and tropical scalar multiplication.

A *tropical cone* in \mathbb{T}^n is a nonempty subset C which contains each tropical linear combination $\lambda \odot x \oplus \mu \odot y$ for $\lambda, \mu \in \mathbb{T}$ and $x, y \in C$. Each tropical cone contains the point $-\infty \mathbf{1}$ and the entire set $\mathbb{R} \mathbf{1}$, where $\mathbf{1}$ is the all-ones vector. Therefore it is convenient to pass to the quotient $\mathbb{TP}^{n-1} := (\mathbb{T}^n \setminus \{-\infty \mathbf{1}\}) / \mathbb{R} \mathbf{1}$, which is called the *tropical projective space*. A subset of \mathbb{TP}^{n-1} is *tropically convex* if it arises as the image of a tropical cone under the canonical projection. A *tropical polytope* is a finitely generated tropically convex set. The *tropical projective torus* $\mathbb{R}^n / \mathbb{R} \mathbf{1}$ is the subset of points in \mathbb{TP}^{n-1} with finite coordinates. We say that a tropical polytope is *bounded* if it lies in $\mathbb{R}^n / \mathbb{R} \mathbf{1}$.

Tropical convexity is intimately related to ordinary convexity, polyhedral geometry and (linear) optimization. For instance, tropical polytopes arise as the images of ordinary convex polytopes over ordered fields of real Puiseux series under the valuation map. Yet, here the following less algebraic description is more relevant.




FIGURE 1. Five sites in $\mathbb{R}^3/\mathbb{R}\mathbb{1}$, their max-tropical convex hull, the induced min-tropical hyperplane arrangement, and the unique Fermat–Weber point (marked white)

We consider an arbitrary $m \times n$ -matrix $V = (v_{ij})$ with real coefficients. Each row $v_i = (v_{i1}, \dots, v_{in})$ is a point in \mathbb{R}^n (or $\mathbb{R}^n/\mathbb{R}\mathbb{1}$, if we ignore the tropical scaling). So V may be viewed as a configuration of m labeled points in $\mathbb{R}^n/\mathbb{R}\mathbb{1}$. Technically, it is convenient to assume that each ordinary row sum equals zero, i.e., each row lies in the set

$$\mathcal{H} = \{x \in \mathbb{R}^n \mid x_1 + x_2 + \dots + x_n = 0\} .$$

Observe that each point in $\mathbb{R}^n/\mathbb{R}\mathbb{1}$ has a unique representative in \mathbb{R}^n which lies in \mathcal{H} . So we can identify the tropical projective torus $\mathbb{R}^n/\mathbb{R}\mathbb{1}$ with the ordinary linear hyperplane \mathcal{H} in \mathbb{R}^n . This also works topologically, since the quotient vector space $\mathbb{R}^n/\mathbb{R}\mathbb{1}$ is homeomorphic to \mathbb{R}^{n-1} (and thus with \mathcal{H} considered as a subset of \mathbb{R}^n). The tropical projective space \mathbb{TP}^{n-1} is a compactification of $\mathbb{R}^n/\mathbb{R}\mathbb{1}$, where the boundary comprises those points which have at least one infinite coordinate.

Adding vectors tropically works coefficient-wise, and there is also tropical multiplication by scalars. With these notions, the *max-tropical convex hull* of (the rows of) V is

$$\text{tconv}^{\max}(V) := \{\lambda_1 \odot v_1 \oplus \dots \oplus \lambda_m \odot v_m \mid \lambda_i \in \mathbb{R}\} + \mathbb{R}\mathbb{1} ,$$

which is a subset of $\mathbb{R}^n/\mathbb{R}\mathbb{1}$. The rows of the matrix V also define an arrangement of m tropical hyperplanes in $\mathbb{R}^n/\mathbb{R}\mathbb{1}$, with respect to min as the tropical addition, and we denote this $\mathcal{T}(V)$. In this context each row arises as the *apex* of a min-tropical hyperplane. Everything that we explained above also works for the min-tropical semiring $\mathbb{T}^{\min} = (\mathbb{R} \cup \{\infty\}, \min, +)$, which is isomorphic to \mathbb{T}^{\max} as a semiring via $x \mapsto -x$. Observe that this involution leaves the hyperplane \mathcal{H} invariant. The min-tropical hyperplane arrangement with the rows of V as their apices induces an ordinary polyhedral subdivision, $\text{CovDec}(V)$, of $\mathbb{R}^n/\mathbb{R}\mathbb{1}$ (or, equivalently, \mathcal{H}) into cells which are convex in three different senses: they are max-tropically convex, min-tropically convex and convex in the ordinary sense (as subsets of \mathcal{H}). Such polyhedra are known as *polytropes*, and they may be bounded or




FIGURE 2. Mixed subdivision $\mathcal{S}(V)$ of $5 \cdot \Delta_2$ for V as in Example 1. The 21 lattice points of $5 \cdot \Delta_2$ are marked with their coordinates. The min-tropical hyperplane arrangement $\mathcal{T}(V)$ admits a piecewise-linear embedding

unbounded. As ordinary polyhedra, the polytopes are characterized by the property that their facet normal directions are $e_i - e_j$ for $i, j \in [n]$ distinct. The subdivision $\text{CovDec}(V)$ is the *covector subdivision* induced by (the rows of) V .

Example 1. We illustrate the various concepts from tropical convexity for the matrix $V \in \mathbb{R}^{5 \times 3}$ whose transpose reads

$$V^\top = \begin{pmatrix} 14 & 13 & 11 & 10 & 3 \\ -7 & -14 & -13 & 1 & -3 \\ -7 & 1 & 2 & -11 & 0 \end{pmatrix}.$$

The rows of V (equivalently, the columns of V^\top) encode five points in $\mathbb{R}^3/\mathbb{R}\mathbf{1}$; see Figure 1. The covector decomposition has 15 regions of maximal dimension 2, and six of them are bounded. The max-tropical convex hull is a non-pure two-dimensional ordinary polytopal complex, which is shaded gray; it also contains the green line segment extending from $(3, -3, 0)$ to the lower left. Each covector cell is a polytrope, and it is bounded if and only if it is contained in the max-tropical convex hull of the rows of V .

We consider the *envelope*

$$(1) \quad \mathcal{E}(V) := \{(t, x) \in \mathbb{R}^m \times \mathbb{R}^n \mid t_i + x_j \geq v_{ij}\},$$

which is an unbounded ordinary polyhedron. By [16, Theorem 6.14] the cells of $\text{CovDec}(V)$ arise as the images of faces of $\mathcal{E}(V)$ under the coordinate projection $(t, x) \mapsto x$. Moreover, $\text{CovDec}(V)$ is dual to the regular subdivision, $\Sigma(V)$, of the product of simplices $\Delta_{m-1} \times \Delta_{n-1} = \text{conv}\{(e_i, e_j) \mid i \in [m], j \in [n]\}$ obtained from lifting (e_i, e_j) to the height v_{ij} . Here we take regular subdivisions induced by upper convex hulls since max is our tropical addition. For the same reason the inequality sign “ \geq ” is reversed in comparison with the min-tropical version in [16, (6.1)]. Via the Cayley trick the subdivision $\Sigma(V)$ of $\Delta_{m-1} \times \Delta_{n-1}$ corresponds to a mixed subdivision, $\mathcal{S}(V)$, of the dilated simplex $m \cdot \Delta_{n-1}$; see [13, §9.2] and [16, §4.5]. For instance, this is convenient for properly visualizing $\Sigma(V)$, which is a polyhedral complex of dimension $(m-1)(n-1)$. Figure 2 shows the mixed subdivision $\mathcal{S}(V)$ of $5 \cdot \Delta_2$ for the matrix V from Example 1.

3. FERMAT–WEBER SETS

We examine the Fermat–Weber problem through tropical combinatorics and polyhedral geometry. As our first key observation we show that asymmetric tropical Fermat–Weber sets arise as specific cells in the covector subdivisions induced by the sites.

The *asymmetric tropical distance* in \mathbb{R}^n is given by

$$(2) \quad d_{\Delta}(a, b) = \sum_{i \in [n]} (b_i - a_i) - n \min_{i \in [n]} (b_i - a_i) = \sum_{i \in [n]} (b_i - a_i) + n \max_{i \in [n]} (a_i - b_i) ,$$

where $a, b \in \mathbb{R}^n$. Since $d_{\Delta}(a', b') = d_{\Delta}(a, b)$ for $a - a' \in \mathbb{R}\mathbf{1}$ and $b - b' \in \mathbb{R}\mathbf{1}$, this induces a directed distance function in the $(n-1)$ -dimensional quotient vector space $\mathbb{R}^n/\mathbb{R}\mathbf{1}$. We do not distinguish between the function $d_{\Delta} : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$ and the induced function on $\mathbb{R}^n/\mathbb{R}\mathbf{1}$. The asymmetric tropical distance is a “polyhedral norm” with respect to the standard simplex $\Delta_{n-1} := \text{conv}\{e_1, \dots, e_n\} + \mathbb{R}\mathbf{1}$ in $\mathbb{R}^n/\mathbb{R}\mathbf{1}$; see [7, §20]. This may also be seen as a rescaled version of the “tropical Funk metric” discussed in [1, §3.3]. More common in tropical geometry is the *symmetric tropical distance* between $a, b \in \mathbb{R}^n$ (or $\mathbb{R}^n/\mathbb{R}\mathbf{1}$). It is defined as

$$\text{dist}(a, b) = \max_{i \in [n]} (a_i - b_i) - \min_{j \in [n]} (a_j - b_j) = \max_{i, j \in [n]} (a_i - b_i - a_j + b_j) ;$$

see [16, §5.3]. We have $\text{dist}(a, b) = \frac{1}{n}(d_{\Delta}(a, b) + d_{\Delta}(b, a))$. Throughout this section we fix a finite set of sites $V = \{v_1, v_2, \dots, v_m\}$ in \mathcal{H} , which we identify with $\mathbb{R}^n/\mathbb{R}\mathbf{1}$.

Definition 2. An (*asymmetric tropical*) *Fermat–Weber point* with respect to V is a point in \mathcal{H} which minimizes the sum of the asymmetric tropical distances from these sites.

In general such a point is not unique. Hence, we let $\text{FW}(V)$ denote the set of all asymmetric tropical Fermat–Weber points and call it the (*asymmetric tropical*) *Fermat–Weber set* with respect to V . This is the asymmetric analog of the symmetric tropical Fermat–Weber set studied in [21].

Fixing the site $v_i \in V$, the distance function from v_i , which reads

$$(3) \quad d_{\Delta}(v_i, x) = n \max_{j \in [n]} (v_{ij} - x_j) \quad \text{for } x \in \mathcal{H} ,$$

is convex in the ordinary sense and piecewise linear. Its regions of linearity are precisely the n closed sectors of the min-tropical hyperplane with apex v_i ; see [16, §5.5]. Consequently, the common subdivision of the regions of linearity of the sites is exactly the covector decomposition $\text{CovDec}(V)$. Our first main theorem shows that the Fermat–Weber set $\text{FW}(V)$ is a bounded cell of that subdivision.

For the sake of a precise formulation of that result, we pass to the regular triangulation, $\Sigma(V)$, of $\Delta_{m-1} \times \Delta_{n-1}$ which is dual to the covector subdivision $\text{CovDec}(V)$. The relatively open cells of $\Sigma(V)$ partition the product of the ordinary polytope $\Delta_{m-1} \times \Delta_{n-1}$. The point $(\frac{1}{m}\mathbf{1}, \frac{1}{n}\mathbf{1})$ is the vertex barycenter of $\Delta_{m-1} \times \Delta_{n-1}$. So there is a unique cell, C_V , which contains that vertex barycenter in its relatively interior. We call C_V the *central cell* of $\Sigma(V)$ and its dual in $\text{CovDec}(C)$ the *central covector cell*.

Theorem 3. *The Fermat–Weber set $\text{FW}(V)$ agrees with the central covector cell in $\text{CovDec}(V)$. In particular, $\text{FW}(V)$ is a bounded polytrope in \mathcal{H} , and it is contained in the tropical polytope $\text{tconv}^{\max}(V)$.*

Proof. Consider the linear program

$$(4) \quad \begin{array}{ll} \text{minimize} & n \cdot (t_1 + \dots + t_m) \\ \text{subject to} & v_{ij} - x_j \leq t_i, \quad \text{for } i \in [m] \text{ and } j \in [n] \\ & x_1 + \dots + x_n = 0 \end{array}$$

with $mn+1$ constraints in the $m+n$ variables $t_1, t_2, \dots, t_m, x_1, x_2, \dots, x_n$. The coefficients v_{ij} are the coordinates of the sites. The constant factor n in the objective function is not relevant here, but it does make the dual linear program (5) studied below look more natural. If (t^*, x^*) is an optimal solution of (4), then $x^* \in \text{FW}(V)$, and $t_i^* = \frac{1}{n}d_\Delta(v_i, x^*)$. Conversely, each Fermat–Weber point arises in this way. The constraints $v_{ij} \leq t_i + x_j$ describe the max-tropical version of the envelope $\mathcal{E}(V)$; see [16, §6.1]. In loc. cit. that inequality would look like “ $v_{ij} \leq -t_i + x_j$ ”. Yet the linear substitution $t_i \mapsto -t_i$ is natural here, because (4) is a minimization problem; see also [16, Remark 6.28]. The cells of $\text{CovDec}(V)$ are precisely the projections of the faces of the unbounded ordinary polyhedron $\mathcal{E}(V) \subset \mathbb{R}^m \times \mathbb{R}^n$ onto the x -coordinates; see [16, Proposition 6.11]. Let F be the optimal face of the linear program (4). The set $\text{FW}(V)$ is the projection of F .

The face F is gotten by minimizing $t_1 + \dots + t_m$, or equivalently $t_1 + \dots + t_m + x_1 + \dots + x_n$, as the points x are restricted to the hyperplane \mathcal{H} . Let D be the cell in the triangulation of $\Delta_{m-1} \times \Delta_{n-1}$ which is dual to F . Then D contains the vector $(\frac{1}{m}\mathbf{1}, \frac{1}{n}\mathbf{1}) \in \mathbb{R}^m \times \mathbb{R}^n$ in its relative interior. In other words, $\text{FW}(V)$ is dual to the central cell C_V .

The sublevel sets of $d_\Delta(v_i, \cdot)$ are simplices if non-empty; in particular, they are bounded. Consequently, the function $x \mapsto \sum_{i \in [m]} d_\Delta(v_i, x)$ has bounded sublevel sets. The latter implies that $\text{FW}(V)$ is bounded, as it is the set of minimizers of the aforementioned function. The max-tropical convex hull of the rows of V equals the union of the bounded covector cells in $\text{CovDec}(V)$; see [16, Corollary 6.17]. \square

Example 4. For the matrix V from Example 1 the unique optimal solution of the primal linear program (4) reads

$$t^* = (5, 4, 5, 7, 3) \quad \text{and} \quad x^* = (9, -6, -3) ,$$

with optimal value $3 \cdot 24 = 72$. We have $\text{FW}(V) = \{x^*\}$; see Figure 1. As $\gcd(5, 3) = 1$, the uniqueness is implied by Theorem 6 below. The point x^* , which is a pseudovortex of $\text{CovDec}(V)$, is dual to the central cell $C_V = \text{conv}\{113, 122, 212, 221\}$; see Figure 2.

Remark 5. Theorem 3 reveals a similarity to the Euclidean case: by [7, Proposition 19.1] any Fermat–Weber point with respect to the Euclidean distance is contained in the convex hull of the sites. The analogous result to Theorem 3 for the symmetric tropical distance is [20, Proposition 26]. As shown in [20, Example 27], in general, the symmetric tropical Fermat–Weber set is not contained in the tropical convex hull.

Via the Cayley trick, the central cell C_V in $\Sigma(V)$ corresponds to the central covector cell in $\text{CovDec}(V)$, which is $\text{FW}(V)$. The dimension of the latter is the codimension of the former.

Theorem 6. *The dimension of $\text{FW}(V)$ is at most $\gcd(m, n) - 1$. In particular, if m and n are relatively prime, the Fermat–Weber point is unique.*

Proof. We consider the regular subdivision, $\Sigma(V)$, of $\Delta_{m-1} \times \Delta_{n-1}$ induced by the matrix V . Let C_V be the central cell. By [13, §6.2] the vertices of C_V are in bijection with the edges in a subgraph $G(C)$ of the complete bipartite graph $K_{m,n}$ with $c := \text{codim}(C) + 1$ connected components; see also [16, §4.7]. We will show that $c \leq \gcd(m, n)$. Here we may assume that V is generic, whence C_V is a simplex; note that any refinement of $\Sigma(V)$ can only increase the codimension of the cell containing a specific point in its relative interior. If $c = 1$, there is nothing to prove. Hence, we assume that $c \geq 2$.

We consider a maximal simplex, U , of $\Sigma(V)$ which contains C_V , and let T be the subtree of $K_{m,n}$ corresponding to U . A facet, F , of U corresponds to the removal of some edge e of T , yielding disjoint unions $[m] = I' \sqcup I''$ and $[n] = J' \sqcup J''$ such that the

connected components of $T \setminus e$ are the restrictions on $I' \sqcup J'$ and $I'' \sqcup J''$ and e is incident to a vertex in I'' and one in J'' . The hyperplane $\sum_{i' \in I'} x_{i'} + \sum_{j'' \in J''} y_{j''} = 1$ contains the facet F . If I' were empty, then $\sum_{j' \in J'} y_{j'} = 0$ for every $(x, y) \in C$. But C_V contains the point $(\frac{1}{m}\mathbb{1}, \frac{1}{n}\mathbb{1})$, so J' must be empty as well. This contradicts the fact that J' contains a vertex incident to e . Hence, $I' \neq \emptyset$ and, similarly, $J'' \neq \emptyset$.

Consequently, there exist two proper partitions $[m] = I_1 \sqcup I_2 \sqcup \cdots \sqcup I_c$ and $[n] = J_1 \sqcup J_2 \sqcup \cdots \sqcup J_c$ such that the restriction of $G(C)$ on $I_i \sqcup J_i$ is a tree, and there exist edges e_2, \dots, e_c in $K_{m,n}$ between a vertex of I_i and J_{i-1} such that adding these edges we obtain the tree T .

A supporting hyperplane for the facet corresponding to $T \setminus \{e_i\}$ is given by the equation

$$\sum_{i \in I_1 \sqcup \cdots \sqcup I_{i-1}} x_i + \sum_{j \in J_i \sqcup \cdots \sqcup J_c} y_j = 1 ,$$

where $2 \leq i \leq c$. Now the point $(\frac{1}{m}\mathbb{1}, \frac{1}{n}\mathbb{1})$ is contained in these hyperplanes, yielding the $c - 1$ equalities

$$\frac{1}{m} \sum_{k < i} |I_k| + \frac{1}{n} \sum_{\ell \geq i} |J_\ell| = 1, \quad \text{for } 2 \leq i \leq c .$$

Multiplying with the least common multiple of m and n we obtain

$$\frac{n}{\gcd(m,n)} \sum_{k < i} |I_k| + \frac{m}{\gcd(m,n)} \sum_{\ell \geq i} |J_\ell| = \text{lcm}(m, n), \quad \text{for } 2 \leq i \leq c .$$

Putting those relations together with $\sum_{k \in [c]} |I_k| = m$ and $\sum_{\ell \in [c]} |J_\ell| = n$, we yield that $|I_k|$ is a multiple of $m/\gcd(m, n)$ for every $k \in [c]$ and $|J_\ell|$ is a multiple of $n/\gcd(m, n)$ for all $\ell \in [c]$. Further $I_k \neq \emptyset$ for every $k \in [c]$, and so $|I_k| \geq m/\gcd(m, n)$ for all $k \in [c]$. Hence

$$m = \sum_{k \in [c]} |I_k| \geq c \cdot \frac{m}{\gcd(m,n)} ,$$

which implies $c \leq \gcd(m, n)$ as claimed. \square

Restricting to the one-dimensional case (i.e., $n = 2$), we recover the known fact that an odd number of points have a unique median, while the median can be selected from an interval for an even number of points. The following example shows that the inequality in Theorem 6 is tight for all m and n ; see also Example 1.

Example 7. Our example employs the matrix $V = (v_{ij}) \in \mathbb{R}^{m \times n}$ with $v_{ij} = (i - 1)(j - 1)$. The rows are points on the tropical moment curve, and their (max-)tropical convex hull is a tropical cyclic polytope; see [6, §4] and [16, Example 5.18]. The dual triangulation $\Sigma(V)$ of $\Delta_{m-1} \times \Delta_{n-1}$ is the *staircase triangulation*; see [13, §6.2.3]. We explain the construction.

In this case, we represent the simplices as points in an $m \times n$ grid instead of subgraphs of $K_{m,n}$. The staircase triangulation consists of all the paths in an $m \times n$ grid starting at $(1, 1)$ and ending at (m, n) obtained by going right or down. Figure 3 portrays such a path. Due to its shape, it is called a *staircase*. We will show that in the staircase triangulation the simplex containing $(\frac{1}{m}\mathbb{1}, \frac{1}{n}\mathbb{1})$ lies in a cell of codimension $\gcd(m, n) - 1$. For improved readability, we abbreviate $d = \gcd(m, n)$, $a = m/d$, and $b = n/d$.

If $d = 1$, then there is a unique maximal simplex in the staircase triangulation containing $(\frac{1}{m}\mathbb{1}, \frac{1}{n}\mathbb{1})$ in its interior. To find the precise staircase, we use the Northwest Corner Rule [11, §8.3.1] in the standard transportation array with marginal column $\frac{1}{m}\mathbb{1}$ and marginal row $\frac{1}{n}\mathbb{1}$. The visited cells form the staircase, which we call the *central staircase*. Note that this method provides also barycentric coordinates for $(\frac{1}{m}\mathbb{1}, \frac{1}{n}\mathbb{1})$ in this simplex.




FIGURE 3. Staircase in a 6×9 grid

When $d \geq 2$, consider the partitions in d subsets $[m] = I_1 \sqcup \dots \sqcup I_d$ and $[n] = J_1 \sqcup \dots \sqcup J_d$ where $I_i = \{(i-1)a+1, (i-1)a+2, \dots, ia\}$ and $J_i = \{(i-1)b+1, (i-1)b+2, \dots, ib\}$. Consider on $I_i \times J_i$ the central staircase on an $a \times b$ grid and add the points $(ia, ib+1)$ for $i = 1, \dots, d-1$: in Figure 3 the blocks on $I_i \times J_i$ correspond to the gray areas whereas the added points are those in the white area. This staircase corresponds to a maximal simplex U in the staircase triangulation, which contains $(\frac{1}{m}\mathbb{1}, \frac{1}{n}\mathbb{1})$.

The removal of the grid point $(ia, ib+1)$ yields the facet defined by

$$\sum_{k \leq ia} x_k + \sum_{\ell > ib} y_\ell = 1 .$$

Using $d = m/a = n/b$, we see that $(\frac{1}{m}\mathbb{1}, \frac{1}{n}\mathbb{1})$ is contained in this facet. In total, there are $d-1$ facets of this form.

Each remaining facet of U corresponds to the removal of a grid point from some block $I_k \times J_k$. In particular, each facet induces a partition $[m] = I' \sqcup I''$ such that $I_1 \sqcup \dots \sqcup I_{k-1} \subset I'$, $I_{k+1} \sqcup \dots \sqcup I_d \subset I''$ for some $k \in [d]$; similarly, there is a partition $J' \sqcup J''$ on $[n]$. Moreover, at least one of the intersections $I_k \cap I' \cap I''$ and $J_k \cap J' \cap J''$ is nonempty, which implies that at least one of $|I'|$ and $|J''|$ is not a multiple of d . As in the proof of Theorem 6, the facet defining equation

$$\sum_{i' \in I'} x_{i'} + \sum_{j'' \in J''} y_{j''} = 1$$

is not satisfied by $(\frac{1}{m}\mathbb{1}, \frac{1}{n}\mathbb{1})$. Since $(\frac{1}{m}\mathbb{1}, \frac{1}{n}\mathbb{1})$ is contained in precisely $d-1$ facets of U , the dimension of $\text{FW}(V)$ equals $d-1$.

The staircase for U is obtained by using the Northwest Corner Rule with breaking ties by going East. If we break the ties randomly, then 2^{d-1} staircases appear with nonzero probability. These staircases are in bijection with the ordinary vertices of $\text{FW}(V)$, which is a $(d-1)$ -dimensional cube, seen as an ordinary polytope.

Remark 8. In the special case $m = n$ computing tropical Fermat–Weber points reduces to the tropical Cramer rule [16, §4.9]. This also agrees with results of Gärtner and Jaggi [15, §4.1] in the context of “tropical support vector machines” on computing a “separating hyperplane for n points”.

Remark 9. As a consequence of [16, Theorem 6.14] the Fermat–Weber set $\text{FW}(V^\top)$ in $\mathbb{R}^m/\mathbb{R}\mathbb{1}$ of the n columns of V is affinely isomorphic to the Fermat–Weber set $\text{FW}(V)$ in $\mathbb{R}^n/\mathbb{R}\mathbb{1}$ of the m rows.

Remark 10. Our analysis rests on the decision, in (3), to look at the distances from the sites to the Fermat–Weber points. This leads to min-tropical hyperplane arrangements and max-tropical convex hulls. Reversing the direction, i.e., considering distances from the Fermat–Weber points to the sites, amounts to exchanging min and max throughout. Conceptually, the results remain the same; cf. [16, §1.3].

4. TRANSPORTATION

This section comprises the algorithmic core of this paper. The basic ingredient is the transportation problem which already occurred in Example 7.

Again let us fix a matrix $V = (v_{ij}) \in \mathbb{R}^{m \times n}$. Whenever it will suit us, we may also view the rows of V as a m labeled points v_1, \dots, v_m in \mathcal{H} or $\mathbb{R}^n/\mathbb{R}\mathbf{1}$. The following is the dual of the linear program (4) with variables λ and y_{ij} for $i \in [m]$ and $j \in [n]$:

$$(5) \quad \begin{aligned} & \text{maximize} && \sum_{i \in [m]} \sum_{j \in [n]} v_{ij} \cdot y_{ij} \\ & \text{subject to} && \sum_{j \in [n]} y_{ij} = n, && \text{for } i \in [m] \\ & && \lambda + \sum_{i \in [m]} y_{ij} = 0, && \text{for } j \in [n] \\ & && y_{ij} \geq 0, && \text{for } i \in [m] \text{ and } j \in [n]. \end{aligned}$$

From $0 = n \cdot \lambda + \sum_{i,j} y_{ij} = n \cdot (\lambda + m)$, we get $\lambda = -m$ for every feasible point. By substituting that value in (5) we obtain the standard linear programming formulation of a transportation problem; see [27, §21.6]. The primal linear program (4) and the envelope $\mathcal{E}(V)$ are related to transportation via Hitchcock’s theorem [27, Theorem 21.13].

As the right hand sides are the integral constants m and n , the feasible region of (5) is a *central transportation polytope*, and we denote it $T(m, n)$. The polytope $T(m, n)$ is known to be integral [12, Lemma 2.13]. The nonzero entries of any vertex, which is an $m \times n$ -matrix, defines a subgraph of $K_{m,n}$ by picking edges [12, Lemma 2.9]. This is the *support graph* of that vertex, and this is a forest; see [27, Theorem 21.15].

Example 11. For the 5×3 -matrix V from Example 1 the transpose of the unique optimal solution of (5) is

$$(6) \quad (y_{ij}^*)^\top = \begin{pmatrix} 3 & 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 3 & 2 \\ 0 & 1 & 3 & 0 & 1 \end{pmatrix}.$$

The support graph is a spanning tree; see Figure 4. By Theorem 14 below that tree encodes the covector of the unique Fermat–Weber point x^* . The degree sequence for the column nodes is (223). This is the coarse type of x^* and also the componentwise maximum of the four vertices of the central cell in the mixed subdivision; cf. Example 4 and [16, §4.5].




FIGURE 4. Spanning subtree of $K_{5,3}$ encoding the covector of the unique Fermat–Weber point from Example 1. Column nodes at the bottom

Assuming $m \geq n$, Tokuyama and Nakano gave an algorithm (which they called “splitter finding”) to solve a transportation problem like (5) in $O(n^2 m \log^2 m)$ time [30, Theorem 3.1]. We borrow some of their ideas. For $J \subseteq [n]$ and $u \in \mathcal{H}$ consider the set

$$S_J(u) := \left\{ x \in \mathcal{H} \mid \max_{j \in J} (x_j - u_j) \geq \max_{i \notin J} (x_i - u_i) \right\} .$$

We have $S_\emptyset(u) = \emptyset$, and $S_{[n]}(u) = \mathcal{H}$, where we use the convention that the maximum of the empty set is $-\infty$, the neutral element of the tropical addition. If J is a nonempty proper subset of $[n]$, then $S_J(u)$ is a max-tropical halfspace with apex u ; see [16, §7.1]. In the special case where $J = \{j\}$ is a singleton that tropical halfspace is a closed sector; in general, $S_J(u) = \bigcup_{j \in J} S_{\{j\}}(u)$.

Definition 12. We say that $u \in \mathcal{H}$ *evenly splits* V if for every subset J of $[n]$ we have $n \cdot |V \cap S_J(u)| \geq m \cdot |J|$.

Tokuyama and Nakano [30] call the point u a “1-splitter”, and the sectors are the “regions split by u ”. Theorem 14 below may be seen as our interpretation of their results in the setting of tropical convexity.

Example 13. Consider the points V from Example 1. The Fermat–Weber point $u = (9, -6, -3)$ evenly splits them. This is illustrated in Figure 5, where we have drawn the max-tropical hyperplane based at u with dotted lines. In particular, we see the subdivision of $\mathbb{R}^3/\mathbb{R}\mathbf{1}$ in three convex regions.

Theorem 14. *A point $u \in \mathcal{H}$ evenly splits V if and only if $u \in \text{FW}(V)$. The support graph of an optimal dual solution y^* is the covector of the Fermat–Weber point $u(y^*)$.*

Proof. Assume that u evenly splits V and consider $t_i = \max_{j \in [n]} (v_{ij} - u_j)$ for $i \in [m]$. Thus (u, t) is a feasible solution of the primal linear program (4). Moreover, $t_i + u_j = v_{ij}$ if and only if $v_i \in S_{\{j\}}(u)$.

Now [30, Theorem 2.2] says that there exists a solution (y_{ij}) of the transportation problem (5), which is dual to (4), such that (u, t) and (y_{ij}) satisfy the complementary slackness condition. Indeed, if $t_i + u_j \neq v_{ij}$, then $v_i \notin S_{\{j\}}(u)$, so the aforementioned result gives $y_{ij} = 0$. So it follows from linear programming duality that (u, t) is an optimal solution of (4). In particular, u is a Fermat–Weber point for V .

For the converse, we denote by ϕ the convex function $\frac{1}{n} \sum_{s \in V} d_\Delta(s, \cdot)$. Also, for every subset J of $[n]$, denote by σ_J the cardinality of $V \cap S_J(u)$. Abbreviating $f_J := (n - |J|) \sum_{j \in J} e_j - |J| \sum_{i \in [n] \setminus J} e_i$, which is a point in \mathcal{H} , we obtain:

- if $s \in S_J(u)$, then $s \in S_J(u - \delta f_J)$ for any $\delta \geq 0$;
- if $s \notin S_J(u)$, then $s \notin S_J(u - \delta f_J)$ for any $\delta \geq 0$ sufficiently small.

The condition $s \in S_J(u)$ implies $d_\Delta(s, u) = n(s_j - u_j)$ for some $j \in J$. Therefore, for $\delta > 0$ sufficiently small, we obtain

- $d_\Delta(s, u - \delta f_J) = d_\Delta(s, u) + n \cdot \delta(n - |J|)$, when $s \in S_J(u)$;
- $d_\Delta(s, u - \delta f_J) = d_\Delta(s, u) - n \cdot \delta |J|$, when $s \notin S_J(u)$.

Summing up and dividing by n yields

$$(7) \quad \phi(u - \delta f_J) = \phi(u) + \delta(n - |J|)\sigma_J - \delta|J|(m - \sigma_J) = \phi(u) + \delta(n\sigma_J - m|J|)$$

for any $\delta > 0$ sufficiently small. If $u \in \text{FW}(V)$, then u is a minimizer, so $\phi(u - \delta f_J) \geq \phi(u)$ for any $\delta > 0$ and $J \subseteq [n]$. Equation (7) implies $n\sigma_J \geq m|J|$ for every $J \subseteq [n]$, under this assumption. Consequently, u evenly splits V . \square




FIGURE 5. Five sites in $\mathbb{R}^3/\mathbb{R}\mathbb{1}$ and the unique point that evenly splits them. Each closed sector of the max-tropical halfspace with apex $(9, -6, -3)$ contains at least two sites

The argument in the proof of Theorem 14 leads to the following algorithm for obtaining a tropical Fermat–Weber point from an optimal solution y^* of (5). By complementary slackness each edge (i, j) of the support graph, $B(y^*)$, imposes the equality

$$(8) \quad t_i + x_j = v_{ij}$$

for any optimal dual solution. If we assume $x_j = 0$ for some column node j , then we can perform a depth-first search from j and recover all the other values of (t, x) using the equalities in (8). There may be more connected components, in which case we start a depth-first search from every unvisited column node. In this way all the values are recovered, as $B(y^*)$ is spanning—no row or column of y^* can be zero as its elements must sum up to a positive number. Note that the point x obtained this way may not lie in \mathcal{H} . Yet, by adding $(x_1 + \dots + x_n)/n$ to every entry of x and subtracting the same value from every entry of t , we obtain a feasible solution (t^*, x^*) of (4) that satisfies the equations (8). In particular, that solution is optimal, whence $x^* \in \text{FW}(V)$. So the method of [30] gives the following complexity result.

Corollary 15. *Assuming $m \geq n$, one tropical Fermat–Weber point can be computed in $O(n^2 m \log^2 m)$ time.*

The complexity bounds of the algorithms by Kleinschmidt and Schannath [18] and Brenner [8] are similar, but slightly different. Naturally, they carry over as well.

It is also natural to ask for explicit representations of the entire Fermat–Weber set. A first idea could be to list the vertices of $\text{FW}(V)$, seen as an ordinary polytope. Example 7 shows that cubes occur as Fermat–Weber sets, and their number of vertices is exponential in the dimension. Yet, the polytropical structure allows for more efficient choices. Namely, a polytope in $\mathbb{R}^n/\mathbb{R}\mathbb{1}$ has at most n tropical vertices and at most $n^2 - n$ ordinary facets.

Let us start with the latter. Since we know the possible directions of the (outer) facet normal vectors, $e_k - e_\ell$ for $k \neq \ell$, we can find the ordinary facets by solving $n^2 - n$ linear

programs like:

$$(9) \quad \begin{array}{ll} \text{maximize} & x_k - x_\ell \\ \text{subject to} & v_{ij} - x_j \leq t_i, \quad \text{for } i \in [m] \text{ and } j \in [n] \\ & x_1 + \cdots + x_n = 0 \\ & t_1 + \cdots + t_m = p^*/n \end{array}$$

where p^* is the optimal value of (4). The constraint matrix has only 0 and ± 1 entries, and so a linear program of the form (9) can be solved in strongly polynomial time [28, Corollary 15.3a].

Each such linear program yields one tight inequality $x_k - x_\ell \leq a_{k,\ell}$ of $\text{FW}(V)$, where $a_{k,\ell}$ is the optimal value. Then, the tropical vertices can be found as $A_k = (-a_{k,1}, \dots, -a_{k,n}) \in \mathbb{R}^n / \mathbb{R}\mathbf{1}$, where $a_{k,k} = 0$. From Corollary 15 we thus infer the following result.

Corollary 16. *The ordinary facet description and the tropical vertices of $\text{FW}(V)$ can be found in strongly polynomial time.*

If the tropical vertices are known, then we can check if a given point lies in $\text{FW}(V)$ in $O(n^2)$ time by checking the criterion [16, Proposition 5.37].

5. TROPICAL MEDIAN CONSENSUS TREES

Phylogenetics is a branch of (computational) biology that seeks to associate trees to mark ancestral relations among given *taxa*, e.g., species [29]. The taxa correspond to the leaves. Here we show how asymmetric tropical Fermat–Weber sets give rise to a new algorithm for finding consensus trees. That method is efficient, and it exhibits other nice features.

A *dissimilarity map* is a symmetric $n \times n$ -matrix $D = (d_{ij})$ with zero diagonal. It is called an *ultrametric* if D is nonnegative, and the *ultrametric inequality*

$$(10) \quad d_{ik} \leq \max(d_{ij}, d_{jk})$$

holds for all $i, j, k \in [n]$. Since the diagonal is implicit, and the matrix is required to be symmetric, we may view a dissimilarity map as an element of $\mathbb{R}^{\binom{n}{2}}$.

A rooted metric tree with n labeled leaves is *equidistant* if the distance from any leaf to the root is the same. It is known that the ultrametries are precisely the distance functions among the leaves in an equidistant tree; see [29, Theorem 7.2.5]. Note that the all-ones vector $\mathbf{1}$ of length $\binom{n}{2}$ is an ultrametric. The corresponding equidistant tree has interior edges of length zero, while each leaf edge has length $\frac{1}{2}$.

Ardila and Klivans [3, Theorem 3] showed that a dissimilarity map is an ultrametric if and only if it corresponds to a point on the Bergman fan $\tilde{B}(K_n)$ of the complete graph K_n . The Bergman fan of a matroid is a special case of a tropical linear space, i.e., with constant coefficients. The ultrametric inequality (10) is a tropical convexity condition with respect to \max . From a dissimilarity map D and any real constant c we can construct the dissimilarity map $D + c\mathbf{1}$. Moreover, if D is an ultrametric, and $D + c\mathbf{1}$ is nonnegative, then it is an ultrametric, too. In this way, we may view $\tilde{B}(K_n)$ as a max-tropical linear space in the tropical projective torus $\mathbb{R}^{\binom{n}{2}} / \mathbb{R}\mathbf{1}$; see [20, Proposition 16] and [31, Theorem 3]. We abbreviate $\mathcal{T}_n := \tilde{B}(K_n) / \mathbb{R}\mathbf{1}$ and call it the *space of equidistant trees* on n labeled leaves. The following result says that Fermat–Weber points of equidistant trees are again equidistant trees.

Theorem 17. *Let $V \subset \mathcal{T}_n$ be a finite set of equidistant trees on n leaves. Then the tropical polytope $\text{FW}(V)$ is contained in \mathcal{T}_n . Moreover, any two trees in $\text{FW}(V)$ share the same tree topology.*




FIGURE 6. Three trees (a),(b),(c) and their tropical median consensus tree (d)

Proof. According to Theorem 3 the set $\text{FW}(V)$ is contained in the max-tropical convex hull of V . The space of equidistant trees \mathcal{T}_n is a max-tropical linear space and thus max-tropically convex; see [16, Proposition 10.33]. This is the first claim.

Page, Yoshida and Zhang showed in [25, Theorem 3.2] that the trees in any relatively open cell of the covector decomposition of V in \mathcal{T}_n share the same tree topology. Now the second claim follows from Theorem 3. \square

The previous result is in contrast with the situation of symmetric Fermat–Weber points, which may lie outside the max-tropical convex hull; see [20, Example 27].

Corollary 18. *Let $V \subset \mathcal{T}_n$ be set of m equidistant trees on n leaves. Then*

$$\dim \text{FW}(V) \leq \min(n - 1, \gcd(m, \binom{n}{2})) - 1 .$$

Proof. This follows from Theorem 6 and the fact that the graphic matroid of K_n has rank $n - 1$, so $\dim \mathcal{T}_n = n - 2$. \square

Given multiple trees, their Fermat–Weber set may have positive dimension. To devise a consensus method, we need to consistently select one point from the relative interior of this set. We exploit the fact that the Fermat–Weber set is a polytrope, whose tropical vertices

can be found in strongly polynomial time, by Corollary 16. Our consensus tree will be the ordinary average of the tropical vertices and we call it the *tropical median consensus tree*. Taking the average tropically, we would arrive at the “tropical barycenter” defined in [1, §3.2]; that always lies on the boundary and is thus undesirable in this context.

Example 19. The first three equidistant trees on $n = 9$ taxa in Figure 6 are taken from [17, Chapter 7]. Since, in loc. cit., there are no edge lengths, here we choose heights that are compatible with the graphical representation in [17, Fig. 7.1]. The tropical median consensus tree is depicted in Figure 6 as (d). This is the unique Fermat–Weber point of the three input trees in \mathcal{T}_9 .

Let $i, j, k \in [n]$ be pairwise distinct taxa in an equidistant tree such that the lowest common ancestor of i and j is a proper descendant of the lowest common ancestor of i, j , and k . Then we say that these taxa form a *rooted triplet*, and we denote it by $ij|k$. If $D = (d_{ij})$ is its ultrametric distance, then $ij|k$ is a rooted triplet if and only if $d_{ij} < d_{ik}$. Note that the ultrametric property implies $d_{jk} = d_{ik}$, so we also have $d_{ij} < d_{jk}$. We denote by $r(D)$ the set of rooted triplets of the tree. A consensus method is called

- *Pareto on rooted triplets* if $\bigcap_{\ell \in [m]} r(D_\ell) \subseteq r(D)$;
- *co-Pareto on rooted triplets* if $r(D) \subseteq \bigcup_{\ell \in [m]} r(D_\ell)$;

where D_1, \dots, D_m corresponds to the input trees and D represents the consensus tree. These are desirable properties for consensus methods; see [9, §3]. The set of equidistant trees containing the rooted triplet $ij|k$ is

$$\mathcal{T}_n(ij|k) := \{D \in \mathcal{T}_n \mid d_{ij} < d_{ik}\} .$$

This set is tropically convex as it arises as the intersection of \mathcal{T}_n with an open tropical halfspace. This argument also shows that its complement in \mathcal{T}_n is tropically convex.

Proposition 20. *The tropical median method is Pareto and co-Pareto on rooted triplets.*

Proof. Let D_1, \dots, D_m be the input ultrametrics and D any Fermat–Weber point.

If $ij|k$ belongs to $\bigcap_{\ell} r(D_\ell)$, then $D_\ell \in \mathcal{T}_n(ij|k)$ for every $\ell \in [m]$. As $\mathcal{T}_n(ij|k)$ is tropically convex, Theorem 3 implies that D also belongs to this set. Thus, $ij|k$ also belongs to $r(D)$, showing that our method is Pareto on rooted triplets.

If $ij|k$ does not belong to $\bigcup_{\ell \in [m]} r(D_\ell)$, then the input ultrametrics are all contained in the complement of $\mathcal{T}_n(ij|k)$. Again, Theorem 3 yields that $D \notin \mathcal{T}_n(ij|k)$, because $\mathcal{T}_n \setminus \mathcal{T}_n(ij|k)$ is tropically convex. This is equivalent to $ij|k \notin r(D)$. We conclude that our method is co-Pareto on rooted triplets. \square

A survey of consensus methods for trees can be found in [9]. Most methods deal with unweighted phylogenetic trees, so they are discrete analogues of our approach. Unweighted *median consensus trees* were defined in [4], and the *asymmetric* case is analyzed in [26]. Bryant [9] presents only one consensus method that uses the edge lengths in the computation. The result is the *average consensus tree* [19]; one of its drawbacks being that there is no efficient algorithm known for its construction. The difficulty arises from the necessity to add the ultrametric inequality as an additional constraint. The same issue occurs for consensus trees via the symmetric tropical distance [21]. However, Page, Yoshida and Zhang showed that a consensus tree can be selected from a symmetric tropical Fermat–Weber set of ultrametrics [25, Lemma 3.5]. The advantage of our method is that the ultrametric property is automatic, even if it is not required explicitly in the problem formulation.

REFERENCES

- [1] Xavier Allamigeon, Pascal Benchimol, Stéphane Gaubert, and Michael Joswig. “Log-barrier interior point methods are not strongly polynomial”, *SIAM J. Appl. Algebra Geom.* **2.1** (2018), pp. 140–178. DOI: [10.1137/17M1142132](https://doi.org/10.1137/17M1142132).
- [2] Omid Amini and Madhusudan Manjunath. “Riemann-Roch for sub-lattices of the root lattice A_n ”, *Electron. J. Combin.* **17.1** (2010), Research Paper 124, 50.
- [3] Federico Ardila and Caroline J. Klivans. “The Bergman complex of a matroid and phylogenetic trees”, *J. Combin. Theory Ser. B* **96.1** (2006), pp. 38–49. DOI: [10.1016/j.jctb.2005.06.004](https://doi.org/10.1016/j.jctb.2005.06.004).
- [4] Jean-Pierre Barthélemy and Frederick R. McMorris. “The median procedure for n-trees”. English, *J. Classif.* **3** (1986), pp. 329–334. DOI: [10.1007/BF01894194](https://doi.org/10.1007/BF01894194).
- [5] Louis J. Billera, Susan P. Holmes, and Karen Vogtmann. “Geometry of the space of phylogenetic trees”, *Adv. in Appl. Math.* **27.4** (2001), pp. 733–767. DOI: [10.1006/aama.2001.0759](https://doi.org/10.1006/aama.2001.0759).
- [6] Florian Block and Josephine Yu. “Tropical convexity via cellular resolutions”, *J. Algebraic Combin.* **24.1** (2006), pp. 103–114. DOI: [10.1007/s10801-006-9104-9](https://doi.org/10.1007/s10801-006-9104-9).
- [7] Vladimir Boltyanski, Horst Martini, and Valeriu Soltan. *Geometric methods and optimization problems*. Vol. 4. Combinatorial Optimization. Kluwer Academic Publishers, Dordrecht, 1999. DOI: [10.1007/978-1-4615-5319-9](https://doi.org/10.1007/978-1-4615-5319-9).
- [8] Ulrich Brenner. “A faster polynomial algorithm for the unbalanced Hitchcock transportation problem”, *Oper. Res. Lett.* **36.4** (2008), pp. 408–413. DOI: [10.1016/j.orl.2008.01.011](https://doi.org/10.1016/j.orl.2008.01.011).
- [9] David Bryant. “A classification of consensus methods for phylogenetics”, *Bioconsensus (Piscataway, NJ, 2000/2001)*. Vol. 61. DIMACS Ser. Discrete Math. Theoret. Comput. Sci. Amer. Math. Soc., Providence, RI, 2003, pp. 163–183.
- [10] Peter Butkovič. *Max-linear systems: theory and algorithms*. Springer Monographs in Mathematics. Springer-Verlag London, Ltd., London, 2010, pp. xviii+272. DOI: [10.1007/978-1-84996-299-5](https://doi.org/10.1007/978-1-84996-299-5).
- [11] George B. Dantzig and Mukund N. Thapa. *Linear programming. 1*. Springer Series in Operations Research. Springer-Verlag, New York, 1997.
- [12] Jesús A. De Loera and Edward D. Kim. “Combinatorics and geometry of transportation polytopes: an update”, *Discrete geometry and algebraic combinatorics*. Vol. 625. Contemp. Math. Amer. Math. Soc., Providence, RI, 2014, pp. 37–76. DOI: [10.1090/conm/625/12491](https://doi.org/10.1090/conm/625/12491).
- [13] Jesús A. De Loera, Jörg Rambau, and Francisco Santos. *Triangulations*. Vol. 25. Algorithms and Computation in Mathematics. Structures for algorithms and applications. Springer-Verlag, Berlin, 2010. DOI: [10.1007/978-3-642-12971-1](https://doi.org/10.1007/978-3-642-12971-1).
- [14] Mike Develin and Bernd Sturmfels. “Tropical convexity”, *Doc. Math.* **9** (2004). correction: *ibid.*, pp. 205–206, 1–27 (electronic).
- [15] Bernd Gärtner and Martin Jaggi. “Tropical support vector machines”. ACS Technical Report No.: ACS-TR-362502-01. 2006.
- [16] Michael Joswig. *Essentials of tropical combinatorics*. Vol. 219. Graduate Studies in Mathematics. Providence, RI: American Mathematical Society, 2021.
- [17] Ian J. Kitching, Peter L. Forey, Christopher J. Humphries, and David M. Williams. *Cladistics: The Theory and Practice of Parsimony Analysis*. 2nd edition. Oxford University Press, 1998.

- [18] Peter Kleinschmidt and Heinz Schannath. “A strongly polynomial algorithm for the transportation problem”, *Math. Programming* **68.1**, Ser. A (1995), pp. 1–13. DOI: [10.1007/BF01585755](https://doi.org/10.1007/BF01585755).
- [19] François-Joseph Lapointe and Guy Cucumel. “The Average Consensus Procedure: Combination of Weighted Trees Containing Identical or Overlapping Sets of Taxa”, *Systematic Biology* **46.2** (1997), pp. 306–312.
- [20] Bo Lin, Bernd Sturmfels, Xiaoxian Tang, and Ruriko Yoshida. “Convexity in tree spaces”, *SIAM J. Discrete Math.* **31.3** (2017), pp. 2015–2038. DOI: [10.1137/16M1079841](https://doi.org/10.1137/16M1079841).
- [21] Bo Lin and Ruriko Yoshida. “Tropical Fermat-Weber points”, *SIAM J. Discrete Math.* **32.2** (2018), pp. 1229–1245. DOI: [10.1137/16M1071122](https://doi.org/10.1137/16M1071122).
- [22] Diane Maclagan and Bernd Sturmfels. *Introduction to tropical geometry*. Vol. 161. Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2015, pp. xii+363.
- [23] Madhusudan Manjunath. “The Laplacian lattice of a graph under a simplicial distance function”, *European J. Combin.* **34.6** (2013), pp. 1051–1070. DOI: [10.1016/j.ejc.2013.01.010](https://doi.org/10.1016/j.ejc.2013.01.010).
- [24] Lior Pachter and Bernd Sturmfels, eds. *Algebraic statistics for computational biology*. New York: Cambridge University Press, 2005.
- [25] Robert Page, Ruriko Yoshida, and Leon Zhang. “Tropical principal component analysis on the space of phylogenetic trees”, *Bioinformatics* **36.17** (June 2020), pp. 4590–4598. DOI: [10.1093/bioinformatics/btaa564](https://doi.org/10.1093/bioinformatics/btaa564).
- [26] Cynthia Phillips and Tandy J. Warnow. “The asymmetric median tree—a new model for building consensus trees”, *Discrete Appl. Math.* **71.1-3** (1996), pp. 311–335. DOI: [10.1016/S0166-218X\(96\)00071-6](https://doi.org/10.1016/S0166-218X(96)00071-6).
- [27] Alexander Schrijver. *Combinatorial optimization. Polyhedra and efficiency. Vol. A*. Vol. 24. Algorithms and Combinatorics. Paths, flows, matchings, Chapters 1–38. Berlin: Springer-Verlag, 2003.
- [28] Alexander Schrijver. *Theory of linear and integer programming*. Wiley-Interscience Series in Discrete Mathematics. A Wiley-Interscience Publication. John Wiley & Sons, Ltd., Chichester, 1986.
- [29] Charles Semple and Mike Steel. *Phylogenetics*. Vol. 24. Oxford Lecture Series in Mathematics and its Applications. Oxford University Press, Oxford, 2003.
- [30] Takeshi Tokuyama and Jun Nakano. “Efficient algorithms for the Hitchcock transportation problem”, *SIAM J. Comput.* **24.3** (1995), pp. 563–578. DOI: [10.1137/S0097539792236729](https://doi.org/10.1137/S0097539792236729).
- [31] Ruriko Yoshida, Leon Zhang, and Xu Zhang. “Tropical principal component analysis and its application to phylogenetics”, *Bull. Math. Biol.* **81.2** (2019), pp. 568–597. DOI: [10.1007/s11538-018-0493-4](https://doi.org/10.1007/s11538-018-0493-4).

(Andrei Comăneci) TECHNISCHE UNIVERSITÄT BERLIN, CHAIR OF DISCRETE MATHEMATICS/GEOMETRY,
COMANECI@MATH.TU-BERLIN.DE

(Michael Joswig) TECHNISCHE UNIVERSITÄT BERLIN, CHAIR OF DISCRETE MATHEMATICS/GEOMETRY,
MAX-PLANCK INSTITUTE FOR MATHEMATICS IN THE SCIENCES, LEIPZIG, JOSWIG@MATH.TU-BERLIN.DE