

EXAMPLES OF COMPACT EMBEDDED λ -HYPERSURFACES

QING-MING CHENG, JUNQI LAI AND GUOXIN WEI

ABSTRACT. In the paper, we construct compact embedded λ -hypersurfaces which are diffeomorphic to a sphere and are not isometric to a standard sphere. Hence, one can not expect to have Alexandrov type theorem for λ -hypersurfaces.

1. INTRODUCTION

A hypersurface $\Sigma^n \subset \mathbb{R}^{n+1}$ is called a λ -hypersurface if it satisfies

$$(1.1) \quad H + \langle X, \nu \rangle = \lambda,$$

where λ is a constant, X is the position vector, ν is a unit normal vector and H is the mean curvature. The notation of λ -hypersurfaces were first introduced by Cheng and Wei in [4] (also see [11]). Cheng and Wei [4] proved that λ -hypersurfaces are critical points of the weighted area functional with respect to weighted volume-preserving variations. λ -hypersurfaces can also be viewed as stationary solutions to the isoperimetric problem in the Gaussian space. For more informations on λ -hypersurfaces, one can see [4] and [11]. It is well known that there are several special complete embedded solutions to (1.1):

- hyperplanes with a distance of $|\lambda|$ from the origin,
- sphere with radius $\frac{-\lambda + \sqrt{\lambda^2 + 4n}}{2}$ centered at origin,
- cylinders with an axis through the origin and radius $\frac{-\lambda + \sqrt{\lambda^2 + 4(n-1)}}{2}$.

Remark 1.1. *If $\lambda = 0$, $\langle X, \nu \rangle + H = \lambda = 0$, then $X : \Sigma^n \rightarrow \mathbb{R}^{n+1}$ is a self-shrinker. Hence, one knows that λ -hypersurfaces is a natural generalization of self-shrinkers of mean curvature flow, which plays an important role for study on singularities of the mean curvature flow.*

In [5], Cheng and Wei constructed the first nontrivial example of a λ -hypersurface which is diffeomorphic to $\mathbb{S}^{n-1} \times \mathbb{S}^1$ using techniques similar to Angenent [1]. In [12], using a similar method to McGrath [10], Ross constructed a λ -hypersurface in \mathbb{R}^{2n+2} which is diffeomorphic to $\mathbb{S}^n \times \mathbb{S}^n \times \mathbb{S}^1$ and exhibits a $SO(n) \times SO(n)$ rotational symmetry. In [9], Li and Wei constructed an immersed S^n λ -hypersurface using a similar method to [7]. It is quite interesting to find other nontrivial examples of λ -hypersurfaces, in this paper, motivated by [7, 8, 9], we construct a nontrivial embedded λ -hypersurface which is diffeomorphic to \mathbb{S}^n and they are not isometric to a standard sphere. In fact, we obtain the following theorem:

Theorem 1.1. *For $n \geq 2$ and $-\frac{2}{\sqrt{n+2}} < \lambda < 0$, there exists an embedded convex λ -hypersurface $\Sigma^n \subset \mathbb{R}^{n+1}$ which is diffeomorphic to \mathbb{S}^n and is not isometric to a standard sphere.*

The first author was partially supported by JSPS Grant-in-Aid for Scientific Research (B): No.16H03937 and No.22K03303. The third author was partly supported by NSFC Grant No.12171164, GDUPS (2018).

Remark 1.2. *It is well-known that for compact embedded hypersurfaces with constant mean curvature in \mathbb{R}^{n+1} , Alexandrov theorem holds, that is, a compact embedded hypersurface with constant mean curvature in \mathbb{R}^{n+1} is isometric to a round sphere. But for λ -hypersurfaces, one can not expect to have Alexandrov type theorem for λ -hypersurfaces according to the above theorem 1.1.*

Remark 1.3. *For self-shrinkers, there is a well-known conjecture asserts that the round sphere should be the only embedded self-shrinker which is diffeomorphic to a sphere. Brendle [2] proved the above conjecture for 2-dimension self-shrinker. For the higher dimensional self-shrinker, the conjecture is still open. But for λ -hypersurfaces, we can construct compact embedded λ -hypersurface which is diffeomorphic to a sphere and is not isometric to a round sphere.*

Remark 1.4. *For $-\frac{2}{\sqrt{3}} < \lambda < 0$, Chang [3] proved that there exists a compact embedded λ -curve with 2-symmetry in \mathbb{R}^2 , which is not a circle.*

2. PRELIMINARIES

Let $SO(n)$ denote the special orthogonal group and act on $\mathbb{R}^{n+1} = \{(x, y) : x \in \mathbb{R}, y \in \mathbb{R}^n\}$ in the usual way, then we can identify the space of orbits $\mathbb{R}^{n+1}/SO(n)$ with the half plane $\mathbb{H} = \{(x, r) \in \mathbb{R}^2 : x \in \mathbb{R}, r \geq 0\}$ under the projection(see [12])

$$\Pi(x, y) = (x, |y|) = (x, r).$$

If a hypersurface Σ is invariant under the action $SO(n)$, then the projection $\Pi(\Sigma)$ will give us a profile curve in the half plane, which can be parametrized by Euclidean arc length and write as $\gamma(s) = (x(s), r(s))$. Conversely, if we have a curve $\gamma(s) = (x(s), r(s))$, $s \in (a, b)$ parametrized by Euclidean arc length in the half plane, then we can reconstruct the hypersurface by

$$(2.1) \quad \begin{aligned} X : (a, b) \times S^{n-1}(1) &\hookrightarrow \mathbb{R}^{n+1}, \\ (s, \alpha) &\mapsto (x(s), r(s)\alpha). \end{aligned}$$

Let

$$(2.2) \quad \nu = (-\dot{r}, \dot{x}\alpha),$$

where the dot denotes taking derivative with respect to arc length s . A direct calculation shows that ν is a unit normal vector for the hypersurface. Then we can calculate that the principal curvatures of hypersurface(see [5, 6]):

$$(2.3) \quad \begin{aligned} \lambda_i &= -\frac{\dot{x}}{r}, \quad i = 1, 2, \dots, n-1 \\ \lambda_n &= \dot{x}\ddot{r} - \ddot{x}\dot{r} \end{aligned}$$

Hence the mean curvature equals to

$$(2.4) \quad H = \dot{x}\ddot{r} - \ddot{x}\dot{r} - (n-1)\frac{\dot{x}}{r},$$

and then, by (2.1), (2.2) and (2.4), equation (1.1) reduces to(see also [5, 8, 12])

$$(2.5) \quad \dot{x}\ddot{r} - \ddot{x}\dot{r} = \left(\frac{n-1}{r} - r\right)\dot{x} + x\dot{r} + \lambda,$$

where $(\dot{x})^2 + (\dot{r})^2 = 1$. Let $\theta(s)$ denote the angles between the tangent vectors of the profile curve and x -axis, then (2.5) can be written as the following system of differential equation:

$$(2.6) \quad \begin{cases} \dot{x} = \cos \theta, \\ \dot{r} = \sin \theta, \\ \dot{\theta} = \left(\frac{n-1}{r} - r\right) \cos \theta + x \sin \theta + \lambda. \end{cases}$$

Let P denote the projection from $\mathbb{H} \times \mathbb{R}$ to \mathbb{H} . Obviously, if $\gamma(s)$ is a solution of (2.6), then the curve $P(\gamma(s))$ will generate a λ -hypersurface by (2.1). If we can, by this way, find a curve that starts and ends on the x -axis and is perpendicular to the x -axis at both ends, then we obtain an embedded λ -hypersurface, which is diffeomorphic to a sphere. Hence, Theorem 1.1 will be proved. By some symmetry of (2.6), a curve that starts perpendicularly on the x -axis and ends perpendicularly on the r -axis will also solve the problem. This paper's main goal is to find the latter.

Letting (x_0, r_0, θ_0) be a point in $\{(x, r, \theta) : x \in \mathbb{R}, r > 0, \theta \in \mathbb{R}\}$, by an existence and uniqueness theorem of the solutions for first order ordinary differential equations, there is a unique solution $\Gamma(x_0, r_0, \theta_0)(s)$ to (2.6) satisfying initial conditions $\Gamma(x_0, r_0, \theta_0)(0) = (x_0, r_0, \theta_0)$. Moreover, the solution depends smoothly on the initial conditions. Note that (2.6) has a singularity at $r = 0$ due to the term $\frac{1}{r}$. But there is also an existence and uniqueness theorem for (2.6) with initial conditions at singularity and $\theta_0 = \pi/2$. And the previously mentioned smooth dependence can also extend to singularity smoothly (see [7]).

When the profile curve can be written in the form $(x, u(x))$, by (2.6), the function $u(x)$ satisfies the differential equation

$$(2.7) \quad \frac{u''}{1 + (u')^2} = x u' - u + \frac{n-1}{u} + \lambda \sqrt{1 + (u')^2}.$$

When the profile curve can be written in the form $(f(r), r)$, by (2.6), the function $f(r)$ satisfies the differential equation

$$(2.8) \quad \frac{f''}{1 + (f')^2} = \left(r - \frac{n-1}{r}\right) f' - f - \lambda \sqrt{1 + (f')^2}.$$

Next we consider equations (2.6) in polar coordinates, when the profile curve can be written in the form $\rho = \rho(\phi)$, where $\rho = \sqrt{x^2 + r^2}$ and $\phi = \arctan(r/x)$, the function $\rho(\phi)$ satisfies the differential equation

$$(2.9) \quad \rho'' = \frac{1}{\rho} \left\{ \rho'^2 + (\rho^2 + \rho'^2) \left[n - \rho^2 - (n-1) \frac{\rho'}{\rho} \cot \phi - \lambda \sqrt{\rho^2 + \rho'^2} \right] \right\}.$$

Note that (2.8) has a solution $f = -\lambda$, which corresponds to hyperplane, (2.9) has a solution $\rho = \frac{-\lambda + \sqrt{\lambda^2 + 4n}}{2}$, which corresponds to the round sphere.

We conclude this section with a lemma about the solutions of (2.8), this lemma can be found in [9].

Lemma 2.1 ([9]). *Let f be a solution of (2.8) with $f(0) > -\lambda$ and $f'(0) = 0$. Then we have $f'' < 0$ and there exists a point $r_* < \infty$ so that $\lim_{r \rightarrow r_*} f'(r) = -\infty$ and $\lim_{r \rightarrow r_*} f(r) > -\infty$, i.e., f blows-up at r_* .*

3. BEHAVIOR NEAR THE KNOWN SOLUTIONS

In order to solve the problem, we need to study the behavior near two known solutions first, and use continuity argument to find the curve that we want to have.

3.1. Behavior near the plane. Let $f_\epsilon(r)$ be the solution of (2.8) with $f_\epsilon(0) = -\lambda + \epsilon$ and $f'_\epsilon(0) = 0$ where $'$ denotes $\frac{d}{dr}$. For $\epsilon > 0$, by Lemma 2.1, let r_*^ϵ denote the point where f_ϵ blows-up, and let $x_*^\epsilon = f_\epsilon(r_*^\epsilon)$. Let $h_\epsilon(r) = f_\epsilon(r) + \lambda$, we may write $h_\epsilon(r)$ to $h(r)$, $f_\epsilon(r)$ to $f(r)$, r_*^ϵ to r_* and x_*^ϵ to x_* when there is no ambiguity. We introduce the following proposition which gives a description of the behavior near the plane $f_0 = -\lambda$.

Proposition 3.1. *For any fixed $n \geq 2$, $-\min\left\{\frac{23n+4}{28\sqrt{n}}, \frac{25n-6}{30\sqrt{n}}\right\} \leq \lambda < 0$, there exists $\bar{\epsilon} > 0$ so that*

$$r_*^\epsilon \geq \sqrt{\log \frac{1}{\sqrt{\pi\epsilon}}}, \quad -\frac{30n+4}{\sqrt{\log \frac{1}{\sqrt{\pi\epsilon}}}} - \lambda \leq x_*^\epsilon < -\lambda$$

for $\epsilon \in (0, \bar{\epsilon}]$.

To prove the proposition, we need several informations on $f_\epsilon(r)$ under small ϵ .

Lemma 3.1. *If $\lambda \leq 0$, $0 < \epsilon \leq \frac{1}{\sqrt{\pi}}$, then $r_* > \sqrt{\log \frac{1}{\sqrt{\pi\epsilon}}}$.*

Proof. Since $f'(0) = 0$, $f''(r) < 0$, and $\lim_{r \rightarrow r_*} f'(r) = -\infty$, there exists $r' \in (0, r_*)$ so that $f'(r') = -1$. Then $h'(r')$ also equals to -1 . For $r \in (0, r')$, we have

$$\begin{aligned} \frac{d}{dr}(e^{-r^2}h'(r)) &= e^{-r^2}h''(r) - 2re^{-r^2}h'(r) \\ &\geq \frac{2}{1+h'(r)^2}e^{-r^2}h''(r) - 2re^{-r^2}h'(r) \\ &= 2e^{-r^2} \left[\left(r - \frac{n-1}{r}\right)h'(r) - h(r) + \lambda(1 - \sqrt{1+h'(r)^2}) \right] - 2re^{-r^2}h'(r) \\ &\geq -2e^{-r^2}h(r), \end{aligned}$$

where we have used that $\lambda \leq 0$ in the last inequality. Integrating from 0 to r' ,

$$-e^{-(r')^2} \geq -2 \int_0^{r'} e^{-r^2}h(r)dr \geq -2\epsilon \int_0^{r'} e^{-r^2}dr \geq -\sqrt{\pi}\epsilon.$$

Hence, we obtain $r_* > r' \geq \sqrt{\log \frac{1}{\sqrt{\pi\epsilon}}}$ for $\lambda \leq 0$, $0 < \epsilon \leq \frac{1}{\sqrt{\pi}}$. \square

Lemma 3.2. *For any fixed $n \geq 2$, $\lambda \in \mathbb{R}$, there exists $\epsilon_1 > 0$ so that $f_\epsilon(r) = -\lambda$ (i.e., $h_\epsilon(r) = 0$) has a solution in $[\sqrt{n}, \sqrt{2n}]$ for $|\epsilon| \leq \epsilon_1$.*

Proof. We adopt an argument from the appendix B of [8]. In order to understand the behavior of $f_\epsilon(r)$ when ϵ is close to 0, we study the linearization of the rotational λ -hypersurface differential equation (2.8) near the plane $f_0(r) = -\lambda$. We define w by

$$w(r) = \frac{d}{d\epsilon} \Big|_{\epsilon=0} f_\epsilon(r).$$

Since $f_\epsilon(r)$ satisfies equation

$$\frac{f''_\epsilon}{1+(f'_\epsilon)^2} = \left(r - \frac{n-1}{r}\right)f'_\epsilon - f_\epsilon - \lambda\sqrt{1+(f'_\epsilon)^2},$$

by differentiating the above equation with respect to ϵ and letting $\epsilon = 0$, we obtain a differential equation for w :

$$(3.1) \quad w'' = \left(r - \frac{n-1}{r}\right)w' - w,$$

with $w(0) = 1$ and $w'(0) = 0$, where we have used $f'_0(r) = 0$. We will show that $w(\sqrt{n}) > 0$ and $w(\sqrt{2n}) < 0$ which lead to the lemma.

Defining $\xi = r^2$, the (3.1) becomes into the following differential equation:

$$(3.2) \quad 4\xi \frac{d^2w}{d\xi^2} = 2(\xi - n) \frac{dw}{d\xi} - w$$

with the initial conditions at $\xi = 0$:

$$w(0) = 1, \quad \frac{dw}{d\xi}(0) = -\frac{1}{2n}.$$

Taking derivatives of (3.2), we have the following second order differential equation:

$$(3.3) \quad 4\xi \frac{d^3w}{d\xi^3} = 2(\xi - n - 2) \frac{d^2w}{d\xi^2} + \frac{dw}{d\xi}$$

with the initial conditions at $\xi = 0$:

$$\frac{dw}{d\xi}(0) = -\frac{1}{2n}, \quad \frac{d^2w}{d\xi^2}(0) = -\frac{1}{4n(n+2)}.$$

We also note that the differential equation (3.3) for $\frac{dw}{d\xi}$ satisfies a maximum principle, which yields that $\frac{d^2w}{d\xi^2} < 0$ and then $\frac{dw}{d\xi} \leq \frac{dw}{d\xi}(0) < 0$. Hence w is strictly concave and strictly decreasing on $[0, \infty)$. Combining this fact with $w|_{\xi=0} = 1$ and $\frac{dw}{d\xi}(0) = -\frac{1}{2n}$ we obtain $w|_{\xi=2n} < 0$. Let $\xi = n$ in (3.2), we get $w|_{\xi=n} > 0$. Now we have proved that $w|_{\xi=n} > 0$ and $w|_{\xi=2n} < 0$, i.e., $w|_{r=\sqrt{n}} > 0$ and $w|_{r=\sqrt{2n}} < 0$. \square

Lemma 3.3. *Suppose $n \geq 2$, $-\frac{25n-6}{30\sqrt{n}} \leq \lambda < 0$. If there exists a point $r_0 \in [\sqrt{n}, \infty)$ so that $h(r_0) = 0$, we have*

$$h(r) > \frac{30n}{r} h'(r)$$

for $r \in [r_0, r_*)$.

Proof. Let $\Phi(r) = \frac{1}{30n} r h(r) - h'(r)$. We want to show that $\Phi(r) > 0$ for $r \in [r_0, r_*)$. If $r = r_0$, then $\Phi(r_0) = -h'(r_0) > 0$. If $r > r_0$, by a direct computation, we obtain

$$\begin{aligned} \Phi(r) &= \frac{1}{30n} r h(r) - h'(r) \\ &= \frac{1}{30n} r \int_{r_0}^r h'(\xi) d\xi - h'(r) \\ &> \frac{1}{30n} r(r - r_0) h'(r) - h'(r) \\ &= \frac{1}{30n} h'(r) (r^2 - r_0 r - 30n) \\ &\geq \frac{1}{30n} h'(r) (r^2 - \sqrt{nr} - 30n) \end{aligned}$$

that is, $\Phi(r) > 0$ for $r \leq 6\sqrt{n}$.

Suppose $\Phi(r) = 0$ for some $r \in [r_0, r_*)$. Then $r > 6\sqrt{n}$ and there exists a point $\bar{r} \in (6\sqrt{n}, r_*)$ so that $\Phi(\bar{r}) = 0$ and $\Phi(r) > 0$ for $r \in [r_0, \bar{r})$. This implies that $\Phi'(\bar{r}) \leq 0$ and $\frac{1}{30n} \bar{r} h(\bar{r}) = h'(\bar{r})$. On the other hand, since

$$\Phi'(\bar{r}) = \frac{1}{30n} h(\bar{r}) + \frac{1}{30n} \bar{r} h'(\bar{r}) - h''(\bar{r})$$

$$\begin{aligned}
 &\geq \frac{1}{30n}h(\bar{r}) + \frac{1}{30n}\bar{r}h'(\bar{r}) - \frac{h''(\bar{r})}{1+h'(\bar{r})^2} \\
 &= \frac{1}{30n}h(\bar{r}) + \frac{1}{30n}\bar{r}h'(\bar{r}) - \left[\left(\bar{r} - \frac{n-1}{\bar{r}} \right) h'(\bar{r}) - f(\bar{r}) - \lambda \sqrt{1+h'(\bar{r})^2} \right] \\
 &= \frac{1}{30n}h(\bar{r}) + \frac{1}{30n}\bar{r}h'(\bar{r}) - \left[\left(\bar{r} - \frac{n-1}{\bar{r}} \right) h'(\bar{r}) - h(\bar{r}) + \lambda(1 - \sqrt{1+h'(\bar{r})^2}) \right] \\
 &\geq \frac{1}{30n}h(\bar{r}) + \frac{1}{30n}\bar{r}h'(\bar{r}) - \left[\left(\bar{r} - \frac{n-1}{\bar{r}} \right) h'(\bar{r}) - h(\bar{r}) + \lambda h'(\bar{r}) \right] \\
 &= \frac{1}{30n}h(\bar{r}) + \frac{1}{900n^2}\bar{r}^2h(\bar{r}) - \left[\frac{1}{30n}\bar{r}(\bar{r} - \frac{n-1}{\bar{r}} + \lambda)h(\bar{r}) - h(\bar{r}) \right] \\
 &= \frac{1}{900n^2}h(\bar{r}) [(1-30n)\bar{r}^2 - 30n\lambda\bar{r} + 930n^2] \\
 &> 0.
 \end{aligned}$$

where we have used that $n \geq 2$, $-\frac{25n-6}{30\sqrt{n}} \leq \lambda < 0$ and $\bar{r} > 6\sqrt{n}$ in the last inequality, this contradicts $\Phi'(\bar{r}) \leq 0$. Hence, we complete the proof. \square

Proof of Proposition 3.1. The first part of the proposition 3.1 has been shown in the lemma 3.1. We know that $r' \geq \sqrt{\log \frac{1}{\sqrt{\pi}\epsilon}}$ for $0 < \epsilon \leq \frac{1}{\sqrt{\pi}}$ from the lemma 3.1. In particular, $r' \geq \sqrt{\log \frac{1}{\sqrt{\pi}\epsilon}} \geq 7\sqrt{n}$ for $0 < \epsilon \leq \frac{1}{\sqrt{\pi}e^{49n}}$. Choosing $\bar{\epsilon} = \min \left\{ \epsilon_1, \frac{1}{\sqrt{\pi}e^{49n}} \right\}$ and assuming $0 < \epsilon \leq \bar{\epsilon}$, by the lemma 3.2 and lemma 3.3, we have $h(r_*) < h(r') < h(r_0) = 0$ (that is, $x_* < -\lambda$) and

$$h(r') > -\frac{30n}{r'} \geq -\frac{30n}{\sqrt{\log \frac{1}{\sqrt{\pi}\epsilon}}}.$$

We will extend this estimate for $h(r')$ to an estimate for $h(r_*) = x_* + \lambda$. For $r \geq r'$, we have

$$\begin{aligned}
 h''(r) &< h'(r)^2 \frac{h''(r)}{1+h'(r)^2} \\
 &= h'(r)^2 \left[\left(r - \frac{n-1}{r} \right) h'(r) - h(r) + \lambda(1 - \sqrt{1+h'(r)^2}) \right] \\
 &= h'(r)^2 \left[\left(r - \frac{n-1}{r} \right) h'(r) - h(r) + \lambda h'(r) \right] \\
 &< h'(r)^3 \left(r - \frac{31n-1}{r} + \lambda \right) \\
 &\leq \frac{1}{4} r h'(r)^3,
 \end{aligned}$$

where we have used that $n \geq 2$, $r \geq 7\sqrt{n}$ and $-\frac{23n+4}{28\sqrt{n}} \leq \lambda < 0$ in the last inequality. Integrating the previous inequality from r to r_* , implies

$$h'(r)^2 \leq \frac{4}{r_*^2 - r^2}$$

for $r \geq r'$. Since $h'(r) < 0$, we have

$$(3.4) \quad h'(r) \geq -\frac{2}{\sqrt{r_*^2 - r^2}} \geq -\frac{1}{\sqrt{r_* + r'}} \frac{2}{\sqrt{r_* - r}}$$

for $r \in [r', r_*)$. At r' , this tells us that

$$-\frac{\sqrt{r_* - r'}}{\sqrt{r_* + r'}} \geq -\frac{2}{r_* + r'}.$$

Finally, integrating (3.4) from r' to r_* , we have

$$h(r_*) - h(r') \geq -\frac{4}{\sqrt{r_* + r'}} \sqrt{r_* - r'},$$

and therefore

$$\begin{aligned} h(r_*) &\geq h(r') - \frac{4}{\sqrt{r_* + r'}} \sqrt{r_* - r'} \\ &\geq -\frac{30n}{r'} - \frac{8}{r_* + r'} \\ &\geq -\frac{30n + 4}{r'} \\ &\geq -\frac{30n + 4}{\sqrt{\log \frac{1}{\sqrt{\pi}\epsilon}}}. \end{aligned}$$

Hence $x_* \geq -\frac{30n+4}{\sqrt{\log \frac{1}{\sqrt{\pi}\epsilon}}} - \lambda$, this completes the proof. \square

3.2. Behavior near the round sphere. As in the proof of the lemma 3.2, we also make use of the method from the appendix B of [8]. Recall equation (2.9), if we can write the profile curve in the form $\rho = \rho(\phi)$ in polar coordinates, where $\rho = \sqrt{x^2 + r^2}$ and $\phi = \arctan(r/x)$, then (2.9) tells us that $\rho = \rho(\phi)$ satisfies

$$(2.8) \quad \rho'' = \frac{1}{\rho} \left\{ \rho'^2 + (\rho^2 + \rho'^2) \left[n - \rho^2 - (n-1) \frac{\rho'}{\rho} \cot \phi - \lambda \sqrt{\rho^2 + \rho'^2} \right] \right\}.$$

This ordinary differential equation has a constant solution

$$\rho = \frac{-\lambda + \sqrt{\lambda^2 + 4n}}{2}$$

which corresponds to the round sphere. We note that this equation has a singularity when $\phi = 0$ due to the $\cot \phi$ term. Let $\rho(\phi, \epsilon)$ be the solution to (2.8) with initial conditions $\rho(0, \epsilon) = \frac{-\lambda + \sqrt{\lambda^2 + 4n}}{2} + \epsilon$ and $\frac{d\rho}{d\phi}(0, \epsilon) = 0$. As is Drugan and Kleene said in [8], for $\lambda = 0$, the solutions $\rho(\phi, \epsilon)$ is smooth when $\phi \in [0, \pi/2]$ and ϵ is close to 0. This is also true for $\lambda \in \mathbb{R}$.

In order to understand the behavior of $\rho(\phi, \epsilon)$ when ϵ is close to 0, we study the linearization of the rotational self-shrinker differential equation near the round sphere $\rho(\phi, 0) = \frac{-\lambda + \sqrt{\lambda^2 + 4n}}{2}$. We define w by

$$w(\phi) = \frac{d}{d\phi} \Big|_{\epsilon=0} \rho(\phi, \epsilon).$$

Then w satisfies the (singular) linear differential equation:

$$(3.5) \quad w'' = -(n-1) \cot \phi w' - \frac{-\lambda + \sqrt{\lambda^2 + 4n}}{2} \sqrt{\lambda^2 + 4n} w$$

with $w(0) = 1$ and $w'(0) = 0$. Letting $A(\lambda) = \frac{-\lambda + \sqrt{\lambda^2 + 4n}}{2} \sqrt{\lambda^2 + 4n}$, we may write $A(\lambda)$ for A when there is no ambiguity. Note that the sign of A is positive for $\lambda \in \mathbb{R}$. In the following lemma, we claim that $w(\pi/2) < 0$ and $w'(\pi/2) < 0$ which yields that, for $\epsilon < 0$

and ϵ closed to 0, $r > \frac{-\lambda + \sqrt{\lambda^2 + 4n}}{2}$ and $\pi/2 < \theta < \pi$ at the first point where the profile curve meets r -axis.

Lemma 3.4. *Let $-\frac{2}{\sqrt{n+2}} < \lambda \leq 0$, w be the solution to (3.5) with $w(0) = 1$ and $w'(0) = 0$. Then $w(\pi/2) < 0$ and $w'(\pi/2) < 0$.*

Proof. By making use of the substitution $\xi = \cos \phi$, (3.5) becomes into the following Legendre type differential equation:

$$(3.6) \quad (1 - \xi^2) \frac{d^2 w}{d\xi^2} = n \xi \frac{dw}{d\xi} - A w$$

with the initial conditions at $\xi = 1$:

$$w(1) = 1, \quad \frac{dw}{d\xi}(1) = \frac{A}{n}.$$

To prove the lemma, we need to show that $w = w(\xi)$ satisfies $w(0) < 0$ and $\frac{dw}{d\xi}(0) > 0$. Taking derivatives of (3.6), we have the following second order differential equations:

$$(3.7) \quad (1 - \xi^2) \frac{d^3 w}{d\xi^3} = (n + 2) \xi \frac{d^2 w}{d\xi^2} + (n - A) \frac{dw}{d\xi},$$

$$(3.8) \quad (1 - \xi^2) \frac{d^4 w}{d\xi^4} = (n + 4) \xi \frac{d^3 w}{d\xi^3} + (2n + 2 - A) \frac{d^2 w}{d\xi^2}.$$

It follows from (3.7) and (3.8) that

$$\frac{d^2 w}{d\xi^2}(1) = -\frac{(n - A)A}{n(n + 2)}, \quad \frac{d^3 w}{d\xi^3}(1) = \frac{(2n + 2 - A)(n - A)A}{n(n + 2)(n + 4)}.$$

By a direct calculation, we get that $n - A = \frac{1}{2}\lambda(\sqrt{\lambda^2 + 4n} - \lambda) - n < 0$ for $\lambda \leq 0$, and we also get $2n + 2 - A = -\frac{1}{2}(\lambda^2 - 4 - \lambda\sqrt{\lambda^2 + 4n})$. Then one can obtain $\lambda > -\frac{2}{\sqrt{n+2}}$ by solving inequality $2n + 2 - A > 0$. Summing up, we have $n - A < 0$ and $2n + 2 - A > 0$ for $-\frac{2}{\sqrt{n+2}} < \lambda \leq 0$, which tell us that $\frac{d^2 w}{d\xi^2}(1) > 0$ and $\frac{d^3 w}{d\xi^3}(1) < 0$. We also note that the differential equation (3.8) for $\frac{d^2 w}{d\xi^2}$ satisfies a maximum principle since $2n + 2 - A > 0$.

Using the same method as in [8], we can obtain $\frac{d^2 w}{d\xi^2}(0) > 0$ and $\frac{d^3 w}{d\xi^3}(0) < 0$, which follows from (3.6) and (3.7) that $w(0) < 0$ and $\frac{dw}{d\xi}(0) > 0$. Regarding $w = w(\phi)$ as a function of ϕ , this says that $w(\pi/2) < 0$ and $w'(\pi/2) < 0$, which proves the lemma. \square

4. PROOF OF THE THEOREM

Proof of Theorem 1.1. Let $S[x]$ denote the solution to (2.6) with initial conditions $S[x](0) = (x, 0, \pi/2)$. According to the lemma 2.1 we know that, for $x > -\lambda$, the first component of $P(S[x])$ written as a graph over the r -axis is concave down and decreasing before it blows-up at the point $B_x = (f_{x+\lambda}(r_*^{x+\lambda}), r_*^{x+\lambda}) = (x_*^{x+\lambda}, r_*^{x+\lambda})$. The proposition 3.1 tells us that when $x > -\lambda$ and x closed to $-\lambda$, B_x is in the first quadrant. From the lemma 3.4 we know that when $-\lambda < x < \frac{-\lambda + \sqrt{\lambda^2 + 4n}}{2}$ and x closed to $\frac{-\lambda + \sqrt{\lambda^2 + 4n}}{2}$, B_x is in the second quadrant. Since the mapping $(-\lambda, \infty) \rightarrow \mathbb{H}, x \mapsto B_x$ is continuous, the previous results imply that there exists $\hat{x} \in (-\lambda, \frac{-\lambda + \sqrt{\lambda^2 + 4n}}{2})$ such that $B_{\hat{x}}$ lies in r -axis. Suppose $P(S[\hat{x}] (\hat{s})) = B_{\hat{x}}$, then the curve $[0, 2\hat{s}] \rightarrow \mathbb{H}, s \mapsto P(S[\hat{x}] (s))$ will generate an embedded λ -hypersurface by (2.1), which is diffeomorphic to a sphere \mathbb{S}^n and is not isometric to the sphere $S^n(\frac{-\lambda + \sqrt{\lambda^2 + 4n}}{2})$. Moreover, by (2.3) and lemma 2.1, all the principal curvatures of

the hypersurface we construct is positive. In other words, the hypersurface we construct is convex. \square

Remark 4.1. According to some sketches, we know, for $n \geq 2$ and some $\lambda \leq -\frac{2}{\sqrt{n+2}}$, there exists an embedded λ -hypersurface $\Sigma^n \subset \mathbb{R}^{n+1}$ which is diffeomorphic to \mathbb{S}^n and is not $S^n(\frac{-\lambda + \sqrt{\lambda^2 + 4n}}{2})$ (see Figure 4.1, Figure 4.2, Figure 4.3 and Figure 4.4).




Figure 4.1 $n = 2, \lambda = -1, \hat{x} \approx 1.31$.




Figure 4.2 $n = 3, \lambda = -0.9, \hat{x} \approx 1.11$.




Figure 4.3 The graph of the λ -hypersurface generated by the profile curve in Figure 4.1




Figure 4.4 The graph of half of the λ -hypersurface generated by the profile curve in Figure 4.1

Remark 4.2. In the proof of Theorem 1.1, we can not assure that the point \hat{x} that we find in $(-\lambda, \frac{-\lambda + \sqrt{\lambda^2 + 4n}}{2})$ is unique. According to some sketches, for some n and $\lambda < 0$ (for example, $n = 3$ and $\lambda = -1$), there may be more than one points $\hat{x}_1, \hat{x}_2, \dots$ in $(-\lambda, \frac{-\lambda + \sqrt{\lambda^2 + 4n}}{2})$ so that

$$P(S[\hat{x}_1](s)), P(S[\hat{x}_2](s)), \dots$$

will generate embedded λ -hypersurfaces by (2.1) (see Figure 4.5 and Figure 4.6).




Figure 4.5 $n = 3$, $\lambda = -1$, $\hat{x} \approx 1.57$.




Figure 4.6 $n = 3$, $\lambda = -1$, $\hat{x} \approx 1.69$.

REFERENCES

- [1] S. B. Angenent, *Shrinking doughnuts* Birkhäuser, Boston-Basel-Berlin, **7**, 21-38, 1992.
- [2] S. Brendle, *Embedded self-similar shrinkers of genus 0*, Ann. of Math. **183** (2016), 715-728.
- [3] J. -E. Chang, *1-dimensional solutions of the λ -self shrinkers*, Geom. Dedicata **189** (2017), 97-112.
- [4] Q. -M. Cheng and G. Wei, *Complete λ -hypersurfaces of weighted volume-preserving mean curvature flow*, Cal. Var. Partial Differential Equations **57** (2018), 32.
- [5] Q. -M. Cheng and G. Wei, *Examples of compact λ -hypersurfaces in Euclidean spaces*, Sci. China Math. **64** (2021), 155-166.
- [6] M. do Carmo and M. Dajczer, *Hypersurfaces in space of constant curvature*, Trans. Amer. Math. Soc. **277** (1983), 685-709.
- [7] G. Drugan, *An immersed S^2 self-shrinker*. Trans. Amer. Math. Soc. **367** (2015), 3139-3159.
- [8] G. Drugan and S. J. Kleene, *Immersed self-shrinkers*, Trans. Amer. Math. Soc. **369** (2017), 7213-7250.
- [9] Z. Li and G. Wei, *An immersed S^n λ -hypersurface*, arXiv:2204.11390.
- [10] P. McGrath, *Closed mean curvature self-shrinking surfaces of generalized rotational type*, arXiv: 1507.00681. 2015.
- [11] M. McGonagle and J. Ross, *The hyperplane is the only stable, smooth solution to the isoperimetric problem in Gaussian space*, Geom. Dedicata **178** (2015), 277-296.
- [12] J. Ross, *On the existence of a closed, embedded, rotational λ -hypersurface*, J. Geom. **110** (2019), 1-12.

QING-MING CHENG, DEPARTMENT OF APPLIED MATHEMATICS, FACULTY OF SCIENCE, FUKUOKA UNIVERSITY, 814-0180, FUKUOKA, JAPAN, CHENG@FUKUOKA-U.AC.JP

JUNQI LAI, SCHOOL OF MATHEMATICAL SCIENCES, SOUTH CHINA NORMAL UNIVERSITY, 510631, GUANGZHOU, CHINA, 2019021668@M.SCNU.EDU.CN

GUOXIN WEI, SCHOOL OF MATHEMATICAL SCIENCES, SOUTH CHINA NORMAL UNIVERSITY, 510631, GUANGZHOU, CHINA, WEIGUOXIN@TSINGHUA.ORG.CN