

# $W$ -core inverses in rings with involution

Huihui Zhu<sup>a</sup>, Liyun Wu<sup>a</sup>, Jianlong Chen<sup>b</sup>

<sup>a</sup>*School of Mathematics, Hefei University of Technology, Hefei 230009, China.*

<sup>b</sup>*School of Mathematics, Southeast University, Nanjing 210096, China.*

---

## Abstract

Let  $R$  be a unital  $*$ -ring and let  $a, w, v \in R$ . The initial goal of this work is to introduce two new classes of generalized inverses, called the  $w$ -core inverse and the dual  $v$ -core inverse. It is shown that the core inverse and the pseudo core inverse belongs to the defined  $w$ -core inverse, and the dual core inverse and the dual pseudo core inverse are instances of the dual  $v$ -core inverse. An element  $a \in R$  is  $w$ -core invertible if there exists some  $x \in R$  such that  $awx^2 = x$ ,  $xawa = a$  and  $(awx)^* = awx$ . Such an  $x$  is called a  $w$ -core inverse of  $a$ . The dual  $v$ -core inverse of  $a$  is defined by the existence of  $y \in R$  satisfying  $y^2va = y$ ,  $avay = a$  and  $(yva)^* = yva$ . Several characterizations of them are given, and the expression of the  $w$ -core inverse (resp. the dual  $v$ -core inverse) of  $a$  is given by the inverse of  $w$  along  $a$  and  $\{1, 3\}$ -inverses of  $a$  (resp. the inverse of  $v$  along  $a$  and  $\{1, 4\}$ -inverses of  $a$ ). Also, the characterization for both  $w$ -core invertible and dual  $v$ -core invertible elements is given by units. Finally, the relationships among the  $w$ -core inverse, the dual  $v$ -core inverse and other generalized inverses are given.

*Keywords:* Jacobson pairs, core inverses, dual-core inverses, the inverse along an element,  $\{1, 3\}$ -inverses,  $\{1, 4\}$ -inverses, Moore-Penrose inverses, Rings with involution

*2010 MSC:* 15A09, 16W10, 16E50

---

*Email addresses:* hhzhu@hfut.edu.cn (Huihui Zhu), wlymath@163.com (Liyun Wu), jlchen@seu.edu.cn (Jianlong Chen)

## 1. Introduction

Let  $R$  be an associative ring with unity 1. Following [6], an element  $a \in R$  is Drazin invertible if there exists some  $x \in R$  such that

- (i)  $ax = xa$ ,
- (ii)  $xax = x$ ,
- (iii)  $a^k = a^{k+1}x$  for some nonnegative integer  $k$ .

Such an  $x$  is called a Drazin inverse of  $a$ . It is unique if it exists, and is denoted by  $a^D$ . The smallest nonnegative integer  $k$  in the conditions above is called the Drazin index of  $a$ , and is denoted by  $\text{ind}(a)$ . If  $\text{ind}(a) = 1$ ,  $a$  is called group invertible and the group inverse of  $a$  is denoted by  $a^\#$ . We denote by  $R^D$  and  $R^\#$  the sets of all Drazin invertible and group invertible elements in  $R$ , respectively. It is known that  $a \in R^D$  if and only if  $a^n \in a^{n+1}R \cap Ra^{n+1}$  for some positive integer  $n$ , and that  $a \in R^\#$  if and only if  $a \in a^2R \cap Ra^2$ . In particular, if  $a = a^2x = ya^2$  for some  $x, y \in R$ , then  $a^\# = yax = y^2a = ax^2$ .

Let  $a, d \in R$ . The element  $a$  is called invertible along  $d$  [12] if there exists some  $b \in R$  such that  $bad = d = dab$  and  $b \in dR \cap Rd$ . Such an element  $b$  is called the inverse of  $a$  along  $d$ . It is unique if it exists, and is denoted by  $a^{\parallel d}$ . By  $R^{\parallel d}$  we denote the set of all invertible elements along  $d$ . The inverse along an element encompasses the classical invertibility, the Drazin inverse and the group inverse. Mary in [12, Theorem 11] illustrated that (i)  $a$  is invertible if and only if  $a^{\parallel 1}$  exists, (ii)  $a \in R^D$  if and only if  $a^{\parallel a^n}$  exists for some positive integer  $n$ , (iii)  $a \in R^\#$  if and only if  $a^{\parallel a}$  exists. In these cases,  $a^{-1} = a^{\parallel 1}$ ,  $a^D = a^{\parallel a^n}$  and  $a^\# = a^{\parallel a}$ . One also knows from [13, Corollary 3.4] that  $a \in R^\#$  if and only if  $1^{\parallel a}$  exists. Moreover,  $1^{\parallel a} = aa^\#$ .

Suppose that  $R$  is a unital  $*$ -ring, that is a ring with unity 1 and an involution  $*$  satisfying  $(x^*)^* = x$ ,  $(xy)^* = y^*x^*$  and  $(x + y)^* = x^* + y^*$  for all  $x, y \in R$ . Throughout this paper, we assume that  $R$  is a unital  $*$ -ring.

Recall that an element  $a \in R$  is said to have a Moore-Penrose inverse [15] if there exists some  $x \in R$  such that

- (i)  $axa = a$ , (ii)  $xax = x$ , (iii)  $(ax)^* = ax$ , (iv)  $(xa)^* = xa$ .

Such an  $x$  is called a Moore-Penrose inverse of  $a$ . It is unique if it exists, and is denoted by  $a^\dagger$ . By  $R^\dagger$  we denote the set of all Moore-Penrose invertible elements in  $R$ . More generally, if  $a$  and  $x$  satisfy the equations (i)  $axa = a$  and (iii)  $(ax)^* = ax$ , then  $x$  is called a  $\{1, 3\}$ -inverse of  $a$ , and is denoted by

$a^{(1,3)}$ . If  $a$  and  $x$  satisfy the equations (i)  $axa = a$  and (iv)  $(xa)^* = xa$ , then  $x$  is called a  $\{1, 4\}$ -inverse of  $a$ , and is denoted by  $a^{(1,4)}$ . As usual, we denote by  $R^{(1,3)}$  and  $R^{(1,4)}$  the sets of all  $\{1, 3\}$ -invertible and  $\{1, 4\}$ -invertible elements in  $R$ , respectively. It is known that  $a \in R^{(1,3)}$  if and only if  $a \in Ra^*a$ , and  $a \in R^{(1,4)}$  if and only if  $a \in aa^*R$ . In particular, If  $a = xa^*a$  for some  $x \in R$ , then  $x^*$  is a  $\{1, 3\}$ -inverse of  $a$ . Mary [12, Theorem 11] proved also that  $a \in R^\dagger$  if and only if  $a^{\parallel a^*}$  exists. Moreover,  $a^\dagger = a^{\parallel a^*}$ .

The core inverse and the dual-core inverse of complex matrices were first introduced by Baksalary and Trenkler in their paper [1]. Let  $M_n(\mathbb{C})$  be the ring of all  $n$  by  $n$  complex matrices. Suppose  $A \in M_n(\mathbb{C})$ . A matrix  $X \in M_n(\mathbb{C})$  is called a core inverse of  $A$  if it satisfies  $AX = P_A$  and  $\mathcal{R}(X) \subseteq \mathcal{R}(A)$ , where  $\mathcal{R}(A)$  denotes the column space of  $A$ , and  $P_A$  is the orthogonal projector onto  $\mathcal{R}(A)$ . Such a matrix  $X$  is unique if it exists, and is denoted by  $A^\oplus$ . The dual-core inverse, when it exists, is the unique  $A_{\oplus}$  satisfying  $A_{\oplus}A = P_{A^*}$  and  $\mathcal{R}(A_{\oplus}) \subseteq \mathcal{R}(A^*)$ .

In 2014, Rakić et al. [16] extended the core inverse and the dual-core inverse of a complex matrix to the case of a unital  $*$ -ring  $R$ . They proved that the core inverse of  $a \in R$  is the solution of the following five equations

$$(1) \ axa = a, (2) \ xax = x, (3) \ ax^2 = x, (4) \ xa^2 = a, (5) \ (ax)^* = ax.$$

Also, they showed that the dual-core inverse of  $a \in R$  is the solution of the following five equations

$$(1') \ axa = a, (2') \ xax = x, (3') \ x^2a = x, (4') \ a^2x = a, (5') \ (xa)^* = xa.$$

As usual, by  $R^\oplus$  and  $R_{\oplus}$  we denote the sets of all core invertible and dual-core invertible elements in  $R$ , respectively.

In 2017, Xu et al. [17] found that the equations (1) and (2) above can be dropped, more precisely, they characterized the core inverse of  $a \in R$  by the solution of the following three equations

$$(3) \ ax^2 = x, (4) \ xa^2 = a, (5) \ (ax)^* = ax.$$

The dual-core inverse can also be expressed by the solution of

$$(3') \ x^2a = x, (4') \ a^2x = a, (5') \ (xa)^* = xa.$$

In addition, they derived that (i)  $a \in R^\oplus$  if and only if  $a \in R^\# \cap R^{(1,3)}$ , and (ii)  $a \in R_{\oplus}$  if and only if  $a \in R^\# \cap R^{(1,4)}$ . Moreover,  $a^\oplus = a^\#aa^{(1,3)}$  and  $a_{\oplus} = a^{(1,4)}aa^\#$ . Through the aspect of Mary's inverse along an element,

the existence criteria of the core inverse and the dual-core inverse can be correspondingly stated: (i)  $a \in R^{\oplus}$  if and only if  $1 \in R^{\parallel a}$  and  $a \in R^{(1,3)}$ , and (ii)  $a \in R_{\oplus}$  if and only if  $1 \in R^{\parallel a}$  and  $a \in R^{(1,4)}$ . Moreover,  $a^{\oplus} = 1^{\parallel a} a^{(1,3)}$  and  $a_{\oplus} = a^{(1,4)} 1^{\parallel a}$ . More results on core inverses and dual-core inverses can be referred to [21].

In [7], Gao and Chen defined the pseudo core inverse (a.k.a. the core-EP inverse [11]) by three equations in  $\ast$ -rings. An element  $a \in R$  is pseudo core invertible if there exists an  $x \in R$  such that  $xa^{m+1} = a^m$ ,  $ax^2 = x$  and  $(ax)^{\ast} = ax$  for some positive integer  $m$ . Such an  $x$  is called a pseudo core inverse of  $a$ . It is unique if it exists, and is denoted by  $a^{\odot}$ . The smallest positive integer  $m$  is called the pseudo core index of  $a$ , and is denoted by  $I(a) = m$ . We will use the symbol  $R^{\odot}$  to denote all pseudo core invertible elements in  $R$ . One knows from [7] that  $a \in R^{\odot}$  with  $I(a) = m$  if and only if  $a \in R^D$  with  $\text{ind}(a) = m$  and  $a^k \in R^{(1,3)}$ , for any integer  $k \geq m$ .

As is stated above, the core inverse (pseudo core inverse) can be explained by the intersection of the group inverse (the Drazin inverse) and  $\{1,3\}$ -inverse. It is well known that the inverse along an element extends the group inverse and the Drazin inverse. So, it is of interest to ask what is the intersection of the inverse along an element and  $\{1,3\}$ -inverse.

The paper is organized as follows. In Section 2, the  $w$ -core inverse and the dual  $v$ -core inverse are defined. Then, several existence criteria of them are given. In particular, we show in Theorem 2.9 that  $a$  is  $w$ -core invertible if and only if  $w$  is invertible along  $a$  and  $a$  is  $\{1,3\}$ -invertible. Moreover,  $w^{\parallel a} a^{(1,3)}$  is the  $w$ -core inverse of  $a$ . Also, we show in Theorem 2.20 that  $a$  is dual  $v$ -core invertible if and only if  $v$  is invertible along  $a$  and  $a$  is  $\{1,4\}$ -invertible. Moreover,  $a^{(1,4)} v^{\parallel a}$  is the dual  $v$ -core inverse of  $a$ . It is also shown that  $a$  is  $w$ -core invertible if and only if  $a^{\ast}$  is dual  $w^{\ast}$ -core invertible. In particular,  $a$  is  $a^{\ast}$ -core invertible if and only if it is Moore-Penrose invertible if and only if it is dual  $a^{\ast}$ -core invertible. Further, it is shown that  $a$  is  $w$ -core invertible if and only if there exists a (unique) projection  $p \in R$  such that  $pa = 0$  and  $u = p + aw \in R^{-1}$ . In Section 3, we in Theorem 3.4 characterize both  $w$ -core invertible and dual  $v$ -core invertible elements by units, under certain assumptions. Also, a counterexample is given to show that the assumption can not be removed in a general  $\ast$ -ring  $R$ . Specially, if  $R$  is a Dedekind-finite ring, the hypothesis can be dropped (see Theorem 3.7). In Section 4, we show that the  $w$ -core inverse and the dual  $v$ -core inverse are both instances of Mary's inverses along an element and Drazin's hybrid  $(b, c)$ -inverses.

## 2. The $w$ -core inverse and the dual $v$ -core inverse

Our main result in this section is to give several characterizations for the  $w$ -core inverse and the dual  $v$ -core inverse. Theorem 2.9 establishes the equivalence that  $a$  is  $w$ -core invertible if and only if  $w \in R^{\parallel a}$  and  $a \in R^{(1,3)}$ . Theorem 2.20 presents the equivalence that  $a$  is dual  $v$ -core invertible if and only if  $v \in R^{\parallel a}$  and  $a \in R^{(1,4)}$ .

**Definition 2.1.** *Let  $a, w \in R$ . An element  $a$  is called  $w$ -core invertible if there exists some  $x \in R$  such that  $awx^2 = x$ ,  $xawa = a$  and  $awx = (awx)^*$ . Such an  $x$  is called a  $w$ -core inverse of  $a$ .*

**Lemma 2.2.** *For any  $a, w \in R$ , if  $x \in R$  is the  $w$ -core inverse of  $a$ , then  $awxa = a$  and  $xawx = x$ .*

PROOF. As  $a$  is  $w$ -core invertible with the  $w$ -core inverse  $x$ , then  $awx^2 = x$ ,  $xawa = a$  and  $(awx)^* = awx$ . Hence,  $a = xawa = (awx^2)awa = (awx)xawa = awxa$ , and  $x = awx^2 = (xawa)wx^2 = xaw(awx^2) = xawx$ .  $\square$

In what follows, we establish a few facts about the  $w$ -core inverse in a  $*$ -ring not subjected to any additional conditions. Most fundamentally, we have

**Theorem 2.3.** *Let  $a, w \in R$ . Then  $a$  has at most one  $w$ -core inverse in  $R$ .*

PROOF. Suppose that  $x, y \in R$  are any two  $w$ -core inverses of  $a$ . Then  $awx^2 = x$ ,  $awy^2 = y$ ,  $xawa = a = yawa$ ,  $(awx)^* = awx$  and  $(awy)^* = awy$ . It follows from Lemma 2.2 that  $awxa = a = awya$ ,  $xawx = x$  and  $yawy = y$ .

So, we have

$$\begin{aligned}
 x &= xawx = x(awya)wx = x(awy)(awx) \\
 &= x(awy)^*(awx)^* = x(awxawy)^* \\
 &= x(awy)^* = xawy = xaw(awy^2) \\
 &= (xawa)wy^2 = awy^2 \\
 &= y.
 \end{aligned}$$

The proof is complete.  $\square$

In view of Theorem 2.3, it is known that the  $w$ -core inverse is unique if it exists. The  $w$ -core inverse of  $a$  is denoted by  $a_w^\oplus$ . We denote by  $R_w^\oplus$  all  $w$ -core invertible elements in  $R$ .

It is not difficult to observe that the 1-core inverse is just the core inverse. So, core invertible elements are  $w$ -core invertible. However,  $w$ -core invertible elements may not be core invertible as the following example shows.

**Example 2.4.** Let  $R$  be the ring of all  $2 \times 2$  complex matrices and let the involution  $*$  be the conjugate transpose. Suppose  $a = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ ,  $w = \begin{bmatrix} 3 & 6 \\ 1 & 0 \end{bmatrix} \in R$ . Then  $a$  is  $w$ -core invertible and  $a_w^\oplus = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ . Clearly,  $a \notin a^2R \cap Ra^2$ , i.e.,  $a \notin R^\#$ , and hence  $a \notin R^\oplus$ .

**Remark 2.5.** For any  $a, w_1, w_2 \in R$ ,  $a_{w_1}^\oplus$  and  $a_{w_2}^\oplus$  are not equal in general provided that  $a_{w_1}^\oplus$  and  $a_{w_2}^\oplus$  both exist. There are, of course, lots of examples to illustrate this fact. However, we can construct a counterexample such that  $a_{w_1}^\oplus = a_{w_2}^\oplus = \cdots = a_{w_n}^\oplus = \cdots$ , for different  $w_i$  ( $i = 1, 2, \dots, n, \dots$ ). Such as, let  $R$  and the involution  $*$  be the same as that of the previous Example 2.4. Take  $a = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \in R$ , then  $a_{w_i}^\oplus = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$  for any  $w_i$  of the form  $\begin{bmatrix} x & y \\ 1 & 0 \end{bmatrix}$ , where  $x, y \in \mathbb{C}$ .

Given any  $a \in R$ , the right annihilator of  $a$  is denoted by  $a^0 = \{x \in R : ax = 0\}$  and the left annihilator of  $a$  is denoted by  ${}^0a = \{x \in R : xa = 0\}$ . Recall that (see e.g. [16, Lemmas 2.5 and 2.6]) for any  $a, b \in R$ , if  $aR = bR$  then  ${}^0a = {}^0b$ , and if  $Ra = Rb$  then  $a^0 = b^0$ .

We next give several characterizations of the  $w$ -core inverse in a ring  $R$ .

**Theorem 2.6.** *Let  $a, w \in R$ . Then the following conditions are equivalent:*

- (i)  $a$  is  $w$ -core invertible.
- (ii) There exists some  $x \in R$  such that (1)  $awxa = a$ , (2)  $xawx = x$ , (3)  $(awx)^* = awx$ , (4)  $xawa = a$  and (5)  $awx^2 = x$ .
- (iii) There exists some  $x \in R$  such that  $awxa = a$ ,  $xR = aR$  and  $Rx = Ra^*$ .
- (iv) There exists some  $x \in R$  such that  $awxa = a$ ,  $xR = x^*R = aR$ .
- (v) There exists some  $x \in R$  such that  $awxa = a$ ,  ${}^0x = {}^0a$  and  $x^0 = (a^*)^0$ .
- (vi) There exists some  $x \in R$  such that  $awxa = a$ ,  ${}^0x = {}^0a$  and  $(a^*)^0 \subseteq x^0$ .

PROOF. (i)  $\Rightarrow$  (ii) by Lemma 2.2.

(ii)  $\Rightarrow$  (iii) Given (ii), we have (4)  $xawa = a$  and (5)  $awx^2 = x$ , which imply  $aR = xawaR \subseteq xR$  and  $xR = awx^2R \subseteq aR$ . So,  $xR = aR$ . Also, by (2)  $xawx = x$  and (3)  $(awx)^* = awx$ , we have  $Rx = Rxawx = Rx(awx)^* = Rxx^*w^*a^* \subseteq Ra^*$ . Note that (1)  $awxa = a$  and (3)  $(awx)^* = awx$ . Then  $Ra^* = R(awxa)^* = Ra^*(awx)^* = Ra^*awx \subseteq Rx$ . Therefore,  $Rx = Ra^*$ .

(iii)  $\Rightarrow$  (iv), (iv)  $\Rightarrow$  (v) and (v)  $\Rightarrow$  (vi) are clear.

(vi)  $\Rightarrow$  (i) From  $awxa = a$ , it follows that  $(1-awx)a = 0$  and  $1-awx \in {}^0a$ . Since  ${}^0x = {}^0a$ , we have  $(1-awx)x = 0$ , i.e.,  $awx^2 = x$ . Note that  $a^* = a^*(awx)^*$  and  $(a^*)^0 \subseteq x^0$ . Then  $x(1-(awx)^*) = 0$ , namely  $x = x(awx)^*$ . Then  $awx = awx(awx)^*$ , which implies  $awx = (awx)^*$ . So,  $a^* = a^*awx$ , which together with  $(a^*)^0 \subseteq x^0$  ensures  $(1-xaw)x = 0$ . Again by  ${}^0x = {}^0a$ , we have  $(1-xaw)a = 0$ , i.e.,  $xawa = a$ . So,  $a$  is  $w$ -core invertible.  $\square$

Set  $w = 1$  in Theorem 2.6, we get the characterization for the core inverse in a ring  $R$ , some of which were given in [16].

**Corollary 2.7.** *Let  $a \in R$ . Then the following conditions are equivalent:*

- (i)  $a$  is core invertible.
- (ii) There exists some  $x \in R$  such that (1)  $axa = a$ , (2)  $xax = x$ , (3)  $(ax)^* = ax$ , (4)  $xa^2 = a$  and (5)  $ax^2 = x$ .
- (iii) There exists some  $x \in R$  such that  $axa = a$ ,  $xR = aR$  and  $Rx = Ra^*$ .
- (iv) There exists some  $x \in R$  such that  $axa = a$ ,  $xR = x^*R = aR$ .
- (v) There exists some  $x \in R$  such that  $axa = a$ ,  ${}^0x = {}^0a$  and  $x^0 = (a^*)^0$ .
- (vi) There exists some  $x \in R$  such that  $axa = a$ ,  ${}^0x = {}^0a$  and  $(a^*)^0 \subseteq x^0$ .

The following theorem, a main result of this paper, presents the representation of the  $w$ -core inverse by the inverse along an element and  $\{1, 3\}$ -inverses. First, an auxiliary lemma about the existence criterion of the inverse along an element is given.

**Lemma 2.8.** [13, Theorem 2.2] *Let  $a, d \in R$ . Then  $a \in R^{\parallel d}$  if and only if  $d \in dadR \cap Rdad$ . In this case,  $a^{\parallel d} = dx = yd$ , where  $x, y \in R$  satisfy  $d = dadx = ydad$ .*

**Theorem 2.9.** *Let  $a, w \in R$ . Then  $a \in R_w^{\oplus}$  if and only if  $w^{\parallel a}$  and  $a^{(1,3)}$  both exist. In this case,  $a_w^{\oplus} = w^{\parallel a}a^{(1,3)}$ .*

PROOF. First suppose  $x \in R$  is the  $w$ -core inverse of  $a$ . Then, by Theorem 2.6, we have  $a = xawa \in Rawa$  and  $a = awxa = aw(awx^2)a \in awaR$ , which give  $a \in awaR \cap Rawa$ , and hence  $w \in R^{\parallel a}$  by Lemma 2.8.

Again, by Theorem 2.6, we have  $(awx)^* = awx$  and  $awxa = a$ , and hence  $a \in R^{(1,3)}$ .

Conversely, if  $w^{\parallel a}$  and  $a^{(1,3)}$  exist, then  $x = w^{\parallel a}a^{(1,3)}$  is the  $w$ -core inverse of  $a$ . Indeed, we have

(1) As  $w^{\parallel a} \in aR$ , there exists some  $y \in R$  such that  $w^{\parallel a} = ay$  and hence  $aa^{(1,3)}w^{\parallel a} = aa^{(1,3)}ay = ay = w^{\parallel a}$ . So,  $awx^2 = (aww^{\parallel a})a^{(1,3)}w^{\parallel a}a^{(1,3)} = aa^{(1,3)}w^{\parallel a}a^{(1,3)} = w^{\parallel a}a^{(1,3)} = x$ .

(2) Note also that  $w^{\parallel a} \in Ra$ . Then  $w^{\parallel a}a^{(1,3)}a = w^{\parallel a}$  and  $xawa = w^{\parallel a}a^{(1,3)}awa = w^{\parallel a}wa = a$ .

(3)  $(awx)^* = awx$  since  $awx = aww^{\parallel a}a^{(1,3)} = aa^{(1,3)}$ .

**Remark 2.10.** In Theorem 2.9 above, the  $w$ -core inverse of  $a \in R$  is expressed by the product of  $w^{\parallel a}$  and  $a^{(1,3)}$ . It is well known that an element can have different  $\{1, 3\}$ -inverses. However, the product of  $w^{\parallel a}$  and  $a^{(1,3)}$ , i.e.,  $w^{\parallel a}a^{(1,3)}$  is unique, that is for  $x, y \in a\{1, 3\}$ , we have  $w^{\parallel a}x = w^{\parallel a}y$ . Indeed, the equality  $ax = aya x = (ay)^*(ax)^* = (axay)^* = (ay)^* = ay$  implies  $w^{\parallel a}x = w^{\parallel a}y$  since  $w^{\parallel a} \in Ra$ .

Applying Lemma 2.11 below, we can obtain another representation of the  $w$ -core inverse in  $R$ .

**Lemma 2.11.** [13, Theorem 2.1] *Let  $a, w \in R$ . Then the following conditions are equivalent:*

- (i)  $w \in R^{\parallel a}$ .
- (ii)  $a \in awR$  and  $aw \in R^\#$ .
- (iii)  $a \in Rwa$  and  $wa \in R^\#$ .

*In this case,  $w^{\parallel a} = a(wa)^\# = (aw)^\#a$ .*

**Proposition 2.12.** *Let  $a, w \in R$ . Then  $a \in R_w^\oplus$  if and only if  $w^{\parallel a}$  and  $a^{(1,3)}$  both exist. In this case, we have  $a_w^\oplus = a(wa)^\#a^{(1,3)} = (aw)^\#aa^{(1,3)}$ .*

If we particularize  $R$  to  $M_n(\mathbb{C})$ , the ring of all  $n \times n$  complex matrices, then we can obtain the concise existence criteria for the  $W$ -core inverse of  $A$ . It should be mentioned that the notion of the  $W$ -core inverse was not considered so far in the context of complex matrices.



**Theorem 2.13.** *Let  $A, W, X \in M_n(\mathbb{C})$ . Then the following conditions are equivalent:*

- (i)  $X$  is the  $W$ -core inverse of  $A$ .
- (ii)  $AWX = P_A$  and  $\mathcal{R}(X) = \mathcal{R}(A)$ .
- (iii)  $AWX = P_A$  and  $\mathcal{R}(X) \subseteq \mathcal{R}(A)$ .

PROOF. (i)  $\Rightarrow$  (ii) As  $X$  is the  $W$ -core inverse of  $A$ , then, by the implication (i)  $\Rightarrow$  (iii) of Theorem 2.6, we have  $XM_n = AM_n$ . So,  $\mathcal{R}(X) = \mathcal{R}(A)$ . It follows from Theorem 2.9 that  $X = W^{\parallel A}A^\dagger$  and hence  $AWX = AWW^{\parallel A}A^\dagger = AA^\dagger = P_A$ .

(ii)  $\Rightarrow$  (iii) is a tautology.

(iii)  $\Rightarrow$  (i) Given  $AWX = P_A$ , then  $AWX = (AWX)^*$ . There exists some  $Y \in M_n(\mathbb{C})$  such that  $X = AY$  since  $\mathcal{R}(X) \subseteq \mathcal{R}(A)$ . Also, it follows from  $AWX = P_A$  that  $AWXA = P_AA = A$  and hence  $AWX^2 = AWXAY = AY = X$ . Note that  $A^* = (AWXA)^* = A^*(AWX)^* = A^*AWX$  and  $\mathcal{R}(X) \subseteq \mathcal{R}(A)$  (i.e.,  $\mathcal{C}(A^*)$ , where  $\mathcal{C}(A^*)$  denotes the row space of  $A^*$ ). Then there is some  $Z \in M_n(\mathbb{C})$  such that  $X = ZA^* = ZA^*AWX = XAWX$ . Observe that  $\text{rank}(A) = \text{rank}(AWXA) \leq \text{rank}(X)$ . Then  $\mathcal{R}(X) = \mathcal{R}(A)$  and  $A = XN$  for some  $N \in M_n(\mathbb{C})$ . We hence have  $A = XN = XAWXN = XAWA$ , as required.  $\square$

As a consequence of Theorem 2.9, we have the following result.

**Corollary 2.14.** *Let  $A, W \in M_n(\mathbb{C})$ . Then*

- (i)  $A$  is  $W$ -core invertible if and only if  $W$  is invertible along  $A$ . In this case,  $A_W^\oplus = W^{\parallel A}A^\dagger$ .
- (ii) If  $W^{\parallel A}$  exists and  $\lambda \in \mathbb{C}/\{0\}$ , then  $(\lambda A)_W^\oplus = \frac{1}{\lambda}A_W^\oplus$ .

In terms of Lemma 2.8 and Corollary 2.14, the existence criterion for the  $W$ -core inverse of  $A$  is given below in the ring of all  $n \times n$  complex matrices.

**Theorem 2.15.** *Let  $A, W \in M_n(\mathbb{C})$ . Then  $A$  is  $W$ -core invertible if and only if  $\text{rank}(A) = \text{rank}(AWA)$ .*

Let us now back to a general  $*$ -ring. In Proposition 2.17 below, we show that  $a \in R$  is core invertible if and only if it is  $a$ -core invertible, and that  $a$  is pseudo core invertible with  $I(a) = n$  if and only if  $a^n$  is  $a$ -core invertible. Also, we prove in Proposition 2.25 that  $a \in R$  is Moore-Penrose invertible if and only if it is  $a^*$ -core invertible if and only if it is dual  $a^*$ -core invertible.

**Proposition 2.16.** *Let  $a \in R$ . Then the following conditions are equivalent:*

- (i)  $a \in R^{\oplus}$ .
- (ii)  $a \in R^{\#} \cap R^{(1,3)}$ .
- (iii)  $a \in R_a^{\oplus}$ .
- (iv) *There exists some  $x \in R$  such that  $a^2x^2 = x$ ,  $xa^3 = a$  and  $(a^2x)^* = a^2x$ .*

*In this case,  $a^{\oplus} = aa_a^{\oplus}$  and  $a_a^{\oplus} = a^{\#}a^{\oplus}$ .*

PROOF. (i)  $\Leftrightarrow$  (ii) by taking  $w = 1$  in Proposition 2.12.

(ii)  $\Leftrightarrow$  (iii) by Theorem 2.9 and  $a \in R^{\#} \Leftrightarrow a \in R^{\parallel a}$ .

(iii)  $\Leftrightarrow$  (iv) by taking  $w = a$ .

It is known that  $x \in R$  satisfying the condition (iv) is the  $a$ -core inverse of  $a$ , and  $x = a^{\parallel a}a^{(1,3)} = a^{\#}a^{(1,3)}$ .

So,  $a^{\oplus} = ax = aa_a^{\oplus}$  and  $x = a_a^{\oplus} = a^{\#}a^{\oplus}$ .  $\square$

**Proposition 2.17.** *Let  $a \in R$  and let  $n$  be a positive integer. Then  $a \in R^{\mathfrak{D}}$  if and only if  $a^n \in R_a^{\oplus}$ .*

PROOF. Suppose  $a \in R^{\mathfrak{D}}$  with  $I(a) = n$ . Then, by [7, Theorem 2.3],  $a \in R^D$  with  $\text{ind}(a) = n$  and  $a^n \in R^{(1,3)}$ . Again, it follows from [12, Theorem 11](2) that  $a \in R^D$  if and only if  $a$  is invertible along  $a^n$ , for some positive integer  $n$ . So,  $a \in R^{\mathfrak{D}}$  implies the fact that  $a$  is invertible along  $a^n$  and  $a^n \in R^{(1,3)}$ , and hence  $a^n \in R_a^{\oplus}$  by Theorem 2.9.

For the converse part, given  $a^n \in R_a^{\oplus}$ , then  $a^{\parallel a^n}$  exists and  $a^n \in R^{(1,3)}$ . As  $a^{\parallel a^n}$  exists, then  $a \in R^D$  with  $\text{ind}(a) = n$  in terms of [12, Theorem 11](2), so that  $a \in R^{\mathfrak{D}}$ .  $\square$

Dually, we can give the definition of the dual  $v$ -core inverse in a ring  $R$ .

**Definition 2.18.** *For any  $a, v \in R$ ,  $a$  is called dual  $v$ -core invertible if there exists some  $y \in R$  such that  $avay = a$ ,  $y^2va = y$  and  $yva = (yva)^*$ . Such an  $y$  is called a dual  $v$ -core inverse of  $a$ .*

The dual  $v$ -core inverse of  $a$  is unique if it exists, and is denoted by  $a_{v, \oplus}$ . By  $R_{v, \oplus}$  we denote all dual  $v$ -core invertible elements of  $R$ . A characterization of the dual  $v$ -core inverse is given below.

**Theorem 2.19.** *Let  $a, v \in R$ . Then the following conditions are equivalent:*

- (i)  *$a$  is dual  $v$ -core invertible.*
- (ii) *There exists some  $y \in R$  such that (1')  $ayva = a$ , (2')  $yvay = y$ , (3')  $(yva)^* = yva$ , (4')  $avay = a$  and (5')  $y^2va = y$ .*
- (iii) *There exists some  $y \in R$  such that  $ayva = a$ ,  $yR = a^*R$  and  $Ry = Ra$ .*
- (iv) *There exists some  $y \in R$  such that  $ayva = a$ ,  $yR = y^*R = a^*R$ .*
- (v) *There exists some  $y \in R$  such that  $ayva = a$ ,  ${}^0y = {}^0(a^*)$  and  $y^0 = a^0$ .*
- (vi) *There exists some  $y \in R$  such that  $ayva = a$ ,  ${}^0(a^*) \subseteq {}^0y$  and  $a^0 = y^0$ .*

Dually, we can give the formula of the dual  $v$ -core inverse of  $a$  by  $v^{\parallel a}$  and  $a^{(1,4)}$  in a ring  $R$ .

**Theorem 2.20.** *Let  $a, v \in R$ . Then  $a \in R_{v, \oplus}$  if and only if  $v^{\parallel a}$  and  $a^{(1,4)}$  both exist. In this case,  $a_{v, \oplus} = a^{(1,4)}v^{\parallel a} = a^{(1,4)}a(va)^{\#} = a^{(1,4)}(av)^{\#}a$ .*

**Corollary 2.21.** [17, Theorem 2.8] *Let  $a \in R$ . Then  $a \in R_{\oplus}$  if and only if  $a \in R^{\#} \cap R^{(1,4)}$ . In this case,  $a_{\oplus} = a^{(1,4)}aa^{\#}$ .*

Combining with Theorems 2.9 and 2.20, we have the following result.

**Proposition 2.22.** *Let  $a, w, v \in R$ . Then  $a \in R_w^{\oplus} \cap R_{v, \oplus}$  if and only if  $w, v \in R^{\parallel a}$  and  $a \in R^{\dagger}$ .*

For any  $a, w \in R$ , it is known from Lemma 2.8 that  $w$  is invertible along  $a$  if and only if  $a \in awaR \cap Raw$  if and only if  $a^* \in a^*w^*a^*R \cap Ra^*w^*a^*$  if and only if  $w^*$  is invertible along  $a^*$ . Moreover,  $(w^*)^{\parallel a^*} = (w^{\parallel a})^*$ . One also knows that  $a \in R^{(1,3)}$  if and only if  $a^* \in R^{(1,4)}$ . Moreover,  $(a^*)^{(1,4)} = (a^{(1,3)})^*$ . We hence have the following result.

**Proposition 2.23.** *Let  $a, w \in R$ . Then  $a$  is  $w$ -core invertible if and only if  $a^*$  is dual  $w^*$ -core invertible. In this case,  $(a^*)_{w^*, \oplus} = (a_w^{\oplus})^*$ .*

PROOF. By Theorems 2.9 and 2.20, we have  $(a^*)_{w^*, \oplus} = (a^*)^{(1,4)}(w^*)^{\parallel a^*} = (a^{(1,3)})^*(w^{\parallel a})^* = (w^{\parallel a}a^{(1,3)})^* = (a_w^{\oplus})^*$ .  $\square$

**Lemma 2.24.** [20, Theorem 3.12] *Let  $a \in R$ . The following conditions are equivalent:*

- (i)  *$a \in R^{\dagger}$ .*

(ii)  $a \in aa^*aR$ .

(iii)  $a \in Raa^*a$ .

In this case,  $a^\dagger = (ax)^* = (ya)^*$ , where  $x, y \in R$  satisfy  $a = aa^*ax = yaa^*a$ .

The following result shows that the existence of the  $a^*$ -core inverse of  $a$  coincides with the existence of its dual  $a^*$ -core inverse, which are indeed equivalent to the existence of its Moore-Penrose inverse.

**Proposition 2.25.** *Let  $a \in R$ . Then the following conditions are equivalent:*

(i)  $a \in R_{a^*}^{\oplus}$ .

(ii)  $a \in R^\dagger$ .

(iii)  $a \in R_{a^*, \oplus}$ .

In this case,  $a^\dagger = (a_{a^*}^{\oplus})^* = (aa_{a^*, \oplus})^*$ ,  $a_{a^*}^{\oplus} = (a^\dagger)^*a^\dagger$  and  $a_{a^*, \oplus} = a^\dagger(a^\dagger)^*$ .

PROOF. (i)  $\Rightarrow$  (ii) Suppose  $x \in R$  is the  $a^*$ -core inverse of  $a$ . Then, by taking  $w = a^*$ , we have  $a = xaa^*a \in Raa^*a$ , and hence  $a \in R^\dagger$  by Lemma 2.24. Moreover,  $(xa)^*$  is the Moore-Penrose inverse of  $a$ .

(ii)  $\Rightarrow$  (iii) To show  $a \in R_{a^*, \oplus}$ , it suffices to prove  $(a^*)^{\parallel a}$  (i.e.,  $a \in R^\dagger$ ) and  $a^{(1,4)}$  both exist. The condition (ii) immediately gives this.

(iii)  $\Rightarrow$  (i) From Theorems 2.9 and 2.20, it is known that  $a \in R_{a^*, \oplus}$  if and only if both  $(a^*)^{\parallel a}$  and  $a^{(1,4)}$  exist if and only if  $(a^*)^{\parallel a}$  exists if and only if  $(a^*)^{\parallel a}$  and  $a^{(1,3)}$  exist if and only if  $a \in R_{a^*}^{\oplus}$ .

By a direct calculation, we get that  $(a^\dagger)^*a^\dagger$  is the  $a^*$ -core inverse of  $a$ , and  $a^\dagger(a^\dagger)^*$  is the dual  $a^*$ -core inverse of  $a$ .  $\square$

Given any  $a, w \in R$ , the  $w$ -core inverse of  $a$  can also be equivalently described in terms of the intersection of some ideals and the core inverse of  $aw$ .

**Theorem 2.26.** *Let  $a, w \in R$  and let  $n \geq 2$  be an integer. Then the following conditions are equivalent:*

(i)  $a \in R_w^{\oplus}$ .

(ii)  $a \in R[(aw)^*]^n a \cap R(aw)^{n-1}a$ .

(iii)  $a \in awR$  and  $aw \in R^{\oplus}$ .

In this case,  $a_w^{\oplus} = (aw)^{\oplus} = w^{\parallel a}w(aw)^{(1,3)}$ .

PROOF. (i)  $\Rightarrow$  (ii) As  $a$  is  $w$ -core invertible, then there exists some  $x \in R$  such that  $xawa = a$ ,  $awx^2 = x$  and  $(awx)^* = awx$ , which guarantee  $xawaw = aw$ ,  $awx^2 = x$  and  $(awx)^* = awx$ . So  $aw \in R^\oplus$  by Corollary 2.7. Applying [10, Theorem 2.10], we know that  $aw \in R^\oplus$  implies that  $aw \in R[(aw)^*]^n aw \cap R(aw)^n$  for all integers  $n \geq 2$ . It follows from Theorem 2.9 that  $a \in R_w^\oplus$  implies  $w \in R^{\parallel a}$  and hence  $a \in awaR \subseteq awR$ . Then  $a \in R[(aw)^*]^n a \cap R(aw)^{n-1}a$  for any integer  $n \geq 2$ .

(ii)  $\Rightarrow$  (iii) Given  $a \in R[(aw)^*]^n a \cap R(aw)^{n-1}a$ , then  $aw \in R[(aw)^*]^n aw \cap R(aw)^n$  and so  $aw \in R^\oplus$  by [10, Theorem 2.10]. Since  $a \in R[(aw)^*]^n a$ , there exists some  $x \in R$  such that  $a = x[(aw)^*]^n a = x[(aw)^*]^{n-1} w^* a^* a \subseteq Ra^*a$ . So,  $a \in R^{(1,3)}$  and  $w(aw)^{n-1}x^*$  is a  $\{1,3\}$ -inverse of  $a$ . We have at once  $a = aw(aw)^{n-1}x^*a \in awR$ .

(iii)  $\Rightarrow$  (i) As  $aw \in R^\oplus$ , then there exists some  $y \in R$  satisfying  $awyaw = aw$ ,  $yawy = y$ ,  $y(aw)^2 = aw$ ,  $awy^2 = y$  and  $awy = (awy)^*$  by Corollary 2.7. Since  $a \in awR$ , there exists some  $t \in R$  such that  $a = awt = (awyaw)t = awya = (awy)^*a = (wy)^*a^*a \in Ra^*a$ , i.e.,  $a \in R^{(1,3)}$ .

Note also that  $a = awya = aw(awy^2)a \in awaR$  and  $a = awt = y(aw)^2t = yaw(awt) = yawa \in Rawa$ . Then  $w \in R^{\parallel a}$ , which combines with  $a \in R^{(1,3)}$  to ensure that  $a \in R_w^\oplus$  by Theorem 2.6.

It follows from Lemma 2.11 and Proposition 2.17 that

$$(aw)^\oplus = (aw)^\# aw(aw)^{(1,3)} = w^{\parallel a} w(aw)^{(1,3)}.$$

To show that  $w^{\parallel a} w(aw)^{(1,3)}$  is the  $w$ -core inverse of  $a$ , it suffices to prove that  $z = w(aw)^{(1,3)}$  is a  $\{1,3\}$ -inverse of  $a$  by Theorem 2.9. Since  $a \in awR$ , it follows that  $a = aw(aw)^{(1,3)} awt = aw(aw)^{(1,3)} a = aza$ . Also,  $az = aw(aw)^{(1,3)} = (az)^*$ , as required.  $\square$

**Remark 2.27.** In Theorem 2.26, (i)  $\Leftrightarrow$  (ii) does not hold for  $n = 1$ , i.e.,  $a \in R(aw)^*a \cap Ra$  can not imply  $a \in R_w^\oplus$ . Such as, let  $R$  be an infinite complex matrix ring whose rows and columns are both finite and let the involution  $*$  be the conjugate transpose. Suppose  $a = \sum_{i=1}^{\infty} e_{i+1,i}$  and  $w = 1$ . Then  $a^*a = 1$  and  $a \in R(aw)^*a \cap Ra$ . However,  $w^{\parallel a} = 1^{\parallel a}$  does not exist, so that  $a$  is not  $w$ -core invertible by Theorem 2.9. In fact, if  $1^{\parallel a}$  exists, then, by Lemma 2.8,  $a \in (a \cdot 1 \cdot a)R \cap R(a \cdot 1 \cdot a)$ , and consequently  $a = a^2s$  for some  $s \in R$ . So,  $1 = a^*a = a^*a^2s = as$ , which together with  $a^*a = 1$  to guarantee that  $a$  is invertible. A contradiction.

For the group inverse, the Drazin inverse and the core inverse, one knows that  $(a^\#)^\# = a^2a^\#$ ,  $(a^D)^D = a^2a^D$  and  $(a^\oplus)^\oplus = a^2a^\oplus$ .

It is natural to ask that whether the  $w$ -core inverse also shares a similar property, i.e., whether  $w$ -core invertible elements are  $w$ -core invertible. In fact, the answer to this question is negative. See the following example.

**Example 2.28.** Let  $R$  and the involution  $*$  be the same as that of the previous Example 2.4. Take  $a = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ ,  $w = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \in R$ , then  $a_w^\oplus = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$  by Remark 2.5. However,  $a_w^\oplus$  is not  $w$ -core invertible as there exists no  $x \in R$  such that  $xa_w^\oplus wa_w^\oplus = a_w^\oplus$  since  $a_w^\oplus wa_w^\oplus = 0$ .

It is of interest to study that whether  $w$ -core invertible elements are core invertible. The following result provides a positive answer.

**Theorem 2.29.** *Let  $a, w \in R$  and let  $a \in R_w^\oplus$ . Then  $a_w^\oplus \in R^\oplus$  and  $(a_w^\oplus)^\oplus = (aw)^2 a_w^\oplus$ .*

PROOF. Let  $x = (aw)^2 a_w^\oplus$ . It is routine to check that  $x$  is the core inverse of  $a_w^\oplus$  by the following three steps.

- (1)  $a_w^\oplus x = a_w^\oplus (aw)^2 a_w^\oplus = aw a_w^\oplus = (a_w^\oplus x)^*$ .
  - (2)  $a_w^\oplus x^2 = aw a_w^\oplus (aw)^2 a_w^\oplus = (aw)^2 a_w^\oplus = x$ .
  - (3)  $x(a_w^\oplus)^2 = (aw)^2 a_w^\oplus (a_w^\oplus)^2 = (aw)^2 (a_w^\oplus)^2 a_w^\oplus = a_w^\oplus$ .
- So,  $a_w^\oplus \in R^\oplus$  and  $(a_w^\oplus)^\oplus = (aw)^2 a_w^\oplus$ . □

An element  $p \in R$  is called a Hermitian element if  $p^* = p$ . If in addition,  $p = p^2$ , then  $p$  is called a projection. By  $R^{-1}$  we denote the set of all invertible elements in  $R$ . The classical generalized inverses, including the group inverse, the Moore-Penrose inverse and the core inverse are characterized as follows, respectively. (i)  $a \in R^\#$  if and only if there exists an idempotent  $e \in R$  such that  $ea = ae = 0$  and  $a + e \in R^{-1}$  (see [2, Proposition 8.24]). (ii)  $a \in R^\dagger$  if and only if there exists a projection  $p \in R$  such that  $pa = 0$  and  $aa^* + p \in R^{-1}$  (see [14, Theorem 1.2]). (iii)  $a \in R^\oplus$  if and only if there exists a projection  $q \in R$  such that  $qa = 0$  and  $a^n + q \in R^{-1}$  for any integer  $n \geq 1$  (see [10, Theorems 3.3 and 3.4]). Inspired by these, we aim to the characterization and the representation of the  $w$ -core inverse by projections and units.

**Theorem 2.30.** *Let  $a, w \in R$ . The following conditions are equivalent:*

- (i)  $a \in R_w^\oplus$ .
- (ii) *There exists a unique projection  $p \in R$  such that  $pa = 0$  and  $u = p + aw \in R^{-1}$ .*

(iii) *There exists a projection  $p \in R$  such that  $pa = 0$  and  $u = p + aw \in R^{-1}$ .*

*In this case,  $a_w^\oplus = u^{-1}(1 - p)$ .*

PROOF. (i)  $\Rightarrow$  (ii) Given  $a \in R_w^\oplus$ , set  $p = 1 - awa_w^\oplus$ , then  $p^2 = p = p^*$  and  $pa = (1 - awa_w^\oplus)a = 0$ . Set  $x = a_w^\oplus + 1 - a_w^\oplus aw$ , then  $xu = 1$ . Indeed, we have

$$\begin{aligned} xu &= (a_w^\oplus + 1 - a_w^\oplus aw)(aw + 1 - awa_w^\oplus) \\ &= a_w^\oplus aw + a_w^\oplus(1 - awa_w^\oplus) + (1 - a_w^\oplus aw)aw \\ &\quad + (1 - a_w^\oplus aw)(1 - awa_w^\oplus) \\ &= a_w^\oplus aw + 0 + 0 + 1 - a_w^\oplus aw \\ &= 1. \end{aligned}$$

It is routine to check that  $ux = 1$  and hence  $u = aw + p \in R^{-1}$ .

We next show that such  $p$  is unique. Let  $p, q$  satisfy  $pa = 0 = qa$ ,  $p + aw \in R^{-1}$  and  $q + aw \in R^{-1}$ . As  $(1 - p)(p + aw) = aw$  and  $p + aw \in R^{-1}$ , then  $1 - p = aw(p + aw)^{-1}$  and consequently  $q(1 - p) = qaw(p + aw)^{-1} = 0$ , i.e.,  $q = qp$ .

Similarly, we can prove  $1 - q = aw(q + aw)^{-1}$  and  $p = pq$ . So,  $p = (pq)^* = q^*p^* = qp = q$ .

(ii)  $\Rightarrow$  (iii) is obvious.

(iii)  $\Rightarrow$  (i) Note that  $(1 - p)(p + aw) = aw \Rightarrow 1 - p = aw(p + aw)^{-1}$  and the implication  $(p + aw)a = awa \Rightarrow (p + aw)^{-1}awa = a$ . Then  $x = u^{-1}(1 - p) = (p + aw)^{-1}(1 - p)$  is the  $w$ -core inverse of  $a$ . Indeed, we have

$$(1) \quad xawa = (p + aw)^{-1}(1 - p)awa = (p + aw)^{-1}awa = a.$$

$$(2) \quad awx = aw(p + aw)^{-1}(1 - p) = (1 - p)^2 = 1 - p = (awx)^*.$$

(3) As  $p(p + aw) = p$ , then  $p = p(p + aw)^{-1}$  and hence  $(1 - p)(p + aw)^{-1} = (p + aw)^{-1} - p$ , so that  $awx^2 = (1 - p)(p + aw)^{-1}(1 - p) = ((p + aw)^{-1} - p)(1 - p) = (p + aw)^{-1}(1 - p) = x$ .  $\square$

### 3. Characterizations of $w$ -core inverses and dual $v$ -core inverses by units

In this section, we mainly characterize the existence criteria of both  $w$ -core invertible and dual  $v$ -core invertible elements by units. To make an approach to our main result, we begin with several auxiliary lemmas.

**Lemma 3.1.** *For any  $a, b \in R$ , if  $\alpha = 1 - ab$  is invertible, then so is  $\beta = 1 - ba$ . Moreover,  $\beta^{-1} = 1 + ba^{-1}a$ .*

The formula above in Lemma 3.1 is well known as Jacobson's Lemma. Two elements  $\alpha = 1 - ab$  and  $\beta = 1 - ba$  are called Jacobson pair [8]. More results on Jacobson pairs can be referred to [8, 9].

Recall that an element  $a \in R$  is (von Neumann) regular if there exists an  $x \in R$  such that  $a = axa$ . Such an  $x$  is called an inner inverse or  $\{1\}$ -inverse of  $a$ , and is denoted by  $a^-$ . We herein remark that  $a \in R^{\parallel d}$  implies that  $d$  is regular. As usual, by  $a\{1\}$  we denote the set of all  $\{1\}$ -inverses of  $a$ .

**Lemma 3.2.** [13, Theorem 3.2] *Let  $d \in R$  be regular with  $d^- \in d\{1\}$ . The following conditions are equivalent:*

- (i)  $a$  is invertible along  $d$ .
  - (ii)  $u = da + 1 - dd^- \in R^{-1}$ .
  - (iii)  $v = ad + 1 - d^-d \in R^{-1}$ .
- In this case,  $a^{\parallel d} = u^{-1}d = dv^{-1}$ .*

We here remind the reader that  $u = da + 1 - dd^-$  and  $v = ad + 1 - d^-d$  are Jacobson pairs as  $u$  and  $v$  can be written as  $u = 1 + d(a - d^-)$  and  $v = 1 + (a - d^-)d$ , respectively.

**Lemma 3.3.** [20, Corollary 3.17] *Let  $a \in R$  be regular with  $a^- \in a\{1\}$ . Then the following conditions are equivalent:*

- (i)  $a \in R^\dagger$ .
  - (ii)  $u = aa^* + 1 - aa^- \in R^{-1}$ .
  - (iii)  $v = a^*a + 1 - a^-a \in R^{-1}$ .
- In this case,  $a^\dagger = (u^{-1}a)^* = (av^{-1})^*$ .*

**Theorem 3.4.** *Let  $a, w, v \in R$  and let  $v \in R^{\parallel a}$  with  $a^- \in a\{1\}$ . Then the following conditions are equivalent:*

- (i)  $a \in R_w^{\oplus} \cap R_{v, \oplus}$ .
  - (ii)  $w \in R^{\parallel a}$  and  $a \in R^\dagger$ .
  - (iii)  $u = awavaa^* + 1 - aa^- \in R^{-1}$ .
  - (iv)  $r = avawaa^* + 1 - aa^- \in R^{-1}$ .
  - (v)  $s = wavaa^*a + 1 - a^-a \in R^{-1}$ .
  - (vi)  $t = vawaa^*a + 1 - a^-a \in R^{-1}$ .
- In this case,  $a_w^{\oplus} = avaa^*as^{-1}(u^{-1}awava)^*$  and  $a_{v, \oplus} = (u^{-1}awava)^*(awaa^*at^{-1})$ .*



PROOF. (i)  $\Leftrightarrow$  (ii) follows from Theorems 2.9 and 2.20. (iii)  $\Leftrightarrow$  (v) and (iv)  $\Leftrightarrow$  (vi) by Lemma 3.1. We next just show (ii)  $\Leftrightarrow$  (iii)  $\Leftrightarrow$  (iv).

(ii)  $\Rightarrow$  (iii) Since  $v \in R^{\parallel a}$ , one can get  $av + 1 - aa^- \in R^{-1}$  by Lemma 3.2. Also,  $w \in R^{\parallel a}$  gives  $aw + 1 - aa^- \in R^{-1}$  and hence  $awaa^- + 1 - aa^- \in R^{-1}$  by Lemma 3.1. Thus, we have  $(awaa^- + 1 - aa^-)(av + 1 - aa^-) = awav + 1 - aa^- \in R^{-1}$ . Lemma 3.3 ensures that  $aa^* + 1 - aa^- \in R^{-1}$  as  $a \in R^\dagger$ . Note that  $awav + 1 - aa^-$  and  $awavaa^- + 1 - aa^-$  are Jacobson pairs. Then  $u = awavaa^* + 1 - aa^- = (awavaa^- + 1 - aa^-)(aa^* + 1 - aa^-) \in R^{-1}$ .

(iii)  $\Rightarrow$  (ii) Given (iii), i.e.,  $u = awavaa^* + 1 - aa^- \in R^{-1}$  for some  $a^- \in a\{1\}$ , then we have  $ua = awavaa^*a$  and whence  $a = u^{-1}awavaa^*a \in Raa^*a$ . By Lemma 2.24, we have at once  $a \in R^\dagger$  and  $a^\dagger = (u^{-1}awava)^*$ . This in turn gives  $aa^* + 1 - aa^- \in R^{-1}$  from Lemma 3.3. We get that  $awavaa^- + 1 - aa^- = u(aa^* + 1 - aa^-)^{-1} \in R^{-1}$ , and consequently  $awav + 1 - aa^- \in R^{-1}$ . Note that the assumption  $v \in R^{\parallel a}$  implies  $av + 1 - aa^- \in R^{-1}$ . Then we also can get  $awaa^- + 1 - aa^- = (awav + 1 - aa^-)(av + 1 - aa^-)^{-1} \in R^{-1}$ , then  $aw + 1 - aa^- \in R^{-1}$ , which guarantees that  $w \in R^{\parallel a}$  by Lemma 3.2.

(ii)  $\Leftrightarrow$  (iv) can be proved by a similar way of (ii)  $\Leftrightarrow$  (iii).

We next give the formulae of  $a_w^{\oplus}$  and  $a_{v,\oplus}$ , respectively. Since  $s = wavaa^*a + 1 - a^-a \in R^{-1}$ , we have  $as = awavaa^*a$  and  $a = awavaa^*as^{-1}$ . As  $w^{\parallel a}$  exists, then  $w^{\parallel a} = avaa^*as^{-1}$  by Lemma 2.8. So,  $a_w^{\oplus} = w^{\parallel a}a^{(1,3)} = w^{\parallel a}a^\dagger = avaa^*as^{-1}(u^{-1}awava)^*$ .

Similarly,  $a_{v,\oplus} = a^{(1,4)}v^{\parallel a} = (u^{-1}awava)^*(awaa^*at^{-1})$ .  $\square$

Let us now pause to make some remarks.

**Remark 3.5.** (1) In Theorem 3.4 above, the assumption “ $v \in R^{\parallel a}$ ” cannot be dropped. Such as, let  $R$  and the involution  $*$  be the same as that of Remark 2.27 above. Take  $w = \sum_{i=1}^{\infty} e_{i,i+1}$ ,  $v = \sum_{i=1}^{\infty} e_{i+1,i}$  and  $a = 1$ , then  $v^{\parallel a} = v^{\parallel 1}$  do not exist as  $v \notin R^{-1}$ . By a direct check, the condition (iii)  $awavaa^* + 1 - aa^- = wv = 1 \in R^{-1}$  cannot imply that  $w^{\parallel 1}$  exists (in the item (ii)) as  $w \notin R^{-1}$ .

(2) The equivalences of the items (i), (iii), (iv), (v) and (vi) in Theorem 3.4 also hold without the assumption when  $R$  is a Dedekind-finite ring satisfying the property  $xy = 1 \Rightarrow yx = 1$  for any  $x, y \in R$  (see Theorem 3.7 below).

(3) The formula of  $a_w^{\oplus}$  can also be given by Proposition 2.12. From  $s = wavaa^*a + 1 - a^-a \in R^{-1}$ , it follows that  $was = (wa)^2vaa^*a$  and hence  $wa = (wa)^2vaa^*as^{-1}$ . As  $wa \in R^\#$ , then  $(wa)^\# = wa(vaa^*as^{-1})^2 = (wavaa^*as^{-1})vaa^*as^{-1}$ . So,  $a(wa)^\# = (awavaa^*as^{-1})vaa^*as^{-1} = avaa^*as^{-1}$ . As a consequence,  $a_w^{\oplus} = a(wa)^\#a^{(1,3)} = avaa^*as^{-1}(u^{-1}awava)^*$ .

Combining with Lemma 3.2 and Theorem 3.4, we get the characterization of both  $w$ -core invertible and dual  $v$ -core invertible elements by Mary's inverse along an element in  $R$ .

**Corollary 3.6.** *Let  $a, w, v \in R$  and let  $v \in R^{\parallel a}$ . Then the following conditions are equivalent:*

- (i)  $a \in R_w^{\oplus} \cap R_{v, \oplus}$ .
- (ii)  $w \in R^{\parallel a}$  and  $a^* \in R^{\parallel a}$ .
- (iii)  $wavaa^* \in R^{\parallel a}$ .
- (iv)  $vavaa^* \in R^{\parallel a}$ .

*In this case,  $a_w^{\oplus} = avaa^*ax(yawava)^*$  and  $a_{v, \oplus} = (tavawa)^*awaa^*as$ , where  $x, y \in R$  satisfy  $a = awavaa^*ax = yawavaa^*a$ , and  $s, t \in R$  satisfy  $a = avavaa^*as = tavavaa^*a$ .*

PROOF. As  $wavaa^* \in R^{\parallel a}$ , then  $a \in awavaa^*aR \cap Rawavaa^*a$ . There exist  $x, y \in R$  such that  $a = awavaa^*ax = yawavaa^*a$ . Since  $w \in R^{\parallel a}$ , we have  $w^{\parallel a} = avaa^*ax$ . Recall that if  $a = xa^*a$  then  $x^*$  is a  $\{1, 3\}$ -inverse of  $a$ . From  $a = yawavaa^*a$ , we have  $a^{(1,3)} = (yawava)^*$ . So,  $a_w^{\oplus} = w^{\parallel a}a^{(1,3)} = avaa^*ax(yawava)^*$ .

Similarly, we have  $a_{v, \oplus} = (tavawa)^*awaa^*as$ , where  $s, t \in R$  satisfy  $a = avavaa^*as = tavavaa^*a$ .  $\square$

**Theorem 3.7.** *Let  $R$  be a Dedekind-finite ring and let  $a, w, v \in R$ . Suppose  $a$  is regular and  $a^- \in a\{1\}$ . Then the following conditions are equivalent:*

- (i)  $a \in R_w^{\oplus} \cap R_{v, \oplus}$ .
- (ii)  $wav \in R^{\parallel a}$  and  $a \in R^\dagger$ .
- (iii)  $vaw \in R^{\parallel a}$  and  $a \in R^\dagger$ .
- (iv)  $u = awavaa^* + 1 - aa^- \in R^{-1}$ .
- (v)  $r = avavaa^* + 1 - aa^- \in R^{-1}$ .
- (vi)  $s = wavaa^*a + 1 - a^-a \in R^{-1}$ .
- (vii)  $t = vavaa^*a + 1 - a^-a \in R^{-1}$ .

*In this case,  $a_w^{\oplus} = avaa^*as^{-1}(u^{-1}awava)^*$  and  $a_{v, \oplus} = (u^{-1}awava)^*(awaa^*at^{-1})$ .*

PROOF. (i)  $\Rightarrow$  (ii) Given  $a \in R_w^{\oplus} \cap R_{v, \oplus}$ , it follows from Proposition 2.22 that  $w, v \in R^{\parallel a}$  and  $a \in R^\dagger$ . From Lemma 2.8, we know that  $w, v \in R^{\parallel a}$  imply that  $a \in awaR \cap Rawa$  and  $a \in avaR \cap Rava$ . Hence,  $a \in awavaR \cap Rawava$ , i.e.,  $wav \in R^{\parallel a}$ .

(ii)  $\Rightarrow$  (i) As  $wav \in R^{\parallel a}$ , then  $a \in awavaR \cap Rawava$  by Lemma 2.8. In addition,  $a \in awavaR \subseteq awaR$  shows that  $w$  is right invertible along  $a$  (see

[19, Definition 2.2 and Theorem 2.4]). Note the fact that  $w$  is right invertible along  $a$  implies that  $w$  is left invertible along  $a$  in a Dedekind-finite ring  $R$ . So,  $w \in R^{\parallel a}$ . Similarly,  $v \in R^{\parallel a}$ .

- (i)  $\Leftrightarrow$  (iii) can be proved similarly.
- (ii)  $\Leftrightarrow$  (iv) and (iii)  $\Leftrightarrow$  (v) are similar to the proof of Theorem 3.4.
- (iv)  $\Leftrightarrow$  (vi) and (v)  $\Leftrightarrow$  (vii) follow from Lemma 3.1.

The proof is complete.  $\square$

As a consequence of Proposition 2.22 and Theorem 3.7, the characterization about the product along an element can be obtained.

**Corollary 3.8.** *Let  $R$  be a Dedekind-finite ring and let  $a, w, v \in R$ . Then the following conditions are equivalent:*

- (i)  $w, v \in R^{\parallel a}$ .
- (ii)  $wav \in R^{\parallel a}$ .
- (iii)  $vaw \in R^{\parallel a}$ .

In Theorem 3.4 above, set  $v = w$ , then we can get the following characterization of the  $w$ -core inverse and the dual  $w$ -core inverse by units, whose proof is left to the reader.

**Theorem 3.9.** *Let  $a, w \in R$  and let  $a$  be regular with  $a^- \in a\{1\}$ . Then the following conditions are equivalent:*

- (i)  $a \in R_w^{\oplus} \cap R_{w, \oplus}$ .
- (ii)  $w \in R^{\parallel a}$  and  $a \in R^\dagger$ .
- (iii)  $u = awaa^* + 1 - aa^- \in R^{-1}$ .
- (iv)  $r = a^*awa + 1 - a^-a \in R^{-1}$ .
- (v)  $s = waa^*a + 1 - a^-a \in R^{-1}$ .
- (vi)  $t = aa^*aw + 1 - aa^- \in R^{-1}$ .

*In this case,  $a_w^{\oplus} = t^{-1}aa^*$  and  $a_{w, \oplus} = a^*as^{-1}$ .*

**Corollary 3.10.** [3, Theorem 5.6] *Let  $a \in R$  be regular with  $a^- \in a\{1\}$ . Then the following conditions are equivalent:*

- (i)  $a \in R^{\oplus} \cap R_{\oplus}$ .
- (ii)  $a \in R^\# \cap R^\dagger$ .
- (iii)  $u = a^2a^* + 1 - aa^- \in R^{-1}$ .
- (iv)  $v = a^*a^2 + 1 - a^-a \in R^{-1}$ .
- (v)  $s = aa^*a + 1 - a^-a \in R^{-1}$ .
- (vi)  $t = aa^*a + 1 - aa^- \in R^{-1}$ .

*In this case,  $a^{\oplus} = t^{-1}aa^*$  and  $a_{\oplus} = a^*as^{-1}$ .*

#### 4. Relations with two classes of generalized inverses

In this short section, we aim to show that our defined  $w$ -core inverses and dual  $v$ -core inverses are instances of two classes of outer generalized inverses.

**Theorem 4.1.** *Let  $a, w, v \in R$  and  $a \in R^\dagger$ . Then*

- (i)  $a \in R_w^\oplus$  if and only if  $aw$  is invertible along  $aa^*$ . In this case, the  $w$ -core inverse of  $a$  coincides with the inverse of  $aw$  along  $aa^*$ .
- (ii)  $a \in R_{v,\oplus}$  if and only if  $va$  is invertible along  $a^*a$ . In this case, the dual  $v$ -core inverse of  $a$  coincides with the inverse of  $va$  along  $a^*a$ .

PROOF. (i) Suppose  $a \in R_w^\oplus$  and  $x$  is the  $w$ -core inverse of  $a$ . Then,  $awxa = a$ ,  $xawx = x$ ,  $xawa = a$ ,  $awx^2 = x$  and  $awx = (awx)^*$  by Theorem 2.6. We now show that  $x = a_w^\oplus$  is the inverse of  $aw$  along  $d = aa^*$ . Indeed,

$$(1) \quad xawd = xawaa^* = aa^* = d = (aa^*)^* = (awxaa^*)^* = aa^*(awx)^* = aa^*awx = dawx.$$

$$(2) \quad x = awx^2 = a(a^\dagger a)^*wx^2 = aa^*(a^\dagger)^*wx^2 = d(a^\dagger)^*wx^2 \in dR.$$

$$(3) \quad x = xawx = x(awx)^* = x(wx)^*a^* = x(wx)^*a^\dagger aa^* = x(wx)^*a^\dagger d \in Rd.$$

So,  $x = a_w^\oplus$  is the inverse of  $aw$  along  $aa^*$ .

Conversely, let  $x = (aw)^{\parallel aa^*}$  be the inverse of  $aw$  along  $aa^*$ . Then  $xawaa^* = aa^* = aa^*awx$  and  $x \in aa^*R \cap Raa^*$ . We next show that  $x$  is the  $w$ -core inverse of  $a$ .

As  $a \in R^\dagger$ , then  $awx = (aa^\dagger)^*awx = (a^\dagger)^*a^*awx = (a^\dagger aa^\dagger)^*a^*awx = (a^\dagger)^*a^\dagger aa^*awx = (a^\dagger)^*a^\dagger aa^* = (a^\dagger)^*a^* = aa^\dagger$ . This means  $awx = (awx)^*$  and  $awxa = a$ .

Since  $xawaa^* = aa^*$ , we have  $xawa = a$  by the implication  $a \in R^\dagger \Rightarrow a \in aa^*R$ .

It follows from  $x \in aa^*R$  that  $x = aa^*s$  for some  $s \in R$ , and consequently  $awx^2 = awx(aa^*s) = (awxa)a^*s = aa^*s = x$ .

So,  $x$  is the  $w$ -core inverse of  $a$ .

(ii) can be proved similarly. □

In terms of Proposition 2.17 and Theorem 4.1, we have the following corollary, among them, (i) and (ii) were given in [16, Theorem 4.3].

**Corollary 4.2.** *Let  $a \in R^\dagger$ . Then the following conditions are equivalent:*

- (i)  $a \in R^\oplus$ .
- (ii)  $a$  is invertible along  $aa^*$ .
- (iii)  $a \in R_a^\oplus$ .

- (iv)  $a^2$  is invertible along  $aa^*$ .  
 In this case,  $a^{\oplus} = a^{\parallel aa^*} = aa_a^{\oplus} = a(a^2)^{\parallel aa^*}$ .

In 2012, Drazin [4] defined an outer generalized inverse, called hybrid  $(b, c)$ -inverse, now renamed right hybrid  $(b, c)$ -inverse in [5], which encompasses his  $(b, c)$ -inverses and Mary's inverse along an element. For any  $a, b, c \in R$ , an element  $a$  is called right hybrid  $(b, c)$ -invertible [4] if there exists some  $y \in R$  such that  $yay = y$ ,  $yR = bR$  and  $y^0 = c^0$ . Such an  $y$  is called the right hybrid  $(b, c)$ -inverse of  $a$ . It is unique if it exists. Also, it is shown in [18, Theorem 2.11] that the right hybrid  $(d, d)$ -inverse of  $a$  is indeed Mary's inverse of  $a$  along  $d$ . More results on right hybrid  $(b, c)$ -inverses can be referred to [22].

**Lemma 4.3.** [18, Theorem 2.11] *Let  $a, b, c, y \in R$ . Then the following conditions are equivalent:*

- (i)  $y$  is the right hybrid  $(b, c)$ -inverse of  $a$ .
- (ii)  $yab = b$ ,  $cay = c$ ,  $yR \subseteq bR$  and  $c^0 \subseteq y^0$ .

**Theorem 4.4.** *Let  $a, w, v \in R$ . Then*

- (i)  $a \in R_w^{\oplus}$  if and only if  $aw$  is right hybrid  $(a, a^*)$ -invertible. In this case, the  $w$ -core inverse of  $a$  coincides with the right hybrid  $(a, a^*)$ -inverse of  $aw$ .
- (ii)  $a \in R_{v, \oplus}$  if and only if  $va$  is right hybrid  $(a^*, a)$ -invertible. In this case, the dual  $v$ -core inverse of  $a$  coincides with the right hybrid  $(a^*, a)$ -inverse of  $va$ .

**PROOF.** We only prove the statement (i) as the statement (ii) can be proved similarly.

Suppose that  $a \in R_w^{\oplus}$  and  $x$  is the  $w$ -core inverse of  $a$ . Then, by Theorem 2.6, we have  $xawx = x$ ,  $xR = aR$  and  $Rx = Ra^*$ . Clearly,  $Rx = Ra^*$  gives  $x^0 = (a^*)^0$ . So,  $x$  is the right hybrid  $(a, a^*)$ -inverse of  $aw$ .

Conversely, suppose that  $aw$  is right hybrid  $(a, a^*)$ -invertible and  $x$  is the right hybrid  $(a, a^*)$ -inverse of  $aw$ . Then, by Lemma 4.3, we have  $xawa = a$ ,  $a^*awx = a^*$ ,  $xR \subseteq aR$  and  $(a^*)^0 \subseteq x^0$ .

Pre-multiplying  $a^*awx = a^*$  by  $(wx)^*$  gives  $(awx)^*awx = (wx)^*a^*awx = (wx)^*a^* = (awx)^*$  and  $awx = (awx)^*$ .

The equation  $a^*awx = a^*$  implies  $a^* = a^*(awx)^* = (awxa)^*$ , i.e.,  $a = awxa$ . As  $xR \subseteq aR$ , then  $x = at$  for some  $t \in R$ . Post-multiplying  $a = awxa$  by  $t$  implies  $awx^2 = x$ .

Therefore,  $x$  is the  $w$ -core inverse of  $a$ . □

Given any  $a, b, c \in R$ , an element  $a$  is  $(b, c)$ -invertible [4] if there some  $y \in R$  such that  $yab = b$ ,  $cay = c$  and  $y \in bRy \cap yRc$ , or equivalently,  $yay = y$ ,  $yR = bR$  and  $Ry = Rc$ . Such an  $y$  is called a  $(b, c)$ -inverse of  $a$ .

By a similar proof of Theorem 4.4 above, the relations about  $w$ -core inverses and  $(b, c)$ -inverses can also be obtained as follows.

**Theorem 4.5.** *Let  $a, w, v \in R$ . Then*

(i)  $a \in R_w^{\oplus}$  if and only if  $aw$  is  $(a, a^*)$ -invertible. In this case, the  $w$ -core inverse of  $a$  coincides with the  $(a, a^*)$ -inverse of  $aw$ .

(ii)  $a \in R_{v, \oplus}$  if and only if  $va$  is  $(a^*, a)$ -invertible. In this case, the dual  $v$ -core inverse of  $a$  coincides with the  $(a^*, a)$ -inverse of  $va$ .

Combining with Theorems 4.4 and 4.5, we have the following result.

**Corollary 4.6.** *Let  $a, w, v \in R$ . Then*

(i)  $aw$  is  $(a, a^*)$ -invertible if and only if it is right hybrid  $(a, a^*)$ -invertible. In this case, the  $(a, a^*)$ -inverse of  $aw$  coincides with its right hybrid  $(a, a^*)$ -inverse.

(ii)  $va$  is  $(a^*, a)$ -invertible if and only if it is right hybrid  $(a^*, a)$ -invertible. In this case, the  $(a^*, a)$ -inverse of  $va$  coincides with its right hybrid  $(a^*, a)$ -inverse.

## ACKNOWLEDGMENTS

This research is supported by the National Natural Science Foundation of China (No. 11771076, No. 11801124) and China Postdoctoral Science Foundation (No. 2020M671068).

## References

## References

- [1] O.M. Baksalary, G. Trenkler, Core inverse of matrices, Linear Multilinear Algebra 58 (2010) 681-697.
- [2] K.P.S. Bhaskara Rao, Theory of Generalized Inverses over Commutative Rings, London and New York: Taylor and Francis; 2002.

- [3] J.L. Chen, H.H. Zhu, P. Patrício, Y.L. Zhang, Characterizations and representations of core and dual core inverses, *Canad. Math. Bull.* 60 (2017) 269-282.
- [4] M.P. Drazin, A class of outer generalized inverses, *Linear Algebra Appl.* 436 (2012) 1909-1923.
- [5] M.P. Drazin, Hybrid  $(b, c)$ -inverses and three finiteness properties in rings, semigroups and categories, *Comm. Algebra* 49 (2021) 2265-2277.
- [6] M.P. Drazin, Pseudo-inverses in associative rings and semigroups, *Amer. Math. Monthly* 65 (1958) 506-514.
- [7] Y.F. Gao, J.L. Chen, Pseudo core inverses in rings with involution, *Comm. Algebra* 64 (2018) 38-50.
- [8] T.Y. Lam, Pace P. Nielsen, Inner inverses and inner annihilators in rings, *J. Algebra* 397 (2014) 91-110.
- [9] T.Y. Lam, Pace P. Nielsen, Jacobson's lemma for Drazin inverses, *Contemp. Math.* 609 (2014) 185-195.
- [10] T.T. Li, J.L. Chen, Characterizations of core and dual core inverses in rings with involution, *Linear Multilinear Algebra* 66 (2018) 717-730.
- [11] K. Manjunatha Prasad, K.S. Mohana, Core-EP inverse, *Linear Multilinear Algebra* 62 (2014) 792-802.
- [12] X. Mary, On generalized inverses and Green's relations, *Linear Algebra Appl.* 434 (2011) 1836-1844.
- [13] X. Mary, P. Patrício, Generalized inverses modulo  $\mathcal{H}$  in semigroups and rings, *Linear Multilinear Algebra* 61 (2013) 886-891.
- [14] P. Patrício, C.M. Araújo, Moore-Penrose invertibility in involutory rings:  $aa^\dagger = bb^\dagger$ , *Linear Multilinear Algebra* 58 (2010) 445-452.
- [15] R. Penrose, A generalized inverse for matrices, *Proc. Camb. Phil. Soc.* 51 (1955) 406-413.
- [16] D.S. Rakić, N.C. Dinčić, D.S. Djordjević, Group, Moore-Penrose, core and dual core inverse in rings with involution, *Linear Algebra Appl.* 463 (2014) 115-133.

- [17] S.Z. Xu, J.L. Chen, X.X. Zhang, New characterizations for core inverses in rings with involution, *Front. Math. China* 12 (2017) 231-246.
- [18] H.H. Zhu, Further results on several generalized inverses, *Comm. Algebra* 46 (2018) 3388-3396.
- [19] H.H. Zhu, J.L. Chen, P. Patrício, Further results on the inverse along an element in semigroups and rings, *Linear Multilinear Algebra* 64 (2016) 393-403.
- [20] H.H. Zhu, J.L. Chen, P. Patrício, X. Mary, Centralizer's applications to the inverse along an element, *Appl. Math. Comput.* 315 (2017) 27-33.
- [21] H.H. Zhu, Q.W. Wang, Weighted pseudo core inverses in rings, *Linear Multilinear Algebra* 68 (2020) 2434-2447.
- [22] H.H. Zhu, L.Y. Wu, F. Peng, P. Patrício, Characterizations and representations of left and right hybrid  $(b, c)$ - inverses in rings, *RACSAM* 115 (2021) 8 pages.