

QUADRATURE FORMULAS FOR BESSEL POLYNOMIALS

HIDEKI MATSUMURA*

CONTENTS

1. Introduction	1
2. Quadrature formulas for Bessel polynomials	4
3. Proof of Theorem 1.3	6
3.1. Proof of Theorem 1.3 (1), (2)	6
3.2. Proof of Theorem 1.3 (3)	7
4. Proof of Theorem 1.4	10
5. Proof of Theorem 1.5	12
References	13

ABSTRACT. A quadrature formula is a formula computing a definite integration by evaluation at finite points. Existence of certain quadrature formulas for orthogonal polynomials is related to interesting problems such as Waring's problem in number theory and spherical designs in algebraic combinatorics. Sawa and Uchida proved the existence and the non-existence of certain rational quadrature formulas for the weight functions of certain classical orthogonal polynomials. Classical orthogonal polynomials belong to the Askey-scheme, which is a hierarchy of hypergeometric orthogonal polynomials. Thus, it is natural to extend the work of Sawa and Uchida to other polynomials in the Askey-scheme. In this article, we extend the work of Sawa and Uchida to the weight function of the Bessel polynomials. In the proofs, we use the Riesz-Shohat theorem and Newton polygons. It is also of number theoretic interest that proofs of some results are reduced to determining the sets of rational points on elliptic curves.

1. INTRODUCTION

A quadrature formula of degree n for a weight function $w(t)$ is an integration formula of the form

$$\sum_{i=1}^m x_i f(y_i) = \int_{\gamma} f(t)w(t)dt,$$

where γ is a path in \mathbb{R} or \mathbb{C} and $f(x)$ is any polynomial such that $\deg(f) \leq n$. The points y_i 's are called the nodes of the quadrature formula. For example, for a real interval $\gamma = [a, b]$, $w(t) = 1$ and $n = 1$, the trapezoidal rule states that

$$\frac{b-a}{2}(f(a) + f(b)) = \int_a^b f(t)dt$$

Date: June 17, 2022.

2010 Mathematics Subject Classification. primary 33C45; secondary 65D32; tertiary 14G05.

Key words and phrases. quadrature formula, Bessel polynomial, Riesz-Shohat theorem, Newton polygon, Christoffel-Darboux kernel, elliptic curve, rational points.

This research is supported by KAKENHI 18H05233.

is a quadrature formula of degree 1 for $w(t) = 1$ with 2 nodes. For $\gamma = [a, b]$, $w(t) = 1$ and $n = 3$, the Simpson rule states that

$$\frac{b-a}{6} \left(f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right) = \int_a^b f(t) dt$$

is a quadrature formula of degree 3 for $w(t) = 1$ with 3 nodes.

A quadrature formula is called rational if all nodes are rational numbers. Existence of certain rational quadrature formulas for orthogonal polynomials is related to interesting problems such as Waring's problem in number theory (cf. [5, 8]) and spherical designs in algebraic combinatorics (cf. [1, 4]). See [5, 10] for applications.

Quadrature formulas are determined by the weight function $w(t)$, the degree n and the number m of nodes. The following problem is fundamental:

Problem 1.1. *For a fixed weight function $w(t)$ and a natural number m , determine the range of n such that there exists a rational quadrature formula of degree n for $w(t)$ with m nodes.*

By the Stroud-type bound (cf. [9, Proposition 4.3], [10, Proposition 2.7]), if there exists a quadrature formula of degree n for $w(t)$ with m nodes, then $n \leq 2m - 1$. The case $n = 2m - 1$ is called the tight case. Sawa and Uchida [9, 10] considered the "almost tight case" and proved that there exists no quadrature formula of degree $2r$ for certain classical orthogonal polynomials (Hermite polynomials $H_n(t)$, Legendre polynomials $P_n(t)$ and Laguerre polynomials $L_n^{(0)}(t)$) with $r + 1$ nodes on \mathbb{Q} ([10, Theorem 1.4]). They also proved that there exist quadrature formulas of degree $r + 1$ for certain classical orthogonal polynomials with $r + 1$ nodes on \mathbb{Q} ([10, Theorem 5.5]). They took algebro-geometric approaches involving the Riesz-Shohat theorem, Christoffel-Darboux kernels and Newton polygons.

Classical orthogonal polynomials belong to the Askey-scheme [7], which is a hierarchy of hypergeometric orthogonal polynomials. Thus, it is natural to extend the work of [9, 10] to more general polynomials in the Askey-scheme. In this note, we extend the work of Sawa and Uchida [9, 10] for the weight function of Bessel polynomials [6, p. 101]

$$y_n(x) := \sum_{k=0}^n \frac{(n+k)!}{(n-k)!k!} \left(\frac{x}{2}\right)^k.$$

The Bessel polynomials belong to the Askey-scheme and are orthogonal with respect to the weight function

$$w(z) = -\frac{1}{4\pi\sqrt{-1}} e^{-\frac{2}{z}},$$

where the path of integration is $S^1 \subset \mathbb{C}$, i.e.,

$$\int_{S^1} y_m(z) y_n(z) w(z) dz = 0$$

for $m \neq n$ (cf. [6, p. 104]).

Orthogonal polynomials are defined from certain inner products expressed by integrations. The inner products for the classical orthogonal polynomials in [9, 10] are expressed by real integrations. On the other hand, in our case, note that the inner product is expressed by complex integrations. As a generalization of "quadrature formulas with nodes on \mathbb{Q} " ([9, 10]), we consider "quadrature formulas with the real part and the imaginary parts of nodes on \mathbb{Q} ." Since the path of integration is S^1 , we consider the following problems:

Problem 1.2. (1) *Does there exist a quadrature formula of degree $2r$ for Bessel polynomials with $r + 1$ nodes on $\mathbb{Q}(\sqrt{-1}) \cap S^1$?*

- (2) *Does there exist a quadrature formula of degree $r + 1$ for Bessel polynomials with $r + 1$ nodes on $\mathbb{Q}(\sqrt{-1}) \cap S^1$?*

As far as the author knows, there are few works considering quadrature formulas whose nodes lie in a quadratic extension of \mathbb{Q} . Note that [9, 10] do not consider quadrature formulas with nodes on S^1 . We give negative answers to Problem 1.2 (1) for all $r \in \mathbb{Z}_{\geq 1}$ and Problem 1.2 (2) for $r = 2$.

- Theorem 1.3.** (1) *There exist quadrature formulas of degree 2 for Bessel polynomials with 2 nodes on $\mathbb{Q}(\sqrt{-1}) \cap S^1$.*
 (2) *There exists no quadrature formula of degree 4 for Bessel polynomials with 3 nodes on $\mathbb{Q}(\sqrt{-1}) \cap S^1$.*
 (3) *There exists no quadrature formula of degree $2r$ for Bessel polynomials with $r + 1$ nodes on $\mathbb{Q}(\sqrt{-1})$ for all $r \in \mathbb{Z}_{\geq 3}$.*

The proof of (1) is straightforward. (2) follows from Theorem 1.4. In the proof of (3), we use the Riesz–Shohat theorem (Theorem 2.2) and a fact on Newton polygons (Lemma 2.4). An advantage of Bessel polynomials is that the computation (or estimate) of p -adic valuation is relatively easy. On the other hand, to prove (2), we cannot use Lemma 2.4 since we can construct infinitely many quadrature formulas of degree 4 for Bessel polynomials with 3 nodes on $\mathbb{Q}(\sqrt{-1})$ by using the Christoffel–Darboux kernel (cf. Example 3.2). By contrast, there exists no quadrature formula of degree 4 for Hermite polynomials with 3 nodes on \mathbb{Q} by [10, Theorem 1.4].

Theorem 1.4. *There exists no quadrature formula of degree 3 for Bessel polynomials with 3 nodes on $\mathbb{Q}(\sqrt{-1}) \cap S^1$.*

Note that it is an immediate consequence of the Riesz–Shohat theorem that there exists a quadrature formula of degree r for Bessel polynomials with $r + 1$ nodes on $\mathbb{Q}(\sqrt{-1}) \cap S^1$ (Remark 2.3). Thus, increasing the degree is a problem. In the proof of Theorem 1.4, we use the Riesz–Shohat theorem (Theorem 2.2) to reduce the problem to determining the set of rational points on an elliptic curve. We do not know whether we can take the nodes y_1, \dots, y_{r+1} on $\mathbb{Q}(\sqrt{-1}) \cap S^1$ for $r \geq 3$. On the other hand, by the Riesz–Shohat theorem (Theorem 2.2), we can prove that there exist quadrature formulas of degree $r + 1$ with $r + 1$ nodes on \mathbb{Q} for all $r \in \mathbb{Z}_{\geq 0}$.

Theorem 1.5. *There exist quadrature formulas of degree $r + 1$ for Bessel polynomials with $r + 1$ nodes on \mathbb{Q} for all $r \in \mathbb{Z}_{\geq 0}$.*

This article is organized as follows:

In §2, we formulate a quadrature formula on \mathbb{C} and introduce three key tools: the Riesz–Shohat theorem (Theorem 2.2), Newton polygons and Christoffel–Darboux kernels. In §3, we prove Theorem 1.3 by using the Riesz–Shohat theorem and Newton polygons. We also construct an example of quadrature formulas of degree 4 for Bessel polynomials with 3 nodes on $\mathbb{Q}(\sqrt{-1})$ by using the Christoffel–Darboux kernel (Example 3.2). In §4, we prove Theorem 1.4 by using the Riesz–Shohat theorem. It is also of number theoretic interest that this proof is reduced to determining the set of rational points on an elliptic curve. Then, in §5, we prove Theorem 1.5 by using the Riesz–Shohat theorem.

2. QUADRATURE FORMULAS FOR BESSEL POLYNOMIALS

We consider the “quadrature formulas” for Bessel polynomials. Since

$$\frac{1}{2^k} \frac{(n+k)!}{(n-k)!k!} = \frac{1}{2^k} \frac{(n+k)!}{(n-k)!(2k)!} \frac{(2k)!}{k!} = \binom{n+k}{2k} (2k-1)!!,$$

note that $y_n(x) \in \mathbb{Z}[x]$.

Unlike the classical orthogonal polynomials in [10], the path of integration is $S^1 \subset \mathbb{C}$. We consider “quadrature formulas” in the following sense:

Definition 2.1. Let $\gamma \subset \mathbb{C}$ be a path.¹ An integration formula of the form

$$(1) \quad \sum_{i=1}^m x_i f(z_i) = \int_{\gamma} f(z) w(z) dz$$

with $x_1, \dots, x_m, z_1, \dots, z_m \in \mathbb{C}$ is called a quadrature formula of degree n for a weight function $w(z)$ with nodes z_i and weights x_i if it holds for all $f(z) \in \mathbb{C}[z]$ such that $\deg(f) \leq n$. A quadrature formula on a subset $S \subset \mathbb{C}$ is a quadrature formula such that $z_i \in S$ for all i .

For Bessel polynomials, we take

$$w(z) = -\frac{1}{4\pi\sqrt{-1}} e^{-\frac{2}{z}}.$$

so that

$$\int_{S^1} w(z) dz = 1.$$

We introduce key tools: the Riesz–Shohat theorem [11] and a lemma on Newton polygons. They are key tools in the proof of [10, Theorem 1.4, Theorem 5.5], and also work over \mathbb{C} .

In the proof of Theorems 1.3 to 1.5, we use the Riesz–Shohat theorem.

Theorem 2.2 (Cf. Riesz–Shohat theorem, [10, Proposition 2.4], [11, Theorem I]). *Let $\gamma \subset \mathbb{C}$ be a path. Suppose that $1 \leq k \leq r+2$, $z_1, \dots, z_{r+1} \in \mathbb{C}$ are distinct and let*

$$\omega_{r+1}(z) := \prod_{i=1}^{r+1} (z - z_i).$$

Then, the following are equivalent:

- (1) *There exist $x_1, \dots, x_{r+1} \in \mathbb{C}$ such that*

$$\sum_{i=1}^{r+1} x_i f(z_i) = \int_{\gamma} f(z) w(z) dz$$

is a quadrature formula of degree $2(r+1) - k$.

- (2) *ω_{r+1} is a quasi-orthogonal polynomial of degree $r+1$ and order $k-1$, i.e., there exist $b_1, \dots, b_{k-1} \in \mathbb{C}$ such that*

$$\omega_{r+1}(x) = \phi_{r+1}(x) + b_1 \phi_r(x) + \dots + b_{k-1} \phi_{r+2-k}(x).$$

Here, $\{\phi_l\}$ is the system of monic orthogonal polynomial with respect to $w(z)$.

Furthermore, if the above equivalent conditions hold, then we have

$$x_i = \int_{\gamma} \frac{\omega_{r+1}(z)}{(z - z_i) \omega'_{r+1}(z_i)} w(z) dz.$$

¹Later, we consider $\gamma = S^1$. For the formulation over \mathbb{R} , see [10].

Remark 2.3. Let $\{\Phi_n\}$ be a system of orthogonal polynomials with respect to a weight function $w(z)$. Then, there exists a quadrature formula of degree r for $\{\Phi_n\}$ with $r + 1$ nodes on any subset $A \subset \mathbb{C}$ with at least $r + 1$ elements. Indeed, this is an immediate consequence of Theorem 2.2. Note that this is the case $2r + 2 - k = r$. i.e., $k = r + 2$ in Theorem 2.2. Let $\{\phi_n\}$ be the system of monic orthogonal polynomials with respect to $w(z)$. Then, since $\{\phi_0, \dots, \phi_{r+1}\}$ is a basis of the space of polynomials of degree $\leq r + 1$ and $\omega_{r+1}(x)$ is monic, there exist $b_1, \dots, b_{r+1} \in \mathbb{C}$ such that

$$\omega_{r+1}(x) = \phi_{r+1}(x) + b_1\phi_r(x) + \dots + b_{r+1}\phi_0(x).$$

Therefore, by Theorem 2.2, for all $z_1, \dots, z_{r+1} \in A$, there exist $x_1, \dots, x_{r+1} \in \mathbb{C}$ such that

$$\sum_{i=1}^{r+1} x_i f(z_i) = \int_{\gamma} f(z) w(z) dz$$

is a quadrature formula of degree r . In particular, there exists a quadrature formula of degree r for Bessel polynomials with $r + 1$ nodes on $\mathbb{Q}(\sqrt{-1}) \cap S^1$.²

In the proof of Theorem 1.3 (3), we use a fact on Newton polygons.

Lemma 2.4 (Cf. [10, Lemma 4.2]). *Let K be a discrete valuation field and $f(x) \in \mathcal{O}_K[x]$. If all zeros of $f(x)$ are K -rational, then all edges of the Newton polygon of $f(x)$ have integral slopes.*

We also use the following version of Bertrand's postulate:

Theorem 2.5 ([3, p. 505]). *For $n \geq 4$, there exists a prime number p such that $n < p \leq 2n$ and $p \equiv 3 \pmod{4}$.*

As in the proof of [10, Lemma 4.5], we obtain the following lemma:

Lemma 2.6. *For all $l \in \mathbb{Z}_{\geq 7}$, there exists a prime number p such that $(l + 1)/2 < p \leq l$ and $p \equiv 3 \pmod{4}$.*

In Example 3.2, we use the Christoffel–Darboux kernel.

Definition 2.7 ([10, p. 1243]). Let $\{\Phi_l\}$ be a sequence of orthogonal polynomials with respect to a weight function $w(z)$ over $\gamma \subset \mathbb{C}$. The Christoffel–Darboux kernel for polynomials of degree at most l is defined by

$$K_l(x, y) := \sum_{k=0}^l h_k h_k^{-1} \Phi_k(x) \Phi_k(y).$$

Here, $h_k := \int_{\gamma} \Phi_k(z)^2 dz$.

The Christoffel–Darboux kernel is computed by the following formula:

Proposition 2.8 (Christoffel–Darboux formula (cf. [10, Proposition 2.1], [12, Theorem 3.2.2])). *For $l \in \mathbb{Z}_{\geq 0}$, let k_l be the leading coefficient of $\Phi_l(x)$. Then,*

$$K_l(x, y) = \frac{k_l}{k_{l+1}} \frac{\Phi_{l+1}(x)\Phi_l(y) - \Phi_l(x)\Phi_{l+1}(y)}{x - y}.$$

Definition 2.9.

$$f_r(x, y) := \frac{\Phi_{l+1}(x)\Phi_l(y) - \Phi_l(x)\Phi_{l+1}(y)}{x - y}.$$

²We can take arbitrary $r + 1$ points on $\mathbb{Q}(\sqrt{-1}) \cap S^1$ as nodes.

By Theorem 2.2 and Proposition 2.8,

Lemma 2.10 (Cf. [10, Lemma 2.5]). *Suppose that $x_1, \dots, x_{r+1}, z_1, \dots, z_{r+1} \in \mathbb{C}$ satisfy eq. (1) for $(m, n) = (r+1, 2r)$. Assume that z_1, \dots, z_{r+1} are distinct. Then $f_r(z_i, z_j) = 0$ for every distinct i, j .*

3. PROOF OF THEOREM 1.3

In this section, we prove Theorem 1.3. We also construct an example of quadrature formulas of degree 4 for Bessel polynomials with 3 nodes on $\mathbb{Q}(\sqrt{-1})$.

3.1. Proof of Theorem 1.3 (1), (2). In this subsection, we prove Theorem 1.3 (1) and (2).

Proof of Theorem 1.3 (1), (2). (1) Suppose that there exist $x_1, x_2 \in \mathbb{C}, z_1, z_2 \in \mathbb{Q}(\sqrt{-1})$ such that $|z_1| = |z_2| = 1$ and

$$\begin{aligned} x_1 + x_2 &= 1, \\ x_1 z_1 + x_2 z_2 &= -1, \\ x_1 z_1^2 + x_2 z_2^2 &= \frac{2}{3}. \end{aligned}$$

By the first and the second equality, we obtain

$$(z_1 - z_2)x_1 = -z_2 - 1.$$

If $z_1 = z_2$, then

$$z_1 = z_2 = -\frac{1}{2}$$

by the first and the second equality. Thus,

$$x_1 + x_2 = \frac{8}{3},$$

which contradicts the first equality. Since $z_1 \neq z_2$,

$$x_1 = \frac{-z_2 - 1}{z_1 - z_2}.$$

We substitute it and the first equality into the third equality to obtain

$$-z_1 - z_1 z_2 - z_2 = \frac{2}{3}.$$

If $z_1 = -1$, then

$$1 + z_2 - z_2 = \frac{2}{3},$$

which is a contradiction. Thus, $z_1 \neq -1$. Therefore,

$$z_2 = \frac{-z_1 - \frac{2}{3}}{z_1 + 1}.$$

Since $|z_1| = |z_2| = 1$,

$$\operatorname{Re}(z_1) = -\frac{5}{6}.$$

Thus, we have

$$\operatorname{Im}(z_1) = \pm \frac{\sqrt{11}}{6} \notin \mathbb{Q},$$

which is a contradiction.

(2) This is an immediate consequence of Theorem 1.4. □

Remark 3.1. There exists a unique quadrature formula of degree 2 for Bessel polynomials with 2 nodes on $S^1 \subset \mathbb{C}$. Indeed, if there exists such a quadrature formula on S^1 , we have

$$z_1 = \frac{-5 + \sqrt{-11}}{6},$$

$$z_2 = \frac{-5 - \sqrt{-11}}{6}$$

by the proof of Theorem 1.3 (1). Then, by Theorem 2.2 and the residue theorem,

$$x_1 = \int_{S^1} \frac{z - z_2}{z_2 - z_1} w(z) dz = \frac{11 + \sqrt{-11}}{22},$$

$$x_2 = \int_{S^1} \frac{z - z_1}{z_1 - z_2} w(z) dz = \frac{11 - \sqrt{-11}}{22}.$$

Since

$$-\frac{1}{4\pi\sqrt{-1}} \int_{S^1} z^j e^{-\frac{2}{z}} dz = \frac{(-2)^j}{(j+1)!}$$

for $j \in \mathbb{Z}_{\geq 0}$ by the residue theorem, we can check that

$$x_1 f(z_1) + x_2 f(z_2) = -\frac{1}{4\pi\sqrt{-1}} \int_{S^1} f(z) e^{-\frac{2}{z}} dz$$

is a quadrature formula of degree 2 for Bessel polynomials.

3.2. Proof of Theorem 1.3 (3). In this subsection, we prove Theorem 1.3 (3). We also construct an example of quadrature formulas of degree 4 for Bessel polynomials with 3 nodes on $\mathbb{Q}(\sqrt{-1})$.

Proof of Theorem 1.3 (3). Assume that there exist $x_1, \dots, x_{r+1} \in \mathbb{C}, z_1, \dots, z_{r+1} \in \mathbb{Q}(\sqrt{-1})$ such that

$$\sum_{i=1}^{r+1} x_i z_i^j = -\frac{1}{4\pi\sqrt{-1}} \int_{S^1} z^j e^{-\frac{2}{z}} dz \quad (j = 0, 1, \dots, 2r).$$

By Theorem 2.2, there exists $c \in \mathbb{Q}(\sqrt{-1})$ such that the zeros of the quasi-Bessel polynomial $y_{r+1;c}(x)$ are z_1, \dots, z_{r+1} . Write $c = s/t$, with $s, t \in \mathbb{Z}[\sqrt{-1}]$ and $\gcd(s, t) = 1$. Let

$$f(x) := t y_{r+1;c}(x) = t y_{r+1}(x) + s y_r(x) \in \mathbb{Z}[x].$$

Write

$$f(x) = \sum_{k=0}^{r+1} a_k x^{r+1-k}.$$

We have

$$(2) \quad a_k = \begin{cases} t(2r+1)!! & (k=0), \\ (t(2r+1)+s)(2r-1)!! & (k=1), \\ \left(t \binom{2(r+1)-k}{2(r+1-k)} + s \binom{2r+1-k}{2(r+1-k)} \right) (2(r+1-k)-1)!! & (2 \leq k \leq r+1). \end{cases}$$

First, assume that $r \geq 7$, $r = 3$ or 4 . For $r \geq 7$, by Lemma 2.6, we can take a prime number p such that $(r+1)/2 < p \leq r$ and $p \equiv 3 \pmod{4}$ (i.e., p is inert in $\mathbb{Q}(\sqrt{-1})$). For $r = 3, 4$, let $p = 3$. We consider the Newton polygon of $f(x)$ with respect to p . Let $P_k := (k, v_p(a_k)) \in \mathbb{R}^2$.

Let

$$j := \begin{cases} r & (t + s \equiv 0 \pmod{p}), \\ r + 1 & (t + s \not\equiv 0 \pmod{p}). \end{cases}$$

Then, $v_p(a_j) = 0$ by eq. (2). Indeed, if $t + s \equiv 0 \pmod{p}$, then $v_p(s) = v_p(t) = 0$ by $\gcd(s, t) = 1$. Since $p < r + 1 < 2p$ and p is odd,

$$v_p(a_r) = v_p\left(\frac{(r+1)((t+s)r+2t)}{2}\right) = 0.$$

If $t + s \not\equiv 0 \pmod{p}$, then $v_p(a_{r+1}) = v_p(t + s) = 0$.

Therefore, there exists $j_0 := \min\{j \mid v_p(a_j) = 0\}$.

If $v_p(t) = 0$, then

$$v_p(a_0) = v_p((2r+1)!) = \begin{cases} 1 & (2r+1 < 3p), \\ 2 & (2r+1 \geq 3p, p > 3), \\ 3 & (2r+1 \geq 3p, p = 3). \end{cases}$$

by eq. (2) and $p < 2r + 1 < 4p$. If $v_p(t) \geq 1$, then $v_p(s) = 0$ by $\gcd(s, t) = 1$. Thus,

$$v_p(a_1) = v_p((2r-1)!) = \begin{cases} 1 & (2r-1 < 3p),^3 \\ 2 & (2r-1 \geq 3p). \end{cases}$$

by eq. (2) and $p < 2r - 1 < 4p$.

(1) Suppose that $v_p(t) = 0$.

(a) If $2r + 1 < 3p$,⁴ then $v_p(a_0) = 1$. Let $j_0 := \min\{j \mid v_p(a_j) = 0\}$. In this case, the Newton polygon of $f(x)$ has the edge $P_0P_{j_0}$, whose slope is $-1/j_0 \notin \mathbb{Z}$ since $j_0 > r - (p-1)/2 \geq p - (p-1)/2 = (p+1)/2 \geq 2$ by $p \geq 3$. For the first inequality, note that if p is an odd prime and $p \leq 2(r+1-k) - 1$, i.e., $k \leq r - (p-1)/2$, then $v_p(a_k) \geq 1$.

(b) If $r \geq 7$ and $2r + 1 \geq 3p$, then $v_p(a_0) = 2$ and $v_p(a_1) \geq 1$.

(i) If $v_p(a_1) = 1$, then the Newton polygon of $f(x)$ has the edge $P_1P_{j_0}$, whose slope is $-1/(j_0 - 1) \notin \mathbb{Z}$ since $j_0 > 2$.

(ii) If $v_p(a_1) \geq 2$ and $v_p(a_i) \neq 1$ for all i , then the Newton polygon of $f(x)$ has the edge $P_0P_{j_0}$, whose slope is $-2/j_0 \notin \mathbb{Z}$ since $j_0 > 2$.

(iii) If $v_p(a_1) \geq 2$ and $v_p(a_i) = 1$ for some $i \geq 2$, then let $i_0 := \min\{i \mid v_p(a_i) = 1\}$. The Newton polygon of $f(x)$ has the edge $P_0P_{i_0}$, whose slope is $-1/i_0 \notin \mathbb{Z}$ since $i_0 > 1$.

(c) If $r = 4$, then

$$\begin{aligned} a_0 &= 945t, \\ a_1 &= 3(315t + 35s), \\ a_2 &= 3(140t + 35s), \\ a_3 &= 3(35t + 15s), \\ a_4 &= 15t + 10s, \\ a_5 &= t + s. \end{aligned}$$

³If $p = 3$, then $r < 5$ by $(r+1)/2 < p$. Thus, $2r - 1 < 9 = 3p$.

⁴This includes the case $r = 3$.

- (i) If $v_3(s) > 0$, then $v_3(a_0) = 3$, $v_3(a_1) \geq 2$, $v_3(a_2) = 1$, $v_3(a_3), v_3(a_4) \geq 1$ and $v_3(a_5) = 0$. Thus, the Newton polygon of $f(x)$ has the edge P_2P_5 , whose slope is $-1/3 \notin \mathbb{Z}$.
 - (ii) If $v_3(s) = 0$, then $v_3(a_0) = 3$, $v_3(a_1) = 1$, $v_3(a_2), v_3(a_3) \geq 1$ and $v_3(a_4) = 0$. Thus, the Newton polygon of $f(x)$ has the edge P_1P_4 , whose slope is $-1/3 \notin \mathbb{Z}$.
- (2) Suppose that $v_p(t) > 0$. Since $\gcd(s, t) = 1$, $v_p(s) = v_p(t+s) = 0$. We have $v_p(a_0) \geq 2$.
- (a) If $2r - 1 < 3p$, then $v_p(a_1) = 1$ and the Newton polygon of $f(x)$ has the edge $P_1P_{j_0}$, whose slope is $-1/(j_0 - 1) \notin \mathbb{Z}$ since $j_0 > 2$.
 - (b) If $2r - 1 \geq 3p$, then $v_p(a_0) \geq 3$ and $v_p(a_1) = 2$. We have $v_p(a_2) \geq 1$ by $j_0 > (p+1)/2 \geq 3$.
 - (i) If $v_p(a_2) = 1$, then the Newton polygon of $f(x)$ has the edge $P_2P_{j_0}$, whose slope is $-1/(j_0 - 2) \notin \mathbb{Z}$ since $j_0 > r - (p-1)/2 \geq (3p+1)/2 - (p-1)/2 \geq p+1 \geq 4$ by $2r - 1 \geq 3p$. The first inequality follows from the argument in Case (1-a).
 - (ii) If $v_p(a_2) \geq 2$ and $v_p(a_i) \neq 1$ for all i , then the Newton polygon of $f(x)$ has the edge $P_1P_{j_0}$, whose slope is $-2/(j_0 - 1) \notin \mathbb{Z}$ since $j_0 > 4$.
 - (iii) If $v_p(a_2) \geq 2$ and $v_p(a_i) = 1$ for some $i \geq 3$, then the Newton polygon of $f(x)$ has the edge $P_1P_{i_0}$, whose slope is $-1/(i_0 - 1) \notin \mathbb{Z}$ since $i_0 > 2$.

For $r = 5, 6$, we consider $(2 + \sqrt{-1})$ -adic valuation and we can prove the theorem similarly. □

Next, we construct an example of quadrature formulas of degree 4 for Bessel polynomials with 3 nodes on $\mathbb{Q}(\sqrt{-1})$.

Since $y_2(z) = 3z^2 + 3z + 1$ and $y_3(z) = 15z^3 + 15z^2 + 6z + 1$,

$$f_2(x, y) = 3((15y^2 + 15y + 5)x^2 + (15y^2 + 14y + 4)x + (5y^2 + 4y + 1)).$$

by Definition 2.9. Multiplying both sides of $f_2(x, y) = 0$ by $4(15y^2 + 15y + 5)$, we have

$$(2(15y^2 + 15y + 5)x + (15y^2 + 14y + 4))^2 + 75y^4 + 120y^3 + 84y^2 + 28y + 4 = 0.$$

Let $w := 2(15y^2 + 15y + 5)x + (15y^2 + 14y + 4)$, then we have a curve

$$C : w^2 = -75y^4 - 120y^3 - 84y^2 - 28y - 4.$$

The problem is reduced to determining the set $C(\mathbb{Q}(\sqrt{-1}))$ of $\mathbb{Q}(\sqrt{-1})$ -rational points on C . Let J be the Jacobian variety of C . Then, by MAGMA [2], we can check that $\text{rank } J(\mathbb{Q}(\sqrt{-1})) = 1$.

Example 3.2. From the rational points on the above elliptic curve C , we can construct an example of quadrature formulas of degree 4 for Bessel polynomials with 3 nodes on $\mathbb{Q}(\sqrt{-1})$. For example, by MAGMA [2],

$$P := (-264/743, \pm 377866\sqrt{-1}/552049) \in C(\mathbb{Q}(\sqrt{-1})).$$

Then, by Mathematica [13], for any weight x_1 , other weights x_2, x_3 and nodes z_1, z_2, z_3 are expressed by rational functions of x_1 and $\sqrt{-75x_1^4 - 120x_1^3 - 84x_1^2 - 28x_1 - 4}$. Substituting

⁵Note that for $n \in \mathbb{Z}$, n is divisible by $2 + \sqrt{-1}$ in $\mathbb{Z}[\sqrt{-1}]$ if and only if n is divisible by 5 in \mathbb{Z} .

$x_1 = -264/743$, we can check that

$$\begin{aligned} & \frac{304758098401}{73863713379} f\left(\frac{-264}{743}\right) \\ & + \left(-\frac{115447192511}{73863713379} + \frac{28870417761487643\sqrt{-1}}{27910585919669214}\right) f\left(\frac{-253754 + 188933\sqrt{-1}}{863405}\right) \\ & + \left(-\frac{115447192511}{73863713379} - \frac{28870417761487643\sqrt{-1}}{27910585919669214}\right) f\left(\frac{-253754 - 188933\sqrt{-1}}{863405}\right) \\ & = -\frac{1}{4\pi\sqrt{-1}} \int_{S^1} f(z)e^{-\frac{2}{z}} dz \end{aligned}$$

is a quadrature formula of degree 4 for Bessel polynomials with nodes on $\mathbb{Q}(\sqrt{-1})$ corresponding to P .

Remark 3.3. (1) By the same argument as in Theorem 1.3, we can prove that there exists no quadrature formula of degree $2r$ for Bessel polynomials with $r + 1$ nodes on $\mathbb{Q}(\sqrt{-1})$ for all $r \in \mathbb{Z}_{\geq 3}$.

(2) We cannot prove Theorem 1.3 (2) by using Lemma 2.4 since there exist quadrature formulas of degree 4 for Bessel polynomials with 3 nodes on $\mathbb{Q}(\sqrt{-1})$ by Example 3.2.

4. PROOF OF THEOREM 1.4

In this section, we prove Theorem 1.4

Proof of Theorem 1.4. Note that this is the case $2r + 2 - k = r + 1$. i.e., $k = r + 1 = 3$ in the Riesz-Shohat theorem Theorem 2.2. In this case, we prove the following by contradiction:

There exist no $z_1, z_2, z_3 \in \mathbb{Q}(\sqrt{-1}) \cap S^1$, $b_1, b_2 \in \mathbb{C}$ such that

$$(x - z_1)(x - z_2)(x - z_3) = \phi_3(x) + b_1\phi_2(x) + b_2\phi_1(x).$$

Here,

$$\phi_n(x) := \frac{y_n(x)}{\frac{(2n)!}{2^n n!}}.$$

Let

$$A := \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ \frac{2}{5} & 1 & 1 & 0 \\ \frac{1}{15} & \frac{1}{3} & 1 & 1 \end{pmatrix},$$

and suppose that

$$A \begin{pmatrix} 1 \\ b_1 \\ b_2 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ -z_1 - z_2 - z_3 \\ z_1 z_2 + z_2 z_3 + z_3 z_1 \\ -z_1 z_2 z_3 \end{pmatrix}.$$

Comparing the second to the fourth row,

$$\begin{aligned} b_1 &= -1 - z_1 - z_2 - z_3, \\ b_2 &= \frac{3}{5} + (z_1 + z_2 + z_3) + (z_1 z_2 + z_2 z_3 + z_3 z_1), \\ \left(z_1 + z_2 + \frac{2}{3} + z_1 z_2\right) z_3 &= -\frac{1}{3} - \frac{2}{3}(z_1 + z_2) - z_1 z_2. \end{aligned}$$

By the third equality, if

$$z_1 + z_2 + \frac{2}{3} + z_1 z_2 = 0,$$

then

$$z_1 z_2 = -\frac{2}{3}(z_1 + z_2) - \frac{1}{3}.$$

By the above two equalities,

$$z_1 + z_2 + 1 = 0.$$

Write $z_1 = s + t\sqrt{-1}$, $z_2 = u + v\sqrt{-1}$ with $s, t, u, v \in \mathbb{Q}$ and $s^2 + t^2 = u^2 + v^2 = 1$. Then, $s + u + 1 = 0$ and $t + v = 0$. Thus, $u = -s - 1$ and $v = -t$. Since $s^2 + t^2 = u^2 + v^2 = 1$,

$$1 = (-s - 1)^2 + t^2 = 2s + 2.$$

Thus, $s = -1/2$ and $t = \pm\sqrt{3}/2 \notin \mathbb{Q}$, which is a contradiction. Therefore,

$$z_1 + z_2 + \frac{2}{3} + z_1 z_2 \neq 0$$

and we have

$$z_3 = \frac{-\frac{1}{3} - \frac{2}{3}(z_1 + z_2) - z_1 z_2}{z_1 + z_2 + \frac{2}{3} + z_1 z_2}.$$

We solve $|z_1|^2 = |z_2|^2 = |z_3|^2 = 1$ by Mathematica [13]. Here are outputs:

$$(t, u) = \left(\sqrt{1 - s^2}, \frac{-91 - 202s - 112s^2 \pm 2\sqrt{43 + 84s + s^2 - 84s^3 - 44s^4}}{106 + 224s + 120s^2} \right)^6$$

$$(t, u) = \left(-\sqrt{1 - s^2}, \frac{-91 - 202s - 112s^2 \pm 2\sqrt{43 + 84s + s^2 - 84s^3 - 44s^4}}{106 + 224s + 120s^2} \right)$$

$$(s, t, u, v) = \left(-1, 0, -\frac{1}{2}, \pm\frac{\sqrt{3}}{2} \right),$$

$$(s, t, u, v) = \left(1, 0, -\frac{9}{10}, \pm\frac{\sqrt{19}}{10} \right),$$

$$(s, t, u, v) = \left(\frac{-28 - \sqrt{-11}}{30}, -\frac{\sqrt{127 - 56\sqrt{-11}}}{30}, \frac{-28 + \sqrt{-11}}{30}, \frac{\sqrt{127 + 56\sqrt{-11}}}{30} \right),$$

$$(s, t, u, v) = \left(\frac{-28 - \sqrt{-11}}{30}, \frac{\sqrt{127 - 56\sqrt{-11}}}{30}, \frac{-28 + \sqrt{-11}}{30}, -\frac{\sqrt{127 + 56\sqrt{-11}}}{30} \right),$$

$$(s, t, u, v) = \left(\frac{-28 + \sqrt{-11}}{30}, -\frac{\sqrt{127 + 56\sqrt{-11}}}{30}, \frac{-28 - \sqrt{-11}}{30}, \frac{\sqrt{127 - 56\sqrt{-11}}}{30} \right),$$

$$(s, t, u, v) = \left(\frac{-28 + \sqrt{-11}}{30}, \frac{\sqrt{127 + 56\sqrt{-11}}}{30}, \frac{-28 - \sqrt{-11}}{30}, -\frac{\sqrt{127 - 56\sqrt{-11}}}{30} \right).$$

For the third to the eighth solution, $s \notin \mathbb{Q}$ or $v \notin \mathbb{Q}$, which are impossible.

For the first and the second solution, we claim that

$$43 + 84s + s^2 - 84s^3 - 44s^4$$

⁶ v can be expressed by a rational function of s , $\sqrt{43 + 84s + s^2 - 84s^3 - 44s^4}$ and $\sqrt{1 - s^2}$.

is a rational square if and only if $s = \pm 1$ or $-1/2$ or $-9/10$, which contradict $t \in \mathbb{Q}$ or $v \in \mathbb{Q}$.
Indeed, let C be an elliptic curve over \mathbb{Q} defined by

$$y^2 = -44x^4 - 84x^3 + x^2 + 84x + 43.$$

By MAGMA [2], $\text{rank } C(\mathbb{Q}) = 0$. We also know that $(\pm 1, 0), (-\frac{1}{2}, \pm 3), (-\frac{9}{10}, \pm \frac{19}{25}) \in C(\mathbb{Q})$ and the torsion subgroup is isomorphic to $\mathbb{Z}/6\mathbb{Z}$. Therefore, we have

$$C(\mathbb{Q}) = \left\{ (\pm 1, 0), \left(-\frac{1}{2}, \pm 3\right), \left(-\frac{9}{10}, \pm \frac{19}{25}\right) \right\}.$$

□

Remark 4.1. (1) Remark 3.1 implies that there exists a unique quadrature formulas of degree 2 for Bessel polynomials with 2 nodes on $\mathbb{Q}(\sqrt{-11}) \cap S^1$. On the other hand, there exists no quadrature formula of degree 3 for Bessel polynomials with 3 nodes on $\mathbb{Q}(\sqrt{-11}) \cap S^1$. The proof is exactly similar to Theorem 1.4.

(2) There exists quadrature formulas of degree 3 for Bessel polynomials with 3 nodes on $\mathbb{Q}(\sqrt{-3}) \cap S^1$. For example,

$$\frac{2}{3}f(-1) + \frac{3 + \sqrt{-3}}{18}f\left(\frac{-1 + \sqrt{-3}}{2}\right) + \frac{3 - \sqrt{-3}}{18}f\left(\frac{-1 - \sqrt{-3}}{2}\right) = -\frac{1}{4\pi\sqrt{-1}} \int_{S^1} f(z)e^{-\frac{z}{2}} dz$$

is a quadrature formula of degree 3 for Bessel polynomials with 3 nodes on $\mathbb{Q}(\sqrt{-3}) \cap S^1$.

5. PROOF OF THEOREM 1.5

In this section, we prove Theorem 1.5.

Proof of Theorem 1.5. By Theorem 2.2, it is enough to show that there exist $z_1, \dots, z_{r+1} \in \mathbb{Q}$, $b_1, \dots, b_r \in \mathbb{C}$ such that

$$(3) \quad \omega_{r+1}(x) := (x - z_1) \cdots (x - z_{r+1}) = \phi_{r+1}(x) + b_1 \phi_r(x) + \dots + b_r \phi_1(x) + 0 \cdot \phi_0(x).$$

Here,

$$\phi_n(x) := \frac{y_n(x)}{2^n n!}.$$

Write

$$(x - z_1) \cdots (x - z_{r+1}) = 1 \cdot x^{r+1} + Z_1(z_1, \dots, z_{r+1})x^r + \dots + Z_{r+1}(z_1, \dots, z_{r+1}),$$

$$\phi_{r+1}(x) + b_1 \phi_r(x) + \dots + b_r \phi_1(x) = 1 \cdot x^{r+1} + B_1(b_1, \dots, b_r)x^r + \dots + B_{r+1}(b_1, \dots, b_r).$$

Let

$$A := \begin{pmatrix} 1 & 0 & 0 & 0 & \dots & 0 & 0 & 0 \\ a_{(r+1)1} & 1 & 0 & 0 & \dots & 0 & 0 & 0 \\ a_{(r+1)2} & a_{r1} & 1 & 0 & \dots & 0 & 0 & 0 \\ a_{(r+1)3} & a_{r2} & a_{(r-1)1} & 1 & \dots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots & \vdots & \vdots \\ a_{(r+1)(r-1)} & a_{r(r-2)} & a_{(r-1)(r-3)} & a_{(r-2)(r-4)} & \dots & 1 & 0 & 0 \\ a_{(r+1)r} & a_{r(r-1)} & a_{(r-1)(r-2)} & a_{(r-2)(r-3)} & \dots & a_{21} & 1 & 0 \\ a_{(r+1)(r+1)} & a_{rr} & a_{(r-1)(r-1)} & a_{(r-2)(r-2)} & \dots & a_{22} & a_{11} & 1 \end{pmatrix},$$

where we write

$$\phi_k(x) := \sum_{i=0}^k a_{ki} x^{k-i} \quad (a_{k0} = 1).$$

Then, by comparing the coefficients of x^k in (2), eq. (3) is equivalent to

$$A \begin{pmatrix} 1 \\ b_1 \\ \vdots \\ b_r \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ B_1(b_1, \dots, b_r) \\ \vdots \\ B_{r+1}(b_1, \dots, b_r) \end{pmatrix} = \begin{pmatrix} 1 \\ Z_1(z_1, \dots, z_{r+1}) \\ \vdots \\ Z_{r+1}(z_1, \dots, z_{r+1}) \end{pmatrix}.$$

Since $\det(A) \neq 0$, this is equivalent to

$$(4) \quad \begin{pmatrix} 1 \\ b_1 \\ \vdots \\ b_r \\ 0 \end{pmatrix} = A^{-1} \begin{pmatrix} 1 \\ Z_1(z_1, \dots, z_{r+1}) \\ \vdots \\ Z_{r+1}(z_1, \dots, z_{r+1}) \end{pmatrix}.$$

$z_1, \dots, z_r, b_1, \dots, b_r$ in eq. (3) can be taken as follows:

First, take $z_1, \dots, z_r \in \mathbb{Q}$ arbitrarily. By comparing the last row of eq. (4), we have a relation on z_1, \dots, z_{r+1} . For each i , note that Z_i is an elementary symmetric polynomial in z_1, \dots, z_{r+1} and is linear in z_{r+1} . Thus, $z_{r+1} \in \mathbb{Q}$ can be expressed by a rational function of z_1, \dots, z_r . Then, by eq. (4), b_1, \dots, b_{r+1} can be expressed in terms of z_1, \dots, z_{r+1} .⁷ \square

Example 5.1.

$$\frac{172}{441} f\left(\frac{1}{2}\right) - \frac{1625}{1611} f\left(\frac{1}{5}\right) + \frac{42592}{26313} f\left(-\frac{27}{44}\right) = -\frac{1}{4\pi\sqrt{-1}} \int_{S^1} f(z) e^{-\frac{2}{z}} dz$$

is a quadrature formula of degree 3 with 3 nodes on \mathbb{Q} .

Acknowledgements. The author thanks his advisor Professor Ken-ichi Bannai for reading the draft and giving helpful comments. The author also thanks him for warm and constant encouragement. The author gratefully thanks Professor Yukihiro Uchida for helpful comments and discussions. The author also thanks Professors Shuji Yamamoto, Yoshinosuke Hirakawa, Naoto Dainobu, Yoshinori Kanamura for helpful comments and discussions.

REFERENCES

- [1] Eiichi Bannai and Etsuko Bannai, *Tight Gaussian 4-designs*, J. Algebraic Combin. **22** (2005), no. 1, 39–63, DOI 10.1007/s10801-005-2505-3. MR2163709
- [2] Wieb Bosma, John Cannon, and Catherine Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), no. 3-4, 235–265, DOI 10.1006/jsco.1996.0125. Computational algebra and number theory (London, 1993). MR1484478
- [3] Robert Breusch, *Zur Verallgemeinerung des Bertrandschen Postulates, daß zwischen x und $2x$ stets Primzahlen liegen*, Math. Z. **34** (1932), no. 1, 505–526, DOI 10.1007/BF01180606 (German). MR1545270
- [4] P. Delsarte, J. M. Goethals, and J. J. Seidel, *Spherical codes and designs*, Geometriae Dedicata **6** (1977), no. 3, 363–388, DOI 10.1007/bf03187604. MR485471
- [5] F. Hausdorff, *Zur Hilbertschen Lösung des Waringschen Problems*, Math. Ann. **67** (1909), no. 3, 301–305, DOI 10.1007/BF01450406 (German). MR1511531

⁷Thus, $b_1, \dots, b_r \in \mathbb{Q}$.

- [6] H. L. Krall and Orrin Frink, *A new class of orthogonal polynomials: The Bessel polynomials*, Trans. Amer. Math. Soc. **65** (1949), 100–115, DOI 10.2307/1990516. MR28473
- [7] Roelof Koekoek, Peter A. Lesky, and René F. Swarttouw, *Hypergeometric orthogonal polynomials and their q -analogues*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2010. With a foreword by Tom H. Koornwinder. MR2656096
- [8] Yu. V. Nesterenko, *On Waring’s problem (elementary methods)*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **322** (2005), no. Trudy po Teorii Chisel, 149–175, 254, DOI 10.1007/s10958-006-0266-8 (Russian, with English and Russian summaries); English transl., J. Math. Sci. (N.Y.) **137** (2006), no. 2, 4699–4715. MR2138457
- [9] Masanori Sawa and Yukihiro Uchida, *Discriminants of classical quasi-orthogonal polynomials with application to Diophantine equations*, J. Math. Soc. Japan **71** (2019), no. 3, 831–860, DOI 10.2969/jmsj/79877987. MR3984244
- [10] ———, *Algebro-geometric aspects of the Christoffel-Darboux kernels for classical orthogonal polynomials*, Trans. Amer. Math. Soc. **373** (2020), no. 2, 1243–1264, DOI 10.1090/tran/7998. MR4068263
- [11] J. Shohat, *On mechanical quadratures, in particular, with positive coefficients*, Trans. Amer. Math. Soc. **42** (1937), no. 3, 461–496, DOI 10.2307/1989740. MR1501930
- [12] Gábor Szegő, *Orthogonal polynomials*, 4th ed., American Mathematical Society Colloquium Publications, Vol. XXIII, American Mathematical Society, Providence, R.I., 1975. MR0372517
- [13] Wolfram Research, Inc., *Mathematica, Version 11.1.1.0*, Champaign, IL (2017).

Email address: `hidekimatsumura@keio.jp`

*DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND TECHNOLOGY, KEIO UNIVERSITY, 3-14-1 HIYOSHI, KOHOKU, YOKOHAMA, KANAGAWA, JAPAN