

Generalized Fisher-Darmois-Koopman-Pitman Theorem and Rao-Blackwell Type Estimators for Power-Law Distributions

Atin Gayen and M. Ashok Kumar

Department of Mathematics

Indian Institute of Technology Palakkad

Kerala 678557, India

Email: atinfordst@gmail.com; ashokm@iitpkd.ac.in

Abstract

This paper generalizes the notion of sufficiency for estimation problems beyond maximum likelihood. In particular, we consider estimation problems based on Jones et al. and Basu et al. likelihood functions that are popular among distance-based robust inference methods. We first characterize the probability distributions that always have a fixed number of sufficient statistics with respect to these likelihood functions. These distributions are power-law extensions of the usual exponential family and contain Student distributions as special case. We then extend the notion of minimal sufficient statistics and compute it for these power-law families. Finally, we establish a Rao-Blackwell type theorem for finding best estimators for a power-law family. This enables us to derive certain generalized Cramér-Rao lower bounds for power-law families.

Index Terms

Basu et al. and Jones et al. likelihood functions, Cramér-Rao inequality, Fisher-Darmois-Koopman-Pitman theorem, Power-law distributions, Rao-Blackwell theorem, Student distribution, Sufficient Statistics.

I. INTRODUCTION

Consider a parametric family of probability distributions, parameterized by θ , $\Pi = \{p_\theta : \theta \in \Theta\}$, where Θ is an open subset of \mathbb{R}^k . We shall assume (unless explicitly specified) that all distributions in the family have a common support $\mathbb{S} \subset \mathbb{R}$. Indeed, if there is no prior information about θ , an experimenter who wishes to estimate the parameter θ , uses only the given samples to estimate θ . Thus an estimator of θ is a function of the given samples, called a *statistic*. However, if the estimators are functions of some specific statistics of the samples, then the experimenter can use these statistics to estimate θ without knowing the entire sample. Fisher introduced this concept in his seminal paper titled ‘‘On the mathematical foundations of theoretical statistics’’ and termed it as a *sufficient statistic* for the parameter [20, Sec. 7]. He designated a statistic T *sufficient for estimating* θ if the conditional distribution of the samples, given T , is independent of θ . Formally, if $X_1^n := X_1, \dots, X_n$ are n independent and identically distributed (i.i.d.) random variables from some $p_\theta \in \Pi$ and $T := T(X_1^n)$ is a statistic, then T is sufficient (for estimating θ) if the conditional distribution

$$p_{\theta_{X_1^n|T}}(x_1^n|t) \text{ is independent of } \theta \quad (1)$$

for every $t \in \mathcal{T} := \{t : t = T(y_1^n) \text{ for some } y_1^n \in \mathbb{S}^n\}$. This definition, however, is helpful only if one has a guess of a suitable candidate for T . Observe also that the distribution of T is needed to find the conditional distribution in (1). Later Koopman proposed the following equivalent definition, where the distribution of T is not needed [32]. He called a statistic T sufficient for θ , if the following holds. If x_1^n and y_1^n are two sets of i.i.d. samples from some $p_\theta \in \Pi$ such that they are same under T , that is, $T(x_1^n) = T(y_1^n)$, then the difference of their respective likelihoods, $[L(x_1^n; \theta) - L(y_1^n; \theta)]$ is independent of θ , where $L(x_1^n; \theta) := \sum_{i=1}^n \log p_\theta(x_i)$ denotes the log-likelihood function of the sample x_1^n . Observe that, to use this definition also, one needs to have a guess for T . Let us now suppose that a sufficient

statistic T is known. A natural question that arises is, what kind of estimators that can be found using T . Observe that both Fisher's and Koopman's definitions of sufficiency make use of the likelihood function L . However, it is still not clear if Fisher or Koopman had maximum likelihood (ML)-estimation in mind while proposing the above definitions. The following result, known as *factorization theorem* [27], [39], which is equivalent to both the above definitions, seems to answer this. A statistic T is sufficient for θ if and only if there exist two functions $u : \Theta \times \mathcal{F} \rightarrow \mathbb{R}$ and $v : \mathbb{S}^n \rightarrow \mathbb{R}$ such that

$$L(x_1^n; \theta) = u(\theta, T(x_1^n)) + v(x_1^n) \quad (2)$$

for all x_1^n and all $\theta \in \Theta$ [10, Th. 6.2.6]. Observe that, neither any prior guess for T nor its distribution is needed in this case. If θ_{MLE} denotes the maximum likelihood estimator (MLE) of θ , then from (2), we have

$$\theta_{\text{MLE}} = \arg \max_{\theta} L(X_1^n; \theta) = \arg \max_{\theta} [u(\theta, T(X_1^n)) + v(X_1^n)] = \arg \max_{\theta} u(\theta, T(X_1^n)).$$

This implies that MLE of θ depends on the sample only through statistic T (that is, knowledge of T is sufficient for the estimation of θ). This makes it clear that the classical notion of sufficiency is based on ML-estimation.

Observe that the entire sample is always a sufficient statistic with respect to any estimation. However, the notion of sufficient statistics is helpful only when it exists in fixed number, independent of the sample size. *Fisher-Darmois-Koopman-Pitman theorem* addresses this problem [1], [16], [21], [32], [42]. It states that, under certain regularity conditions, existence of such a set of sufficient statistics is possible if and only if the underlying family is exponential. A set of probability distributions \mathcal{E} is said to be an *exponential family* if there exist functions $w = [w_1, \dots, w_s]^\top$ and $f = [f_1, \dots, f_s]^\top$ with $w_i : \Theta \rightarrow \mathbb{R}$, $f_i : \mathbb{S} \rightarrow \mathbb{R}$, and $h : \mathbb{R} \rightarrow \mathbb{R}$ such that each $p_\theta \in \mathcal{E}$ is given by

$$p_\theta(x) = \begin{cases} Z(\theta) \exp [h(x) + w(\theta)^\top f(x)] & \text{if } x \in \mathbb{S} \\ 0 & \text{otherwise,} \end{cases} \quad (3)$$

where $Z(\theta)$ is the normalizing factor [7]. Bernoulli, Binomial, Poisson and Normal distributions are some examples of exponential family. However, a statistic that is sufficient for ML-estimation might not be sufficient for some other estimation. For example, consider the family of binomial distributions with parameters, say (m, θ) , where m is a positive integer denoting the number of trials (known) and $\theta \in (0, 1)$ is the success probability which is unknown. It is well-known that the sum of the samples is always a sufficient statistic for θ (with respect to the usual likelihood function) (see, for example, [10]).

Let us now suppose that we want to estimate θ by maximizing the following likelihood function, called *Cauchy-Schwarz likelihood function* (c.f. [9, Eq. (2.90)]):

$$L_{cs}(x_1^n; \theta) := \log \left[\frac{\sum_{x=0}^m p_n(x) p_\theta(x)}{\sum_{x=0}^m p_\theta(x)^2} \right], \quad (4)$$

where $p_\theta(x) = \binom{m}{x} \theta^x (1 - \theta)^{m-x}$ for $x = 0, \dots, m$ and p_n is the empirical distribution of the observed samples. Maximizing Cauchy-Schwarz likelihood function is same as minimizing the well-known Cauchy-Schwarz divergence. (This divergence is popular in machine learning applications such as blind source separation, independent component analysis, manifold learning, image segmentation and so on [28], [29], [31], [40], [41], [46].)

It is easy to check that the solutions of the estimating equation, in this case, are the roots of a polynomial in θ whose coefficients involve some of $p_n(x)$, $x = 0, \dots, m$. For example, when $m = 2$, it is a polynomial of degree five and the following are, respectively, the coefficients in descending order of the exponent of θ :

$$8p_n(1), \quad 8p_n(0) - 20p_n(1) - 6p_n(2), \quad -20p_n(0) + 16p_n(1) + 12p_n(2), \quad 18p_n(0) - 4p_n(1) - 6p_n(2), \\ -7p_n(0) - 2p_n(1) + p_n(2), \quad \text{and } p_n(0) + p_n(1).$$

This shows that at least two among $p_n(0), p_n(1)$ and $p_n(2)$ are needed to find the estimator. Hence sum of the samples is no longer sufficient for estimating θ .

On the other hand, let us consider the following family of compactly-supported Student distributions given by

$$p_\theta(x) = \begin{cases} \frac{3}{4\sqrt{5}\sigma} \left[1 - \frac{(x-\mu)^2}{5\sigma^2} \right] & \text{if } x \in \mathbb{S} \\ 0 & \text{otherwise,} \end{cases} \quad (5)$$

where $\mathbb{S} = [\mu - \sqrt{3}\sigma, \mu + \sqrt{3}\sigma]$, $\theta = (\mu, \sigma^2)$, $\mu \in (-\infty, \infty)$ and $\sigma \in (0, \infty)$. Observe that (5) does not fall in an exponential family as defined in (3). Hence, by Fisher-Darmois-Koopman-Pitman theorem, there does not exist any fixed number of sufficient statistics (independent of sample size) for μ and σ . Indeed, it can be shown that, the knowledge of entire sample is needed for estimating μ and σ in this case (See [25]). However, Eguchi et al. considered an estimation based on Cauchy-Schwarz likelihood function (4). They found closed-form solution for the estimators under the assumption that $(X_i - \mu)^2 \leq 3\sigma^2$ for all $i = 1, \dots, n$ (See [17, Eq. (38)]). The estimators are functions of the statistics $\sum_{i=1}^n X_i$ and $\sum_{i=1}^n X_i^2$ [17, Th. 3] (c.f. [26, Sec. 4.2]). This motivates us to generalize the concept of sufficiency based on an estimation method. Let us consider the difference of Cauchy-Schwarz likelihood function L_{cs} evaluated at samples x_1^n and y_1^n :

$$L_{cs}(x_1^n; \theta) - L_{cs}(y_1^n; \theta) = \log \sum_{i=1}^n p_\theta(x_i) - \log \sum_{i=1}^n p_\theta(y_i).$$

For the family of Student distributions (5), we have

$$L_{cs}(x_1^n; \theta) - L_{cs}(y_1^n; \theta) = \log \frac{1 - (1/(5\sigma^2 - \mu^2)) \sum_{i=1}^n x_i^2 + (2\mu/(5\sigma^2 - \mu^2)) \sum_{i=1}^n x_i}{1 - (1/(5\sigma^2 - \mu^2)) \sum_{i=1}^n y_i^2 + (2\mu/(5\sigma^2 - \mu^2)) \sum_{i=1}^n y_i}. \quad (6)$$

(6) is independent of θ if

$$\sum_{i=1}^n x_i^2 = \sum_{i=1}^n y_i^2 \quad \text{and} \quad \sum_{i=1}^n x_i = \sum_{i=1}^n y_i.$$

Notice that the Cauchy-Schwarz estimators for μ and σ depend on the sample only through the statistics $\sum_{i=1}^n X_i$ and $\sum_{i=1}^n X_i^2$. This implies that, when the estimation is based on the likelihood function L_{cs} , it is reasonable to call the pair $(\sum X_i, \sum X_i^2)$ sufficient for μ and σ .

It should also be noted that Cauchy-Schwarz likelihood function is a special case ($\alpha = 2$) of the more general class of likelihood functions, namely *Jones et al. likelihood function*:

$$L_J^{(\alpha)}(x_1^n; \theta) := \frac{1}{\alpha - 1} \log \left[\frac{1}{n} \sum_{i=1}^n p_\theta(x_i)^{\alpha-1} \right] - \frac{1}{\alpha} \log \left[\int p_\theta(x)^\alpha dx \right], \quad (7)$$

where $\alpha > 0$, $\alpha \neq 1$. Jones et al. likelihood function is popular in the context of robust estimation [30]. Observe that (7) coincides with the usual likelihood function $L(\theta)$ as $\alpha \rightarrow 1$. Hence Jones et al. likelihood function is considered as a robust alternative to ML-estimation. Motivated by the works of Windham [50] and Field and Smith [19], Jones et al. proposed this method in search of an estimating equation, alternative to the one of MLE, that uses a suitable power of the model density to down-weight the effect of *outliers* (possibly) present in the data [17], [22], [26], [30]. It is well-known that ML-estimation is closely related to the minimization of the well-known Kullback-Leibler divergence or relative entropy [15, Lem. 3.1]. Similarly, Jones et al. estimation is related to the minimization of the so-called Jones et al. divergence [23], [26], [36], [47], [48] (also known as γ -divergence [11], [23], projective power divergence [17], logarithmic density power divergence [37], relative α -entropy [33], [34], Rényi pseudo-distance [5], [6]). While Jones et al. divergence for $\alpha > 1$ corresponds to robust inference, the same for $\alpha < 1$ has connections to problems in information theory, such as mismatched guessing [47], source coding [34], encoding of tasks

[8] and so on. Though Jones et al. estimation for $\alpha < 1$ does not have a meaning in terms of robustness, it can be used to approximate MLEs of Student distributions (c.f. [26, Rem. 7]).

Another likelihood function that is popular in the context of robust inference is the so-called Basu et al. likelihood function given by

$$L_B^{(\alpha)}(x_1^n; \theta) = \frac{1}{n} \sum_{i=1}^n \left[\frac{\alpha \cdot p_\theta(x_i)^{\alpha-1} - 1}{\alpha - 1} \right] - \int p_\theta(x)^\alpha dx, \quad (8)$$

where $\alpha > 0$, $\alpha \neq 1$ [3, Eq. (2.2)], maximization of which results in a non-normalized version of Jones et al. estimating equation (see [26, Sec. 1]). Basu et al. estimation is closely related to the minimization of the so-called Basu et al. divergence which belongs to the Bregman class [13]. $\alpha = 2$ corresponds to the squared Euclidean distance. A comparison of Basu et al. and Jones et al. estimations was studied in [30]. Basu et al. and Jones et al. likelihood functions are special, unlike other likelihood functions such as the Hellinger likelihood function, in the sense that smoothing of the empirical probability measure is not needed (see [26, Sec. 1] for details). In this paper, we propose a generalized notion of sufficiency based on the likelihood function associated with an estimation problem. Specifically, we will focus on Jones et al. and Basu et al. likelihood functions. We will characterize the family of probability distributions that has a fixed number of sufficient statistics independent of the sample size with respect to these likelihood functions. It turns out that these are the power-law families studied in [2], [14], [26], [34], [38]. Student distributions fall in this family.

Sufficient statistics not only help in data reduction but also in finding a better estimator than the given one. Indeed, if one has an unbiased estimator along with sufficient statistics, it is always possible to find another unbiased estimator (usually a function of the sufficient statistics) whose variance is not more than the given one. This result is known as the *Rao-Blackwell theorem* [4], [43]. It takes the conditional expectation of an unbiased estimator given the sufficient statistics which results in a uniform improvement on the variance of the estimator. Hence this suggests one to consider only such conditional expectations while searching for *best unbiased estimators*.

In this paper, we find Rao-Blackwell type estimators for power-law families of the form (5) with respect to Jones et al. and Basu et al. likelihood functions. We also find a lower bound for the variance of these estimators. This generalizes the usual Cramér-Rao Lower Bound (CRLB) and results in a sharper bound for the power-law families. We also apply these results, in particular, to the Student distributions.

Our main contributions in this paper are the following.

- (i) Propose a generalized notion of principle of sufficiency that is appropriate when the underlying estimation is not necessarily ML-estimation.
- (ii) Characterize the families of probability distributions having a fixed number of sufficient statistics independent of sample size with respect to Basu et al. and Jones et al. likelihood functions.
- (iii) Generalize the notion of minimal sufficiency with respect to a likelihood function and find generalized minimal sufficient statistics for power-law family.
- (iv) Extend Rao-Blackwell theorem to both Basu et al. and Jones et al. estimations and find best estimators for power-law family.
- (v) Show that variance of any unbiased estimator of a power-law family is bounded below by the asymptotic variance of Jones et al. estimator.

The rest of the paper is organized as follows. In Section II, we introduce the generalized notion of sufficiency. In Section III, Fisher-Darmois-Koopman-Pitman theorem for Jones et al. and Basu et al. likelihood functions are established. In this section, we also find minimal sufficient statistics for the mean and variance parameters of Student distributions. In Section IV, we find generalized Rao-Blackwell type

estimators for power-law family. In Section V, generalized CRLB for Basu et al. and Jones et al. estimators of power-law family are derived. In this section, we also find best estimators for the mean and variance parameters of Student distributions. We end the paper with a summary and concluding remarks in Section VI.

II. A GENERALIZED NOTION OF PRINCIPLE OF SUFFICIENCY

In this section, we first propose a generalized notion of sufficiency analogous to Koopman's definition of sufficiency. We then derive a factorization theorem associated with this generalized notion. We also show that this is equivalent to generalized Fisher's definition of sufficiency.

Definition 1: Let $\Pi = \{p_\theta : \theta \in \Theta\}$ be a family of probability distributions. Suppose the underlying problem is to estimate θ by maximizing some generalized likelihood function L_G . A set of statistics T is said to be a *sufficient statistic for θ with respect to L_G* if $[L_G(x_1^n; \theta) - L_G(y_1^n, \theta)]$ is independent of θ whenever $T(x_1^n) = T(y_1^n)$.

Observe that, when L_G is the log-likelihood function, above definition coincides with the usual Koopman's definition of sufficiency. We now derive a generalized factorization theorem.

Proposition 2 (Generalized Factorization Theorem): Let $\Pi = \{p_\theta : \theta \in \Theta\}$ be a family of probability distributions. Suppose the underlying problem is to estimate θ by maximizing some generalized likelihood function L_G . Then a set of statistics T is a sufficient statistic for θ with respect to L_G if and only if there exist functions $u : \Theta \times \mathcal{T} \rightarrow \mathbb{R}$ and $v : \mathbb{S}^n \rightarrow \mathbb{R}$ such that

$$L_G(x_1^n; \theta) = u(\theta, T(x_1^n)) + v(x_1^n),$$

for all sample points x_1^n and for all $\theta \in \Theta$.

Proof: See Sec-A.1 of Appendix. ■

Observe that, when $L_G = L$, the above coincides with the usual factorization theorem [27], [39]. We now show that this generalized notion of sufficiency is also equivalent to a generalized version of Fisher's definition of sufficiency. Let X_1^n be i.i.d. random variables according to some $p_\theta \in \Pi$. Let $x_1^n := x_1, \dots, x_n$ be a sample. Let us define the deformed probability distribution associated with the likelihood function L_G by

$$p_\theta^*(x_1^n) := \frac{\exp[L_G(x_1^n, \theta)]}{\int \exp[L_G(y_1^n, \theta)] dy_1^n}.$$

Observe that, if $L_G(x_1^n; \theta) = L(x_1^n; \theta)$, then

$$p_\theta^*(x_1^n) = \frac{\exp[\log p_\theta(x_1^n)]}{\int \exp[\log p_\theta(y_1^n)] dy_1^n} = \frac{p_\theta(x_1^n)}{\int p_\theta(y_1^n) dy_1^n} = p_\theta(x_1^n). \quad (9)$$

Let T be a statistic of the sample x_1^n . For any value t of T , let $q_\theta^*(t) := \int p_\theta^*(y_1^n) \mathbf{1}(T(y_1^n) = t) dy_1^n$, where $\mathbf{1}(\cdot)$ denotes the indicator function. This can be re-written as

$$q_\theta^*(t) = \int_{A_t} p_\theta^*(y_1^n) dy_1^n,$$

where $A_t = \{x_1^n : T(x_1^n) = t\}$. The associated conditional distribution is given by

$$p_{\theta, x_1^n | T}^*(x_1^n | t) := \frac{p_\theta^*(x_1^n) \mathbf{1}(T(x_1^n) = t)}{q_\theta^*(t)} = \begin{cases} \frac{p_\theta^*(x_1^n)}{q_\theta^*(t)} & \text{if } x_1^n \in A_t \\ 0 & \text{otherwise.} \end{cases} \quad (10)$$

Then

$$\int p_{\theta_{x_1^n|T}}^*(x_1^n|t) dx_1^n = \int_{A_t} \frac{p_{\theta}^*(x_1^n)}{q_{\theta}^*(t)} dx_1^n = \int_{A_t} \frac{p_{\theta}^*(x_1^n)}{q_{\theta}^*(T(x_1^n))} dx_1^n = \frac{\int p_{\theta}^*(x_1^n) dx_1^n}{\int_{A_t} p_{\theta}^*(x_1^n) dx_1^n} = 1.$$

Observe that, when $L_G = L$, we have

$$p_{\theta_{x_1^n|T}}^*(x_1^n|t) = p_{\theta_{x_1^n|T}}(x_1^n|t),$$

the usual conditional distribution of x_1^n given $T = t$.

Proposition 3: T is sufficient for θ with respect to a likelihood function L_G if and only if $p_{\theta_{x_1^n|T}}^*(x_1^n|t)$ is independent of θ for any value t of T .

Proof: See Sec-A.2 of Appendix. ■

While sufficient statistics help in data compression, *minimal sufficient statistics* help maximum such compression. We now define the notion of minimal sufficiency in a general set-up. Let us first recall the definition of minimal sufficiency in the classical sense.

Definition 4: [10, Def. 6.2.11] A set of statistics T is said to be a *minimal sufficient statistic* if T is a function of any other sufficient statistic. In other words, T is minimal if the following holds: For any sufficient statistic \tilde{T} and for any two i.i.d. samples x_1^n and y_1^n , $\tilde{T}(x_1^n) = \tilde{T}(y_1^n)$ implies $T(x_1^n) = T(y_1^n)$.

There is an easy criterion to find minimal sufficient statistics for MLE due to Lehmann and Scheffé [10, Th. 6.2.13]. Here we first generalize this criterion to a generalized likelihood function. Later we will use it to find minimal sufficient statistics for power-law families.

Proposition 5: Let $\Pi = \{p_{\theta} : \theta \in \Theta\}$ be a family of probability distributions. Suppose the underlying problem is to estimate θ by maximizing some generalized likelihood function L_G . Assume that the following condition holds for a statistic T . For any two i.i.d. samples x_1^n and y_1^n , $T(x_1^n) = T(y_1^n)$ if and only if $[L_G(x_1^n; \theta) - L_G(y_1^n; \theta)]$ is independent of θ for $\theta \in \Theta$. Then T is a minimal sufficient statistic for θ with respect to L_G .

Proof: See Sec-A.3 of Appendix. ■

III. GENERALIZED FISHER-DARMOIS-KOOPMAN-PITMAN THEOREM

In this section, we first characterize the power-law families that have fixed number of sufficient statistics independent of sample size with respect to Jones et al. and Basu et al. likelihood functions. We then find sufficient statistics for the power-law families. We start by defining the family associated with Jones et al. likelihood function.

Definition 6: A k -parameter α -power-law family characterized by h, w, f, Θ and \mathbb{S} , where Θ is an open subset of \mathbb{R}^k , $w = [w_1, \dots, w_s]^\top$, $f = [f_1, \dots, f_s]^\top$, with $w_i : \Theta \rightarrow \mathbb{R}$ is differentiable for $i = 1, \dots, s$, $f_i : \mathbb{R} \rightarrow \mathbb{R}$ for $i = 1, \dots, s$, $h : \mathbb{R} \rightarrow \mathbb{R}$, and $\mathbb{S} \subseteq \mathbb{R}$, is defined by $\mathbb{M}^{(\alpha)} = \{p_{\theta} : \theta \in \Theta\}$, where

$$p_{\theta}(x) = \begin{cases} Z(\theta) [h(x) + w(\theta)^\top f(x)]^{\frac{1}{\alpha-1}} & \text{for } x \in \mathbb{S} \\ 0 & \text{otherwise,} \end{cases} \quad (11)$$

with $Z(\theta)^{-1} = \int_{\mathbb{S}} [h(x) + w(\theta)^\top f(x)]^{\frac{1}{\alpha-1}} dx$.

Remark 1:

- (i) $\mathbb{M}^{(\alpha)}$ -family is a generalization of the usual exponential family \mathcal{E} . This can be seen by the following way. Consider the α -exponential function $\exp_{\alpha} : [-\infty, \infty] \rightarrow (-\infty, \infty]$ defined by

$$\exp_{\alpha}(x) = \begin{cases} [\max\{0, 1 + (1 - \alpha)x\}]^{\frac{1}{1-\alpha}} & \text{for } \alpha \neq 1, \\ \exp(x) & \text{for } \alpha = 1 \end{cases}$$

[49]. Using this, (11) can be re-written as

$$p_\theta(x)^{-1} = Z(\theta)^{-1} \exp_\alpha \left[-h(x) - w(\theta)^\top f(x) \right], \quad (12)$$

for $x \in \mathbb{S}$. Observe that, the above coincides with an exponential family \mathcal{E} when $\alpha = 1$.

(ii) In [34], this family was derived as projections of Jones et al. divergence on a set of probability distributions determined by linear constraints. More about $\mathbb{M}^{(\alpha)}$ -family can be found in [2], [24], [26].

(iii) Student distributions fall in $\mathbb{M}^{(\alpha)}$ -family [26, Ex. 3].

We now state the main result of this section.

Theorem 7: Let $\Pi = \{p_\theta : \theta \in \Theta \subset \mathbb{R}^k\}$ be a family of probability distributions with common support \mathbb{S} , an open subset of \mathbb{R} . Let x_1^n be an i.i.d. sample from some $p_\theta \in \Pi$. For some s ($k \leq s < n$), let $T_j(x_1^n)$ for $j = 1, \dots, s$ be differentiable functions on \mathbb{S}^n . If $[T_1, \dots, T_s]^\top$ is a sufficient statistic for θ with respect to Jones et al. likelihood function $L_J^{(\alpha)}$, then there exist functions $h : \mathbb{S} \rightarrow \mathbb{R}$, $f_i : \mathbb{S} \rightarrow \mathbb{R}$ and $w_i : \Theta \rightarrow \mathbb{R}$, $i = 1, \dots, s$ such that $p_\theta(x)$ can be expressed as (11) for $x \in \mathbb{S}$.

Proof: We only give the proof when $k = s = 1$. Other cases can be proved similarly. Let us first consider the Jones et al. likelihood function (7).

$$\frac{\partial}{\partial \theta} L_J^{(\alpha)}(x_1^n; \theta) = \frac{1}{\alpha - 1} \frac{\sum_{i=1}^n \frac{\partial}{\partial \theta} [p_\theta(x_i)^{\alpha-1}]}{\sum_{i=1}^n p_\theta(x_i)^{\alpha-1}} - \frac{1}{\alpha} \frac{\partial}{\partial \theta} \log \int p_\theta(x)^\alpha dx. \quad (13)$$

If T is a sufficient statistic for θ with respect to $L_J^{(\alpha)}$, then from Proposition 2, we have

$$L_J^{(\alpha)}(x_1^n; \theta) = u(\theta, T) + v(x_1^n) \quad (14)$$

for some functions u and v , where v is independent of θ . This implies that

$$\frac{\partial}{\partial \theta} L_J^{(\alpha)}(x_1^n; \theta) = u_1(\theta, T), \quad (15)$$

where $u_1(\theta, T) := \frac{\partial}{\partial \theta} [u(\theta, T)]$. Comparing (13) and (15), we have

$$u_1(\theta, T) = \frac{1}{\alpha - 1} \frac{\sum_{i=1}^n \frac{\partial}{\partial \theta} [p_\theta(x_i)^{\alpha-1}]}{\sum_{i=1}^n p_\theta(x_i)^{\alpha-1}} - \frac{1}{\alpha} \frac{\partial}{\partial \theta} \log \int p_\theta(x)^\alpha dx. \quad (16)$$

Now, for a particular value of $\theta \in \Theta$, say θ_0 , (16) can be re-written as

$$\begin{aligned} u_1(\theta_0, T) &= \frac{1}{\alpha - 1} \frac{\sum_{i=1}^n \left[\frac{\partial}{\partial \theta} \{p_\theta(x_i)^{\alpha-1}\} \right]_{\theta_0}}{\sum_{i=1}^n p_{\theta_0}(x_i)^{\alpha-1}} - \left[\frac{\partial}{\partial \theta} \log \|p_\theta\| \right]_{\theta_0} \\ &= \frac{1}{\alpha - 1} \frac{\sum_{i=1}^n g(x_i)}{\sum_{i=1}^n h(x_i)} - \left[\frac{\partial}{\partial \theta} \log \|p_\theta\| \right]_{\theta_0}, \end{aligned} \quad (17)$$

where $\|p_\theta\| := \left[\int p_\theta(x)^\alpha dx \right]^{1/\alpha}$, $g(x_i) = \left[\frac{\partial}{\partial \theta} \{p_\theta(x_i)^{\alpha-1}\} \right]_{\theta_0}$, and $h(x_i) = p_{\theta_0}(x_i)^{\alpha-1}$.

Let us denote $T_1 := \sum_{i=1}^n g(x_i)$ and $T_2 := \sum_{i=1}^n h(x_i)$. Then from (17), there exists a function Φ such that

$$T = \Phi(T_1, T_2).$$

Using this, (15) can be re-written as

$$\frac{\partial}{\partial \theta} L_J^{(\alpha)}(x_1^n; \theta) = u_2(\theta, T_1, T_2) \quad (18)$$

for some function u_2 . Observe also that, (7) can be re-written as

$$L_J^{(\alpha)}(x_1^n; \theta) = \frac{1}{\alpha - 1} \log \left[\frac{1}{n} \sum_{i=1}^n p_\theta(x_i)^{\alpha-1} \right] - \frac{1}{\alpha - 1} \log \|p_\theta\|^{\alpha-1} = \frac{1}{\alpha - 1} \log \left[\frac{1}{n} \sum_{i=1}^n \frac{p_\theta(x_i)^{\alpha-1}}{\|p_\theta\|^{\alpha-1}} \right].$$

This implies that

$$\exp \left[(\alpha - 1) L_J^{(\alpha)}(x_1^n; \theta) \right] = \frac{1}{n} \sum_{i=1}^n \frac{p_\theta(x_i)^{\alpha-1}}{\|p_\theta\|^{\alpha-1}}.$$

Further (14) implies that

$$\exp \left[(\alpha - 1) L_J^{(\alpha)}(x_1^n; \theta) \right] = \exp \left[(\alpha - 1) u(T, \theta) \right] \cdot \exp \left[(\alpha - 1) v(x_1^n) \right].$$

Thus we have

$$\frac{1}{n} \sum_{i=1}^n \frac{p_\theta(x_i)^{\alpha-1}}{\|p_\theta\|^{\alpha-1}} = \exp \left[(\alpha - 1) u(T, \theta) \right] \cdot \exp \left[(\alpha - 1) v(x_1^n) \right]. \quad (19)$$

Fixing $\theta = \theta_0$, the above equation becomes

$$T_2 = n \|p_{\theta_0}\|^{\alpha-1} \exp \left[(\alpha - 1) u(T, \theta_0) \right] \cdot \exp \left[(\alpha - 1) v(x_1^n) \right].$$

Since $T = \Phi(T_1, T_2)$, this implies that $v(x_1^n)$ must be a function of T_1 and T_2 only. Therefore, (19) can be re-written as

$$\sum_{i=1}^n p_\theta(x_i)^{\alpha-1} = u_3(\theta, T_1, T_2), \quad (20)$$

where $u_3(\theta, T_1, T_2) := n \|p_\theta\|^{\alpha-1} \exp \left[(\alpha - 1) u(\Phi(T_1, T_2), \theta) \right] \exp \left[(\alpha - 1) v(x_1^n) \right]$. Now, taking partial derivative with respect to x_j on both sides of (20), we have

$$\begin{aligned} \frac{\partial}{\partial x_j} p_\theta(x_j)^{\alpha-1} &= \frac{\partial}{\partial T_1} u_3(\theta, T_1, T_2) \frac{\partial T_1}{\partial x_j} + \frac{\partial}{\partial T_2} u_3(\theta, T_1, T_2) \frac{\partial T_2}{\partial x_j} \\ &= \frac{\partial}{\partial T_1} u_3(\theta, T_1, T_2) \frac{\partial}{\partial x_j} g(x_j) + \frac{\partial}{\partial T_2} u_3(\theta, T_1, T_2) \frac{\partial}{\partial x_j} h(x_j). \end{aligned} \quad (21)$$

Since the left hand side of (21) is a function of x_j and θ only, we shall denote it by $f(x_j, \theta)$. Thus taking partial derivative with respect to x_i for $i \neq j$ on both sides of (21), we get

$$\frac{\partial}{\partial x_i} \left[\frac{\partial}{\partial T_1} u_3(\theta, T_1, T_2) \right] \frac{\partial}{\partial x_j} [g(x_j)] + \frac{\partial}{\partial x_i} \left[\frac{\partial}{\partial T_2} u_3(\theta, T_1, T_2) \right] \frac{\partial}{\partial x_j} [h(x_j)] = 0. \quad (22)$$

From (22), it is clear that

$$\frac{\partial}{\partial x_i} \left[\frac{\partial}{\partial T_1} u_3(\theta, T_1, T_2) \right] = 0 \text{ if and only if } \frac{\partial}{\partial x_i} \left[\frac{\partial}{\partial T_2} u_3(\theta, T_1, T_2) \right] = 0, \quad (23)$$

since neither $\frac{\partial}{\partial x_j} [g(x_j)]$ nor $\frac{\partial}{\partial x_j} [h(x_j)]$ can be zero.

Equivalently,

$$\frac{\partial}{\partial x_i} \left[\frac{\partial}{\partial T_1} u_3(\theta, T_1, T_2) \right] \neq 0 \text{ if and only if } \frac{\partial}{\partial x_i} \left[\frac{\partial}{\partial T_2} u_3(\theta, T_1, T_2) \right] \neq 0.$$

Suppose that $\frac{\partial}{\partial x_i} \left[\frac{\partial}{\partial T_1} u_3(\theta, T_1, T_2) \right] \neq 0$. Then (22) can be re-written as

$$\frac{\frac{\partial}{\partial x_i} \left[\frac{\partial}{\partial T_1} u_3(\theta, T_1, T_2) \right]}{\frac{\partial}{\partial x_i} \left[\frac{\partial}{\partial T_2} u_3(\theta, T_1, T_2) \right]} = - \frac{\frac{\partial}{\partial x_j} [h(x_j)]}{\frac{\partial}{\partial x_j} [g(x_j)]} =: \xi(x_j). \quad (24)$$

Since the right hand side of (24) is a function of x_j alone, both $\frac{\partial}{\partial T_1}[u_3(\theta, T_1, T_2)]$ and $\frac{\partial}{\partial T_2}[u_3(\theta, T_1, T_2)]$ must be of the following form

$$\begin{aligned}\frac{\partial}{\partial T_1}u_3(\theta, T_1, T_2) &= \Psi(x_1^n, \theta)S_1(x_j) + V_1(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n, \theta), \\ \frac{\partial}{\partial T_2}u_3(\theta, T_1, T_2) &= \Psi(x_1^n, \theta)S_2(x_j) + V_2(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n, \theta),\end{aligned}$$

for some real-valued functions Ψ, S_1, S_2, V_1 , and V_2 . Observe that (24) holds for any $i \neq j$. This implies V_1 and V_2 must be a function of x_j and θ only. Using these in (21), we get

$$f(x_j, \theta) = \Psi(x_1^n, \theta)K_1(x_j) + K_2(x_j, \theta), \quad (25)$$

where

$$\begin{aligned}K_1(x_j) &:= S_1(x_j) \frac{\partial}{\partial x_j}[g(x_j)] - S_2(x_j) \frac{\partial}{\partial x_j}[h(x_j)], \quad \text{and} \\ K_2(x_j, \theta) &:= V_1(x_j, \theta) \frac{\partial}{\partial x_j}[g(x_j)] - V_2(x_j, \theta) \frac{\partial}{\partial x_j}[h(x_j)].\end{aligned}$$

In order for (25) to hold, Ψ must be a function of x_j and θ only. This implies that both

$$\frac{\partial}{\partial T_1}u_3(\theta, T_1, T_2) \quad \text{and} \quad \frac{\partial}{\partial T_2}u_3(\theta, T_1, T_2) \quad (26)$$

must be a function of x_j and θ only.

On the other hand, let $\frac{\partial}{\partial x_i} \left[\frac{\partial}{\partial T_1}u_3(\theta, T_1, T_2) \right] = 0$ for some $i \neq j$. Then from (23), we have

$$\frac{\partial}{\partial x_i} \left[\frac{\partial}{\partial T_2}u_3(\theta, T_1, T_2) \right] = 0.$$

This implies that both $\frac{\partial}{\partial T_1}u_3(\theta, T_1, T_2)$ and $\frac{\partial}{\partial T_2}u_3(\theta, T_1, T_2)$ are independent of x_i 's where $i \neq j$. Thus, in this case also, (26) holds.

Since $u_3(\theta, T_1, T_2)$ is a function of T_1, T_2 and θ only, both $\frac{\partial}{\partial T_1}u_3(\theta, T_1, T_2)$ and $\frac{\partial}{\partial T_2}u_3(\theta, T_1, T_2)$ are also functions of T_1, T_2 and θ only. But both T_1 and T_2 are symmetric functions of the x_i 's, $i = 1, \dots, n$. This implies that both the quantities in (26) must be a function of θ only. Let

$$\frac{\partial}{\partial T_1}u_3(\theta, T_1, T_2) =: w_1(\theta) \quad \text{and} \quad \frac{\partial}{\partial T_2}u_3(\theta, T_1, T_2) =: w_2(\theta)$$

for some w_1 and w_2 which are functions of θ only. Then we have

$$u_3(\theta, T_1, T_2) = w_1(\theta)T_1 + w_2(\theta)T_2 + w_3(\theta).$$

Using (20), we get

$$p_\theta(x_i)^{\alpha-1} = w_1(\theta)g(x_i) + w_2(\theta)h(x_i) + w_3(\theta)$$

for $i = 1, \dots, n$. This completes the proof. ■

The converse of the above theorem is also true. Indeed, in the following, we show that an $\mathbb{M}^{(\alpha)}$ -family always has a fixed number of sufficient statistics with respect to the Jones et al. likelihood function $L_J^{(\alpha)}$.

Theorem 8: Consider a parametric family of probability distributions $\Pi = \{p_\theta : \theta \in \Theta \subset \mathbb{R}^k\}$ having a common support \mathbb{S} . Let $T = [T_1, \dots, T_s]^\top$ be s statistics defined on \mathbb{S}^n where $s \geq k$. T is a sufficient statistic for θ with respect to Jones et al. likelihood function $L_J^{(\alpha)}$ if the following hold.

- (a) Π is a k -parameter $\mathbb{M}^{(\alpha)}$ -family as in Definition 6,
- (b) there exist real valued functions ψ_1, \dots, ψ_s defined on \mathcal{T} such that for any i.i.d. sample x_1^n

$$\psi_i(T(x_1^n)) = \bar{f}_i / \bar{h} \quad (27)$$

for $i = 1, \dots, s$, where $\bar{f}_i = \frac{1}{n} \sum_{j=1}^n f_i(x_j)$ and $\bar{h} := \frac{1}{n} \sum_{j=1}^n h(x_j)$.

Proof: See Sec-B.1 of Appendix. ■

Remark 2: It is easy to see from the above theorem that $\bar{f}/\bar{h} := [\bar{f}_1/\bar{h}, \dots, \bar{f}_s/\bar{h}]^\top$ is a sufficient statistic for the k -parameter $\mathbb{M}^{(\alpha)}$ -family with respect to Jones et al. likelihood function.

Example 1: Consider the *Student distributions* with location parameter μ , scale parameter σ and degrees of freedom parameter $\nu \in (2, \infty)$:

$$p_{\mu,\sigma}(x) = N_{\sigma,\nu} \left[1 + \frac{(x - \mu)^2}{\nu\sigma^2} \right]^{-\frac{\nu+1}{2}}, \quad (28)$$

for $x \in \mathbb{R}$, where $N_{\sigma,\nu} := \frac{\Gamma(\frac{\nu+1}{2})}{\sqrt{\nu\pi\sigma^2} \Gamma(\frac{\nu}{2})}$ is the normalizing factor. Let ν be known, and μ and σ^2 be unknown. Letting $\alpha = \frac{\nu - 1}{\nu + 1}$ and $\theta = (\mu, \sigma)$, (28) can be re-written as

$$p_\theta(x) = N_{\theta,\alpha} \left(1 + \frac{\mu^2 b_\alpha}{\sigma^2} \right)^{\frac{1}{\alpha-1}} \left[1 + \frac{b_\alpha x^2}{\sigma^2 + b_\alpha \mu^2} - \frac{2\mu b_\alpha x}{\sigma^2 + b_\alpha \mu^2} \right]^{\frac{1}{\alpha-1}}, \quad (29)$$

where $b_\alpha = \frac{1}{\nu} = \frac{1 - \alpha}{1 + \alpha}$ and $N_{\theta,\alpha} = N_{\sigma,\nu}$. (29) implies that Student distributions form an $\mathbb{M}^{(\alpha)}$ -family with

$$\begin{aligned} h(x) &\equiv 1, & Z(\theta) &= N_{\theta,\alpha} \left(1 + \frac{\mu^2 b_\alpha}{\sigma^2} \right)^{\frac{1}{\alpha-1}}, & w(\theta) &= [w_1(\theta), w_2(\theta)]^\top, & f(x) &= [f_1(x), f_2(x)]^\top, \\ w_1(\theta) &= b_\alpha / (\sigma^2 + b_\alpha \mu^2), & f_1(x) &= x^2, & w_2(\theta) &= -2\mu b_\alpha / (\sigma^2 + b_\alpha \mu^2), & \text{and } f_2(x) &= x. \end{aligned}$$

By Theorem 8, $\bar{f}_1 = \sum_{i=1}^n X_i^2$ and $\bar{f}_2 = \sum_{i=1}^n X_i$ are sufficient statistics for μ and σ with respect to Jones et al. likelihood function.

A. Generalized Fisher-Darmois-Koopman-Pitman theorem for Basu et al. likelihood function

In [25], we characterized the probability distributions that have a fixed number of sufficient statistics with respect to Basu et al. likelihood function (only for the case $\Theta \subset \mathbb{R}$ though). We termed the associated family of probability distributions as $\mathbb{B}^{(\alpha)}$ -family. In the following, we first define the $\mathbb{B}^{(\alpha)}$ -family and then state the results concerning sufficient statistics for the same.

Definition 9: Let h, w, f, Θ and \mathbb{S} be as in Definition 6. A k -parameter $\mathbb{B}^{(\alpha)}$ -family characterized by h, w, f, Θ and \mathbb{S} is defined by $\mathbb{B}^{(\alpha)} = \{p_\theta : \theta \in \Theta\}$, where

$$p_\theta(x) = \begin{cases} [h(x) + F(\theta) + w(\theta)^\top f(x)]^{\frac{1}{\alpha-1}} & \text{for } x \in \mathbb{S} \\ 0 & \text{otherwise} \end{cases} \quad (30)$$

for some $F : \Theta \rightarrow \mathbb{R}$, the normalizing factor.

Remark 3:

- (i) Student distributions fall in $\mathbb{B}^{(\alpha)}$ -family [26, Expl. 1].
- (ii) $\mathbb{B}^{(\alpha)}$ -family is also a generalization of the usual exponential family [25], [26]. A more general form of this family can be found in [14] in connection to the projection theorem of Bregman divergences (c.f. [26]). This family was also studied in [2], [38]. A comparison between $\mathbb{B}^{(\alpha)}$ and $\mathbb{M}^{(\alpha)}$ families can be found in [26, Rem. 2].

The following is the generalized Fisher-Darmois-Koopman-Pitman theorem with respect to Basu et al. likelihood function $L_B^{(\alpha)}$.

Theorem 10: Let $k \leq s < n$. If $[T_1, \dots, T_s]^\top$ is a sufficient statistic for θ with respect to the Basu et al. likelihood function $L_B^{(\alpha)}$, then for $x \in \mathbb{S}$, there exist functions $h : \mathbb{S} \rightarrow \mathbb{R}$, $f_i : \mathbb{S} \rightarrow \mathbb{R}$ and $w_i : \Theta \rightarrow \mathbb{R}$, $i = 1, \dots, s$ such that $p_\theta(x)$ can be expressed as (30).

Proof: The proof for the case $s = k = 1$ can be found in [25, Th. 2]. On the other hand, if $k < s$, say $k = 1$ and $s = 2$, following the same steps as in Theorem 7, one gets (18) using $L_B^{(\alpha)}(x_1^n, \theta)$ in place of $L_J^{(\alpha)}(x_1^n, \theta)$. This, upon taking partial derivative with respect to x_j first, and then with respect to some x_i , $i \neq j$, implies (22). Then, proceeding as in Theorem 7, one can prove the result. ■

The converse of the above theorem is also true.

Theorem 11: Consider a k -parameter $\mathbb{B}^{(\alpha)}$ -family as in Definition 9. Let $T = [T_1, \dots, T_s]^\top$ be s statistics defined on \mathbb{S}^n where $s \geq k$. Then T is sufficient statistic for θ with respect to $L_B^{(\alpha)}$ if there exist real valued functions ψ_1, \dots, ψ_s defined on \mathcal{T} such that $\psi_i(T(x_1^n)) = \bar{f}_i$ for any i.i.d. sample x_1^n and $i = 1, \dots, s$. In particular, $\bar{f} := [\bar{f}_1, \dots, \bar{f}_s]^\top$ is a sufficient statistic for θ with respect to $L_B^{(\alpha)}$.

Proof: See [25, Th. 4]. ■

Example 2: Consider the Student distributions as in Example 1. In [26], we showed that these also form a 2-parameter $\mathbb{B}^{(\alpha)}$ -family. Using Theorem 11, the same statistics as in Example 1 form sufficient statistics for their parameters with respect to Basu et al. likelihood function.

B. Minimal sufficiency with respect to Basu et al. and Jones et al. likelihood functions

In the following, we show that \bar{f}/\bar{h} and \bar{f} are also minimal, respectively, for regular $\mathbb{M}^{(\alpha)}$ and $\mathbb{B}^{(\alpha)}$ families. An $\mathbb{M}^{(\alpha)}$ -family as in Definition 6 is said to be *regular* if the following are satisfied (c.f. [26, Def. 3]).

- (i) The support \mathbb{S} does not depend on the parameter θ .
- (ii) The number of unknown parameters is same as the number of w_i 's, that is, $s = k$.
- (iii) $1, w_1, \dots, w_k$ are linearly independent.
- (iv) f_1, \dots, f_k are linearly independent.

Similarly, a $\mathbb{B}^{(\alpha)}$ -family as in Definition 9 is said to be *regular* if $1, f_1, \dots, f_k$ are linearly independent along with conditions (i)-(iii) above (c.f. [26, Def. 1]).

Theorem 12: The statistics $\bar{f}/\bar{h} = [\bar{f}_1/\bar{h}, \dots, \bar{f}_k/\bar{h}]^\top$ and $\bar{f} = [\bar{f}_1, \dots, \bar{f}_k]^\top$ are minimal sufficient statistics, respectively, for a k -parameter regular $\mathbb{M}^{(\alpha)}$ -family and a k -parameter regular $\mathbb{B}^{(\alpha)}$ -family with respect to Jones et al. and Basu et al. likelihood functions.

Proof: See Sec-B.2 of Appendix. ■

Remark 4: In view of Theorem 12, the sufficient statistics derived in Example 1 for Student distributions are minimal too.

IV. RAO-BLACKWELL THEOREM FOR JONES ET AL. ESTIMATION

Let X_1^n be an i.i.d. random sample drawn according to some $p_\theta \in \Pi$ where $\theta \in \Theta \subset \mathbb{R}$. Let $\hat{\theta} := \hat{\theta}(X_1^n)$ be an unbiased estimator of θ . Let $T := T(X_1^n)$ be a sufficient statistic for θ with respect to the usual likelihood function. Let us consider the conditional expectation $\mathbb{E}_\theta[\hat{\theta}|T]$. It is easy to show from Fisher's definition of sufficiency that $\mathbb{E}_\theta[\hat{\theta}|T]$ is independent of θ [4]. Hence $\mathbb{E}_\theta[\hat{\theta}|T]$ is a function of T alone. Let $\phi(T) := \mathbb{E}_\theta[\hat{\theta}|T]$. Then by Rao-Blackwell theorem, $\phi(T)$ is a *uniformly better* unbiased estimator of θ than $\hat{\theta}$ (see [4], [43]). In other words,

1. $\mathbb{E}_\theta[\phi(T)] = \mathbb{E}_\theta[\hat{\theta}] = \theta$ for all $\theta \in \Theta$, and
2. $\text{Var}_\theta[\phi(T)] \leq \text{Var}_\theta[\hat{\theta}]$ for all $\theta \in \Theta$, and equality holds if and only if $\hat{\theta}$ is a function of the sufficient statistic T .

This implies that the ‘best’ unbiased estimator of θ (the one with least variance) is a function of the sufficient statistic T only, that is, one from the family $\{\psi(T) : \mathbb{E}_\theta[\psi(T)] = \tau(\theta)\}$.

Let us consider an exponential family \mathcal{E} as in (3). $\bar{f} = \frac{1}{n} \sum_{i=1}^n f(X_i)$ is a sufficient statistic for θ with respect to the usual likelihood function [35] and $\mathbb{E}_\theta[\bar{f}] = \mathbb{E}_\theta[f(X)] =: \tau(\theta)$, called the *mean value parameter*. Let $\hat{\theta}$ be any unbiased estimator of $\tau(\theta)$. Then by Rao-Blackwell theorem, we have $\phi(\bar{f}) := \mathbb{E}_\theta[\hat{\theta} | \bar{f}]$ is also an unbiased estimator of $\tau(\theta)$ and has variance not more than that of $\hat{\theta}$. That is,

$$\mathbb{E}_\theta[\phi(\bar{f})] = \tau(\theta) \quad \text{and} \quad \text{Var}_\theta[\phi(\bar{f})] \leq \text{Var}_\theta[\hat{\theta}]. \quad (31)$$

Now consider $\psi(\bar{f}) = \phi(\bar{f}) - \bar{f}$. Then $\mathbb{E}_\theta[\psi(\bar{f})] = \mathbb{E}_\theta[\phi(\bar{f})] - \mathbb{E}_\theta[\bar{f}] = \tau(\theta) - \tau(\theta) = 0$ and

$$\text{Var}_\theta[\phi(\bar{f})] = \text{Var}_\theta[\psi(\bar{f}) + \bar{f}] = \text{Var}_\theta[\psi(\bar{f})] + \text{Var}_\theta[\bar{f}] + 2 \text{Cov}_\theta[\psi(\bar{f}), \bar{f}]. \quad (32)$$

It can be shown, under some mild regularity conditions, that $\text{Cov}_\theta[\psi(\bar{f}), \bar{f}] = 0$. See Sec-C.1 of Appendix for the detailed derivation. Hence (32) reduces to

$$\text{Var}_\theta[\phi(\bar{f})] = \text{Var}_\theta[\psi(\bar{f})] + \text{Var}_\theta[\bar{f}] \geq \text{Var}_\theta[\bar{f}].$$

Thus from (31), for any $\hat{\theta}$ with $\mathbb{E}_\theta[\hat{\theta}] = \mathbb{E}_\theta[\bar{f}]$, we have

$$\text{Var}_\theta[\hat{\theta}] \geq \text{Var}_\theta[\phi(\bar{f})] \geq \text{Var}_\theta[\bar{f}].$$

This shows that the variance of \bar{f} is not more than that of any unbiased estimator of $\tau(\theta)$. Thus we have the following (c.f. [10], [45]).

Proposition 13: Consider a regular exponential family as in (3). Let $\mathbb{E}_\theta[f(X)] = \tau(\theta)$. Then there does not exist any unbiased estimator of $\tau(\theta)$ that has lesser variance than that of \bar{f} . That is, \bar{f} is the best estimator of its expected value.

Further, it can be easily shown that (c.f. [44], [45])

$$\text{Var}_\theta[\bar{f}] = \frac{1}{n} \text{Var}[f(X)] = \frac{[\tau'(\theta)]^2}{n I(\theta)},$$

where $I(\theta)$ is the Fisher information, that is, $I(\theta) = \mathbb{E}_\theta \left[\left(\frac{\partial}{\partial \theta} \log p_\theta(X) \right)^2 \right]$. This and Proposition 13 imply that

$$\text{Var}_\theta[\hat{\theta}] \geq \frac{[\tau'(\theta)]^2}{n I(\theta)}. \quad (33)$$

Equality holds in (33) if and only if $\hat{\theta} = \bar{f}$ almost surely. Notice that (33) is the well-known *Cramér-Rao inequality* [12], [43].

Observe that, in Rao-Blackwell theorem, instead of a sufficient statistic T , if one conditions on some other statistic, then also the resulting estimator has variance not more than that of the given estimator. However, conditioning on any other statistic does not guarantee that the conditional expectation is independent of θ . Also by Fisher-Darmois-Koopman-Pitman theorem, a sufficient statistic with respect to the usual likelihood function exists if and only if the underlying model is exponential [16], [32], [42]. Hence Rao-Blackwell theorem is not applicable outside the exponential family. However in Section III, we showed that there are models that have a fixed number of sufficient statistics with respect to likelihood functions other than log-likelihood. In particular, we showed that the $\mathbb{M}^{(\alpha)}$ -family is one such model which has a fixed number of sufficient statistics with respect to Jones et al. likelihood function (see Theorem 7 and Theorem 8). In the following, we generalize the Rao-Blackwell theorem for Jones et al. likelihood function. Later we will apply this to $\mathbb{M}^{(\alpha)}$ -family and find Rao-Blackwell type ‘‘best’’ estimator. This also helps us find a lower bound for the variance of the estimators of $\mathbb{M}^{(\alpha)}$ -family as an alternative to the usual CRLB.

The Jones et al. likelihood function $L_J^{(\alpha)}$ as in (7) can be re-written as

$$L_J^{(\alpha)}(x_1^n, \theta) = \log \left[\frac{\left(\frac{1}{n} \sum_{i=1}^n p_\theta(x_i)^{\alpha-1} \right)^{\frac{1}{\alpha-1}}}{\|p_\theta\|} \right].$$

Then the deformed probability distribution associated with $L_J^{(\alpha)}$ is given by

$$p_\theta^*(x_1^n) := \frac{\exp[L_J^{(\alpha)}(x_1^n, \theta)]}{\int \exp[L_J^{(\alpha)}(y_1^n, \theta)] dy_1^n} = \frac{\left(\frac{1}{n} \sum_{i=1}^n p_\theta(x_i)^{\alpha-1} \right)^{\frac{1}{\alpha-1}}}{\int \left(\frac{1}{n} \sum_{i=1}^n p_\theta(y_i)^{\alpha-1} \right)^{\frac{1}{\alpha-1}} dy_1^n}. \quad (34)$$

Observe that when $\alpha \rightarrow 1$, $L_J^{(\alpha)}(x_1^n; \theta)$ coincides with the log-likelihood function $L(x_1^n; \theta)$. Hence, in this case, $p_\theta^*(x_1^n) = p_\theta(x_1^n)$.

Let T be a statistic of the sample x_1^n . Then, from (10), we have

$$p_{\theta_{X_1^n|T}}^*(x_1^n|t) = \begin{cases} \frac{p_\theta^*(x_1^n)}{q_\theta^*(t)} & \text{if } x_1^n \in A_t \\ 0 & \text{otherwise.} \end{cases}$$

Observe that, as $\alpha \rightarrow 1$, the above coincides with the usual conditional distribution of x_1^n given $T = t$.

Let $\hat{\theta} := \hat{\theta}(X_1^n)$ be an unbiased estimator of $\tau^*(\theta) := \mathbb{E}_\theta^*[\hat{\theta}]$, where $\mathbb{E}_\theta^*[\cdot \cdot \cdot]$ denotes the expectation with respect to p_θ^* . Let $T := T(X_1^n)$ be a sufficient statistic for θ with respect to Jones et al. likelihood function $L_J^{(\alpha)}$. Let us consider the conditional expectation

$$\mathbb{E}_\theta^*[\hat{\theta}|T = T(x_1^n)] = \int \hat{\theta}(y_1^n) p_\theta^*(y_1^n|T(x_1^n)) dy_1^n. \quad (35)$$

In the foregoing, we first establish some properties of (35).

Proposition 14: The conditional expectation $\mathbb{E}_\theta^*[\hat{\theta}|T]$ is independent of θ .

Proof: See Sec-C.2 of Appendix. ■

Let

$$\mathbb{E}_\theta^*[\hat{\theta}|T = T(x_1^n)] =: \phi^*(T(x_1^n)).$$

Let us consider the expectation of $\phi^*(T)$ with respect to p_θ^* . That is,

$$\mathbb{E}_\theta^*[\phi^*(T(X_1^n))] = \int p_\theta^*(x_1^n) \phi^*(T(x_1^n)) dx_1^n.$$

Then we have the following.

Proposition 15: $\phi^*(T)$ is unbiased for $\tau^*(\theta)$, that is, $\mathbb{E}_\theta^*[\phi^*(T(X_1^n))] = \mathbb{E}_\theta^*[\hat{\theta}(X_1^n)] = \tau^*(\theta)$.

Proof: See Sec-C.3 of Appendix. ■

Thus, if we have an unbiased estimator of $\tau^*(\theta)$ and a sufficient statistic with respect to Jones et al. likelihood function, we can always get another unbiased estimator for $\tau^*(\theta)$ that is based only on the sufficient statistic. Let us now consider the following two classes of estimators based on the sufficient statistic T :

$$\begin{aligned} \mathcal{A} &= \{\mathbb{E}_\theta^*[\hat{\theta}|T] : \hat{\theta} \text{ is an unbiased estimator of } \tau^*(\theta), \text{ that is, } \mathbb{E}_\theta^*[\hat{\theta}] = \tau^*(\theta)\}, \\ \mathcal{B} &= \{\psi(T) : \mathbb{E}_\theta^*[\psi(T)] = \tau^*(\theta)\}. \end{aligned} \quad (36)$$

Proposition 16: The above two classes of estimators are the same, that is $\mathcal{A} = \mathcal{B}$.

Proof: See Sec-C.4 of Appendix. ■

The above proposition suggests that any unbiased estimator of $\tau^*(\theta)$ that is a function of the sufficient statistic T is of the form $\mathbb{E}_\theta^*[\hat{\theta}|T]$, where $\hat{\theta}$ is an unbiased estimator of $\tau^*(\theta)$. In the following, we compare the variance of an estimator of the form (36) and any other unbiased estimator of $\tau^*(\theta)$.

Lemma 17: Let $\hat{\theta}$ be any unbiased estimator of $\tau^*(\theta)$ and $\phi^*(T) = \mathbb{E}_\theta^*[\hat{\theta}|T]$. Let $\psi^*(T) \in \mathcal{B}$. Then

$$\text{Var}_\theta^*[\hat{\theta}] = \mathbb{E}_\theta^*[(\hat{\theta} - \psi^*(T))^2] + \text{Var}_\theta^*[\psi^*(T)] + 2\mathbb{E}_\theta^*[\psi^*(T)\{\phi^*(T) - \psi^*(T)\}]. \quad (37)$$

Proof: See Sec-C.5 of Appendix. ■

The following (also the main result of this section) is a generalization of Rao-Blackwell theorem for Jones et al. likelihood function.

Theorem 18: Let $\Pi = \{p_\theta : \theta \in \Theta \subset \mathbb{R}\}$ be a family of probability distributions with common support $\mathbb{S} \subset \mathbb{R}$. Consider the deformed family of probability distributions $\Pi^* = \{p_\theta^*\}$ of Π , where p_θ^* is as in (34). Let $\hat{\theta}$ be an unbiased estimator for its expected value with respect to p_θ^* . Let $\tau^*(\theta) := \mathbb{E}_\theta^*[\hat{\theta}]$. Suppose T is a sufficient statistic for θ with respect to the Jones et al. likelihood function. Let $\phi^*(T) := \mathbb{E}_\theta^*[\hat{\theta}|T]$. Then $\phi^*(T)$ is a uniformly better estimator of $\tau^*(\theta)$ than $\hat{\theta}$, that is,

- (a) $\mathbb{E}_\theta^*[\phi^*(T)] = \tau^*(\theta)$,
- (b) $\text{Var}_\theta^*[\phi^*(T(X_1^n))] \leq \text{Var}_\theta^*[\hat{\theta}(X_1^n)]$, and equality holds if and only if $\hat{\theta}$ is a function of the sufficient statistic T .

Proof: See Sec-C.6 of Appendix. ■

Theorem 18 implies that $\phi^*(T)$ is *uniformly a better estimator* of $\tau^*(\theta)$ than $\hat{\theta}$. That means the *best unbiased estimator* of $\tau^*(\theta)$ is a function of the sufficient statistic T . The following result asserts that the estimator with minimum variance (best estimator) is unique.

Proposition 19: The best unbiased estimator for $\tau^*(\theta)$ with respect to p_θ^* is unique.

Proof: See Sec-C.7 of Appendix. ■

In the following, we apply the above results to the family of Bernoulli distributions.

A. *Example:*

Observe that the family of Bernoulli distributions can be written as

$$p_\theta(x) = (1 - \theta)[1 + x \cdot (2\theta - 1)/(1 - \theta)] \quad \text{for } x = 0, 1,$$

where $\theta \in (0, 1)$ is the unknown parameter. Thus Bernoulli distributions form an $\mathbb{M}^{(\alpha)}$ -family with $\alpha = 2$, $h(x) \equiv 1$, $Z(\theta) = 1 - \theta$, $w(\theta) = (2\theta - 1)/(1 - \theta)$ and $f(x) = x$. By Theorem 8, $\sum_{i=1}^n X_i$ is a sufficient statistic for θ with respect to the Jones et al. likelihood function.

The Jones et al. likelihood function for Bernoulli distributions is

$$L_{cs}(x_1^n, \theta) = \log \left[\frac{\frac{1}{n} \sum_{i=1}^n p_\theta(x_i)}{(\mathbb{E}_\theta[p_\theta(X)])^{1/2}} \right] = \log \left[\frac{1 + \bar{x} \cdot (2\theta - 1)/(1 - \theta)}{(1 + \theta(2\theta - 1)/(1 - \theta))^{1/2}} \right].$$

The associated deformed probability distribution is given by

$$p_\theta^*(x_1^n) = \frac{\exp[L_{cs}(x_1^n, \theta)]}{\sum_{y_1^n} \exp[L_{cs}(y_1^n, \theta)]} = \frac{1 + \bar{x} \cdot (2\theta - 1)/(1 - \theta)}{\sum_{y_1^n} [1 + \bar{y} \cdot (2\theta - 1)/(1 - \theta)]} = N(\theta) \left[1 + \bar{x} \cdot \frac{2\theta - 1}{1 - \theta} \right],$$

where

$$\begin{aligned}
N(\theta)^{-1} &:= \sum_{y_1^n} \left[1 + \bar{y} \cdot \frac{2\theta - 1}{1 - \theta} \right] = 2^n + \left[\frac{(2\theta - 1)}{(1 - \theta)} \cdot \frac{1}{n} \sum_{i=1}^n i \cdot \binom{n}{i} \right] \\
&= 2^n + \left[\frac{(2\theta - 1)}{n(1 - \theta)} \cdot (n2^{n-1}) \right] \\
&= \frac{2^{n-1}}{1 - \theta}.
\end{aligned}$$

Also

$$p_\theta^* \left(X_1^n = x_1^n \mid T = \sum_{i=1}^n x_i \right) = \frac{1 + \bar{x} \cdot (2\theta - 1)/(1 - \theta)}{\sum_{y_1^n \in A_{\sum x_i}} [1 + \bar{y} \cdot (2\theta - 1)/(1 - \theta)]} = \frac{1}{|A_{\sum x_i}|} = \frac{1}{\binom{n}{\sum x_i}}.$$

Let $\hat{\theta} = X_1$. Then

$$\begin{aligned}
\tau^*(\theta) = \mathbb{E}_\theta^*[\hat{\theta}] &= \sum_{x_1^n} x_1 \cdot p_\theta^*(x_1^n) = \sum_{x_1} N(\theta) \left[2^{n-1} + \frac{2\theta - 1}{n(1 - \theta)} \sum_{i=1}^n (x_1 + i) \binom{n-1}{i} \right] \\
&= N(\theta) \cdot 2^{n-1} \left[1 + \frac{2\theta - 1}{n(1 - \theta)} \cdot \left(1 + \frac{n+1}{2} \right) \right] \\
&= \frac{\theta}{n} + \frac{n-1}{2n}.
\end{aligned}$$

The variance of $\hat{\theta}$ with respect to p_θ^* is given by

$$\text{Var}_\theta^*[\hat{\theta}] = \mathbb{E}_\theta^*[X_1^2] - (\mathbb{E}_\theta^*[X_1])^2 = \tau^*(\theta) - \tau^*(\theta)^2.$$

Let us consider the generalized Rao-Blackwell estimator of $\tau^*(\theta)$. Recall that $\sum_{i=1}^n X_i$ is a sufficient statistic for θ with respect to Jones et al. likelihood function.

$$\begin{aligned}
\phi^*(T(x_1^n)) &= \phi^* \left(\sum_{i=1}^n x_i \right) = \sum_{y_1^n} \hat{\theta}(y_1^n) \cdot p_\theta^* \left(y_1^n \mid T = \sum_{i=1}^n x_i \right) \\
&= \sum_{y_1^n \in A_{\sum x_i}} y_1 \cdot p_\theta^* \left(y_1^n \mid T = \sum_{i=1}^n x_i \right) \\
&= \binom{n}{\sum x_i}^{-1} \cdot \sum_{y_1^n \in A_{\sum x_i}} y_1 \\
&= \binom{n}{\sum x_i}^{-1} \cdot \sum_{y_1^n \in A_{(\sum_{i=1}^n x_i - 1)}} 1 \\
&= \binom{n-1}{\sum x_i - 1} / \binom{n}{\sum x_i} \\
&= \frac{1}{n} \sum_{i=1}^n x_i.
\end{aligned}$$

Thus $\phi^*(T) = \frac{1}{n} \sum_{i=1}^n X_i$ is uniformly better estimator for $\tau^*(\theta)$ than X_1 . It can also be shown that

$$\text{Var}_\theta^*[\phi^*(T)] = \frac{\left[\frac{\partial}{\partial \theta} \tau^*(\theta) \right]^2}{I_{\alpha, n}^*(\theta)},$$

where $I_{\alpha,n}^*(\theta)$ is the inverse of the asymptotic variance of Jones et al. estimator for $p_\theta^*(X_1^n)$ [30]. This shows that the Rao-Blackwell estimator derived from sufficient statistic attains asymptotic variance of Jones et al. estimation. In the following section we elucidate this for a general $\mathbb{M}^{(\alpha)}$ -family.

V. EFFICIENCY OF RAO-BLACKWELL ESTIMATORS OF POWER-LAW FAMILY

In this section, we shall show the efficiency of Rao-Blackwell estimators for a power-law family. For simplicity, we shall consider a regular $\mathbb{M}^{(\alpha)}$ -family with $h \equiv 1$. That is, for $x \in \mathbb{S}$,

$$p_\theta(x) = Z(\theta)[1 + w(\theta)f(x)]^{\frac{1}{\alpha-1}}. \quad (38)$$

Then

$$\left(\frac{1}{n} \sum_{i=1}^n p_\theta(x_i)^{\alpha-1}\right)^{\frac{1}{\alpha-1}} = \left(\frac{1}{n} \sum_{i=1}^n Z(\theta)^{\alpha-1}[1 + w(\theta)f(x_i)]\right)^{\frac{1}{\alpha-1}} = Z(\theta)[1 + w(\theta)\bar{f}]^{\frac{1}{\alpha-1}}.$$

Thus, using (34), we get

$$p_\theta^*(x_1^n) = \frac{[1 + w(\theta)\bar{f}]^{\frac{1}{\alpha-1}}}{\int [1 + w(\theta)\bar{f}]^{\frac{1}{\alpha-1}} dx_1^n} = N(\theta)[1 + w(\theta)\bar{f}]^{\frac{1}{\alpha-1}}, \quad (39)$$

where $\bar{f} := \frac{1}{n} \sum_{i=1}^n f(x_i)$ and $N(\theta)^{-1} := \int [1 + w(\theta)\bar{f}]^{\frac{1}{\alpha-1}} dx_1^n$ is the normalizing constant. Comparing this with (11), we see that (39) also forms an $\mathbb{M}^{(\alpha)}$ -family with $h \equiv 1$ and $f(x_1^n) = \bar{f}$. Recall that, \bar{f} is a sufficient statistic for $\mathbb{M}^{(\alpha)}$ with respect to Jones et al. likelihood function (Theorem 8). Let $\tau^*(\theta) := \mathbb{E}_\theta^*[\bar{f}]$. In the following, we show that \bar{f} is the best estimator of $\tau^*(\theta)$. That is, variance of any unbiased estimator of $\tau^*(\theta)$ is bounded below by the variance of \bar{f} with respect to p_θ^* .

Let $\hat{\theta}$ be an unbiased estimator of $\tau^*(\theta)$ with respect to p_θ^* . Then by Theorem 18, $\phi^*(\bar{f}) = \mathbb{E}_\theta^*[\hat{\theta}|\bar{f}]$ is also an unbiased estimator of $\tau^*(\theta)$ and

$$\text{Var}_\theta^*[\hat{\theta}] \geq \text{Var}_\theta^*[\phi^*(\bar{f})]. \quad (40)$$

Let us consider $\psi^*(\bar{f}) := \phi^*(\bar{f}) - \bar{f}$. Then $\mathbb{E}_\theta^*[\psi^*(\bar{f})] = \tau^*(\theta) - \tau^*(\theta) = 0$. Also

$$\text{Var}_\theta^*[\phi^*(\bar{f})] = \text{Var}_\theta^*[\psi^*(\bar{f}) + \bar{f}] = \text{Var}_\theta^*[\psi^*(\bar{f})] + \text{Var}_\theta^*[\bar{f}] + 2 \text{Cov}_\theta^*[\psi^*(\bar{f}), \bar{f}]. \quad (41)$$

In Sec-D.1 of Appendix, we have shown that, if $w(\theta) > 0$ for $\theta \in \Theta$, then $\text{Cov}_\theta^*[\psi^*(\bar{f}), \bar{f}] \geq 0$. Using this in (41), we get

$$\text{Var}_\theta^*[\phi^*(\bar{f})] = \text{Var}_\theta^*[\psi^*(\bar{f})] + \text{Var}_\theta^*[\bar{f}] + 2 \text{Cov}_\theta^*[\psi^*(\bar{f}), \bar{f}] \geq \text{Var}_\theta^*[\bar{f}]. \quad (42)$$

Using (40) and (42), we get

$$\text{Var}_\theta^*[\hat{\theta}] \geq \text{Var}_\theta^*[\bar{f}], \quad (43)$$

where equality holds if and only if $\hat{\theta} = \bar{f}$ almost surely. Thus we have the following.

Theorem 20: Consider an $\mathbb{M}^{(\alpha)}$ -family as in (38) with $w(\theta) > 0$ for $\theta \in \Theta$. Let $\tau^*(\theta) = \mathbb{E}_\theta^*[\bar{f}]$. Then the variance of any unbiased estimator of $\tau^*(\theta)$ with respect to p_θ^* is not lower than that of \bar{f} . That is, \bar{f} is the best estimator of its expected value with respect to p_θ^* .

We now show that $\text{Var}_\theta^*[\bar{f}]$, indeed, equals to the asymptotic variance of Jones et al. estimator for the model p_θ^* . The Jones et al. asymptotic variance for p_θ^* is given by (c.f. [30])

$$\frac{1}{I_{\alpha,n}^*(\theta)} := \frac{\text{Var}_\theta^*[p_\theta^*(X_1^n)^{\alpha-1} s^*(X_1^n, \theta)]}{\text{Cov}_\theta^*[s^*(X_1^n, \theta), p_\theta^*(X_1^n)^{\alpha-1} s^*(X_1^n, \theta)]^2}, \quad (44)$$

where $s^*(x_1^n, \theta) = \frac{\partial}{\partial \theta} \log p_\theta^*(x_1^n)$. It can be easily seen that $I_{\alpha, n}^*(\theta)$ coincides with the usual Fisher information $I_n^*(\theta)$ for $\alpha = 1$, where $I_n^*(\theta) = \text{Var}_\theta^*[\frac{\partial}{\partial \theta} \log p_\theta^*(X_1^n)]$. Observe that (39) can be re-written as

$$p_\theta^*(x_1^n) = N(\theta)[1 + w(\theta)\bar{f}]^{\frac{1}{\alpha-1}} = [N(\theta)^{\alpha-1} + N(\theta)^{\alpha-1}w(\theta)\bar{f}]^{\frac{1}{\alpha-1}}.$$

For convenience, let $F(\theta) := N(\theta)^{\alpha-1}$ and $\tilde{w}(\theta) := N(\theta)^{\alpha-1}w(\theta)$. Then

$$p_\theta^*(x_1^n) = [F(\theta) + \tilde{w}(\theta)\bar{f}]^{\frac{1}{\alpha-1}}$$

and hence

$$\frac{\partial}{\partial \theta} p_\theta^*(x_1^n) = \frac{1}{\alpha-1} p_\theta^*(x_1^n)^{2-\alpha} [F'(\theta) + \tilde{w}'(\theta)\bar{f}].$$

Then

$$\text{Var}_\theta^*[p_\theta^*(X_1^n) s^*(X_1^n, \theta)] = \left(\frac{1}{\alpha-1}\right)^2 [\tilde{w}'(\theta)]^2 \text{Var}_\theta^*[\bar{f}], \quad (45)$$

and

$$\text{Cov}_\theta^*[s^*(X_1^n, \theta), p_\theta^*(X_1^n)^{\alpha-1} s^*(X_1^n, \theta)] = \frac{1}{\alpha-1} \tilde{w}'(\theta) \frac{\partial}{\partial \theta} \tau^*(\theta). \quad (46)$$

The detailed derivations of (45) and (46) can be found in Sec-D.2 of the Appendix. Using these values, (44) becomes

$$\frac{1}{I_{\alpha, n}^*(\theta)} = \frac{\text{Var}_\theta^*[\bar{f}]}{\left[\frac{\partial}{\partial \theta} \mathbb{E}_\theta^*[\bar{f}]\right]^2},$$

that is,

$$\frac{\left[\frac{\partial}{\partial \theta} \mathbb{E}_\theta^*[\bar{f}]\right]^2}{I_{\alpha, n}^*(\theta)} = \frac{\left[\frac{\partial}{\partial \theta} \tau^*(\theta)\right]^2}{I_{\alpha, n}^*(\theta)} = \text{Var}_\theta^*[\bar{f}].$$

This, together with (43), yields the following result.

Corollary 21: Consider an $\mathbb{M}^{(\alpha)}$ -family as in Theorem 20. Let $\tau^*(\theta) = \mathbb{E}_\theta^*[\bar{f}]$. Let $\hat{\theta}$ be any unbiased estimator of $\tau^*(\theta)$ with respect to p_θ^* . Then

$$\text{Var}_\theta^*[\hat{\theta}] \geq \frac{\left[\frac{\partial}{\partial \theta} \tau^*(\theta)\right]^2}{I_{\alpha, n}^*(\theta)}. \quad (47)$$

Equality holds in (47) if and only if $\hat{\theta} = \bar{f}$ almost surely.

Remark 5:

(i) Observe that

$$\text{Cov}_\theta^*[s^*(X_1^n, \theta), p_\theta^*(X_1^n)^{\alpha-1} s^*(X_1^n, \theta)]^2 \leq \text{Var}_\theta^*[p_\theta^*(X_1^n)^{\alpha-1} s^*(X_1^n, \theta)] \text{Var}_\theta^*[s^*(X_1^n, \theta)],$$

by Cauchy-Schwarz inequality. This implies that

$$\frac{1}{I_n^*(\theta)} \leq \frac{1}{I_{\alpha, n}^*(\theta)}.$$

That means the bound given by Corollary 21 is sharper than the usual CRLB for any unbiased estimator.

(ii) In view of (9) and Remark 1(i), we see that (47) in Corollary 21 is essentially a generalization of the usual Cramér-Rao inequality for an exponential family.

Example 3: Consider the Student distributions with location parameter $\mu = 0$, scale parameter σ^2 (unknown) and degrees of freedom parameter $\nu \in (2, \infty)$:

$$p_\sigma(x) = N_{\nu,\sigma} \left[1 + \frac{x^2}{\nu\sigma^2} \right]^{-\frac{\nu+1}{2}}, \quad (48)$$

where $N_{\nu,\sigma}$ is the normalizing constant. Let $\alpha = 1 - \frac{2}{\nu+1}$, $b_\alpha = 1/\nu$ and $N_{\alpha,\sigma} = N_{\nu,\sigma}$. Then (48) can be re-written as a one parameter regular $\mathbb{M}^{(\alpha)}$ -family with $h \equiv 1$, $w(\sigma) = b_\alpha/\sigma^2 > 0$ and $f(x) = x^2$ [26]. By Theorem 20, $\overline{X^2}$ is the best estimator for $\mathbb{E}_\sigma^*[\overline{X^2}]$. We shall now find $\mathbb{E}_\sigma^*[\overline{X^2}]$ explicitly. Using (39), we have

$$p_\sigma^*(x_1^n) = Z_{\alpha,\sigma} \left[1 + (b_\alpha/\sigma^2)\overline{x^2} \right]^{\frac{1}{\alpha-1}},$$

where $Z_{\alpha,\sigma}^{-1} := \int [1 + (b_\alpha/\sigma^2)y^2]^{\frac{1}{\alpha-1}} dy_1^n = \sigma^n \cdot H_n(\alpha)$, provided $\alpha > 1 - \frac{2}{n}$. Here

$$H_n(\alpha) = \begin{cases} \frac{(\pi b_\alpha)^{\frac{1}{2}} (n\pi/b_\alpha)^{\frac{n}{2}} \Gamma\left(\frac{1+\alpha}{2(1-\alpha)}\right) A_n(\alpha)^k}{\Gamma(n/2)\Gamma\left(\frac{1}{1-\alpha}\right)} & \text{if } n = 2k + 1, k \geq 0, \\ \frac{(n\pi/b_\alpha)^{\frac{n}{2}} (1-\alpha) A_n(\alpha)^{k-1}}{\alpha \Gamma(n/2)} & \text{if } n = 2k, k \geq 1, \end{cases}$$

where $A_n(\alpha) := \frac{(1-\alpha)(n-2)}{2-n(1-\alpha)}$. Let us now denote $B_n(\alpha) = \frac{n(1-\alpha)}{2\alpha-n(1-\alpha)}$. Then $\mathbb{E}_\sigma^*[\overline{X^2}]$ equals to $\frac{A_n(\alpha)^{2-k} B_n(\alpha)^{k-1}}{b_\alpha} \sigma^2$ when $n = 2k$, and $\frac{A_n(\alpha)^{1-k} B_n(\alpha)^k}{b_\alpha} \sigma^2$ when $n = 2k + 1$, provided $\alpha > \frac{n}{n+2}$. Hence the generalized CRLB is given by

$$\begin{cases} [\{A_n(\alpha)^{2-k}(C_n(\alpha)^k - A_n(\alpha)^{-k} B_n(\alpha)^{2k})\}/b_\alpha^2] \sigma^4 & \text{if } n = 2k + 1, \\ [\{A_n(\alpha)^{3-k}(C_n(\alpha)^{k-1} - A_n(\alpha)^{1-k} B_n(\alpha)^{(2k-1)})\}/b_\alpha^2] \sigma^4 & \text{if } n = 2k, \end{cases}$$

where $C_n(\alpha) := \frac{(1-\alpha)(n+2)}{2\alpha+(n+2)(\alpha-1)}$ and $\alpha > \frac{n+2}{n+4}$.

Remark 6: Observe that Theorem 20 can not be applied, for example, to Student distributions with unknown location parameter (scale parameter known), since they do not form a regular $\mathbb{M}^{(\alpha)}$ -family [26, Rem. 2(c)]. However, in the following, we address this problem by deriving a Rao-Blackwell type theorem for Basu et al. likelihood function (8). Note that, for this model, Basu et al. estimator and Jones et al. estimator are the same [24], [26].

A. Analogous results for Basu et al. likelihood function

Consider the Basu et al. likelihood function $L_B^{(\alpha)}$ as in (8). The associated deformed probability distribution is given by

$$p_\theta^*(x_1^n) := \frac{\exp[L_B^{(\alpha)}(x_1^n; \theta)]}{\int \exp[L_B^{(\alpha)}(y_1^n; \theta)] dy_1^n} = \frac{\exp\left[\frac{\alpha}{n(\alpha-1)} \sum_{i=1}^n p_\theta(x_i)^{\alpha-1}\right]}{\int \exp\left[\frac{\alpha}{n(\alpha-1)} \sum_{i=1}^n p_\theta(y_i)^{\alpha-1}\right] dy_1^n}. \quad (49)$$

Recall that, $\mathbb{B}^{(\alpha)}$ -families are the ones that have a fixed number of sufficient statistics with respect to Basu et al. likelihood function (Theorem 10). Consider a $\mathbb{B}^{(\alpha)}$ -family as in (30). Then, using (49), we have

$$p_\theta^*(x_1^n) = \exp\left[\frac{\alpha}{\alpha-1} \bar{h} + \tilde{F}(\theta) + \frac{\alpha}{\alpha-1} w(\theta) \bar{f}\right], \quad (50)$$

where $\tilde{F}(\theta) = -\log \int \exp\left[\frac{\alpha}{\alpha-1} (\bar{h} + w(\theta) \bar{f})\right] dx_1^n$, is the normalizing constant. Observe that p_θ^* forms a 1-parameter exponential family. Then we can derive the following result concerning the best estimator for the parameter, analogous to Theorem 20 and Corollary 21.

Theorem 22: Consider a regular $\mathbb{B}^{(\alpha)}$ -family as in (30). Let $\mathbb{E}_\theta^*[f] = \tau(\theta)$. Then \bar{f} is the best estimator of its expected value with respect to p_θ^* . Furthermore, if $\hat{\theta}$ is any estimator such that $\mathbb{E}_\theta^*[\hat{\theta}] = \tau(\theta)$, then

$$\text{Var}_\theta^*[\hat{\theta}] \geq \frac{[\tau'(\theta)]^2}{I_n^*(\theta)}. \quad (51)$$

Proof: See Sec-D.3 of Appendix. ■

Remark 7: Notice that the bound in (51) is given by the usual Fisher information (of course, applied to the model p_θ^*) unlike the one in Corollary 21 where the bound is given by the asymptotic variance of Jones et al. estimators.

We now apply the above result to find best estimator for the mean parameter of a Student distribution. Consider the Student distributions with degrees of freedom parameter $\nu \in (2, \infty)$, location parameter μ and scale parameter $\sigma^2 = 1$:

$$p_\mu(x) = N_\nu \left[1 + \frac{1}{\nu}(x - \mu)^2 \right]^{-\frac{\nu+1}{2}}, \quad (52)$$

where N_ν is the normalizing constant. Letting $\alpha = 1 - \frac{2}{\nu+1}$, $b_\alpha = 1/\nu$ and $N_\alpha = N_\nu$, (52) forms a regular $\mathbb{B}^{(\alpha)}$ -family as in (30) with $h(x) = N_\alpha^{\alpha-1} b_\alpha x^2$, $F(\mu) = (1 + b_\alpha \mu^2) N_\alpha^{\alpha-1}$, $w(\mu) = -2\mu N_\alpha^{\alpha-1} b_\alpha$ and $f(x) = x$. This implies that \bar{X} is a sufficient statistic for μ with respect to the Basu et al. likelihood function [25, Th. 4] and \bar{X} is the best estimator for $\mathbb{E}_\mu^*[\bar{X}]$ (Theorem 22).

It can also be shown, in this case, that

$$p_\mu^*(x_1^n) = \exp \left[\frac{\alpha}{\alpha-1} N_\alpha^{\alpha-1} b_\alpha \bar{x}^2 + F_\alpha(\mu) - \frac{2\alpha}{\alpha-1} N_\alpha^{\alpha-1} b_\alpha \mu \bar{x} \right],$$

where $F_\alpha(\mu) := \frac{\alpha}{\alpha-1} N_\alpha^{\alpha-1} b_\alpha \mu^2 - \frac{n}{2} \log \left[\pi - \frac{\alpha}{n(\alpha-1)} N_\alpha^{\alpha-1} b_\alpha \right]$,

$$\mathbb{E}_\mu^*[\bar{X}] = - \left(\frac{\alpha-1}{\alpha} \right) \frac{\frac{2\alpha}{(1-\alpha)} N_\alpha^{\alpha-1} b_\alpha \mu}{w'(\mu)} = \mu \quad \text{and} \quad \text{Var}_\mu^*[\bar{X}] = \frac{[\tau'(\mu)]^2}{I_n^*(\mu)} = \frac{(1-\alpha)}{2\alpha b_\alpha N_\alpha^{\alpha-1}}.$$

VI. SUMMARY AND CONCLUDING REMARKS

Fisher introduced the concept of sufficient statistics in connection with parameter estimation problems using which one can estimate the unknown parameters of a population without knowing the entire sample [20]. The classical notion of sufficiency is based on the usual likelihood function and is useful only when the underlying probability distribution is from a (regular) exponential family. If the distribution is not from an exponential family, it may not have sufficient statistics (fixed in number independent of sample size) [16], [32], [42]. However, such distributions may also have sufficient statistics with respect to a likelihood function different from the usual one [17], [18]. In this paper, we considered two families of likelihood functions, namely Basu et al. and Jones et al. likelihood functions, that arise in distance-based robust inference methods and are also generalizations of the usual likelihood function. We characterized the family of probability distributions that have a fixed number of sufficient statistics with respect to these likelihood functions. This family generalizes the usual exponential family and has a power law. Student distributions fall in this family. We also extended the notion of minimal sufficiency in this connection and found minimal sufficient statistics for the power-law families.

An important application of sufficient statistics is that it helps find an estimator uniformly better than the given one. This result is known as Rao-Blackwell theorem. It essentially proves that a sufficient statistic is the best estimator for its expected value. In this paper, we generalized Rao-Blackwell theorem when one has sufficient statistics with respect to Basu et al. or Jones et al. likelihood function. This helps us find the ‘best’ estimator for the parameters of a power-law family and hence for Student distributions. Consequently, we derived a lower bound for the variance of estimators of a deformed form of power-law family, p_θ^* . In case of Jones et al. likelihood function, the bound turns out to be its asymptotic variance

[30] (with respect to p_θ^*). This bound is sharper than the usual CRLB (See Corollary 21). Recall that the usual CRLB equals the asymptotic variance of MLE.

We now make some remarks on the deformed probability distribution p_θ^* . Recall that, it is the factorization theorem (Halmos and Savage [27] in the classical case and Proposition 2 in the general case) that makes it clear that the notion of sufficiency is with respect to a likelihood function. However, the generalized notion of sufficiency is defined using the deformed probability distribution p_θ^* . The equivalence of analogous factorization theorem and Fisher's definition of sufficiency, shows the uniqueness of the usage of p_θ^* in the definition of generalized Fisher sufficiency. This fact was further evidenced in the case of Jones et al. likelihood function, where the lower bound of variance of estimators for power-law families equals the asymptotic variance of the Jones et al. estimator for p_θ^* . However, for Basu et al. likelihood function though, the bound is given by the usual CRLB, computed with respect to p_θ^* , not the asymptotic variance of the Basu et al. estimator (Theorem 22). This is surprising as one would expect the bound to equal the Basu et al. asymptotic variance ([3, Th. 2]), similar to the case of Jones et al. likelihood function. This will be the focus of one of our forthcoming works. In any case, such a distinct behavior of Basu et al. likelihood function (from Jones et al. likelihood function) was also noticed in the study of geometry of associated divergences in connection with projection theorems for families of probability distributions determined by linear constraints. While the geometry of the Basu et al. divergence is similar to that of the Kullback-Leibler divergence, the same of Jones et al. divergence was vastly different from either of these two [34].

ACKNOWLEDGEMENTS

Atin Gayen is supported by an INSPIRE fellowship of the Department of Science and Technology, Govt. of India.

APPENDIX

A. PROOF OF RESULTS OF SECTION 2

A.1. Proof of Proposition 2:

Let T be a sufficient statistic for θ with respect to L_G . Let x_1^n and y_1^n be two samples such that $T(x_1^n) = T(y_1^n)$. Then from generalized Koopman's definition of sufficiency, we have $[L_G(x_1^n; \theta) - L_G(y_1^n; \theta)]$ is independent of θ . Let us define a relation ' \sim ' on the set of all sample points of length n by $x_1^n \sim y_1^n$ if and only if $T(x_1^n) = T(y_1^n)$. Then ' \sim ' is an equivalence relation. Let us denote the equivalence classes by \mathbb{S}^t for each $t \in \mathcal{T} = \{r : r = T(x_1^n) \text{ for some } x_1^n \in \mathbb{S}^n\}$. For each equivalence class \mathbb{S}^t , designate an element $x_{1,t}^n \in \mathbb{S}^t$.

Let x_1^n be a sample point such that $T(x_1^n) = t^* \in \mathcal{T}$. Then $x_1^n \in \mathbb{S}^{t^*}$ and $T(x_1^n) = T(x_{1,t^*}^n)$. This implies that $[L_G(x_1^n; \theta) - L_G(x_{1,t^*}^n; \theta)]$ is independent of θ by hypothesis. Let $v(x_1^n) := [L_G(x_1^n; \theta) - L_G(x_{1,t^*}^n; \theta)]$. Then

$$L_G(x_1^n; \theta) = L_G(X_{1,t^*}^n; \theta) + v(x_1^n) = u(\theta, t^*) + v(x_1^n),$$

where $u(\theta, t) := L_G(X_{1,t}^n; \theta)$ for each $t \in \mathcal{T}$.

Conversely, let us assume that there exist two functions u and v such that

$$L_G(x_1^n; \theta) = u(\theta, T(x_1^n)) + v(x_1^n)$$

for all sample points x_1^n and for all $\theta \in \Theta$. Thus for any two sample points x_1^n and y_1^n with $T(x_1^n) = T(y_1^n)$, we have

$$\begin{aligned} L_G(x_1^n; \theta) - L_G(y_1^n; \theta) &= u(\theta, T(x_1^n)) + v(x_1^n) - u(\theta, T(y_1^n)) - v(y_1^n) \\ &= v(x_1^n) - v(y_1^n). \end{aligned}$$

This implies that $[L_G(x_1^n; \theta) - L_G(y_1^n; \theta)]$ is independent of θ . By generalized Koopman's definition of sufficiency, T is a sufficient statistic for θ with respect to L_G . This completes the proof. ■

A.2. Proof of Proposition 3:

Let T be sufficient for θ with respect to L_G . Then, from Proposition 2.2, there exist functions u and v such that

$$L_G(x_1^n, \theta) = u(T(x_1^n), \theta) + v(x_1^n), \quad (53)$$

for all $x_1^n \in \mathbb{S}^n$. We need to only show $p_{\theta_{x_1^n|T}}^*(x_1^n|t)$ is independent of θ when $t = T(x_1^n)$. Observe that if $t \neq T(x_1^n)$, $p_{\theta_{x_1^n|T}}^*(x_1^n|t) = 0$ which is independent of θ . Using (53), we have

$$\begin{aligned} p_{\theta_{x_1^n|T}}^*(x_1^n|t) &= \frac{p_{\theta}^*(x_1^n)}{\int_{A_T(x_1^n)} p_{\theta}^*(y_1^n) dy_1^n} = \frac{\exp[L_G(x_1^n; \theta)]}{\int_{A_T(x_1^n)} \exp[L_G(y_1^n; \theta)] dy_1^n} \\ &= \frac{\exp[u(T(x_1^n), \theta)] \exp[v(x_1^n)]}{\int_{A_T(x_1^n)} \exp[u(T(x_1^n), \theta)] \exp[v(y_1^n)] dy_1^n} \\ &= \frac{\exp[v(x_1^n)]}{\int_{A_T(x_1^n)} \exp[v(y_1^n)] dy_1^n}, \end{aligned}$$

which is independent of θ , where the first equality is by the definition of generalized conditional distribution.

Conversely, let $p_{\theta_{x_1^n|T}}^*(x_1^n|t)$ be independent of θ . Let $p_{\theta_{x_1^n|T}}^*(x_1^n|t) := v_1(x_1^n)$. Then

$$\frac{\exp[L_G(x_1^n; \theta)]}{\int_{A_T(x_1^n)} \exp[L_G(y_1^n; \theta)] dy_1^n} = v_1(x_1^n).$$

This implies

$$L_G(x_1^n; \theta) = u(T(x_1^n), \theta) + v(x_1^n),$$

where $u(T(x_1^n), \theta) = \log \int_{A_T(x_1^n)} \exp[L_G(y_1^n; \theta)] dy_1^n$ and $v(x_1^n) = \log v_1(x_1^n)$. Since the above is true for any $x_1^n \in \mathbb{S}^n$, we have the result. ■

A.3. Proof of Proposition 5:

The condition that if $T(x_1^n) = T(y_1^n)$ then $[L_G(x_1^n; \theta) - L_G(y_1^n; \theta)]$ is independent of θ , for all $\theta \in \Theta$ implies that T is a sufficient statistic for θ with respect to L_G , by Proposition 2.2. Thus the only thing we need to prove is that T is further minimal. Let \tilde{T} be a sufficient statistics such that $\tilde{T}(x_1^n) = \tilde{T}(y_1^n)$. Then by Proposition 2.2, $[L_G(x_1^n; \theta) - L_G(y_1^n; \theta)]$ is independent of θ . Hence by hypothesis, we have $T(x_1^n) = T(y_1^n)$. Therefore T is minimal, by definition. ■

B. PROOF OF RESULTS OF SECTION 3

B.1. Proof of Theorem 8:

Let $\Pi = \mathbb{M}^{(\alpha)}$ as in Definition 3.1. Let x_1^n and y_1^n be two i.i.d. samples from some $p_{\theta} \in \mathbb{M}^{(\alpha)}$ such that $T(x_1^n) = T(y_1^n)$. Assume that, for $i = 1, \dots, s$,

$$\overline{f_i(x_1^n)/h(x_1^n)} = \psi_i(T(x_1^n)) \text{ and } \overline{f_i(y_1^n)/h(y_1^n)} = \psi_i(T(y_1^n)).$$

Then

$$\begin{aligned}
& L_J^{(\alpha)}(x_1^n; \theta) - L_J^{(\alpha)}(y_1^n; \theta) \\
&= \frac{1}{\alpha - 1} \log \left[\frac{1}{n} \sum_{i=1}^n p_\theta(x_i)^{\alpha-1} \right] - \frac{1}{\alpha - 1} \log \left[\frac{1}{n} \sum_{i=1}^n p_\theta(y_i)^{\alpha-1} \right] \\
&= \frac{1}{\alpha - 1} \log \frac{Z(\theta)^{\alpha-1} \{ \overline{h(x_1^n)} + w(\theta)^T \overline{f(x_1^n)} \}}{Z(\theta)^{\alpha-1} \{ \overline{h(y_1^n)} + w(\theta)^T \overline{f(y_1^n)} \}} \\
&= \frac{1}{\alpha - 1} \log \frac{\overline{h(x_1^n)}}{\overline{h(y_1^n)}} + \frac{1}{\alpha - 1} \log \frac{1 + w(\theta)^T \{ \overline{f(x_1^n)} / \overline{h(x_1^n)} \}}{1 + w(\theta)^T \{ \overline{f(y_1^n)} / \overline{h(y_1^n)} \}} \\
&= \frac{1}{\alpha - 1} \log \left[\overline{h(x_1^n)} / \overline{h(y_1^n)} \right],
\end{aligned} \tag{54}$$

which is independent of θ . The last equality holds since $T(x_1^n) = T(y_1^n)$ and thus $\overline{f_i(x_1^n)} / \overline{h(x_1^n)} = \overline{f_i(y_1^n)} / \overline{h(y_1^n)}$. Hence T is a sufficient statistic for θ with respect to the Jones et al. likelihood function $L_J^{(\alpha)}$. ■

B.2. Proof of Theorem 12:

We only give the proof for $\mathbb{M}^{(\alpha)}$ -family. The result for $\mathbb{B}^{(\alpha)}$ -family can be proved similarly.

Let us consider a regular $\mathbb{M}^{(\alpha)}$ -family as in Definition 3.1. We will only show that the statistic

$$\overline{f(x_1^n)} / \overline{h(x_1^n)} = (\overline{f_1(x_1^n)} / \overline{h(x_1^n)}, \dots, \overline{f_k(x_1^n)} / \overline{h(x_1^n)})$$

is minimal. Let us consider two sample points x_1^n and y_1^n such that $[L_J^{(\alpha)}(x_1^n; \theta) - L_J^{(\alpha)}(y_1^n; \theta)]$ is independent of θ . Our aim is to prove that $\overline{f(x_1^n)} / \overline{h(x_1^n)} = \overline{f(y_1^n)} / \overline{h(y_1^n)}$.

Consider $[L_J^{(\alpha)}(x_1^n; \theta) - L_J^{(\alpha)}(y_1^n; \theta)]$. Using (54), we have

$$L_J^{(\alpha)}(x_1^n; \theta) - L_J^{(\alpha)}(y_1^n; \theta) = \frac{1}{\alpha - 1} \log \frac{\overline{h(x_1^n)}}{\overline{h(y_1^n)}} + \frac{1}{\alpha - 1} \log \frac{1 + w(\theta)^T \{ \overline{f(x_1^n)} / \overline{h(x_1^n)} \}}{1 + w(\theta)^T \{ \overline{f(y_1^n)} / \overline{h(y_1^n)} \}}.$$

Since the above is independent of θ , we must have

$$\log \frac{1 + w(\theta)^T \{ \overline{f(x_1^n)} / \overline{h(x_1^n)} \}}{1 + w(\theta)^T \{ \overline{f(y_1^n)} / \overline{h(y_1^n)} \}} = C,$$

where C is a constant independent of θ (may depend on x_1^n and y_1^n). This implies

$$(1 - \exp(C)) + w(\theta)^T \left[\overline{f(x_1^n)} / \overline{h(x_1^n)} - \exp(C) \{ \overline{f(y_1^n)} / \overline{h(y_1^n)} \} \right] = 0.$$

Since the given family is regular, the functions $1, w_1, \dots, w_k$ are linearly independent. Hence we have $\exp(C) = 1$ and therefore

$$\left[\overline{f(x_1^n)} / \overline{h(x_1^n)} - \overline{f(y_1^n)} / \overline{h(y_1^n)} \right] = 0.$$

That is,

$$\overline{f(x_1^n)} / \overline{h(x_1^n)} = \overline{f(y_1^n)} / \overline{h(y_1^n)}.$$

Thus $\overline{f}/\overline{h}$ is a minimal sufficient statistic for a regular $\mathbb{M}^{(\alpha)}$ -family with respect to $L_J^{(\alpha)}$. ■

C. PROOF OF RESULTS OF SECTION 4

C.1. *Proof of $\text{Cov}_\theta(\psi(\bar{f}), \bar{f}) = 0$ for regular exponential family:*

Recall that $p_\theta(x) = [h(x) + Z(\theta) + w(\theta)f(x)]$ for $x \in \mathbb{S}$ and $\mathbb{E}_\theta[\psi(\bar{f})] = 0$. That is

$$\int \psi(\bar{f})p_\theta(x_1^n)dx_1^n = 0.$$

Taking derivative with respect to θ on both sides, we get

$$\int \psi(\bar{f}) \frac{\partial}{\partial \theta} \left\{ \prod_{i=1}^n \exp[h(x_i) + w(\theta)f(x_i)] \right\} dx_1^n = 0.$$

This implies

$$\int \psi(\bar{f})p_\theta(x_1^n)w'(\theta)n\bar{f}dx_1^n = 0.$$

Since $w'(\theta) \neq 0$ for all $\theta \in \Theta$, we have $\mathbb{E}_\theta[\psi(\bar{f})\bar{f}] = 0$. Therefore

$$\text{Cov}_\theta[\psi(\bar{f}), \bar{f}] = \mathbb{E}_\theta[\psi(\bar{f})\bar{f}] - \mathbb{E}_\theta[\psi(\bar{f})]\mathbb{E}_\theta[\bar{f}] = 0.$$

■

C.2. *Proof of Proposition 14:*

Since T is a sufficient statistic with respect to the Jones et al. likelihood function, by Proposition 2.3, the conditional distribution of x_1^n given T with respect to p_θ^* is independent of θ . Hence the result follows as $\hat{\theta}$ is a function of x_1^n only. ■

C.3. *Proof of Proposition 15:*

For every partition set $A_{T(y_1^n)}$ of \mathbb{S}^n , the space of all possible n -length samples, we have

$$\mathbb{E}_\theta^* \left[\phi^*(T(X_1^n)) \cdot \mathbf{1}_{A_{T(y_1^n)}} \right] := \int_{A_{T(y_1^n)}} p_\theta^*(x_1^n) \phi^*(T(x_1^n)) dx_1^n = \int_{A_{T(y_1^n)}} p_\theta^*(x_1^n) \phi^*(T(y_1^n)) dx_1^n, \quad (55)$$

since $x_1^n \in A_{T(y_1^n)}$ implies $T(x_1^n) = T(y_1^n)$.

Observe that

$$\begin{aligned} \phi^*(T(x_1^n)) &= \int \hat{\theta}(y_1^n) p_\theta^*(y_1^n | T(x_1^n)) dy_1^n \\ &= \int_{A_{T(x_1^n)}} \hat{\theta}(y_1^n) p_\theta^*(y_1^n | T(x_1^n)) dy_1^n + \int_{A_{T(x_1^n)}^c} \hat{\theta}(y_1^n) p_\theta^*(y_1^n | T(x_1^n)) dy_1^n \\ &= \int_{A_{T(x_1^n)}} \hat{\theta}(y_1^n) p_\theta^*(y_1^n | T(x_1^n)) dy_1^n + 0 \\ &= \int_{A_{T(x_1^n)}} \hat{\theta}(y_1^n) \cdot \left[\frac{p_\theta^*(y_1^n)}{q_\theta^*(T(x_1^n))} \right] dy_1^n \\ &= \frac{\int_{A_{T(x_1^n)}} \hat{\theta}(y_1^n) p_\theta^*(y_1^n) dy_1^n}{\int_{A_{T(x_1^n)}} p_\theta^*(y_1^n) dy_1^n}. \end{aligned}$$

Thus (55) implies

$$\begin{aligned}
\mathbb{E}_\theta^* \left[\phi^*(T(X_1^n)) \cdot \mathbf{1}_{A_{T(y_1^n)}} \right] &= \int_{A_{T(y_1^n)}} p_\theta^*(x_1^n) \phi^*(T(y_1^n)) dx_1^n \\
&= \phi^*(T(y_1^n)) \int_{A_{T(y_1^n)}} p_\theta^*(x_1^n) dx_1^n \\
&= \frac{\int_{A_{T(y_1^n)}} \widehat{\theta}(x_1^n) p_\theta^*(x_1^n) dx_1^n}{\int_{A_{T(y_1^n)}} p_\theta^*(x_1^n) dx_1^n} \left[\int_{A_{T(y_1^n)}} p_\theta^*(x_1^n) dx_1^n \right] \\
&= \int_{A_{T(y_1^n)}} \widehat{\theta}(x_1^n) p_\theta^*(x_1^n) dx_1^n \\
&= \mathbb{E}_\theta^* \left[\widehat{\theta}(x_1^n) \cdot \mathbf{1}_{A_{T(y_1^n)}} \right]. \tag{56}
\end{aligned}$$

Observe that (56) holds for every partition set $A_{T(y_1^n)}$ of \mathbb{S}^n . This implies both the estimators $\widehat{\theta}$ and $\phi^*(T)$ have the same expected value with respect to p_θ^* . That is,

$$\mathbb{E}_\theta^*[\phi^*(T(X_1^n))] = \mathbb{E}_\theta^*[\widehat{\theta}(X_1^n)] = \tau^*(\theta).$$

■

C.4. Proof of Proposition 16:

$\mathcal{A} \subseteq \mathcal{B}$ is immediate from Propositions 4.2 and 4.3. Conversely, let $\psi(T) \in \mathcal{B}$. Since $\mathbb{E}_\theta^*[\psi(T)] = \tau^*(\theta)$, we have $\mathbb{E}_\theta^*[\psi^*(T)|T] \in \mathcal{A}$. But $\mathbb{E}_\theta^*[\psi^*(T)|T] = \psi^*(T)$. This implies that $\psi^*(T) \in \mathcal{A}$. Hence $\mathcal{B} \subseteq \mathcal{A}$. This completes the proof. ■

C.5. Proof of Lemma 17:

We have

$$\begin{aligned}
\text{Var}_\theta^*[\widehat{\theta}] &= \mathbb{E}_\theta^*[(\widehat{\theta} - \tau^*(\theta))^2] \\
&= \mathbb{E}_\theta^*[(\widehat{\theta} - \psi^*(T) + \psi^*(T) - \tau^*(\theta))^2] \\
&= \mathbb{E}_\theta^*[(\widehat{\theta} - \psi^*(T))^2] + \mathbb{E}_\theta^*[(\psi^*(T) - \tau^*(\theta))^2] + 2\mathbb{E}_\theta^*[(\widehat{\theta} - \psi^*(T))(\psi^*(T) - \tau^*(\theta))] \\
&= \mathbb{E}_\theta^*[(\widehat{\theta} - \psi^*(T))^2] + \text{Var}_\theta^*[\psi^*(T)] + 2\mathbb{E}_\theta^*[(\widehat{\theta} - \psi^*(T))(\psi^*(T) - \tau^*(\theta))].
\end{aligned}$$

Now

$$\begin{aligned}
\mathbb{E}_\theta^*[(\widehat{\theta} - \psi^*(T))(\psi^*(T) - \tau^*(\theta))] &= \mathbb{E}_\theta^*[(\widehat{\theta} - \psi^*(T))\psi^*(T)] - \tau^*(\theta) \mathbb{E}_\theta^*[(\widehat{\theta} - \psi^*(T))] \\
&\stackrel{(a)}{=} \mathbb{E}_\theta^*[(\widehat{\theta} - \psi^*(T))\psi^*(T)] \\
&= \mathbb{E}_\theta^*[\widehat{\theta} \psi^*(T)] - \mathbb{E}_\theta^*[(\psi^*(T))^2] \\
&\stackrel{(b)}{=} \mathbb{E}_\theta^*[\mathbb{E}_\theta^*[\widehat{\theta} \psi^*(T)|T]] - \mathbb{E}_\theta^*[(\psi^*(T))^2] \\
&= \mathbb{E}_\theta^*[\psi^*(T) \mathbb{E}_\theta^*[\widehat{\theta}|T]] - \mathbb{E}_\theta^*[(\psi^*(T))^2] \\
&= \mathbb{E}_\theta^*[\psi^*(T) \{ \mathbb{E}_\theta^*[\widehat{\theta}|T] - \psi^*(T) \}] \\
&= \mathbb{E}_\theta^*[\psi^*(T) \{ \phi^*(T) - \psi^*(T) \}],
\end{aligned}$$

where (a) follows from the fact that both $\widehat{\theta}$ and $\psi^*(T)$ are unbiased estimators of $\tau^*(\theta)$ and (b) follows from law of total expectation. Thus we have

$$\text{Var}_\theta^*[\widehat{\theta}] = \mathbb{E}_\theta^*[(\widehat{\theta} - \psi^*(T))^2] + \text{Var}_\theta^*[\psi^*(T)] + 2\mathbb{E}_\theta^*[\psi^*(T)\{\phi^*(T) - \psi^*(T)\}].$$

■

C.6. Proof of Theorem 18:

From Proposition 4.3, we have $\mathbb{E}_\theta^*[\phi^*(T)] = \mathbb{E}_\theta^*[\widehat{\theta}] = \tau^*(\theta)$. Now in Lemma 4.5, if we take $\psi^*(T) = \phi^*(T)$, we have

$$\text{Var}_\theta^*[\widehat{\theta}] = \mathbb{E}_\theta^*[(\widehat{\theta} - \phi^*(T))^2] + \text{Var}_\theta^*[\phi^*(T)] \geq \text{Var}_\theta^*[\phi^*(T)].$$

Equality holds if and only if $\mathbb{E}_\theta^*[(\widehat{\theta} - \phi^*(T))^2] = 0$, that is, if and only if $\widehat{\theta} = \phi^*(T) = \mathbb{E}_\theta^*[\widehat{\theta}|T]$ almost surely. This happens if and only if $\widehat{\theta}$ is a function of T only. This completes the proof. ■

Proof of Proposition 19:

Let $\phi_1(T)$ and $\phi_2(T)$ be two unbiased estimators for $\tau^*(\theta)$ with respect to p_θ^* . Let σ_*^2 be the minimum variance that any unbiased estimator for $\tau^*(\theta)$ can achieve. Suppose $\text{Var}_\theta^*[\phi_1(T)] = \text{Var}_\theta^*[\phi_2(T)] = \sigma_*^2$. We show that $\phi_1(T) = \phi_2(T)$ almost surely.

Consider $\xi(T) := \frac{\phi_1(T) + \phi_2(T)}{2}$. Then $\mathbb{E}_\theta^*[\xi(T)] = \tau^*(\theta)$ and

$$\text{Var}_\theta^*[\xi(T)] \leq \sigma_*^2,$$

since $\text{Cov}_\theta^*[\phi_1(T), \phi_2(T)] \leq \sigma_*^2$, by Cauchy-Schwarz inequality. As σ_*^2 is the minimum variance that one unbiased estimator of $\tau^*(\theta)$ can achieve, we have $\text{Var}_\theta^*[\xi(T)] \geq \sigma_*^2$. Therefore $\text{Var}_\theta^*[\xi(T)] = \sigma_*^2$. This implies

$$\text{Cov}_\theta^*[\phi_1(T), \phi_2(T)] = \sigma_*^2.$$

That means $\phi_1(T)$ and $\phi_2(T)$ are positively correlated. Thus there exist constants, say $\alpha > 0$ and β , such that

$$\phi_1(T) = \alpha\phi_2(T) + \beta \quad \text{almost surely.}$$

Taking expectation and variance with respect to p_θ^* on the both sides, respectively, we get

$$\tau^*(\theta) = \alpha\tau^*(\theta) + \beta \quad \text{and} \quad \sigma_*^2 = \alpha^2\sigma_*^2.$$

These together imply $\alpha = 1$ and $\beta = 0$. That is, $\phi_1(T) = \phi_2(T)$ almost surely. ■

D. PROOF OF RESULTS OF SECTION 5

D.1. Proof of $\text{Cov}_\theta^*(\psi^*(\bar{f}), \bar{f}) \geq 0$ in a regular $\mathbb{M}^{(\alpha)}$ -family:

Since $\mathbb{E}_\theta^*[\psi^*(\bar{f})] = 0$,

$$\int \psi^*(\bar{f})N(\theta)[1 + w(\theta)\bar{f}]^{\frac{1}{\alpha-1}} dx_1^n = 0.$$

That is,

$$\int \psi^*(\bar{f})[1 + w(\theta)\bar{f}]^{\frac{1}{\alpha-1}} dx_1^n = 0.$$

Differentiating both sides with respect to θ ,

$$\int \psi^*(\bar{f})\frac{1}{\alpha-1}[1 + w(\theta)\bar{f}]^{\frac{2-\alpha}{\alpha-1}}w'(\theta)\bar{f} dx_1^n = 0.$$

This implies

$$\int \frac{\psi^*(\bar{f})\bar{f}}{[1+w(\theta)\bar{f}]} p_\theta^*(x_1^n) dx_1^n = 0.$$

That is,

$$\mathbb{E}_\theta^* \left[\frac{\psi^*(\bar{f})\bar{f}}{1+w(\theta)\bar{f}} \right] = 0. \quad (57)$$

We now claim that $\mathbb{E}_\theta^*[\psi^*(\bar{f})\bar{f}] \geq 0$. Let

$$k(\theta) := \mathbb{E}_\theta^*[\psi^*(\bar{f})\bar{f}]. \quad (58)$$

Suppose that $k(\theta) \neq 0$ for all $\theta \in \Theta$. From (58),

$$\int \psi^*(\bar{f})\bar{f} N(\theta) [1+w(\theta)\bar{f}]^{\frac{1}{\alpha-1}} dx_1^n = k(\theta).$$

That is,

$$\int \psi^*(\bar{f})\bar{f} [1+w(\theta)\bar{f}]^{\frac{1}{\alpha-1}} dx_1^n = \frac{k(\theta)}{N(\theta)}.$$

Differentiating both sides with respect to θ , we get

$$\int \psi^*(\bar{f})\bar{f} \frac{1}{\alpha-1} [1+w(\theta)\bar{f}]^{\frac{2-\alpha}{\alpha-1}} w'(\theta)\bar{f} dx_1^n = \frac{\partial}{\partial \theta} \left[\frac{k(\theta)}{N(\theta)} \right].$$

Hence

$$\int \psi^*(\bar{f}) \frac{\bar{f}^2}{1+w(\theta)\bar{f}} p_\theta^*(x_1^n) dx_1^n = (\alpha-1) \frac{N(\theta)}{w'(\theta)} \frac{\partial}{\partial \theta} \left[\frac{k(\theta)}{N(\theta)} \right].$$

Hence

$$\int \psi^*(\bar{f})\bar{f} \left[\frac{1+w(\theta)\bar{f}-1}{1+w(\theta)\bar{f}} \right] p_\theta^*(x_1^n) = (\alpha-1) \frac{w(\theta)N(\theta)}{w'(\theta)} \frac{\partial}{\partial \theta} \left[\frac{k(\theta)}{N(\theta)} \right].$$

This implies

$$\mathbb{E}_\theta^*[\psi^*(\bar{f})\bar{f}] - \mathbb{E}_\theta^* \left[\frac{\psi^*(\bar{f})\bar{f}}{1+w(\theta)\bar{f}} \right] = (\alpha-1) \frac{w(\theta)N(\theta)}{w'(\theta)} \frac{\partial}{\partial \theta} \left[\frac{k(\theta)}{N(\theta)} \right].$$

Using (57), we get

$$k(\theta) = (\alpha-1) \frac{w(\theta)N(\theta)}{w'(\theta)} \frac{\partial}{\partial \theta} \left[\frac{k(\theta)}{N(\theta)} \right].$$

Since $k(\theta) \neq 0$, we have

$$(\alpha-1) \frac{\partial}{\partial \theta} \left[\frac{k(\theta)}{N(\theta)} \right] / \left[\frac{k(\theta)}{N(\theta)} \right] = \frac{w'(\theta)}{w(\theta)}.$$

That is,

$$(\alpha-1) \frac{\partial}{\partial \theta} \log \left[\frac{k(\theta)}{N(\theta)} \right] = \frac{\partial}{\partial \theta} \log w(\theta).$$

This reduces to

$$\frac{\partial}{\partial \theta} \log \left[\frac{k(\theta)}{N(\theta)} \right]^{\alpha-1} = \frac{\partial}{\partial \theta} \log w(\theta).$$

Integrating both sides with respect to θ , we get

$$\log \left[\frac{k(\theta)}{N(\theta)} \right]^{\alpha-1} = \log w(\theta) + \log C$$

where $\log C$ is the constant of integration. This implies

$$k(\theta)^{\alpha-1} = Cw(\theta)N(\theta)^{\alpha-1}.$$

Observe that $C > 0$ and $N(\theta) > 0$ for all $\theta \in \Theta$. Hence, if $w(\theta) > 0$ for all $\theta \in \Theta$, we get $k(\theta) > 0$. This, together with the case $k(\theta) = 0$, implies $k(\theta) \geq 0$. That means, in any case, $\mathbb{E}_\theta^*[\psi^*(\bar{f})\bar{f}] \geq 0$. Therefore

$$\text{Cov}_\theta^*[\psi^*(\bar{f}), \bar{f}] = \mathbb{E}_\theta^*[\psi^*(\bar{f})\bar{f}] - \mathbb{E}_\theta^*[\psi^*(\bar{f})]\mathbb{E}_\theta^*[\bar{f}] = \mathbb{E}_\theta^*[\psi^*(\bar{f})\bar{f}] - 0 = \mathbb{E}_\theta^*[\psi^*(\bar{f})\bar{f}] \geq 0.$$

■

D.2. Derivation of generalized Cramér-Rao bound for $\mathbb{M}^{(\alpha)}$:

Consider

$$p_\theta^*(x_1^n) = N(\theta)[1 + w(\theta)\bar{f}]^{\frac{1}{\alpha-1}} = [N(\theta)^{\alpha-1} + N(\theta)^{\alpha-1}w(\theta)\bar{f}]^{\frac{1}{\alpha-1}}.$$

For convenience, let $F(\theta) := N(\theta)^{\alpha-1}$ and $\tilde{w}(\theta) := N(\theta)^{\alpha-1}w(\theta)$. Then

$$p_\theta^*(x_1^n) = [F(\theta) + \tilde{w}(\theta)\bar{f}]^{\frac{1}{\alpha-1}},$$

and thus

$$\frac{\partial}{\partial \theta} p_\theta^*(x_1^n) = \frac{1}{\alpha-1} p_\theta^*(x_1^n)^{2-\alpha} [F'(\theta) + \tilde{w}'(\theta)\bar{f}]. \quad (59)$$

Then using (59), we get

$$\begin{aligned} \int s^*(x_1^n, \theta) p_\theta^*(x_1^n)^\alpha dx_1^n &= \int \frac{\partial}{\partial \theta} \log p_\theta^*(x_1^n) p_\theta^*(x_1^n)^\alpha dx_1^n \\ &= \frac{1}{\alpha-1} \int p_\theta^*(x_1^n) [F'(\theta) + \tilde{w}'(\theta)\bar{f}] dx_1^n \end{aligned}$$

and

$$\begin{aligned} \int s^*(x_1^n, \theta)^2 p_\theta^*(x_1^n)^{2\alpha-1} dx_1^n &= \int \left[\frac{\partial}{\partial \theta} \log p_\theta^*(x_1^n) \right]^2 p_\theta^*(x_1^n)^{2\alpha-1} dx_1^n \\ &= \left(\frac{1}{\alpha-1} \right)^2 \int p_\theta^*(x_1^n) [F'(\theta) + \tilde{w}'(\theta)\bar{f}]^2 dx_1^n. \end{aligned}$$

Thus

$$\begin{aligned} \text{Var}_\theta^*[p_\theta^*(X_1^n) s^*(X_1^n, \theta)] &= \int s^*(x_1^n, \theta)^2 p_\theta^*(x_1^n)^{2\alpha-1} dx_1^n - \left[\int s^*(x_1^n, \theta) p_\theta^*(x_1^n)^\alpha dx_1^n \right]^2 \\ &= \left(\frac{1}{\alpha-1} \right)^2 \{ \mathbb{E}_\theta^*[(F'(\theta) + \tilde{w}'(\theta)\bar{f})^2] - (\mathbb{E}_\theta^*[F'(\theta) + \tilde{w}'(\theta)\bar{f}])^2 \} \\ &= \left(\frac{1}{\alpha-1} \right)^2 \text{Var}_\theta^*[F'(\theta) + \tilde{w}'(\theta)\bar{f}] \\ &= \left(\frac{1}{\alpha-1} \right)^2 [\tilde{w}'(\theta)]^2 \text{Var}_\theta^*[\bar{f}]. \end{aligned}$$

Also, using (59), we get

$$\begin{aligned}
& \text{Cov}_\theta^*[s^*(X_1^n, \theta), p_\theta^*(X_1^n)^{\alpha-1} s^*(X_1^n, \theta)] \\
&= \int s^*(x_1^n, \theta)^2 p_\theta^*(x_1^n)^\alpha dx_1^n \\
&= \int s^*(x_1^n, \theta) \left[\frac{1}{p_\theta^*(x_1^n)} \frac{\partial}{\partial \theta} p_\theta^*(x_1^n) \right] p_\theta^*(x_1^n)^\alpha dx_1^n \\
&= \int \frac{1}{\alpha-1} p_\theta^*(x_1^n) s^*(x_1^n, \theta) [F'(\theta) + \tilde{w}'(\theta) f(x_1^n)] dx_1^n \\
&= \frac{1}{\alpha-1} \int p_\theta^*(x_1^n) s^*(x_1^n, \theta) F'(\theta) dx_1^n + \frac{1}{\alpha-1} \int p_\theta^*(x_1^n) s^*(x_1^n, \theta) \tilde{w}'(\theta) \bar{f} dx_1^n \\
&= 0 + \frac{1}{\alpha-1} \tilde{w}'(\theta) \int p_\theta^*(x_1^n) \frac{1}{p_\theta^*(x_1^n)} \frac{\partial}{\partial \theta} p_\theta^*(x_1^n) \bar{f} dx_1^n \\
&= \frac{1}{\alpha-1} \tilde{w}'(\theta) \frac{\partial}{\partial \theta} \int p_\theta^*(x_1^n) \bar{f} dx_1^n \\
&= \frac{1}{\alpha-1} \tilde{w}'(\theta) \frac{\partial}{\partial \theta} \tau^*(\theta).
\end{aligned}$$

■

D.3. Proof of Theorem 51:

The deformed probability distribution associated with Basu et al. likelihood function is given by

$$p_\theta^*(x_1^n) = \frac{\exp\left[\frac{\alpha}{n(\alpha-1)} \sum_{i=1}^n p_\theta(x_i)^{\alpha-1}\right]}{\int \exp\left[\frac{\alpha}{n(\alpha-1)} \sum_{i=1}^n p_\theta(y_i)^{\alpha-1}\right] dy_1^n}. \quad (60)$$

$T := T(X_1^n)$ is a sufficient statistic for θ with respect to Basu et al. likelihood function. Let $\hat{\theta} := \hat{\theta}(X_1^n)$ be an unbiased estimator of $\tau(\theta)$ with respect to p_θ^* . Let us define

$$\phi(T(x_1^n)) := \mathbb{E}^*[\hat{\theta}|T = T(x_1^n)] = \int \hat{\theta}(y_1^n) p_\theta^*(y_1^n | T(x_1^n)) dy_1^n,$$

where the conditional distribution is as defined in (10) of the main article. Then proceeding similar steps as in Lemma 4.5 and Theorem 4.6, we have the following.

1. $\mathbb{E}^*[\phi(T)] = \mathbb{E}_\theta^*[\hat{\theta}] = \tau(\theta)$,
2. $\text{Var}_\theta^*[\phi(T)] \leq \text{Var}_\theta^*[\hat{\theta}]$, and equality holds if and only if $\hat{\theta}$ is a function of T only.

Now for the given $\mathbb{B}^{(\alpha)}$ -family, using (60), we have

$$p_\theta^*(x_1^n) = \exp\left[\frac{\alpha}{\alpha-1} \bar{h} + \tilde{F}(\theta) + \frac{\alpha}{\alpha-1} w(\theta) \bar{f}\right].$$

Recall that $T = \bar{f}$ is a sufficient statistic for θ with respect to Basu et al. likelihood function. Using generalized Rao-Blackwell theorem for Basu et al. likelihood function, we get

$$\mathbb{E}_\theta^*[\phi(\bar{f})] = \mathbb{E}_\theta^*[\hat{\theta}] = \tau(\theta) \quad \text{and} \quad \text{Var}_\theta^*[\phi(\bar{f})] \leq \text{Var}_\theta^*[\hat{\theta}]. \quad (61)$$

Let us define $\psi(\bar{f}) := \phi(\bar{f}) - \bar{f}$. Then $\mathbb{E}_\theta^*[\psi(\bar{f})] = 0$, since $\mathbb{E}_\theta^*[\bar{f}] = \tau(\theta)$ by hypothesis. Also

$$\text{Var}_\theta^*[\phi(\bar{f})] = \text{Var}_\theta^*[\psi(\bar{f}) + \bar{f}] = \text{Var}_\theta^*[\psi(\bar{f})] + \text{Var}_\theta^*[\bar{f}] + 2 \text{Cov}^*(\psi(\bar{f}), \bar{f}).$$

Proceeding as in Section III(A), one can show that $\text{Cov}^*(\psi(\bar{f}), \bar{f}) = 0$. Thus we get

$$\text{Var}_\theta^*[\phi(\bar{f})] = \text{Var}_\theta^*[\psi(\bar{f})] + \text{Var}_\theta^*[\bar{f}] \geq \text{Var}_\theta^*[\bar{f}].$$

Hence the inequality in (61) reduces to

$$\text{Var}_\theta^*[\hat{\theta}] \geq \text{Var}_\theta^*[\bar{f}], \quad (62)$$

that is, \bar{f} is the best estimator of its expected value with respect to p_θ^* .

We next show that $\text{Var}_\theta^*[\bar{f}] = \frac{[\tau'(\theta)]^2}{I_n^*(\theta)}$. Observe that p_θ^* forms a 1-parameter exponential family and

$$\begin{aligned} I_n^*(\theta) &= \text{Var}_\theta^*\left[\frac{\partial}{\partial\theta} \log p_\theta^*(X_1^n)\right] \\ &= \text{Var}_\theta^*\left[\tilde{F}'(\theta) + \frac{\alpha}{\alpha-1} w'(\theta) \bar{f}\right] \\ &= \left(\frac{\alpha}{\alpha-1}\right)^2 (w'(\theta))^2 \text{Var}_\theta^*[\bar{f}]. \end{aligned} \quad (63)$$

Using the fact $\mathbb{E}_\theta^*\left[\frac{\partial}{\partial\theta} \log p_\theta^*(X_1^n)\right] = 0$ and $\mathbb{E}_\theta^*[\bar{f}] = \tau(\theta)$, we get

$$\tau(\theta) = -\frac{(\alpha-1)}{\alpha} \frac{\tilde{F}'(\theta)}{w'(\theta)}. \quad (64)$$

Further taking derivative on both sides of $\mathbb{E}_\theta^*\left[\frac{\partial}{\partial\theta} \log p_\theta^*(X_1^n)\right] = 0$, we get

$$\tau'(\theta) = \frac{\alpha}{\alpha-1} w'(\theta) \text{Var}_\theta^*[\bar{f}]. \quad (65)$$

Thus, using (63), (64) and (65), we get

$$\text{Var}_\theta^*[\bar{f}] = \frac{[\tau'(\theta)]^2}{I_n^*(\theta)}.$$

This completes the proof. ■

REFERENCES

- [1] E. W. Barankin and A. P. Maitra, "Generalization of the fisher-darmois-koopman-pitman theorem on sufficient statistics," *Sankhya: The Indian Journal of Statistics, Series A*, vol. 25(3), pp. 217–244, 1963.
- [2] A. G. Bashkirov, "On maximum entropy principle, superstatistics, power-law distribution and rényi parameter," *Phys. A*, vol. 340, pp. 153–162, 2004.
- [3] A. Basu, I. R. Harris, N. L. Hjort, and M. C. Jones, "Robust and efficient estimation by minimizing a density power divergence," *Biometrika*, vol. 85, pp. 549–559, 1998.
- [4] D. Blackwell, "Conditional expectation and unbiased sequential estimation," *Ann. Math. Statist.*, vol. 18(1), pp. 105–110, 1947.
- [5] M. Broniatowski, A. Toma, and I. Vajda, "Decomposable pseudo-distances and applications in statistical estimation," *J. Statist. Plann. Inference*, vol. 142, pp. 2574–2585, 2012.
- [6] M. Broniatowski and I. Vajda, "Several applications of divergence criteria in continuous families," *Kybernetika (Prague)*, vol. 48, pp. 600–636, 2012.
- [7] L. D. Brown, *Fundamentals of Statistical Exponential Families: with Applications in Statistical Decision Theory*. Institute of Mathematical Statistics, Hayward, CA., 1986.
- [8] C. Bunte and A. Lapidoth, "Encoding tasks and rényi entropy," *IEEE Trans. Inform. Theory*, vol. 60, pp. 5065–5076, 2014.
- [9] J. c Principe, *Information Theoretic Learning: Renyi's Entropy and Kernel Perspectives*. Springer Science and Business Media, New York, USA., 2010.
- [10] G. Casella and R. L. Berger, *Statistical Inference*. 2nd ed. Duxbury Press, 2002.
- [11] A. Cichocki and S. Amari, "Families of alpha-beta-and gamma-divergences: Flexible and robust measure of similarities," *Entropy*, vol. 12, pp. 1532–1568, 2010.
- [12] H. Cramér, *Mathematical methods of statistics*. Princeton: Princeton University Press, 1946.
- [13] I. Csiszár, "Generalized cutoff rates and rényi's information measures," *IEEE Trans. Inform. Theory*, vol. 41, pp. 26–34, 1995.
- [14] I. Csiszár and F. Matúš, "Generalized minimizers of convex integral functionals, bergman distance, pythagorean identities," *Kybernetika (Prague)*, vol. 48, pp. 637–689, 2012.
- [15] I. Csiszár and P. C. Shields, *Information Theory and Statistics: A Tutorial*. Foundations and Trends in Communications and Information Theory, 2004.
- [16] G. Darmois, "Sur les lois de probabilités a estimation exhaustive," *C. R. Acad. Sci. Paris (in French)*, vol. 200, pp. 1265–1266, 1935.
- [17] S. Eguchi, O. Komori, and S. Kato, "Projective power entropy and maximum tsallis entropy distributions," *Entropy*, vol. 13, pp. 1746–1764, 2011.

- [18] S. Eguchi, O. Komori, and A. Ohara, “Duality of maximum entropy and minimum divergence,” *Phys. Rev. E.*, vol. 16, pp. 3552–3572, 2014.
- [19] C. Field and B. Smith, “Robust estimation: A weighted maximum likelihood approach,” *Int. Stat. Rev.*, vol. 62, pp. 405–424, 1994.
- [20] R. A. Fisher, “On the mathematical foundations of theoretical statistics,” *Philos. Trans. Roy. Soc. A.*, vol. 222, pp. 309–368, 1922.
- [21] —, “Two new properties of mathematical likelihood,” *Proceedings of the Royal Society, Series A*, vol. 144, pp. 285–307, 1934.
- [22] H. Fujisawa, “Normalized estimating equation for robust parameter estimation,” *Electron. J. Stat.*, vol. 7, pp. 1587–1606, 2013.
- [23] H. Fujisawa and S. Eguchi, “Robust parameter estimation with a small bias against heavy contamination,” *J. Multivariate Anal.*, vol. 99, pp. 2053–2081, 2008.
- [24] A. Gayen and M. A. Kumar, “Generalized estimating equation for the student-t distributions,” in *2018 IEEE International Symposium on Information Theory (ISIT)*, 2018, pp. 571–575.
- [25] —, “A generalized notion of sufficiency for power-law distributions,” in *2021 IEEE International Symposium on Information Theory (ISIT)*, 2021, pp. 2185–2190.
- [26] —, “Projection theorems and estimating equations for power-law models,” *Journal of Multivariate Analysis*, vol. 184, p. 104734, 2021.
- [27] P. R. Halmos and L. J. Savage, “Application of the radon-nikodym theorem to the theory of sufficient statistics,” *The Annals of Mathematical Statistics*, vol. 20, pp. 225 – 241, 1949.
- [28] H. G. Hoang, B. Vo, B. T. Vo, and R. Mahler, “The cauchy–schwarz divergence for poisson point processes,” *IEEE Transactions on Information Theory*, vol. 61, pp. 4475–4485, 2015.
- [29] R. Jenssen, J. C. Principe, D. Erdogmus, and T. Eltoft, “The cauchy–schwarz divergence and parzen windowing: Connections to graph theory and mercer kernels,” *Journal of the Franklin Institute*, vol. 343, pp. 614–629, 2006.
- [30] M. C. Jones, N. L. Hjort, I. R. Harris, and A. Basu, “A comparison of related density based minimum divergence estimators,” *Biometrika*, vol. 88, pp. 865–873, 2001.
- [31] K. Kampa, E. Hasanbelliu, and J. C. Principe, “Closed-form cauchy-schwarz pdf divergence for mixture of gaussians,” *Proceedings of International Joint Conference on Neural Networks, San Jose, California, USA*, pp. 2578–2585, 2011.
- [32] B. O. Koopman, “On distributions admitting a sufficient statistic,” *Trans. Amer. Math. Soc.*, vol. 39, pp. 399–409, 1936.
- [33] M. A. Kumar and R. Sundaresan, “Minimization problems based on relative α -entropy i: Forward projection,” *IEEE Trans. Inform. Theory.*, vol. 61, pp. 5063–5080, 2015.
- [34] —, “Minimization problems based on relative α -entropy ii: Reverse projection,” *IEEE Trans. Inform. Theory.*, vol. 61, pp. 5081–5095, 2015.
- [35] E. L. Lehmann and G. Casella, *Theory of point estimation*. Springer, 2nd ed., 1998.
- [36] E. Lutwak, D. Yang, and G. Zhang, “Cramér-rao and moment-entropy inequalities for rényi entropy and generalized fisher information,” *IEEE Trans. Inform. Theory.*, vol. 51, pp. 473–478, 2005.
- [37] A. Maji, A. Ghosh, and A. Basu, “The logarithmic super divergence and asymptotic inference properties,” *ASTA Adv. Stat. Anal.*, vol. 100, pp. 99–131, 2016.
- [38] J. Naudts, “Estimators, escort probabilities, and ϕ -exponential families in statistical physics,” *J. Inequal. Pure. Appl. Math.*, vol. 5, p. 102, 2004.
- [39] J. Neyman, “Sur un teorema concernente la cosidette statisticle sufficienti,” *Giorn. Ist. Ital. Attuari*, vol. 6, pp. 320–334, 1935.
- [40] F. Nielsen, K. Sun, and S. Marchand-Maillet, “k-means clustering with hölder divergences,” *Geometric Science of Information*, pp. 856–863, 2017.
- [41] —, “On hölder projective divergences,” *Entropy*, vol. 19, pp. 1–28, 2017.
- [42] E. J. G. Pitman, “Sufficient statistics and intrinsic accuracy,” *Mathematical Proceedings of the Cambridge Phil. Soc.*, vol. 32, pp. 567–579, 1936.
- [43] C. R. Rao, “Information and accuracy attainable in the estimation of statistical parameters,” *Bulletin of the Calcutta Mathematical Society*, vol. 37(3), pp. 81–91, 1945.
- [44] —, “Minimum variance and the estimation of several parameters,” *Mathematical Proceedings of the Cambridge Philosophical Society*, vol. 43, pp. 280–283, 1947.
- [45] —, “Sufficient statistics and minimum variance estimates,” *Mathematical Proceedings of the Cambridge Philosophical Society*, vol. 45(2), pp. 213–218, 1949.
- [46] Z. Shanqing, Z. Kunlong, and X. Weibin, “Application of cauchy-schwarz divergence in image segmentation,” *Computer Engineering and Applications*, vol. 49, pp. 129–131, 2013.
- [47] R. Sundaresan, “A measure of discrimination and its geometric properties,” in *2002 IEEE International Symposium on Information Theory (ISIT)*, 2002, p. 264.
- [48] —, “Guessing under source uncertainty,” *IEEE Trans. Inform. Theory.*, vol. 53, pp. 269–287, 2007.
- [49] C. Tsallis, “What are the numbers that experiments provide?” *Quim. Nova*, vol. 17(6), pp. 468–471, 1994.
- [50] M. P. Windham, “Robustifying model fitting,” *J. R. Stat. Soc. Ser. B. Stat. Methodol.*, vol. 57, pp. 599–609, 1995.