

DISTINGUISHED VARIETIES IN THE POLYDISC

SOURAV PAL

ABSTRACT. We characterize all distinguished varieties in the polydisc \mathbb{D}^n and thus generalize the seminal work due to Agler and McCarthy on distinguished varieties in the bidisc. We show that a distinguished variety in \mathbb{D}^n is a part of an affine algebraic curve which is a set-theoretic complete intersection and can be represented by Taylor joint spectrum of $n - 1$ commuting linear complex matrix pencils satisfying certain conditions.

1. INTRODUCTION

In 2005, Agler and McCarthy published a very influential article [1], where they established the fact that for any pair of commuting contractive matrices (T_1, T_2) , there is a one-dimensional complex algebraic variety V in \mathbb{C}^2 such that V intersects \mathbb{D}^2 and that the von-Neumann inequality holds on $V \cap \mathbb{D}^2$ for any polynomial p in $\mathbb{C}[z_1, z_2]$, i.e.

$$\|p(T_1, T_2)\| \leq \sup_{(z_1, z_2) \in V \cap \mathbb{D}^2} |p(z_1, z_2)|, \quad (1.1)$$

provided that T_1, T_2 do not have eigenvalues of unit modulus. The variety V exits the bidisc through its distinguished boundary, the 2-torus \mathbb{T}^2 , without intersecting any other part of its topological boundary. Such a set $V \cap \mathbb{D}^2$ is called a distinguished variety in \mathbb{D}^2 and has the following determinantal representation

$$V \cap \mathbb{D}^2 = \{(z, w) \in \mathbb{D}^2 : \det(\Psi(z) - wI) = 0\} \quad (1.2)$$

for some matrix-valued rational function Ψ on the unit disc \mathbb{D} that is unitary on the unit circle \mathbb{T} . The novel idea of Agler and McCarthy refines Ando's inequality for commuting matrices T_1, T_2 . Ando proved in [3] that for any pair of commuting contractions T_1, T_2

$$\|p(T_1, T_2)\| \leq \sup_{(z_1, z_2) \in \overline{\mathbb{D}^2}} |p(z_1, z_2)|, \quad p \in \mathbb{C}[z_1, z_2]. \quad (1.3)$$

So in particular when T_1, T_2 are commuting matrices, Ando's inequality (1.3), which holds on the two-dimensional bidisc, also holds on $V \cap \mathbb{D}^2$ of dimension one. In geometric language, a pair of commuting matrices which live inside the bidisc, actually live on a curve in \mathbb{D}^2 . Recently, these path-breaking results for \mathbb{D}^2 have been reformulated and extended for a wider class of operators in [4] and [7] respectively.

2010 *Mathematics Subject Classification.* 14H50, 14M10, 14M12, 32A60.

Key words and phrases. Distinguished Varieties, Polydisc, Taylor Joint Spectrum.

The author is supported by the Seed Grant of IIT Bombay, the CPDA of Govt. of India and the MATRICS Award of SERB, (Award No. SERB/F/10165/2019-2020) of DST, India.

Recall that for any positive integer n , $\mathbb{A}_{\mathbb{C}}^n$ denotes the *affine n -space* \mathbb{C}^n over \mathbb{C} . If T is any subset of the polynomial ring $\mathbb{C}[z_1, \dots, z_n]$, the *zero set* of T is defined to be the set of common zeros of all elements of T , namely

$$Z(T) = \{P \in \mathbb{A}_{\mathbb{C}}^n : f(P) = 0 \text{ for all } f \in T\}.$$

We shall follow standard terminologies from the literature of basic algebraic geometry (e.g. [9]) and define the following.

Definition 1.1. A subset W of $\mathbb{A}_{\mathbb{C}}^n$ is called an *affine algebraic set* or simply an *algebraic set* if there is a subset T of $\mathbb{C}[z_1, \dots, z_n]$ such that $W = Z(T)$. An *affine algebraic variety* or simply an *algebraic variety* in $\mathbb{A}_{\mathbb{C}}^n$ is an irreducible algebraic set in $\mathbb{A}_{\mathbb{C}}^n$.

Definition 1.2. A *distinguished set* Ω in a domain $G \subset \mathbb{C}^n$ is the intersection of G with an algebraic set $W \subset \mathbb{A}_{\mathbb{C}}^n$ such that the complex-dimension of W is greater than zero and that W exits the domain G through the distinguished boundary $b\overline{G}$ of G , that is,

$$\dim W > 0, \quad \Omega = G \cap W \quad \text{and} \quad \overline{\Omega} \cap \partial \overline{G} = \overline{\Omega} \cap b\overline{G},$$

where $\partial \overline{G}$ is the topological boundary of G . A *distinguished variety* in G is an irreducible distinguished set. Also for a distinguished set Ω , that is defined by an algebraic set W , we denote by $\overline{\Omega}$ the set $W \cap \overline{G}$, that is, $\overline{\Omega} = W \cap \overline{G}$.

According to our definitions, what we mean by a distinguished set in \mathbb{D}^n here, was called an *inner variety* in [13]. Also, note that the distinguished sets in \mathbb{D}^2 were studied in [1], not only the distinguished varieties. Nevertheless, a distinguished set is nothing but a union of distinguished varieties. Only the distinguished sets or varieties with dimension greater than zero are considered here in order to avoid triviality as the same was done in [1].

In this article we study the distinguished varieties in the polydisc

$$\mathbb{D}^n = \{(z_1, \dots, z_n) : |z_i| < 1, i = 1, \dots, n\}.$$

In Theorem 3.1 and Corollary 3.2 of this paper, we show that all distinguished sets are one-dimensional. The second main result of this paper, Theorem 3.5, finds a representation for all distinguished varieties in \mathbb{D}^n in terms of Taylor joint spectrum of $n - 1$ commuting complex square matrices satisfying certain conditions. We obtain this representation via the distinguished varieties in the *symmetrized polydisc* \mathbb{G}_n , where

$$\mathbb{G}_n = \left\{ \left(\sum_{1 \leq i \leq n} z_i, \sum_{1 \leq i < j \leq n} z_i z_j, \dots, \prod_{i=1}^n z_i \right) : |z_i| < 1, i = 1, \dots, n \right\}.$$

Note that the Taylor joint spectrum of finitely many commuting complex square matrices A_1, \dots, A_n of same size is just the set of joint eigenvalues of A_1, \dots, A_n , but this is not true in general for bounded operators. One can see Curto's well-written survey article [6] or Taylor's original works [14, 15] to have a detailed discussion on Taylor joint spectrum for commuting bounded operators.

2. DISTINGUISHED VARIETIES IN THE SYMMETRIZED POLYDISC

We begin with a few definitions from basic algebraic geometry which will be useful for the rest of the paper.

Definition 2.1. An *affine algebraic curve* or simply an *algebraic curve* in $\mathbb{A}_{\mathbb{C}}^n$ is an algebraic set that has dimension one as a complex manifold.

Definition 2.2. An affine algebraic set or variety V of dimension $k (\leq n)$ in $\mathbb{A}_{\mathbb{C}}^n$ is called a *set-theoretic complete intersection* or simply a *complete intersection* if V is defined by $n - k$ independent polynomials in $\mathbb{C}[z_1, \dots, z_n]$. Thus an algebraic curve C in $\mathbb{A}_{\mathbb{C}}^n$ is a complete intersection if C is defined by $n - 1$ independent polynomials in $\mathbb{C}[z_1, \dots, z_n]$.

Theorem 2.3 ([9], CH-I, Corollary 1.6). *Every algebraic set in $\mathbb{A}_{\mathbb{C}}^n$ can be uniquely expressed as a union of affine algebraic varieties, no one containing another.*

Lemma 2.4. *Let X be an affine algebraic variety in $\mathbb{A}_{\mathbb{C}}^n$ such that $X \cap V(z_n - b)$ is a finite set for every $b \in \mathbb{C}$. Then X is an algebraic curve.*

Proof. The collection $\{V(z_n - b) : b \in \mathbb{C}\}$ defines a one parameter family over \mathbb{C} and by hypothesis $X \cap V(z_n - b)$ is finite for every $b \in \mathbb{C}$. Let $A = \bigcap_{b \in \mathbb{C}} V(z_n - b)$. Then we can define a regular morphism $\phi : X \setminus A \rightarrow \mathbb{A}_{\mathbb{C}}^1$ and hence a proper flat map $\phi^* : X^* \rightarrow \mathbb{A}_{\mathbb{C}}^n$, where X^* is the blow up of X along A such that the fibre

$$X_b^* = \phi^*^{-1}(\{b\}) = X \cap V(z_n - b).$$

Since by hypothesis $X \cap V(z_n - b)$ is a finite set, it follows that $\dim(X^*) = \{0\}$. Since ϕ^* is proper and flat, we also have

$$\dim(X_b^*) = \dim(X^*) - \dim(\mathbb{A}_{\mathbb{C}}^n).$$

So we have $\dim(X^*) = \dim(\mathbb{A}_{\mathbb{C}}^n)$. Since X^* is birational with X , it follows that

$$\dim(X) = \dim(X^*) = 1.$$

Hence X is an algebraic curve. □

Recall from the literature (see [5, 10] and the references therein) that the *symmetrized n -disc* or simply the *symmetrized polydisc* \mathbb{G}_n is the image of the polydisc \mathbb{D}^n under the symmetrization map $\pi_n : \mathbb{C}^n \rightarrow \mathbb{C}^n$ defined by

$$\pi_n(z) = (s_1(z), \dots, s_{n-1}(z), p(z)), \quad z = (z_1, \dots, z_n),$$

where

$$s_i(z) = \sum_{1 \leq k_1 \leq k_2 \leq \dots \leq k_i \leq n} z_{k_1} \dots z_{k_i}, \quad i = 1, \dots, n-1 \quad \text{and} \quad p(z) = \prod_{i=1}^n z_i.$$

In [12], the author and Shalit have given an explicit description of distinguished varieties in the symmetrized bidisc

$$\mathbb{G}_2 = \{(z_1 + z_2, z_1 z_2) : |z_i| < 1, i = 1, 2\}.$$

The author further generalized the result for the symmetrized polydisc in [11] and the following representation for a distinguished variety in \mathbb{G}^n was obtained.

Theorem 2.5 ([11], Theorem 3.14). *Let*

$$\Lambda = \{(s_1, \dots, s_{n-1}, p) \in \mathbb{G}_n : (s_1, \dots, s_{n-1}) \in \sigma_T(F_1^* + pF_{n-1}, F_2^* + pF_{n-2}, \dots, F_{n-1}^* + pF_1)\}, \quad (2.1)$$

where σ_T denotes the Taylor joint spectrum and F_1, \dots, F_{n-1} are complex square matrices of same order that satisfy the following conditions:

- (i) $[F_i, F_j] = 0$ and $[F_i^*, F_{n-j}] = [F_j^*, F_{n-i}]$, for $1 \leq i < j \leq n-1$;
- (ii) $\sigma_T(F_1^* + zF_{n-1}, F_2^* + zF_{n-2}, \dots, F_{n-1}^* + zF_1, zI) \subseteq \mathbb{G}_n$, $\forall z \in \mathbb{D}$;
- (iii) the polynomials $\{f_1, \dots, f_{n-1}\}$, where $f_i = \det(F_i^* + pF_{n-i} - s_i I)$, form a regular sequence;
- (iv) either $\Lambda = V_S \cap \mathbb{G}_n$ or Λ is an irreducible component of $V_S \cap \mathbb{G}_n$, where V_S is the complex algebraic set generated by the set of polynomials $S = \{f_1, \dots, f_{n-1}\}$.

Then, Λ is a distinguished variety in \mathbb{G}_n , which further is an affine algebraic curve lying in \mathbb{G}_n and a complete intersection.

Conversely, every distinguished variety Λ in \mathbb{G}_n is an affine algebraic curve in \mathbb{G}_n , a complete intersection and has representation as in (2.1), where F_1, \dots, F_{n-1} are complex square matrices of same order satisfying the above conditions (i) – (iv).

3. DISTINGUISHED VARIETIES IN THE POLYDISC

In this section, we find characterization of the distinguished varieties in the polydisc \mathbb{D}^n by an application of Theorem 2.5. We first prove that every distinguished variety in the polydisc is an affine algebraic curve. Then we show that a distinguished variety in \mathbb{D}^n gives rise to a distinguished variety in \mathbb{G}_n and vice-versa. It will follow as a corollary that a distinguished variety in \mathbb{D}^n is a complete intersection. Combining all these together, we obtain the desired representation of a distinguished variety in \mathbb{D}^n . Also, we show that every distinguished variety in \mathbb{D}^n is polynomially convex.

Theorem 3.1. *Every distinguished variety in the polydisc \mathbb{D}^n is a part of an affine algebraic curve in \mathbb{C}^n .*

Proof. Let Ω be a distinguished variety in \mathbb{D}^n . Then $\Omega = V_S \cap \mathbb{D}^n$ for some complex algebraic variety V_S in \mathbb{C}^n , generated by a set of polynomials $S \subseteq \mathbb{C}[z_1, \dots, z_n]$. By Eisenbud-Evans theorem (see [8]), there exist n polynomials say f_1, \dots, f_n in $\mathbb{C}[z_1, \dots, z_n]$ such that V_S is generated by f_1, \dots, f_n . So, without loss of generality let us assume that

$$\Omega = V_S \cap \mathbb{D}^n = \{(z_1, \dots, z_n) \in \mathbb{D}^n : f_i(z_1, \dots, z_n) = 0, 1 \leq i \leq n\}.$$

Let if possible Ω has complex dimension u and $u > 1$. Let $(t_1, \dots, t_n) \in \Omega$. Then $|t_i| < 1$ for $i = 1, \dots, n$. Let

$$\Omega_{t_n} = \{(z_1, \dots, z_n) \in \Omega : z_n = t_n\}$$

and let

$$V_{S_n} = \{(z_1, \dots, z_{n-1}, t_n) \in \mathbb{C}^n : f_i(z_1, \dots, z_{n-1}, t_n) = 0, i = 1, \dots, n\} \subseteq V_S.$$

Then $\Omega_{t_n} = V_{S_n} \cap \mathbb{D}^n \subset V_S \cap \mathbb{D}^n = \Omega$. Now if every such set Ω_{t_n} is finite whenever $(t_1, \dots, t_n) \in \Omega$, then by an argument similar to that in Lemma 2.4, it follows that Ω has complex dimension 1, a contradiction as $u > 1$. Thus, we choose such a point $(t_1, \dots, t_n) \in \Omega$ for which Ω_{t_n} has complex dimension at least 1. Set

$$V_{S'} = \{(z_1, \dots, z_{n-1}) \in \mathbb{C}^{n-1} : f_i(z_1, \dots, z_{n-1}, t_n) = 0, 1 \leq i \leq n\}.$$

Then $V_{S'}$ is a complex algebraic variety in \mathbb{C}^{n-1} with complex dimension at least $u - 1$ and $u - 1 \geq 1$. The intersection of $V_{S'}$ and \mathbb{D}^{n-1} is not empty as $(t_1, \dots, t_{n-1}) \in V_{S'} \cap \mathbb{D}^{n-1}$. Note that $V_{S'}$, being a complex algebraic variety, is an unbounded connected subset of \mathbb{C}^{n-1} . Therefore, $V_{S'}$ must exit through the topological boundary $\partial \overline{\mathbb{D}^{n-1}}$ of \mathbb{D}^{n-1} . Let $(q_1, \dots, q_{n-1}) \in V_{S'} \cap \partial \overline{\mathbb{D}^{n-1}}$. Then at least one of q_1, \dots, q_{n-1} has modulus equal to 1. Let $|q_t| = 1$. Clearly $(q_1, \dots, q_{n-1}, t_n) \in V_{S_n} \subseteq V_S$ as $(q_1, \dots, q_{n-1}) \in V_{S'} \cap \partial \overline{\mathbb{D}^{n-1}}$. Since $|q_t| = 1$ and $|t_n| < 1$, $(q_1, \dots, q_{n-1}, t_n)$ belongs to $\overline{\mathbb{D}^n} \setminus \mathbb{D}^n$ but does not belong to \mathbb{T}^n . Thus, $(q_1, \dots, q_{n-1}, t_n) \in V_S \cap (\partial \overline{\mathbb{D}^n} \setminus \mathbb{T}^n)$ and hence Ω does not exit through the distinguished boundary of \mathbb{D}^n , a contradiction. Hence Ω has complex dimension 1 and consequently it is an affine algebraic curve lying in \mathbb{D}^n . □

Corollary 3.2. *Every distinguished set in \mathbb{D}^n (i.e. an inner variety) has complex dimension 1.*

Proof. An algebraic set W is nothing but the union of its irreducible components which are algebraic varieties and the dimension of W is the maximum of the dimensions of its irreducible components. Hence a distinguished set in \mathbb{D}^n has complex dimension 1. □

Note that Theorem 3.1 and Corollary 3.2 contradict Theorem 1.4 in [13], which ensures the existence of a distinguished set in \mathbb{D}^n of dimension k for every $k \leq n$.

Theorem 3.3. *Let $\Lambda \subset \mathbb{G}_n$. Then Λ is a distinguished variety in \mathbb{G}_n if and only if $\Lambda = \pi_n(\Omega)$ for some distinguished variety Ω in \mathbb{D}^n .*

Proof. Let Λ be a distinguished variety in \mathbb{G}_n . Then by Theorem 2.5, Λ is an affine algebraic curve and is a complete intersection. Let Λ be generated by $S = \{f_1, \dots, f_{n-1}\} \subset \mathbb{C}[z_1, \dots, z_{n-1}]$, that is,

$$\Lambda = \{(s_1, \dots, s_{n-1}, p) \in \mathbb{G}_n : f_i(s_1, \dots, s_{n-1}, p) = 0, 1 \leq i \leq n-1\}.$$

Let

$$\Omega = \{(z_1, \dots, z_n) \in \mathbb{D}^n : f_i \circ \pi_n(z_1, \dots, z_n) = 0, 1 \leq i \leq n-1\}.$$

Then $\Lambda = \pi_n(\Omega)$ and thus Ω is a one-dimensional complex algebraic variety in \mathbb{D}^n . Since $\pi_n(\mathbb{T}^n) = b\Gamma_n$, it follows that Ω exits through the distinguished boundary \mathbb{T}^n of \mathbb{D}^n . Hence Ω is a distinguished variety in \mathbb{D}^n .

Conversely, suppose Ω is a distinguished variety in \mathbb{D}^n . Then by Theorem 3.1, Ω has complex dimension 1 and by Eisenbud-Evans theorem (see [8]), there exist n polynomials

g_1, \dots, g_n in $\mathbb{C}[z_1, \dots, z_n]$ that generate Ω . For $i = 1, \dots, n$ let

$$\tilde{g}_i(z_1, \dots, z_n) = \prod_{(\sigma(z_1), \dots, \sigma(z_n)) \in S_n} g_i(\sigma(z_1), \dots, \sigma(z_n)),$$

where S_n is the group of permutation of $\{1, \dots, n\}$. If V_i and \tilde{V}_i are the algebraic sets generated by g_i and \tilde{g}_i respectively, then $\pi_n(\tilde{V}_i) = \pi_n(V_i)$ for each i . Thus if $S = \{g_1, \dots, g_n\}$ and $S' = \{\tilde{g}_1, \dots, \tilde{g}_n\}$, then $\Omega = V_S \cap \mathbb{D}^n = V_{S'} \cap \mathbb{D}^n$. Again since \tilde{g}_i is a symmetric polynomial, $\tilde{g}_i(z_1, \dots, z_n) = f_i \circ \pi_n(z_1, \dots, z_n)$ for some polynomial f_i . Let

$$\Lambda = \{(s_1, \dots, s_{n-1}, p) \in \mathbb{G}_n : f_i(s_1, \dots, s_{n-1}, p) = 0, 1 \leq i \leq n\}.$$

Then $\Lambda = \pi_n(\Omega)$ and Λ is a one-dimensional complex algebraic variety in \mathbb{G}_n . Since $\pi_n(\mathbb{T}_n) = b\Gamma_n$, Λ exits through the distinguished boundary $b\Gamma$. Hence $\Lambda = \pi_n(\Omega)$ is a distinguished variety in \mathbb{G}_n and the proof is complete. \square

Corollary 3.4. *Every distinguished variety in the polydisc \mathbb{D}^n is a complete intersection.*

This is straight-forward because for any distinguished variety Ω in \mathbb{D}^n , if $\Lambda = \pi_n(\Omega)$ then $\{f_1 \circ \pi_n, \dots, f_{n-1} \circ \pi_n\}$ is a set of generators for Ω while $\{f_1, \dots, f_{n-1}\}$ is a generating set for Λ .

Thus combining Theorem 2.5, Theorem 3.1, Theorem 3.3 and Corollary 3.4, we obtain the following representation of distinguished varieties in the polydisc.

Theorem 3.5. *A set $\Omega \subset \mathbb{D}^n$ is a distinguished variety in \mathbb{D}^n if and only if Ω is an affine algebraic curve in \mathbb{D}^n , a complete intersection and there are $n - 1$ complex square matrices F_1, \dots, F_{n-1} of same order such that the following conditions are satisfied:*

- (i) $[F_i, F_j] = 0$ and $[F_i^*, F_{n-j}] = [F_j^*, F_{n-i}]$, for $1 \leq i < j \leq n - 1$;
- (ii) $\sigma_T(F_1^* + zF_{n-1}, F_2^* + zF_{n-2}, \dots, F_{n-1}^* + zF_1, zI) \subseteq \mathbb{G}_n, \forall z \in \mathbb{D}$;
- (iii) *the set of polynomials $S = \{f_1, \dots, f_{n-1}\} \subset \mathbb{C}[z_1, \dots, z_n]$, where $f_i = \det(F_i^* + pF_{n-i} - s_i I)$, forms a regular sequence ;*
- (iv) *either the set*

$$\Lambda = \{(s_1, \dots, s_{n-1}, p) \in \mathbb{G}_n : (s_1, \dots, s_{n-1}) \in \sigma_T(F_1^* + pF_{n-1}, F_2^* + pF_{n-2}, \dots, F_{n-1}^* + pF_1)\},$$

coincides with $V_S \cap \mathbb{G}_n$, or Λ is an irreducible component of $V_S \cap \mathbb{G}_n$;

- (v) *the set $S' = \{g_i \in \mathbb{C}[z_1, \dots, z_n] : g_i = f_i \circ \pi_n, i = 1, \dots, n - 1\}$ generates Ω , that is, either $\Omega = V_{S'} \cap \mathbb{D}^n$, or Ω is an irreducible component of $V_{S'} \cap \mathbb{D}^n$.*

Proof. Let there be $n - 1$ complex square matrices of same order such that the conditions (i) – (iv) are satisfied. Then Λ is a distinguished variety in \mathbb{G}_n , by Theorem 2.5. Evidently condition-(v) implies that $\Lambda = \pi_n(\Omega)$ and thus by Theorem 3.3, Ω is a distinguished variety in \mathbb{D}^n .

Conversely, suppose that Ω is a distinguished variety in \mathbb{D}^n . Then by Theorem 3.1 and Corollary 3.4, Ω is an affine algebraic curve in \mathbb{D}^n and is a complete intersection. Also by Theorem 3.3, there exists a distinguished variety Λ in \mathbb{G}_n such that $\Lambda = \pi_n(\Omega)$. Again since Λ is a distinguished variety in \mathbb{G}_n , by Theorem 2.5, there are $n - 1$ complex square matrices

F_1, \dots, F_{n-1} of same order such that the conditions (i) – (iv) are satisfied. Condition-(v) follows from the proof of Theorem 3.3. Hence the proof is complete. \square

Remark 3.6. Needless to mention that the preceding theorem is a generalization of the famous result due to Agler and McCarthy ([1], Theorem 1.12) describing the distinguished sets in the bidisc \mathbb{D}^2 . Their description of a distinguished set V in \mathbb{D}^2 is given in terms of a matrix-valued rational function in the following way:

$$V = \{(z, w) \in \mathbb{D}^2 : \det(\psi(z) - wI) = 0\},$$

where $\psi(z) = A + zB(I - zD)^{-1}C$ is an $m \times m$ -matrix-valued rational function defined on \mathbb{D} by the entries of a unitary $(m+n) \times (m+n)$ -matrix U given by

$$U = \begin{pmatrix} A & B \\ C & D \end{pmatrix} : \mathbb{C}^m \oplus \mathbb{C}^n \rightarrow \mathbb{C}^m \oplus \mathbb{C}^n.$$

For $n = 2$, our representation of a distinguished variety as in Theorem 3.5 is simpler and is given by determinant of a matrix-valued polynomial in the following way:

$$V = \{(z, w) \in \mathbb{D}^2 : \det(A^* + Azw - (z+w)I) = 0\},$$

where A is any square matrix such that each joint eigenvalue (s, p) of $(A^* + Ap, sI)$ is in the symmetrized bidisc \mathbb{G}_2 .

We conclude this article by proving that the closure of every distinguished variety is polynomially convex.

Proposition 3.7. *The closure of any distinguished variety in \mathbb{D}^n is polynomially convex.*

Proof. Suppose Ω is a distinguished variety in \mathbb{D}^n and suppose $\Omega = V \cap \mathbb{D}^n$ for some variety V . Without loss of generality let us assume that $\{f_1, \dots, f_k\}$ is a set of generators for V . Let $y = (y_1, \dots, y_n) \in \mathbb{C}^n \setminus \overline{\mathbb{D}^n}$. Since $\overline{\mathbb{D}^n}$ is polynomially convex, there is a polynomial say $f \in \mathbb{C}[z_1, \dots, z_n]$ such that

$$|f(y)| > \sup_{z \in \overline{\mathbb{D}^n}} |f(z)| = \|f\|_{\infty, \overline{\mathbb{D}^n}} \geq \|f\|_{\infty, \overline{\Omega}}.$$

Now let $z = (z_1, \dots, z_n) \in \overline{\mathbb{D}^n} \setminus \overline{\Omega}$. Then, there is at least one of f_1, \dots, f_k say f_t such that $f_t(z) \neq 0$. It follows that

$$|f_t(z)| > \sup_{x \in \overline{\Omega}} |f_t(x)| = \|f_t\|_{\infty, \overline{\Omega}} = 0.$$

Hence $\overline{\Omega}$ is polynomially convex. \square

Acknowledgement. The author is grateful to Sudhir Ghorpade and Souvik Goswami for stimulating conversations on basic algebraic geometry.

REFERENCES

- [1] J. Agler and J.E. McCarthy, Distinguished varieties, *Acta Math.*, 194 (2005), no. 2, 133 – 153.
- [2] J. Agler and J.E. McCarthy, Parametrizing distinguished varieties, *Contemp. Math.*, 393 (2006), 29 – 34.
- [3] T. Ando, On a pair of commutative contractions, *Acta Sci Math* 24 (1963), 88 – 90.
- [4] T. Bhattacharyya, P. Kumar and H. Sau, Distinguished varieties through the Berger-Coburn-Lebow theorem, To appear in *Anal. PDE*, available at <https://arxiv.org/abs/2001.01410>.
- [5] B. Bisai and S. Pal, A certain rational function and operator theory on the symmetrized polydisc, *J. Math. Anal. Appl.*, 495 (2021), Paper No. 124714, 15 pp.
- [6] R. E. Curto, Applications of several complex variables to multiparameter spectral theory, Surveys of Some Recent Results in Operator Theory, Vol. II, *Pitman Res. Notes Math. Ser., Longman Sci. Tech., Harlow*, 192 (1988), 25-90.
- [7] B. K. Das and J. Sarkar, Ando dilations, von Neumann inequality, and distinguished varieties, *J. Funct. Anal.*, 272 (2017), 2114 – 2131.
- [8] D. Eisenbud and E. Graham Evans, Jr., Every algebraic set in n -space is the intersection of n hypersurfaces, *Inventiones Math.*, 19 (1973), 107 – 112.
- [9] R. Hartshorne, Algebraic Geometry, *Graduate Texts in Mathematics; Springer-Verlag New York*, ISBN 0-387-90244-9, 1997.
- [10] S. Pal, Canonical decomposition of operators associated with the symmetrized polydisc, *Complex Anal. Oper. Theory*, 12 (2018), 931 – 943.
- [11] S. Pal, Distinguished varieties in a families of domains associated with spectral interpolation and operator theory, *Preprint*, available at <https://arxiv.org/abs/2010.16077>.
- [12] S. Pal and O. M. Shalit, Spectral sets and distinguished varieties in the symmetrized bidisc, *J. Funct. Anal.*, 266 (2014), 5779 – 5800.
- [13] D. Scheinker, Hilbert function spaces and the Nevanlinna-Pick problem on the polydisc, *J. Funct. Anal.*, 261 (2011), 2238 – 2249.
- [14] J. L. Taylor, The analytic-functional calculus for several commuting operators, *Acta Math.* 125 (1970), 1 – 38.
- [15] J. L. Taylor, A joint spectrum for several commuting operators, *J. Funct. Anal.*, 6 (1970), 172 – 191.

(Sourav Pal) MATHEMATICS DEPARTMENT, INDIAN INSTITUTE OF TECHNOLOGY BOMBAY, POWAI, MUMBAI - 400076, INDIA.

Email address: sourav@math.iitb.ac.in, souravmaths@gmail.com