

FULL EXCEPTIONAL COLLECTIONS FOR ANTICANONICAL LOG DEL PEZZO SURFACES

GIULIA GUGLIATTI AND FRANCO ROTA

ABSTRACT. Motivated by homological mirror symmetry, this paper constructs explicit full exceptional collections for the canonical stacks associated with the series of log del Pezzo surfaces constructed by Johnson and Kollár in [JK01]. These surfaces have cyclic quotient, non-Gorenstein, singularities. The construction involves both the $\mathrm{GL}(2, \mathbb{C})$ McKay correspondence, and the study of the minimal resolutions of the surfaces, which are birational to degree two del Pezzo surfaces. We show that a degree two del Pezzo surface arises in this way if and only if it admits a generalized Eckardt point, and in the course of the paper we classify the blow-ups of \mathbb{P}^2 giving rise to them. Our result on the adjoints of the functor of Ishii–Ueda [IU15] applies to any finite small subgroup of $\mathrm{GL}(2, \mathbb{C})$.

CONTENTS

1. Introduction	1
2. Preliminaries	4
3. Computation of left adjoint functor	10
4. The minimal resolution of X	15
5. The derived category of \mathcal{X}	22
References	28

1. INTRODUCTION

For each positive integer k , consider the family of anticanonical del Pezzo surfaces

$$(1) \quad X = X_{8k+4} \subset \mathbb{P}(2, 2k+1, 2k+1, 4k+1).$$

With the exception of 22 sporadic cases, these surfaces are all the quasismooth wellformed anticanonical del Pezzo weighted hypersurfaces [JK01]. They have isolated, noncanonical singularities which are locally qG-rigid in the sense of [ACC⁺16]. Motivated by homological mirror symmetry, summarized in Section 1.1 below, we study the derived category of coherent sheaves of the canonical stack of X . The main result is the following.

Theorem 1.1 (= (5.2)). *Let \mathcal{X} be the canonical stack associated with X . Then $D^b(\mathrm{Coh}(\mathcal{X}))$ admits full exceptional collections, given explicitly in Section 5.*

The exceptional collections of Theorem 1.1 are formed by $10 + 12k$ objects. In Section 5.2 we also exhibit a collection whose nontrivial morphisms are only in degree 0 and 1, which in turn implies the existence of a tilting complex on \mathcal{X} (see Section 5.3).

The starting point of our construction is the $\mathrm{GL}(2, \mathbb{C})$ McKay correspondence, as stated in [IU15]: let X be a surface with finite quotient singularities, let \tilde{X} be its minimal resolution, and let \mathcal{X} be the canonical stack associated with X . Then, there is a fully faithful functor

2020 *Mathematics Subject Classification*. Primary 14F08, 14A20, 14E16, 14J45; secondary 14J33.

Key words and phrases. Derived categories, canonical smooth Deligne–Mumford stacks, McKay correspondence, log del Pezzo surfaces, mirror symmetry.

$\Phi: D^b(\text{Coh}(\tilde{X})) \rightarrow D^b(\text{Coh}(\mathcal{X}))$. Moreover, if X has only cyclic singularities, there exists a semiorthogonal decomposition (SOD)

$$(2) \quad D^b(\text{Coh}(\mathcal{X})) = \left\langle \mathbf{e}_1, \dots, \mathbf{e}_N, \Phi \left(D^b(\text{Coh}(\tilde{X})) \right) \right\rangle,$$

where $(\mathbf{e}_1, \dots, \mathbf{e}_N)$ is an exceptional collection determined by the singularities of X .

The exceptional sequences of Theorem 1.1 are constructed by applying the above decomposition applied to the surfaces (1), together with:

- (i) a detailed description of $D^b(\text{Coh}(\tilde{X}))$;
- (ii) a computation of the images of $\mathbf{e}_1, \dots, \mathbf{e}_N$ through the left adjoint functor Ψ to Φ .

To obtain (i) and (ii), it is necessary to prove two results which may be of independent interest. We give a brief overview here.

The first result describes the geometry of the minimal resolution \tilde{X} of the surfaces (1).

Theorem 1.2 (= (4.3)). *Let \tilde{X} be the minimal resolution of X . Then \tilde{X} is obtained from a degree 2 del Pezzo surface X' via a sequence of $1 + 4k$ blow-ups.*

As an immediate consequence, full exceptional collections for \tilde{X} , and hence \mathcal{X} , exist. The surface X' appearing in Theorem 4.3 does not depend on the integer k and it admits a generalized Eckardt point, i.e. a point of intersection of four (-1) -curves. Del Pezzo surfaces of degree 2 with a generalized Eckardt point form a stratum of codimension two in the moduli space of all del Pezzo surfaces [Dol12, Table 8.9]. Since every such surface appears as one of the X' , the following geometric corollary holds.

Corollary 1.3 (= (4.6)). *For all $k > 0$ integer, the moduli stack of hypersurfaces of degree $8k + 4$ in $\mathbb{P}(2, 2k + 1, 2k + 1, 4k + 1)$ is birational to the moduli space of del Pezzo surfaces with a generalised Eckardt point.*

In Proposition 4.9 we classify all configurations of seven points on \mathbb{P}^2 which, once blown up, give rise to a del Pezzo surface with a generalized Eckardt point. This classification is then used to describe the Hom-spaces of the exceptional collections of \mathcal{X} .

Our second result computes the image via Ψ ¹ of simple sheaves $\mathcal{O}_0 \otimes \rho \in D([\mathbb{C}^2/G])$, where G is any finite small subgroup of $\text{GL}(2, \mathbb{C})$ and ρ is an irreducible representation of G . Consider the minimal resolution of \mathbb{C}^2/G , let E_1, \dots, E_m be the exceptional curves and let E be the fundamental cycle. Write ρ_i , with $i = 0, \dots, n$, for the irreducible representations of G , and assume that ρ_i , with $i = 1, \dots, m$, is the special representation corresponding to E_i via the McKay correspondence (see Section 2.3).

Theorem 1.4 (= (3.1)). *Let ρ_{nat} be the natural representation of $G \subset \text{GL}(2, \mathbb{C})$ and $\rho_{\text{det}} := \Lambda^2 \rho_{\text{nat}}$. For $i \in \{1, \dots, m\}$ let $-\alpha_i$ be the self intersection of the exceptional curve E_i . Then we have:*

$$(3) \quad \Psi(\mathcal{O}_0 \otimes \rho) \simeq \begin{cases} \mathcal{O}_E(E)[1] & \text{if } \rho = \rho_{\text{det}}^* \\ \mathcal{O}_{E_i}(-\alpha_i + 1) & \text{if } \rho \otimes \rho_{\text{det}} = \rho_i \text{ for some } 1 \leq i \leq m \\ 0 & \text{otherwise.} \end{cases}$$

In Section 3.1, we apply Theorem 1.4 to give an alternative proof of [IU15, Proposition 1.1], which states that the essential image of Φ is generated by $\{\mathcal{O}_{\mathbb{C}^2} \otimes \rho_i\}_{i=1}^m$, and its right orthogonal by $\{\mathcal{O}_{\mathbb{C}^2} \otimes \rho_i\}_{i=m+1}^n$.

To describe morphism spaces of the SOD (2), one can compute either the images through Φ of objects in $D(\tilde{X})$, or the images $\Psi(\mathbf{e}_i)$ and use adjunction. The latter approach is simpler and gives a more general answer. It is simpler since the computations take place on the smooth surface \tilde{X} and not on \mathcal{X} . On the other hand, our Theorem 1.4 shows that the objects $\Psi(\mathcal{O}_0 \otimes \rho)$ do not depend on the global properties of \tilde{X} . In fact, the formula (3) determines $\Psi(\mathbf{e}_i)$ for all

¹In the rest of the paper, we will distinguish local and global functors by denoting them with lower-case and upper-case letters.

surfaces with quotient singularities by finite small groups, since the \mathbf{e}_i all have a composition series whose factors are simple sheaves $\mathcal{O}_0 \otimes \rho$ [IU15, Section 3].

1.1. Related works and further problems. The main motivation for this work comes from mirror symmetry for Fano varieties – also known as Fano/Landau–Ginzburg (LG) correspondence – from the point of view of Kontsevich’s homological mirror symmetry (HMS) [Kon95a]. The correspondence predicts that the mirror of a n -dimensional Fano orbifold \mathcal{X} is a LG model (Y, w) of the same dimension², and can be interpreted at different levels.

Its Hodge-theoretic formulation identifies the quantum differential operator of \mathcal{X} and the Picard–Fuchs operator of (Y, w) [Gol07, Gol08, Prz07]. Recent works suggest that the Hodge-theoretic mirrors of Fano orbifolds with \mathbb{Q} -Gorenstein (qG) toric degenerations should be LG models restricting to specific classes of Laurent polynomials. This perspective, first laid out in [Gol07, Prz11, CCG⁺13], agrees with the Givental/Hori–Vafa mirror construction [Giv98, HV00] for toric complete intersections and the intrinsic mirror symmetry program [GS18] for log Calabi–Yau pairs.

One formulation of HMS³ predicts an equivalence $D^b(\mathrm{Coh}(\mathcal{X})) \cong \mathrm{DLag}_{\mathrm{vc}}(Y, w)$ between the bounded derived category of coherent sheaves of \mathcal{X} and the derived category of Lagrangian vanishing cycles of (Y, w) [Sei08]⁴. It is expected that HMS implies Hodge-theoretic mirror symmetry (see [Kon95b], and [GPS15]), even though the relation between these two formulations is not proven in full generality. Unlike Hodge-theoretic mirror symmetry, HMS has been proven systematically only for few classes of Fano varieties; among them, smooth del Pezzo surfaces [AKO06], and weighted projective planes [AKO08].

The focus of our work is on the surfaces (1), which we think of as canonical stacks. These surfaces have isolated log terminal singularities which are locally qG-rigid, hence they fit into the mirror symmetry context of [ACC⁺16]. The work [CG21] shows that their quantum differential operator is hypergeometric, and uses hypergeometric motives to build their Hodge-theoretic mirrors: these are pencils of hyperelliptic curves which do not restrict to Laurent polynomials. The mirror construction in [CG21] is the only one known for the surfaces (1). Indeed, since their anticanonical linear system is empty, none of the standard methods apply (see [CG21, Remark 2.7]).

This paper is the first step towards the study of HMS for the surfaces (1), as it describes exceptional collections of $D^b(\mathrm{Coh}(\mathcal{X}))$. A construction of other exceptional collections for \mathcal{X} using other methods appears to be challenging: for example, neither X nor \tilde{X} are toric, so the Kawamata construction [Kaw06] does not apply. Nevertheless, the surfaces X are hypersurfaces in weighted projective spaces, so they inherit a – not full – exceptional collection from the ambient space. This collection appears in Elagin [Ela07]. There, the author uses it to build a full exceptional collection for the stack of a surface with one log terminal singularity via an *ad hoc* mutation of a set of generators for the residual category. There seems to be no systematic way of extending the construction [Ela07] to other cases.

Structure of the paper. Section 2 recalls definitions and basic properties. We set some toric geometry notation in 2.1. Then we recall properties of log del Pezzo surfaces in 2.2 and the $\mathrm{GL}(2, \mathbb{C})$ McKay correspondence in 2.3.

Section 3 contains the computation of the adjoint functor Ψ , and the alternative proof of [Ish02, Prop. 1.1] in 3.1. In Section 3.2, we rephrase Theorem 1.4 in the cyclic singularities setting, and reprove it using toric geometry.

Section 4 is dedicated to the minimal resolutions and contains the proof of Theorem 1.2. In 4.3 we classify degree 2 del Pezzo surfaces admitting a generalized Eckardt point.

²In mirror symmetry, it is important to think of a Fano variety X with quotient singularities as a canonical smooth Deligne–Mumford stack \mathcal{X} . This allows one to define Gromov–Witten invariants for \mathcal{X} . See [OP18, Section 3] for a concise summary of the quantum period of a Fano orbifold.

³A parallel formulation, which translates the Hodge-theoretic version of mirror symmetry mentioned above, is an equivalence between a suitable Fukaya category of \mathcal{X} and the category of matrix factorisations of (Y, w) .

⁴A rigorous definition of this category has been proposed by Seidel in the case where (Y, w) is a Lefschetz fibration.

In Section 5 we show the existence of full exceptional collections for \tilde{X} and \mathcal{X} and describe them explicitly. We discuss the existence of a tilting complex for \tilde{X} and \mathcal{X} in 5.3.

Notation and conventions. We work over the field of complex numbers. We use $D(-) = D^b(\text{Coh}(-))$ to denote the bounded derived category of coherent sheaves and $D_G(-)$ to denote the bounded derived category of G -equivariant coherent sheaves. We write $\mathcal{O}(n)$ to denote the line bundle of degree n in the (weighted) projective space \mathbb{P} . For $C \subset \mathbb{P}$ a rational curve, we write $\mathcal{O}_C(1)$ for its line bundle of degree one, so that $\mathcal{O}(1)|_C = \mathcal{O}_C(\deg C)$.

Acknowledgements. We are thankful to Arend Bayer, Alessio Corti, and Michael Wemyss for helpful discussions and advice. We thank Akira Ishii for his remarks on Section 3.1. The second author was partially supported by EPSRC grant EP/R034826/1.

2. PRELIMINARIES

This section is organised as follows: in 2.1 we set up our notation for toric varieties and we state some well-known facts which we will use repeatedly; in 2.2 we recall the notion of anticanonical log del Pezzo surface, the classification [JK01], and some basic facts about smooth del Pezzo surfaces of degree two; 2.3 is a summary of the $\text{GL}(2, \mathbb{C})$ McKay correspondence, with a focus on the case of cyclic quotient singularities.

2.1. Toric setup. For a lattice L , we denote by $L_{\mathbb{R}} = L \otimes_{\mathbb{Z}} \mathbb{R}$ the associated real vector space. For a torus \mathbb{T} , we denote by $M = \text{Hom}(\mathbb{T}, \mathbb{C}^{\times})$ the character lattice and we write $N = \text{Hom}(M, \mathbb{Z})$. Given vectors $v_1, \dots, v_k \in L_{\mathbb{R}}$, we write $\langle v_1, \dots, v_k \rangle$ for the cone they generate.

For a fan $\Sigma \subset N_{\mathbb{R}}$, we denote by F_{Σ} the associated toric variety. The toric variety is proper if and only if Σ is a complete fan, that is, its support $|\Sigma|$ is all $N_{\mathbb{R}}$. The toric variety has finite quotient singularities if and only if the fan is simplicial. We write $\Sigma(r)$ for the set of r -dimensional cones of the fan Σ . We denote by ρ_i the primitive generators of the 1-dimensional cones (or rays) of Σ , and by $D_{\rho_i} \subset F_{\Sigma}$ the corresponding divisors under the toric orbit-cone correspondence.

Below we recall the construction of star subdivision of a fan.

Lemma/Definition 2.1. *Let $\Sigma \subset N_{\mathbb{R}}$ be a fan with rays ρ_1, \dots, ρ_n and let $v \in |\Sigma| \cap N$ be a primitive element. Define the star subdivision of Σ at v , denoted by $\Sigma^*(v)$, as the set of the cones:*

- (1) σ , where $v \notin \sigma \in \Sigma$;
- (2) $\text{cone}(\tau, v)$, where $v \notin \tau \in \Sigma$ and $\{v\} \cup \tau \subset \sigma \in \Sigma$.

The set $\Sigma^*(v)$ is a refinement of Σ , thus induces a toric morphism $X_{\Sigma^*(v)} \rightarrow X_{\Sigma}$.

In Section 4 we will use toric morphisms coming from star subdivisions to resolve singularities of toric varieties.

In particular, we will consider situations where Σ contains a cone τ with rays $\rho_{i_1}, \dots, \rho_{i_r}$ such that the corresponding affine chart U_{τ} is of the form:

$$U_{\tau} \simeq \mathbb{C}^r / \mathbb{Z}_m(a_1, \dots, a_r) \times (\mathbb{C}^{\times})^{n-r} \subset X_{\Sigma}$$

where m, a_i are positive integers such that $a_i < m$ for all i . Then the toric variety $X_{\Sigma^*(v)}$ associated to the star subdivision $\Sigma^*(v)$, with:

$$(4) \quad v = \frac{1}{m} \sum_{i=1}^r a_j \rho_{i_j}$$

is the weighted blow-up of X_{Σ} along $V(\tau)$ with weights $\frac{1}{m}(a_1, \dots, a_r)$, where we write $V(\tau)$ for the closure of the \mathbb{T}_N -orbit in X_{Σ} corresponding to τ . As such, the variety $X_{\Sigma^*(v)}$ fits in a fiber square:

$$\begin{array}{ccc} \mathcal{E} & \hookrightarrow & X_{\Sigma^*(v)} \\ \downarrow & & \downarrow \\ V(\tau) & \hookrightarrow & X_{\Sigma} \end{array}$$

where the exceptional divisor \mathcal{E} is a $\mathbb{P}(a_1, \dots, a_r)$ -bundle over $V(\tau)$.

2.2. Log del Pezzo weighted hypersurfaces. For positive integers $a_i, i = 1, \dots, m$, we denote by $S = S(a_1, \dots, a_m)$ the graded polynomial ring $\mathbb{C}[x_1, \dots, x_m]$ where $\deg x_i = a_i$, and by $\mathbb{P}(a_1, \dots, a_m)$ the weighted projective space with weights a_1, \dots, a_m . We simply write \mathbb{P} if the weights and m are clear from the context. We write \mathbb{P}_{sing} for the singular locus of \mathbb{P} . The space \mathbb{P} is said *wellformed* if the quotient stack $\mathbf{P} = [\text{Spec } S \setminus \{0\}/\mathbb{C}^\times]$ is canonical, that is, the natural morphism $s: \mathbf{P} \rightarrow \mathbb{P}$ is an isomorphism in codimension 1.

A subvariety $X \subset \mathbb{P}$ is called *quasismooth* of dimension m if its affine cone $C_X \subset \mathbb{C}^{m+1}$ is smooth of dimension $m + 1$ outside of its vertex $\underline{0}$. This implies that the singularities of X are only due to the \mathbb{C}^\times -action defining \mathbb{P} , hence are cyclic quotient singularities. A subvariety $X \subset \mathbb{P}$ of codimension c is called *wellformed* if \mathbb{P} is wellformed and $\text{codim}_X(\mathbb{P}_{\text{sing}} \cap X) \geq 2$. Observe that, if $X = \text{Proj}(S/I) \subset \mathbb{P}$, where I is a homogeneous ideal of S , is quasismooth and wellformed, then X is the coarse moduli space of the canonical stack $[\text{Spec}(S/I) \setminus \{0\}/\mathbb{C}^\times]$.

We denote by $X_d \subset \mathbb{P}(a_1, \dots, a_m)$ a sufficiently general member of the family of all hypersurfaces of weighted degree d . For $X = X_d \subset \mathbb{P}(a_1, \dots, a_m)$ a quasismooth and wellformed hypersurface of degree d , the adjunction formula for the canonical sheaf K_X [IF00, Section 6.14] states that

$$K_X = \mathcal{O}_X(d - \sum_{i=1}^m a_i)$$

By definition X is Fano if $-K_X$ is ample, i.e., if and only if $\sum_{i=1}^m a_i - d > 0$. X is called *anticanonical* if $-K_X = \mathcal{O}_X(1)$, i.e., if and only if $\sum_{i=1}^m a_i - d = 1$.

A Fano surface with quotient singularities is called a *log del Pezzo surface*. Johnson and Kollár [JK01] classify all anticanonical quasismooth wellformed log del Pezzo surfaces in weighted projective 3-spaces $X_d \subset \mathbb{P}(a_1, a_2, a_3, a_4)$. The classification consists of the series (1), where $k \in \mathbb{N}$, $k > 0$, and 22 sporadic cases. The 22 sporadic cases are all listed in [JK01, Theorem 8].

For all $k > 0$ the surfaces (1) have isolated qG-rigid, log-terminal singularities (see Section 4.1). Among the sporadic cases, three are smooth surfaces; precisely: cubic surfaces $X_3 \subset \mathbb{P}^3$, degree-two del Pezzo surfaces $X_4 \subset \mathbb{P}(1, 1, 1, 2)$, and degree-one del Pezzo surfaces $X_6 \subset \mathbb{P}(1, 1, 2, 3)$.

2.2.1. Degree-two del Pezzo surfaces. We will prove in Section 4 that the geometry of the series (1) is related to the geometry of some special degree-two smooth del Pezzo surfaces. Below we recall some basic facts about degree-two smooth del Pezzo surfaces which we will use repeatedly.

Let S be a smooth del Pezzo surface of degree two. Then S is isomorphic to the blow-up of \mathbb{P}^2 at 7 distinct points in general position [Man86, IV.24.4][Dem80, Thm 1]. Conversely, the choice of a set $\mathcal{L} = \{l_1, \dots, l_7\}$ of 7 disjoint (-1) -curves on S is equivalent to the choice of a birational morphism $\pi: S \rightarrow \mathbb{P}^2$ which expresses S as the blowup of \mathbb{P}^2 at the 7 points $x_i = \pi(l_i)$. We refer to such a set \mathcal{L} as to an *exceptional set* for S .

On the other hand, S is expressed by its anticanonical series as a double cover $\sigma: S \rightarrow \mathbb{P}^2$ branched along a quartic curve $Q = (f_4(x_1, x_2, x_3) = 0) \subset \mathbb{P}^2$ [Kol96, III.3.5] Hence, S can be defined by an equation of the form

$$(x_4^2 - f_4(x_1, x_2, x_3) = 0) \subset \mathbb{P}(1, 1, 1, 2).$$

The surface S contains 56 (-1) -curves, which can be described in terms of the morphisms σ and π . On the one hand, the 56 (-1) -curves are the preimages through σ of the 28 lines bitangent to Q . Each preimage is a reducible curve of the form $m \cup m'$ where m, m' are (-1) -curves on S satisfying $m + m' \sim -K_S$ (we refer to m and m' as *dual curves*). Thus any two of the 56 (-1) -curves are either disjoint, intersect at one point, or intersect at two points precisely if they are dual to each other. On the other hand, given an exceptional set $\mathcal{L} = \{l_1, \dots, l_7\}$ for S and the corresponding blow-down morphism $\pi: S \rightarrow \mathbb{P}^2$, the 28 pairs of (-1) -curves described above are the strict transforms through π of:

- the line through two points x_i, x_j and the conic through the remaining 5 points (21 pairs);

- the cubic with a node at one of the x_i and passing through the remaining 6 points, paired with the exceptional divisor l_i (7 pairs).

By the proof of [TVAV09, Lemma 4.1], at most four bitangents to Q can meet at one point, thus at most four (-1) -curves on S can meet at one point of S . Such points on S are called *generalized Eckardt points*.

Denote by $I^{1,7}$ the lattice \mathbb{Z}^8 together with the symmetric bilinear form defined by the diagonal matrix with diagonal entries $(1, -1, \dots, -1)$ with respect to the standard basis

$$e_0 = (1, 0, \dots, 0)^T, \quad e_1 = (1, 0, \dots, 0)^T, \quad \dots, \quad e_7 = (0, 0, \dots, 1)^T.$$

The Picard lattice $\text{Pic}(S)$ is isomorphic to the lattice $I^{1,7}$ [Dol12, Section 8.2.6]. The set of (-1) -curves on S is in bijection with the set of exceptional vectors in $I^{1,7}$, that is, the vectors $v \in I^{1,7}$ such that $v \cdot k = -1$, where $k = (3, -1, \dots, -1)^T$, and $v^2 = -1$. These vectors are listed in [Dol12, Proposition 8.2.19].

2.3. $\text{GL}(2, \mathbb{C})$ McKay correspondence. Suppose G is a finite small subgroup of $\text{GL}(2, \mathbb{C})$ acting naturally on $S = \mathbb{C}[x, y]$ and let $R = S^G$. We can associate two resolutions to the singularity $X = \text{Spec } R$: its minimal resolution $f: Y \rightarrow X$, and a *non-commutative crepant resolution* ([vdB04a, Definition 4.1]) given by the skew group algebra $S * G$. The category of finitely generated $S * G$ -modules is equivalent to the category of G -equivariant sheaves on \mathbb{C}^2 , and therefore to the category of coherent sheaves over the quotient stack $[\mathbb{C}^2/G]$:

$$\text{mod-}S * G \simeq \text{Coh}[\mathbb{C}^2/G].$$

There are bijective correspondences of irreducible representations of G , indecomposable Cohen-Macaulay R -modules, indecomposable projective $S * G$ -modules, and indecomposable full sheaves on Y [Esn85, Remark 1.4, Lemma 2.2]. More explicitly, to an irreducible representation ρ of G , we associate a free S -module⁵ $S \otimes \rho$, the CM R -module

$$M_\rho := (S \otimes \rho^*)^G$$

and the coherent sheaf $\widetilde{M}_\rho := f^*(M_\rho)/\text{tors.}$ on Y (sheaves arising this way are called *full*).

Moreover, irreducible exceptional curves of Y are in bijection with *special* representations of G , i.e. irreducible representations ρ for which $H^1(\widetilde{M}_\rho^\vee) = 0$. Let E_1, \dots, E_m be the exceptional curves on Y : Wunram [Wun88, Theorem 1.2] shows that there are exactly $m + 1$ special representations $\rho_0, \rho_1, \dots, \rho_m$ of G , where ρ_0 is the trivial representation and the other satisfy the *multiplication formula*

$$c_1(\widetilde{M}_{\rho_i}) \cdot E_j = \delta_{ij}.$$

A refined version of the formula is given by Ishii in [Ish02], and we briefly recall it here.

Consider the equivariant Koszul complex of $0 \in \mathbb{C}^2$:

$$(5) \quad 0 \rightarrow S \otimes \rho_{\det} \xrightarrow{(y, -x)^T} S \otimes \rho_{\text{nat}} \xrightarrow{(x, y)} S \rightarrow \mathcal{O}_0 \rightarrow 0,$$

where ρ_{nat} is the representation induced by $G \subset \text{GL}(2, \mathbb{C})$ and $\rho_{\det} := \Lambda^2 \rho_{\text{nat}}$. Tensoring with ρ^* and taking invariants we obtain the Auslander-Reiten (AR) sequences

$$(6) \quad A_\rho := [0 \rightarrow \tau(M_\rho) \rightarrow N_\rho \rightarrow M_\rho \rightarrow 0],$$

where $N_\rho := (S \otimes \rho_{\text{nat}} \otimes \rho^*)^G$ and $\tau(M_\rho) := (S \otimes \rho_{\det} \otimes \rho^*)^G$. The A_ρ are exact except for $\rho = \rho_0$, in which case

$$0 \rightarrow \tau(R) \rightarrow N_{\rho_0} \rightarrow R \rightarrow \mathcal{O}_0 \rightarrow 0$$

is exact.

Recall that the fundamental cycle of $Y \rightarrow \text{Spec } R$ is $E = \sum_{i=1}^m r_i E_i$, where $r_i \geq 1$ are the unique smallest integers satisfying $E \cdot E_i \leq 0$ for all $i = 1, \dots, m$ [Art66]. An immediate consequence of the proof of [Ish02, Theorem 5.1] is:

⁵By [Aus86, Lemma 1.1], a $S * G$ -module is projective precisely if it is free as a S -module. Then, the $S \otimes \rho$ for $\rho \in \text{Irrep}(G)$ are all the indecomposable projectives.

Proposition 2.2. *Let $\rho \in \text{Irrep}(G)$. Denote by E the fundamental cycle of $f: Y \rightarrow X$ and by $\widetilde{A}_\rho \in D^b(Y)$ the torsion-free pull-back of A_ρ . Then*

$$(7) \quad \widetilde{A}_\rho \simeq \begin{cases} \mathcal{O}_E & \text{if } \rho = \rho_0, \\ \mathcal{O}_{E_i}(-1)[1] & \text{if } \rho = \rho_i \text{ for } 1 \leq i \leq m, \\ 0 & \text{otherwise.} \end{cases}$$

2.3.1. *Derived McKay correspondence.* In the above setting, there is a full and faithful functor $\phi: D(Y) \rightarrow D[\mathbb{C}^2/G]$ constructed as follows. One observes that Y can be constructed as the G -Hilbert scheme parametrizing G -clusters of \mathbb{C}^2 (i.e. finite, G -invariant subschemes $Z \subset \mathbb{C}^2$ such that $H^0(\mathcal{O}_Z)$ is isomorphic as a $\mathbb{C}[G]$ -module to the regular representation of G) [Ish02, Theorem 3.1, Theorem 8.1]. Thus, $Y \times \mathbb{C}^2$ admits a universal family Z , and one can define the diagram

$$(8) \quad \begin{array}{ccc} Z & \subseteq & Y \times \mathbb{A}^2 \xrightarrow{\pi_{\mathbb{C}^2}} \mathbb{A}^2 \\ & & \downarrow \pi_Y \qquad \qquad \downarrow \\ & & Y \xrightarrow{f} X \end{array}$$

and the Fourier-Mukai functor

$$(9) \quad \phi(-) := \pi_{\mathbb{C}^2*}(\pi_Y^*(-) \otimes^{\mathbf{L}} \mathcal{O}_Z).$$

The functor ϕ is an equivalence if and only if Y is a crepant resolution, which happens if and only if $G \subset \text{SL}(2, \mathbb{C})$, equivalently R is Gorenstein. Even in the $G \subset \text{GL}(2, \mathbb{C})$ case, ϕ admits left and right adjoints. So its essential image is admissible and there is a semiorthogonal decomposition:

$$(10) \quad D([\mathbb{C}^2/G]) = \langle \phi(D(Y))^\perp, \phi(D(Y)) \rangle.$$

The essential image of ϕ and its right orthogonal are described as follows. Since the functor $(-)^G$ of taking invariant part defines an equivalence of categories between projective $S * G$ -modules and CM R -modules [Aus86, Sec. 2], the $\{S \otimes \rho\}_{\rho \in \text{Irrep}(G)}$ are projective generators of $\text{mod-}S * G$ and hence they generate $D([\mathbb{C}^2/G])$.

Proposition 2.3. [IU15, Prop. 1.1] ⁶ *The essential image of ϕ in $D([\mathbb{C}^2/G])$ is generated by the modules $\{S \otimes \rho\}$ where ρ is a special representation. The right orthogonal of $\phi(D(Y))$ in $D([\mathbb{C}^2/G])$ is generated by sheaves $\mathcal{O}_0 \otimes \rho$ where $\mathcal{O}_0 = S/(x, y)$ and ρ is not special.*

Remark 2.4. Let μ be an irreducible representation of G and write $a_{\mu\nu}$ for the multiplicity of $\nu \in \text{Irrep}(G)$ in the decomposition of $\mu \otimes \rho_{\text{nat}}$ into irreducible representations. Then by (5) we have

$$\begin{aligned} \dim \text{Hom}(\mathcal{O}_0 \otimes \mu, \mathcal{O}_0 \otimes \nu) &= \delta_{\mu\nu} \\ \dim \text{Ext}^1(\mathcal{O}_0 \otimes \mu, \mathcal{O}_0 \otimes \nu) &= a_{\mu\nu} \\ \dim \text{Ext}^2(\mathcal{O}_0 \otimes \mu, \mathcal{O}_0 \otimes \nu) &= \dim \text{Hom}(\mathcal{O}_0 \otimes \nu, \mathcal{O}_0 \otimes \mu \otimes \rho_{\text{det}}) \end{aligned}$$

This information is summarised in the *McKay quiver* of G , whose vertices represent the irreducible representation of G , and whose solid (resp. dashed) arrows from μ to ν denote a basis of $\text{Ext}^1(\mathcal{O}_0 \otimes \mu, \mathcal{O}_0 \otimes \nu)$ (resp. $\text{Ext}^2(\mathcal{O}_0 \otimes \mu, \mathcal{O}_0 \otimes \nu)$). We give two examples in Figure 1 and Figure 2.

2.3.2. *Cyclic singularities.* For relatively prime integers $1 \leq a < r$, define the group $G = \frac{1}{r}(1, a)$ as the cyclic small subgroup of $\text{GL}(2, \mathbb{C})$ generated by

$$\zeta := \begin{pmatrix} \epsilon & 0 \\ 0 & \epsilon^a \end{pmatrix}$$

where ϵ is a primitive r -th root of unity. We use the same notation $\frac{1}{r}(1, a)$ for the corresponding cyclic quotient singularity $\text{Spec } R$.

⁶In Section 3.1 we provide an alternative proof of Proposition 2.3 based on our Theorem 3.1.

Definition 2.5. For integers $1 \leq a < r$ as above, set $i_0 = r$, $i_1 = a$, and define i_t inductively by

$$(11) \quad i_t := \alpha_{t-1} i_{t-1} - i_{t-2}$$

where $\alpha_{t-1} \geq 2$ and $0 \leq i_t < i_{t-1}$, until $i_n = 1$ and $i_{n+1} = 0$. The I -series $I(r, a)$ is the sequence $I(r, a) := \{i_0 = r, \dots, i_{n+1} = 0\}$.

The equations (11) can be rearranged and spliced together to give the Hirzebruch–Jung continued fraction expansion of $\frac{r}{a}$, i.e.:

$$(12) \quad \frac{r}{a} = \alpha_1 - \frac{1}{\alpha_2 - \frac{1}{\alpha_3 - \frac{1}{\dots}}} =: [\alpha_1, \dots, \alpha_n]$$

It is well-known that the dual graph of the minimal resolution Y of $\frac{1}{r}(1, a)$ is the labeled diagram

$$\begin{array}{ccccccc} \bullet & \text{---} & \bullet & \text{---} & \dots & \text{---} & \bullet & \text{---} & \bullet \\ -\alpha_1 & & -\alpha_2 & & & & -\alpha_{n-1} & & -\alpha_n \end{array}$$

For $l \in \mathbb{Z}$, denote by ρ_l the irreducible representation of G acting with weight l . Then the modules

$$M_l := (\mathbb{C}[x, y] \otimes \rho_l^*)^G = \left\{ f \in \mathbb{C}[x, y] \mid \zeta \cdot f = \epsilon^l f \right\}$$

are the indecomposable CM R -modules⁷, and M_l is special if $l \in I(r, a)$ [Wun87]. In [Wun88, Appendix A1]) it is shown that their torsion free pull-backs \widetilde{M}_l satisfy

$$(13) \quad c_1(\widetilde{M}_l) \cdot E_t = \delta_{l, i_t}.$$

Moreover, since in this case $\rho_{\text{nat}} = \rho_1 \oplus \rho_a$ and $\rho_{\text{det}} = \rho_{a+1}$, the AR-sequences (6) specialise as follows:

$$(14) \quad A_l := [0 \rightarrow M_{l-a-1} \rightarrow M_{l-1} \oplus M_{l-a} \rightarrow M_l \rightarrow 0],$$

and by (13) Proposition 2.2 rewrites as:

Proposition 2.6. Let $l \in \{0, \dots, r-1\}$. Denote by E the fundamental cycle of $f: Y \rightarrow X$ and by $\widetilde{A}_l \in D^b(Y)$ the torsion-free pull-back of the AR-sequence ending in M_l . Then we have

$$(15) \quad \widetilde{A}_l \simeq \begin{cases} \mathcal{O}_E & \text{if } l = 0 \\ \mathcal{O}_{E_t}(-1)[1] & \text{if } l = i_t \in I(r, a) \setminus \{0\} \\ 0 & \text{otherwise.} \end{cases}$$

In the cyclic case, the right orthogonal $\phi(D(Y))^\perp$ admits a full exceptional collection. Every non-special representation ρ contributes one exceptional sheaf, which has finite length and socle $\mathcal{O}_0 \otimes \rho$ [IU15, Sec. 3]. We do not recall here the construction of [IU15], but we point out below that the exceptional sheaves are simple in the two cases we consider in this paper, namely $G = \frac{1}{n}(1, 1)$, $n > 1$ integer, and $G = \frac{1}{2k+1}(1, k)$, $k \geq 1$ integer.

Example 2.7. Let $G = \frac{1}{n}(1, 1)$, for $n > 1$ integer. Then $I(n, 1) = \{n, 1, 0\}$ and $\frac{n}{1} = [n]$. It follows that the minimal resolution $Y = \text{Tot}(\mathcal{O}_{\mathbb{P}^1}(-n))$ has a single exceptional $(-n)$ -curve, ρ_0 and ρ_1 are the only special representations, and the exceptional collection computed in [IU15] reads

$$(16) \quad \phi(D(Y))^\perp = (\mathcal{O}_0 \otimes \rho_2, \dots, \mathcal{O}_0 \otimes \rho_{n-1}).$$

For all i , the McKay quiver of G has two solid arrows from ρ_i to ρ_{i+1} , and one dashed arrow from ρ_i to ρ_{i+1} (see Figure 1).

⁷Clearly, $\rho_l = \rho_m$ and $M_l = M_m$ if $l \equiv m \pmod{r}$.

Example 2.8. Let $G = \frac{1}{2k+1}(1, k)$, for $k \geq 1$ integer. One computes

$$\frac{2k+1}{k} = [3, \underbrace{2, \dots, 2}_{k-1 \text{ times}}],$$

and the relevant terms of the i -series are $i_t = k + 1 - t$ for $t = 1, \dots, k + 1$. It follows that the minimal resolution contains one (-3) -curve E_1 and $k - 1 \times (-2)$ -curves $E_i, i \in \{2, \dots, k\}$, and the special representations are $\rho_{i_t} = \rho_{k+1-t}, t = 1, \dots, k + 1$. The exceptional collection computed in [IU15] reads

$$\phi(D(Y))^\perp = (\mathcal{O}_0 \otimes \rho_{k+1}, \dots, \mathcal{O}_0 \otimes \rho_{2k}).$$

For all i , the McKay quiver of G has a solid arrow from ρ_i to ρ_{i+1} , a solid arrow from ρ_i to ρ_{i+k} , and a dashed arrow from ρ_i to ρ_{i+k+1} .

Figure 1 and 2 depict the McKay quiver of $\frac{1}{5}(1, 1)$ and of $\frac{1}{7}(1, 3)$, with objects and morphisms in $\phi(D(Y))^\perp$ colored in blue.




Figure 1. The case $\frac{1}{5}(1, 1)$.




Figure 2. The case $\frac{1}{7}(1, 3)$.

2.3.3. *Global case.* Suppose X is a complex quasi-projective surface with isolated quotient singularities, i.e. every singular point of X is étale-locally isomorphic to a quotient \mathbb{C}^2/G , where G is a finite subgroup of $\mathrm{GL}(2, \mathbb{C})$. Let \tilde{X} be the minimal resolution of X , and denote by \mathcal{X} the canonical stack associated to X , that is the unique smooth Deligne-Mumford stack \mathcal{X} such that the natural morphism $s: \mathcal{X} \rightarrow X$ is an isomorphism over $X \setminus \mathrm{Sing}(X)$. By [IU15, Theorem 1.4], there is a fully faithful functor $\Phi: D(\tilde{X}) \rightarrow D^b(\mathcal{X})$.

Suppose that X has only cyclic quotient singularities and consider the diagram

$$\begin{array}{ccc} \mathcal{Z} & \subseteq & \tilde{X} \times \mathcal{X} \xrightarrow{\pi_{\mathcal{X}}} \mathcal{X} \\ & & \downarrow \pi_{\tilde{X}} \quad \quad \downarrow s \\ & & \tilde{X} \xrightarrow{f} X \end{array}$$

where \mathcal{Z} is the substack $(\tilde{X} \times_X \mathcal{X})_{\mathrm{red}}$. In this case, [IU15, Prop. 8.1] shows that the Fourier-Mukai functor $\Phi = \pi_{\mathcal{X}*}(\pi_{\tilde{X}}^*(-) \otimes \mathcal{O}_{\mathcal{Z}})$ is fully faithful. Over the smooth locus of X , \mathcal{Z} is the graph of s . On a singular chart of X , \mathcal{W} restricts to the quotient $[Z/G]$ of the universal subscheme of diagram (8). This yields a semiorthogonal decomposition

$$D(\mathcal{X}) = \left\langle \mathbf{e}_1, \dots, \mathbf{e}_N, \Phi(D(\tilde{X})) \right\rangle,$$

where $(\mathbf{e}_1, \dots, \mathbf{e}_n)$ is an exceptional collection, containing the local contributions described in Section 2.3.2.

3. COMPUTATION OF LEFT ADJOINT FUNCTOR

Consider the fully faithful functor ϕ of Equation (9) and the semiorthogonal decomposition (10). By Proposition 2.3, $\phi(D(Y))^\perp$ is generated by simple sheaves of the form $\mathcal{O}_0 \otimes \rho$, where ρ is nonspecial. In this Section we introduce the functor ψ left adjoint to ϕ , and compute the images $\psi(\mathcal{O}_0 \otimes \rho)$ for all ρ .

We follow the same setup and notation of Section 2.3. The left adjoint of ϕ is:

$$(17) \quad \psi(-) = [\mathbf{R}\pi_{Y*} (\pi_{\mathbb{C}^2}^*(- \otimes K_{\mathbb{C}^2}) \otimes^{\mathbf{L}} \mathcal{O}_{\mathbb{Z}}^\vee)]^G[2]$$

where $K_{\mathbb{C}^2} = \mathcal{O}_{\mathbb{C}^2} \otimes \rho_{\det}$ is the G -equivariant canonical sheaf of \mathbb{C}^2 .

Theorem 3.1. *For $i \in \{1, \dots, m\}$ let $-\alpha_i$ be the self intersection of the exceptional curve E_i . Then we have:*

$$\psi(\mathcal{O}_0 \otimes \rho) \simeq \begin{cases} \mathcal{O}_E(E)[1] & \text{if } \rho = \rho_{\det}^* \\ \mathcal{O}_{E_i}(-\alpha_i + 1) & \text{if } \rho \otimes \rho_{\det} = \rho_i \text{ for some } 1 \leq i \leq m \\ 0 & \text{otherwise.} \end{cases}$$

Proof. By definition of ψ we have:

$$(18) \quad \begin{aligned} \psi(\mathcal{O}_0 \otimes \rho) &= [\mathbf{R}\pi_{Y*} (\pi_{\mathbb{A}^2}^*(\mathcal{O}_0) \otimes^{\mathbf{L}} \mathcal{O}_{\mathbb{Z}}^\vee \otimes \rho \otimes \rho_{\det}[2])]^G \\ &= [\mathbf{R}\pi_{Y*} (\mathcal{O}_{Y \times \{0\}} \otimes^{\mathbf{L}} \mathcal{O}_{\mathbb{Z}}^\vee \otimes \rho \otimes \rho_{\det}[2])]^G \\ &= [\mathbf{R}\pi_{Y*} (\mathbf{R}\mathcal{H}om(\mathcal{O}_{\mathbb{Z}}, \mathcal{O}_{Y \times \{0\}} \otimes \rho \otimes \rho_{\det}[2]))]^G \end{aligned}$$

Consider the resolution obtained from pulling back (5) to $Y \times \mathbb{C}^2$:

$$0 \rightarrow \mathcal{O}_{Y \times \mathbb{A}^2} \otimes \rho_{\det} \xrightarrow{(y, -x)^T} \mathcal{O}_{Y \times \mathbb{A}^2} \otimes \rho_{\text{nat}} \xrightarrow{(x, y)} \mathcal{O}_{Y \times \mathbb{A}^2} \rightarrow \mathcal{O}_{Y \times \{0\}} \rightarrow 0.$$

By Grothendieck–Verdier duality

$$(19) \quad \mathbf{R}\pi_{Y*} \mathbf{R}\mathcal{H}om(\mathcal{O}_{\mathbb{Z}}, \mathcal{O}_{Y \times \mathbb{A}^2} \otimes \rho_{\det}[2]) = \mathbf{R}\mathcal{H}om(\mathbf{R}\pi_{Y*} \mathcal{O}_{\mathbb{Z}}, \mathcal{O}_Y)$$

Denote by \mathcal{R} the vector bundle $\mathcal{R} = \mathbf{R}\pi_{Y*} \mathcal{O}_{\mathbb{Z}}$. Then the right hand side of the equation above is $\mathcal{R}^\vee = \mathbf{R}\mathcal{H}om(\mathcal{R}, \mathcal{O}_Y)$. Combining (18) and (19) we find that

$$(20) \quad \psi(\mathcal{O}_0 \otimes \rho) = [\mathcal{R}^\vee \otimes (\rho_{\det} \otimes \rho) \rightarrow \mathcal{R}^\vee \otimes (\rho_{\text{nat}} \otimes \rho) \rightarrow \mathcal{R}^\vee \otimes \rho]^G.$$

The vector bundle \mathcal{R} decomposes as a sum

$$(21) \quad \mathcal{R} = \bigoplus_{\rho} \mathcal{R}_{\rho} \otimes \rho$$

according to the decomposition of the regular representation. Moreover, for all $\rho \in \text{Irrep}(G)$, we have $(\mathcal{R}^\vee \otimes \rho)^G = (\mathcal{R}_{\rho}^\vee \otimes \rho^* \otimes \rho) = \mathcal{R}_{\rho}^\vee$. Then by exactness of the functor $(-)^G$:

$$(22) \quad \psi(\mathcal{O}_0 \otimes \rho) = [\mathcal{R}_{\rho_{\det} \otimes \rho}^\vee \rightarrow \mathcal{R}_{\rho_{\text{nat}} \otimes \rho}^\vee \rightarrow \mathcal{R}_{\rho}^\vee].$$

By [Ish02, Corollary 3.2], \mathcal{R}_{ρ} is the torsion free pull-back $\mathcal{R}_{\rho} = \widetilde{M}_{\rho}$. Therefore, $\psi(\mathcal{O}_0 \otimes \rho)$ is dual to the torsion-free pull-back of $\widetilde{A}_{\rho \otimes \rho_{\det}}$:

$$\mathbf{R}\mathcal{H}om(\widetilde{A}_{\rho \otimes \rho_{\det}}, \mathcal{O}_Y)[2] = \psi(\mathcal{O}_0 \otimes \rho),$$

since all summands of \mathcal{R} are locally free. The statement then follows from Proposition 2.2. \square

Remark 3.2. Let $\text{Coh}_G(\mathbb{C}^2)_0$ be the finite length category of G -equivariant sheaves supported at $0 \in \mathbb{C}^2$, and let ${}^0\text{Per}(Y/X)$ be the category of 0-perverse sheaves on Y (as defined in [Bri02]).

The complexes \widetilde{A}_{ρ} , for ρ special, are quasi-isomorphic to simple (-1) -perverse sheaves. Dualizing them (with the functor $\mathbf{R}\mathcal{H}om(-, K_Y)[2]$), we obtain simple 0-perverse sheaves ([VdB04b, Prop. 3.5.8]). Theorem 3.1 shows that ψ sends simple sheaves supported at 0 to simple 0-perverse

sheaves twisted by $-K_Y$. It follows that $\psi \otimes K_Y$ restricts to an exact functor of abelian categories $\text{Coh}_G(\mathbb{C}^2)_0 \rightarrow {}^0\text{Per}(Y/X)$.

3.1. Alternative proof of Prop. 2.3 (or [Ish02, Prop. 1.1]). As an application of Theorem 3.1, we give an alternative proof of [IU15, Prop. 1.1].

Let θ be the right adjoint to ϕ :

$$\begin{aligned} \theta(-) &:= [\mathbf{R}\pi_{Y*}(\pi_{\mathbb{C}^2}^*(-) \otimes^{\mathbf{L}} \mathcal{O}_{\mathbb{Z}}^{\vee} \otimes \pi_Y^*(K_Y)[2])]^G \\ &= [\mathbf{R}\pi_{Y*}(\pi_{\mathbb{C}^2}^*(-) \otimes^{\mathbf{L}} \mathcal{O}_{\mathbb{Z}}^{\vee})]^G \otimes K_Y[2]. \end{aligned}$$

Note that

$$(23) \quad \theta(- \otimes K_{\mathbb{C}^2}) = \psi(-) \otimes K_Y.$$

Proof of Prop. 2.3. Consider the semiorthogonal decomposition (10). Let $\mathcal{C} = \ker(\theta)$, i.e. the full triangulated subcategory of $D_G^b(\mathbb{C}^2)$ containing objects C such that $\theta(C) = 0$. By the adjunction between ϕ and θ one has $\mathcal{C} = \phi(D(Y))^\perp$, thus ${}^\perp\mathcal{C} \simeq \phi(D(Y))$.

Step 1: we show that the category \mathcal{C} is generated by the simple sheaves $\mathcal{O}_0 \otimes \rho$, with ρ not special.

Suppose E is a complex of G -equivariant sheaves on \mathbb{C}^2 which is not supported on the origin. Then by [BM02, Lemma 5.3] there exists a point in $X \setminus \{0\}$, equivalently a free orbit Gx in \mathbb{C}^2 , and an integer i such that

$$\text{Ext}_{\mathbb{C}^2}^i(\mathcal{O}_{Gx}, E)^G \neq 0.$$

The functor ϕ maps points of $Y \setminus F$ to free orbits, so we may write $\mathcal{O}_{Gx} \simeq \phi(\mathcal{O}_p)$ for some $p \in Y \setminus F$. Then,

$$\text{Ext}^i(\phi(\mathcal{O}_p), E)^G = \text{Ext}_{[\mathbb{C}^2/G]}^i(\phi(\mathcal{O}_p), E) = \text{Ext}_Y^i(\mathcal{O}_p, \theta(E)) \neq 0,$$

thus $E \notin \mathcal{C}$. Therefore, all objects of \mathcal{C} are supported at the origin.

Theorem (3.1), combined with (23), implies that $\mathcal{O}_0 \otimes \rho \in \mathcal{C}$ if and only if ρ is not special. Then sheaves in \mathcal{C} have a composition series with factors the non-special simples. Since θ is exact on $\text{Coh}_G(\mathbb{C}^2)_0$ by Remark 3.2 and (23), a complex $E \in D_G(\mathbb{C}^2)$ lies in \mathcal{C} precisely if its cohomologies do, which proves the claim.

Step 2: we show that the sheaf $\mathcal{O}_{\mathbb{C}^2} \otimes \rho$ is in the essential image of Φ for ρ special.

One readily checks that, for two irreducible representations σ and τ ,

$$\mathbf{R}\text{Hom}(\mathcal{O}_{\mathbb{C}^2} \otimes \sigma, \mathcal{O}_0 \otimes \tau) = 0$$

unless $\sigma = \tau$ (see Remark 2.4). Then by Step 1 we have that $\mathcal{O}_{\mathbb{C}^2} \otimes \rho \in {}^\perp\mathcal{C} = \text{Im } \Phi$ for ρ special.

Step 3: we show that the sheaves $\mathcal{O}_{\mathbb{C}^2} \otimes \rho$ for ρ special generate the essential image of Φ .

We have

$$\begin{aligned} \psi(\mathcal{O}_{\mathbb{C}^2} \otimes \rho) &= [\mathbf{R}\pi_{Y*} \mathbf{R}\text{Hom}(\mathcal{O}_{\mathbb{Z}}, \mathcal{O}_{Y \times \mathbb{A}^2} \otimes \rho \otimes \rho_{\det}[2])]^G \\ (24) \quad &= [\mathbf{R}\text{Hom}(\mathbf{R}\pi_{Y*} \mathcal{O}_{\mathbb{Z}}, \mathcal{O}_Y) \otimes \rho]^G \\ &= [\mathcal{R}^\vee \otimes \rho]^G = \mathcal{R}_\rho^\vee \end{aligned}$$

by (19) and (21). Then by Step 2, Equation (24), and the fact that

$$\phi \circ \psi|_{\phi(D(\text{Coh}(Y)))} = \text{id}_{\phi(D(\text{Coh}(Y)))}$$

we get that $\Phi(\mathcal{R}_\rho^\vee) = \mathcal{O}_{\mathbb{C}^2} \otimes \rho$ for all ρ special. Since the \mathcal{R}_ρ^\vee are generators for $D^b(Y)$ by [VdB04b, Theorem B], the special $\{\mathcal{O}_{\mathbb{C}^2} \otimes \rho\}$ generate $\Phi(D^b(\text{Coh}(Y)))$. \square

3.2. Cyclic singularities case. Suppose we are in the cyclic setting of Section 2.3.2. Then we can give a more precise statement of Theorem 3.1:

Proposition 3.3. *We have:*

$$(25) \quad \psi(\mathcal{O}_0 \otimes \rho_i) \simeq \begin{cases} \mathcal{O}_E(E)[1] & \text{if } a + 1 + i = 0 \\ \mathcal{O}_{E_t}(-\alpha_t + 1) & \text{if } a + 1 + i = i_t \in I(r, a) \setminus \{0, r\} \\ 0 & \text{otherwise.} \end{cases}$$

Proof. The proof goes as the proof of Theorem 3.1. Recall that in this case $\rho_{\det} = \rho_{a+1}$, and that Proposition 2.6 computes the torsion-free pullbacks \widetilde{A}_l . \square

Example 3.4. Suppose $G = \frac{1}{5}(1, 3)$. We have $\frac{5}{3} = [2, 3]$ and $I(5, 3) = \{5, 3, 1, 0\}$. The SOD (2) from [IU15] reads

$$D^b(\mathcal{X}) = \left\langle G, \mathcal{O}_0 \otimes \rho_4, \phi(D^b(\text{Coh}(Y))) \right\rangle$$

where $G = \mathbb{C}[x, y]/(x, y^2) \otimes \rho_4$ fits in an extension $\mathcal{O}_0 \otimes \rho_2 \xrightarrow{y} G \rightarrow \mathcal{O}_0 \otimes \rho_4$. By Proposition 3.3 there is a short exact sequence

$$\mathcal{O}_{E_2}(-2) \rightarrow \psi(G) \rightarrow \mathcal{O}_{E_1}(-1).$$

It is straightforward to check that $\psi(G)$ can also be written as an extension of \mathcal{O}_p by $\mathcal{O}_E(E)$, where $p \in E_2$.

Example 3.5. Let $G = \frac{1}{n}(1, 1)$, $n > 1$ integer. Let E the single exceptional $(-n)$ -curve in the minimal resolution of X (see Example 2.7). Then Theorem 3.3 reads

$$(26) \quad \psi(\mathcal{O}_0 \otimes \rho_i) \simeq \begin{cases} 0 & \text{if } i = 0, \dots, n-3 \\ \mathcal{O}_E(-n)[1] & \text{if } i = n-2 \\ \mathcal{O}_E(-n+1) & \text{if } i = n-1. \end{cases}$$

Example 3.6. Let $G = \frac{1}{2k+1}(1, k)$, with $k \geq 1$. Let E_1, \dots, E_k be the k exceptional curves in Y , where $E_1^2 = -3$ and $E_i^2 = -2$ for $i = 2, \dots, k$ (see Example 2.8). The fundamental cycle is $E = \sum_{i=1}^k E_i$. Theorem 3.3 reads

$$(27) \quad \psi(\mathcal{O}_0 \otimes \rho_i) \simeq \begin{cases} 0 & \text{if } i = 0, \dots, k-1 \\ \mathcal{O}_E(E)[1] & \text{if } i = k \\ \mathcal{O}_{E_{2k+1-i}}(-\alpha_{2k+1-i} + 1) & \text{if } i = k+1, \dots, 2n. \end{cases}$$

3.2.1. *Alternative (toric) approach to Proposition 3.3.* Here we present another possible approach, based on toric geometry, to the computation in Proposition 3.3.

First, we briefly recall how to describe the singularity $X = \frac{1}{r}(1, a)$ and its minimal resolution torically. The singularity X is the affine toric surface U_σ associated to the cone

$$(28) \quad \sigma = \langle re_1 - ae_2, e_2 \rangle \subset N_{\mathbb{R}} \simeq \mathbb{R}^2$$

Its minimal resolution can be obtained by performing n consecutive weighted blow-ups with weights $\frac{1}{i_{t-1}}(1, i_t)$, where $t = 1, \dots, n$. More explicitly, write $\rho_0 = e_2$ and $\rho_{n+1} = re_1 - ae_2$. Then Y is the smooth affine toric surface $Y = F_\Sigma$, where Σ is the refinement of σ obtained by adding the rays generated by the vectors

$$(29) \quad \rho_t = \frac{1}{i_{t-1}}(\rho_{n+1} + i_t \cdot \rho_{t-1}) \quad t = 1, \dots, n$$

Indeed, if $\sigma_{t-1} = \langle \rho_{n+1}, \rho_{t-1} \rangle$, the associated surface is the cyclic singularity $U_{\sigma_{t-1}} = \frac{1}{i_{t-1}}(1, i_t)$. The surface associated to the refinement of σ_{t-1} obtained by adding the ray ρ_t has two charts: the chart corresponding to $\tau_t = \langle \rho_{t-1}, \rho_t \rangle$ is $U_{\tau_t} = \mathbb{C}^2$, and the chart corresponding to $\sigma_t = \langle \rho_{n+1}, \rho_t \rangle$ is $U_{\sigma_t} = \frac{1}{i_t}(1, i_{t+1})$ if $t < n$, while is \mathbb{C}^2 if $t = n$. Note that the vectors ρ_t satisfy the relation $\rho_{t-1} + \rho_{t+1} = \alpha_t \cdot \rho_t$, which implies that the self-intersection of the exceptional curve E_t associated to the ray ρ_t is $-\alpha_t$ (see [CLS11, Theorem 10.4.4]).

The map $f: Y \rightarrow X$ can be described locally as follows. Let u_t, v_t be the coordinates on U_{τ_t} , and, for $t < n$, let x_t, y_t be the coordinates on \mathbb{C}^2 such that $U_{\sigma_t} = \text{Spec } \mathbb{C}[x_t, y_t]^G$, $G = \frac{1}{i_t}(1, i_{t+1})$. Then:

$$(30) \quad \begin{cases} x_{t-1}^{i_{t-1}} = u_t \\ y_{t-1}^{i_{t-1}} = u_t^{i_t} v_t^{i_{t-1}} \end{cases} \quad \text{and} \quad \begin{cases} x_{t-1}^{i_{t-1}} = x_t^{i_{t-1}} y_t \\ y_{t-1}^{i_{t-1}} = y_t^{i_t} \end{cases}$$

For $t = n$, the same expressions hold, x_n, y_n being the coordinates on $U_{\sigma_n} = \mathbb{C}^2$.

Remark 3.7. Any 2-dimensional strictly convex cone can be written in the form (28), called *normal form*. There is an ambiguity for normal forms of 2-dimensional cones: if $\sigma = \langle re_1 - ae_2, e_2 \rangle$ and $\sigma' = \langle r'e_1 - a'e_2, e_2 \rangle$ are lattice equivalent cones, then $r = r'$ and either $a = a'$ or $aa' = 1 \pmod r$. In terms of the associated singularity, this corresponds to writing X as either $\frac{1}{r}(1, a)$ or $\frac{1}{r}(a', 1)$. Note that, if $r/a = [\alpha_1, \dots, \alpha_n]$, then $r/a' = [\alpha_n, \dots, \alpha_1]$, that is, E_1, \dots, E_n appear in reverse order when resolving $\frac{1}{r}(a', 1)$ as indicated above. Define the j -series $J(r, a)$ to be the sequence $J(r, a) := \{j_0, j_1, \dots, j_n, j_{n+1}\}$, where $j_0 = 0, j_1 = 1$, and $j_t = j_{t-1}\alpha_{t-1} - j_{t-2}$. Then, $I(r, a') = \{j_{n+1}, j_n, \dots, j_1, j_0\}$.

The Picard group of the minimal resolution Y is generated by divisors D_1, \dots, D_n determined by the condition $D_i \cdot E_j = \delta_{ij}$. For $t = 1, \dots, n$, formula (13) implies $\widetilde{M}_{\rho_{i_t}} = \mathcal{O}_Y(D_t)$. Now let $l \in \{0, \dots, r-1\}$. By [Wun88, §A.1, Theorem a)] the summand $\mathcal{R}_{\rho_l} = \widetilde{M}_{\rho_l}$ appearing in (21) is the line bundle $\mathcal{O}_Y(\sum_i d_i^{(l)} D_i)$, where the $d_i^{(l)} \geq 0$ are uniquely determined by the I-series $I(r, a)$ by requiring

$$l = d_1 i_1 + \dots + d_n i_n \quad \text{and} \quad 0 \leq \sum_{t>t_0} d_t i_t < i_{t_0} \text{ for any } t_0.$$

We can then interpret the sequences (22) by studying the D_i 's and the divisors associated to the invariant coordinates x^r and y^r on X , without recurring to Proposition 2.2.

Consider the divisors on X defined by x^r and y^r . By (30), their strict transforms under $f: Y \rightarrow X$ are nonreduced curves $A' = rA, B' = rB$; by construction, the curve A (resp. B) meets the curve E_n (resp. E_1) at exactly one point (see top of Figure 3). In what follows, we write the pull-backs $f^*(x^r), f^*(y^r)$ and the D_i 's as linear combinations of A, B , and E_1, \dots, E_n .

For $s \in \{0, 1, \dots, n, n+1\}$ define a sequence $H(s) := \{h_0^{(s)}, \dots, h_{n+1}^{(s)}\}$ by letting $h_s^{(s)} = 0, h_{s-1}^{(s)} = h_{s+1}^{(s)} = 1$ and

$$\begin{aligned} h_t^{(s)} &= h_{t+1}^{(s)} \alpha_{t+1} - h_{t+2}^{(s)} \quad \text{for } 0 \leq t < s-1, \\ h_t^{(s)} &= h_{t-1}^{(s)} \alpha_{t-1} - h_{t-2}^{(s)} \quad \text{for } s+1 < t \leq n+1. \end{aligned}$$

Observe that $H(0) = J(r, a)$ and $H(n+1) = I(r, a)$. More generally, the truncation $H(s)_{\geq s}$ is the j -series $J(i_s, i_{s+1})$ and the truncation $H(s)_{\leq s}$ is the i -series $I(j_s, j'_{s-1})$, with $j_{s-1} j'_{s-1} = 1 \pmod j_s$.

By (30), the terms of $H(0)$ and $H(n+1)$ are the coefficients of the pull-back of $(x^r = 0)$ and $(y^r = 0)$ to Y :

$$(f^*(x^r) = 0) = \sum_{t=1}^n h_t^{(0)} E_t + h_{n+1}^{(0)} A \quad (f^*(y^r) = 0) = h_0^{(n+1)} B + \sum_{t=1}^n h_t^{(n+1)} E_t.$$

Moreover, one checks that there are numerical equivalences⁸

$$(31) \quad D_s \equiv \sum_{t=s+1}^n h_t^{(s)} E_t + h_{n+1}^{(s)} A \equiv h_0^{(s)} B + \sum_{t=1}^{s-1} h_t^{(s)} E_t \quad s = 1, \dots, n.$$

An alternative way to obtain the expressions (31) is to look at the fibers of maps $Y \rightarrow \mathbb{P}^1$ constructed as follows. For each s , contract all the exceptional curves on Y except for E_s , to obtain a morphism $\tau_s: Y \rightarrow T_s$, where T_s is the total space of an orbibundle over the projective line $\mathbb{P}(j_{s-1}, i_{s+1}) \simeq \mathbb{P}^1$ (see Figure 3). Project further $\sigma_s: T_s \rightarrow \mathbb{P}^1$. Then, by (30), the two expressions for D_s written in (31) are the fibers of $\sigma_s \circ \tau_s: Y \rightarrow \mathbb{P}^1$ over the points $(0:1)$ and $(1:0)$ in \mathbb{P}^1 .

⁸Since we are in a toric setting, linear and numerical equivalence coincide. For the same reason, it is not surprising that working up to equivalence does not suffice to compute the cohomologies of the sequences (22), which is why the expressions (31) are relevant to our computation.




Figure 3. A picture of the surface Y and the maps $f: Y \rightarrow X$, $p_3 \circ \tau_3: Y \rightarrow \mathbb{P}^1$. The toric surface T_3 has two singular points of type $\frac{1}{i_3}(1, i_4)$ and $\frac{1}{j_3}(j_2, 1)$.

This setup can be used to interpret the maps in the complexes (22). Indeed, invariant functions pull-back to maps:

$$\mathcal{O}_Y \xrightarrow{f^*(x^r)} \mathcal{O}_Y \quad \mathcal{O}_Y \xrightarrow{f^*(y^r)} \mathcal{O}_Y$$

which must factor through rational maps

$$(32) \quad x_{(l)}: \mathcal{O}_Y(\sum_i d_i^{(l)} D_i) \simeq \mathcal{R}_l \rightarrow \mathcal{R}_{l+1} \simeq \mathcal{O}_Y(\sum_i d_i^{(l+1)} D_i)$$

(and similarly for y) satisfying $(x_{(l)}) \leq (f^*(x^r))$. This last condition uniquely determines a divisor in the linear series $|\sum_i d_i^{(l)} D_i|$, which allows one to compute the cohomologies of (22) without ambiguity: the maps $x_{(l)}$ and $y_{(l)}$ are precisely those appearing in (22) (after taking duals). We illustrate this in an example.

Example 3.8. Suppose $G = \frac{1}{5}(1, 2)$. The continued fraction expansion is $\frac{7}{5} = [3, 2]$ and the i -series $I(7, 3) = \{5, 2, 1, 0\}$. Equation (27) reads

$$(33) \quad \psi(\mathcal{O}_0 \otimes \rho_i) \simeq \begin{cases} 0 & \text{if } i = 0, 1, \\ \mathcal{O}_F(F)[1] & \text{if } i = 2 \\ \mathcal{O}_{E_{5-i}(-\alpha_{5-i}+1)} & \text{if } i = 3, 4. \end{cases}$$

We verify it using the approach of Section 3.2.1. Applying [Wun88, §A.1] we have

$$\mathcal{R}_i = \begin{cases} \mathcal{O}_Y & i = 0 \\ \mathcal{O}_Y(D_{3-i}) & i = 1, 2 \\ \mathcal{O}_Y(D_1 + D_{5-i}) & i = 3, 4 \end{cases}$$

From Section 3.2.1 we get the following equalities:

$$\begin{aligned} (f^*x^5 = 0) &= E_1 + 3E_2 + 5A, & D_1 &\equiv B \equiv E_2 + 2A, \\ (f^*y^5 = 0) &= 5B + 2E_1 + E_2, & D_2 &\equiv 3B + E_1 \equiv A. \end{aligned}$$

The maps $x_{(l)}$ are uniquely determined by the above. Indeed, denote by a, b, e_1, e_2 the sections defining the divisors A, B, E_1, E_2 . Then we have:

$$\begin{array}{ll} x_{(0)}: \mathcal{R}_0 \xrightarrow{a} \mathcal{R}_1 & y_{(0)}: \mathcal{R}_0 \xrightarrow{b} \mathcal{R}_2 \\ x_{(1)}: \mathcal{R}_1 \xrightarrow{e_2 a} \mathcal{R}_2 & y_{(2)}: \mathcal{R}_2 \xrightarrow{b} \mathcal{R}_4 \\ x_{(2)}: \mathcal{R}_2 \xrightarrow{a} \mathcal{R}_3 & y_{(4)}: \mathcal{R}_4 \xrightarrow{be_1} \mathcal{R}_1 \\ x_{(3)}: \mathcal{R}_3 \xrightarrow{e_2 a} \mathcal{R}_4 & y_{(1)}: \mathcal{R}_1 \xrightarrow{b} \mathcal{R}_3 \\ x_{(4)}: \mathcal{R}_4 \xrightarrow{e_1 e_2 a} \mathcal{R}_0 & y_{(3)}: \mathcal{R}_3 \xrightarrow{be_1 e_2} \mathcal{R}_0, \end{array}$$

so that $f^* x^5 = x_{(4)} x_{(3)} x_{(2)} x_{(1)} x_{(0)}$ (and similarly for y). The dual maps $x_{(l)}^\vee$ (resp. $y_{(l)}^\vee$) to $x_{(l)}$ (resp. $y_{(l)}$) correspond to multiplication by the same sections. Then, consider for example the sequence:

$$\psi(\mathcal{O}_0 \otimes \rho_2) = \left[\mathcal{R}_0^\vee \xrightarrow{(-y_{(3)}^\vee, x_{(4)}^\vee)^T} \mathcal{R}_3^\vee \oplus \mathcal{R}_4^\vee \xrightarrow{(x_{(2)}^\vee, y_{(2)}^\vee)} \mathcal{R}_2^\vee \right].$$

The kernel of the map $(x_{(2)}^\vee, y_{(2)}^\vee) = (a, b)$ is

$$\mathcal{O}_Y(-2D_1 - D_2) \xrightarrow{(-b, a)^T} \mathcal{R}_3^\vee \oplus \mathcal{R}_4^\vee.$$

We have $(-y_{(3)}^\vee, x_{(4)}^\vee)^T = e_1 e_2 (-b, a)^T$, so

$$\psi(\mathcal{O}_0 \otimes \rho_2) = \text{coker} \left(\mathcal{R}_0 \xrightarrow{e_1 e_2} \mathcal{O}(-2D_1 - D_2) \right) [1] \simeq \mathcal{O}_F(F)[1].$$

Similarly, consider:

$$\psi(\mathcal{O}_0 \otimes \rho_3) = \left[\mathcal{R}_1^\vee \xrightarrow{(-y_{(4)}^\vee, x_{(0)}^\vee)^T} \mathcal{R}_4^\vee \oplus \mathcal{R}_0^\vee \xrightarrow{(x_{(3)}^\vee, y_{(3)}^\vee)} \mathcal{R}_3^\vee \right].$$

with $(-y_{(4)}^\vee, x_{(0)}^\vee)^T = (-be_1, a)^T$ and $(x_{(3)}^\vee, y_{(3)}^\vee) = (e_2 a, be_1 e_2)$. This shows that the complex is exact in the middle. However, the cokernel of $(-be_1, a)^T$ is (a, be_1) , and hence

$$\psi(\mathcal{O}_0 \otimes \rho_3) = \text{coker} \left(\mathcal{O}(-2D_1 + D_2) \xrightarrow{e_2} \mathcal{R}_3^\vee \right) \simeq \mathcal{O}_{E_2}(-1).$$

One proves the remaining cases in the same way.

4. THE MINIMAL RESOLUTION OF X

For $k > 0$ integer, let $X^{(k)}$ be a sufficiently general log del Pezzo surface of the family (1). The surface $X^{(k)}$ is the vanishing locus of a generic section of the sheaf $\mathcal{O}_{\mathbb{P}}(8k + 4)$. Let $f^{(k)}: \tilde{X}^{(k)} \rightarrow X^{(k)}$ be the minimal resolution. In this Section we describe in details the geometry of the surface $\tilde{X}^{(k)}$. We will omit the superscript $^{(k)}$ from our notation whenever the context allows it.

4.1. Basic properties. Fix $k > 0$ integer. We denote by x, y_1, y_2, z the weighted homogeneous coordinates of $\mathbb{P} = \mathbb{P}(2, 2k+1, 2k+1, 4k+1)$. The singular locus \mathbb{P}_{sing} of \mathbb{P} has three components:

1. the point $(1 : 0 : 0 : 0)$, which is a singularity of type $1/2(1, 1, 1)$;
2. the line $(x = z = 0)$, whose points all have a neighbourhood analytically isomorphic to $\mathbb{A}^1 \times 1/(2k+1)(1, k)$;
3. the point $(0 : 0 : 0 : 1)$, which is a singularity of type $1/(4k+1)(4, 1, 1)$.

It is easy to check that a section of the sheaf $\mathcal{O}_{\mathbb{P}}(8k+4)$ is of the form:

$$(34) \quad h^{(k)}(x, y_1, y_2, z) = ax^{4k+2} + h_4(y_1, y_2) + bxz^2 + x^{2k+1}h_2(y_1, y_2) + x^{k+1}h_1(y_1, y_2)z$$

where $a, b \in \mathbb{C}$ and $h_i(y_1, y_2)$ is a homogeneous polynomial of degree i in the variables y_1, y_2 .⁹ It follows that, for a generic choice of h , $(1 : 0 : 0 : 0) \notin X$, and $(0 : 0 : 0 : 1) \in X$ is a singularity of type $1/(4k+1)(1, 1)$. Moreover, since $h(0, y_1, y_2, 0) = h_4(y_1, y_2)$, X intersects the line $(x = z = 0)$ in four points, each of which is a singularity of type $1/(2k+1)(1, k)$. We write $p = (0 : 0 : 0 : 1)$ and denote by p_1, p_2, p_3, p_4 the four points in $X \cap (x = z = 0)$.

By Equation (34) the curve $(x = 0) \cap X$ is the union of four lines $c_i \in |\mathcal{O}_{\mathbb{P}}(2)| \oplus |\mathcal{O}_{\mathbb{P}}(2k+1)|$, $i = 1, 2, 3, 4$; for each i , the line c_i passes through the points p_i and p (see the bottom left of Figure 6).

Remark 4.1. By the general theory of surface cyclic quotient singularities, the surface \tilde{X} is obtained from X by a sequence of $1 + 4k$ weighted blow-ups: a single weighted blow-up at p , and a sequence of k weighted blow-ups centered at each p_i (see Examples 2.7 and 2.8).

We denote by F the $-(4k+1)$ -exceptional curve over p , and for $i = 1, 2, 3, 4$ we denote by $C_{i,0}$ the (-3) -exceptional curve, and by $C_{i,j}$, $j = 1, \dots, k-1$, the (-2) -exceptional curves over p_i . Moreover, for $i = 1, 2, 3, 4$ we denote by $C_{i,k}$ the strict transform in \tilde{X} of the line $c_i \subset X$ (see the top of Figure 6).

Remark 4.2. The minimal resolution $f: \tilde{X} \rightarrow X$ is noncrepant. More precisely, the singularities of X are strictly log-terminal:

$$(35) \quad K_{\tilde{X}} = \tau^*(K_X) + d \cdot F + \sum_{i=1}^4 \sum_{j=1}^k d_j \cdot C_{ij}$$

where

$$(36) \quad d = -\frac{4k-1}{4k+1} \quad \text{and} \quad d_j = -\frac{k-(j-1)}{2k+1}$$

4.2. The minimal resolution. Fix a generic polynomial $h = h^{(k)}$, equivalently, $a, b \in \mathbb{C} \setminus \{0\}$, and polynomials h_4, h_2, h_1 as in Equation (34). Let X be the corresponding log del Pezzo surface and \tilde{X} its minimal resolution. Let $X' \subset \mathbb{P}(2, 1, 1, 1)$ be the smooth del Pezzo surface of degree two defined by the polynomial $h^{(0)}$. The plane $(x = 0) \subset \mathbb{P}(2, 1, 1, 1)$ intersects X' in four lines $c_{i,0}$, $i = 1, 2, 3, 4$, which meet at the point $q := (0 : 0 : 0 : 1)$. The next Theorem shows that \tilde{X} is obtained from X' by a sequence of $4k+1$ smooth blow-ups.

Theorem 4.3. *There is a birational morphism $\tau: \tilde{X} \rightarrow X'$, which contracts the $4k+1$ curves $C_{i,k}, \dots, C_{i,1}$, $i = 1, 2, 3, 4$, and F to the point q , as pictured in Figure 4.*

Proof. Let $\Delta \subset N_{\mathbb{R}}$ be the fan of \mathbb{P} and let $\rho_1, \rho_2, \rho_3, \rho_4$ be the primitive generators of the rays of Δ . One may choose for instance:

$$\rho_1 = (k, k, 4k+1)^T \quad \rho_2 = (1, 0, 0)^T \quad \rho_3 = (0, 1, 0)^T \quad \rho_4 = (-1, -1, -2)^T$$

Consider the star subdivision $\Sigma = \Delta^*(v)$, where

$$(37) \quad v = \frac{1}{4k+1} (4\rho_1 + \rho_2 + \rho_3)$$

The toric variety $G = G_{\Sigma}$ is the weighted blow-up of \mathbb{P} at p with weights $1/(4k+1)(4, 1, 1)$. Note that the ray ρ_1 belongs to the cone generated by ρ_4 and v (see Figure 5); more precisely,

$$\rho_1 = \frac{1}{2}(\rho_4 + (2k+1)v)$$

⁹Note that by [IF00, Corollary 8.5] the vanishing locus of h is quasismooth if and only if $a \neq 0$, $b \neq 0$, and for $i = 1, 2$ the polynomial $h_i(y_1, y_2)$ has a monomial of the form $y_i^m y_j$, $m > 0$, $j \in \{1, 2\}$.




Figure 4. A picture of the morphism $\tau: \tilde{X} \rightarrow X'$. The surface \tilde{X} is obtained from the surface X' by a sequence of $4k + 1$ smooth blow-ups.

Let $Y \subset G$ be the strict transform of X . The exceptional divisor of the morphism $G \rightarrow \mathbb{P}$ is the torus-invariant divisor $D_v \subset G$. By construction, the exceptional curve $D_v \cap Y \subset Y$ over the point $p \in X$ is a $-(4k + 1)$ -curve. The line $D_{\rho_1} \cap D_{\rho_4} \subset G$ is a line of $1/(2k + 1)(1, k)$ points, and the surface Y intersects this line in four points (see the left of Figure 6).

Now consider the sequence of k star-subdivisions of the fan Σ defined by setting $\Sigma_0 = \Sigma^*(v_0)$, where

$$(38) \quad v_0 = \frac{1}{2k + 1}(\rho_1 + k\rho_4)$$

and, for $i = 1, \dots, k - 1$, $\Sigma_i = \Sigma_{i-1}^*(v_i)$, where

$$(39) \quad v_i = \frac{1}{k - (i - 1)}(\rho_1 + (k - i)v_{i-1})$$

and let $G_i = G_{\Sigma_i}$ (see Figure 5). The variety $G_0 = G_{\Sigma_0}$ is the weighted blow-up of G along the line $D_{\rho_1} \cap D_{\rho_4}$ with weights $1/(2k + 1)(1, k)$. For $i = 1, \dots, k - 1$, the line $D_{\rho_1} \cap D_{v_{i-1}} \subset G_{i-1}$ is a line of $\frac{1}{k - (i - 1)}(1, k - i)$ points, and the variety G_i is the weighted blow-up of G_{i-1} along this line.

Write $\tilde{G} = G_{k-1}$ and let \tilde{X} be the strict transform of X in \tilde{G} . By construction, the surface \tilde{X} is the minimal resolution of X . The curve $D_v \cap \tilde{X} \subset \tilde{X}$ is the $-(4k + 1)$ -exceptional curve F over the point $p \in X$, and the intersection $\tilde{X} \cap (D_{v_0} \cup D_{v_1} \cup \dots \cup D_{v_{k-1}})$ is given by the union of the four chains

$$C_i = \bigcup_{j=0}^{k-1} C_{i,j} \quad i = 1, 2, 3, 4$$

where $C_{i,j}$ is the j th exceptional curve over the point $p_i \in X$ (see Remark 4.1 and the top of Figure 6). The intersection $\tilde{X} \cap D_{\rho_1} \subset \tilde{X}$ is the union of the proper transforms $C_{i,k}$ of the lines $c_i \subset X$, $i = 1, 2, 3, 4$. Each $C_{i,k}$ has self-intersection -1 ; indeed $K_X \cdot c_i = -\frac{1}{(2k+1)(4k+1)}$ since $c_i \in |\mathcal{O}_{\mathbb{P}(2)}| \oplus |\mathcal{O}_{\mathbb{P}(2k+1)}|$, thus by Remark 4.2

$$K_{\tilde{X}} \cdot C_{i,k} = K_X \cdot c_i - \frac{4k - 1}{4k + 1} - \frac{1}{2k + 1} = -1$$

and by adjunction $C_{i,k}^2 = -1$.

Now, observe that the vectors $v_1, \dots, v_{k-1}, \rho_1$ all belong to the cone $\langle v_0, v \rangle$ (see Figure 5). In particular, it follows from (37) and (38) that $\rho_1 = v_0 + kv$; combining this and (39) one finds that:

$$(40) \quad v_i = v_{i-1} + v \quad i = 1, \dots, k - 1 \quad \text{and} \quad \rho_1 = v_{k-1} + v$$




Figure 5. A picture of the fan $\tilde{\Sigma}$ for $k = 2$. The segments colored in blue all belong to the same plane. The three star-subdivisions of the fan Δ are represented by the segments in red. (For sake of clarity, the picture is out of scale.)

This implies that the fan $\tilde{\Sigma}$ is obtained from the fan Σ'_0 with rays $\rho_2, \rho_3, \rho_4, v_0, v$ and maximal cones

$$\langle \rho_2, \rho_3, \rho_4 \rangle \quad \langle \rho_2, v_0, v \rangle \quad \langle \rho_2, \rho_4, v_0 \rangle \quad \langle \rho_3, v_0, v \rangle \quad \langle \rho_3, \rho_4, v_0 \rangle \quad \langle \rho_2, \rho_3, v \rangle$$

by the sequence of k star subdivisions:

$$\Sigma'_i = (\Sigma'_{i-1})^*(v_{i-1} + v) \quad i = 1, \dots, k$$

where $\tilde{\Sigma} = \Sigma'_k$. For $i = 1, \dots, k$ the toric variety $G'_i = G_{\Sigma'_i}$ is the blow up of the toric variety G'_{i-1} along $D_{v_{i-1}} \cap D_v$.

Let Y' be the proper transform of \tilde{X} in G'_0 . The morphism $\tilde{G} \rightarrow G'_0$ contracts the divisors $D_{\rho_1}, D_{v_{k-1}}, \dots, D_{v_1}$ in \tilde{G} , thus contracts the curves $C_{i,k}, C_{i,k-1}, \dots, C_{i,1}$, $i = 1, 2, 3, 4$, in \tilde{X} . For $i = 1, 2, 3, 4$ the proper transform of $C_{i,0}$ in Y' is a (-2) -curve, and the proper transform of F in Y' is a (-1) -curve (see the right of Figure 6).

Now, note that the vector v can be written in terms of ρ_2, ρ_3, v_0 as:

$$(41) \quad v = \rho_2 + \rho_3 + 4v_0$$

and that $\rho_2, \rho_3, \rho_4, v_0$ span the lattice N and satisfy:

$$(42) \quad 2v_0 + \rho_2 + \rho_3 + \rho_4 = 0$$

This implies that G'_0 is the weighted blow-up of $\mathbb{P}(2, 1, 1, 1)$ at the smooth point $q = (0 : 0 : 0 : 1)$ with weights $(4, 1, 1)$.

Let X' be the proper transform of Y' in $\mathbb{P}(2, 1, 1, 1)$. The morphism $G'_0 \rightarrow \mathbb{P}(2, 1, 1, 1)$ contracts the divisor $D_v \subset G'_0$ to the point q , thus contracts the proper transform of F in Y' to q . The proper transforms $c_{i,0} \subset X'$ of the curves $C_{i,0} \subset \tilde{X}$, $i = 1, 2, 3, 4$, are (-1) -curves meeting at the point q (see the bottom right of Figure 6).




Figure 6. A picture of the surfaces X, Y, \tilde{X}, Y' , and X' . The surface Y is the weighted blow-up of X at p with weights $1/(4k+1)(1, 1)$; the surface Y' is the blow-up of X' at the smooth point q . The curves $F, C_{i,0}, \dots, C_{i,k-1}, i = 1, 2, 3, 4$, are the exceptional curves of the minimal resolution $\tilde{X} \rightarrow X$, while the curves $F, C_{i,1}, \dots, C_{i,k}, i = 1, 2, 3, 4$, are the exceptional curves of the morphism $\tilde{X} \rightarrow X'$. We use the same color for a curve and its proper transforms under the maps considered.

It follows from (34) and (42) that the surface X' is the smooth del Pezzo surface of degree two cut out of $\mathbb{P}(2, 1, 1, 1)$ by the section of $\mathcal{O}(4)$ given by the polynomial $h^{(0)}$. The union of the four curves $c_{i,0}$ is given by $(x = f_4(y_1, y_2) = 0) \subset \mathbb{P}(2, 1, 1, 1)$. \square

An immediate consequence of Theorem 4.3 is:

Corollary 4.4. *For $k > 0$ integer, let $X = X^{(k)}$ be a generic log del Pezzo surface (1) and let $\tilde{X}^{(k)}$ be its minimal resolution. Then $\tilde{X}^{(k)}$ is the blow-up of \mathbb{P}^2 in $8 + 4k$ points. Moreover, for all $k > 0$, $X^{(k)}$ is a rational surface of Picard rank $\rho(X^{(k)}) = 8$.*

Theorem 4.3 highlights a relation between the surfaces of the series (1) as k varies. Fix a generic choice of $a, b \in \mathbb{C} \setminus \{0\}$, and of polynomials h_4, h_2, h_1 . This defines a sequence of polynomials $\{h^{(k)}\}_{k>0}$, and with it sequences $\{X^{(k)}\}$ and $\{\tilde{X}^{(k)}\}$ of log del Pezzo surfaces and their minimal resolutions. By Theorem 4.3, every surface $\tilde{X}^{(k)}$ admits a birational morphism $\tau^{(k)}: \tilde{X}^{(k)} \rightarrow X'$.

Corollary 4.5. *For $0 < k_1 < k_2$, the map $\tau^{(k_2)}$ factors through the map $\tau^{(k_1)}$, as illustrated in Figure 7.*

Any del Pezzo surface of degree two as in Theorem 4.3 has a generalized Eckardt point, that is, a point where four (-1) -curves of X' meet.




Figure 7. The morphisms $f^{(k)}: \tilde{X}^{(k)} \rightarrow X^{(k)}$ and $\tau^{(k)}: \tilde{X}^{(k)} \rightarrow X'$. For all $k > 0$, the surface $\tilde{X}^{(k+1)}$ is the blow-up at four points of the surface $\tilde{X}^{(k)}$. We use dashed arrows to denote rational maps and solid arrows to denote morphisms.

By [Kuw05, Theorem 5.1] a del Pezzo surface S of degree two admits a generalized Eckardt point if and only if it satisfies any of the following equivalent properties:

(i) by a suitable change of coordinates, S can be defined by an equation of the form

$$(43) \quad (x_4^2 + x_3^4 + g_2(x_1, x_2)x_3^2 + g_4(x_1, x_2) = 0) \subset \mathbb{P}(1, 1, 1, 2)$$

where x_1, x_2, x_3, x_4 are weighted homogeneous coordinates on $\mathbb{P}(1, 1, 1, 2)$ and $g_i(x_1, x_2)$ is a polynomial of degree i in the two variables x_1, x_2 ;

(ii) the quartic $Q \subset \mathbb{P}^2$ along which the natural double cover $\sigma: S \rightarrow \mathbb{P}^2$ is branched admits an involution;

(iii) the automorphism group $\text{Aut}(S)$ contains $\mathbb{Z}/2 \times \mathbb{Z}/2$ as a subgroup.

The moduli space of smooth del Pezzo surfaces of degree two whose automorphism group contains $\mathbb{Z}/2 \times \mathbb{Z}/2$ as a subgroup is a stratum of dimension 4 of the moduli space of smooth del Pezzo surfaces of degree 2. Surfaces in this stratum are called *of type XII* in [Dol12, Table 8.9].

It is easy to see that by a change of coordinates any polynomial of the form $h^{(0)}$ can be rewritten in the form (43). Then, it follows that:

Corollary 4.6. *For all $k > 0$ integer, the moduli stack of hypersurfaces of degree $8k + 4$ in the weighted projective 3-space $\mathbb{P}(2, 2k + 1, 2k + 1, 4k + 1)$ is birational to the moduli space of smooth del Pezzo surfaces of type XII.*

4.3. Classification of generalized Eckardt points. Suppose that S is a del Pezzo surface of degree two with a generalized Eckardt point q . Denote by m_l , $l = 1, 2, 3, 4$, the four (-1) -curves meeting at the point q . In this Section we classify all the possible configurations $n_l = \pi(m_l)$, where $\pi: S \rightarrow \mathbb{P}^2$ is the blow-down of an exceptional set. Keeping this notation:

Lemma 4.7. *Let $\pi: S \rightarrow \mathbb{P}^2$ be a blowup of \mathbb{P}^2 at seven points x_i in general position (equivalently, a choice of an exceptional set for S). Up to reindexing, we have the following possibilities for the curves n_l :*

- n_1 and n_2 are lines, and n_3 and n_4 are conics;
- n_1 is a point, and the remaining are conics, or n_1 is a cubic and the remaining are lines;
- n_1 is a point, and the remaining are a line, a conic, and a cubic.

Proof. Observe first of all that n_k is a point for at most one k , since the m_k are not disjoint.

If $n_1 = x_1$ is a point, then at most one of the other n_k is a line: two lines in \mathbb{P}^2 containing x_1 are disjoint in the blowup, unless they coincide.

Suppose $n_1 = x_1$ and n_2 is a line (containing x_2). At most one between n_3, n_4 is a conic: otherwise, n_3 and n_4 pass through x_1 with the same tangent direction of n_2 , and therefore they intersect in x_1 (with multiplicity 2). Each conic must contain four points among (x_3, \dots, x_7) , but if they share three they coincide, a contradiction.

If n_1, n_2, n_3 are lines, n_4 cannot be a line: each line contains $\pi(q)$ and two of the x_i , and no point x_i can be common to distinct lines. Similarly, n_4 is not a conic: if it was, it would

have to share $\pi(q)$ and two of the x_i with one among n_1, n_2, n_3 , say n_1 . But then $n_1 \subset n_4$, a contradiction.

At this point, we may argue dually on the curves $m'_k = -K_S - m_k$. In fact, they also intersect at a generalized Eckardt point q' , and since the image of m_k determines that of m'_k and conversely, we can exclude all the configurations dual to those excluded above. For example, no more than one n_k is a cubic, otherwise more than one n'_k would be a point. The only possibilities left are those in the statement: Example 4.8 shows that these do occur. \square

Example 4.8 (Construction of generalized Eckardt points). We present three constructions of seven points $x_1, \dots, x_7 \in \mathbb{P}^2$ in general position. The blowup $\text{Bl}_{\{x_i\}} \mathbb{P}^2$ is a del Pezzo surface of degree two with a generalized Eckardt point (which is the preimage of q in the first example, and lies in the preimage of x_1 in the other cases). In particular, these constructions are examples of the configurations listed in Lemma 4.7.

- Let t_1, t_2 be two lines in \mathbb{P}^2 intersecting at a point q . Pick general points $x_1, x_2 \in t_1, x_3, x_4 \in t_2$. Let s_1 be a generic conic through q, x_1, x_3 and s_2 a generic conic through q, x_2, x_4 . Then $s_1 \cap s_2 = \{q, x_5, x_6, x_7\}$ (Figure 8).
- Let $x_1 \in \mathbb{P}^2$, and fix a conic s_1 through x_1 . Define two general conics s_2 and s_3 which pass through x_1 with the same tangent direction of s_1 . We have $s_1 \cap s_2 = \{x_2, x_3\}$, $s_1 \cap s_3 = \{x_4, x_5\}$, and $s_2 \cap s_3 = \{x_6, x_7\}$. For generic choices, the points $x_i, i = 1, \dots, 7$ are in general position, and the blowup $\text{Bl}_{\{x_i\}} \mathbb{P}^2$ is a del Pezzo surface of degree two with a generalized Eckardt point at the intersection of the strict transforms of the s_k with the exceptional divisor of x_1 (Figure 9).
- Fix a point $x_1 \in \mathbb{P}^2$, a line t through it and a conic s tangent to t at x_1 . Now pick generic points $x_2 \in t$, and $x_3, \dots, x_6 \in s$. Then, pick a nodal cubic in the 2-dimensional linear system of cubics containing x_2, \dots, x_6 and passing through x_1 with tangent direction t (Figure 10).




Figure 8




Figure 9




Figure 10

In fact, the configurations of Example 4.8 are the only possibility for generalized Eckardt points. Keeping the same notation as Lemma 4.7, we have:

Proposition 4.9. *The map $\pi : S \rightarrow \mathbb{P}^2$ is one of the blowups described in Example 4.8.*

Proof. By Lemma 4.7, there are only three possibilities for the n_l (after possibly considering the dual curves n'_l). We only show one case, the others being similar.

Suppose n_1, n_2 are lines and n_3, n_4 are conics. Since n_1 and n_2 intersect at $\pi(q)$, they cannot share the x_i 's they contain, so we may assume $x_1, x_2 \in n_1$ and $x_3, x_4 \in n_2$. The conic n_3 must contain 5 of the x_i and pass through q , moreover it intersects n_1 and n_2 at two points (n_3 is smooth, otherwise the x_i are not in general position). The only possibility is that n_3 contains, say, x_1 and x_3 , whereas $x_2, x_4 \in n_4$. Then, n_3 and n_4 must meet at q and at x_5, x_6, x_7 . This is the situation of Example 4.8 (a). \square

Later it will be useful to know that it is always possible to choose π as in Example 4.8 (a). This is Corollary 4.13 below, which follows from more general statements about *cliques on S*.

In graph theory, a clique is a complete subgraph of an undirected graph. Given a degree two del Pezzo surface S , by a clique on S we mean a set of four (-1) -curves on S whose dual graph is

a clique in the dual graph of all (-1) -curves. The (-1) -curves meeting at a generalized Eckardt point form a clique, but the converse does not hold.

Remark 4.10. Lemma 4.7 holds if $\{m_1, \dots, m_4\}$ is only a clique on S , with a nearly identical proof.

Lemma 4.11. *Let S be a del Pezzo surface of degree two and let $\{m_1, \dots, m_4\}$ be a clique on S . Then there exists an exceptional set \mathcal{L} on S such that $\mathcal{L} \cap \{m_1, \dots, m_4\} = \emptyset$.*

Proof. The set of (-1) -curves on S is in bijection with the set of exceptional vectors in $I^{1,7}$ (see Section 2.2.1). Identifying the two sets, one checks via computer algebra that on S there are 576 exceptional sets and 630 cliques, and that every clique admits a disjoint exceptional set. \square

Lemma 4.12. *Let S be a del Pezzo surface of degree two, let $\{m_1, \dots, m_4\}$ be a clique on S and let $\{m'_1, \dots, m'_4\}$ be the dual clique. Then there exists an exceptional set \mathcal{L} on S such that $\mathcal{L} \cap \{m_1, \dots, m_4, m'_1, \dots, m'_4\} = \emptyset$.*

Proof. Via computer algebra one checks that, for each of the 630 cliques on S , there is an exceptional set which is disjoint from both the clique and its dual. \square

Corollary 4.13. *Suppose that S is a del Pezzo surface of degree two with a generalized Eckardt point q . Then S can always be realized as the blow-up of \mathbb{P}^2 described in Example 4.8 (a).*

Proof. An immediate consequence of Proposition 4.9 and Lemma 4.12. \square

5. THE DERIVED CATEGORY OF \mathcal{X}

We construct full exceptional collections for $D(\tilde{X})$ in Section 5.1; this directly implies the existence of full exceptional collections for $D(\mathcal{X})$. In Section 5.2, after some additional choices, we compute the morphism spaces between objects in the above collections. The only non-vanishing spaces are Hom and Ext^1 . In Section 5.3, we observe that this is sufficient to produce tilting complexes for $D(\tilde{X})$ and $D(\mathcal{X})$.

5.1. Full exceptional collections for \tilde{X} . Several choices are involved in the construction of full exceptional collections for \tilde{X} . Our guiding principle is to get exceptional collections whose Hom-spaces are as simple as possible, and we will make a choice explicit when it is made.

We start by letting $\pi: X' \rightarrow \mathbb{P}^2$ be a blow-down map which does not contract any of the lines $c_{i,0}$: such a π always exists by Lemma 4.11. Let $\tau: \tilde{X} \rightarrow X'$ be the morphism in Theorem 4.3 and write $\alpha = \tau \circ \pi$. For $i = 1, 2, 3, 4$ and for $1 \leq j \leq k$, write $b_{i,j} = \mathcal{O}_{\sum_{j=1}^k C_{ij}}$ and $\mathcal{B}_i = (b_{i,1}, \dots, b_{i,k})$. Let $D := F + \sum_{i=1}^4 \sum_{j=1}^k j C_{ij}$.

Theorem 5.1. *Let (A_0, A_1, A_2) be a full exceptional collection on \mathbb{P}^2 and let L_1, \dots, L_7 be the strict transforms in \tilde{X} of the exceptional curves of $\pi: X' \rightarrow \mathbb{P}^2$. Then*

$$(44) \quad \sigma_{\tilde{X}} = (\alpha^*(A_1), \alpha^*(A_1), \alpha^*(A_2), \mathcal{O}_{L_1}, \dots, \mathcal{O}_{L_7}, \mathcal{O}_D, \mathcal{B}_1, \mathcal{B}_2, \mathcal{B}_3, \mathcal{B}_4)$$

is a full exceptional collection on \tilde{X} .

Proof. Let l_1, \dots, l_7 be the exceptional curves of $\pi: X' \rightarrow \mathbb{P}^2$. Then, because of our choice of the map π , we have that for all i, j $c_{i,0} \neq l_j$, thus none of the l_j contains the point q . This implies that $\tau^* \mathcal{O}_{l_i} = \mathcal{O}_{L_i}$. Then the statement follows immediately by an iterative application of Orlov's blow-up formula [Orl92] (as stated in [Kuz14]). \square

Let p, p_1, \dots, p_4 be the five singular points on X , as in Section 4.1. Let $G = \frac{1}{4k+1}(1, 1)$ and $H = \frac{1}{2k+1}(1, k)$. For $j \in \mathbb{Z}/4k+1$ let ρ_j be the irreducible representation of G of weight j and write $e_j = \mathcal{O}_p \otimes \rho_j$. For $j \in \mathbb{Z}/2k+1$ let η_j be the irreducible representation of H of weight j and write $e_{i,j} = \mathcal{O}_{p_i} \otimes \eta_j$. Then:

Corollary 5.2. *The derived category $D(\mathcal{X})$ admits a full exceptional collection of the form*

$$(45) \quad \sigma_{\mathcal{X}} = (\mathcal{E}_p, \mathcal{E}_{p_1}, \mathcal{E}_{p_2}, \mathcal{E}_{p_3}, \mathcal{E}_{p_4}, \Phi(\sigma_{\tilde{\mathcal{X}}})) ,$$

where $\mathcal{E}_p = (e_2, \dots, e_{4k})$, $\mathcal{E}_{p_i} = (e_{i,k+1}, \dots, e_{i,2k})$, $i = 1, 2, 3, 4$. Here, $\sigma_{\tilde{\mathcal{X}}}$ is the full exceptional collection in (44), and Φ is the fully faithful functor from Section 2.3.3.

Proof. An immediate consequence of the global $\mathrm{GL}(2, \mathbb{C})$ McKay correspondence 2.3.3, Theorem 5.1, Example 2.7 and Example 2.8. \square

5.2. Hom-spaces. There are various standard full exceptional collections (A_0, A_1, A_2) on \mathbb{P}^2 . We choose $(A_0, A_1, A_2) = (\mathcal{O}, \mathcal{T}(-1), \mathcal{O}(1))$, where \mathcal{T} is the tangent bundle on \mathbb{P}^2 , following [AKO06]. Moreover, we choose the map $\pi: X' \rightarrow \mathbb{P}^2$ to be a blow-up morphism as described in Example 4.8 (a): this is always possible by Corollary 4.13. This choice implies that the images $\alpha(C_{i,0}) \subset \mathbb{P}^2$ of the curves $C_{i,0}$ are two lines and two conics. Without loss of generality, we assume that $\alpha(C_{i,0})$ is a line if $i = 1, 2$, is a conic if $i = 3, 4$. Moreover, setting $x_j = \alpha(L_j)$, we also assume that $x_1, x_2 \in \alpha(C_{1,0})$, $x_3, x_4 \in \alpha(C_{2,0})$, $x_1, x_3 \in \alpha(C_{3,0})$, and $x_2, x_4 \in \alpha(C_{4,0})$ (see Figure 8).

With these choices, we describe below the spaces of morphisms of the exceptional collections $\sigma_{\tilde{\mathcal{X}}}$ ¹⁰ and

$$(46) \quad \sigma'_{\mathcal{X}} = (\mathcal{E}'_p, \mathcal{E}_{p_1}, \mathcal{E}_{p_2}, \mathcal{E}_{p_3}, \mathcal{E}_{p_4}, \Phi(\sigma_{\tilde{\mathcal{X}}}))$$

where $\mathcal{E}'_p = (e_2[-4k+1], \dots, e_{4k-1}[-2], e_{4k}[-1])$.

More precisely, in Lemma 5.6 and Lemma 5.7 we describe the spaces of morphisms of $\sigma_{\tilde{\mathcal{X}}}$. Since Φ is a fully faithful functor, the same two lemmas also describe all morphisms between objects of $\sigma'_{\mathcal{X}}$ of the form $\Phi(E)$, where $E \in \sigma_{\tilde{\mathcal{X}}}$. Lemma 5.8 and Lemma 5.9 complete the description of the spaces of morphisms of $\sigma'_{\mathcal{X}}$. We omit the computations of the morphisms between objects in \mathcal{E}'_p and \mathcal{E}_{p_i} since they follow directly from the McKay quivers in Example 2.7 and Example 2.8.

Our computations show that the collections $\sigma_{\tilde{\mathcal{X}}}$ and $\sigma'_{\mathcal{X}}$ are not strong. However, they have non-trivial morphisms only in degree 0 and 1, which implies the existence of a tilting complex in $D(\tilde{X})$ and $D(\mathcal{X})$ (see Section 5.3).

In fact, one can change the exceptional collection $\sigma_{\tilde{\mathcal{X}}}$ to another one with simpler spaces of morphisms. The exceptional collection for \tilde{X} obtained by the left mutation of $\sigma_{\tilde{\mathcal{X}}}$ in the pair $(\alpha^*\mathcal{O}(1), \mathcal{O}_D)$ ¹¹, that is,

$$(47) \quad L\sigma_{\tilde{\mathcal{X}}} = (\mathcal{O}_{\tilde{X}}, \alpha^*\mathcal{T}(-1), M, \alpha^*\mathcal{O}(1), \mathcal{O}_{L_1}, \dots, \mathcal{O}_{L_7}, \mathcal{B}_1, \mathcal{B}_2, \mathcal{B}_3, \mathcal{B}_4)$$

has only non-trivial morphisms in degree 0, except for the morphisms from $\mathrm{Ext}^*(b_{i,j}, b_{i,l}) = \mathbb{C} \oplus \mathbb{C}[-1]$ when $j < l$. Here, M denotes the left mutation $M = L_{\alpha^*\mathcal{O}(1)}\mathcal{O}_D$, defined by the short exact sequence

$$(48) \quad 0 \rightarrow M \rightarrow \alpha^*\mathcal{O}(1) \xrightarrow{r} \mathcal{O}_D \rightarrow 0.$$

Moreover, the corresponding exceptional collection for \mathcal{X}

$$(49) \quad L\sigma'_{\mathcal{X}} = (\mathcal{E}'_p, \mathcal{E}_{p_1}, \mathcal{E}_{p_2}, \mathcal{E}_{p_3}, \mathcal{E}_{p_4}, \Phi(L\sigma_{\tilde{\mathcal{X}}}))$$

has only non-trivial morphism in degree 0, except for the morphisms from $\Phi(b_{i,j})$ to $\Phi(b_{i,l})$ when $j < l$ (Lemma 5.10), and the morphisms starting from $e_{i,j}$, which have degree 0 or 1 (Lemma 5.11).

We collect the data of the dimensions of the Hom-spaces between objects of $L\sigma_{\tilde{\mathcal{X}}}$ and $L\sigma'_{\mathcal{X}}$ in Table 1, Table 2 and Table 3. In each table, the entry in position (i, j) is either the integer $\dim \mathrm{Hom}(E_i, E_j)$, or a pair of integers $(\dim \mathrm{Hom}(E_i, E_j), \dim \mathrm{Ext}^1(E_i, E_j))$.

¹⁰The dimension of the Hom spaces of $\sigma_{\tilde{\mathcal{X}}}$ are computed in Section 5.2.2 for arbitrary vector bundles A_0, A_1, A_2 . For $\sigma'_{\mathcal{X}}$ they depend on our choices of A_1, A_2, A_3 and π .

¹¹More precisely, one left-mutates \mathcal{O}_D through $\alpha^*\mathcal{O}(1), \mathcal{O}_{L_1}, \dots, \mathcal{O}_{L_7}$. However, the left mutation $L_{\mathcal{O}_{L_i}}(\mathcal{O}_D) = \mathcal{O}_D$ since \mathcal{O}_D and \mathcal{O}_{L_i} are orthogonal.

Remark 5.3. If $k = 1$, the exceptional collection $L\sigma_{\tilde{\mathcal{X}}}$ is in fact strong. Moreover, by shifting by -1 any element of \mathcal{E}_{p_i} , $i = 1, 2, 3, 4$, one obtains a full strong exceptional collection also for the stack \mathcal{X} . The dimensions of the Hom-spaces of this collection are summarised in Table 4.

It may be interesting to find out whether, for all $k > 0$ integer, one could obtain a full strong exceptional collection for \mathcal{X} through a sequence of mutations and shifts of $L\sigma'_{\mathcal{X}}$.

5.2.1. *Basic facts.* We recall some well-known facts about sheaves of pure dimension 1 on surfaces which we use repeatedly in the proofs of the lemmas. The first one is a Riemann-Roch computation. Here, by a curve on S we mean a connected effective 1-cycle.

Lemma 5.4. *Let S be a smooth projective surface, and let V_1, V_2 be sheaves of pure dimension one on S . For $i = 1, 2$, let D_i be curves on S such that $\text{ch}_1(V_i) = D_i \in H^*(S)$. Then*

$$\chi(V_1, V_2) = \sum_{i \geq 0} (-1)^i \dim \text{Ext}^i(V_1, V_2) = -D_1 \cdot D_2.$$

Proof. Since V_1 and V_2 are torsion, their Chern characters satisfy $\text{ch}_0(V_i) = 0$. Then the Hirzebruch–Riemann–Roch theorem shows the statement. \square

The next Lemma follows directly from the definition of pure sheaf and simple homological algebra:

Lemma 5.5. *Let V, W be sheaves of pure dimension one on S , supported on curves C and D , respectively. Then:*

- (i) *if C and D have no common irreducible component, then $\text{Hom}(V, W) = 0$;*
- (ii) *every map $V \rightarrow W$ factors through the restriction map $V \rightarrow V|_{C \cap D}$;*
- (iii) *Let $D' = \overline{D \setminus (C \cap D)}$. Every map $V \rightarrow W$ factors through the kernel of the restriction map $W \rightarrow W|_{D'}$.*

For future reference, we also list the intersection numbers of $K_{\tilde{\mathcal{X}}}$ with the curves appearing in Figure 6:

$$(50) \quad F \cdot K_{\tilde{\mathcal{X}}} = 4k - 1 \quad C_{i,0} \cdot K_{\tilde{\mathcal{X}}} = 1 \quad C_{i,j} \cdot K_{\tilde{\mathcal{X}}} = 0 \quad C_{i,k} \cdot K_{\tilde{\mathcal{X}}} = -1$$

for $i = 1, \dots, 4$ and $j = 1, \dots, k - 1$.

5.2.2. *Morphisms of the collections $\sigma_{\tilde{\mathcal{X}}}$ and $\sigma'_{\mathcal{X}}$.*

Lemma 5.6. *Let V be a vector bundle on \mathbb{P}^2 . Then $\text{Ext}^*(\alpha^*V, E) = \mathbb{C}^{\text{rank}(V)}$, where E is any object of $(\mathcal{O}_{L_1}, \dots, \mathcal{O}_{L_7}, \mathcal{O}_D, \mathcal{B}_1, \mathcal{B}_2, \mathcal{B}_3, \mathcal{B}_4)$. Moreover, \mathcal{O}_{L_i} is mutually orthogonal to \mathcal{O}_{L_j} for all $i \neq j$, and to any object of $(\mathcal{O}_D, \mathcal{B}_1, \mathcal{B}_2, \mathcal{B}_3, \mathcal{B}_4)$.*

Proof. To prove the first part of the lemma note that, by adjunction,

$$\text{Ext}^*(\alpha^*V, E) = \text{Ext}^*(V, \mathbf{R}\alpha_*E)$$

and $\mathbf{R}\alpha_*E$ is the structure sheaf of a point. The second part of the lemma follows from the fact that the objects involved have disjoint supports. \square

Lemma 5.7. *Let $i = 1, 2, 3, 4$. Then:*

- 1. *for $j = 1, \dots, k$, $\text{Ext}^*(\mathcal{O}_D, b_{i,j}) = \mathbb{C} \oplus \mathbb{C}[-1]$;*
- 2. *\mathcal{B}_i and \mathcal{B}_j , $j \neq i$, are mutually orthogonal;*
- 3. *for $1 \leq j < l \leq k$, $\text{Ext}^*(b_{i,j}, b_{i,l}) = \mathbb{C} \oplus \mathbb{C}[-1]$.*

Proof. To prove (1) and (3) use Lemma 5.4 and Lemma 5.5. We show for example the proof of (1) for $j = k$. It follows from Lemma 5.5 (ii) that there is only one map $\mathcal{O}_D \rightarrow b_{i,k}$. Moreover, by Lemma 5.4

$$\chi(\mathcal{O}_D, b_{i,k}) = -D \cdot C_{i,k} = -(F + (k - 1) \cdot C_{i,k-1} + k \cdot C_{i,k}) \cdot C_{i,k} = 0$$

On the other hand, $\text{Ext}^2(\mathcal{O}_D, b_{i,j}) = \text{Hom}(\mathcal{O}_{C_{i,k}}(1), \mathcal{O}_D)$ by Serre duality and (50), and the latter is trivial by Lemma 5.5 (iii). Assertion (2) is immediate since \mathcal{B}_i and \mathcal{B}_j have disjoint supports. \square

To carry out the rest of the computations, we will combine our result on the left adjoint functor Ψ to Φ with the configuration of curves described in Theorem 4.3. In particular, the local description from Examples 3.5 and 3.6 gives

$$(51) \quad \Psi(e_i) \simeq \begin{cases} 0 & \text{if } i \in 0, \dots, 4k-2 \\ \mathcal{O}_F(-4k-1)[1] & \text{if } i = 4k-1 \\ \mathcal{O}_F(-4k) & \text{if } i = 4k \end{cases}$$

and

$$(52) \quad \Psi(e_{ij}) \simeq \begin{cases} 0 & \text{if } j \in 0, \dots, k-1 \\ \mathcal{O}_{C_i}(C_i)[1] & \text{if } j = k \\ \mathcal{O}_{C_{i,2k-j}}(-1) & \text{if } j \in \{k+1, \dots, 2k-1\} \\ \mathcal{O}_{C_{i,0}}(-2) & \text{if } j = 2k \end{cases}$$

Lemma 5.8. *We have that:*

1. for $2 \leq l < 4k-1$, $\text{Ext}^*(e_l, \Phi(E)) = 0$ for all $E \in D^b(\text{Coh}(\tilde{X}))$;
2. for V a vector bundle on \mathbb{P}^2 , $\text{Ext}^*(e_{4k-1}, \Phi(\alpha^*(V))) = \mathbb{C}^{\text{rank}(V)}[-2]$;
3. $\text{Ext}^*(e_{4k-1}, \Phi(\mathcal{O}_D)) = \mathbb{C}^2[-1] \oplus \mathbb{C}[-2]$;
4. for $i = 1, 2, 3, 4$, $j = 1, \dots, k$, $\text{Ext}^*(e_{4k-1}, \Phi(b_{i,j})) = \mathbb{C}[-1]$;
5. for V a vector bundle on \mathbb{P}^2 , $\text{Ext}^*(e_{4k}, \Phi(\alpha^*(V))) = 0$;
6. $\text{Ext}^*(e_{4k}, \Phi(\mathcal{O}_D)) = \mathbb{C}$;
7. for $i = 1, 2, 3, 4$, $j = 1, \dots, k$, $\text{Ext}^*(e_{4k}, \Phi(b_{i,j})) = \mathbb{C}[-1]$;
8. \mathcal{E}_p and $(\Phi(\mathcal{O}_{L_1}), \dots, \Phi(\mathcal{O}_{L_7}))$ are mutually orthogonal.

Proof. In all the statements above, we use that Ψ is left adjoint to Φ , and conclude with techniques similar to the previous lemmas. We show (3) as an illustration:

$$\text{Ext}^*(e_{4k-1}, \Phi(\mathcal{O}_D)) = \text{Ext}_{\tilde{X}}^*(\mathcal{O}_F(-4k-1)[1], \mathcal{O}_D)$$

by (51). The restriction of \mathcal{O}_D to $\overline{D \setminus F}$ has kernel $\mathcal{O}_F(-4k)$, so we have

$$\text{Hom}_{\tilde{X}}(\mathcal{O}_F(-4k-1), \mathcal{O}_D) = \text{Hom}_{\tilde{X}}(\mathcal{O}_F(-4k-1), \mathcal{O}_F(-4k)) = \mathbb{C}^2$$

by Lemma 5.5 (iii). On the other hand,

$$\text{Ext}_{\tilde{X}}^2(\mathcal{O}_F(-4k-1), \mathcal{O}_D) \simeq \text{Hom}(\mathcal{O}_D, \mathcal{O}_F(-2))^* = \text{Hom}(\mathcal{O}_F, \mathcal{O}_F(-2)) = 0$$

by Serre duality, (50), and Lemma 5.5 (ii). Since $\chi(\mathcal{O}_F(-4k-1), \mathcal{O}_D) = -F \cdot D = 1$ (using Lemma 5.4), we show (3). □

Lemma 5.9. *Let $i = 1, 2, 3, 4$. Then:*

1. for V a vector bundle on \mathbb{P}^2 , $\text{Ext}^*(e_{i,j}, \Phi(\alpha^*(V))) = 0$ for all $j \neq 2k$;
2. $e_{i,j}$ and $\Phi(\mathcal{O}_{L_h})$ are mutually orthogonal for all $j \neq 2k$;
3. $\text{Ext}^*(e_{i,2k}, \Phi(\mathcal{O}_{\tilde{X}})) = 0$;
4. $\text{Ext}^*(e_{i,2k}, \Phi(\alpha^*(\mathcal{T}(-1)))) = \text{Ext}^*(e_{i,2k}, \Phi(\alpha^*(\mathcal{O}(1)))) = \begin{cases} \mathbb{C}[-1] & \text{if } i = 1, 2 \\ \mathbb{C}^2[-1] & \text{if } i = 3, 4 \end{cases}$
5. $\text{Ext}^*(e_{i,2k}, \Phi(\mathcal{O}_{L_h})) = \mathbb{C}^{a_{i,j}}[-1]$, where $a_{i,j}$ is the entry (i, j) of the matrix:

$$A = \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 & 1 & 1 & 1 \end{pmatrix}$$

6. $\text{Ext}^*(e_{i,j}, \Phi(\mathcal{O}_D)) = \begin{cases} 0 & \text{if } j < 2k \\ \mathbb{C}[-1] & \text{if } j = 2k \end{cases}$

$$7. \text{Ext}^*(e_{i,j}, \Phi(b_{i,l})) = \begin{cases} \mathbb{C} & \text{if } l = 2k - j \\ \mathbb{C}[-1] & \text{if } l = 2k - j + 1 \\ 0 & \text{otherwise} \end{cases}$$

8. \mathcal{E}_i and $\Phi(\mathcal{B}_j)$, $j \neq i$, are mutually orthogonal.

Proof. Again, in all the statements above, we use that Ψ is left adjoint to Φ and we conclude using ideas similar to those of the previous lemmas. As an example, we show (4).

By (52), Serre duality, (50), and adjunction,

$$\begin{aligned} \text{Ext}^*(e_{i,2k}, \Phi(\alpha^*(\mathcal{T}(-1)))) &= \text{Ext}^*(\mathcal{O}_{C_{i,0}}(-2), \alpha^*(\mathcal{T}(-1))) \\ &= \text{Ext}^{2-*}(\alpha^*(\mathcal{T}(-1)), \mathcal{O}_{C_{i,0}}(-1)) \\ &= \text{Ext}^{2-*}(\mathcal{T}(-1), \mathcal{O}_{\alpha(C_{i,0})}(-1)) \end{aligned}$$

Now, by the long exact sequence for the Hom functor applied to the Euler sequence, one finds that

$$\text{Ext}^*(\mathcal{T}(-1), \mathcal{O}_{\alpha(C_{i,0})}(-1)) = \text{Ext}^{*-1}(\mathcal{O}(-1), \mathcal{O}_{\alpha(C_{i,0})}(-1))$$

since $H^*(\mathbb{P}^2, \mathcal{O}_{\alpha(C_{i,0})}(-1)) = 0$ because $C_{i,0}$ is rational. One concludes by observing that

$$\text{Ext}^*(\mathcal{O}(-1), \mathcal{O}_{\alpha(C_{i,0})}(-1)) = \begin{cases} \text{Ext}^*(\mathcal{O}, \alpha(C_{i,0})) = \mathbb{C} & \text{if } \alpha(C_{i,0}) \text{ is a line,} \\ \text{Ext}^*(\mathcal{O}, \alpha(C_{i,0})(1)) = \mathbb{C}^2 & \text{if } \alpha(C_{i,0}) \text{ is a conic.} \end{cases}$$

Analogously,

$$\text{Ext}^*(e_{i,2k}, \Phi(\alpha^*(\mathcal{O}(1)))) = \text{Ext}^*(\mathcal{O}_{C_{i,0}}(-2), \alpha^*(\mathcal{O}(1))) = \text{Ext}^{2-*}(\mathcal{O}(1), \mathcal{O}_{\alpha(C_{i,0})}(-1))$$

Hence, $\text{Ext}^*(e_{i,2k}, \Phi(\alpha^*(\mathcal{O}(1))))$ is $\mathbb{C}[-1]$ if $\alpha(C_{i,0})$ is a line, $\mathbb{C}^2[-1]$ if $\alpha(C_{i,0})$ is a conic. \square

5.2.3. Morphisms of the mutated collections.

Lemma 5.10. *Let M be as in (48). Then:*

1. $\text{Ext}^*(\mathcal{O}_{\tilde{X}}, M) = \mathbb{C}^2$
2. $\text{Ext}^*(\alpha^*(\mathcal{T}(-1)), M) = \mathbb{C}$
3. $\text{Ext}^*(M, \alpha^*(\mathcal{O}(1))) = \mathbb{C}$
4. for $i = 1, \dots, 7$, $\text{Ext}^*(M, \mathcal{O}_{L_i}) = \mathbb{C}$
5. for $i = 1, 2, 3, 4$, $j = 1, \dots, k$, $\text{Ext}^*(M, b_{i,j}) = \mathbb{C}$

Proof. All the computations follow from the long exact sequence of the Hom functor applied to (48), combined with Lemmas 5.6 and 5.7. For example, composition with r gives a surjection

$$\text{Hom}(\alpha^*(\mathcal{T}(-1)), \alpha^*(\mathcal{O}(1))) \twoheadrightarrow \text{Hom}(\alpha^*(\mathcal{T}(-1)), \mathcal{O}_D),$$

which immediately implies (2). \square

Lemma 5.11. *Let M be as in (48). Then:*

1. $\text{Ext}^*(e_l, \Phi(M)) = \begin{cases} 0 & \text{if } 2 \leq l < 4k - 1 \\ \mathbb{C}^2[-2] & \text{if } l = 4k - 1 \\ \mathbb{C}[-1] & \text{if } l = 4k \end{cases}$
2. for $i = 1, 2, 3, 4$ and for $k < j < 2k$, $\text{Ext}^*(e_{i,j}, \Phi(M)) = 0$;
3. $\text{Ext}^*(e_{i,2k}, \Phi(M)) = \begin{cases} 0 & \text{if } i = 1, 2 \\ \mathbb{C}[-1] & \text{if } i = 3, 4 \end{cases}$

Proof. All the computations follow from the long exact sequence of the Hom functor applied to (48), together with Lemma 5.8 and Lemma 5.9. For example, assertions (1) and (6) in Lemma 5.9 immediately imply (2). \square

5.3. Exceptional collections and tilting. Let \mathcal{D} be a triangulated category. We say that $T \in \mathcal{D}$ *generates* \mathcal{D} if, for every $E \in \mathcal{D}$, we have that $\mathrm{Hom}(T, E) = 0$ implies $E \simeq 0$, and that T *classically generates* \mathcal{D} if the smallest full triangulated subcategory containing T , closed under isomorphisms and direct summands, is \mathcal{D} itself. An object $T \in \mathcal{D}$ is *partial tilting* if $\mathrm{Hom}(T, T[l]) = 0$ for all $l \neq 0$. We say that T is a *tilting object* if, moreover, it generates \mathcal{D} .

Generally speaking, tilting objects produce equivalences between \mathcal{D} and derived categories of modules over their endomorphism algebra. More precisely, suppose $\mathcal{D} = D^b(\mathrm{Coh}(Z))$ where Z is a smooth Deligne-Mumford stack with finite stabilizers and projective coarse moduli space. Then we have:

Proposition 5.12. *Let T be a tilting complex in \mathcal{D} and denote by $\Lambda = \mathrm{Hom}(T, T)$ its endomorphism algebra. Then, the functor $\mathbf{R}\mathrm{Hom}_{\mathcal{O}_Z}(T, -)$ defines an equivalence of triangulated categories*

$$\mathcal{D} \xrightarrow{\sim} D^b(\mathrm{mod}\text{-}\Lambda)$$

with inverse $-\otimes_{\Lambda}^{\mathbf{L}} T$.

Proof. For schemes, this is [HVdB07, Theorem 7.6]. If \mathcal{X} is a smooth DM stack, $D(\mathrm{QCoh}(Z))$ is compactly generated (this is shown for the more general class of *perfect* stacks in [BZFN10, Prop. 3.9]¹²). Then, T generates $D(\mathrm{QCoh}(Z))$ if and only if it classically generates \mathcal{D} [BvdB03, Theorem 2.1.2], and the standard proof (see e.g. [Bae88, §3]) suffices to conclude. \square

If (E_1, \dots, E_n) is a strong full exceptional collection in \mathcal{D} , then $\mathcal{E} = \bigoplus_{i=1}^n E_i$ is a tilting complex. If the collection is not strong, $\Lambda_{\mathcal{E}} = \mathrm{Hom}(\mathcal{E}, \mathcal{E})$ is a dg-algebra, and one may still obtain an equivalence, analogous to Prop. 5.12, between \mathcal{D} and the derived category of dg-modules over $\Lambda_{\mathcal{E}}$. Alternatively, if the exceptional collection only admits maps in degrees 0 and 1, we obtain an explicit tilting complex from the so called *universal extension trick*, which we briefly recall here:

Lemma 5.13. *Assume that (E_1, \dots, E_n) is a collection of objects of \mathcal{D} such that:*

- (i) *each E_i is partial tilting;*
- (ii) *the sum $\bigoplus_{i=1}^n E_i$ generates $D^b(\mathrm{Coh}(\mathcal{X}))$;*
- (iii) *if $i < j$, then $\mathrm{Hom}(E_i, E_j[l]) = 0$ for all $l \neq 0, 1$;*
- (iv) *if $i < j$, then $\mathrm{Hom}(E_j, E_i[l]) = 0$ for all $l \in \mathbb{Z}$.*

Then \mathcal{D} admits a tilting complex.

Proof. We direct the reader to [Har21, Lemma 2.4] for the details of the proof, we only recall the explicit construction here.

Proceed by induction on n : the case $n = 1$ is clear. Assume now that $n > 1$. Let r denote the dimension of $\mathrm{Ext}^1(E_1, E_2)$, then the universal extension map produces a triangle

$$(53) \quad E_2 \rightarrow F \rightarrow E_1^{\oplus r}.$$

Put $E'_2 = E_2 \oplus F$ and $E'_i = E_i$ for $i = 3, \dots, n$. Then, $\{E'_2, E'_3, \dots, E'_n\}$ still satisfies the assumptions. This claim is proven as in [Har21]. The only significant difference is that we do not assume the E_i to be sheaves, and therefore need more care to control negative Ext groups. \square

Corollary 5.14. *Suppose (E_1, \dots, E_n) is a full exceptional collection of \mathcal{D} , such that $\mathrm{Hom}(E_i, E_j[l]) = 0$ for all $l \neq 0, 1$. Then \mathcal{D} admits a tilting complex.*

Proof. This is immediate from Lemma 5.13. \square

Applying this to the case at hand:

Proposition 5.15. *The exceptional collections $L\sigma_{\tilde{X}}$ and $L\sigma'_{\mathcal{X}}$ from Section 5.2 give rise to tilting complexes in $D(\tilde{X})$ and $D(\mathcal{X})$.*

¹²Smooth DM stacks with finite stabilizers have affine diagonal, and hence are perfect. Our setting is analogous to that of [AU15, Lemma 4.1], which follows a similar argument.

	$\mathcal{O}_{\tilde{X}}$	$\alpha^*\mathcal{T}(-1)$	M	$\alpha^*\mathcal{O}(1)$	\mathcal{O}_{L_t}	$b_{i,j}$	$b_{i,l}$
$\mathcal{O}_{\tilde{X}}$	1	3	2	3	1	1	1
$\alpha^*\mathcal{T}(-1)$		1	1	3	2	2	2
M			1	1	1	1	1
$\alpha^*\mathcal{O}(1)$				1	1	1	1
\mathcal{O}_{L_t}					1	0	0
$b_{i,j}$						1	(1, 1)
$b_{i,l}$							1

Table 1. The Hom-spaces of the collection $L\sigma_{\tilde{X}}$.

	$e_{4k-j}[-j-1]$	$\Phi(\mathcal{O}_{\tilde{X}})$	$\Phi\alpha^*\mathcal{T}(-1)$	ΦM	$\Phi\alpha^*\mathcal{O}(1)$	$\Phi\mathcal{O}_{L_t}$	$\Phi b_{i,1}$
\vdots							
$e_{4k-2}[-3]$	McKay quiver (with shifts)	0	0	0	0	0	0
$e_{4k-1}[-2]$	Example 2.7	1	2	2	1	0	1
$e_{4k}[-1]$		0	0	1	0	0	1

Table 2. The Hom-spaces between objects in \mathcal{E}'_p and from \mathcal{E}'_p to $\Phi(L\sigma_{\tilde{X}})$. The shifts of the objects in \mathcal{E}'_p make the arrows from the McKay quiver into maps of degree 0.

	$e_{i,2k-j}$	$\Phi(\mathcal{O}_{\tilde{X}})$	$\Phi\alpha^*\mathcal{T}(-1)$	ΦM	$\Phi\alpha^*\mathcal{O}(1)$	$\Phi\mathcal{O}_{L_t}$	$\Phi b_{i,2k-j}$	$\Phi b_{i,2k-j+1}$
$e_{i,k+1}$		0	0	0	0	0
\vdots								
$e_{i,j}$	McKay quiver	0	0	0	0	0	(1, 0)	(0, 1)
\vdots	Example 2.8							
$e_{i,2k-1}$		0	0	0	0	0
$e_{1,2k}$		0	(0, 1)	0	(0, 1)	(0, $a_{1,t}$)
$e_{3,2k}$		0	(0, 2)	(0, 1)	(0, 2)	(0, $a_{3,t}$)

Table 3. The Hom-spaces from between objects in \mathcal{E}_{p_i} and from \mathcal{E}_{p_i} to $\Phi(L\sigma_{\tilde{X}})$. The arrows of the McKay quiver involved in the first column are morphisms of degree 1. Thus every $e_{i,j}$ admits both Hom's and Ext¹'s to other objects of the collection.

	$e_2[-3]$	$e_3[-2]$	$e_4[-1]$	$e_{i,2}[-1]$	$\Phi\alpha^*\mathcal{O}_{\tilde{X}}$	$\Phi\alpha^*\mathcal{T}(-1)$	ΦM	$\Phi\alpha^*\mathcal{O}(1)$	$\Phi\mathcal{O}_{L_t}$	$\Phi\mathcal{O}_{C_{i,1}}$
$e_2[-3]$	1	2	1	0	0	0	0	0	0	0
$e_3[-2]$		1	2	$O_{2 \times 4}$	1	2	2	1	$O_{2 \times 7}$	1
$e_4[-1]$			1		0	0	1	0		1
$e_{1,2}[-1]$				Id_4	0	1	0	1	A	Id_4
$e_{2,2}[-1]$					0	1	0	1		
$e_{3,2}[-1]$					0	2	1	2		
$e_{4,2}[-1]$					0	2	1	2		

Table 4. The strong exceptional collection for \mathcal{X} when $k = 1$.

REFERENCES

- [ACC⁺16] Mohammad Akhtar, Tom Coates, Alessio Corti, Liana Heuberger, Alexander Kasprzyk, Alessandro Oneto, Andrea Petracci, Thomas Prince, and Ketil Tveiten, *Mirror symmetry and the classification of orbifold del Pezzo surfaces*, Proc. Amer. Math. Soc. **144** (2016), no. 2, 513–527. MR 3430830
- [AKO06] Denis Auroux, Ludmil Katzarkov, and Dmitri Orlov, *Mirror symmetry for del Pezzo surfaces: vanishing cycles and coherent sheaves*, Invent. Math. **166** (2006), no. 3, 537–582. MR 2257391
- [AKO08] ———, *Mirror symmetry for weighted projective planes and their noncommutative deformations*, Ann. of Math. (2) **167** (2008), no. 3, 867–943. MR 2415388
- [Art66] Michael Artin, *On isolated rational singularities of surfaces*, Amer. J. Math. **88** (1966), 129–136. MR 199191

- [AU15] Tarig Abdelgadir and Kazushi Ueda, *Weighted projective lines as fine moduli spaces of quiver representations*, *Comm. Algebra* **43** (2015), no. 2, 636–649. MR 3274027
- [Aus86] Maurice Auslander, *Rational singularities and almost split sequences*, *Trans. Amer. Math. Soc.* **293** (1986), no. 2, 511–531. MR 816307
- [Bae88] Dagmar Baer, *Tilting sheaves in representation theory of algebras*, *Manuscripta Math.* **60** (1988), no. 3, 323–347. MR 928291
- [BM02] Tom Bridgeland and Antony Maciocia, *Fourier-Mukai transforms for K3 and elliptic fibrations*, *J. Algebraic Geom.* **11** (2002), no. 4, 629–657. MR 1910263
- [Bri02] Tom Bridgeland, *Flops and derived categories*, *Invent. Math.* **147** (2002), no. 3, 613–632. MR 1893007
- [BvdB03] A. Bondal and M. van den Bergh, *Generators and representability of functors in commutative and noncommutative geometry*, *Mosc. Math. J.* **3** (2003), no. 1, 1–36, 258. MR 1996800
- [BZFN10] David Ben-Zvi, John Francis, and David Nadler, *Integral transforms and Drinfeld centers in derived algebraic geometry*, *J. Amer. Math. Soc.* **23** (2010), no. 4, 909–966. MR 2669705
- [CCG⁺13] Tom Coates, Alessio Corti, Sergey Galkin, Vasily Golyshev, and Alexander Kasprzyk, *Mirror symmetry and Fano manifolds*, *European Congress of Mathematics*, *Eur. Math. Soc.*, Zürich, 2013, pp. 285–300. MR 3469127
- [CG21] Alessio Corti and Giulia Gugiatti, *Hyperelliptic integrals and mirrors of the Johnson-Kollár del Pezzo surfaces*, *Trans. Amer. Math. Soc.* **374** (2021), no. 12, 8603–8637. MR 4337923
- [CLS11] David A Cox, John B Little, and Henry K Schenck, *Toric varieties*, vol. 124, *American Mathematical Soc.*, 2011.
- [Dem80] Michel Demazure, *Surfaces de del Pezzo II, III, IV, V*, *Séminaire sur les singularités des surfaces*, *Lecture notes in Mathematics*, vol. 777, Springer, Berlin, 1980, pp. 23–69.
- [Dol12] Igor V. Dolgachev, *Classical algebraic geometry*, Cambridge University Press, Cambridge, 2012, A modern view. MR 2964027
- [Ela07] A. D. Elagin, *Exceptional sets on some del Pezzo surfaces with one log-terminal singularity*, *Mat. Zametki* **82** (2007), no. 1, 36–51. MR 2374881
- [Esn85] Hélène Esnault, *Reflexive modules on quotient surface singularities*, *J. Reine Angew. Math.* **362** (1985), 63–71. MR 809966
- [Giv98] Alexander Givental, *A mirror theorem for toric complete intersections*, *Topological field theory, primitive forms and related topics (Kyoto, 1996)*, *Progr. Math.*, vol. 160, Birkhäuser Boston, Boston, MA, 1998, pp. 141–175. MR 1653024
- [Gol07] Vasily V. Golyshev, *Classification problems and mirror duality*, *Surveys in geometry and number theory: reports on contemporary Russian mathematics*, *London Math. Soc. Lecture Note Ser.*, vol. 338, Cambridge Univ. Press, Cambridge, 2007, pp. 88–121. MR 2306141
- [Gol08] V. Golyshev, *Spectra and strains*, arXiv e-prints (2008), arXiv:0801.0432.
- [GPS15] Sheel Ganatra, Timothy Perutz, and Nick Sheridan, *Mirror symmetry: from categories to curve counts*, arXiv e-prints (2015), arXiv:1510.03839.
- [GS18] Mark Gross and Bernd Siebert, *Intrinsic mirror symmetry and punctured Gromov–Witten invariants*, *Algebraic geometry: Salt Lake City 2015*, *Proc. Sympos. Pure Math.*, vol. 97, Amer. Math. Soc., Providence, RI, 2018, pp. 199–230. MR 3821173
- [Har21] Wahei Hara, *On the Abuaf-Ueda flop via non-commutative crepant resolutions*, *SIGMA Symmetry Integrability Geom. Methods Appl.* **17** (2021), Paper No. 044, 22. MR 4251444
- [HV00] Kentaro Hori and Cumrun Vafa, *Mirror Symmetry*, arXiv e-prints (2000).
- [HVdB07] Lutz Hille and Michel Van den Bergh, *Fourier-Mukai transforms*, *Handbook of tilting theory*, *London Math. Soc. Lecture Note Ser.*, vol. 332, Cambridge Univ. Press, Cambridge, 2007, pp. 147–177. MR 2384610
- [IF00] A. R. Iano-Fletcher, *Working with weighted complete intersections*, *Explicit birational geometry of 3-folds*, *London Math. Soc. Lecture Note Ser.*, vol. 281, Cambridge Univ. Press, Cambridge, 2000, pp. 101–173. MR 1798982
- [Ish02] Akira Ishii, *On the McKay correspondence for a finite small subgroup of $GL(2, \mathbb{C})$* , *J. Reine Angew. Math.* **549** (2002), 221–233. MR 1916656
- [IU15] Akira Ishii and Kazushi Ueda, *The special McKay correspondence and exceptional collections*, *Tohoku Math. J. (2)* **67** (2015), no. 4, 585–609. MR 3436544
- [JK01] Jennifer M. Johnson and János Kollár, *Kähler–Einstein metrics on log del Pezzo surfaces in weighted projective 3-spaces*, *Ann. Inst. Fourier (Grenoble)* **51** (2001), no. 1, 69–79. MR 1821068
- [Kaw06] Yujiro Kawamata, *Derived categories of toric varieties*, *Michigan Math. J.* **54** (2006), no. 3, 517–535. MR 2280493
- [Kol96] János Kollár, *Rational curves on algebraic varieties*, *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*, vol. 32, Springer-Verlag, Berlin, 1996. MR 1440180
- [Kon95a] Maxim Kontsevich, *Homological algebra of mirror symmetry*, *Proceedings of the International Congress of Mathematicians*, Vol. 1, 2 (Zürich, 1994), Birkhäuser, Basel, 1995, pp. 120–139. MR 1403918

- [Kon95b] ———, *Homological algebra of mirror symmetry*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994), Birkhäuser, Basel, 1995, pp. 120–139. MR 1403918
- [Kuw05] Masato Kuwata, *Twenty-eight double tangent lines of a plane quartic curve with an involution and the Mordell-Weil lattices*, Comment. Math. Univ. St. Pauli **54** (2005), no. 1, 17–32. MR 2153953
- [Kuz14] Alexander Kuznetsov, *Semiorthogonal decompositions in algebraic geometry*, Proceedings of the International Congress of Mathematicians—Seoul 2014. Vol. II, Kyung Moon Sa, Seoul, 2014, pp. 635–660. MR 3728631
- [Man86] Yu. I. Manin, *Cubic forms: algebra, geometry, arithmetic*, North-Holland Mathematical Library, Vol. 4, North-Holland Publishing Co., Amsterdam-London; American Elsevier Publishing Co., Inc., New York, 1986, Translated from the Russian by M. Hazewinkel. MR 0460349
- [OP18] Alessandro Oneto and Andrea Petracci, *On the quantum periods of del Pezzo surfaces with $\frac{1}{3}(1, 1)$ singularities*, Adv. Geom. **18** (2018), no. 3, 303–336. MR 3830183
- [Orl92] D. O. Orlov, *Projective bundles, monoidal transformations, and derived categories of coherent sheaves*, Izv. Ross. Akad. Nauk Ser. Mat. **56** (1992), no. 4, 852–862. MR 1208153
- [Prz07] Victor Przyjalkowski, *On Landau-Ginzburg models for Fano varieties*, Commun. Number Theory Phys. **1** (2007), no. 4, 713–728. MR 2412270
- [Prz11] ———, *Hori-Vafa mirror models for complete intersections in weighted projective spaces and weak Landau-Ginzburg models*, Cent. Eur. J. Math. **9** (2011), no. 5, 972–977. MR 2824440
- [Sei08] Paul Seidel, *Fukaya categories and Picard-Lefschetz theory*, Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2008. MR 2441780
- [TVAV09] Damiano Testa, Anthony Várilly-Alvarado, and Mauricio Velasco, *Cox rings of degree one del Pezzo surfaces*, Algebra Number Theory **3** (2009), no. 7, 729–761. MR 2579393
- [vdB04a] Michel van den Bergh, *Non-commutative crepant resolutions*, The legacy of Niels Henrik Abel, Springer, Berlin, 2004, pp. 749–770. MR 2077594
- [VdB04b] Michel Van den Bergh, *Three-dimensional flops and noncommutative rings*, Duke Math. J. **122** (2004), no. 3, 423–455. MR 2057015
- [Wun87] J. Wunram, *Reflexive modules on cyclic quotient surface singularities*, Singularities, representation of algebras, and vector bundles (Lambrecht, 1985), Lecture Notes in Math., vol. 1273, Springer, Berlin, 1987, pp. 221–231. MR 915177
- [Wun88] Jürgen Wunram, *Reflexive modules on quotient surface singularities*, Math. Ann. **279** (1988), no. 4, 583–598. MR 926422

GG: MATH SECTION, ICTP, LEONARDO DA VINCI BUILDING, STRADA COSTIERA 11, 34151 TRIESTE, ITALY
 Email address: ggugiatt@ictp.it

FR: SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF GLASGOW, GLASGOW G12 8QQ, UNITED KINGDOM
 Email address: franco.rota@glasgow.ac.uk