

# FINITE ABELIAN SUBGROUPS IN THE GROUPS OF BIRATIONAL AND BIMEROMORPHIC SELFMAPS

ALEKSEI GOLOTA

ABSTRACT. Let  $X$  be a complex projective variety. Suppose that the group of birational automorphisms of  $X$  contains finite subgroups isomorphic to  $(\mathbb{Z}/N_i\mathbb{Z})^r$  for  $r$  fixed and  $N_i$  arbitrarily large. We show that  $r$  does not exceed  $2 \dim(X)$ . We also show that the same result holds for groups of bimeromorphic automorphisms of compact Kähler threefolds.

## 1. INTRODUCTION

This paper studies finite abelian subgroups in the groups of birational automorphisms of projective varieties over a field of zero characteristic, or in the groups of bimeromorphic automorphisms of compact Kähler spaces. The starting point for us is the following question, posed in [Xu18, Question 1.2].

*Question 1.1.* Let  $X$  be a projective variety over an algebraically closed field of characteristic zero. Suppose that for some  $r \in \mathbb{N}$  there exists a subgroup  $G \subseteq \text{Bir}(X)$  isomorphic to  $(\mathbb{Z}/p\mathbb{Z})^r$ , for a sufficiently large prime number  $p$ . Is  $X$  birational to a variety having an effective action of an algebraic torus or an abelian variety?

The answer to this question in full generality is currently unknown. However, in [Xu18, Theorem 2.9] Jinsong Xu gave an affirmative answer to this question for non-uniruled projective varieties.

**Theorem 1.2.** *Let  $X$  be a non-uniruled variety over an algebraically closed field of characteristic zero. There exists a constant  $b(X)$  such that the group  $\text{Bir}(X)$  contains an element of order greater than  $b(X)$  if and only if  $X$  is birational to a variety  $X'$  which admits an effective action of an abelian variety.*

Moreover, J. Xu proved a rationality criterion for rationally connected varieties admitting an action of  $(\mathbb{Z}/p\mathbb{Z})^r$  in terms of  $r$  and  $p$  (see [Xu18, Theorem 4.5]).

**Theorem 1.3.** *Let  $X$  be a rationally connected variety of dimension  $n$  over an algebraically closed field of characteristic zero. Then there exists a constant  $R(n)$  such that if  $\text{Bir}(X)$  contains a subgroup isomorphic to  $(\mathbb{Z}/p\mathbb{Z})^r$  for some  $p > R(n)$  then  $X$  is rational.*

Informally speaking, these results suggest that finite abelian subgroups in  $\text{Bir}(X)$  of unbounded orders should come from certain algebraic groups acting on  $X$  by birational automorphisms.

Recently, I. Mundet i Riera [Mun21] studied finite abelian groups of unbounded orders acting on topological and differentiable manifolds. Among many other results, he proved a result similar to Theorem 1.2 for groups of biholomorphic automorphisms of compact Kähler manifolds [Mun21, Theorem 1.15].

**Theorem 1.4.** *Let  $X$  be a compact Kähler manifold. Suppose that there exists  $r \in \mathbb{N}$  such that for infinitely many positive integers  $N_i$  the group  $\text{Aut}(X)$  contains a subgroup isomorphic to  $(\mathbb{Z}/N_i\mathbb{Z})^r$ . Then  $\text{Aut}(X)$  contains a subgroup isomorphic to a compact real torus of dimension  $r$ . Moreover, one has  $r \leq 2 \dim(X)$ , and if  $r = 2 \dim(X)$  then  $X$  is biholomorphic to a compact complex torus.*

In this paper we study Question 1.1 in the case when the value of  $r$  is maximal. In particular, we show that  $r$  is always bounded from above by  $2 \dim(X)$ . Our main theorem is as follows.

**Theorem 1.5.** *Let  $X$  be a projective variety over an algebraically closed field of zero characteristic. Suppose that the group  $\text{Bir}(X)$  contains subgroups isomorphic to  $(\mathbb{Z}/N_i\mathbb{Z})^r$  for some fixed  $r$  and arbitrarily large  $N_i$ . Then  $r \leq 2 \dim(X)$  and in case of equality  $X$  is birational to an abelian variety.*

---

The author was supported by the Basic Research Program of the National Research University Higher School of Economics.

Compared to Theorem 1.2, in our theorem we allow  $X$  to be uniruled. Also the orders  $N_i$  of generators of the finite groups are not necessarily prime. The main ideas of the proof are, however, already present in J. Xu's work (see e. g. the proof of [Xu18, Proposition 2.12]). Combining these ideas with some technical results from our paper [Gol21], we prove the same result for groups of bimeromorphic selfmaps of compact Kähler spaces of dimension 3.

**Theorem 1.6.** *Let  $X$  be a compact Kähler space of dimension 3. Suppose that the group  $\text{Bim}(X)$  contains subgroups isomorphic to  $(\mathbb{Z}/N_i\mathbb{Z})^r$  for some fixed  $r$  and arbitrarily large  $N_i$ . Then  $r \leq 6$  and in case of equality  $X$  is bimeromorphic to a compact complex torus.*

Let us outline the structure of the paper. In Section 2 we gather some technical results. Section 3 is devoted to the proof of our main theorem. First, in subsection 3.1 we use techniques from our previous paper [Gol21] to generalize Theorem 1.4 to pseudoautomorphisms of compact Kähler spaces with rational singularities (see Theorem 3.5). Then, in subsection 3.2 we prove the main theorem for non-uniruled projective varieties (Theorem 3.10), following the ideas from [Xu18, Section 2]. In subsection 3.3 we use the results of Prokhorov and Shramov from [PS16] to derive the bound on  $r$  for abelian groups acting on rationally connected varieties. Finally, in subsection 3.4 we derive Theorem 1.5 from Theorems 3.10 and 3.14, using the maximal rationally connected fibration of  $X$ . We also prove Theorem 1.6 in this section.

**Acknowledgement.** The author thanks Constantin Shramov for suggesting this problem and Ignasi Mundet i Riera for valuable remarks.

## 2. PRELIMINARIES

**2.1. Structure of abelian subgroups.** In this subsection we collect a few technical statements about subgroups of finite abelian groups.

**Lemma 2.1.** *Let  $G \simeq (\mathbb{Z}/N\mathbb{Z})^r$  be a finite abelian group. Let  $H \subset G$  be a subgroup of index  $I_H \leq N - 1$ . Then there exists a subgroup  $H' \subseteq H$  such that  $H' \simeq (\mathbb{Z}/N'\mathbb{Z})^r$  for some  $N' \geq N/I_H$ .*

*Proof.* Let  $H \subseteq G$  be a subgroup of index  $I_H \leq N - 1$ . Then  $H$  is a finite abelian group of order at least  $N(r - 1) + 1$ . The orders of generators of  $H$  do not exceed  $N$ , so the group  $H$  is generated by at least  $r$  elements. By the structure theorem for finite abelian groups we have

$$H \simeq \bigoplus_{1 \leq i \leq r} \mathbb{Z}/N_i\mathbb{Z}$$

where  $N_i | N_{i+1}$  for all  $i \in \{1, \dots, r - 1\}$ . Moreover, from the equality

$$|H| = N^r / I_H = N_1 \cdots N_r$$

we obtain that  $N_1 \geq N/I_H$ . Then we can take  $H'$  to be a subgroup of the form  $(\mathbb{Z}/N_1\mathbb{Z})^r \subseteq H$ .  $\square$

**Lemma 2.2.** *Let  $G \simeq (\mathbb{Z}/N\mathbb{Z})^r$  be a finite abelian group and let  $H \subset G$  be a subgroup. There exists a set of generators  $\{b_1, \dots, b_r\}$  for  $H$  and a set of generators  $\{a'_1, \dots, a'_r\}$  for  $G$  such that the embedding  $H \rightarrow G$  can be written as the direct sum of homomorphisms*

$$\mathbb{Z}/N_i\mathbb{Z} \rightarrow \mathbb{Z}/N\mathbb{Z}.$$

*Proof.* Choose a set of generators  $a_1, \dots, a_r$  for  $\mathbb{Z}^r$  such that their images under the projection

$$\mathbb{Z}^r \rightarrow \mathbb{Z}^r / (N\mathbb{Z})^r$$

give an isomorphism  $(\mathbb{Z}/N\mathbb{Z})^r \simeq G$ . Let us denote by  $\tilde{H}$  the preimage of  $H$  under the map  $\mathbb{Z}^r \rightarrow G$ . Then  $\tilde{H}$  is a free abelian group of rank  $r$  containing the subgroup  $(\mathbb{Z}/N\mathbb{Z})^r$ . Let  $A$  be a presentation matrix for  $\tilde{H}$ . Then (see e. g. [Art91, Theorem (4.3)]) there exists the Smith normal form

$$A' = QAP^{-1}$$

where  $Q, P \in \text{GL}_r(\mathbb{Z})$  and the matrix  $A'$  is diagonal with entries  $N_1, \dots, N_r$  such that  $N_i$  divides  $N_{i+1}$  for any  $i \in \{1, \dots, r - 1\}$ . So there exists a basis  $a'_1, \dots, a'_r$  of  $\mathbb{Z}^r$  such that  $\tilde{H}$  is generated by

$$b_1 = a'_1 N_1, \dots, b_r = a'_r N_r.$$

Since multiplication by invertible matrices preserves the sublattice  $(N\mathbb{Z})^r \subset \mathbb{Z}^r$ , we can take the images of  $b_1, \dots, b_r$  under the projection  $\mathbb{Z}^r \rightarrow \mathbb{Z}^r / (n\mathbb{Z})^r \simeq G$ .  $\square$

It will be useful to introduce the following definition.

**Definition 2.3.** Let  $\{G_i\}, i \in \mathbb{N}$  be a sequence of finite groups. We define the *unbounded rank* of the sequence  $\{G_i\}$  to be the minimal number  $r$  such that the following condition is satisfied. There exists a constant  $L$  such that for infinitely many  $i \in \mathbb{N}$  we can find an abelian subgroup  $H_i \subset G_i$  such that

- $H_i$  is generated by  $r$  elements;
- $\frac{|G_i|}{|H_i|} \leq L$ .

*Example 2.4.* If the orders of the groups  $G_i$  are bounded by a constant then the unbounded rank of the sequence  $\{G_i\}$  is equal to zero.

*Example 2.5.* The unbounded rank of the sequence  $G_i = (\mathbb{Z}/2\mathbb{Z})^i$  is infinite.

*Example 2.6.* The unbounded rank of the sequence  $G_i = (\mathbb{Z}/i\mathbb{Z})^r \times (\mathbb{Z}/2\mathbb{Z})^{10}$  is equal to  $r$ .

*Remark 2.7.* Let  $\{G_i\}$  be a sequence of finite groups. Then we can always find a sequence of abelian subgroups  $H_i \subset G_i$  such that the unbounded rank of  $\{G_i\}$  is equal to the unbounded rank of  $\{H_i\}$ . Moreover, if  $\{G'_i \subset G_i\}$  is a sequence of subgroups of bounded index then the unbounded ranks of  $\{G'_i\}$  and  $\{G_i\}$  are equal.

For a sequence of finite abelian groups we can compute the unbounded rank in terms of the direct sum decomposition provided by the structure theorem.

**Proposition 2.8.** *Let  $\{G_i\}$  be a sequence of finite abelian groups. Suppose that for every  $i \in \mathbb{N}$  the group  $G_i$  can be generated by  $r$  elements. Consider the decomposition*

$$G_i \simeq \bigoplus_{1 \leq k \leq r} \mathbb{Z}/N_{i,k}\mathbb{Z},$$

where  $N_{i,k} | N_{i,k+1}$  for every  $k \in \{1, \dots, r-1\}$ . Then the unbounded rank of the sequence  $\{G_i\}$  is equal to

$$r - \max\{k \mid \text{the sequence } \{N_{i,k}\}, i \in \mathbb{N} \text{ is bounded as } i \rightarrow \infty\}.$$

*Proof.* Let us denote

$$k_{\max} = \max\{k \mid \text{the sequence } \{N_{i,k}\}, i \in \mathbb{N} \text{ is bounded as } i \rightarrow \infty\}.$$

Then we may take a sequence of subgroups

$$H_i = \bigoplus_{k_{\max}+1 \leq k \leq r} \mathbb{Z}/N_{i,k}\mathbb{Z} \subset G_i,$$

so the unbounded rank of  $\{G_i\}$  is at most  $r - k_{\max}$ . We denote by  $L$  a constant such that  $|G_i|/|H_i| \leq L$  for all  $i \in \mathbb{N}$ .

Suppose that the unbounded rank  $r'$  of  $\{G_i\}$  is less than  $r - k_{\max}$ . Then there exists a sequence of subgroups  $\{H'_i \subset G_i\}$  such that for every  $i \in \mathbb{N}$  the group  $H'_i$  is generated by  $r' < r - k_{\max}$  elements and the indices  $|G_i|/|H'_i|$  are bounded as  $i \rightarrow \infty$ . We have

$$|H'_i| = |H'_i/(H_i \cap H'_i)| \cdot |H_i \cap H'_i| \leq L \cdot |H_i \cap H'_i|.$$

Then we have an inequality

$$\frac{|G_i|}{|H'_i|} \geq \frac{|G_i|}{L \cdot |H_i \cap H'_i|} = \frac{|G_i|}{L|H_i|} \cdot \frac{|H_i|}{|H_i \cap H'_i|}.$$

On the other hand, since  $H_i \cap H'_i$  is generated by  $r' < r - k_{\max}$  elements, the indices  $|H_i|/|H_i \cap H'_i|$  are unbounded as  $i \rightarrow \infty$ . This contradiction shows that the unbounded rank of  $\{G_i\}$  is equal to  $r - k_{\max}$ .  $\square$

Another way to express the unbounded rank of a sequence of abelian groups is given by the corollary below.

**Corollary 2.9.** *Let  $\{G_i\}$  be a sequence of finite abelian groups. Suppose that for every  $i \in \mathbb{N}$  the group  $G_i$  can be generated by  $r$  elements. Then the unbounded rank of  $\{G_i\}$  is equal to*

$$\max\{r \mid G_i \supset (\mathbb{Z}/M_i\mathbb{Z})^r \text{ for infinitely many } i \in \mathbb{N} \text{ and } M_i \rightarrow \infty\}.$$

*Proof.* By Proposition 2.8 the unbounded rank of the sequence  $\{G_i\}$  is equal to  $r - k_{\max}$ , where

$$k_{\max} = \max\{k \mid \text{the sequence } \{N_{i,k}\}, i \in \mathbb{N} \text{ is bounded as } i \rightarrow \infty\}.$$

Consider the sequence of subgroups

$$H_i = \bigoplus_{k_{\max}+1 \leq k \leq r} \mathbb{Z}/N_{i,k}\mathbb{Z} \subset G_i.$$

Then each  $H_i$  contains a subgroup isomorphic to  $(\mathbb{Z}/M_i\mathbb{Z})^{r-k_{\max}}$  where  $M_i = N_{i,k_{\max}+1}$ . Suppose that for infinitely many  $i \in \mathbb{N}$  we can find subgroups  $H'_i \subset G_i$  such that  $H'_i \simeq (\mathbb{Z}/M'_i\mathbb{Z})^s$  for  $M_i \rightarrow \infty$ . Consider the images of  $H'_i$  under the quotient homomorphisms  $G_i \rightarrow G_i/H_i$ . Since the indices  $|G_i|/|H_i|$  are bounded by a constant  $L$  independent of  $i \in \mathbb{N}$ , the number  $s$  does not exceed  $r - k_{\max}$ .  $\square$

We deduce the following subadditivity property for the unbounded ranks.

**Lemma 2.10.** *Let  $G_i \simeq (\mathbb{Z}/N_i\mathbb{Z})^r$  be a sequence of abelian groups such that  $N_i$  tend to infinity. Consider a sequence of subgroups  $G'_i \subset G_i$  for  $i \in \mathbb{N}$  and denote the quotient groups by  $G''_i$ . Suppose that the unbounded rank of the sequence  $\{G'_i\}$  is at most  $r'$  and that the unbounded rank of the sequence  $\{G''_i\}$  is at most  $r''$ . Then the unbounded rank  $r$  of the sequence  $\{G_i\}$  is at most  $r' + r''$ .*

*Proof.* By Lemma 2.2 we may choose compatible systems of generators in  $G'_i$  and  $G_i$  for every  $i \in \mathbb{N}$ . Let us denote by

$$N'_{i,1}|N'_{i,2}| \cdots |N'_{i,r}$$

the divisors in the decomposition of  $G'_i$  given by the structure theorem. Then the quotient groups  $G''_i$  are isomorphic to

$$\bigoplus_{1 \leq j \leq r} \mathbb{Z}/\frac{N_i}{N'_{i,j}}\mathbb{Z}.$$

By assumption, the unbounded rank of the sequence  $\{G'_i\}$  is at most  $r'$ . By Proposition 2.8 we obtain

$$r - \max\{j \mid \text{the sequence } \{N_{i,j}\} \text{ is bounded as } i \rightarrow \infty\} \leq r'.$$

Also, since the unbounded rank of the sequence  $\{G''_i\}$  is at most  $r''$ , by Proposition 2.8 we obtain the inequality

$$r - \min\{j \mid \text{the sequence } \{\frac{N_i}{N'_{i,j}}\} \text{ is bounded as } i \rightarrow \infty\} \leq r''.$$

However, since  $N_i$  tend to infinity, the sequences  $\{N'_{i,j}\}$  and  $\{N_i/N'_{i,j}\}$  for a fixed  $j \in \{1, \dots, r\}$  cannot be bounded simultaneously. Therefore, adding the above inequalities we obtain  $r \leq r' + r''$ , as it was to be shown.  $\square$

**2.2. The MRC fibration.** In this subsection we briefly recall the construction of the maximal rationally connected fibration (the MRC fibration) for a compact Kähler space  $X$ . In this generality existence of the MRC fibration was established in [Cam81, Cam04]. We refer to [Cam04] for details, including the definition of the cycle space  $\mathcal{C}(X)$  for a compact complex space  $X$ . For a purely algebraic proof of this result in the case of algebraic varieties, see [KMM92].

**Definition 2.11.** A covering family of cycles on  $X$  is a subvariety  $S \subset \mathcal{C}(X)$  such that

- $S$  is a countable union of compact irreducible subvarieties;
- For  $s \in S$  a general point the cycle  $Z_s$  is irreducible and reduced;
- $X$  is a union of  $\text{Supp}(Z_s)$  for  $s \in S$ .

A covering family of cycles induces an equivalence relation  $R(S)$  on points of  $X$ . Namely, two points  $x, y \in X$  are equivalent if and only if they are contained in a connected union of a finite number of cycles parameterized by  $S$ .

**Theorem 2.12.** *Let  $X$  be a normal compact connected complex space. Let  $S \subset \mathcal{C}(X)$  be a covering family of cycles on  $X$ . Denote by  $R(S)$  the equivalence relation on  $X$  induced by  $S$ . Then there exists a meromorphic fibration  $q_S: X \dashrightarrow B_S$  such that a typical fiber of  $q_S$  is an equivalence class for  $R(S)$ .*

**Definition 2.13.** Let  $X$  be a compact complex space and let  $S$  be a family of all rational curves on  $X$ . The fibration  $f: X \dashrightarrow B$  corresponding to  $S$  by Theorem 2.12 is called the maximal rationally connected (MRC) fibration of  $X$ .

The MRC fibration is known to exist if  $X$  is a compact Kähler space (see [Cam92, Remarque 1.3]).

Obviously, if  $X$  is not covered by rational curves then  $f$  is the identity map. A crucial property of the MRC fibration is that its base  $B$  is not covered by rational curves. This statement was shown in [GHS03, Corollary 1.4] for  $X$  an algebraic variety. The same argument generalizes to the Kähler case (see, for instance, [HP15, Remark 3.2] or [PS20, Proposition 3.8]).

**Proposition 2.14.** *Let  $X$  be a compact Kähler space. Consider the MRC fibration  $f: X \dashrightarrow B$ . Then the base  $B$  is not uniruled.*

Moreover, the fibers of the MRC fibration of a compact Kähler space  $X$  are in fact projective.

**Proposition 2.15.** *Let  $X$  be a rationally connected compact Kähler space. Then  $X$  is projective.*

**2.3. Finite group actions.** We will need a well-known result (see e. g. [PS14, Lemma-Definition 3.1]) on existence of regularizations of birational  $G$ -actions.

**Proposition 2.16.** *Let  $X$  be a normal projective variety and let  $G \subset \text{Bir}(X)$  be a finite group. Then there exists a smooth projective variety  $\overline{X}$  with a regular action of  $G$  and a  $G$ -equivariant birational map*

$$\varphi: \overline{X} \dashrightarrow X.$$

*Proof.* Replacing  $X$  by an affine open subset, we may assume that the action of  $G$  on  $X$  is regular. Then by [Sum74, Theorem 3] there exists a  $G$ -equivariant projective completion  $\varphi: \overline{X} \dashrightarrow X$ . Replacing  $\overline{X}$  by a  $G$ -equivariant resolution of singularities of  $\overline{X}$  (see e. g. [BM97]), we may assume  $\overline{X}$  to be smooth.  $\square$

The following proposition was shown e. g. in [Akh95, p. 38].

**Proposition 2.17.** *Let  $G$  be a finite group acting on a variety  $X$  with a fixed point  $p \in X$ . Then the induced action of  $G$  on the tangent space  $T_p(X)$  is faithful.*

### 3. MAIN RESULTS

**3.1. Groups of pseudoautomorphisms.** We begin this section by a straightforward generalization of Theorem 1.4 to automorphism groups of singular compact Kähler spaces. First of all, we prove the following simple lemma (cf. [Ue75, Lemma 9.11]).

**Lemma 3.1.** *Let  $X$  be a normal compact Kähler space. Suppose that there exists a bimeromorphic morphism*

$$\varphi: T \rightarrow X$$

*where  $T$  is a compact complex torus. Then  $\varphi$  is an isomorphism.*

*Proof.* Let  $E$  be an irreducible component of the exceptional locus of  $\varphi$  of dimension  $c > 0$ . Take a Kähler class  $\omega$  on  $X$ . By the projection formula we have

$$(\varphi^*\omega)^c \cdot E = \omega^c \cdot (\varphi_*E) = 0.$$

On the other hand, we can choose a general translation  $\tau: T \rightarrow T$  such that the image of  $\tau^*(E)$  under the map  $\varphi$  is not contained in the singular locus of  $X$ . Therefore we obtain

$$(\varphi^*\omega)^c \cdot \tau^*E = \omega^c \cdot (\varphi_*\tau^*E) > 0.$$

However, since  $\tau$  is an automorphism of  $T$ , we must have  $(\varphi^*\omega)^c \cdot E = (\varphi^*\omega)^c \cdot (\tau^*E)$ . This contradiction shows that the exceptional locus of  $\varphi$  is empty, so  $\varphi$  is an isomorphism.  $\square$

We also state another theorem by Mundet i Riera, see [Mun21, Theorem 1.16].

**Theorem 3.2.** *Let  $G$  be a Lie group with finitely many connected components. For every natural number  $r$  the following properties are equivalent:*

- $G$  contains subgroups of the form  $(\mathbb{Z}/N_i\mathbb{Z})^r$  for arbitrarily large positive integers  $N_i$ ;
- $G$  contains a subgroup isomorphic to a compact real torus  $T^r$  of real dimension  $r$ .

We can deduce the following corollary from Theorem 3.2 by an argument similar to the proof of [Mun21, Theorem 1.15].

**Corollary 3.3.** *Let  $X$  be a (possibly singular) normal compact Kähler space. For every natural number  $r$  the following properties are equivalent:*

- $\text{Aut}(X)$  contains subgroups of the form  $(\mathbb{Z}/N_i\mathbb{Z})^r$  for arbitrarily large positive integers  $N_i$ ;
- $\text{Aut}(X)$  contains a subgroup isomorphic to  $T^r$ .

Moreover,  $r \leq 2 \dim(X)$  and if  $r = 2 \dim(X)$  then  $X$  is biholomorphic to a compact complex torus.

*Proof.* By [Kim18, Lemma 3.1] the quotient group  $\text{Aut}(X)/\text{Aut}^0(X)$  has bounded finite subgroups. So by Lemma 2.1 we may assume that the finite subgroups in question lie in  $\text{Aut}^0(X)$ . The group  $\text{Aut}^0(X)$  is a connected complex Lie group, see e. g. [Akh95, Theorem on p. 40]. So the first statement of the corollary follows from Theorem 3.2. Suppose that there is an effective action of  $T^r$  by holomorphic automorphisms. Then by [BM97] we can take a  $T^r$ -equivariant Kähler resolution of singularities  $\varphi: X' \rightarrow X$ . We apply Theorem 1.4 to  $X'$  and obtain that  $r \leq 2 \dim(X') = 2 \dim(X)$ . Moreover, if  $r = 2 \dim(X')$  then  $X'$  is biholomorphic to a compact complex torus. By Lemma 3.1 we obtain that  $\varphi$  is an isomorphism, so  $X$  is nonsingular and biholomorphic to a compact complex torus.  $\square$

The next step is to extend the above result to groups of pseudoautomorphisms of singular compact Kähler spaces. Recall that a bimeromorphic map  $f: X \dashrightarrow X$  is a pseudoautomorphism if both  $f$  and  $f^{-1}$  do not contract divisors. The group of pseudoautomorphisms of  $X$  is denoted by  $\text{PsAut}(X)$ .

The following proposition was proved in [Gol21, Corollary 4.6].

**Proposition 3.4.** *Let  $X$  be a compact Kähler space with rational singularities. Let  $f: X \dashrightarrow X$  be a pseudoautomorphism. Suppose that there exists a Kähler class  $\omega$  such that  $f_*\omega$  is also a Kähler class. Then  $f$  is a biholomorphic automorphism of  $X$ .*

Now we apply this proposition to generalize the statement of Corollary 3.3 to the group  $\text{PsAut}(X)$ .

**Theorem 3.5.** *Let  $X$  be a compact Kähler space with rational singularities. Suppose that there exists  $r \in \mathbb{N}$  such that the group  $\text{PsAut}(X)$  contains a subgroup isomorphic to  $(\mathbb{Z}/N_i\mathbb{Z})^r$  for arbitrarily large  $N_i$ . Then  $r \leq 2 \dim(X)$  and  $\text{PsAut}(X)$  contains a subgroup isomorphic to a compact complex torus of dimension  $r$ . Moreover, if  $r = 2 \dim(X)$  then  $X$  is biholomorphic to a compact complex torus.*

*Proof.* Consider the action of  $\text{PsAut}(X)$  on  $H^{1,1}(X, \mathbb{Q})$  by pushforward. We obtain an exact sequence of groups

$$1 \rightarrow \text{PsAut}(X)_\tau \rightarrow \text{PsAut}(X) \rightarrow \text{PsAut}(X)/\text{PsAut}(X)_\tau \rightarrow 1,$$

where we denote  $\text{PsAut}(X)_\tau = \{f \in \text{PsAut}(X) \mid f_*|_{H^{1,1}(X, \mathbb{Q})} = \text{Id}\}$ . Note that the quotient group  $\text{PsAut}(X)/\text{PsAut}(X)_\tau$  embeds into  $\text{GL}(H^{1,1}(X, \mathbb{Q}))$ , therefore, by Minkowski's theorem (see e.g. [Ser07, Theorem 1]), the orders of finite subgroups of  $\text{PsAut}(X)/\text{PsAut}(X)_\tau$  are bounded by a constant  $M(X)$  depending on  $h^{1,1}(X, \mathbb{Q})$  only. Therefore, the group  $\text{PsAut}(X)_\tau$  contains finite abelian subgroups generated by at least  $r$  elements of arbitrarily large orders; moreover, by Lemma 2.1 we may assume that these subgroups are of the form  $(\mathbb{Z}/N\mathbb{Z})^r$ , where  $N$  go to infinity. The group  $\text{PsAut}(X)_\tau$  acts trivially on  $H^{1,1}(X, \mathbb{R}) = H^{1,1}(X, \mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{R}$  and, in particular, preserves every Kähler class on  $X$ . Thus by Proposition 3.4 the group  $\text{PsAut}(X)_\tau$  is contained in  $\text{Aut}(X)$ . The conclusion of the theorem now follows from Corollary 3.3.  $\square$

**3.2. Non-uniruled varieties.** In this subsection we derive a slightly more precise version of Theorem 1.2 from Theorem 3.5.

**Definition 3.6.** A compact Kähler space (or a projective variety)  $X$  with terminal  $\mathbb{Q}$ -factorial singularities is called

- *minimal* (or a *minimal model*) if the canonical class  $K_X$  is nef;
- *quasi-minimal* (or a *quasi-minimal model*) if  $K_X$  is modified nef.

Note that a minimal model is also quasi-minimal. Existence of quasi-minimal models for non-uniruled projective varieties was shown in [PS14, Lemma 4.4].

**Proposition 3.7.** *Let  $X$  be a non-uniruled projective variety. Then there exists a quasi-minimal model of  $X$ , that is, a quasi-minimal variety  $X'$  birational to  $X$ .*

In the case of non-uniruled compact Kähler spaces of dimension 3, minimal models exist by [HP16, Theorem 1.1].

**Theorem 3.8.** *Let  $X$  be a compact Kähler space of dimension 3. Then there exists a minimal compact Kähler space  $X'$  bimeromorphic to  $X$ .*

The reason why we need quasi-minimal models is the following description of their bimeromorphic (or birational) automorphisms. The case when  $X$  is a projective variety was settled in [PS14, Corollary 4.7]; for the general case of compact Kähler spaces see [Gol21, Proposition 4.2].

**Proposition 3.9.** *Let  $X$  be a quasi-minimal compact Kähler space. Let  $f: X \dashrightarrow X$  be a bimeromorphic map. Then  $f$  is a pseudoautomorphism.*

**Theorem 3.10.** *Let  $X$  be a non-uniruled projective variety. Suppose that there exists  $r \in \mathbb{N}$  such that the group  $\text{Bir}(X)$  contains a subgroup isomorphic to  $(\mathbb{Z}/N_i\mathbb{Z})^r$  where  $N_i$  tend to infinity. Then  $r \leq 2 \dim(X)$ , and the group  $\text{Bir}(X)$  contains a subgroup isomorphic to an abelian variety of dimension  $\lceil r/2 \rceil$ . Moreover, if  $r = 2 \dim(X)$  then  $X$  is birational to an abelian variety.*

*Proof.* Since  $X$  is non-uniruled, by Proposition 3.7 there exists a quasi-minimal projective variety  $X'$  birational to  $X$ . By Proposition 3.9, we have  $\text{Bir}(X) \simeq \text{Bir}(X') = \text{PsAut}(X')$ . Then the result follows from Theorem 3.5.  $\square$

**Proposition 3.11.** *Let  $X$  be a non-uniruled compact Kähler space of dimension 3. Suppose that there exists  $r \in \mathbb{N}$  such that the group  $\text{Bim}(X)$  contains a subgroup isomorphic to  $(\mathbb{Z}/N_i\mathbb{Z})^r$  where  $N_i$  tend to infinity. Then  $r \leq 6$ , and the group  $\text{Bim}(X)$  contains a subgroup isomorphic to a compact complex torus of dimension  $\lceil r/2 \rceil$ . Moreover, if  $r = 6$  then  $X$  is bimeromorphic to a compact complex torus.*

*Proof.* By Theorem 3.8 there exists a minimal compact Kähler space  $X'$  bimeromorphic to  $X$ . By Proposition 3.9 we have  $\text{Bim}(X) \simeq \text{Bim}(X') = \text{PsAut}(X')$ . The statement now follows from Theorem 3.5.  $\square$

### 3.3. Rationally connected varieties.

**Proposition 3.12.** *Let  $X$  be a rationally connected variety of dimension  $n$  over an algebraically closed field  $k$  of zero characteristic. Then there exists a constant  $J(n)$ , depending on  $n$  only, such that for any finite subgroup  $G \subseteq \text{Aut}(X)$  there exists a subgroup  $H \subseteq G$  of index at most  $J(n)$  acting on  $X$  with a fixed point.*

*Proof.* The statement follows from [PS16, Theorem 4.2] and [Bir21, Theorem 1.1].  $\square$

**Corollary 3.13.** *Let  $X$  be a rationally connected variety of dimension  $n$  over a field  $k$  of zero characteristic. There exists a constant  $M = M(n)$  such that for any finite subgroup  $G \subset \text{Bir}(X)$  there exists an abelian subgroup  $H \subset G$  of index at most  $M(n)$  and such that  $H$  is generated by at most  $n$  elements.*

*Proof.* We pass to the algebraic closure  $\bar{k}$  of  $k$  and replace  $X$  by  $X \times_k \bar{k}$ . Let  $\varphi: X' \dashrightarrow X$  be a smooth birational regularization of the action of  $G$ , which exists by Proposition 2.16. Note that  $X'$  is rationally connected. Therefore by Proposition 3.12 there exists a constant  $J'(n)$  such that  $G$  contains a subgroup  $H$  of index at most  $J'(n)$  acting on  $X'$  with a fixed point. By Proposition 2.17 the group  $H$  embeds into  $\text{GL}_n(\bar{k})$ . Therefore by Jordan's theorem (see [Jor78] or [Rob90, Theorem A]) the group has an abelian subgroup  $A \subset H$  of index at most  $J''(n)$ . Then  $A \subset G$  is a subgroup of index at most

$$M(n) = J'(n) \cdot J''(n),$$

moreover,  $A$  is generated by at most  $n$  elements.  $\square$

**Theorem 3.14.** *Let  $X$  be a rationally connected variety of dimension  $n$  over a field  $k$  of zero characteristic. Suppose that there exists  $r \in \mathbb{N}$  such that for arbitrarily large  $N_i \in \mathbb{N}$  the group  $\text{Bir}(X)$  contains a subgroup  $G_i \simeq (\mathbb{Z}/N_i\mathbb{Z})^r$ . Then  $r \leq n$ .*

*Proof.* By Proposition 3.12 for every  $G_N$  we can find a subgroup  $H_N \subseteq G_N$  of index at most  $J(n)$  such that  $H_N$  acts on  $X$  with a fixed point. Moreover, by Lemma 2.1 we may replace each  $H_N$  for  $N > J(n)$  by a subgroup  $H_{N'}$  isomorphic to  $(\mathbb{Z}/N'\mathbb{Z})^r$ , where  $N' \geq N/J(n)$ . Then by Proposition 2.17 the induced action of  $H_{N'}$  on the tangent space is faithful. Since all subgroups  $H_{N'}$  map injectively to  $\text{GL}_n(\bar{k})$ , we obtain that  $r \leq n$ .  $\square$

**3.4. The general case.** Now we can prove Theorems 1.5 and 1.6.

*Proof of Theorem 1.5.* If  $X$  is not uniruled then the result follows from Theorem 3.10. Suppose that  $X$  is uniruled and consider its MRC fibration  $f: X \dashrightarrow B$  where  $\dim(B) < \dim(X)$ . Then for every  $i \in \mathbb{N}$  we obtain an exact sequence

$$1 \rightarrow G'_i \rightarrow G_i \rightarrow G''_i \rightarrow 1$$

where the action of  $G'_i$  is fiberwise with respect to  $f$  (that is, every element  $g \in G'_i$  maps a point in a fiber of  $f$  where  $g$  is defined to a point in the same fiber) and  $G''_i$  acts faithfully on the base  $B$ . Let  $X_\eta$  be the scheme-theoretic generic fiber of  $f$ . Then for every  $i \in \mathbb{N}$  we have  $G'_i \subset \text{Bir}(X_\eta)$ . We denote  $n' = \dim(X_\eta)$ . Then by Theorem 3.14 the unbounded rank of the sequence  $\{G'_i\}$  is at most  $n'$ . Since  $B$  is not uniruled by Proposition 2.14, we apply Theorem 3.10 and obtain that the unbounded rank of the sequence  $\{G''_i\}$  is at most  $2 \dim(B)$ . Therefore by Lemma 2.10 the unbounded rank  $r$  of the sequence  $\{G_i\}$  is at most  $n' + 2 \dim(B)$ . In particular, one has  $r \leq 2 \dim(B) + n' < 2n$ , as it was to be shown.  $\square$

*Proof of Theorem 1.6.* If  $X$  is not uniruled then the result follows from Proposition 3.11.

Suppose that  $X$  is uniruled. Then we consider the MRC fibration  $f: X \dashrightarrow B$  with  $B$  non-uniruled and  $\dim(B) < \dim(X)$ . Then for every  $i \in \mathbb{N}$  there exists an exact sequence of groups

$$1 \rightarrow G'_i \rightarrow G_i \rightarrow G''_i \rightarrow 1,$$

where the action of  $G'_i$  is fiberwise with respect to  $f$  and  $G''_i$  acts faithfully on  $B$ . Let  $X_t$  be a very typical fiber of  $f$ , that is,  $t$  is a point in a complement of at most countable union of proper analytic subsets of  $B$ . Since the set of finite groups  $\{G_i\}$  is countable, by [PS21, Lemma 3.1] we may assume that

$$G'_i \subset \text{Bim}(X_t)$$

for a very typical fiber  $X_t$ . Note that by Proposition 2.15 the fibers of  $f$  are projective. Then by Theorem 3.14 the unbounded rank of the sequence  $\{G'_i\}$  is at most  $\dim(X_t)$ . Moreover, by Theorem 3.10 the unbounded rank of the sequence  $\{G''_i\}$  is at most  $2 \dim(B)$ . By Lemma 2.10 we obtain that the unbounded rank of the sequence  $\{G_i\}$  is at most  $2 \dim(B) + \dim(X_t)$ ; in particular, one has

$$r \leq \dim(X_t) + 2 \dim(B) < 6.$$

$\square$

## REFERENCES

- [Akh95] D. N. Akhiezer. Lie group actions in complex analysis. *Aspects of Mathematics, E27*. Friedr. Vieweg & Sohn, Braunschweig, 1995. viii+201 pp.
- [Art91] M. Artin. Algebra. Prentice Hall, Inc., Englewood Cliffs, NJ, 1991. xviii+618 pp.
- [Bir21] C. Birkar. Singularities of linear systems and boundedness of Fano varieties. *Ann. of Math. (2)* 193 (2021), no. 2, 347–405.
- [BM97] E. Bierstone, P. Milman. Canonical desingularization in characteristic zero by blowing up the maximum strata of a local invariant. *Invent. Math.* 128, 207–302 (1997).
- [Cam81] F. Campana. Coréduction algébrique d'un espace analytique faiblement Kählérien compact. *Invent. Math.*, 63 (1981), 187–223.
- [Cam92] F. Campana. Connexité rationnelle des variétés de Fano. *Annales scientifiques de l'École Normale Supérieure, Série 4, Tome 25* (1992) no. 5, 539–545.
- [Cam04] F. Campana. Orbifolds, special varieties and classification theory: an appendix. *Annales de l'Institut Fourier, Tome 54* (2004) no. 3, 631–665.
- [Gol21] A. Golota. Jordan property for groups of bimeromorphic automorphisms of compact Kähler threefolds. arXiv:2112.02673.
- [GHS03] T. Graber, J. Harris, J. Starr. Families of rationally connected varieties. *J. Amer. Math. Soc.* 16 (2003), no. 1, 57–67.
- [HP15] A. Höring, T. Peternell. Mori fibre spaces for Kähler threefolds. *J. Math. Sci. Univ. Tokyo*, 22:1–28, 2015.
- [HP16] A. Höring, T. Peternell. Minimal models for Kähler threefolds. *Invent. Math.* 203 (2016), no. 1, 217–264.
- [Jor78] C. Jordan. Mémoire sur les équations différentielles linéaires à intégrale algébrique. *Journal für die reine und angewandte Mathematik (Crelles Journal)*, vol. 1878, no. 84, 1878, pp. 89–215.
- [Kim18] J. H. Kim. Jordan property and automorphism groups of normal compact Kähler varieties. *Commun. Contemp. Math.* 20 (2018), no. 3, 1750024, 9 pp.
- [KMM92] J. Kollar, Y. Miyaoka, S. Mori. Rationally connected varieties. *J. Alg. Geom.* 1, 1992, 429–448.
- [MZ18] S. Meng, D.-Q. Zhang. Jordan property for non-linear algebraic groups and projective varieties. *Amer. J. Math.* 140 (2018), no. 4, 1133–1145.
- [Mun21] I. Mundet i Riera. Topological rigidity of tori and finite group actions on hypertoral manifolds. arXiv:2112.05599.

- [PS14] Yu. Prokhorov, C. Shramov. Jordan property for groups of birational selfmaps. *Compositio Mathematica*, 150(12):2054–2072, 2014.
- [PS16] Yu. Prokhorov, C. Shramov. Jordan property for Cremona groups. *Amer. J. Math.*, 138(2):403–418, 2016.
- [PS20] Yu. Prokhorov, C. Shramov. Finite groups of bimeromorphic selfmaps of uniruled Kähler threefolds. *Izv. Ross. Akad. Nauk Ser. Mat.*, 84(5):169–196, 2020.
- [PS21] Yu. Prokhorov, C. Shramov. Finite groups of bimeromorphic selfmaps of non-uniruled Kähler threefolds. arXiv:2110.05825.
- [Rob90] G. Robinson. On linear groups. *J. Algebra* 131 (1990), no. 2, 527–534.
- [Ser07] J.-P. Serre. Bounds for the orders of the finite subgroups of  $G(k)$ . In *Group representation theory*, pages 405–450. EPFL Press, Lausanne, 2007.
- [Sum74] H. Sumihiro. Equivariant completion. *J. Math. Kyoto Univ.* 14-1 (1974) 1–28.
- [Ue75] K. Ueno. Classification theory of algebraic varieties and compact complex spaces. Notes written in collaboration with P. Cherenack. *Lecture Notes in Mathematics*, Vol. 439. Springer-Verlag, Berlin-New York, 1975. xix+278 pp.
- [Xu18] J. Xu. Finite  $p$ -groups of birational automorphisms and characterizations of rational varieties. arXiv:1809.09506.

National Research University Higher School of Economics, Russian Federation  
Laboratory of Mirror Symmetry, NRU HSE, 6 Usacheva str., Moscow, Russia, 119048  
golota.g.a.s@mail.ru, agolota@hse.ru