

GRAPHING, HOMOTOPY GROUPS OF SPHERES, AND SPACES OF LINKS AND KNOTS

ROBIN KOYTCHEFF

ABSTRACT. We study homotopy groups of spaces of links, focusing on long links of codimension at least three. In the case of multiple components, they admit split injections from homotopy groups of spheres. We calculate them, up to knotting, in a range depending on the dimensions of the source manifolds and target manifold which roughly generalizes the triple-point-free range for isotopy classes. At the edge of this range, joining components sends both a parametrized long Borromean rings class and a Hopf fibration to a generator of the first nontrivial homotopy group of the space of long knots. For spaces of two-component long links, we give generators of the homotopy group in this dimension in terms of this class from the Borromean rings and homotopy groups of spheres. A key ingredient in most of our results is a graphing map that increases source and target dimensions by one.

CONTENTS

1. Introduction	2
1.1. Previous related results	2
1.2. Main results and organization	2
1.3. Acknowledgments	5
2. Key definitions	6
2.1. Spaces of embeddings, link maps, and pseudoisotopy embeddings	6
2.2. Maps between spaces of links	7
3. Injectivity of graphing	10
3.1. Injectivity for spaces of embeddings via the λ -invariant	10
3.2. Injectivity for spaces of link maps in a range via the α -invariant	13
4. Bijectivity of graphing in a range	15
4.1. Lemmas on pseudoisotopy embedding spaces	15
4.2. Bijectivity of graphing and homotopy groups of spaces of long links in a range	17
4.3. Spherical links and a further calculation for long links	20
5. Homotopy classes in spaces of long knots and links from joining pure braids	22
5.1. Configuration space integrals for cohomology of spaces of long 1-dimensional links	22
5.2. Some low-dimensional graph cocycles	24
5.3. Homology classes dual to configuration space integrals	26
5.4. Long Borromean rings, the Hopf map, and classes of long links and knots	27
5.5. Questions and future directions	34
References	35

2020 *Mathematics Subject Classification.* Primary: 57K45, 55Q40, 57R40, 55R80; Secondary: 58D10, 81Q30, 55P35.

Key words and phrases. Spaces of embeddings, long knots, long links, pure braids, graphing, homotopy groups of spheres, configuration spaces, the λ -invariant, the α -invariant, bordism, pseudoisotopy embedding spaces, configuration space integrals, immersions.

1. INTRODUCTION

This paper concerns homotopy groups of spaces of links of various dimensions, where the source and target manifolds are either Euclidean spaces or spheres. We focus mainly, though not exclusively, on spaces $\mathcal{L}_{p,q}^n = \text{Emb}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n)$ of 2-component long links, meaning smooth embeddings $\mathbb{R}^p \sqcup \mathbb{R}^q \hookrightarrow \mathbb{R}^n$ with fixed compact support. Usually we will assume $p, q \leq n - 3$. Most of our results use a graphing map that increases both source and target dimensions by one. Our main results are the following:

- Theorems [A](#) and [B](#) concern injectivity on homotopy groups of graphing maps from spheres.
- Theorems [C](#), [D](#), and [E](#) calculate, modulo knotting, homotopy groups of spaces of long links, in roughly a “metastable” or “triple-point-free” range, namely up to degree $2n - 3p - 3$ if $p = q$. Most of the classes in these groups come from homotopy groups of spheres.
- Theorem [F](#) gives explicit generators of homotopy groups of spaces of long links with $p = q$ and long knots in the above-mentioned degree. It also establishes that a certain homotopy group of spheres gives rise to nontrivial classes in spaces of long knots.

Each set of results is proven using a different method. Before describing them in more detail, we survey some earlier related work.

1.1. Previous related results. In all of the following previous results, the codimensions of the embeddings are assumed to be at least 3. Haefliger [[Hae67](#)] determined the group $\pi_0 \text{Emb}(S^p \sqcup S^q, S^n)$ of isotopy classes of spherical links in a range of values of p, q , and n . This range was improved slightly by M. Skopenkov [[Sko09](#)]. Their result applies in the 2-metastable range. This is roughly the quadruple-point-free range, meaning that if $p = q$, then roughly $p < 3n/4$. More precisely their range is $3n - 2p - 2q \geq 6$. Crowley, Ferry, and M. Skopenkov [[CFS14](#)] computed rational isotopy classes of spherical links with an arbitrary number of components. Songhafouo Tsopméné and Turchin [[STT18b](#), Theorem 3.2] described rational isotopy classes of long links in terms of trivalent trees. They conjectured an extension to all rational homotopy groups of spaces of long links in terms of the homology of a graph complex, which was proven by Fresse, Turchin, and Willwacher [[FTW](#), Section 5.5].

Haefliger [[Hae66](#)] showed that the group $\pi_0 \text{Emb}(S^p, S^n)$ of isotopy classes of knots $S^p \hookrightarrow S^n$ is trivial in the metastable range, which is roughly the triple-point-free range $p < 2n/3$. More precisely, there is no knotting if $2n - 3p \geq 4$. He also computed this group for the first family of dimensions (p, n) where they are nontrivial, namely $2n - 3p = 3$. Budney [[Bud08](#)] generalized these results on knots from isotopy classes to the i -th homotopy groups of spaces $\text{Emb}_c(\mathbb{R}^p, \mathbb{R}^n)$ of long knots, i.e., supported embeddings $\mathbb{R}^p \hookrightarrow \mathbb{R}^n$ with fixed compact support. He used iterates of graphing maps which increase the dimensions of both source and target manifolds by one. The range of dimensions where his results apply, $0 \leq i \leq 2n - 3p - 3$, generalizes the range for Haefliger’s results from isotopy classes to families of knots.

1.2. Main results and organization. Most of our results rely on the same iterated graphing construction, adapted easily from long knots to long links. To get nontrivial classes for knots from iterated graphing, one must start with at least 1-dimensional knots, whereas in our setting of links, we can start with links where one or both components have dimension 0. The space of such links has a subspace homeomorphic to a sphere, and most of our results involve classes from homotopy groups of spheres. Though some of our results hold in codimension less than 3, they say nothing new in these cases.

Our first result concerns nontrivial homotopy classes in spaces of 2-component links from homotopy groups of spheres, with not much restriction on the dimensions involved.

Theorem A. *If $0 \leq p \leq q \leq n - 1$ and $i \geq 0$, then $\pi_i \text{Emb}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n)$ contains a direct summand $\pi_{i+p} S^{n-q-1}$, the inclusion of which can be realized by graphing.*

Theorem A appears in the main body as Theorem 3.1. Theorem 3.4 is the analogue of it for spherical links in a sphere. The proof is not difficult, and putting $i = 0$ gives a result known at least as far back as work of Haefliger [Hae62]. The case $i = 0$ and $p + q = n - 1$ corresponds to classes dual to the generalized linking number, including 2-strand pure braids when $p = q = 1$ and $n = 3$. See Example 3.5. Theorem A yields many nontrivial torsion classes in spaces of links. Applying it in some low-dimensional examples, one obtains a split injection of $\pi_4 S^2 \cong \mathbb{Z}/2$ into $\pi_3 \text{Emb}_c(\mathbb{R} \sqcup \mathbb{R}, \mathbb{R}^4)$, as well as into $\pi_3 \text{Emb}_c(\mathbb{R} \sqcup \mathbb{R}^2, \mathbb{R}^5)$. For more details on the graphing map, see Definition 2.4 or the statement of Theorem C below.

The next result is an analogue of the previous one for link maps rather than embeddings. Recall that a link map of one manifold into another is a smooth map such that the images of the components are pairwise disjoint. We write $\text{Link}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n)$ for the space of long links maps, i.e., link maps $\mathbb{R}^p \sqcup \mathbb{R}^q \rightarrow \mathbb{R}^n$ with fixed compact support.

Theorem B. *If $1 \leq p \leq q \leq n - 3$ and $0 \leq i \leq 2n - p - 2q - 4$, then $\pi_i \text{Link}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n)$ contains a direct summand $\pi_{i+p} S^{n-q-1}$, the inclusion of which can be realized by graphing.*

The condition on i is precisely that $\pi_{i+p} S^{n-q-1}$ lies in the stable range of the Freudenthal suspension theorem. Theorem B is the special case of Theorem 3.7 applied to this stable range. The statement is also true for based spherical link maps in Euclidean space; see Remark 3.8. Putting $i = 0$ gives a result of Scott [Sco68], which was improved upon by Massey and Rolfsen [MR85], Koschorke [Kos90], and Habegger and Kaiser [HK98].

The proofs of Theorems A and B and their related results are fairly elementary, using respectively the λ -invariant (or an analogue of it) and the α -invariant, as well as the Pontryagin–Thom theorem. The λ -invariant comes from the topology of an unknot complement, while the α -invariant is akin to a Gauss map. See Definitions 3.2 and 3.6 for more details.

Our third main result is a mutual extension to $\pi_i \text{Emb}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n)$ of both Budney’s result [Bud08, Proposition 3.9] on $\pi_i \text{Emb}_c(\mathbb{R}^p, \mathbb{R}^n)$ and the result of Haefliger [Hae67] and M. Skopenkov [Sko09] on $\pi_0 \text{Emb}(S^p \sqcup S^q, \mathbb{R}^n)$. Indeed, Lemma 4.8 identifies spherical isotopy classes with long isotopy classes. The next theorem applies in a range that generalizes the triple-point-free range for isotopy classes to i -parameter families. There are however many nontrivial groups for links in this range, in contrast to the setting of knots. For the remaining results, we abbreviate $\mathcal{L}_{p,q}^n := \text{Emb}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n)$ and $\mathcal{K}_p^n := \text{Emb}_c(\mathbb{R}^p, \mathbb{R}^n)$. Theorem C is a slightly condensed version of Theorem 4.3.

Theorem C. *If $1 \leq p \leq q \leq n - 3$ and $i \leq 2n - p - 2q - 4$, then*

$$\pi_i \mathcal{L}_{p,q}^n \cong \pi_{i+p} S^{n-q-1} \oplus \pi_0 \mathcal{K}_{i+q}^{i+n}.$$

The inclusion of the first summand can be realized as a map $\pi_{i+p} S^{n-q-1} \rightarrow \pi_{i+p} \mathcal{L}_{0,q-p}^{n-p}$ followed by a composition of the first p graphing maps below, while the projection onto the second summand can be realized as a composition of the last i graphing maps below followed by restriction to the second component $\pi_0 \mathcal{L}_{i+p,i+q}^{i+n} \rightarrow \pi_0 \mathcal{K}_{i+q}^{i+n}$:

$$\pi_{i+p} \mathcal{L}_{0,q-p}^{n-p} \xrightarrow{G_*} \pi_{i+p-1} \mathcal{L}_{1,q-p+1}^{n-p+1} \xrightarrow{G_*} \dots \xrightarrow{G_*} \pi_i \mathcal{L}_{p,q}^n \xrightarrow{G_*} \dots \xrightarrow{G_*} \pi_0 \mathcal{L}_{i+p,i+q}^{i+n}$$

In this sequence, each map is an isomorphism, except possibly the first, which is always surjective.

If $n - q \geq 2$, then we can identify $\pi_0 \mathcal{K}_{i+q}^{i+n}$ with isotopy classes of spherical knots in the sphere or Euclidean space. This group vanishes if $2n - 3q \geq 4$, hence it vanishes in the assumed range if $p = q$. It is also known if $2n - 3q$ is 3 or 2, by work of Haefliger [Hae66] and Milgram [Mil72]. That work of Milgram furthermore identifies the 2-primary part of this group for some smaller values of this quantity.

See also the exposition by A. Skopenkov [Skob]. The proof of Theorem C uses connectivity results on pseudoisotopy embedding spaces due to Goodwillie, much like the result of Budney's that it generalizes.

The statements of Theorems A, B, and C, as well as their proofs, become slightly simpler when $p = q$. In that case, the space $\mathcal{L}_{0,q-p}^{n-p}$ from which we start the iteration of graphing maps in each theorem is just the configuration space of two points in \mathbb{R}^{n-p} . With $p = q$ in Theorem C, in addition to the vanishing of the group of knots, one gets that every graphing map G_* shown is an isomorphism; see Theorem 4.3.

Theorem D appears as Theorem 4.6 in the main body, and its proof is similar to that of Theorem C. It says that in a range similar to the one in Theorem C, all elements of those homotopy groups come from links with at most 2 components.

Theorem D. *Suppose that $1 \leq p_1 \leq \dots \leq p_m \leq n-3$ and $0 \leq i \leq 2n-2p_{m-1}-p_m-4$. Then every class in $\pi_i \mathcal{L}_{p_1, \dots, p_m}^n$ is in the image of $(\iota_S)_*$ for some $S \subset \{1, \dots, m\}$ with $|S| \leq 2$, where ι_S is the inclusion of the space of links with components labeled by S .*

In the slightly more restrictive range $i \leq 2n-p_{m-1}-2p_m-4$, we can thus describe all classes in $\pi_i \mathcal{L}_{p_1, \dots, p_m}^n$ as in Theorem C.

Methods like those used to prove Theorem C yield a calculation of homotopy groups of equidimensional 2-component links of dimension at least 2 in the degree where graphing from spheres ceases to be surjective. Theorem E appears as 4.10 in the main body.

Theorem E. *Suppose $1 \leq p \leq n-3$, and consider the sequence of maps induced by graphing:*

$$\pi_{2n-3p-3} \mathcal{L}_{p,p}^n \rightarrow \pi_{2n-3p-4} \mathcal{L}_{p+1,p+1}^{n+1} \rightarrow \dots \rightarrow \pi_0 \mathcal{L}_{2n-2p-3, 2n-2p-3}^{3n-3p-3}$$

If $p \geq 2$, then all the maps are isomorphisms. If $p = 1$, then the first map is surjective, and the remaining maps are isomorphisms. If $p \geq 2$, these groups are isomorphic to

$$\begin{aligned} \mathbb{Z}^3 \oplus \pi_{2n-2p-3} S^{n-p-1} & \quad \text{if } n-p \text{ is odd} \\ (\mathbb{Z}/2)^3 \oplus \pi_{2n-2p-3} S^{n-p-1} & \quad \text{if } n-p \text{ is even} \end{aligned}$$

Conjecture 5.7 concerns the left-hand group for $p = 1$, that is, $\pi_{2n-6} \mathcal{L}_{1,1}^n$. It is known rationally [STT18a], and we expect that all the torsion comes from the graphing map from a sphere.

Our last main result concerns equidimensional long links. It connects spaces of 2- and 3-component pure braids to spaces of long knots and 2-component long links, in the homotopy group just outside the ranges of Theorems C and D, i.e., the group $\pi_{2n-3p-3}$, which appeared in Theorem E.

It is well known for (spherical) 1-dimensional links in \mathbb{R}^3 that one can obtain the trefoil knot by joining all three components of the Borromean rings. One can obtain the Whitehead link by joining just two of the three components. A generalization of these facts to isotopy classes of higher-dimensional (spherical) links and knots is also known [Skoa]. Theorem F generalizes this fact to higher homotopy groups. The classes in it can be viewed as analogues of the trefoil, Borromean rings, and Whitehead link, though there is also a class coming from Hopf fibrations or Whitehead squares which has no analogue for links in \mathbb{R}^3 . Theorem F appears as Theorem 5.3 in the main body of the paper.

Theorem F. *Suppose $1 \leq p \leq n-3$ and $2n-3p-3 \geq 0$.*

- (a) *If $n-p$ is odd, then the map $\pi_{2n-2p-3} S^{n-p-1} \rightarrow \pi_{2n-3p-3} \mathcal{K}_p^n$ given by p -fold graphing followed by joining the two link components is nontrivial. It sends each Hopf fibration to a generator. More generally, it sends $[\mathbb{1}_{n-p-1}, \mathbb{1}_{n-p-1}]$, the Whitehead square of the identity, to twice a generator.*
- (b) *The map $\pi_{2n-3p-3} \Omega^p \text{Conf}(3, \mathbb{R}^{n-p}) \rightarrow \pi_{2n-3p-3} \mathcal{K}_p^n$ given by p -fold graphing followed by joining the three components together maps the “parametrized long Borromean rings” class $[b_{21}, b_{31}]$ to a generator.*

- (c) *There is a basis for $\pi_{2n-6}\mathcal{L}_{1,1}^n$ modulo torsion consisting of*
- *the images of a generator of $\pi_{2n-6}\mathcal{K}_1^n$ under the inclusions $\iota_1, \iota_2 : \mathcal{K}_1^n \rightarrow \mathcal{L}_{1,1}^n$,*
 - *the result of joining two components of $[b_{21}, b_{31}]$, and*
 - *for $n - p$ odd, the result of graphing $[\mathbb{1}_{n-2}, \mathbb{1}_{n-2}]$.*

If $p \geq 2$, then there is a minimal generating set for $\pi_{2n-3p-3}\mathcal{L}_{p,p}^n$ given by the two inclusions of a generator of $\pi_{2n-3p-3}\mathcal{K}_p^n$, the result of joining two components of $[b_{21}, b_{31}]$, and the image under graphing of a minimal generating set of $\pi_{2n-2p-3}S^{n-p-1}$.

By the connectivity result of Budney [Bud08], $2n - 3p - 3$ is the lowest dimension where such a result is possible for spaces of long knots. It is also the lowest dimension in which a class detected by the third stage of the Goodwillie–Weiss Taylor tower can appear. Equivalently, it is the lowest dimension of classes dual to analogues of type-2 Vassiliev invariants. Conjecture 5.8 is that classes in this dimension in a space of long links with any number of components must come from the classes described in Theorem F. It is true rationally [FTW].

Theorem F is proven via cohomology classes from configuration space integrals and dual homology classes obtained by resolving singular links. Specifically, we represent all the homotopy classes appearing in it by such resolutions or their images under graphing. It hints at the possibility of obtaining arbitrary classes of long links from pure braids, generalizing known results on obtaining isotopy classes and n -equivalence classes of knots via pure braids.

The paper is organized as follows. Section 2 sets the main definitions and notation. We define spaces of embeddings and link maps, configuration spaces, and pseudoisotopy embedding spaces in Section 2.1. We define graphing and closure maps, as well as maps that join components in Section 2.2. Readers familiar with the subject matter may wish to initially skip Section 2 and refer back to it as needed.

The remaining Sections are essentially independent of each other and can be read in any order.

In Section 3, we prove results about the injectivity of homotopy groups of spheres into homotopy groups of spaces of 2-component links. Section 3.1 contains Theorem 3.1 (Theorem A) on long embeddings and Theorem 3.4, which is a spherical analogue of it. Both use a map called the λ -invariant or an analogue of it. In Section 3.2, we prove Theorem 3.7 (and hence Theorem B) on link maps, by using a map called the α -invariant. These two Subsections are mostly independent of each other.

Section 4 contains Theorem 4.3 (Theorem C) and Theorem 4.6 (Theorem D), which determine homotopy groups of all spaces of long links, at least up to knotting, in a certain range. We first establish some lemmas about pseudoisotopy embedding spaces in Section 4.1, which are then used to prove those Theorems in Section 4.2. At the end of this Section, we prove Theorem 4.10 (Theorem E), which improves upon the calculation in Theorem D for equidimensional links of dimension at least 2.

In Section 5, we prove Theorem 5.3 (Theorem F) about nontrivial classes in spaces of long knots and links from classes in spaces of pure braids, including classes from spheres. We review configuration space integrals in Section 5.1, including some specific cohomology classes from graph cocycles in Section 5.2. In Section 5.3, we review how to construct dual homology classes by resolving singular links. In Section 5.4, we use this material to prove the Theorem. We conclude with Section 5.5, where we pose questions to explore in future work, on graphing and joining components, as well as subspaces of split links and cables.

1.3. Acknowledgments. This work was supported by the Louisiana Board of Regents Support Fund, contract number LEQSF(2019-22)-RD-A-22. We thank Ryan Budney, Danica Kosanović, and Victor Turchin for useful conversations.

2. KEY DEFINITIONS

In Section 2.1, we define a handful of spaces of smooth embeddings or smooth maps, and in Section 2.2, we define several maps involving these spaces. We begin with some basic conventions and notation.

For any $k \geq 0$, let D^k denote the closed k -dimensional unit disk in \mathbb{R}^k . Let $I := D^1 = [-1, 1]$. For a smooth manifold X with a basepoint x , we write $\Omega^k X$ for the space of smooth based k -fold loops in X . We take these to be smooth maps $\mathbb{R}^k \rightarrow X$ which are constant at x outside I^k . We may sometimes allow the role of I^k to be played by D^k , or by a product of disks whose dimensions sum to k , via a homeomorphism that is a diffeomorphism between the interiors. One can define $\pi_k X$ as $\pi_0 \Omega^k X$, and $\pi_i \Omega^j X \cong \pi_{i+j} X$ for any non-negative integers i and j . We may also represent a k -fold loop in X by a based map $S^k \rightarrow X$. We write π_i^S for the i -th stable homotopy group of spheres, i.e., $\pi_i^S := \operatorname{colim}_{k \rightarrow \infty} \pi_{k+i} S^k$.

2.1. Spaces of embeddings, link maps, and pseudoisotopy embeddings. For smooth manifolds P and N , we write $\operatorname{Emb}(P, N)$ for the space of smooth embeddings of P into N . Let P_1, \dots, P_m be the path components of P . A **link map** is a smooth map $(f_1, \dots, f_m) : \coprod_1^m P_i \rightarrow N$ such that the images of the f_i are pairwise disjoint. We denote the space of such link maps by $\operatorname{Link}(\coprod_1^m P_i, N)$. We will usually use the term **links** for embeddings $\coprod_1^m P_i \rightarrow N$, especially when $m > 1$, though we may occasionally use it for link maps when there is no risk of confusion.

If the components of P have basepoints b_1, \dots, b_m and y_1, \dots, y_m are fixed distinct points in N , we define $\operatorname{Emb}_*(P, N)$ as the space of **based embeddings**, meaning embeddings $f : P \rightarrow N$ such that $f(b_1) = y_1, \dots, f(b_m) = y_m$. We write $\operatorname{Emb}_c(P, N)$ for a space of compactly supported embeddings of P into N , which we will define more precisely in the setting of interest (long links) in Definition 2.1. We use the notation $\operatorname{Link}_*(P, N)$ and $\operatorname{Link}_c(P, N)$ for similarly defined spaces of based link maps and compactly supported link maps respectively. For spaces X and Y , we write $\operatorname{Map}_*(X, Y)$ for the space of based continuous maps $X \rightarrow Y$.

All spaces of smooth maps are equipped with the Whitney C^∞ -topology, while spaces of continuous maps have the compact-open topology. In this paper, N will often be a sphere or Euclidean space, while P will often be a disjoint union of finitely many spheres or Euclidean spaces.

Definition 2.1. Let m and n be positive integers and p_1, \dots, p_m be nonnegative integers less than n . Fix real numbers t_1^*, \dots, t_m^* such that $-1 < t_1^* < t_2^* < \dots < t_m^* < 1$. We define the space $\operatorname{Emb}_c(\mathbb{R}^{p_1} \sqcup \dots \sqcup \mathbb{R}^{p_m}, \mathbb{R}^n)$ of **long links** in \mathbb{R}^n as follows. An element f of this embedding space is required to map $\coprod_1^m (-1, 1)^{p_i}$ into $(-1, 1)^n$. Outside of $\coprod_1^m (-1, 1)^{p_i}$, f must agree with a fixed embedding

$$e = (e_1, \dots, e_m) : \coprod_1^m \mathbb{R}^{p_i} \hookrightarrow \mathbb{R}^n$$

$$e_i : (t_1, \dots, t_{p_i}) \mapsto (t_i^*, 0, \dots, 0, t_1, \dots, t_{p_i}).$$

We take e to be the basepoint of $\operatorname{Emb}_c(\coprod_1^m \mathbb{R}^{p_i}, \mathbb{R}^n)$. A **long knot** is a long link with one component. We define the space $\operatorname{Link}_c(\coprod_1^m \mathbb{R}^{p_i}, \mathbb{R}^n)$ of **long link maps** similarly, just replacing embeddings by link maps.

We sometimes abbreviate $\mathcal{L}_{p_1, \dots, p_m}^n := \operatorname{Emb}_c(\coprod_1^m \mathbb{R}^{p_i}, \mathbb{R}^n)$ and write $\mathcal{L}_{m \cdot p}^n$ when $p_1 = \dots = p_m = p$. Some authors use the term “string links” instead of “long links.”

Our convention of using the last rather than first p_i coordinates in \mathbb{R}^n in Definition 2.1 is motivated by the convention of braids flowing vertically rather than horizontally, and it informs the convention we will use in Definition 2.4 of the graphing map. Any other choice of standard long link e produces a space of long links that can be related to the one in Definition 2.1 by a homeomorphism.

If the dimensions p_1, \dots, p_m of the source manifolds are all at least 1, then the space of long links has a homotopy-associative H-space structure given by stacking. Given $f, g \in \operatorname{Emb}_c(\coprod_1^m \mathbb{R}^{p_i}, \mathbb{R}^n)$, their

product is defined on $\coprod_1^m [-1, 1]^{p_i}$ by setting

$$(f \# g)_i(t_1, \dots, t_{p_i}) := \begin{cases} (\mathbb{1}^{n-1} \times \rho_-^{-1}) \circ f_i \circ (\mathbb{1}^{p_i-1} \times \rho_-) & \text{on } [-1, 1]^{p_i-1} \times [-1, 0] \\ (\mathbb{1}^{n-1} \times \rho_+^{-1}) \circ f_i \circ (\mathbb{1}^{p_i-1} \times \rho_+) & \text{on } [-1, 1]^{p_i-1} \times [0, 1] \end{cases}$$

for each $i = 1, \dots, m$, where $\rho_{\pm}(t) := 2t \mp 1$ and where $\mathbb{1}^k$ denotes the identity map on \mathbb{R}^k . (By abuse of notation, we also use $\mathbb{1}^k$ for the identity on S^k in a context where no ambiguity should arise.) The same operation gives a homotopy associative H-space structure on $\text{Link}_c(\coprod_1^m \mathbb{R}^{p_i}, \mathbb{R}^n)$.

If $p_1 = \dots = p_m = 0$ in Definition 2.1, one obtains the **configuration space** of m points in \mathbb{R}^n , which we also denote $\text{Conf}(m, \mathbb{R}^n)$. Alternatively, it can be defined as

$$\text{Conf}(m, \mathbb{R}^n) := \{(x_1, \dots, x_m) \in (\mathbb{R}^n)^m : x_i \neq x_j \ \forall i \neq j\}.$$

Applying Definition 2.1, we obtain $((t_1^*, 0, \dots, 0), \dots, (t_m^*, 0, \dots, 0))$ as the basepoint of $\text{Conf}(m, \mathbb{R}^n)$. For a finite set S , we will also write $\text{Conf}(S, \mathbb{R}^n)$ for $\text{Emb}(S, \mathbb{R}^n)$. We call an element of the space $\Omega^p \text{Conf}(m, \mathbb{R}^{n-p})$ a p -dimensional, m -component **pure braid** in \mathbb{R}^n . Using the graphing map (see Definition 2.4 below), we may sometimes view such an element as lying in $\mathcal{L}_{m,p}^n$.

For any i and j with $1 \leq i \neq j \leq m$, let $b_{ij} \in \pi_{n-1} \text{Conf}(m, \mathbb{R}^n)$ be the cycle obtained from the map

$$\begin{aligned} S^{n-1} &\rightarrow \text{Conf}(m, \mathbb{R}^n) \\ v &\mapsto (x_1, \dots, x_{i-1}, x_j + \varepsilon v, x_{i+1}, \dots, x_m) \end{aligned}$$

where $\varepsilon < \min_i (t_i^* - t_{i-1}^*)$. More precisely, we get a based map by conjugating by a path which takes the i -th point from its position at the basepoint to the image of the sphere above. For $m = 2$, the map b_{12} (or b_{21}) is a homotopy equivalence.

Definition 2.2. Define the space of **Brunnian long links** $\text{BrEmb}_c(\coprod_1^m \mathbb{R}^{p_i}, \mathbb{R}^n)$ as the subspace of links f in $\text{Emb}_c(\coprod_1^m \mathbb{R}^{p_i}, \mathbb{R}^n)$ such that the restriction of f to any $m-1$ of its components is isotopic to a standard link (e_1, \dots, e_{m-1}) . Define the space $\text{BrEmb}(\coprod_1^m S^{p_i}, S^n)$ of **Brunnian spherical links in a sphere** as the subspace of spherical links f in $\text{Emb}(\coprod_1^m S^{p_i}, S^n)$ such that f is isotopic to a trivial link, where a spherical link is trivial if each component S^{p_i} bounds a disk D^{p_i+1} in S^n .

Thus $\text{BrEmb}_c(\coprod_1^m \mathbb{R}^{p_i}, \mathbb{R}^n)$ is a union of path components of $\text{Emb}_c(\coprod_1^m \mathbb{R}^{p_i}, \mathbb{R}^n)$. In many of the cases we consider, the latter space is path-connected, in which case all embeddings are Brunnian.

Definition 2.3. Define the space $\text{PEmb}(\coprod_{i=1}^m \mathbb{R}^{p_i}, \mathbb{R}^n)$ of **pseudoisotopy embeddings** as the subspace of embeddings $f = (f_1, \dots, f_m) : \coprod_{i=1}^m \mathbb{R}^{p_i} \hookrightarrow \mathbb{R}^n$ such that

- f agrees with the standard embedding $e = (e_1, \dots, e_m)$ outside of $\coprod_{i=1}^m I^{p_i-1} \times [-1, \infty)$ and
- there is a long link $g = (g_1, \dots, g_m) \in \mathcal{L}_{p_1-1, \dots, p_m-1}^{n-1}$ such that if $t_{p_i} \in [1, \infty)$, then $f_i(t_1, \dots, t_{p_i}) = (g_i(t_1, \dots, t_{p_i-1}), t_{p_i})$.

We will often abbreviate this space by $\mathcal{P}_{p_1, \dots, p_m}^n := \text{PEmb}(\coprod_{i=1}^m \mathbb{R}^{p_i}, \mathbb{R}^n)$.

Some authors write $\text{PE}(I^{p-1}, I^{n-1})$ or $\text{CE}(I^{p-1}, I^{n-1})$ to denote a similarly defined space of embeddings $I^p \hookrightarrow I^n$. This space is homeomorphic to \mathcal{P}_p^n via restriction and extension maps. One can think of a pseudoisotopy embedding as an embedding that is fixed on the bottom and sides of the domain cubes and takes the top faces of the domain into the top face of the codomain.

2.2. Maps between spaces of links. The next map appears in most of our main results.

Definition 2.4. For any $m \geq 1$, $0 \leq p_1, \dots, p_m < n$, define the **graphing map**

$$G : \Omega \text{Emb}_c \left(\prod_1^m \mathbb{R}^{p_i}, \mathbb{R}^n \right) \rightarrow \text{Emb}_c \left(\prod_1^m \mathbb{R}^{p_i+1}, \mathbb{R}^{n+1} \right)$$

$$(t \mapsto f(t) = (f_1(t), \dots, f_m(t))) \mapsto (G(f)_1, \dots, G(f)_m)$$

by

$$G(f)_i(t_1, \dots, t_{p_i+1}) := (f_i(t_{p_i+1})(t_1, \dots, t_{p_i}), t_{p_i+1})$$

for $i = 1, \dots, m$.

Compared to the usual conventions for graphs, the coordinates of the domain and range are swapped because for $n = 2$ and $p_1 = \dots = p_m = 0$, braids are more often drawn as flowing vertically rather than horizontally.

Iterating such graphing maps gives rise to maps of the form

$$G^k : \Omega^k \text{Emb}_c \left(\prod_1^m \mathbb{R}^{p_i}, \mathbb{R}^n \right) \rightarrow \text{Emb}_c \left(\prod_1^m \mathbb{R}^{p_i+k}, \mathbb{R}^{n+k} \right).$$

More precisely, G^k is the composition of the maps induced on various iterated loop spaces by various graphing maps, all of which by abuse of notation we will denote G .

The next map lets us pass from long links to spherical (based) links in a Euclidean space. It will be used in results on injectivity of graphing in Section 3. It is roughly given by fixing disjoint embeddings of disks into the complement of $[-1, 1]^n$ and gluing these to the parts of the long link in $[-1, 1]^n$.

To give the precise definition, we introduce some more notation. For any $k \geq 1$, let rD^k be the closed disk in \mathbb{R}^k centered at the origin and with radius r . For $i = 1, \dots, m$, let D_i be the image of $\sqrt{n}D^{p_i}$ under e_i , the i -th component of the standard long link. Thus $e_i([-1, 1]^{p_i})$ is a cube inscribed in the disk D_i . Fix disjoint open intervals $I_1, \dots, I_m \subset \mathbb{R}$ such that $t_i^* \in I_i$, where t_i^* are the basepoints in $[-1, 1]$ from Definition 2.1.

Definition 2.5. For $m \geq 1$ and $1 \leq p_1, \dots, p_m \leq n-2$, define the **closure map**

$$\hat{\cdot} : \text{Emb}_c \left(\prod_1^m \mathbb{R}^{p_i}, \mathbb{R}^n \right) \rightarrow \text{Emb}_* \left(\prod_1^m S^{p_i}, \mathbb{R}^n \right) \tag{2.1}$$

$$f = (f_1, \dots, f_m) \mapsto \hat{f} = (\hat{f}_1, \dots, \hat{f}_m)$$

by fixing for each $i = 1, \dots, m$ a proper smooth embedding

$$g_i : D^{p_i} \rightarrow I_i \times \mathbb{R}^{n-1} - \sqrt{n}D^n$$

which maps ∂D^{p_i} to $\{(t_i^*, 0^{n-p_i-1})\} \times \partial D_i \subset \mathbb{R}^{n-p_i} \times \mathbb{R}^{p_i}$ and setting

$$\hat{f}_i := f_i|_{D^{p_i}} \cup g_i : D^{p_i} \cup D^{p_i} = S^{p_i} \rightarrow \mathbb{R}^n. \tag{2.2}$$

The codomain of the map $\hat{\cdot}$ is defined as the space of based embeddings where we take the basepoint b_i in each copy of S^{p_i} to be any point in the image of g_i , and we take the image of b_i in \mathbb{R}^n to be the base value y_i .

An embedding g_i as in Definition 2.5 exists because its target is homeomorphic to $(0, 1) \times (\mathbb{R}^{n-1} - D^{n-1})$ and $n - p_i \geq 2$. For example, one could construct g_i by joining a small outward perturbation of a hemisphere of S^{p_i} in $\mathbb{R}^{p_i+1} \subset \mathbb{R}^{n-1}$ to the equator of S^{p_i} . The closure map depends on the choice of embeddings g_i , but these can be made independently of f , so that it is well defined on the space level, not just on isotopy classes. In Section 3, we will further specify these g_i for $m = 2$.

Because the intervals I_1, \dots, I_m are disjoint, the closure \widehat{e} of the standard long link e is a trivial link, meaning that its components bound disjoint $(p_i + 1)$ -dimensional disks in \mathbb{R}^n . One could define a closure map where one allows $n - p_i = 1$, but it cannot take a standard long link to a trivial link if $n - p_i = 1$ for three or more values of i . We take \widehat{e} to be the basepoint of $\text{Emb}_* \left(\coprod_{i=1}^m S^{p_i}, \mathbb{R}^n \right)$. We can also use it as the basepoint of the space $\text{Emb} \left(\coprod_{i=1}^m S^{p_i}, \mathbb{R}^n \right)$ of unbased embeddings.

We will use similarly defined closure maps denoted by the same symbol:

$$\begin{aligned} \widehat{\cdot} : \text{Link}_c \left(\coprod_{i=1}^m \mathbb{R}^{p_i}, \mathbb{R}^n \right) &\rightarrow \text{Link}_* \left(\coprod_{i=1}^m S^{p_i}, \mathbb{R}^n \right) \\ \widehat{\cdot} : \text{Emb}_c \left(\coprod_{i=1}^m \mathbb{R}^{p_i}, \mathbb{R}^n \right) &\rightarrow \text{Emb} \left(\coprod_{i=1}^m S^{p_i}, S^n \right) \end{aligned}$$

The second one is obtained by post-composing the map (2.1) by the map induced by a fixed inclusion $\mathbb{R}^n \hookrightarrow S^n$, and forgetting that the resulting embeddings are based.

The following map will be used in Section 5 to study long knots and 2-component links via 2- and 3-component links. Recall that $\mathcal{L}_{m,p}^n$ denotes the space $\mathcal{L}_{p_1, \dots, p_m}^n = \text{Emb}_c \left(\coprod_{i=1}^m \mathbb{R}^{p_i}, \mathbb{R}^n \right)$ where each $p_i = p$.

Definition 2.6. Define the **joining map**

$$J : \mathcal{L}_{m,p}^n \rightarrow \mathcal{L}_{(m-1),p}^n$$

for any $m \geq 2$, as follows. For $f = (f_1, \dots, f_m)$ in $\mathcal{L}_{m,p}^n$, take the closure \widehat{f}_m , as defined in (2.2), of just the last component. Then form a connected sum of the images of f_{m-1} and \widehat{f}_m , where the connected sum preserves orientation and occurs outside $[-1, 1]^n$. The result is diffeomorphic to \mathbb{R}^p . By an appropriate parametrization, an isotopy supported outside of $[-1, 1]^n$, and a linear diffeomorphism of \mathbb{R}^n , we can ensure that the result $J(f)$ is indeed an element of $\mathcal{L}_{(m-1),p}^n$.

Some choices must be made to define J , but they can all be made independently of f so that J is well defined on the space level, not just on isotopy classes. In Section 5.4, we will further specify some of these choices. We will also use an alternative description for $p = 1$, where one can delete an open ray from each component and join the remaining closed rays by an arc. The joining map J behaves with respect to standard links much like the closure map does with respect to standard and trivial links. Namely, $J(e_1, \dots, e_m)$ is isotopic to (e_1, \dots, e_{m-1}) . One could generalize J to maps that join different pairs of components, but we do not need such a generalization.

Finally, we consider **restriction** maps ρ_S and **inclusion** maps ι_S between spaces of links of different numbers of components. As a special case, if $0 \leq p, q < n$, there is a fibration

$$\text{Emb}_c(\mathbb{R}^p, \mathbb{R}^n - \mathbb{R}^q) \rightarrow \mathcal{L}_{p,q}^n \xrightarrow{\rho_2} \mathcal{K}_q^n \quad (2.3)$$

which restricts a 2-component link to the second component. It is a fibration by work of Palais or Lima [Pal60, Lim64]. Taking the fiber over the standard long link $e = (e_1, e_2)$, the space $\mathbb{R}^n - \mathbb{R}^q$ is more accurately $\mathbb{R}^n - e_2(\mathbb{R}^q)$. One can define an inclusion $\iota_2 : \mathcal{K}_q^n \rightarrow \mathcal{L}_{p,q}^n$ which is a section of ρ_2 up to homotopy. Let $t^* \in (-1, 1)$ and $t_1^*, t_2^* \in (-1, 1)$ be the basepoint values for \mathcal{K}_q^n and $\mathcal{L}_{p,q}^n$ respectively. Fix a real number a such that $t_1^* < a < t_2^*$. Define ι_2 by first mapping the ambient space \mathbb{R}^n homeomorphically onto $(a, \infty) \times \mathbb{R}^{n-1}$ so that $\{t^*\} \times \mathbb{R}^{n-1}$ is mapped to $\{t_2^*\} \times \mathbb{R}^{n-1}$. Then embed a standard first component $\mathbb{R}^p \hookrightarrow (\infty, a) \times \mathbb{R}^{n-1}$. Because of this section, the long exact sequence in homotopy of the fibration (2.3) splits as a direct sum.

There is similarly a restriction ρ_1 to the first component with a section ι_1 . We leave it to the reader to generalize this definition to maps ρ_S (respectively ι_S) with domain (respectively codomain) $\mathcal{L}_{p_1, \dots, p_m}^n$,

where S is any subset of $\{1, \dots, m\}$. In the general case, one may use an isotopy, a reparametrization, and a diffeomorphism as in the definition of the joining maps to ensure that the resulting embeddings have the required behavior outside of $[-1, 1]^n$.

We will use the fact that the graphing and closure maps preserve the Brunnian property. It is easy to see that the graphing, restriction, and inclusion maps are maps of H-spaces, whereas the maps that join components are not.

3. INJECTIVITY OF GRAPHING

In Section 3.1, we prove injectivity results for graphing into spaces of long and spherical links using the λ -invariant or an analogue of it. These results concern homotopy groups in arbitrarily high degrees. In Section 3.2, we prove an analogous result of spaces of long link maps, using the α -invariant. Injectivity in that setting applies only to homotopy groups in a range. The proofs of all the results in this Section are somewhat similar.

Throughout this Section, we consider only 2-component links. In this setting, we first impose some further conditions on the data used in the definitions of long links and the closure map.

Fix $t_1^* = -1/2$ and $t_2^* = 0$ so that the standard long link $e = (e_1, e_2)$ for $\text{Emb}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n)$ has $e_1(t) = (-1/2, 0^{n-p-1}, t)$ and $e_2(u) = (0^{n-q}, u)$. Take e as the standard long link map for $\text{Link}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n)$.

For the closure map $\hat{\cdot} : \text{Emb}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n) \rightarrow \text{Emb}(S^p \sqcup S^q, \mathbb{R}^n)$, recall that $\hat{f}_i := f_i \cup g_i$ for some fixed embeddings g_1 and g_2 of D^p and D^q respectively. Choose the g_i so that

$$\begin{aligned} \text{im}(g_1) &\subset \{(t_1, \dots, t_n) : t_1 \leq -1/2\} \quad \text{and} \\ \text{im}(g_2) &\subset \{(t_1, \dots, t_n) : t_1 \geq 0\}. \end{aligned} \tag{3.1}$$

We impose this same condition on the g_i for the closure map $\text{Link}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n) \rightarrow \text{Link}(S^p \sqcup S^q, S^n)$.

We will use diffeomorphisms of the form

$$\begin{aligned} \mathbb{R}^n - \mathbb{R}^q &\rightarrow S^{n-q-1} \times \mathbb{R}^{q+1} \\ (t_1, \dots, t_n) &\mapsto \left(\frac{(t_1, \dots, t_{n-q})}{|(t_1, \dots, t_{n-q})|}, \sigma(|(t_1, \dots, t_{n-q})|), t_{n-q+1}, \dots, t_n \right) \end{aligned} \tag{3.2}$$

where $\sigma : (0, \infty) \times \mathbb{R}^q \rightarrow \mathbb{R}^{q+1}$ is the homeomorphism given by $\sigma(r, \vec{t}) := (\ln r, \vec{t})$. By taking the one-point compactification of \mathbb{R}^n , we get a diffeomorphism

$$S^n - S^q \rightarrow S^{n-q-1} \times \mathbb{R}^{q+1} \tag{3.3}$$

The exact formula for this diffeomorphism is not as important as the fact that it can be written as the product of two maps, where the second is the identity on the space \mathbb{R}^q given by the last q coordinates.

3.1. Injectivity for spaces of embeddings via the λ -invariant. We begin with long links, where we will define a space-level retract from a space of embeddings to a sphere that can be viewed as an adaptation to long links of the λ -invariant (defined in Definition 3.2 below).

Theorem 3.1. *Suppose $0 \leq p \leq q \leq n-1$ and $i \geq 0$. Then $\pi_i \text{Emb}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n)$ contains a direct summand isomorphic to $\pi_{i+p} S^{n-q-1}$, the inclusion of which can be realized by a graphing map.*

Proof. To prove the theorem for $\text{Emb}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n)$, it suffices to prove it instead for $\text{Emb}_c(\mathbb{R}^p, \mathbb{R}^n - \mathbb{R}^q)$. Indeed, the fibration (2.3) and its section (up to homotopy) ensure that $\pi_i \text{Emb}_c(\mathbb{R}^p, \mathbb{R}^n - \mathbb{R}^q)$ is a direct summand of $\pi_i \mathcal{L}_{p,q}^n = \pi_i \text{Emb}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n)$.

To prove the theorem for $\text{Emb}_c(\mathbb{R}^p, \mathbb{R}^n - \mathbb{R}^q)$, we show that it has $\Omega^p S^{n-q-1}$ as a retract, by showing that the following composition (of maps of H-spaces) is the identity:

$$\begin{aligned}
 \Omega^p S^{n-q-1} &\xrightarrow{\cong} \Omega^p(S^{n-q-1} \times \mathbb{R}^{q-p+1}) \xrightarrow{\cong} \Omega^p(\mathbb{R}^{n-p} - \mathbb{R}^{q-p}) \\
 &\xrightarrow{\cong} \Omega^p \text{Emb}_c(\{*\}, \mathbb{R}^{n-p} - \mathbb{R}^{q-p}) \\
 &\xrightarrow{G^p} \text{Emb}_c(\mathbb{R}^p, \mathbb{R}^n - \mathbb{R}^q) \\
 &\longrightarrow \Omega^p(\mathbb{R}^n - \mathbb{R}^q) \\
 &\xrightarrow{\cong} \Omega^p(S^{n-q-1} \times \mathbb{R}^{q+1}) \\
 &\xrightarrow{\cong} \Omega^p S^{n-q-1}
 \end{aligned} \tag{3.4}$$

The first map comes the inclusion $i : S^{n-q-1} \rightarrow S^{n-q-1} \times \mathbb{R}^{q-p+1}$ defined by $i(s) = (s, 0)$. The second map comes from a diffeomorphism $h : \mathbb{R}^{n-p} - \mathbb{R}^{q-p} \rightarrow S^{n-q-1} \times \mathbb{R}^{q-p+1}$ as in (3.2). The unlabeled arrow is a forgetful map. The penultimate map comes from a diffeomorphism $\mathbb{R}^n - \mathbb{R}^q \rightarrow S^{n-q-1} \times \mathbb{R}^{q+1}$, as in formula (3.2), which is $h \times \text{id}_{\mathbb{R}^p}$. The last map comes the projection $\pi : S^{n-q-1} \times \mathbb{R}^{i+q+1} \rightarrow S^{n-q-1}$. The composition of the last three arrows then maps a link g to $\pi \circ (h \times \text{id}_{\mathbb{R}^{i+p}}) \circ g$.

The result of applying the first three maps to $f : \mathbb{R}^p \rightarrow S^{n-q-1}$ is the composite $h^{-1}if$. Thus the composite (3.4) sends f to the composite

$$\begin{array}{ccccccc}
 \mathbb{R}^p & \longrightarrow & (\mathbb{R}^{n-p} - \mathbb{R}^{q-p}) \times \mathbb{R}^p & \xrightarrow{h \times \text{id}_{\mathbb{R}^p}} & (S^{n-q-1} \times \mathbb{R}^{q-p+1}) \times \mathbb{R}^p & \xrightarrow{\pi} & S^{n-q-1} \\
 t & \longmapsto & (h^{-1}if(t), t) & \longmapsto & (if(t), t) & \longmapsto & f(t)
 \end{array}$$

Hence the composite (3.4) it is the identity, as desired. \square

For Theorem 3.4, the analogue of Theorem 3.1 for spherical links in a sphere, we use the subspace of Brunnian links; see Definition 2.2. This is because for spherical links, the restriction to one component does not in general admit a section. (In fact, for $p = q = 1$ and $n = 3$, the component of the Hopf link is homotopy equivalent to $SO(4)$ [HK21, Corollary 4.4(e)], while the component of the unknot is homotopy equivalent to $SO(4)/SO(2)$ [Hat83]. One can see that the projection $SO(4) \rightarrow SO(4)/SO(2)$ admits no section by considering the long exact sequence in homotopy at π_2 and π_1 .) We will use the λ -invariant, which appears at least as far back as work of Zeeman [Zee60, Zee62] and Haefliger [Hae62].

Definition 3.2. For $p \leq q < n$, define a map

$$\pi_0 \text{BrEmb}(S^p \sqcup S^q, S^n) \xrightarrow{\lambda} \pi_p S^{n-q-1}$$

as follows. Denote an element of the domain by $b = (b_1, b_2)$. Take any isotopy of b_2 to the standard embedding \widehat{e}_2 , and extend it to an isotopy from b to a link $b' = (b'_1, \widehat{e}_2)$. Define $\lambda(b)$ as the composite

$$S^p \xrightarrow{b'_1} S^n - S^q \longrightarrow S^{n-q-1} \times \mathbb{R}^{q+1} \longrightarrow S^{n-q-1}$$

where the second map is the diffeomorphism (3.3) and the third map is projection onto the first factor.

The proof of the next lemma is essentially from Haefliger's work [Hae67, Theorem 7.1] but is included for the reader's convenience.

Lemma 3.3. *The map λ is well defined.*

Proof. We need to show that $\lambda(b)$ is independent of the choice of isotopy from b_2 to \widehat{e}_2 and the choice of an extension of it to the first component. Suppose b' and b'' are the endpoints of two such isotopies

starting at b . Then b' and b'' are isotopic links, each restricting to \widehat{e}_2 on S^q (though the links throughout the isotopy need not satisfy this property).

An isotopy from b' to b'' can be extended to a path of diffeomorphisms of S^n starting at the identity. Restrict this path to a disk D^n which contains $b'_1(S^p)$ and which intersects $\widehat{e}_2(S^q)$ in a disk $D^q \subset D^n$. The endpoints g_0 and g_1 of this path lie in the space $\text{Emb}^+(D^n, S^n; D^q)$ of orientation-preserving embeddings $D^n \hookrightarrow S^n$ which agree with the fixed inclusion on D^q . By a shrinking and linearization argument, $\text{Emb}^+(D^n, S^n; D^q) \simeq SO(n-q)$. In particular, it is path-connected, so there is a path in $\text{Emb}^+(D^n, S^n; D^q)$ from g_0 to g_1 .

Lift this path to the space $\text{Diff}^+(S^n; \widehat{e}_2(S^q))$ of diffeomorphisms of S^n which fix $\widehat{e}_2(S^q)$ pointwise. We can do so by the isotopy extension theorem [Hir76, Theorem 8.1.3] or more generally the fact that the restriction $\text{Diff}^+(S^n; \widehat{e}_2(S^q)) \rightarrow \text{Emb}^+(D^n, S^n; D^q)$ is a fibration [Pal60, Lim64]. Restricting the latter path to the image of b' gives rise to an isotopy from b' to b'' which leaves the second component fixed throughout. \square

Theorem 3.4. *Let $1 \leq p \leq q \leq n-2$, and let $i \geq 0$. Then $\pi_i \text{Emb}(S^p \sqcup S^q, S^n)$ contains a direct summand isomorphic to $\pi_{i+p} S^{n-q-1}$, the inclusion of which can be realized by a graphing map.*

Theorem 3.4 is valid for $q = n-1$ but trivial because we assume $p \geq 1$. We impose that hypothesis because we defined the closure map only for codimension at least 2.

Proof. To prove the theorem for $\text{Emb}(S^p \sqcup S^q, S^n)$, it suffices to prove the statement for $\text{BrEmb}(S^p \sqcup S^q, S^n)$ because the latter space is a union of path components of the former space. We will show that the following composition of maps is the identity:

$$\begin{aligned}
\pi_{i+p} S^{n-q-1} &\cong \dots \cong \pi_i \Omega^p(\mathbb{R}^{n-p} - \mathbb{R}^{q-p}) \\
&\cong \pi_i \Omega^p \text{BrEmb}_c(\{*\} \sqcup \mathbb{R}^{q-p}, \mathbb{R}^{n-p}) \\
&\xrightarrow{G_*^p} \pi_i \text{BrEmb}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n) \\
&\xrightarrow{\widehat{}} \pi_i \text{BrEmb}(S^p \sqcup S^q, S^n) \\
&\xrightarrow{G_*^i} \pi_0 \text{BrEmb}(S^{i+p} \sqcup S^{i+q}, S^{i+n}) \\
&\xrightarrow{\lambda} \pi_{i+p} S^{n-q-1}
\end{aligned} \tag{3.5}$$

The ellipsis stands for isomorphisms coming from homotopy equivalences or homeomorphisms like those in the first line of the diagram (3.4). The map denoted G^i (by abusing notation from the setting of long links) is the composition of i maps, starting with

$$\Omega \text{BrEmb}(S^p \sqcup S^q, S^n) \rightarrow \text{BrEmb}(S^{p+1} \sqcup S^{q+1}, S^{n+1}).$$

To obtain this first map, start by graphing a loop restricted to I to get an embedding $S^p \times I \sqcup S^q \times I \rightarrow S^n \times I$. Then attach two disks D^{n+1} to $S^n \times I$ using diffeomorphisms $\partial D^{n+1} \rightarrow S^n \times \{\pm 1\}$. In each copy of D^{n+1} , fix smooth proper embeddings of D^{p+1} and D^{q+1} and glue two copies of each of these to the embeddings of $S^p \times I$ and $S^q \times I$ to obtain smooth embeddings of S^{p+1} and S^{q+1} . We can use fixed embeddings of D^{p+1} and D^{q+1} independent of the embeddings of S^p and S^q because the loop of embeddings is based at the standard trivial link \widehat{e} . The remaining $i-1$ maps needed to obtain G^i are defined similarly.

From the proof of Theorem 3.1 and the relationship between λ and the map $\text{Emb}_c(\mathbb{R}^p, \mathbb{R}^n - \mathbb{R}^q) \rightarrow \Omega^p(\mathbb{R}^n - \mathbb{R}^q)$, we can see that the composite (3.5) would be the identity if we applied λ to the restriction $D^{i+p} \sqcup D^{i+q} \rightarrow D^{i+n}$ of an element of $\pi_0 \text{BrEmb}(S^{i+p} \sqcup S^{i+q}, S^{i+n})$.

The proof will now be completed by identifying elements of $\pi_{i+p}S^{n-q-1}$ with bordism classes of framed manifolds, via the Pontryagin–Thom correspondence. A homotopy class is identified with a bordism class by taking the preimage of a regular value of a smooth representative $\mathbb{R}^{i+p} \rightarrow S^{n-q-1}$. Recall the conditions assumed on the closure map in (3.1). By choosing the regular value $y \in S^{n-q-1}$ sufficiently far from the first coordinate axis in \mathbb{R}^{n-q} , we can ensure that the framed submanifold representing a bordism class in the target copy of $\pi_{i+p}S^{n-q-1}$ lies in $\mathbb{R}^i \times D^p \subset \mathbb{R}^{i+p}$. For example, we can take y to lie in an ε -neighborhood of $(0, 1, 0, \dots, 0)$, where exactly how small ε must be depends on the fixed embeddings g_i used in the definition of the closure map. (A similar argument will be used in the proof of Theorem 3.7.)

The fact that λ yields the identity when restricted to embeddings of $D^{i+p} \sqcup D^{i+q}$ therefore guarantees that the composite (3.5) is the identity. \square

Example 3.5 (Generalized linking numbers). One of the simplest cases of Theorem 3.1 and 3.4 is when $p = q = 1$, $n = 3$, and $i = 0$, where the theorem reduces to an easily proven injection of $\pi_1(S^1) \cong \mathbb{Z}$, the group of 2-strand classical pure braids, into the monoid of isotopy classes of classical 2-component long, or closed, links. More generally, if $i + p + q = n - 1$, then a generator of the resulting copy of \mathbb{Z} in $\pi_i \text{Emb}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^{i+p+q+1})$ is dual to the linking number of manifolds of dimensions say $i + p$ and q in $\mathbb{R}^{i+p+q+1}$, by combining the domain of an element of π_i and the source manifold \mathbb{R}^p into an embedded $(i + p)$ -dimensional manifold.

3.2. Injectivity for spaces of link maps in a range via the α -invariant. Our next main result is proven using the α -invariant of spherical links in Euclidean space. It is essentially a Gauss map and appears at least as early as the work of Massey and Rolfsen [MR85]. Further developments on it include work of Koschorke [Kos88, Kos97] and A. Skopenkov [Sko00].

Definition 3.6. We define α as the map

$$\begin{aligned} \alpha : \text{Link}_*(S^p \sqcup S^q, \mathbb{R}^n) &\longrightarrow \text{Map}_*(S^p \times S^q, \text{Conf}(2, \mathbb{R}^n)) \\ g = (g_1, g_2) &\longmapsto ((t, u) \mapsto (g_1(t), g_2(u))). \end{aligned}$$

Now $\text{Conf}(2, \mathbb{R}^n) \simeq S^{n-1}$, so if $p, q \leq n - 2$, then the restriction of any map $S^p \times S^q \rightarrow \text{Conf}(2, \mathbb{R}^n)$ to $S^p \vee S^q$ is nullhomotopic. Thus the map induced by α on π_i for any $i \geq 0$ can be written as

$$\begin{aligned} \alpha_* : \pi_i \text{Link}_*(S^p \sqcup S^q, \mathbb{R}^n) &\rightarrow \pi_i \text{Map}_*((S^p \times S^q)/(S^p \vee S^q), \text{Conf}(2, \mathbb{R}^n)) \\ &\cong \pi_{i+p+q} \text{Conf}(2, \mathbb{R}^n) \\ &\cong \pi_{i+p+q} S^{n-1} \end{aligned}$$

Theorem 3.7. *Suppose $1 \leq p \leq q \leq n - 2$ and $i \geq 0$. Let $G_*^p : \pi_{i+p}S^{n-q-1} \rightarrow \pi_i \text{Link}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n)$ be the map induced by graphing. Let $E^q : \pi_{i+p}S^{n-q-1} \rightarrow \pi_{i+p+q}S^{n-1}$ be the q -fold suspension map. Then G_p^* is injective, nonzero, or split injective if the q -fold suspension map is respectively injective, nonzero, or an isomorphism.*

Like Theorem 3.4, Theorem 3.7 is valid for $q = n - 1$ but trivial because we assume $p \geq 1$. Again, we impose that hypothesis because we defined the closure map only for codimension at least 2.

Proof. The main idea is to check that the composite $\pi_{i+p}S^{n-q-1} \rightarrow \pi_{i+p+q}S^{n-1}$, shown below in the composite (3.6), agrees with the q -fold iterated suspension E^q . For notational ease, we write \widehat{G}^p to denote the

p -fold graphing G^p followed by the closure map $\hat{\cdot}$. Thus we write the composite below as $[f] \mapsto \alpha_* \widehat{G}_*^p [f]$.

$$\begin{aligned}
\pi_{i+p} S^{n-q-1} &\cong \pi_i \Omega^p S^{n-q-1} \cong \pi_i \Omega^p (S^{n-q-1} \times \mathbb{R}^{q-p+1}) \\
&\cong \pi_i \Omega^p (\mathbb{R}^{n-p} - \mathbb{R}^{q-p}) \\
&\longrightarrow \pi_i \Omega^p \text{Link}_c(\{*\} \sqcup \mathbb{R}^{q-p}, \mathbb{R}^{n-p}) \\
&\xrightarrow{G_*^p} \pi_i \text{Link}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n) \\
&\xrightarrow{(\hat{\cdot})_*} \pi_i \text{Link}_*(S^p \sqcup S^q, \mathbb{R}^n) \\
&\xrightarrow{\alpha_*} \pi_{i+p+q} S^{n-1}
\end{aligned} \tag{3.6}$$

All of the above maps are homomorphisms, even if $i = 0$, because G^p is a map of H-spaces.

We will use the Pontryagin–Thom Theorem to identify elements of the first and last homotopy groups in the composite (3.6) with bordism classes of manifolds. To do so, one can take the preimage of any regular value under a smooth representative of a homotopy class. In particular, we will identify $[f]$ with a bordism class of framed submanifold of \mathbb{R}^{i+p} of dimension $i + p + q - n + 1$, and we will identify $\alpha_* \widehat{G}_*^p [f]$ with a class of framed submanifold of \mathbb{R}^{i+p+q} , also of dimension $i + p + q - n + 1$. Let $y = (y_1, \dots, y_{n-q}) \in S^{n-q-1}$ be a regular value of f that lies in a small neighborhood of $(0, 1, 0, \dots, 0)$. The latter assumption will simplify a consideration below related to the closure map. Let $X := f^{-1}\{y\}$ be the framed bordism class corresponding to $[f]$.

The result of applying the first three isomorphisms in the composite (3.6) to a map $f : \mathbb{R}^{i+p} \rightarrow S^{n-q-1}$ is more explicitly written as the composition

$$\mathbb{R}^i \times \mathbb{R}^p \xrightarrow{f} S^{n-q-1} \xrightarrow{i} S^{n-q-1} \times \mathbb{R}^{q-p+1} \xrightarrow{h^{-1}} \mathbb{R}^{n-p} - \mathbb{R}^{q-p}$$

where i maps S^{n-q-1} to $S^{n-q-1} \times \{0^{q-p+1}\}$. Above h is a diffeomorphism as in the proof of Theorem 3.1. Note that by identifying $S^{n-q-1} \times \mathbb{R}$ with $\mathbb{R}^{n-q} - \{0^{n-q}\}$, we can easily choose h so that $h^{-1}i(S^{n-q-1})$ is the unit sphere in $\mathbb{R}^{n-q} \times \{0^{q-p}\} \subset \mathbb{R}^{n-p}$.

The first arrow in the composite (3.6) comes from mapping $\mathbb{R}^{n-p} - \mathbb{R}^{q-p}$ homeomorphically onto the subspace of $\text{Link}_c(\{*\} \sqcup \mathbb{R}^{q-p}, \mathbb{R}^{n-p})$ where the \mathbb{R}^{q-p} component is standard. Thus the restriction to the first component of the image of $[f]$ in $\pi_i \text{Link}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n)$ is represented by the map

$$\begin{aligned}
\mathbb{R}^i \times \mathbb{R}^p &\longrightarrow (\mathbb{R}^{n-p} - \mathbb{R}^{q-p}) \times \mathbb{R}^p \\
(s, t) &\longmapsto (h^{-1}if(s, t), t).
\end{aligned}$$

Hence on the subspace $D^p \times D^q \subset S^p \times S^q$, the class $\alpha_*(\widehat{G}_*^p [f])$ is represented by the composite

$$\begin{aligned}
\mathbb{R}^i \times D^p \times D^q &\longrightarrow \text{Conf}(2, \mathbb{R}^n) \xrightarrow{\cong} S^{n-1} \\
(s, t, u) &\longmapsto ((h^{-1}if(s, t), t), (0^{n-q}, u))
\end{aligned} \tag{3.7}$$

Using the formula in (3.7), one can check that $(y, 0^q) \in S^{n-1} \subset \mathbb{R}^n$ is a regular value of the composite shown there. Because of the conditions on the closure map assumed at the beginning of this Section, taking the regular value y in a small enough neighborhood of $(0, 1, 0, \dots, 0)$ guarantees that the preimage of $(y, 0^q)$ under $\alpha \widehat{G}^p f$ is contained in $\mathbb{R}^i \times D^p \times D^q \subset \mathbb{R}^i \times S^q \times S^q$. Let $Z := (\alpha \widehat{G}^p f)^{-1}\{(y, 0^q)\}$. From the formula in (3.7) and the fact that $h^{-1}i(S^{n-q-1})$ is the unit sphere in $\mathbb{R}^{n-q} \times \{0^{q-p}\} \subset \mathbb{R}^{n-p}$, we deduce that

$$\begin{aligned}
Z &= \{(s, t, 0^{q-p}, t) : h^{-1}if(s, t) = h^{-1}i(y)\} \\
&= \{(s, t, 0^{q-p}, t) : f(s, t) = y\}
\end{aligned}$$

where the second equality comes from the fact that $h^{-1}i$ is injective.

The iterated suspension $E^q[f]$ corresponds to the image of X under the inclusion $\iota_{\mathbb{R}^{i+p}, \mathbb{R}^{i+p+q}} : \mathbb{R}^{i+p} \hookrightarrow \mathbb{R}^{i+p+q}$ of the first $i+p$ coordinates. Now

$$\iota_{\mathbb{R}^{i+p}, \mathbb{R}^{i+p+q}}(X) = \{(s, t, 0^q) : f(s, t) = y\}.$$

This submanifold (with its induced framing) is bordant to Z (with its induced framing) via the ambient isotopy

$$\begin{aligned} \mathbb{R}^{i+p+q} \times [0, 1] &\rightarrow \mathbb{R}^{i+p+q} \\ ((t_1, \dots, t_{i+p+q}), r) &\mapsto (t_1, \dots, t_{i+q}, t_{i+q+1} + r t_{i+1}, \dots, t_{i+q+p} + r t_{i+p}). \end{aligned}$$

Thus the composition $\pi_{i+p} S^{n-q-1} \rightarrow \pi_{i+p+q} S^{n-1}$ shown in (3.6) is the q -fold suspension E^q , so the map $G_*^p : \pi_{i+p} S^{n-q-1} \rightarrow \pi_i \text{Link}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n)$ is injective or nonzero if E^q is, and if E^q is an isomorphism, G_*^p maps isomorphically onto a direct summand. \square

Imposing the condition $i \leq 2n - p - 2q - 4$ and applying the Freudenthal suspension theorem yields Theorem B from Theorem 3.7.

One can immediately adapt Example 3.5 on linking number classes from Theorem 3.1 to Theorem 3.7, i.e., from embeddings to link maps.

Remark 3.8 (Spaces of based spherical link maps in Euclidean space). An analogue of Theorem 3.7 also holds for the space $\text{Link}_*(S^p \sqcup S^q, \mathbb{R}^n)$. Indeed, the proof applies to \widehat{G}_*^p just as well as it does to G_*^p .

Remark 3.9 (Spaces of embeddings and the α -invariant). Theorem 3.7 and its proof are equally valid for spaces of embeddings rather than link maps. However, for spaces of embeddings, Theorem 3.1 already gives a stronger result.

4. BIJECTIVITY OF GRAPHING IN A RANGE

We will now give a complete calculation, at least up to knotting, of homotopy groups of spaces of long links in a certain range. We will consider spaces of long (knots and) links, such as \mathcal{K}_p^n and $\mathcal{L}_{p,q}^n$, as well as spaces of pseudoisotopy embeddings, such as \mathcal{P}_p^n and $\mathcal{P}_{p,q}^n$. We will not consider spaces of link maps any further. The key ingredients, given in Section 4.1, are a fibration sequence involving spaces of long links and a space of pseudoisotopy embeddings, together with a lower bound on the connectivity of the latter space. In Section 4.2, we first establish bijectivity in a range for graphing on spaces of 2-component links (Theorem 4.3). In this range, this determines the homotopy groups of any space of long links (Theorem 4.6). In Section 4.3, we extend the bijectivity result to one degree higher for equidimensional links by starting with 2-dimensional rather than 0-dimensional links (Theorem 4.10). This requires an identification of isotopy classes between spherical links and long links (Lemma 4.8). Many of the results and the methods in this Section are extensions to long links of Budney's work [Bud08, Proposition 3.9] on spaces of long knots.

4.1. Lemmas on pseudoisotopy embedding spaces. Restriction of a 2-component pseudoisotopy embedding $\mathbb{R}^p \sqcup \mathbb{R}^q \hookrightarrow \mathbb{R}^n$ to the hyperplanes $\{t_p = 1\} \sqcup \{t_q = 1\}$ gives the following fibration:

$$\mathcal{L}_{p,q}^n \xrightarrow{i} \mathcal{P}_{p,q}^n \xrightarrow{\rho} \mathcal{L}_{p-1,q-1}^{n-1}$$

In turn, this leads to the sequence

$$\Omega \mathcal{L}_{p-1,q-1}^{n-1} \xrightarrow{\partial} \mathcal{L}_{p,q}^n \xrightarrow{i} \mathcal{P}_{p,q}^n$$

which up to homotopy is a fibration. To be somewhat explicit, we review the general construction of the connecting map ∂ for a fibration in this special case. It comes from the following homotopy-commutative diagram:

$$\begin{array}{ccccc} \Omega \mathcal{L}_{p-1, q-1}^{n-1} & \xrightarrow{\partial} & \mathcal{L}_{p, q}^n & \xrightarrow{i} & \mathcal{P}_{p, q}^n \\ p \uparrow \simeq & & \pi \uparrow \simeq & & \parallel \\ \text{hofib}(i) & \xrightarrow{j} & \widetilde{\mathcal{L}}_{p, q}^n & \xrightarrow{k} & \mathcal{P}_{p, q}^n \end{array} \quad (4.1)$$

The space $\widetilde{\mathcal{L}}_{p, q}^n$ is defined by

$$\widetilde{\mathcal{L}}_{p, q}^n := \left\{ (f, \gamma) : f \in \mathcal{L}_{p, q}^n, \gamma : [-1, 1] \rightarrow \mathcal{P}_{p, q}^n, \gamma(1) = i(f) \right\}.$$

The homotopy equivalence $\pi : \widetilde{\mathcal{L}}_{p, q}^n \rightarrow \mathcal{L}_{p, q}^n$ is the projection $\pi : (f, \gamma) \mapsto f$. The map $k : \widetilde{\mathcal{L}}_{p, q}^n \rightarrow \mathcal{P}_{p, q}^n$ is given by $k : (f, \gamma) \mapsto \gamma(-1)$, and it is homotopic to $i \circ \pi$. One defines $\text{hofib}(i)$ as its fiber, i.e.,

$$\text{hofib}(i) := \left\{ (f, \gamma) : f \in \mathcal{L}_{p, q}^n, \gamma : [-1, 1] \rightarrow \mathcal{P}_{p, q}^n, \gamma(-1) = e, \gamma(1) = i(f) \right\}.$$

The homotopy equivalence $p : \text{hofib}(i) \rightarrow \Omega \mathcal{L}_{p-1, q-1}^{n-1}$ is given by $(f, \gamma) \mapsto \rho \circ \gamma$, and for h homotopy inverse to p , we can define $\partial := \pi \circ j \circ h$.

Lemma 4.1. *The graphing map $G : \Omega \mathcal{L}_{p-1, q-1}^{n-1} \rightarrow \mathcal{L}_{p, q}^n$ agrees with ∂ .*

Proof. We will specify a map $h : \Omega \mathcal{L}_{p-1, q-1}^{n-1} \rightarrow \text{hofib}(i)$ homotopy inverse to p so that with $\partial = \pi \circ j \circ h$, we get $G = \partial$. We thus need to construct out of a loop of long links a path of pseudoisotopy embeddings in one dimension higher, which starts at the standard embedding e and ends at a long link. Write a map $f \in \Omega \mathcal{L}_{p-1, q-1}^{n-1}$ as

$$\begin{aligned} f &= (f_1, f_2) : \mathbb{R} \times (\mathbb{R}^{p-1} \sqcup \mathbb{R}^{q-1}) \rightarrow \mathbb{R}^{n-1} \\ f_1 &: (s, (t_1, \dots, t_p)) \mapsto f_1(s)(t_1, \dots, t_{p-1}) \\ f_2 &: (s, (t_1, \dots, t_q)) \mapsto f_2(s)(t_1, \dots, t_{q-1}) \end{aligned}$$

by identifying $\mathbb{R} \times (\mathbb{R}^{p-1} \sqcup \mathbb{R}^{q-1})$ with $(\mathbb{R} \times \mathbb{R}^{p-1}) \sqcup (\mathbb{R} \times \mathbb{R}^{q-1})$. We define h by the formula

$$\begin{aligned} h(f) &= (h(f)_1, h(f)_2) : [-1, 1] \times (\mathbb{R}^p \sqcup \mathbb{R}^q) \rightarrow \mathbb{R}^n \\ h(f)_1 &: (s, (t_1, \dots, t_p)) \mapsto (f_1(g(s, t_p))(t_1, \dots, t_{p-1}), t_p) \\ h(f)_2 &: (s, (t_1, \dots, t_q)) \mapsto (f_2(g(s, t_q))(t_1, \dots, t_{q-1}), t_q) \end{aligned}$$

where $g(s, t) = (s+1)\rho(t) - 1$ and $\rho : \mathbb{R} \rightarrow \mathbb{R}$ is a smooth cutoff function that is -1 for $t \leq -1/2$ and 1 for $t \geq 1/2$. We conclude the proof by noting that the function $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ satisfies the following properties:

- g is a smooth function of t for all $s \in [-1, 1]$,
- $g(-1, t) \leq -1$, so that $h(f)$ starts at e ,
- $g(s, t) \leq -1$ if $t \leq -1$, so that at every time s , $h(f)$ is a pseudoisotopy embedding (standard on the bottom face of $I^p \sqcup I^q$ and all slices of $(I^{p-1} \sqcup I^{q-1}) \times \mathbb{R}$ below it),
- $g(1, t) \geq 1$ if $t \geq 1$, so that $h(f)$ ends at a long knot,
- $g(s, t)$ is independent of t for all $t \geq 1$, so that at every time s , $h(f)$ is a pseudoisotopy embedding (given by the same long knot on the top face of $I^p \sqcup I^q$ and all slices of $(I^{p-1} \sqcup I^{q-1}) \times \mathbb{R}$ above it), and
- $g(s, 1) = s$, so that $\pi \circ j \circ h = G$.

□

Lemma 4.2. *If $0 \leq p \leq q \leq n - 3$, then the space $\mathcal{P}_{p,q}^n$ is $(2n - 2q - 5)$ -connected.*

Proof. Consider the square

$$\begin{array}{ccc} \mathcal{P}_{p,q}^n & \longrightarrow & \mathcal{P}_p^n \\ \downarrow & & \downarrow \\ \mathcal{P}_q^n & \longrightarrow & * \end{array} \quad (4.2)$$

where the maps with domain $\mathcal{P}_{p,q}^n$ are obtained by restricting an embedding to each of the two components. This square is $(2n - p - q - 4)$ -cartesian. This fact is the improved version of Morlet's disjunction lemma due to Goodwillie [Goo90b, p. 6], who further generalized this result [Goo90b, p. 12, Theorem D] [GK15, Lemma 7.4]. One can see that the statement about the square (4.2) is equivalent to Morlet's disjunction lemma by considering the connectivity of the map between the fibers of the rows or columns. Thus the map from $\mathcal{P}_{p,q}^n$ to the homotopy limit of the rest of the square is $(2n - p - q - 4)$ -connected. This homotopy limit is simply $\mathcal{P}_p^n \times \mathcal{P}_q^n$. Another result of Goodwillie implies that the space \mathcal{P}_p^n is $(2n - 2p - 5)$ -connected [Goo90b, pp. 9-10] [Goo90a, Lemma 3.16]. Thus $\mathcal{P}_{p,q}^n$ has connectivity at least $\min\{2n - p - q - 5, 2n - 2p - 5, 2n - 2q - 5\} = 2n - 2q - 5$. \square

4.2. Bijectivity of graphing and homotopy groups of spaces of long links in a range. We now calculate homotopy groups of 2-component long links, at least up to knotting, in a certain range.

Theorem 4.3. *Assume $1 \leq p \leq q \leq n - 3$ and $0 \leq i \leq 2n - p - 2q - 4$.*

(a) *We have*

$$\pi_i \mathcal{L}_{p,q}^n \cong \pi_{i+p} S^{n-q-1} \oplus \pi_0 \mathcal{K}_{i+q}^{i+n}. \quad (4.3)$$

(b) *In the sequence of maps*

$$\pi_{i+p} \mathcal{L}_{0,q-p}^{n-p} \xrightarrow{G_*} \pi_{i+p-1} \mathcal{L}_{1,q-p+1}^{n-p+1} \xrightarrow{G_*} \dots \xrightarrow{G_*} \pi_i \mathcal{L}_{p,q}^n \xrightarrow{G_*} \dots \xrightarrow{G_*} \pi_0 \mathcal{L}_{i+p,i+q}^{i+n} \quad (4.4)$$

induced by graphing, each map except possibly the first is an isomorphism. The first map is always a surjection, and it is an isomorphism if $i \leq 2n - p - 2q - 5$ or $p = q$.

(c) *The inclusion of the summand $\pi_{i+p} S^{n-q-1}$ in the direct sum (4.3) is equal to the composite*

$$\pi_{i+p} S^{n-q-1} \cong \pi_{i+p} (\mathbb{R}^{n-p} - \mathbb{R}^{q-p}) \hookrightarrow \pi_{i+p} \mathcal{L}_{0,q-p}^{n-p} \xrightarrow{G_*^p} \pi_i \mathcal{L}_{p,q}^n. \quad (4.5)$$

The projection onto the summand $\pi_0 \mathcal{K}_{i+q}^{i+n}$ in the direct sum (4.3) is equal to the i -fold graphing map followed by restriction to the second component:

$$\pi_i \mathcal{L}_{p,q}^n \cong \pi_0 \Omega^i \mathcal{L}_{p,q}^n \xrightarrow{G_*^i} \pi_0 \mathcal{L}_{i+p,i+q}^{i+n} \longrightarrow \pi_0 \mathcal{K}_{i+q}^{i+n}$$

Naively, one might expect a term coming from the knotting of the \mathbb{R}^p component in the direct sum (4.3), but the assumed inequalities imply that $i \geq 2n - 3p - 4$, so a result of Budney [Bud08, Proposition 3.9] implies that $\pi_i \mathcal{K}_p^n = 0$ and likewise, an earlier result of Haefliger [Hae66] which it generalizes implies that $\pi_0 \mathcal{K}_{i+q}^{i+n} = 0$. For the same reason, $p = q$ is a sufficient condition to guarantee that $\pi_0 \mathcal{K}_{i+q}^{i+n} = 0$.

The inequality involving i , p , q , and n is precisely the condition that $\pi_{i+p} S^{n-q-1}$ lies within the stable range of the homotopy groups of S^{n-q-1} . Of course, i must satisfy $n - p - q - 1 \leq i$ for $\pi_{i+p} S^{n-q-1} \neq 0$.

Proof. We will first prove the connectivity of the graphing maps G claimed in part (b), except for the case where $p = q$ and $i = 2n - p - 2q - 4$, which we will address at the end of the proof. The maps of spaces underlying the sequence (4.4) are

$$\Omega^{i+p} \mathcal{L}_{0,q-p}^{n-p} \rightarrow \Omega^{i+p-1} \mathcal{L}_{1,q-p+1}^{n-p+1} \rightarrow \dots \rightarrow \Omega^i \mathcal{L}_{p,q}^n \rightarrow \dots \rightarrow \Omega \mathcal{L}_{i+p-1,i+q-1}^{i+n-1} \rightarrow \mathcal{L}_{i+p,i+q}^{i+n}. \quad (4.6)$$

We are interested in the maps they induce on π_0 . By Lemma 4.1, each such graphing map fits into a fibration

$$\Omega^{j+1} \mathcal{L}_{i+p-j-1, i+q-j-1}^{i+n-j-1} \rightarrow \Omega^j \mathcal{L}_{i+p-j, i+q-j}^{i+n-j} \rightarrow \Omega^j \mathcal{D}_{i+p-j, i+q-j}^{i+n-j} \quad (4.7)$$

where $0 \leq j \leq i+p-1$. By Lemma 4.2, the connectivity c of the base space in this case satisfies

$$c \geq (2(i+n-j) - 2(i+q-j) - 5) - j = 2n - 2q - j - 5 \geq 2n - i - p - 2q - 4$$

If $i \leq 2n - p - 2q - 5$, then $c \geq 1$. Thus each map in the sequence (4.6) is 1-connected and hence induces isomorphisms on π_0 .

If $i = 2n - p - 2q - 4$, then $j = i + p - 1$ yields $c \geq 0$, while $j \leq i + p - 2$ yields $c \geq 1$. Hence in this case, the first map in the sequence (4.6) induces a surjection in π_0 and the remaining maps induce isomorphisms on π_0 . Thus part (b) is proven, apart from the exceptional case of equidimensional source manifolds.

For part (a), the restriction fibration (2.3) from long links to long knots and its section (up to homotopy) yield the decomposition

$$\pi_i \mathcal{L}_{p,q}^n \cong \pi_i \text{Emb}_c(\mathbb{R}^p, \mathbb{R}^n - \mathbb{R}^q) \oplus \pi_i \mathcal{K}_q^n. \quad (4.8)$$

There are similar direct sum decomposition of each homotopy group in the sequence (4.4) obtained by replacing p, q , and n by the appropriate dimensions. In particular, the first group is

$$\begin{aligned} \pi_{i+p} \mathcal{L}_{0, q-p}^{n-p} &\cong \pi_{i+p} \text{Emb}_c(\{*\}, \mathbb{R}^{n-p} - \mathbb{R}^{q-p}) \oplus \pi_{i+p} \mathcal{K}_{q-p}^{n-p} \\ &\cong \pi_{i+p} S^{n-q-1} \oplus \pi_{i+p} \mathcal{K}_{q-p}^{n-p} \end{aligned}$$

Each graphing map G_* preserves the direct sum decomposition and hence can be written as $G_* = (G'_*, G''_*)$. The properties established for G_* in part (b) also apply to G'_* and G''_* . So for all values of i under consideration, the first instance of G'_* is surjective and the remaining ones are isomorphisms. By Theorem 3.1, the first instance of G'_* is injective, so every map G'_* is an isomorphism. The composition of the first p instances of G'_* map $\pi_{i+p} S^{n-q-1}$ isomorphically onto the first summand of the direct sum (4.8), thus establishing a decomposition

$$\pi_i \mathcal{L}_{p,q}^n \cong \pi_{i+p} S^{n-q-1} \oplus \pi_i \mathcal{K}_q^n. \quad (4.9)$$

where the inclusion of the first summand into $\pi_i \mathcal{L}_{p,q}^n$ is equal to the composite (4.5).

The last group in the sequence (4.4) is

$$\pi_0 \mathcal{L}_{i+p, i+q}^{i+n} \cong \pi_0 \text{Emb}_c(\mathbb{R}^{i+p}, \mathbb{R}^{i+n} - \mathbb{R}^{i+q}) \oplus \pi_0 \mathcal{K}_{i+q}^{i+n}. \quad (4.10)$$

Therefore, to prove part (a), it suffices to check that the composition $\pi_i \mathcal{K}_q^n \rightarrow \pi_0 \mathcal{K}_{i+q}^{i+n}$ of the last i instances of G''_* is an isomorphism. From the direct sum decompositions (4.8), (4.9), and (4.10) and the fact that graphing preserves them, part (c) will be proven as well. In turn, it suffices to check that each of the last i instances of G_* in the sequence (4.4) are isomorphisms. But because $p \geq 1$, the last i instances of G_* do not include the first instance, hence by part (b) are all isomorphisms.

Finally, if $p = q$, then the first instance of G_* is G'_* , hence an isomorphism by Theorem 3.1 as previously noted. This completes the proof of the exceptional equidimensional case in part (b). \square

One can easily extend Theorem 4.3 to the case where $0 = p \leq q \leq n - 2$, while still assuming $0 \leq i \leq 2n - p - 2q - 4$. In that case, the fibration $\mathcal{L}_{0,q}^n \rightarrow \mathcal{K}_q^n$ with a section, as discussed after the map (2.3), implies that $\pi_i \mathcal{L}_{0,q}^n \cong \pi_i S^{n-q-1} \oplus \pi_i \mathcal{K}_q^n$.

We will see in Remark 5.6 that if $p = q$, the range of values for i in Theorem 4.3 is sharp, even if one ignores knotting. That is, the map $\pi_{2n-3p-3} \Omega^p S^{n-p-1} \rightarrow \pi_{2n-3p-3} \mathcal{L}_{p,p}^n$ does not surject even onto the factor that does not come from knotting. However, the sequence of graphing maps involving $\pi_{2n-3p-3} \mathcal{L}_{p,p}^n$

consists of isomorphisms if one starts with 2-dimensional instead of 0-dimensional links, as we will see in Theorem 4.10.

Remark 4.4 (Isotopy classes of links). Lemma 4.8 below implies that for $i = 0$, Theorem 4.3 reduces to the result of Haefliger [Hae67] and M. Skopenkov [Sko09]. Their result applies in a larger range, namely $3n - 2p - 2q - 6 \geq 0$ (roughly, the quadruple-point-free range), in which the group is a direct sum of four factors. There is one factor for knotting of each of the components, a third factor $\pi_p S^{n-q-1}$, and a fourth factor $\pi_{p+q+2-n} V_M(\mathbb{R}^{M+n-p-1})$ where $V_M(\mathbb{R}^{M+k})$ is the Stiefel manifold of M -frames in \mathbb{R}^{M+k} and M is large enough for this group to be stable (or more precisely, $M \geq 2p + q + 5 - 2n$). In the range $2n - p - 2q - 4 \geq 0$ of Theorem 4.3, there is no knotting of the \mathbb{R}^p component, and the homotopy group of the Stiefel manifold is zero.

Remark 4.5 (Spaces of spherical links). One could ask whether Theorem 4.3 describes homotopy groups of the space of spherical links in the range $i \leq 2n - p - 2q - 4$, or perhaps some smaller range, via the closure map $\text{Emb}_c(\mathbb{R}^p \sqcup \mathbb{R}^q, \mathbb{R}^n) \rightarrow \text{Emb}(S^p \sqcup S^q, S^n)$. A result of Budney [Bud08] suggests that the closure map has connectivity at most $n - q - 2$, though the lack of a section $\text{Emb}(S^p, S^n) \rightarrow \text{Emb}(S^q \sqcup S^q, S^n)$ (as discussed before Definition 3.2) makes a proof via that result currently elusive. The proof of Lemma 4.8 suggests that this connectivity could be even lower. We do not pursue this question beyond isotopy classes, which are addressed in Lemma 4.8.

The next result implies that in the range assumed in Theorem 4.3, all elements of homotopy groups of spaces of long links come from 2-component links and knots. Recall that ι_S is a section (up to homotopy) of the restriction map ρ_S , described after the map (2.3).

Theorem 4.6. *Suppose that $1 \leq p_1 \leq \dots \leq p_m \leq n - 3$ and $0 \leq i \leq 2n - 2p_{m-1} - p_m - 4$. Then every class in $\pi_i \mathcal{L}_{p_1, \dots, p_m}^n$ is in the image of $(\iota_S)_*$ for some $S \subset \{1, \dots, m\}$ with $|S| \leq 2$.*

Proof. We proceed by induction on m , with $m = 2$ as the basis case. Suppose we know the theorem to be true for links with $m - 1$ components. The restriction fibration

$$\text{Emb}_c \left(\mathbb{R}^{p_1}, \mathbb{R}^n - \left(\prod_{k=2}^m \mathbb{R}^{p_k} \right) \right) \rightarrow \mathcal{L}_{p_1, \dots, p_m}^n \rightarrow \mathcal{L}_{p_2, \dots, p_m}^n$$

has a section up to homotopy, so by the induction hypothesis, it suffices to prove the theorem for the fiber. We use the sequence of graphing maps

$$\Omega^{p_1} \text{Emb}_c \left(\mathbb{R}^0, \mathbb{R}^{n-p_1} - \left(\prod_{k=2}^m \mathbb{R}^{p_k-p_1} \right) \right) \rightarrow \dots \rightarrow \text{Emb}_c \left(\mathbb{R}^{p_1}, \mathbb{R}^n - \left(\prod_{k=2}^m \mathbb{R}^{p_k} \right) \right). \quad (4.11)$$

Now $\mathbb{R}^{n-p_1} - \left(\prod_{k=2}^m \mathbb{R}^{p_k-p_1} \right)$ is homeomorphic to a boundary connected sum

$$(S^{n-p_2-1} \times \mathbb{R}^{p_2-p_1+1}) \natural \dots \natural (S^{n-p_m-1} \times \mathbb{R}^{p_m-p_1+1})$$

so the leftmost space in the sequence (4.11) is equivalent to $\Omega^{p_1} \left(\bigvee_{k=2}^m S^{n-p_k-1} \right)$. By the Hilton–Milnor theorem [Hil55] and the upper bound on i , any element of π_i of this space is in the image of the inclusion of some summand S^{n-p_k-1} . (A Whitehead product of length 2 lies in π_i with $i \geq 2n - 2p_m - 3 - p_1$.)

It remains to show that the maps in the sequence (4.11) induce isomorphisms on π_i for any i in the assumed range. An analogue of Lemma 4.1 shows that each map is the inclusion of the fiber in a fibration of the following form, analogous to the fibration (4.7), with $0 \leq j \leq p_1 - 1$:

$$\Omega^j \text{Emb}_c \left(\mathbb{R}^{p_1-j}, \mathbb{R}^{n-j} - \left(\prod_{k=2}^m \mathbb{R}^{p_k-j} \right) \right) \rightarrow \Omega^j \text{PEmb} \left(\mathbb{R}^{p_1-j}, \mathbb{R}^{n-j} - \left(\prod_{k=2}^m \mathbb{R}^{p_k-j} \right) \right) \quad (4.12)$$

The space $\text{PEmb}(\dots)$ above is defined as in Definition 2.3, replacing \mathbb{R}^n by the complement of the standard long components (e_2, \dots, e_m) . A result of Goodwillie [Goo90b, pp. 9-10] says that if $i \leq 2n - 2p_1 - 5$, then π_i of this pseudoisotopy embedding space is $\pi_{i+p_1+2-n}^S(\Omega N)$, where $N = \mathbb{R}^{n-j} - (\coprod_{k=2}^m \mathbb{R}^{p_k-j})$. (This result was applied to $N = \mathbb{R}^n$ in the proof of Lemma 4.2.) Since N is homotopy equivalent to a wedge of spheres of dimension at least $n - p_m - 1$, we get that ΩN is $(n - p_m - 3)$ -connected. Thus the connectivity of the space $\Omega^j \text{PEmb}(\dots)$ in (4.12) is at least $c = 2n - p_1 - p_m - 5 - j$.

For a graphing map with $j \leq p_1 - 2$, we have $c \geq 2n - 2p_1 - p_m - 3$, so we get isomorphisms on π_i for all i as assumed. For the first graphing map where $j = p_1 - 1$, we have $c \geq 2n - 2p_1 - p_m - 4$, so we get isomorphisms on π_i for $i \leq 2n - 2p_1 - p_m - 5$ and a surjection for $i = 2n - 2p_1 - p_m - 4$. But using Theorem 3.1, an appropriate projection, and the fact that a class in this dimension i must come from a single sphere, we deduce that the first graphing map is also injective on this homotopy group, hence an isomorphism. (Note that even though these last few inequalities feature p_1 , we needed p_{m-1} rather than p_1 in the inequality assumed on i to be able to apply the induction hypothesis.) \square

Remark 4.7. One can approach Theorem 4.3 by arguing as in the proof of Theorem 4.6, first splitting off knots and then considering pseudoisotopy fibrations, rather than vice-versa. One then deduces that on the factor complementary to knots, the graphing maps in the sequence (4.4) are isomorphisms for all i at most $2n - 2p - q - 4$, rather than $2n - p - 2q - 4$.

4.3. Spherical links and a further calculation for long links. Theorem 4.10, the last result of this Section, extends the range of Theorem 4.3, provided we start with 2-dimensional links. The graphing sequence ends with isotopy classes of long links, so we first digress to relate those to isotopy classes of spherical links, which are known in the range we will consider.

Lemma 4.8. *If $1 \leq p \leq n - 3$, then the closure map*

$$\hat{\cdot} : \mathcal{L}_{m,p}^n \rightarrow \text{Emb}\left(\coprod_1^m S^p, S^n\right)$$

induces a bijection on path components.

Proof. We first check that closure induces a surjection on path components. Given a spherical embedding f , let f_+ be its restriction to $\coprod_1^m D^p$, where D^p is the upper hemisphere of S^p . Find an isotopy $F_+ = F_+(x, t)$ of f_+ to the embedding used to construct the closure of a long link (called $g = (g_1, \dots, g_m)$ in Definition 2.5). By the isotopy extension theorem, F_+ extends to an isotopy F of spherical embeddings starting at f . Then $F(-, 1)$ is in the image of the closure map and is isotopic to f , so $\hat{\cdot}$ is surjective on π_0 .

For injectivity, we will use an intermediate space, which we will show has the same path components as $\mathcal{L}_{m,p}^n$. Fix a diffeomorphism $\varphi : \coprod_1^m D^p \rightarrow (\coprod_{i=1}^m \{(t_i^*, 0^{n-p-1})\} \times D^p)$, where each summand of D^p in the domain is the upper hemisphere of S^p . We define the space $\text{DbEmb}(\coprod_1^m S^p, S^n)$ of **disk-based embeddings** as the space of pairs (f, g) where $f \in \text{Emb}(\coprod_1^m S^p, S^n)$ and $g : D^n \rightarrow S^n$ is a smooth, orientation-preserving embedding such that

- $g\varphi$ is the restriction of f to $\coprod_1^m D^p$, and
- $g(D^n) \cap \text{im} f = f(\coprod_1^m D_i^p)$.

We topologize it as a subspace of $\text{Emb}(\coprod_1^m S^p, S^n) \times \text{Emb}(D^n, S^n)$. This definition is motivated by the d -based links used by Habegger and Lin [HL90, Definition 2.1] for link maps with $p = 1$ and $n = 3$.

We start by establishing a bijection on isotopy classes between disk-based embeddings and long links. The projection $(f, g) \mapsto g$ to the second factor is a fibration

$$\mathcal{L}_{m,p}^n \rightarrow \text{DbEmb}\left(\coprod_1^m S^p, S^n\right) \rightarrow \text{Emb}^+(D^n, S^n) \quad (4.13)$$

where $\text{Emb}^+(D^n, S^n)$ is the space of orientation-preserving embeddings $D^n \rightarrow S^n$. (The fact that this projection and (4.14) below satisfy the homotopy lifting property can be seen by lifting a homotopy to the space of embeddings of the n -manifold with boundary obtained from a neighborhood of $g(D^n) \cup f(\coprod_1^m S^p)$.) The inclusion of the fiber over g is $f \mapsto (\hat{f}, g)$, where \hat{f} is the closure obtained via the images of D^p under g . The base space is homotopy equivalent to $SO(n+1)$ by shrinking and linearizing. In particular, it is path-connected.

It now suffices to check that the boundary map $\pi_1 SO(n+1) \rightarrow \pi_0 \mathcal{L}_{p,m}^n$ in the long exact sequence of the fibration (4.13) for is trivial. Find a neighborhood of $g(D^n)$ homeomorphic to $D^{n-1} \times I$ such that the induced inclusion $D^n \rightarrow D^{n-1} \times I$ is the standard one. A generator of $\pi_1 SO(n+1)$ in this context can be represented by a loop of rotations of the D^{n-1} factor through an angle of 2π . The effect on a long link f in the fiber is to send it to (the isotopy class of) the long link obtained by rotating the factor of I^n in $I^n = I^{n-1} \times I$ by 2π , that is, to f itself. Thus the action is trivial on π_0 , so the inclusion of the fiber in (4.13) is a bijection on π_0 .

We now connect to isotopy classes of spherical links. The projection $(f, g) \mapsto f$ of a disk-based embedding to the first factor is a fibration

$$\text{DbEmb}\left(\coprod_1^m S^p, S^n\right) \rightarrow \text{Emb}\left(\coprod_1^m S^p, S^n\right). \quad (4.14)$$

We will show that it is injective on π_0 by showing that its fibers are path connected. Since the closure map factors through the map (4.14), this will establish injectivity on π_0 of the closure map.

Because $n - p \geq 3$, we know that π_0 of the base space in (4.14) is a group [Hae67], so its path components are homotopy equivalent to each other. Therefore it suffices to consider the fiber F over a standard trivial link. By considering a small neighborhood of the interval $[t_1^*, t_m^*] \times \{0^{n-1}\} = \bigcup_{i=2}^m [t_{i-1}^*, t_i^*] \times \{0^{n-1}\}$ in D^n , we see that F is homotopy equivalent to the space of framed embeddings of $m-1$ intervals with interiors lying in $M := D^n - (\coprod_1^m S^p)$ and prescribed values and p -frames at the endpoints. In turn, F fibers over the space E of unframed such embeddings in M , with fiber \tilde{F} given by the space of such framings. By a linearization argument again, $\pi_0 \tilde{F}$ is $(\pi_1(O(n-1), O(n-p-1)))^m$, which is trivial since $n-p \geq 3$.

It remains to show that E is path-connected. Note that M has a handle decomposition with one n -disk and m handles of index $n-p-1$. Let $A \in E$. Since $n-p-1 \geq 2$, we can perform an isotopy of each sub-arc of A that lies in a handle, fixing its endpoints, so that it ends up in the n -disk. Since $n \geq 4$, we can take each such isotopy to be an isotopy of A , i.e., we can avoid self-intersections. By a similar argument using that $n \geq 4$, we can find a further isotopy to a fixed standard arc in the n -disk. Thus E is path-connected. Hence so is F , so the map (4.14) is injective on π_0 , and the closure map is bijective on π_0 . \square

The work of Budney [Bud08, Proposition 3.9] gives a stronger analogue of Lemma 4.8 for \mathcal{K}_p^n , namely that the closure map is $(n-p-2)$ -connected.

Remark 4.9 (Closure of classical long links). For $p = 1$ and $n = 3$, the closure map is not injective on isotopy classes for $m \geq 2$. For example, long links differing by conjugation by a pure braid have the same closure. In the proof above, we would accordingly have $\pi_0 E \neq \{*\}$, i.e., multiple isotopy classes of based embeddings of arcs in $S^3 - (\coprod_1^m S^1)$, including for $m \geq 3$ distinct classes lying in the image of a 2-disk. (For such an embedding of arcs $A_1 \cup \dots \cup A_{m-1} = A$ in $D^2 - \{q_1, \dots, q_m\}$ with $m \geq 3$, a corresponding pure braid β can be found by “thickening” A_i to an unbased loop L_i enclosing both of its endpoints. Then take β to be any preimage in $\text{Aut}(F_m)$ of the element of $\text{Out}(F_m)$ that sends $x_i x_{i+1}$ to L_i .) For $m = 2$, conjugation by pure braids is trivial, but the failure of injectivity of closure can be seen from the fact that $\pi_0 \tilde{F}$ would be nontrivial. One need only look at long links with 2 and 3 crossings to find an example of two non-isotopic long links with isotopic closure that arises in this way.

Theorem 4.10. *Suppose $1 \leq p \leq n - 3$, and consider the sequence of maps induced by graphing:*

$$\pi_{2n-3p-3} \mathcal{L}_{p,p}^n \rightarrow \pi_{2n-3p-4} \mathcal{L}_{p+1,p+1}^{n+1} \rightarrow \dots \rightarrow \pi_0 \mathcal{L}_{2n-2p-3, 2n-2p-3}^{3n-3p-3}$$

If $p \geq 2$, then all the maps are isomorphisms. If $p = 1$, then the first map is surjective, and the remaining maps are isomorphisms. If $p \geq 2$, these groups are isomorphic to

$$\begin{aligned} \mathbb{Z}^3 \oplus \pi_{2n-2p-3} S^{n-p-1} & \quad \text{if } n-p \text{ is odd} \\ (\mathbb{Z}/2)^3 \oplus \pi_{2n-2p-3} S^{n-p-1} & \quad \text{if } n-p \text{ is even} \end{aligned}$$

Conjecture 5.7 is that one gets isomorphism in this set of degrees if one starts with 1-dimensional links in an even-dimensional Euclidean space.

Proof. We use the fibration

$$\Omega \mathcal{L}_{p+j-1, p+j-1}^{n+j-1} \rightarrow \mathcal{L}_{p+j, p+j}^{n+j} \rightarrow \mathcal{D}_{p+j, p+j}^{n+j}$$

where $1 \leq j \leq 2n - 3p - 3$. The pseudoisotopy embedding space $\mathcal{D}_{p+j, p+j}^{n+j}$ is $(2n - 2p - 5)$ -connected. Thus

$$\pi_i \mathcal{L}_{p+j-1, p+j-1}^{n+j-1} \rightarrow \pi_{i-1} \mathcal{L}_{p+j, p+j}^{n+j}$$

is an isomorphism if $i \leq 2n - 2p - 5$, which is at most $2n - 3p - 3$ if $p \geq 2$ and equal to $2n - 3p - 4$ if $p = 1$. If $p = 1$, the first map in the sequence is surjective because $2n - 3p - 3 = 2n - 2p - 4$. By Lemma 4.8, we have $\pi_0 \mathcal{L}_{2n-2p-3, 2n-2p-3}^{3n-3p-3} \cong \pi_0 \text{Emb}(\coprod_1^2 S^{2n-2p-3}, S^{3n-3p-3})$. The proof is completed using the calculation of the latter group from Haefliger's result [Hae67, Hae66]. Each 3-fold direct sum comes from two summands of $\pi_0 \text{Emb}(S^{2n-2p-3}, S^{3n-3p-3}) \cong \pi_0 \mathcal{K}_{2n-2p-3}^{3n-3p-3}$ and one summand of $\pi_{n-p-1} V_M(\mathbb{R}^{M+n-p-1})$ where $M \geq 2$. \square

5. HOMOTOPY CLASSES IN SPACES OF LONG KNOTS AND LINKS FROM JOINING PURE BRAIDS

We now seek nontrivial elements and spanning sets of homotopy groups of long knots and links obtained by applying the joining map J to pure braids. Although Theorem 3.1 guarantees that many nontrivial torsion classes of 2-component pure braids survive as classes of 2-component long links, the proof of our last main result uses real cohomology classes. Thus we can apply it only to classes that either are or can be related to classes of infinite order.

Classes of 2-component p -dimensional pure braids in \mathbb{R}^n are elements in $\pi_i \Omega^p S^{n-p-1}$. The class of the identity (with $i = n - 2p - 1$) cannot yield a nontrivial class in the space of long knots, since \mathcal{K}_p^n is $(2n - 3p - 4)$ -connected [Bud08]. We focus on the dimension of the only other (possibly) nontrivial rational homotopy group of a sphere, namely $\pi_{2n-3p-3} \Omega^p S^{n-p-1}$. We also consider the class $[b_{21}, b_{31}] \in \pi_{2n-3p-3} \Omega^p \text{Conf}(3, \mathbb{R}^n)$, which we view as the (parametrized) long Borromean rings.

In Section 5.1, we briefly review configuration space integrals, which produce cohomology classes in spaces of links from a certain cochain complex of graphs (diagrams). We specify a handful of graph cocycles in Section 5.2. In Section 5.3, we describe how to produce dual homology classes by resolving singular links. The proof of Theorem 5.3 (a.k.a. Theorem F), given in Section 5.4, involves identifying certain such homology classes with the homotopy classes in the Theorem statement, or more precisely their images under the Hurewicz map. We discuss possible future directions in Section 5.5.

5.1. Configuration space integrals for cohomology of spaces of long 1-dimensional links. The proof of the last result will use several cohomology classes in spaces of long links that we will describe via configuration space integrals. We now digress for a brief overview of these integrals for long m -component, 1-dimensional links. Full details are given in our previous joint work [KMV13], which extended work of Cattaneo, Cotta-Ramusino, and Longoni [CCRL02] and Bott and Taubes [BT94] from knots to long links.

(Sakai and Watanabe [SW12] extended the construction to higher-dimensional long knots, but we will not use this extension.)

The classes produced by this construction are indexed by linear combinations of link diagrams, which are graphs on m oriented line segments, for example as in formulas (5.1) through (5.8) below. Such a graph Γ consists of a vertex set $V(\Gamma)$ and an edge set $E(\Gamma)$. The vertices are partitioned into sets $V_{\text{seg}}(\Gamma)$ and $V_{\text{free}}(\Gamma)$ of segment vertices and free vertices. An edge joining two segment vertices is called a chord. Diagrams where all the edges are chords, which play a somewhat special role, are called a chord diagrams. A part of one of the m segments between two vertices is called an arc. All vertices are required to have valence at least 3, where both edges and arcs count towards valence.

For a smooth manifold X , the configuration space $\text{Conf}(m, X)$ has a compactification $\text{Conf}[m, X]$ due to Fulton and Macpherson. It is a smooth manifold with corners homotopy equivalent to $\text{Conf}(m, X)$. For any graph Γ , one defines a bundle ξ_Γ over the space \mathcal{L}_{m-1}^n of m -component 1-dimensional links. To be able to compactify at infinity, one uses a version of \mathcal{L}_{m-1}^n where the standard link has all m components approach infinity in different directions. The total space of ξ_Γ is the pullback of the diagram

$$\mathcal{L}_{m-1}^n \times \text{Conf}_0 \left[V_{\text{seg}}(\Gamma), \prod_1^m \mathbb{R} \right] \rightarrow \text{Conf} [V_{\text{seg}}(\Gamma), \mathbb{R}^n] \leftarrow \text{Conf} [V(\Gamma), \mathbb{R}^n].$$

In the left-hand configuration space, the subscript 0 indicates that the segment vertices are required lie on the components and in the order given by the graph Γ . Thus the fiber of ξ_Γ over a link f is the subspace Conf_Γ of $\text{Conf}[V(\Gamma), \mathbb{R}^n]$ where the points in $V_{\text{seg}}(\Gamma)$ lie in the image of f , on certain components and in the order determined by Γ . Each pair of vertices i, j determines a map $\varphi_{ij} : \text{Conf}(V(\Gamma), \mathbb{R}^n) \rightarrow S^{n-1}$ given by the unit vector from point i to point j . Let ω be a unit volume form on S^{n-1} . One sends a graph Γ to a differential form via the map

$$I : \Gamma \mapsto \int_{\text{Conf}_\Gamma} \bigwedge_{e \in E(\Gamma)} \varphi_{ij}^* \omega$$

where the integration is along the fiber Conf_Γ of the bundle ξ_Γ over the space of links.

To orient the configuration space over which one integrates and determine the sign of the form to be integrated, one needs certain labelings on the graphs Γ , which depend on the parity of n . For n odd, one needs an ordering of the vertices and an orientation of each edge. For n even, one needs an ordering of the segment vertices and an ordering of the edges. Changing the labeling changes the integral only by a sign, so if two labeled graphs Γ and Γ' differ by a permutation σ of labels and (for n odd) r edge reversals, one sets $\Gamma = (-1)^{\text{sign}(\sigma) + r} \Gamma'$. An equivalence class of labeling is viewed as an orientation of a graph.

There is a coboundary operator δ on such graphs that makes the space of graphs into a cochain complex. On a graph Γ , it is defined as the signed sum of edge contractions

$$\delta\Gamma := \sum_e \varepsilon(e)\Gamma/e$$

over all arcs e and all edges e that are not chords or self-loops. We first define the sign $\varepsilon(e)$ for n odd. If e is an edge or arc with endpoints $i < j$, then

$$\varepsilon(e) := \begin{cases} (-1)^j & \text{if } e = (i \rightarrow j) \\ (-1)^{j+1} & \text{if } e = (i \leftarrow j) \end{cases}$$

If n is even and e is an arc with endpoints $i < j$, define $\varepsilon(e)$ as above (where the arc orientation comes from the orientation of the segments). If n is even and e is an edge, set

$$\varepsilon(e) = (-1)^{e+1+|V_{\text{seg}}(\Gamma)|}$$

where by abuse of notation e also denotes the label on this edge.

With this coboundary operator, the integration map I is a cochain map if $n \geq 4$. At the level of cohomology, I is known to be injective on the subspace of trivalent graphs [CCRL02]. It produces all of the real cohomology of the space of braids in dimension at least 4 [KKV20].

5.2. Some low-dimensional graph cocycles. We will focus on a handful of graph cocycles, which for $m = 1, 2, 3$ yield cohomology classes in dimension $2n - 6$ in the space of m -component 1-dimensional long links in \mathbb{R}^n for any $n \geq 4$.

The graphs below are oriented using the following shorthand: segment vertices are ordered first by their order on the components (with e.g. all those on the first component coming first), a free vertex is last, edges are oriented from smaller to larger label, and edges are ordered by the smallest label of an endpoint. With this convention, some of the formulas giving cocycles are the same in both parities of n except for some signs; for these classes (κ , μ , and the v_i), we list the sign for n odd above the sign for n even. For the other classes (η and λ), we separately list the formulas for the two parities (indicated by subscripts), but we later use these symbols without the subscripts to refer to both parities at once.

$$\kappa := \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} \mp \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} \quad (5.1)$$

$$\eta_{\text{odd}} := \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} - \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} \quad (5.2)$$

$$\lambda_{\text{odd}} := \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} - \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} - \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} + \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} + \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} \quad (5.3)$$

$$\eta_{\text{even}} := \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} + \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} + \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} + \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} + \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} \quad (5.4)$$

$$\lambda_{\text{even}} := \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} + \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} - \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} + \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} - \begin{array}{c} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \\ \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \end{array} \quad (5.5)$$

$$\mu := \begin{array}{c} \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \end{array} + \begin{array}{c} \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \end{array} + \begin{array}{c} \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \end{array} \mp \begin{array}{c} \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \end{array} \quad (5.6)$$

$$v_1 := \begin{array}{c} \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \end{array} \mp \begin{array}{c} \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \end{array} \quad v_2 := \begin{array}{c} \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \end{array} \mp \begin{array}{c} \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \end{array} \quad (5.7)$$

$$v_3 := \begin{array}{c} \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \end{array} \mp \begin{array}{c} \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \quad \uparrow \end{array} \quad (5.8)$$

We will use these classes to identify families of 1-, 2-, and 3-component long links. Recall that $\rho_i : \mathcal{L}_{1,1}^n \rightarrow \mathcal{K}_1^n$, $i = 1, 2$ is the restriction to the i -th component. Let $\rho_{ij} : \mathcal{L}_{1,1,1}^n \rightarrow \mathcal{L}_{1,1}^n$, $1 \leq i < j \leq 3$ be the restriction to the i -th and j -th components.

Lemma 5.1.

- (a) The space $H^{2n-6}(\mathcal{K}_1^n; \mathbb{R})$ is spanned by $I(\kappa)$.
- (b) The class $I(\eta)$ spans a subspace of $H^{2n-6}(\mathcal{L}_{1,1}^n; \mathbb{R})$ which pulls back via the graphing map $\Omega S^{n-2} \simeq \Omega \text{Conf}(2, \mathbb{R}^{n-1}) \rightarrow \mathcal{L}_{1,1}^n$ isomorphically onto $H^{2n-6}(\Omega \text{Conf}(2, \mathbb{R}^{n-1}); \mathbb{R})$. If n is even, the evaluation of $I(\eta)$ on the image of this graphing map (pre-composed with the Hurewicz map) is up to a sign the Hopf invariant of a class in $\pi_{2n-5} S^{n-2}$.
- (c) The set $\{I(\eta), I(\lambda), \rho_1^* I(\kappa), \rho_2^* I(\kappa)\}$ is a basis for $H^{2n-6}(\mathcal{L}_{1,1}^n; \mathbb{R})$.
- (d) The set $\{I(\mu), I(v_1), I(v_2), I(v_3), \rho_{12}^* I(\eta), \rho_{13}^* I(\eta), \rho_{23}^* I(\eta)\}$ is a basis for a subspace of $H^{2n-6}(\mathcal{L}_{1,1,1}^n; \mathbb{R})$ that maps isomorphically onto $H^{2n-6}(\Omega \text{Conf}(3, \mathbb{R}^{n-1}); \mathbb{R})$ via the graphing map $\Omega \text{Conf}(3, \mathbb{R}^{n-1}) \rightarrow \mathcal{L}_{1,1,1}^n$.

Proof. The definition of the coboundary operator ensures that with our conventions for graph orientations, these cochains are cocycles. They lie in a dimension which admits no nonzero coboundary map, so they represent nontrivial and moreover linearly independent cohomology classes.

Part (a) then follows from the fact that $H^{2n-6}(\mathcal{K}_1^n; \mathbb{Z}) \cong \mathbb{Z}$, which was known at least as far back as work of Turchin [Tur07].

For parts (b) and (d), Cohen and Gitler [CG02, Theorem 2.3] described $H_*(\Omega \text{Conf}(m, \mathbb{R}^{n-1}); \mathbb{Z})$ as the (free abelian) groups of horizontal chord diagrams modulo the 4T relations, where each chord has degree $n-3$. (The cohomology thus corresponds to Vassiliev invariants of pure braids.) Consider the linear combinations of link diagrams obtained by remembering only the horizontal chord diagrams in the cocycles in question. One can check by directly computing with the 4T relations that in degree $2n-6$, these chord diagrams form bases for $H^{2n-6}(\Omega \text{Conf}(m, \mathbb{R}^{n-1}))$ for $m = 2, 3$, where by abuse of notation, an element dual to a diagram is represented by the same diagram. (These classes correspond to the triple linking number and 2-fold products of pairwise linking numbers.)

From our previous joint work [KKV20, Corollary 5.21], the graphing map $\Omega \text{Conf}(m, \mathbb{R}^{n-1}) \rightarrow \mathcal{L}_{m,1}^n$ induces a surjection in cohomology, and the cocycles in question are precisely the link diagrams for classes

in $\mathcal{L}_{m,1}^n$ corresponding to elements in these bases of chord diagrams. (Roughly, the dual injection in homology is that of horizontal chord diagrams modulo 4T relations into uni-trivalent diagrams modulo STU and IHX relations. The inclusion of *all* chord diagrams modulo 4T into uni-trivalent diagrams modulo STU and IHX is an isomorphism, by work of Bar-Natan [BN95].) The fact that $I(\eta)$ is the Hopf invariant as claimed in part (b) is established in our recent joint work [KKV, Example B.2].

For part (c), work of Songhafouo Tsopméné and Turchin [STT18a] implies that

$$\dim \pi_{n-3} \mathcal{L}_{1,1}^n \otimes \mathbb{Q} = 1$$

$$\dim \pi_{2n-6} \mathcal{L}_{1,1}^n \otimes \mathbb{Q} = \begin{cases} 3 & n \text{ odd} \\ 4 & n \text{ even} \end{cases}$$

and that these are the only nonzero groups up to degree $2n - 6$.

In more detail, they describe these groups via a complex of graphs with leaves labeled by the link components. It is roughly dual the complex described in Section 5.1, except that there are no link components, only labels by them, and the graphs are required to be connected. It is bigraded by quantities called complexity (which is the Vassiliev order for trivalent graphs) and Hodge (multi-)degree. It generalizes work of Arone and Turchin [AT15] from one component to multiple components and can be used for various source and target dimensions. For source dimension 1, a class of complexity t has degree at least $t(n - 3)$ in $\mathcal{L}_{m,1}^n$. Arone and Turchin [AT15, Section 3.2, Tables A and B] list classes of complexity $t \leq 2$ for knots, and generalizing to m -component links requires simply considering all possible labelings of leaves by $\{1, \dots, m\}$. By [AT15, Theorem 3.1], some of these graphs involving only one component appear for spaces of embeddings modulo immersions, but not for spaces of embeddings. Ultimately, in $\mathcal{L}_{1,1}^n$, there is a graph of degree $n - 3$ consisting of a single edge with different labels on the endpoints, for either parity of n ; the “2-hair hedgehog” graphs in degree $2n - 6$ with the 3 possible leaf-labelings, for n odd; and the “tripod” graphs in degree $2n - 6$ with the 4 possible leaf-labelings, for n even.

(Work of Fresse, Turchin, and Willwacher [FTW] similarly implies that for $p \geq 2$, $\pi_{2n-3p-3} \mathcal{L}_{p,p}^n \otimes \mathbb{Q}$ is 4-dimensional for $n - p$ even, generated by the same tripods, but 0 for $n - p$ odd. We do not however need to use this result anywhere.)

To complete the proof of (c), since $\mathcal{L}_{1,1}^n$ is a homotopy-associative H-space, the Milnor–Moore theorem [MM65] says that its rational homology is the universal enveloping algebra on its rational homotopy. Let a denote a generator of $\pi_{n-3} \mathcal{L}_{1,1}^n \otimes \mathbb{Q}$ as well as its image in rational homology under the Hurewicz map. Under this map, the Whitehead product in homotopy corresponds, up to a sign, to the graded commutator $[-, -]$ under the Pontryagin product \cdot in homology. If n is even, then $[a, a] = \pm 2a \cdot a$, while if n is odd, $[a, a] = 0$ so $a \cdot a$ is not primitive. Thus in either parity, the rational homology (or cohomology) in degree $2n - 6$ is 4-dimensional. Hence the linearly independent set in question is a basis. \square

Thus $I(\lambda_{\text{odd}})$ must be dual (at least up to a scalar multiple) to the class represented by the 2-hair hedgehog graph with different labels on its hairs; by the STU relation, that graph corresponds to the sum of tripods appearing in λ_{odd} . We expect that each of $I(\eta_{\text{even}})$ and $I(\lambda_{\text{even}})$ is dual to the linear combination of tripods that appear in the corresponding graph cocycle, with leaves labeled by the segments on which the segment vertices lie. However, we neither prove nor need this statement.

5.3. Homology classes dual to configuration space integrals. We now describe a process for constructing a family of 1-dimensional m -component links F out of an immersion $f : \coprod_1^m \mathbb{R} \rightarrow \mathbb{R}^n$ with k double-points. As guaranteed by Lemma 5.2 below, such a family F will be dual to a certain chord diagram Γ with k chords. We assume that the tangent vectors are linearly independent at each double-point of f .

Suppose the double-points are given by $f(s_1) = f(t_1), \dots, f(s_k) = f(t_k)$. The s_i may lie in different components, as may the t_i , and we make no assumptions on the order in which they lie even if some of them do lie on the same component.

The family F is constructed as follows. The unit sphere in the orthogonal complement to the two tangent vectors at each double-point is S^{n-3} . Build a $k(n-3)$ -parameter family F of (nonsingular) m -component braids by varying the strand on which t_i lies through a normal sphere S_i^{n-3} of a small radius within a small neighborhood N_i in \mathbb{R}^n of the i -th double-point for $i = 1, \dots, k$. Somewhat more precisely, parametrize the family by $(D^{n-3})^{\times k}$ and define it using degree-1 maps $(D^{n-3}, \partial D^{n-3}) \rightarrow (S_i^{n-3}, *)$ around t_i for each i . We can arrange for its support to lie in neighborhoods of t_1, \dots, t_k , interpolating between the degree-1 maps and constant maps within these neighborhoods.

Lemma 5.2. *Let F be the family constructed out of a singular link f as just described, and let Γ be a graph. Then $\langle I(\Gamma), F \rangle$ is nonzero if and only if Γ is the chord diagram with chords corresponding exactly to the singularities in f .*

For example, if $f = (f_1, f_2)$ has two double points, one where $f_1(s_1) = f_2(t_1)$ and another where $f_1(s_2) = f_2(t_2)$, with $s_1 < s_2$ and $t_1 < t_2$, then F pairs nontrivially with the first graph in formula (5.2), but trivially with the second graph in that formula. This F would also pair trivially with all the graphs in formula (5.3), including the three chord diagrams in that formula.

Sketch of proof of Lemma 5.2. The proof is based on that of a result of Cattaneo, Cotta-Ramusino, and Longoni [CCRL02, Theorem 6.1]. First, on the space of configurations where fewer than 2 points lie in some neighborhood N_i , any integral vanishes because on that space, the variation of F through the i -th copy of S^{n-3} admits a homotopy back to the original singular link, thus yielding a degenerate family. This implies that if Γ is not a chord diagram, then $\langle I(\Gamma), F \rangle = 0$.

It remains to consider the integrals of chord diagrams on the space \mathcal{C} of configurations where each N_i contains exactly two points. The configuration points in a chord diagram can be identified by their order on the link components, so there is only one way for each N_i to contain a pair of points, i.e., \mathcal{C} is connected. If points j and k do not lie in the same N_i , then φ_{jk} restricted to \mathcal{C} is nullhomotopic. Thus for a chord diagram Γ , $\langle I(\Gamma), F \rangle \neq 0$ only if the chords of Γ correspond exactly to the singularities resolved in F .

Finally, consider a factor $\varphi_{jk}^* \omega$ in $I(\Gamma)$. Its integral over the configuration space $I \times I$ of two points on segments of the link in some N_i produces an $(n-3)$ -form. The pairing of this form with the $(n-3)$ -parameter family coming from the resolution of the i -th double point is the same as the linking number of the sphere S^{n-2} obtained from the $(n-3)$ -parameter family that varies one segment and (a closure of) the other segment. Then by construction, if the chords in Γ correspond to the resolved singularities of F , $\langle I(\Gamma), F \rangle$ is product of factors ± 1 , hence is itself ± 1 .

This argument can be made more precise by taking the limit as the diameters of the neighborhoods N_i and the spheres S^{n-3} inside them approaches zero. \square

5.4. Long Borromean rings, the Hopf map, and classes of long links and knots. In this section, we will use maps that join components of a long link. By abuse of notation, we will denote all of them J .

We first specify a map $J: \mathcal{L}_{1,1,1}^n \rightarrow \mathcal{L}_{1,1}^n$ which joins the second and third components of a 3-component link. Given $f = (f_1, f_2, f_3) \in \mathcal{L}_{1,1,1}^n$, $n \geq 3$, remove $f_2((2, \infty))$ and $f_3((-\infty, -2))$. Then join $f_2(2)$ to $f_3(-2)$ by an embedded arc in $[-1, 1] \times [-3, 3]^{n-1} - [-1, 1]^n$, and parametrize the result so that $[-1, 1]$ is mapped to the segment between $f_2(-1)$ and $f_3(1)$. Next perform isotopies of the strands $f_2((-\infty, 2])$ and $f_3([2, \infty))$, fixing the endpoint of each, so that they agree with the image of the standard component e_2 for $|t| \geq 3$.

The arcs and isotopies can be arranged to lie in $\{0^{n-3}\} \times \mathbb{R}^3$. Finally, perform a scaling by $1/3$ in every coordinate but the first so that $[-1, 1] \times [-3, 3]^{n-1}$ is taken to $[-1, 1]^n$.

We similarly specify a map $J : \mathcal{L}_{1,1}^n \rightarrow \mathcal{K}_1^n$ which joins the components of a 2-component link. Given $f = (f_1, f_2) \in \mathcal{L}_{1,1}^n$, remove $f_1((2, \infty))$ and $f_2((-\infty, -2))$, and join $f_1(2)$ to $f_2(-2)$ by an embedded arc in $[-1, 1] \times [-3, 3]^{n-1} - [-1, 1]^n$. Carry out a parametrization, isotopies away from the knotted part, and a rescaling of Euclidean space as in the previous setting to obtain an element of \mathcal{K}_1^n .

By induction on p , we can require that the joining maps $J : \mathcal{L}_{p,p,p}^n \rightarrow \mathcal{L}_{p,p}^n$ and $J : \mathcal{L}_{p,p}^n \rightarrow \mathcal{K}_p^n$ are compatible with joining maps J on $(p-1)$ -dimensional links in \mathbb{R}^{n-1} and graphing maps G from such links to p -dimensional links in \mathbb{R}^n . That is, we require that $GJ = JG$.

We are now ready to prove our last main result:

Theorem 5.3. *Suppose $1 \leq p \leq n-3$ and $2n-3p-3 \geq 0$.*

- (a) *If $n-p$ is odd, then the map $\pi_{2n-2p-3}S^{n-p-1} \rightarrow \pi_{2n-3p-3}\mathcal{K}_p^n$ given by p -fold graphing followed by joining the two link components is nontrivial. If $n-p-1 = 2, 4$, or 8 , it sends each Hopf fibration to a generator. More generally, it sends $[\mathbb{1}_{n-p-1}, \mathbb{1}_{n-p-1}]$, the Whitehead square of the identity, to twice a generator.*
- (b) *For either parity of $n-p$, the map $\pi_{2n-3p-3}\Omega^p\text{Conf}(3, \mathbb{R}^{n-p}) \rightarrow \pi_{2n-3p-3}\mathcal{K}_p^n$ given by p -fold graphing followed by joining the three components together maps the “parametrized long Borromean rings” class $[b_{21}, b_{31}]$ to a generator.*
- (c) *There is a basis for $\pi_{2n-6}\mathcal{L}_{1,1}^n$ modulo torsion consisting of*
 - *the images of a generator of $\pi_{2n-6}\mathcal{K}_1^n$ under the inclusions $\iota_1, \iota_2 : \mathcal{K}_1^n \rightarrow \mathcal{L}_{1,1}^n$,*
 - *the result of joining two components in the parametrized long Borromean rings $[b_{21}, b_{31}]$, and*
 - *for $n-p$ odd, the result of graphing $[\mathbb{1}_{n-2}, \mathbb{1}_{n-2}]$.*

If $p \geq 2$, then there is a minimal generating set for $\pi_{2n-3p-3}\mathcal{L}_{p,p}^n$ given by the two inclusions of a generator of $\pi_{2n-3p-3}\mathcal{K}_p^n$, the result of joining two components of $[b_{21}, b_{31}]$, and the image under graphing of a minimal generating set of $\pi_{2n-2p-3}S^{n-p-1}$.

Proof. The main idea of the proof is to define families of 1-dimensional m -component long links, for $m = 1, 2, 3$, by resolving singular links and joining components. The families we define will be called F and F' (for $m = 3$), H , L , and L' (for $m = 2$), and K (for $m = 1$). Via maps that join components, we will have $(F \pm F') \mapsto (L \pm L') \mapsto K$ and $H \mapsto K$, mostly by construction. Our first main task is to show using Lemmas 5.1 and 5.2 that at the level of homology, $F - F'$ is (up to a sign) the image of $[b_{21}, b_{31}]$, H is (up to a sign) the image of the Hopf map, and K is a generator of $H_{2n-6}\mathcal{K}_1^n$.

Families F and F' of 3-component 1-dimensional pure braids: Take a singular 3-strand braid $f = (f_1, f_2, f_3) : \coprod_1^3 \mathbb{R} \hookrightarrow \mathbb{R}^n$ with two double-points, given by $f_1(s_1) = f_2(t)$ and $f_1(s_2) = f_3(u)$ with $s_1 < s_2$. One can construct f by starting with a singular braid in \mathbb{R}^3 , for example as in Figure 1 and taking its image under an inclusion of coordinates $i : \mathbb{R}^3 \hookrightarrow \mathbb{R}^n$. Then construct a $(2n-6)$ -parameter family F by resolving the singularities, as described before Lemma 5.2.

Next repeat this process, but with a singular 3-strand braid $f' = (f'_1, f'_2, f'_3)$ with two double-points, given by $f'_1(s_1) = f'_3(u)$ and $f'_1(s_2) = f'_2(t)$ with $s_1 < s_2$. Let F' be the $(2n-6)$ -parameter family obtained by resolving the singularities of f' . We orient it by ordering the parameters as s_1, s_2, t, u and ordering the spheres of resolution in the order that the double-points are listed.

Note that both F and F' can be taken to be families of pure braids, rather than arbitrary long links. As a result, we can use Lemma 5.1 (d). The graphing map $\Omega\text{Conf}(3, \mathbb{R}^{n-1}) \rightarrow \mathcal{L}_{1,1}^n$ induces an injection in homology, and the dual of its image has a basis as specified there. We apply Lemma 5.2 to the cycles F and F' and to the seven graph cocycles γ in that basis. Only the first graphs in the expression for μ pairs




FIGURE 1. (a) The singular 3-strand braid $f = (f_1, f_2, f_3)$ used to build the family F . (b) The singular 3-strand braid $f' = (f'_1, f'_2, f'_3)$ used to build the family F' .

nontrivially with F , and the same is true for v_1 and F' . We conclude that

$$\begin{aligned}
 \langle I(\mu), F \rangle &= 1 & \langle I(\mu), F' \rangle &= 0 \\
 \langle I(v_1), F \rangle &= 1 & \langle I(v_1), F' \rangle &= \mp 1 \\
 \langle I(v_i), F \rangle &= 0 \text{ for } i = 2, 3 & \langle I(v_i), F' \rangle &= 0 \text{ for } i = 2, 3 \\
 \langle \rho_{ij}^* I(\eta), F \rangle &= 0 \text{ for } 1 \leq i < j \leq 3 & \langle \rho_{ij}^* I(\eta), F' \rangle &= 0 \text{ for } 1 \leq i < j \leq 3
 \end{aligned}
 \tag{5.9}$$

where as before, the sign for n odd is shown above the sign for n even. The class $[b_{21}, b_{31}]$ is known to generate $\pi_{2n-6} \Omega \text{Conf}(3, \mathbb{R}^{n-1}) \otimes \mathbb{Q}$. From our previous joint work [KKV20] [KKV], we have $\langle I(\mu), [b_{21}, b_{31}] \rangle = 1$ (possibly up to a sign) and $\langle I(\gamma), [b_{21}, b_{31}] \rangle = 0$ for all the other graph cocycles γ appearing in (5.9). Hence $F \pm F'$ (for n odd, respectively even) is homologous to $[b_{21}, b_{31}]$, as desired.

Families L and L' of 2-component 1-dimensional long links by joining components: We next define $(2n-6)$ -dimensional families L and L' of 2-component long links as the images of F and F' under the map that joins the second and third components, as described just before the Theorem statement. We can also describe L and L' as the resolutions of the singular links ℓ and ℓ' obtained by joining components of the singular links f and f' . (That is, joining components commutes with resolving singularities.) These singular links are shown in Figure 3.




FIGURE 2. The singular 2-component long link $\ell = (\ell_1, \ell_2)$ obtained by joining strand 2 to strand 3 in the singular braid $f = (f_1, f_2, f_3)$. An isotopy takes this singular long link to a singular 2-strand pure braid.




FIGURE 3. The singular 2-component long link $\ell' = (\ell'_1, \ell'_2)$ obtained by joining strand 2 to strand 3 in the singular braid $f' = (f'_1, f'_2, f'_3)$.




FIGURE 4. The singular 2-component braid $h = (h_1, h_2)$ used to construct the family H .

By Lemma 5.2

$$\begin{aligned} \langle I(\eta), L \rangle &= 1 & \langle I(\eta), L' \rangle &= \begin{cases} -1 & \text{if } n \text{ is odd} \\ 0 & \text{if } n \text{ is even} \end{cases} \\ \langle I(\lambda), L \rangle &= 0 & \langle I(\lambda), L' \rangle &= 1 \\ \langle \rho_i^* I(\kappa), L \rangle &= 0 \text{ for } i = 1, 2 & \langle \rho_i^* I(\kappa), L' \rangle &= 0 \text{ for } i = 1, 2 \end{aligned}$$

Indeed, L pairs nontrivially with only the first diagram in formulas (5.2) and (5.4) for η , and L' pairs nontrivially with only the first diagram in formulas (5.3) and (5.5) for λ .

A family H of 2-component 1-dimensional pure braids: Next, we construct a $(2n-6)$ -parameter family H of 2-component 1-dimensional long links. Start with an immersion $h = (h_1, h_2)$ which is an embedding except at two double-points, where the double-points are given by $h_1(s_1) = h_2(t_1)$ and $h_1(s_2) = h_2(t_2)$ where $s_1 < s_2$ and $t_1 < t_2$. See Figure 4. Resolve the singularities to obtain a $(2n-6)$ -parameter family H out of h . We orient it by ordering the parameters as s_1, s_2, t_1, t_2 and ordering the resolution spheres in the order that the double-points are listed. By Lemma 5.2

$$\begin{aligned} \langle I(\eta), H \rangle &= 1 \\ \langle I(\lambda), H \rangle &= 0 \\ \langle \rho_i^* I(\kappa), H \rangle &= 0 \text{ for } i = 1, 2 \end{aligned} \tag{5.10}$$

Thus $L \pm L'$ and H represent linearly independent elements of $H_{2n-6}(\mathcal{L}_{1,1}^n; \mathbb{Z})$. Note however that L and H represent the same element.

We now identify H with the image of a better known element in the homotopy groups of spheres. Recall that if n is even, then the free part of $\pi_{2n-5}(S^{n-2})$ maps isomorphically onto $H_{2n-6}(\Omega S^{n-2})$, which

via graphing injects into $H_{2n-6}(\mathcal{L}_{1,1}^n)$. Lemma 5.1 (b) and the pairings in (5.10) show that if $n - 2 = 2, 4,$ or $8,$ then H represents (up to a sign) the class of the pure braid coming from the Hopf fibration. More generally, this shows that $2H$ and $\pm[\mathbb{1}_{n-2}, \mathbb{1}_{n-2}]$ represent the same class, since the Whitehead square of an even-dimensional identity map has Hopf invariant ± 2 .

A family K of 1-dimensional long knots: Finally, we define a $(2n - 6)$ -dimensional family K of long knots in \mathbb{R}^n as the image of the family H of 2-component pure braids under the map $J : \mathcal{L}_{1,1}^n \rightarrow \mathcal{K}_1^n$ described just before the Theorem statement. As noted earlier, joining commutes with resolving singularities. Thus we can alternatively define K by applying an obvious extension of the map J to the singular link h to get a singular knot k , as shown in Figure 5(b), and then resolving the singularities of k . One could also obtain K without reference to multiple-component links, defining k directly as a singular knot prescribed by the first diagram in formula (5.1). The matches the construction of a generator of $\pi_{2n-6}\mathcal{K}_1^n$ in Budney’s work [Bud08], so K generates that group. (By Lemma 5.2, $\langle I(\kappa), K \rangle = 1$, establishing the nontriviality of K in $H_{2n-6}\mathcal{K}_1^n; \mathbb{Z} \cong \pi_{2n-6}\mathcal{K}_1^n$, but we need Budney’s construction to show that it is a generator.)




FIGURE 5. The singular knot k obtained by joining the two components of the singular 2-component braid h .

Since $H \mapsto K$ and $2H$ was identified with the Whitehead square of the identity, part (a) is proven in the case that $p = 1$. In particular, if the ambient dimension n is $4, 6,$ or $10,$ graphing the pure braid associated to the Hopf fibration gives a generator of the first nontrivial homotopy group of the space of long knots in \mathbb{R}^n .

We next prove part (b) for $p = 1$. We just have to consider the result of joining the two components in the cycle $L \pm L'$. We saw that L is homologous to H and hence maps to K , so it remains to check that the result K' of joining the components of L' pairs trivially with κ . Indeed, the singularities of K' do not coincide with the chords of κ , so by Lemma 5.1 (a) and Lemma 5.2, K' is zero in homology, and part (b) is proven for $p = 1$. A picture of K' is shown in Figure 6.

We now prove part (c) for $p = 1$. Recall that for $i = 1, 2,$ we have sections $\iota_i : \mathcal{K}_1^n \rightarrow \mathcal{L}_{1,1}^n$ of the restrictions $\rho_i : \mathcal{L}_{1,1}^n \rightarrow \mathcal{K}_1^n$ to the i -th component, as described after the map (2.3). The classes $(\iota_1)_*(K), (\iota_2)_*(K),$ and $L \pm L'$, and for n even the class H are all in the image of the Hurewicz map. Indeed, we have




FIGURE 6. The singular knot obtained by joining the two components of the singular link $\ell' = (\ell'_1, \ell'_2)$.

already identified them with (images of) elements of homotopy groups. It is easy to see that

$$\begin{aligned} \langle \rho_j^* I(\kappa), (t_i)_*(K) \rangle &= \delta_{ij} \text{ for } 1 \leq i, j \leq 2 \\ \langle I(\eta), (t_i)_*(K) \rangle &= 0 \\ \langle I(\lambda), (t_i)_*(K) \rangle &= 0. \end{aligned}$$

Thus for n odd (respectively even) these 3 (respectively 4) classes are linearly independent. Recall that $\pi_*(\mathcal{L}_{1,1}^n) \otimes \mathbb{Q}$ is known to be 3-dimensional for n odd and 4-dimensional for n even [STT18a], as discussed in the proof of Lemma 5.1 (c). Furthermore, $(t_1)_*(K)$, $(t_2)_*(K)$, and $L \pm L'$ cannot be proper multiples of other classes because each one maps to a generator of $\pi_{2n-6}\mathcal{K}_1^n$. The same is true for H because by Theorem 3.1, graphing from a sphere is a split injection. So part (c) is proven for $p = 1$.

To prove parts (a), (b), and (c) in the case of arbitrary p , we use diagram (5.11) below. All the horizontal arrows are graphing maps, and the arrows from the groups $\pi_{2n-p-3}\Omega\text{Conf}(m, \mathbb{R}^{n-1})$, $m = 2, 3$, are iterated graphing maps. Each vertical arrow is given by joining two components. As mentioned just before the Theorem statement, we have defined the joining maps for various values of p in such a way that the diagram commutes.

$$\begin{array}{ccccccc}
 \pi_{2n-3p-3}\Omega^p\text{Conf}(3, \mathbb{R}^{n-p}) & & & & & & \\
 \searrow & \xrightarrow{\quad} & & \xrightarrow{\quad} & \xrightarrow{\quad} & \xrightarrow{\quad} & \\
 & \pi_{2n-2p-4}\mathcal{L}_{1,1,1}^{n-p+1} & \rightarrow & \pi_{2n-2p-5}\mathcal{L}_{2,2,2}^{n-p+2} & \rightarrow & \dots & \rightarrow \pi_{2n-3p-3}\mathcal{L}_{p,p,p}^n \\
 & \downarrow & & \downarrow & & & \downarrow \\
 \pi_{2n-3p-3}\Omega^p\text{Conf}(2, \mathbb{R}^{n-p}) & \xrightarrow{\quad} & \pi_{2n-2p-4}\mathcal{L}_{1,1}^{n-p+1} & \rightarrow & \pi_{2n-2p-5}\mathcal{L}_{2,2}^{n+2} & \rightarrow & \dots & \rightarrow \pi_{2n-3p-3}\mathcal{L}_{p,p}^n \\
 & \downarrow & \downarrow & & \downarrow & & \downarrow & \\
 & \pi_{2n-2p-4}\mathcal{K}_1^{n-p+1} & \xrightarrow{\cong} & \pi_{2n-2p-5}\mathcal{K}_2^{n+2} & \xrightarrow{\cong} & \dots & \xrightarrow{\cong} & \pi_{2n-3p-3}\mathcal{K}_p^n
 \end{array} \tag{5.11}$$

By work of Budney [Bud08], the horizontal maps in the bottom row are isomorphisms except possibly the first one. For odd codimension $n - p$, all the groups in the bottom row are \mathbb{Z} and all the maps there

are isomorphisms. For even codimension $n - p$, the first group is \mathbb{Z} , while the remaining groups are $\mathbb{Z}/2$, and the first map is surjective while the rest are isomorphisms. By commutativity, the compositions from the groups $\pi_{2n-3p-3}\Omega^p\text{Conf}(m, \mathbb{R}^{n-p})$ to $\pi_{2n-3p-3}\mathcal{K}_p^n$ through $\pi_{2n-2p-4}\mathcal{K}_1^{n-p+1}$ are equal to the compositions through $\pi_{2n-3p-3}\mathcal{L}_{p,p}^n$ and $\pi_{2n-3p-3}\mathcal{L}_{p,p,p}^n$ respectively. Thus we obtain parts (a) and (b) in the case of arbitrary p from the case of $p = 1$.

To prove part (c) for $p \geq 2$, recall that by Theorem 4.10, $\pi_{2n-3p-3}\mathcal{L}_{p,p}^n$ is $\mathbb{Z}^3 \oplus \pi_{2n-2p-3}S^{n-p-1}$ if $n - p$ is odd and $(\mathbb{Z}/2)^3 \oplus \pi_{2n-2p-3}S^{n-p-1}$ if $n - p$ is even,

We first treat the case of $n - p$ odd. Consider the graphing map from $\pi_{2n-2p-4}\mathcal{L}_{1,1}^{n-p+1}$ to $\pi_{2n-3p-3}\mathcal{L}_{p,p}^n$. Theorem 4.10 says that it is surjective, and the domain and codomain both have rank 4 [STT18a]. The commutativity of diagram (5.11) and part (c) for $p = 1$ imply that the elements in question generate the group $\pi_{2n-3p-3}\mathcal{L}_{p,p}^n$ modulo torsion. By Theorem 3.1, graphing restricted to the torsion subgroup of $\pi_{2n-2p-3}S^{n-p-1}$ is injective, and by the calculation of its codomain in Theorem 4.10, it must surject onto the torsion subgroup of $\pi_{2n-3p-3}\mathcal{L}_{p,p}^n$. Thus the proof is complete for $n - p$ odd.

If $n - p$ is even, graphing sends $\pi_{2n-2p-3}S^{n-p-1}$ isomorphically onto a summand of $\pi_{2n-3p-3}\mathcal{L}_{p,p}^n$ by Theorem 3.1, but to zero in $\pi_{2n-3p-3}\mathcal{K}_p^n$ because it is finite and factors through $\pi_{2n-2p-4}\mathcal{K}_1^{n-p+1} \cong \mathbb{Z}$. Part (b) and diagram (5.11) imply that $[b_{21}, b_{31}]$ maps downward to the generator of $\pi_{2n-3p-3}\mathcal{K}_p^n \cong \mathbb{Z}/2$ in that diagram. The same is true of $(\iota_1)_*(K)$ and $(\iota_2)_*(K)$. So they lie in the summand complementary to the image of $\pi_{2n-2p-3}S^{n-p-1}$. Considering the images of these 3 classes under the restriction maps ρ_1 and ρ_2 shows that they generate distinct summands. A cardinality consideration shows that these 3 classes together with $\pi_{2n-2p-3}S^{n-p-1}$ generate $\pi_{2n-3p-3}\mathcal{L}_{p,p}^n$ and are a minimal generating set. \square

Remark 5.4 (Sharpness of Theorem 5.3 (a)). We cannot extend part (a) of Theorem 5.3 on graphing spheres, joining 2-component links, and knots to the case where $n - p$ is even. In that case, although the target is $\mathbb{Z}/2$ for $p \geq 2$, we deduce the failure of this extension from the facts (noted just above in the proof of part (c)) that $\pi_{2n-2p-3}S^{n-p-1}$ is finite, $\pi_{2n-2p-4}\mathcal{K}_1^{n-p+1} \cong \mathbb{Z}$, and diagram (5.11) commutes.

Remark 5.5 (Other ways of joining Borromean rings). Similar arguments show that joining a pair of the three components of the long 1-dimensional Borromean rings class $[b_{21}, b_{31}] \in \pi_{2n-6}\Omega\text{Conf}(3, \mathbb{R}^{n-1})$ in any other of the six possible ways yields either the same element of $\pi_{2n-6}\mathcal{L}_{1,1}^n$ or its negative.

Remark 5.6 (Sharpness of Theorem 4.3 for $p = q$). Theorem 5.3 (c) shows that Theorem 4.3 does not extend to $\pi_{2n-6}\mathcal{L}_{1,1}^n$, even if one works modulo knotting. Indeed, the class $L \pm L'$, obtained by joining two components of the parametrized long Borromean rings, comes from neither $\pi_{2n-5}S^{n-2}$ nor from knots.

To calculate $\pi_{2n-3p-3}\mathcal{L}_{p,p}^n$ for all p and n with $p \leq n - 3$, one would just need to calculate the torsion subgroup for $p = 1$. Theorem 3.1 gives a lower bound on this subgroup, and we suspect that it is sharp:

Conjecture 5.7. *There is an isomorphism $\pi_{2n-6}\mathcal{L}_{1,1}^n \cong \mathbb{Z}^3 \oplus \pi_{2n-5}S^{n-2}$, where the \mathbb{Z}^3 summand is generated by the three basis elements in Theorem 5.3 (c) that do not come from S^{n-2} . The graphing map $\pi_{2n-6}\mathcal{L}_{1,1}^n \rightarrow \pi_{2n-5}\mathcal{L}_{2,2}^{n+1}$ is an isomorphism if n is even, and it is given by reduction modulo 2 on the \mathbb{Z}^3 summand and the identity on the summand $\pi_{2n-5}S^{n-2}$ if n is odd.*

Conjecture 5.7 would further generalize Budney's work [Bud08] on long knots, which says that it is true for the factors in the sequence that come from knots. The next conjecture generalizes Theorem 4.6 to one degree higher in the case of equidimensional long links.

Conjecture 5.8. *Any class in $\pi_{2n-3p-3}\mathcal{L}_{m,p}^n$ comes from the images under inclusions of classes in the groups $\pi_{2n-3p-3}\mathcal{L}_{k,p}^n$ with $k \leq 2$ and the class $[b_{21}, b_{31}] \in \pi_{2n-3p-3}\mathcal{L}_{3,p}^n$.*

Remark 5.9 (Knots, links, and braids in \mathbb{R}^3 and Vassiliev invariants). The constructions in the proof of Theorem 5.3 can be applied when $n = 3$. In fact, for the specific cocycles considered, the configuration space integral map I gives link invariants in this setting. The dual families are parametrized by $(S^0)^2$, so each of the classes $F + F'$, $L + L'$, H , and K in $H_0\mathcal{L}_{m-1}^3$ is an alternating sum of four links. More generally, one can view a link with i double points as an alternating sum of 2^i links. Now $H_0(\mathcal{L}_{m-1}^3; \mathbb{R}) = \mathbb{R}[\pi_0\mathcal{L}_{m-1}^3]$ fits into the Vassiliev–Goussarov filtration

$$\mathbb{R}[\pi_0\mathcal{L}_{m-1}^3] = J_0 \supset J_1 \supset \dots$$

where J_i is the ideal generated by singular m -component links with at least i double-points. A Vassiliev invariant of type r is a link invariant which vanishes on J_{r+1} . A similar setup applies to pure braids, in which case J_i is the i -th power of the augmentation ideal in the group ring on the pure braid group.

The only invariants of type 1 are linking numbers μ_{ij} of strands i and j . There is one knot invariant of type 2, called the Casson knot invariant (corresponding to κ). The only invariants of type 2 for pure braids are triple linking numbers (corresponding to μ) and 2-fold products of pairwise linking numbers (corresponding to η and the ν_i). There is an invariant of 2-component long links of type 2 corresponding to λ . For any m the dual space $(\mathbb{R}[\pi_0\mathcal{L}_{m-1}^3]/J_3)^*$ can be given a basis consisting of the appropriate collection of these invariants.

One extends link invariants linearly to linear combinations of links. Then for any m , one can obtain a basis for $\mathbb{R}[\pi_0\mathcal{L}_{m-1}^3]/J_3$ by choosing the appropriate elements out of $F + F'$, $L + L'$, H , and K and their images under including components. The classes $F + F'$, $L + L'$ and K are equivalent to classes in $\pi_0\mathcal{L}_{m-1}^3 \subset H_0\mathcal{L}_{m-1}^3$ in this stage of the Goussarov–Vassiliev filtration. (We neglect signs in what follows.) A long trefoil is equivalent to K , as it has Casson invariant 1. A nontrivial 3-component pure braid commutator, whose closure will be the Borromean rings, is equivalent to $F + F'$, since it has the same type-2 invariants. A long Whitehead link is equivalent to $L + L'$. The class of $H \in H_0(\Omega S^1)$ is not primitive because there is of course no 2-strand pure braid (or long link) which has zero pairwise linking number but a nonzero square of its pairwise linking number. This mirrors what happens for other odd values of n .

5.5. Questions and future directions. In future work, we plan to further study the realization of classes of long links and long knots by graphing braids and joining components.

Question 5.10. (a) Can all classes in $\pi_i\mathcal{L}_{m,p}^n$ be realized by graphing and then joining components of elements of $\pi_i\Omega^p\text{Conf}(q, \mathbb{R}^{n-p})$ for some $q \geq m$? (b) If so, how does one find such a representative?

An affirmative answer to part (a) could be seen as a higher-dimensional analogue of a theorem of Alexander [Ale23, CG01] that any knot can be obtained by joining components of a pure braid. At a more refined level, it could also be seen as an analogue of a result of Stanford [Sta]. His result says that knots are equivalent in the Vassiliev filtration quotient J_0/J_r if and only if they differ by a pure braid in the r -th stage of the lower central series of the pure braid group. The result of Fresse, Turchin, and Willwacher [FTW] on embedding spaces and graph complexes suggests that the rational homology groups of spaces of knots and links in higher dimensions with codimension at least 3 can be viewed roughly as generalizations of terms J_{r-1}/J_r in the associated graded of the Vassiliev filtration, with larger values of r corresponding to homology groups in higher degrees and with the homotopy groups consisting of the primitive elements. It could be interesting to compare any potential progress on this question to the work of Kosanović [Kos] on realizing isotopy classes of 1-dimensional knots via gropes.

One could also try to build m -component links out of links with fewer than m components. The simplest such map is an inclusion that is a section to a restriction like the map (2.3). One calls an element in the image of such an inclusion a split link. Another map that increases the number of components is cabling, where parallel components are added to a knot. Clearly such maps cannot produce braids

out of knots. In the classical case, it is not hard to construct a link that is neither split, nor a cable, nor a braid. In higher dimensions, one could consider families of links in $\mathcal{L}_{1,1}^n$ obtained by split links, cables, and braids, where these classes no longer form distinct components but subspaces.

Remark 5.11. Preliminary computations similar to the ones in the proof of Theorem 5.3 suggest that in the bases given in part (c), one can use a cable of the generator in $\pi_{2n-6}\mathcal{K}_1^n$ to replace $J_*[b_{21}, b_{31}]$ for n odd. However, for n even, this cable spans the same subgroup as $[\mathbb{1}_{n-2}, \mathbb{1}_{n-2}]$ so in this parity, $J_*[b_{21}, b_{31}]$ is neither split, nor a cable, nor a braid. We omit further details here, including a precise definition of cabling at the space level, but we may pursue them in future work.

Question 5.12. *Is there a systematic decomposition of a space of links in \mathbb{R}^n , $n \geq 4$, in terms of configuration spaces, split links, cables, and some remaining subspace?*

An example of such a decomposition is the one obtained by Batelier and Ducoulombier [BD] for 2-component long links with $n = 3$ in terms of the Swiss cheese operad. (That result extended results of ours on 2-component long links [BK15, BBK15], which in turn extended a result of Budney on long knots and the little 2-cubes operad [Bud07] and Schubert’s prime decomposition of knots [Sch49].) Such a decomposition in higher dimensions could lead to constructions of explicit geometric representatives of homotopy and homology classes in those spaces of links.

REFERENCES

- [Ale23] J. W. Alexander. A lemma on systems of knotted curves. *Proc. Natl. Acad. Sci. USA*, 9(3), 1923. [34](#)
- [AT15] G. Arone and V. Turchin. Graph-complexes computing the rational homotopy of high dimensional analogues of spaces of long knots. *Ann. Inst. Fourier (Grenoble)*, 65(1):1–62, 2015. [26](#)
- [BBK15] R. Blair, J. Burke, and R. Koytcheff. A prime decomposition theorem for the string link monoid. *J. Knot Theory Ramif.*, 24(2):24 pp., 2015. [35](#)
- [BD] É. Batelier and J. Ducoulombier. On operadic actions of spaces of knots and 2-links. arXiv:2005.06036. [35](#)
- [BK15] J. Burke and R. Koytcheff. A colored operad for string link infection. *Algebr. Geom. Topol.*, 15(6):3371–3408, 2015. [35](#)
- [BN95] D. Bar-Natan. On the Vassiliev invariants. *Topology*, 34(2):423–472, 1995. [26](#)
- [BT94] R. Bott and C. Taubes. On the self-linking of knots. *J. Math. Phys.*, 35(10):5247–5287, 1994. [22](#)
- [Bud07] R. Budney. Little cubes and long knots. *Topology*, 46(1):1–27, 2007. [35](#)
- [Bud08] R. Budney. A family of embedding spaces. In *Groups, homotopy and configuration spaces*, volume 13 of *Geom. Topol. Monogr.*, pages 41–83. Geom. Topol. Publ., Coventry, 2008. [2](#), [3](#), [5](#), [15](#), [17](#), [19](#), [21](#), [22](#), [31](#), [32](#), [33](#)
- [CCRL02] A.S. Cattaneo, P. Cotta-Ramusino, and R. Longoni. Configuration spaces and Vassiliev classes in any dimension. *Algebr. and Geom. Top.*, 2:949–1000, 2002. [22](#), [24](#), [27](#)
- [CFS14] D. Crowley, S. C. Ferry, and M. Skopenkov. The rational classification of links of codimension > 2 . *Forum Math.*, 26(1):239–269, 2014. [2](#)
- [CG01] F. R. Cohen and S. Gitler. Loop spaces of configuration spaces, braid-like groups, and knots. In *Cohomological methods in homotopy theory (Bellaterra, 1998)*, volume 196 of *Progr. Math.*, pages 59–78. Birkhäuser, Basel, 2001. [34](#)
- [CG02] F. Cohen and S. Gitler. On loop spaces of configuration spaces. *Trans. Amer. Math. Soc.*, 354(5):1705–1748, 2002. [25](#)
- [FTW] B. Fresse, V. Turchin, and T. Willwacher. On the rational homotopy type of embedding spaces of manifolds in \mathbb{R}^n . arXiv:2008.08146. [2](#), [5](#), [26](#), [34](#)
- [GK15] T. G. Goodwillie and J. R. Klein. Multiple disjunction for spaces of smooth embeddings. *J. Topol.*, 8(3):651–674, 2015. [17](#)
- [Goo90a] T. G. Goodwillie. Calculus I. The first derivative of pseudoisotopy theory. *K-Theory*, 4(1):1–27, 1990. [17](#)
- [Goo90b] T. G. Goodwillie. A multiple disjunction lemma for smooth concordance embeddings. *Mem. Amer. Math. Soc.*, 86(431):viii+317, 1990. [17](#), [20](#)
- [Hae62] A. Haefliger. Differentiable links. *Topology*, 1:241–244, 1962. [3](#), [11](#)
- [Hae66] A. Haefliger. Differential embeddings of S^n in S^{n+q} for $q > 2$. *Ann. of Math. (2)*, 83:402–436, 1966. [2](#), [3](#), [17](#), [22](#)
- [Hae67] A. Haefliger. Enlacements de sphères en codimension supérieure à 2. *Comment. Math. Helv.*, 41:51–72, 1966/67. [2](#), [3](#), [11](#), [19](#), [21](#), [22](#)
- [Hat83] A. Hatcher. A proof of the Smale conjecture. *Ann. of Math. (2)*, 117(3):553–607, 1983. [11](#)

- [Hil55] P. J. Hilton. On the homotopy groups of the union of spheres. *J. London Math. Soc.*, 30:154–172, 1955. [19](#)
- [Hir76] M. Hirsch. *Differential topology*. Springer-Verlag, 1976. [12](#)
- [HK98] N. Habegger and U. Kaiser. Link homotopy in the 2-metastable range. *Topology*, 37(1):75–94, 1998. [3](#)
- [HK21] A. Havens and R. Koytcheff. Spaces of knots in the solid torus, knots in the thickened torus, and links in the 3-sphere. *Geom. Dedicata*, 214:671–737, 2021. arXiv:2008.03192. [11](#)
- [HL90] N. Habegger and X.-S. Lin. The classification of links up to link-homotopy. *J. Amer. Math. Soc.*, 3(2):389–419, 1990. [20](#)
- [KKV] R. Komendarczyk, R. Koytcheff, and I. Volić. Diagrams for primitive cycles in spaces of pure braids and string links. arXiv:2106.11441. [26](#), [29](#)
- [KKV20] R. Komendarczyk, R. Koytcheff, and I. Volić. Diagram complexes, formality, and configuration space integrals for spaces of braids. *Q. J. Math.*, 71(2):729–779, 2020. [24](#), [25](#), [29](#)
- [KMV13] R. Koytcheff, B. Munson, and I. Volić. Configuration space integrals and the cohomology of the space of homotopy string links. *J. Knot Theory Ramif.*, 22(11):73 pp., 2013. [22](#)
- [Kos] D. Kosanović. Embedding calculus and grope cobordism of knots. arXiv:2010.05120. [34](#)
- [Kos88] U. Koschorke. Link maps and the geometry of their invariants. *Manuscripta Math.*, 61(4):383–415, 1988. [13](#)
- [Kos90] U. Koschorke. On link maps and their homotopy classification. *Math. Ann.*, 286(4):753–782, 1990. [3](#)
- [Kos97] U. Koschorke. A generalization of Milnor’s μ -invariants to higher-dimensional link maps. *Topology*, 36(2):301–324, 1997. [13](#)
- [Lim64] E. L. Lima. On the local triviality of the restriction map for embeddings. *Comment. Math. Helv.*, 38:163–164, 1964. [9](#), [12](#)
- [Mil72] R. J. Milgram. On the Haefliger knot groups. *Bull. Amer. Math. Soc.*, 78:861–865, 1972. [3](#)
- [MM65] J. W. Milnor and J. C. Moore. On the structure of Hopf algebras. *Ann. of Math. (2)*, 81:211–264, 1965. [26](#)
- [MR85] W. S. Massey and D. Rolfsen. Homotopy classification of higher-dimensional links. *Indiana Univ. Math. J.*, 34(2):375–391, 1985. [3](#), [13](#)
- [Pal60] R. S. Palais. Local triviality of the restriction map for embeddings. *Comment. Math. Helv.*, 34:305–312, 1960. [9](#), [12](#)
- [Sch49] H. Schubert. Die eindeutige Zerlegbarkeit eines Knotens in Primknoten. *S.-B. Heidelberger Akad. Wiss. Math.-Nat. Kl.*, (3):57–104, 1949. [35](#)
- [Sco68] G. P. Scott. Homotopy links. *Abh. Math. Sem. Univ. Hamburg*, 32:186–190, 1968. [3](#)
- [Skoa] A. Skopenkov. High codimension links. http://www.map.mpim-bonn.mpg.de/High_codimension_links. [4](#)
- [Skob] A. Skopenkov. Knots, i.e. embeddings of spheres. http://www.map.mpim-bonn.mpg.de/Knots,_i.e._embeddings_of_spheres. [4](#)
- [Sko00] A. Skopenkov. On the generalized Massey-Rolfsen invariant for link maps. *Fund. Math.*, 165(1):1–15, 2000. [13](#)
- [Sko09] M. Skopenkov. Suspension theorems for links and link maps. *Proc. Amer. Math. Soc.*, 137(1):359–369, 2009. [2](#), [3](#), [19](#)
- [Sta] T. Stanford. Vassiliev invariants and knots modulo pure braid subgroups. arXiv:math/9805092. [34](#)
- [STT18a] P.A. Songhafou Tsopméné and V. Turchin. Euler characteristics for spaces of string links and the modular envelope of \mathcal{L}_∞ . *Homology Homotopy Appl.*, 20(2):115–144, 2018. [4](#), [26](#), [32](#), [33](#)
- [STT18b] P.A. Songhafou Tsopméné and V. Turchin. Rational homology and homotopy of high-dimensional string links. *Forum Math.*, 30(5):1209–1235, 2018. [2](#)
- [SW12] K. Sakai and T. Watanabe. 1-loop graphs and configuration space integral for embedding spaces. *Math. Proc. Cambridge Philos. Soc.*, 152(3):497–533, 2012. [23](#)
- [Tur07] V. Turchin. On the other side of the bialgebra of chord diagrams. *J. Knot Theory Ramifications*, 16(5):575–629, 2007. [25](#)
- [Zee60] E. C. Zeeman. Linking spheres. *Abh. Math. Sem. Univ. Hamburg*, 24:149–153, 1960. [11](#)
- [Zee62] E. C. Zeeman. Isotopies and knots in manifolds. In *Topology of 3-manifolds and related topics (Proc. The Univ. of Georgia Institute, 1961)*, pages 187–193. Prentice-Hall, Englewood Cliffs, N.J., 1962. [11](#)

Email address: koytcheff@louisiana.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF LOUISIANA, LAFAYETTE, LA, USA 70504