

DISCRIMINANTS AND TORIC K -THEORY

R. PAUL HORJA AND LUDMIL KATZARKOV

ABSTRACT. We discuss a categorical approach to the theory of discriminants in the combinatorial language introduced by Gelfand, Kapranov and Zelevinsky. Our point of view is inspired by homological mirror symmetry and provides K -theoretic evidence for a conjecture presented by Paul Aspinwall in a conference talk in Banff in March 2016 and later in a joint paper with Plesser and Wang.

1. INTRODUCTION.

In this note, we investigate a conjecture of Aspinwall, Plesser and Wang [APW]. Our calculations offer supporting evidence at the level of toric K -theory and clarify some aspects related to multiplicities of discriminants which play an important implicit role in the statement of the conjecture.

In the projective Calabi–Yau case, homological mirror symmetry [K1] is stated as an equivalence of the bounded derived category of coherent sheaves $D^b(X)$ on a smooth projective Calabi–Yau variety X and the derived Fukaya category $DFuk(Y)$ of the mirror Calabi–Yau variety Y . Mirror symmetry and string theory considerations about D -brane moduli spaces also predict an identification between the global structure of locally trivial families of categories $D^b(X)$ and $DFuk(Y)$ over the Kähler parameters of X and the complex parameters of Y , respectively. As a by-product of his general conjecture, Kontsevich [K2] conjectured that the action on cohomology of the group of self-equivalences of $D^b(X)$ matches the monodromy action on the cohomology of the mirror Calabi–Yau variety Y . In particular, for a smooth projective Calabi–Yau variety X , the action of the spherical twist induced by the structure sheaf of X is ”mirrored” by the monodromy action around a distinguished ”primary” component in the moduli space of complex structures of Y . While such a statement is implicit in the existing proofs of homological mirror symmetry as well as in the statement of the Strominger–Yau–Zaslow conjecture [SYZ] as identifications

at large complex/radius limits, our work is concerned with a similar global identification in the moduli spaces.

In the present work, we consider the simplified case of toric quasi-projective Calabi-Yau Deligne-Mumford stacks. The proposal by Aspinwall, Plesser and Wang generalizes Kontsevich's identification to all the irreducible components of the principal A -determinant E_A . The polynomial E_A has integer coefficients and provides a remarkable generalization of the classical discriminant and was introduced and studied by Gelfand, Kapranov and Zelevinsky [GKZ]. The APW conjecture predicts the spherical functors that correspond to the various components of the principal A -determinant. Our result (Theorem 3.5) lends further support for the conjecture by checking its consistency in terms of the ranks of the K -theory for the various toric Deligne-Mumford stacks that enter the geometrical picture, and local multiplicity of discriminants and their intersecting components. The importance of these multiplicities has been recognized in the original work of Aspinwall, Plesser and Wang [APW], and their relevance is clearly explained in the more recent paper of Kite and Segal [KS]. Such multiplicities are also featured in earlier works [AHK], [HLS]. In the purely categorical sense, they are implicit in the theory of "windows" that began with the string theoretical work of Herbst, Hori and Page [HHP], and was made rigorous by Ballard, Favero, Katzarkov [BFK] and Halpern-Leistner [HL]. Our approach to determining multiplicities is elementary algebraic (see Definition 3.6) and it uses the powerful GKZ toolbox for studying discriminants. There are well known subtleties related to the local topological picture of intersecting discriminants. A categorification of the classical braid factorization technique should offer a truly conceptual picture of the expected semiorthogonal decompositions beyond the K -theoretic point of view. For a glimpse of how such a procedure might work, see [AHK].

In this note, we do not perform any mirror monodromy calculations for the associated GKZ D -module. A true check of the APW conjecture would require such computations, but we leave this discussion for future work. However, by simply combining the results of this work with the analytic monodromy calculations from [H], [BH2], we can check the APW predictions in some cases under certain transversality assumptions, see Corollary 4.2. It is important to note that the principal A -determinant is determined by the characteristic cycle of the GKZ

D -module. We expect that the "stacky" nature of the toric geometrical context will require the use of the better behaved GKZ system [BH] and an appropriately adapted version of the topological mirror map.

2. REVIEW OF TORIC GEOMETRY AND A -DISCRIMINANTS

2.1. Toric Geometry. We briefly review the definitions of reduced toric Deligne-Mumford stacks, as developed in [BCS] and [JT]. A *stacky fan* Σ is defined as the data $(\Sigma, \{v_i\})$ where Σ is a simplicial fan in $\bar{N} = N \otimes_{\mathbb{Z}} \mathbb{Q}$, and $\{v_i\}$ $1 \leq i \leq n$, is a collection of elements in the finitely generated group N . We assume that the rays of the simplicial fan Σ are generated by the possibly non-minimal non-zero integral elements \bar{v}_i .

The set $\{v_i\}$ defines a map $\alpha : \mathbb{Z}^n \rightarrow N$ with finite cokernel. If we dualize, we get an exact sequence

$$0 \rightarrow N^\vee \rightarrow (\mathbb{Z}^n)^\vee \rightarrow N' \rightarrow K \rightarrow 0.$$

defining the Gale dual of α (see [BCS]), with K a finite abelian group. We apply the functor $\text{Hom}(\cdot, \mathbb{C}^\times)$ to get the exact sequence

$$0 \rightarrow K \rightarrow G \rightarrow (\mathbb{C}^\times)^n \rightarrow (\mathbb{C}^\times)^d \rightarrow 1,$$

where G is the algebraic group $\text{Hom}(N', \mathbb{C}^\times)$.

Consider the subset Z of \mathbb{C}^n that consists of all the points $\mathbf{z} = (z_1, \dots, z_n)$ such that the set of v_i for the zero coordinates of \mathbf{z} is contained in a cone of Σ . Then the toric DM stack X_Σ that corresponds to the stacky fan Σ is defined as the stack quotient $[Z/G]$ where Z and G are endowed with the natural reduced scheme structures. It has been shown in [BCS] that X_Σ is a Deligne-Mumford stack whose moduli space is the simplicial toric variety X_Σ . The category of coherent sheaves on $[Z/G]$ is equivalent to that of G -linearized coherent sheaves on Z , see [V, Example 7.21].

To a cone σ in Σ , one can associate a closed substack X_σ of X_Σ by looking at the quotient of N by the subgroup spanned by v_i with $\bar{v}_i \in \sigma$. The new fan Σ/σ is defined as the image of the link of σ in Σ , and the new v_i are the images of the old v_i from the link. The quotient of N may have torsion, even in the case of a torsionfree N . Therefore, the closed substack X_σ may not be reduced even when X_Σ is.

2.2. Combinatorics of A -sets. In this section, we review some concepts discussed in chapter 7 of the book [GKZ]. A finite subset $A = \{v_1, \dots, v_n\}$ of the integral lattice $N = \mathbb{Z}^d$ is called an A -**set** if the set

A generates the lattice N as an abelian group, and there exists a linear function $h : N \rightarrow \mathbb{Z}$ such that $h(v_i) = 1$, for all $i, 1 \leq i \leq n$.

For the rest of the paper, we will assume that the set

$$A = \{v_1, \dots, v_n\} \subset N = \mathbb{Z}^d$$

is an A -set. However, see Remark 3.10 below.

We will use the notations

$$\begin{aligned} Q &:= \text{conv}(A) \subset \mathbb{R}^{d-1}, \\ K &:= \sum_{1 \leq i \leq n} \mathbb{R}_{\geq 0} v_i = \mathbb{R}_{\geq 0} Q \subset N \otimes \mathbb{R} = \mathbb{R}^d, \\ S &:= \sum_{1 \leq i \leq n} \mathbb{Z}_{\geq 0} v_i = \mathbb{Z}_{\geq 0} A \subset N = \mathbb{Z}^d. \end{aligned}$$

The rational polyhedral cone K defines an affine toric variety $Y = \text{Spec } \mathbb{C}[K^\vee \cap N^\vee]$. The hyperplane condition on the set A implies that this variety has Gorenstein singularities. Moreover, any regular (i.e. projective) triangulation of the polytope Q with vertices among the elements of A induces a stacky fan structure Σ supported on the cone K . The notion of a regular triangulation will be explained below. There is a natural crepant birational morphism $\pi : X_\Sigma \rightarrow Y$ from the induced smooth toric Deligne-Mumford stack X_Σ to the toric affine Gorenstein singularity Y .

A *triangulation* of the set $A = \{v_1, \dots, v_n\}$ is a triangulation of the polytope Q such that all vertices are among the the points v_1, \dots, v_n . A marked polytope (P, B) consists of the convex polytope P and a finite subset B of P such that $P = \text{Conv}(B)$.

Definition 2.1. [GKZ, Definition 7.2.1] *Given the set $A \subset N = \mathbb{Z}^d$, a polyhedral subdivision \mathcal{P} of (Q, A) is a family of $(d-1)$ -dimensional marked polytopes $(Q_i, A_i), i = 1, \dots, l, A_i \subset A$, such that any intersection $Q_i \cap Q_j$ is a face (possibly empty) of both Q_i and Q_j ,*

$$A_i \cap (Q_i \cap Q_j) = A_j \cap (Q_i \cap Q_j),$$

and the union of all Q_i coincides with Q .

A function $f : Q \rightarrow \mathbb{R}$ is said to be \mathcal{P} -linear for the polyhedral subdivision $\mathcal{P} = (Q_i, A_i)$ if f is continuous and the restriction of f to each Q_i is affine-linear. A function $\psi : A \rightarrow \mathbb{R}$ is said to be \mathcal{P} -linearizable if there exists a \mathcal{P} -affine function $g_\psi : Q \rightarrow \mathbb{R}$ such that $g_\psi(v) = \psi(v)$ for any $v \in \cup A_i$. A continuous function $g : Q \rightarrow \mathbb{R}$ is *concave*, if for any $x, y \in Q$, we have $g(tx + (1-t)y) \geq tg(x) + (1-t)g(y)$, $0 \leq t \leq 1$. For any polyhedral subdivision $\mathcal{P} = (Q_i, A_i)$, let $C(\mathcal{P})$ denote the cone of \mathcal{P} -linearizable functions $\psi : A \rightarrow \mathbb{R}$ such that the

associated function $g_\psi : Q \rightarrow \mathbb{R}$ is concave and $g_\psi(v) \geq \psi(v)$ for all $v \in A \setminus \cup A_i$. We say that the subdivision \mathcal{P} is *regular* if the cone $C(\mathcal{P})$ has relative non-empty interior inside the linear space of \mathcal{P} -linearizable functions.

For two polyhedral subdivisions $\mathcal{P} = (Q_i, A_i)$ and $\mathcal{P}' = (Q'_j, A'_j)$, we shall say that \mathcal{P} is a *refinement* \mathcal{P}' if, for each j , the collection of (Q_i, A_i) such that $Q_i \subset Q'_j$, forms a polyhedral subdivision of (Q'_j, A'_j) . The set of polyhedral subdivisions is then naturally a partially ordered set (poset) with respect to refinement. Triangulations are minimal elements of this poset. The maximal element is the polyhedral subdivision $(Q = \text{conv}(A), A)$.

The cones $C(\mathcal{T})$ for all the regular triangulations of (Q, A) together with all the faces of these cones form a complete fan $F(A)$ in $\mathbb{R}^A = \mathbb{R}^n$ called the *secondary fan*. We choose a translation invariant volume form vol in \mathbb{R}^{n-1} (or in \mathbb{R}^n) such that elementary simplex in the lattice $N = \mathbb{Z}^n \subset \mathbb{R}^n$ has volume 1. The characteristic function of the triangulation \mathcal{T} is the function $\phi_{\mathcal{T}} : A \rightarrow \mathbb{R}$ defined by

$$\phi_{\mathcal{T}}(v) := \sum_{v \in \text{Vert}(\sigma)} \text{vol}(\sigma),$$

where the summation is over all the maximal simplices of the triangulations \mathcal{T} for which v is vertex. We set $\phi(v) = 0$ if v is not a vertex of simplex in \mathcal{T} . The *secondary polytope* $S(A)$ is the convex hull in $\mathbb{R}^A = \mathbb{R}^n$ of the functions $\phi_{\mathcal{T}}$ for all the triangulations \mathcal{T} of (Q, A) . Theorem 7.1.7 in [GKZ] shows that the secondary polytope is $n - d$ dimensional and its vertices are the characteristics functions $\phi_{\mathcal{T}}$ for all triangulations \mathcal{T} of (Q, A) . Moreover, the secondary fan $F(A)$ is the normal fan of the secondary polytope $S(A)$. We identify $\mathbb{R}^A = \mathbb{R}^n$ with its dual in the canonical way.

Consider two regular triangulations such that the corresponding vertices in the secondary polytope are joined by an edge. The two triangulations differ by a *modification* along a circuit. A *circuit* in A is a minimal dependent subset $\{v_i, i \in I\}$ with $I \subset \{1, \dots, N\}$. In particular any circuit determines a relation of the form

$$(1) \quad \sum_{i \in I_+} l_i v_i + \sum_{i \in I_-} l_i v_i = 0,$$

with $I = I_+ \cup I_-$, where the two subsets $I_+ := \{i : l_i > 0\}$ and $I_- := \{i : l_i < 0\}$ are uniquely defined by the circuit up to replacing I_+ by I_- . We will assume that the relation (1) is *primitive*, i.e. that

the integers l_i have no common prime factor. Given a circuit, one can write a (possibly) non-primitive relation by choosing

$$|l_i| := \text{vol}(\text{conv}(v_i, i \in I \setminus i)).$$

When there is no danger of confusion, we may call the index subset I a circuit. The minimizing condition in the definition of a circuit implies that the sets $\text{conv}(v_i, i \in I_+)$ and $\text{conv}(v_i, i \in I_-)$ intersect in their common interior point. The polytope determined by $v_i, i \in I$ admits exactly two triangulations $\mathcal{T}_+(I)$ and $\mathcal{T}_-(I)$ defined by the simplices $\text{conv}(\{v_j, j \in I \setminus i\})$, for $i \in I_+$, respectively $i \in I_-$.

Suppose that the regular triangulations \mathcal{T} and \mathcal{T}' of A are obtained from each other by a modification along a circuit I . We say that a subset $J \subset A \setminus I$, is *separating* for \mathcal{T} and \mathcal{T}' if, for some $i \in I$, the set of v_i , with $i \in (I \setminus i) \cup J$ is the set of vertices of a simplex (of maximal dimension) of \mathcal{T}_1 . It turns out that a separating set J has the property that $(I \setminus i) \cup J$, for $i \in I_+$, determine simplices in \mathcal{T} and $(I \setminus i) \cup J$, for $i \in I_-$, determine simplices in \mathcal{T}' .

Proposition 2.2. [GKZ, Prop 7.2.12] *The polyhedral subdivision $\mathcal{F} = \mathcal{F}(\mathcal{T}, \mathcal{T}')$ corresponding to the edge of the secondary polytope joining the vertices determined by \mathcal{T} and \mathcal{T}' consists of the simplices $(\text{conv}(K), K)$ which \mathcal{T} and \mathcal{T}' have in common and the polyhedra $(\text{conv}(I \cup J), I \cup J)$ for all separating subsets $J \subset A \setminus I$.*

2.3. Principal A -Determinants. Define ∇_A as the Zariski closure in \mathbb{C}^n of the set of polynomials $f = \sum_{1 \leq j \leq n} a_j x^{v_j}$ in $\mathbb{C}[x_1, \dots, x_d]$ such that there exists some $y \in (\mathbb{C}^\times)^n$ with the property that $f = 0$ is singular at y . By definition, the discriminant $\Delta_A \in \mathbb{Z}[a_1, \dots, a_n]$ is the irreducible polynomial (defined up to a sign) whose zero set is given by the union of the irreducible codimension 1 components of ∇_A . For the case $\text{codim} \nabla_A > 1$, one sets $\nabla_A = 1$.

For any non-empty face Γ of the polytope Q , let $i(\Gamma)$ denote the index of the sublattice $\mathbb{Z}(A \cap \Gamma)$ inside the lattice $N \cap \mathbb{R}\Gamma$,

$$(2) \quad i(\Gamma) := [N \cap \mathbb{R}\Gamma : \mathbb{Z}(A \cap \Gamma)].$$

Recall that $S = \mathbb{Z}_{\geq 0}A$ is the semigroup generated by A . Furthermore, let S/Γ denote the image semigroup of S in the quotient free group $N_{\mathbb{R}}/\mathbb{R}\Gamma$, with $N_{\mathbb{R}} = N \otimes_{\mathbb{Z}} \mathbb{R}$. The semigroup S/Γ generates a pointed cone in $N_{\mathbb{R}}/\mathbb{R}\Gamma$, so we can define

$$(3) \quad u(\Gamma) := \text{vol}(\text{conv}(S/\Gamma) \setminus \text{conv}(S/\Gamma \setminus \{0\})),$$

where the volume form on $N_{\mathbb{R}}/\mathbb{R}\Gamma$ is induced from the standard volume form on N .

Definition 2.3. [GKZ, Theorem 10.1.2] *The principal A -determinant E_A is the polynomial in $\mathbb{Z}[a_1, \dots, a_n]$ defined as*

$$(4) \quad E_A := \prod_{\Gamma} (\Delta_{A \cap \Gamma})^{u(\Gamma) \cdot i(\Gamma)},$$

where the product is taken over all the non-empty faces of the polytope $Q = \text{conv}(A)$.

For the faces Γ such that $\text{codim} \nabla_A > 1$, the corresponding factor in the product is equal to 1. Hence the product is taken over the vertices and some the faces Γ such that $A \cap \Gamma$ is not simplicial.

The principal A -determinant has the following remarkable property.

Theorem 2.4. [GKZ, Theorem 10.1.4] *For a given set A , the Newton polytope of E_A coincides with the secondary polytope $S(A)$. In particular, the vertices of the Newton polytope of E_A are given by the characteristic functions $\phi_{\mathcal{T}}$ for all the regular triangulations \mathcal{T} of (Q, A) .*

3. K -THEORY AND DISCRIMINANT INTERSECTION MULTIPLICITIES

We will now compute the dimensions of the Grothendieck rings of some toric stacks involved in the main result and conjecture discussed in this paper.

Let Γ be a non-empty face of the polytope $Q = \text{conv}(A)$. We will consider a stacky fan $\Sigma_{\Gamma} = (\Sigma_{\Gamma}, \{\pi(v_i)\}_{i \in \mathcal{I}})$ where $\pi : N \rightarrow N/\mathbb{Z}(A \cap \Gamma)$ is the canonical map, and Σ_{Γ} is a fan in the real linear space $(N/\mathbb{Z}(A \cap \Gamma)) \otimes \mathbb{R} = N_{\mathbb{R}}/\mathbb{R}\Gamma$ with rays generated by the images of the vectors v_i for $i \in \mathcal{I}$. The set \mathcal{I} consists of all indices i such that the image of v_i in $N_{\mathbb{R}}/\mathbb{R}\Gamma$ is contained in the closure of the set

$$\text{conv}(S/\Gamma) \setminus \text{conv}(S/\Gamma \setminus \{0\}),$$

where S/Γ is the image semigroup of $S = \mathbb{Z}A$ in $N_{\mathbb{R}}/\mathbb{R}\Gamma$. It is important to note that the set A and the nonempty face Γ uniquely determine the vectors $v_i, i \in \mathcal{I}$, but the cones of the fan Σ_{Γ} are not uniquely determined. However, the following statement is true.

Proposition 3.1. *For any two choices of stacky fans Σ_{Γ} and Σ'_{Γ} as above, the bounded derived categories of coherent categories $D^b(X_{\Sigma_{\Gamma}})$ and $D^b(X_{\Sigma'_{\Gamma}})$ are equivalent.*

Proof. The stacky fans Σ_{Γ} and Σ'_{Γ} are connected by a finite sequence of toric birational flops. Each of them induces an equivalence of the bounded derived categories of coherent sheaves on the corresponding DM stacks [BO], [Ka], so the result follows. \square

The rank of the Grothendieck group of the DM stack Z_Γ is given by the following result.

Proposition 3.2. *The rank of the Grothendieck group of the DM stack Z_Γ is equal to the product $u(\Gamma) \cdot i(\Gamma)$.*

Proof. Since N is a lattice, the quotient group $N/N \cap \mathbb{R}\Gamma$ is torsion free. This implies that the torsion part of the quotient $N/\mathbb{Z}(A \cap \Gamma)$ is isomorphic to the torsion part of the quotient $N \cap \mathbb{R}\Gamma/\mathbb{Z}(A \cap \Gamma)$. The order of the torsion part is given by the index $i(\Gamma)$. The discussion in section 6 of [BH1], as well as the work of Jiang-Tseng [JT], show that the rank of $K_0(Z_\Gamma)$ is equal to $K_0(Z_{\Sigma_\Gamma}) \cdot i(\Gamma)$.

In order to finish the proof, we have to show that the dimension $K_0(Z_{\Sigma_\Gamma})$ is equal to $u(\Gamma)$. Definition 3 shows that $u(\Gamma)$ is equal to the sum of the volumes of the (simplicial) maximal dimensional cones in the fan Σ_Γ . The result follows from [DKK, Proposition 3.20] which shows that the desired rank is equal to the sum of the volumes maximal dimensional cones in the fan Σ_Γ . \square

We now consider two triangulations \mathcal{T} and \mathcal{T}' corresponding to vertices of the secondary polytope which are joined by an edge F . According to Proposition 2.2, there exists a circuit $I = \{v_i\}, i \in I$, associated to this edge. Let $\mathbb{Z}I := \sum_{i \in I} \mathbb{Z}v_i$ be the sublattice generated in N by the vectors of the circuit. We now define a stacky fan $\Sigma_F = (\Sigma_F, \{\pi(v_i)\}_{i \in K})$ where $\pi : N \rightarrow N/\mathbb{Z}I$ is the canonical map, and Σ_F is the fan in the real linear space $N_{\mathbb{R}}/\mathbb{R}I$ whose maximal cones with rays generated by the images in $N_{\mathbb{R}}/\mathbb{R}I$ of $v_i, i \in J$, for all the separating sets J for the circuit I and the pair of triangulations \mathcal{T} and \mathcal{T}' . The set $\{\pi(v_i)\}_{i \in K}$ consists of images of vectors v_i which are not in I but are a vertex for one of the separating sets J . Note that all the separating sets J are simplicial. Denote by Z_F the toric DM stack defined by the stacky fan Σ_F . For the general theory of spherical functors, see [AL].

Proposition 3.3. *There exists a spherical functor $D^b(Z_F) \rightarrow D^b(X)$ where X is the toric DM stack induced by either one of the triangulations \mathcal{T} or \mathcal{T}' .*

Proof. The EZ-twist argument in [EZ] is mostly formal and can be adapted to the toric DM stacky context. The result also follows from the work of Halpern-Leistner-Shipman [HLS]. In the physics literature, the spherical twist induced by this spherical functor is known as the “wall monodromy”. \square

Proposition 3.4. *The rank of the Grothendieck group of the DM stack Z_F is equal to the sum*

$$\sum_{J\text{-sep}} [N : \mathbb{Z}(I \cup J)]$$

over all the separating sets J .

Proof. Since $I \cup J$ is a simplex, and $\mathbb{Z}I \cap \mathbb{Z}J = (0)$, we can write a non-canonical splitting for the torsion group $N/\mathbb{Z}(I \cup J)$ as a direct sum of finite torsion groups $\mathbb{Z}^k/\mathbb{Z}I \oplus \mathbb{Z}^{d-k}/\mathbb{Z}J$. Note that the index $\mathbb{Z}^{d-k}/\mathbb{Z}J$ is the volume of the cone generated by the separating set J in the fan Σ_I in $N_{\mathbb{R}}/\mathbb{R}I$ associated to the toric DM stack Z_F . As in the proof of Proposition 3.2, we conclude that the rank of the Grothendieck group of the toric DM stack is indeed the sum $\sum_J [N : \mathbb{Z}(I \cup J)]$ over all the separating sets J . \square

The main result of this note is the following theorem.

Theorem 3.5. *For any edge $F = [\phi_{\mathcal{T}}, \phi_{\mathcal{T}'}]$ of the secondary polytope, between the triangulations \mathcal{T} and \mathcal{T}' , the following equality holds:*

$$(5) \quad \text{rk}(K_0(D^b(Z_F))) = \sum_{\Gamma \subset Q} n_{\Gamma, F} \cdot \text{rk}(D^b(K_0(Z_{\Gamma}))),$$

for some non-negative integers $n_{\Gamma, F}$.

The precise combinatorial definition of the non-negative integers $n_{\Gamma, F}$ will be given below in Definition 3.6. It arises naturally as a by-product of the proof of the theorem and it is close in spirit to the string theoretical analysis in [AGM]. Geometrically, the index $n_{\Gamma, F}$ coincides in most cases with the intersection multiplicity of the discriminant component $\nabla_{\Gamma, F}$ with the rational curve induced by edge F in the DM toric stack defined by the secondary fan. This point of view been discussed in [HLS], [KS].

Proof. We will prove the result by giving two interpretations to the coefficient restriction of the principal A -determinant E_A to the edge F of the secondary polytope. By [GKZ, Prop 6.1.3], the coefficient restriction of any Laurent polynomial $P(x_1, \dots, x_n)$ to any face F of its Newton polytope has the leading term of the polynomial in t

$$t \mapsto P(t^{\psi_1} x_1, \dots, t^{\psi_n} x_n)$$

equal to $t^{\psi(F)} P_F(x_1, \dots, x_n)$, where ψ is a lineat support function for the face F , and P_F is the coefficient restriction of P to the face F .

Let $\mathcal{P} = (Q_i, A_i)$ be a polyhedral subdivision of the marked polytope (Q, A) . Theorem 1.12' in [GKZ] shows that, up to multiplication by

a constant, the coefficient restriction of E_A to the edge $F(\mathcal{P})$ of the secondary polytope equals

$$\prod_i (E_{A_i})^{[N:\mathbb{Z}A_i]} .$$

In particular, if \mathcal{F} is the polyhedral subdivision corresponding to the circuit I and the edge $[\phi_{\mathcal{T}}, \phi_{\mathcal{T}'}]$ for two triangulations $\mathcal{T}, \mathcal{T}'$ (see Proposition 2.2), then, up to multiplication by a constant, E_A is equal to

$$\prod_{i \in K} (E_{A_i})^{[N:\mathbb{Z}A_i]} \cdot \prod_{J\text{-sep}} (E_{I \cup J})^{[N:\mathbb{Z}(I \cup J)]} ,$$

where $(\text{conv}(A_i), A_i)_{i \in K}$ is the family of simplices that \mathcal{T} and \mathcal{T}' have in common, and the second product is taken over all the separating sets J such that $J \cup I$ has maximal dimension. Note that each such $J \cup I$ contains as a unique circuit the circuit I . In the terminology of [GKZ, pg 309], the set $J \cup I$ is *weakly dependent*. As such, the argument on [GKZ, pg 309] shows that, up to a multiplication by a constant, $E_{I \cup J}$ is in fact equal to the the discriminant $\Delta_{I \cup J}$, which by [GKZ, Prop 9.1.8] is a non-zero scalar multiple of the polynomial

$$(6) \quad \Delta_I := \left(\prod_{i \in I_+} l_i^{l_i} \right) \prod_{i \in I_-} a_i^{-l_i} - \left(\prod_{i \in I_-} l_i^{-l_i} \right) \prod_{i \in I_+} a_i^{l_i} ,$$

where $\sum_{i \in I_+ \cup I_-} l_i v_i = 0$ is a primitive integer relation (unique up to sign) associated to the circuit I . Note [GKZ, Prop 9.1.8] assumes that the circuit generates the ambient lattice so the choice $|l_i| := \text{vol}(\text{conv}(v_i, i \in I \setminus i))$, would work in that case. We do not make that assumption.

Consider a linear form $\psi : \mathbb{R}^A \rightarrow \mathbb{R}$ lying in the interior of cone $C(\mathcal{F})$ (the normal cone to the edge F) of the secondary fan as defined in section 2.2. This means that the edge F is the supporting face of ψ . This means that ψ is equal to a constant $\psi(F)$ along F , and $\psi(v_i) < \psi(F)$ when v_i is not on F . It follows that the coefficient of the leading term in t of the Laurent polynomial $t \mapsto E_A(\sum_i t^{\psi_i} a_i x^{v_i})$ is, up to multiplication by a constant and a monomial in the variables a_i , the product

$$(7) \quad \prod_{J\text{-sep}} (E_{I \cup J})^{[N:\mathbb{Z}(I \cup J)]} = (\Delta_I)^{\sum_J [N:\mathbb{Z}(I \cup J)]}$$

where the sum and product in the formula above are over all the separating sets J , a the discriminant polynomial Δ_I is defined by (6).

On the other hand, according to Definition 2.3 the principal A -determinant E_A is the product

$$E_A := \prod_{\Gamma} (\Delta_{A \cap \Gamma})^{u(\Gamma) \cdot i(\Gamma)},$$

where the product is taken over all the non-empty faces of the polytope $Q = \text{conv}(A)$. Due to the properties of Minkovski sums for Newton polytopes, the linear form Ψ attains its maximum value on the Newton polytope of $\Delta_{A \cap \Gamma}$ at either a vertex or an edge parallel to F . This means that, up to multiplication by a constant and a monomial in the variables a_i , the product the coefficient of the leading term in t of the Laurent polynomial $t \mapsto \Delta_{A \cap \Gamma}(\sum_i t^{\psi_i} a_i x^{v_i})$ is equal to $(\Delta_I)^{n_{\Gamma, F}}$.

Definition 3.6. *The multiplicity $n_{\Gamma, F}$ is the non-negative integer defined by the property that, up to multiplication by a constant and a monomial in the variables a_i , the coefficient of the leading term in t of the Laurent polynomial $t \mapsto \Delta_{A \cap \Gamma}(\sum_i t^{\psi_i} a_i x^{v_i})$ is equal to $(\Delta_I)^{n_{\Gamma, F}}$.*

The theorem follows as a direct consequence of the results of Propositions 3.2 and 3.4. \square

Remark 3.7. Note that $n_{\Gamma, F} = 0$, when $\Gamma \cap F$ is a vertex of Q . Furthermore, the definition shows that, if the circuit I determined by the edge F is not contained in the face Γ , then $n_{\Gamma, F} = 0$.

Remark 3.8. In practice, calculating the multiplicities $n_{\Gamma, F}$ can be computationally daunting. We will provide some examples in the next section. Note that our definition does not use the Horn uniformization of the A -discriminant (proved in this case by Kapranov [Kap]) which is a useful tool but only provides a birational parametrization of the discriminant.

The previous theorem adds to the body of evidence supporting the conjecture of Aspinwall-Plesser-Wang [APW]. The importance of multiplicities and other aspects of this conjecture have been discussed by Kite-Segal [KS].

Conjecture 3.9.

1) *For any edge F of the secondary polytope, the category $D^b(Z_F)$ admits a semiorthogonal decomposition consisting of $n_{\Gamma, F}$ components $D^b(Z_{\Gamma})$ for each face Γ of the polytope Q .*

2) *For each face Γ , there exist spherical functors $D^b(Z_{\Gamma}) \rightarrow D^b(X)$ for any toric DM stack X determined by a triangulation corresponding to a vertex of the secondary polytope.*

The second part is a direct consequence of the first: the wall monodromy functors $D^b(Z_F) \rightarrow D^b(X)$ are spherical, so result of Kuznetsov (unpublished) and Halpern-Leristner-Shipman [HLS] implies that each component of the semiorthogonal decomposition determines a spherical functor.

Remark 3.10. One of the hypotheses in the definition of an A -set is that the set A generates the lattice $N = \mathbb{Z}^d$. In particular this guarantees that the multiplicity of the "primary" component of the discriminant, associated to the face $\Gamma = Q$, has $u(\Gamma) = i(\Gamma) = 1$ and Z_Γ is a point $\text{Spec } \mathbb{C}$. This assumption is likely not essential. We could simply ask that A linearly generates the vector space $N_{\mathbb{R}} = \mathbb{R}^d$. The adapted results of this note should continue to hold, although the multiplicities involved gain the extra factor $N : \mathbb{Z}A$. On the other hand, the technical details in [GKZ] are worked out in the book for A -sets, so one would have to work out and adapt the calculations to the more general situation.

4. APPLICATIONS

Example 4.1. Recall that for a general set A in the lattice $N = \mathbb{Z}^d$, the rational polyhedral cone K generated by A defines a Gorenstein affine toric variety $Y = \text{Spec } \mathbb{C}[K^\vee \cap N^\vee]$. For any stacky resolution $\pi : X_\Sigma \rightarrow Y$ induced by the a regular triangulation of the polytope Q , the structure of the exceptional set is quite complicated. It may contain irreducible components of different dimensions. Consider an edge of the secondary polytope and let $I = I_+ \cup I_-$ be the associated circuit in A . We assume that I is maximal dimension, $\dim I = \dim C$. Let X_+ and X_- denote the coresponding toric stacky resolutions of Y that differ by a flop. In this case Z_F is the stacky point $[\text{Spec } \mathbb{C}/\mathbb{Z}_n]$ where n is the index of the sublattice generated by the circuit I in N , $n = [N : \mathbb{Z}I]$. The sets I_+ and I_- generate cones that determine the exceptional loci $E_+ \subset X_+$, and $E_- \subset X_-$, and, by Proposition 3.3, we have spherical functors $D^b(\text{Spec } \mathbb{C}/\mathbb{Z}_n) \rightarrow D^b(E_\pm) \hookrightarrow D^b(X_\pm)$. The category $D^b([\text{Spec } \mathbb{C}/\mathbb{Z}_n])$ splits into n orthogonal copies of $D^b(\text{Spec } \mathbb{C})$, so we get the spherical functors predicted by Kontsevich's conjecture described in the introduction. Theorem 3.5 implies that $n_{F,Q} = n$, where F is the edge of the secondary polytope associated to the flop. This multiplicity statement is not obvious if one attempts a direct calculation. In the transversal case $n_{F,Q} = 1$, the analytic monodromy calculations from [H], [BH2] for the classical GKZ module are valid, so we get that:

Corollary 4.2. *The wall monodromy spherical functor induced by the edge F is the spherical functor $D^b(\text{Spec } \mathbb{C}) \rightarrow D^b(E_F) \rightarrow D^b(X)$, where $E_F \subset X$ is the exceptional locus corresponding to the circuit I . The K -theory action of this functor matches the analytical continuation functor along a corresponding loop γ_F (see [BH2])*

A similar argument expands the range of cases where Kontsevich's conjecture is true beyond the cases considered in [H]. We hope to return to this issue in future work.

Example 4.3. Let X be the resolution of the A_3 singularity, with $v_0 = (1, 0), v_1 = (1, 1), v_2 = (1, 2), v_3 = (1, 3), v_4 = (1, 4)$. As it is well known, the secondary polytope is combinatorially equivalent to a cube in \mathbb{R}^3 . The principal A -determinant is

$$\begin{aligned} E_A = a_0 a_4 \Delta_Q = & a_0 a_4 (256 a_0^3 a_4^3 - 192 a_0^2 a_1 a_3 a_4^2 - 128 a_0^2 a_2^2 a_4^2 \\ & + 144 a_0^2 a_2 a_3^2 a_4 - 27 a_0^2 a_4^4 + 144 a_0 a_1^2 a_2 a_4^2 - 6 a_0 a_1^2 a_3^2 a_4 - 80 a_0 a_1 a_2^2 a_3 a_4 \\ & + 18 a_0 a_1 a_2 a_3^3 + 16 a_0 a_2^4 a_4 - 4 a_0 a_2^3 a_3^2 - 27 a_1^4 a_4^2 \\ & + 18 a_1^3 a_2 a_3 a_4 - 4 a_1^3 a_3^3 - 4 a_1^2 a_2^3 a_4 + a_1^2 a_2^2 a_3^2). \end{aligned}$$

It is clear that in this case Z_Q is the point $\text{Spec } \mathbb{C}$. Let X denote the DM stack $[\mathbb{C}^2/\mathbb{Z}_4]$ whose stacky fan has one cone and the rays generated by the vectors v_0, v_4 . Consider the edge F_1 corresponding to the birational transformation $X \leftrightarrow X_1$, where X_1 is the toric DM stack with cones determined by the pairs v_0, v_1 and v_1, v_4 . The associated polyhedral subdivision is $(\text{conv}\{0, 4\}, \{0, 1, 4\})$, and the circuit relation I is $3v_0 - 4v_1 + v_4 = 0$. The discriminant Δ_I is $256a_0^3a_4 - 27a_1^4$, and the leading term with respect to the edge F_1 in the quartic discriminant Δ_Q is

$$256a_0^3a_4^3 - 27a_1^4a_4^2 = a_4^2 \cdot \Delta_I.$$

This means that $n_{Q, F_1} = 1$, which is consistent with the fact that $Z_{F_1} = \text{Spec } \mathbb{C}$. The associated spherical functor is

$$D^b(\text{Spec } \mathbb{C}) \rightarrow D^b([\mathbb{C}^2/\mathbb{Z}_4]).$$

Let F_2 denote the edge corresponding to the birational transformation $X \leftrightarrow X_2$, where X_2 is the toric DM stack with cones determined by the pairs v_0, v_2 and v_2, v_4 . The associated polyhedral subdivision is $(\text{conv}\{0, 4\}, \{0, 2, 4\})$, and the circuit relation I is $v_0 - 2v_2 + v_4 = 0$. The discriminant Δ_I is $a_0 a_4 - 4a_2^2$, and the leading term with respect to the edge F_2 in the quartic discriminant Δ_Q is

$$256a_0^3a_4^3 - 128a_0^2a_2^2a_4^2 + 16a_0a_2^4a_4 = 16a_0a_4 \cdot \Delta_I^2.$$

This means that $n_{Q, F_2} = 2$ which is consistent with the fact that Z_{F_2} is the stacky point $[\text{Spec } \mathbb{C}/\mathbb{Z}_2]$. The derived category $D^b([\text{Spec } \mathbb{C}/\mathbb{Z}_2])$ splits into two copies of $D^b(\text{Spec } \mathbb{C})$, the associated wall monodromy spherical functor is

$$D^b([\text{Spec } \mathbb{C}/\mathbb{Z}_2]) \rightarrow D^b([\mathbb{C}^2/\mathbb{Z}_4]).$$

Example 4.4. Let X_1 be the well known quasi-projective Calabi-Yau toric variety defined as the total space of the canonical bundle of \mathbb{P}^2 . The toric structure is given by the vectors $v_0 = (0, 0, 1)$, $v_1 = (1, 0, 1)$, $v_2 = (0, 1, 1)$ and $v_3 = (-1, -1, 1)$. The lattice of linear relations is one dimensional and generated by the vector $(-3, 1, 1, 1)$, and the secondary polytope is the segment $F = [(3, 2, 2, 2), (0, 3, 3, 3)]$. There are two toric birational models in this case, namely X_1 and the stacky quotient $X_2 = [\mathbb{C}^3/\mathbb{Z}^3]$. The principal A -determinant is $E_A = a_1^2 a_2^2 a_3^2 (a_0^3 - 27a_1 a_2 a_3)$. The multiplicity of the discriminant components corresponding to v_1, v_2, v_3 are all equal to 2, and $n_{F, Q} = 1$. The spherical twist of $D^b(X_1)$ is determined by the spherical object $\mathcal{O}_{\mathbb{P}^2} \subset X_1$.

Example 4.5. Let's consider the case the Calabi-Yau toric variety X_1 is the total space of the canonical bundle of the Hirzebruch surface F_2 . The toric structure is given by the vectors $v_0 = (0, 0, 1)$, $v_1 = (1, 0, 1)$, $v_2 = (0, 1, 1)$, $v_3 = (-1, 2, 1)$ and $v_4 = (0, -1, 1)$ with the obvious cones. The lattice of relations is two dimensional and generated by the vectors $(-2, 0, 1, 0, 1)$ and $(0, 1, -2, 1, 0)$. There are four stacky toric birational models for X_1 .

The discriminant picture is obtained by studying the singularities of the Laurent polynomial $\sum a_i x^{v_i}$. Besides the vertices corresponding to v_1, v_3 and v_4 there are two other faces that contain circuits. The discriminant component determined by the face Γ generated by the vectors v_1, v_2 and v_3 is

$$f_\Gamma = a_2^2 - 4a_1 a_3.$$

For the maximal dimensional face Q we obtain the discriminant

$$f_Q = a_0^4 - 8a_0^2 a_2 a_4 + 16a_2^2 a_4^2 - 64a_1 a_3 a_4^2.$$

The principal A -determinant is

$$E_A = a_1^2 a_2^2 a_4^2 \cdot f_\Gamma \cdot f_Q.$$

If we choose the coordinates adapted to the large complex structure point

$$x := \frac{a_2 a_4}{a_0^2}, \quad y = \frac{a_1 a_3}{a_2^2}$$




FIGURE 1. The F_2 example.

the we obtain that the two relevant discriminant components are given by

$$y = \frac{1}{4}\left(1 - \frac{1}{4x}\right)^2 \quad \text{and} \quad y = \frac{1}{4}.$$

Figure 1 is suggestive and intuitively very useful and appeared for the first time in the paper by Dave Morrison [DRM] on the 3-fold octic in $\mathbb{P}^4(1, 1, 1, 2, 2)$ which is a higher dimensional generalization of our example.

The analysis of the neighborhood of the point A is completely analogous to the work [AHK], so we will not discuss here. By analyzing the moduli space picture, we see that a rational curve $x = \text{constant}$ infinitesimally close to the y -axis corresponds to a transition from X_1 , the total space of the canonical bundle of F_2 to the stacky version X_4 of the weighted projective space $\mathbb{P}(2, 1, 1)$. This curve intersects $E_A = 0$ in the neighborhood of the point B . The circuit I associated to the corresponding edge of the secondary quadrilateral is $v_1 - 2v_2 + v_3 = 0$. One of the associated EZ-spherical functor is induced by the diagram

$$\begin{array}{ccc} E = \mathbb{A}^1 \times \mathbb{P}^1 & \hookrightarrow & X_1 \\ \downarrow q & & \\ Z = \mathbb{A}^1 & & \end{array}$$

The birational transformation corresponding a rational curve $x = \text{constant}$ infinitesimally close to the $y = \infty$ corresponds to a transition from X_2 , a different birational model of X_1 to X_4 which is the stacky quotient $[\mathbb{C}^3/\mathbb{Z}_4]$. This curve intersects $E_A = 0$ in the neighborhood of




FIGURE 2. The "mirror" complex discriminant moduli space.

the point C . One of the associated spherical functors is induced by the diagram

$$\begin{array}{ccc} E = [\mathbb{A}^1/\mathbb{Z}_2] \times \mathbb{P}^1 & \hookrightarrow & X_4 = [\mathbb{C}^3/\mathbb{Z}_4] \\ \downarrow q & & \\ Z = [\mathbb{A}^1/\mathbb{Z}_2] & & \end{array}$$

It is interesting to note that the circuit I associated to this transition and determined by corresponding edge of the secondary polytope is also $v_1 - 2v_2 + v_3 = 0$.

Recall that in determining multiplicities, the circuit discriminant $\Delta_I = 4a_1a_3 - a_2^2$ played a crucial role. For the transition $X_1 \leftrightarrow X_4$, we can consider the linear form ψ in the cone $\mathcal{C}(\mathcal{F}_{14})$ whose supporting face is the edge F_{14} given by

$$\psi(v_0) = u, \psi(v_1) = \psi(v_2) = \psi(v_3) = 1, \psi(v_4) = 0,$$

with $u < 1/2$. A similar linear form ψ will work for the transition $X_1 \leftrightarrow X_4$, with $u > 1/2$ in that case. The monomials in f_Q $a_0^4, a_0^2a_2a_4, a_2^2a_4^2, a_1a_3a_4^2$ have the weights $4u, 2u + 1, 2, 2$, respectively. The leading term for the edge F_{14} is a_0^4 . The highest weight terms for the edge F_{23} are $a_2^2a_4^2$ and $a_1a_3a_4^2$, so in this case the leading term for f_Q is

$$16a_2^2a_4^2 - 64a_1a_3a_4^2 = -16a_4^2 \cdot \Delta_I.$$

We conclude that $n_{Q,F_{1,4}} = 0$ and $n_{Q,F_{2,3}} = 1$, as Figure 2 indicates. In this simple example, we do have the true parametrizations of the

components, so the weight calculation is not needed. However, only the birational Horn parametrizations are available in the general case.

5. FINAL COMMENTS AND SOME SPECULATIONS

The proposed constructions of spherical functors appearing in this paper seem to be related of the notion of co-sheaf over the topological space of the cone $K = \sum \mathbb{R}v_i$, in the sense of Bressler-Lunts [BL], section 6.7.1.

Since the defined spherical functors can be thought as categorical maps corresponding to the Bressler-Lunts strata, it is a very intriguing question whether the whole structure can be expressed in the language of schobers introduced by Kapranov and Schetchman [KaS]. Such an abstract realization would be consistent with the fact that our EZ -spherical functors map "stalks" between various strata of a Whitney stratification of the characteristic cycle of the GKZ hypergeometric system. Such a picture would implicitly subsume our multiplicity result. The stalks in this case would be bounded derived categories of coherent sheaves of the form $D(Z_\Gamma)$, indexed by the faces Γ . In particular, if $\Gamma = \emptyset$, the "stalk" is the bounded derived category of coherent sheaves on the ambient toric DM stack. Donovan [Do] studied certain wall crossing phenomena in terms of spherical pairs and schobers. In a combinatorial context similar to ours, Špenko and Van den Bergh [SvB] produced a categorification of the GKZ system in the quasi-symmetric toric case. More work is needed in order to make our speculation rigorous.

REFERENCES

- [AL] R. Anno and T. Logvinenko, *Spherical DG-functors*, J. EMS **19** (9), 2577–2656, [arXiv:1309.5035](#).
- [AGM] P. S. Aspinwall, B. R. Greene and D. R. Morrison, *Calabi–Yau moduli space, mirror manifolds and spacetime topology change in string theory*, Nuclear Phys. B **416** (1994), 414–480, [hep-th/9309097](#).
- [AHK] P. S. Aspinwall, R. P. Horja, and R. L. Karp, *Massless D-Branes on Calabi–Yau Threefolds and Monodromy*, Comm. in Math. Physics, **259** (2005), 45–69, [hep-th/0209161](#)
- [APW] P. S. Aspinwall, M. R. Plesser and K. Wang, *Mirror Symmetry and Discriminants*, preprint (2017), [arXiv:1702.04661](#).
- [BFK] M. Ballard, D. Favero, David, L. Katzarkov, *Variation of geometric invariant theory quotients and derived categories*, J. Reine Angew. Math. **746** (2019), 235–303.
- [BO] A. Bondal and D. Orlov, *Semiorthogonal decompositions for algebraic varieties*, preprint [alg-geom/9506012](#).

- [BL] P. Bressler, V. Lunts, *Intersection cohomology on nonrational polytopes*, *Compositio Math.* **135** (2003), 245–278.
- [BCS] L. A. Borisov, L. Chen, G. Smith, *The orbifold Chow ring of toric Deligne-Mumford stacks*, *J. Amer. Math. Soc.* **18** (2005), no. 1, 193–215, [math.AG/0309229](#).
- [BH2] L. A. Borisov, R. P. Horja, *Mellin-Barnes integrals as Fourier-Mukai transforms*, *Adv. Math.* **207** (2006), no. 2, 876–927, [math.AG/0510486](#).
- [BH1] L. A. Borisov, R. P. Horja, *On the K-theory of smooth toric DM stacks*, *Snowbird lectures on string geometry*, 21–42, *Contemp. Math.*, **401**, Amer. Math. Soc., Providence, RI, 2006, [math.AG/0503277](#).
- [BH] Lev A. Borisov, R. Paul Horja, *On the better behaved version of the GKZ hypergeometric system*, *Math. Annalen*, **357** (2013), 585–603.
- [DKK] C. Diemer, L. Katzarkov, G. Kerr, *Symplectomorphism group relations and degenerations of Landau-Ginzburg models*, *J. Eur. Math. Soc. (JEMS)* **18** (2016), no. 10, 2167–2271, [arXiv:1204.2233 \[math.SG\]](#)
- [Do] W. Donovan, *Perverse schobers and wall crossing*, *Int. Math. Res. Not. IMRN* **18** (2019), 5777–5810, [arXiv:1703.00592](#)
- [GKZ] I. M. Gelfand, A. V. Zelevinsky and M. M. Kapranov, *Hypergeometric functions and toral varieties*, *Funct. Analysis and its Appl.* **23** (1989), 94–106.
- [GKZ] I. M. Gelfand, M. M. Kapranov, A. V. Zelevinsky, *Discriminants, resultants and multidimensional determinants*, Birkhäuser, 1994.
- [HL] D. Halpern-Leistner, *The derived category of a GIT quotient*, *J. AMS* **28** (2015), 872–912, [arXiv:1601.02030](#)
- [HLS] D. Halpern-Leistner, I. Shipman, *Autoequivalences of derived categories via geometric invariant theory*, *Adv. Math.* **303**, 1264–1299, [arXiv:1303.5531](#)
- [HHP] M. Herbst, K. Hori, K., D. Page, *Phases of $N = 2$ Theories in $1+1$ Dimensions with Boundary*, CERN-PH-TH/2008-048 (2008), [arXiv:0803.2045](#)
- [H] R. P. Horja, *Hypergeometric functions and mirror symmetry in toric varieties*, preprint (1999), [math.AG/9912109](#).
- [EZ] R. P. Horja, *Derived category automorphisms from mirror symmetry*, *Duke Mathematical Journal*, **127** (2005), 1–34.
- [JT] Y. Jiang, H-H. Tseng, *On the K-theory of toric stack bundles*, *Asian J. Math.* **14** (2010), no. 1, 1–10. [arXiv:0808.1754 \[math.AG\]](#)
- [KaS] M.M. Kapranov, V. Schechtman, *Perverse Schobers*, preprint (2014), [arXiv:1411.2772 \[math.AG\]](#)
- [Kap] M.M. Kapranov, *A characterization of A -discriminantal hypersurfaces in terms of the logarithmic Gauss map*, *Math. Ann.* **290** (1991), 277–285.
- [Ka] Y. Kawamata, *Log crepant birational maps and derived categories*, *J. Math. Sci. Univ. Tokyo* **12** (2005), no. 2, 211–231, [math.AG/0311139](#).
- [KS] A. Kite, E. Segal, *Discriminants and semi-orthogonal decompositions*, *Comm. in Math. Physics*, **390** (2022), 907–931, [arXiv:2102:08412](#)
- [K1] M. Kontsevich, *Homological algebra of mirror symmetry*, *Proc. Internat. Congr. Math. Zürich 1994*, vol. 1, Birkhäuser Basel, Boston, Berlin, 1995, 120–139, [alg-geom/9411018](#).
- [K2] M. Kontsevich, *Lecture at Rutgers University*, November 11, 1996, (unpublished).

- [DRM] D. R. Morrison, *Through the looking glass*, Mirror symmetry III (D. H. Phong, L. Vinet and S.-T. Yau, eds), American Math. Soc., International Press, 1999, 263–277, [alg-geom/9705028](#).
- [SvB] Špela Špenko and M. Van den Bergh, *Perverse schobers and GKZ systems*, preprint (2020), [arXiv: 2007.04924](#).
- [SYZ] A. Strominger, S.-T. Yau and E. Zaslow, *Mirror symmetry is T-duality*, Nuclear Phys. B **479** (1996), 243–259, [hep-th/9606040](#).
- [V] A. Vistoli, *Intersection theory on algebraic stacks and on their moduli spaces*, Invent. Math. **97** (1989), no. 3, 613–670.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MIAMI, CORAL GABLES,
FL 33146, USA, horja@math.miami.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MIAMI, CORAL GABLES,
FL 33146, USA, l.katzarkov@miami.edu