

SUM OF ELEMENTS IN FINITE SIDON SETS II

YUCHEN DING

ABSTRACT. A set $S \subset \{1, 2, \dots, n\}$ is called a Sidon set if all the sums $a + b$ ($a, b \in S$) are different. Let S_n be the largest cardinality of the Sidon sets in $\{1, 2, \dots, n\}$. In a former article, the author proved the following asymptotic formula

$$\sum_{a \in S, |S|=S_n} a = \frac{1}{2}n^{3/2} + O(n^{111/80+\varepsilon}),$$

where $\varepsilon > 0$ is an arbitrary small constant. In this note, we give an extension of the above formula. We show that

$$\sum_{a \in S, |S|=S_n} a^\ell = \frac{1}{\ell+1}n^{\ell+1/2} + O(n^{\ell+61/160})$$

for any positive integers ℓ . Besides, we also consider the asymptotic formulae of other type summations involving Sidon sets. The proofs are established in a more general setting, namely we obtain the asymptotic formulae of the Sidon sets with t elements when t is near the magnitude $n^{1/2}$.

1. INTRODUCTION

The notion of the Sidon set dates back to a paper of Sidon [23] where he investigated the coefficients of the Fourier series. Since then, the researches of Sidon sets became rich after the past ninety years. Let n be a positive integer and S be a subset of the first n numbers $\{1, 2, \dots, n\}$. We call S a Sidon set if all the sums $a + b$ ($a, b \in S$) are different. Denoting by S_n the largest cardinality of the Sidon sets in $\{1, 2, \dots, n\}$, i.e.,

$$S_n = \max_{S \subset \{1, 2, \dots, n\}, S \text{ Sidon}} |S|.$$

It is a natural question to ask the magnitude of the number S_n . And there are a few literatures involving this magnitude. In 1941, Erdős and Turán [12] proved that

$$\frac{1}{\sqrt{2}} \leq \liminf_{n \rightarrow \infty} \frac{S_n}{\sqrt{n}} \leq \limsup_{n \rightarrow \infty} \frac{S_n}{\sqrt{n}} \leq 1.$$

Actually, they gave more explicit upper bound $S_n \leq n^{1/2} + O(n^{1/4})$. This upper bound was sharpen to $S_n \leq n^{1/2} + n^{1/4} + 1$ by Lindström [17]. Later, Cilleruelo [6] improved slightly this bound by showing that $S_n \leq n^{1/2} + n^{1/4} + 1/2$. In about 2017, Professor Yong-Gao Chen in Nanjing Normal University showed me that the original method of Erdős and Turán with some more elaborate calculations can already lead to the bound $S_n \leq n^{1/2} + n^{1/4} + 5/8$ (private communications). Recently, Balogh, Füredi and Roy [3]

2010 *Mathematics Subject Classification*. Primary 11B75; Secondary 11B83.

Key words and phrases. Sidon sets; asymptotic formula.

proved that $S_n \leq n^{1/2} + 0.998n^{1/4}$ for sufficiently large n by combining the methods of Erdős–Turán and Lindström.

In the other direction, Chowla [5] and Erdős [9] independently found that the constructions of Singer [24] and Bose [4] provided the evidences of $S_n \geq (1 + o(1))n^{1/2}$. Based on the results of gaps between primes (up to the year of 1999), it has been remarked by Ruzsa [22] that $S_n \geq n^{1/2} + O(n^{11/40+\varepsilon})$, where $\varepsilon > 0$ is an arbitrary small constant. This lower bound together with the upper bound give us that

$$S_n = n^{1/2} + O(n^{11/40+\varepsilon}). \quad (1.1)$$

It was conjectured by Erdős [11] that for any $\varepsilon > 0$, we have $S_n = n^{1/2} + O(n^\varepsilon)$. He offered one thousand dollars for the first one to solve this conjecture.

In the famous book ‘*Unsolved Problems in Number Theory*’ of Guy [13, Problem E28], it asked what is the supremum of the sum of reciprocal terms taken from a Sidon set. It has been proved that

$$\max_{\substack{S \subseteq \{1, 2, \dots, n\} \\ \text{Sidon}}} \sum_{a \in S} \frac{1}{a}$$

is near 2 for sufficiently large n . For results of this topic, one can refer to [1, 10, 15, 16, 18, 21, 25, 26, 27, 28].

For the collections of researches about Sidon sets, one can refer to the excellent survey article [19] and P.hD. thesis [20] of O’Bryant.

In the Lindström argument for bounding the maximal size of a Sidon set, it arises naturally the estimate of

$$\sum_{\substack{S = \{a_1 < a_2 < \dots < a_t\} \\ S \subseteq [1, n] \text{ Sidon}}} (n + 1 - i)a_i.$$

In 2021, the author [8] considered the magnitude of the sum of elements in Sidon sets. To be precise, let S be a Sidon set in $\{1, 2, \dots, n\}$ with $|S| = S_n$, using the asymptotic formula of S_n in equation (1.1) the author proved for any $\varepsilon > 0$

$$\sum_{a \in S} a = \frac{1}{2}n^{3/2} + O(n^{111/80+\varepsilon}). \quad (1.2)$$

In this subsequent note, we provide the asymptotic formulae of the following two type summations:

$$\sum_{\substack{S = \{a_1 < a_2 < \dots < a_t\} \\ S \subseteq [1, n] \text{ Sidon}}} a_i^\ell \quad \text{and} \quad \sum_{\substack{S = \{a_1 < a_2 < \dots < a_t\} \\ S \subseteq [1, n] \text{ Sidon}}} ia_i^\ell.$$

Using these estimates with $\ell = 1$, we finally attain the asymptotic formula of

$$\sum_{\substack{S = \{a_1 < a_2 < \dots < a_t\} \\ S \subseteq [1, n] \text{ Sidon}}} (n + 1 - i)a_i$$

which arose in the text of Lindström. Although this formula can not effect the upper bound of the maximal Sidon sets in Lindström's argument, this is of interest itself.

Our first theorem deals with the general case of equation (1.2).

Theorem 1.1. *Let $S = \{a_1 < a_2 < \dots < a_t\}$ be a Sidon set in $\{1, 2, \dots, n\}$. Then for any positive integer ℓ , we have*

$$\sum_{i=1}^t a_i^\ell = \begin{cases} \frac{1}{\ell+1} t n^\ell + O_\ell(n^{\ell+3/8}), & \text{if } t \geq n^{1/2} - n^{1/4} \\ \frac{1}{\ell+1} t n^\ell + O_\ell(n^{\ell+1/4} \sqrt{n^{1/2} - t}), & \text{if } t < n^{1/2} - n^{1/4}. \end{cases}$$

Several corollaries can be deduced from Theorem 1.1. The asymptotic formula mentioned in the abstract is proved by taking $t = S_n = n^{1/2} + O(n^{21/80})$ (see Lemma 2.10).

Taking $\ell = 1$, we have the following corollary, which is another extension of equation (1.2).

Corollary 1.2. *Let $S = \{a_1 < a_2 < \dots < a_t\}$ be a Sidon set in $\{1, 2, \dots, n\}$. Then*

$$\sum_{i=1}^t a_i = \begin{cases} \frac{1}{2} n t + O(n^{11/8}), & \text{if } t \geq n^{1/2} - n^{1/4} \\ \frac{1}{2} n t + O(n^{5/4} \sqrt{n^{1/2} - t}), & \text{if } t < n^{1/2} - n^{1/4}. \end{cases}$$

Taking $t = S_n$ and then Corollary 1.2 reduces to the following Corollary 1.3.

Corollary 1.3. *Let S be a Sidon set in $\{1, 2, \dots, n\}$ with $|S| = S_n$, then we have*

$$\sum_{a \in S} a = \frac{1}{2} n^{3/2} + O(n^{221/160}).$$

It is certainly that $11/8 < 221/160 < 111/80$, so this is a small improvement of the error term in equation (1.2). In the former article, the author pointed out that one can improve the error term in equation (1.2) to $O(n^{11/8})$ under the assumption $S_n = n^{1/2} + O(n^{1/4})$. This assumption is still open at present due to the barrier of the gaps between primes. By some applications of the mean value estimate rather than the direct use of the size of Sidon sets, we can prove the following stronger result comparing with Corollary 1.3.

Theorem 1.4. *Let S be a Sidon set in $\{1, 2, \dots, n\}$ with $|S| = S_n$, then for all $n \leq N$ but at most $O(N/(\log N)^{7/19})$ exceptions, we have*

$$\sum_{a \in S} a = \frac{1}{2} n^{3/2} + O(n^{11/8} \log n).$$

From Theorem 1.4, we immediately have the following corollary.

Corollary 1.5. *Let S be a Sidon set in $\{1, 2, \dots, n\}$ with $|S| = S_n$, then the asymptotic formula $\sum_{a \in S} a = \frac{1}{2}n^{3/2} + O(n^{11/8} \log n)$ is true for almost all integers n .*

As an application of Theorem 1.1, we establish the last theorem in this note.

Theorem 1.6. *Let $S = \{a_1 < a_2 < \dots < a_t\}$ be a Sidon set in $\{1, 2, \dots, n\}$. Then for any positive integer ℓ we have*

$$\sum_{i=1}^t i a_i^\ell = \begin{cases} \frac{1}{\ell+2} t^2 n^\ell + O_\ell(n^{\ell+3/8} t), & \text{if } t \geq n^{1/2} - n^{1/4} \\ \frac{1}{\ell+2} t^2 n^\ell + O_\ell(n^{\ell+1/4} t \sqrt{n^{1/2} - t}), & \text{if } t < n^{1/2} - n^{1/4}. \end{cases}$$

By Theorem 1.6, we finally offer the asymptotic formula of the summation occurred in the Lindström article.

Corollary 1.7. *Let $S = \{a_1 < a_2 < \dots < a_t\}$ be a Sidon set in $\{1, 2, \dots, n\}$. Then*

$$\sum_{i=1}^t (n+1-i) a_i = \begin{cases} \frac{1}{2} n^2 t + O(n^{19/8}), & \text{if } t \geq n^{1/2} - n^{1/4} \\ \frac{1}{2} n^2 t + O(n^{9/4} \sqrt{n^{1/2} - t}), & \text{if } t < n^{1/2} - n^{1/4}. \end{cases}$$

2. LEMMAS

In this section, we provide some lemmas which shall be used in the proofs of our theorems.

Lemma 2.8 (Baker–Harman–Pintz). [2] *Let p_k be the k -th prime, then we have*

$$p_{k+1} - p_k \ll p_k^{0.525}.$$

Lemma 2.9 (Bose). [4] *Let p be a prime, then there are at least p elements between $[1, p^2 - 1]$ such that all the sums of two of these elements are different modulo $p^2 - 1$. In other words, we have $S_{p^2-1} \geq p$ for primes p .*

Lemma 2.10. *For $n \geq 2$, we have $S_n = n^{1/2} + O(n^{21/80})$.*

This is mentioned in [19] without proof and we offer one here for complement.

Proof. Let p_i be the i -th prime. Suppose that p_t is the largest prime such that $p_t^2 - 1 < n$, then we have

$$p_t^2 - 1 < n \leq p_{t+1}^2 - 1. \quad (2.1)$$

By Lemmas 2.8 and 2.9, it follows that

$$S_n \geq S_{p_t^2-1} \geq p_t = p_{t+1} - (p_{t+1} - p_t) \geq \sqrt{n+1} + O(p_t^{0.525}).$$

From the equation (2.1), it is clear that $p_t^{0.525} \ll n^{0.525/2} = n^{21/80}$. Now the lemma follows from the fact that $S_n \leq n^{1/2} + O(n^{1/4})$. \square

Lemma 2.11 (Cilleruelo). [7] *Given a Sidon set $A \subset \{1, 2, \dots, n\}$ with $|A| = n^{1/2} - L$. Then every subinterval $I \subset [1, n]$ with length cn contains $c|A| + E_I$ elements of A , where*

$$|E_I| \leq 52n^{1/4}(1 + c^{1/2}n^{1/8})(1 + L_+^{1/2}n^{-1/8}), \quad L_+ = \max\{0, L\}.$$

Let B_n denote the n -th Bernoulli number, i.e., it is given by

$$\frac{x}{e^x - 1} = \sum_{n=0}^{\infty} \frac{B_n}{n!} x^n.$$

It is well known that (see for example [14])

$$B_0 = 1, \quad B_1 = -\frac{1}{2}, \quad B_2 = \frac{1}{6}, \quad B_4 = -\frac{1}{30}, \quad B_6 = \frac{1}{42}, \quad B_8 = -\frac{1}{30}, \quad B_{10} = \frac{5}{66}, \dots$$

and $B_n = 0$ for odd integers $n \geq 3$. The Bernoulli polynomial $B_n(x)$ ($n = 0, 1, 2, 3, \dots$) is defined by

$$B_n(x) = \sum_{i=0}^n \binom{n}{i} B_i x^{n-i}, \quad (2.2)$$

where $\binom{n}{i} = \frac{n!}{i!(n-i)!}$. We have the following identity.

Lemma 2.12. [14, Page 93] *For natural numbers r and x , we have*

$$\sum_{n=0}^{x-1} n^{r-1} = \frac{1}{r} (B_r(x) - B_r).$$

3. PROOFS

Proof of Theorem 1.1. We start with calculations of the summation $\sum_{k=1}^n k^{\ell-1} S(k)$, where $S(k) = |S \cap [1, k]|$ for any $1 \leq k \leq n$. On one hand,

$$\sum_{k=1}^n k^{\ell-1} S(k) = \sum_{k=1}^n k^{\ell-1} \sum_{\substack{a \in S \\ 1 \leq a \leq k}} 1 = \sum_{a \in S} \sum_{a \leq k \leq n} k^{\ell-1}. \quad (3.1)$$

By Lemma 2.12, we have

$$\sum_{a \leq k \leq n} k^{\ell-1} = \frac{1}{\ell} B_\ell(n+1) - \frac{1}{\ell} B_\ell(a). \quad (3.2)$$

Taking equation (3.2) into equation (3.1) gives

$$\sum_{k=1}^n k^{\ell-1} S(k) = \frac{1}{\ell} B_\ell(n+1)t - \frac{1}{\ell} \sum_{a \in S} B_\ell(a). \quad (3.3)$$

On the other hand, by Lemma 2.11 with $A = S$ and $I = S \cap [1, k]$ and Lemma 2.12,

$$\begin{aligned} \sum_{k=1}^n k^{\ell-1} S(k) &= \sum_{k=1}^n k^{\ell-1} |S \cap [1, k]| = \sum_{k=1}^n k^{\ell-1} \left(\frac{k}{n} t + E_k \right) \\ &= \frac{t}{n(\ell+1)} (B_{\ell+1}(n+1) - B_{\ell+1}) + \sum_{k=1}^n k^{\ell-1} E_k, \end{aligned} \quad (3.4)$$

where $|E_k| \leq 52n^{1/4}(1 + (k/n)^{1/2}n^{1/8})(1 + L_+^{1/2}n^{-1/8})$, $L_+ = \max\{0, n^{1/2} - t\}$. Hence, we get

$$\sum_{a \in S} B_\ell(a) = \left(B_\ell(n+1) - \frac{\ell}{\ell+1} \frac{B_{\ell+1}(n+1)}{n} \right) t + \frac{\ell t}{(\ell+1)n} B_{\ell+1} - \sum_{k=1}^n k^{\ell-1} E_k. \quad (3.5)$$

From the definition of the Bernoulli polynomial (equation (2.2)), we have

$$B_\ell(n+1) - \frac{\ell}{\ell+1} \frac{B_{\ell+1}(n+1)}{n} = \frac{1}{\ell+1} n^\ell + O_\ell(n^{\ell-1}) \quad (3.6)$$

and

$$B_\ell(a) = a^\ell + O_\ell(a^{\ell-1}). \quad (3.7)$$

Therefore,

$$\sum_{i=1}^t a_i^\ell = \sum_{a \in S} a^\ell = \frac{1}{\ell+1} t n^\ell + O_\ell(t n^{\ell-1}) - \sum_{k=1}^n k^{\ell-1} E_k. \quad (3.8)$$

It remains to bound the sum involving E_k . Since

$$\begin{aligned} |E_k| &\ll \begin{cases} n^{1/4} (1 + (k/n)^{1/2} n^{1/8}), & \text{if } t \geq n^{1/2} - n^{1/4} \\ n^{1/8} (1 + (k/n)^{1/2} n^{1/8}) \sqrt{n^{1/2} - t}, & \text{if } t < n^{1/2} - n^{1/4} \end{cases} \\ &\ll \begin{cases} n^{1/4} + k^{1/2} n^{-1/8}, & \text{if } t \geq n^{1/2} - n^{1/4} \\ (n^{1/8} + k^{1/2} n^{-1/4}) \sqrt{n^{1/2} - t}, & \text{if } t < n^{1/2} - n^{1/4}, \end{cases} \end{aligned} \quad (3.9)$$

it follows that

$$\sum_{k=1}^n k^{\ell-1} E_k \ll \begin{cases} n^{\ell+3/8}, & \text{if } t \geq n^{1/2} - n^{1/4} \\ n^{\ell+1/4} \sqrt{n^{1/2} - t}, & \text{if } t < n^{1/2} - n^{1/4}. \end{cases} \quad (3.10)$$

Now the theorem follows from the trivial estimates

$$t n^{\ell-1} \ll n^{\ell+1/8}$$

for $t < n^{1/2} + O(n^{1/4})$ and

$$t n^{\ell-1} \ll n^{\ell+1/4} \sqrt{n^{1/2} - t}$$

provided that $t < n^{1/2} - n^{1/4}$. \square

Proof of Theorem 1.4. The proof is an application of the mean value idea. For any $1 \leq n \leq N$, let \mathcal{S}_n be a Sidon set in $\{1, 2, \dots, n\}$ with $|\mathcal{S}_n| = S_n$. By Lemma 2.10 and equation (3.5) with $\ell = 1$ and $S = \mathcal{S}_n$, we know that

$$\begin{aligned} \sum_{a \in \mathcal{S}_n} a &= \left(B_1(n+1) - \frac{1}{2} \frac{B_2(n+1)}{n} \right) S_n + O\left(\frac{1}{\sqrt{n}}\right) - \sum_{k=1}^n E_{n,k} \\ &= \frac{1}{2} n^{3/2} + O(n^{101/80}) - \sum_{k=1}^n E_{n,k}, \end{aligned} \quad (3.11)$$

where

$$|E_{n,k}| \leq 52n^{1/4}(1 + (k/n)^{1/2}n^{1/8})(1 + L_n^{1/2}n^{-1/8}), \quad L_n = \max\{0, n^{1/2} - S_n\}. \quad (3.12)$$

In another word, we shall have

$$\left| \sum_{a \in \mathcal{S}_n} a - \frac{1}{2}n^{3/2} \right| \ll n^{101/80} + \sum_{k=1}^n |E_{n,k}|. \quad (3.13)$$

By equation (3.13), we immediately obtain that

$$\sum_{n=1}^N \left| \sum_{a \in \mathcal{S}_n} a - \frac{1}{2}n^{3/2} \right| \ll N^{181/80} + \sum_{n=1}^N \sum_{k=1}^n |E_{n,k}|. \quad (3.14)$$

We now treat the summation involving $E_{n,k}$. From equation (3.12), we know that

$$\sum_{k=1}^n |E_{n,k}| \ll (n^{1/4} + L_n^{1/2}n^{1/8}) \sum_{k=1}^n (1 + k^{1/2}n^{-3/8}) \ll n^{11/8} + L_n^{1/2}n^{5/4}. \quad (3.15)$$

From which it can be deduced that

$$\sum_{n=1}^N \sum_{k=1}^n |E_{n,k}| \ll \sum_{n=1}^N (n^{11/8} + L_n^{1/2}n^{5/4}) \ll N^{19/8} + \sum_{n=1}^N L_n^{1/2}n^{5/4}. \quad (3.16)$$

Suppose that p_ℓ is the least prime such that $N \leq p_\ell^2 - 1$, where p_i is the i -th prime and we define p_0 to be 1. Then we have

$$\sum_{n=1}^N L_n^{1/2}n^{5/4} \leq \sum_{m=0}^{\ell-1} \sum_{p_m^2-1 < n \leq p_{m+1}^2-1} L_n^{1/2}n^{5/4}. \quad (3.17)$$

Recall that $S_{p^2-1} \geq p$ from Lemma 2.9, thus by the definition of L_n in equation (3.12), we get

$$\begin{aligned} \sum_{p_m^2-1 < n \leq p_{m+1}^2-1} L_n^{1/2}n^{5/4} &\leq \sum_{p_m^2-1 < n \leq p_{m+1}^2-1} n^{5/4}(n^{1/2} - p_m)^{1/2} \\ &< \sum_{p_m^2 < n \leq p_{m+1}^2} n^{5/4}(n^{1/2} - p_m)^{1/2} \\ &\ll p_{m+1}^{5/4} \sum_{p_m^2 < n \leq p_{m+1}^2} (n^{1/2} - p_m)^{1/2} \\ &\ll p_{m+1}^{5/4} \int_{p_m^2}^{p_{m+1}^2} (x^{1/2} - p_m)^{1/2} dx \\ &= \frac{t=x^{1/2}-p_m}{x=(t+p_m)^2} p_{m+1}^{5/4} \int_0^{p_{m+1}^2-p_m} 2t^{1/2}(t+p_m) dt \\ &\ll p_{m+1}^{5/4} (p_{m+1} - p_m)^{5/2} + p_{m+1}^{5/4} p_m (p_{m+1} - p_m)^{3/2} \\ &\ll p_{m+1}^{9/4} (p_{m+1} - p_m)^{3/2}. \end{aligned} \quad (3.18)$$

Taking this into equation (3.17), we get

$$\begin{aligned}
\sum_{n=1}^N L_n^{1/2} n^{5/4} &\ll \sum_{m=0}^{\ell-1} p_{m+1}^{9/4} (p_{m+1} - p_m)^{3/2} \\
&\leq \sum_{m=0}^{\ell-1} p_{m+1}^{11/4} (p_{m+1} - p_m) \\
&\leq p_\ell^{11/4} \sum_{m=0}^{\ell-1} (p_{m+1} - p_m) \\
&\leq p_\ell^{15/4}.
\end{aligned} \tag{3.19}$$

Recall that $p_{\ell-1}^2 - 1 < N \leq p_\ell^2 - 1$ from the definition of ℓ , so we have

$$p_\ell = p_{\ell-1} + (p_\ell - p_{\ell-1}) \leq \sqrt{N+1} + O(p_{\ell-1}^{0.525}) \ll N^{1/2}. \tag{3.20}$$

It follows that

$$\sum_{n=1}^N L_n^{1/2} n^{5/4} \ll N^{15/8}. \tag{3.21}$$

Thus, we conclude that

$$\sum_{n=1}^N \left| \sum_{a \in \mathcal{S}_n} a - \frac{1}{2} n^{3/2} \right| \ll N^{19/8} \tag{3.22}$$

from equations (3.14), (3.16) and (3.21). Now let \mathcal{A} be the set of integers $n \leq N$ such that

$$\left| \sum_{a \in \mathcal{S}_n} a - \frac{1}{2} n^{3/2} \right| > n^{11/8} \log n. \tag{3.23}$$

If there are more than $N/(\log N)^{7/19}$ integers in \mathcal{A} , we shall have

$$\sum_{n \in \mathcal{A}} \left| \sum_{a \in \mathcal{S}_n} a - \frac{1}{2} n^{3/2} \right| > \sum_{n \leq N/(\log N)^{7/19}} n^{11/8} \log n \gg N^{19/8} (\log N)^{1/8}. \tag{3.24}$$

This a contradiction with equation (3.22) for sufficiently large N , which means that we should have $|\mathcal{A}| < N/(\log N)^{7/19}$ provided that N is large enough. Therefore, for all integers $n \leq N$ but at most $N/(\log N)^{7/19}$ exceptions, we have

$$\left| \sum_{a \in \mathcal{S}_n} a - \frac{1}{2} n^{3/2} \right| \leq n^{11/8} \log n. \tag{3.25}$$

□

Remark 3.1. It is sure that the error term $O(n^{11/8} \log n)$ in Theorem 1.4 can be improved to $O(n^{11/8} f(n))$ for the integers of density one provided that $f(n)$ tends to infinity. It seems that this is the best possible which can be achieved by the present method.

Proof of Theorem 1.6. It is clear that

$$\sum_{i=1}^t ia_i^\ell = \sum_{a \in S} a^\ell \sum_{\substack{b \in S \\ b \leq a}} 1 = \sum_{a \in S} a^\ell \left(\frac{a}{n} t + E_a \right) = \frac{t}{n} \sum_{a \in S} a^{\ell+1} + \sum_{a \in S} a^\ell E_a \quad (3.26)$$

from Lemma 2.11 with $A = S$ and $I = S \cap [1, a]$, where

$$\begin{aligned} |E_a| &\leq 52n^{1/4}(1 + (a/n)^{1/2}n^{1/8})(1 + L_+^{1/2}n^{-1/8}) \quad (\text{with } L_+ = \max\{0, n^{1/2} - t\}) \\ &\ll \begin{cases} n^{1/4} + a^{1/2}n^{-1/8}, & \text{if } t \geq n^{1/2} - n^{1/4} \\ (n^{1/8} + a^{1/2}n^{-1/4})\sqrt{n^{1/2} - t}, & \text{if } t < n^{1/2} - n^{1/4}. \end{cases} \end{aligned} \quad (3.27)$$

For $t \geq n^{1/2} - n^{1/4}$, we have

$$\begin{aligned} \sum_{a \in S} a^\ell E_a &\ll n^{1/4} \sum_{a \in S} a^\ell + n^{-1/8} \sum_{a \in S} a^{\ell+1/2} \\ &\ll n^{1/4} n^\ell t + n^{-1/8} n^{\ell+1/2} t \\ &\ll n^{\ell+3/8} t. \end{aligned} \quad (3.28)$$

For $t < n^{1/2} - n^{1/4}$, we have

$$\begin{aligned} \sum_{a \in S} a^\ell E_a &\ll n^{1/8} \sqrt{n^{1/2} - t} \sum_{a \in S} a^\ell + n^{-1/4} \sqrt{n^{1/2} - t} \sum_{a \in S} a^{\ell+1/2} \\ &\ll n^{1/8} n^\ell t \sqrt{n^{1/2} - t} + n^{-1/4} n^{\ell+1/2} t \sqrt{n^{1/2} - t} \\ &\ll n^{\ell+1/4} t \sqrt{n^{1/2} - t}. \end{aligned} \quad (3.29)$$

Recall that

$$\sum_{a \in S} a^{\ell+1} = \begin{cases} \frac{1}{\ell+2} t n^{\ell+1} + O_\ell(n^{\ell+11/8}), & \text{if } t \geq n^{1/2} - n^{1/4} \\ \frac{1}{\ell+2} t n^{\ell+1} + O_\ell(n^{\ell+5/4} \sqrt{n^{1/2} - t}), & \text{if } t < n^{1/2} - n^{1/4}. \end{cases} \quad (3.30)$$

from Theorem 1.1, thus the theorem follows from equations (3.26), (3.28), (3.29) and (3.30). \square

Proof of Corollary 1.7. From Corollary 1.2 and Theorem 1.6 with $\ell = 1$, we have

$$\begin{aligned} \sum_{i=1}^t (n+1-i)a_i &= (n+1) \sum_{i=1}^t a_i - \sum_{i=1}^t ia_i \\ &= \begin{cases} \frac{1}{2} n^2 t + O(n^{19/8}), & \text{if } t \geq n^{1/2} - n^{1/4} \\ \frac{1}{2} n^2 t + O(n^{9/4} \sqrt{n^{1/2} - t}), & \text{if } t < n^{1/2} - n^{1/4}. \end{cases} \end{aligned} \quad (3.31)$$

\square

ACKNOWLEDGMENTS

The author is supported by the Natural Science Foundation of Jiangsu Province of China, Grant No. BK20210784. He is also supported by the foundations of the projects ‘‘Jiangsu Provincial Double-Innovation Doctor Program’’, Grant No. JSSCBS20211023

and “Golden Phoenix of the Green City–Yang Zhou” to excellent PhD, Grant No. YZLYJF2020PHD051.

REFERENCES

- [1] H. L. Abbott, On sum-free sequences, *Acta Arith.*, **48** (1987), 93–96.
- [2] R. C. Baker, G. Harman, J. Pintz, The difference between consecutive primes, II, *Proc. Lond. Math. Soc.*, **83** (2001), 532–562.
- [3] J. Balogh, Z. Füredi and S. Roy, An upper bound on the size of Sidon set, <http://arxiv.org/abs/2103.15850>.
- [4] R. C. Bose, An affine analogue of Singer’s theorem, *J. Indian Math. Soc.*, **6** (1942), 1–15.
- [5] S. Chowla, Solution of a problem of Erdős and Turán in additive number theory, *Proc. Natl. Acad. Sci. India Sect. A*, **14** (1944), 1–2.
- [6] J. Cilleruelo, Sidon sets in \mathbb{N}^d , *J. Combin. Theory Ser. A*, **117** (2010), 857–871.
- [7] J. Cilleruelo, Gaps in dense Sidon sets, *Integers*, **0** (2000).
- [8] Y. Ding, Sum of elements in finite Sidon sets, *Int. J. Number Theory*, **17** (2021), 991–1001.
- [9] P. Erdős, On a problem of Sidon in additive number theory and on some related problems. Addendum, *J. London Math. Soc.*, **19** (1944), 208.
- [10] P. Erdős, Problems and results in combinatorial analysis and combinatorial number theory, *Graph Theory, Combinatorics and Applications*, Vol. 1 (Kalamazoo MI, 1988) 397–406, Wiley, New York, 1991.
- [11] P. Erdős, Problems and results in discrete mathematics, *Discrete Math.*, **136** (1994), 53–73.
- [12] P. Erdős and P. Turán, On a problem of Sidon in additive number theory, and on some related problems, *J. London Math. Soc.*, **16** (1941), 212–215.
- [13] R. K. Guy, *Unsolved Problems in Number Theory*, Volume 1, 2nd Edition, Springer–Verlag, (1994).
- [14] K. Kato, N. Kurokawa, T. Saito, *Number theory, 1, Fermat’s dream*, Translated from the 1996 Japanese original by Masato Kuwata, *Translations of Mathematical Monographs* 186, Iwanami Series in Modern Mathematics, American Mathematical Society, Providence, RI, 2000.
- [15] E. Levine, An extremal result for sum-free sequences, *J. Number Theory*, **12** (1980), 251–257.
- [16] E. Levine, J. O’Sullivan, An upper estimate for the reciprocal sum of a sum-free sequence, *Acta Arith.*, **34** (1977), 9–24.
- [17] B. Lindström, An inequality for B_2 -sequences, *J. Combin. Theory*, **6** (1969), 211–212.
- [18] A. M. Mian, S. D. Chowla, On the B_2 -sequences of Sidon, *Proc. Nat. Acad. Sci. India Sect. A*, **14** (1944), 3–4.
- [19] K. O’Bryant, A complete annotated bibliography of work related to Sidon sequences, *Electron. J. Combin.*, **11** (2004).
- [20] K. O’Bryant, *Sidon sets and Beatty sequences*. Thesis (Ph.D.)–University of Illinois at Urbana-Champaign. 2002.
- [21] J. O’Sullivan, *On reciprocal sums of sum-free sequences*, PhD thesis, Adelphi University, 1973.
- [22] I. Z. Ruzsa, Erdős and the integers, *J. Number Theory*, **79** (1999), 115–163.
- [23] S. Sidon, Ein Satz über trigonometrische Polynome und seine Anwendung in der Theorie der Fourier-Reihen, *Math. Ann.*, **106**(1932), 536–539.
- [24] J. Singer, A theorem in finite projective geometry and some applications to number theory, *Trans. Amer. math. Soc.*, **43** (1938), 377–385.
- [25] H. Taylor and G. S. Yovanof, B_2 -sequences and the distinct distance constant, *Comput. Math. Appl.*, **39** (2000), 37–42.
- [26] Z.-X. Zhang, A sum-free sequence with larger reciprocal sum, *Discrete Math.*, (1992).
- [27] Z.-X. Zhang, A B_2 -sequence with larger reciprocal sum, *Math. Comput.*, **60** (1993), 835–839.
- [28] Z.-X. Zhang, Finding finite B_2 -sequences with larger $m - a_m^{1/2}$, *Math. Comput.*, **61** (1993).

(YUCHEN DING) SCHOOL OF MATHEMATICAL SCIENCE, YANGZHOU UNIVERSITY, YANGZHOU 225002, PEOPLE’S REPUBLIC OF CHINA

Email address: ycding@yzu.edu.cn