

# WITTEN–RESHETIKHIN–TURAEV INVARIANTS AND HOMOLOGICAL BLOCKS FOR PLUMBED HOMOLOGY SPHERES

YUYA MURAKAMI

ABSTRACT. In this paper, we prove a conjecture by Gukov–Pei–Putrov–Vafa for a wide class of plumbed 3-manifolds. Their conjecture states that Witten–Reshetikhin–Turaev (WRT) invariants are radial limits of homological blocks, which are  $q$ -series introduced by them for plumbed 3-manifolds with negative definite linking matrices. The most difficult point in our proof is to prove the vanishing of weighted Gauss sums that appear in coefficients of negative degree in asymptotic expansions of homological blocks. To deal with it, we develop a new technique for asymptotic expansions, which enables us to compare asymptotic expansions of rational functions and false theta functions related to WRT invariants and homological blocks, respectively. In our technique, our vanishing results follow from holomorphy of such rational functions.

## CONTENTS

1. Introduction	1
Acknowledgement	2
2. Basic notations for plumbing graphs	3
2.1. Notations for graphs	3
2.2. The inverse matrix of the linking matrix	3
2.3. Periodic maps determined by the linking matrix	5
2.4. Rational functions determined by the linking matrix	6
3. Calculations of WRT invariants	7
4. Expressions of homological blocks as false theta functions	9
4.1. Settings and the main result	9
4.2. A proof of Proposition 4.2	10
5. Asymptotic formulas	11
5.1. Asymptotic formulas obtained from Euler–Maclaurin summation formula	11
5.2. An asymptotic formula for infinite series with signatures	13
5.3. Asymptotic expansion of infinite series with weighted Gauss sums	15
6. A proof of the main theorem	16
6.1. The main result	16
6.2. A proof of the main result	17
References	18

## 1. INTRODUCTION

In this paper, we study Witten–Reshetikhin–Turaev (WRT) invariants. They are quantum invariants of 3-manifolds constructed by Witten [Wit89] from a physical viewpoint and constructed by Reshetikhin–Turaev [RT91, Theorem 3.3.2] from a mathematical viewpoint. The Witten’s asymptotic expansion conjecture plays a central role in studying WRT invariants. It claims that the Chern–Simons invariants and the Reidemeister torsions appear in the asymptotic expansion of WRT invariants. This conjecture is proved by Lawrence–Zagier [LZ99] for the Poincaré homology sphere, Hikami [Hik05a] for Brieskorn homology spheres, Hikami [Hik06a] and Matsusaka–Terashima [MT21] independently for Seifert homology spheres. Many other authors have studied this conjecture [AH12, AM20, And13, BW05, Cha16, Chu17, CM15, FIMT21, FG91, GMP16, Hik05b, Hik06b, HT02, Jef92, Roz94, Roz96, Wu21].

Recently, Gukov–Pei–Putrov–Vafa [GPPV20] introduced  $q$ -series invariants called homological blocks or GPPV invariants for plumbed manifolds. They conjectured that homological blocks have good modular transformations and their radial limits yield WRT invariants. These conjectures show a roadmap to prove the Witten’s asymptotic expansion conjecture. The first conjecture, modular transformations of homological blocks, are related to quantum modular forms introduced by Zagier [Zag10]. This conjecture is proved by Matsusaka–Terashima [MT21] for Seifert homology spheres and Bringmann–Mahlburg–Milas [BMM20a] for non-Seifert homology spheres whose surgery diagrams are the H-graphs. Bringmann–Nazaroglu [BN19] and Bringmann–Kasdzian–Milas–Nazaroglu [BKMN21] clarified and proved the modular transformation formulas of false theta functions, which are main tools to study modular transformations of homological blocks.

The second conjecture, which asks the relation between WRT invariants and radial limits of homological blocks, is proved by Fuji–Iwaki–Murakami–Terashima [FIMT21] with a result in Andersen–Mistegård [AM20] for Seifert homology spheres and Mori–Murakami [MM21] for non-Seifert homology spheres whose surgery diagrams are the H-graphs. In this paper, we prove this conjecture for a large class of plumbed manifolds by developing some new techniques.

Let us explain our setting. Let  $\Gamma = (V, E, (w_v)_{v \in V})$  be a plumbing graph, that is, a finite tree with the vertex set  $V$ , the edge set  $E$ , and integral weights  $w_v \in \mathbb{Z}$  for each vertex  $v \in V$ . Following [GPPV20], we assume that the linking matrix  $W$  of  $\Gamma$  is negative definite. Then one can define the homological block  $\widehat{Z}_\Gamma(q)$  with  $q \in \mathbb{C}$  and  $|q| < 1$ . Let  $M(\Gamma)$  be the plumed 3-manifold obtained from  $S^3$  through the surgery along the diagram defined by  $\Gamma$ . We also assume that  $M(\Gamma)$  is an integral homology sphere. This condition is equivalent to  $\det W = \pm 1$  since  $H_1(M(\Gamma), \mathbb{Z}) \cong \mathbb{Z}^V / W(\mathbb{Z}^V)$  by use of the Mayer–Vietoris sequence. For a positive integer  $k$ , let  $\text{WRT}_k(M(\Gamma))$  be the WRT invariant of  $M(\Gamma)$  normalised as  $\text{WRT}_k(S^3) = 1$ . We also denote  $\zeta_k := e^{2\pi\sqrt{-1}/k}$ .

In the above setting, we can state the conjecture of Gukov–Pei–Putrov–Vafa [GPPV20] by the following.

**Conjecture 1.1** ([GPPV20, Conjecture 2.1, Equation (A.28)]).

$$\text{WRT}_k(M(\Gamma)) = \frac{1}{2(\zeta_{2k} - \zeta_{2k}^{-1})} \lim_{q \rightarrow \zeta_k} \widehat{Z}_\Gamma(q).$$

In this paper, we prove this conjecture for a wide class of plumbing graphs.

**Theorem 1.2.** *Conjecture 1.1 is true if  $|\bar{v}| + 2 - \deg(v) > 0$  for any vertex  $v \in V$ , where  $\deg(v)$  is the degree of  $v$  and  $\bar{v} := \{i \in V \mid \{i, v\} \in E, \deg(i) = 1\}$ .*

In a proof of this theorem, the most difficult point is to prove the vanishing of weighted Gauss sums that appear in coefficients of negative degree in asymptotic expansions of homological blocks. The previous works [BMM20a, MM21] deal with this difficulty by direct calculations ([BMM20a, Theorem 4.1], [MM21, Proposition 4.2]). However, we prove it indirectly by using our asymptotic formula (Proposition 5.2 and Corollary 5.7) and holomorphy of a rational function whose radial limits are WRT invariants. In a sense, our technique is a generalisation of the method of Lawrence–Zagier [LZ99, pp.98, Proposition] using  $L$ -functions for periodic maps.

This paper will be organised as follows. In Section 2, we prepare some notations for plumbing graphs  $\Gamma$  which we use throughout this paper. In Section 3, we calculate WRT invariants for plumbed homology spheres  $M(\Gamma)$ . The point of our calculation is to represent WRT invariants as a sum for the submatrix of  $W^{-1}$  with vertices whose degrees are greater than 2. In Section 4, we express homological blocks as false theta functions. Then we can study the asymptotic expansions of homological blocks as  $q \rightarrow \zeta_k$ . To calculate asymptotic expansions, we develop a formula in Section 5.1 by the Euler–Maclaurin summation formula based on Zagier [Zag06, Equation (44)]. In Section 5.2, we rewrite this formula as an appropriate form for false theta functions. In Section 5.3, we develop the important asymptotic formula, which plays a central role in proving our main theorem. This formula asserts that the same factors appear in the asymptotic expansions of homological blocks and rational functions whose radial limits are WRT invariants. Finally, we prove our main theorem in Section 6.

#### ACKNOWLEDGEMENT

The author would like to show the greatest appreciation to Takuya Yamauchi for giving many pieces of advice. The author would like to thank Takuya Yamauchi, Yuji Terashima, Kazuhiro Hikami,

Toshiki Matsusaka for giving many comments. The author is deeply grateful to Akihito Mori for a lot of discussion. The author is supported by JSPS KAKENHI Grant Number JP 20J20308.

## 2. BASIC NOTATIONS FOR PLUMBING GRAPHS

In this section, we list some notations for plumbing graphs and their basic properties, which we use throughout this paper.

**2.1. Notations for graphs.** In this subsection, we prepare settings for graphs. As in Section 1, let  $\Gamma = (V, E, (w_v)_{v \in V})$  be a plumbing graph and  $W$  be its linking matrix such that it is negative definite and  $\det W = \pm 1$ . Here we consider the edge set  $E$  as the subset of  $\{\{v, v'\} \mid v, v' \in V\}$ . We identify  $M_{|V|}(\mathbb{Z})$  and  $\text{End}(\mathbb{Z}^V)$  and consider  $W$  as the element of  $\text{End}(\mathbb{Z}^V)$ . We remark that  $w_v \in \mathbb{Z}_{<0}$  for any vertex  $v \in V$ .

For two plumbing graphs  $\Gamma$  and  $\Gamma'$ , Neumann ([Neu80, Proposition 2.2], [Neu81, Theorem 3.1]) proved that two 3-manifolds  $M(\Gamma)$  and  $M(\Gamma')$  are homeomorphic if and only if  $\Gamma$  and  $\Gamma'$  are related by Neumann moves shown in Figure 1. Thus, we can assume  $w_v \leq -2$  for a vertex  $i$  with  $\deg(i) = 1$ .




FIGURE 1. Neumann moves

For a positive integer  $d$ , let

$$V_d := \{v \in V \mid \deg(v) = d\}, \quad V_{\geq d} := \{v \in V \mid \deg(v) \geq d\}.$$

Let  $W_d \in \text{End}(\mathbb{Z}^{V_d})$  and  $W_{\geq d} \in \text{End}(\mathbb{Z}^{V_{\geq d}})$  be the submatrices of  $W$  with  $V_d \times V_d$  and  $V_{\geq d} \times V_{\geq d}$  components respectively. Moreover, for a positive integer  $e$ , let  $W_{d, \geq e} \in \text{Hom}(\mathbb{Z}^{V_{\geq e}}, \mathbb{Z}^{V_d})$  and  $W_{\geq d, e} \in \text{Hom}(\mathbb{Z}^{V_e}, \mathbb{Z}^{V_{\geq d}})$  be the submatrices of  $W$  with  $V_d \times V_{\geq e}$  and  $V_{\geq d} \times V_e$  components respectively.

We need to focus on vertices with degree 1 to calculate WRT invariants and homological blocks. For this reason, we define

$$\bar{v} := \{i \in V_1 \mid \{i, v\} \in E\}, \quad M_v := \prod_{i \in \bar{v}} w_i$$

for each vertex  $v \in V_{\geq 2}$ . Here we define  $M_v := 1$  if  $\bar{v} = \emptyset$ .

The condition  $\det W = \pm 1$  implies the following lemma, which Akihito Mori told the author.

**Lemma 2.1.** *For a vertex  $v \in V_{\geq 2}$  and distinct vertices  $i, j \in \bar{v}$ , it holds  $\gcd(w_i, w_j) = 1$ .*

*Proof.* Let  $\{e_v\}_{v \in V}$  be the standard basis of  $\mathbb{R}^V$ . Since

$$We_i - We_j = (e_v + w_j e_j) - (e_v + w_i e_i) = w_j e_j - w_i e_i,$$

$\gcd(w_i, w_j)$  divides  $\det W = \pm 1$ . □

**2.2. The inverse matrix of the linking matrix.** In this subsection, we calculate the inverse matrix of the linking matrix.

Let  $S \in \text{Aut}(\mathbb{Q}^{V_{\geq 2}}) \cap \text{End}(\mathbb{Z}^{V_{\geq 2}})$  be the  $V_{\geq 2} \times V_{\geq 2}$  submatrix of  $-W^{-1} \in \text{Aut}(\mathbb{Q}^V)$ . We also denote  $T := -W_1^{-1} W_{1, \geq 2}$ . Here we remark that  $S$  is positive definite since  $W$  is negative definite.

The inverse matrices  $W^{-1}$  and  $S^{-1}$  have the following properties.

**Proposition 2.2.** (i)

$$W^{-1} = \begin{pmatrix} T \\ I \end{pmatrix} S \begin{pmatrix} {}^t T & I \end{pmatrix} + \begin{pmatrix} W_1^{-1} & \\ & O \end{pmatrix}.$$

(ii)

$$\det S = \det W \prod_{i \in V_1} w_i = \prod_{i \in V_1} |w_i|.$$

(iii)

$$S^{-1} = -W_{\geq 2} + \text{diag} \left( \sum_{i \in \bar{v}} \frac{1}{w_i} \right)_{v \in V_{\geq 2}}.$$

Here, we denote  $\text{diag}(a_v)_{v \in V_{\geq 2}}$  the diagonal matrix whose  $(v, v)$ -component is  $a_v$ .

(iv) For distinct vertices  $v, v' \in V_{\geq 2}$ , the  $(v, v)$ -component of  $S$  is in  $M_v \mathbb{Z}$  and the  $(v, v')$ -component of  $S$  is in  $M_v M_{v'} \mathbb{Z}$ .

(v)

$$S^{-1}(\mathbb{Z}^{V_{\geq 2}}) = \bigoplus_{v \in V_{\geq 2}} \frac{1}{M_v} \mathbb{Z} \subset \mathbb{Q}^{V_{\geq 2}}.$$

To prove this proposition, we prepare a lemma for the inverse matrices of symmetric block matrices.

**Lemma 2.3.** For a symmetric block matrix

$$X = \begin{pmatrix} A & B \\ {}^t B & C \end{pmatrix} \in \text{GL}_{m+n}(\mathbb{C})$$

such that  $A \in \text{GL}_m(\mathbb{C})$  and  $C \in \text{GL}_n(\mathbb{C})$  be symmetric matrices and  $B \in M_{m,n}(\mathbb{C})$ , let

$$S := (C - {}^t B A^{-1} B)^{-1} \in \text{GL}_n(\mathbb{C}), \quad T := -A^{-1} B \in M_{m,n}(\mathbb{C}).$$

Then, it holds

$$X^{-1} = \begin{pmatrix} T \\ I \end{pmatrix} S \begin{pmatrix} {}^t T & I \end{pmatrix} + \begin{pmatrix} A^{-1} & \\ & O \end{pmatrix}, \quad \det S = \frac{\det A}{\det X}.$$

In particular,  $S$  is the  $n \times n$  bottom right submatrix of  $X^{-1}$ .

*Proof.* The first equality follows from

$$X \left( \begin{pmatrix} T \\ I \end{pmatrix} S \begin{pmatrix} {}^t T & I \end{pmatrix} + \begin{pmatrix} A^{-1} & \\ & O \end{pmatrix} \right) = \begin{pmatrix} (AT + B)S^t T + I & (AT + B)S \\ ({}^t B T + C)S^t T + {}^t B A^{-1} & ({}^t B T + C)S \end{pmatrix}$$

and the second equality follows from

$$X = \begin{pmatrix} I & \\ -{}^t T & I \end{pmatrix} \begin{pmatrix} A & \\ & S^{-1} \end{pmatrix} \begin{pmatrix} I & -T \\ & I \end{pmatrix}.$$

□

Here we remark that Lemma 2.3 is a generalisation of the completing the square  $ax^2 + bx + c = a(x + b/2a)^2 - (b^2 - 4ac)/4a$  for symmetric block matrices.

*Proof of Proposition 2.2.* We obtain (i) and the equalities

$$S^{-1} = -W_{\geq 2} + W_{\geq 2,1} W_1^{-1} W_{1,\geq 2}, \quad \det S = \frac{\det W_1}{\det W}$$

by applying Lemma 2.3 for the block matrix

$$W = \begin{pmatrix} W_1 & W_{1,\geq 2} \\ W_{\geq 2,1} & W_{\geq 2} \end{pmatrix} \in \text{GL}_{m+n}(\mathbb{Z}).$$

Since  $\det W \in \{\pm 1\}$ ,  $W_1 = \text{diag}(w_i)_{i \in V_1}$ , and  $\det S > 0$ , we get (ii). (iii) follows from the fact that the  $(v, v')$ -component of  $W_{\geq 2,1} W_1^{-1} W_{1,\geq 2}$  is

$$\sum_{i \in V_1, \{i,v\}, \{i,v'\} \in E} \frac{1}{w_i} = \delta_{v,v'} \sum_{i \in V_1, \{i,v\} \in E} \frac{1}{w_i}.$$

We prove (iv). By (iii), the  $(v, v')$ -cofactor of  $S^{-1}$  is an element of  $\left( \prod_{v'' \in V_{\geq 2} \setminus \{v,v'\}} 1/M_{v''} \right) \mathbb{Z}$ . Since  $\det S = \pm \prod_{v \in V_{\geq 2}} M_v$  by (iii), we obtain the claim. Finally, we prove (v). Let  $W'_{\geq 2} := \text{diag}(M_v)_{v \in V_{\geq 2}} \in \text{Aut}(\mathbb{Q}^{|V_{\geq 2}|})$ . It suffices to show  $S^{-1} W'_{\geq 2} \in \text{Aut}(\mathbb{Q}^{|V_{\geq 2}|})$ . Since  $\det S = \pm \det W'_{\geq 2}$ , we need only check that each component of  $S^{-1} W'_{\geq 2}$  is an integer, which follows from (iii). □

**2.3. Periodic maps determined by the linking matrix.** In this subsection, we introduce periodic maps determined by the linking matrix and prove its basic properties.

For each vertex  $v \in V_{\geq 2}$ , let

$$\mathcal{S}_v := \left\{ \frac{1}{2} \deg(v) - 1 + \sum_{i \in \bar{v}} \frac{l_i}{2w_i} \mid l_i \in \{\pm 1\}^{\bar{v}} \right\}$$

Here we set  $\mathcal{S}_v := \{\deg(v)/2 - 1\}$  if  $\bar{v} = \emptyset$ . We also define the periodic map  $\varepsilon_v: \frac{1}{2M_v} \mathbb{Z}/\mathbb{Z} \rightarrow \{0, \pm 1\}$  by

$$\varepsilon_v(\alpha_v) := \begin{cases} \prod_{i \in \bar{v}} l_i & \text{if } \alpha_v \equiv \frac{1}{2} \deg(v) - 1 + \sum_{i \in \bar{v}} \frac{l_i}{2w_i} \pmod{\mathbb{Z}} \text{ for some } (l_i)_{i \in \bar{v}} \in \{\pm 1\}^{\bar{v}}, \\ 0 & \alpha_v \notin \mathcal{S}_v. \end{cases}$$

This map is well-defined by the following lemma.

**Lemma 2.4.** *For a vertex  $v \in V_{\geq 2}$  such that  $\bar{v} \neq \emptyset$ , the followings hold.*

(i) *The map*

$$\begin{aligned} \{\pm 1\}^{\bar{v}} &\longrightarrow \mathcal{S}_v \\ (l_i)_{i \in \bar{v}} &\longmapsto \frac{1}{2} \deg(v) - 1 + \sum_{i \in \bar{v}} \frac{l_i}{2w_i} \end{aligned}$$

*is bijective.*

(ii) *The natural projection  $\mathcal{S}_v \rightarrow \frac{1}{2M_v} \mathbb{Z}/\mathbb{Z}$  is injective. Thus, the map  $\varepsilon_v$  is well-defined.*

(iii) *For any  $\alpha_v \in \mathcal{S}_v$ , it holds  $\gcd(2M_v \alpha_v, M_v) = 1$ .*

*Proof.* By Lemma 2.1, there exists  $i_0 \in \bar{v}$  such that  $w_i$  is odd for any  $i \in \bar{v} \setminus \{i_0\}$ . Then, we have

$$\mathbb{Z}/2M_v \mathbb{Z} \cong \mathbb{Z}/2w_{i_0} \mathbb{Z} \oplus \bigoplus_{i \in \bar{v} \setminus \{i_0\}} \mathbb{Z}/w_i \mathbb{Z}$$

by the Chinese remainder theorem. For each

$$\alpha_v = \frac{1}{2} \deg(v) - 1 + \sum_{i \in \bar{v}} \frac{l_i}{2w_i} \in \mathcal{S}_v, \quad (l_i)_{i \in \bar{v}} \in \{\pm 1\}^{\bar{v}},$$

it holds

$$2M_v \alpha_v \equiv \begin{cases} \frac{M_v}{w_i} l_i \pmod{w_i} & \text{if } i \in \bar{v} \setminus \{i_0\}, \\ M_v \deg(v) + \frac{M_v}{w_{i_0}} l_{i_0} + \sum_{i \in \bar{v} \setminus \{i_0\}} \frac{M_v}{w_i} \pmod{2w_{i_0}} & \text{if } i = i_0. \end{cases}$$

Here  $M_v/w_i \not\equiv -M_v/w_i \pmod{w_i}$  for any vertex  $i \in \bar{v} \setminus \{i_0\}$  since  $w_i$  is odd. We also have  $M_v/w_{i_0} \not\equiv -M_v/w_{i_0} \pmod{w_{i_0}}$  since we assume  $w_{i_0} \leq -2$  in Section 2.1. Thus, we obtain (ii). By this argument, we can prove injectivity of the composition of the map  $\{\pm 1\}^{\bar{v}} \rightarrow \mathcal{S}_v$  in (i) and the map  $\mathcal{S}_v \rightarrow \mathbb{Z}/2M_v \mathbb{Z}$ ,  $\alpha_v \mapsto 2M_v \alpha_v$  which is injective by (ii). Hence, we obtain (i). Moreover, we have (iii) since  $\gcd(M_v/w_i, w_i) = 1$  for each  $i \in \bar{v}$ .  $\square$

In the above lemma, (i) is due to Akihito Mori. Our proof is due to the author.

The following lemma follows from the definition immediately.

**Lemma 2.5.** *For any vertex  $v \in V_{\geq 2}$ , the followings hold.*

(i)

$$\sum_{\alpha_v \in \frac{1}{2M_v} \mathbb{Z}/\mathbb{Z}} \varepsilon_v(\alpha_v) = 0.$$

(ii) *For each  $\alpha_v \in \frac{1}{2M_v} \mathbb{Z}/\mathbb{Z}$  such that  $\varepsilon_v(\alpha_v) \neq 0$ ,  $M_v \alpha_v, M_v \alpha_v^2 \pmod{\mathbb{Z}}$  are independent of  $\alpha_v$  respectively.*

(iii)

$$\sum_{\alpha_v \in \mathcal{S}_v} \varepsilon_v(\alpha_v) q^{\alpha_v} = q^{\deg(v)/2 - 1} \prod_{i \in \bar{v}} \left( q^{1/2w_i} - q^{-1/2w_i} \right).$$

The following lemma is very important for our proof of our main theorem. We use it in a proof of Proposition 6.1 later.

**Lemma 2.6** ([FIMT21, Lemma 8]). *For  $0 \leq n \leq |\bar{v}| - 1$ , it holds*

$$\sum_{\alpha_v \in \mathcal{S}_v} \varepsilon_v(\alpha_v) \alpha_v^n = 0.$$

In the next subsection, we give a proof of this lemma by a different method from [FIMT21].

At the end of this section, we consider products for vertices. Let  $\mathcal{S} := \prod_{v \in V_{\geq 2}} \mathcal{S}_v$  and the map  $\varepsilon: (2S)^{-1}(\mathbb{Z}^{V_{\geq 2}})/\mathbb{Z}^{V_{\geq 2}} \rightarrow \{0, \pm 1\}$  be

$$\varepsilon((\alpha_v)_{v \in V_{\geq 2}}) := \prod_{v \in V_{\geq 2}} \varepsilon_v(\alpha_v).$$

This is well-defined since  $\mathcal{S} \subset (2S)^{-1}(\mathbb{Z}^{V_{\geq 2}})$  by Proposition 2.2 (v).

The following lemma follows from Lemma 2.4.

**Lemma 2.7.** (i) *The map*

$$\begin{aligned} \{\pm 1\}^{V_1} &\longrightarrow \mathcal{S} \\ (l_i)_{i \in V_1} &\longmapsto \left( \frac{1}{2} \deg(v) - 1 + \sum_{i \in \bar{v}} \frac{l_i}{2w_i} \right)_{v \in V_{\geq 2}} \end{aligned}$$

*is bijective.*

(ii) *The natural projection  $\mathcal{S} \rightarrow (2S)^{-1}(\mathbb{Z}^{V_{\geq 2}})/\mathbb{Z}^{V_{\geq 2}}$  is injective.*

**2.4. Rational functions determined by the linking matrix.** In this subsection, we introduce rational functions determined by the linking matrix and prove its basic properties.

For a vertex  $v \in V_{\geq 2}$  and complex variable  $q$ , define a rational function

$$G_v(q) := (q^{M_v} - q^{-M_v})^{2-\deg(v)} \prod_{i \in \bar{v}} \left( q^{M_v/w_i} - q^{-M_v/w_i} \right).$$

It has the following Laurant expansion.

**Lemma 2.8.** *For a vertex  $v \in V_{\geq 2}$ ,  $G_v(q)$  is expanded as*

$$G_v(q) = \sum_{\alpha_v \in \mathcal{S}_v} \varepsilon_v(\alpha_v) \sum_{n_v=0}^{\infty} \binom{n_v + \deg(v) - 3}{n_v} q^{2M_v(n_v + \alpha_v)}$$

for  $0 < |q|^{\text{sgn } M_v} < 1$ . Here we define

$$(2.1) \quad \binom{m}{l} := \begin{cases} \frac{m!}{l!(m-l)!} & \text{if } 0 \leq l \leq m, \\ 1 & \text{if } m = -1, \\ 0 & \text{otherwise.} \end{cases}$$

*Proof.* The claim follows from (iii) and the binomial theorem

$$(1-q)^{-d} = \sum_{n=0}^{\infty} \binom{n+d-1}{n} q^n$$

which holds for any integer  $d \geq 0$ . □

The rational function  $G_v(q)$  is expanded at  $q = 1$  as follows.

**Lemma 2.9.** *For a vertex  $v \in V_{\geq 2}$ , it holds*

$$G_v(q) = 2^{2-\deg(v)+|\bar{v}|} (M_v)^{1-\deg(v)+|\bar{v}|} (q-1)^{2-\deg(v)+|\bar{v}|} + O((q-1)^{3-\deg(v)+|\bar{v}|}).$$

*Proof.* The idea of our proof is due to Akihito Mori. The claim follows from

$$\begin{aligned} G_v(q) &= q^{(\deg(v)-1-\sum_{i \in \bar{v}} 1/w_i)M_v} (q^{2M_v} - 1)^{2-\deg(v)} (q-1)^{\bar{v}} \prod_{i \in \bar{v}} \frac{q^{2M_v/w_i} - 1}{q-1} \\ &= q^{(\deg(v)-1-\sum_{i \in \bar{v}} 1/w_i)M_v} (1+q+\dots+q^{2M_v-1})^{2-\deg(v)} (q-1)^{2-\deg(v)+\bar{v}} \prod_{i \in \bar{v}} \frac{q^{2M_v/w_i} - 1}{q-1} \end{aligned}$$

and

$$\lim_{q \rightarrow 1} \frac{q^{2M_v/w_i} - 1}{q - 1} = \frac{2M_v}{w_i}.$$

□

Lemma 2.6 follows from Lemma 2.9. Thus, Lemma 2.9 is a refinement of Lemma 2.6.

*Proof of Lemma 2.6.* Since

$$G_v(q) = (q^{2M_v} - 1)^{2-\deg(v)} \sum_{\alpha_v \in \mathcal{S}_v} \varepsilon_v(\alpha_v) q^{2M_v \alpha_v},$$

we have

$$G_v(e^{-t}) = \left( t^{2-\deg(v)} + O(t^{3-\deg(v)}) \right) \sum_{n=0}^{\infty} \frac{(-2M_v t)^n}{n!} \sum_{\alpha_v \in \mathcal{S}_v} \varepsilon_v(\alpha_v) \alpha_v^n.$$

Since  $G_v(e^{-t}) = O(t^{2-\deg(v)+|\bar{v}|})$  by Lemma 2.9, we obtain the claim. □

At the end of this section, we consider products for vertices.

For  $n \in \mathbb{Z}^{V_{\geq 3}}$ , we denote

$$(2.2) \quad P(n) := \prod_{v \in V_{\geq 4}} \frac{(n_v + \deg(v) - 3)(n_v + \deg(v) - 4) \cdots (n_v + 1)}{(\deg(v) - 3)!}.$$

We remark that for  $n \in \mathbb{Z}_{\geq -1}^{V_{\geq 3}}$  it holds

$$P(n) = \prod_{v \in V_{\geq 4}} \binom{n_v + \deg(v) - 3}{n_v} = \prod_{v \in V} \binom{n_v + \deg(v) - 3}{n_v}$$

by our definition of binomial coefficients in (2.1).

For each complex number  $z$ , we denote  $e(z) := e^{2\pi\sqrt{-1}z}$ . We fix a positive integer  $k$  and denote  $\zeta_k := e^{2\pi\sqrt{-1}/k}$ .

Then, the following holds by Lemma 2.8.

**Lemma 2.10.** *For  $\mu \in \mathbb{Z}^{V_{\geq 2}}$  and  $t \in \mathbb{R}^{V_{\geq 2}}$ , it holds*

$$\prod_{v \in V_{\geq 2}} G_v \left( \zeta_{2k}^{\mu_v} e^{-t_v/2M_v} \right) = \sum_{\alpha \in \mathcal{S}} \varepsilon(\alpha) \sum_{n \in \mathbb{Z}_{\geq 0}^{V_{\geq 3}}} P(n) e \left( \frac{1}{k} {}^t(n + \alpha)\mu \right) \exp(-{}^t(n + \alpha)t).$$

### 3. CALCULATIONS OF WRT INVARIANTS

In this section, we calculate WRT invariants of the plumbed homology sphere  $M(\Gamma)$ . Our starting point is the following expression by [GPPV20].

**Proposition 3.1** ([GPPV20, Equation A.12]).

$$\text{WRT}_k(M(\Gamma)) = \frac{e(|V|/8)\zeta_{4k}^{-\sum_{v \in V} (w_v+3)}}{2\sqrt{2k}^{|V|}(\zeta_{2k} - \zeta_{2k}^{-1})} \sum_{\mu \in (\mathbb{Z} \setminus k\mathbb{Z})^V / 2k\mathbb{Z}^V} \zeta_{4k}^{t\mu W\mu} \prod_{v \in V} \left( \zeta_{2k}^{\mu_v} - \zeta_{2k}^{-\mu_v} \right)^{2-\deg(v)}.$$

We can calculate WRT invariants as follows.

**Proposition 3.2.**

$$\text{WRT}_k(M(\Gamma)) = \frac{(-1)^{|V_1|} e(|V_{\geq 2}|/8) \zeta_{4k}^{-\sum_{v \in V} (w_v+3) - \sum_{i \in V_1} 1/w_i}}{2\sqrt{2k}^{|V_{\geq 2}|} (\zeta_{2k} - \zeta_{2k}^{-1}) \prod_{i \in V_1} \sqrt{|w_i|}} \sum_{\mu \in (\mathbb{Z} \setminus k\mathbb{Z})^{V_{\geq 2}} / 2k\mathbb{S}(\mathbb{Z}^{V_{\geq 2}})} e \left( -\frac{1}{4k} {}^t\mu S^{-1}\mu \right) \prod_{v \in V_{\geq 2}} G_v(\zeta_{2k}^{\mu_v}).$$

To prove this, we need the following property called ‘‘reciprocity of Gauss sums.’’

**Proposition 3.3** ([DT07, Theorem 1]). *Let  $L$  be a lattice of finite rank  $n$  equipped with a non-degenerated symmetric  $\mathbb{Z}$ -valued bilinear form  $\langle \cdot, \cdot \rangle$ . We write*

$$L' := \{y \in L \otimes \mathbb{R} \mid \langle x, y \rangle \in \mathbb{Z} \text{ for all } x \in L\}$$

for the dual lattice. Let  $0 < k \in |L'/L|\mathbb{Z}$ ,  $u \in \frac{1}{k}L$ , and  $h: L \otimes \mathbb{R} \rightarrow L \otimes \mathbb{R}$  be a self-adjoint automorphism such that  $h(L') \subset L'$  and  $\frac{k}{2} \langle y, h(y) \rangle \in \mathbb{Z}$  for all  $y \in L'$ . Let  $\sigma$  be the signature of the quadratic form  $\langle x, h(y) \rangle$ . Then it holds

$$\sum_{x \in L/kL} e\left(\frac{1}{2k} \langle x, h(x) \rangle + \langle x, u \rangle\right) = \frac{e(\sigma/8)k^{n/2}}{\sqrt{|L'/L| |\det h|}} \sum_{y \in L'/h(L')} e\left(-\frac{k}{2} \langle y + u, h^{-1}(y + u) \rangle\right).$$

*Proof of Proposition 3.2.* Idea of our proof is the same as in [MM21, Proposition 6.1] which deal with the case when  $\Gamma$  is the H-graph. However, our calculation is slightly different and clearer than it. To begin with, since  $\zeta_{2k}^{\mu_v} - \zeta_{2k}^{-\mu_v} = 0$  for  $\mu_v \in k\mathbb{Z}$ , we can write

$$(3.1) \quad \begin{aligned} & \sum_{\mu \in (\mathbb{Z} \setminus k\mathbb{Z})^V / 2k\mathbb{Z}^V} \zeta_{4k}^{t\mu W \mu} \prod_{v \in V} \left( \zeta_{2k}^{\mu_v} - \zeta_{2k}^{-\mu_v} \right)^{2-\deg(v)} \\ &= \sum_{\mu \in (\mathbb{Z}/2k\mathbb{Z})^{V_1} \oplus ((\mathbb{Z} \setminus k\mathbb{Z})/2k\mathbb{Z})^{V_{\geq 2}}} \zeta_{4k}^{t\mu W \mu} \prod_{v \in V} \left( \zeta_{2k}^{\mu_v} - \zeta_{2k}^{-\mu_v} \right)^{2-\deg(v)}. \end{aligned}$$

Since

$$\prod_{i \in V_1} \left( \zeta_{2k}^{\mu_i} - \zeta_{2k}^{-\mu_i} \right) = \prod_{i \in V_1} \sum_{l_i \in \{\pm 1\}} l_i \zeta_{2k}^{l_i \mu_i}$$

and for  $\mu = (\mu_1, \mu_{\geq 2}) \in \mathbb{Z}^V$ ,  $\mu_1 \in \mathbb{Z}^{V_1}$ ,  $\mu_{\geq 2} \in \mathbb{Z}^{V_{\geq 2}}$  it holds

$${}^t\mu W \mu = {}^t\mu_{\geq 2} W \mu_{\geq 2} + \sum_{v \in V_{\geq 2}} \sum_{i \in \bar{v}} (w_i \mu_i^2 + 2\mu_v \mu_i),$$

the right hand side of (3.1) can be written as

$$\begin{aligned} & \sum_{\mu_{\geq 2} \in ((\mathbb{Z} \setminus k\mathbb{Z})/2k\mathbb{Z})^{V_{\geq 2}}} \zeta_{4k}^{t\mu_{\geq 2} W \mu_{\geq 2}} \prod_{v \in V_{\geq 2}} \left( \zeta_{2k}^{\mu_v} - \zeta_{2k}^{-\mu_v} \right)^{2-\deg(v)} \\ & \quad \prod_{i \in \bar{v}} \sum_{l_i \in \{\pm 1\}} l_i \sum_{\mu_i \in \mathbb{Z}/2k\mathbb{Z}} \zeta_{4k}^{w_i \mu_i^2 + 2(\mu_v + l_i) \mu_i}. \end{aligned}$$

Since the last sum for  $\mu_i$  is equal to

$$\frac{e(-1/8)\sqrt{2k}}{\sqrt{|w_i|}} \sum_{\mu_i \in \mathbb{Z}/w_i\mathbb{Z}} \zeta_{4k w_i}^{-(2k\mu_i + \mu_v + l_i)^2}$$

by reciprocity of Gauss sums (Proposition 3.3), the right hand side of (3.1) can be written as

$$(3.2) \quad \begin{aligned} & \frac{e(-|V_1|/8)\sqrt{2k}^{|V_1|}}{\prod_{i \in V_1} \sqrt{|w_i|}} \sum_{\mu_{\geq 2} \in ((\mathbb{Z} \setminus k\mathbb{Z})/2k\mathbb{Z})^{V_{\geq 2}}} \zeta_{4k}^{t\mu_{\geq 2} W \mu_{\geq 2}} \prod_{v \in V_{\geq 2}} \left( \zeta_{2k}^{\mu_v} - \zeta_{2k}^{-\mu_v} \right)^{2-\deg(v)} \\ & \quad \prod_{i \in \bar{v}} \sum_{l_i \in \{\pm 1\}} l_i \sum_{\mu_i \in \mathbb{Z}/w_i\mathbb{Z}} \zeta_{4k w_i}^{-(2k\mu_i + \mu_v + l_i)^2}. \end{aligned}$$

Since  $\bigoplus_{i \in \bar{v}} \mathbb{Z}/w_i\mathbb{Z} \cong \mathbb{Z}/M_v\mathbb{Z}$  by the Chinese remainder theorem and Lemma 2.1, the last line in (3.2) is equal to

$$\sum_{\mu'_v \in \mathbb{Z}/M_v\mathbb{Z}} \prod_{i \in \bar{v}} \sum_{l_i \in \{\pm 1\}} l_i \zeta_{4k w_i}^{-(2k\mu'_v + \mu_v + l_i)^2}.$$



By replacing  $2k\mu'_v + \mu_v$  by  $\mu_v$  and using Proposition 2.2 (v), the sum for  $\mu_{\geq 2}$  in (3.2) is written as

$$\begin{aligned}
& \sum_{\mu_{\geq 2} \in (\mathbb{Z} \setminus k\mathbb{Z})^{V_{\geq 2}} / 2kS(\mathbb{Z}^{V_{\geq 2}})} \zeta_{4k}^{t\mu_{\geq 2} W \mu_{\geq 2}} \prod_{v \in V_{\geq 2}} \left( \zeta_{2k}^{\mu_v} - \zeta_{2k}^{-\mu_v} \right)^{2 - \deg(v)} \prod_{i \in \bar{v}} \sum_{l_i \in \{\pm 1\}} l_i \zeta_{4kw_i}^{-(\mu_v + l_i)^2} \\
&= \sum_{\mu_{\geq 2} \in (\mathbb{Z} \setminus k\mathbb{Z})^{V_{\geq 2}} / 2kS(\mathbb{Z}^{V_{\geq 2}})} e \left( \frac{1}{4k} \left( t\mu_{\geq 2} W \mu_{\geq 2} - \sum_{v \in V_{\geq 2}} \sum_{i \in \bar{v}} \frac{1}{w_i} \mu_v^2 \right) \right) \\
& \quad \prod_{v \in V_{\geq 2}} \left( \zeta_{2k}^{\mu_v} - \zeta_{2k}^{-\mu_v} \right)^{2 - \deg(v)} \prod_{i \in \bar{v}} \sum_{l_i \in \{\pm 1\}} l_i \zeta_{4kw_i}^{-2l_i \mu_v - l_i^2} \\
&= \zeta_{4k}^{-\sum_{i \in V_1} 1/w_i} \sum_{\mu_{\geq 2} \in (\mathbb{Z} \setminus k\mathbb{Z})^{V_{\geq 2}} / 2kS(\mathbb{Z}^{V_{\geq 2}})} e \left( \frac{1}{4k} \left( t\mu_{\geq 2} W \mu_{\geq 2} - \sum_{v \in V_{\geq 2}} \sum_{l_v \in \{\pm 1\}^{\bar{v}}} \frac{1}{w_i} \mu_v^2 \right) \right) \\
& \quad \prod_{v \in V_{\geq 2}} (-1)^{|\bar{v}|} G_v(\zeta_{2k}^{\mu_v} M_v).
\end{aligned}$$

By Proposition 2.2 (iii), it holds

$$t\mu_{\geq 2} W \mu_{\geq 2} - \sum_{v \in V_{\geq 2}} \sum_{l_v \in \{\pm 1\}^{\bar{v}}} \frac{1}{w_i} \mu_v^2 = -t\mu_{\geq 2} S^{-1} \mu_{\geq 2}.$$

Thus, we obtain the claim.  $\square$

#### 4. EXPRESSIONS OF HOMOLOGICAL BLOCKS AS FALSE THETA FUNCTIONS

In this section, we represent the homological block of  $M(\Gamma)$  as a false theta function. Our main result in this section is Proposition 4.2.

**4.1. Settings and the main result.** For the plumbed homology sphere  $M(\Gamma)$ , the homological block  $\widehat{Z}_{\Gamma}(q)$  is defined as follows.

**Definition 4.1** ([GPPV20, Subsection 3.4]). *The homological block of the plumbed homology sphere  $M(\Gamma)$  is defined as*

$$\widehat{Z}_{\Gamma}(q) = 2^{-|V|} q^{-\sum_{v \in V} (w_v + 3)/4} \text{v.p.} \int_{|z_v|=1, v \in V} \Theta_{-W, \delta}(q; z) \prod_{v \in V} (z_v - 1/z_v)^{2 - \deg(v)} \frac{dz_v}{2\pi\sqrt{-1}z_v},$$

where v.p. is the Cauchy principal value defined as

$$\text{v.p.} := \lim_{\varepsilon \rightarrow +0} \left( \int_{|z|=1+\varepsilon} + \int_{|z|=1-\varepsilon} \right),$$

$\delta := (\deg(v))_{v \in V} \in \mathbb{Z}^V$ , and

$$\Theta_{-W, \delta}(q; z) := \sum_{l \in 2\mathbb{Z}^V + \delta} q^{-tW^{-1}l/4} \prod_{v \in V} z_v^{l_v}$$

is the theta function.

Here we remark that our definition of the Cauchy principal value is half of it in [GPPV20] ([GPPV20, pp. 55]). However, our definition of the homological block  $\widehat{Z}_{\Gamma}(q)$  is the same as [GPPV20] since we multiply it by  $2^{-|V|}$ .

To state our main result, we need some notations.

For a real number  $x$ , let

$$\text{sgn}(x) := \begin{cases} 1 & x \geq 0, \\ -1 & x < 0. \end{cases}$$

For  $n \in \mathbb{Z}^{V_{\geq 2}}$ , let  $Q(n) := {}^t n S n$  be the quadratic form and  $\text{sgn}(n) := \prod_{v \in V_{\geq 2}} \text{sgn}(n_v)$ .

Our main result in this section is the following.

**Proposition 4.2.** *It holds*

$$\widehat{Z}_\Gamma(q) = (-1)^{|V_1|} 2^{|V_{\geq 3}|} q^{-(\sum_{v \in V} (w_v + 3) + \sum_{i \in V_1} 1/w_i)/4} \sum_{\alpha \in \mathcal{S}} \varepsilon(\alpha) \sum_{n \in \mathbb{Z}^{V_{\geq 3}}} \operatorname{sgn}(n) P(n) q^{Q(n+\alpha)},$$

where  $P(n)$  is the polynomial defined in (2.2).

**4.2. A proof of Proposition 4.2.** In this subsection, we prove Proposition 4.2.

To begin with, we calculate Cauchy principal values.

**Lemma 4.3.** *For  $d \in \mathbb{Z}_{>0}$  and  $l \in \mathbb{Z}$ , it holds*

$$\operatorname{v.p.} \int_{|z|=1} \frac{z^{l-1}}{(z-1/z)^{d-2}} \frac{dz}{2\pi\sqrt{-1}} = \begin{cases} -2l & d=1, l \in \{\pm 1\}, \\ 2 & d=2, l=0, \\ \operatorname{sgn}(l-d+2) \binom{(l+d)/2-2}{(l-d)/2+1} & d \geq 3, l \equiv d \pmod{2}, \\ 0 & \text{otherwise.} \end{cases}$$

*Proof.* Let  $f(z) := z^{l-1}(z-z^{-1})^{2-d}$ . Then, we have

$$\begin{aligned} \operatorname{v.p.} \int_{|z|=1} f(z) \frac{dz}{2\pi\sqrt{-1}} &= \lim_{\varepsilon \rightarrow +0} \left( \int_{|z|=1+\varepsilon} + \int_{|z|=1-\varepsilon} \right) f(z) \frac{dz}{2\pi\sqrt{-1}} \\ &= 2 \operatorname{Res}_{z=0} f(z) + \operatorname{Res}_{z=1} f(z) + \operatorname{Res}_{z=-1} f(z). \end{aligned}$$

If  $d=1, 2$ , then we obtain the claim by direct calculations since  $f(z)$  has no residues at  $z = \pm 1$ . If  $d \geq 3$ , then for any complex number  $\alpha$  it holds

$$\operatorname{Res}_{z=\alpha} f(z) = \binom{(l+d)/2-2}{(l-d)/2+1} \operatorname{Res}_{z=\alpha} \left( \frac{z^{l-d+2}}{z-z^{-1}} \right)$$

by [BMM20b, Corollary 3.3]. Since

$$\begin{aligned} \operatorname{Res}_{z=0} \left( \frac{z^{l-d+2}}{z-z^{-1}} \right) &= \begin{cases} -1 & d-l-4 \in 2\mathbb{Z}_{\geq 0} \\ 0 & \text{otherwise,} \end{cases} \\ \operatorname{Res}_{z=1} \left( \frac{z^{l-d+2}}{z-z^{-1}} \right) &= \frac{1}{2}, \\ \operatorname{Res}_{z=-1} \left( \frac{z^{l-d+2}}{z-z^{-1}} \right) &= \frac{1}{2} (-1)^{l-d}, \end{aligned}$$

we have

$$\begin{aligned} & 2 \operatorname{Res}_{z=0} \left( \frac{z^{l-d+2}}{z-z^{-1}} \right) + \operatorname{Res}_{z=1} \left( \frac{z^{l-d+2}}{z-z^{-1}} \right) + \operatorname{Res}_{z=-1} \left( \frac{z^{l-d+2}}{z-z^{-1}} \right) \\ &= \begin{cases} -2 + \frac{1}{2} + \frac{1}{2} (-1)^{l-d} & d-l-4 \in 2\mathbb{Z}_{\geq 0}, \\ \frac{1}{2} + \frac{1}{2} (-1)^{l-d} & \text{otherwise} \end{cases} \\ &= \begin{cases} -1 & l \equiv d \pmod{2}, l-d+2 < 0, \\ 1 & l \equiv d \pmod{2}, l-d+2 \geq 0, \\ 0 & l \not\equiv d \pmod{2} \end{cases} \\ &= \begin{cases} \operatorname{sgn}(l-d+2) & l \equiv d \pmod{2}, \\ 0 & l \not\equiv d \pmod{2} \end{cases} \end{aligned}$$

and thus, we obtain the claim.  $\square$

*Proof of Proposition 4.2.* By definition, we have

$$\begin{aligned} & 2^{|V|} q^{\sum_{v \in V} (w_v + 3)/4} \widehat{Z}_\Gamma(q) \\ &= \sum_{l \in 2\mathbb{Z}^V + \delta} q^{-lW^{-1}l/4} \prod_{v \in V} \operatorname{v.p.} \int_{|z_v|=1} \frac{z_v^{l_v-1}}{(z_v-1/z_v)^{\deg(v)-2}} \frac{dz_v}{2\pi\sqrt{-1}}. \end{aligned}$$

By Lemma 4.3, this is equal to

$$\sum_{\substack{l=(l_1, l_2, l_{\geq 3}), \\ l_1 \in \{\pm 1\}^{V_1}, l_2=0, \\ l_{\geq 3} \in 2\mathbb{Z}^{V_{\geq 3}} + \delta_{\geq 3}}} q^{-\iota W^{-1}l/4} \left( \prod_{i \in V_1} (-2l_i) \right) 2^{|V_2|} \\ \prod_{v \in V_{\geq 3}} \operatorname{sgn}(l_v - \deg(v) + 2) \binom{(l_v + \deg(v))/2 - 2}{(l_v - \deg(v))/2 + 1},$$

where  $\delta_{\geq 3} := (\deg(v))_{v \in V_{\geq 3}} \in \mathbb{Z}^{V_{\geq 3}}$ . Let

$$l = {}^t(l_1, l_2, l_{\geq 3}), \quad l_2 = 0, \quad l_{\geq 3} = 2n_{\geq 3} + \delta_{\geq 3} - (2, \dots, 2), \quad n_{\geq 3} \in \mathbb{Z}^{V_{\geq 3}}.$$

By Proposition 2.2 (i), we have

$$-{}^t W^{-1}l = Q(2n_{\geq 3} + \delta_{\geq 3} + {}^t T l_1 - (2, \dots, 2)) - {}^t l_1 W_1^{-1} l_1.$$

Thus, we obtain

$$(-1)^{|V_1|} 2^{|V_{\geq 3}|} q^{(\sum_{v \in V} (w_v + 3) + \sum_{i \in V_1} 1/w_i)/4} \widehat{Z}_\Gamma(q) \\ = \sum_{l_1 \in \{\pm 1\}^{V_1}} \left( \prod_{i \in V_1} l_i \right) \sum_{n_{\geq 3} \in \mathbb{Z}^{V_{\geq 3}}} q^{Q(n_{\geq 3} + \delta_{\geq 3}/2 + {}^t T l_1/2 - (1, \dots, 1))} \prod_{v \in V_{\geq 3}} \operatorname{sgn}(n_v) \binom{n_v + \deg(v) - 3}{n_v}.$$

Since  $W^{-1} \in \operatorname{Aut}(\mathbb{Z}^V)$ , every component of  $S^t T$  is integers by Proposition 2.2 (i). Thus, for any  $l_1 \in \{\pm 1\}^{V_1}$ , we have  ${}^t T l_1/2 \in (2S)^{-1}(\mathbb{Z}^{V_{\geq 2}})$ . Since

$$\frac{1}{2} {}^t T l_1 = \left( \sum_{i \in \bar{v}} \frac{l_i}{2w_i} \right)_{v \in V_{\geq 2}}$$

by the definition of  $T$  in Section 2.2, the map

$$\begin{aligned} \{\pm 1\}^{V_1} &\longrightarrow S \\ l_1 &\longmapsto \delta_{\geq 3}/2 + {}^t T l_1/2 - (1, \dots, 1) \end{aligned}$$

is bijective by Lemma 2.7 (ii). Thus, the claim follows by letting  $\alpha = \delta_{\geq 3}/2 + {}^t T l_1/2 - (1, \dots, 1)$ .  $\square$

## 5. ASYMPTOTIC FORMULAS

In this section, we develop asymptotic formulas which we need to calculate WRT invariants and radial limits of homological blocks.

**5.1. Asymptotic formulas obtained from Euler–Maclaurin summation formula.** Some useful methods have been developed to calculate radial limits of false theta functions. The method of Lawrence–Zagier [LZ99, pp. 98, Proposition] is the beginning of these. Lawrence–Zagier [LZ99] developed it by using  $L$ -functions for periodic maps. Zagier [Zag06] collected many techniques to calculate asymptotic expansions of infinite series. In particular, [Zag06, Equation (44)] is the very powerful method followed by the Euler–Maclaurin summation formula. In our setting, these methods deal with the case when  $|V_{\geq 2}| = 1$  (in this case,  $M(\Gamma)$  is a Seifert 3-manifold). Bringmann–Kasjian–Milas [BKM19, Equation (2.8)] and Bringmann–Mahlburg–Milas [BMM20a, Lemma 2.2] stated the two-variable case of [Zag06, Equation (44)].

In this subsection, we develop a more general formula (Proposition 5.2) to deal with  $n$ -variable case with polynomial weights. Our starting point is the following lemma followed by the Euler–Maclaurin summation formula.

**Lemma 5.1** ([Zag06, Equation (44)]). *Let  $N$  be a positive number and  $f: \mathbb{R}^N \rightarrow \mathbb{C}$  be a  $C^\infty$  function of rapid decay as  $x_1, \dots, x_N \rightarrow \infty$ . Fix  $\alpha \in \mathbb{R}^N$ . Then, for a variable  $t \in \mathbb{R}_{>0}^N$ , an asymptotic expansion as  $t \rightarrow +0$*

$$\sum_{n \in \mathbb{Z}_{\geq 0}^N} f(t(n + \alpha)) \sim \sum_{-1 \leq n_i, 1 \leq i \leq N} \left( \prod_{1 \leq i \leq N} \frac{B_{n_i+1}(\alpha_i)}{(n_i+1)!} \right) f^{(n)}(0) t_1^{n_1} \dots t_N^{n_N}$$

holds. Here,  $F(t) \sim G(t)$  means that  $F(t) = G(t) + O(t^R)$  for any positive number  $R$  and let  $t(n+\alpha) := (t_i(n_i + \alpha_i))_{1 \leq i \leq N}$ ,  $B_i(x)$  be the  $i$ -th Bernoulli polynomial,

$$g^{(-1)}(x) = \frac{d^{-1}}{dx^{-1}}g(x) := - \int_x^\infty g(x')dx', \quad f^{(n)}(x) := \frac{\partial^{n_1+\dots+n_N} f}{\partial x_1^{n_1} \dots \partial x_N^{n_N}}(x).$$

Here we remark that Zagier [Zag06, Equation (44)] stated for the case  $N = 1$  and Bringmann–Kaszián–Milas [BKM19, Equation (2.8)] and Bringmann–Mahlburg–Milas [BMM20a, Lemma 2.2] stated for the case  $N = 2$ .

By Lemma 5.1, we obtain the following asymptotic expansion formula.

**Proposition 5.2.** *Let  $N$  be a positive number,  $f: \mathbb{R}^N \rightarrow \mathbb{C}$  be a  $C^\infty$  function of rapid decay as  $x_1, \dots, x_N \rightarrow \infty$ , and  $P(x) = \sum_{m \in \mathbb{Z}_{\geq 0}^N} p_m x_1^{m_1} \dots x_N^{m_N}$  be a polynomial. Fix  $\alpha, \lambda \in \mathbb{R}^N$ . Then, for a variable  $t \in \mathbb{R}_{>0}^N$ , an asymptotic expansion as  $t \rightarrow +0$*

$$\sum_{n \in \mathbb{Z}_{\geq 0}^N} P(\lambda + n) f(t(\alpha + \lambda + n)) \sim \sum_{n \in \mathbb{Z}^N} t_1^{n_1} \dots t_N^{n_N} f^{(n)}(0) \sum_{m \in \mathbb{Z}_{\geq 0}^N} p_m \mathbb{B}_{m,n}(\alpha, \lambda)$$

holds. Here we define

$$\begin{aligned} \mathbb{B}_{m,n}(\alpha, \lambda) &:= \prod_{1 \leq i \leq N} \mathbb{B}_{m_i, n_i}(\alpha_i, \lambda_i), \\ \mathbb{B}_{m_i, n_i}(\alpha_i, \lambda_i) &:= \begin{cases} \sum_{0 \leq l \leq m_i + n_i + 1} b_{m_i, n_i, l} B_{m_i + n_i + 1 - l}(\lambda_i) \alpha_i^l & \text{if } n_i \geq 0, \\ \frac{m_i!}{(m_i + n_i + 1)!} (-\alpha_i)^{m_i + n_i + 1} & \text{if } -m_i - 1 \leq n_i \leq -1, \\ 0 & \text{if } n_i \leq -m_i - 2, \end{cases} \\ b_{m_i, n_i, l} &:= \frac{m_i!}{(m_i + n_i + 1 - l)!} \sum_{0 \leq k \leq l} \binom{m_i + n_i - k}{n_i} \frac{(-1)^k}{k!(l-k)!}. \end{aligned}$$

*Proof.* It suffices to show the case when  $N = 1$ . Let  $g_\alpha(m, f; x) := (x - \alpha)^m f(x)$ . By Lemma 5.1, we have

$$\begin{aligned} \sum_{n \geq 0} P(\lambda + n) f(t(\alpha + \lambda + n)) &= \sum_{m \geq 0} p_m t^{-m} \sum_{n \geq 0} (t(\alpha + \lambda + n) - t\alpha)^m f(t(\alpha + \lambda + n)) \\ &= \sum_{m \geq 0} p_m t^{-m} \sum_{n \geq 0} g_{t\alpha}(m, f; t(\alpha + \lambda + n)) \\ &\sim \sum_{m \geq 0} p_m t^{-m} \sum_{n \geq -1} \frac{B_{n+1}(\alpha + \lambda)}{(n+1)!} g_{t\alpha}^{(n)}(m, f; 0) t^n \\ &= \sum_{n > -\infty} t^n \sum_{m \geq 0} p_m \frac{B_{m+n+1}(\alpha + \lambda)}{(m+n+1)!} g_{t\alpha}^{(m+n)}(m, f; 0). \end{aligned}$$

We need the following lemma.

**Lemma 5.3.** *For any  $n \geq -1$ , it holds*

$$\begin{aligned} g_\alpha^{(n)}(m, f; 0) &= \sum_{0 \leq l \leq m} \binom{m}{l} \binom{n}{l} l! (-\alpha)^{m-l} f^{(n-l)}(0) \\ &= m! \sum_{0 \leq l \leq m} \binom{n}{l} \frac{(-\alpha)^{m-l}}{(m-l)!} f^{(n-l)}(0). \end{aligned}$$

*Proof.* We recall that  $\binom{n}{l} = 1$  for  $n = -1$  by our definition of binomial coefficients in (2.1). If  $n \geq 0$ , then by the Leibnitz rule, we have

$$g_\alpha^{(n)}(m, f; 0) = \sum_{0 \leq l \leq n} \binom{n}{l} \frac{d^l}{dx^l} (x - \alpha)^m \Big|_{x=0} f^{(n-l)}(0).$$

Thus, we obtain the claim. If  $n = -1$ , then the claim follows by induction and

$$g_\alpha^{(-1)}(m, f; 0) = (-\alpha)^m f^{(-1)}(0) + m g_\alpha^{(-1)}(m-1, f^{(-1)}; 0)$$

which follows from a partial integration.  $\square$

We turn back to the proof of Proposition 5.2. By Lemma 5.3, we have

$$\begin{aligned} & \sum_{n \geq 0} P(\lambda + n) f(t(\alpha + \lambda + n)) \\ & \sim \sum_{n > -\infty} t^n \sum_{m \geq 0} p_m \frac{B_{m+n+1}(\alpha + \lambda)}{(m+n+1)!} m! \sum_{0 \leq l \leq m} \binom{m+n}{l} \frac{(-t\alpha)^{m-l}}{(m-l)!} f^{(m+n-l)}(0). \end{aligned}$$

By replacing  $m+n-l$  by  $n$ , this can be written as

$$\sum_{n > -\infty} t^n f^{(n)}(0) \sum_{m \geq 0} p_m m! \sum_{0 \leq l \leq m} \frac{B_{n+l+1}(\alpha + \lambda)}{(n+l+1)!} \binom{n+l}{l} \frac{(-\alpha)^{m-l}}{(m-l)!}.$$

Thus, it suffices to show that

$$(5.1) \quad \mathbb{B}_{m,n}(\alpha, \lambda) = m! \sum_{0 \leq l \leq m} \frac{B_{n+l+1}(\alpha + \lambda)}{(n+l+1)!} \binom{n+l}{l} \frac{(-\alpha)^{m-l}}{(m-l)!}.$$

By the addition formula of Bernoulli polynomials

$$B_i(\alpha + \lambda) = \sum_{0 \leq j \leq i} \binom{i}{j} B_j(\lambda) \alpha^j,$$

the right hand side of (5.1) is equal to

$$m! \sum_{0 \leq l \leq m} \binom{n+l}{l} \frac{(-\alpha)^{m-l}}{(m-l)!} \sum_{0 \leq k \leq n+l+1} \frac{B_{n+l+1-k}(\lambda)}{(n+l+1-k)!} \frac{\alpha^k}{k!}.$$

By letting  $i := m+k-l, j := m-l$ , this is written as

$$\sum_{0 \leq i \leq m+n+1} B_{m+n+1-i}(\lambda) \alpha^i \frac{m!}{(m+n+1-i)!} \sum_{0 \leq j \leq i} \binom{m+n-j}{n} \frac{(-1)^j}{j!(i-j)!}.$$

Thus, we obtain (5.1) for the case when  $n \geq 0$ . Assume  $n \leq -1$ . Then, the right hand side of (5.1) is 0 unless  $m+n-j = -1$  for some  $0 \leq j \leq i \leq m+n+1$ . It occurs only for the case when  $i = j = m+n+1$ . Thus, if  $m+n+1 < 0$  then the right hand side of (5.1) is 0 and (5.1) holds. If  $m+n+1 \geq 0$ , the right hand side of (5.1) is written as

$$\frac{m!}{(m+n+1)!} (-\alpha)^{m+n+1}$$

and thus, (5.1) also holds in this case.  $\square$

**Remark 5.4.** If  $m_i = 0$ , then since  $b_{0,n_i,l} = 1/l!(n_i+1-l)!$  we have

$$\mathbb{B}_{0,n_i}(\alpha_i, \lambda_i) = B_{n_i+1}(\lambda_i + \alpha_i)$$

by the addition formula of Bernoulli polynomials. Thus, in this case, Proposition 5.2 coincides with [Zag06, Equation (44)], [BKM19, Equation (2.8)], and [BMM20a, Lemma 2.2].

**5.2. An asymptotic formula for infinite series with signatures.** We modify Proposition 5.2 as an appropriate form for false theta functions.

**Proposition 5.5.** *Let  $k$  and  $N$  be positive integers,  $C: \mathbb{Z}^N/k\mathbb{Z}^N \rightarrow \mathbb{C}$  be a periodic map,  $f: \mathbb{R}^N \rightarrow \mathbb{C}$  be a  $C^\infty$  function of rapid decay as  $x_1, \dots, x_N \rightarrow \infty$ , and  $P(x) = \sum_{m \in \mathbb{Z}_{\geq 0}^N} p_m x_1^{m_1} \cdots x_N^{m_N}$  be a*

polynomial. Fix  $\alpha \in \mathbb{R}^N$ . Then, for a variable  $t \in \mathbb{R}_{>0}^N$ , asymptotic expansions as  $t \rightarrow +0$

$$\begin{aligned} & \sum_{n \in \mathbb{Z}^N} \operatorname{sgn}(n) C(n) P(n) f(t(n + \alpha)) \\ & \sim 2^N \sum_{n \in \mathbb{Z}^N} (kt)^{n_1 + \dots + n_N} f^{(n)}(0) \sum_{m \in \mathbb{Z}_{\geq 0}^N} p_m k^{m_1 + \dots + m_N} \sum_{\lambda \in \{0, \dots, k-1\}^N} C(\lambda) \mathbb{B}_{m,n}(k^{-1}\alpha, k^{-1}\lambda) \\ & \sim 2^N \sum_{n=0}^{\infty} C(n) P(n) f(t(n + \alpha)) \end{aligned}$$

holds. Here,  $\mathbb{B}_{m,n}(\alpha, \lambda)$  is the polynomial defined in Proposition 5.2.

The point of a proof is the following lemma.

**Lemma 5.6.** For integers  $m_i \in \mathbb{Z}_{\geq 0}$  and  $n_i \in \mathbb{Z}$ , it holds

$$\mathbb{B}_{m_i, n_i}(-\alpha_i, 1 - \lambda_i) = (-1)^{m_i + n_i + 1} \mathbb{B}_{m_i, n_i}(\alpha_i, \lambda_i).$$

*Proof.* The claim follows from the definition if  $n_i \leq -1$  and the fact  $B_m(1 - \lambda) = (-1)^m B_m(\lambda)$  if  $n \geq 0$ .  $\square$

*Proof of Proposition 5.5.* The second asymptotic follows from Proposition 5.2. In the following, we prove the first asymptotic. Consider two bijections

$$\begin{aligned} \{0, \dots, k-1\} \times \mathbb{Z}_{\geq 0} & \longrightarrow \mathbb{Z}_{\geq 0} & \{0, \dots, k-1\} \times \mathbb{Z}_{\geq 0} & \longrightarrow \mathbb{Z}_{< 0} \\ (\lambda, n) & \longmapsto \lambda + kn, & (\lambda, n) & \longmapsto -(k - \lambda + kn). \end{aligned}$$

By replacing  $n$  by  $e(\lambda_e + kn)$ , we have

$$\begin{aligned} & \sum_{n \in \mathbb{Z}^N} \operatorname{sgn}(n) C(n) P(n) f(t(n + \alpha)) \\ & = \sum_{e \in \{\pm 1\}^N} \left( \prod_{1 \leq i \leq N} e_i \right) \sum_{\lambda \in \{0, \dots, k-1\}^N} C(e(\lambda_e + kn)) \sum_{n \in \mathbb{Z}_{\geq 0}^N} P(e(\lambda_e + kn)) f(t(e(\lambda_e + kn) + \alpha)). \end{aligned}$$

Here, for  $e \in \{\pm 1\}^N$  and  $\lambda \in \mathbb{Z}^N$  let  $e\lambda, \lambda_e \in \mathbb{Z}^N$  be the vector whose  $i$ -th components are

$$(e\lambda)_i := e_i \lambda_i, \quad (\lambda_e)_i := \begin{cases} \lambda_i & e_i = 1, \\ k - \lambda_i & e_i = -1 \end{cases}$$

respectively. Let  $P_e(x) := P(ex)$  and  $f_e(x) := f(ex)$ . Since  $e\lambda_e \equiv \lambda \pmod{k}$ , we can write the above equation as

$$\sum_{e \in \{\pm 1\}^N} \left( \prod_{1 \leq i \leq N} e_i \right) \sum_{\lambda \in \{0, \dots, k-1\}^N} C(\lambda) \sum_{n \in \mathbb{Z}_{\geq 0}^N} P_e(\lambda_e + kn) f_e(t(\lambda_e + kn + e\alpha)).$$

Since  $f_e^{(n)}(x) = e_1^{n_1} \dots e_N^{n_N} f^{(n)}(x)$  for any  $n \in \mathbb{Z}_{\geq 0}^N$  and for  $m \in \mathbb{Z}_{\geq 0}^N$ , the coefficient of  $x_1^{m_1} \dots x_N^{m_N}$  in  $P_e(x)$  is  $e_1^{m_1} \dots e_N^{m_N} p_m$ , we have an asymptotic expansion as  $t \rightarrow +0$

$$\begin{aligned} & \sum_{n \in \mathbb{Z}^N} (kt)^{n_1 + \dots + n_N} f^{(n)}(0) \sum_{m \in \mathbb{Z}_{\geq 0}^N} p_m k^{m_1 + \dots + m_N} \sum_{e \in \{\pm 1\}^N} \left( \prod_{1 \leq i \leq N} e_i^{m_i + n_i + 1} \right) \\ & \sum_{\lambda \in \{0, \dots, k-1\}^N} C(\lambda) \mathbb{B}_{m,n}(k^{-1}e\alpha, k^{-1}\lambda_e) \end{aligned}$$

by Proposition 5.2. Then, we obtain the claim since it holds

$$\mathbb{B}_{m,n}(k^{-1}e\alpha, k^{-1}\lambda_e) = \mathbb{B}_{m,n}(k^{-1}\alpha, k^{-1}\lambda) \prod_{1 \leq i \leq N} e_i^{m_i + n_i + 1}$$

by Lemma 5.6.  $\square$

**5.3. Asymptotic expansion of infinite series with weighted Gauss sums.** In this subsection, we study a family of infinite series  $F(f; t)$  with weighted Gauss sums, which involves both WRT invariants and homological blocks. We start with the following asymptotic formula following from Proposition 5.2.

**Corollary 5.7.** *For a  $C^\infty$  function  $f: \mathbb{R}^{V_{\geq 2}} \rightarrow \mathbb{C}$  of rapid decay as  $x_v \rightarrow \infty$  for each  $v \in V_{\geq 2}$ , let*

$$F(f; t) := \sum_{\alpha \in \mathcal{S}} \varepsilon(\alpha) \sum_{n \in \mathbb{Z}_{\geq 0}^{V_{\geq 3}}} e\left(\frac{1}{k}Q(n + \alpha)\right) P(n) f(t(n + \alpha))$$

be a series for a variable  $t \in \mathbb{R}_{> 0}^{V_{\geq 3}}$ , where  $t(n + \alpha) := (t_v(n_v + \alpha_v))_{v \in V_{\geq 2}}$  and  $t_v = 0$  for  $v \in V_{\geq 2}$ . Then, it holds

$$F(f; t) \sim \sum_{n \in \mathbb{Z}^{V_{\geq 3}}} c_n f^{(n)}(0) \left( \prod_{v \in V_{\geq 3}} t_v^{n_v} \right)$$

as  $t \rightarrow +0$ , where we define

$$c_n = \sum_{m \in \mathbb{Z}_{\geq 0}^{V_{\geq 3}}} p_m \left( \prod_{v \in V_{\geq 3}} k^{m_v + n_v} \right) \sum_{\alpha \in \mathcal{S}} \varepsilon(\alpha) \sum_{\lambda \in \{0, \dots, k-1\}^{V_{\geq 3}}} e\left(\frac{1}{k}Q(\lambda + \alpha)\right) \mathbb{B}_{m, n}(\alpha, \lambda)$$

for  $n \in \mathbb{Z}^{V_{\geq 3}}$ .

We define two  $C^\infty$  function of rapid decay  $f_1, f_2: \mathbb{R}^{V_{\geq 2}} \rightarrow \mathbb{C}$  as  $f_1(x) := \exp(-\sum_{v \in V_{\geq 2}} x_v)$  and  $f_2: \mathbb{R}^{V_{\geq 2}} \rightarrow \mathbb{C}$  respectively. Then, WRT invariants and limit values of homological blocks can be written as limit values of  $F(f_1; t)$  and  $F(f_2; t)$  as  $t \rightarrow 0$  shown later in Propositions 5.8 and 6.3, respectively.

Here, we need to prove that  $F(f; t)$  has a limit as  $t \rightarrow 0$ , that is,  $c_n$  vanishes for  $n \in \mathbb{Z}^{V_{\geq 3}}$  with non-positive components. Such a result is called ‘‘a vanishing result of weighted Gauss sums’’ in [MM21], and it is the most important part in a proof of the conjecture of Gukov–Pei–Putrov–Vafa [GPPV20, Conjecture 2.1, Equation (A.28)]. In the previous works [BMM20a, MM21], it is proved by direct calculations ([BMM20a, Theorem 4.1], [MM21, Proposition 4.2]). On the other hand, our proof is based on the above asymptotic expansion in Corollary 5.7. Since  $c_n$  is independent of  $f$  in this asymptotic expansion, it suffices to consider not for any  $f$ , but for  $f_1$ . Since  $F(f_1; t)$  is a rational function shown later in Lemma 5.10, vanishings results of  $c_n$  is equivalent to holomorphy of  $F(f_1; t)$ , which is proved in Section 6.

The asymptotic expansion of homological blocks is equal to it of  $F(f_2; t)$  as follows.

**Proposition 5.8.** *It holds*

$$\lim_{q \rightarrow \zeta_k} \widehat{Z}_\Gamma(q) = (-1)^{|V_1|} \zeta_{4k}^{-\sum_{v \in V} (w_v + 3) - \sum_{i \in V_1} 1/w_i} \lim_{t \rightarrow +0} F(f_2; t)$$

if the limit values in both sides converge.

This proposition follows from Propositions 4.2 and 5.5.

In the last of this subsection, we prove that  $F(f_1; t)$  is a rational function related to WRT invariants. To begin with, we replace quadratic forms  $Q(n + \alpha)$  as linear forms in the indices of  $\zeta_k$  in the definition of  $F(f; t)$ .

**Lemma 5.9.** *For a  $C^\infty$  function  $f: \mathbb{R}^{V_{\geq 2}} \rightarrow \mathbb{C}$  of rapid decay as  $x_v \rightarrow \infty$  for each  $v \in V_{\geq 2}$ , it holds*

$$F(f; t) = \frac{e(|V_{\geq 2}|/8)}{\sqrt{2k}^{|V_{\geq 2}|} \prod_{i \in V_1} \sqrt{|w_i|}} \sum_{\mu \in \mathbb{Z}^{V_{\geq 2}/2k\mathcal{S}}(\mathbb{Z}^{V_{\geq 2}})} e\left(-\frac{1}{4k}t\mu S^{-1}\mu\right) \sum_{\alpha \in \mathcal{S}} \varepsilon(\alpha) \sum_{n \in \mathbb{Z}_{\geq 0}^{V_{\geq 3}}} e\left(\frac{1}{k}t(n + \alpha)\mu\right) P(n) f(t(n + \alpha)).$$

*Proof.* Let

$$G(2kS) := \sum_{\mu \in \mathbb{Z}^{V_{\geq 2}} / 2kS(\mathbb{Z}^{V_{\geq 2}})} e\left(-\frac{1}{4k} t \mu S^{-1} \mu\right).$$

Then, we have  $G(2kS) = \left(\sqrt{2k} e(-1/8)\right)^{|V_{\geq 2}|} \sqrt{\det S}$  by Proposition 3.3. Here we remark  $\det S = \prod_{i \in V_1} |w_i|$  by Proposition 2.2 (ii). Then, we can write

$$\begin{aligned} & G(2kS)F(f; t) \\ &= \sum_{\mu \in \mathbb{Z}^{V_{\geq 2}} / 2kS(\mathbb{Z}^{V_{\geq 2}})} \sum_{\alpha \in \mathcal{S}} \varepsilon(\alpha) \sum_{n \in \mathbb{Z}_{\geq 0}^{V_{\geq 3}}} e\left(\frac{1}{k} \left(Q(n + \alpha) - \frac{1}{4} t \mu S^{-1} \mu\right)\right) P(n) f(t(n + \alpha)). \end{aligned}$$

By replacing  $\mu$  by  $\mu - 2S(n + \alpha)$ , we obtain the claim.  $\square$

By this lemma, we obtain the following representation.

**Lemma 5.10.**

$$F(f_1; t) = \frac{e(|V_{\geq 2}|/8)}{(2k)^{|V_{\geq 2}|} \prod_{i \in V_1} \sqrt{|w_i|}} \sum_{\mu \in \mathbb{Z}^{V_{\geq 2}} / 2kS(\mathbb{Z}^{V_{\geq 2}})} e\left(-\frac{1}{4k} t \mu S^{-1} \mu\right) \prod_{v \in V_{\geq 2}} G_v\left(\zeta_{2kM_v}^{\mu_v} e^{-t_v/2M_v}\right).$$

In particular,  $F(f_1; t)$  is a rational function.

*Proof.* By Lemma 5.9, we have

$$\begin{aligned} G(2kS)F(f_1; t) &= \sum_{\mu \in \mathbb{Z}^{V_{\geq 2}} / 2kS(\mathbb{Z}^{V_{\geq 2}})} e\left(-\frac{1}{4k} t \mu S^{-1} \mu\right) \\ &\quad \sum_{\alpha \in \mathcal{S}} \varepsilon(\alpha) \sum_{n \in \mathbb{Z}_{\geq 0}^{V_{\geq 3}}} e\left(\frac{1}{k} t(n + \alpha)\mu\right) P(n) \exp\left(-\sum_{v \in V_{\geq 3}} t_v(n_v + \alpha_v)\right). \end{aligned}$$

By applying Lemma 2.10 for this, we obtain the claim.  $\square$

## 6. A PROOF OF THE MAIN THEOREM

In this section, we prove holomorphy of  $F(f_1; t)$  (that is, vanishings of  $c_n$ ) and our main result.

**6.1. The main result.** In this subsection, we state our main result without a proof.

**Proposition 6.1.** (i) For each vertex  $v \in V_{\geq 3}$ , the order of  $F(f_1; t)$  at  $t_v = 0$  satisfies

$$\text{ord}_{t_v=0} F(f_1; t) \geq \min\{0, |\bar{v}| + 2 - \deg(v)\}.$$

(ii) If  $|\bar{v}| + 2 - \deg(v) > 0$  for each vertex  $v \in V_{\geq 3}$ , then

$$F(f_1; 0) = c_0 = \frac{e(|V_{\geq 2}|/8)}{\sqrt{2k}^{|V_{\geq 2}|} \prod_{i \in V_1} \sqrt{|w_i|}} \sum_{\mu \in (\mathbb{Z} \setminus k\mathbb{Z})^{V_{\geq 2}} / 2kS(\mathbb{Z}^{V_{\geq 2}})} e\left(-\frac{1}{4k} t \mu S^{-1} \mu\right) \prod_{v \in V_{\geq 2}} G_v\left(\zeta_{2kM_v}^{\mu_v}\right).$$

We obtain the following corollary by Proposition 6.1 (i).

**Corollary 6.2.** Suppose  $|\bar{v}| + 2 - \deg(v) \geq 0$  for each vertex  $v \in V_{\geq 3}$ . Then,  $c_n = 0$  holds for  $n \in \mathbb{Z}^{V_{\geq 3}}$  such that  $n_v \leq -1$  for some vertex  $v \in V_{\geq 3}$ . In particular, for a  $C^\infty$  function  $f: \mathbb{R}^{V_{\geq 2}} \rightarrow \mathbb{C}$  of rapid decay as  $x_v \rightarrow \infty$  for each  $v \in V_{\geq 2}$ , the limit  $\lim_{t \rightarrow +0} F(f; t)$  converges and it can be written as

$$\lim_{t \rightarrow +0} F(f; t) = c_0,$$

which is independent of  $f$ .

We obtain the following proposition by combining Proposition 6.1 (ii) and Proposition 3.2.

**Proposition 6.3.** If  $|\bar{v}| + 2 - \deg(v) > 0$  for any vertices  $v \in V_{\geq 3}$ , then it holds

$$\text{WRT}_k(M(\Gamma)) = \frac{(-1)^{|V_1|} \zeta_{4k}^{-\sum_{v \in V} (w_v + 3) - \sum_{i \in V_1} 1/w_i}}{2(\zeta_{2k} - \zeta_{2k}^{-1})} F(f_1; 0).$$



We obtain our main result by combining Propositions 5.8 and 6.3.

**Corollary 6.4** (Theorem 1.2). *If  $|\bar{v}| + 2 - \deg(v) > 0$  for any vertices  $v \in V_{\geq 3}$ , then it holds*

$$\text{WRT}_k(M(\Gamma)) = \frac{1}{2(\zeta_{2k} - \zeta_{2k}^{-1})} \lim_{q \rightarrow \zeta_k} \widehat{Z}_\Gamma(q).$$

**6.2. A proof of the main result.** Finally, we prove Proposition 6.1. We need the following lemma.

**Lemma 6.5.** *Let  $k, M \in \mathbb{Z} \setminus \{0\}$  and  $a, b \in \mathbb{Z}$  be integers such that  $\gcd(a, M) = 1$ . For  $\alpha \in \frac{1}{2M}\mathbb{Z}$  such that  $\gcd(2M\alpha, M) = 1$ , the complex number*

$$\sum_{\mu \in \mathbb{Z}/k\mathbb{Z} + \alpha} e\left(\frac{M}{k}(a\mu^2 + b\mu)\right)$$

*depends only on  $M\alpha^2, M\alpha \bmod \mathbb{Z}$ .*

This lemma is a little generalisation of [MM21, Lemma 4.7 and 4.8] which is based on the proof of [BMM20a, Theorem 4.1]. In this lemma, we remove the assumption  $\gcd(2M, 2M\alpha) = 1$  in [MM21, Lemma 4.7 and 4.8]. Although our proof is essentially the same as in [MM21, Lemma 4.7 and 4.8], we give a proof for the convenience of readers.

*Proof. Case 1:* Suppose  $\gcd(M, k) > 1$ . We can write

$$\sum_{\mu \in \mathbb{Z}/k\mathbb{Z} + \alpha} e\left(\frac{M}{k}(a\mu^2 + b\mu)\right) = e\left(\frac{M}{k}(a\alpha^2 + b\alpha)\right) \sum_{\mu \in \mathbb{Z}/k\mathbb{Z}} e\left(\frac{1}{k}(Ma\mu^2 + (2M\alpha a + Mb)\mu)\right).$$

By the assumption, we have  $\gcd(a, M) = 1$ ,  $\gcd(2M\alpha, M) = 1$  and thus,  $\gcd(2M\alpha a + Mb, M) = 1$  holds. By combining with  $\gcd(M, k) \mid \gcd(Ma, k)$  and  $1 < \gcd(M, k) \mid M$ , we obtain  $\gcd(Ma, k) \nmid 2M\alpha a + Mb$ . Thus, it is 0 by the well-known vanishing result of one-variable Gauss sums ([MM21, Lemma 4.3]).

**Case 2:** Suppose  $\gcd(M, k) = 1$ . Let  $M^* \in \mathbb{Z}$  be an integer such that  $MM^* \equiv 1 \pmod{k}$ . For any  $\mu \in \mathbb{Z}$ , we have

$$\begin{aligned} & M(a(\mu + \alpha)^2 + b(\mu + \alpha)) - M(a(\mu + MM^*\alpha)^2 + b(\mu + MM^*\alpha)) \\ &= M((1 - (MM^*)^2)a\alpha^2 + (1 - MM^*)(b + 2a\mu)\alpha) \\ &\equiv (1 - MM^*)((1 + MM^*)a\alpha^2 + b\alpha) \pmod{k\mathbb{Z}}. \end{aligned}$$

Thus, we obtain

$$\begin{aligned} & \sum_{\mu \in \mathbb{Z}/k\mathbb{Z} + \alpha} e\left(\frac{M}{k}(a\mu^2 + b\mu)\right) \\ &= e\left(\frac{1 - MM^*}{k}M((1 + MM^*)a\alpha^2 + b\alpha)\right) \sum_{m \in \mathbb{Z}/k\mathbb{Z} + MM^*\alpha} e\left(\frac{M}{k}(a\mu^2 + b\mu)\right). \end{aligned}$$

Since this sum depends only on  $M\alpha^2 \bmod \mathbb{Z}$  and  $M\alpha \bmod \mathbb{Z}$ , we obtain the claim.  $\square$

*Proof of Proposition 6.1.* For (i), fix a vertex  $v \in V_{\geq 3}$  and let

$$\begin{aligned} F(t) &:= (2ke(-1/8))^{|V_{\geq 2}|} \sqrt{\det SF(f_1; t)} \\ F_v(t) &:= \sum_{\mu \in (\mathbb{Z}^{V_{\geq 2} \setminus \{v\}} \oplus k\mathbb{Z})/2kS(\mathbb{Z}^{V_{\geq 2}})} e\left(-\frac{1}{4k}t\mu S^{-1}\mu\right) \prod_{v' \in V_{\geq 2}} G_{v'}\left(\zeta_{2kM_{v'}}^{\mu_{v'}} e^{-t_{v'}/2M_{v'}}\right). \end{aligned}$$

By Lemma 5.10, we can write

$$\begin{aligned} & F(t) - F_v(t) \\ &= \sum_{\mu \in (\mathbb{Z}^{V_{\geq 2} \setminus \{v\}} \oplus (\mathbb{Z} \setminus k\mathbb{Z})) / 2kS(\mathbb{Z}^{V_{\geq 2}})} e\left(-\frac{1}{4k}t\mu S^{-1}\mu\right) \prod_{v' \in V_{\geq 2}} G_{v'}\left(\zeta_{2kM_{v'}}^{\mu_{v'}} e^{-t_{v'}/2M_{v'}}\right). \end{aligned}$$

Since the rational function

$$G_v(\zeta_{2k}^{\mu_v} e^{-t_v/2}) = (\zeta_{2k}^{\mu_v} e^{-t_v/2} - \zeta_{2k}^{-\mu_v} e^{t_v/2})^{2-\deg(v)} \prod_{i \in \bar{v}} \left( \left( \zeta_{2k}^{\mu_v} e^{-t_v/2} \right)^{M_v/w_i} - \left( \zeta_{2k}^{\mu_v} e^{-t_v/2} \right)^{-M_v/w_i} \right)$$

is holomorphic at  $t_v = 0$  for any  $\mu_v \in \mathbb{Z} \setminus k\mathbb{Z}$ , the rational function  $F(t) - F_v(t)$  is also holomorphic at  $t_v = 0$ . Thus, it suffices to show  $\text{ord}_{t_v=0} F_v(t) \geq |\bar{v}| + 2 - \deg(v)$ . Let

$$a_v := M_v \left( -w_v + \sum_{i \in \bar{v}} \frac{1}{w_i} \right), \quad b_v := \sum_{v' \in V_{\geq 3}, \{v, v'\} \in E} \mu_{v'}$$

for each  $\mu' \in \mathbb{Z}^{V_{\geq 2} \setminus \{v\}}$ . Under this notations, we can write

$$(6.1) \quad F_v(t) := \sum_{\mu' \in \bigoplus_{v' \in V_{\geq 2} \setminus \{v\}} \mathbb{Z}/2kM_{v'}\mathbb{Z}} e \left( -\frac{1}{4k} {}_t\mu' S^{-1} \mu' \right) \left( \prod_{v \in V_{\geq 2} \setminus \{v\}} G_{v'} \left( \zeta_{2kM_{v'}}^{\mu_{v'}} e^{-t_{v'}/2M_{v'}} \right) \right) \\ \sum_{\mu_v \in \mathbb{Z}/2M_v\mathbb{Z}} e \left( -\frac{1}{4k} \left( \frac{ka'_v}{M_v} \mu_v^2 + 2b_v \mu_v \right) \right) G_v \left( \zeta_{2M_v}^{\mu_v} e^{-t_v/2M_v} \right).$$

Since we have

$$G_v(q) = (q^{2M_v} - 1)^{2-\deg(v)} \sum_{\alpha_v \in \mathcal{S}_v} \varepsilon_v(\alpha_v) q^{2M_v \alpha_v}$$

by the definition of  $G_v(q)$ , the last line of (6.1) is equal to

$$(e^{-t_v} - 1)^{2-\deg(v)} \sum_{\alpha_v \in \mathcal{S}_v} \varepsilon_v(\alpha_v) e^{-\alpha_v t_v} \sum_{\mu_v \in \mathbb{Z}/2M_v\mathbb{Z}} e \left( -\frac{ka'_v}{4M_v} \mu_v^2 + \left( \alpha_v - \frac{b_v}{2} \right) \mu_v \right).$$

By applying reciprocity of Gauss sums (Proposition 3.3), this can be written as

$$(6.2) \quad \frac{e(1/8) \sqrt{ka'_v}}{\sqrt{2|M_v|}} (e^{-t_v} - 1)^{2-\deg(v)} \sum_{\alpha_v \in \mathcal{S}_v} \varepsilon_v(\alpha_v) e^{-\alpha_v t_v} \sum_{\mu_v \in \mathbb{Z}/ka_v\mathbb{Z}} e \left( -\frac{M_v}{ka'_v} \left( \mu_v + \alpha_v - \frac{b_v}{2} \right)^2 \right).$$

Since the second summation is independent of  $\alpha_v$  by Lemma 2.5 (ii), Lemma 2.4 (iii), and Lemma 6.5, (6.2) is the multiplication of  $G_v(e^{-t_v/2M_v})$  by a constant independent of  $\alpha_v$ . Thus, we obtain  $\text{ord}_{t_v=0} F_v(t) \geq \text{ord}_{t_v=0} G_v(e^{-t_v/2M_v})$ . Since  $\text{ord}_{t_v=0} G_v(e^{-t_v/2M_v}) = 2 - \deg(v) + |\bar{v}|$  holds by Lemma 2.9, the claim holds.

Finally we prove (ii). If  $|\bar{v}| + 2 - \deg(v) > 0$  for each vertex  $v \in V_{\geq 3}$ , then  $G_v(e^{-t_v/2M_v})$  has a zero at  $t_v = 0$  by Lemma 2.9, thus  $F_v(t)$  also has a zero. Thus, we obtain

$$F(t)|_{t_v=0} = F(t) - F_v(t)|_{t_v=0}.$$

Fix any vertex  $v' \in V_{\geq 2} \setminus \{v\}$  and let

$$F_{v,v'}(t) := F(t) - F_v(t) - \sum_{\mu \in \left( \mathbb{Z}^{V_{\geq 2} \setminus \{v, v'\}} \oplus (\mathbb{Z} \setminus k\mathbb{Z})^{\{v, v'\}} \right) / 2kS(\mathbb{Z}^{V_{\geq 2}})} e \left( -\frac{1}{4k} {}_t\mu S^{-1} \mu \right) \\ \prod_{v' \in V_{\geq 2}} G_{v'} \left( \zeta_{2kM_{v'}}^{\mu_{v'}} e^{-t_{v'}/2M_{v'}} \right).$$

By the same argument in a proof of (i), the rational function  $F_{v,v'}(t)$  has a zero at  $t_{v'} = 0$ . By induction, we obtain the claim.  $\square$

## REFERENCES

- [AH12] Jørgen Ellegaard Andersen and Benjamin Himpel. The Witten-Reshetikhin-Turaev invariants of finite order mapping tori II. *Quantum Topol.*, 3(3-4):377–421, 2012.
- [AM20] J. Andersen and W. Mistegård. Resurgence analysis of quantum invariants of Seifert fibered homology spheres. September 2020. arXiv:1811.05376v3.
- [And13] Jørgen Ellegaard Andersen. The Witten-Reshetikhin-Turaev invariants of finite order mapping tori I. *J. Reine Angew. Math.*, 681:1–38, 2013.
- [BKM19] Kathrin Bringmann, Jonas Kaszian, and Antun Milas. Higher depth quantum modular forms, multiple Eichler integrals, and  $\mathfrak{sl}_3$  false theta functions. *Res. Math. Sci.*, 6(2):Paper No. 20, 41, 2019.

- [BKMN21] Kathrin Bringmann, Jonas Kaszian, Antun Milas, and Caner Nazaroglu. Higher depth false modular forms. *arXiv:2109.00394*, 2021.
- [BMM20a] Kathrin Bringmann, Karl Mahlburg, and Antun Milas. Higher depth quantum modular forms and plumbed 3-manifolds. *Lett. Math. Phys.*, 110(10):2675–2702, 2020.
- [BMM20b] Kathrin Bringmann, Karl Mahlburg, and Antun Milas. Quantum modular forms and plumbing graphs of 3-manifolds. *J. Combin. Theory Ser. A*, 170:105145, 32, 2020.
- [BN19] Kathrin Bringmann and Caner Nazaroglu. A framework for modular properties of false theta functions. *Res. Math. Sci.*, 6(3):Paper No. 30, 23, 2019.
- [BW05] Chris Beasley and Edward Witten. Non-abelian localization for Chern-Simons theory. *J. Differential Geom.*, 70(2):183–323, 2005.
- [Cha16] L. Charles. On the Witten asymptotic conjecture for Seifert manifolds. *arXiv:1605.04124*, 2016.
- [Chu17] Sungbong Chun. A resurgence analysis of the  $SU(2)$  Chern-Simons partition functions on a Brieskorn homology sphere  $\Sigma(2, 5, 7)$ . *arXiv:1701.03528v1*, 2017.
- [CM15] L. Charles and J. Marché. Knot state asymptotics II: Witten conjecture and irreducible representations. *Publ. Math. Inst. Hautes Études Sci.*, 121:323–361, 2015.
- [DT07] Florian Deloup and Vladimir Turaev. On reciprocity. *J. Pure Appl. Algebra*, 208(1):153–158, 2007.
- [FG91] Daniel S. Freed and Robert E. Gompf. Computer calculation of Witten’s 3-manifold invariant. *Comm. Math. Phys.*, 141(1):79–117, 1991.
- [FIMT21] Hiroyuki Fuji, Kohei Iwaki, Hitoshi Murakami, and Yuji Terashima. Witten–Reshetikhin–Turaev function for a knot in Seifert manifolds. *Communications in Mathematical Physics*, 2021.
- [GMP16] Sergei Gukov, Marcos Marino, and Pavel Putrov. Resurgence in complex Chern-Simons theory. 2016. *arXiv:1605.07615v2*.
- [GPPV20] Sergei Gukov, Du Pei, Pavel Putrov, and Cumrun Vafa. BPS spectra and 3-manifold invariants. *J. Knot Theory Ramifications*, 29(2):2040003, 85, 2020.
- [Hik05a] Kazuhiro Hikami. On the quantum invariant for the Brieskorn homology spheres. *Internat. J. Math.*, 16(6):661–685, 2005.
- [Hik05b] Kazuhiro Hikami. Quantum invariant, modular form, and lattice points. *Int. Math. Res. Not.*, (3):121–154, 2005.
- [Hik06a] Kazuhiro Hikami. On the quantum invariants for the spherical Seifert manifolds. *Comm. Math. Phys.*, 268(2):285–319, 2006.
- [Hik06b] Kazuhiro Hikami. Quantum invariants, modular forms, and lattice points. II. *J. Math. Phys.*, 47(10):102301, 32, 2006.
- [HT02] Søren Kold Hansen and Toshie Takata. Quantum invariants of Seifert 3-manifolds and their asymptotic expansions. In *Invariants of knots and 3-manifolds (Kyoto, 2001)*, volume 4 of *Geom. Topol. Monogr.*, pages 69–87. Geom. Topol. Publ., Coventry, 2002.
- [Jef92] Lisa C. Jeffrey. Chern-Simons-Witten invariants of lens spaces and torus bundles, and the semiclassical approximation. *Comm. Math. Phys.*, 147(3):563–604, 1992.
- [LZ99] Ruth Lawrence and Don Zagier. Modular forms and quantum invariants of 3-manifolds. volume 3, pages 93–107. 1999. Sir Michael Atiyah: a great mathematician of the twentieth century.
- [MM21] Akihito Mori and Yuya Murakami. Witten-Reshetikhin-Turaev invariants, homological blocks, and quantum modular forms. *arXiv:2110.10958*, 2021.
- [MT21] Toshiki Matsusaka and Yuji Terashima. Modular transformations of homological blocks for Seifert fibered homology 3-spheres. *arXiv:2112.06210*, 2021.
- [Neu80] Walter D. Neumann. An invariant of plumbed homology spheres. In *Topology Symposium, Siegen 1979 (Proc. Sympos., Univ. Siegen, Siegen, 1979)*, volume 788 of *Lecture Notes in Math.*, pages 125–144. Springer, Berlin, 1980.
- [Neu81] Walter D. Neumann. A calculus for plumbing applied to the topology of complex surface singularities and degenerating complex curves. *Trans. Amer. Math. Soc.*, 268(2):299–344, 1981.
- [Roz94] L. Rozansky. A large  $k$  asymptotics of Witten’s invariant of Seifert manifolds. In *Proceedings of the Conference on Quantum Topology (Manhattan, KS, 1993)*, pages 307–354. World Sci. Publ., River Edge, NJ, 1994.
- [Roz96] L. Rozansky. Residue formulas for the large  $k$  asymptotics of Witten’s invariants of Seifert manifolds. The case of  $SU(2)$ . *Comm. Math. Phys.*, 178(1):27–60, 1996.
- [RT91] N. Reshetikhin and V. G. Turaev. Invariants of 3-manifolds via link polynomials and quantum groups. *Invent. Math.*, 103(3):547–597, 1991.
- [Wit89] Edward Witten. Quantum field theory and the Jones polynomial. *Comm. Math. Phys.*, 121(3):351–399, 1989.
- [Wu21] David H. Wu. Resurgent analysis of  $SU(2)$  Chern-Simons partition function on Brieskorn spheres  $\Sigma(2, 3, 6n + 5)$ . *J. High Energy Phys.*, (2):Paper No. 008, 18, 2021.
- [Zag06] D. Zagier. *The Mellin transform and other useful analytic techniques*. In Appendix of E. Zeidler, Quantum Field Theory I: Basics in Mathematics and Physics. A Bridge Between Mathematicians and Physicists, pages 305–323. Springer, Berlin, 2006.
- [Zag10] Don Zagier. Quantum modular forms. In *Quanta of maths*, volume 11 of *Clay Math. Proc.*, pages 659–675. Amer. Math. Soc., Providence, RI, 2010.