

An R Package AZIAD for Analyzing Zero-Inflated and Zero-Altered Data

Niloufar Dousti Mousavi¹, Hani Aldirawi² and Jie Yang¹

¹University of Illinois at Chicago and

²California State University, San Bernardino

May 4, 2022

Abstract

Sparse data with a large portion of zeros arise in many scientific disciplines. Modeling sparse data is very challenging due to the skewness of the distribution. We adopt bootstrapped Monte Carlo method to estimate the p-value of the Kolmogorov-Smirnov test, as well as bootstrapped likelihood ratio tests for zero-inflated and zero-altered (or hurdle) model selection. Our new package AZIAD provides miscellaneous functions to simulate zero-inflated or zero-altered data and calculate maximum likelihood estimates of unknown parameters for a large class of discrete or continuous distributions. In addition, we calculate the Fisher information matrix and the confidence intervals of unknown parameters. Compared with other R packages available so far, our package covers many more types of zero-inflated and zero-altered distributions, provides more accurate estimates for unknown parameters, and achieves higher power for model selection. To facilitate the potential users, in this paper we provide theoretical justifications and detailed formulae for functions in AZIAD and illustrate the use of them with executable R code and real dataset.

Key words and phrases: Kolmogorov-Smirnov (KS) test, Zero-inflated model, Hurdle model, Model selection, Fisher information matrix

1 Introduction

Sparse or zero-inflated data arise frequently from a rich variety of scientific disciplines including microbiome (Xia et al., 2018; Metwally et al., 2018), gene expression (McDavid et al., 2019), health care (Majo and van Soest, 2011), insurance claim (Boucher et al., 2009), security (Chen et al., 2018), and more. Modeling sparse data is very challenging due to the high proportion of zero values and the skewness of the distribution. Zero-inflated Poisson (ZIP), zero-inflated negative binomial (ZINB), Poisson hurdle (PH), and negative binomial hurdle (NBH) models have been widely used to model sparse data (Metwally et al., 2018; Aldirawi et al., 2019; Metwally et al., 2019).

On one hand, more and more zero-inflated models have been proposed under different circumstances. On the other hand, it becomes more and more difficult for researchers

and practitioners to choose the most appropriate model for their sparse data. Some partial comparisons among existing models have been done for certain sparse data. For examples, Xu et al. (2015) recommended ZINB and NBH models for microbiome data after comparing the performance of Poisson, ZIP, PH, NB (negative binomial), ZINB, and NBH models; while Aldirawi et al. (2019) indicated that two new models, zero-inflated beta negative binomial (ZIBNB) and beta negative binomial hurdle (BNBH) models, are more appropriate for microbiome data examples.

For analyzing zero-inflated data, there are some available R packages from the Comprehensive R Archive Network (CRAN, <https://cran.r-project.org/>) including `bzinb` (Cho et al., 2018), `hurdlr` (Balderama and Trippe, 2017), `iZID` (Wang et al., 2019), `mazeinda` (Albasi, 2018), `mhurdle` (Croissant et al., 2021), `rbtt` (Waudby-Smith and Li, 2017), `ZIBBSeqDiscovery` (Hu and Zhou, 2018), `ZIBseq` (Peng et al., 2017), `zic` (Jochmann, 2017), `ZIM` (Yang et al., 2018), `ziphsmm` (Xu and Liu, 2018), etc. For example, `iZID` covers 12 discrete distributions including Poisson, NB, BB (beta binomial), BNB (beta negative binomial) and their zero-inflated and hurdle versions. It implemented the bootstrapped Monte Carlo p value estimates proposed by Aldirawi et al. (2019) for identifying a discrete distribution. For packages other than `iZID`, please see Wang et al. (2020) for a good review.

The existing literature and packages on analyzing zero-inflated data, including the very recent work Aldirawi et al. (2019) and Wang et al. (2020), are still limited for three reasons. First, only a small number of zero-inflated models were under consideration at the same time. Discrete and continuous baseline distributions were typically considered separately. Secondly, the accuracy of the parameter estimates and the power of the tests still have room for improvement. Thirdly, the confidence intervals of the parameter values were seldom provided along with their estimates, which is critical for model diagnostics, such as, testing whether the inflation or deflation of zeros exists in the data.

In this paper, we introduce our newly developed R package named `AZIAD` for Analyzing Zero-Inflated and Zero-Altered Data, available from the Comprehensive R Archive Network (CRAN, <https://cran.r-project.org/package=AZIAD>). Compared with the existing R packages for similar purposes, our `AZIAD` achieves the following significant improvements: (1) We not only cover discrete baseline distributions including Poisson, geometric, NB, BB, and BNB, but also cover commonly used continuous baseline distributions including normal (or Gaussian), log-normal, half-normal, and exponential distributions along with their zero-inflated and zero-altered (also known as hurdle) models; (2) By more precise specifications on solution forms under different situations and lower/upper bounds needed for numerical optimizations, our R functions provide more accurate maximum likelihood estimates (MLE) even under extreme circumstances, which further gains more power when identifying the most appropriate zero-inflated or hurdle models for a given data set; (3) Following Aldirawi and Yang (2022), we provide not only MLEs for parameters, but also the Fisher information matrix and the corresponding confidence intervals for estimated parameters, which will facilitate the potential users from biological sciences, insurance, health studies, security, ecology, etc, to identify the most appropriate probabilistic model for their dataset and make robust statistical inference based on it.

The rest of this paper is organized as follows. In Section 2, we not only review relevant theoretical results from Aldirawi and Yang (2022), but also provide new formulae

for Fisher information matrices of the zero-inflated (ZI) and zero-altered (ZA or hurdle) models with geometric, BB, BNB, normal, log-normal, half-normal and exponential distributions, as well as ZINB. In Section 3, we summarize and use numerical examples to illustrate the improvements by using our package over existing R packages, as well as identifying the most appropriate models for a real dataset. We interpret and discuss indistinguishable pairs of distributions in Section 4.

2 MLE and Fisher Information for ZI and ZA Models

2.1 MLE and Fisher information for zero-altered or hurdle models

Zero-altered models or hurdle models have been widely used for modeling data with an excess or deficit of zeros (see, for example, Metwally et al. (2018), for a good review). A general hurdle model consists a baseline distribution and a component generating the zeros. The baseline distribution could be fairly general with the distribution function $f_{\theta}(y)$ and its zero-truncated version $f_{\text{tr}}(y | \theta) = [1 - p_0(\theta)]^{-1}f_{\theta}(y)$, $y \neq 0$, where θ is the model parameter(s), and $p_0(\theta) = P_{\theta}(Y = 0)$ is the probability that $Y = 0$ under the baseline distribution. Following Aldirawi and Yang (2022), the distribution function of the corresponding hurdle model can written as:

$$f_{\text{ZA}}(y | \phi, \theta) = \phi \mathbf{1}_{\{y=0\}} + (1 - \phi) f_{\text{tr}}(y | \theta) \mathbf{1}_{\{y \neq 0\}} \quad (1)$$

where $\phi \in [0, 1]$ is the weight parameter of zeros. Actually, $\phi = P(Y = 0)$ if $Y \sim f_{\text{ZA}}$.

If the baseline distribution is discrete with a probability mass function (pmf) $f_{\theta}(y)$, such as Poisson, negative binomial (NB), geometric (Ge), beta binomial (BB) and beta negative binomial (BNB) distributions, then both $f_{\text{tr}}(y | \theta)$ and $f_{\text{ZA}}(y | \phi, \theta)$ are pmfs as well. The corresponding zero-altered or hurdle models may be called as zero-altered Poisson (ZAP) or Poisson hurdle (PH), zero-altered negative binomial (ZANB) or negative binomial hurdle (NBH), zero-altered geometric (ZAGe) or geometric hurdle (GeH), zero-altered beta binomial (ZABB) or beta binomial hurdle (BBH), zero-altered beta negative binomial (ZABNB) or beta negative binomial hurdle (BNBH) models, respectively.

If the baseline distribution is continuous with a probability density function (pdf) $f_{\theta}(y)$, such as Gaussian (or normal), log-normal, half-normal and exponential distributions, then $p_0(\theta) = 0$ and $f_{\text{tr}}(y | \theta) = f_{\theta}(y)$. In this case, $f_{\text{ZA}}(y | \phi, \theta)$ in (1) represents a mixture distribution consisting of a discrete component at 0 with probability ϕ and a continuous component with density function $(1 - \phi)f_{\theta}(y)$. We will revisit these continuous cases in Section 2.3. In this section, we focus on discrete cases.

The parameters of the hurdle model (1) include both ϕ and θ . In this paper, we adopt the maximum likelihood estimate (MLE) for estimating the parameters (see, for example, Section 1.3.1 in Agresti (2013), for justifications on adopting MLE). Let Y_1, \dots, Y_n be a random sample from model (1). Then the likelihood function of (ϕ, θ) is

$$L(\phi, \theta) = \phi^{n-m} (1 - \phi)^m \cdot \prod_{i: Y_i \neq 0} f_{\text{tr}}(Y_i | \theta) \quad (2)$$

where $m = \#\{i : Y_i \neq 0\}$ is the number of nonzero observations. According to Theorem 1 in Aldirawi and Yang (2022), the maximum likelihood estimate (MLE) of $(\phi, \boldsymbol{\theta})$ that maximizes (2) is

$$\hat{\phi} = 1 - \frac{m}{n}, \quad \hat{\boldsymbol{\theta}} = \operatorname{argmax}_{\boldsymbol{\theta}} \prod_{i:Y_i \neq 0} f_{\text{tr}}(Y_i | \boldsymbol{\theta}) \quad (3)$$

That is, $\hat{\boldsymbol{\theta}}$ is simply the MLE for the truncated model with distribution function $f_{\text{tr}}(y | \boldsymbol{\theta}) = f_{\boldsymbol{\theta}}(y)/[1 - p_0(\boldsymbol{\theta})], y \neq 0$.

According to Theorem 3 in Aldirawi and Yang (2022), under some regularity conditions, the Fisher information matrix of the zero-altered distribution is

$$\mathbf{F}_{\text{ZA}} = \begin{bmatrix} \phi^{-1}(1 - \phi)^{-1} & \mathbf{0}^T \\ \mathbf{0} & \mathbf{F}_{\text{ZA}\boldsymbol{\theta}} \end{bmatrix} \quad (4)$$

where

$$\mathbf{F}_{\text{ZA}\boldsymbol{\theta}} = -\frac{1 - \phi}{1 - p_0(\boldsymbol{\theta})} \left(E \left[\frac{\partial^2 \log f_{\boldsymbol{\theta}}(Y')}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} \right] + \frac{p_0(\boldsymbol{\theta})}{1 - p_0(\boldsymbol{\theta})} \cdot \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} \cdot \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}^T} \right)$$

and Y' follows the baseline distribution $f_{\boldsymbol{\theta}}(y)$. Note that the sample size n does not show up in (4) since the Fisher information here is of the distribution, not of the sample.

Major advantages for adopting MLE (3) and calculating the Fisher information matrix (4) include: (i) $\hat{\phi}$ and $\hat{\boldsymbol{\theta}}$ are consistent estimators of ϕ and $\boldsymbol{\theta}$, respectively (Theorem 2 in Aldirawi and Yang (2022)); (ii) $\sqrt{n}(\hat{\phi} - \phi) \xrightarrow{\mathcal{L}} N(0, \phi(1 - \phi))$ and $\sqrt{n}(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}) \xrightarrow{\mathcal{L}} N(\mathbf{0}, \mathbf{F}_{\text{ZA}\boldsymbol{\theta}}^{-1})$, which provides the formulae for building up approximate confidence intervals and relevant hypothesis tests for ϕ and $\boldsymbol{\theta}$ (see, for example, Sections 1.3.3 and 1.3.4 in Agresti (2013)). For example, an approximate $(1 - \alpha)100\%$ confidence interval for ϕ is

$$\phi \in \left(\hat{\phi} - \frac{z_{\frac{\alpha}{2}}}{\sqrt{n}} \sqrt{\hat{\phi}(1 - \hat{\phi})}, \hat{\phi} + \frac{z_{\frac{\alpha}{2}}}{\sqrt{n}} \sqrt{\hat{\phi}(1 - \hat{\phi})} \right) \quad (5)$$

where $z_{\frac{\alpha}{2}} = \Phi^{-1}(1 - \frac{\alpha}{2})$ is the upper $\frac{\alpha}{2}$ th quantile of the standard normal distribution, and $\alpha \in (0, 1)$ is the desired significance level, such as 0.05. If $p_0(\boldsymbol{\theta})$ does not belong to the confidence interval (5), then there is a significant evidence for zero-inflation or deflation.

Therefore, calculating MLE and Fisher information accurately and efficiently is the basis of sound and thorough statistical inferences.

Explicit formulae of the gradients $(\partial \log f_{\boldsymbol{\theta}}(y)/\partial \boldsymbol{\theta})$ and $\partial \log p_0(\boldsymbol{\theta})/\partial \boldsymbol{\theta}$, need for finding MLEs) and the Fisher information matrices of zero-altered Poisson (ZAP) or Poisson hurdle (PH) model, zero-altered negative binomial (ZANB) or negative binomial hurdle (NBH) model, have been provided in Examples 2 and 3 of Aldirawi and Yang (2022), respectively. In this section, we will provide explicit formulae for zero-altered geometric, beta binomial, and beta negative binomial models.

Example 2.1. Zero-altered geometric (ZAGe) or geometric hurdle (GeH) model has been used, for example, in Mullahy (1986) for modeling count data in econometrics. The pmf of the baseline distribution can be written as $f_p(y) = p(1 - p)^y$ with

$y \in \{0, 1, 2, \dots\}$ and parameter $p \in (0, 1)$. In this case, $p_0(p) = p$, $\frac{\partial \log f_p(y)}{\partial p} = \frac{1}{p} - \frac{y}{1-p}$, and $\frac{\partial \log p_0(p)}{\partial p} = \frac{1}{p}$.

According to Theorem 3 in Aldirawi and Yang (2022), the Fisher information matrix of the ZAGe or GeH distribution can be written as

$$\mathbf{F}_{\text{ZAGe}} = \begin{bmatrix} \frac{1}{\phi(1-\phi)} & 0 \\ 0 & \frac{1-\phi}{p^2(1-p)} \end{bmatrix}$$

□

Example 2.2. Zero-altered beta-binomial (ZABB) or beta-binomial hurdle (BBH) model has been used by Aldirawi et al. (2019) for modeling microbiome data. The pmf of the baseline distribution with parameters $\boldsymbol{\theta} = (n, \alpha, \beta) \in \mathbb{N} \times (0, \infty) \times (0, \infty)$ is given by $f_{\boldsymbol{\theta}}(y) = \binom{n}{y} \frac{\text{Beta}(y+\alpha, n-y+\beta)}{\text{Beta}(\alpha, \beta)}$, $y \in \{0, 1, 2, \dots, n\}$. In this case, $p_0(\boldsymbol{\theta}) = \frac{\Gamma(n+\beta)\Gamma(\alpha+\beta)}{\Gamma(n+\alpha+\beta)\Gamma(\beta)}$. Then

$$\begin{aligned} \frac{\partial \log f_{\boldsymbol{\theta}}(y)}{\partial n} &= \Psi(n+1) - \Psi(n-y+1) + \Psi(n-y+\beta) - \Psi(n+\alpha+\beta) \\ \frac{\partial \log f_{\boldsymbol{\theta}}(y)}{\partial \alpha} &= \Psi(y+\alpha) - \Psi(n+\alpha+\beta) + \Psi(\alpha+\beta) - \Psi(\alpha) \\ \frac{\partial \log f_{\boldsymbol{\theta}}(y)}{\partial \beta} &= \Psi(n-y+\beta) - \Psi(n+\alpha+\beta) + \Psi(\alpha+\beta) - \Psi(\beta) \\ \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial n} &= \Psi(n+\beta) - \Psi(n+\alpha+\beta) \\ \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \alpha} &= \Psi(\alpha+\beta) - \Psi(n+\alpha+\beta) \\ \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \beta} &= \Psi(n+\beta) + \Psi(\alpha+\beta) - \Psi(n+\alpha+\beta) - \Psi(\beta) \end{aligned}$$

where $\Psi(\cdot) = \Gamma'(\cdot)/\Gamma(\cdot)$ is known as the *digamma* function. Note that the range of parameter n can be extended to positive real numbers. According to Theorem 3 in Aldirawi and Yang (2022), the Fisher information matrix of the ZABB or BBH distribution is

$$\mathbf{F}_{\text{ZABB}} = \begin{bmatrix} \frac{1}{\phi(1-\phi)} & \mathbf{0}^T \\ \mathbf{0} & \mathbf{F}_{\text{BB}\boldsymbol{\theta}} \end{bmatrix}$$

where

$$\begin{aligned} \mathbf{F}_{\text{BB}\boldsymbol{\theta}} &= -\frac{(1-\phi)\Gamma(n+\alpha+\beta)\Gamma(\beta)}{\Gamma(n+\alpha+\beta)\Gamma(\beta) - \Gamma(n+\beta)\Gamma(\alpha+\beta)} \left(\begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{12} & A_{22} & A_{23} \\ A_{13} & A_{23} & A_{33} \end{bmatrix} \right. \\ &\quad \left. + \frac{\Gamma(n+\beta)\Gamma(\alpha+\beta)}{\Gamma(n+\alpha+\beta)\Gamma(\beta) - \Gamma(n+\beta)\Gamma(\alpha+\beta)} \begin{bmatrix} B_{11} & B_{12} & B_{13} \\ B_{12} & B_{22} & B_{23} \\ B_{13} & B_{23} & B_{33} \end{bmatrix} \right) \end{aligned}$$

with

$$\begin{aligned}
A_{11} &= \Psi_1(n+1) - \Psi_1(n+\alpha+\beta) + E\Psi_1(n-Y'+\beta) - E\Psi_1(n-Y'+1) \\
A_{12} &= -\Psi_1(n+\alpha+\beta) \\
A_{13} &= E\Psi_1(n-Y'+\beta) - \Psi_1(n+\alpha+\beta) \\
A_{22} &= \Psi_1(\alpha+\beta) - \Psi_1(n+\alpha+\beta) - \Psi_1(\alpha) + E\Psi_1(Y'+\alpha) \\
A_{23} &= \Psi_1(\alpha+\beta) - \Psi_1(n+\alpha+\beta) \\
A_{33} &= \Psi_1(\alpha+\beta) - \Psi_1(\beta) - \Psi_1(n+\alpha+\beta) + E\Psi_1(n-Y'+\beta) \\
B_{11} &= [\Psi(n+\beta) - \Psi(n+\alpha+\beta)]^2 \\
B_{12} &= [\Psi(n+\beta) - \Psi(n+\alpha+\beta)] \cdot [\Psi(\alpha+\beta) - \Psi(n+\alpha+\beta)] \\
B_{13} &= [\Psi(n+\beta) - \Psi(n+\alpha+\beta)] \cdot [\Psi(n+\beta) + \Psi(\alpha+\beta) - \Psi(n+\alpha+\beta) - \Psi(\beta)] \\
B_{22} &= [\Psi(\alpha+\beta) - \Psi(n+\alpha+\beta)]^2 \\
B_{23} &= [\Psi(\alpha+\beta) - \Psi(n+\alpha+\beta)] \cdot [\Psi(n+\beta) + \Psi(\alpha+\beta) - \Psi(n+\alpha+\beta) - \Psi(\beta)] \\
B_{33} &= [\Psi(n+\beta) + \Psi(\alpha+\beta) - \Psi(n+\alpha+\beta) - \Psi(\beta)]^2
\end{aligned}$$

where $\Psi_1(\cdot) = \Psi'(\cdot)$ is known as the *trigamma* function, and Y' follows the baseline distribution $f_{\boldsymbol{\theta}}(y)$. \square

Due to the lack of simple forms, we may use Monte Carlo estimates for calculating $E\Psi_1(\cdot)$ in Example 2.2 and others approximately. For example, $E\Psi_1(n-Y'+\beta)$ can be approximated by $N^{-1} \sum_{i=1}^N \Psi_1(n-Y'_i+\beta)$ with simulated Y'_1, \dots, Y'_N from $f_{\boldsymbol{\theta}}(y)$.

Another computational issue involved in Example 2.2 and others is that $\log[\Gamma(n+\alpha+\beta)\Gamma(\beta) - \Gamma(n+\beta)\Gamma(\alpha+\beta)]$, which is relevant to $\log p_0(\boldsymbol{\theta})$, may be undefined numerically for large n since both $\Gamma(n+\alpha+\beta)$ and $\Gamma(n+\beta)$ are numerical infinity. To overcome this kind of issues, we use the fact $\log(A-B) = \log(1 - \exp(\log B - \log A)) + \log A$ if $A \geq B$. It can be verified that $\Gamma(n+\alpha+\beta)\Gamma(\beta) > \Gamma(n+\beta)\Gamma(\alpha+\beta)$.

Example 2.3. Zero-altered beta negative binomial (ZABNB) or beta negative binomial hurdle (BNBH) model has been recommended by Aldirawi et al. (2019) and Aldirawi and Yang (2022) for modeling microbiome data. The pmf of the baseline distribution with parameters $\boldsymbol{\theta} = (r, \alpha, \beta) \in \mathbb{N} \times (0, \infty) \times (0, \infty)$ is given by $f_{\boldsymbol{\theta}}(y) = \binom{r+y-1}{y} \frac{\text{Beta}(r+\alpha, y+\beta)}{\text{Beta}(\alpha, \beta)}$, $y \in \{0, 1, 2, \dots\}$. Then $p_0(\boldsymbol{\theta}) = \frac{\Gamma(r+\alpha)\Gamma(\alpha+\beta)}{\Gamma(r+\alpha+\beta)\Gamma(\alpha)}$. Note that r can also be extended to positive real numbers.

$$\begin{aligned}
\frac{\partial \log f_{\boldsymbol{\theta}}(y)}{\partial r} &= \Psi(r+y) - \Psi(r) + \Psi(r+\alpha) - \Psi(r+y+\alpha+\beta) \\
\frac{\partial \log f_{\boldsymbol{\theta}}(y)}{\partial \alpha} &= \Psi(r+\alpha) - \Psi(r+y+\alpha+\beta) + \Psi(\alpha+\beta) - \Psi(\alpha) \\
\frac{\partial \log f_{\boldsymbol{\theta}}(y)}{\partial \beta} &= \Psi(y+\beta) - \Psi(r+y+\alpha+\beta) + \Psi(\alpha+\beta) - \Psi(\beta) \\
\frac{\partial \log p_0(\boldsymbol{\theta})}{\partial r} &= \Psi(r+\alpha) - \Psi(r+\alpha+\beta) \\
\frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \alpha} &= \Psi(r+\alpha) + \Psi(\alpha+\beta) - \Psi(r+\alpha+\beta) - \Psi(\alpha) \\
\frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \beta} &= \Psi(\alpha+\beta) - \Psi(r+\alpha+\beta)
\end{aligned}$$

According to Theorem 3 in Aldirawi and Yang (2022), the Fisher information matrix of the ZABNB or BNBH distribution is

$$\mathbf{F}_{\text{ZABNB}} = \begin{bmatrix} \frac{1}{\phi(1-\phi)} & \mathbf{0}^T \\ \mathbf{0} & \mathbf{F}_{\text{BNBH}\boldsymbol{\theta}} \end{bmatrix}$$

where

$$\mathbf{F}_{\text{BNBH}\boldsymbol{\theta}} = -\frac{(1-\phi)\Gamma(r+\alpha+\beta)\Gamma(\alpha)}{\Gamma(r+\alpha+\beta)\Gamma(\alpha) - \Gamma(r+\alpha)\Gamma(\alpha+\beta)} \begin{pmatrix} \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{12} & A_{22} & A_{23} \\ A_{13} & A_{23} & A_{33} \end{bmatrix} \\ + \frac{\Gamma(r+\alpha)\Gamma(\alpha+\beta)}{\Gamma(r+\alpha+\beta)\Gamma(\alpha) - \Gamma(r+\alpha)\Gamma(\alpha+\beta)} \begin{bmatrix} B_{11} & B_{12} & B_{13} \\ B_{12} & B_{22} & B_{23} \\ B_{13} & B_{23} & B_{33} \end{bmatrix} \end{pmatrix}$$

with

$$\begin{aligned} A_{11} &= E\Psi_1(r+Y') - \Psi_1(r) + \Psi_1(r+\alpha) - E\Psi_1(r+Y'+\alpha+\beta) \\ A_{12} &= \Psi_1(r+\alpha) - E\Psi_1(r+Y'+\alpha+\beta) \\ A_{13} &= -E\Psi_1(r+Y'+\alpha+\beta) \\ A_{22} &= \Psi_1(r+\alpha) - E\Psi_1(r+Y'+\alpha+\beta) + \Psi_1(\alpha+\beta) - \Psi_1(\alpha) \\ A_{23} &= -E\Psi_1(r+Y'+\alpha+\beta) + \Psi_1(\alpha+\beta) \\ A_{33} &= E\Psi_1(Y'+\beta) - E\Psi_1(r+Y'+\alpha+\beta) + \Psi_1(\alpha+\beta) - \Psi_1(\beta) \\ B_{11} &= [\Psi(r+\alpha) - \Psi(r+\alpha+\beta)]^2 \\ B_{12} &= [\Psi(r+\alpha) - \Psi(r+\alpha+\beta)] \cdot [\Psi(r+\alpha) + \Psi(\alpha+\beta) - \Psi(r+\alpha+\beta) - \Psi(\alpha)] \\ B_{13} &= [\Psi(r+\alpha) - \Psi(r+\alpha+\beta)] \cdot [\Psi(r+\alpha) - \Psi(r+\alpha+\beta)] \\ B_{22} &= [\Psi(r+\alpha) + \Psi(\alpha+\beta) - \Psi(r+\alpha+\beta) - \Psi(\alpha)]^2 \\ B_{23} &= [\Psi(r+\alpha) + \Psi(\alpha+\beta) - \Psi(r+\alpha+\beta) - \Psi(\alpha)] \cdot [\Psi(r+\alpha) - \Psi(r+\alpha+\beta)] \\ B_{33} &= [\Psi(r+\alpha) - \Psi(r+\alpha+\beta)]^2 \end{aligned}$$

As for Ψ_1 and relevant calculations, please see the arguments right after Example 2.2.

□

2.2 MLE and Fisher information for zero-inflated models

When data is sparse, a zero-inflated (ZI) model is more commonly used in practice, which assumes an excess of zeros (see, for example, Metwally et al. (2018) for a good review). Similar as the zero-altered (ZA) models in Section 2.1, there is a baseline distribution with distribution function $f_{\boldsymbol{\theta}}(y)$ and parameter(s) $\boldsymbol{\theta}$. We also denote $p_0(\boldsymbol{\theta}) = P_{\boldsymbol{\theta}}(Y = 0)$, if $Y \sim f_{\boldsymbol{\theta}}$. Different from ZA models, a zero-weighting parameter $\phi \in [0, 1]$ adds additional probability of zeros to the ZI model. Following Aldirawi and Yang (2022), we write the distribution function of the corresponding ZI model as

$$f_{\text{ZI}}(y | \phi, \boldsymbol{\theta}) = [\phi + (1-\phi)p_0(\boldsymbol{\theta})]\mathbf{1}_{\{y=0\}} + (1-\phi)f_{\boldsymbol{\theta}}(y)\mathbf{1}_{\{y \neq 0\}} \quad (6)$$

Note that if $Y \sim f_{\text{ZI}}$, then $P(Y = 0) = \phi + (1 - \phi)p_0(\boldsymbol{\theta})$, which is larger than ϕ in general.

When the baseline distribution is either continuous with a pdf $f_{\boldsymbol{\theta}}(y)$ or discrete but with $p_0(\boldsymbol{\theta}) = 0$, the corresponding zero-inflated model (6) is essentially the same as the corresponding zero-altered model (1). Examples include Gaussian or normal, log-normal, half-normal, and exponential distributions. We will revisit them in Section 2.3.

When the baseline distribution is discrete, such as Poisson (P), negative binomial (NB), geometric (Ge), beta binomial (BB) and beta negative binomial (BNB), the corresponding zero-inflated models can be written as ZIP, ZINB, ZIGe, ZIBB, and ZIBNB, respectively.

Given a random sample Y_1, \dots, Y_n from the zero-inflated model $f_{\text{ZI}}(y|\phi, \boldsymbol{\theta})$, the likelihood function of $(\phi, \boldsymbol{\theta})$ can be written as

$$\begin{aligned} L(\phi, \boldsymbol{\theta}) &= [\phi + p_0(\boldsymbol{\theta})(1 - \phi)]^{n-m} \cdot (1 - \phi)^m \prod_{i:Y_i \neq 0} f_{\boldsymbol{\theta}}(Y_i) \\ &= [\phi + p_0(\boldsymbol{\theta})(1 - \phi)]^{n-m} \cdot (1 - \phi)^m (1 - p_0(\boldsymbol{\theta}))^m \cdot \prod_{i:Y_i \neq 0} f_{\text{tr}}(Y_i | \boldsymbol{\theta}) \end{aligned} \quad (7)$$

where $m = \#\{i : Y_i \neq 0\}$, and $f_{\text{tr}}(y; \boldsymbol{\theta}) = f_{\boldsymbol{\theta}}(y)/[1 - p_0(\boldsymbol{\theta})]$, $y \neq 0$.

According to Theorem 4 in Aldirawi and Yang (2022), the maximum likelihood estimate $(\hat{\phi}, \hat{\boldsymbol{\theta}})$ maximizing (7) can be obtained as follows:

- (0) Determine $\boldsymbol{\theta}_* = \operatorname{argmax}_{\boldsymbol{\theta}} L_{\text{tr}}(\boldsymbol{\theta})$, where $L_{\text{tr}}(\boldsymbol{\theta}) = \prod_{i:Y_i \neq 0} f_{\text{tr}}(Y_i; \boldsymbol{\theta})$.
- (1) If $m/n \leq 1 - p_0(\boldsymbol{\theta}_*)$, then $\hat{\boldsymbol{\theta}} = \boldsymbol{\theta}_*$ and $\hat{\phi} = 1 - [1 - p_0(\boldsymbol{\theta}_*)]^{-1} \cdot m/n$.
- (2) Otherwise, $\hat{\boldsymbol{\theta}} = \operatorname{argmax}_{\boldsymbol{\theta}} L(\psi(\boldsymbol{\theta}), \boldsymbol{\theta})$ and $\hat{\phi} = 1 - \psi(\hat{\boldsymbol{\theta}}) \cdot [1 - p_0(\hat{\boldsymbol{\theta}})]^{-1}$, where $\psi(\boldsymbol{\theta}) = \min\{m/n, 1 - p_0(\boldsymbol{\theta})\}$, and $L(\psi, \boldsymbol{\theta}) = (1 - \psi)^{n-m} \psi^m \prod_{i:Y_i \neq 0} f_{\text{tr}}(Y_i; \boldsymbol{\theta})$.

Solving for the MLE $(\hat{\phi}, \hat{\boldsymbol{\theta}})$ may involve two maximization problems, $\boldsymbol{\theta}_* = \operatorname{argmax}_{\boldsymbol{\theta}} L_{\text{tr}}(\boldsymbol{\theta})$ and $\hat{\boldsymbol{\theta}} = \operatorname{argmax}_{\boldsymbol{\theta}} L(\psi(\boldsymbol{\theta}), \boldsymbol{\theta})$. The following first order derivatives may be needed by, for example, quasi-Newton algorithms:

$$\begin{aligned} \frac{\partial \log L_{\text{tr}}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} &= \sum_{i:Y_i \neq 0} \frac{\partial \log f_{\boldsymbol{\theta}}(Y_i)}{\partial \boldsymbol{\theta}} - m \frac{\partial \log[1 - p_0(\boldsymbol{\theta})]}{\partial \boldsymbol{\theta}} \\ \frac{\partial \log L(\psi(\boldsymbol{\theta}), \boldsymbol{\theta})}{\partial \boldsymbol{\theta}} &= \begin{cases} \sum_{i:Y_i \neq 0} \frac{\partial \log f_{\boldsymbol{\theta}}(Y_i)}{\partial \boldsymbol{\theta}} - m \frac{\partial \log[1 - p_0(\boldsymbol{\theta})]}{\partial \boldsymbol{\theta}} & \text{if } 1 - p_0(\boldsymbol{\theta}) > \frac{m}{n} \\ \sum_{i:Y_i \neq 0} \frac{\partial \log f_{\boldsymbol{\theta}}(Y_i)}{\partial \boldsymbol{\theta}} + (n - m) \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} & \text{if } 1 - p_0(\boldsymbol{\theta}) < \frac{m}{n} \end{cases} \end{aligned}$$

Note that

$$\frac{\partial \log[1 - p_0(\boldsymbol{\theta})]}{\partial \boldsymbol{\theta}} = -\frac{p_0(\boldsymbol{\theta})}{1 - p_0(\boldsymbol{\theta})} \cdot \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}}$$

Thus for different models, we only need to prepare specific formulae of $\partial \log f_{\boldsymbol{\theta}}(y)/\partial \boldsymbol{\theta}$ and $\partial \log p_0(\boldsymbol{\theta})/\partial \boldsymbol{\theta}$ for numerical calculations.

According to Theorem 5 in Aldirawi and Yang (2022), under some regularity conditions, the Fisher information matrix of a general zero-inflated distribution is

$$\mathbf{F}_{\text{ZI}} = \begin{bmatrix} \frac{1 - p_0(\boldsymbol{\theta})}{[\phi + (1 - \phi)p_0(\boldsymbol{\theta})](1 - \phi)} & \frac{p_0(\boldsymbol{\theta})}{\phi + (1 - \phi)p_0(\boldsymbol{\theta})} \cdot \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}^T} \\ \frac{p_0(\boldsymbol{\theta})}{\phi + (1 - \phi)p_0(\boldsymbol{\theta})} \cdot \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} & \mathbf{F}_{\text{ZI}\boldsymbol{\theta}} \end{bmatrix} \quad (8)$$

where

$$\mathbf{F}_{\text{ZI}\boldsymbol{\theta}} = -(1 - \phi) \left(E \left[\frac{\partial^2 \log f_{\boldsymbol{\theta}}(Y')}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} \right] + \frac{\phi p_0(\boldsymbol{\theta})}{\phi + (1 - \phi)p_0(\boldsymbol{\theta})} \cdot \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} \cdot \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}^T} \right)$$

and Y' follows the baseline distribution $f_{\boldsymbol{\theta}}(y)$. We denote $\mathbf{F}_{\boldsymbol{\theta}} = -E \left[\frac{\partial^2 \log f_{\boldsymbol{\theta}}(Y')}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} \right]$, which is essentially the Fisher information matrix of the baseline distribution.

The Fisher information matrix (8) is more complicated than (4) for zero-altered models. To obtain the asymptotic distributions of MLEs, it is more convenient to obtain the following theorem by applying the formulae (see, for example, Formulae 4.33 and 14.13(b) in Seber (2008)) of block matrices and the Sherman-Morrison formula (see, for example, Section 2.1.4 in Golub and Loan (2013)).

Theorem 2.1. *Suppose $\phi \in (0, 1)$, $|\mathbf{F}_{\boldsymbol{\theta}}| \neq 0$ and $|\mathbf{F}_{\text{ZI}}| \neq 0$. Then*

$$\mathbf{F}_{\text{ZI}}^{-1} = \begin{bmatrix} \phi(1 - \phi) \cdot \frac{d_{\boldsymbol{\theta}} \delta_{\boldsymbol{\theta}}}{d_{\boldsymbol{\theta}} \delta_{\boldsymbol{\theta}} - p_0(\boldsymbol{\theta})} & -\frac{\phi p_0(\boldsymbol{\theta})}{d_{\boldsymbol{\theta}} \delta_{\boldsymbol{\theta}} - p_0(\boldsymbol{\theta})} \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}^T} \mathbf{F}_{\boldsymbol{\theta}}^{-1} \\ -\frac{\phi p_0(\boldsymbol{\theta})}{d_{\boldsymbol{\theta}} \delta_{\boldsymbol{\theta}} - p_0(\boldsymbol{\theta})} \mathbf{F}_{\boldsymbol{\theta}}^{-1} \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} & \mathbf{D}_{\boldsymbol{\theta}} \end{bmatrix}$$

where $d_{\boldsymbol{\theta}} = \phi + (1 - \phi)p_0(\boldsymbol{\theta}) > 0$, $\delta_{\boldsymbol{\theta}} = 1 - \frac{\phi p_0(\boldsymbol{\theta})}{d_{\boldsymbol{\theta}}} \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}^T} \mathbf{F}_{\boldsymbol{\theta}}^{-1} \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} \neq 0$, and

$$\mathbf{D}_{\boldsymbol{\theta}} = \frac{1}{1 - \phi} \left[\mathbf{F}_{\boldsymbol{\theta}}^{-1} + \frac{\phi p_0(\boldsymbol{\theta})}{d_{\boldsymbol{\theta}} \delta_{\boldsymbol{\theta}} - p_0(\boldsymbol{\theta})} \mathbf{F}_{\boldsymbol{\theta}}^{-1} \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}^T} \mathbf{F}_{\boldsymbol{\theta}}^{-1} \right]$$

□

The proof of Theorem 2.1 is relegated to the Appendix. With Theorem 2.1, we have $\sqrt{n}(\hat{\phi} - \phi) \xrightarrow{\mathcal{L}} N\left(0, \phi(1 - \phi) \frac{d_{\boldsymbol{\theta}} \delta_{\boldsymbol{\theta}}}{d_{\boldsymbol{\theta}} \delta_{\boldsymbol{\theta}} - p_0(\boldsymbol{\theta})}\right)$ and $\sqrt{n}(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}) \xrightarrow{\mathcal{L}} N(\mathbf{0}, \mathbf{D}_{\boldsymbol{\theta}})$. The relevant confidence intervals and hypothesis tests can be performed similarly as for zero-altered models.

As mentioned in Section 2.1, we need to calculating MLE and Fisher information accurately and efficiently. Explicit formulae of the Fisher information matrix of zero-inflated Poisson (ZIP) has been provided in Example 5 of Aldirawi and Yang (2022). In this section, we will provide explicit formulae of the gradients and Fisher information matrices for zero-inflated geometric, negative binomial, beta binomial, and beta negative binomial models.

Example 2.4. Zero-inflated geometric model (ZIGe) Same as in Example 2.1, the pmf of the baseline distribution can be written as $f_p(y) = p(1 - p)^y$ with $y \in \{0, 1, 2, \dots\}$ and parameter $p \in (0, 1)$. According to Theorem 5 in Aldirawi and Yang (2022), the Fisher information matrix of the corresponding ZIGe distribution is

$$\mathbf{F}_{\text{ZIGe}} = \frac{1}{\phi + (1 - \phi)p} \begin{bmatrix} \frac{1-p}{1-\phi} & 1 \\ 1 & \frac{(1-\phi)[\phi(1-p)+p(1-\phi)+\phi p^2]}{p^2(1-p)} \end{bmatrix}$$

□

Example 2.5. Zero-inflated negative binomial model (ZINB) Different from Example 3 of Aldirawi and Yang (2022), we take the form of the pmf of the baseline NB distribution as $f_{\theta}(y) = \frac{\Gamma(y+r)}{\Gamma(y+1)\Gamma(r)}p^r(1-p)^y$ with parameters $\theta = (r, p) \in (0, \infty) \times [0, 1]$, $y \in \{0, 1, 2, \dots\}$, which is more popular in the statistical literature (see, for example, Hogg et al. (2019)). In short, the p in Example 3 of Aldirawi and Yang (2022) is replaced by $1-p$ here. Then $p_0(\theta) = p^r$ and

$$\begin{aligned}\frac{\partial \log f_{\theta}(y)}{\partial r} &= \Psi(y+r) - \Psi(r) + \log p \\ \frac{\partial \log f_{\theta}(y)}{\partial p} &= \frac{r}{p} - \frac{y}{1-p} \\ \frac{\partial \log p_0(\theta)}{\partial r} &= \log p \\ \frac{\partial \log p_0(\theta)}{\partial p} &= \frac{r}{p}\end{aligned}$$

where $\Psi(\cdot)$ is the digamma function. According to Theorem 5 in Aldirawi and Yang (2022), the Fisher information matrix of the ZINB distribution is

$$\mathbf{F}_{\text{ZINB}} = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{12} & & \mathbf{F}_{\text{ZINB}\theta} \\ A_{13} & & \end{bmatrix}$$

where $A_{11} = \frac{1-p^r}{[\phi+(1-\phi)p^r](1-\phi)}$, $A_{12} = \frac{p^r \log p}{\phi+(1-\phi)p^r}$, $A_{13} = \frac{rp^{r-1}}{\phi+(1-\phi)p^r}$, and

$$\mathbf{F}_{\text{ZINB}\theta} = -(1-\phi) \left(\begin{bmatrix} B_{11} & B_{12} \\ B_{12} & B_{22} \end{bmatrix} + \frac{\phi p^r}{\phi+(1-\phi)p^r} \begin{bmatrix} C_{11} & C_{12} \\ C_{12} & C_{22} \end{bmatrix} \right)$$

with $B_{11} = E\Psi_1(Y'+r) - \Psi_1(r)$, $B_{12} = \frac{1}{p}$, $B_{22} = -\frac{r}{p^2(1-p)}$, $C_{11} = (\log p)^2$, $C_{12} = \frac{r \log p}{p}$, and $C_{22} = \frac{r^2}{p^2}$. Here $\Psi_1(\cdot)$ is the trigamma function (see Example 2.2). \square

Example 2.6. Zero-inflated beta-binomial model (ZIBB) Same as in Example 2.2, the pmf of the baseline distribution is $f_{\theta}(y) = \binom{n}{y} \frac{\text{Beta}(y+\alpha, n-y+\beta)}{\text{Beta}(\alpha, \beta)}$ with parameters $\theta = (n, \alpha, \beta)$ and $y = 0, 1, \dots, n$. According to Theorem 5 in Aldirawi and Yang (2022), the Fisher information matrix of the ZIBB distribution is

$$\mathbf{F}_{\text{ZIBB}} = \begin{bmatrix} C_{11} & C_{12} & C_{13} & C_{14} \\ C_{12} & & & \\ C_{13} & & \mathbf{F}_{\text{ZIBB}\theta} & \\ C_{14} & & & \end{bmatrix}$$

where

$$\begin{aligned}
C_{11} &= \frac{\Gamma(n + \alpha + \beta)\Gamma(\beta) - \Gamma(n + \beta)\Gamma(\alpha + \beta)}{[\phi\Gamma(n + \alpha + \beta)\Gamma(\beta) + (1 - \phi)\Gamma(n + \beta)\Gamma(\alpha + \beta)](1 - \phi)} \\
C_{12} &= \frac{\Gamma(n + \beta)\Gamma(\alpha + \beta)[\Psi(n + \beta) - \Psi(n + \alpha + \beta)]}{\phi\Gamma(n + \alpha + \beta)\Gamma(\beta) + (1 - \phi)\Gamma(n + \beta)\Gamma(\alpha + \beta)} \\
C_{13} &= \frac{\Gamma(n + \beta)\Gamma(\alpha + \beta)[\Psi(\alpha + \beta) - \Psi(n + \alpha + \beta)]}{\phi\Gamma(n + \alpha + \beta)\Gamma(\beta) + (1 - \phi)\Gamma(n + \beta)\Gamma(\alpha + \beta)} \\
C_{14} &= \frac{\Gamma(n + \beta)\Gamma(\alpha + \beta)[\Psi(n + \beta) + \Psi(\alpha + \beta) - \Psi(n + \alpha + \beta) - \Psi(\beta)]}{\phi\Gamma(n + \alpha + \beta)\Gamma(\beta) + (1 - \phi)\Gamma(n + \beta)\Gamma(\alpha + \beta)} \\
\mathbf{F}_{\text{ZIBB}\boldsymbol{\theta}} &= -(1 - \phi) \left(\mathbf{A} + \frac{\phi\Gamma(n + \beta)\Gamma(\alpha + \beta)}{\phi\Gamma(n + \alpha + \beta)\Gamma(\beta) + (1 - \phi)\Gamma(n + \beta)\Gamma(\alpha + \beta)} \cdot \mathbf{B} \right)
\end{aligned}$$

Note that

$$\mathbf{A} = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{12} & A_{22} & A_{23} \\ A_{13} & A_{23} & A_{33} \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} B_{11} & B_{12} & B_{13} \\ B_{21} & B_{22} & B_{23} \\ B_{31} & B_{32} & B_{33} \end{bmatrix}$$

here are the same as in Example 2.2. \square

Example 2.7. Zero-inflated beta negative binomial model (ZIBNB) Same as in Example 2.3, the pmf of the baseline distribution is given by $f_{\boldsymbol{\theta}}(y) = \binom{r+y-1}{y} \frac{\text{Beta}(r+\alpha, y+\beta)}{\text{Beta}(\alpha, \beta)}$ with parameters $\boldsymbol{\theta} = (r, \alpha, \beta)$, $y \in \{0, 1, 2, \dots\}$. According to Theorem 5 in Aldirawi and Yang (2022), the Fisher information matrix of the ZIBNB distribution is

$$\mathbf{F}_{\text{ZIBNB}} = \begin{bmatrix} C_{11} & C_{12} & C_{13} & C_{14} \\ C_{12} & & & \\ C_{13} & & \mathbf{F}_{\text{ZIBNB}\boldsymbol{\theta}} & \\ C_{14} & & & \end{bmatrix}$$

where

$$\begin{aligned}
C_{11} &= \frac{\Gamma(r + \alpha + \beta)\Gamma(\alpha) - \Gamma(r + \alpha)\Gamma(\alpha + \beta)}{[\phi\Gamma(r + \alpha + \beta)\Gamma(\alpha) + (1 - \phi)\Gamma(r + \alpha)\Gamma(\alpha + \beta)](1 - \phi)} \\
C_{12} &= \frac{\Gamma(r + \alpha)\Gamma(\alpha + \beta)[\Psi(r + \alpha) - \Psi(r + \alpha + \beta)]}{\phi\Gamma(r + \alpha + \beta)\Gamma(\alpha) + (1 - \phi)\Gamma(r + \alpha)\Gamma(\alpha + \beta)} \\
C_{13} &= \frac{\Gamma(r + \alpha)\Gamma(\alpha + \beta)[\Psi(r + \alpha) + \Psi(\alpha + \beta) - \Psi(r + \alpha + \beta) - \Psi(\alpha)]}{\phi\Gamma(r + \alpha + \beta)\Gamma(\alpha) + (1 - \phi)\Gamma(r + \alpha)\Gamma(\alpha + \beta)} \\
C_{14} &= \frac{\Gamma(r + \alpha)\Gamma(\alpha + \beta)[\Psi(\alpha + \beta) - \Psi(r + \alpha + \beta)]}{\phi\Gamma(r + \alpha + \beta)\Gamma(\alpha) + (1 - \phi)\Gamma(r + \alpha)\Gamma(\alpha + \beta)} \\
\mathbf{F}_{\text{ZIBNB}\boldsymbol{\theta}} &= -(1 - \phi) \left(\mathbf{A} + \frac{\phi\Gamma(r + \alpha)\Gamma(\alpha + \beta)}{\phi\Gamma(r + \alpha + \beta)\Gamma(\alpha) + (1 - \phi)\Gamma(r + \alpha)\Gamma(\alpha + \beta)} \cdot \mathbf{B} \right)
\end{aligned}$$

Note that

$$\mathbf{A} = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{12} & A_{22} & A_{23} \\ A_{13} & A_{23} & A_{33} \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} B_{11} & B_{12} & B_{13} \\ B_{12} & B_{22} & B_{23} \\ B_{13} & B_{23} & B_{33} \end{bmatrix}$$

are the same as in Example 2.3. \square

2.3 Zero-altered and zero-inflated models with continuous baseline distributions

As mentioned in Section 2.2, if the baseline distribution $f_{\boldsymbol{\theta}}(y)$ satisfies $p_0(\boldsymbol{\theta}) = P_{\boldsymbol{\theta}}(Y = 0) = 0$ given $Y \sim f_{\boldsymbol{\theta}}(y)$, then the zero-altered model (1) and the zero-inflated model (6) are the same. We call such kind of models the *zero-altered-zero-inflated* (ZAZI) models. Examples include all continuous baseline distributions, as well as discrete or mixture baseline distributions satisfying $p_0(\boldsymbol{\theta}) = 0$. For ZAZI models with a baseline distribution $f_{\boldsymbol{\theta}}(y)$, its distribution function can be written as

$$f_{\text{ZAZI}}(y \mid \phi, \boldsymbol{\theta}) = \phi \mathbf{1}_{\{y=0\}} + (1 - \phi) f_{\boldsymbol{\theta}}(y) \mathbf{1}_{\{y \neq 0\}} \quad (9)$$

Given a random sample $Y_1, \dots, Y_n \sim f_{\text{ZAZI}}(y \mid \phi, \boldsymbol{\theta})$, the likelihood function of $(\phi, \boldsymbol{\theta})$ is

$$L(\phi, \boldsymbol{\theta}) = \phi^{n-m} (1 - \phi)^m \cdot \prod_{i: Y_i \neq 0} f_{\boldsymbol{\theta}}(Y_i) \quad (10)$$

where $m = \#\{i : Y_i \neq 0\}$. Then the MLEs maximizing (10) are $\hat{\phi} = 1 - m/n$ and $\hat{\boldsymbol{\theta}} = \operatorname{argmax}_{\boldsymbol{\theta}} \prod_{i: Y_i \neq 0} f_{\boldsymbol{\theta}}(Y_i)$, which are similar as (3) for zero-altered models. As a special case of Theorem 3 in Aldirawi and Yang (2022), we have the following formulae for the Fisher information matrix of ZAZI distributions.

Theorem 2.2. Under regularity conditions, the Fisher information matrix of the ZAZI distribution (9) is

$$\mathbf{F}_{\text{ZAZI}} = \begin{bmatrix} \phi^{-1}(1 - \phi)^{-1} & \mathbf{0}^T \\ \mathbf{0} & \mathbf{F}_{\text{ZAZI}\boldsymbol{\theta}} \end{bmatrix}$$

where

$$\mathbf{F}_{\text{ZAZI}\boldsymbol{\theta}} = -(1 - \phi) \cdot E \left[\frac{\partial^2 \log f_{\boldsymbol{\theta}}(Y')}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} \right]$$

and Y' follows the baseline distribution $f_{\boldsymbol{\theta}}(y)$. \square

In this section, we provide explicit formulae of gradients and Fisher information matrices for commonly used ZAZI models with continuous baseline distributions including Gaussian or normal, log-normal, half-normal, and exponential distributions.

Example 2.8. Zero-altered-zero-inflated Gaussian model (ZAZIG) This model has been used by, for example, Zhang et al. (2020) for analyzing longitudinal microbiome data, known as a zero-inflated Gaussian (ZIG) model. The pdf of the baseline distribution with parameters $\boldsymbol{\theta} = (\mu, \sigma) \in \mathbb{R} \times (0, \infty)$ is given by $f_{\boldsymbol{\theta}}(y) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\{-\frac{1}{2\sigma^2}(y - \mu)^2\}$. Then

$$\begin{aligned} \frac{\partial \log f_{\boldsymbol{\theta}}(y)}{\partial \mu} &= \frac{y - \mu}{\sigma^2} \\ \frac{\partial \log f_{\boldsymbol{\theta}}(y)}{\partial \sigma} &= -\frac{1}{\sigma} + \frac{(y - \mu)^2}{\sigma^3} \end{aligned}$$

According to Theorem 2.2, the Fisher information matrix of the ZAZIG distribution is

$$\mathbf{F}_{\text{ZAZIG}} = \begin{bmatrix} \phi^{-1}(1-\phi)^{-1} & 0 & 0 \\ 0 & \frac{1-\phi}{\sigma^2} & 0 \\ 0 & 0 & \frac{2(1-\phi)}{\sigma^2} \end{bmatrix}$$

Note that in this case, given a random sample Y_1, \dots, Y_n from ZAZIG, $\hat{\mu} = \frac{1}{m} \sum_{i:Y_i \neq 0} Y_i$ and $\hat{\sigma} = \left(\frac{1}{m} \sum_{i:Y_i \neq 0} (Y_i - \hat{\mu})^2 \right)^{1/2}$ have explicit formulae, where $m = \#\{i : Y_i \neq 0\}$. \square

Example 2.9. Zero-altered-zero-inflated log-normal model (ZAZILN) This model, also known as zero-inflated log-normal (ZILN) model, has been used by, for example, Zhou and Tu (1999) to study the effects of a prospective DUR intervention program for randomized clinical trials. The pdf of the baseline distribution with parameters $\boldsymbol{\theta} = (\mu, \sigma) \in \mathbb{R} \times (0, \infty)$ is given by $f_{\boldsymbol{\theta}}(y) = \frac{1}{y\sqrt{2\pi\sigma^2}} \exp\{-\frac{1}{2\sigma^2}(\log y - \mu)^2\}$, $y > 0$. Then

$$\begin{aligned} \frac{\partial \log f_{\boldsymbol{\theta}}(y)}{\partial \mu} &= \frac{\log y - \mu}{\sigma^2} \\ \frac{\partial \log f_{\boldsymbol{\theta}}(y)}{\partial \sigma} &= -\frac{1}{\sigma} + \frac{(\log y - \mu)^2}{\sigma^3} \end{aligned}$$

According to Theorem 2.2, the Fisher information matrix of the ZAZILN distribution is

$$\mathbf{F}_{\text{ZAZILN}} = \begin{bmatrix} \phi^{-1}(1-\phi)^{-1} & 0 & 0 \\ 0 & \frac{1-\phi}{\sigma^2} & 0 \\ 0 & 0 & \frac{2(1-\phi)}{\sigma^2} \end{bmatrix}$$

which is exactly the same as the one in Example 2.8. \square

Example 2.10. Zero-altered-zero-inflated half-normal model (ZAZIHN) Also known as a zero-inflated half-normal model (ZIHN), this model has been used by, for example, Reyna-Hurtado et al. (2012) as a candidate distribution for modeling the animal movement distances in the wild. In our notations, the pdf of its baseline distribution with parameter $\boldsymbol{\theta} = \sigma \in (0, \infty)$, known as a standard half-normal distribution, is given by $f_{\boldsymbol{\theta}}(y) = \frac{\sqrt{2}}{\sqrt{\pi}\sigma^2} \exp\{-\frac{y^2}{2\sigma^2}\}$, $y > 0$. Then $\frac{\partial \log f_{\boldsymbol{\theta}}(y)}{\partial \sigma} = -\frac{1}{\sigma} + \frac{y^2}{\sigma^3}$. According to Theorem 2.2, the Fisher information matrix of the ZAZIHN distribution is

$$\mathbf{F}_{\text{ZAZIHN}} = \begin{bmatrix} \phi^{-1}(1-\phi)^{-1} & 0 \\ 0 & \frac{2(1-\phi)}{\sigma^2} \end{bmatrix}$$

It should be noted that other forms of ZIHN models have also been used in the literature. For example, Chen and Dunson (2003) utilized a more general ZIHN distribution as a prior for a hierarchical Bayesian model. The continuous component of their ZIHN is a general normal distribution $N(\mu, \sigma^2)$ truncated below by zero. In that case, $\boldsymbol{\theta} = (\mu, \sigma)$. \square

Example 2.11. Zero-altered-zero-inflated exponential model (ZAZIE) Also known as zero-inflated exponential model (ZIE), this model has been used by, for example, Huang

et al. (2019) for modeling casualty rates in ship collision. The pdf of its baseline distribution with parameter $\theta = \lambda \in (0, \infty)$ can be written as $f_{\theta}(y) = \lambda e^{-\lambda y}$, $y > 0$. Then $\frac{\partial \log f_{\theta}(y)}{\partial \lambda} = \frac{1}{\lambda} - y$. According to Theorem 2.2, the Fisher information matrix of the ZAZIE distribution is

$$\mathbf{F}_{\text{ZAZIE}} = \begin{bmatrix} \phi^{-1}(1 - \phi)^{-1} & 0 \\ 0 & \frac{1 - \phi}{\lambda^2} \end{bmatrix}$$

Note that the MLE of λ has an explicit form $\hat{\lambda} = \frac{m}{\sum_{i: Y_i \neq 0} Y_i}$ given a random sample Y_1, \dots, Y_n from ZAZIE, where $m = \#\{i : Y_i \neq 0\}$. \square

2.4 Model selection based on KS and likelihood ratio tests

Given so many zero-altered or zero-inflated models listed in Sections 2.1, 2.2 and 2.3, a critical question is which model is the most appropriate one for a given dataset.

The Kolmogorov-Smirnov (KS) test has been commonly used for testing whether a random sample $\{Y_1, \dots, Y_n\}$ comes from a continuous cumulative distribution function $F_{\theta}(y)$ with specified model parameter(s) θ (Massey Jr, 1951). It is based on the KS statistic $D_n = \sup_y |F_n(y) - F_{\theta}(y)|$, where $F_n(y) = n^{-1} \sum_{i=1}^n \mathbf{1}_{(-\infty, y]}(Y_i)$ is known as the *empirical distribution function*. Dimitrova et al. (2020) extended the KS test for general distributions $F_{\theta}(y)$ with known parameter(s) θ , including discrete and mixed ones. For typical applications, θ is unknown and an estimate $\hat{\theta}$ is plugged in when calculating D_n , which tends to overestimate the corresponding p value (Lilliefors, 1967, 1969; Aldirawi et al., 2019). To overcome the biasedness of estimated p value due to the plugged-in estimated parameters, Aldirawi et al. (2019) proposed a bootstrapped Monte Carlo estimate $\frac{\#\{D_n^{(b)} > D_n\} + 1}{B + 1}$ for the p value in their Algorithm 1, where $D_n^{(b)} = \sup_y |F_n^{(c)}(y) - F_{\hat{\theta}^{(b)}}(y)|$, $F_n^{(c)}(y)$ is the empirical distribution function of a random sample $\mathbf{Y}^{(c)} = \{Y_1^{(c)}, \dots, Y_n^{(c)}\}$ from $F_{\hat{\theta}^{(b)}}$, $\hat{\theta}^{(b)}$ is the MLE based on a bootstrapped sample $\mathbf{Y}^{(b)} = \{Y_1^{(b)}, \dots, Y_n^{(b)}\}$ of $\{Y_1, \dots, Y_n\}$, $b = 1, \dots, B$, and B is a predetermined large number, typically $B = 1000$. If the estimated p value is larger than 0.05, we say that the specified distribution passes the KS test for the given dataset.

Since $D_n = \sup_y |F_n(y) - F_{\hat{\theta}}(y)|$ when θ is unknown, where both F_n and $\hat{\theta}$ are based on the same data $\{Y_1, \dots, Y_n\}$, a more reasonable bootstrapped version of D_n would be $D_n^{(b)'} = \sup_y |F_n^{(c)}(y) - F_{\hat{\theta}^{(c)}}(y)|$, where $\hat{\theta}^{(c)}$ is the MLE based on $\mathbf{Y}^{(c)} = \{Y_1^{(c)}, \dots, Y_n^{(c)}\}$ instead of $\mathbf{Y}^{(b)}$. In this paper, we propose the following nested bootstrap estimate for KS test p value based on the above argument.

To make a distinction, we call Aldirawi et al. (2019)'s Algorithm 1 as Algorithm 1A and our Algorithm 1 as Algorithm 1B. The corresponding R functions are named `kstest.A` and `kstest.B`, respectively. According to our simulation studies in Section 3.4, we recommend `kstest.B` for small sample sizes such as $n = 30, 50$. For larger sample sizes, since the difference in terms of test power is negligible, we recommend `kstest.A` for less computational cost.

In practice, it is not uncommon that two or more distributions pass the KS test for the same dataset, especially when the sample size is moderate or small (see, for example, Aldirawi et al. (2019)). In this situation, likelihood ratio tests may be used for pairwise

Algorithm 1 Nested Bootstrap Algorithm for Estimating p value of KS Test

- 1: Given data $\mathbf{Y} = \{Y_1, Y_2, \dots, Y_n\}$, calculate the MLE $\hat{\boldsymbol{\theta}}$ of $\boldsymbol{\theta}$ and the KS statistic $D_n = \sup_y |F_n(y) - F_{\hat{\boldsymbol{\theta}}}(y)|$.
 - 2: For $b = 1, \dots, B$, do steps 3~7.
 - 3: Resample \mathbf{Y} with replacement to get a bootstrapped sample $\mathbf{Y}^{(b)} = \{Y_1^{(b)}, \dots, Y_n^{(b)}\}$.
 - 4: Calculate the MLE $\hat{\boldsymbol{\theta}}^{(b)}$ of $\boldsymbol{\theta}$ based on $\mathbf{Y}^{(b)}$.
 - 5: Simulate a random sample $\mathbf{Y}^{(c)} = \{Y_1^{(c)}, \dots, Y_n^{(c)}\}$ from $F_{\hat{\boldsymbol{\theta}}^{(b)}}$.
 - 6: Calculate the MLE $\hat{\boldsymbol{\theta}}^{(c)}$ of $\boldsymbol{\theta}$ based on $\mathbf{Y}^{(c)}$.
 - 7: Calculate the bootstrapped KS statistic $D_n^{(b)'} = \sup_y |F_n^{(c)}(y) - F_{\hat{\boldsymbol{\theta}}^{(c)}}(y)|$, where $F_n^{(c)}(y)$ is the empirical distribution function of $\mathbf{Y}^{(c)}$.
 - 8: Estimate the p value of the KS test by $\frac{\#\{b|D_n^{(b)'} > D_n\}}{B}$.
-

comparisons. More specifically, for testing $H_0 : Y_1, \dots, Y_n \text{ iid } \sim f(y; \boldsymbol{\theta})$ with unknown parameter(s) $\boldsymbol{\theta}$ against $H_1 : Y_1, \dots, Y_n \text{ iid } \sim g(y; \boldsymbol{\delta})$ with unknown parameter(s) $\boldsymbol{\delta}$, the likelihood ratio test statistic in log scale (see, for example, Aldirawi et al. (2019)) can be defined as

$$\Lambda = \log \frac{\prod_{i=1}^n f(Y_i; \hat{\boldsymbol{\theta}})}{\prod_{i=1}^n g(Y_i; \hat{\boldsymbol{\delta}})}$$

where $\hat{\boldsymbol{\theta}}$ and $\hat{\boldsymbol{\delta}}$ are the corresponding maximum likelihood estimates. Smaller Λ values are in favor of the alternative distribution $g(y; \boldsymbol{\delta})$.

To overcome the possible biasedness due to plugged-in estimated parameters, we adopt the bootstrapped estimate of p value described by Algorithm 2 of Aldirawi et al. (2019). More specifically, (i) bootstrap samples $\mathbf{Y}^{(b)} = \{Y_1^{(b)}, \dots, Y_n^{(b)}\}$, $b = 1, \dots, B$ are obtained from the original data $\mathbf{Y} = \{Y_1, \dots, Y_n\}$; (ii) MLEs $\hat{\boldsymbol{\theta}}^{(b)}$ and $\hat{\boldsymbol{\delta}}^{(b)}$ are calculated based on $\mathbf{Y}^{(b)}$; (iii) a random sample $\mathbf{Y}^{(c)} = \{Y_1^{(c)}, \dots, Y_n^{(c)}\}$ is simulated from $f(y; \hat{\boldsymbol{\theta}}^{(b)})$; (iv) bootstrapped test statistic $\Lambda^{(b)}$ is calculated based on $\mathbf{Y}^{(c)}$; and (v) the estimated p value is $\frac{\#\{b|\Lambda^{(b)} < \Lambda\}}{B}$. If the p value is less than 0.05, we claim that H_1 is significantly better; otherwise, we stick to H_0 .

2.5 Zero-altered model versus zero-inflated model

Clearly a zero-altered model (1) and its corresponding zero-inflated model (6) are connected by sharing the same baseline distribution $f_{\boldsymbol{\theta}}(y)$. Under some conditions, they are actually equivalent due to the following theorem.

Theorem 2.3. Let $f_{ZA}(y | \phi_{ZA}, \boldsymbol{\theta})$ be a zero-altered model as in (1) and $f_{ZI}(y | \phi_{ZI}, \boldsymbol{\theta})$ be the corresponding zero-inflated model as in (6) with the same baseline distribution $f_{\boldsymbol{\theta}}(y)$.

- (i) Given $f_{ZI}(y | \phi_{ZI}, \boldsymbol{\theta})$, we let $\phi_{ZA} = \phi_{ZI} + (1 - \phi_{ZI})p_0(\boldsymbol{\theta})$ and then $f_{ZA}(y | \phi_{ZA}, \boldsymbol{\theta}) = f_{ZI}(y | \phi_{ZI}, \boldsymbol{\theta})$ for all y .
- (ii) Given $f_{ZA}(y | \phi_{ZA}, \boldsymbol{\theta})$ with $\phi_{ZA} \geq p_0(\boldsymbol{\theta})$, we let $\phi_{ZI} = [\phi_{ZA} - p_0(\boldsymbol{\theta})]/[1 - p_0(\boldsymbol{\theta})]$ and then $f_{ZI}(y | \phi_{ZI}, \boldsymbol{\theta}) = f_{ZA}(y | \phi_{ZA}, \boldsymbol{\theta})$ for all y .

The proof of Theorem 2.3 is relegated to the Appendix. Theorem 2.3 implies that if $\phi_{ZA} \geq p_0(\boldsymbol{\theta})$, which indicates that the data is zero-inflated, then a ZA model is equivalent to the corresponding ZI model. Given a random sample from a zero-inflated model $f_{ZI}(y | \phi_{ZI}, \boldsymbol{\theta})$, a KS test will conclude that some zero-altered model $f_{ZA}(y | \phi_{ZA}, \boldsymbol{\theta})$ seems fine with the data as well. The other direction is slightly different though. That is, if the true model is a zero-altered one, only when ϕ_{ZA} is at least as large as $p_0(\boldsymbol{\theta})$, a KS test will conclude that some zero-inflated model seems to be true as well. In Section 3.5, we will provide a numerical example such that both ZIBNB and BNBH fit the data well.

3 Software Implementation and Numerical Analysis

In this section, the R implementation of the proposed package AZIZD is introduced and its numerical analysis is performed. A real data example is used to illustrate how our package can be used in practice.

Compared with existing R packages each covering only a limited number of baseline distributions, our package covers many more discrete and continuous distributions including Poisson, geometric, negative binomial, beta binomial, beta negative binomial, normal (or Gaussian), log-normal, half-normal, and exponential distributions along with their zero-altered and zero-inflated models, which facilitates the potential users to choose the most appropriate model from a large class of candidates for their dataset. For all the models mentioned above, we provide the corresponding Fisher information matrix and confidence intervals for estimated parameters, which allows users to run hypothesis tests and make further inference.

3.1 Improvements over existing packages on finding MLEs

From Section 2.4, we can see that accurate MLE is a critical component for both KS test and likelihood ratio test. In this section, we first summarize the improvements of the proposed AZIAD over existing R packages on finding MLEs for zero-altered and zero-inflated models.

- (1) Compared with other packages, AZIAD provides MLEs well even for extreme cases including $\phi = 0$ or $\phi = 1$ (please see the comparison analysis in Section 3.3).
- (2) Some packages encountered error message `L-BFGS-B needs finite values of "fn"` for some zero-inflated data when using, for example, the function `dis.kstest` in package `iZID`. This issue is solved in AZIAD by specifying appropriate lower

bound and upper bound for optimizations (please see Section 3.3 for a comparison).

- (3) Compared with package `iZID` which also covered MLEs for ZIBNB, BNBH, ZIBB, and BBH models, our results are more reliable (see Example 3.2 for a comparison) by applying Theorem 4 in Aldirawi and Yang (2022). More specifically, we separate the two situations (1) $m/n \leq 1 - p_0(\boldsymbol{\theta}_*)$ and (2) $m/n > 1 - p_0(\boldsymbol{\theta}_*)$ and calculate the MLEs in each case.
- (4) Some parameters are originally defined as positive integers, such as n in Examples 2.2 and 2.6 and r in Examples 2.3 and 2.5, while in practice they could be extended to positive real numbers. In AZIAD, we seek for real-valued MLEs by default. In the mean time, we keep the option of integer-valued n or r in response to users' call.

In the rest part of this section, we use examples to illustrate how to use our package to find the MLEs and the improved accuracy by using our package.

Example 3.1. In order to find the MLE for the parameter(s) of zero-inflated or hurdle models, the main function built in AZIAD is

```
zih.mle(x, r, p, alpha1, alpha2, n, lambda, mean, sigma, type = c("zi", "h"),
dist, lowerbound = 0.01, upperbound = 10000)
```

where `x` is a sequence of numbers, which could be integers for discrete cases or real numbers for continuous cases; the arguments `r`, `p`, `alpha1`, `alpha2`, `n`, `lambda`, `mean`, and `sigma` are initial values of the corresponding parameters; `dist` could be chosen as `poisson.zihmle`, `geometric.zihmle`, `nb.zihmle`, `nb1.zihmle`, `bb.zihmle`, `bb1.zihmle`, `bnb.zihmle`, `bnb1.zihmle`, `normal.zihmle`, `halfnorm.zihmle`, `lognorm.zimle`, and `exp.zihmle`, which correspond to zero-inflated or zero-altered Poisson, geometric, negative binomial, negative binomial with integer-valued r , beta binomial, beta binomial with integer-valued n , beta negative binomial, beta negative binomial with integer-valued r , normal, log-normal, half-normal, and exponential distributions; option `type` indicates the type of distribution is zero-inflated (`zi`) or hurdle/zero-altered (`h`); `lowerbound` and `upperbound` specify the searching range of parameters when maximizing the likelihood function. For instance, in order to calculate the MLE of zero-inflated geometric distribution (see Example 2.4), one may use the following R code:

```
R> set.seed(008)
R> x22=sample.h1(2000,phi=0.3,dist='geometric',p=0.3)
R> zih.mle(x22,p=0.2,dist="geometric.zihmle",type="h")
      p      phi    loglik
0.2942292 0.3015 -4101.05
```

Our estimates $\hat{p} = 0.2942292$ and $\hat{\phi} = 0.3015$ are fairly close to the true parameter values $p = 0.3$ and $\phi = 0.3$. □

Example 3.2. To compare the performance of our AZIAD package and the existing iZID on finding MLEs, we generate random samples from a BNBH model (see Example 2.3) with parameters $(\phi, r, \alpha_1, \alpha_2) = (0.3, 5, 8, 3)$, and a BBH model (see Example 2.2) with parameters $(\phi, n, \alpha_1, \alpha_2) = (0.6, 5, 8, 3)$, each with increasing sample sizes $N = 10^4, 5 \times 10^4, 20 \times 10^4$ and 100×10^4 . To see the converging pattern more clearly, we generate nested datasets such that any dataset with a smaller N is a subset of the corresponding datasets with bigger N s, if they are simulated from the same distribution.

For each sample, we find the MLEs $(\hat{\phi}, \hat{\theta}_1, \hat{\theta}_2, \hat{\theta}_3)$ of the parameters and calculate the aggregated L_1 relative distance (L1RD) $L_1 = |\hat{\phi} - \phi|/|\phi| + \sum_{i=1}^3 |\hat{\theta}_i - \theta_i|/|\theta_i|$. The implemented R codes are as follows:

```
R> set.seed(167)
R> hi1=sample.h1(N=1000000,phi=0.3,dist="bnb",r=5,alpha1=8,alpha2=3)
R> hi2=hi1[1:200000]
R> hi3=hi1[1:50000]
R> hi4=hi1[1:10000]
R> mle13=zih.mle(hi4,type="h",r=6,alpha1=9,alpha2=4,dist="bnb.zihmle")
R> mle13=zih.mle(hi4,type="h",r=6,alpha1=9,alpha2=4,dist="bnb1.zihmle")
R> mle14=bnb.zihmle(hi4,type="h",r=6,alpha1=9,alpha2=4)

R> set.seed(171)
R> hi1=sample.h1(N=1000000,phi=0.6,dist="bb",n=5,alpha1 = 8,alpha2=3)
R> hi2=hi1[1:200000]
R> hi3=hi1[1:50000]
R> hi4=hi1[1:10000]
R> mle23=zih.mle(hi1,n=6,alpha1=9,alpha2=4,type="h",dist="bb.zihmle")
R> mle23=zih.mle(hi4,n=6,alpha1=9,alpha2=4,type="h",dist="bb1.zihmle")
R> mle24=bb.zihmle(hi1,n=6,alpha1=9,alpha2=3,type="h")
```

In Tables 1 and 2, we list and compare the MLEs, log-likelihood, and L1RD obtained by our package (R function `zih.mle` with real-valued or integer-valued MLEs) and the iZID package (R function `xxx.zihmle`) for BNBH and BBH models. As displayed in these tables, the estimates based on our functions are more accurate as indicated by smaller L1RD. As the sample size increases, the L1RD based on our MLEs shows an overall decreasing pattern which indicates the convergence of MLEs towards the true parameter values. Similar results are collected for ZIBNB and ZIBB as well but not shown here. \square

3.2 Fisher information, confidence interval, and test on zero-inflation

As mentioned in Sections 2.1, 2.2 and 2.3, the Fisher information matrix \mathbf{F}_{ZA} , \mathbf{F}_{ZI} or \mathbf{F}_{ZAZI} can be calculated for zero-altered/hurdle, zero-inflated, or ZAZI models, respectively. Its inverse matrix is approximately the variance-covariance matrix of parameter estimates $(\hat{\phi}, \hat{\boldsymbol{\theta}})$. For example, for zero-altered or hurdle models (see Section 2.1),

$$\sqrt{n}(\hat{\phi} - \phi) \sim N(0, \phi(1 - \phi)), \quad \sqrt{n}(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}) \sim N(\mathbf{0}, \mathbf{F}_{ZA\boldsymbol{\theta}}^{-1})$$

N	R function	r	α_1	α_2	ϕ	loglike	L1RD
1×10^4	zih.mle(bnb)	3.81	7.69	3.81	0.303	-19232.34	0.555
	zih.mle(bnb1)	4	7.70	3.64	0.303	-19232.35	0.460
	bnb.zihmle	5.83	20.68	3.89	0	777027.7	3.05
5×10^4	zih.mle(bnb)	5.35	7.69	2.67	0.301	-96201.68	0.223
	zih.mle(bnb1)	5	7.59	2.82	0.301	-96201.71	0.113
	bnb.zihmle	30.09	453.25	28.96	0	134618737	70.32
20×10^4	zih.mle(bnb)	5.36	7.88	2.74	0.299	-384542.1	0.175
	zih.mle(bnb1)	5	7.78	2.90	0.299	-384542.2	0.061
	bnb.zihmle	92.90	5492.33	119.79	0	8730091135	743.05
1×10^6	zih.mle(bnb)	5.46	8.01	2.78	0.299	-1921040	0.181
	zih.mle(bnb1)	5	7.97	2.98	0.299	-1921041	0.009
	bnb.zihmle	101.23	6842.71	138.64	0	55719787227	919.80

Table 1: Comparison of MLE estimates, log-likelihood, and L_1 relative distance (L1RD) for BNB Hurdle distribution with true parameters ($r = 5$, $\alpha_1 = 8$, $\alpha_2 = 3$, $\phi = 0.3$) using `bnb.zihmle` in `iZID` and `zih.mle` in `AZIAD` with `dist=bnb.zihmle` (real-valued r) and `dist=bnb1.zihmle` (integer-valued r) with various sample sizes N

N	R function	n	α_1	α_2	ϕ	loglike	L1RD
1×10^4	zih.mle(bb)	4.99	7.58	2.84	0.597	-12572.96	0.110
	zih.mle(bb1)	5	7.86	2.96	0.597	-12579.31	0.033
	bb.zihmle	6	9	3	0.597	28775.56	0.33
5×10^4	zih.mle(bb)	4.99	7.57	2.78	0.597	-62688.44	0.129
	zih.mle(bb1)	5	7.85	2.91	0.597	-62721.29	0.051
	bb.zihmle	6	9	3	0.597	143733.9	0.328
20×10^4	zih.mle(bb)	4.99	7.67	2.84	0.600	-249962	0.094
	zih.mle(bb1)	5	7.95	2.97	0.600	-250089.3	0.015
	bb.zihmle	6	9	3	0.600	570424.6	0.325
1×10^6	zih.mle(bb)	4.99	7.66	2.85	0.600	-1249976	0.093
	zih.mle(bb1)	5	7.94	2.97	0.600	-1250610	0.015
	bnb.zihmle	6	9	3	0.600	2850793	0.325

Table 2: Comparison of MLE estimates, log-likelihood, and L_1 relative distance (L1RD) for BB Hurdle distribution with true parameters ($n = 5$, $\alpha_1 = 8$, $\alpha_2 = 3$, $\phi = 0.6$) using `bb.zihmle` in `iZID` and `zih.mle` in `AZIAD` with `dist=bb.zihmle` (real-valued r) and `dist=bb1.zihmle` (integer-valued r) with various sample sizes N

for large n . Approximate confidence intervals can be constructed for ϕ and θ . For example, Equation (5) provides an approximate confidence interval for ϕ in ZA models.

In our package AZIAD, we use the function

```
FI.ZI(x, dist="poisson", r = NULL, p = NULL, alpha1 = NULL,
alpha2 = NULL, n = NULL, lambda=NULL, mean=NULL, sigma=NULL,
lowerbound = 0.01, upperbound = 10000)
```

to calculate (the inverse of) the Fisher information matrix at the MLE and 95% approximate confidence intervals for all parameters, where x is the data in its vector form; `dist` can be `poisson`, `geometric`, `nb`, `bb`, `bnb`, `normal`, `halfnormal`, `lognormal`, `exponential`, `zip`, `zigeom`, `zinb`, `zibb`, `zibnb`, `zinormal`, `zilognorm`, `zihalfnorm`, `ziexp`, `ph`, `geomh`, `nbh`, `bbh`, and `bnbh` for Poisson, geometric, negative binomial, beta binomial, beta negative binomial, normal/Gaussian, log-normal, half-normal, exponential, their zero-inflated versions and hurdle versions, respectively; `r`, `p`, `alpha1`, `alpha2`, `n`, `lambda`, `mean`, `sigma` provide initial values of distribution parameters; `lowerbound` and `upperbound` are predetermined ranges for MLEs, which could be extended if attained by any of the estimated parameter values.

Example 3.3. Confidence intervals for ZINB As an illustration, we apply the FI.ZI function to a simulated ZINB dataset with true parameters $\phi = 0.4$, $r = 10$ and $p = 0.2$.

```
R> set.seed(117)
R> N=1000;r=10;p=0.2;phi=0.4;
R> x<-sample.zi1(N,phi=phi,dist="nb",r=r,p=p)
R> FI.ZI(x,r=3,p=0.1, dist="zinb")
```

```
$inversefisher
      [,1]      [,2]      [,3]
[1,] 0.32302489 0.0661767 0.02116832
[2,] 0.06617670 35.0781124 -0.53660185
[3,] 0.02116832 -0.5366019 0.03658915
```

```
$ConfidenceIntervals
      [,1]      [,2]
CI of phi 0.3937737 0.4642262
CI of r   9.8952803 10.6294496
CI of p   0.1923346 0.2160458
```

With sample size $N = 1000$, the derived 95% confidence intervals cover the true parameter values fairly well. \square

Example 3.4. Test zero-inflation in Poisson data As an illustration, we simulate a ZIP data with $N = 1000$, $\phi = 0$ and $\lambda = 0.8$, which is actually a regular Poisson data with $\lambda = 0.8$ without zero-inflation. There are 459 zeros which roughly matches the probability of zero $p_0(\lambda) = 0.449$. Then we use the FI.ZI function with distribution Poisson Hurdle (PH) to allow both zero-inflation and deflation.

```

R> set.seed(337)
R> x<-sample.zi1(N=1000,phi=0,dist="poisson",lambda=0.8)
R> table(x)
x
0  1  2  3  4  5
459 334 153 41 10 3
R> dpois(0,lambda=0.8)
[1] 0.449329
R> FI.ZI(x,lambda=1, dist="ph")
$inversefisher
      [,1]      [,2]
[1,] 0.248319 0.000000
[2,] 0.000000 2.558941

$ConfidenceIntervals
      [,1]      [,2]
CI of Phi  0.4281146 0.4898854
CI of lambda 0.7937409 0.9920343

R> FI.ZI(x,lambda=1, dist="poisson")
$inversefisher
lambda
[1,] 0.818

$ConfidenceIntervals
[1] 0.7619437 0.8740563

```

It turns out that the 95% approximate confidence interval (0.4281, 0.4899) of ϕ based on PH model (`dist="ph"`) covers the baseline zero probability $p_0(\lambda) = 0.4493$, which indicates that neither zero-inflation nor zero-deflation is significant at the 5% level. Actually, the 95% approximate confidence interval (0.7619, 0.8741) of λ based on the regular Poisson model (`dist="poisson"`) covers the true value 0.8 roughly at the center, which is better than (0.7937, 0.9920) based on the PH model. \square

3.3 Comparison study in terms of type I error

In this section and Section 3.4, we use simulation studies to compare the performance of our algorithms for model identification with existing R functions for the same purposes. In this section we focus on type I errors. That is, given the true distribution, say ZINB, what is the chance that the test erroneously concludes that the distribution is not ZINB due to a p value less than 0.05. Such an error is known as a *type I error* (Lehmann and Romano, 2005). Ideally, such a chance is no more than 0.05, known as the size of the test.

The R functions under comparison in this section include the basic function `ks.test`, function `disc_ks_test` in package `KSgeneral` (Dimitrova et al., 2020), function `dis.kstest`

in package `iZID` (Wang et al., 2020), and two of our functions in `AZIAD`, `kstest.A` based on Aldirawi et al. (2019)'s Algorithm 1 with $(\#b + 1)/(B + 1)$ replaced by $\#b/B$, and `kstest.B` based on our Algorithm 1 described in Section 2.4.

For illustration purposes, we consider four zero-inflated models, ZIP, ZINB, ZIBNB, and ZIBB. For each model, we consider five different samples sizes, $N = 30, 50, 100, 200, 500$, and simulate $B = 1000$ independent datasets for each sample size. For each simulated dataset, we run the five R functions under comparison individually targeting the corresponding true model. If the p value of the test is less than 0.05, we count it as a type I error. The ratios of type I errors (that is, the number of type I errors divided by $B = 1000$) are listed in Tables 3, 4, 5 and 6. For readers' reference, we provide the R code for generating Table 4 below.

```
R> set.seed(007)
R> NoIterations <- 1000
R> listnb1 <- numeric(NoIterations)
R> listnb2 <- numeric(NoIterations)
R> listnb3 <- numeric(NoIterations)
R> listnb4 <- numeric(NoIterations)
R> listnb5 <- numeric(NoIterations)
R> for(i in 1:NoIterations){
R>   sampled <- sample.zi(N=500,phi=0.3,dist = 'nb',r=5,p=.2)
R>   listnb1[i]<-dist.ksnew(sampled,nsim=100,bootstrap = TRUE,dist= 'zinb',
R>     lowerbound = 1e-10, upperbound = 100000)$pvalue
R>   listnb2[i]<-dist.ksmle(sampled,nsim=100,bootstrap = TRUE,dist= 'zinb',
R>     lowerbound = 1e-10, upperbound = 100000)$pvalue
R>   listnb3[i]<-try(dis.kstest(sampled,nsim=100,bootstrap = TRUE,distri='zinb',
R>     lowerbound = 1e-10, upperbound = 100000)$pvalue }
R> for(i in 1:NoIterations) {
R>   sampled <-sample.zi(N=30,phi=0.3,dist = 'nb',r=5,p=.2)
R>   a=zih.mle(sampled,r=10,p=.3,type="zi",dist = "nb1.zihmle")
R>   sampled1<- sample.zi(N=30,r=a[1],p=a[2],phi=a[3],dist="nb")
R>   listnb4[i]<-ks.test(sampled,sampled1)$p.value
R>   x.unique=sort(unique(sampled))
R>   probs_ori = a[3] + (1 - a[3]) * stats::pnbinom(x.unique,
R>     size = a[1], prob = a[2])
R>   step_ori = stats::stepfun(x.unique, c(0, probs_ori))
R>   listnb5[i]<-disc_ks_test(x=sampled,y=step_ori,exact = TRUE)$p.value }
```

According to tables 3 to 6, `disc_ks_test` (package `KSgeneral`) seems to have larger type I error rates than the nominal level 0.05. One concern about function `dis.kstest` (`iZID`) for ZINB, ZIBNB and ZIBB distributions is its significant portions of NA's due to errors. Our two functions, `kstest.A` and `kstest.B`, and R basic function `ks.test` have type I error rates nearly zero, which are satisfactory.

Sample Size N	30	50	100	200	500
ks.test	0	0	0	0.001	0.005
disc_ks.test	0.074	0.116	0.081	0.112	0.08
dis.kstest	0	0	0	0	0
kstest.A	0	0	0.001	0	0
kstest.B	0	0	0	0	0

Table 3: Type I error rates of KS tests on whether the data comes from ZIP, based on $B = 1000$ simulated ZIP datasets with parameters $\phi = 0.3$, $\lambda = 10$ for each sample size N

Sample Size N	30	50	100	200	500
ks.test	0.001	0.002	0.001	0.004	0.003
disc_ks.test	0.062	0.065	0.099	0.089	0.107
dis.kstest	0.003 (536NA)	0.004 (638NA)	0.003 (903NA)	0.001 (996NA)	All NA
kstest.A	0	0	0	0	0
kstest.B	0	0	0	0	0

Table 4: Type I error rates of KS tests on whether the data comes from ZINB, based on $B = 1000$ simulated ZINB datasets with parameters $\phi = 0.3$, $r = 5$, $p = 0.2$ for each sample size N

Sample Size N	30	50	100	200	500
ks.test	0.058	0	0	0.001	0
disc_ks.test	0.062	0.049	0.067	0.07	0.084
dis.kstest	0.788 (176NA)	0.889 (89NA)	0.921 (87NA)	0.832 (168NA)	0.671 (383NA)
kstest.A	0	0	0	0.002	0.002
kstest.B	0.002	0	0	0.003	0.001

Table 5: Type I error rates of KS tests on whether the data comes from ZIBNB, based on $B = 1000$ simulated ZIBNB datasets with parameters $\phi = 0.3$, $r = 3$, $\alpha_1 = 3$, $\alpha_2 = 5$ for each sample size N

Sample Size N	30	50	100	200	500
ks.test	0	0	0.001	0	0
disc_ks.test	0.054	0.06	0.06	0.058	0.066
dis.kstest	0.039 (24NA)	0.05 (34NA)	0.084 (47NA)	0.645 (64NA)	0.947 (53NA)
kstest.A	0.001	0	0	0	0
kstest.B	0.001	0	0	0	0

Table 6: Type I error rates of KS tests on whether the data comes from ZIBB, based on $B = 1000$ simulated ZIBB datasets with parameters $\phi = 0.3$, $n = 5$, $\alpha_1 = 8$, $\alpha_2 = 3$ for each sample size N

3.4 Comparison study in terms of test power

In this section, we continue the simulation studies in Section 3.3 and compare the power of the tests based on four different R functions. More specifically, given the $B = 1000$

datasets simulated from, for example, the ZIP distribution, we run a KS test on whether the data comes from a ZINB distribution, and denote the test as ZIP versus ZINB. If we reject ZINB distribution for 800 times, then the empirical power of the KS test at ZINB is $800/1000 = 0.80$. Given that the type I error rate does not go beyond the nominal level 0.05, the bigger the power is, the better the test performs (see, for example, Lehmann and Romano (2005)). We remove function `dis.kstest` of package `iZID` from power analysis since it generates too many NA's for ZINB (see Table 4) or has a type I error rate much higher than 0.05 for ZIBNB (see Table 5) and ZIBB (see Table 6).

Table 7 shows that our functions, `kstest.A` and `kstest.B`, have larger power at ZIP versus ZIBNB. All tests fail at ZIP versus ZINB and ZIP versus ZIBB. It often happens for a test of a simpler model versus a more flexible model, especially when the flexible model could approximate the simpler model well (see Section 4 for a discussion on it). Table 8 shows that our functions have comparable power at ZINB versus ZIP. All tests fail at ZINB versus ZIBNB. All tests lose power gradually at ZINB versus ZIBB, as sample size N increases. Table 9 shows that our functions have the largest powers at ZIBNB versus ZIP. For ZIBNB versus ZINB, our functions have larger power than `ks.test` but less power than `disc_ks.test`, which tends to reject more for ZINB (see Table 4). All tests lose power quickly at ZIBNB versus ZIBB, while `disc_ks.test` shows a slower decreasing rate. Table 10 shows that our functions have significantly larger power at ZIBB versus ZIBNB. Function `disc_ks.test` has greater power at ZIBB versus ZINB and ZIBB versus ZIP when $N \leq 100$. Note that both `ks.test` and `disc_ks.test` here rely on the MLEs for ZIBB and ZIBNB provided by our package `AZIAD`.

Overall our two functions `kstest.A` and `ks.test.B` are most reliable for KS tests involving zero-inflated models. In practice, we recommend `ks.testB` for small sample sizes such as $N = 30, 50$ (see ZIP versus ZIBNB in Table 7, ZINB versus ZIP in Table 9, and ZIBNB versus ZIP in Table 9) and `ks.testA` for large sample size such as $N = 100, 200, 500$, which is faster.

Test	Function	$N = 30$	$N = 50$	$N = 100$	$N = 200$	$N = 500$
ZIPvsZINB	<code>ks.test</code>	0.002	0	0.001	0.001	0
	<code>disc_ks.test</code>	0.079	0.087	0.076	0.096	0.093
	<code>kstest.A</code>	0	0	0	0	0
	<code>kstest.B</code>	0	0	0	0	0
ZIPvsZIBNB	<code>ks.test</code>	0.132	0.441	0.844	0.978	0.984
	<code>disc_ks.test</code>	0.115	0.096	0.084	0.099	0.101
	<code>kstest.A</code>	0.749	0.892	0.954	0.973	0.992
	<code>kstest.B</code>	0.73	0.984	0.954	0.973	0.992
ZIPvsZIBB	<code>ks.test</code>	0.002	0.001	0.001	0.005	0.007
	<code>disc_ks.test</code>	0.079	0.084	0.09	0.074	0.091
	<code>kstest.A</code>	0	0	0	0.001	0
	<code>kstest.B</code>	0	0	0	0	0

Table 7: Empirical power of KS tests on whether the data comes from ZINB, ZIBNB or ZIBB, based on $B = 1000$ simulated ZIP datasets with parameters $\phi = 0.3$, $\lambda = 10$ for each sample size N

Test	Function	$N = 30$	$N = 50$	$N = 100$	$N = 200$	$N = 500$
ZINBvsZIP	ks.test	0.05	0.111	0.325	0.801	1
	disc_ks.test	0.316	0.54	0.89	1	1
	kstest.A	0.2	0.501	0.942	1	1
	kstest.B	0.199	0.512	0.942	1	1
ZINBvsZIBNB	ks.test	0	0	0	0	0
	disc_ks.test	0.054	0.045	0.045	0.041	0.056
	kstest.A	0	0	0	0	0
	kstest.B	0	0	0	0	0
ZINBvsZIBB	ks.test	0.297	0.529	0.6	0.567	0.429
	disc_ks.test	0.634	0.663	0.646	0.57	0.457
	kstest.A	0.632	0.641	0.59	0.535	0.368
	kstest.B	0.632	0.642	0.59	0.535	0.368

Table 8: Empirical power of KS tests on whether the data comes from ZIP, ZIBNB or ZIBB, based on $B = 1000$ simulated ZINB datasets with parameters $\phi = 0.3$, $r = 3$, $\alpha_1 = 3$, $\alpha_2 = 5$ for each sample size N

Test	Function	$N = 30$	$N = 50$	$N = 100$	$N = 200$	$N = 500$
ZIBNBvsZIP	ks.test	0.183	0.413	0.751	0.975	1
	disc_ks.test	0.552	0.764	0.981	1	1
	kstest.A	0.569	0.866	0.996	1	1
	kstest.B	0.573	0.871	0.996	1	1
ZIBNBvsZINB	ks.test	0.006	0.011	0.022	0.021	0.099
	disc_ks.test	0.106	0.14	0.145	0.253	0.437
	kstest.A	0.019	0.016	0.043	0.097	0.248
	kstest.B	0.018	0.019	0.046	0.102	0.307
ZIBNBvsZIBB	ks.test	0.422	0.266	0.167	0.069	0.046
	disc_ks.test	0.473	0.337	0.251	0.209	0.285
	kstest.A	0.413	0.282	0.144	0.084	0.056
	kstest.B	0.413	0.282	0.144	0.084	0.056

Table 9: Empirical power of KS tests on whether the data comes from ZIP, ZINB or ZIBB, based on $B = 1000$ simulated ZIBNB datasets with parameters $\phi = 0.3$, $r = 3$, $\alpha_1 = 3$, $\alpha_2 = 5$ for each sample size N

3.5 Real data analysis

In this section, we use a real data `DebTrivedi` from R package `MixAll` to illustrate how to use our package `AZIAD` to identify the most appropriate zero-inflated model. The `DebTrivedi` data was obtained from the US National Medical Expenditure Survey (Deb and Trivedi, 1997; Zeileis et al., 2008). It contains 19 variables from 4,406 individuals aged 66 and over. For illustration purpose, we select the variable `ofp`, which is the number of physician office visits.

In order to analyze the data, we first use KS tests to check each distribution covered by our package. Since the sample size 4,406 is fairly large, we use our function `kstest.A` instead of `kstest.B`. Out of 19 distributions under test, we have 7 models with p value larger than 0.05, including geometric, NB (with real-valued r), NB1 (with integer-valued

Test	Function	$N = 30$	$N = 50$	$N = 100$	$N = 200$	$N = 500$
ZIBBvsZIP	ks.test	0.01	0.017	0.038	0.235	0.983
	disc_ks.test	0.241	0.299	0.714	0.963	1
	kstest.A	0.001	0.037	0.459	0.986	1
	kstest.B	0.01	0.05	0.455	0.992	1
ZIBBvsZINB	ks.test	0.008	0.016	0.052	0.246	0.989
	disc_ks.test	0.237	0.316	0.688	0.965	1
	kstest.A	0.015	0.058	0.494	0.99	1
	kstest.B	0.015	0.063	0.494	0.99	1
ZIBBvsZIBNB	ks.test	0.011	0.039	0.144	0.548	1
	disc_ks.test	0.86	0.075	0.085	0.126	0.262
	kstest.A	0.275	0.75	0.999	1	1
	kstest.B	0.265	0.739	0.998	1	1

Table 10: Empirical power of KS tests on whether the data comes from ZIP, ZINB or ZIBNB, based on $B = 1000$ simulated ZIBB datasets with parameters $\phi = 0.3$, $n = 5$, $\alpha_1 = 8$, $\alpha_2 = 3$ for each sample size N

r), ZIBB, ZIBNB, BBH, and BNBH. The relevant R code is displayed below:

```

set.seed(1008)
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="poisson")$pvalue      #0
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="geometric")$pvalue  #0.06
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="nb")$pvalue         #0.08
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="nb1")$pvalue       #0.105
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="bb")$pvalue        #0.02
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="bb1")$pvalue      #0.04
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="bnb")$pvalue       #0.015
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="bnb")$pvalue       #0.025
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="bnb1")$pvalue     #0.01
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="zip")$pvalue       #0
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="zigeom")$pvalue    #0.015
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="zinb")$pvalue     #0.005
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="zibb")$pvalue     #0.085
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="zibnb")$pvalue    #0.85
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="ph")$pvalue       #0
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="geomh")$pvalue    #0.005
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="nbh")$pvalue      #0.005
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="bbh")$pvalue     #0.08
kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="bnbh")$pvalue     #0.825

```

Since there are seven distributions passing the KS test, we further run our likelihood ratio test function `lrt.A` to compare each pair of the candidate distributions. For example, if `d1` represents geometric distribution and `d2` represents NB distribution, we run `lrt.A(d1, d2)`. A p value less than 0.05 indicates that `d2` is significantly better than `d1`. The relevant R code is displayed below:

$H_0 \backslash H_1$	Geometric	NB	NB1	ZIBB	ZIBNB	BBH	BNBH
Geometric	1	0.88	0.79	0.9	0	0.81	0
NB	0.07	1	0.165	0.835	0	0.49	0
NB1	0.84	0.82	1	0.895	0	0.8	0
ZIBB	0	0.04	0.015	1	0	0.96	0
ZIBNB	0.66	0.725	0.69	0.81	1	0.83	1
BBH	0.005	0.09	0.015	1	0	1	0
BNBH	0.705	0.73	0.67	0.81	0.995	0.735	1

Table 11: The p values of pairwise comparisons of the seven candidate models based on `lrt.A` for data `DebTrivedi`, a p value less than 0.05 indicating that the corresponding column model is significantly better than the corresponding row model

```
#Best Model By using LRT
d1=kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="geometric")
d2=kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="nb")
d3=kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="nb1")
d4=kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="zibb")
d5=kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="zibnb")
d6=kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="bbh")
d7=kstest.A(ofp,nsim=200,bootstrap=TRUE,dist="bnbh")
pmat = matrix(, nrow=7, ncol=7)
dvec = list(d1,d2,d3,d4,d5,d6,d7)
for(i in 1:7) for(j in 1:7) pmat[i,j] = lrt.A(dvec[[i]],dvec[[j]]);
```

Table 11 shows the results of pairwise comparisons of the seven candidate distributions based on `lrt.A`(H_0, H_1) with row names indicating H_0 and column names indicating H_1 . A small p value implies that the column distribution is significantly better than the row distribution. It should be noted that in general the p values of `lrt.A`(d_1, d_2) and `lrt.A`(d_2, d_1) are not equal. Based on the p values in Table 11, we conclude that ZIBNB and BNBH are significantly better than geometric, NB, NB1, ZIBB and BBH, while there is no significant difference between ZIBNB and BNBH, which confirms our conclusion in Section 2.5. Note that neither ZIBNB nor BNBH was considered by Deb and Trivedi (1997) and Zeileis et al. (2008). Both distributions were recommended by Aldirawi et al. (2019) and Aldirawi and Yang (2022) for microbiome data analysis.

We run the same procedure based on our function `lrt.B` as well, which is based on `kstest.B`. The results are consistent with `lrt.A`'s. Similarly as in Section 3.4, we recommend `lrt.B` for cases with smaller sample sizes and `lrt.A` for cases with a sample size larger than 100.

N	30	50	100	200	500	1000	5000
$\sup_y F_{\text{ZIP}}(y) - F_{\text{ZINB}}(y) $	0.166	0.12	0.08	0.03	0.024	0.04	0.0158

Table 12: Maximum difference between the CDF of ZIP and the CDF of fitted ZINB based on random samples from $\text{ZIP}(\phi = 0.3, \lambda = 10)$ with various sample sizes N

4 Discussion

A major goal targeted in this paper is to identify the underlying distribution F_0 given a random sample $\{Y_1, \dots, Y_n\}$ from it. Given the data $\{Y_1, \dots, Y_n\}$, ideally our tests could accomplish two tasks, (i) do not reject F_0 itself; (ii) reject any F_1 which is not F_0 .

According to the Glivenko-Cantelli theorem (see, for example, Theorem 19.1 in Vaart (2000)), the empirical distribution function F_n converges to F_0 uniformly and then Task (i) can be achieved by proper KS tests up to a type I error, which is confirmed by our simulation studies in Section 3.3.

Task (ii) is much more complicated. First of all, in Section 2.5, we show the equivalence of a zero-inflated model and its corresponding hurdle model given that $\phi_{\text{ZA}} \geq p_0(\boldsymbol{\theta})$. Therefore, one could not make a distinction between a ZI model and its corresponding ZA or hurdle model when zero-inflation exists. We have such a real data example in Section 3.5.

Secondly, as shown by simulation studies in Section 3.4, one may not be able to reject F_1 if F_1 is a more flexible model than F_0 , especially when F_1 with fitted parameters could approximate F_0 well. In this section, we use ZIP versus ZINB as an example to illustrate why one may not be able to reject F_1 given that F_0 is the true model.

It is known that $\text{Poisson}(\lambda)$ can be approximated by $\text{NB}(r, p)$ with a large r and a p close to 1 such that $\lambda = r(1 - p)$ (see, for example, Teerapabolarn (2014)). To illustrate how well ZINB could approximate a ZIP distribution, we simulate random samples $\{Y_1, \dots, Y_N\}$ from $\text{ZIP}(\phi = 0.3, \lambda = 10)$ with various sample sizes N , then fit a ZINB model. In Table 12, we list the maximum difference between the cumulative distribution function (CDF) $F_{\text{ZIP}}(y)$ of ZIP and the CDF $F_{\text{ZINB}}(y)$ of the fitted ZINB. As the sample size N increases, the maximum distance $\sup_y |F_{\text{ZIP}}(y) - F_{\text{ZINB}}(y)|$ decreases, which indicates ZINB approximates ZIP better and better.

Another example is the KS test of ZIBNB versus ZIBB, whose empirical power shows a decreasing pattern as the sample size N increases (see Table 9 in Section 3.4). We perform a similar numerical study here for ZIBNB versus ZIBB as well. More specifically, we simulate random samples $\{Y_1, \dots, Y_N\}$ from $\text{ZIBNB}(\phi = 0.3, r = 15, \alpha_1 = 19, \alpha_2 = 10)$ with various sample sizes N , then fit a ZIBB model. Table 13 shows a decreasing pattern of the maximum difference between the two CDFs as the sample size N increases, which indicates ZIBB can approximate ZIBNB better and better.

Acknowledgments

This work was supported in part by the U.S. NSF grant DMS-1924859, and CSUSB research fellowship.

N	30	50	100	200	500	1000	5000
$\sup_y F_{\text{ZIBNB}}(y) - F_{\text{ZIBB}}(y) $	0.166	0.1	0.1	0.11	0.04	0.024	0.014

Table 13: Maximum difference between the CDF of ZIBNB and the CDF of fitted ZIBB based on random samples from $\text{ZIBNB}(\phi = 0.3, r = 15, \alpha_1 = 19, \alpha_2 = 10)$ with various sample sizes N

Appendix

Proof of Theorem 2.1:

According to formulae 14.13(b) in Seber (2008), if $\beta = a - \mathbf{b}^T \mathbf{D}^{-1} \mathbf{b} \neq 0$, then

$$\begin{bmatrix} a & \mathbf{b}^T \\ \mathbf{b} & \mathbf{D} \end{bmatrix}^{-1} = \begin{bmatrix} \frac{1}{\beta} & -\frac{1}{\beta} \mathbf{b}^T \mathbf{D}^{-1} \\ -\frac{1}{\beta} \mathbf{D}^{-1} \mathbf{b} & \mathbf{D}^{-1} + \frac{1}{\beta} \mathbf{D}^{-1} \mathbf{b} \mathbf{b}^T \mathbf{D}^{-1} \end{bmatrix} = \begin{bmatrix} \frac{1}{\beta} & -\frac{1}{\beta} \boldsymbol{\gamma}^T \\ -\frac{1}{\beta} \boldsymbol{\gamma} & \mathbf{D}^{-1} + \frac{1}{\beta} \boldsymbol{\gamma} \boldsymbol{\gamma}^T \end{bmatrix} \quad (11)$$

where $a \in \mathbb{R}$, \mathbf{b} is a vector, \mathbf{D} is a symmetric nonsingular matrix, $\boldsymbol{\gamma} = \mathbf{D}^{-1} \mathbf{b}$ is a vector. For our case \mathbf{F}_{ZI} as in (8), $a = \frac{1-p_0(\boldsymbol{\theta})}{[\phi+(1-\phi)p_0(\boldsymbol{\theta})](1-\phi)}$, $\mathbf{b} = \frac{p_0(\boldsymbol{\theta})}{\phi+(1-\phi)p_0(\boldsymbol{\theta})} \cdot \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}}$, $\mathbf{D} = \mathbf{F}_{\text{ZI}\boldsymbol{\theta}} = (1-\phi)(\mathbf{F}_{\boldsymbol{\theta}} - \frac{\phi p_0(\boldsymbol{\theta})}{\phi+(1-\phi)p_0(\boldsymbol{\theta})} \cdot \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} \cdot \frac{\partial \log p_0(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}^T})$. Since $|\mathbf{F}_{\text{ZI}}| \neq 0$, then \mathbf{F}_{ZI} is positive definite and so is \mathbf{D} . On the other hand, $|\mathbf{F}_{\text{ZI}}| = |\mathbf{D}| \cdot (a - \mathbf{b}^T \mathbf{D}^{-1} \mathbf{b}) = |\mathbf{D}| \cdot \beta$ (see, for example, formula 14.21(b) in Seber (2008)), then $|\mathbf{F}_{\text{ZI}}| \neq 0$ implies $\beta \neq 0$ in our case.

To simplify the notations, we write in this proof only $p = p_0(\boldsymbol{\theta})$, $d = d_{\boldsymbol{\theta}} = \phi + (1-\phi)p_0(\boldsymbol{\theta})$, and $\mathbf{F} = \mathbf{F}_{\boldsymbol{\theta}}$. We first derive the formula of \mathbf{D}^{-1} . By the Sherman-Morrison-Woodbury formula (see for example, 15.3(a)(iii) in Seber (2008)), $(A+UV)^{-1} = A^{-1} - A^{-1}U(I+VA^{-1}U)^{-1}VA^{-1}$ if $|A| \neq 0$ and $|I+VA^{-1}U| \neq 0$. In our case, $A = \mathbf{F}$ satisfies $|A| = |\mathbf{F}| = |\mathbf{F}_{\boldsymbol{\theta}}| \neq 0$, $U = -\frac{\phi p}{d} \frac{\partial \log p}{\partial \boldsymbol{\theta}}$, $V = \frac{\partial \log p}{\partial \boldsymbol{\theta}^T}$. On the other hand, $|\mathbf{D}| \neq 0$ implies $\phi < 1$ and $|\mathbf{F} - \frac{\phi p}{d} \frac{\partial \log p}{\partial \boldsymbol{\theta}} \cdot \frac{\partial \log p}{\partial \boldsymbol{\theta}^T}| \neq 0$. Since $|\mathbf{F}| \neq 0$, then \mathbf{F} is positive definite and $\mathbf{F}^{-1/2}$ exists. According to formulae 4.33 in Seber (2008), $|I - \mathbf{a}\mathbf{a}^T| = 1 - \mathbf{a}^T \mathbf{a}$ for any vector \mathbf{a} . Then

$$\begin{aligned} 0 &\neq |\mathbf{F}^{-1/2}| \cdot \left| \mathbf{F} - \frac{\phi p}{d} \frac{\partial \log p}{\partial \boldsymbol{\theta}} \cdot \frac{\partial \log p}{\partial \boldsymbol{\theta}^T} \right| \cdot |\mathbf{F}^{-1/2}| \\ &= \left| I - \left(\sqrt{\frac{\phi p}{d}} \mathbf{F}^{-1/2} \frac{\partial \log p}{\partial \boldsymbol{\theta}} \right) \cdot \left(\sqrt{\frac{\phi p}{d}} \mathbf{F}^{-1/2} \frac{\partial \log p}{\partial \boldsymbol{\theta}} \right)^T \right| \\ &= 1 - \left(\sqrt{\frac{\phi p}{d}} \mathbf{F}^{-1/2} \frac{\partial \log p}{\partial \boldsymbol{\theta}} \right)^T \cdot \left(\sqrt{\frac{\phi p}{d}} \mathbf{F}^{-1/2} \frac{\partial \log p}{\partial \boldsymbol{\theta}} \right) \\ &= 1 - \frac{\phi p}{d} \frac{\partial \log p}{\partial \boldsymbol{\theta}^T} \mathbf{F}^{-1} \frac{\partial \log p}{\partial \boldsymbol{\theta}} \\ &= |I + VA^{-1}U| \end{aligned}$$

That is, $|I + VA^{-1}U| \neq 0$ in our case. By applying the Sherman-Morrison-Woodbury

formula,

$$\mathbf{D}^{-1} = \frac{1}{1-\phi} \left(\mathbf{F} - \frac{\phi p}{d} \cdot \frac{\partial \log p}{\partial \boldsymbol{\theta}} \cdot \frac{\partial \log p}{\partial \boldsymbol{\theta}^T} \right)^{-1} = \frac{1}{1-\phi} \left(\mathbf{F}^{-1} - \frac{\phi p}{d\delta} \mathbf{c} \mathbf{c}^T \right)$$

where $\delta = \delta_{\boldsymbol{\theta}} = 1 - \frac{\phi p}{d} \frac{\partial \log p}{\partial \boldsymbol{\theta}^T} \mathbf{F}^{-1} \frac{\partial \log p}{\partial \boldsymbol{\theta}} \neq 0$ and $\mathbf{c} = \mathbf{F}^{-1} \frac{\partial \log p}{\partial \boldsymbol{\theta}}$. It can be verified that $\boldsymbol{\gamma} = \mathbf{D}^{-1} \mathbf{b} = \frac{p}{d\delta(1-\phi)} \cdot \mathbf{c}$ and $\beta = a - \mathbf{b}^T \mathbf{D}^{-1} \mathbf{b} = \frac{1}{\phi(1-\phi)} \cdot \frac{d\delta - p}{d\delta} \neq 0$. Then the final expression of $\mathbf{F}_{\text{ZI}}^{-1}$ can be obtained from (11). \square

Proof of Theorem 2.3:

Given $f_{\text{ZI}}(y \mid \phi_{\text{ZI}}, \boldsymbol{\theta})$ and $\phi_{\text{ZA}} = \phi_{\text{ZI}} + (1 - \phi_{\text{ZI}})p_0(\boldsymbol{\theta})$, we have $1 - \phi_{\text{ZA}} = (1 - \phi_{\text{ZI}})[1 - p_0(\boldsymbol{\theta})]$ and $f_{\text{ZA}}(y \mid \phi_{\text{ZA}}, \boldsymbol{\theta}) = \phi_{\text{ZA}} \mathbf{1}_{\{y=0\}} + (1 - \phi_{\text{ZA}}) \frac{f_{\boldsymbol{\theta}}(y)}{1-p_0(\boldsymbol{\theta})} \mathbf{1}_{\{y \neq 0\}} = [\phi_{\text{ZI}} + (1 - \phi_{\text{ZI}})p_0(\boldsymbol{\theta})] \mathbf{1}_{\{y=0\}} + (1 - \phi_{\text{ZI}}) f_{\boldsymbol{\theta}}(y) \mathbf{1}_{\{y \neq 0\}} = f_{\text{ZI}}(y \mid \phi_{\text{ZI}}, \boldsymbol{\theta})$.

Given $f_{\text{ZA}}(y \mid \phi_{\text{ZA}}, \boldsymbol{\theta})$ and $\phi_{\text{ZI}} = [\phi_{\text{ZA}} - p_0(\boldsymbol{\theta})]/[1 - p_0(\boldsymbol{\theta})]$, if $\phi_{\text{ZA}} \geq p_0(\boldsymbol{\theta})$, then $\phi_{\text{ZI}} \in [0, 1]$, $\phi_{\text{ZI}} + (1 - \phi_{\text{ZI}})p_0(\boldsymbol{\theta}) = \phi_{\text{ZA}}$, $1 - \phi_{\text{ZI}} = \frac{1 - \phi_{\text{ZA}}}{1 - p_0(\boldsymbol{\theta})}$, and $f_{\text{ZI}}(y \mid \phi_{\text{ZI}}, \boldsymbol{\theta}) = \phi_{\text{ZA}} \mathbf{1}_{\{y=0\}} + \frac{1 - \phi_{\text{ZA}}}{1 - p_0(\boldsymbol{\theta})} f_{\boldsymbol{\theta}}(y) \mathbf{1}_{\{y \neq 0\}} = \phi_{\text{ZA}} \mathbf{1}_{\{y=0\}} + (1 - \phi_{\text{ZA}}) f_{\text{tr}}(y \mid \boldsymbol{\theta}) \mathbf{1}_{\{y \neq 0\}} = f_{\text{ZA}}(y \mid \phi_{\text{ZA}}, \boldsymbol{\theta})$. \square

References

- Agresti, A., 2013: *Categorical Data Analysis*. Wiley, 3rd ed.
- Albasi, A., 2018: *mazeinda: Monotonic Association on Zero-Inflated Data*. R package version 0.0.1.
- Aldirawi, H. and J. Yang, 2022: Modeling sparse data using mle with applications to microbiome data. *Journal of Statistical Theory and Practice*, **16(1)**, Article 13.
- Aldirawi, H., J. Yang, and A. A. Metwally, 2019: Identifying appropriate probabilistic models for sparse discrete omics data. In *2019 IEEE EMBS International Conference on Biomedical & Health Informatics (BHI)*, IEEE, pp. 1–4.
- Balderama, E. and T. Trippe, 2017: *hurdlr: Zero-Inflated and Hurdle Modelling Using Bayesian Inference*. R package version 0.1.
- Boucher, J.-P., M. Denuit, and M. Guillen, 2009: Number of accidents or number of claims? an approach with zero-inflated poisson models for panel data. *Journal of Risk and Insurance*, **76(4)**, 821–846.
- Chen, P., Q. Liu, and F. Sun, 2018: Bicycle parking security and built environments. *Transportation Research Part D: Transport and Environment*, **62**, 169–178.
- Chen, Z. and D. B. Dunson, 2003: Random effects selection in linear mixed models. *Biometrics*, **59(4)**, 762–769.
- Cho, H., C. Liu, J. Park, and D. Wu, 2018: *binzb: Bivariate Zero-Inflated Negative Binomial Model Estimator*. R package version 1.0.4.

- Croissant, Y., F. Carlevaro, and S. Hoareau, 2021: *mhurdle: Multiple Hurdle Tobit Models*. R package version 1.3.0.
- Deb, P. and P. K. Trivedi, 1997: Demand for medical care by the elderly: a finite mixture approach. *Journal of Applied Econometrics*, **12(3)**, 313–336.
- Dimitrova, D. S., V. K. Kaishev, and S. Tan, 2020: Computing the kolmogorov-smirnov distribution when the underlying cdf is purely discrete, mixed, or continuous. *Journal of Statistical Software*, **95(1)**, 1–42.
- Golub, G. and C. V. Loan, 2013: *Matrix Computations*. Johns Hopkins University Press, 4th ed.
- Hogg, R. V., J. W. McKean, and A. T. Craig, 2019: *Introduction to Mathematical Statistics*. Pearson Education, eighth ed.
- Hu, T. and Y. Zhou, 2018: *ZIBBSeqDiscovery: Zero-Inflated Beta-Binomial Modeling of Microbiome Count Data*. R package version 1.0.
- Huang, D., H. Hu, and Y. Li, 2019: Zero-inflated exponential distribution of casualty rate in ship collision. *Journal of Shanghai Jiaotong University (Science)*, **24(6)**, 739–744.
- Jochmann, M., 2017: *zic: Bayesian Inference for Zero-Inflated Count Models*. R package version 0.9.1.
- Lehmann, E. L. and J. P. Romano, 2005: *Testing Statistical Hypotheses*. Springer, 3rd ed.
- Lilliefors, H. W., 1967: On the kolmogorov-smirnov test for normality with mean and variance unknown. *Journal of the American statistical Association*, **62(318)**, 399–402.
- , 1969: On the kolmogorov-smirnov test for the exponential distribution with mean unknown. *Journal of the American Statistical Association*, **64(325)**, 387–389.
- Majo, M. C. and A. van Soest, 2011: The fixed-effects zero-inflated poisson model with an application to health care utilization. <https://pure.uvt.nl/ws/portalfiles/portal/1341748/2011-083.pdf>.
- Massey Jr, F. J., 1951: The kolmogorov-smirnov test for goodness of fit. *Journal of the American statistical Association*, **46(253)**, 68–78.
- McDavid, A., R. Gottardo, N. Simon, and M. Drton, 2019: Graphical models for zero-inflated single cell gene expression. *The Annals of Applied Statistics*, **13(2)**, 848–873.
- Metwally, A. A., H. Aldirawi, and J. Yang, 2018: A review on probabilistic models used in microbiome studies. *Communications in Information and Systems*, **18(3)**, 173–191.
- Metwally, A. A., Y. Dai, P. W. Finn, and D. L. Perkins, 2019: *MetaLonDA: a flexible R package for identifying time intervals of differentially abundant features in metagenomic longitudinal studies*. R package version 1.1.8.

- Mullahy, J., 1986: Specification and testing of some modified count data models. *Journal of econometrics*, **33(3)**, 341–365.
- Peng, X., G. Li, Z. Liu, and H. Chen, 2017: *ZIBseq: Differential Abundance Analysis for Metagenomic Data via Zero-Inflated Beta Regression*. R package version 1.2.
- Reyna-Hurtado, R., C. A. Chapman, S. Calme, and E. J. Pedersen, 2012: Searching in heterogeneous and limiting environments: foraging strategies of white-lipped peccaries (*tayassu pecari*). *Journal of Mammalogy*, **93(1)**, 124–133.
- Seber, G., 2008: *A Matrix Handbook for Statisticians*. Wiley.
- Teerapabolarn, K., 2014: An improved poisson to approximate the negative binomial distribution. *International Journal of Pure and Applied Mathematics*, **91(3)**, 369–373.
- Vaart, A. W. v. d., 2000: *Asymptotic Statistics*. Cambridge university press.
- Wang, L., H. Aldirawi, and J. Yang, 2019: *iZID: Identify Zero-Inflated Distributions*. R package version 0.0.1.
- , 2020: Identifying zero-inflated distributions with a new r package izid. *Communications in Information and Systems*, **20(1)**, 23–44.
- Waudby-Smith, I. and P. Li, 2017: *rbtt: Alternative Bootstrap-Based t-Test Aiming to Reduce Type-I Error for Non-Negative, Zero-Inflated Data*. R package version 0.1.0.
- Xia, Y., J. Sun, and D.-G. Chen, 2018: *Statistical Analysis of Microbiome Data with R*. Springer.
- Xu, L., A. D. Paterson, W. Turpin, and W. Xu, 2015: Assessment and selection of competing models for zero-inflated microbiome data. *PLoS ONE*, **10(7)**.
- Xu, Z. J. and Y. Liu, 2018: *ziphsmm: Zero-Inflated Poisson Hidden (Semi-)Markov Models*. R package version 2.0.6.
- Yang, M., G. Zamba, and J. Cavanaugh, 2018: *ZIM: Zero-Inflated Models (ZIM) for Count Time Series with Excess Zeros*. R package version 1.1.0.
- Zeileis, A., C. Kleiber, and S. Jackman, 2008: Regression models for count data in r. *Journal of Statistical Software*, **27(8)**, 1–25.
- Zhang, X., B. Guo, and N. Yi, 2020: Zero-inflated gaussian mixed models for analyzing longitudinal microbiome data. *PLoS ONE*, **15(11)**, e0242073.
- Zhou, X.-H. and W. Tu, 1999: Comparison of several independent population means when their samples contain log-normal and possibly zero observations. *Biometrics*, **55(2)**, 645–651.