

Riemann-Roch for $\overline{\text{Spec } \mathbb{Z}}$

Alain Connes and Caterina Consani

Abstract

We prove a Riemann-Roch formula for Arakelov divisors on $\overline{\text{Spec } \mathbb{Z}}$ equating the integer valued Euler characteristic with a simple modification of the traditional expression (*i.e.* the degree of the divisor plus $\log 2$). The integer valued topological side involves besides the ceiling function, the division by $\log 3$. The notion of dimension and the definition of the cohomologies are based on the universal arithmetic over the sphere spectrum which we developed using Segal's Gamma rings. This new perspective allows us to parallel Weil's adelic proof of the Riemann-Roch formula for function fields including the use of Pontryagin duality.

Keywords: Riemann-Roch, Arakelov compactification, Adeles, Segal's Gamma-ring.

MSC Classification: 14C40; 14G40; 14H05; 11R56; 13F35; 18N60; 19D55.

1 Introduction

The main result of this paper is the following arithmetic Riemann-Roch theorem for $\overline{\text{Spec } \mathbb{Z}}$ (see Theorem 5.3)

Theorem 1 *Let D be an Arakelov divisor on $\overline{\text{Spec } \mathbb{Z}}$. Then*

$$\dim_{\mathbb{S}[\pm 1]} H^0(D) - \dim_{\mathbb{S}[\pm 1]} H^1(D) = \left[\frac{\deg D + \log 2}{\log 3} \right]' - \mathbf{1}_L. \quad (1)$$

On the (topological) right-hand side of (1), $[x]'$ denotes the odd function on \mathbb{R} that agrees with the ceiling function on positive reals. The set $L \subset \mathbb{R}$ is the union, for $k \geq 0$, of the intervals $(\log \frac{3^k}{2}, \log \frac{3^k+1}{2})$: L has finite Lebesgue measure. The symbol $\mathbf{1}_L$ denotes the (degree) indicator function: this is zero unless $\deg D \in L$ in which case it is 1. The numbers $\dim_{\mathbb{S}[\pm 1]} H^i(D)$ on the (cohomological) left-hand side are integers providing the dimension of the cohomology of D , as a module over the spherical extension $\mathbb{S}[\pm 1]$ of the absolute base

\mathbb{S} (see section 2). Before to introduce the definitions of the cohomology and their dimension, we comment on the new geometric perspective introduced by Theorem 1 in comparison with the asymptotic Riemann-Roch formula of S. Lang [8] (foreseen by A. Weil in [10]) comparing $\log \#H^0(D)$ and $\deg D + \log 2$, when $\deg D \rightarrow \infty$. The first novelty is that unlike $\log \#H^0(D)$, $\dim_{\mathbb{S}[\pm 1]} H^i(D)$ are non-negative integers. Secondly, the cohomological side of (1) implements explicitly (the dimension of) $H^1(D)$, thus it goes well beyond the arithmetic Riemann-Roch formula for \mathbb{Q} , where $H^1(D)$ is either undefined or set to equate $H^0(K - D)$, for an expected Serre duality. All the more: (1) is an exact formula and not an asymptotic statement. All these advancements derive from the new understanding of $\overline{\text{Spec } \mathbb{Z}}$ as a curve over the absolute ring $\mathbb{S}[\pm 1]$. The third relevant fact is that on the topological side of the Riemann-Roch formula the traditional sum $\deg D + \log 2$ is now divided by $\log 3 \sim 1.09861$, and there also appears an exceptional set $L \subset \mathbb{R}$ of finite Lebesgue measure $|L| = \log \prod (1 + 3^{-n}) \sim 1.14099$. The Euler characteristic of D is a non-decreasing function of $\deg D$, while the topological side in (1) drops by 1 if $\deg D \in L$. Except for the set L , this topological side is invariant under Serre's duality (Theorem 2) which replaces D with $K - D$ ($\deg K = -2 \log 2$), since the function $[x]'$ is odd.

In [1, 4] we have developed the fundamentals of a geometric theory for Segal's Γ -rings : this means that the initial ring \mathbb{Z} of the category of rings is replaced by the sphere spectrum \mathbb{S} . In particular, we have defined the structure sheaf of the one-point compactification $\overline{\text{Spec } \mathbb{Z}}$ of the algebraic spectrum of the integers by suitably extending, at the archimedean place, the structure sheaf of $\text{Spec } \mathbb{Z}$ as a subsheaf of the constant sheaf \mathbb{Q} . The global sections of this sheaf determine the \mathbb{S} -algebra $\mathbb{S}[\pm 1]$. In [2] we extended the notion of homology for simplicial complexes, when group coefficients are replaced by \mathbb{S} -modules. In [3] we implemented homological algebra in this context by showing a generalization of the Dold-Kan correspondence which associates a Γ -space to a short complex of \mathbb{S} -modules. In this paper we apply all these results to adapt, for $\overline{\text{Spec } \mathbb{Z}}$, the adelic proof of the Riemann-Roch formula for function fields given by Weil in [11].

Next, we introduce the cohomologies and their dimensions. An Arakelov divisor $D = \sum_j a_j \{p_j\} + a\{\infty\}$ on $\overline{\text{Spec } \mathbb{Z}}$ determines a compact subset $\mathcal{O}(D) \subset \mathbb{A}_{\mathbb{Q}}$ of the adeles of \mathbb{Q} . This is the product, indexed by the places of \mathbb{Q} , of the abelian group $\mathbb{Z}_p \subset \mathbb{Q}_p$ for each finite $p \notin \{p_j\}_j$, the abelian groups $p_j^{-a_j} \mathbb{Z}_{p_j}$, and the interval $[-e^a, e^a] \subset \mathbb{R}$. This last component is implemented by a sub-module of the Eilenberg-MacLane $\mathbb{S}[\pm 1]$ -module $H\mathbb{R}$ which functorially encodes the additive structure of the group \mathbb{R} . The morphisms of abelian groups used by Weil in the development of the adelic geometry of function fields are meaningful in this absolute set-up since they are viewed as morphisms of $\mathbb{S}[\pm 1]$ -modules. In particular, we shall focus on the morphism

$$\psi : \mathbb{Q} \times \mathcal{O}(D) \rightarrow \mathbb{A}_{\mathbb{Q}} \quad \psi(q, a) = q + a \quad \forall q \in \mathbb{Q}, a \in \mathcal{O}(D), \quad (2)$$

where \mathbb{Q} is embedded diagonally in the adèles. The Γ -space $\mathbf{H}(D)$ associated by the Dold-Kan correspondence to (2) (ψ is viewed here as a short complex of $\mathbb{S}[\pm 1]$ -modules: see Appendix A) provides the absolute incarnation of the Riemann-Roch problem for the divisor D . In fact, by implementing the multiplicative action of \mathbb{Q}^\times on $\mathbb{A}_{\mathbb{Q}}$, one is reduced to consider only the case $D = a\{\infty\}$, for which $\ker(\psi)$ is the $\mathbb{S}[\pm 1]$ -module $H^0(D) = \|H\mathbb{Z}\|_{e^a}$ that associates to a finite pointed set X the \mathbb{Z} -valued divisors $\sum_i n_i x_i$, $x_i \in X$, which fulfill the condition $\sum_i |n_i| \leq e^a$ ($|\cdot|$ = euclidean absolute value). By definition, $H^0(D)$ is a covariant functor $\Gamma^o \rightarrow \mathfrak{Sets}_*$ from the small category Γ^o (a skeleton of the category of finite pointed sets) to pointed sets that keeps track of the partially defined addition in $I = \|H\mathbb{Z}\|_{e^a}(1_+) = [-e^a, e^a] \cap \mathbb{Z}$, so that for $n_i \in I$, $\sum n_i$ is meaningful provided $\sum_i |n_i| \leq e^a$. Its dimension $\dim_{\mathbb{S}[\pm 1]} H^0(D)$ is defined to be the smallest number of generators of I , where by definition, a subset $F \subset I$ generates if and only if for every $m \in I$ there exist coefficients $\alpha(f) \in \{-1, 0, 1\} = \mathbb{S}[\pm 1](1_+)$ such that $m = \sum \alpha(f)f$, $f \in F$, and $\sum |\alpha(f)f| \leq e^a$ (see section 4). The $\mathbb{S}[\pm 1]$ -module $H^0(D)$ only depends upon the integer part $n = \lfloor e^a \rfloor$ of e^a and Proposition 4.3 determines its dimension to be¹

$$\dim_{\mathbb{S}[\pm 1]}(\|H\mathbb{Z}\|_n) = \left\lceil \frac{\log(2n+1)}{\log 3} \right\rceil. \quad (3)$$

The cokernel of ψ , that is $H^1(D)$, is the couple of the $\mathbb{S}[\pm 1]$ -module $H\mathbb{A}_{\mathbb{Q}}$ and a suitable tolerance relation \mathcal{R} on it (Proposition 2.2). This relation is defined by pairs whose difference belongs to the image of ψ . More specifically, $H^1(D)$ defines an object of the category $\Gamma\mathcal{T}_*$ of tolerance \mathbb{S} -modules (see section 3). Its dimension $\dim_{\mathbb{S}[\pm 1]} H^1(D)$ is defined to be the smallest number of generators in a generating set, where a subset $F \subset H\mathbb{A}_{\mathbb{Q}}(1_+) = \mathbb{A}_{\mathbb{Q}}$ generates if and only if for every $x \in H\mathbb{A}_{\mathbb{Q}}(1_+)$ there exists coefficients $\alpha(f) \in \{-1, 0, 1\}$ such that the pair $(x, \sum \alpha(f)f)$ belongs to the relation \mathcal{R} (see section 5). For a divisor $D = a\{\infty\}$, the dimension of $H^1(D)$ is the same as the dimension of the pair $(H(\mathbb{R}/\mathbb{Z}), \mathcal{R})$, where the relation \mathcal{R} on $H(\mathbb{R}/\mathbb{Z})(1_+) = \mathbb{R}/\mathbb{Z}$ is given by

$$(x, y) \in \mathcal{R} \iff d(x, y) \leq e^a, \quad (4)$$

for d the translation invariant Riemannian metric of length 1 on \mathbb{R}/\mathbb{Z} . In Proposition 5.1 we show that this dimension is

$$\dim_{\mathbb{S}[\pm 1]}(H(\mathbb{R}/\mathbb{Z}), \mathcal{R}) = \left\lceil \frac{-a - \log 2}{\log 3} \right\rceil. \quad (5)$$

Theorem 1 follows from (3) when $\deg D \geq -\log 2$, since in this case $\dim_{\mathbb{S}[\pm 1]} H^1(D) = 0$. The case $\deg D < -\log 2$ is proved using (5).

The symmetry of the term $\lceil \frac{\deg D + \log 2}{\log 3} \rceil'$ under the replacement of D by the divisor $K - D$, with $K = -2\{2\}$, is the numerical evidence of a geometric

¹For the values $n = 4, 13, 40, \dots, (3^k - 1)/2$, every element of I is uniquely written in terms of the generating set $F = \{3^i \mid 0 \leq i < k\}$ (Proposition 4.3 (iii)).

duality holding on the curve $\overline{\text{Spec } \mathbb{Z}}$ over $\mathbb{S}[\pm 1]$ (see section 6) which is precisely stated by the following (see Theorem 6.3)

Theorem 2 *Let D be an Arakelov divisor on $\overline{\text{Spec } \mathbb{Z}}$ and $K = -2\{2\}$, with $\deg K = -2 \log 2$. Then there is an isomorphism of $\mathbb{S}[\pm 1]$ -modules*

$$H^0(K - D) \simeq \underline{\text{Hom}}_{\Gamma \mathcal{T}_*}(H^1(D), U(1)_{\frac{1}{4}}), \quad (6)$$

where $U(1)_{\frac{1}{4}}$ coincides with $H^1(K)$, playing the role of the dualizing module in Pontryagin duality.

The isomorphism $H^0(K - D) \simeq \underline{\text{Hom}}_{\Gamma \mathcal{T}_*}(H^1(D), H^1(K))$ is the analogue of Grothendieck's version of Serre's duality [6] (12.15.1955). The duality (6) derives from Pontryagin's duality holding for tolerant $\mathbb{S}[\pm 1]$ -modules (Proposition 6.2), in analogy with the corresponding result proven by Weil for curves over fields of positive characteristic.

2 The cohomology $H^\bullet(D)$

We recall from [3] the construction of the Γ -space $\mathbf{H}(D)$ naturally associated to an Arakelov divisor D on $\overline{\text{Spec } \mathbb{Z}}$. This space is the absolute homological incarnation of the Riemann-Roch problem for the divisor D . We formulate this construction in terms of the adeles of \mathbb{Q} .

Let $D = \sum_j a_j \{p_j\} + a\{\infty\}$ be an Arakelov divisor on $\overline{\text{Spec } \mathbb{Z}}$. This is a formal finite sum with $a_j \in \mathbb{Z}$ and $a \in \mathbb{R}$, where p_j are rational primes in \mathbb{Z} and the symbol ∞ stands for the restriction $|\cdot| : \mathbb{Q} \rightarrow \mathbb{R}$ of the euclidean absolute value $|\cdot|_\infty$. Let $\Sigma_{\mathbb{Q}}$ denote the full set of places ν of \mathbb{Q} . To D is naturally associated the following idele $\exp(D)$ of \mathbb{Q}

$$\exp(D)_\nu := \begin{cases} p_j^{-a_j} & \text{if } \nu = \nu_j \\ 1 & \forall \nu \neq \nu_j, \forall j \\ e^a & \text{if } \nu = \infty. \end{cases}$$

The equality $\exp(D)(\nu) = |\exp(D)_\nu|_\nu$ defines a map $\exp(D) : \Sigma_{\mathbb{Q}} \rightarrow \mathbb{R}_+^*$ such that $\exp(D)(\nu) \in \text{mod}(\mathbb{Q}_\nu)$, $\forall \nu \in \Sigma_{\mathbb{Q}}$, and $\exp(D)(\nu) = 1$, for $\nu \neq \infty, \nu_j, \forall j$. To the divisor D corresponds the compact subset of the adeles of \mathbb{Q}

$$\mathcal{O}(D) := \{(a_\nu) \in \mathbb{A}_{\mathbb{Q}} \mid |a_\nu|_\nu \leq \exp(D)(\nu), \forall \nu \in \Sigma_{\mathbb{Q}}\} \subset \mathbb{A}_{\mathbb{Q}}.$$

By definition $\mathcal{O}(D) = \mathcal{O}(D)_f \times \mathcal{O}(D)_\infty = \prod_\nu \mathcal{O}(D)_\nu$ with

$$\mathcal{O}(D)_\nu = \begin{cases} p_j^{-a_j} \mathbb{Z}_{p_j} & \text{if } \nu = \nu_j \\ \mathbb{Z}_\nu & \forall \nu \neq \nu_j, \forall j, \nu < \infty \\ [-e^a, e^a] & \text{if } \nu = \infty. \end{cases}$$

In particular, the product $\mathcal{O}(D)_f = \prod_{\nu \neq \infty} \mathcal{O}(D)_\nu$ is a compact abelian group. The archimedean component on the other hand, *i.e.* the real interval $[-e^a, e^a]$, is not, and for this reason we implement the functorial viewpoint to suitably understand $\mathcal{O}(D)$. Indeed, to $\mathcal{O}(D)_\infty$ we associate the covariant functor

$$\|H\mathbb{R}\|_{e^a} : \Gamma^\circ \longrightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_* \quad \|H\mathbb{R}\|_{e^a}(F) := \left\{ \phi \in H\mathbb{R}(F) \mid \sum_{F \setminus \{*\}} |\phi(x)| \leq e^a \right\}$$

from the opposite of the Segal category (see *e.g.* [5] Chpt. 2 and [1]) to pointed sets, which maps a finite pointed set F to the pointed set of \mathbb{R} -valued divisors on F vanishing on the base point and whose total mass $\sum_{F \setminus \{*\}} |\phi(x)|$ is bounded by e^a . Covariant functors $\Gamma^\circ \longrightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$ and their natural transformations determine the category $\Gamma\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$ of Γ -sets (aka \mathbb{S} -modules). In particular, the Eilenberg-MacLane functor H encodes an abelian group A as the covariant functor $HA : \Gamma^\circ \longrightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$ that associates to F the pointed set of A -valued divisors on F vanishing on the base point. Functoriality holds by taking sums on the inverse images of a point. The functor HA encodes the addition on $A = HA(1_+)$. In general, let $\sigma \in \text{Hom}_{\Gamma^\circ}(k_+, 1_+)$ with $\sigma(\ell) = 1 \ \forall \ell \neq *$ and

$$\delta(j, k) \in \text{Hom}_{\Gamma^\circ}(k_+, 1_+), \quad \delta(j, k)(\ell) := \begin{cases} 1 & \text{if } \ell = j \\ * & \text{if } \ell \neq j. \end{cases} \quad (7)$$

Given an \mathbb{S} -module \mathcal{F} and elements $x, x_j \in \mathcal{F}(1_+)$, $j = 1, \dots, k$, one writes

$$x = \sum_j x_j \iff \exists z \in \mathcal{F}(k_+) \text{ s.t. } \mathcal{F}(\sigma)(z) = x, \mathcal{F}(\delta(j, k))(z) = x_j, \forall j. \quad (8)$$

For $A = \mathbb{R}$ the key point is that this general construction respects the bound on the total mass while encoding the addition. One easily sees that $\|H\mathbb{R}\|_{e^a}$ is an $\mathbb{S}[\pm 1]$ -module (an “ $(\mathbb{S}[\pm 1] \wedge -)$ -algebra” as in [5] 2.1.5), where $\mathbb{S}[\pm 1] : \Gamma^\circ \longrightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$, $\mathbb{S}[\pm 1](F) := \{-1, 0, 1\} \wedge F$ is the spherical monoid algebra of the multiplicative monoid $\{\pm 1\}$. In particular, the inclusion functor $\|H\mathbb{R}\|_{e^a} \longrightarrow H\mathbb{R}$ determines $\|H\mathbb{R}\|_{e^a}$ as a sub $\mathbb{S}[\pm 1]$ -module of $H\mathbb{R}$, where $\mathbb{S} : \Gamma^\circ \longrightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$ is the identity functor.

Given two \mathbb{S} -modules \mathcal{F}_j , $j = 1, 2$, one defines their product as the functor

$$\mathcal{F}_1 \times \mathcal{F}_2 : \Gamma^\circ \longrightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*, \quad (\mathcal{F}_1 \times \mathcal{F}_2)(F) = \mathcal{F}_1(F) \times \mathcal{F}_2(F)$$

with the base point of $\mathcal{F}_1(F) \times \mathcal{F}_2(F)$ taken to be $(*, *)$. For abelian groups A, B one has a canonical isomorphism $H(A \times B) \simeq HA \times HB$. In particular, the morphism of addition in adèles $\alpha : \mathbb{Q} \times \mathbb{A}_{\mathbb{Q}} \rightarrow \mathbb{A}_{\mathbb{Q}}$, $\alpha(q, a) = q + a$ (using the diagonal embedding of \mathbb{Q} in $\mathbb{A}_{\mathbb{Q}}$) determines a morphism of $\mathbb{S}[\pm 1]$ -modules $H\alpha : H\mathbb{Q} \times H\mathbb{A}_{\mathbb{Q}} \longrightarrow H\mathbb{A}_{\mathbb{Q}}$. Next proposition shows how this functorial construction is well combined with the adelic formalism.

Proposition 2.1 (i) *The $\mathbb{S}[\pm 1]$ -module*

$$H\mathcal{O}(D) := H(\mathcal{O}(D)_f) \times \|H\mathbb{R}\|_{e^a}$$

determines, canonically, a sub $\mathbb{S}[\pm 1]$ -module $\iota : H\mathcal{O}(D) \rightarrow H\mathbb{A}_{\mathbb{Q}}$.

(ii) *The restriction of $H\alpha$ to $H\mathbb{Q} \times H\mathcal{O}(D)$ defines a morphism of $\mathbb{S}[\pm 1]$ -modules*

$$\psi : H\mathbb{Q} \times H\mathcal{O}(D) \rightarrow H\mathbb{A}_{\mathbb{Q}}. \quad (9)$$

Proof (i) With $\mathbb{A}_{\mathbb{Q}} = \mathbb{A}_f \times \mathbb{R}$, let $\iota_f : \mathcal{O}(D)_f \rightarrow \mathbb{A}_f$ be the inclusion of abelian groups. Then $\iota : H\mathcal{O}(D) \rightarrow H\mathbb{A}_{\mathbb{Q}}$ is the product of $H\iota_f$ and the inclusion $\|H\mathbb{R}\|_{e^a} \rightarrow H\mathbb{R}$.

(ii) By composition one obtains $\psi = H\alpha \circ (\text{Hid}_{\mathbb{Q}} \times \iota)$. \square

The morphism ψ in (9) is obtained by restricting $H\alpha$ to the (adelic) divisor D , where α is a morphism of abelian groups. The theory developed in [3] associates to ψ a Γ -space $\mathbf{H}(D)$, by implementing the Dold-Kan correspondence in the special case of a short complex of abelian groups and its restriction to a sub- \mathbb{S} -module. The Γ -space $\mathbf{H}(D)$ supplies the absolute cohomology of the divisor D . We recall this construction in Appendix A. Moreover, we review how the generalized homotopy $\pi_*^{\mathcal{T}}(\mathbf{H}(D))$, where \mathcal{T} is the category of tolerance relations defined in 3.1, is meaningful in this context supplying, with $\pi_1^{\mathcal{T}}(\mathbf{H}(D))$ and $\pi_0^{\mathcal{T}}(\mathbf{H}(D))$, the (tolerant) \mathbb{S} -modules descriptions of *resp.* the kernel and the cokernel of ψ . In other words, one obtains the following result

Proposition 2.2 (i) *The kernel of ψ is the $\mathbb{S}[\pm 1]$ -module:*

$$H^0(D) = (\text{Hid}_{\mathbb{Q}} \times \iota)^{-1}(H \ker(\alpha)) \simeq \iota^{-1}H\mathbb{Q} = \|H^0(\text{Spec } \mathbb{Z}, \mathcal{O}(D)_f)\|_{e^a}. \quad (10)$$

(ii) *The cokernel of ψ is the $\mathbb{S}[\pm 1]$ -module $H^1(D) = (H\mathbb{A}_{\mathbb{Q}}, \mathcal{R})$ endowed with the relations*

$$\mathcal{R}_k := \{(x, y) \in H\mathbb{A}_{\mathbb{Q}}(k_+) \times H\mathbb{A}_{\mathbb{Q}}(k_+) \mid x - y \in \text{Image } \psi(k_+)\}. \quad (11)$$

(iii) *Both $H^0(D)$ and $H^1(D)$ only depend, up to isomorphism, on the linear equivalence class of the divisor D .*

Proof We refer to Appendix A.2. \square

3 Tolerance \mathbb{S} -modules

The construction of the category $\Gamma\mathcal{S}_*$ of Γ -spaces (see Appendix A) can be broadly generalized by considering in place of the category \mathcal{S}_* of simplicial pointed sets any pointed category \mathcal{C} with initial and final object $*$. In this way, one obtains a category $\Gamma\mathcal{C}$ of pointed covariant functor $\Gamma^{\circ} \rightarrow \mathcal{C}$. We shall apply this formal construction

to the category of tolerance relations and introduce the notion of tolerant $\mathbb{S}[\pm 1]$ -modules which plays a central role in the development of the absolute Riemann-Roch problem. We start with the following general fact

Lemma 3.1 *Let \mathcal{C} be a pointed category with initial and final object $*$. Then $\Gamma\mathcal{C}$ is naturally enriched in \mathbb{S} -modules. More precisely, the following formula endows the internal $\underline{\text{Hom}}_{\Gamma\mathcal{C}}(A, B)$ with a structure of \mathbb{S} -module defined by*

$$\underline{\text{Hom}}_{\Gamma\mathcal{C}}(A, B)(k_+) := \text{Hom}_{\Gamma\mathcal{C}}(A, B(k_+ \wedge -)) \quad k \in \mathbb{N}. \quad (12)$$

Proof Let $\phi \in \text{Hom}_{\Gamma^\circ}(k_+, \ell_+)$. For every object F of Γ° the morphism $\phi \wedge \text{id} \in \text{Hom}_{\Gamma^\circ}(k_+ \wedge F, \ell_+ \wedge F)$ gives, by functoriality of $B : \Gamma^\circ \rightarrow \mathcal{C}$, a morphism $B(\phi \wedge \text{id}) \in \text{Hom}_{\mathcal{C}}(B(k_+ \wedge F), B(\ell_+ \wedge F))$. These morphisms define a natural transformation of functors $B(\phi \wedge -) \in \text{Hom}_{\Gamma\mathcal{C}}(B(k_+ \wedge -), B(\ell_+ \wedge -))$, and one obtains the functoriality on the right hand side of (12) using the left composition

$$\psi \in \text{Hom}_{\Gamma\mathcal{C}}(A, B(k_+ \wedge -)) \mapsto B(\phi \wedge -) \circ \psi \in \text{Hom}_{\Gamma\mathcal{C}}(A, B(\ell_+ \wedge -)).$$

□

3.1 The category $\Gamma\mathcal{T}_*$

A tolerance relation \mathcal{R} on a set X is a reflexive and symmetric relation on X . Equivalently, \mathcal{R} is a subset $\mathcal{R} \subset X \times X$ which is symmetric and containing the diagonal. We shall denote by \mathcal{T} the category of tolerance relations (X, \mathcal{R}) . Morphisms in \mathcal{T} are defined by

$$\text{Hom}_{\mathcal{T}}((X, \mathcal{R}), (X', \mathcal{R}')) := \{\phi : X \rightarrow X', \phi(\mathcal{R}) \subset \mathcal{R}'\}.$$

We denote \mathcal{T}_* the pointed category under the object $\{*\}$ endowed with the trivial relation. One has the following

Definition 3.2 *A tolerant \mathbb{S} -module is a pointed covariant functor $\Gamma^\circ \rightarrow \mathcal{T}_*$. We denote by $\Gamma\mathcal{T}_*$ the category of tolerant \mathbb{S} -modules.*

Next statement is an easy but useful fact

Proposition 3.3 (i) *The functor $\mathfrak{S}\text{ets} \rightarrow \mathcal{T}$ which endows a set with the diagonal relation, embeds the category of sets as a full subcategory of \mathcal{T} , and consequently the category of \mathbb{S} -modules as a full subcategory of the category $\Gamma\mathcal{T}_*$.*

(ii) *The forgetful functor is the right adjoint of the inclusion in (i).*

3.2 The tolerant \mathbb{S} -module $(A, d)_\lambda$

A relevant example of tolerant \mathbb{S} -module is given by the following construction. Let A be an additive abelian group. A translation invariant metric d on A is a metric on

A that satisfies $d(x, y) = d(x - y, 0)$, so that the triangle inequality can be read as $d(x + y, 0) \leq d(x, 0) + d(y, 0) \forall x, y \in A$. This fact implies that the inequality

$$d\left(\sum_I x_i, \sum_I y_i\right) \leq \sum_I d(x_i, y_i) \quad (13)$$

holds for any finite index set I and maps $x, y : I \rightarrow A$.

Proposition 3.4 *Let (A, d) be an abelian group endowed with a translation invariant metric d and let $\lambda \in \mathbb{R}_{>0}$. The following relations turn the Eilenberg-MacLane \mathbb{S} -module HA into a tolerant $\mathbb{S}[\pm 1]$ -module $(A, d)_\lambda : \Gamma^\circ \rightarrow \mathcal{T}_*$:*

$$\mathcal{R}_k = \left\{ ((x_i), (y_i)) \in A^{\times k} \times A^{\times k} \mid \sum_{I=k_+} d(x_i, y_i) \leq \lambda \right\} \quad \forall k_+, k \in \mathbb{N}. \quad (14)$$

Proof For any $\phi \in \text{Hom}_{\Gamma^\circ}(k_+, \ell_+)$, the map $HA(\phi) : HA(k_+) \rightarrow HA(\ell_+)$ fulfills $HA(\phi)^{\times 2}(\mathcal{R}_k) \subset \mathcal{R}_\ell$. Indeed, this follows from (13) applied to the finite sets $I_j = \{i \in k_+ \mid \phi(i) = j\}$ which label pairs of elements of A . \square

Let $U(1)$ be the abelian group \mathbb{R}/\mathbb{Z} endowed with the canonical metric d of length 1. We shall denote by $U(1)_\lambda$ the tolerant \mathbb{S} -module $(U(1), d)_\lambda$, ($\lambda \in \mathbb{R}_{>0}$).

3.3 The dimension of a tolerant $\mathbb{S}[\pm 1]$ -module

In this part we introduce a notion of dimension for a tolerant $\mathbb{S}[\pm 1]$ -module that naturally generalizes, in the absolute context, the definition of dimension of a vector space.

Definition 3.5 *Let (E, \mathcal{R}) be a tolerant $\mathbb{S}[\pm 1]$ -module. A subset $F \subset E(1_+)$ generates $E(1_+)$ if the following two conditions hold*

1. For $x, y \in F$, with $x \neq y \implies (x, y) \notin \mathcal{R}$
2. For every $x \in E(1_+)$ there exists $\alpha_j \in \{-1, 0, 1\}$, $j \in F$ and $y \in E(1_+)$ such that $y = \sum_F \alpha_j j \in E(1_+)$ in the sense of (8), and $(x, y) \in \mathcal{R}$.

The dimension $\dim_{\mathbb{S}[\pm 1]}(E, \mathcal{R})$ is defined as the minimal cardinality of a generating set F .

To familiarize with this notion we prove the following

Proposition 3.6 *Let $p : A \rightarrow B$ be a morphism of abelian groups and (HB, \mathcal{R}) a tolerant $\mathbb{S}[\pm 1]$ -module.*

- (i) *Consider the relation $p^*(\mathcal{R}_k) := \{(x, y) \in HA(k_+) \times HA(k_+) \mid (Hp(x), Hp(y)) \in \mathcal{R}_k\}$. Then the pair $p^*(HB, \mathcal{R}) := (HA, p^*(\mathcal{R}))$ is a tolerant $\mathbb{S}[\pm 1]$ module.*
- (ii) *If p is surjective: $\dim_{\mathbb{S}[\pm 1]}(p^*(HB, \mathcal{R})) = \dim_{\mathbb{S}[\pm 1]}(HB, \mathcal{R})$.*

Proof (i) Since (HB, \mathcal{R}) is a tolerant $\mathbb{S}[\pm 1]$ -module, for any $\phi \in \text{Hom}_{\Gamma^\circ}(k_+, \ell_+)$ one has $HB(\phi)^{\times 2}(\mathcal{R}_k) \subset \mathcal{R}_\ell$. Also

$$HA(\phi)^{\times 2}(p^*\mathcal{R}_k) \subset p^*\mathcal{R}_\ell \quad \forall \phi \in \text{Hom}_{\Gamma^\circ}(k_+, \ell_+),$$

which shows that $p^*(HB, \mathcal{R}) = (HA, p^*(\mathcal{R}))$ is a tolerant $\mathbb{S}[\pm 1]$ module.

(ii) Let $F \subset HB(1_+) = B$ be a generating set for (HB, \mathcal{R}) , and let $F' \subset HA(1_+) = A$ be a lift of F , with $\#F' = \#F$. Let us show that F' is a generating set for $p^*(HB, \mathcal{R}) = (HA, p^*(\mathcal{R}))$. Let $a \in HA(1_+) = A$, then there exists coefficients $\alpha_j \in \{-1, 0, 1\}$, $j \in F$, such that $(p(a), \sum_F \alpha_j j) \in \mathcal{R}$. It follows that using the lifts $j' \in F'$ of $j \in F$ one has $(a, \sum_{F'} \alpha_j j') \in p^*(\mathcal{R})$, hence F' is a generating set for $p^*(HB, \mathcal{R})$. Thus $\dim_{\mathbb{S}[\pm 1]}(p^*(HB, \mathcal{R})) \leq \dim_{\mathbb{S}[\pm 1]}(HB, \mathcal{R})$. Conversely, let $F' \subset A$ be a generating set for $p^*(HB, \mathcal{R})$ and $F := p(F')$. Then condition 1. of Definition 3.5 for F' implies the same condition for F , thus one has $\#F' = \#F$. Let $b \in HB(1_+) = B$ and $a \in A$ with $p(a) = b$. Then there exists coefficients $\alpha_j \in \{-1, 0, 1\}$, $j \in F'$, such that $(a, \sum_{F'} \alpha_j j') \in p^*(\mathcal{R})$. This implies $(b, \sum \alpha_j j) \in \mathcal{R}$ so that F is a generating set for (HB, \mathcal{R}) . \square

Next, we apply this functorial machinery to the geometry of $\overline{\text{Spec } \mathbb{Z}}$. We retain the notations of section 2.

Proposition 3.7 *Let $D = \sum_j a_j \{p_j\} + a\{\infty\}$ be an Arakelov divisor on $\overline{\text{Spec } \mathbb{Z}}$ and let $\pi : \mathbb{A}_{\mathbb{Q}} \rightarrow \mathbb{R}/L$ be the projection of the adèles on their archimedean component modulo the lattice*

$$L = H^0(\text{Spec } \mathbb{Z}, \mathcal{O}(D)_f) := \{q \in \mathbb{Q} \mid |q|_\nu \leq \exp(D)(\nu), \forall \nu \neq \infty\}. \quad (15)$$

- (i) *Let d be the metric on \mathbb{R}/L induced by the standard metric on \mathbb{R} and set $\lambda = \exp a$. Then one has $H^1(D) = \pi^*((\mathbb{R}/L, d)_\lambda)$.*
 (ii) $\dim_{\mathbb{S}[\pm 1]} H^1(D) = \dim_{\mathbb{S}[\pm 1]}((\mathbb{R}/L, d)_\lambda)$.

Proof (i) Let $j : \mathbb{A}_{\mathbb{Q}}^f \rightarrow \mathbb{A}_{\mathbb{Q}}$, $j(x) := (x, 0)$, be the embedding of finite adèles in adèles. Using the ultrametric property of the local norms at the finite places one sees that $j(\mathcal{O}(D)_f) = \mathcal{O}(D) \cap j(\mathbb{A}_{\mathbb{Q}}^f) \subset \mathbb{A}_{\mathbb{Q}}$ is a compact subgroup. Set $G = \mathbb{Q} \times j(\mathcal{O}(D)_f)$: one has $j(\mathcal{O}(D)_f) \cap \mathbb{Q} = \{0\}$ since all the adèles in $j(\mathcal{O}(D)_f)$ have archimedean component equal to 0. Thus, the restriction of the morphism of addition, $\alpha : \mathbb{Q} \times \mathbb{A}_{\mathbb{Q}} \rightarrow \mathbb{A}_{\mathbb{Q}}$, $\alpha(q, a) = q + a$, to G determines an isomorphism of G with the subgroup $\alpha(G) = \mathbb{Q} + j(\mathcal{O}(D)_f)$ of $\mathbb{A}_{\mathbb{Q}}$. Note, in particular, that $\alpha(G)$ is closed in $\mathbb{A}_{\mathbb{Q}}$, since \mathbb{Q} is discrete (hence closed) and $j(\mathcal{O}(D)_f)$ is compact. In the following, we identify (set-theoretically) $\mathbb{A}_{\mathbb{Q}}$ with the product $\mathbb{A}_{\mathbb{Q}}^f \times \mathbb{R}$ endowed with the two projection morphisms $p_f : \mathbb{A}_{\mathbb{Q}} \rightarrow \mathbb{A}_{\mathbb{Q}}^f$ and $p_\infty : \mathbb{A}_{\mathbb{Q}} \rightarrow \mathbb{R}$. The subgroup $p_f(\mathbb{Q}) \subset \mathbb{A}_{\mathbb{Q}}^f$ is dense and the subgroup $p_f(j(\mathcal{O}(D)_f)) = \mathcal{O}(D)_f \subset \mathbb{A}_{\mathbb{Q}}^f$ is open. Hence $p_f(\alpha(G)) = \mathbb{A}_{\mathbb{Q}}^f$. Thus the projection p_∞ induces the isomorphism of groups $p : \mathbb{A}_{\mathbb{Q}}/\alpha(G) \xrightarrow{\sim} \mathbb{R}/L$, where $L = p_\infty(\ker(p_f \circ \alpha))$. The kernel of the composite $p_f \circ \alpha : G \rightarrow \mathbb{A}_{\mathbb{Q}}^f$ is the group of pairs $(q, a) \in \mathbb{Q} \times j(\mathcal{O}(D)_f)$ such that $p_f(q) + p_f(a) = 0$. Such pairs are determined by the value of q and thus

$$L = p_\infty(H^0(\text{Spec } \mathbb{Z}, \mathcal{O}(D)_f)) = \{q \in \mathbb{Q} \mid |q|_\nu \leq \exp(D)(\nu), \forall \nu \neq \infty\}. \quad (16)$$

By Proposition 2.2, $H^1(D)$ is the tolerant $\mathbb{S}[\pm 1]$ -module $H^1(D) = (H\mathbb{A}_{\mathbb{Q}}, \mathcal{R})$ where the relations are given by (11), *i.e.*

$$\mathcal{R}_k := \{(x, y) \in H\mathbb{A}_{\mathbb{Q}}(k_+) \times H\mathbb{A}_{\mathbb{Q}}(k_+) \mid x - y \in \text{Image } \psi(k_+)\},$$

and where ψ is as in (9). By construction one has $\mathbb{Q} \times \mathcal{O}(D) \simeq G \times [-e^a, e^a]$. After quotienting both sides of (9) by $G = \mathbb{Q} \times j(\mathcal{O}(D)_f)$, the map ψ becomes

$$\phi : \|H\mathbb{R}\|_{e^a} \rightarrow H(\mathbb{R}/L), \quad \phi(x) = x + L \in \mathbb{R}/L \quad \forall x \in [-e^a, e^a] \subset \mathbb{R}, \quad (17)$$

where $L \subset \mathbb{Q} \subset \mathbb{R}$ is the lattice (16). Thus one obtains that the relation \mathcal{R} is equal to the inverse image by the map $\pi : \mathbb{A}_{\mathbb{Q}} \rightarrow \mathbb{R}/L$ of the relation (14).

(ii) Follows from Proposition 3.6 (ii). \square

4 The dimension of $H^0(D)$

In this section we consider, for each integer $n > 0$, the $\mathbb{S}[\pm 1]$ -module $\|HZ\|_n$

$$\|HZ\|_n : \Gamma^{\circ} \longrightarrow \mathfrak{Sets}_* \quad \|HZ\|_n(F) := \{\phi \in HZ(F) \mid \sum_{F \setminus \{*\}} |\phi(x)| \leq n\}$$

and evaluate its dimension. It follows from the $\mathbb{S}[\pm 1]$ -module structure, that (8) holds for a sum of the form

$$\sum_F \alpha_j j \in [-n, n] \cap \mathbb{Z} = \|HZ\|_n(1_+) \quad \forall \alpha_j \in \{-1, 0, 1\} \quad \text{s.t.} \quad \sum_F |\alpha_j j| \leq n.$$

By applying Definition 3.5, a subset $F \subset [-n, n] \cap \mathbb{Z} = \|HZ\|_n(1_+)$ generates $\|HZ\|_n(1_+)$ if and only if for every element $x \in \|HZ\|_n(1_+)$ there exists coefficients $\alpha_j \in \{-1, 0, 1\}$, $j \in F$, such that $x = \sum_F \alpha_j j$ and $\sum_F |\alpha_j j| \leq n$. The *dimension* $\dim_{\mathbb{S}[\pm 1]} \|HZ\|_n$ is the minimal cardinality of a generating set.

The explicit computation of $\dim_{\mathbb{S}[\pm 1]} \|HZ\|_n$ follows from the next two statements.

Lemma 4.1 *For $k \in \mathbb{N}$, let $n = (3^k - 1)/2$. Consider the subset $F := \{3^i, 0 \leq i \leq k-1\} \subset [-n, n]$. Then F is a generating set and the map*

$$\theta : \{-1, 0, 1\}^{\times k} \rightarrow [-n, n] \cap \mathbb{Z}, \quad \theta((\alpha_i)) := \sum_{i=0}^{k-1} \alpha_i 3^i \quad (18)$$

is bijective. Moreover $\sum_{i=0}^{k-1} |\alpha_i 3^i| \leq n$, for all $\alpha_i \in \{-1, 0, 1\}^{\times k}$.

Proof First of all, note that for all $\alpha \in \{-1, 0, 1\}^{\times k}$ one has

$$\sum_{i=0}^{k-1} |\alpha_i 3^i| \leq \sum_{i=0}^{k-1} 3^i = n.$$

This shows that for any $\alpha \in \{-1, 0, 1\}^{\times k}$ one has $z = (\alpha_i 3^i) \in \|HZ\|_n(k_+)$, and, as in (8), $\|HZ\|_n(\sigma)(z) = \theta((\alpha_i))$. We prove the bijectivity of θ by induction on k . It is clear for $k = 1$. For $k = 2$, $n = 4$ and one writes the set $[-4, 4] \cap \mathbb{Z}$ as the disjoint union of three subsets as follows

$$[-4, 4] \cap \mathbb{Z} = \{-4, -3, -2\} \cup \{-1, 0, 1\} \cup \{2, 3, 4\} =$$

$$= (-3 + \{-1, 0, 1\}) \cup \{-1, 0, 1\} \cup (3 + \{-1, 0, 1\}).$$

This shows that θ is bijective. Next, by assuming to have shown the bijectivity up to $k-1$, we deduce it for k . Indeed, with $n = (3^k - 1)/2$ let $m = (3^{k-1} - 1)/2$. The induction hypothesis ensures that every element of the set $[-m, m] \cap \mathbb{Z}$ can be uniquely written as a sum $\sum_{i=0}^{k-2} \alpha_i 3^i$. We divide the set $[-n, n] \cap \mathbb{Z}$ into three disjoint parts of the form

$$J_{-1} := [-n, -n + 2m] \cap \mathbb{Z}, \quad J_0 := [-m, m] \cap \mathbb{Z}, \quad J_1 := [n - 2m, n] \cap \mathbb{Z}.$$

One has $n - 2m = m + 1$ and thus the union of these three disjoint sets is $[-n, n] \cap \mathbb{Z}$. Moreover one has $J_1 = (n - m) + J_0$ with $n - m = 3^{k-1}$, and similarly $J_{-1} = J_0 - 3^{k-1}$. Thus, given $x \in [-n, n] \cap \mathbb{Z}$ one determines uniquely the coefficients $\alpha_i \in \{-1, 0, 1\}$ such that $\theta((\alpha_i)) = x$. Indeed, one uses the above partition in three intervals to determine the coefficient α_{k-1} and then one applies the induction hypothesis to determine the others. \square

Remark 4.2 1. *The conceptual explanation of Lemma 4.1 derives from the following peculiar property of the Teichmüller lift $\tau : \mathbb{F}_3 \rightarrow \mathbb{Z}_3$. One has $\tau(\mathbb{F}_3) = \{-1, 0, 1\} \subset \mathbb{Z} \subset \mathbb{Z}_3$ and τ extends to a canonical map $\tilde{\tau} : W_k(\mathbb{F}_3) \rightarrow \mathbb{Z}$ from Witt vectors to \mathbb{Z}*

$$\xi = (\xi_j) \in W_k(\mathbb{F}_3) \mapsto \tilde{\tau}(\xi) := \sum_0^{k-1} \tau(\xi_j) 3^j \in \mathbb{Z}.$$

This map coincides with θ in (18) and one can compute the partially defined sums (8) in $\|HZ\|_n$ using the addition in $W_k(\mathbb{F}_3)$. The prime $p = 3$ is the only prime such that the subring $\mathbb{Z} \subset \mathbb{Z}_p = W(\mathbb{F}_p)$ coincides with the ring of Witt vectors with finitely many non-zero components. When $p > 3$ one has $\tau(\mathbb{F}_p) \not\subset \mathbb{Z}$, while for $p = 2$ the integer $-1 \in \mathbb{Z} \subset \mathbb{Z}_2$ is the Witt vector whose components are all equal to 1. In general the Witt vectors with finitely many non-zero components do not even form a subgroup of the additive group of Witt vectors.

2. *The Teichmüller lift τ , as a morphism of multiplicative pointed monoids, induces a morphism $\mathbb{S}[\pm 1] \rightarrow H\mathbb{Z}$ of \mathbb{S} -algebras.*
3. *The ordering of the natural numbers encoded by θ coincides with the lexicographic ordering of the coefficients $(\alpha_i) \in \{-1, 0, 1\}^{\times k}$*

$$\theta((\alpha_i)) < \theta((\beta_i)) \iff \exists j \text{ s.t. } \alpha_j < \beta_j \text{ and } \alpha_\ell = \beta_\ell \quad \forall \ell > j.$$

We recall that the ceiling function $\lceil x \rceil$ associates to a positive real number x the smallest integer $n \geq x$.

Proposition 4.3 (i) *For any integer $n \geq 0$ one has*

$$\dim_{\mathbb{S}[\pm 1]} \|HZ\|_n = \left\lceil \frac{\log(2n+1)}{\log 3} \right\rceil. \quad (19)$$

(ii) For $n \notin \{2, 5\}$, there exists a generating subset $F \subset \{1, \dots, n\}$ of cardinality $\#F = \lceil \frac{\log(2n+1)}{\log 3} \rceil$, with $\sum_{j \in F} j = n$, such that the following map surjects onto $[-n, n] \cap \mathbb{Z}$

$$\theta : \{-1, 0, 1\}^{\times(\#F)} \rightarrow [-n, n] \cap \mathbb{Z}, \quad \theta((\alpha_i)) := \sum_{j \in F} \alpha_j j.$$

Proof (i) Set $\kappa(n) := \dim_{\mathbb{S}[\pm 1]} \|HZ\|_n$. Since the cardinality of $[-n, n] \cap \mathbb{Z}$ is $2n + 1$ and the cardinality of $\{-1, 0, 1\}^{\times \#F}$ is $3^{\#F}$, it follows that if F is a generating set one must have $3^{\#F} \geq 2n + 1$. Thus one has $\#F \geq \frac{\log(2n+1)}{\log 3}$, and since $\#F$ is an integer one gets $\#F \geq \lceil \frac{\log(2n+1)}{\log 3} \rceil$. This shows that $\kappa(n) \geq \lceil \frac{\log(2n+1)}{\log 3} \rceil$. It remains to prove that for each $n \geq 0$ one can find a generating set F with $\#F = \lceil \frac{\log(2n+1)}{\log 3} \rceil$. One uses Lemma 4.1, and distinguishes several cases starting with the easiest one. Set $F(k) := \{3^i, 0 \leq i \leq k - 1\}$ and let $E \subset \mathbb{N}$ be the subset

$$E = \left\{ 3^\ell + \frac{1}{2} (3^m - 1) \mid m \in \mathbb{N}, \ell < m \right\}. \quad (20)$$

We list the first elements of E as follows

2, 5, 7, 14, 16, 22, 41, 43, 49, 67, 122, 124, 130, 148, 202, 365, 367, 373, 391, 445, 607 ...

First, assume that $n \notin E$. Then let m be the largest integer such that $3^m \leq 2n + 1$. If $3^m = 2n + 1$, then Lemma 4.1 provides one with a generating set of cardinality $m = \frac{\log(2n+1)}{\log 3}$. Assume now that $3^m < 2n + 1$. Then, the map θ for $F(m)$ covers the set $[-q, q] \cap \mathbb{Z}$, where $q = \frac{1}{2} (3^m - 1)$. Thus by adjoining to $F(m)$ the element $n - q$ one covers $[-n, n] \cap \mathbb{Z}$. More precisely, let $F = F(m) \cup \{n - \frac{1}{2} (3^m - 1)\}$, then we show that F fulfills the conditions in (ii). One has $\#F = \#F(m) + 1$ since the additional element $n - \frac{1}{2} (3^m - 1)$ does not belong to $F(m)$ as, by hypothesis, $n \notin E$. Since $\#F(m) = m$, $\#F = m + 1 = \lceil \frac{\log(2n+1)}{\log 3} \rceil$. The sum of elements in $F(m)$ is $\frac{1}{2} (3^m - 1)$, so the sum of elements in F is equal to n . Next, we show that θ is surjective. Its image contains the three sets $\theta(F(m))$, $\theta(F(m)) + n - \frac{1}{2} (3^m - 1)$, and $\theta(F(m)) - (n - \frac{1}{2} (3^m - 1))$. Using Lemma 4.1 one obtains, with $q = \frac{1}{2} (3^m - 1)$

$$\theta F(m) = [-q, q] \cap \mathbb{Z}, \quad \theta F(m) + n - q = [n - 2q, n] \cap \mathbb{Z},$$

$$\theta F(m) - (n - q) = [-n, -n + 2q] \cap \mathbb{Z}.$$

One has $3(2q + 1) = 3^{m+1} > 2n + 1$. This inequality prevents the three subsets from being disjoint, thus the upper limit q of $\theta F(m)$ is greater or equal to the lower limit $n - 2q$ of the interval $\theta F(m) + n - q$. Thus θ is surjective.

The next case is for $n \in E$ of the form $n = 1 + \frac{1}{2} (3^m - 1)$, for some m ($\ell = 0$ in (20)). We assume that $m > 2$ (this excludes the cases $n = 2$ for $m = 1$, and $n = 5$ for $m = 2$). Let

$$F := \{3^k \mid 0 \leq k \leq m - 2\} \cup \{2, 3^{m-1} - 1\}.$$

(This choice avoids the repetition of the element $1 = n - q$ while keeping the sum of elements of F equal to n). We show that F fulfills the conditions in (ii). We have $\#F = m - 1 + 2 = m + 1$ since $3^{m-1} - 1 \notin \{1, 2\}$. Moreover $2n + 1 = 3^m + 2$, thus $\#F = m + 1 = \lceil \frac{\log(2n+1)}{\log 3} \rceil$. The sum of terms in F is equal to one plus the sum of

the powers of 3 up to 3^{m-1} and thus is equal to n . Let us now show that the map θ is surjective. By construction $\theta(F)$ is the union of nine intervals obtained, with $q' = \frac{1}{2}(3^{m-1} - 1)$, by adding to the set $[-q', q'] \cap \mathbb{Z}$ the terms in $\{2, 3^{m-1} - 1\}$ with coefficients in $\{-1, 0, 1\}$. With the term $3^{m-1} - 1$ one covers the set $[-q+1, q-1] \cap \mathbb{Z}$, where $q = \frac{1}{2}(3^m - 1)$ as the union of the three sets

$$[-q', q'] \cap \mathbb{Z}, \quad ([-q', q'] + 3^{m-1} - 1) \cap \mathbb{Z}, \quad ([-q', q'] - (3^{m-1} - 1)) \cap \mathbb{Z}.$$

One has $q' + 3^{m-1} - 1 = n - 2$, and thus it remains to cover the two elements $n - 1, n$ and their opposite. This is done using the set $([-q', q'] + 3^{m-1} - 1 + 2) \cap \mathbb{Z}$ whose upper limit is n , and appealing to the fact that since $q' > 1$ this interval contains $n - 1$.

Finally, we consider the case $n \in E$ of the form $n = 3^\ell + \frac{1}{2}(3^m - 1)$, with $m \in \mathbb{N}$, $0 < \ell < m$. Here, the additional element $n - \frac{1}{2}(3^m - 1)$ does belong to $F(m)$ so the proof of the first case considered above does not apply since 3^ℓ appears twice in $F(m) \cup \{n - \frac{1}{2}(3^m - 1)\}$. We replace this double occurrence of 3^ℓ by the two distinct elements $3^\ell \pm 1$. This gives

$$F = (F(m) \setminus \{3^\ell\}) \cup \{3^\ell - 1, 3^\ell + 1\}.$$

By construction $\#F(m) + 1 = m + 1$. Furthermore one has: $n = 3^\ell + \frac{1}{2}(3^m - 1)$, $2n + 1 = 3^m + 2 \times 3^\ell$ and since $0 < \ell < m$ one gets $\#F = m + 1 = \lceil \frac{\log(2n+1)}{\log 3} \rceil$. The sum of elements of F is the same as the sum of elements of $F(m)$ plus 3^ℓ which is equal to n . It remains to show that, for such F , θ is surjective. We first deal with the set $G(\ell) := F(\ell) \cup \{3^\ell - 1, 3^\ell + 1\}$ and show that the map $\theta_{G(\ell)}$, for this set, surjects onto the interval $[-t(\ell), t(\ell)] \cap \mathbb{Z}$, where $t(\ell) = \sum_{G(\ell)} j = 3^\ell + \frac{1}{2}(3^{\ell+1} - 1)$. Lemma 4.1 shows that the image of $\theta_{F(\ell)}$ is the set $J \cap \mathbb{Z}$, with $J = [-\frac{1}{2}(3^\ell - 1), \frac{1}{2}(3^\ell - 1)]$. To show the surjectivity of $\theta_{G(\ell)}$ it is enough to show that the following intervals cover $[0, t(\ell)]$:

$$J, \quad J + 3^\ell - 1, \quad J + 3^\ell + 1, \quad J + 2 \times 3^\ell.$$

The upper limit of $J + 2 \times 3^\ell$ is $\frac{1}{2}(3^\ell - 1) + 2 \times 3^\ell = t(\ell)$, its lower limit is equal to the upper limit of $J + 3^\ell + 1$. The lower limit of $J + 3^\ell - 1$ is $3^\ell - 1 - \frac{1}{2}(3^\ell - 1) = \frac{1}{2}(3^\ell - 1)$ which is the upper limit of J . Moreover, the length of J is $3^\ell - 1 \geq 2$ so that two translates of J of the form $J + a, J + a + 2$ necessarily overlap. This shows that the map $\theta_{G(\ell)}$ surjects onto the interval $[-t(\ell), t(\ell)] \cap \mathbb{Z}$. Thus $\theta_{G(\ell)}$ is the symmetric enlargement of the interval $\theta(F(\ell + 1))$ obtained by adding 3^ℓ to the upper limit. Then, as in the proof of Lemma 4.1, one obtains by induction that when one adjoins the higher powers $3^{\ell'}$, where $\ell < \ell' < m$, one achieves the enlargement of the interval $\theta(F(m))$ obtained by adding 3^ℓ to the upper limit. This shows that the map θ for F is surjective. In fact the above arguments prove (ii) and (i), except that one has to take care of the special cases $n = 2, 5$. For $n = 2$, $\{1, 2\}$ is a generating set but the sum of its terms is > 2 . For $n = 5$, $\{1, 2, 3\}$ is a generating set but the sum of its terms is > 5 . \square

We are now ready to determine the dimension of the $\mathbb{S}[\pm 1]$ -module $H^0(D)$, for any Arakelov divisor D . It is a non-decreasing function of $\deg D$ but the formula for this dimension drops by 1 on an exceptional set L . It is the open set defined as the union of open intervals as follows

$$L + \log 2 = \bigcup_{k \in \mathbb{N}} (k \log 3, k \log 3 + \epsilon_k), \quad \epsilon_k := \log(1 + 3^{-k}). \quad (21)$$

It has finite Lebesgue measure

$$|L| = \sum_{k \geq 0} \epsilon_k = \log \left(-1; \frac{1}{3} \right)_{\infty} = 1.14099 \dots$$

where $(-1; \frac{1}{3})_{\infty} = \prod_{n=0}^{\infty} (1 + 3^{-n})$ is the -1 Pochhammer symbol at $\frac{1}{3}$.

Theorem 4.4 *Let D be an Arakelov divisor on $\overline{\text{Spec } \mathbb{Z}}$. If $\deg D \geq -\log 2$, then*

$$\dim_{\mathbb{S}[\pm 1]} H^0(D) = \left\lceil \frac{\deg D + \log 2}{\log 3} \right\rceil - \mathbf{1}_L \quad (22)$$

where $\mathbf{1}_L$ is the characteristic function of the open set L .

Proof Let $a = \deg D$ and $n = \lfloor \exp(\deg(D)) \rfloor$, then $H^0(D) = \|HZ\|_n$. By Proposition 4.3 one gets

$$\dim_{\mathbb{S}[\pm 1]} H^0(D) = \left\lceil \frac{\log(2n+1)}{\log 3} \right\rceil, \quad n := \lfloor \exp(a) \rfloor.$$

Let us compare this formula with (22). Assume first that $a \in L$, thus there exists $k \in \mathbb{N}$ such that $a + \log 2 \in (k \log 3, k \log 3 + \epsilon_k)$. Then, $2 \exp a \in (3^k, 3^k + 1)$ and with $n = \lfloor \exp a \rfloor$, one has $n = \frac{1}{2}(3^k - 1)$. Thus $2n+1 = 3^k$ and $\lceil \frac{\log(2n+1)}{\log 3} \rceil = k$. Moreover, $\frac{a+\log 2}{\log 3} \in (k, k + \frac{\epsilon_k}{\log 3})$, where $0 < \frac{\epsilon_k}{\log 3} < 1$, hence one has $\lceil \frac{a+\log 2}{\log 3} \rceil = k+1$, so that (22) holds since $\mathbf{1}_L(a) = 1$. Now, assume that $a \notin L$, equivalently that $\frac{a+\log 2}{\log 3} \in [k + \frac{\epsilon_k}{\log 3}, k+1]$ for some integer $k \geq 0$ (since $a \geq -\log 2$ by hypothesis). Then $\lceil \frac{a+\log 2}{\log 3} \rceil = k+1$. One has

$$2 \exp a \in [\exp(k \log 3)(1 + 3^{-k}), \exp((k+1) \log 3)] = [3^k + 1, 3^{k+1}].$$

In this case $n = \lfloor \exp a \rfloor \in [\frac{1}{2}(3^k + 1), \frac{1}{2}(3^{k+1} - 1)]$ and thus $2n+1 \in [3^k + 2, 3^{k+1}]$. This implies that $\frac{\log(2n+1)}{\log 3} \in (k, k+1]$ and hence that $\lceil \frac{\log(2n+1)}{\log 3} \rceil = k+1$. In this case (22) holds since $\mathbf{1}_L(a) = 0$. \square

5 The dimension of $H^1(D)$

Let (A, d) be an abelian group endowed with a translation invariant metric d . For $\lambda \in \mathbb{R}_{>0}$, we shall refer to $(A, d)_{\lambda}$ as the associated object of the category $\Gamma\mathcal{T}_{*}$ as in Proposition 3.4. The \mathbb{S} -module structure determines the addition in $A = (A, d)_{\lambda}(1_+)$, as well as the action of $\mathbb{S}[\pm 1]$ on it, where for $x \in A = (A, d)_{\lambda}(1_+)$, $-x$ is the additive inverse of x . The metric d determines the tolerance relation \mathcal{R} on $A = (A, d)_{\lambda}(1_+)$. Next result computes the dimension of the tolerant $\mathbb{S}[\pm 1]$ -module $(\mathbb{R}/\mathbb{Z}, d)_{\lambda}$.

Proposition 5.1 *Let $\lambda \in \mathbb{R}_{>0}$, and $U(1)_{\lambda} = (U(1), d)_{\lambda}$ where $U(1)$ is the abelian group \mathbb{R}/\mathbb{Z} endowed with the canonical metric d of length 1. Then*

$$\dim_{\mathbb{S}[\pm 1]} U(1)_{\lambda} = \begin{cases} \left\lceil \frac{-\log \lambda - \log 2}{\log 3} \right\rceil & \text{if } \lambda < \frac{1}{2} \\ 0 & \text{if } \lambda \geq \frac{1}{2}. \end{cases} \quad (23)$$

Proof For $\lambda \geq \frac{1}{2}$, any element of $U(1)_\lambda = (\mathbb{R}/\mathbb{Z}, d)_\lambda$ is at distance $\leq \lambda$ from 0, thus one can take $F = \emptyset$ as generating set since, by convention, $\sum_\emptyset = 0$. Thus $\dim_{\mathbb{S}[\pm 1]} U(1)_\lambda = 0$. Next, we assume $\lambda < \frac{1}{2}$. Let $F \subset U(1)$ be a generating set and let $k = \#F$. One easily sees that there are at most 3^k elements of the form $\sum_F \alpha_j j$. The subsets $\{x \in U(1) \mid d(x, \alpha_j j) \leq \lambda\}$ cover $U(1)$, and since each of them has measure 2λ one gets the inequality $2\lambda \cdot 3^k \geq 1$. Thus $k \geq \frac{-\log \lambda - \log 2}{\log 3}$. For $\lambda = \frac{1}{6}$ one has $k \geq 1$ and the subset $F = \{\frac{1}{3}\}$ generates, thus $\dim_{\mathbb{S}[\pm 1]} U(1)_\lambda = 1$. When $\frac{-\log \lambda - \log 2}{\log 3} = m$ is an integer, one has $\lambda = \frac{1}{2}3^{-m}$. Let $F(m) = \{\frac{1}{3}, \dots, \frac{1}{3^m}\}$. The minimal distance between two elements of $F(m)$ is the distance between 3^{-m} and $3^{-m+1} = 3 \cdot 3^{-m}$ which is $2 \cdot 3^{-m} = 4\lambda$. Let us show that $F(m)$ is a generating set. By Lemma 4.1 any integer q in the interval $[-N, N]$, for $N = \frac{1}{2}(3^m - 1)$ can be written as $q = \sum_{i=0}^{m-1} \alpha_i 3^i$, with $\alpha_i \in \{-1, 0, 1\}$. One then gets

$$q \cdot 3^{-m} = \sum_{i=0}^{m-1} \alpha_i 3^{i-m} = \sum_{j=1}^m \alpha_{m-j} 3^{-j}.$$

Let $x \in [-\frac{1}{2}, \frac{1}{2}]$, then $3^m x \in [-\frac{3^m}{2}, \frac{3^m}{2}]$ and there exists an integer $q \in [-N, N]$ such that $|3^m x - q| \leq \frac{1}{2}$. Hence $d(x, q \cdot 3^{-m}) \leq \lambda$. This proves that $F(m)$ is a generating set (see Definition 3.5) and one derives $\dim_{\mathbb{S}[\pm 1]} U(1)_\lambda = m$. Assume now that $\frac{-\log \lambda - \log 2}{\log 3} \in (m, m+1)$, where m is an integer. For any generating set F of cardinality k one has $k \geq \frac{-\log \lambda - \log 2}{\log 3} > m$ so that $k \geq m+1$. The subset $F(m+1) = \{\frac{1}{3}, \dots, \frac{1}{3^{m+1}}\}$ fulfills the first condition of Definition 3.5 since the minimal distance between two elements of $F(m+1)$ is $2 \cdot 3^{-m-1}$ which is larger than $\lambda < \frac{1}{2}3^{-m}$. As shown above, the subset $F(m+1)$ is generating for $\lambda = \frac{1}{2}3^{-m-1}$ and a fortiori for $\lambda > \frac{1}{2}3^{-m-1}$ (as by assumption $\frac{-\log \lambda - \log 2}{\log 3} < m+1$). Thus one obtains $\dim_{\mathbb{S}[\pm 1]} U(1)_\lambda = m+1$ and (23) is proven. \square

Remark 5.2 *The proof of Proposition 5.1 is related to a definite advantage of the triadic expansion of numbers [12]: truncating a number is identical to rounding it. This property does not hold in the decimal system where rounding a number requires the knowledge of the next digit.*

We extend the ceiling function to negative values of the variable as follows

$$\lceil x \rceil' = \begin{cases} \lceil x \rceil & \text{if } x \geq 0 \\ -\lceil -x \rceil & \text{if } x \leq 0. \end{cases} \quad (24)$$

We can now state and prove the absolute Riemann-Roch theorem for $\overline{\text{Spec } \mathbb{Z}}$ over $\mathbb{S}[\pm 1]$. With L the exceptional set defined in (21) one has

Theorem 5.3 *Let D be an Arakelov divisor on $\overline{\text{Spec } \mathbb{Z}}$. Then*

$$\dim_{\mathbb{S}[\pm 1]} H^0(D) - \dim_{\mathbb{S}[\pm 1]} H^1(D) = \left\lceil \frac{\deg D + \log 2}{\log 3} \right\rceil' - \mathbf{1}_L. \quad (25)$$

Proof By appealing to the invariance under linear equivalence (Proposition 2.2 (iii)), one may assume that $D = a\{\infty\}$. Then it follows from Proposition 3.7 (ii) that $\dim_{\mathbb{S}[\pm 1]} H^1(D) = \dim_{\mathbb{S}[\pm 1]} U(1)_\lambda$ for $\lambda = \exp(\deg(D))$. Assume first that $\exp(\deg(D)) \geq \frac{1}{2}$, then by Proposition 5.1: $\dim_{\mathbb{S}[\pm 1]} H^1(D) = 0$, thus for $\deg(D) \geq -\log 2$ (25) follows from (22). Let us now assume that $\deg(D) < -\log 2$. Then $\dim_{\mathbb{S}[\pm 1]} H^0(D) = 0$ since $H^0(D) = \{*\}$ when $\deg D < 0$. Moreover $\deg D \notin L$ since L is lower bounded by $-\log 2$. Thus (25) follows from (23) which gives, using (24)

$$-\dim_{\mathbb{S}[\pm 1]} H^1(D) = -\dim_{\mathbb{S}[\pm 1]} U(1)_\lambda = -\left\lceil \frac{-\log \lambda - \log 2}{\log 3} \right\rceil = \left\lceil \frac{\deg D + \log 2}{\log 3} \right\rceil'.$$

This ends the proof of (25). \square

6 Duality

In this section we prove an absolute analogue of Serre's duality, namely the following isomorphism of $\mathbb{S}[\pm 1]$ -modules, for any divisor D on $\overline{\text{Spec } \mathbb{Z}}$:

$$H^0(D) \simeq \underline{\text{Hom}}_{\Gamma\mathcal{T}^*}(H^1(K - D), U(1)_{\frac{1}{4}}).$$

Here, the divisor $K = -2\{2\}$ plays the role of the canonical divisor. The choice of the tolerant $\mathbb{S}[\pm 1]$ -module $(U(1), d)_{\frac{1}{4}}$ as dualizing module is motivated by Pontryagin duality (see 6.2). One has $\dim_{\mathbb{S}[\pm 1]}(U(1)_{\frac{1}{4}}) = 1$. This equality, in fact, also holds for the tolerant $\mathbb{S}[\pm 1]$ -module $U(1)_\lambda$ for $\frac{1}{6} \leq \lambda < \frac{1}{2}$: the specific choice $\lambda = \frac{1}{4}$ is dictated by the invariance of the Riemann-Roch formula (25) when one switches H^0 and H^1 and replaces D by $K - D$ (ignoring the exceptional set L).

6.1 $\underline{\text{Hom}}_{\mathbb{S}}(\|H\mathbb{R}\|_\lambda, \|H\mathbb{R}\|_\mu)$

We start with the following general statement.

Lemma 6.1 *Let $\lambda, \mu \in \mathbb{R}_{>0}$. The \mathbb{S} -algebra structure of $H\mathbb{R}$ induces an isomorphism of \mathbb{S} -modules*

$$\|H\mathbb{R}\|_{\mu/\lambda} \simeq \underline{\text{Hom}}_{\mathbb{S}}(\|H\mathbb{R}\|_\lambda, \|H\mathbb{R}\|_\mu). \quad (26)$$

Proof One starts by defining the morphism of \mathbb{S} -modules

$$\eta : \|H\mathbb{R}\|_{\mu/\lambda} \rightarrow \underline{\text{Hom}}_{\mathbb{S}}(\|H\mathbb{R}\|_\lambda, \|H\mathbb{R}\|_\mu). \quad (27)$$

Precisely, the multiplication in the \mathbb{S} -algebra $H\mathbb{R}$ determines natural maps

$$m : H\mathbb{R}(X) \wedge H\mathbb{R}(Y) \rightarrow H\mathbb{R}(X \wedge Y)$$

inducing, for a fixed pointed set X , the map $m_X : H\mathbb{R}(X) \rightarrow \text{Hom}_{\mathbb{S}}(H\mathbb{R}, H\mathbb{R}(X \wedge -))$. The morphism η is defined as the natural transformation of functors which associates to $X = k_+$, the map $\eta_X : \|H\mathbb{R}\|_{\mu/\lambda}(X) \rightarrow \text{Hom}_{\mathbb{S}}(\|H\mathbb{R}\|_\lambda, \|H\mathbb{R}\|_\mu(X \wedge -))$ defined as the restriction of m_X . This restriction is meaningful in view of [1] (Proposition 6.1). Next, we show that η is an isomorphism. First, we determine the \mathbb{S} -module $\text{Hom}_{\mathbb{S}}(\|H\mathbb{R}\|_\lambda, \|H\mathbb{R}\|_\mu)$. For integers $1 \leq j \leq k$ we let as in (7)

$$\delta(j, k) \in \text{Hom}_{\Gamma^0}(k_+, 1_+), \quad \delta(j, k)(\ell) := \begin{cases} 1 & \text{if } \ell = j \\ * & \text{if } \ell \neq j. \end{cases}$$

Let $\phi \in \text{Hom}_{\mathbb{S}}(\|H\mathbb{R}\|_{\lambda}, \|H\mathbb{R}\|_{\mu})$. By construction, the natural transformation ϕ reads, for each k , as a map $\phi(k_+) : \|H\mathbb{R}\|_{\lambda}(k_+) \rightarrow \|H\mathbb{R}\|_{\mu}(k_+)$ and by naturality we have

$$\|H\mathbb{R}\|_{\mu}(\delta(j, k)) \circ \phi(k_+) = \phi(1_+) \circ \|H\mathbb{R}\|_{\lambda}(\delta(j, k)). \quad (28)$$

Since an element $y \in \|H\mathbb{R}\|_{\mu}(k_+)$ is determined by its components $y_j \in \mathbb{R}$, $1 \leq j \leq k$, with $y_j = \|H\mathbb{R}\|_{\mu}(\delta(j, k))(y)$, (28) shows that $\phi(k_+)$ is uniquely determined by the map $\phi(1_+)$ acting componentwise, *i.e.*

$$\phi(k_+)((x_j)) = (\phi(1_+)(x_j)). \quad (29)$$

Moreover, the map $f = \phi(1_+) : [-\lambda, \lambda] \rightarrow [-\mu, \mu]$ fulfills

$$f(x + y) = f(x) + f(y) \quad \forall x, y \text{ s.t. } |x| + |y| \leq \lambda, \quad (30)$$

as one sees using the naturality of ϕ for the map $\sigma \in \text{Hom}_{\Gamma \circ} (2_+, 1_+)$, *i.e.* using

$$(\|H\mathbb{R}\|_{\mu}(\sigma) \circ \phi(2_+))(x, y) = (\phi(1_+) \circ \|H\mathbb{R}\|_{\lambda}(\sigma))(x, y) \implies f(x) + f(y) = f(x + y).$$

By (30) one has $f(x2^{-n}) = 2^{-n}f(x)$, for any $x \in [-\lambda, \lambda]$ and $n \geq 0$. Thus the ‘germ of map’ f uniquely extends to a map $\tilde{f} : \mathbb{R} \rightarrow \mathbb{R}$ defined by $\tilde{f}(x) := 2^n f(x2^{-n})$ for any n such that $|x2^{-n}| \leq \lambda$. Moreover, again by (30), the map \tilde{f} is additive and since $\tilde{f}([-2^{-n}\lambda, 2^{-n}\lambda]) \subset [-2^{-n}\mu, 2^{-n}\mu]$, it is also continuous and hence determined by the multiplication by a real number r . One has $|r\lambda| \leq \mu$ and hence $r \in \|H\mathbb{R}\|_{\mu/\lambda}(1_+)$. Thus one gets $\phi(1_+)(x) = rx$, $\forall x \in [-\lambda, \lambda]$. By (29) one obtains $\phi(k_+)((x_j)) = (rx_j)$, $\forall (x_j) \in \|H\mathbb{R}\|_{\lambda}(k_+)$. This shows that for $X = 1_+$, $\eta_X : \|H\mathbb{R}\|_{\mu/\lambda}(X) \rightarrow \text{Hom}_{\mathbb{S}}(\|H\mathbb{R}\|_{\lambda}, \|H\mathbb{R}\|_{\mu}(X \wedge -))$ is bijective. The next step is to determine $\text{Hom}_{\mathbb{S}}(\|H\mathbb{R}\|_{\lambda}, \|H\mathbb{R}\|_{\mu}(X \wedge -))$ for $X = \ell_+$. Let $\phi \in \text{Hom}_{\mathbb{S}}(\|H\mathbb{R}\|_{\lambda}, \|H\mathbb{R}\|_{\mu}(\ell_+ \wedge -))$. One has $\ell_+ \wedge k_+ = (\ell k)_+$ and an element $z \in \|H\mathbb{R}\|_{\mu}(\ell_+ \wedge k_+)$ is determined by its components $z(i, j) \in \mathbb{R}$ for $1 \leq i \leq \ell$, $1 \leq j \leq k$ such that $\sum |z(i, j)| \leq \mu$. In particular, for each j , the $z(i, j)$, $1 \leq i \leq \ell$, are the components of $z_j = \|H\mathbb{R}\|_{\mu}(\text{id} \wedge \delta(j, k))(z) \in \|H\mathbb{R}\|_{\mu}(\ell_+)$. This implies by applying the naturality of ϕ , that

$$\phi(k_+)(z) = (\phi(1_+)(z_j)). \quad (31)$$

As shown above the map $f = \phi(1_+) : [-\lambda, \lambda] \rightarrow \|H\mathbb{R}\|_{\mu}(\ell_+)$ fulfills

$$f(x + y) = f(x) + f(y) \quad \forall x, y \text{ s.t. } |x| + |y| \leq \lambda \quad (32)$$

and it extends to an additive map $\tilde{f} : \mathbb{R} \rightarrow \mathbb{R}^{\ell}$ which is continuous since \tilde{f} maps the interval $[-\lambda, \lambda]$ inside $\|H\mathbb{R}\|_{\mu}(\ell_+)$. Thus there exists real numbers r_i , $1 \leq i \leq \ell$, such that $\tilde{f}(x) = (r_i x) \forall x \in \mathbb{R}$. One has $\sum |r_i \lambda| \leq \mu$ and thus $r = (r_i) \in \|H\mathbb{R}\|_{\mu/\lambda}(\ell_+)$. Finally, using (31) it follows that $\phi = \eta_X(r)$. This proves that η as in (27) is an isomorphism. \square

6.2 Pontryagin duality

In order to formulate Pontryagin duality in this context we consider, for $\lambda > 0$, the $\mathbb{S}[\pm 1]$ -module $U(1)_{\lambda} = (U(1), d)_{\lambda}$, where $U(1)$ is the abelian group \mathbb{R}/\mathbb{Z} endowed with its canonical metric d of length 1. For a metric abelian group (A, d) , we denote by \widehat{A} the abelian group of continuous characters, *i.e.* of continuous group homomorphisms $A \rightarrow U(1)$, where A is endowed with the topology associated to the metric d . We retain the notations of section 3. Next statement is motivated by Lemma 6.1.

Proposition 6.2 *Let (A, d) be an abelian group endowed with a translation invariant metric d .*

- (i) For $\lambda, \mu \in \mathbb{R}_{>0}$, the $\mathbb{S}[\pm 1]$ -module $\underline{\text{Hom}}_{\Gamma\mathcal{T}_*}((A, d)_\lambda, U(1)_\mu)$ is isomorphic to the sub $\mathbb{S}[\pm 1]$ -module of $H\widehat{A}$ which, on the set k_+ , is given by k -tuples (χ_j) , $1 \leq j \leq k$ of continuous characters $\chi_j \in \widehat{A}$ such that, with $|x| := d(x, 0)$, and for all finite collections $\{x_i\} \subset A$, fulfill

$$\sum_i |x_i| \leq \lambda \Rightarrow \sum_{i,j} |\chi_j(x_i)| \leq \mu. \quad (33)$$

- (ii) Let $p : A' \rightarrow A$ be a surjective morphism of abelian groups and let $p^*(A, d)_\lambda$ be the pullback as in Proposition 3.6. One has the following canonical isomorphism

$$\underline{\text{Hom}}_{\Gamma\mathcal{T}_*}(p^*(A, d)_\lambda, U(1)_\mu) \simeq \underline{\text{Hom}}_{\Gamma\mathcal{T}_*}((A, d)_\lambda, U(1)_\mu). \quad (34)$$

Proof (i) Let $\phi \in \underline{\text{Hom}}_{\Gamma\mathcal{T}_*}((A, d)_\lambda, U(1)_\mu)(1_+) = \text{Hom}_{\Gamma\mathcal{T}_*}((A, d)_\lambda, U(1)_\mu)$. By applying the forgetful functor $\mathcal{F} : \mathcal{T} \rightarrow \mathfrak{Sets}$ which associates to (X, \mathcal{R}) the set X (see Proposition 3.3), one obtains an element $\mathcal{F}(\phi) \in \text{Hom}_{\mathbb{S}}(HA, HU(1))$. Since the Eilenberg-MacLane functor H determines a full and faithful embedding of the category of abelian groups inside the category of \mathbb{S} -modules, there exists a unique group homomorphism $\chi : A \rightarrow U(1)$ such that $H(\chi) = \mathcal{F}(\phi)$. The condition that ϕ preserves the relation \mathcal{R}_k on the set k_+ means that for any $x_i, y_i \in A$, $1 \leq i \leq k$ such that $\sum d(x_i, y_i) \leq \lambda$ one has $\sum d(\chi(x_i), \chi(y_i)) \leq \mu$. By translation invariance of the metrics this condition is equivalent to

$$\sum_i |x_i| \leq \lambda \Rightarrow \sum_i |\chi(x_i)| \leq \mu. \quad (35)$$

This shows that $\text{Hom}_{\Gamma\mathcal{T}_*}((A, d)_\lambda, U(1)_\mu)$ consists exactly of the group homomorphisms $\chi : A \rightarrow U(1)$ fulfilling (33). Specializing (35) to the case where all $x_i = x$, $i = 1, \dots, n$, one obtains the implication $|x| \leq \lambda/n \Rightarrow |\chi(x)| \leq \mu/n$, and hence that χ is uniformly continuous. Let $\phi \in \underline{\text{Hom}}_{\Gamma\mathcal{T}_*}((A, d)_\lambda, U(1)_\mu)(k_+) = \text{Hom}_{\Gamma\mathcal{T}_*}((A, d)_\lambda, U(1)_\mu(k_+ \wedge -))$. The object $U(1)_\mu(k_+ \wedge -)$ of $\Gamma\mathcal{T}_*$ is $(U(1)^{\times k}, d_k)_\mu$, where the metric d_k on the product group $(U(1)^{\times k})$ is defined by

$$d_k((x_i), (y_i)) := \sum_1^k d(x_i, y_i) \quad \forall x_i, y_i \in U(1).$$

Replacing $U(1)_\mu$ with $(U(1)^{\times k}, d_k)_\mu$ in the first part of the proof one obtains that $\mathcal{F}(\phi) \in \text{Hom}_{\mathbb{S}}(HA, HU(1)^{\times k}) = H((\chi_j))$, where (χ_j) , $1 \leq j \leq k$ is a k -tuple of characters of A fulfilling (33). It follows that $\chi_j \in \widehat{A}$.

(ii) Let $\phi \in \underline{\text{Hom}}_{\Gamma\mathcal{T}_*}(p^*(A, d)_\lambda, U(1)_\mu)(1_+) = \text{Hom}_{\Gamma\mathcal{T}_*}(p^*(A, d)_\lambda, U(1)_\mu)$. As in the proof of (i), there exists a group homomorphism $\chi' : A' \rightarrow U(1)$ such that $H(\chi') = \mathcal{F}(\phi)$. Moreover χ' preserves the relation \mathcal{R}_k for any k , and this implies

$$\sum_{i=1}^k |p(x'_i)| \leq \lambda \Rightarrow \sum_{i=1}^k |\chi'(x'_i)| \leq \mu, \quad \forall (x'_i) \in (A')^{\times k}.$$

In particular, taking all $x'_i = x' \in \ker(p)$ one obtains $|\chi'(x')| \leq \mu/k \forall k$, and hence $\chi'(\ker(p)) = \{1\}$. This implies that there exists a group homomorphism $\chi : A \rightarrow U(1)$ such that $\chi' = \chi \circ p$. \square

We can now state and prove Serre's duality.

Theorem 6.3 *Let $D = \sum_j a_j \{p_j\} + a\{\infty\}$ be an Arakelov divisor on $\overline{\text{Spec } \mathbb{Z}}$. There is a canonical isomorphism of $\mathbb{S}[\pm 1]$ -modules*

$$H^0(K - D) \simeq \underline{\text{Hom}}_{\Gamma\mathcal{T}_*}(H^1(D), U(1)_{\frac{1}{4}}), \quad (36)$$

where K is the divisor $K = -2\{2\}$.

Proof By Proposition 3.7, with $\lambda = \exp a$, one has $H^1(D) = \pi^*((\mathbb{R}/L, d)_\lambda)$ and by Proposition 6.2, (ii), one gets the isomorphism

$$\underline{\text{Hom}}_{\Gamma\mathcal{T}_*}(H^1(D), U(1)_{\frac{1}{4}}) \simeq \underline{\text{Hom}}_{\Gamma\mathcal{T}_*}((\mathbb{R}/L, d)_\lambda, U(1)_{\frac{1}{4}}).$$

In fact, we can assume that $D = a\{\infty\}$, with $a = \deg D$ so that $L = \mathbb{Z}$. Then we apply Proposition 6.2 (i), with $A = \mathbb{R}/\mathbb{Z}$ and $\mu = \frac{1}{4}$. One has $\widehat{A} = \mathbb{Z}$ and the characters $\chi_n \in \widehat{A}$ are given by multiplication by n , i.e. $\chi_n(s) := ns \in \mathbb{R}/\mathbb{Z}$, $\forall s \in \mathbb{R}/\mathbb{Z}$. Next, we need to determine the $\mathbb{S}[\pm 1]$ -submodule of $H\widehat{A} = H\mathbb{Z}$ which, on the set k_+ , is given by k -tuples (n_j) , $1 \leq j \leq k$ of characters $n_j \in \mathbb{Z}$ such that (33) holds. This means, using the distance d on \mathbb{R}/\mathbb{Z} , that

$$\sum_i d(x_i, 0) \leq \lambda \Rightarrow \sum_{i,j} d(n_j x_i, 0) \leq \frac{1}{4}. \quad (37)$$

The distance $d(x, 0)$ is given, for any $x' \in \mathbb{R}$ in the class of x by the distance between x' and the closed subset $\mathbb{Z} \subset \mathbb{R}$. Thus for any integer n one has: $d(nx, 0) \leq |n|d(x, 0)$. Assume that $\sum_j |n_j| \leq \frac{1}{4\lambda}$, then (37) follows since

$$\sum_{i,j} d(n_j x_i, 0) \leq \sum_{i,j} |n_j| d(x_i, 0) \leq \frac{1}{4\lambda} \sum_i d(x_i, 0).$$

Conversely, assume (37). Then repeating m times the same x , gives

$$d(x, 0) \leq \frac{\lambda}{m} \Rightarrow \sum_j d(n_j x, 0) \leq \frac{1}{4m}.$$

Taking m large enough and $x = \frac{\lambda}{m}$ one obtains $\sum_j |n_j| x \leq \frac{1}{4m}$, and hence $\sum_j |n_j| \leq \frac{1}{4\lambda}$. This proves that the $\mathbb{S}[\pm 1]$ -submodule of $H\widehat{A} = H\mathbb{Z}$ determined by (33) is equal to $\|H\mathbb{Z}\|_{\frac{1}{4\lambda}}$ which gives (36). \square

Appendix A The Γ -space $\mathbf{H}(D)$

Let $\phi : A \rightarrow B$ be a morphism of abelian groups. To ϕ one associates the following (short) complex $\mathfrak{C} = \{C_n, \phi_n\}$ of abelian groups indexed in non-negative degrees

$$\mathfrak{C} = \{C_1 \xrightarrow{\phi_1} C_0\}; \quad C_0 = B, \quad C_1 = A, \quad \phi_1(x) := \phi(x). \quad (A1)$$

The Dold-Kan correspondence associates to \mathfrak{C} the simplicial abelian group (see [7] III.2, Proposition 2.2)

$$\mathcal{A}_n = \bigoplus_{\mathcal{F}(n,k)} C_k, \quad \mathcal{F}(n, k) := \{\sigma \in \text{Hom}_\Delta([n], [k]) \mid \sigma([n]) = [k]\} \quad (A2)$$

where Δ denotes the simplicial category. The direct sum in (A2) repeats the term C_k of the chain complex as many times as the number of elements of the set $\mathcal{F}(n, k)$ of surjective morphisms $\sigma \in \text{Hom}_\Delta([n], [k])$. For the short complex \mathfrak{C} , the allowed values of k are $k = 0, 1$. Therefore the set $\mathcal{F}(n, 0)$ is reduced to $\text{Hom}_\Delta([n], [0]) =: \Delta_n^0$, *i.e.* to a single point. A morphism $\xi \in \text{Hom}_\Delta([n], [1]) =: \Delta_n^1$ is characterized by $\xi^{-1}(\{1\})$ which is an hereditary subset of $[n]$. It follows that the vertices in Δ_n^1 are labelled by the

$$\xi_j, \quad 0 \leq j \leq n+1, \quad \xi_j(k) = 1 \iff k \geq j. \quad (\text{A3})$$

For each integer $n \geq 0$ the finite set $F(n) := \mathcal{F}(n, 1)$ of surjective elements $\xi \in \Delta_n^1$ excludes ξ_0 and ξ_{n+1} , thus the set $F(n) = \{\xi_j, 1 \leq j \leq n\}$ has n elements. This gives the identification

$$\mathcal{A}_n = HB(1_+) \oplus HA(F(n)_+) \quad (\text{A4})$$

We refer to [7] (III.2, pp. 160-161) for a detailed description of the simplicial structure, namely the definition for each $\theta \in \text{Hom}_\Delta([m], [n])$ of a map of sets $\mathcal{A}(\theta) : \mathcal{A}_n \rightarrow \mathcal{A}_m$. Next, we introduce some notations.

We identify the opposite Δ^o of the simplicial category with (the skeleton of) the category of finite intervals. An interval is a totally ordered set with the smallest element distinct from the largest one. A morphism of intervals is a non-decreasing map that preserves the smallest and largest elements. The canonical contravariant functor $\Delta \rightarrow \Delta^o$ which identifies the opposite category of Δ with Δ^o (described by intervals as above), maps the finite ordinal object $[n] = \{0, 1, \dots, n\}$ in Δ to the interval $[n]^* = \{0, \dots, n+1\}$.

We denote by $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_{2,*}$ the category of pairs of pointed sets (X, Y) , with $X \supset Y$. The morphisms are maps of pairs of pointed sets. We let $c : \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_{2,*} \rightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_*$ be the functor $(X, Y) \rightarrow X/Y$ of collapsing Y to the base point $*$.

Let $\partial : \Delta^o \rightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_{2,*}$ be the functor that replaces an interval I with the pair $\partial I = (X, Y)$, where X is the set I pointed by its smallest element, and $Y \subset X$ is the subset formed by the smallest and largest elements of I .

Finally, we denote by $\gamma : \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_{2,*} \rightarrow \Gamma\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_{2,*}$ (see section 3) the functor that associates to an object (X, Y) of $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_{2,*}$ the covariant functor

$$\gamma(X, Y) : \Gamma^o \rightarrow \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_{2,*}, \quad k_+ := \{0, \dots, k\} \mapsto (X \wedge k_+, Y \wedge k_+).$$

The following formula defines a covariant functor that associates to a pair of pointed sets an abelian group directly related to the morphism $\phi : A \rightarrow B$

$$H_\phi : \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_{2,*} \rightarrow \mathfrak{A}\mathfrak{b}, \quad H_\phi(X, Y) := HB(Y) \times HA(X/Y). \quad (\text{A5})$$

On morphisms $f : (X, Y) \rightarrow (X', Y')$ in $\mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_{2,*}$ with $\alpha = (\alpha_Y, \alpha_{X/Y}) \in H_\phi(X, Y)$, the functor H_ϕ acts as follows

$$\begin{aligned} H_\phi(f)(\alpha) &= \left(\alpha_{Y'}, HA(f)(\alpha_{X/Y}) \right) \\ \alpha_{Y'}(y') &:= HB(f)(\alpha_Y)(y') + \sum_{x \in X \setminus Y, f(x)=y'} \phi(\alpha_{X/Y}(x)), \quad y' \neq *. \end{aligned} \quad (\text{A6})$$

By [3] (Proposition 4.5), the Dold-Kan correspondence for the short complex (A1), *i.e.* the simplicial abelian group \mathcal{A}_n in (A4) is canonically isomorphic to the composite functor $H_\phi \circ \partial : \Delta^o \rightarrow \mathfrak{A}\mathfrak{b}$, with H_ϕ as in (A5). By composing H_ϕ with the Eilenberg-MacLane functor H one obtains a covariant functor $HH_\phi = H \circ H_\phi : \mathfrak{S}\mathfrak{e}\mathfrak{t}\mathfrak{s}_{2,*} \rightarrow \mathbb{S}\text{-Mod}$ to the category of \mathbb{S} -modules, which is naturally isomorphic to the functor

$((U \circ H_\phi) \times \text{id}) \circ \gamma$, where $U : \mathfrak{Ab} \rightarrow \mathfrak{S}\mathfrak{ets}_*$ is the forgetful functor (see *op.cit.* Lemma 4.6). Moreover, as shown in *op.cit.* (Theorem 4.7) the Γ -space associated by the Dold-Kan correspondence to the complex \mathfrak{C} is canonically isomorphic to the functor

$$((U \circ H_\phi) \times \text{id}) \circ \gamma \circ \partial : \Delta^o \rightarrow \mathbb{S} - \text{Mod}. \quad (\text{A7})$$

Notice that the above construction involves the morphism $\phi : A \rightarrow B$ *only* through the composite functor $U \circ H_\phi : \mathfrak{S}\mathfrak{ets}_{2,*} \rightarrow \mathfrak{S}\mathfrak{ets}_*$. This latter functor is still meaningful when one restricts ϕ to a sub- \mathbb{S} -module E of the \mathbb{S} -module HA and it is given by

$$H_\phi|_E : \mathfrak{S}\mathfrak{ets}_{2,*} \rightarrow \mathfrak{S}\mathfrak{ets}_*, \quad H_\phi(X, Y) := HB(Y) \times E(X/Y). \quad (\text{A8})$$

This provides the following construction

Definition A.1 *Let $\phi : A \rightarrow B$ be a morphism of abelian groups and E a sub \mathbb{S} -module of the \mathbb{S} -module HA . We denote by $\Gamma(\phi|_E)$ the Γ -space obtained as a sub-functor of (A7)*

$$\Gamma(\phi|_E) := (H_\phi|_E \times \text{id}) \circ \gamma \circ \partial : \Delta^o \rightarrow \mathbb{S} - \text{Mod}. \quad (\text{A9})$$

When evaluated on the object $1_+ = \{0, 1\}$ of Γ^o , the Γ -space $\Gamma(\phi|_E)$ defines a sub-simplicial set of the Kan simplicial set \mathcal{A}_n in (A4). However it is not in general a Kan simplicial set, thus one needs to exert care when considering its homotopy. We refer to [2] 2.1, for the definition of the homotopies used there. The relation \mathcal{R} of homotopy between n -simplices $(x, y) \in X_n \times X_n$ is defined as follows

$$(x, y) \in \mathcal{R} \iff \partial_j x = \partial_j y \forall j \ \& \ \exists z \text{ s.t. } \partial_j z = s_{n-1} \partial_j x \forall j < n, \ \partial_n z = x, \ \partial_{n+1} z = y. \quad (\text{A10})$$

In general, the relation \mathcal{R} in (A10) fails to be an equivalence relation. In place of the quotient we consider pairs of sets and relations. We define $\pi_n^{\mathcal{T}}$ to be the set of spherical elements in X_n (*i.e.* of n -simplices x , with $\partial_j x = * \forall j$), endowed with the relation \mathcal{R} . Then we have the following result (we refer to section 3 for the notion of tolerant module)

Proposition A.2 *Let $\phi : A \rightarrow B$ be a morphism of abelian groups and let E be a sub- $\mathbb{S}[\pm 1]$ -module of the Eilenberg-MacLane $\mathbb{S}[\pm 1]$ -module HA (where $(\pm 1)x := \pm x$).*

(i) *The homotopy $\pi_1^{\mathcal{T}}(\Gamma(\phi|_E))$ is the sub- $\mathbb{S}[\pm 1]$ -module $\ker(H\phi|_E)$ of E*

$$\ker(H\phi|_E)(k_+) = \{x \in E(k_+) \mid H\phi(x) = *\}. \quad (\text{A11})$$

(ii) *The homotopy $\pi_0^{\mathcal{T}}(\Gamma(\phi|_E))$ is the tolerant $\mathbb{S}[\pm 1]$ -module HB endowed with the relations*

$$\mathcal{R}_k = \{(x, y) \in HB(k_+) \times HB(k_+) \mid x - y \in H\phi(E(k_+))\} \quad k \in \mathbb{N}. \quad (\text{A12})$$

Proof By construction, $\Gamma(\phi|_E)$ is the composite of the functors $H_\phi|_E : \mathfrak{S}\mathfrak{ets}_{2,*} \rightarrow \mathfrak{S}\mathfrak{ets}_*$ of (A8) and $\gamma \circ \partial : \Delta^o \rightarrow \mathfrak{S}\mathfrak{ets}_{2,*}$. One has $\partial[0]^* = (X, Y)$ where $X = Y =$

$\{0, 1\}$ with base point 0. Thus $\gamma \circ \partial[0]^*(k_+) = (X \wedge k_+, Y \wedge k_+) = (k_+, k_+)$ and by (A8)

$$H_\phi|_E(\gamma \circ \partial[0]^*)(k_+) = H_\phi|_E(X \wedge k_+, Y \wedge k_+) = HB(k_+).$$

One has $\partial[1]^* = (X, Y)$, where $X = \{0, 1, 2\}$, $Y = \{0, 2\}$ with base point 0. Thus one has $\gamma \circ \partial[1]^*(k_+) = (X \wedge k_+, Y \wedge k_+)$ and $Y \wedge k_+ = k_+$, while $(X \wedge k_+)/ (Y \wedge k_+) = \{0, 1\} \wedge k_+ = k_+$ and by (A8) it follows that

$$H_\phi|_E(\gamma \circ \partial[1]^*)(k_+) = E(k_+) \times HB(k_+). \quad (\text{A13})$$

The boundaries $\partial_j : H_\phi|_E(\gamma \circ \partial[1]^*) \rightarrow H_\phi|_E(\gamma \circ \partial[0]^*)$ are obtained as in [3] Proposition 4.11

$$\partial_0(\psi) = \psi_2, \quad \partial_1(\psi) = H\phi(\psi_1) + \psi_2, \quad \forall \psi = (\psi_1, \psi_2) \in H_\phi|_E(\gamma \circ \partial[1]^*). \quad (\text{A14})$$

(i) The spherical condition $\partial_j(\psi) = *$ on $\psi \in H_\phi|_E(\gamma \circ \partial[1]^*)(k_+)$ means that $\psi_2 = 0$ and $H\phi(\psi_1) = 0$. Thus the solutions correspond to $\ker(H\phi|_E)(k_+)$ as in (A11). One shows as in [3] Proposition 4.11 (iii), that the relation of homotopy is the identity.

(ii) The relation \mathcal{R}_k on elements of $H_\phi|_E(\gamma \circ \partial[0]^*)(k_+)$ is defined as follows

$$(\alpha, \beta) \in \mathcal{R}_k \iff \exists \psi \in H_\phi|_E(\gamma \circ \partial[1]^*)(k_+) \text{ s.t. } \partial_0(\psi) = \alpha \ \& \ \partial_1(\psi) = \beta. \quad (\text{A15})$$

By (A14) it coincides with (A12).

Note that this relation is a tolerance relation *i.e.* is symmetric since E is a sub $\mathbb{S}[\pm 1]$ -module of the $\mathbb{S}[\pm 1]$ -module HA , so that

$$x - y \in H\phi(E(k_+)) \iff y - x \in H\phi(E(k_+)).$$

□

A.1 The Γ -space $\mathbf{H}(D)$

The above general construction applies in the geometric adelic context and by implementing the action of \mathbb{Q}^\times on adeles, we obtain the following variant of Proposition 4.9 in [3].

Proposition A.3 *Let D be an Arakelov divisor on $\overline{\text{Spec } \mathbb{Z}}$. Let $A = \mathbb{Q} \times \mathbb{A}_{\mathbb{Q}}$, $B = \mathbb{A}_{\mathbb{Q}}$ and $\alpha : A \rightarrow B$, $\alpha(q, a) = q + a$. Let $E = H\mathbb{Q} \times H\mathcal{O}(D)$ be the sub $\mathbb{S}[\pm 1]$ -module of HA as in Proposition 2.1. Then the functor*

$$\mathbf{H}(D) := \Gamma(\alpha|_E) : \Delta^\circ \longrightarrow \mathbb{S} - \text{Mod} \quad (\text{A16})$$

defines a Γ -space that depends only on the linear equivalence class of D .

A.2 Proof of Proposition 2.2

With the notations of section 3 and Proposition A.3, one has $H^0(D) := \pi_1^T(\mathbf{H}(D)) = \pi_1^T(\Gamma(\alpha|_E))$. Proposition A.2, (i), gives

$$H^0(D)(k_+) = \{x \in E(k_+) \mid H\alpha(x) = *\}.$$

An element $x \in E(k_+)$ is a k -tuple (q_j, a_j) with $q_j \in \mathbb{Q}$ for all j , and $(a_j) \in H\mathcal{O}(D)(k_+)$, where $H\mathcal{O}(D) := H\mathcal{O}(D)_f \times \|H\mathbb{R}\|_{e^\alpha}$. The condition $H\alpha(x) = *$ means $q_j = -a_j$ for all j , so that x is uniquely determined by the k -tuple (q_j) . Moreover, the allowed k -tuples are those for which $\sum |q_j| \leq e^\alpha$ and $q_j \in \mathcal{O}(D)_f$ for all j . One has

$$\mathbb{Q} \cap \mathcal{O}(D)_f = H^0(\text{Spec } \mathbb{Z}, \mathcal{O}(D)_f).$$

This proves (i) of Proposition 2.2. The statement (ii) follows from Proposition A.2 (ii). Finally, (iii) follows from Proposition A.3.

Acknowledgments. The second author is partially supported by the Simons Foundation collaboration grant n. 691493 and thanks IHES for the hospitality, where part of this research was done.

References

- [1] A. Connes, C. Consani, *Absolute algebra and Segal's Gamma sets*, J. Number Theory 162 (2016), 518–551.
- [2] A. Connes, C. Consani, $\overline{\text{Spec } \mathbb{Z}}$ and the Gromov norm, Theory and Applications of Categories, 35, No. 6, (2020), pp. 155–178.
- [3] A. Connes, C. Consani, *Segal's Gamma rings and universal arithmetic*, The Quarterly Journal of Mathematics, 72, 1-2, (2021), pp. 1–29
- [4] A. Connes, C. Consani, *On Absolute Algebraic Geometry, the affine case*, Advances in Mathematics, 390, Paper No. 107909 (2021), 44 pp.
- [5] Dundas, Bjorn Ian; Goodwillie, Thomas G.; McCarthy, Randy *The local structure of algebraic K-theory*, Algebra and Applications, 18. Springer-Verlag London, Ltd., London, 2013.
- [6] *Correspondance Grothendieck-Serre*. Edited by Pierre Colmez and Jean-Pierre Serre. Documents Mathématiques (Paris), 2. Société Mathématique de France, Paris, 2001. xii+288 pp.
- [7] Goerss, Paul G.; Jardine, John F. *Simplicial homotopy theory*. Progress in Mathematics, 174. Birkhauser Verlag, Basel, 1999.
- [8] S. Lang, *Algebraic Number Theory*. Addison Wesley, 1970.
- [9] J. Neukirch, *Algebraic number theory*. Translated from the 1992 German original and with a note by Norbert Schappacher. With a foreword by G. Harder. Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 322. Springer-Verlag, Berlin, 1999.
- [10] A. Weil *Sur l'analogie entre les corps de nombres algébriques et les corps de fonctions algébriques*, Oeuvres scientifiques/Collected papers I. 1926–1951. Springer, Heidelberg, 2014.
- [11] A. Weil *Basic number theory*. Reprint of the second (1973) edition. Classics in Mathematics. Springer-Verlag, Berlin, 1995
- [12] <http://solbakkn.com/math/triadic-nums.htm>

Alain Connes
Collège de France and IHES
alain@connes.org

Caterina Consani
Johns Hopkins University
cconsan1@jhu.edu