

# D-FINITENESS, RATIONALITY, AND HEIGHT II: LOWER BOUNDS OVER A SET OF POSITIVE DENSITY

JASON P. BELL, KHOA D. NGUYEN, AND UMBERTO ZANNIER

ABSTRACT. We consider D-finite power series  $f(z) = \sum a_n z^n$  with coefficients in a number field  $K$ . We show that there is a dichotomy governing the behaviour of  $h(a_n)$  as a function of  $n$ , where  $h$  is the absolute logarithmic Weil height. As an immediate consequence of our results, we have that either  $f(z)$  is rational or  $h(a_n) > [K : \mathbb{Q}]^{-1} \cdot \log(n) + O(1)$  for  $n$  in a set of positive upper density and this is best possible when  $K = \mathbb{Q}$ .

## 1. INTRODUCTION

Within number theory, combinatorics, and other disciplines, it is often of use to understand the asymptotics of sequence of complex numbers; that is, to describe how the sequence is growing. This approach is especially fruitful when trying to gain insight into the behaviour of integer-valued sequences satisfying recurrences. When considering non-integer-valued sequences, however, knowing the actual size (or, more precisely, modulus) is typically of less importance and one often uses other measures that serve as proxies for how the values of a sequence  $(a_n)$  increase in complexity with  $n$ . For number fields, one generally uses the notion of *heights* (see §2.1 for relevant definitions) as a measure of the complexity of elements of the field, and interprets numbers with smaller height as being simpler, or less complex, than numbers with larger height.

In the paper [BNZ20], improving results of van der Poorten-Shparlinski [vdPS96] and Bell-Chen [BC17], we considered exactly these sorts of questions, looking at the growth of heights of coefficients of multivariate D-finite power series with coefficients in a number field  $K$ . These are power series  $F(z_1, \dots, z_d) \in K[[z_1, \dots, z_d]]$  in variables  $z_1, \dots, z_d$  that satisfy non-trivial homogeneous linear differential equations with polynomial coefficients with respect to each differential operator  $\partial/\partial z_i$ . In this case we found that if the heights of the coefficients of  $F$  do not grow too quickly then  $F$  is necessarily a rational function of a special form. In simpler terms, if the complexity of the coefficients of  $F$  grows sufficiently slowly then  $F$  is well behaved. More precisely, we showed that if

$$f(z_1, \dots, z_d) = \sum a_{n_1, \dots, n_d} z_1^{n_1} \cdots z_d^{n_d} \in K[[x]]$$

and the logarithmic Weil height of  $a_{n_1, \dots, n_d}$  is  $o(\log \|\mathbf{n}\|)$ , with  $\|\mathbf{n}\| := n_1 + \cdots + n_d$ , as  $\|\mathbf{n}\| \rightarrow \infty$  then  $f(z_1, \dots, z_d)$  is the power series expansion of a rational function and the coefficients  $a_{n_1, \dots, n_d}$  belong to a finite set. This result suggests, then, that

---

*Date:* April 2022.

*2010 Mathematics Subject Classification.* Primary: 13F25, 12H05. Secondary: 11G50.

*Key words and phrases.* D-finite power series, heights, rational functions,

Jason Bell was supported by NSERC grant RGPIN-2016-03632; Khoa Nguyen was supported by NSERC grant RGPIN-2018-03770 and CRC tier-2 research stipend 950-231716.

there is a type of “height boundary” for coefficients of multivariate D-finite series, given by the function  $\log \|\mathbf{n}\|$ , where if the coefficients have heights that are small compared to this boundary function then the power series is rational and the height growth of the coefficients falls into the next level, namely  $h(a_{n_1, \dots, n_d}) = O(1)$ .

This paper provides a more refined result toward a complete classification of the height growth of the coefficients of a D-finite univariate power series in  $\bar{\mathbb{Q}}[z]$ . These are exactly power series solutions of linear homogeneous differential equations with coefficients in  $\bar{\mathbb{Q}}(z)$  and play an important role in many areas of mathematics. While the coefficients of the power series of a rational function in  $\bar{\mathbb{Q}}(z)$  satisfy a linear recurrence relation with constant coefficients and are rather well understood, the coefficients of a D-finite series satisfy a linear recurrence relation with polynomial coefficients and remain quite mysterious (see the questions in Section 4). Although the authors obtained most of the results in this paper in the fall of 2019 shortly after [BNZ20], its release has been delayed due to various reasons. In the meantime, [BNZ20] motivates several highly interesting results in number theory and dynamics. In arithmetic dynamics, [BNZ20] motivates a height gap conjecture in work of Bell, Hu, Ghioca, and Satriano [BHS20, BGS21] concerning the quantity  $\limsup_{n \rightarrow \infty} h(f(\Phi^n(x)))/\log n$ , where  $\Phi : X \dashrightarrow X$  and  $f : X \dashrightarrow \mathbb{P}^1$  are rational maps on quasi-projective varieties defined over  $\bar{\mathbb{Q}}$  and  $x \in X(\bar{\mathbb{Q}})$  is such that the forward orbit of  $x$  under  $\Phi$  is well defined and avoids the indeterminacy locus of  $f$ . In the theory of Mahler functions, Adamczewski, Bell, and Smertnig [ABS] provide a complete classification of the possible height growth of the coefficients of a Mahler function. The paper [BNZ20] together with earlier work of Perelli and Zannier [PZ84, Zan96] motivate [BN21] in which an analogue of a conjecture of Ruzsa for polynomials over finite fields is established. Finally [BNZ20, BN21] and the discussion related to results in this paper inspire Dimitrov’s spectacular solution [Dim] of the Schinzel-Zassenhaus conjecture from the 1960s.

For an algebraic number  $\alpha$ , we define its denominator  $\text{den}(\alpha)$  to be the smallest positive integer  $d$  such that  $d\alpha$  is an algebraic integer. We recall that a subset  $S$  of  $\mathbb{N}$  is said to have positive (upper) density if

$$\limsup_{n \rightarrow \infty} \frac{|S \cap [1, n]|}{n} > 0;$$

otherwise  $S$  is said to have zero density. Finally, we use Vinogradov notation  $\gg$  and  $\ll$ , where  $a_n \gg b_n$  for  $n$  in a set  $T$  means that there is a positive constant  $C$  such that  $a_n \geq Cb_n$  for  $n \in T$ , with  $\ll$  defined analogously. Our main result gives a coarse classification of the growth of  $h(a_n)$ , where

$$f(z) = \sum_{n=0}^{\infty} a_n z^n$$

is a D-finite power series with algebraic coefficients,  $n$  belongs to a set of positive density, and where  $h : K \rightarrow [0, \infty)$  is the absolute logarithmic Weil height.

**Theorem 1.1.** *Let  $K$  be a number field and let  $f(z) = \sum_n a_n z^n \in K[[z]]$  be a D-finite power series. Let  $r$  be the radius of convergence of  $f$ , we have:*

- (a) *If  $r \in \{0, \infty\}$  and  $f$  is not a polynomial then  $h(a_n) = O(n \log n)$  as  $n$  tends to infinity and  $h(a_n) \gg n \log n$  on a set of positive upper density.*
- (b) *If  $r \notin \{0, \infty\}$  then at least one of the following holds:*
  - (i)  *$h(a_n) \gg n$  on a set of positive upper density;*

- (ii)  $\text{den}(a_n) \gg n$ , and hence  $h(a_n) > \frac{1}{[K : \mathbb{Q}]} \log n + O(1)$ , on a set of positive upper density;
- (iii)  $f(z)$  is a rational function whose denominator divides  $(1 - z^M)^N$  for some nonnegative integers  $M, N$  (i.e. all of the finite poles of  $f$  are roots of unity).

*Remark 1.2.* In fact, we prove a more precise version of Theorem 1.1 (see Theorem 3.9 and Lemma 3.4). When (iii) holds, we have  $a_n = P(n)$  for all large  $n$  in an appropriate arithmetic progression where  $P$  is a polynomial. In this case, we still have  $h(a_n) > \log n + O(1)$  on a set of positive density unless  $P$  is constant, in which case the sequence  $(a_n)_{n \geq 0}$  is eventually periodic.

Theorem 1.1 immediately yields the following result.

**Corollary 1.3.** *Let  $K$  and  $f$  be as in Theorem 1.1. Suppose that  $f$  is not a rational function, then*

$$h(a_n) > \frac{1}{[K : \mathbb{Q}]} \log n + O(1)$$

on a set of positive upper density.

*Remark 1.4.* It follows from [BNZ20] that  $\limsup h(a_n)/\log n > 0$  or in other words there exist  $C > 0$  and an infinite  $S \subseteq \mathbb{N}$  depending on  $f$  such that  $h(a_n) > C \log n$  for  $n \in S$ . Corollary 1.3 improves this in two aspects: one may require that  $S$  has positive density and take the uniform constant  $C = 1/[K : \mathbb{Q}]$  up to the additional  $O(1)$  term. When  $K = \mathbb{Q}$ , we have the lower bound  $h(a_n) > \log n + O(1)$  on a set of positive density. This is best possible thanks to examples such as

$$\log(1 + z^m) = z^m - \frac{z^{2m}}{2} + \frac{z^{3m}}{3} - \dots,$$

which show both that we cannot improve the constant and that the density of the set  $S$  can be arbitrarily close to zero. For a general number field  $K$ , it is plausible that the lower bound  $\log n + O(1)$  remains valid (i.e., the factor  $1/[K : \mathbb{Q}]$  in the statement of Corollary 1.3 (ii) can be improved to 1) but our method has not been able to yield this. We leave this as a question for future work (see Question 4.2).

The outline of this paper is as follows. In §2, we give an overview of Weil heights and the technical ingredients in the proof of Theorem 1.1. This includes the use of Hankel determinants, Pólya's inequality, and diophantine approximation results when approximating a D-finite series by a rational function. In §3, we first prove Theorem 3.9 which is a slightly more precise variant of Theorem 1.1 then obtain the proof of the latter. Besides the technical results introduced in §2, our method involves the construction of an auxiliary polynomial  $P(z)$  and work with the power series  $\sum P(n)a_n z^n$  instead of the original series  $f(z) = \sum a_n z^n$ . Right after the completion of this paper, a variant of our method has been extended to obtain a general criterion for the Pólya-Carlson dichotomy and its application to the Artin-Mazur zeta function associated to endomorphisms on positive characteristic tori [BGNS]. In §4 we provide some comments to results in the paper. This includes a demonstration that one cannot, in a certain sense, strengthen Theorem 1.1 in the style of Theorem 3.9 and several open problems for future work. These problems include a complete classification of the possible growth of  $h(a_n)$  for a D-finite series  $\sum a_n z^n \in \mathbb{Q}[[z]]$  as well as a similar long standing open problem in the theory of Siegel E-functions.

## 2. PRELIMINARY RESULTS

**2.1. Heights.** We give a quick overview of heights along with the results we will need. For a number field  $K$ , we let  $M_K^\infty$  denote the set of archimedean places (equivalence classes of archimedean absolute values) of  $K$  and we let  $M_K^0$  denote the set of finite places. We write  $M_K = M_K^\infty \cup M_K^0$ . For every place  $w \in M_K$ , let  $K_w$  denote the completion of  $K$  with respect to  $w$  and we let  $d(w)$  denote the quantity  $[K_w : \mathbb{Q}_v]$  where  $v$  is the restriction of  $w$  to  $\mathbb{Q}$ .

We now follow the treatment given in [BG06, Chapter 1]: for every  $w \in M_K$  with restriction  $v$  on  $\mathbb{Q}$ , we may always take  $|\cdot|_v$  to be either the ordinary Euclidean absolute value or the  $p$ -adic absolute value for some prime  $p$ . We can then normalize  $|\cdot|_w$  by defining

$$|x|_w = |\mathrm{N}_{K_w/\mathbb{Q}_v}(x)|_v^{1/[K:\mathbb{Q}]}.$$

Let  $m \in \mathbb{N}$ . Given a vector  $\mathbf{u} = (u_0, \dots, u_m) \in K^{m+1} \setminus \{\mathbf{0}\}$  and  $w \in M_K$ , we can now define

$$|\mathbf{u}|_w := \max_{0 \leq i \leq m} |u_i|_w.$$

In particular, we can now define heights for points in projective varieties. For  $P \in \mathbb{P}^m(\bar{\mathbb{Q}})$ , let  $K$  be a number field such that  $P$  has a representative  $\mathbf{u} \in K^{m+1} \setminus \{\mathbf{0}\}$  and define:

$$H(P) = \prod_{w \in M_K} |\mathbf{u}|_w.$$

Define  $h(P) = \log(H(P))$ . Finally, for  $\alpha \in \bar{\mathbb{Q}}$ , we write  $H(\alpha) = H([\alpha : 1])$  and  $h(\alpha) = \log(H(\alpha))$ .

Then the height function defined has the following properties (see [HS00, Part B] or [BG06, Chapters 1–2] for proofs of these facts):

- (i) For  $a \in \bar{\mathbb{Q}}^*$  and  $m \in \mathbb{Z}$ ,  $h(a^m) = |m|h(a)$ .
- (ii) For  $r \in \mathbb{N}$  and  $a_1, \dots, a_r \in \bar{\mathbb{Q}}$ ,  $h(a_1 + \dots + a_r) \leq h(a_1) + \dots + h(a_r) + \log r$ .
- (iii) For every  $a, b \in \bar{\mathbb{Q}}$ ,  $h(ab) \leq h(a) + h(b) + \log 2$ . Hence if  $b \neq 0$  then  $h(ab) \geq h(a) - h(b) - \log 2$ .

**2.2. Hankel determinants.** Throughout this subsection, let  $k$  be a field, let  $(a_n)_{n \in \mathbb{N}_0}$  be a  $k$ -valued sequence, and let

$$f(z) = \sum_{n=0}^{\infty} a_n z^n \in k[[z]]$$

be the associated formal power series. We assume tacitly that  $f$  is nonzero. We let  $\mathrm{ord}$  denote the order function on  $k[[z]]$ . For a power series  $g \in k[[z]]$  we use the notation  $g = O(z^m)$  to mean that  $\mathrm{ord} g \geq m$ .

For integers  $\ell, m \geq 0$  we define the Hankel matrix  $H_{\ell, m}(f)$  and Hankel determinant  $\Delta_{\ell, m}(f)$  by:

$$H_{\ell, m}(f) = \begin{pmatrix} a_\ell & a_{\ell+1} & \cdots & a_{\ell+m} \\ a_{\ell+1} & a_{\ell+2} & \cdots & a_{\ell+m+1} \\ \cdots & & & \\ a_{\ell+m} & a_{\ell+m+1} & \cdots & a_{\ell+2m} \end{pmatrix}, \quad \Delta_{\ell, m}(f) := \det H_{\ell, m}(f).$$

Hankel determinants are of importance due to their connection with rationality of power series. We begin with a lemma, which illustrates this connection in terms of approximation by rational power series.

**Lemma 2.1.** *Let  $\ell$  and  $m$  be nonnegative integers. Then  $\Delta_{\ell,m}(f) = 0$  if and only if there exist polynomials  $P(z), Q(z) \in k[z]$  with  $\deg P \leq \ell + m - 1$ ,  $Q(z) \neq 0$ ,  $\deg Q \leq m$  and  $P(z) - Q(z)f(z) = O(z^{\ell+2m+1})$ .*

*Proof.* Suppose  $\Delta_{\ell,m} = 0$ , so the columns of  $H_{\ell,m}$  are linearly dependent over  $k$ ; i.e., there exist  $q_0, \dots, q_m \in k$ , not all 0, and with

$$q_0 a_{\ell+r+m} + q_1 a_{\ell+r+m-1} + \dots + q_m a_{\ell+r} = 0$$

for  $r = 0, 1, \dots, m$ .

Setting  $Q(z) := q_0 + q_1 z + \dots + q_m z^m$ , we have  $Q(z) \neq 0$  and

$$Q(z)f(z) = \sum_{s=0}^{\infty} c_s z^s, \quad c_s = q_0 a_s + q_1 a_{s-1} + \dots + q_m a_{s-m},$$

with the convention that  $a_n = 0$  for  $n < 0$ .

Note now that the above equations give that  $c_s = 0$  for  $s = \ell + m, \dots, \ell + 2m$ , hence setting  $P(z) = c_0 + c_1 z + \dots + c_{\ell+m-1} z^{\ell+m-1}$ , we have  $\deg P \leq \ell + m - 1$ ,  $\deg Q \leq m$  and  $\text{ord}(P(z) - Q(z)f(z)) \geq \ell + 2m + 1$ , as required. The converse assertion is proved by simply reversing the argument.  $\square$

In the sequel we shall let  $\Delta_m(f)$  denote the Hankel determinant  $\Delta_{0,m}(f)$ . We have the following consequence.

**Corollary 2.2.** *Let  $m$  and  $d$  be nonnegative integers. If  $\Delta_{m+i}(f) = 0$  for  $0 \leq i \leq d$  then there exist  $P(z), Q(z) \in k[z]$  with  $\deg P \leq m - 1$ ,  $Q(0) \neq 0$ ,  $\deg Q \leq m$ ,  $\gcd(P, Q) = 1$ , and  $f(z) - P(z)/Q(z) = O(z^{m+d+1})$ .*

*Proof.* For  $0 \leq i \leq d$ , Lemma 2.1 gives  $P_i(z), Q_i(z) \in k[z]$  with  $\deg P_i \leq m + i - 1$ ,  $Q_i \neq 0$ ,  $\deg Q_i \leq m + i$  and  $P_i(z) - Q_i(z)f(z) = O(z^{2m+2i+1})$ . Therefore:

$$P_i Q_{i+1} - P_{i+1} Q_i = O(z^{2m+2i+1}).$$

Since the left-hand side has degree at most  $2m + 2i$ , we have  $P_i Q_{i+1} = P_{i+1} Q_i$  for  $0 \leq i \leq d - 1$ . We have

(1)

$$f(z) - P_0(z)/Q_0(z) = f(z) - P_d(z)/Q_d(z) = \frac{1}{Q_d(z)} O(z^{2m+2d+1}) = O(z^{m+d+1}).$$

Express  $\frac{P_0}{Q_0} = \frac{P}{Q}$  with  $\gcd(P, Q) = 1$ . Let  $a = \text{ord } Q_0$  then we have  $a \leq \text{ord } P_0$ ; otherwise the order function of the LHS of (1) has a negative value. This implies  $Q(z)$  divides  $Q_0(z)/z^a$ . Hence  $Q(0) \neq 0$  and this finishes the proof.  $\square$

By letting  $d \rightarrow \infty$  in Corollary 2.2, we have the following well-known criterion for rationality by Kronecker:

**Corollary 2.3** (Kronecker's criterion).  *$f(z) \in k(z)$  if and only if  $\Delta_n(f) = 0$  for all sufficiently large  $n$ .*

**2.3. D-finite series and rational approximation.** Throughout this subsection, we let  $k$  be a field of characteristic zero and let  $f(z) \in k[[z]]$  be a (nonzero) D-finite power series over  $k(z)$ ; explicitly:

$$(2) \quad \mathcal{L}f = 0, \quad \mathcal{L} = A_0(z)D^p + A_1(z)D^{p-1} + \dots + A_p(z), \quad D = \frac{d}{dz},$$

where  $p = \dim_{k(z)} \text{Span}\{f, f', f'', \dots\}$  and  $A_i \in k[z]$  for  $0 \leq i \leq p$  with  $A_0 \neq 0$ . Let  $\delta = \max\{\deg(A_i) : 0 \leq i \leq p\}$ .

**Lemma 2.4.** *Let  $n$  be an integer with  $n \geq p-1$ , let  $P_0(z), \dots, P_n(z)$  be polynomials in  $k[z]$  of degree at most  $n$ , and let*

$$g(z) = \sum_{i=0}^n P_i(z) f^{(i)}(z).$$

If

$$\dim_{k(z)} \text{Span}\{g, g', g'', \dots\} = p,$$

then there is a  $p \times p$  matrix  $B(z)$  with rational function entries whose numerators and denominators are bounded in degree by  $(2\delta + 1)np$  such that

$$B(z)[g(z), \dots, g^{(p-1)}(z)]^T = [f(z), f'(z), \dots, f^{(p-1)}(z)]^T.$$

*Proof.* We have  $f^{(p)}(z) = -\sum_{i=0}^{p-1} (A_{p-i}(z)/A_0(z))f^{(i)}(z)$ . Then a straightforward induction shows that for  $j \geq 0$  we have

$$f^{(p+j)}(z) = \sum_{i=0}^{p-1} (A_{p-i,j}(z)/A_0(z)^{j+1})f^{(i)}(z),$$

where the  $A_{p-i,j}$ 's are polynomials of degree at most  $\delta(j+1)$ . Then for  $j = 0, \dots, p-1$  we have

$$g^{(j)}(z) = \sum_{i=0}^{p-1} C_{p-i,j}(z)/A_0(z)^{n+j-p+1} f^{(i)}(z),$$

where  $C_{p-i,j}$  has degree at most  $n + \delta(n+j-p+1)$ . Then

$$[g(z), \dots, g^{(p-1)}(z)]^T = C(z)[f(z), f'(z), \dots, f^{(p-1)}(z)]^T,$$

where  $C(z)$  is the  $p \times p$  matrix whose  $(i, j)$ -entry is  $C_{p-j,i}(z)/A_0(z)^{n+i-p+1}$  for  $0 \leq i, j \leq p-1$ . Observe that  $C(z)$  is invertible, since otherwise there would be a nonzero row vector  $w(z) \in k(z)^p$  such that  $w(z)C(z) = 0$ , which would give that  $g, g', \dots, g^{(p-1)}$  are linearly dependent over  $k(z)$ , contradicting the assumption that

$$\dim_{k(z)} \text{Span}\{g, g', g'', \dots\} = p.$$

The determinant of  $C(z)$  has the form:

$$\det(C(z)) = \frac{P(z)}{A_0(z)^{p(n-p+1)+(p(p-1)/2)}}$$

where  $P(z) \in k[z]$  with  $\deg P \leq np + \delta p(n-p+1) + \delta p(p-1)/2$ . For  $0 \leq i, j \leq p-1$ , we remove the row  $(C_{i,k})_{0 \leq k \leq p-1}$  and the column  $(C_{k,j})_{0 \leq k \leq p-1}^T$  from  $C(z)$  then the determinant of the resulting matrix has the form:

$$\frac{P_{i,j}(z)}{A_0(z)^{(p-1)(n-p+1)+(p(p-1)/2)-i}}$$

where  $P_{i,j}(z) \in k[z]$  with  $\deg P_{i,j} \leq n(p-1) + \delta(p-1)(n-p+1) + \delta((p(p-1)/2) - i)$ . Let  $B(z)$  be the inverse of the matrix  $C(z)$ . We have that each entry of  $B(z)$  is a rational function whose numerator and denominator are bounded in degree by:

$$np + \delta p(n-p+1) + \delta p(p-1)/2 \leq (2\delta + 1)np.$$

□

We conclude this subsection with a rational approximation principle. We recall that we are assuming our base field  $k$  has characteristic zero, which is a needed hypothesis in the following result.

**Lemma 2.5.** *Suppose that  $P, Q \in k[z]$  are polynomials of degree  $\leq \ell$  such that  $Q(0) \neq 0$  and  $P - Qf = O(z^N)$ , where  $N > (p + 2)\ell + \delta + p$ . Then we have  $\mathcal{L}(P/Q) = 0$ . Moreover, there is an effectively computable subset  $T$  of  $\mathbb{N}$  with  $|T| \leq p$ , depending only on  $\mathcal{L}$ , such that if  $\text{ord}(P - Qf) \notin T$  then we must have in fact  $f = P/Q$ .*

*Proof.* If  $Qf = P$  we are done, so assume this is not the case. Since  $Q(0) \neq 0$  we may write  $f - P/Q = z^m R$  with  $m = \text{ord}(P - Qf)$  and  $R \in k[[z]]$  satisfying  $R(0) \neq 0$ .

We have  $\mathcal{L}(P/Q) = -\mathcal{L}(z^m R)$ . The right-hand side is a power series of order  $\geq m - p$ . The left-hand side is a rational function whose numerator degree is bounded above by  $(p + 2)\ell + \delta$ . This numerator must be divisible by  $z^{m-p}$ , thus must vanish since  $m \geq N > (p+2)\ell + \delta + p$  by assumption. This proves  $\mathcal{L}(P/Q) = 0$ .

For the second assertion, we use that  $\mathcal{L}(z^m R) = 0$  as follows. Note that the term of lowest order in  $A_i(z)D^{p-i}(z^m R(z))$  is of the form

$$R(0)\alpha_i(p-i)! \binom{m}{p-i} z^{m-p+i+e_i},$$

where  $\alpha_i z^{e_i}$  is the term of lowest order in  $A_i(z)$  (i.e.  $e_i = \text{ord } A_i$ ). Since the sum of the terms of lowest order in

$$\sum_{i=0}^p A_i(z)D^{p-i}(z^m R)$$

vanishes, the sum of such terms with the property that  $m - p + i + e_i$  assumes the minimum value, say equal to  $\mu$ , must also vanish.

Let  $I \subseteq \{0, 1, \dots, p\}$  be the subset of indices  $i$  such that  $m - p + i + e_i = \mu$ ; note that, for  $m \geq p$ ,  $I$  does not depend on  $m$  and corresponds to  $i + e_i$  being minimal. We then have

$$\sum_{i \in I} \alpha_i(p-i)! \binom{m}{p-i} = 0.$$

Since  $\text{char}(k) = 0$ , the left side is a nonzero polynomial in  $m$ , of degree at most  $p$ , depending explicitly on the  $A_i(x)$ . This polynomial has at most  $p$  roots, and if  $m$  is outside the set of these roots we have a contradiction, which in fact proves that  $f = P/Q$ .  $\square$

**2.4. Pólya's inequality.** In this subsection, we let  $G \subsetneq \mathbb{C}$  be a simply connected domain containing 0 with conformal radius  $\rho > 1$  from the origin. This means we have a (unique) conformal map  $\varphi$  from  $G$  onto the disk  $D(0, \rho)$  with  $\varphi(0) = 0$  and  $\varphi'(0) = 1$ . Let  $r \in (1, \rho)$  and let  $\Gamma = \varphi^{-1}(\partial D(0, r))$ . A more general version of Pólya's inequality treats functions with isolated singularities, the following version for analytic functions is sufficient for our purpose:

**Theorem 2.6** (Pólya's inequality). *Let  $f(z)$  be an analytic function on  $G$  with the Taylor series  $f(z) = \sum_{n=0}^{\infty} a_n z^n$  at 0. Then there exists a constant  $C > 0$  depending only on the data  $(G, r)$  such that*

$$|\Delta_n(f)| \leq (n+1)!(C\|f\|_{\Gamma})^{n+1} r^{-n(n+1)} \text{ for every } n \in \mathbb{N}_0,$$

where  $\|f\|_\Gamma := \max\{|f(z)| : z \in \Gamma\}$ .

*Proof.* The more general version in which  $f$  can have isolated singularities is proved in [Bie55, pp. 121–124].  $\square$

For the proof of Theorem 1.1, we need the following estimate when applying Pólya's inequality for certain auxiliary functions  $g$  constructed from  $f$ :

**Corollary 2.7.** *Let  $f(z)$  be an analytic function on  $G$  with the Taylor series  $f(z) = \sum_{n=0}^{\infty} a_n z^n$  at 0. There exist constants  $C_1 > 0$  depending on  $(G, r, f)$  and  $C_2 \geq 1$  depending on  $(G, r)$  such that for every  $d \geq 0$  and  $c_0, \dots, c_d \in \mathbb{C}$ , we have*

$$|\Delta_n(g)| \leq (n+1)! C_1^{n+1} \left( \sum_{j=0}^d |c_j| j! C_2^j \right)^{n+1} r^{-n(n+1)}$$

for every  $n \in \mathbb{N}_0$ , where  $g(z) := \sum_{j=0}^d c_j z^j f^{(j)}(z)$ .

*Proof.* In order to apply Theorem 2.6, we need an upper bound for  $\|g\|_\Gamma$  as follows. Let  $r' = (r + \rho)/2 \in (r, \rho)$  and let  $\Gamma' = \varphi^{-1}(\partial D(0, r'))$ . Let

$$\kappa := \min\{|z - z'| : z \in \Gamma, z' \in \Gamma'\}$$

denote the distance between  $\Gamma$  and  $\Gamma'$ . For  $z_0 \in \Gamma$ , Cauchy's formula gives:

$$f^{(j)}(z_0) = \frac{j!}{2\pi i} \int_{\Gamma'} f(z)/(z - z_0)^{j+1} dz$$

and so

$$|f^{(j)}(z_0)| \leq (2\pi)^{-1} j! \|f\|_{\Gamma'} \kappa^{-j-1} \text{length}(\Gamma').$$

Consequently,

$$\|g\|_\Gamma \leq C_3 \left( \sum_{j=0}^d |c_j| j! M^j \kappa^{-j-1} \right)$$

for some constant  $C_3$  depending only on  $(G, r, f)$ , where  $M = \|z\|_\Gamma$ . Let  $C$  be the resulting constant in Theorem 2.6, we now take  $C_1 = CC_3 \kappa^{-1}$  and  $C_2 = \max(1, M \kappa^{-1})$  then apply Theorem 2.6 to finish the proof.  $\square$

### 3. PROOF OF THEOREM 1.1

In this section we prove our main result. We begin with a simple lemma.

**Lemma 3.1.** *Let  $L$  and  $n$  be natural numbers with  $n > 4L$  and let  $i_1, \dots, i_L$  be integers with*

$$1 \leq i_1 < i_2 < \dots < i_L \leq n.$$

*Then there exist  $c_0, \dots, c_{2L} \in \mathbb{Z}$ , not all zero, with  $|c_i| < n \binom{n}{2L}$  for  $i = 0, \dots, 2L$  such that*

$$\sum_{i=0}^{2L} c_i \binom{m}{i} = 0$$

for  $m = i_1, \dots, i_L$ .



*Proof.* Since  $n > 4L$ , the maximum of  $\binom{m}{i}$  with  $i \leq 2L$  and  $m \in \{i_1, \dots, i_L\}$  is  $\binom{n}{2L}$ . Then by Siegel's lemma [BG06, p. 72], there exist  $c_0, \dots, c_{2L} \in \mathbb{Z}$  not all zero, with

$$|c_i| < \left( (2L+1) \cdot \binom{n}{2L} \right)^{L/(L+1)} < n \binom{n}{2L}$$

such that

$$\sum_{i=0}^{2L} c_i \binom{m}{i} = 0$$

for  $m = i_1, \dots, i_L$ . □

**Definition 3.2.** We say that a polynomial  $P(z) \in \mathbb{Q}[z]$  is *integer-valued* if  $P(n)$  is an integer for every  $n \in \mathbb{Z}$ .

**Theorem 3.3.** Let  $f(z) = \sum a_n z^n \in \mathbb{Q}[[z]]$  be a power series that converges on the open unit disk and write  $d_n = \text{den}(a_n)$ . Then one of the following three statements must hold:

- (i)  $f(z)$  admits the unit circle as a natural boundary;
- (ii) there is some  $\beta > 1$ , depending only upon  $f(z)$ , such that for every set  $S \subseteq \mathbb{N}$  of zero density, we have  $\text{lcm}\{d_i : i \leq n, i \notin S\} > \beta^n$  for all sufficiently large  $n$ ;
- (iii) for every positive constant  $C$ , there exist a positive integer  $n$ , an integer-valued polynomial  $P(z)$  of degree at most  $n$ , and co-prime polynomials  $u(z)$  and  $v(z)$  of degree at most  $n$  with  $v(0) \neq 0$  such that

$$\text{ord} \left( \sum_{j \geq 0} P(j) a_j z^j - u(z)/v(z) \right) > Cn.$$

*Proof.* Suppose that  $f(z)$  does not admit the unit circle as a natural boundary. Let  $D$  be a simply connected region of the plane that properly contains the open unit disc such that  $f(z)$  can be extended to an analytic function in  $D$ . Let  $r \in (1, \rho)$ , where  $\rho > 1$  is the conformal radius (from the origin) of  $D$ . For the moment, let  $C$  be a fixed positive integer.

Let  $S$  be a subset of the natural numbers of zero density and for  $m \geq 1$ , let  $S(m)$  denote the set of natural numbers  $j \leq m$  with  $j \in S$ . Let  $n$  be a natural number and let  $i_1 < \dots < i_L \leq Cn$  denote the elements of  $S(Cn)$ . Then by assumption  $L/n \rightarrow 0$  as  $n \rightarrow \infty$  and we assume that  $n$  is large enough that  $2L < n$ .

By Lemma 3.1 there exist  $c_0, \dots, c_{2L} \in \mathbb{Z}$ , not all zero, with  $|c_i| < Cn \binom{Cn}{2L}$  for  $i = 0, \dots, 2L$  such that

$$\sum_{i=0}^{2L} c_i \binom{m}{i} = 0$$

for  $m = i_1, \dots, i_L$ . Then we let  $g_n(z) = \sum_{j=0}^{2L} c_j z^j f^{(j)}(z)/j!$ . We can express

$$g_n(z) = \sum_j P_n(j) a_j z^j$$

for some nonzero integer-valued polynomial  $P_n(z)$  of degree at most  $2L < n$  and  $P_n(i_1) = \dots = P_n(i_L) = 0$ . Then by Corollary 2.7, there exist constants  $A > 0$

and  $\alpha \geq 1$ , depending only on the data  $(f, D, r)$ , such that for  $m \in \{n, \dots, Cn\}$  we have

$$\begin{aligned} |\Delta_m(g_n(z))| &\leq (m+1)!A^{m+1} \left( \sum_{j=0}^{2L} |c_j|\alpha^j \right)^{m+1} r^{-m(m+1)} \\ &< (m+1)!A^{m+1} \left( Cn^2 \binom{Cn}{2L} \alpha^{2L} \right)^{m+1} r^{-m(m+1)}. \end{aligned}$$

Let  $d(m) = \text{lcm}\{d_i : i \leq m, i \notin S\}$ . By definition,  $d(m)^{m+1}\Delta_m(g_n(z))$  is an integer for  $m = n, \dots, Cn$  and hence we must have either  $\Delta_m(g_n(z)) = 0$  for all  $m$  in this range, or there is some  $m \in \{n, \dots, Cn\}$  such that

$$1/d(m)^{m+1} \leq |\Delta_m(g_n(z))| < (m+1)!A^{m+1} \left( Cn^2 \binom{Cn}{2L} \alpha^{2L} \right)^{m+1} r^{-m(m+1)}.$$

Stirling's estimate now shows that if  $1 < r_0 < r$  there is some fixed  $\epsilon > 0$  such that if  $L < \epsilon n$  then

$$CAn^2 \binom{Cn}{2L} \alpha^{2L} < r_0^n$$

for  $n$  sufficiently large. In particular, since  $L < \epsilon n$  for  $n$  sufficiently large, we see that

$$1/d(m)^{m+1} < (m+1)!(r_0/r)^{m(m+1)}$$

for all large  $n$  with  $m \in \{n, \dots, Cn\}$  such that  $\Delta_m(g_n(z)) \neq 0$ .

There are now two cases. First, if for every positive integer  $C$ , there is a set  $S \subset \mathbb{N}$  of zero density and a large integer  $n$  such that  $\Delta_m(g_n(z)) = 0$  for all  $m \in \{n, \dots, Cn\}$  where  $g_n$  is constructed as above. In this case we see that (iii) holds by Corollary 2.2. Second, if there is some  $C$  such that for every set  $S \subset \mathbb{N}$  of zero density and for all large  $n$  we have  $\Delta_m(g_n(z)) \neq 0$  for some  $m \in \{n, \dots, Cn\}$  then we have that

$$1/d(m)^{m+1} < (m+1)!(r_0/r)^{m(m+1)}.$$

But this implies:

$$d(Cn) \geq d(m) > (r/r_0)^m (m+1)!^{-1/(m+1)} > \frac{(r/r_0)^m}{m+1} \geq \frac{(r/r_0)^n}{Cn+1}$$

for all sufficiently large  $n$ . Since we may begin with any  $r_0 \in (1, r)$ , we may take any  $\beta \in (1, r)$  and have that  $d(n) > \beta^n$  for all sufficiently large  $n$ . This means (ii) holds and we finish the proof.  $\square$

We have an elementary result relating a lower bound on  $\text{lcm}\{\text{den}(a_i) : i \leq n\}$  to a lower bound on  $\text{den}(a_i)$ :

**Lemma 3.4.** *Let  $(d_n)_{n \geq 0}$  be a sequence of positive integers. Suppose there exists  $\beta > 1$  such that for every set  $S \subset \mathbb{N}$  of zero density, we have:*

$$\text{lcm}\{d_i : i \leq n, i \notin S\} > \beta^n$$

*for all sufficiently large  $n$ . Let  $\kappa \in (0, \log \beta)$ , then the set  $\{n : d_n \geq \kappa n\}$  has positive density.*

*Proof.* Put  $S = \{n : d_n \geq \kappa n\}$  and suppose that  $S$  has zero density. Then

$$\text{lcm}\{d_i : i \leq n, i \notin S\} \leq \text{lcm}\{m : 1 \leq m \leq \kappa n\} = e^{\kappa n + o(n)},$$

where the second equality follows from the prime number theorem. This yields a contradiction when  $n$  is large.  $\square$

**Proposition 3.5.** *Let  $f(z) = \sum a_n z^n \in \bar{\mathbb{Q}}[[z]]$  with the property that there exists a nonzero  $P(z) \in \bar{\mathbb{Q}}[z]$  such that  $g(z) = \sum P(n)a_n z^n$  is the power series expansion of a rational function. Then at least one of the following holds:*

- (i)  $h(a_n) \gg n$  on a set of positive density.
- (ii) *There exists  $s \in \mathbb{N}$  and polynomials  $Q_0, \dots, Q_{s-1} \in \bar{\mathbb{Q}}[z]$  such that  $P(sn + i)a_{sn+i} = Q_i(n)$  for every  $i \in \{0, \dots, s-1\}$  and sufficiently large  $n$ .*

*Proof.* Suppose that (i) does not hold. We may assume that  $g(z)$  is not a polynomial since otherwise  $a_n = 0$  for all sufficiently large  $n$  and (ii) holds trivially. Write  $g(z) = u(z)/v(z)$  for coprime polynomials  $u, v \in \bar{\mathbb{Q}}[z]$  with  $v(0) = 1$  and  $v$  is non-constant. Write  $v(z) = (1 - \alpha_1 z)^{m_1} \cdots (1 - \alpha_q z)^{m_q}$  with  $q \geq 1$  and  $\alpha_1, \dots, \alpha_q$  are distinct nonzero algebraic numbers. For all sufficiently large  $n$ , we can express

$$P(n)a_n = R_1(n)\alpha_1^n + \dots + R_q(n)\alpha_q^n$$

where the  $R_i$ 's are nonzero elements of  $\bar{\mathbb{Q}}[z]$ . First, we prove that all the  $\alpha_i$ 's are roots of unity. Restricting  $n$  to appropriate arithmetic progressions if necessary, we may assume that  $\alpha_i/\alpha_j$  is not a root of unity for  $i \neq j$ . Suppose there is a place  $w$  of  $\bar{\mathbb{Q}}$  such that

$$M := \max_{1 \leq i \leq q} |\alpha_i|_w > 1.$$

Then for every  $M_0 \in (1, M)$ , we apply results in [KMN19, Section 2] to conclude that  $|a_n|_w > M_0^n$  for all but finitely many  $n$ . This gives (i), contradicting the assumption at the beginning of the proof. Hence we have  $|\alpha_i|_w \leq 1$  for every place  $w$  and for  $0 \leq i \leq q$ . Therefore all the  $\alpha_i$ 's are roots of unity. Let  $s \in \mathbb{N}$  such that  $\alpha_i^s = 1$  for every  $i$ . Then we have polynomials  $Q_i(z)$ 's for  $i \in \{0, \dots, s-1\}$  satisfying (ii).  $\square$

**Theorem 3.6.** *Let  $f(z) = \sum a_n z^n \in \mathbb{Q}[[z]]$  be a  $D$ -finite power series that converges on the open unit disk such that for every nonzero  $P(z) \in \mathbb{Q}[z]$  the series  $\sum P(n)a_n z^n$  is not a rational function. Put  $d_n = \text{den}(a_n)$  for every  $n$ . There exists  $\beta > 1$  such that for every set  $S \subset \mathbb{N}$  of zero density, we have*

$$\text{lcm}\{d_i : i \leq n, i \notin S\} > \beta^n$$

for all sufficiently large  $n$ .

*Proof.* Among all the nonzero integer-valued polynomials  $P(z)$ , we pick one such that

$$\dim_{\mathbb{C}(z)} \text{Span}\{t(z), t'(z), t''(z), \dots\} \text{ is minimal,}$$

where  $t(z) = \sum_{n \geq 0} a_n P(n) z^n$ . Then it suffices to prove the result for  $t(z)$  since  $\text{den}(P(n)a_n)$  divides  $\text{den}(a_n)$  for every  $n$ . Thus for the remainder of the proof, we replace  $f(z)$  by  $t(z)$  and let

$$p = \dim_{\mathbb{C}(z)} \text{Span}\{f(z), f'(z), f''(z), \dots\}.$$

Then by construction,  $f(z) = \sum a_n z^n$  has the property that for every nonzero integer-valued  $P(z)$  we have  $p = \dim_{\mathbb{C}(z)} \text{Span}\{t(z), t'(z), t''(z), \dots\} > 1$ , where  $t(z) = \sum a_n P(n) z^n$ . Let

$$\mathcal{L} = \sum_{i=0}^p A_i D^{p-i}$$

be the differential operator as in (2) such that  $\mathcal{L}(f) = 0$  and let

$$\delta = \max_{0 \leq i \leq p} \deg(A_i),$$

as before. Since  $f(z)$  is D-finite it does not admit the unit circle as a natural boundary. We assume that the conclusion of this theorem does not hold and arrive at a contradiction.

By Theorem 3.3, we have that for every positive integer  $C$ , there is a natural number  $n$ , a nonzero integer-valued polynomial  $P(z)$  of degree at most  $n$  such that

$$g(z) = \sum a_n P(n) z^n$$

has the property that there exist co-prime polynomials  $u(z), v(z)$  of degree at most  $n$  with  $v(0) \neq 0$  such that

$$(3) \quad \text{ord}(g(z) - u(z)/v(z)) > Cn.$$

By Lemma 2.4, there are polynomial  $r_0(z), \dots, r_{p-1}(z), s(z)$  of degree at most  $(2\delta + 1)np$ , with  $s(z)$  nonzero, such that

$$s(z)f(z) = \sum_{i=0}^{p-1} r_i(z)g^{(i)}(z).$$

Now  $\text{ord}(g^{(i)}(z) - (u(z)/v(z))^{(i)}) > Cn - i$  by (3) and so

$$\text{ord} \left( s(z)f(z) - \sum_{i=0}^{p-1} r_i(z)(u/v)^{(i)}(z) \right) > Cn - p.$$

This gives  $\text{ord}(f(z) - \Theta(z)) > Cn - p - (2\delta + 1)pn$ , where

$$\Theta(z) = s(z)^{-1} \sum_{i=0}^{p-1} r_i(z)(u/v)^{(i)}(z).$$

Notice that  $\Theta(z)$  is a rational function whose numerator and denominator have degree bounded by  $(2\delta + 1)np + np$ . Taking  $C$  large enough, we see that  $f(z)$  is necessarily rational by Lemma 2.5, contradiction.  $\square$

**Lemma 3.7.** *If  $f(z)$  has radius of convergence  $r \in (0, \infty)$  and if  $f(z) = \sum_{n \in S} a_n z^n + g(z)$  where  $g(z)$  has radius of convergence strictly greater than  $r$  and  $S$  has zero density, then  $f(z)$  admits the circle of radius  $r$  as a natural boundary.*

*Proof.* Let  $f_0(z) = \sum_{n \in S} a_n z^n$ . Then  $f_0(z)$  has radius of convergence  $r$  and since  $S$  has zero density, by Fabry's gap theorem (see [Tur47]),  $f_0(z)$  admits the circle of radius  $r$  as its natural boundary. Thus  $f(z)$  does too, since  $g(z)$  is analytic in a circle of radius  $r + \epsilon$  centred at zero, for some  $\epsilon > 0$ .  $\square$

**Proposition 3.8.** *Let  $K$  be a number field, let  $f(z) = \sum a_n z^n \in K[[z]]$ , and let  $r$  be the radius of convergence of  $f(z)$ . Then the following hold:*

- (a) if  $r \in (1, \infty)$  and  $f(z)$  does not admit the circle of radius  $r$  as a natural boundary then there is a positive constant  $\kappa$  such that  $h(a_n) \geq \kappa n$  for  $n$  in a set of positive density;
- (b) if  $r \in (0, 1)$  and  $f(z)$  does not admit the circle of radius  $r$  as a natural boundary then there is a positive constant  $\kappa'$  such that  $h(a_n) \geq \kappa' n$  for  $n$  in a set of positive density;
- (c) if  $r \in \{0, \infty\}$ ,  $f(z)$  is  $D$ -finite and  $f(z)$  is not a polynomial then  $h(a_n) = O(n \log n)$  and there is a positive constant  $\kappa$  such that  $h(a_n) > \kappa n \log n$  on a set of positive density.

*Proof.* If  $r \in (1, \infty)$ , then there is some  $c < 1$  such that  $|a_n| < c^n$  when  $n$  is sufficiently large. In particular, there is a positive constant  $\kappa$  such that for  $n$  sufficiently large we have  $h(a_n) \geq \kappa n$  whenever  $a_n$  is nonzero. Since  $f(z)$  does not admit the circle of radius  $r$  as a natural boundary, by Lemma 3.7 we have that the set of  $n$  for which  $a_n \neq 0$  has positive density.

If  $r \in (0, 1)$ , there is some  $c > 1$  such that  $|a_n| > c^n$  for infinitely many  $n$ . Pick  $b \in (1, c)$  with  $1/b > r$  and let  $S$  denote the set of  $n$  for which  $|a_n| > b^n$ . We claim that  $S$  has positive density. To see this, suppose that  $S$  has density zero. Then we can write  $f(z) = f_0(z) + f_1(z)$ , where  $f_0(z) = \sum_{n \in S} a_n z^n$  and  $f_1(z) = \sum_{n \notin S} a_n z^n$ . Then by construction  $f_1$  is convergent in the disc  $D(0, 1/b)$ . Then  $f_0(z)$  must have radius of convergence  $r$  since  $f(z)$  has radius of convergence  $r$  and  $f_1$  has radius of convergence strictly greater than  $r$ . Since  $S$  has zero density, by Lemma 3.7 we have that  $f_0(z)$  admits the circle of radius  $r$  as its natural boundary and so  $f(z)$  must too since  $f_1(z)$  is analytic in  $B(0, 1/b)$  and  $1/b > r$ . This is a contradiction. Thus we obtain (b).

It remains to consider the case  $r \in \{0, \infty\}$ . The property  $h(a_n) = O(n \log n)$  has been proved in [BNZ20]. The power series  $f(z)$  is a Gevrey series whose precise Gevrey order is a rational number: there is  $s \in \mathbb{Q}$  such that  $\sum a_n (n!)^s z^n$  has positive finite radius of convergence (see, for example, [Ram78]). Since  $r \in \{\infty, 0\}$ , we have  $s \neq 0$ . We write  $s = a/b$  with  $a, b$  nonzero integers and  $b > 0$ . Then by construction  $g(z) := \sum_{n \geq 0} a_n^b n!^a z^n$  is  $D$ -finite and has positive nonzero radius of convergence. In particular, by using Lemma 3.7 as above there exist  $r_0, r_1 \in (0, \infty)$  with  $r_0 < r_1$  such that

$$r_0^n < a_n^b n!^a < r_1^n$$

for  $n$  in a set of natural numbers of positive density. In particular, if  $a > 0$  then we see that  $a_n^{-b} > n!^a / r_1^n$ , which gives that  $h(a_n) > \kappa n \log n$  for some positive constant  $\kappa$  for  $n$  in a set of positive density. Similarly, if  $a < 0$  then  $a_n^b > r_0^n n!^{-a}$  and so the result follows similarly in this case.  $\square$

**Theorem 3.9.** *Let  $f(z) = \sum_{n=0}^{\infty} a_n z^n \in \bar{\mathbb{Q}}[[z]]$  be a  $D$ -finite power series. Let  $r$  be the radius of convergence of  $f$ , we have:*

- (a) If  $r \in \{0, \infty\}$  and  $f$  is not a polynomial then  $h(a_n) = O(n \log n)$  for every large  $n$  and  $h(a_n) \gg n \log n$  on a set of positive density.
- (b) If  $r \notin \{0, \infty\}$  then at least one of the following holds:
  - (i)  $h(a_n) \gg n$  on a set of positive density.
  - (ii) There is a constant  $\beta > 1$  such that for every set  $S \subset \mathbb{N}$  of zero density, we have:

$$\text{lcm}\{\text{den}(a_i) : i \leq n, i \notin S\} > \beta^n$$

for all sufficiently large  $n$ .

- (iii) *There exist  $s \in \mathbb{N}$  and rational functions  $Q_0(z), \dots, Q_{s-1}(z) \in \bar{\mathbb{Q}}(z)$  such that  $a_{sn+i} = Q_i(n)$  for every  $i \in \{0, \dots, s-1\}$  and sufficiently large  $n$ .*

*Proof.* Part (a) is from Proposition 3.8, we now assume  $r \notin \{0, \infty\}$ . Suppose neither (i) nor (ii) hold. Let  $K$  be a number field such that  $f(z) \in K[[z]]$  and let  $\mathcal{O}$  be its ring of integers. Let  $m := [K : \mathbb{Q}]$  and let  $\{\gamma_1, \dots, \gamma_m\}$  be an integral basis of  $K$ . Write  $\text{Tr} := \text{Trace}_{K/\mathbb{Q}}$ . If there exist a nonzero  $P(z) \in \bar{\mathbb{Q}}[z]$  such that  $\sum P(n)a_n z^n$  is a rational function then we have (iii) thanks to Proposition 3.5. From now on, we assume that there does not exist such a  $P$ .

For  $1 \leq i \leq m$ , write

$$g_i(z) = \text{Tr}(\gamma_i f(z)) = \sum_n a_{i,n} z^n \in \mathbb{Q}[[z]],$$

where  $a_{i,n} = \text{Tr}(\gamma_i a_n)$ . From the basic height inequalities in Subsection 2.1 and the assumption that (i) does not hold, for each  $i \in \{1, \dots, m\}$  we do *not* have that  $h(a_{i,n}) \gg n$  on a set of positive density. By Proposition 3.8 and the fact that  $g_i$  is D-finite, we have that  $g_i$  has radius of convergence 1 for  $i \in \{1, \dots, m\}$ .

First, consider the case that for every  $i \in \{1, \dots, m\}$  there exists a nonzero  $P_i(z) \in \bar{\mathbb{Q}}[z]$  such that  $\sum_n P_i(n)a_{i,n}z^n$  is a rational function. For every  $i \in \{1, \dots, m\}$ , Proposition 3.5 implies the existence of  $s_i \in \mathbb{N}$  and  $R_{i,j}(z) \in \bar{\mathbb{Q}}(z)$  such that  $a_{i,s_i n+j} = R_{i,j}(n)$  for  $j \in \{0, \dots, s_i - 1\}$  and sufficiently large  $n$ . From elementary field theory, there exist  $c_1, \dots, c_m \in K$  such that for every  $x \in K$  we have

$$x = c_1 \text{Tr}(\gamma_1 x) + \dots + c_m \text{Tr}(\gamma_m x);$$

therefore  $a_n = c_1 a_{1,n} + \dots + c_m a_{m,n}$  for every  $n$ . It now follows that (iii) holds with  $s = \text{lcm}\{s_1, \dots, s_m\}$ .

We now consider the case when there is  $i_0 \in \{1, \dots, m\}$  such that for every nonzero  $P(z) \in \bar{\mathbb{Q}}[z]$  the series  $\sum_n P(n)a_{i_0,n}z^n$  is not a rational function. Theorem 3.6 yields  $\beta > 1$  such that for every set  $S \subset \mathbb{N}$  of zero density we have

$$\text{lcm}\{\text{den}(a_{i_0,k}) : k \leq n, k \notin S\} > \beta^n$$

for all sufficiently large  $n$ . This implies

$$\text{lcm}\{\text{den}(a_k) : k \leq n, k \notin S\} > \beta^n$$

for all sufficiently large  $n$  since  $\text{den}(a_{i_0,k}) \mid \text{den}(a_k)$  for every  $k$ . But since we are assuming that (ii) does not hold, the current case cannot happen and we finish the proof.  $\square$

*Proof of Theorem 1.1.* By Theorem 3.9, it remains to prove part (b) of Theorem 1.1. Assume that neither (i) nor (ii) of Theorem 1.1(b) hold. By Lemma 3.4, we have that neither (i) nor (ii) of Theorem 3.9 hold. Hence there exist  $s \in \mathbb{N}$  and rational functions  $Q_0, \dots, Q_{s-1} \in K(z)$  such that  $a_{sn+i} = Q_i(n)$  for every  $i \in \{0, \dots, s-1\}$  and sufficiently large  $n$ . First, we consider the case when  $Q_i(z)$  is a polynomial for every  $i \in \{0, \dots, s-1\}$ . Then it follows that  $f(z)$  is a rational function whose finite poles must be roots of unity.

We now consider the case when some  $Q_{i_0}$  is not a polynomial. Let  $\mathcal{O}$  be the ring of integers of  $K$ , write  $Q_{i_0}(z) = A(z)/B(z)$ , where  $A$  and  $B$  are coprime

polynomials with coefficients in  $\mathcal{O}$  and  $B$  is non-constant. It is an easy fact that  $\text{den}(A(n)/B(n)) \gg n^{\deg(B)}$  when  $n$  is sufficiently large and we include the proof here for the sake of completeness. Write  $\delta_n = \text{den}(A(n)/B(n))$ . There exist  $U(z), V(z) \in \mathcal{O}[z]$  and a nonzero  $C \in \mathcal{O}$  such that:

$$U(z)A(z) + V(z)B(z) = C.$$

Then for every  $v \in M_K^0$  and  $n \in \mathbb{N}$  we have:

$$(4) \quad |C|_v \leq \max\{|A(n)|_v, |B(n)|_v\}.$$

Let  $v \in M_K^0$  such that  $|A(n)|_v \geq |B(n)|_v$ . From  $|\delta_n A(n)/B(n)|_v \leq 1$ , we have:

$$|C\delta_n|_v \leq |C|_v \cdot |B(n)/A(n)|_v \leq |B(n)|_v,$$

where the rightmost inequality follows from (4) and the assumption on  $v$ .

Let  $v \in M_K^0$  such that  $|A(n)|_v < |B(n)|_v$ . We have:

$$|C\delta_n|_v \leq |C|_v \leq |B(n)|_v,$$

where the rightmost inequality follows from (4) and the assumption on  $v$ .

Therefore we have  $|C\delta_n|_v \leq |B(n)|_v$  for every  $v \in M_K^0$ . Combining this with the product formula and the fact that  $\delta_n \in \mathbb{Q}$ , we have:

$$\delta_n \prod_{v \in M_K^\infty} |C|_v = \frac{1}{\prod_{v \in M_K^0} |C\delta_n|_v} \geq \frac{1}{\prod_{v \in M_K^0} |B(n)|_v} = \prod_{v \in M_K^\infty} |B(n)|_v \gg n^{\deg(B)}.$$

Since  $\deg(B) \geq 1$ , we have that  $\text{den}(a_n) \gg n$  on a set of positive density. Since we are assuming that (ii) does not hold, the current case cannot happen and we finish the proof.  $\square$

*Remark 3.10.* Throughout this paper, we obtain lower bounds for  $h(a_n)$  for  $n$  in a set of positive *upper* density. We explain how to slightly improve some of the previous results. For a subset  $S$  of  $\mathbb{N}$ , its lower density is

$$\liminf_{n \rightarrow \infty} \frac{|S \cap [1, n]|}{n}.$$

We can strengthen Theorem 3.3(ii), Theorem 3.6, and Theorem 3.9(b)(ii) by replacing “zero density” by “zero lower density” using similar arguments to the earlier proofs. Finally, we can strengthen Theorem 1.1(b)(ii) by replacing “positive upper density” by “positive lower density”.

#### 4. CONCLUDING REMARKS

We start with a comment on the subtlety of Theorem 1.1. In view of Theorem 1.1, Theorem 3.9, and Lemma 3.4, it is natural to ask whether one can strengthen Theorem 1.1 by replacing the property “ $\text{den}(a_n) \gg n$  on a set of positive density” in (ii) by the property “ $\text{lcm}\{\text{den}(a_k) : k \leq n, k \notin S\} > \beta^n$  for all sufficiently large  $n$  for every  $S$  of zero density”. The answer is negative even for the simple example

$$\log(1+x) = \frac{1}{x} - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \dots$$

in which  $\text{den}(a_n) = n$  for every  $n$ . A version of the prime number theorem gives  $\text{lcm}\{1, \dots, n\} = e^{n+o(n)}$  for all large  $n$ . On the other hand, even when we relax the condition “for all large  $n$ ” to the condition “for infinitely many  $n$ ”, it is still possible to construct a set  $S$  of zero density such that  $\text{lcm}\{i : i \leq n, i \notin S\}$  fails to have an exponential lower bound:

**Proposition 4.1.** *There exists  $S \subset \mathbb{N}$  of zero density such that for every  $\beta > 1$  the inequality*

$$\text{lcm}\{i : i \leq n, i \notin S\} > \beta^n$$

*holds for only finitely many  $n$ .*

*Proof.* For  $k \in \mathbb{N}_0$ , let  $n_k = k!$ . Let  $(c_k)_{k \in \mathbb{N}}$  be a sequence of positive integers such that

$$(5) \quad \lim_{k \rightarrow \infty} \frac{c_k}{k} = \infty, \quad \lim_{k \rightarrow \infty} \frac{\log c_k}{\log k} = 1, \quad \text{and} \quad \lim_{k \rightarrow \infty} \frac{c_k}{k \log k} = 0;$$

for example one may take  $c \in (0, 1)$  and let  $c_k = \lfloor k(\log k)^c \rfloor$  for large  $k$ . For  $k \in \mathbb{N}$ , let  $S_k$  be the set of integers in  $(n_{k-1}, n_k]$  that are divisible by a prime in  $(n_k/c_k, n_k]$ . Let  $S = \bigcup_{k \in \mathbb{N}} S_k$ .

Fix  $\beta > 1$ . Let  $n$  denote a sufficiently large integer and let  $k$  be such that  $n_{k-1} < n \leq n_k$ . From the construction of  $S$ , we have that  $\text{lcm}\{i : i \leq n, i \notin S\}$  is not divisible by any prime in  $(n_k/c_k, n]$ . Therefore:

$$(6) \quad \text{lcm}\{i : i \leq n, i \notin S\} \leq \frac{\text{lcm}\{1, \dots, n\}}{\prod_{p \in (n_k/c_k, n]} p} = e^{n_k/c_k + o(n)}$$

thanks to the prime number theorem. From  $n_k/c_k < nk/c_k$  and (5), it follows that the right side of (6) is  $e^{o(n)}$ . Therefore

$$\text{lcm}\{i : i \leq n, i \notin S\} \leq \beta^n$$

for all sufficiently large  $n$ .

It remains to prove that  $S$  has zero density. In fact, for the purpose of [BGNS, Section 1], we prove the stronger property that

$$(7) \quad |S \cap [1, n]| = O(n \log \log n / \log n).$$

For  $I \subseteq \mathbb{R}$  let  $\pi(I)$  denote the number of primes in  $I$  and for  $x \geq 1$  let  $\pi(x) = \pi([1, x])$ . We will use the following well-known estimates repeatedly:

$$(8) \quad \begin{aligned} \pi(x) &= \frac{x}{\log x} + O(x/(\log x)^2), \quad \sum_{m \leq x} \frac{1}{m} = \log x + O(1), \\ \log n_k &\sim k \log k, \quad \text{and} \quad \frac{1}{k} \sim \frac{\log \log n_k}{\log n_k}. \end{aligned}$$

First we give an upper bound of  $|S_k|$  for large  $k$ . Since  $S_k$  consists of numbers of the form  $jp$ , where  $1 \leq j \leq m$  and  $p$  is a prime in  $(n_k/(m+1), n_k/m]$  for  $1 \leq m < c_k$ , we have

$$(9) \quad |S_k| = \sum_{m=1}^{c_k-1} m \pi((n_k/(m+1), n_k/m]).$$

Since  $c_k = n_k^{o(1)}$ , we have  $\log(n_k/m) \gg \log n_k$  for  $1 \leq m \leq c_k$  and then

$$\frac{n_k}{m \log(n_k/m)} - \frac{n_k}{m \log n_k} = O\left(\frac{n_k \log m}{(\log n_k)^2 \cdot m}\right).$$

Therefore

$$\pi((n_k/(m+1), n_k/m]) = \frac{n_k}{m \log n_k} - \frac{n_k}{(m+1) \log n_k} + O\left(\frac{n_k \log m}{(\log n_k)^2 \cdot m}\right).$$



Combining this with (9), we have:

$$\begin{aligned} |S_k| &= \frac{n_k}{\log n_k} \sum_{m=1}^{c_k-1} \frac{1}{m+1} + \sum_{m=1}^{c_k-1} O\left(\frac{n_k \log m}{(\log n_k)^2}\right) \\ &= \frac{n_k \log c_k}{\log n_k} + O(n_k / \log n_k) + O(n_k c_k \log c_k / (\log n_k)^2). \end{aligned}$$

Since  $\log n_k \sim k \log k$ ,  $\log c_k \sim \log k$ , and  $c_k = o(k \log k)$ , we have  $\frac{n_k \log c_k}{\log n_k} \sim \frac{n_k}{k}$ ,  $n_k / \log n_k = o(n_k / k)$ , and  $n_k c_k \log c_k / (\log n_k)^2 = o(n_k / k)$ . Therefore

$$(10) \quad |S_k| = \frac{n_k}{k} + o(n_k / k) \sim (k-1)!$$

and hence

$$(11) \quad \sum_{j=1}^k |S_j| = O((k-1)!).$$

Let  $n$  denote a large integer and let  $k$  be such that  $(k-1)! < n \leq k!$ . From (11), we have:

$$\left| \bigcup_{j=1}^{k-1} S_j \right| \ll (k-2)! < n / (k-1) = O(n \log \log n / \log n).$$

Therefore it remains to show that  $|S_k \cap [1, n]| = O(n \log \log n / \log n)$ . Let  $\lambda$  be the integer in  $[1, k-1]$  such that  $n \in (n_k / (\lambda+1), n_k / \lambda]$ . Every element in  $S_k \cap [1, n]$  must have the form  $jp$  where:

- $p$  is a prime in the set  $(n_k / (\lambda+m), n_k / (\lambda+m-1)] \cap [1, n]$  for  $1 \leq m \leq c_k - \lambda$  (when  $m \geq 2$ , the above set is simply  $(n_k / (\lambda+m), n_k / (\lambda+m-1)]$ ).
- $j \in \mathbb{N}$  with  $j \leq \frac{n}{p} < \frac{n_k}{\lambda} \cdot \frac{\lambda+m}{n_k} = \frac{m}{\lambda} + 1$

By using similar estimates as before and the above properties of  $p$  and  $j$ , we have the following upper bound for  $|S_k \cap [1, n]|$ :

$$\begin{aligned} & \sum_{m=1}^{c_k-\lambda} \left(\frac{m}{\lambda} + 1\right) \pi((n_k / (\lambda+m), n_k / (\lambda+m-1)] \cap [1, n]) \\ & \leq \sum_{m=1}^{c_k-\lambda} \frac{m}{\lambda} \left( \frac{n_k}{(m+\lambda-1) \log n_k} - \frac{n_k}{(m+\lambda) \log n_k} + O\left(\frac{n_k \log(m+\lambda)}{(\log n_k)^2 \cdot (m+\lambda)}\right) \right) \\ & \quad + \pi(n) \\ & = \frac{n_k}{\lambda \log n_k} \sum_{m=1}^{c_k-\lambda} m \left( \frac{1}{m+\lambda-1} - \frac{1}{m+\lambda} \right) + O\left(\frac{n_k}{\lambda (\log n_k)^2} \sum_{j=\lambda+1}^{c_k} \log j\right) \\ & \quad + O(n / \log n). \end{aligned}$$

The first  $O$ -term is  $O\left(\frac{n_k c_k \log c_k}{\lambda (\log n_k)^2}\right) = o\left(\frac{n_k}{\lambda k}\right) = o\left(\frac{n \log \log n}{\log n}\right)$  by the same reasoning as before, and the second  $O$ -term is also  $o\left(\frac{n \log \log n}{\log n}\right)$

$$\begin{aligned} \sum_{m=1}^{c_k-\lambda} m \left( \frac{1}{m+\lambda-1} - \frac{1}{m+\lambda} \right) &= \frac{1}{\lambda} + \frac{1}{\lambda+1} + \cdots + \frac{1}{c_k-1} - \frac{c_k-\lambda}{c_k} \\ &= \log(c_k/\lambda) + O(1). \end{aligned}$$

Combining all the above, we have:

$$|S_k \cap [1, n]| \leq \frac{n_k}{\lambda \log n_k} \log(c_k) + o(n \log \log n / \log n) = O(n \log \log n / \log n)$$

where the last equality follows from  $n_k/\lambda = O(n)$ ,  $\log(n_k) \sim \log n$  and  $\log(c_k) \sim \log k \sim \log \log n$ . This finishes the proof.  $\square$

We now pose some questions related to the work we have obtained. In Corollary 1.3 we show that if  $f(z) = \sum a_n z^n \in K[[z]]$  is D-finite and not rational, with  $K$  a number field, then we have the inequality

$$h(a_n) > \frac{1}{[K:\mathbb{Q}]} \log n + O(1)$$

for  $n$  in a set of positive density. Moreover, we showed that there are examples which show this is essentially best possible when  $K = \mathbb{Q}$ . In the case that  $[K:\mathbb{Q}] > 1$ , however, the bound appears to be suboptimal and so we ask if one can improve the bounds in this case.

**Question 4.2.**  *$K$  and  $f$  as in Corollary 1.3, is it true that  $h(a_n) > \log n + O(1)$  on a set of positive density?*

Another natural question is whether one can give a complete classification of the types of growth rates of heights of coefficients of D-finite series that can occur. To the best of our knowledge, there are really only four types of possible behaviours, and the question below asks if this gives a taxonomy of possible ‘‘height gaps’’ of D-finite series.

**Question 4.3.** *Let  $f(z) \in \bar{\mathbb{Q}}[[z]]$  be D-finite, is it true that one of the following holds:*

- (i)  $h(a_n) = O(n \log n)$  for every  $n$  and  $h(a_n) \gg n \log n$  on a set of positive density.
- (ii)  $h(a_n) = O(n)$  for every  $n$  and  $h(a_n) \gg n$  on a set of positive density.
- (iii)  $h(a_n) = O(\log n)$  for every  $n$  and  $h(a_n) \gg \log n$  on a set of positive density.
- (iv)  $h(a_n) = O(1)$  for every  $n$ .

A weaker version of (the affirmative answer to) Question 4.3 is the following collection of four statements concerning a D-finite power series  $\sum_n a_n z^n \in \bar{\mathbb{Q}}[[z]]$ :

- (i')  $h(a_n) = O(n \log n)$ .
- (ii') If  $h(a_n) = o(n \log n)$  then  $h(a_n) = O(n)$ .
- (iii') If  $h(a_n) = o(n)$  then  $h(a_n) = O(\log n)$ .
- (iv') If  $h(a_n) = o(\log n)$  then  $h(a_n) = O(1)$ .

Statements (i') and (iv') have been established in [BNZ20]. Statement (ii') is similar to the following long standing open problem in the theory of Siegel E-functions [Sie29], [Beu, pp. 11–12], [FR, Section 2.1]:

**Question 4.4.** *Let  $f(z) = \sum_n a_n z^n \in \bar{\mathbb{Q}}[[z]]$  be  $D$ -finite. Suppose that  $h(a_0, \dots, a_n) = o(n \log n)$ , is it true that  $h(a_0, \dots, a_n) = O(n)$ ? Here  $h(a_0, \dots, a_n)$  is taken to be  $h([1 : a_0 : \dots : a_n])$ .*

## REFERENCES

- [ABS] B. Adamczewski, J. Bell, and D. Smertnig, *A height gap theorem for coefficients of Mahler functions*, to appear in J. Eur. Math. Soc. (JEMS), 2020, arXiv:2003.03429.
- [BC17] J. P. Bell and S. Chen, *Power series with coefficients from a finite set*, J. Combin. Theory Ser. A **151** (2017), 241–253.
- [Beu] F. Beukers, *E-functions and G-functions*, lecture notes available at <https://www.math.arizona.edu/~swc/aws/2008/08BeukersNotesDraft.pdf>.
- [BG06] E. Bombieri and W. Gubler, *Heights in Diophantine geometry*, New Mathematical Monographs, vol. 4, Cambridge University Press, Cambridge, 2006.
- [BGNS] J. P. Bell, K. Gunn, K. D. Nguyen, and J. C. Saunders, *A general criterion for the Pólya-Carlson dichotomy and application*, available on the arXiv, 2022.
- [BGS21] J. P. Bell, D. Ghioca, and M. Satriano, *Dynamical uniform bound for fibers and a gap conjecture*, Int. Math. Res. Not. **2021** (2021), 7932–7946.
- [BHS20] J. P. Bell, F. Hu, and M. Satriano, *Height gap conjectures, D-finiteness, and a weak dynamical Mordell-Lang conjecture*, Math. Ann. **378** (2020), 971–992.
- [Bie55] L. Bieberbach, *Analytische Fortsetzung*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 2. Folge, Springer, Berlin, 1955.
- [BN21] Jason P. Bell and Khoa D. Nguyen, *An analogue of Ruzsa's conjecture for polynomials over finite fields*, J. Combin. Theory Ser. A **178** (2021).
- [BNZ20] J. P. Bell, K. D. Nguyen, and U. Zannier, *D-finiteness, rationality, and height*, Trans. Amer. Math. Soc. **373** (2020), 4889–4906.
- [Dim] V. Dimitrov, *A proof of the Schinzel-Zassenhaus conjecture on polynomials*, arXiv:1912.12545.
- [FR] S. Fischler and T. Rivoal, *On Siegel's problem for E-functions*, to appear in Rendiconti del Seminario Matematico della Università di Padova, 2022.
- [HS00] M. Hindry and J. Silverman, *Diophantine Geometry. An introduction*, Graduate Texts in Mathematics, vol. 201, Springer, New York, 2000.
- [KMN19] A. Kulkarni, N. M. Mavraki, and K. D. Nguyen, *Algebraic approximations to linear combinations of powers: an extension of results by Mahler and Corvaja-Zannier*, Trans. Amer. Math. Soc. **371** (2019), 3787–3804.
- [PZ84] A. Perelli and U. Zannier, *On recurrent mod  $p$  sequences*, J. Reine Angew. Math. **348** (1984), 135–146.
- [Ram78] J.-P. Ramis, *Déviage Gevrey*, Journées singulières de Dijon, Astérisque, no. 59-60, Société mathématique de France, 1978.
- [Sie29] C.L. Siegel, *Über einige Anwendungen diophantischer Approximationen*, Abhandlungen der preuss. vol. 1, Akademie der Wissenschaften, 1929.
- [Tur47] P. Turán, *On the gap-theorem of Fabry*, Hungarica Acta Math. **1** (1947), 21–29. MR 22905
- [vdPS96] A. J. van der Poorten and I. Shparlinski, *On linear recurrence sequences with polynomial coefficients*, Glasg. Math. J. **38** (1996), 147–155.
- [Zan96] U. Zannier, *On periodic mod  $p$  sequences and G-functions*, Manuscripta Math. **90** (1996), 391–402.

JASON P. BELL, UNIVERSITY OF WATERLOO, DEPARTMENT OF PURE MATHEMATICS, WATERLOO,  
ONTARIO, CANADA N2L 3G1

*Email address:* `jpbell@uwaterloo.ca`

KHOA D. NGUYEN, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF CALGARY,  
AB T2N 1N4, CANADA

*Email address:* `dangkhoa.nguyen@ucalgary.ca`

UMBERTO ZANNIER, SCUOLA NORMALE SUPERIORE, CLASSE DI SCIENZE MATEMATICHE E NAT-  
URALI, PISA, ITALY

*Email address:* `umberto.zannier@sns.it`