

VARIANCE OF THE k -FOLD DIVISOR FUNCTION IN ARITHMETIC PROGRESSIONS FOR INDIVIDUAL MODULUS

DAVID T. NGUYEN

ABSTRACT. In this paper, we confirm a smoothed version of a recent conjecture on the variance of the k -fold divisor function in arithmetic progressions to individual composite moduli, in a restricted range. In contrast to a previous result of Rodgers and Soundararajan [25], we do not require averaging over the moduli. Our proof adapts a technique of S. Lester [20] who treated in the same range the variance of the k -fold divisor function in the short intervals setting, and is based on a smoothed Voronoï summation formula but twisted by multiplicative characters. The use of Dirichlet characters allows us to extend to a wider range from previous result of Kowalski and Ricotta [19] who used additive characters. Smoothing also permits us to treat all k unconditionally. This result is closely related to moments of Dirichlet L -functions.

CONTENTS

1. Introduction	2
2. Statement of results and a conjecture	4
3. Outline of the proofs	6
4. Notations, parameters, and conventions	7
5. Preparatory Lemmas	9
6. Proof of Theorem 1 Full main term with power-saving error term	12
6.1. Step 1: Applying the functional equation and reduction in q	12
6.2. Step 2: Reduction to the dual sum	13
6.3. Step 3: Off-diagonal analysis	15
6.4. Step 4: Diagonal analysis	16
7. Proof of Corollary 1 Asymptotic for leading order main term	18
7.1. Step a: Full main term as a residue plus an error term	18
7.2. Step b: The error term and $\gamma_k(c)$	19
7.3. Step c: Bounding lower order main terms	19
7.4. Step d: Evaluating the matching arithmetic constant $a_k(d)$	20
8. Acknowledgments	21
References	21

Date: Friday 6th May, 2022

2020 Mathematics Subject Classification: [11B25](#); [11N37](#); [11M50](#).

Key words and phrases: Divisor sums; variance in arithmetic progressions; summation formulae .

1. INTRODUCTION

One form of the celebrated Bombieri-Vinogradov Theorem [3] [27] (1965) asserts that

$$(1.1) \quad \sum_{1 \leq q \leq N^{1/2}(\log N)^{-B}} \max_{y \leq N} \max_{(a,q)=1} \left| \sum_{\substack{1 \leq n \leq y \\ n \equiv a(q)}} \Lambda(n) - \frac{y}{\varphi(q)} \right| \ll N(\log N)^{-A}$$

where $\Lambda(n)$ is the von Mangoldt function and $B = 4A + 40$ with $A > 0$ arbitrary. Analogues of (1.1) has been found for $\tau_k(n)$ [24] and $\tau_2(n)^2$ [22, Lemma 8], where $\tau_k(n)$ is the k -fold divisor function: $\zeta^k(s) = \sum_{n=1}^{\infty} \tau_k(n)n^{-s}$, ($\sigma > 1$).

Around the same time, Barban [1] [2] (1963-1964), Davenport-Halberstam [12] (1966), and Gallagher [15] (1967) found the following related inequality in which the absolute value is being squared:

$$(1.2) \quad \sum_{1 \leq q \leq N(\log N)^{-B}} \sum_{\substack{1 \leq a \leq q \\ (a,q)=1}} \left| \sum_{\substack{1 \leq n \leq N \\ n \equiv a(q)}} \Lambda(n) - \frac{N}{\varphi(q)} \right|^2 \ll N^2(\log N)^{-A},$$

giving a much wider range for q . Davenport and Halberstam proved a slightly stronger result than Barban's, while Gallagher gave a simplified elegant proof. For this reason, this type of inequalities are often referred to as Barban-Davenport-Halberstam type inequalities. Barban-Davenport-Halberstam type inequalities have many applications in number theory. For instance, a version of this inequality (with $\Lambda(n)$ replaced by related convolutions over primes) was skillfully used by Zhang [29, Lemma 10] (2014) in his spectacular work on bounded gaps between primes.

Shortly after, in 1970 Montgomery [21] succeeded in replacing the inequality in (1.2) by an asymptotic equality. One of his results is

$$(1.3) \quad \sum_{1 \leq q \leq Q} \sum_{\substack{1 \leq a \leq q \\ (a,q)=1}} \left| \sum_{\substack{1 \leq n \leq N \\ n \equiv a(q)}} \Lambda(n) - \frac{N}{\varphi(q)} \right|^2 \\ = QN \log N + O(QN \log(2N/Q)) + O(N^2(\log N)^{-A})$$

for $Q \leq N$ and $A > 0$ arbitrary. A few years latter, Hooley [17] (1975), by introducing new ideas in treatment of the off-diagonal terms specific to primes, sharpened the right side of (1.3) to

$$QN \log N + O(QN) + O(N^2(\log N)^{-A})$$

with $\Lambda(n)$ replaced by the Chebyshev function $\theta(n)$.

Motohashi [23] (1973), by using an approach similar to Montgomery, elaborately established a more precise asymptotic with lower order and power saving error terms for the divisor function $\tau(n)$. Recently (2018), in a function field variant, the work of Keating, Rodgers, Roditty-Gershon, and Rudnick in [18] leads to the following conjecture (in our notation) over the integers for the variance of $\tau_k(n)$ in arithmetic progressions, with general form for composite moduli put forward by Rodgers and Soundararajan in [25] (c.f. [18, Conjecture 3.3] and [25, Conjecture 1]).

Conjecture 1 (Keating–Rodgers–Roditty–Gershon–Rudnick–Soundararajan). *For $X, d \rightarrow \infty$ such that $\log X / \log d \rightarrow c \in (0, k)$, we have*

$$(1.4) \quad \sum_{\substack{a=1 \\ (a,d)=1}}^d \left| \sum_{\substack{1 \leq n \leq X \\ n \equiv a \pmod{d}}} \tau_k(n) - \frac{1}{\varphi(d)} \sum_{\substack{1 \leq n \leq X \\ (n,d)=1}} \tau_k(n) \right|^2 \sim a_k(d) \gamma_k(c) X (\log d)^{k^2-1},$$

where $a_k(d)$ is the arithmetic constant

$$(1.5) \quad a_k(d) = \lim_{s \rightarrow 1^+} (s-1)^{k^2} \sum_{\substack{n=1 \\ (n,d)=1}}^{\infty} \frac{\tau_k(n)^2}{n^s},$$

and $\gamma_k(c)$ is a piecewise polynomial of degree $k^2 - 1$ defined by

$$(1.6) \quad \gamma_k(c) = \frac{1}{k! G(k+1)^2} \int_{[0,1]^k} \delta_c(w_1 + \cdots + w_k) \Delta(w)^2 d^k w,$$

where $\delta_c(x) = \delta(x - c)$ is a Dirac delta function centered at c , $\Delta(w) = \prod_{i < j} (w_i - w_j)$ is a Vandermonde determinant, and G is the Barnes G -function, so that in particular $G(k+1) = (k-1)!(k-2)! \cdots 1!$.

This conjecture is closely related to the problem of moments of Dirichlet L -functions [5] and correlations of divisor sums [7–11]. In [25], Rodgers and Soundararajan confirmed an averaged version of this conjecture in a restricted range over smooth cutoffs. Harper and Soundararajan in [16] obtained a lower bound of the right order of magnitude for the average of this variance. Nguyen [24, Theorem 3], by using the large sieve inequality, obtained a matching upper bound of the same order of magnitude for this variance when averaged over the moduli d .

More is known about the variance (1.4) when the modulus d is prime. For instance, when $k \geq 3$ and d is prime, Kowalski and Ricotta in 2014 have computed, as one of the many results, [19, Theorem C, pg. 1235] a smoothed version of the variance (1.4), for $c \in (k - \frac{1}{2}, k)$. Their proof uses deep equidistribution result of products of hyper-Kloosterman sums and is based on a Voronoï summation formula for $GL(N)$ twisted by additive characters. This theme of using additive characters to study moments of arithmetic sequences in progressions has also been pursued by de la Bretèche and Fiorilli in [13].

In KRRR ([18]), an analogous conjecture to the variance of divisor sums in arithmetic progressions was also made for divisor sums in short intervals. Lester [20] in 2016 has succeeded in evaluating this variance in the short intervals setting, for $c \in (k-1, k)$, unconditionally for $k=3$ and conditionally on the Lindelöf hypothesis for all $k > 3$. His proof combines two previous methods of Selberg and Jutila, and is based in part on a summation formula for coefficients of L -functions developed in [14] by Friedlander and Iwaniec in 2005.

In this paper, we confirm a smoothed version of the variance (1.4) of $\tau_k(n)$ in arithmetic progressions to any composite moduli d , for $c \in (k-1, k)$. Our procedure follows, with some modifications, that of Lester in his work [20]. We briefly explain here the wider range for c in our result and how we handled the off-diagonal and error terms; see Section 3 below for more detailed discussions.

For ease of exposition, suppose χ is a primitive Dirichlet character modulo a prime p . The functional equation for $L^k(s, \chi)$ reduces a sum of length X to a dual sum of length

X/p^k , which will be less than p when $c > k - 1$ if we write $X = p^c$. So if $n \equiv m \pmod{p}$, and $n, m < p$, then n and m are equal and there are no off-diagonal terms. Thus, since the modulus is prime, all primitive characters modulo p are non-principal, and we have, by orthogonality of all characters, that

$$\frac{1}{\varphi(p)} \sum_{\chi \pmod{p}}^* \chi(n) \overline{\chi}(m) = 1 - \frac{1}{\varphi(p)} = 1 + O(p^{-1}).$$

Hence, roughly speaking, the saving p^{-1} in the error term of the above allows us to take c to be as small as $k - 1$, where the -1 corresponds to the -1 in the exponent of p . Moreover, direct use of summation formulas from [14] when applied to the variance (1.4) yields an error term that is larger than the main term—this is because the absolute value squared in (1.4) requires better than square-root cancellation in the error term, say, E in order for $|E|^2 \ll X^{1-\epsilon}$. We circumvent this by introducing smooth weights in (1.4), and combine this with the methods of Friedlander and Iwaniec in [14] in order to get better control on the error terms. These two ingredients in part lead to the full asymptotic equality, with a power-saving error term, for this smoothed version of the variance of $\tau_k(n)$ in arithmetic progressions unconditionally for all $k \geq 3$, in a restricted range.

Below is the organization of the paper. In the next section, we state this main result as Corollary 1 to Theorem 1. The key ideas of the proofs are outlined in Section 3. Section 4 lists notations, parameters, and conventions used in this paper, which largely follows those in [24]. Preparatory lemmas are collected in Section 5, and the proofs are given in Sections 6 and 7.

2. STATEMENT OF RESULTS AND A CONJECTURE

Let $g(n)$ be an arithmetic function, and $w(y)$ a smooth weight. For $(a, d) = 1$, define

$$\Delta_w(g; X, d, a) := \sum_{n \equiv a \pmod{d}} g(n) w\left(\frac{n}{X}\right) - \frac{1}{\varphi(d)} \sum_{(n, d) = 1} g(n) w\left(\frac{n}{X}\right).$$

Corollary 1 (Asymptotic for leading order main term). *Fix $k \geq 3$. Let $w(y)$ be a smooth function with support in $[1, 2]$ such that*

$$(2.1) \quad \int w(y)^2 dy = 1.$$

Then, uniformly in c for all

$$c \in (k - 1, k),$$

we have the asymptotic

$$(2.2) \quad \sum_{\substack{1 \leq a \leq d \\ (a, d) = 1}} |\Delta_w(\tau_k; X, d, a)|^2 \sim a_k(d) \gamma_k(c) X (\log d)^{k^2 - 1},$$

as $d \rightarrow \infty$, where

$$(2.3) \quad X = d^c,$$

$$a_k(d) = \lim_{s \rightarrow 1^+} (s - 1)^{k^2} \sum_{\substack{n=1 \\ (n, d)=1}}^{\infty} \frac{\tau_k(n)^2}{n^s},$$

and

$$(2.4) \quad \gamma_k(c) = \frac{1}{(k^2 - 1)!} (k - c)^{k^2 - 1}, \quad (k - 1 \leq c < k).$$

Remark 1. By Theorem 1.5 [18, p. 173], Lemma 4.1 [18, p. 180], and the results in Section 4.4.3 [18, pg. 194] of KRRR, the complicated expression for $\gamma_k(c)$ given in (1.6) agrees with the expression (2.4), when $k - 1 \leq c < k$.

Remark 2. We believe that the asymptotic for the smoothed variance (2.2) should hold for all $c \in (0, k)$, with the expression for the piece-wise polynomial $\gamma_k(c)$ given in (1.6). The asymptotic (2.2) for $0 < c < 1$ was essentially established in [25, Lemma 2, pg. 13]. This, together with Corollary 1 above, therefore suggests the following conjecture over smooth cutoffs for the variance of $\tau_k(n)$ in AP's:

Conjecture 2. Let $w(y)$ be a smooth function supported in $[1, 2]$ with

$$\int w(y)^2 dy = 1,$$

and

$$\mathcal{M}[w](\sigma + it) \ll_\ell \frac{1}{1 + |t|^\ell}$$

for all positive integers ℓ , where $\mathcal{M}[w]$ denotes the Mellin transform of w . Then, for $X, d \rightarrow \infty$ such that $\frac{\log X}{\log d} \rightarrow c \in (0, k)$, we have

$$\sum_{\substack{1 \leq a \leq d \\ (a, d) = 1}} \left| \sum_{n \equiv a(d)} \tau_k(n) w\left(\frac{n}{X}\right) - \frac{1}{\varphi(d)} \sum_{(n, d) = 1} \tau_k(n) w\left(\frac{n}{X}\right) \right|^2 \sim a_k(d) \gamma_k(c) X (\log d)^{k^2 - 1},$$

where the arithmetic constant $a_k(d)$ and the piecewise polynomial $\gamma_k(c)$ are given in (1.5) and (1.6), respectively.

The leading order main term of (2.2) is extracted from the full main term (2.7) below, given as a contour integral with a power saving error term. All other lower order main terms of the smoothed variance (2.2) can, in principal, be written down from (2.7).

Theorem 1 (Full main term with power-saving error term). *Fix parameters*

$$(2.5) \quad k \geq 3, \quad \delta > 0,$$

and

$$(2.6) \quad c \in [k - 1 + \delta, k - \delta].$$

For $d \geq 1$, put

$$X = d^c.$$

Let $w(y)$ be a smooth function with support in $[1, 2]$ such that $\int w(y)^2 dy = 1$. Then, we have the asymptotic equality, as $d \rightarrow \infty$,

$$(2.7) \quad \sum_{\substack{1 \leq a \leq d \\ (a,d)=1}} |\Delta_w(\tau_k; X, d, a)|^2 \\ = X \frac{1}{2\pi i} \int_{(2)} \sum_{\substack{d=q^r \\ q>1}} \frac{\varphi^*(q)}{\varphi(d)} \left(\frac{(q/\pi)^k}{X} \right)^{s-1} \sum_{\substack{n=1 \\ (n,r)=1}}^{\infty} \frac{\tau_k(n)^2}{n^s} \mathcal{M}[|f_r|^2](s) ds \\ + O_{\epsilon, \delta, k} \left(X^{1+\epsilon} d^{-\frac{\delta}{2k^2}} \right),$$

for some $\epsilon > 0$, where $\mathcal{M}[|f_r|^2]$ denotes the Mellin transform of $|f_r|^2$, with f_r a complicated inverse Mellin transform of the Gamma factors and $w(y)$, given explicitly in (6.17). The implied constant in the error term of (2.7) is effective, dependent on ϵ, δ, k , and uniform in c .

Remark 3. Because of the ϵ in the error term of (2.7), our method of proof fails to handle the endpoint $c = k - 1$, corresponding to $\delta = 0$. More specifically, this restriction came from bounding the off-diagonal terms; see the bound (6.15). Thus, when $c \leq k - 1$, the off-diagonal terms start contributing to the main term and can no longer be ignored.

Remark 4. It may be possible, by using the asymptotic large sieve of Conrey-Iwaniec-Soundararajan [6] together with the functional equation, to handle the off-diagonal terms and extend the range for c to a slightly wider range

$$c \in \left[k - 1 + \delta - \frac{2}{k}, k - \delta \right]$$

for the smoothed averaged variance

$$\sum_d W \left(\frac{d}{D} \right) \sum_{\substack{1 \leq a \leq d \\ (a,d)=1}} |\Delta_w(\tau_k; X, d, a)|^2$$

with

$$D = X^{1/c}.$$

We hope to return to this question in a future article.

Remark 5. It remains an interesting challenge to remove the smooth weight $w(n/X)$ in (2.7).

We will prove (2.7) first, then derive (2.2) from this.

3. OUTLINE OF THE PROOFS

Since the proofs are quite lengthy and technical in details, we shall divide the proofs into smaller steps and outline in this section the key ideas in each step.

Outline of the proof of Theorem 1

The main idea in the proof of (2.7) is to develop a smoothed version of Voronoï summation twisted by multiplicative characters, by applying the functional equation to transform the original sum of length X , which larger than d^{k-1} , to a shorter (dual) sum of length less than

d . The variance (2.7) is easily handled when $X < d$, corresponding to $c < 1$, because, in this range, only the diagonal terms contribute to the leading main term. When $X > d$, the functional equation converts a sum of length X to a dual sum of length d^k/X , which will be less than d when $c > k - 1$. To carry this out, we first rewrite the congruence relation $n \equiv a \pmod{d}$ as a sum over non-principal characters $\chi \pmod{d}$. We next need to reduce non-principal characters to primitive ones, in order to apply the functional equation. This key reduction introduces extra coprime conditions that makes the details more tricky, but still manageable. The functional equation is also crucial for bounding the error terms, since, roughly speaking, it converts the factor X^σ , which is huge when $\sigma > 1$, into a large power saving $X^{1-\sigma}$. (!) Moreover, since we allow ourselves with smooth weights, which have rapid decay in the vertical direction, we need not strictly work inside the critical strip, and, thus, dispensing with the Lindelöf hypothesis, making the proofs unconditional.

We break the proof of (2.7) into several steps.

Step 1: Applying the functional equation and reduction in q . We first rewrite the variance (2.7) as a sum over primitive characters and apply the functional equation. We show that contributions from characters with modulus $q < Q$ are negligible. It is here that the parameter Q is chosen, so that the error from removing these characters are small.

Step 2: Reduction to the dual sum. Assume $q \geq Q$. Introduce parameter N . Show that contributions from terms with $n > N$ are acceptable by pushing the line of integration far to the right.

Step 3: Off-diagonal analysis. Assume $q \geq Q$ and $n \leq N$. Estimate the off-diagonal contributions and show they do not exceed the main term. The key tool in this step is Shiu's bound (5.1). It is here that sets the lower bound for the range of c in Theorem 1.

Step 4: Diagonal analysis. Assume $q \geq Q$ and $n \leq N$. We apply Mellin inversion to extract the contribution from the diagonal terms as a contour integral.

Outline of the proof of Corollary 1

We extract the leading order term in the full main term from Theorem 1 and also need to show that the lower order main terms are indeed of lower order than the leading main term. A certain arithmetic function arises which is then evaluated to match the arithmetic factor $a_k(d)$. Below are the steps in the proof of Corollary 1.

Step a: Full main term as a residue plus an error term. Write the contour integral from (2.7) as a residue plus an error term. This is fairly straightforward.

Step b: The error term and $\gamma_k(c)$. Show that contribution from the error term in **Step a** is small. It is here that the condition $q \geq Q$ is needed.

Step c: Bounding lower order main terms. Write the residue from **Step a** as a polynomial, and show that contributions from lower order terms are of order of magnitude smaller than the leading term.

Step d: Evaluating the matching arithmetic constant $a_k(d)$. Evaluate the contribution from the leading term of the residue from **Step a** and verifying that the leading constants match.

4. NOTATIONS, PARAMETERS, AND CONVENTIONS

i, j, ℓ —non-negative integers.

p, p_i, p_j —are prime numbers.

$d, a, n, m, k, q, r, s, \alpha_i, \beta_j, Q, T, N$ —positive integers.

X —a large real number.

$\Lambda(n)$ —the von Mangoldt function.

$\tau_k(n)$ —the k -fold divisor function; $\tau_2(n) = \tau(n)$, the usual divisor function.

$\varphi(n)$ —the Euler’s totient function.

$\mu(n)$ —the Möbius function.

$s = \sigma + it$

$\Gamma(s)$ —the Gamma function.

$\int_{(\sigma)}$ —means $\int_{\sigma-i\infty}^{\sigma+i\infty}$.

$\mathcal{M}[f](s)$ —the Mellin transform of a suitable function f , i.e.,

$$\mathcal{M}[f](s) = \int_0^\infty f(x)x^{s-1}dx.$$

$\mathcal{M}^{-1}[F](x)$ —the inverse Mellin transform of a suitable function $F(s)$, i.e.,

$$\mathcal{M}^{-1}[F](x) = \frac{1}{2\pi i} \int_{(\sigma)} F(s)x^{-s}ds, \quad (\sigma > 1).$$

$\chi(n)$ —a Dirichlet character.

$\bar{\chi}(n)$ —means the complex conjugate of $\chi(n)$.

$\tau(\chi)$ —the Gauss sum of the character χ .

$m \equiv a(d)$ means $m \equiv a \pmod{d}$.

$\alpha \star \beta$ means the Dirichlet convolution of α and β , i.e.,

$$(\alpha \star \beta)(d) = \sum_{d=qr} \alpha(q)\beta(r).$$

$\sum'_{\chi(\bmod d)}$ —means a summation over nonprincipal characters $\chi(\bmod d)$.

$\sum^*_{\chi(\bmod d)}$ —means a summation over primitive characters $\chi(\bmod d)$.

$\sum_{b(\bmod d)}$ —means $\sum_{b=1}^d$.

$\sum^*_{b(\bmod d)}$ —means $\sum_{\substack{b=1 \\ (b,d)=1}}^d$.

ϵ —any sufficiently small, positive constant, not necessarily the same in each occurrence.

δ —a fixed small positive number.

A —any sufficiently large, positive constant, not necessarily the same in each occurrence.

B —some large positive constant, not necessarily the same in each occurrence.

We follow standard notations and write $f(X) = O(g(X))$ or $f(X) \ll g(X)$ to mean that $|f(X)| \leq Cg(X)$ for some fixed constant C , and $f(X) = o(g(X))$ if $|f(X)| \leq c(X)g(X)$ for some function $c(X)$ that goes to zero as X goes to infinity.

We begin the proofs with some preliminary lemmata.

TABLE 1. Table of parameters and their first appearance.

Parameters	First appearance
$k \geq 3$	
$\delta > 0$	(2.5)
$c \in [k - 1 + \delta, k - \delta]$	(2.6)
$X = d^c$	(2.3)
$Q = d^{\frac{c}{k} + \frac{\delta}{k^3}}$	(6.3)
$N = d$	(6.9)
$Y = d^{k-c}$	(6.12)
$P = (\log d)^2$	(7.9)

5. PREPARATORY LEMMAS

Lemma 1 (Shiu's bound). *Suppose that $1 \leq N < N' < 2X$, $N' - N > X^\epsilon d$, and $(a, d) = 1$. Then for $j, \nu \geq 1$ we have*

$$(5.1) \quad \sum_{\substack{N \leq n \leq N' \\ n \equiv a(d)}} \tau_j(n)^\nu \ll \frac{N' - N}{\varphi(d)} (\log X)^{j\nu - 1}.$$

The implied constants depending on ϵ, j , and ν at most.

Proof. See [26, Theorem 2]. □

Lemma 2. *We have*

$$(5.2) \quad \sum_{\substack{1 \leq a \leq d \\ (a, d) = 1}} |\Delta_w(\tau_k; X, d, a)|^2 = \frac{1}{\varphi(d)} \sum'_{\chi(\bmod d)} \left| \sum_n \tau_k(n) \chi(n) w\left(\frac{n}{X}\right) \right|^2.$$

Proof. By the orthogonality relation

$$\frac{1}{\varphi(d)} \sum_{\chi(\bmod d)} \chi(n) \bar{\chi}(a) = \begin{cases} 1, & \text{if } n \equiv a(d), \\ 0, & \text{otherwise,} \end{cases}$$

we have

$$\Delta_w(\tau_k; X, d, a) = \frac{1}{\varphi(d)} \sum'_{\chi(\bmod d)} \bar{\chi}(a) \sum_n \tau_k(n) \chi(n) w\left(\frac{n}{X}\right).$$

Thus, the left side of (5.2) can be written as

$$(5.3) \quad \sum_{\substack{1 \leq a \leq d \\ (a, d) = 1}} \frac{1}{\varphi(d)^2} \sum'_{\chi_1(\bmod d)} \bar{\chi}_1(a) \sum_n \tau_k(n) \chi_1(n) w\left(\frac{n}{X}\right) \sum'_{\chi_2(\bmod d)} \chi_2(a) \sum_m \tau_k(m) \bar{\chi}_2(m) w\left(\frac{m}{X}\right) \\ = \frac{1}{\varphi(d)^2} \sum'_{\chi_1, \chi_2(\bmod d)} \sum_{n, m} \tau_k(n) \chi_1(n) w\left(\frac{n}{X}\right) \tau_k(m) \bar{\chi}_2(m) w\left(\frac{m}{X}\right) \sum_{\substack{1 \leq a \leq d \\ (a, d) = 1}} \bar{\chi}_1(a) \chi_2(a).$$

By the relation

$$\sum_{\substack{1 \leq a \leq d \\ (a,d)=1}} \bar{\chi}_1(a) \chi_2(a) = \begin{cases} \varphi(d), & \text{if } \chi_1 = \chi_2, \\ 0, & \text{otherwise,} \end{cases}$$

the right side of (5.3) is equal to

$$\frac{1}{\varphi(d)} \sum'_{\chi(\bmod d)} \sum_{n,m} \tau_k(n) \chi(n) w\left(\frac{n}{X}\right) \tau_k(m) \bar{\chi}(m) w\left(\frac{m}{X}\right)$$

which is equal to the right side of (5.2). \square

We next reduce non-principal to primitive characters.

Lemma 3. *We have*

$$(5.4) \quad \sum'_{\chi(\bmod d)} \left| \sum_n \tau_k(n) \chi(n) w\left(\frac{n}{X}\right) \right|^2 = \sum_{\substack{d=qr \\ q>1}} \sum^*_{\chi_1(\bmod q)} \left| \sum_{(n,r)=1} \tau_k(n) \chi_1(n) w\left(\frac{n}{X}\right) \right|^2.$$

Proof. If $\chi(\bmod d)$ is non-principal, then there is a unique $q \mid d$ with $q > 1$, and a unique primitive character $\chi_1(\bmod q)$, such that, with $r = d/q$,

$$(5.5) \quad \chi(n) = \chi_1(n) \chi_{0,r}(n),$$

where $\chi_{0,r}$ is the principal character modulo r . Thus, replacing the sum $\sum'_{\chi(\bmod d)}$ by $\sum_{\substack{d=qr \\ q>1}} \sum^*_{\chi_1(\bmod q)}$

and substituting (5.5) into the left side of (5.4) give the right side of (5.4). \square

Lemma 4. *For $(mn, q_1) = 1$, we have*

$$(5.6) \quad \sum^*_{\chi(\bmod q_1)} \chi(m) \bar{\chi}(n) = \sum_{\substack{q_1=q_2 r_2 \\ r_2 \mid m-n}} \mu(q_2) \varphi(r_2).$$

Proof. See, e.g., [4, Lemma 2.7, pg. 348]. \square

The next is the well-known functional equation for primitive Dirichlet L -functions.

Theorem 2. *Let $\chi(\bmod q)$ be a primitive character. Then, for $\sigma > 1$, we have*

$$(5.7) \quad L^k(1-s, \chi) = \gamma(s, \chi) L^k(s, \bar{\chi}),$$

where

$$\gamma(s, \chi) = \varepsilon \left(\frac{q}{\pi}\right)^{s-\frac{1}{2}} \frac{\Gamma\left(\frac{s+a}{2}\right)}{\Gamma\left(\frac{1-s+a}{2}\right)}, \quad |\varepsilon| = 1,$$

with

$$a = \begin{cases} 1, & \text{if } \chi(-1) = -1, \\ 0, & \text{if } \chi(-1) = 1. \end{cases}$$

Lemma 5. *We have, for any $k \geq 2$,*

$$(5.8) \quad \tau_k(p^j) = \binom{k+j-1}{k-1}.$$

Proof. By the relation

$$\left(\frac{1}{1-z}\right)^k = \sum_{j=0}^{\infty} \binom{k+j-1}{j} z^j, \quad (|z| < 1),$$

and the Euler product

$$\zeta(s) = \prod_p \frac{1}{1-p^{-s}}, \quad (\sigma > 1),$$

we have

$$(5.9) \quad \zeta^k(s) = \prod_p \left(\frac{1}{1-p^{-s}}\right)^k = \prod_p \sum_{j=0}^{\infty} \binom{k+j-1}{j} p^{-js}, \quad (\sigma > 1).$$

We also have

$$(5.10) \quad \zeta^k(s) = \sum_{n=1}^{\infty} \frac{\tau_k(n)}{n^s} = \prod_p \sum_{j=0}^{\infty} \tau_k(p^j) p^{-js}, \quad (\sigma > 1).$$

Thus, by (5.9) and (5.10),

$$\tau_k(p^j) = \binom{k+j-1}{k-1}.$$

□

Lemma 6. *We have*

$$(5.11) \quad \sum_{j=0}^{\infty} \frac{\tau_k(n)^2}{p^{js}} = \left(1 - \frac{1}{p^s}\right)^{-(2k-1)} \sum_{j=0}^{k-1} \frac{\binom{k-1}{j}^2}{p^{js}}.$$

Proof. We have the identity

$$(5.12) \quad \sum_{j=0}^{\infty} \frac{\binom{k+j-1}{k-1}^2}{p^{js}} = \left(1 - \frac{1}{p^s}\right)^{-(2k-1)} \sum_{j=0}^{k-1} \frac{\binom{k-1}{j}^2}{p^{js}}.$$

This identity (5.12) follows from a hypergeometric relation; see, e.g., [19, Proposition C.2, pg. 1295]. By (5.8), the left side of (5.12) is that of (5.11). □

The next lemma is based on Parseval formula for Mellin transforms, see, e.g., [28, Theorem 1.17, pg. 33], and makes use of the condition (2.1) in statement of Corollary 1.

Lemma 7. *We have, with f_r defined as in (6.16),*

$$(5.13) \quad \mathcal{M}[|f_r|^2](1) = 1.$$

Proof. By definition of the Mellin transform and Parseval formula for the Mellin transform, we have

$$\mathcal{M}[|f_r|^2](1) = \int_0^{\infty} f_r(y, \chi_1) \overline{f_r(y, \chi_1)} dy = \frac{1}{2\pi i} \int_{(2)} \mathcal{M}[f_r](s) \mathcal{M}[\overline{f_r}](1-s) ds.$$

By definition of the Mellin transform again and by (6.16), the above is equal to

$$\begin{aligned} & \frac{1}{2\pi i} \int_{(2)} \mathcal{M}[w](1-s) \frac{\Gamma^k\left(\frac{s+a}{2}\right)}{\Gamma^k\left(\frac{1-s+a}{2}\right)} \prod_{p|r} \left(\frac{1 - \frac{\chi_1(p)}{p^{1-s}}}{1 - \frac{\bar{\chi}_1(p)}{p^s}} \right)^k \mathcal{M}[w](s) \frac{\Gamma^k\left(\frac{1-s+a}{2}\right)}{\Gamma^k\left(\frac{s+a}{2}\right)} \prod_{p|r} \left(\frac{1 - \frac{\bar{\chi}_1(p)}{p^s}}{1 - \frac{\chi_1(p)}{p^{1-s}}} \right)^k ds \\ &= \frac{1}{2\pi i} \int_{(2)} \mathcal{M}[w](1-s) \mathcal{M}[w](s) ds = \int_0^\infty |w(x)|^2 dx, \end{aligned}$$

where the last equality follows by Parseval formula for Mellin transforms once more time. By (2.1), this gives the right side of (5.13). \square

6. PROOF OF THEOREM 1 FULL MAIN TERM WITH POWER-SAVING ERROR TERM

6.1. Step 1: Applying the functional equation and reduction in q . We first rewrite the variance (2.7) as a sum over primitive characters, in order to apply the functional equation. By Lemmas 2 and 3, we have

$$\sum_{\substack{1 \leq a \leq d \\ (a,d)=1}} |\Delta_w(\tau_k; X, d, a)|^2 = \frac{1}{\varphi(d)} \sum_{\substack{d=qr \\ q>1}} \sum_{\chi_1 \pmod{q}}^* \left| \sum_{(n,r)=1} \tau_k(n) \chi_1(n) w\left(\frac{n}{X}\right) \right|^2.$$

Substituting

$$w\left(\frac{n}{X}\right) = \frac{1}{2\pi i} \int_{(\sigma)} \mathcal{M}[w](s) \left(\frac{n}{X}\right)^{-s} ds, \quad (\sigma > 1),$$

into the above and interchanging the order of summation and integration, the left side of (2.7) can be written as

$$(6.1) \quad \frac{1}{\varphi(d)} \sum_{\substack{d=qr \\ q>1}} \sum_{\chi_1 \pmod{q}}^* \left| \frac{1}{2\pi i} \int_{(\sigma)} \mathcal{M}[w](s) X^s \sum_{(n,r)=1} \frac{\tau_k(n) \chi_1(n)}{n^s} ds \right|^2, \quad (\sigma > 1).$$

Since $q > 1$, χ_1 modulo q is non-principal, thus, the function $\sum_{(n,r)=1} \tau_k(n) \chi_1(n) n^{-s}$ has no poles. Moreover, since zero is not in the support of $w(y)$, the Mellin transform $\mathcal{M}[w](s)$ also has no poles. Hence, the integrand in (6.1) is entire and we may shift the line of integration to the left of the one-line.

Shifting the line of integration to $\sigma = -\epsilon$, applying the functional equation (5.7), and making a change of variables s to $1-s$, (6.1) is equal to

$$(6.2) \quad \frac{1}{\varphi(d)} \sum_{\substack{d=qr \\ q>1}} \sum_{\chi_1 \pmod{q}}^* \left| \frac{1}{2\pi i} \int_{(1+\epsilon)} \mathcal{M}[w](1-s) X^{1-s} \gamma^k(s, \chi_1) L^k(s, \bar{\chi}_1) \prod_{p|r} \left(1 - \frac{\chi_1(p)}{p^{1-s}} \right)^k ds \right|^2.$$

Let

$$(6.3) \quad Q := d^{\frac{\epsilon}{k} + \frac{\delta}{k^3}}.$$

We will now show that contributions from characters with modulus $q < Q$ are negligible. Suppose

$$(6.4) \quad q < Q.$$

By repeated integration by parts and rapid decay of derivatives of the smooth weight $w(y)$, we have

$$\mathcal{M}[w](1-s) \ll \frac{1}{(1+|t|)^A}$$

for any $A > 0$. Also, for $s = 1 + \epsilon + it$, we have

$$\gamma^k(s, \chi_1) \ll q^{k(\frac{1}{2} + \epsilon)} \ll d^{\frac{c}{2} + \frac{\delta}{2k^2} + \frac{\epsilon\delta}{k^2} + \epsilon c},$$

$$L^k(s, \bar{\chi}_1) \ll 1,$$

$$\prod_{p|r} \left(1 - \frac{\chi_1(p)}{p^{1-s}}\right)^k \ll r^\epsilon.$$

Thus, by the above estimates, the integral in (6.2) is

$$\ll d^{\frac{c}{2} + \frac{\delta}{2k^2} + \frac{\epsilon\delta}{k^2} + \epsilon}.$$

Hence,

$$\begin{aligned} (6.5) \quad & \frac{1}{\varphi(d)} \sum_{\substack{d=qr \\ 1 < q < Q}} \sum_{\chi_1 \pmod{q}}^* \left| \frac{1}{2\pi i} \int_{(1+\epsilon)} \mathcal{M}[w](1-s) X^{1-s} \gamma^k(s, \chi_1) L^k(s, \bar{\chi}_1) \prod_{p|r} \left(1 - \frac{\chi_1(p)}{p^{1-s}}\right)^k ds \right|^2 \\ & \ll X d^{\frac{\delta}{k^2} + \frac{2\epsilon\delta}{k^2} + \epsilon} \frac{1}{\varphi(d)} \sum_{\substack{d=qr \\ 1 < q < Q}} \varphi^*(q). \end{aligned}$$

By (6.4), the crude bounds $\tau(d) = d^\epsilon$, $\frac{1}{\varphi(d)} \ll d^{-1+\epsilon}$, and $\varphi^*(q) \ll q$, we have

$$\frac{1}{\varphi(d)} \sum_{\substack{d=qr \\ 1 < q < Q}} \varphi^*(q) \ll d^{-\frac{\delta}{k} + \frac{\delta}{k^2} + \epsilon}$$

for all $c \leq k - \delta$. With this, (6.5) is

$$\ll X d^{-\frac{\delta}{k} + \frac{2\delta}{k^2} + \frac{2\epsilon\delta}{k^2} + \epsilon},$$

which is acceptable for (2.7). It thus suffices to assume $q \geq Q$.

6.2. Step 2: Reduction to the dual sum. Suppose $q \geq Q$. We now introduce the parameter N and truncate the n sum in $L^k(s, \bar{\chi}_1)$. Terms with $n \leq N$ account for the main term, and terms with $n > N$ will be shown in this step to be small. By Euler products, we have

$$\sum_{\substack{n \leq N \\ (n,r)=1}} \frac{\tau_k(n) \bar{\chi}_1(n)}{n^s} = \prod_{p|r} \left(1 - \frac{\bar{\chi}_1(p)}{p^s}\right)^k L^k(s, \bar{\chi}_1), \quad (\sigma > 1).$$

Thus, writing

$$(6.6) \quad L^k(s, \bar{\chi}_1) = \prod_{p|r} \left(1 - \frac{\bar{\chi}_1(p)}{p^s}\right)^{-k} \left(\sum_{\substack{n \leq N \\ (n,r)=1}} \frac{\tau_k(n) \bar{\chi}_1(n)}{n^s} + \sum_{\substack{n > N \\ (n,r)=1}} \frac{\tau_k(n) \bar{\chi}_1(n)}{n^s} \right),$$

for $L^k(s, \bar{\chi}_1)$ in (6.2), with N to be chosen below, we have, by (6.6), (6.2), and (6.1),

(6.7)

$$\begin{aligned} \sum_{\substack{1 \leq a \leq d \\ (a,d)=1}} |\Delta_w(\tau_k; X, d, a)|^2 &= \frac{1}{\varphi(d)} \sum_{\substack{d=qr \\ q \geq Q}} \sum_{\chi_1 \pmod{q}}^* \left| \frac{1}{2\pi i} \int_{(1+B)} \left(\sum_{\substack{n \leq N \\ (n,r)=1}} + \sum_{\substack{n > N \\ (n,r)=1}} \right) \frac{\tau_k(n) \bar{\chi}_1(n)}{n^s} \right. \\ &\quad \left. \times \mathcal{M}[w](1-s) X^{1-s} \gamma^k(s, \chi_1) \prod_{p|r} \left(\frac{1 - \frac{\chi_1(p)}{p^{1-s}}}{1 - \frac{\bar{\chi}_1(p)}{p^s}} \right)^k ds \right|^2 + O\left(X^{1+\epsilon} d^{-\frac{\delta}{2k^2}}\right), \end{aligned}$$

where we have moved the line of integration to $\sigma = 1 + B$, $B > 0$. Analogous to **Step 1**, we will show contributions from terms with $n > N$ in the above are negligible. The main difference in this step is that there is an additional saving from the sum $\sum_{n > N} n^{-\sigma+\epsilon}$. We have, for $s = 1 + B + it$,

$$\sum_{\substack{n > N \\ (n,r)=1}} \frac{\tau_k(n) \bar{\chi}_1(n)}{n^s} \ll N^{-B+\epsilon},$$

$$X^{1-s} \ll X^{-B},$$

$$\gamma^k(s, \chi_1) \ll q^{k(B+\frac{1}{2})},$$

$$L^k(s, \bar{\chi}_1) \ll 1,$$

$$\prod_{p|r} \left(1 - \frac{\chi_1(p)}{p^{1-s}}\right)^k \left(1 - \frac{\bar{\chi}_1(p)}{p^s}\right)^{-k} \ll r^{kB+\epsilon}.$$

Thus, by the above estimates, the contribution from terms with $n > N$ in the integral on the right side of (6.7) is

$$(6.8) \quad \ll N^{-B+\epsilon} X^{-B} q^{k(B+\frac{1}{2})} r^{kB+\epsilon}.$$

We choose

$$(6.9) \quad N = d$$

and

$$(6.10) \quad B = \frac{A}{2\delta} + \frac{k}{4\delta}.$$

With the choices (6.9) and (6.10), the estimate (6.8) is

$$\ll d^{-A/2},$$

for any $A > 0$. This bound shows that the contribution from terms with $n > N$ on the right side of (6.7) can be absorbed into the big-oh term.

We are now reduced to terms with $q \geq Q$ and $n \leq N$. To summarize, we now have

(6.11)

$$\begin{aligned} & \sum_{\substack{1 \leq a \leq d \\ (a,d)=1}} |\Delta_w(\tau_k; X, d, a)|^2 \\ &= \frac{1}{\varphi(d)} \sum_{\substack{d=qr \\ q \geq Q}} \sum_{\chi_1 \pmod{q}}^* \left| \sum_{\substack{n \leq N \\ (n,r)=1}} \tau_k(n) \bar{\chi}_1(n) \frac{1}{2\pi i} \int_{(1+B)} \mathcal{M}[w](1-s) \frac{X}{(nX)^s} \gamma^k(s, \chi_1) \prod_{p|r} \left(\frac{1 - \frac{\chi_1(p)}{p^{1-s}}}{1 - \frac{\bar{\chi}_1(p)}{p^s}} \right)^k ds \right|^2 \\ &+ O\left(X^{1+\epsilon} d^{-\frac{\delta}{2k^2}}\right). \end{aligned}$$

For the evaluation of the main term, we will move the line of integration to the critical line $s = 1/2 + it$.

6.3. Step 3: Off-diagonal analysis. Let $s = 1/2 + it$. We expand out the square on the right side of (6.11) and first treat the off-diagonal terms. Let

$$(6.12) \quad XY = d^k.$$

We write

$$\frac{X}{(nX)^s} \gamma^k(s, \chi_1) = \varepsilon^k \frac{X^{1/2} r^{k/2} \pi^{k/2}}{Y^{1/2}} \left(\frac{nr^k \pi^k}{Y} \right)^{-s} \frac{\Gamma\left(\frac{s+a}{2}\right)^k}{\Gamma\left(\frac{1-s+a}{2}\right)^k},$$

and

$$\begin{aligned} \prod_{p|r} \left(\frac{1 - \frac{\chi_1(p)}{p^{1-s}}}{1 - \frac{\bar{\chi}_1(p)}{p^s}} \right)^k &= \sum_{i_1, \dots, i_\ell=0}^k \sum_{j_1, \dots, j_\ell=0}^\infty \binom{k}{i_1} \cdots \binom{k}{i_\ell} \binom{-k}{j_1} \cdots \binom{-k}{j_\ell} (-1)^{i_1 + \dots + j_\ell} \\ &\times \frac{\chi_1(p_1^{i_1} \cdots p_\ell^{i_\ell}) \bar{\chi}_1(p_1^{j_1} \cdots p_\ell^{j_\ell})}{p_1^{i_1(1-s)} \cdots p_\ell^{i_\ell(1-s)} p_1^{j_1 s} \cdots p_\ell^{j_\ell s}}. \end{aligned}$$

Denote

$$\begin{aligned} f_r(y) &= \sum_{i_1, \dots, i_\ell=0}^k \sum_{j_1, \dots, j_\ell=0}^\infty \binom{k}{i_1} \cdots \binom{k}{i_\ell} \binom{-k}{j_1} \cdots \binom{-k}{j_\ell} (-1)^{i_1 + \dots + j_\ell} \frac{p_1^{i_1(s-1)} \cdots p_\ell^{i_\ell(s-1)}}{p_1^{j_1 s} \cdots p_\ell^{j_\ell s}} \\ &\times \frac{1}{2\pi i} \int_{(\frac{1}{2})} \mathcal{M}[w](1-s) y^{-s} \frac{\Gamma\left(\frac{s+a}{2}\right)^k}{\Gamma\left(\frac{1-s+a}{2}\right)^k} ds. \end{aligned}$$

We write the off-diagonal terms $n \neq m$ on the right side of (6.11) as

$$(6.13) \quad \frac{X}{Y} \frac{1}{\varphi(d)} \sum_{\substack{d=qr \\ q \geq Q}} \pi^k r^k \sum_{\substack{n \leq N \\ (n,r)=1}} \tau_k(n) f_r \left(\frac{nr^k \pi^k}{Y} \right) \sum_{\substack{m \leq N \\ (m,r)=1 \\ m \neq n}} \overline{\tau_k(m) f_r \left(\frac{mr^k \pi^k}{Y} \right)} \\ \times \sum_{\chi_1(\bmod q)}^* \bar{\chi}_1(np_1^{j_1} \cdots p_\ell^{j_\ell}) \chi_1(mp_1^{i_1} \cdots p_\ell^{i_\ell}).$$

We have $(np_1^{j_1} \cdots p_\ell^{j_\ell}, q) = 1$ and $(mp_1^{i_1} \cdots p_\ell^{i_\ell}, q) = 1$. Thus, by (5.6),

$$\sum_{\chi_1(\bmod q)}^* \bar{\chi}_1(np_1^{j_1} \cdots p_\ell^{j_\ell}) \chi_1(mp_1^{i_1} \cdots p_\ell^{i_\ell}) = \sum_{\delta | (q, np_1^{j_1} \cdots p_\ell^{j_\ell} - mp_1^{i_1} \cdots p_\ell^{i_\ell})} \varphi(\delta) \mu \left(\frac{q}{\delta} \right).$$

Let

$$b := p_1^{j_1} \cdots p_\ell^{j_\ell} \overline{p_1^{i_1} \cdots p_\ell^{i_\ell}},$$

where the bar denotes inverse modulo δ . Writing the condition $\delta | np_1^{j_1} \cdots p_\ell^{j_\ell} - mp_1^{i_1} \cdots p_\ell^{i_\ell}$ as $m \equiv nb(\bmod \delta)$, (6.13) can be rewritten as

$$(6.14) \quad \frac{X}{Y} \frac{1}{\varphi(d)} \sum_{\substack{d=qr \\ q \geq Q}} \pi^k r^k \sum_{\delta | q} \varphi(\delta) \mu \left(\frac{q}{\delta} \right) \sum_{\substack{n \leq N \\ (n,r)=1}} \tau_k(n) f_r \left(\frac{nr^k \pi^k}{Y} \right) \sum_{\substack{m \leq N \\ (m,r)=1 \\ m \neq n \\ m \equiv nb(\bmod \delta)}} \overline{\tau_k(m) f_r \left(\frac{mr^k \pi^k}{Y} \right)}.$$

The m sum in the above is bounded by

$$\sum_{\substack{m \leq N \\ (m,r)=1 \\ m \equiv nb(\bmod \delta)}} \tau_k(m) \left| \overline{f_r \left(\frac{mr^k \pi^k}{Y} \right)} \right| \ll r^\epsilon Y^\epsilon \sum_{\substack{m \leq N \\ m \equiv nb(\bmod \delta)}} \tau_k(m) \frac{Y^{1/2}}{m^{1/2} r^{k/2}} \ll \frac{1}{\varphi(\delta)} \frac{Y^{1+\epsilon} r^\epsilon}{r^{k/2}},$$

by Shiu's bound (5.1). With this estimate, (6.14) is bounded by

$$(6.15) \quad \frac{X}{Y} \frac{1}{\varphi(d)} \sum_{\substack{d=qr \\ q \geq Q}} r^k \tau(q) \sum_{\substack{n \leq N \\ (n,r)=1}} \tau_k(n) f_r \left(\frac{nr^k \pi^k}{Y} \right) \frac{Y^{1+\epsilon} r^\epsilon}{r^{k/2}} \ll X \frac{Y^{1+\epsilon}}{\varphi(d)} \ll X d^{-\delta+\epsilon}.$$

This estimate for the off-diagonal terms on the right side of (6.11) is acceptable for (2.7). We last treat the diagonal terms in (6.11).

6.4. Step 4: Diagonal analysis. Let

$$(6.16) \quad f_r(y, \chi_1) = \frac{1}{2\pi i} \int_{(\sigma)} \mathcal{M}[w](1-s) y^{-s} \frac{\Gamma\left(\frac{s+a}{2}\right)^k}{\Gamma\left(\frac{1-s+a}{2}\right)^k} \prod_{p|r} \left(\frac{1 - \frac{\chi_1(p)}{p^{1-s}}}{1 - \frac{\bar{\chi}_1(p)}{p^s}} \right)^k ds, \quad (\sigma > 1).$$

This function can also be written as an inverse Mellin transform as

$$(6.17) \quad f_r(y, \chi_1) = \mathcal{M}^{-1} \left[s \mapsto \mathcal{M}[w](1-s) \frac{\Gamma\left(\frac{s+a}{2}\right)^k}{\Gamma\left(\frac{1-s+a}{2}\right)^k} \prod_{p|r} \left(\frac{1 - \frac{\chi_1(p)}{p^{1-s}}}{1 - \frac{\bar{\chi}_1(p)}{p^s}} \right)^k \right].$$

We thus have

$$\begin{aligned} & \sum_{\substack{1 \leq a \leq d \\ (a,d)=1}} \left| \sum_{n \equiv a(d)} \tau_k(n) w\left(\frac{n}{X}\right) - \frac{1}{\varphi(d)} \sum_{(n,d)=1} \tau_k(n) w\left(\frac{n}{X}\right) \right|^2 \\ &= X^2 \frac{1}{\varphi(d)} \sum_{\substack{d=qr \\ q \geq Q}} \frac{\pi^k}{q^k} \sum_{\chi_1 \pmod{q}}^* \sum_{\substack{n \leq N \\ (n,r)=1}} \tau_k(n)^2 \left| f_r\left(\frac{nX\pi^k}{q^k}, \chi_1\right) \right|^2 + O\left(X^{1+\epsilon} d^{-\frac{\delta}{2k^2}}\right). \end{aligned}$$

We also have

$$(6.18) \quad \left| f_r\left(\frac{nX\pi^k}{q^k}, \chi_1\right) \right|^2 \ll \left(\frac{q^k}{nX}\right)^{2B+2} r^{2kB+\epsilon},$$

for any $B > 0$. We now write

$$\sum_{\substack{n \leq N \\ (n,r)=1}} \tau_k(n)^2 \left| f_r\left(\frac{nX\pi^k}{q^k}, \chi_1\right) \right|^2 = \left(\sum_{\substack{n=1 \\ (n,r)=1}}^{\infty} - \sum_{\substack{n > N \\ (n,r)=1}} \right) \tau_k(n)^2 \left| f_r\left(\frac{nX\pi^k}{q^k}, \chi_1\right) \right|^2.$$

For terms $n > N$ in the above, we have, by (6.18),

$$\begin{aligned} & X^2 \frac{1}{\varphi(d)} \sum_{\substack{d=qr \\ q \geq Q}} \frac{\pi^k}{q^k} \sum_{\chi_1 \pmod{q}}^* \sum_{\substack{n > N \\ (n,r)=1}} \tau_k(n)^2 \left| f_r\left(\frac{nX\pi^k}{q^k}, \chi_1\right) \right|^2 \\ & \ll d^{2kB+k} N^{-(2B+1)} X^{-2B} \ll d^{-2B\delta+k}, \end{aligned}$$

for any $B > 0$. This amount is admissible for (2.7).

Finally, we have, by inverse Mellin transform of f_r ,

$$\sum_{\substack{n=1 \\ (n,r)=1}}^{\infty} \tau_k(n)^2 \left| f_r\left(\frac{nX\pi^k}{q^k}, \chi_1\right) \right|^2 = \frac{1}{2\pi i} \int_{(\sigma)} \mathcal{M}[|f_r|^2](s) \left(\frac{q^k}{X\pi^k}\right)^s \sum_{\substack{n=1 \\ (n,r)=1}}^{\infty} \frac{\tau_k(n)^2}{n^s} ds,$$

for $\sigma > 1$. Therefore,

$$(6.19) \quad \begin{aligned} & \sum_{\substack{1 \leq a \leq d \\ (a,d)=1}} |\Delta_w(\tau_k; X, d, a)|^2 \\ &= X^2 \frac{1}{2\pi i} \int_{(\sigma)} \sum_{\substack{d=qr \\ q \geq Q}} \frac{\varphi^*(q)}{\varphi(d)} \frac{\pi^k}{q^k} \left(\frac{q^k}{X\pi^k}\right)^s \sum_{\substack{n=1 \\ (n,r)=1}}^{\infty} \frac{\tau_k(n)^2}{n^s} \mathcal{M}[|f_r|^2](s) ds + O\left(X^{1+\epsilon} d^{-\frac{\delta}{2k^2}}\right). \end{aligned}$$

This proves (2.7). □

7. PROOF OF COROLLARY 1 ASYMPTOTIC FOR LEADING ORDER MAIN TERM

7.1. Step a: Full main term as a residue plus an error term. We start with (6.19), which we rewrite as

$$(7.1) \quad \sum_{\substack{1 \leq a \leq d \\ (a,d)=1}} |\Delta_w(\tau_k; X, d, a)|^2 \\ = X \sum_{\substack{d=qr \\ q \geq Q}} \frac{\varphi^*(q)}{\varphi(d)} \frac{1}{2\pi i} \int_{(2)} \left(\frac{(q/\pi)^k}{X} \right)^{s-1} \sum_{\substack{n=1 \\ (n,r)=1}}^{\infty} \frac{\tau_k(n)^2}{n^s} \mathcal{M}[|f_r|^2](s) ds + O_{\epsilon, \delta, k} \left(X^{1+\epsilon} d^{-\frac{\delta}{2k^2}} \right).$$

The integrand in (7.1) has a pole of order k^2 at $s = 1$. The residue of this pole is equal to

$$P_{k^2-1} \left(\log \left(\frac{(q/\pi)^k}{X} \right), r \right) \mathcal{M}[|f_r|^2](1) = P_{k^2-1} \left(\log \left(\frac{(q/\pi)^k}{X} \right), r \right),$$

by (5.13), where

$$(7.2) \quad P_{k^2-1}(y, r) = \sum_{j=0}^{k^2-1} b_j(r) y^j$$

is an explicit polynomial in y of degree $k^2 - 1$. The leading coefficient $b_{k^2-1}(r)$ in (7.2) is equal to

$$\frac{a_k(r)}{(k^2-1)!} = \frac{1}{(k^2-1)!} \lim_{s \rightarrow 1^+} (s-1)^{k^2} \sum_{\substack{n=1 \\ (n,r)=1}}^{\infty} \frac{\tau_k(n)^2}{n^s}.$$

We thus move the line of integration of the integral in (7.1) to $\sigma = 1/2$, passing the pole at $s = 1$, pick up this residue and obtain

$$(7.3) \quad P_{k^2-1} \left(\log \left(\frac{(q/\pi)^k}{X} \right), r \right) + O \left(\left(\frac{(q/\pi)^k}{X} \right)^{\frac{1}{2}} \right)$$

for this integral. Thus, we have, from (7.1) and (7.3),

$$(7.4) \quad \sum_{\substack{1 \leq a \leq d \\ (a,d)=1}} |\Delta_w(\tau_k; X, d, a)|^2 \\ = X \frac{1}{\varphi(d)} \sum_{\substack{d=qr \\ q \geq Q}} \varphi^*(q) \left[P_{k^2-1} \left(\log \left(\frac{(q/\pi)^k}{X} \right), r \right) + O \left(\left(\frac{(q/\pi)^k}{X} \right)^{\frac{1}{2}} \right) \right] \\ + O_{\epsilon, \delta, k} \left(X^{1+\epsilon} d^{-\frac{\delta}{2k^2}} \right).$$

7.2. Step b: The error term and $\gamma_k(c)$. The first big-oh term in (7.4) contributes to the variance an amount

$$\ll X \sum_{\substack{d=qr \\ q \geq Q}} \frac{\varphi^*(q)}{\varphi(d)} \left(\frac{q^k}{X}\right)^{-\frac{1}{2}} \ll X^{\frac{3}{2}} \sum_{\substack{d=qr \\ q \geq Q}} d^{-\frac{k}{2}\left(\frac{c}{k} + \frac{\delta}{k^3}\right)} \ll X^{1+\epsilon} d^{-\frac{\delta}{2k^2}}.$$

This is equal to the second big-oh term in (7.4), and so the two error terms can be absorbed into one.

We now write

$$\frac{(q/\pi)^k}{X} = \frac{d^{k-c}}{(\pi r)^k}.$$

With this, the leading order term in (7.3) is equal to

$$\begin{aligned} \frac{a_k(r)}{(k^2-1)!} \log \left(\frac{(q/\pi)^k}{X} \right)^{k^2-1} &= \frac{a_k(r)}{(k^2-1)!} ((k-c) \log d - k \log \pi r)^{k^2-1} \\ &= \frac{a_k(r)}{(k^2-1)!} (k-c)^{k^2-1} (\log d)^{k^2-1} \\ &\quad - \frac{a_k(r)}{(k^2-1)!} \binom{k^2-1}{1} (k-c)^{k^2-2} (\log d)^{k^2-2} k \log \pi r \\ &\quad + \cdots + \frac{a_k(r)}{(k^2-1)!} (-k)^{k^2-1} (\log \pi r)^{k^2-1}. \end{aligned}$$

We first show

$$(7.5) \quad \sum_{\substack{d=qr \\ q \geq Q}} \frac{\varphi^*(q)}{\varphi(d)} a_k(r) \log r = o(\log d).$$

With this, together with induction, we will have, as $d \rightarrow \infty$,

$$(7.6) \quad \sum_{\substack{d=qr \\ q \geq Q}} \frac{\varphi^*(q)}{\varphi(d)} P_{k^2-1} \left(\log \left(\frac{(q/\pi)^k}{X} \right), r \right) \sim \frac{(k-c)^{k^2-1}}{(k^2-1)!} (\log d)^{k^2-1} \sum_{\substack{d=qr \\ q \geq Q}} \frac{\varphi^*(q)}{\varphi(d)} a_k(r) \log r.$$

We now prove (7.5).

7.3. Step c: Bounding lower order main terms. To help with the analysis in this step, we add back terms with $q < Q$. This added an amount

$$\sum_{\substack{d=qr \\ q < Q}} \frac{\varphi^*(q)}{\varphi(d)} a_k(r) \log r \ll \sum_{\substack{d=qr \\ q < Q}} \frac{d^{\frac{c}{k} + \frac{\delta}{k^3}} \log \log d}{d} \log r = o(1),$$

which is acceptable for the asymptotic. Thus, we may add in terms with $q < Q$ to the left side of (7.6). Writing

$$\varphi^* = \mu \star \varphi,$$

we have

$$(7.7) \quad \sum_{d=qr} \frac{\varphi^*(q)}{\varphi(d)} a_k(r) \log r = \sum_{d=qr} \frac{\varphi(q)}{\varphi(d)} a_k(r) \Lambda(r).$$

Write

$$(7.8) \quad d = p_1^{\alpha_1} \cdots p_v^{\alpha_v},$$

with p_j distinct. The r sum in (7.7) is restricted to prime powers of the form

$$r = p_j^{\beta_j}, \quad 1 \leq j \leq v, \quad 1 \leq \beta_j \leq \alpha_j.$$

Thus, (7.7) is equal to

$$\sum_{j=1}^v \sum_{\beta_j=1}^{\alpha_j} \frac{\varphi(p_j^{\alpha_j - \beta_j})}{\varphi(p_j^{\alpha_j})} a_k(p_j^{\beta_j}) \log p_j = \sum_{j=1}^v \frac{1}{p_j} \frac{1 - p_j^{-\alpha_j}}{1 - p_j^{-1}} a_k(p_j) \log p_j.$$

Introduce

$$(7.9) \quad P = (\log d)^2$$

to the j sum, the above is equal to

$$\begin{aligned} \left(\sum_{p_j \leq P} + \sum_{p_j > P} \right) \frac{1}{p_j} \frac{1 - p_j^{-\alpha_j}}{1 - p_j^{-1}} a_k(p_j) \log p_j &\ll \sum_{p_j \leq P} \frac{1}{p_j} \log \log d + \sum_{p_j > P} \frac{1}{(\log d)^2} \log p_j \\ &\ll \log \log d \log \log P = o(\log d). \end{aligned}$$

This leads to (7.5), and, thus, to (7.6). Hence, by (7.6), the left side of (2.2) is asymptotic to

$$X \sum_{\substack{d=qr \\ q \geq Q}} \frac{\varphi^*(q)}{\varphi(d)} P_{k^2-1} \left(\log \left(\frac{(q/\pi)^k}{X} \right), r \right) \sim \frac{(k-c)^{k^2-1}}{(k^2-1)!} X (\log d)^{k^2-1} \sum_{d=qr} \frac{\varphi^*(q)}{\varphi(d)} a_k(r).$$

To complete the proof of (2.2), it remains to show that

$$(7.10) \quad \frac{1}{\varphi(d)} \sum_{d=qr} \varphi^*(q) a_k(r) = a_k(d),$$

which we will do in the next, which is the last, step.

7.4. Step d: Evaluating the matching arithmetic constant $a_k(d)$. We rewrite the left side of (7.10) as

$$\frac{1}{\varphi(d)} (\mu \star \varphi \star a_k)(d).$$

We compute the Dirichlet series for $a_k(r)$. By Euler products and the definition of $a_k(r)$, we have

$$(7.11) \quad \begin{aligned} \sum_{r=1}^{\infty} \frac{a_k(r)}{r^s} &= \prod_p \sum_{j=0}^{\infty} \frac{a_k(p^j)}{p^{js}} = a_k \prod_p \frac{(1 - \frac{1}{p})^{2k-1}}{\sum_{i=0}^{k-1} \binom{k-1}{i}^2 p^{-i}} \sum_{j=0}^{\infty} p^{-js} \\ &= a_k \prod_p \frac{(1 - \frac{1}{p})^{2k-1}}{\sum_{i=0}^{k-1} \binom{k-1}{i}^2 p^{-i}} \zeta(s). \end{aligned}$$

Next, we have the well known series

$$(7.12) \quad \sum_{n=1}^{\infty} \frac{\mu(n)}{n^s} = \frac{1}{\zeta(s)}$$

and Euler product

$$(7.13) \quad \sum_{n=1}^{\infty} \frac{\varphi(n)}{n^s} = \prod_p \sum_{j=0}^{\infty} \frac{\varphi(p^j)}{p^{js}}.$$

Hence, by (7.11), (7.12), and (7.13),

$$\begin{aligned} \sum_{d=1}^{\infty} \frac{(\mu \star \varphi \star a_k)(d)}{d^s} &= \frac{1}{\zeta(s)} \left(\prod_p \sum_{j=0}^{\infty} \frac{\varphi(p^j)}{p^{js}} \right) \left(a_k \zeta(s) \prod_p \frac{(1 - \frac{1}{p})^{2k-1}}{\sum_{i=0}^{k-1} \binom{k-1}{i}^2 p^{-i}} \right) \\ &= a_k \prod_p \sum_{j=0}^{\infty} \frac{\varphi(p^j) (1 - \frac{1}{p})^{2k-1}}{\sum_{i=0}^{k-1} \binom{k-1}{i}^2 p^{-i}} \frac{1}{p^{js}}. \end{aligned}$$

Thus, at prime powers,

$$(7.14) \quad (\mu \star \varphi \star a_k)(p^j) = a_k \frac{\varphi(p^j) (1 - \frac{1}{p})^{2k-1}}{\sum_{i=0}^{k-1} \binom{k-1}{i}^2 p^{-i}}.$$

Hence, if d is factored as in (7.8), then (7.14) implies that

$$\begin{aligned} (\mu \star \varphi \star a_k)(d) &= a_k \cdot (\mu \star \varphi \star a_k)(p_1^{\alpha_1}) \cdots (\mu \star \varphi \star a_k)(p_v^{\alpha_v}) \\ &= a_k \varphi(d) \prod_{p|d} \frac{(1 - \frac{1}{p})^{2k-1}}{\sum_{i=0}^{k-1} \binom{k-1}{i}^2 p^{-i}} = \varphi(d) a_k(d). \end{aligned}$$

Thus, by the above, we have that

$$\frac{(\mu \star \varphi \star a_k)(d)}{\varphi(d)} = a_k(d).$$

This proves (7.10), and, therefore, the asymptotic (2.2), as d goes to infinity. \square

8. ACKNOWLEDGMENTS

I am indebted to Brad Rodgers for suggesting this problem and for extensive discussions around this topic. I also benefited from the many discussions with J. Brian Conrey and am grateful for all his support and generous accommodation. Special thanks to Zeev Rudnick and Stephen Lester for their useful comments. This research is supported in part by the National Science Foundation (NSF) grant Focus Research Group DMS-1854398 and the American Institute of Mathematics.

REFERENCES

- [1] M. B. Barban, Analogues of the divisor problem of Titchmarsh (in Russian), Vestnik Leningrad Univ. Ser. Ma. Meh. Astronom. **18** (4) (1963), pp. 5-13.
- [2] M. B. Barban, On the average error term in the generalized prime number theorem (in Russian), Dokl. Akad. Nauk USSR **5** (1964), pp. 5-7.
- [3] E. Bombieri, On the large sieve, Mathematika **12** (1965), pp. 201-225. <https://doi.org/10.1112/S0025579300005313>
- [4] V. Chandee, X. Li, The eighth moment of Dirichlet L -functions, Adv. Math., **259** (2014), pp. 339-375. <https://doi.org/10.1016/j.aim.2014.03.020>
- [5] J. B. Conrey, S. M. Gonek, High moments of the Riemann zeta-function, Duke Math. J. **107** (3) pp. 577-604. <https://doi.org/10.1215/S0012-7094-01-10737-0>

- [6] J. B. Conrey, H. Iwaniec, K. Soundararajan, The sixth power moment of Dirichlet L -functions, *Geom. Funct. Anal.*, **22** (5), pp. 1257-1288 (2012). <https://doi.org/10.1007/s00039-012-0191-6>
- [7] J. B. Conrey and J. P. Keating, Moments of zeta and correlations of divisor-sums: I, *Phil. Trans. R. Soc. A* **373**:20140313 (2015). <http://dx.doi.org/10.1098/rsta.2014.0313>
- [8] J. B. Conrey, J. P. Keating, Moments of zeta and correlations of divisor-sums: II, In *Advances in the Theory of Numbers Proceedings of the Thirteenth Conference of the Canadian Number Theory Association*, Fields Institute Communications, 75-85 (2015). https://doi.org/10.1007/978-1-4939-3201-6_3
- [9] J. B. Conrey, J. P. Keating, Moments of zeta and correlations of divisor-sums: III, *Indagationes Mathematicae*, **26**(5): 736-747 (2015). <https://doi.org/10.1016/j.indag.2015.04.005>
- [10] J. B. Conrey, J. P. Keating, Moments of zeta and correlations of divisor-sums: IV, *Research in Number Theory*, **2**(24): (2016). <https://doi.org/10.1007/s40993-016-0056-4>
- [11] J. B. Conrey, J. P. Keating, Moments of zeta and correlations of divisor-sums: V, *Proc. London Math. Soc.*, **118**: 729-752 (2018). <https://doi.org/10.1112/plms.12196>
- [12] H. Davenport and H. Halberstam, Primes in arithmetic progressions, *Michigan Math. J.* **13** (1966), pp. 485-489. <https://doi.org/10.1307/mmj/1028999608>
- [13] R. de la Bretèche, D. Fiorilli, Major arcs and moments of arithmetical sequences, *Amer. J. Math.* **142**, no. 1, Feb. 2020, 45-77. <https://doi.org/10.1353/ajm.2020.0001>
- [14] J. Friedlander, H. Iwaniec, Summation formulae for coefficients of L -functions, *Canad. J. Math.* Vol **57** (3), 2005, pp. 494-505. <https://doi.org/10.4153/CJM-2005-021-5>
- [15] P. X. Gallagher, The large sieve, *Mathematika* **14** (1967), pp. 14-20. <https://doi.org/10.1112/S0025579300007968>
- [16] A. Harper, K. Soundararajan, Lower bounds for the variance of sequences in arithmetic progressions: Primes and divisor functions, *The Quarterly Journal of Mathematics*, Volume **68**, Issue 1 (2017), pp. 97-123. <https://doi.org/10.1093/qmath/haw005>
- [17] C. Hooley, On the Barban-Davenport-Halberstam theorem. I, *J. Reine Angew. Math.*, **274/275**, pp. 206-223 (1975). <https://doi.org/10.1515/crll.1975.274-275.206>
- [18] J. Keating, B. Rodgers, E. Roditty-Gershon, Z. Rudnick, Sums of divisor functions in $F_q[t]$ and matrix integrals, *Math. Z.* **288**, 167-198 (2018). <https://doi.org/10.1007/s00209-017-1884-1>
- [19] E. Kowalski, G. Ricotta, Fourier coefficients of $GL(N)$ automorphic forms in arithmetic progressions, *Geom. Funct. Anal.* **24**, 1229-1297 (2014). <https://doi.org/10.1007/s00039-014-0296-1>
- [20] S. Lester, On the variance of sums of divisor functions in short intervals, *Proc. Amer. Math. Soc.* **144** (2016), 5015-5027. <https://doi.org/10.1090/proc/12914>
- [21] H. L. Montgomery, Primes in arithmetic progressions, *Michigan Math. J.* **17**, pp. 33-39 (1970). <https://doi.org/10.1307/mmj/1029000373>
- [22] Y. Motohashi, An asymptotic formula in the theory of numbers, *Acta Arith.* **17** (1970), pp. 255-264. <https://doi.org/10.1112/plms/s3-7.1.396>
- [23] Y. Motohashi, On the distribution of the divisor function in arithmetic progressions, *Acta Arithetica XXII* (1973), pp. 175-199. <https://doi.org/10.4064/aa-22-2-175-199>
- [24] D. T. Nguyen, Generalized divisor functions in arithmetic progressions: I, *J. Number Theory* **227** (2021), pp. 30-93. <https://doi.org/10.1016/j.jnt.2021.03.021>
- [25] B. Rodgers, K. Soundararajan, The variance of divisor sums in arithmetic progressions, *Forum Math.* **30** (2018), no. 2, pp. 269-293. <https://doi.org/10.1515/forum-2016-0227>
- [26] P. Shiu, A Brun-Titchmarsh theorem for multiplicative functions, *J. Reine Angew. Math.* **313** (1980), 161-170. <https://doi.org/10.1515/crll.1980.313.161>
- [27] A. I. Vinogradov, The density Hypothesis for Dirichlet L -series, *Izv. Akad. Nauk SSSR Ser. Mat.* **29** (1965), pp. 903-934. https://doi.org/10.1142/9789814542487_0017
- [28] S. B. Yakubovich, *Index transforms*, World Scientific Publishing Co. Inc., River Edge (1996). With a foreword by H. M. Srivastava.
- [29] Y.-T. Zhang, Bounded gaps between primes, *Ann. of Math.* **179** (2014), no. **3**, pp. 1121-1174. <http://doi.org/10.4007/annals.2014.179.3.7>

AMERICAN INSTITUTE OF MATHEMATICS, 600 E. BROKAW RD., SAN JOSE, CA 95112, USA.
Email address: dtn@aimath.org