
NP-HARDNESS OF COMPUTING UNIFORM NASH EQUILIBRIA ON PLANAR BIMATRIX GAMES

Takashi Ishizuka

Graduate School of Mathematics,
Kyushu University,
744 Motooka, Nishi-ku, Fukuoka, Japan
ishizuka.takashi.664@s.kyushu-u.ac.jp

Naoyuki Kamiyama

Institute of Mathematics for Industry,
Kyushu University,
744 Motooka, Nishi-ku, Fukuoka, Japan
kamiyama@imi.kyushu-u.ac.jp

August 23, 2022

ABSTRACT

We study the complexity of computing a uniform Nash equilibrium on a non-win-lose bimatrix game. It is known that such a problem is NP-complete even if a bimatrix game is win-lose (Bonifaci et al., 2008). Fortunately, if a win-lose bimatrix game is planar, then uniform Nash equilibria always exist. We have a polynomial-time algorithm for finding a uniform Nash equilibrium of a planar win-lose bimatrix game (Addario-Berry et al., 2007). The following question is left: How hard to compute a uniform Nash equilibrium on a planar non-win-lose bimatrix game? This paper resolves this issue. We prove that the problem of deciding whether a non-win-lose planar bimatrix game has uniform Nash equilibrium is also NP-complete.

Keywords Computational Complexity · NP-completeness · Bimatrix Game · Uniform Nash equilibrium

1 Introduction

The research interaction between Game Theory and Theoretical Computer Science has helped to study the computational issues underlying fundamental game-theoretical notions. A seminal topic of these research streams is to clarify the complexity of computing Nash equilibria in multi-player non-cooperative games.

The results of the last two decades unravel the complexity of computing Nash equilibria. It has been known that the complexity of the problem of finding one Nash equilibrium on a finite strategic-form game belongs between P and NP. In particular, Papadimitriou [Pap94] introduced the complexity class PPAD and proved that such a problem is PPAD-complete. Unfortunately, it is also PPAD-complete to compute a Nash equilibrium even if the number of players is constant. Daskalakis et al. [DGP09] have shown the PPAD-completeness of Nash equilibrium computation on a three-player game. Chen

and Deng [CD06] have proven that the problem of computing a Nash equilibrium for a two-player strategic-form game is PPAD-complete.

The most important result is the intractability for computing Nash equilibria for two-player strategic-form games such that both players gain only a unit reward at most [AKV05]. Although the Lemke-Howson algorithm [LH64] is known as the best algorithm for finding a Nash equilibrium, it has been known that there is an example that that algorithm has an exponential worst-case running time [SS04].

Motivated from the above negative facts, some studies have concentrated on the complexity of computing specific classes of equilibria, such as pure, mixed, or correlated equilibria [FPT04; Pap05]. This paper focuses on the uniform Nash equilibria, i.e., Nash equilibria consisting of strategies in which strategies are played according to a uniform distribution. A uniform Nash equilibrium is a kind of mixed equilibrium. However, unlike mixed equilibria, uniform Nash equilibria do not always exist. Bonifaci et al. [BIL08] have shown that it is NP-complete to decide whether a given two-player game has a uniform Nash equilibrium even if every element of payoff matrices is either 0 or 1. Note that we call such a game a win-lose bimatrix game.

These hardness results were based on a graph-theoretical approach. There is a one-to-one correspondence between a win-lose bimatrix game and a bipartite digraph. Addario-Berry et al. [AOV07] have shown that a win-lose bimatrix game corresponding to a planar bipartite digraph always has a uniform Nash equilibrium, and we can find it in polynomial time.

1.1 Our Results

This paper sharpens the tractability-intractability boundary for the problem of computing uniform Nash equilibria on bimatrix games (not necessarily win-lose). We focus on a bimatrix game that corresponds to an edge-weighted planar bipartite digraph; we call such a game a planar bimatrix game. Furthermore, we discuss the following two unsolved issues:

- (1) Does a planar bimatrix game always have a uniform Nash equilibrium?
- (2) Can we efficiently compute a uniform Nash equilibrium on a planar bimatrix game (find out if it exists)?

This paper presents the negative answers to the above issues. We prove that the problem of deciding whether a planar bimatrix game has uniform Nash equilibria is NP-complete — this result implies that uniform Nash equilibria do not always exist. More precisely, we state that the problem of computing a uniform Nash equilibrium on a non-win-lose planar bimatrix game is also NP-complete.

Technical Overview Our proof shown in Section 3 is inspired by the proof techniques used in [AKV05; BIL08; CS05]. We extend their graph-theoretical approach to one on edge-weighted digraphs; we present a one-to-one correspondence between bimatrix games and edge-weighted bipartite digraphs and characterize a uniform Nash equilibrium by a graph-theoretical notion. The key technique in proving the main result is the construction of the clause-variable gadget, shown in Figure 4.

2 Preliminaries

Firstly, we define basic notations. We denote by \mathbb{N} , \mathbb{Q} , $\mathbb{Z}_{\geq 0}$, $\mathbb{R}_{\geq 0}$, and $\mathbb{R}_{>0}$ the sets of positive integers, rational numbers, non-negative integers, non-negative real numbers, and positive real numbers, respec-

tively. We use $[n] := \{1, 2, \dots, n\}$ for $n \in \mathbb{N}$. For a given digraph G , we denote by $V(G)$ and $E(G)$ the sets of vertices and directed edges on G , respectively. For sets A and B , we denote by $A \sqcup B$ the disjoint union of A and B .

2.1 Bimatrix Games

A bimatrix game is a two-player strategic form game where both players have a finite set of strategies. We call the first (resp. second) player the *row* (resp. *column*) player. Here, we denote by R and C the sets of strategies for the row and column player, respectively. A bimatrix game is specified by non-negative real matrices $M_R, M_C \in \mathbb{R}_{\geq 0}^{R \times C}$; the rows and columns of both matrices are indexed by the pure strategies of the players.

Let ρ_R and ρ_C be positive integers. We call a bimatrix game (M_R, M_C) a $\langle \rho_R, \rho_C \rangle$ -bimatrix game if there are subsets $\mathcal{A} \subseteq \mathbb{Q} \setminus \{0\}$ with $|\mathcal{A}| = \rho_R$ and $\mathcal{B} \subseteq \mathbb{Q} \setminus \{0\}$ with $|\mathcal{B}| = \rho_C$ such that $M_R \in (\mathcal{A} \sqcup \{0\})^{R \times C}$, and $M_C \in (\mathcal{B} \sqcup \{0\})^{R \times C}$. Namely, a win-lose bimatrix game is a $\langle 1, 1 \rangle$ -bimatrix game.

A mixed strategy is a probability distribution over pure strategies, i.e., a vector $x_R \in \mathbb{R}_{\geq 0}^R$ such that $\sum_{r_i \in R} x_R(r_i) = 1$ is a mixed strategy of the row player. Similarly, a vector $x_C \in \mathbb{R}_{\geq 0}^C$ such that $\sum_{c_j \in C} x_C(c_j) = 1$ is a mixed strategy of the column player. For a mixed strategy x , the support $\text{supp}(x)$ of x is the set of pure strategies i such that $x(i) > 0$. A mixed strategy x is said to be *uniform* if for every $i \in \text{supp}(x)$, $x(i) = 1/|\text{supp}(x)|$.

When the row player plays a mixed strategy x_R and the column player plays a mixed strategy x_C , the expected payoffs for the row player and the column player is $x_R^\top M_R x_C$ and $x_R^\top M_C x_C$, respectively. A Nash equilibrium of the bimatrix game (M_R, M_C) is a pair of mixed strategies (x_R^*, x_C^*) satisfying for all mixed strategies $x_R \in \mathbb{R}_{\geq 0}^R$ of the row player, $(x_R^*)^\top M_R x_C^* \geq x_R^\top M_R x_C^*$, and for all mixed strategies $x_C \in \mathbb{R}_{\geq 0}^C$ of the column player, $(x_R^*)^\top M_C x_C^* \geq (x_R^*)^\top M_C x_C$. We call a Nash equilibrium (x_R^*, x_C^*) a uniform Nash equilibrium if both mixed strategies are uniform.

2.2 Edge-Weighted Digraphs Induced by Bimatrix Games

We now describe the one-to-one correspondence between bimatrix games and edge-weighted bipartite digraphs. Note that our notation and definition used in this paper are based on [AOV07; BIL08]; these papers only dealt with unweighted digraphs. We generalize their notion to edge-weighted digraphs.

One-to-One Correspondence Let (M_R, M_C) be a bimatrix game, where the sets of strategies for the row player and the column player are R and C , respectively. Now, we define the edge-weighted bipartite digraph $G = (V, E, w)$ induced by the bimatrix game (M_R, M_C) . Here, $w : E \rightarrow \mathbb{R}_{> 0}$ represents weights on edges. The set V of vertices is $R \sqcup C$. We have an arc (r_i, c_j) if $M_R(r_i, c_j) > 0$, and the weight on the arc (r_i, c_j) is equal to $M_R(r_i, c_j)$. Similarly, we have an arc (c_j, r_i) if $M_C(r_i, c_j) > 0$, and the weight on the arc (c_j, r_i) is equal to $M_C(r_i, c_j)$.

The bimatrix game (M_R, M_C) is a planar bimatrix game if the edge-weighted digraph G is induced by (M_R, M_C) is a planar graph. Furthermore, we call an edge-weighted bipartite digraph $G = (R \sqcup C, E, w)$ a $\langle \rho_r, \rho_c \rangle$ -edge-weighted bipartite digraph if there are two sets $\mathcal{A} \subseteq \mathbb{Q} \setminus \{0\}$ with $|\mathcal{A}| = \rho_r$ and $\mathcal{B} \subseteq \mathbb{Q} \setminus \{0\}$ with $|\mathcal{B}| = \rho_c$ such that each edge from R to C has the weight within \mathcal{A} and each edge from C to R has the weight within \mathcal{B} .

Remark that it is possible that the reverse transformation. Thus, we can construct a bimatrix game from an edge-weighted bipartite digraph.

Let $G = (V, E, w)$ be an edge-weighted bipartite digraph. Since G is bipartite, we can naturally separate the set of vertices V into two disjoint subsets R and C . We regard R and C as the sets of pure strategies for the row and column players, respectively. The payoff matrix of the row player M_R is defined as follows: For each $r_i \in R$, $M_R(r_i, c_j) = w(r_i, c_j)$ if $(r_i, c_j) \in E$; otherwise, $M_R(r_i, c_j) = 0$. The definition of the payoff matrix of the column player M_C is similar, i.e., for each $c_j \in C$, $M_C(r_i, c_j) = w(c_j, r_i)$ if $(c_j, r_i) \in E$; otherwise, $M_C(r_i, c_j) = 0$.

From the above observation, we obtain the following statement:

Lemma 2.1. *For every bimatrix game, there is a corresponding edge-weighted bipartite digraph; vice versa.*

Undominated Out-regular Subgraphs For a finite subset $S \subseteq V$, let $G[S]$ be the subgraph induced by S . We denote by $\Gamma^+(v)$ the set of out-neighbors of a vertex $v \in V$, and by $\Gamma_S^+(v)$ the set of out-neighbors of a vertex $v \in V$ that are in $S \subseteq V$.

Fix arbitrary non-empty subset of strategies $S = S_R \sqcup S_C \subseteq V$, where $\emptyset \neq S_R \subseteq R$ and $\emptyset \neq S_C \subseteq C$. The induced subgraph $G[S]$ is (α, β) out-regular if the following two conditions (i) for every $r_i \in S_R$, $\sum_{c_j \in \Gamma_S^+(r_i)} w(r_i, c_j) = \alpha$; (ii) for every $c_j \in S_C$, $\sum_{r_i \in \Gamma_S^+(c_j)} w(c_j, r_i) = \beta$. We say that G has an out-regular subgraph if there is a triple (α, β, S) such that $G[S]$ is (α, β) out-regular.

A vertex $v \in V \setminus S$ dominates an (α, β) out-regular subgraph $G[S]$ if it satisfies that either $\sum_{c_j \in \Gamma_S^+(v)} w(v, c_j) > \alpha$ if $v \in R$; or $\sum_{r_i \in \Gamma_S^+(v)} w(v, r_i) > \beta$ if $v \in C$. We say that an (α, β) out-regular subgraph $G[S]$ is undominated if there are no vertices that dominate $G[S]$. We say that G has an undominated out-regular subgraph if there is a triple (α, β, S) such that $G[S]$ is an undominated (α, β) out-regular subgraph.

Theorem 2.2. *A bimatrix game has a uniform Nash equilibrium if and only if the corresponding edge-weighted bipartite digraph has an undominated out-regular subgraph; vice versa.*

Proof. Let (M_R, M_C) be a bimatrix game, where the sets of strategies for the row player and the column player are R and C , respectively. We denote by G the edge-weighted bipartite digraph corresponding to (M_R, M_C) . Fix a subset of pure strategies $S = S_R \sqcup S_C$, where $\emptyset \neq S_R \subseteq R$ and $\emptyset \neq S_C \subseteq C$.

Now, we show that $G[S]$ is an undominated out-regular subgraph if and only if the pair of uniform strategies (x_R, x_C) such that $\text{supp}(x_R) = S_R$ and $\text{supp}(x_C) = S_C$ is a Nash equilibrium of (M_R, M_C) .

First, we assume that $G[S]$ is an undominated (α, β) out-regular subgraph form some positive reals α and β . Then, we show that the pair of uniform strategies (x_R, x_C) defined as $\text{supp}(x_R) = S_R$ and $\text{supp}(x_C) = S_C$ is a Nash equilibrium. When the column player plays the strategy x_C , the row player gains the expected payoff $\alpha/|S_C|$ by playing $r_i \in S_R$, but gains the expected payoff at most $\alpha/|S_C|$ by playing $r_i \in R \setminus S_R$. Therefore, the uniform strategy is a best response to x_C for the row player. Similarly, we can see that the uniform strategy x_C is a best response to x_R for the column player. Thus, (x_R, x_C) is a uniform Nash equilibrium.

Next, we assume that the pair of uniform strategies (x_R, x_C) such that $\text{supp}(x_R) = S_R$ and $\text{supp}(x_C) = S_C$ is a Nash equilibrium. For the sake of contradiction, we suppose that there are strategies $r_i, r_k \in S_R$ such that $\sum_{c_j \in \Gamma_S^+(r_i)} w(r_i, c_j) > \sum_{c_\ell \in \Gamma_S^+(r_k)} w(r_k, c_\ell)$ when the column player plays x_C . In this case, the row player gains the expected payoff $(1/|S_C|) \sum_{c_j \in \Gamma_S^+(r_i)} w(r_i, c_j)$ by playing the

strategy r_i . On the other hand, she gains the expected payoff $(1/|S_C|) \sum_{c_\ell \in \Gamma_S^+(r_k)} w(r_k, c_\ell)$ by playing r_k . From our assumption, the row player will get a larger expected payoff for playing r_i than for playing r_k , which contradicts that (x_R, x_C) is a Nash equilibrium. Therefore, $G[S]$ is an (α, β) out-regular subgraph for some positive reals α and β . Using a similar technique, it is easy to see that $G[S]$ is undominated. \square

Now, we show an important property of undominated out-regular subgraphs, which we use to prove the main result. If the edge-weighted bipartite digraph $G = (V, E, w)$ is strongly connected, and a subset S induces an undominated out-regular subgraph, then we have $\sum_{u \in \Gamma_S^+(v)} w(v, u) > 0$ for each $v \in S$.

Lemma 2.3. *Assume that the edge-weighted bipartite digraph $G = (V, E, w)$ is strongly connected, and a subset $S \subseteq V$ induces the undominated out-regular subgraph $G[S]$. Then, every vertex $v \in S$ satisfies that $\sum_{u \in \Gamma_S^+(v)} w(v, u) > 0$.*

Proof. For the sake of contradiction, we assume that there is a vertex $v \in S$ such that $\sum_{u \in \Gamma_S^+(v)} w(v, u) = 0$. We take a vertex $u \in S$ that belongs to the different part from v . Since the digraph G is strongly connected, there is an arc $(v', u) \in E$. If $v' \in S$, it contradicts that S induces an out-regular subgraph. On the other hand, if $v' \in V \setminus S$, it contradicts that $G[S]$ has no dominating vertices. \square

2.3 A Brief Introduction to Computational Complexity

We begin with a brief overview of various computational complexity-theoretical notions that will be needed for the remainder of this paper. Let $\Sigma = \{0, 1\}$ be a binary alphabet. We denote by Σ^* the set of finite strings. A decision problem $L \subseteq \Sigma^*$ is defined as follows: given a string x , decide whether x is in L . A complexity class \mathcal{C} is a set of decision problems. The complexity class NP is the set of all decision problems that are decidable by a non-deterministic Turing machine.

A reduction is a key concept of Computational Complexity Theory. A reduction from the problem A to the problem B is a function $f : \Sigma^* \rightarrow \Sigma^*$ such that a string $x \in A$ if and only if $f(x) \in B$. When such a function f is computable by a deterministic Turing machine, we say that such a reduction is a polynomial-time reduction. Furthermore, we say that two decision problems A and B are polynomially equivalent if there are polynomial-time reductions from A to B and from B to A . A decision problem L is NP-hard if all decision problems belonging to NP are polynomial-time reducible to L . A decision problem L is NP-complete if L belongs to NP and is NP-hard. Note that the problem 3-SAT (see Definition 2.4) is the well-known NP-complete problem [Coo71; Lev73].

The List of Decision Problems We conclude this brief introduction by resending a list of decision problems that will be used in this paper.

Definition 2.4. 3-SAT:

Input:

- a 3-CNF formula $\phi : \{0, 1\}^n \rightarrow \{0, 1\}$.

Task: Decide whether

- there exists an assignment $x \in \{0, 1\}^n$ such that $\phi(x) = 1$.

Definition 2.5. $\langle \rho_r, \rho_c \rangle$ -BIMATRIX GAME

Input:

- a $\langle \rho_r, \rho_c \rangle$ -bimatrix game (M_R, M_C)

Task: Decide whether

- there is a uniform Nash equilibrium (x_R, x_C) of (M_R, M_C) .

Definition 2.6. $\langle \rho_r, \rho_c \rangle$ -PLANAR BIMATRIX GAME

Input:

- a $\langle \rho_r, \rho_c \rangle$ -planar bimatrix game (M_R, M_C)

Task: Decide whether

- there is a uniform Nash equilibrium (x_R, x_C) of (M_R, M_C) .

Definition 2.7. $\langle \rho_r, \rho_c \rangle$ -UNDOMINATED OUTREGULAR SUBGRAPH

Input:

- a $\langle \rho_r, \rho_c \rangle$ edge-weighted bipartite digraph $G = (V, E, w)$

Task: Decide whether

- G has an undominated out-regular subgraph.

Definition 2.8. $\langle \rho_r, \rho_c \rangle$ -UNDOMINATED OUTREGULAR SUBGRAPH ON PLANAR GRAPH

Input:

- a $\langle \rho_r, \rho_c \rangle$ edge-weighted bipartite planar digraph $G = (V, E, w)$

Task: Decide whether

- G has an undominated out-regular subgraph.

3 On the Complexity of $\langle 1, 2 \rangle$ -Planar Bimatrix Games

The purpose of this section is to prove the NP-hardness of the problem of deciding whether a given $\langle 2, 2 \rangle$ -planar bimatrix game has a unique Nash equilibrium. So, we prove the next theorem.

Theorem 3.1. *The problem $\langle 1, 2 \rangle$ -PLANAR BIMATRIX GAME is NP-complete.*

It is trivial that the problem $\langle 1, 2 \rangle$ -PLANAR BIMATRIX GAME belongs to the class NP. Thus, it suffices to prove that this problem is NP-hard. For this, we present a polynomial-time reduction from 3-SAT to $\langle 1, 2 \rangle$ -PLANAR BIMATRIX GAME.

Recall that Theorem 2.2 states that the existence of uniform Nash equilibria for a bimatrix game is equal to the existence of undominated out-regular subgraphs of the corresponding edge-weighted bipartite digraph. Moreover, the reduction between a bimatrix game and an edge-weighted bipartite digraph

is polynomial-time computable and preserves planar. Therefore, these two problems $\langle 1, 2 \rangle$ -PLANAR BIMATRIX GAME and $\langle 1, 2 \rangle$ -UNDOMINATED OUTREGULAR SUBGRAPH ON PLANAR GRAPH are polynomially equivalent.

To complete the proof of Theorem 3.1, we present a polynomial-time reduction from 3-SAT to $\langle 1, 2 \rangle$ -UNDOMINATED OUTREGULAR SUBGRAPH, i.e., we show the following lemma:

Lemma 3.2. *3-SAT is polynomial-time reducible to $\langle 1, 2 \rangle$ -UNDOMINATED OUTREGULAR SUBGRAPH ON PLANAR GRAPH.*

Here, we sketch the proof of Lemma 3.2; the full proof of this lemma can be found in Section 4. In the first step, we reduce the problem 3-SAT to $\langle 1, 2 \rangle$ -UNDOMINATED OUTREGULAR SUBGRAPH in Section 4.1. For each 3-SAT instance $\phi : \{0, 1\}^n \rightarrow \{0, 1\}$, we construct the $\langle 1, 2 \rangle$ -edge-weighted bipartite digraph G_ϕ . Note that G_ϕ is not planar; it may have some crossing points. Hence, we eliminate all crossing points from the graph G_ϕ in the second step. The key technique to eliminate crossing points is the construction of the clause-variable gadget, shown in Figure 4. Denote by H_ϕ the planar digraph formed by transforming G_ϕ . To complete the proof of Lemma 3.2, we prove that G_ϕ has an undominated out-regular subgraph if and only if H_ϕ also has it.

4 Proof of Lemma 3.2

4.1 Reduction from 3-SAT to $\langle 1, 2 \rangle$ -UNDOMINATED OUTREGULAR SUBGRAPH

Let $\phi : \{0, 1\}^n \rightarrow \{0, 1\}$ be a 3-CNF formula. Without loss of generality, we assume that the 3-CNF formula ϕ holds the following three properties:

- There are m clauses.
- Each clause has exactly three literals.
- Each literal is contained in at least one clause.

This subsection constructs an edge-weighted bipartite digraph G_ϕ such that every undominated out-regular subgraph leads to a satisfying assignment of ϕ . We show how to embed the digraph G_ϕ into a plane in Section 4.2.

The set of vertices of G_ϕ consists of the followings:

- for each $i \in [n]$, variable vertices x_i, \bar{x}_i and variable coordinating vertices $y_i, \bar{y}_i, z_{i_1}, z_{i_2}$,
- for each $j \in [m]$, clause vertex C_j and clause coordinating vertices $v_{j_1}, v_{j_2}, v_{j_3}, u_{j_1}, u_{j_2}, u_{j_3}$, and
- an additional vertex a .

We separate these vertices into two parts:

- the first part $\mathcal{V}_R = \{y_i, \bar{y}_i, z_{i_1} ; i \in [n]\} \cup \{C_j, u_{j_1}, u_{j_2}, u_{j_3} ; j \in [m]\}$ and
- the second part $\mathcal{V}_C = \{x_i, \bar{x}_i, z_{i_2} ; i \in [n]\} \cup \{v_{j_1}, v_{j_2}, v_{j_3} ; j \in [m]\} \cup \{a\}$.




Figure 1: The Construction of arcs for every $j \in [m]$. Here, weights on **dashed** and **thick** edges are $m + n$ and 1 , respectively.

First, we describe the arcs going out from clause vertices and clause coordinating vertices. For a $j \in [m]$, we denote by $\ell_{j_1}, \ell_{j_2}, \ell_{j_3}$ the three literals contained in the clause C_j . For each $k \in \{1, 2, 3\}$, we create the following three arcs: (C_j, v_{j_k}) , (v_{j_k}, u_{j_k}) , and (u_{j_k}, ℓ_{j_k}) . The weights of these arcs are defined as $w(C_j, v_{j_k}) = 1$, $w(v_{j_k}, u_{j_k}) = m + n$, and $w(u_{j_k}, \ell_{j_k}) = 1$. We illustrate an example of our construction in Figure 1.

Second, we describe the arcs going out from the additional vertex a . For every $j \in [m]$, we create the arc (a, C_j) , and define its weight as $w(a, C_j) = 1$. Furthermore, for each $i \in [n]$, we create the arc (a, z_{i_1}) , and define its weight as $w(a, z_{i_1}) = 1$.

Third, we describe the arcs going out from the variable vertices and variable coordinating vertices. For each $i \in [n]$, we create the arcs (x_i, y_i) , (y_i, a) , (\bar{x}_i, \bar{y}_i) , (\bar{y}_i, a) , (a, z_{i_1}) , (z_{i_1}, z_{i_2}) , (z_{i_2}, y_i) , and (z_{i_2}, \bar{y}_i) . The weights of these arcs are defined as $w(x_i, y_i) = m + n$, $w(y_i, a) = 1$, $w(\bar{x}_i, \bar{y}_i) = m + n$, $w(\bar{y}_i, a) = 1$, $w(a, z_{i_1}) = 1$, $w(z_{i_1}, z_{i_2}) = 1$, $w(z_{i_2}, y_i) = m + n$, and $w(z_{i_2}, \bar{y}_i) = m + n$. We illustrate the construction of arcs for every $i \in [n]$ in Figure 3; we call this construction a variable gadget.

Finally, we embed the graph G_ϕ into a $3m \times 2mn$ grid graph. We arrange all clause coordinating vertices (the type of u_{j_k}) to the left and all variable vertices to the bottom. Each variable vertex has m points to connect clause coordinating vertices. At the variable vertices, we label each point to connect the adjacent edge with $1, 2, \dots, m$. Suppose that the arc u_{j_k} to x_i exists. This arc extends horizontally from u_{j_k} to just above the j -th connection point of x_i . The arc is then lowered vertically to the j -th point of x_i . We repeat this procedure to connect every directed edge between clause coordinating vertices and variable vertices. We lexicographically chose a connecting point of a clause vertex without loss of generality. We illustrate an example of our construction in Figure 2.

We complete the construction of G_ϕ . From the construction, we can straightforwardly see the following three facts:

- (i) Every arc is either going from \mathcal{V}_R to \mathcal{V}_C or from \mathcal{V}_C to \mathcal{V}_R .
- (ii) The weight on an arc outgoing from a vertex in \mathcal{V}_R is one.
- (iii) The weight on an arc outgoing from a vertex in \mathcal{V}_C is either one or $m + n$.

Thus, the digraph G_ϕ is a $\langle 1, 2 \rangle$ -edge-weighted bipartite digraph. Hence, the corresponding two-player game is a $\langle 1, 2 \rangle$ -bimatrix game. Now, we prove the following theorem.




Figure 2: The digraph constructed from a given 3-SAT instance $\phi(x) = (x_1 \vee \bar{x}_2 \vee x_3) \wedge (x_2 \vee \bar{x}_3 \vee x_4) \wedge (\bar{x}_1 \vee x_3 \vee \bar{x}_4)$, where $C_1 = x_1 \vee \bar{x}_2 \vee x_3$, $C_2 = x_2 \vee \bar{x}_3 \vee x_4$, and $C_3 = \bar{x}_1 \vee x_3 \vee \bar{x}_4$.




Figure 3: The construction of the variable gadget. Here, weights on **dashed** and **thick** edges are $m + n$ and 1 , respectively.

Theorem 4.1. *A Boolean formula ϕ has a satisfying assignment if and only if the graph G_ϕ has an undominated out-regular subgraph.*

Proof. First, we assume that ϕ is satisfiable, i.e., there exists an assignment $\xi \in \{0, 1\}^n$ such that $\phi(\xi) = 1$. Then, we construct the subset of vertices $S \subseteq V(G_\phi)$ such that

1. for each $i \in [n]$, $x_i, y_i \in S$ if $\xi_i = 1$; otherwise, $\bar{x}_i, \bar{y}_i \in S$
2. for each $i \in [n]$, $z_{i_1}, z_{i_2} \in S$
3. for each $j \in [m]$, $C_j \in S$
4. for each $j \in [m]$, we select exactly one literal $\ell_{j_k} \in S$ contained in C_j , and put clause coordinating vertices v_{j_k} and u_{j_k} into S .
5. the additional vertex a in S .

What remains is to prove that the subset S induces an undominated out-regular subgraph of G_ϕ .

First, we show that $G_\phi[S]$ is out-regular. Each vertex in $S \cap \mathcal{V}_R$ has exactly one out-neighbor, and the corresponding arc has the weight of one. Each vertex in $S \cap \mathcal{V}_C$ except the additional vertex a has exactly one out-neighbor, and the corresponding arc has the weight of $m + n$. The additional vertex a has $m + n$ out-neighbors, and each arc has the weight of one, which implies that the total weight is equal to $m + n$. Therefore, $G[S]$ is a $(1, m + n)$ out-regular induced subgraph.

Next, we prove that $G_\phi[S]$ is undominated. Note that the only vertex that may dominate $G_\phi[S]$ is z_{i_2} for any $i \in [n]$. For each $i \in [n]$, the variable coordinating vertex z_{i_2} does not dominate $G_\phi[S]$ since S contains at most one of x_i and \bar{x}_i that are out-neighbors of z_{i_2} . Hence, $G_\phi[S]$ is an undominated $(1, m + n)$ out-regular induced subgraph.

Therefore, the graph G_ϕ has an undominated out-regular induced subgraph if the 3-CNF formula ϕ has a satisfiable assignment.

Next, we assume that the subset S of the vertices on the digraph G_ϕ induces an undominated out-regular subgraph. Before we describe the construction of a satisfiable assignment of ϕ , we observe the properties of the subset S . More specifically, we prove the following properties:

- the additional vertex $a \in S$,
- for each $j \in [m]$, $C_j \in S$,
- for each $i \in [n]$, $z_{i_1}, z_{i_2} \in S$,
- for each $i \in [n]$, at most one of x_i and \bar{x}_i is contained in S .

Note that the digraph G_ϕ is strongly connected, and every directed edge has a weight of at least one. Therefore, we have $\sum_{u \in \Gamma_S^+(v)} w(v, u) > 0$ for every vertex $v \in S$ from Lemma 2.3. Furthermore, we can see that the additional vertex a is contained in S . The graph given by removing the additional vertex a from the digraph G_ϕ is an acyclic digraph. Note that any induced subgraph $H[S]$ on an acyclic digraph H has a vertex v such that $\Gamma_S^+(v) = \emptyset$, which implies that $\sum_{u \in \Gamma_S^+(v)} w(v, u) = 0$.

The subset S contains at least one out-neighbor of the additional vertex a . When a clause vertex C_j is contained in S , there exists exactly one $k \in \{1, 2, 3\}$ such that S contains the set of clause coordinating vertices $\{v_{j_k}, u_{j_k}\}$. We write x_i that is the out-neighbor of u_{j_k} , without loss of generality.

From Lemma 2.3, x_i and the variable coordinating vertex y_i are also contained in S . On the other hand, when a variable coordinating vertex z_{i_1} is contained in S , the variable coordinating vertex z_{i_1} and at least one of its out-neighbor are also contained in S from Lemma 2.3. These facts implies that all out-neighbors of a are contained in S since the number of a 's out-neighbors is $m + n$, which is equals to the weight on arcs (x_i, y_i) and (z_{i_2}, y_i) .

Note that for each $i \in [n]$, the subset S contains at most one of variable coordinating vertices y_i and \bar{y}_i . For the sake of contradiction, we assume that there is $i \in [n]$ such that both y_i and \bar{y}_i are contained in S . In this case, the variable coordinating vertex z_{i_2} dominates the subset S because the total weight of x_i is $m + n$ but the total weight of z_{i_2} is $2m + 2n$. Hence, for each $i \in [n]$, at most one of x_i and \bar{x}_i is contained in S . If the variable vertex x_i is in S , the variable coordinating vertex y_i is also in S from Lemma 2.3. Similarly, if \bar{x}_i is in S , \bar{y}_i is also in S .

From now on, we describe how to construct a satisfiable assignment of ϕ . We can easily find such an assignment $\xi \in \{0, 1\}^n$. For each $i \in [n]$, we define $\xi_i = 1$ if and only if $x_i \in S$.

What remains is to prove that the assignment ξ holds that $\phi(\xi) = 1$. It is easy to see that every clause vertex is contained in the subset S . For each clause vertex C_j , exactly one literal ℓ_{j_k} is in S . By the definition of ξ , if the literal ℓ_{j_k} corresponds to a variable vertex x_i , then $\xi_i = 1$; otherwise, $\xi_i = 0$. This implies that the assignment ξ is a satisfiable assignment of ϕ since every clause contains at least one TRUE assignment. \square

4.2 Embedding into plane

In this section, we embed the graph G_ϕ constructed in Section 4.1 into a plane. Thus, we eliminate crossing points. From simple observation, the only directed edges that may not be planar are those between the variable vertices and the clause coordinating vertices. To eliminate every crossing point, we construct a clause-variable gadget shown in Figure 4 and insert this gadget into each crossing point.

We now describe the insertion of the clause-variable gadget in more detail¹. Suppose that arcs from v_{j_k} to ℓ_{j_k} and from $v_{j'_k}$ to $\ell_{j'_k}$ intersect. Without loss of generality, we assume that the arc from v_{j_k} to ℓ_{j_k} crosses vertically. Then, in addition to the arcs on the clause-variable gadgets, we create the following four arcs (v_{j_k}, α_2) , $(v_{j'_k}, \beta_2)$, (γ_3, ℓ_{j_k}) , and $(\delta_3, \ell_{j'_k})$.

By the construction, it is obvious that the clause-variable gadget is planar. Hence, we can convert the digraph G_ϕ to the planar digraph by inserting the clause variable gadget into each crossing point. Naturally, our reconstruction can be done in polynomial time since the clause-variable gadget has at most $4m + 4n + 6$ vertices, and the number of crossing points is bounded by some polynomial.

We denote by H_ϕ the planar digraph constructed from G_ϕ by the above procedure. What remains is to prove that H_ϕ has an undominated out-regular subgraph if and only if the Boolean formula ϕ has a satisfying assignment.

Lemma 4.2. *The graph G_ϕ has an undominated out-regular subgraph if and only if the graph H_ϕ has an undominated out-regular subgraph.*

Proof. First, we assume that G_ϕ has an undominated out-regular subgraph $G_\phi[S]$, i.e., the subset of vertices $S \subseteq V(G_\phi)$ induces an undominated out-regular subgraph. Then, we show how to construct a subset T of vertices on H_ϕ that induces the undominated out-regular subgraph H_ϕ . We define the subset T by performing the procedure shown in Algorithm 1.

¹When an arc has more than one crossing point, we repeat the same operation, replacing v_{j_k} and $v_{j'_k}$ as γ_3 and δ_3 appropriately




Figure 4: The Construction of the Clause-Variable Gadget. Here, weights on **dashed** and **thick** edges are $m + n$ and 1 , respectively.

Algorithm 1 A procedure to construct the subset T

```

1: We initialize  $T$  with  $S$ .
2: for all an arc  $(v_{j_k}, \ell_i) \in E(G_\phi[S])$  that crosses other arcs on  $E(G_\phi[S])$  do
3:     we add some vertices as follows:
4:     if the path passes through the clause-variable gadget horizontally then
5:         we add the vertices  $\beta_2, \beta_1, \epsilon, \delta_1^{m+n}, \delta_2^{m+n}$ , and  $\delta_3$  to  $T$ 
6:     end if
7:     if the path passes through the clause-variable gadget vertically then
8:         we add the vertices  $\alpha_2, \alpha_1, \epsilon, \gamma_1^{m+n}, \gamma_2^{m+n}$ , and  $\gamma_3$  to  $T$ 
9:     end if
10: end for
11: for all  $\epsilon \in T$  do
12:     Set  $\kappa = 1$ 
13:     while  $\sum_{u \in \Gamma_T^+(\epsilon)} w(\epsilon, u) < m + n$  do
14:         if  $\beta_1 \in T$  then
15:             we add two vertices  $\delta_1^{m+n-\kappa}$  and  $\delta_2^{m+n-\kappa}$  to  $T$ 
16:         end if
17:         if  $\alpha_1 \in T$  and  $\sum_{u \in \Gamma_T^+(\epsilon)} w(\epsilon, u) < m + n$  then
18:             we add two vertices  $\gamma_1^{m+n-\kappa}$  and  $\gamma_2^{m+n-\kappa}$  to  $T$ 
19:         end if
20:          $\kappa \leftarrow \kappa + 1$ 
21:     end while
22: end for

```

What remains is to prove that the digraph $H_\phi[T]$ is an undominated out-regular subgraph. By the construction of H_ϕ , the vertices that can dominate $H_\phi[T]$ are the variable coordinating vertices z_{i_2} for $i \in [n]$, the vertices α_2 and β_2 that are on the clause-variable gadget. As mentioned in the proof of Theorem 4.1, any variable coordinating vertex z_{i_2} does not dominate the digraph $H_\phi[T]$ since at most one of variable vertices x_i and \bar{x}_i belongs to S . The vertex α_2 does not dominate $H_\phi[T]$ since at most one of the vertices α_1 and δ_1^1 belongs to T . Similarly, we see that the vertex β_2 is not a dominating vertex. Hence, $H_\phi[T]$ is an undominated subgraph. It is straightforward to see that the total weight on out-neighbors is equal to 1 or $m + n$ for every vertex in T . Therefore, $H_\phi[T]$ is out-regular. From the above argument, the digraph H_ϕ has an undominated out-regular subgraph.

Next, we assume that the graph H_ϕ has an undominated out-regular subgraph $H_\phi[T]$, i.e., the subset T induces an undominated out-regular subgraph. Then, we show how to construct the subset S of vertices on G_ϕ that induces an undominated out-regular subgraph. We construct the subgraph S by performing the procedure shown in Algorithm 2.

Algorithm 2 A procedure to construct the subset S

```

1: We initialize  $S$  with  $\{a\} \cup \{C_j ; j \in [m]\}$ .
2: for all  $i \in [n]$  do
3:   if  $x_i \in T$  then
4:     we add the vertices  $x_i$  and  $y_i$  to  $S$ 
5:   end if
6:   if  $\bar{x}_i \in T$  then
7:     we add the vertices  $\bar{x}_i$  and  $\bar{y}_i$  to  $S$ 
8:   end if
9: end for
10: for all  $j \in [m]$  do
11:   if  $u_{j_k}, v_{j_k} \in T$  then
12:     we add the vertices  $u_{j_k}$  and  $v_{j_k}$  to  $S$ 
13:   end if
14: end for
    
```

What remains is to prove that the graph $G_\phi[S]$ is an undominated out-regular subgraph. First, we observe the next fact:

- If a clause coordinating vertex v_{j_k} belongs to T , there is a variable vertex $x_i \in T$ such that the literal ℓ_{j_k} that is contained in the clause C_j corresponds to x_i .

Since the subset T induces the undominated out-regular subgraph $H_\phi[T]$, the out-neighbor of v_{j_k} is contained in T whenever v_{j_k} belongs to T . Note that a clause coordinating vertex v_{j_k} has only one out-neighbor. If the arc outgoing from v_{j_k} does not cross any other arcs, the above fact immediately follows. Suppose that the arc outgoing from v_{j_k} crosses other arcs. Thus, the out-neighbor of v_{j_k} is either α_2 or β_2 on the clause-variable gadget. Without loss of generality, we assume that the out-neighbor of v_{j_k} is α_2 (the other case follows from the same argument). Since again the subset T induces the undominated out-regular subgraph $H_\phi[T]$, the vertices α_1 and ϵ are contained in T . Then, we can see that at least one of vertices $\gamma_1^2, \gamma_1^3, \dots, \gamma_1^{m+n}$ belongs to T because the vertex δ_1^1 is not in the subset T . If δ_1^1 is contained in T , then the total weight on out-neighbors of α_2 is not equal to $m + n$, which contradicts

the out-regularity of $H_\phi[T]$. The vertex γ_3 is also contained in T since the subset T contains the vertices γ_1^κ and γ_2^κ for some κ . By repeating this argument, we see that the above fact follows.

Next, we show that the digraph $G_\phi[S]$ is out-regular. For each vertex in $S \cap \mathcal{V}_R$, the total weight on its out-neighbors is equal to 1. Furthermore, for each vertex in $S \cap \mathcal{V}_C$, the total weight on its out-neighbors is equal to $m + n$. Hence, the induced subgraph $G_\phi[S]$ is out-regular.

Finally, we show that the digraph $G_\phi[S]$ is undominated. For each $i \in [n]$, at most one of variable vertices x_i and \bar{x}_i is contained in T . If not, then the variable coordinating vertex z_{i_2} dominates the digraph $G_\phi[S]$. This fact implies that the digraph $G_\phi[S]$ is an undominated subgraph. From the above argument, the subset S induces an undominated out-regular subgraph. \square

Acknowledgments

This work was supported by JSPS KAKENHI Grant Numbers JP21J10845 and JP20H05795.

References

- [AKV05] Timothy G. Abbott, Daniel M. Kane, and Paul Valiant. “On the Complexity of Two-Player Win-Lose Games.” In: *Proceedings of the 46th Annual IEEE Symposium on Foundations of Computer Science*. IEEE Computer Society, 2005, pp. 113–122. DOI: 10.1109/SFCS.2005.59.
- [AOV07] Louigi Addario-Berry, Neil Olver, and Adrian Vetta. “A Polynomial Time Algorithm for Finding Nash Equilibria in Planar Win-Lose Games.” In: *Journal of Graph Algorithms and Applications* 11.1 (2007), pp. 309–319. DOI: 10.7155/jgaa.00147.
- [BIL08] Vincenzo Bonifaci, Ugo Di Iorio, and Luigi Laura. “The complexity of uniform Nash equilibria and related regular subgraph problems.” In: *Theoretical Computer Science* 401.1-3 (2008), pp. 144–152. DOI: 10.1016/j.tcs.2008.03.036.
- [CD06] Xi Chen and Xiaotie Deng. “Settling the Complexity of Two-Player Nash Equilibrium.” In: *Proceedings of the 47th Annual IEEE Symposium on Foundations of Computer Science*. IEEE Computer Society, 2006, pp. 261–272. DOI: 10.1109/F0CS.2006.69.
- [Coo71] Stephen A. Cook. “The Complexity of Theorem-Proving Procedures.” In: *Proceedings of the 3rd Annual ACM Symposium on Theory of Computing*. ACM, 1971, pp. 151–158. DOI: 10.1145/800157.805047.
- [CS05] Bruno Codenotti and Daniel Stefankovic. “On the computational complexity of Nash equilibria for $(0, 1)$ bimatrix games.” In: *Information Processing Letters* 94.3 (2005), pp. 145–150. DOI: 10.1016/j.ip1.2005.01.010.
- [DGP09] Constantinos Daskalakis, Paul W. Goldberg, and Christos H. Papadimitriou. “The complexity of computing a Nash equilibrium.” In: *Communications of the ACM* 52.2 (2009), pp. 89–97. DOI: 10.1145/1461928.1461951.
- [FPT04] Alex Fabrikant, Christos H. Papadimitriou, and Kunal Talwar. “The complexity of pure Nash equilibria.” In: *Proceedings of the 36th Annual ACM Symposium on Theory of Computing*. ACM, 2004, pp. 604–612. DOI: 10.1145/1007352.1007445.

- [Lev73] Leonid A. Levin. “Universal sequential search problems.” In: *Problems of Information Transmission* 9.3 (1973), pp. 265–266.
- [LH64] C. E. Lemke and J. T. Howson Jr. “Equilibrium Points of Bimatrix Games”. In: *Journal of the Society for Industrial and Applied Mathematics* 12.2 (1964), pp. 413–423. DOI: 10.1137/0112033.
- [Pap05] Christos H. Papadimitriou. “Computing correlated equilibria in multi-player games.” In: *Proceedings of the 37th Annual ACM Symposium on Theory of Computing*. ACM, 2005, pp. 49–56. DOI: 10.1145/1060590.1060598.
- [Pap94] Christos H. Papadimitriou. “On the Complexity of the Parity Argument and Other Inefficient Proofs of Existence.” In: *Journal of Computer and System Sciences* 48.3 (1994), pp. 498–532. DOI: 10.1016/S0022-0000(05)80063-7.
- [SS04] Rahul Savani and Bernhard von Stengel. “Exponentially Many Steps for Finding a Nash Equilibrium in a Bimatrix Game”. In: *Proceedings of the 45th Symposium on Foundations of Computer Science*. IEEE Computer Society, 2004, pp. 258–267. DOI: 10.1109/FOCS.2004.28.