

An Accelerated Failure Time Regression Model for Illness-Death Data: A Frailty Approach

Lea Kats, leatal@mail.tau.ac.il

and

Malka Gorfine, gorfinem@tauex.tau.ac.il

Department of Statistics and Operations Research

Tel Aviv University, Israel

Abstract

This work presents a new model and estimation procedure for the illness-death survival data where the hazard functions follow accelerated failure time (AFT) models. A shared frailty variate induces positive dependence among failure times of a subject for handling the unobserved dependency between the non-terminal and the terminal failure times given the observed covariates. Semi-parametric maximum likelihood estimation procedure is developed via a kernel smoothed-aided EM algorithm, and variances are estimated by weighted bootstrap. The model is presented in the context of existing frailty-based illness-death models, emphasizing the contribution of the current work. The breast cancer data of the Rotterdam tumor bank are analyzed using the proposed and existing illness-death models. The results are contrasted and evaluated based on a new graphical goodness-of-fit procedure. Simulation results and data analysis nicely demonstrate the practical utility of the shared frailty variate with the AFT regression model under the illness-death framework.

Keywords: goodness of fit; illness-death model; kernel method; semicompeting risks; shared frailty;

1 Introduction

The accelerated failure time (AFT) model (Kalbfleisch and Prentice, 2011) is a well-known alternative to the popular Cox proportional hazard (PH) model (Cox, 1972). The major advantage of AFT over Cox PH is the former being more intuitively interpretable (Wei, 1972; Cox, 1997 among others). This work focuses on the illness-death model which is specified by AFT models for the involved transitions. In illness-death model subjects start at State 0 (e.g. healthy) and then move to State 2 (e.g. death) directly or transit first to State 1 (e.g. the age at diagnosis of the disease under study) and then to State 2. See Fig. 1.

Xu et al. (2010) suggested an illness-death model with three Cox-based hazard functions. One of their major contributions was the inclusion of a shared gamma-frailty variate, that acts multiplicatively on each of the hazard function, for incorporating unobserved dependence between the time to disease diagnosis and time to death. Lee et al. (2015) adopted the model of Xu et al. (2010) but replaced their semi-parametric maximum likelihood estimation procedure with a semi-parametric Bayesian estimation approach. Jiang and Haneuse (2017) developed a class of transformation models for illness-death models that permits a non-parametric specification of the frailty distribution, but to insure identifiability, their model is restricted to parametric transformation and error distribution. Recently, for a simpler interpretation, Gorfine et al. (2020) proposed a frailty-based illness-death model with Cox-type marginalized hazards that also accommodates delayed entry.

Various estimation procedures have been developed for shared-frailty AFT models for clustered data (without competing or semi-competing risks). Pan (2001) considered clustered survival data with an AFT model and a gamma

frailty to characterize the unobserved dependence among cluster members. It is assumed that the shared frailty acts multiplicatively on the hazard function of the error term. [Zhang and Peng \(2007\)](#), [Xu and Zhang \(2010\)](#), [Johnson and Strawderman \(2012\)](#) and [Liu et al. \(2013\)](#) adopted Pan's model and provided various estimation methods. All these AFT estimation procedure for clustered data are not directly applicable to our illness-death setting, due to the differences in the likelihood functions, as will be elaborated in Section 2.3.

Unlike Cox-type models with illness-death framework, AFT models are not well developed. The only work which provides an AFT frailty-based model with illness-death setting is of [Lee et al. \(2017\)](#). [Lee et al. \(2017\)](#), in contrast to [Pan \(2001\)](#), used an additive frailty variate in the log-transformed failure time model, and their parametric and semi-parametric estimation methods are based on a Bayesian approach. Table 1 summarizes the available frailty-based Cox or AFT models for the illness-death setting.

The current work fills the gap and provides a gamma-frailty illness-death AFT model where the frailty acts multiplicatively on the hazards of the error terms, in the spirit of [Pan \(2001\)](#). We extended the estimation approaches of [Zeng and Lin \(2007\)](#) and [Liu et al. \(2013\)](#), and developed a semi-parametric maximum likelihood estimators (MLEs) based on a kernel-smoothing technique combined with an EM algorithm. Conceptual differences between our model and that of [Lee et al. \(2017\)](#) will be demonstrated in Section 2.2.

The proposed model and estimation method along with existing methods were applied to the Rotterdam tumor bank of 1,546 breast cancer patients, whom had node-positive disease and underwent a tumor removal surgery between the years 1978–1993. In this example, date at tumor removal surgery is the entrance time to State 0; date at relapse and date at death are the respective entry times to states 1 and 2. Prognostic variables are age at primary surgery, menopausal status, tumor size, tumor grade, number of positive lymph nodes, estrogen and progesterone receptors in the initial biopsy, hormonal therapy and chemotherapy. For a comparison of the various models, we extended the goodness of fit procedure of [Li et al. \(2021\)](#) to any illness-death model. The results of our proposed goodness of fit visualizing procedure nicely demonstrate the utility of the proposed model and estimation procedure.

The remainder of this article is organized as follows. Section 2 describes the proposed gamma frailty-based AFT illness-death regression model, and the estimation method for the regression coefficients, the hazard functions and the parameter of the gamma-frailty distribution. The illness-death goodness of fit procedure is provided in Section 3. In Section 4 we report the results of extensive simulation study. Section 5 summarises the analyses of the breast cancer data from the Rotterdam tumor bank, while comparing the proposed AFT approach and various existing AFT and Cox models, with and without frailty. A concluding discussion is provided in Section 6. R code for performing the simulations and data analysis in this paper is posted on the following webpage: <https://github.com/LeaKats/sempicompAFT>.

2 The Model and Methods

2.1 The Proposed Multiplicative Frailty-Based Model

Assume a sample of n independent observations. Let T_{1i} and T_{2i} be the respective times to the non-terminal and the terminal events of subject i , $i = 1, \dots, n$. Let X_i be a time-independent vector of covariates. The illness-death model (Fig. 1) is defined by

$$\log(T_{1i}) = \beta_{01}^T X_i + \epsilon_{01i}, T_{1i} > 0 \quad (1)$$

$$\log(T_{2i}) = \beta_{02}^T X_i + \epsilon_{02i}, T_{2i} > 0, \text{ given subject } i \text{ is free of the disease} \quad (2)$$

$$\log(T_{2i}) = \beta_{12}^T X_i + \epsilon_{12i}, T_{2i} > t_1 > 0, \text{ given subject } i \text{ was diagnosed at age } T_{1i} = t_1 \quad (3)$$

where β_{jk} , $jk \in \{01, 02, 12\}$, is a vector of regression coefficients of transition jk , and ϵ_{jki} are random errors with an unspecified distributions. Given that subject i was diagnosed at age $T_{1i} = t_1$, the support of T_{2i} is restricted by t_1 , so the conditional distribution of T_{2i} is truncated by t_1 . Model (3) above is not including age at diagnosis, t_1 , as an additional covariate, but instead, the dependence between T_{1i} and T_{2i} is incorporated via a shared-frailty model. Given the frailty variate of subject i , denoted by γ_i , it is assumed that the respective conditional baseline hazard

functions of $\exp(\epsilon_{jki})$, $jk \in \{01, 02, 12\}$, are given by

$$\lambda_{01}^o(t|\gamma_i) = \gamma_i h_{01}^o(t), t > 0 \quad (4)$$

$$\lambda_{02}^o(t|\gamma_i) = \gamma_i h_{02}^o(t), t > 0, \text{ given subject } i \text{ is free of the disease} \quad (5)$$

$$\lambda_{12}^o(t|t_1, \gamma_i) = \gamma_i h_{12}^o(t), t > t_1 > 0, \text{ given subject } i \text{ was diagnosed at age } T_{1i} = t_1 \quad (6)$$

where each $h_{jk}^o(\cdot)$ is an unspecified baseline hazard function of $\exp(\epsilon_{jk})$ and γ_i is an unobservable non-negative random effect, taken to be independent of X_i . It is assumed that γ_i are gamma distribution with mean 1, unknown variance $\sigma > 0$, and thus with density $f(\gamma; \sigma) = \sigma^{-1/\sigma} \gamma^{1/\sigma-1} e^{-\gamma/\sigma} / \Gamma(\sigma^{-1})$. We also assume that $\epsilon_{01i}, \epsilon_{02i}$, and ϵ_{12i} are independent given (X_i, γ_i) .

Based on Eq.'s (1)–(6), the conditional hazard functions of the three transitions, given (X_i, γ_i) , can be written as

$$\begin{aligned} \lambda_{0k}(t|X_i, \gamma_i) &= \lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(t \leq T_{ki} < t + \Delta | T_{1i} \geq t, T_{2i} \geq t, X_i, \gamma_i) \\ &= \gamma_i h_{0k}^0 \left(t e^{-\beta_{0k}^T X_i} \right) e^{-\beta_{0k}^T X_i}, \quad t > 0, \quad k = 1, 2, \end{aligned} \quad (7)$$

$$\begin{aligned} \lambda_{12}(t|t_1, X_i, \gamma_i) &= \lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(t \leq T_{2i} < t + \Delta | T_{1i} = t_1, T_{2i} \geq t, X_i, \gamma_i) \\ &= \gamma_i h_{12}^0 \left(t e^{-\beta_{12}^T X_i} \right) e^{-\beta_{12}^T X_i}, \quad t > t_1 > 0. \end{aligned} \quad (8)$$

For details, see Section S1 of the Web Supplementary Material (WSM).

2.2 Comparison with the Additive Frailty-Based Model

[Lee et al. \(2017\)](#) proposed to directly model the times of the events via the following AFT model specification:

$$\begin{aligned} \log(T_{1i}) &= \tilde{\beta}_{01}^T X_i + \tilde{\gamma}_i + \tilde{\epsilon}_{01i}, \quad T_{1i} > 0 \\ \log(T_{2i}) &= \tilde{\beta}_{02}^T X_i + \tilde{\gamma}_i + \tilde{\epsilon}_{02i}, \quad T_{2i} > 0, \text{ given subject } i \text{ is free of the disease} \\ \log(T_{2i}) &= \tilde{\beta}_{12}^T X_i + \tilde{\gamma}_i + \tilde{\epsilon}_{12i}, \quad T_{2i} > t_1 > 0, \text{ given subject } i \text{ was diagnosed at age } T_{1i} = t_1 \end{aligned}$$

where $\tilde{\beta}_{jk}^T$, $jk \in \{01, 02, 12\}$, are vectors of transition-specific regression coefficients. The errors $\tilde{\epsilon}_{jki}$ are transition-specific random variable with unspecified distributions in the semi-parametric setting, or with a normal distribution in the parametric setting. Also, $\tilde{\gamma}_i$, $i = 1, \dots, n$, are the unobserved normally distributed frailty variates with mean zero and variance θ , and are assumed to be independent of X_i .

The above additive frailty approach includes the observed covariates X_i and the unobservable $\tilde{\gamma}_i$ in a similar fashion in the models. In contrast, the popular multiplicative frailty approach of [Pan \(2001\)](#) in the context of clustered data and in Section 2.1 above, separates the observed covariates X_i and the unobservable component γ_i . The observed covariates directly affect time to event, and the unobserved affects the hazard functions of the error terms.

Assume, for example, that $\exp(\epsilon_{01i})$ and $\tilde{\epsilon}_{01i}$ are normally distributed with mean μ_{01} and variance ω_{01}^2 . Then, the respective conditional hazard functions of the multiplicative and additive models are

$$\lambda_{01}(t|X_i, \gamma_i) = \frac{\gamma_i e^{-\beta_{01}^T X_i}}{\omega_{01}} \phi \left(\frac{te^{-\beta_{01}^T X_i} - \mu_{01}}{\omega_{01}} \right) \left\{ 1 - \Phi \left(\frac{te^{-\beta_{01}^T X_i} - \mu_{01}}{\omega_{01}} \right) \right\}^{-1}$$

and

$$\tilde{\lambda}_{01}(t|X_i, \tilde{\gamma}_i) = \frac{1}{\omega_{01} t} \phi \left(\frac{\log t - \mu_{01} - \tilde{\beta}_1^T X_i - \tilde{\gamma}_i}{\omega_{01}} \right) \left\{ 1 - \Phi \left(\frac{\log t - \mu_{01} - \tilde{\beta}_1^T X_i - \tilde{\gamma}_i}{\omega_{01}} \right) \right\}^{-1}$$

where $\phi(\cdot)$ and $\Phi(\cdot)$ are the density and CDF of the standard normal distribution, respectively. Evidently, λ_{01} is of a simpler interpretation than that of $\tilde{\lambda}_{01}$ in terms of the unobserved frailty effect. Fig. 2 demonstrates $\tilde{\lambda}_{01}(t|X_i, \tilde{\gamma}_i)$ as

a function of t for $X_i = 0$ and various combinations of $(\tilde{\gamma}_i, \mu_{01}, \omega_{01})$. The top-left (bottom) plot of Fig. 2 shows that $\tilde{\lambda}_{01}$ decreases (increases) as a function of γ_i for any given value of t . The top-right plot indicates that $\tilde{\lambda}_{01}$ could be a non-monotone function of γ_i for some values of t . In contrast, λ_{01} of the multiplicative frailty model, is a monotonic increasing function of the frailty variate γ_i for any error distribution (see Eq. 7), and thus the multiplicative-frailty model could be viewed as a model with a simpler interpretation for the unobserved frailty effect.

2.3 The Proposed Estimation Method

Our goal is to estimate the unknown set of parameters of the illness death model $\Omega = \{\beta_{01}^T, \beta_{02}^T, \beta_{12}^T, h_{01}^o, h_{02}^o, h_{12}^o, \sigma\}$. Let C_i denote the right censoring time of subject i , $i = 1, \dots, n$. Then, the observed data consists of n independent observations $\mathcal{O}_i = \{V_i, W_i, \delta_{1i}, \delta_{2i}, \delta_{3i}, X_i\}$, where $V_i = \min(T_{1i}, T_{2i}, C_i)$, $\delta_{1i} = I(T_{1i} \leq \min(T_{2i}, C_i))$, $\delta_{2i} = I(T_{2i} \leq \min(T_{1i}, C_i))$, $W_i = \delta_{1i} \min(T_{2i}, C_i)$, and $\delta_{3i} = \delta_{1i} I(T_{2i} \leq C_i)$. V_i refers to the first observed time, δ_{1i} and δ_{2i} indicate whether the first observed time was age at disease diagnosis (T_{1i}), age at death (T_{2i}) or age at censoring; W_i is age at death or age at censoring after diagnosis, and δ_{3i} indicates whether death was observed after diagnosis. It is assumed that the censoring and the failure times are conditionally independent and non-informative, given (X_i, γ_i) , and observations are identically distributed.

Then, the likelihood function for Ω is proportional to $L(\Omega) = \prod_{i=1}^n \int L_i(\Omega) d\gamma$ where

$$\begin{aligned} L_i(\Omega) &= \left\{ \gamma h_{01}^o(V_i e^{-\beta_{01}^T X_i}) e^{-\beta_{01}^T X_i} \right\}^{\delta_{1i}} \left\{ \gamma h_{02}^o(W_i e^{-\beta_{02}^T X_i}) e^{-\beta_{02}^T X_i} \right\}^{\delta_{2i}} \exp \left\{ -\gamma H_{01}^o(V_i e^{-\beta_{01}^T X_i}) \right\} \\ &\quad \exp \left\{ -\gamma H_{02}^o(W_i e^{-\beta_{02}^T X_i}) \right\} \left\{ \gamma h_{12}^o(W_i e^{-\beta_{12}^T X_i}) e^{-\beta_{12}^T X_i} \right\}^{\delta_{3i}} \\ &\quad \exp \left\{ -\delta_{1i}\gamma \left(H_{12}^o(W_i e^{-\beta_{12}^T X_i}) - H_{12}^o(V_i e^{-\beta_{12}^T X_i}) \right) \right\} f(\gamma; \sigma). \end{aligned}$$

where $H_{jk}^o(t) = \int_0^t h_{jk}^o(u) du$, $jk \in \{01, 02, 12\}$. For details see Section S2 of WSM. Treating the unobservable frailties as a missing-data problem, calls for the Expectation-Maximization (EM) algorithm for obtaining semi-parametric maximum likelihood estimators (Dempster et al., 1977). It can be verified (see S3 of WSM) that the conditional expectation of the complete log-likelihood given the observed data $\mathcal{O} = \{\mathcal{O}_i, i = 1, \dots, n\}$ and the parameters' values at the m th step, $\widehat{\Omega}^{(m)}$, equals

$$E(l(\sigma)|\mathcal{O}, \widehat{\Omega}^{(m)}) + E(l(\beta_{01}, h_{01}^o)|\mathcal{O}, \widehat{\Omega}^{(m)}) + E(l(\beta_{02}, h_{02}^o)|\mathcal{O}, \widehat{\Omega}^{(m)}) + E(l(\beta_{12}, h_{12}^o)|\mathcal{O}, \widehat{\Omega}^{(m)}) \quad (9)$$

where

$$E(l(\sigma)|\mathcal{O}, \widehat{\Omega}^{(m)}) = \frac{1}{n} \sum_{i=1}^n \left(D_i + \frac{1}{\sigma} \right) \mathcal{E}_{2i}^{(m)} - \frac{1}{n\sigma} \sum_{i=1}^n \mathcal{E}_{1i}^{(m)} - \frac{1}{\sigma} \log \sigma - \log \Gamma \left(\frac{1}{\sigma} \right), \quad (10)$$

$$E(l(\beta_{0k}, h_{0k}^o)|\mathcal{O}, \widehat{\Omega}^{(m)}) = \frac{1}{n} \sum_{i=1}^n \left[\delta_{ki} \left\{ \log h_{0k}^o(V_i e^{-\beta_{0k}^T X_i}) - \beta_{0k}^T X_i \right\} - \mathcal{E}_{1i}^{(m)} H_{0k}^o(V_i e^{-\beta_{0k}^T X_i}) \right] k = 1, 2,$$

$$\begin{aligned} E(l(\beta_{12}, h_{12}^o)|\mathcal{O}, \widehat{\Omega}^{(m)}) &= \frac{1}{n} \sum_{i=1}^n \left[\delta_{3i} \left\{ \log h_{12}^o(W_i e^{-\beta_{12}^T X_i}) - \beta_{12}^T X_i \right\} \right. \\ &\quad \left. - \delta_{1i} \mathcal{E}_{1i}^{(m)} \left\{ H_{12}^o(W_i e^{-\beta_{12}^T X_i}) - H_{12}^o(V_i e^{-\beta_{12}^T X_i}) \right\} \right], \end{aligned}$$

$$\mathcal{E}_{1i}^{(m)} = E(\gamma_i|\mathcal{O}, \widehat{\Omega}^{(m)}) = \left(D_i + 1/\widehat{\sigma}^{(m)} \right) \left\{ 1/\widehat{\sigma}^{(m)} + \widehat{H}^{o(m)}(V_i, W_i, \widehat{\beta}^{(m)}) \right\}^{-1},$$

$$\mathcal{E}_{2i}^{(m)} = E(\log \gamma_i|\mathcal{O}, \widehat{\Omega}^{(m)}) = \Psi(D_i + 1/\widehat{\sigma}^{(m)}) - \log \left\{ 1/\widehat{\sigma}^{(m)} + \widehat{H}^{o(m)}(V_i, W_i, \widehat{\beta}^{(m)}) \right\},$$

$$\widehat{H}^{0(m)}(V_i, W_i, \widehat{\beta}^{(m)}) = \sum_{k=1,2} \widehat{H}_{0k}^{o(m)}(V_i e^{-\widehat{\beta}_{0k}^{(m)T} X_i}) + \delta_{1i} \left\{ \widehat{H}_{12}^{o(m)}(W_i e^{-\widehat{\beta}_{12}^{(m)T} X_i}) - \widehat{H}_{12}^{0(m)}(V_i e^{-\widehat{\beta}_{12}^{(m)T} X_i}) \right\}$$

and finally, $D_i = \sum_{k=1}^3 \delta_{ji}$, $\Gamma(x)$ is the Gamma function and $\Psi(x) = \Gamma'(x)/\Gamma(x)$ is the digamma function.

The M-step consists of maximization of Eq. (9). While maximizing $E(l(\sigma)|\mathcal{O}, \widehat{\Omega}^{(m)})$ as a function of σ can be done by gradient-based optimization algorithms, maximizing the other three expectations cannot be done directly with respect to (β_{jk}, h_{jk}^o) , $jk \in \{01, 02, 12\}$, due to a very non-smooth estimator of the cumulative hazard functions; see Zeng and Lin (2007) for more details under the standard univariate AFT model. Therefore, we aimed to find a smooth alternative and thus we extended the kernel-smoothing approach of Zeng and Lin (2007) and Liu et al. (2013) to accommodate our semi-competing risks setting.

We start with a simple case of piecewise constant hazard functions

$$\tilde{h}_{jk}^o(t) = \sum_{l=1}^{J_{jk,n}} c_{jk,l} I(t_{jk,l-1} \leq t < t_{jk,l})$$

where $0 = t_{jk,0} < t_{jk,1} < \dots < t_{jk,J_{jk,n}} = M_{jk}$, $jk \in \{01, 02, 12\}$, are equally spaced, M_{0k} are the respective upper bounds for $V_i \exp\{-\beta_{0k}^T X_i\}$, $k \in \{1, 2\}$, and M_{12} is the upper bound for $W_i \exp\{-\beta_{12}^T X_i\}$. Then, the cumulative baseline hazard functions are

$$\tilde{H}_{jk}^o(t) = \sum_{l=1}^{J_{jk,n}} c_{jk,l} (t - t_{jk,l-1}) I(t_{jk,l-1} \leq t < t_{jk,l}) + (M_{jk}/J_{jk,n}) \sum_{l=1}^{J_{jk,n}} c_{jk,l} I(t \geq t_{jk,l}).$$

The functions \tilde{h}_{jk}^o and \tilde{H}_{jk}^o are plugged in $E(l(\beta_{jk}, h_{jk}^o)|\mathcal{O}, \widehat{\Omega}^{(m)})$ and by maximizing the resulting expression with respect to $c_{jk,l}$, $l = 1, \dots, J_{jk,n}$, $jk \in \{01, 02, 12\}$, for a given β_{jk} , we left with a closed-form estimator of $c_{jk,l}$, $\widehat{c}_{jk,l}^{(m)}$. Plugging $\widehat{c}_{jk,l}^{(m)}$ in $E(l(\beta_{jk}, h_{jk}^o)|\mathcal{O}, \widehat{\Omega}^{(m)})$ provides an approximated profile-likelihood function of β_{jk} . However, even these profile-likelihood functions are not smooth and have multiple local maxima, and thus additional smoothing step is required. To this end, it can be shown that each of the profile likelihood converges to limiting function of β_{jk} , $jk \in \{01, 02, 12\}$, as $n \rightarrow \infty$, $J_{jk,n} \rightarrow \infty$ and $J_{jk,n}/n \rightarrow 0$. Then, for a given kernel function K with bandwidths $a_{jk,n}$, the estimators of β_{jk} , $jk \in \{01, 02, 12\}$, are obtained by maximizing a smoothed approximation of the limiting function. In the illness-death setting, the age of death after disease diagnosis is truncated by the age at diagnosis. Therefore, the kernel-smoothing approach is adopted to accommodate left truncation. Given the estimators of the regression coefficients, the proposed smoothing approach also yields estimators of the baseline hazard functions. Details of the above summary can be found in Section S4 of the WSM. Here we provide the resulting estimation procedure.

Define $R_i^V(\beta) = \log V_i - \beta^T X_i$, $R_i^W(\beta) = \log W_j - \beta^T X_i$. Then, β_{0k} is estimated by maximization of $l_{0k}^s(\beta_{0k})$, $k = 1, 2$, where

$$\begin{aligned} l_{0k}^s(\beta_{0k}) &= -\frac{1}{n} \sum_{i=1}^n \delta_{ki} \log V_i + \frac{1}{n} \sum_{i=1}^n \delta_{ki} \log \left\{ \frac{1}{na_{0k,n}} \sum_{j=1}^n \delta_{kj} K \left(\frac{R_j^V(\beta_{0k}) - R_i^V(\beta_{0k})}{a_{0k,n}} \right) \right\} \\ &\quad - \frac{1}{n} \sum_{i=1}^n \delta_{ki} \log \left\{ \frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \int_{-\infty}^{\{R_j^V(\beta_{0k}) - R_i^V(\beta_{0k})\}/a_{0k,n}} K(s) ds \right\}, \end{aligned} \tag{11}$$

β_{12} is estimated by maximization of

$$\begin{aligned} l_{12}^s(\beta_{12}) &= -\frac{1}{n} \sum_{i=1}^n \delta_{3i} \log W_i + \frac{1}{n} \sum_{i=1}^n \delta_{3i} \log \left\{ \frac{1}{na_{12,n}} \sum_{j=1}^n \delta_{3j} K \left(\frac{R_j^W(\beta_{12}) - R_i^W(\beta_{12})}{a_{12,n}} \right) \right\} \\ &\quad - \frac{1}{n} \sum_{i=1}^n \delta_{3i} \log \left\{ \frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \delta_{1j} \int_{\{R_j^V(\beta_{12}) - R_i^V(\beta_{12})\}/a_{12,n}}^{\{R_j^W(\beta_{12}) - R_i^W(\beta_{12})\}/a_{12,n}} K(s) ds \right\} \end{aligned} \tag{12}$$

and given $\widehat{\beta}_{jk}^{(m)}$, $jk \in \{01, 02, 12\}$, the baseline hazard functions are estimated by

$$\widehat{h}_{0k}^{o(m)}(t) = \frac{(na_{0k,n}t)^{-1} \sum_{i=1}^n \delta_{ki} K\left(\{R_i^V(\widehat{\beta}_{0k}^{(m)}) - \log t\}/a_{0k,n}\right)}{n^{-1} \sum_{i=1}^n \mathcal{E}_{1i}^{(m)} \int_{-\infty}^{\{R_i^V(\widehat{\beta}_{0k}^{(m)}) - \log t\}/a_{0k,n}} K(s) ds} \quad k = 1, 2, \quad (13)$$

$$\widehat{h}_{12}^{o(m)}(t) = \frac{(na_{12,n}t)^{-1} \sum_{i=1}^n \delta_{3i} K\left(\{R_i^W(\widehat{\beta}_{12}^{(m)}) - \log t\}/a_{12,n}\right)}{n^{-1} \sum_{i=1}^n \mathcal{E}_{1i}^{(m)} \delta_{1i} \int_{\{R_i^V(\widehat{\beta}_{12}^{(m)}) - \log t\}/a_{12,n}}^{\{R_i^W(\widehat{\beta}_{12}^{(m)}) - \log t\}/a_{12,n}} K(s) ds}, \quad (14)$$

and $\widehat{H}_{jk}^{o(m)}(t) = \int_0^t \widehat{h}_{jk}^{o(m)}(s) ds$. The following is a summary of our EM-based estimation algorithm:

Step 0 (Initial values): Set $\mathcal{E}_{1i}^{(0)} = 1$, $i = 1, \dots, n$ and $\beta_{12}^{(0)} = 0$. Get $\widehat{\beta}_{0k}^{(0)}$, $k \in \{1, 2\}$ by the rank-based estimator of the R package `aftgee` (Chiou et al., 2014). Get $\widehat{h}_{jk}^{o(0)}$ using Eq.'s (13)–(14) and $\widehat{H}_{jk}^{o(0)}(t)$ by adaptive quadrature (e.g. `integrate` function of R), $jk \in \{01, 02, 21\}$. Finally, set $\widehat{\sigma}^{(0)}$ between 2 to 5 and $m = 0$.

Step 1 (E-step): Set $m = m + 1$ and get $\mathcal{E}_{1i}^{(m)}$ and $\mathcal{E}_{2i}^{(m)}$, $i = 1, \dots, n$.

Step 2 (M-step): Obtain $\widehat{\sigma}^{(m)}$ by maximizing Eq. (10), and get $\widehat{\beta}_{jk}^{(m)}$ by maximizing Eq.'s (11)–(12). Obtain $\widehat{h}_{jk}^{0(m)}(t)$ and $\widehat{H}_{jk}^{0(m)}(t)$, $jk \in \{01, 02, 12\}$.

Step 3: Repeat Steps 1-2 until convergence is reached.

The asymptotic results of Liu et al. (2013, Theorem 1) for clustered data can be extended to establish the asymptotic properties of the proposed illness-death model estimators. Thus, it can be shown that under some regularity conditions, as $na_{jk,n}^2 \rightarrow \infty$, $na_{jk,n}^4 \rightarrow 0$ and $n \rightarrow \infty$, $\sup_{t \in [0, \tau]} |\widehat{H}_{jk}^o(t) - H_{jk}^o(t)| \rightarrow 0$, $\widehat{\beta}_{jk} \rightarrow \beta_{jk}$ almost surely, and $n^{-1/2}(\widehat{\beta}_{jk} - \beta_{jk})$, $jk \in \{01, 02, 12\}$, converges to a mean-zero multivariate normal distribution.

2.3.1 Bandwidth Selection

For the bandwidth parameters, $a_{jk,n}$, we recommend a modified version of the optimal bandwidths of Jones (1990) and Jones and Sheather (1991), in the spirit of Liu et al. (2013). The smoothed profile-likelihood function involves the kernel density for uncensored subjects and the cumulative kernel for all subjects. The recommended bandwidth for the kernel density is $\zeta \widehat{\tau}_{jk} (8\sqrt{2}/3)^{1/5} n_{jk}^{-1/5}$, $jk \in \{01, 02, 12\}$, where $\widehat{\tau}_{jk}$ is the sample standard deviation of $\log(T_{jki})$ among the subjects with observed event time of transition $j \rightarrow k$, denoted by n_{jk} . The recommended bandwidth for the cumulative kernel is $\zeta \widehat{v}_{jk} 4^{1/3} n_{jk}^{-1/3}$, where \widehat{v}_j is the sample standard deviation of $\log(V_i)$ or $\log(W_i)$ among all subjects in state j , denoted by n_j , $j = 0, 1$. Based on our extensive simulation study, $\zeta = 0.5$ is recommended for β_{jk} , and $\zeta = 0.01$ for h_{jk}^o .

2.3.2 Variance Estimation

In highly censored data, standard bootstrap could produce a low number of distinct event times, which often causes convergence failure. Alternatively, the weighted bootstrap approach can be used (Kosorok et al., 2004). At each bootstrap step, a random weight from the standard exponential distribution is assigned for each observation, and the estimators are derived by the respective weighted log-likelihood functions or weighted sums (see Section S5 of the WSM for details).

2.3.3 The Proposed Model and Estimation Method Without Frailty

The proposed estimation method can also be implemented under the model described by Eq.'s (1)–(6) where $\gamma_i \equiv 1$ for all $i = 1, \dots, n$, i.e., with no frailty effect. Namely, AFT models with competing risks, while model (3) is an

AFT model adjusted for left truncated data, where age at death after disease diagnosis is truncated by the age at diagnosis. If one believes that the functional form of age at diagnosis that effects the time to death after the diagnosis is known, and that frailty is not required, then age at diagnosis should be added as one of the covariates in model (3). In this case, the estimation process is simplified to one step for estimating the vectors of the regression coefficients and to one additional step for the baseline hazard functions' estimators, without the need of iterative process (see Section S6 of the WSM).

3 Visualizing Goodness of Fit

A goodness-of-fit procedure aimed to evaluate how closely observed data mirrors expected data under the assumed model. Recently, [Li et al. \(2021\)](#) proposed a goodness of fit (GOF) method for an arbitrary univariate survival model with right censored data which is based on the randomized survival probabilities (RSP). Their key idea is to replace the survival probability of a censored failure time with a uniform random number between 0 and the survival probability at the censored time. They showed that RSPs always uniformly distributed on (0, 1) under the true model. Then, graphical methods for comparing the distribution of the RSPs with the standard uniform distribution could be used for detecting a lack of model fit. In contrast, the distributions of well-known residuals (e.g. Cox-Snell) under the true model are complicated due to censoring and cannot be characterized clearly with a known distribution since their distribution depends on the censoring distribution. Hence, there is a lack of reference distributions for conducting GOF procedure, and the most widely used diagnostic tool is to apply the Kaplan-Meier estimator on the residual; see [Li et al. \(2021\)](#) and references therein. Here, we extend the RSP approach to the illness-death model.

In our setting of frailty-based AFT illness-death model, the marginal survival functions should be used since the frailties are unobserved. The illness-death model will be examined by two sets of RSPs: (i) the probability of remaining at Stat 0; and (ii) the probability of remaining at State 1 among those who diagnosed with the disease. In particular (for a detailed derivation, see Section S9 of the WSM),

$$S_{0.}^M(t|X_i) = \Pr(T_{1i} > t, T_{2i} > t|X_i) = \left[1 + \sigma \sum_{j=1,2} H_{0j}^o(te^{-\beta_{0j}^T X_i}) \right]^{-1/\sigma}$$

and for $t > t_1$,

$$\begin{aligned} S_{12}^M(t|t_1, X_i) &= \Pr(T_{2i} > t|T_{1i} = t_1, T_{2i} > t_1, X_i) \\ &= \left(\frac{1 + \sigma \sum_{j=1,2} H_{0j}^o(t_1 e^{-\beta_{0j}^T X_i})}{1 + \sigma \{ \sum_{j=1,2} H_{0j}^o(t_1 e^{-\beta_{0j}^T X_i}) + H_{12}^o(t_1 e^{-\beta_{12}^T X_i}) - H_{12}^o(t_1 e^{-\beta_{12}^T X_i}) \}} \right)^{1/\sigma+1}, \end{aligned}$$

where the superscript M denotes the marginal distribution with respect to the frailty variate. Clearly, in the absence of frailty (i.e. $\gamma_i \equiv 1$ for all $i = 1, \dots, n$) we get

$$S_{0.}^M(t|X_i) = \exp\{-H_{01}^o(te^{-\beta_{01}^T X_i}) - H_{02}^o(te^{-\beta_{02}^T X_i})\}$$

and

$$S_{12}^M(t|t_1, X_i) = \exp\{-H_{12}^o(te^{-\beta_{12}^T X_i}) + H_{12}^o(t_1 e^{-\beta_{12}^T X_i})\}, \quad t > t_1.$$

Now we are in a position to define the RSPs:

$$S_{0.}^M(V_i, \delta_{1i}, \delta_{2i}, U_{1i}) = (\delta_{1i} + \delta_{2i}) S_{0.}^M(V_i|X_i) + (1 - \delta_{1i} - \delta_{2i}) U_{1i} S_{0.}^M(V_i|X_i),$$

and

$$S_{12}^M(W_i, V_i, \delta_{3i}, U_{2i}) = \delta_{3i} S_{12}^M(W_i|V_i, X_i) + (1 - \delta_{3i}) U_{2i} S_{12}^M(W_i|V_i, X_i),$$

where U_{1i} and U_{2i} are independent random samples from the standard uniform distribution $U(0, 1)$. Based on [Li et al. \(2021\)](#) it can be shown that if the censoring and failure times are independent, given the observed covariates and the frailty variate, $S_{12}^M(t|t_1, X_i)$ and $S_{0.}^M(V_i, \delta_{1i}, \delta_{2i}, U_{1i})$ are uniformly distributed on (0, 1). See the WSM Section S9 for a detailed proof.

Finally, a visualized GOF procedure for any illness-death model could be accomplished by comparing the histograms of $\widehat{S}_0^M(V_i, \delta_{1i}, \delta_{2i}, U_{1i})$ and of $\widehat{S}_{12}^M(W_i, V_i, \delta_{3i}, U_{2i})$ with the expected values under the standard uniform distribution. This procedure will be demonstrated in Section 5.

4 Simulation Study

4.1 Simulation Setup

To demonstrate the finite sample properties of the proposed estimation method, we conducted an extensive simulation study. Failure times were generated from models (1)-(3) with $X_i = (X_{1i}, X_{2i}, X_{3i}, X_{4i})^T$, $\beta_{01} = (1, 0.5, 0, 0)^T$, $\beta_{02} = (0, 1, 1, 0)^T$ and $\beta_{12} = (0.5, 0.5, 0, 1)^T$, and a sample size of $n = 1000$. The baseline hazard functions of $\exp(\epsilon_{jki})$, $jk \in \{01, 02, 12\}$, were $h_{01}^0(t) = 2t$, $h_{02}^0(t) = 3t$, $h_{12}^0(t) = 2t$ and $X_{1i}, X_{2i}, X_{3i}, X_{4i}$ were sampled independently such that $X_{1i}, X_{3i}, X_{4i} \sim \text{Uniform}(-1, 1)$ and $X_{2i} \sim \text{Bernoulli}(0.5)$. Frailty variates γ_i were generated from gamma distribution with various dependence magnitudes $\sigma = 0.5, 1$ and 2 . Failure times, T_1 and T_2 , were generated by solving $U = \exp\{-\gamma H_{0k}^0(T e^{-\beta_{0k}^T X})\}$, $k \in \{1, 2\}$, for T , where U is uniformly distributed over $(0, 1)$. For those who diagnosed with the disease (i.e. $T_1 < T_2$) new values of T_2 were generated from the appropriate left-truncated distribution, by solving $U = \exp[-\gamma\{H_{12}^0(T e^{-\beta_{12}^T X}) - H_{12}^0(T_1 e^{\beta_{12}^T X})\}]$ for T and a new random U . Censoring times were sampled from $U(0, 15)$, such that for $\sigma = 2$ about 16% of observations were censored prior to disease diagnosis or death; and among those who diagnosed, about 13% were censored before death. Under $\sigma = 1$ the corresponding censoring rates were 9% and 10%, and for $\sigma = 0.5$, 7% and 8%. The analysis in this work uses the Gaussian kernel with bandwidths values according to Section 2.3.1. A range of values for ζ were studied, see Section S8 of the WSM. We set $\widehat{\sigma}^{[0]} = 2$. Finally, the convergence criteria were $\max_{1 \leq q \leq p_{jk}} |\widehat{\beta}_{jkq}^{(m+1)} - \widehat{\beta}_{jkq}^{(m)}| < 0.00001$, $\frac{1}{n_{jk}} \sum_{i=1}^{n_{jk}} |\widehat{H}_{jk}^{0(m+1)}(\tilde{t}_{jki}) - \widehat{H}_{jk}^{0(m)}(\tilde{t}_{jki})| < 0.0001$, $jk \in \{01, 02, 12\}$, and $|\widehat{\sigma}^{(m+1)} - \widehat{\sigma}^{(m)}| < 0.0001$, where p_{jk} is the number of components in β_{jk} , β_{jkq} denotes the q th component of β_{jk} , n_{jk} is the number of subjects relevant to transition $j \rightarrow k$, $\tilde{t}_{jki} = V_i e^{-\beta_{jk}^T X_i}$, $jk \in \{01, 02\}$ and $\tilde{t}_{12i} = W_i e^{-\beta_{12}^T X_i}$.

4.2 Simulation Results

Tables 2–3 present the performance of the proposed estimation method with frailty. The tables show the empirical mean, empirical standard deviations (SDs), estimated standard errors (SEs) and the empirical coverage rate of 95% Wald-type confidence interval (CR) of the dependence parameter, the regression coefficients and the cumulative baseline hazard functions at selected time points. Results are based on 100 repetitions. Tables 2–3 indicate that the proposed approach performs well in terms of bias and coverage rates. Moreover, the empirical standard deviations and the estimated standard errors are reasonably close.

Tables 4–5 present simulation results of model misspecification. The data were generated with gamma frailty effect but we applied our estimation procedure without frailty effect provided in Section 2.3.3. Clearly, under high dependence, ignoring the frailty, yield biased estimates and poor coverage rates. For example, with $\sigma = 2$ and true coefficients $\beta_{01} = (1, 0.5)^T$, $\beta_{02} = (1, 1)^T$ and $\beta_{12} = (0.5, 0.5, 1)^T$, the respective mean estimates were $\widehat{\beta}_{01} = (1.27, 0.36)^T$, $\widehat{\beta}_{02} = (1.15, 1.27)^T$ and $\widehat{\beta}_{12} = (0.25, 0.41, 2.09)^T$ with terribly poor coverage rates. In Section S7 and Table S1 of the WSM it is demonstrated that when the true model is free of frailty, our proposed approach of Section 2.3.3 performs very well in term of bias and coverage rate.

5 Rotterdam Tumor Bank Data

5.1 Data and Models

We analyzed the Rotterdam tumor bank of 1,546 breast cancer patients, whom had node-positive disease and underwent a tumor removal surgery between the years 1978-1993; the dataset is available in the **survival** R package ([Therneau, 2021](#)). T_1 is the time from surgery to relapse and T_2 is the time from surgery or relapse to death. Of the 1,546 patients, 924 patients showed a relapse of the disease (63%), 106 died without a relapse evidence (7%) and 771 patients died after a relapse (79% of the patients who showed a relapse of the cancer). Prognostic variables are age at primary surgery (in years), menopausal status (0 = premenopausal, 1 = postmenopausal), tumor size (≤ 20 mm, 20 – 50 mm and > 50 mm), tumor grade (2 = moderately differentiated, 3 = poorly differentiated), number of positive lymph nodes, estrogen and progesterone receptors in the initial biopsy (fmol/l), and chemotherapy (0 = no, 1 = yes).

The following methods were applied: the proposed AFT model and estimation procedure with gamma frailty (SEs are based on 500 bootstrap samples and starting value of σ was set to 1), the gamma-frailty Cox model of [Lee et al. \(2015\)](#), and the marginalized gamma-frailty of [Gorfine et al. \(2020\)](#). The AFT additive-frailty model of [Lee et al. \(2017\)](#) is implemented in the R package **SemicompRisks** ([Alvares et al., 2019](#)) with sojourn time, $T_2 - T_1$, when death occurs after the disease. Applying it to the current data, resulted in convergence failure (the potential scale reduction factors that should be less than 1.05 are much larger for most of the parameters). We hypothesize that the convergence failure is due to the use of sojourn time which could be negatively correlated with time from surgery to relapse, while a gamma frailty model assumes a positive dependence. Additionally, the data were analyzed with the proposed AFT model without frailty and a Cox illness-death model (for details, see section S11 in the WSM).

5.2 Results

Table 6 presents the estimates of the frailty-based methods with Holm adjusted p-values correction for multiplicity [Holm \(1979\)](#). Hereafter, a result is considered significant based on the adjusted p-value and a significance level of 0.05. The proposed model suggests that when holding the other covariates constant, higher age, higher progesterone, having chemotherapy, and having hormone therapy, each goes with a longer time to relapse after surgery. Also, higher number of positive lymph nodes, larger tumor size and poorly differentiated tumor are related to earlier relapse after the surgery. Hormonal treatment and chemotherapy after the surgery are associated with longer time to relapse after surgery.

In transition from surgery to death, higher age and higher number of positive lymph nodes are related with earlier death after surgery. For the transition from relapse to death, the proposed AFT model implies that lower number of positive lymph nodes, higher progesterone level associate with longer post-surgery survival time, given that the patient experienced relapse. Finally, the proposed AFT model indicates a strong dependence between time to relapse and time to death. The marginalized Cox model also indicates high level dependence between relapse and death times, while the conditional Cox model shows somewhat lower dependence. The directions of covariates' effect under the Cox models are similar to each other and to those of the proposed AFT model, but inference results based on these three models somewhat differ.

GOF assessment is done by a visual inspection of the histograms of the unmodified estimated survival probabilities and the RSPs histograms, shown in Figure 3. As expected, the unmodified histograms are far from that of a uniform distribution. However, based on the RSPs histogram, it is evident that the proposed AFT model fits well the data and is a better fit to the data, especially in comparison with the marginalized Cox model. The GOF plots of the models without frailty are shown in Figure S1 of the WSM, which indicate that the corresponding models with frailty fit the data much better.

6 Discussion

The main contribution of this work is twofold: (i) New semiparametric model and estimation method for a frailty-based AFT regression within the illness-death framework are provided. (ii) An exploratory method for diagnosis the appropriateness of any illness-death model is presented.

The proposed model includes covariates and allows to handle a possible residual dependency between the non-terminal the terminal failure times by incorporating a shared frailty. The estimation method can be applied both with or without frailty, and the simulation study indicates a good finite sample performances of the estimators in terms of bias and variance.

The proposed model and methods can be extended to other types of multi-state models, for example with multiple non-terminal events and a vector of random effects (frailties) capturing multiple levels of dependence among the event. Furthermore, additional work is required to extend the proposed estimation method for time-dependent covariates.

Acknowledgements

This work was supported by the Israel Science Foundation (ISF) grant number 767/21.

References

- Alvares, D., S. Haneuse, C. Lee, and K. H. Lee (2019). Semicomprisks: an r package for the analysis of independent and cluster-correlated semi-competing risks data. *The R journal* 11(1), 376.
- Chiou, S. H., S. Kang, and J. Yan (2014). Fitting accelerated failure time models in routine survival analysis with R package aftgee. *Journal of Statistical Software* 61(11), 1–23.
- Dempster, A. P., N. M. Laird, and D. B. Rubin (1977). Maximum likelihood from incomplete data via the em algorithm. *Journal of the Royal Statistical Society: Series B (Methodological)* 39(1), 1–22.
- Gorfine, M., N. Keret, A. Ben Arie, D. Zucker, and L. Hsu (2020). Marginalized frailty-based illness-death model: Application to the uk-biobank survival data. *Journal of the American Statistical Association* (just-accepted), 1–26.
- Holm, S. (1979). A simple sequentially rejective multiple test procedure. *Scandinavian journal of statistics*, 65–70.
- Jiang, F. and S. Haneuse (2017). A semi-parametric transformation frailty model for semi-competing risks survival data. *Scandinavian Journal of Statistics* 44(1), 112–129.
- Johnson, L. M. and R. L. Strawderman (2012). A smoothing expectation and substitution algorithm for the semi-parametric accelerated failure time frailty model. *Statistics in medicine* 31(21), 2335–2358.
- Jones, M. and S. Sheather (1991). Using non-stochastic terms to advantage in kernel-based estimation of integrated squared density derivatives. *Statistics & Probability Letters* 11(6), 511–514.
- Jones, M. C. (1990). The performance of kernel density functions in kernel distribution function estimation. *Statistics & Probability Letters* 9(2), 129–132.
- Kalbfleisch, J. D. and R. L. Prentice (2011). *The statistical analysis of failure time data*, Volume 360. John Wiley & Sons.
- Kosorok, M. R., B. L. Lee, J. P. Fine, et al. (2004). Robust inference for univariate proportional hazards frailty regression models. *The Annals of Statistics* 32(4), 1448–1491.
- Lee, K. H., S. Haneuse, D. Schrag, and F. Dominici (2015). Bayesian semi-parametric analysis of semi-competing risks data: Investigating hospital readmission after a pancreatic cancer diagnosis. *Journal of the Royal Statistical Society. Series C, Applied statistics* 64(2), 253.

- Lee, K. H., V. Rondeau, and S. Haneuse (2017). Accelerated failure time models for semi-competing risks data in the presence of complex censoring. *Biometrics* 73(4), 1401–1412.
- Li, L., T. Wu, and C. Feng (2021). Model diagnostics for censored regression via randomized survival probabilities. *Statistics in Medicine* 40(6), 1482–1497.
- Liu, B., W. Lu, and J. Zhang (2013). Kernel smoothed profile likelihood estimation in the accelerated failure time frailty model for clustered survival data. *Biometrika* 100(3), 741–755.
- Pan, W. (2001). Using frailties in the accelerated failure time model. *Lifetime Data Analysis* 7(1), 55–64.
- Therneau, T. M. (2021). *A Package for Survival Analysis in R*. R package version 3.2-13.
- Xu, J., J. D. Kalbfleisch, and B. Tai (2010). Statistical analysis of illness-death processes and semicompeting risks data. *Biometrics* 66(3), 716–725.
- Xu, L. and J. Zhang (2010). An em-like algorithm for the semiparametric accelerated failure time gamma frailty model. *Computational statistics & data analysis* 54(6), 1467–1474.
- Zeng, D. and D. Lin (2007). Efficient estimation for the accelerated failure time model. *Journal of the American Statistical Association* 102(480), 1387–1396.
- Zhang, J. and Y. Peng (2007). An alternative estimation method for the accelerated failure time frailty model. *Computational statistics & data analysis* 51(9), 4413–4423.




Figure 1: An illness-death model

Table 1: Illness-death Cox and AFT models, methods and software availability.

Authors	Model	Estimation Procedure	Software
Xu et al. (2010)	Cox, gamma frailty, semiparametric	Semi-parametric MLE	none
Lee et al. (2015)	Cox, gamma frailty, semiparametric	Bayesian	R package SemicompRisks
Jiang and Haneuse (2017)	Transformation model, known transformation function, nonparametric frailty at the price of known error distribution.	Semiparametric efficient score	none
Lee et al. (2017)	AFT, additive normal frailty, parametric and semiparametric	Baysian	R package SemicompRisks
Gorfine et al. (2020)	Cox, marginalized gamma frailty, semiparametric	Pseudo-likelihood approach	https://github.com/nirkeret/frailty-LTRC
Current work	AFT, multiplicative gamma frailty, semiparametric	Semi-parametric MLE	https://github.com/LeaKats/semicompAFT




Figure 2: The additive frailty-based AFT mode of Lee et al. (2017): conditional hazard of transition from healthy state to disease diagnosis.

Table 2: Simulation results with frailty: means of estimates, empirical standard deviations, bootstrap standard errors and empirical coverage rates.

σ		$\hat{\sigma}$	$\beta_{01,1}$	$\beta_{01,2}$	$\beta_{02,1}$	$\beta_{02,2}$	$\beta_{12,1}$	$\beta_{12,2}$	$\beta_{12,3}$
	True values		1	0.5	1	1	0.5	0.5	1
2	mean	1.84	1.05	0.48	1.03	1.05	0.51	0.49	1.06
	empirical SD	0.27	0.10	0.13	0.12	0.10	0.11	0.12	0.10
	bootstrap SE	0.29	0.11	0.12	0.12	0.11	0.13	0.14	0.15
	CR	0.95	0.95	0.94	0.93	0.96	0.99	0.99	0.96
1	mean	0.90	1.05	0.48	1.00	1.03	0.50	0.50	1.06
	empirical SD	0.16	0.08	0.08	0.08	0.08	0.08	0.10	0.11
	bootstrap SE	0.16	0.09	0.10	0.10	0.09	0.10	0.11	0.12
	CR	0.89	0.93	0.97	0.97	0.97	0.99	0.99	0.93
0.5	mean	0.44	1.02	0.51	1.02	1.02	0.51	0.51	1.05
	empirical SD	0.10	0.06	0.07	0.07	0.07	0.07	0.08	0.08
	bootstrap SE	0.10	0.08	0.08	0.09	0.08	0.09	0.09	0.10
	CR	0.93	0.98	0.98	0.96	0.97	0.96	0.96	0.99

Table 3: Simulation results of the cumulative baseline hazard functions with frailty: means of estimates, empirical standard deviations, bootstrap standard errors and empirical coverage rates.

σ	t	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	1.00
2	$H_{01}^0(t)$	0.01	0.04	0.09	0.16	0.25	0.36	0.49	0.64	0.81	1.00
	mean	0.01	0.04	0.09	0.15	0.24	0.34	0.45	0.59	0.75	0.91
	empirical SD	0.00	0.01	0.02	0.03	0.05	0.07	0.09	0.12	0.15	0.19
	bootstrap SE	0.00	0.01	0.02	0.03	0.04	0.07	0.09	0.13	0.17	0.21
	CR	0.96	0.93	0.95	0.93	0.97	0.96	0.95	0.97	0.98	0.97
	$H_{02}^0(t)$	0.02	0.06	0.14	0.24	0.38	0.54	0.74	0.96	1.22	1.50
	mean	0.02	0.06	0.13	0.24	0.37	0.52	0.70	0.90	1.13	1.38
	empirical SD	0.00	0.01	0.02	0.04	0.06	0.08	0.11	0.15	0.21	0.28
	bootstrap SE	0.01	0.01	0.02	0.04	0.06	0.09	0.13	0.19	0.25	0.32
	CR	0.99	0.97	0.96	0.97	0.98	0.97	0.99	0.99	0.98	0.98
	$H_{12}^0(t)$	0.01	0.04	0.09	0.16	0.25	0.36	0.49	0.64	0.81	1.00
	mean	0.01	0.05	0.10	0.17	0.27	0.38	0.51	0.66	0.82	1.00
	empirical SD	0.03	0.04	0.05	0.05	0.07	0.09	0.11	0.13	0.15	0.18
	bootstrap SE	0.03	0.05	0.07	0.09	0.11	0.13	0.15	0.18	0.20	0.24
	CR	0.96	0.98	0.98	0.99	0.99	0.98	0.98	0.98	0.98	0.99
1	$H_{01}^0(t)$	0.01	0.04	0.09	0.16	0.25	0.36	0.49	0.64	0.81	1.00
	mean	0.01	0.04	0.09	0.15	0.24	0.34	0.46	0.60	0.75	0.91
	empirical SD	0.00	0.01	0.01	0.02	0.03	0.04	0.06	0.08	0.10	0.13
	bootstrap SE	0.00	0.01	0.01	0.02	0.04	0.05	0.07	0.10	0.13	0.17
	CR	0.97	0.99	0.98	0.97	1.00	0.99	0.98	1.00	0.99	
	$H_{02}^0(t)$	0.02	0.06	0.14	0.24	0.38	0.54	0.74	0.96	1.22	1.50
	mean	0.02	0.06	0.13	0.23	0.36	0.52	0.71	0.92	1.15	1.40
	empirical SD	0.00	0.01	0.02	0.03	0.04	0.07	0.10	0.13	0.17	0.22
	bootstrap SE	0.00	0.01	0.02	0.03	0.05	0.07	0.10	0.14	0.19	0.25
	CR	0.96	0.98	0.98	0.98	0.95	0.96	0.97	0.98	0.95	
	$H_{12}^0(t)$	0.01	0.04	0.09	0.16	0.25	0.36	0.49	0.64	0.81	1.00
	mean	0.01	0.04	0.10	0.17	0.27	0.38	0.51	0.66	0.83	1.02
	empirical SD	0.03	0.03	0.04	0.05	0.07	0.08	0.10	0.11	0.13	0.15
	bootstrap SE	0.02	0.04	0.06	0.07	0.09	0.11	0.13	0.15	0.17	0.20
	CR	0.95	0.97	0.98	0.97	0.96	0.96	0.96	0.99	0.99	
0.5	$H_{01}^0(t)$	0.01	0.04	0.09	0.16	0.25	0.36	0.49	0.64	0.81	1.00
	mean	0.01	0.04	0.09	0.16	0.25	0.35	0.48	0.63	0.79	0.97
	empirical SD	0.00	0.01	0.01	0.02	0.03	0.04	0.05	0.08	0.10	0.13
	bootstrap SE	0.00	0.01	0.01	0.02	0.03	0.05	0.06	0.08	0.11	0.14
	CR	0.92	0.98	0.97	0.98	0.97	0.98	0.97	0.96	0.97	0.95
	$H_{02}^0(t)$	0.02	0.06	0.14	0.24	0.38	0.54	0.74	0.96	1.22	1.50
	mean	0.02	0.06	0.13	0.24	0.37	0.53	0.72	0.93	1.18	1.44
	empirical SD	0.00	0.01	0.02	0.03	0.04	0.06	0.08	0.11	0.15	0.20
	bootstrap SE	0.00	0.01	0.02	0.03	0.04	0.06	0.09	0.12	0.15	0.20
	CR	0.96	1.00	0.98	0.97	0.97	0.95	0.94	0.94	0.93	0.96
	$H_{12}^0(t)$	0.01	0.04	0.09	0.16	0.25	0.36	0.49	0.64	0.81	1.00
	mean	0.01	0.04	0.09	0.17	0.27	0.38	0.52	0.67	0.85	1.04
	empirical SD	0.02	0.03	0.04	0.05	0.06	0.07	0.09	0.11	0.13	0.14
	bootstrap SE	0.02	0.04	0.05	0.06	0.08	0.09	0.11	0.13	0.14	0.16
	CR	0.89	0.96	0.98	0.99	0.98	0.98	0.98	0.96	0.95	0.96

Table 4: Simulation results of model misspecification where frailty is ignored: means of estimates, empirical standard deviations and empirical coverage rates.

σ		$\beta_{01,1}$	$\beta_{01,2}$	$\beta_{02,1}$	$\beta_{02,2}$	$\beta_{12,1}$	$\beta_{12,2}$	$\beta_{12,3}$
	true values	1	0.5	1	1	0.5	0.5	1
2	mean	1.27	0.36	1.15	1.27	0.25	0.41	2.09
	empirical SD	0.10	0.14	0.15	0.15	0.32	0.31	0.50
	CR	0.25	0.86	0.82	0.53	0.90	0.94	0.45
1	mean	1.19	0.42	1.08	1.17	0.37	0.46	1.63
	empirical SD	0.09	0.09	0.10	0.09	0.16	0.16	0.19
	CR	0.48	0.86	0.87	0.56	0.86	0.93	0.08
0.5	mean	1.09	0.48	1.06	1.10	0.44	0.47	1.33
	empirical SD	0.07	0.08	0.08	0.08	0.11	0.12	0.14
	CR	0.70	0.94	0.89	0.72	0.91	0.96	0.37

Table 5: Simulation results of the cumulative baseline hazard functions under model misspecification where frailty is ignored: means of estimates, empirical standard deviations, and empirical coverage rates.

σ	t	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	1.00
2	$H_{01}^0(t)$	0.01	0.04	0.09	0.16	0.25	0.36	0.49	0.64	0.81	1.00
	mean	0.01	0.03	0.06	0.10	0.14	0.19	0.23	0.28	0.32	0.36
	empirical SD	0.00	0.01	0.01	0.02	0.02	0.03	0.03	0.04	0.04	0.04
	CR	0.96	0.73	0.31	0.06	0.00	0.00	0.00	0.00	0.00	0.00
	$H_{02}^0(t)$	0.02	0.06	0.14	0.24	0.38	0.54	0.74	0.96	1.22	1.50
	mean	0.02	0.06	0.11	0.17	0.23	0.29	0.35	0.40	0.46	0.51
	empirical SD	0.00	0.01	0.02	0.02	0.03	0.03	0.03	0.04	0.04	0.04
	CR	0.91	0.92	0.67	0.12	0.00	0.00	0.00	0.00	0.00	0.00
	$H_{12}^0(t)$	0.01	0.04	0.09	0.16	0.25	0.36	0.49	0.64	0.81	1.00
	mean	0.19	0.44	0.66	0.88	1.07	1.26	1.43	1.58	1.73	1.89
	empirical SD	0.24	0.36	0.42	0.45	0.48	0.50	0.52	0.52	0.55	0.56
	CR	0.90	0.86	0.82	0.75	0.72	0.69	0.68	0.68	0.72	0.77
1	$H_{01}^0(t)$	0.01	0.04	0.09	0.16	0.25	0.36	0.49	0.64	0.81	1.00
	mean	0.01	0.03	0.07	0.12	0.18	0.24	0.31	0.37	0.44	0.51
	empirical SD	0.00	0.01	0.01	0.02	0.02	0.03	0.03	0.04	0.04	0.05
	CR	0.92	0.84	0.63	0.28	0.07	0.00	0.00	0.00	0.00	0.00
	$H_{02}^0(t)$	0.02	0.06	0.14	0.24	0.38	0.54	0.74	0.96	1.22	1.50
	mean	0.02	0.06	0.12	0.19	0.28	0.37	0.46	0.55	0.64	0.73
	empirical SD	0.00	0.01	0.01	0.02	0.03	0.04	0.04	0.05	0.05	0.06
	CR	0.93	0.94	0.79	0.38	0.09	0.00	0.00	0.00	0.00	0.00
	$H_{12}^0(t)$	0.01	0.04	0.09	0.16	0.25	0.36	0.49	0.64	0.81	1.00
	mean	0.06	0.17	0.32	0.47	0.65	0.81	0.97	1.14	1.31	1.46
	empirical SD	0.09	0.11	0.14	0.16	0.17	0.19	0.20	0.22	0.23	0.24
	CR	0.96	0.85	0.71	0.53	0.44	0.37	0.33	0.37	0.47	0.58
0.5	$H_{01}^0(t)$	0.01	0.04	0.09	0.16	0.25	0.36	0.49	0.64	0.81	1.00
	mean	0.01	0.04	0.08	0.14	0.21	0.29	0.38	0.48	0.58	0.69
	empirical SD	0.00	0.01	0.01	0.02	0.02	0.03	0.04	0.05	0.06	0.07
	CR	0.92	0.92	0.91	0.82	0.58	0.40	0.21	0.11	0.01	0.01
	$H_{02}^0(t)$	0.02	0.06	0.14	0.24	0.38	0.54	0.74	0.96	1.22	1.50
	mean	0.02	0.06	0.13	0.22	0.32	0.44	0.56	0.70	0.83	0.97
	empirical SD	0.00	0.01	0.02	0.02	0.03	0.04	0.05	0.07	0.07	0.08
	CR	0.92	0.96	0.94	0.81	0.57	0.36	0.14	0.03	0.00	0.00
	$H_{12}^0(t)$	0.01	0.04	0.09	0.16	0.25	0.36	0.49	0.64	0.81	1.00
	mean	0.02	0.09	0.18	0.29	0.43	0.58	0.73	0.89	1.06	1.24
	empirical SD	0.04	0.06	0.08	0.10	0.12	0.14	0.16	0.17	0.18	0.20
	CR	0.92	0.91	0.84	0.75	0.72	0.70	0.73	0.72	0.74	0.78

Table 6: Rotterdam Tumor Bank Data: Estimates (Est), exponent of estimated regression coefficients (exp), standard errors (SE), p-values and Holm's adjusted p-values. Bold results are significant at 0.05 based on Holm's adjusted p-value.

σ	Proposed Model ($\zeta=65$)						Marginalized Cox (Gorfine et al. 2021)						Conditional Cox (Lee et al. 2015)					
	Est	exp	SE	p-value	Holm	Est	exp	SE	p-value	Holm	Est	exp	SE	p-value	Holm	Est	exp	SE
Transition: surgery \rightarrow relapse																		
Age at surgery (divided by 10)	0.14	1.15	0.06	0.012	0.185	-0.15	0.86	0.06	0.014	0.262	-0.22	0.80	0.08	0.003	0.048			
log of lymph nodes	-0.40	0.67	0.05	0.000	0.000	0.42	1.53	0.04	0.000	0.000	0.71	2.03	0.07	0.000	0.000			
log of estrogen+1	0.07	1.07	0.03	0.030	0.390	-0.03	0.97	0.02	0.186	1.000	-0.10	0.90	0.04	0.003	0.047			
log of progesterone+1	0.09	1.09	0.02	0.000	0.005	-0.04	0.96	0.02	0.065	1.000	-0.11	0.90	0.03	0.001	0.020			
Postmenopausal (vs. premenopausal)	-0.34	0.71	0.15	0.023	0.328	0.13	1.14	0.13	0.296	1.000	0.34	1.40	0.19	0.071	0.785			
Tumor size (ref. < 20mm)	-0.32	0.73	0.09	0.001	0.015	0.20	1.22	0.07	0.006	0.116	0.40	1.49	0.12	0.001	0.019			
> 50mm	-0.49	0.61	0.11	0.000	0.000	0.38	1.46	0.11	0.001	0.020	0.79	2.19	0.16	0.000	0.000			
Hormone therapy	0.60	1.83	0.13	0.000	0.000	-0.38	0.68	0.08	0.000	0.000	-0.88	0.41	0.15	0.000	0.000			
Chemotherapy	0.49	1.64	0.11	0.000	0.000	-0.37	0.69	0.11	0.001	0.023	-0.79	0.46	0.16	0.000	0.000			
Tumor grade 3 (vs. 2)	-0.25	0.78	0.09	0.004	0.081	0.21	1.23	0.08	0.008	0.155	0.44	1.56	0.13	0.000	0.010			
Transition: surgery \rightarrow death																		
Age at surgery (divided by 10)	-0.43	0.65	0.14	0.002	0.051	1.32	3.74	0.37	0.000	0.009	1.43	4.20	0.18	0.000	0.000			
log of lymph nodes	-0.14	0.87	0.08	0.091	1.000	0.13	1.14	0.12	0.298	1.000	0.44	1.54	0.15	0.004	0.049			
log of estrogen+1	0.04	1.04	0.04	0.287	1.000	-0.01	0.99	0.06	0.816	1.000	-0.11	0.89	0.08	0.149	1.000			
log of progesterone+1	0.01	1.01	0.04	0.827	1.000	0.08	1.08	0.06	0.205	1.000	0.01	1.01	0.07	0.857	1.000			
Postmenopausal (vs. premenopausal)	-0.15	0.86	0.34	0.647	1.000	-0.30	0.74	0.50	0.554	1.000	-0.35	0.70	0.70	0.613	1.000			
Tumor size (ref. < 20mm)	-0.13	0.88	0.15	0.376	1.000	-0.16	0.85	0.25	0.526	1.000	-0.04	0.96	0.28	0.874	1.000			
> 50mm	-0.19	0.82	0.18	0.275	1.000	0.15	1.16	0.31	0.634	1.000	0.58	1.79	0.35	0.096	0.956			
Hormone therapy	0.41	1.51	0.18	0.019	0.290	-0.21	0.81	0.25	0.389	1.000	-0.69	0.50	0.29	0.018	0.220			
Chemotherapy	1.13	3.09	0.30	0.000	0.005	-0.22	0.81	0.81	0.789	1.000	-0.78	0.46	0.63	0.215	1.000			
Tumor grade 3 (vs. 2)	-0.06	0.94	0.13	0.641	1.000	-0.01	0.99	0.28	0.961	1.000	0.21	1.23	0.27	0.446	1.000			
Transition: relapse \rightarrow death																		
Age at surgery (divided by 10)	0.00	1.00	0.07	0.956	1.000	0.03	1.03	0.08	0.700	1.000	0.08	1.08	0.07	0.278	1.000			
log of lymph nodes	-0.25	0.78	0.07	0.000	0.010	0.25	1.28	0.05	0.000	0.000	0.38	1.47	0.07	0.000	0.000			
log of estrogen+1	0.04	1.04	0.05	0.341	1.000	-0.03	0.97	0.02	0.193	1.000	-0.10	0.90	0.04	0.008	0.102			
log of progesterone+1	0.13	1.14	0.04	0.001	0.021	-0.08	0.92	0.02	0.000	0.003	-0.19	0.83	0.04	0.000	0.000			
Postmenopausal (vs. premenopausal)	-0.21	0.81	0.17	0.203	1.000	-0.05	0.95	0.13	0.731	1.000	0.04	1.04	0.20	0.843	1.000			
Tumor size (ref. < 20mm)	-0.37	0.69	0.14	0.008	0.131	0.23	1.26	0.07	0.001	0.024	0.46	1.58	0.14	0.001	0.015			
> 50mm	-0.52	0.60	0.17	0.002	0.044	0.40	1.49	0.10	0.000	0.002	0.67	1.96	0.18	0.000	0.003			
Hormone therapy	0.39	1.48	0.14	0.005	0.090	-0.18	0.84	0.09	0.037	0.633	-0.48	0.62	0.16	0.002	0.037			
Chemotherapy	0.23	1.25	0.18	0.205	1.000	-0.16	0.85	0.13	0.227	1.000	-0.18	0.84	0.17	0.287	1.000			
Tumor grade 3 (vs. 2)	-0.26	0.77	0.13	0.047	0.569	0.21	1.23	0.09	0.024	0.440	0.43	1.54	0.14	0.002	0.032			




Figure 3: Goodness of fit plots for the illness-death models. Histograms of $\widehat{S}_0^M(V_i|X_i)$ (left of (a), (c), (e)) $\widehat{S}_0^M(V_i, \delta_{1i}, \delta_{2i}, U_{1i})$ (right of (a), (c), (e)), $\widehat{S}_{12}^M(W_i|V_i, X_i)$ (left of (b), (d), (f)) and $\widehat{S}_{12}^M(W_i, V_i, \delta_{3i}, U_{2i})$ (right of (b), (d), (f)). The dashed lines are the expected values under the uniform distribution.

Web Supplementary Material
An Accelerated Failure Time Regression Model for
Illness-Death Data:
A Frailty Approach

Lea Kats and Malka Gorfine
 Department of Statistics and Operations Research
 Tel Aviv University, Israel

S1 Conditional Hazards

Based on Eq.'s(1)-(6) of the main text, we derive the conditional hazard functions of the three transitions, given (X_i, γ_i) , namely, following conditional hazard functions

$$\begin{aligned}\lambda_{01}(t|X_i, \gamma_i) &= \lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(t \leq T_{1i} < t + \Delta | T_{1i} \geq t, T_{2i} \geq t, X_i, \gamma_i), \quad t > 0, \\ \lambda_{02}(t|X_i, \gamma_i) &= \lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(t \leq T_{2i} < t + \Delta | T_{1i} \geq t, T_{2i} \geq t, X_i, \gamma_i), \quad t > 0, \\ \lambda_{12}(t|t_1, X_i, \gamma_i) &= \lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(t \leq T_{2i} < t + \Delta | T_{1i} = t_1, T_{2i} \geq t, X_i, \gamma_i), \quad t > t_1 > 0.\end{aligned}$$

To this end, write

$$\begin{aligned}\lambda_{01}(t|X_i, \gamma_i) &= \frac{\lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(T_{1i} \in [t, t + \Delta], T_{2i} \geq t | X_i, \gamma_i)}{\Pr(T_{1i} \geq t, T_{2i} \geq t | X_i, \gamma_i)} \\ &= \frac{\lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(e^{\epsilon_{01i}} \in [te^{-\beta_{01}^T X_i}, (t + \Delta)e^{-\beta_{01}^T X_i}], e^{\epsilon_{02i}} \geq te^{-\beta_{02}^T X_i} | X_i, \gamma_i)}{\Pr(e^{\epsilon_{01i}} \geq te^{-\beta_{01}^T X_i}, e^{\epsilon_{02i}} \geq te^{-\beta_{02}^T X_i} | X_i, \gamma_i)} \\ &= \frac{\lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(e^{\epsilon_{01i}} \in [te^{-\beta_{01}^T X_i}, (t + \Delta)e^{-\beta_{01}^T X_i}] | X_i, \gamma_i) \Pr(e^{\epsilon_{02i}} \geq te^{-\beta_{02}^T X_i} | X_i, \gamma_i)}{\Pr(e^{\epsilon_{01i}} \geq te^{-\beta_{01}^T X_i} | X_i, \gamma_i) \Pr(e^{\epsilon_{02i}} \geq te^{-\beta_{02}^T X_i} | X_i, \gamma_i)} \\ &= -\frac{\partial}{\partial t} \log S_{\epsilon_{01}}(te^{-\beta_{01}^T X_i} | \gamma_i) = \frac{f_{\epsilon_{01}}(te^{-\beta_{01}^T X_i} | \gamma_i) e^{-\beta_{01}^T X_i}}{S_{\epsilon_{01}}(te^{-\beta_{01}^T X_i} | \gamma_i)} = \lambda_{01}(te^{-\beta_{01}^T X_i} | \gamma_i) e^{-\beta_{01}^T X_i} \\ &= \gamma_i h_{01}^o(te^{-\beta_{01}^T X_i}) e^{-\beta_{01}^T X_i}, \quad t > 0.\end{aligned}$$

Due to the similarity between T_{1i} and T_{2i} at transitions 0 → 1 and 0 → 2,

$$\lambda_{02}(t|X_i, \gamma_i) = \frac{\lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(T_{2i} \in [t, t + \Delta], T_{1i} \geq t | X_i, \gamma_i)}{\Pr(T_{1i} \geq t, T_{2i} \geq t | X_i, \gamma_i)} = \gamma_i h_{02}^o(te^{-\beta_{02}^T X_i}) e^{-\beta_{02}^T X_i}, \quad t > 0.$$

Finally, since

$$\lambda_{12}(t|t_1, X_i, \gamma_i) = -\frac{\partial}{\partial t} \log \Pr(T_{2i} > t | T_{1i} = t_1, T_{2i} > t_1, X_i, \gamma_i),$$

we first derive $\Pr(T_{2i} > t | T_{1i} = t_1, T_{2i} > t_1, X_i, \gamma_i)$, the probability of staying in State 1 until time t , after diagnosed with the disease at time t_1 . Then, for $t > t_1 > 0$,

$$\begin{aligned}\Pr(T_{2i} > t | T_{1i} = t_1, T_{2i} > t_1, X_i, \gamma_i) &= \frac{\lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(T_{2i} > t, T_{1i} \in [t_1, t_1 + \Delta], T_{2i} > t_1 | X_i, \gamma_i)}{\lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(T_{1i} \in [t_1, t_1 + \Delta], T_{2i} > t_1 | X_i, \gamma_i)}\end{aligned}$$

$$\begin{aligned}
&= \frac{\lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(T_{2i} > t, T_{1i} \in [t_1, t_1 + \Delta) | X_i, \gamma_i)}{\lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(T_{1i} \in [t_1, t_1 + \Delta), T_{2i} > t_1 | X_i, \gamma_i)} \\
&= \frac{\Pr(T_{2i} > t | X_i, \gamma_i) \lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(T_{1i} \in [t_1, t_1 + \Delta) | X_i, \gamma_i)}{\Pr(T_{2i} > t_1 | X_i, \gamma_i) \lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(T_{1i} \in [t_1, t_1 + \Delta) | X_i, \gamma_i)} \\
&= \frac{\Pr(e^{\epsilon_{12}} > te^{-\beta_{12}^T X_i} | X_i, \gamma_i)}{\Pr(e^{\epsilon_{12}} > t_1 e^{-\beta_{12}^T X_i} | X_i, \gamma_i)} = \frac{S_{\epsilon_{12}}(te^{-\beta_{12}^T X_i} | \gamma_i)}{S_{\epsilon_{12}}(t_1 e^{-\beta_{12}^T X_i} | \gamma_i)} \\
&= \frac{\exp \left\{ -\gamma_i \int_0^{te^{-\beta_{12}^T X_i}} h_{12}^o(v) dv \right\}}{\exp \left\{ -\gamma_i \int_0^{t_1 e^{-\beta_{12}^T X_i}} h_{12}^o(v) dv \right\}} \\
&= \exp \left\{ -\gamma_i \left[H_{12}^o(te^{-\beta_{12}^T X_i}) - H_{12}^o(t_1 e^{-\beta_{12}^T X_i}) \right] \right\}.
\end{aligned}$$

Hence,

$$\begin{aligned}
\lambda_{12}(t | t_1, X_i, \gamma_i) &= -\frac{\partial}{\partial t} \log \left(\exp \left\{ -\gamma_i \left[H_{12}^o(te^{-\beta_{12}^T X_i}) - H_{12}^o(t_1 e^{-\beta_{12}^T X_i}) \right] \right\} \right) \\
&= \gamma_i h_{12}^o(te^{-\beta_{12}^T X_i}) e^{-\beta_{12}^T X_i}, \quad t > t_1 > 0.
\end{aligned}$$

S2 Likelihood Function

Based on Eq.'s (1)-(6) of the main text, the contribution of individual i to the complete-data likelihood is given by

$$\begin{aligned}
&f_{V,W,\delta_1,\delta_2,\delta_3,\gamma}(V_i, W_i, \delta_{1i}, \delta_{2i}, \delta_{3i}, \gamma_i | X_i) \\
&= f_{V,W,\delta_1,\delta_2,\delta_3}(V_i, W_i, \delta_{1i}, \delta_{2i}, \delta_{3i} | X_i, \gamma_i) f_{\gamma|X}(\gamma_i | X_i) \\
&= f_{V,W,\delta_1,\delta_2,\delta_3}(V_i, W_i, \delta_{1i}, \delta_{2i}, \delta_{3i} | X_i, \gamma_i) f(\gamma_i; \sigma) \\
&= f(V_i, \delta_{1i}, \delta_{2i} | X_i, \gamma_i) f(W_i, \delta_{3i} | V_i, W_i > V_i, \delta_{1i} = 1, X_i, \gamma_i)^{\delta_{1i}} f(\gamma_i; \sigma) \\
&\propto \lambda_{01}(V_i | X_i, \gamma_i)^{\delta_{1i}} \lambda_{02}(V_i | X_i, \gamma_i)^{\delta_{2i}} \exp \left\{ -\gamma_i \left[H_{01}^o(V_i e^{-\beta_{01}^T X_i}) + H_{02}^o(V_i e^{-\beta_{02}^T X_i}) \right] \right\} \\
&\quad \left(\lambda_{12}(W_i | V_i, X_i, \gamma_i)^{\delta_{3i}} \exp \left\{ -\gamma_i \left[H_{12}^o(W_i e^{-\beta_{12}^T X_i}) - H_{12}^o(V_i e^{-\beta_{12}^T X_i}) \right] \right\} \right)^{\delta_{1i}} f(\gamma_i; \sigma) \\
&= \left\{ \gamma_i h_{01}^o(V_i e^{-\beta_{01}^T X_i}) e^{-\beta_{01}^T X_i} \right\}^{\delta_{1i}} \left\{ \gamma_i h_{02}^o(V_i e^{-\beta_{02}^T X_i}) e^{-\beta_{02}^T X_i} \right\}^{\delta_{2i}} \exp \left\{ -\gamma_i H_{01}^o(V_i e^{-\beta_{01}^T X_i}) \right\} \\
&\quad \exp \left\{ -\gamma_i H_{02}^o(V_i e^{-\beta_{02}^T X_i}) \right\} \left[\left\{ \gamma_i h_{12}^o(W_i e^{-\beta_{12}^T X_i}) e^{-\beta_{12}^T X_i} \right\}^{\delta_{3i}} \right. \\
&\quad \left. \exp \left\{ -\gamma_i \left(H_{12}^o(W_i e^{-\beta_{12}^T X_i}) - H_{12}^o(V_i e^{-\beta_{12}^T X_i}) \right) \right\} \right]^{\delta_{1i}} f(\gamma_i; \sigma). \tag{15}
\end{aligned}$$

S3 Conditional Expectations in the EM Algorithm

Given Eq. (15), the complete-data log-likelihood function can be written as

$$l(\Omega) = \log L(\Omega) = l(\sigma) + l(\beta_{01}, h_{01}^o) + l(\beta_{02}, h_{02}^o) + l(\beta_{12}, h_{12}^o)$$

where

$$l(\sigma) = \frac{1}{n} \sum_{i=1}^n \log f(\gamma_i; \sigma) + \frac{1}{n} \sum_{i=1}^n (\delta_{1i} + \delta_{2i} + \delta_{3i}) \log \gamma_i,$$

$$l(\beta_{0k}, h_{0k}^o) = \frac{1}{n} \sum_{i=1}^n \left\{ \delta_{ki} \left[\log h_{0k}^o(V_i e^{-\beta_{0k}^T X_i}) - \beta_{0k}^T X_i \right] - \gamma_i H_{0k1}^o(V_i e^{-\beta_{0k}^T X_i}) \right\}, \quad k = 1, 2,$$

and

$$\begin{aligned} l(\beta_{12}, h_{12}^o) &= \frac{1}{n} \sum_{i=1}^n \left\{ \delta_{3i} \left[\log h_{12}^o(W_i e^{-\beta_{12}^T X_i}) - \beta_{12}^T X_i \right] \right. \\ &\quad \left. - \delta_{1i} \gamma_i \left[H_{12}^o(W_i e^{-\beta_{12}^T X_i}) - H_{12}^o(V_i e^{-\beta_{12}^T X_i}) \right] \right\}. \end{aligned}$$

In the E-step of the EM algorithm, the expectation of the complete log-likelihood function given the observed data, $\mathcal{O} = \{(V_i, W_i, \delta_{1i}, \delta_{2i}, \delta_{3i}, X_i) : i = 1, \dots, n\}$, and the current values of the estimated parameters is required. The conditional density of γ_i , given the observed data $\mathcal{O}_i = (V_i, W_i, \delta_{1i}, \delta_{2i}, \delta_{3i}, X_i)$ and the parameter estimates at step m , denoted by $\widehat{\Omega}^{(m)} = (\widehat{\beta}_{01}^{(m)}, \widehat{\beta}_{12}^{(m)}, \widehat{\beta}_{12}^{(m)}, \widehat{h}_{01}^{(m)}, \widehat{h}_{02}^{(m)}, \widehat{h}_{12}^{(m)}, \widehat{\sigma}^{(m)})$, is obtained by

$$f(\gamma_i | \mathcal{O}_i, \widehat{\Omega}^{(m)}) = \frac{f(\mathcal{O}_i | \gamma_i, \widehat{\Omega}^{(m)}) f(\gamma_i; \widehat{\sigma}^{(m)})}{\int_0^\infty f(\mathcal{O}_i | \gamma_i, \widehat{\Omega}^{(m)}) f(\gamma_i; \widehat{\sigma}^{(m)}) d\gamma_i},$$

where

$$\begin{aligned} f(\mathcal{O}_i | \gamma_i, \widehat{\Omega}^{(m)}) &= \left\{ \gamma_i \widehat{h}_{01}^{(m)} (V_i e^{-\widehat{\beta}_{01}^{(m)T} X_i}) e^{-\widehat{\beta}_{01}^{(m)T} X_i} \right\}^{\delta_{1i}} \left\{ \gamma_i \widehat{h}_{02}^{(m)} (V_i e^{-\widehat{\beta}_{02}^{(m)T} X_i}) e^{-\widehat{\beta}_{02}^{(m)T} X_i} \right\}^{\delta_{2i}} \\ &\quad \exp \left\{ -\gamma_i [\widehat{H}_{01}^{(m)}(V_i e^{-\widehat{\beta}_{01}^{(m)T} X_i}) - \widehat{H}_{02}^{(m)}(V_i e^{-\widehat{\beta}_{02}^{(m)T} X_i})] \right\} \\ &\quad [\left\{ \gamma_i \widehat{h}_{12}^{(m)} (W_i e^{-\widehat{\beta}_{12}^{(m)T} X_i}) e^{-\widehat{\beta}_{12}^{(m)T} X_i} \right\}^{\delta_{3i}} \\ &\quad \exp \left\{ -\gamma_i [\widehat{H}_{12}^{(m)}(W_i e^{-\widehat{\beta}_{12}^{(m)T} X_i}) - \widehat{H}_{12}^{(m)}(V_i e^{-\widehat{\beta}_{12}^{(m)T} X_i})] \right\}]^{\delta_{1i}}. \end{aligned}$$

Under the gamma frailty model, the respective conditional expectations of γ_i , and $\log \gamma_i$ at m th step are

$$\mathcal{E}_{1i}^{(m)} = E(\gamma_i | \mathcal{O}, \widehat{\Omega}^{(m)}) = \left(D_i + 1/\widehat{\sigma}^{(m)} \right) \left\{ 1/\widehat{\sigma}^{(m)} + \widehat{H}^{(m)}(V_i, W_i, \widehat{\beta}^{(m)}) \right\}^{-1}$$

and

$$\mathcal{E}_{2i}^{(m)} = E(\log \gamma_i | \mathcal{O}, \widehat{\Omega}^{(m)}) = \Psi(D_i + 1/\widehat{\sigma}^{(m)}) - \log \left\{ 1/\widehat{\sigma}^{(m)} + \widehat{H}^{(m)}(V_i, W_i, \widehat{\beta}^{(m)}) \right\},$$

where

$$\widehat{H}^{(m)}(V_i, W_i, \widehat{\beta}^{(m)}) = \sum_{k=1,2} \widehat{H}_{0k}^{(m)}(V_i e^{-\widehat{\beta}_{0k}^{(m)T} X_i}) + \delta_{1i} \left\{ \widehat{H}_{12}^{(m)}(W_i e^{-\widehat{\beta}_{12}^{(m)T} X_i}) - \widehat{H}_{12}^{(m)}(V_i e^{-\widehat{\beta}_{12}^{(m)T} X_i}) \right\},$$

$D_i = \sum_{j=1}^3 \delta_{ji}$, $\Gamma(x)$ is the Gamma function and $\Psi(x) = \Gamma'(x)/\Gamma(x)$ is the digamma function. Therefore, the conditional expectation of the complete-data log-likelihood equals

$$E(l(\sigma) | \mathcal{O}, \widehat{\Omega}^{(m)}) + E(l(\beta_{01}, h_{01}^o) | \mathcal{O}, \widehat{\Omega}^{(m)}) + E(l(\beta_{02}, h_{02}^o) | \mathcal{O}, \widehat{\Omega}^{(m)}) + E(l(\beta_{12}, h_{12}^o) | \mathcal{O}, \widehat{\Omega}^{(m)})$$

where

$$\begin{aligned} E(l(\sigma) | \mathcal{O}, \widehat{\Omega}^{(m)}) &= \frac{1}{n} \sum_{i=1}^n \left(D_i + \frac{1}{\sigma} \right) \mathcal{E}_{2i}^{(m)} - \frac{1}{n\sigma} \sum_{i=1}^n \mathcal{E}_{1i}^{(m)} - \frac{1}{\sigma} \log \sigma - \log \Gamma\left(\frac{1}{\sigma}\right), \\ E(l(\beta_{0k}, h_{0k}^o) | \mathcal{O}, \widehat{\Omega}^{(m)}) &= \frac{1}{n} \sum_{i=1}^n \left[\delta_{ki} \left\{ \log h_{0k}^o(V_i e^{-\beta_{0k}^T X_i}) - \beta_{0k}^T X_i \right\} - \mathcal{E}_{1i}^{(m)} H_{0k}^o(V_i e^{-\beta_{0k}^T X_i}) \right], \quad k = 1, 2, \end{aligned}$$

and

$$\begin{aligned} E(l(\beta_{12}, h_{12}^o) | \mathcal{O}, \widehat{\Omega}^{(m)}) &= \frac{1}{n} \sum_{i=1}^n \left[\delta_{3i} \left\{ \log h_{12}^o(W_i e^{-\beta_{12}^T X_i}) - \beta_{12}^T X_i \right\} \right. \\ &\quad \left. - \delta_{1i} \mathcal{E}_{1i}^{(m)} \left\{ H_{12}^o(W_i e^{-\beta_{12}^T X_i}) - H_{12}^o(V_i e^{-\beta_{12}^T X_i}) \right\} \right]. \end{aligned}$$

S4 Derivation of the Estimators

We start with a simple case of piecewise constant hazard function,

$$\tilde{h}_{jk}^o(t) = \sum_{l=1}^{J_{jk,n}} c_{jk,l} I(t_{jk,l-1} \leq t < t_{jk,l})$$

on $[0, M_{jk}]$, where $0 = t_{jk,0} < t_{jk,1} < \dots < t_{jk,J_{jk,n}} = M_{jk}$ are equally spaced, M_{0k} denotes an upper bound for $V_i e^{-\beta_{jk}^T X_i}$, $k \in \{1, 2\}$, and M_{12} denotes an upper bounds for $W_i e^{-\beta_{12}^T X_i}$. Then, the cumulative piecewise hazard function are

$$\tilde{H}_{jk}^o(t) = \sum_{l=1}^{J_{jk,n}} c_{jk,l} (t - t_{jk,l-1}) I(t_{jk,l-1} \leq t < t_{jk,l}) + (M_{jk}/J_{jk,n}) \sum_{l=1}^{J_{jk,n}} c_{jk,l} I(t \geq t_{jk,l}).$$

In the following, these piecewise functions will be used to acquire an approximated profile-likelihood function of β_{jk} , which converges to a limiting function of β_{jk} , $jk \in \{01, 02, 12\}$, as $n \rightarrow \infty$, $J_{jk,n} \rightarrow \infty$ and $J_{jk,n}/n \rightarrow 0$. The estimators of β_{jk} , $jk \in \{01, 02, 12\}$, are defined as the maximizing values of respective smooth approximations of the limiting functions, with respect to β_{jk} . Given the estimates of the regression coefficients, the proposed smoothing approach also provides estimators of the baseline hazard functions. In the next subsections we provide a detailed derivation of the proposed estimations.

S4.1 Estimating (β_{01}^T, h_{01}^o) and (β_{02}^T, h_{02}^o)

We begin with the estimation method of (β_{01}^T, h_{01}^o) by substituting $\tilde{h}_{01}^o(t)$ and $\tilde{H}_{01}^o(t)$ in $E(l(\beta_{01}, h_{01}^o) | \mathcal{O}, \hat{\Omega}^{(m)})$:

$$\begin{aligned} \tilde{E}\left(l(\beta_{01}, h_{01}^o) | \mathcal{O}, \hat{\Omega}^{(m)}\right) &= \\ &= \frac{1}{n} \sum_{i=1}^n \left\{ \delta_{1i} \left[\log \sum_{l=1}^{J_{01,n}} c_l^{01} I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l}) - \beta_{01}^T X_i \right] \right. \\ &\quad - \mathcal{E}_{1,i}^{(m)} \left[\sum_{l=1}^{J_{01,n}} c_{01,l} (V_i e^{-\beta_{01}^T X_i} - t_{01,l-1}) I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l}) \right. \\ &\quad \left. \left. + \frac{M_{01}}{J_{01,n}} \sum_{l=1}^{J_{01,n}} c_{01,l} I(V_i e^{-\beta_{01}^T X_i} \geq t_{01,l}) \right] \right\} \\ &= -\frac{1}{n} \sum_{i=1}^n \delta_{1i} \beta_{01}^T X_i + \frac{1}{n} \sum_{i=1}^n \delta_{1i} \log \sum_{l=1}^{J_{01,n}} c_{01,l} I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l}) \\ &\quad - \frac{1}{n} \sum_{i=1}^n \mathcal{E}_{1,i}^{(m)} \left[\sum_{l=1}^{J_{01,n}} c_{01,l} (V_i e^{-\beta_{01}^T X_i} - t_{01,l-1}) I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l}) \right. \\ &\quad \left. + \frac{M_{01}}{J_{01,n}} \sum_{l=1}^{J_{01,n}} c_{01,l} I(V_i e^{-\beta_{01}^T X_i} \geq t_{01,l}) \right] \\ &= -\frac{1}{n} \sum_{i=1}^n \delta_{1i} \beta_{01}^T X_i + \frac{1}{n} \sum_{l=1}^{J_{01,n}} \log c_{01,l} \sum_{i=1}^n \delta_{1i} I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l}) \\ &\quad - \frac{1}{n} \sum_{l=1}^{J_{01,n}} c_{01,l} \left[\sum_{i=1}^n \mathcal{E}_{1,i}^{(m)} (V_i e^{-\beta_{01}^T X_i} - t_{01,l-1}) I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l}) \right. \end{aligned}$$

$$+ \frac{M_{01}}{J_{01,n}} \sum_{i=1}^n \mathcal{E}_{1,i}^{(m)} I(V_i e^{-\beta_{01}^T X_i} \geq t_{01,l}) \Big]. \quad (16)$$

For a given value of β_{01} , $c_{01,l}^{(m)}$ that maximizes Eq. (16), $l = 1, 2, \dots, J_{01,n}$, is given by

$$\hat{c}_{01,l}^{(m)} = \frac{\sum_{i=1}^n \delta_{1i} I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l})}{\sum_{i=1}^n \mathcal{E}_{1,i}^{(m)} (V_i e^{-\beta_{01}^T X_i} - t_{01,l-1}) I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l}) + \frac{M_{01}}{J_{01,n}} \sum_{i=1}^n \mathcal{E}_{1,i}^{(m)} I(V_i e^{-\beta_{01}^T X_i} \geq t_{01,l})}. \quad (17)$$

An approximated profile-likelihood function of β_{01} , based on the piecewise hazard, is obtained by plugging (17) into (16),

$$\begin{aligned} l_{01}^{*p}(\beta_{01}) &= \\ &= -\frac{1}{n} \sum_{i=1}^n \delta_{1i} \beta_{01}^T X_i + \frac{1}{n} \sum_{l=1}^{J_{01,n}} \log(\hat{c}_{01,l}) \sum_{i=1}^n \delta_{1i} I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l}) \\ &\quad - \frac{1}{n} \sum_{l=1}^{J_{01,n}} \hat{c}_{01,l} \left[\sum_{i=1}^n \mathcal{E}_{1,i}^{(m)} (V_i e^{-\beta_{01}^T X_i} - t_{01,l-1}) I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l}) \right. \\ &\quad \left. + \frac{M_{01}}{J_{01,n}} \sum_{i=1}^n \mathcal{E}_{1,i}^{(m)} I(V_i e^{-\beta_{01}^T X_i} \geq t_{01,l}) \right] \\ &= -\frac{1}{n} \sum_{i=1}^n \delta_{1i} \beta_{01}^T X_i - \frac{1}{n} \sum_{l=1}^{J_{01,n}} \sum_{i=1}^n \delta_{1i} I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l}) \\ &\quad + \frac{1}{n} \sum_{l=1}^{J_{01,n}} \sum_{i=1}^n \delta_{1i} I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l}) \left[\log \left\{ \sum_{j=1}^n \delta_{1j} I(t_{01,l-1} \leq V_j e^{-\beta_{01}^T X_j} < t_{01,l}) \right\} \right. \\ &\quad \left. - \log \left\{ \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} (V_j e^{-\beta_{01}^T X_j} - t_{01,l-1}) I(t_{01,l-1} \leq V_j e^{-\beta_{01}^T X_j} < t_{01,l}) \right. \right. \\ &\quad \left. \left. + \frac{M_{01}}{J_{01,n}} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} I(V_j e^{-\beta_{01}^T X_j} \geq t_{01,l}) \right\} \right]. \end{aligned}$$

By multiplying the numerator and the denominator by $\frac{J_{01,n}}{n M_{01}}$ we obtain

$$\begin{aligned} l_{01}^{*p}(\beta_{01}) &= \\ &= -\frac{1}{n} \sum_{i=1}^n \delta_{1i} \beta_{01}^T X_i - \frac{1}{n} \sum_{l=1}^{J_{01,n}} \sum_{i=1}^n \delta_{1i} I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l}) \\ &\quad + \frac{1}{n} \sum_{l=1}^{J_{01,n}} \left(\sum_{i=1}^n \delta_{1i} I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l}) \right. \\ &\quad \left. \left[\log \left\{ \frac{J_{01,n}}{n M_{01}} \sum_{j=1}^n \delta_{1j} I(t_{01,l-1} \leq V_j e^{-\beta_{01}^T X_j} < t_{01,l}) \right\} \right. \right. \\ &\quad \left. \left. - \log \left\{ \frac{J_n}{n M_{01}} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} (V_j e^{-\beta_{01}^T X_j} - t_{01,l-1}) I(t_{01,l-1} \leq V_j e^{-\beta_{01}^T X_j} < t_{01,l}) \right. \right. \right. \\ &\quad \left. \left. \left. + \frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} I(V_j e^{-\beta_{01}^T X_j} \geq t_{01,l}) \right\} \right] \right) \\ &= -\frac{1}{n} \sum_{i=1}^n \delta_{1i} \beta_{01}^T X_i - \frac{1}{n} \sum_{l=1}^{J_{01,n}} \sum_{i=1}^n \delta_{1i} I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l}) \end{aligned}$$

$$\begin{aligned}
& + \sum_{l=1}^{J_{01,n}} \left[\frac{1}{n} \sum_{i=1}^n \delta_{1i} I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l}) \log \left\{ \frac{J_{01,n}}{n M_{01}} \sum_{j=1}^n \delta_{1j} I(t_{01,l-1} \leq V_j e^{-\beta_{01}^T X_j} < t_{01,l}) \right\} \right. \\
& \quad - \frac{1}{n} \sum_{i=1}^n \delta_{1i} I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l}) \log \left\{ \frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} I(V_j e^{-\beta_{01}^T X_j} \geq t_{01,l}) \right. \\
& \quad \left. \left. + \frac{J_{01,n}}{n M_{01}} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} (V_j e^{-\beta_{01}^T X_j} - t_{01,l-1}) I(t_{01,l-1} \leq V_j e^{-\beta_{01}^T X_j} < t_{01,l}) \right\} \right].
\end{aligned}$$

The term $-\frac{1}{n} \sum_{l=1}^{J_{01,n}} \sum_{i=1}^n \delta_{1i} I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l})$ equals $-\frac{1}{n} \sum_{i=1}^n \delta_{1i}$, because for subject i , $V_i e^{-\beta_{01}^T X_i}$ must belong to one of the intervals. Therefore, for maximizing $l_{01}^{*p}(\beta_{01})$ with respect to β_{01} it is enough to consider

$$\begin{aligned}
l_{01}^p(\beta_{01}) &= \\
&= -\frac{1}{n} \sum_{i=1}^n \delta_{1i} \beta_{01}^T X_i \\
&+ \sum_{l=1}^{J_{01,n}} \left[\frac{1}{n} \sum_{i=1}^n \delta_{1i} I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l}) \log \left\{ \frac{J_{01,n} n}{n M_{01}} \sum_{j=1}^n \delta_{1j} I(t_{01,l-1} \leq V_j e^{-\beta_{01}^T X_j} < t_{01,l}) \right\} \right. \\
&\quad - \frac{1}{n} \sum_{i=1}^n \delta_{1i} I(t_{01,l-1} \leq V_i e^{-\beta_{01}^T X_i} < t_{01,l}) \log \left\{ \frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} I(V_j e^{-\beta_{01}^T X_j} \geq t_{01,l}) \right. \\
&\quad \left. \left. + \frac{J_{01,n}}{n M_{01}} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} (V_j e^{-\beta_{01}^T X_j} - t_{01,l-1}) I(t_{01,l-1} \leq V_j e^{-\beta_{01}^T X_j} < t_{01,l}) \right\} \right]. \quad (18)
\end{aligned}$$

In the spirit of [Zeng and Lin \(2007\)](#) and [Liu et al. \(2013\)](#), as $n \rightarrow \infty$, $J_{01,n} \rightarrow \infty$ and $J_{01,n}/n \rightarrow 0$, it can be shown that $l_{01}^p(\beta_{01})$ converges uniformly in β_{01} to a limiting function $l_{01}(\beta_{01})$, where

$$l_{01}(\beta_{01}) = E \left[-\delta_1 \beta_{01}^T X + \delta_1 \log \left\{ \frac{\frac{\partial}{\partial t} P(\delta_1 = 1, V \exp\{-\beta_{01}^T X\} \leq t) |_{t=v \exp\{-\beta_{01}^T X\}}}{E[\hat{\gamma} I(V \exp\{-\beta_{01}^T X\} \geq v \exp\{-\beta_{01}^T X\})]} \right\} \right],$$

and $\hat{\gamma} = E(\gamma_1 | \mathcal{O}_1, \hat{\Omega})$.

Moreover, $\frac{\partial}{\partial t} P(\delta_1 = 1, V \exp\{-\beta_{01}^T X\} \leq t)$ can be approximated by

$$\frac{1}{tna_{01}} \sum_{j=1}^n \delta_{1j} K \left(\frac{\log\{V_j\} - \beta_{01}^T X_j - \log t}{a_{01}} \right),$$

and $E[\hat{\gamma} I(V \exp\{-\beta_{01}^T X\} \geq t)]$ can be approximated by

$$\frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \int_{\log t}^{\infty} \frac{1}{a_{01}} K \left(\frac{\log\{V_j\} - \beta_{01}^T X_j - s}{a_{01}} \right) ds,$$

where $K(\cdot)$ is a kernel function and a_{01} is the bandwidth. By setting $x = \frac{\log\{V_j\} - \beta_{01}^T X_j - s}{a_{01}}$ we get

$$\frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \int_{-\infty}^{(\log\{V_j\} - \beta_{01}^T X_j - \log t)/a_{01}} K(x) dx.$$

Thus, the term $E \left[\delta_1 \log \left\{ \frac{\frac{\partial}{\partial t} P(\delta_1 = 1, V \exp\{-\beta_{01}^T X\} \leq t)}{E[\hat{\gamma} I(V \exp\{-\beta_{01}^T X\} \geq v \exp\{-\beta_{01}^T X\})]} \right\} \right]$ can be approximated by

$$\frac{1}{n} \sum_{i=1}^n \delta_{1i} \log \left\{ \frac{\left[V_i \exp\{-\beta_{01}^T X_i\} n a_{01} \right]^{-1} \sum_{j=1}^n \delta_{1j} K \left(\frac{\log\{V_j\} - \beta_{01}^T X_j - (\log\{V_i\} - \beta_{01}^T X_i)}{a_{01}} \right)}{(n)^{-1} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \int_{-\infty}^{[\log\{V_j\} - \beta_{01}^T X_j - (\log\{V_i\} - \beta_{01}^T X_i)]/a_{01}} K(x) dx} \right\}.$$

Based on the approximations above, $l_{01}^p(\beta_{01})$ can be approximated by

$$\begin{aligned} l_{01}^s(\beta_{01}) &= -\frac{1}{n} \sum_{i=1}^n \delta_{1i} \beta_{01}^T X_i \\ &\quad + \frac{1}{n} \sum_{i=1}^n \delta_{1i} \log \left\{ \frac{\left[\exp(\log\{V_i\} - \beta_{01}^T X_i) n a_{01} \right]^{-1} \sum_{j=1}^n \delta_{1j} K \left(\frac{\log\{V_j\} - \beta_{01}^T X_j - \log\{V_i\} + \beta_{01}^T X_i}{a_{01}} \right)}{(n)^{-1} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \int_{-\infty}^{[\log\{V_j\} - \beta_{01}^T X_j - \log\{V_i\} + \beta_{01}^T X_i]/a_{01}} K(x) dx} \right\} \\ &= -\frac{1}{n} \sum_{i=1}^n \delta_{1i} \log\{V_i\} + \frac{1}{n} \sum_{i=1}^n \delta_{1i} \log \left\{ \frac{1}{n a_{01}} \sum_{j=1}^n \delta_{1j} K \left(\frac{\log\{V_j\} - \beta_{01}^T X_j - \log\{V_i\} + \beta_{01}^T X_i}{a_{01}} \right) \right\} \\ &\quad - \frac{1}{n} \sum_{i=1}^n \delta_{1i} \log \left\{ \frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \int_{-\infty}^{[\log\{V_j\} - \beta_{01}^T X_j - \log\{V_i\} + \beta_{01}^T X_i]/a_{01}} K(s) ds \right\}. \end{aligned}$$

Since $K(\cdot)$ is a smooth function, the maximizer of $l_{01}^s(\beta_{01})$, denoted by $\hat{\beta}_{01}$, can be found by gradient-based algorithms. Let $\hat{\beta}_{01}^{(m)}$ denote the maximizer of $l_{01}^s(\beta_{01})$ at step m . Given $\hat{\beta}_{01}^{(m)}$, a smooth estimator of $h_{01}^o(t)$ can be obtained by

$$\hat{h}_{01}^{o(m)}(t; \hat{\beta}_{01}^{(m)}) = \frac{(n a_{01} t)^{-1} \sum_{j=1}^n \delta_{1j} K \left(\frac{\log\{V_j\} - \hat{\beta}_{01}^{T(m)} X_j - \log(t)}{a_{01}} \right)}{(n)^{-1} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \int_{-\infty}^{[\log\{V_j\} - \hat{\beta}_{01}^{T(m)} X_j - \log(t)]/a_{01}} K(u) du}, \quad t > 0,$$

and the estimator of $H_{01}^o(t)$, is obtained by

$$\hat{H}_{01}^{o(m)}(t; \hat{\beta}_{01}^{(m)}) = \int_0^t \hat{h}_{01}^{o(m)}(s; \hat{\beta}_{01}^{(m)}) ds.$$

Similarly, due to the symmetry between transition $0 \rightarrow 1$ and $0 \rightarrow 2$, we get

$$\hat{h}_{02}^{o(m)}(t; \hat{\beta}_{02}^{(m)}) = \frac{(n a_{02} t)^{-1} \sum_{j=1}^n \delta_{2j} K \left(\frac{\log\{V_j\} - \hat{\beta}_{02}^{T(m)} X_j - \log(t)}{a_{02}} \right)}{(n)^{-1} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \int_{-\infty}^{[\log\{V_j\} - \hat{\beta}_{02}^{T(m)} X_j - \log(t)]/a_{02}} K(u) du}, \quad t > 0,$$

where $\hat{\beta}_{02}^{(m)}$ is the maximizer of $l_{02}^s(\beta_{02})$ at step m and

$$\begin{aligned} l_{02}^s(\beta_{02}) &= -\frac{1}{n} \sum_{i=1}^n \delta_{2i} \log\{V_i\} + \frac{1}{n} \sum_{i=1}^n \delta_{2i} \log \left\{ \frac{1}{n a_{02}} \sum_{j=1}^n \delta_{2j} K \left(\frac{\log\{V_j\} - \beta_{02}^T X_j - \log\{V_i\} + \beta_{02}^T X_i}{a_{02}} \right) \right\} \\ &\quad - \frac{1}{n} \sum_{i=1}^n \delta_{2i} \log \left\{ \frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \int_{-\infty}^{[\log\{V_j\} - \beta_{02}^T X_j - \log\{V_i\} + \beta_{02}^T X_i]/a_{02}} K(s) ds \right\}. \end{aligned}$$

S4.2 Estimating (β_{12}^T, h_{12}^o)

In the illness-death setting, the death time after disease diagnosis is truncated by the age at diagnosis. Therefore, the kernel smoothing method described previously should be adapted to accommodate left truncation. To this end, we start with substituting $\tilde{h}_{12}^o(t)$ and $\tilde{H}_{12}^o(t)$ into $E(l(\beta_{12}, h_{12}^o)|\mathcal{O}, \widehat{\Omega}^{(m)})$ and get

$$\begin{aligned}
& \tilde{E}\left(l_n(\beta_{12}, h_{12}^o)|\mathcal{O}, \widehat{\Omega}^{(m)}\right) \\
= & -\frac{1}{n} \sum_{i=1}^n \delta_{3i} \beta_{12}^T X_i + \frac{1}{n} \sum_{i=1}^n \delta_{3i} \log \left\{ \sum_{l=1}^{J_{12,n}} c_{12,l} I(t_{12,l-1} \leq W_i e^{-\beta_{12}^T X_i} < t_{12,l}) \right\} \\
& -\frac{1}{n} \sum_{i=1}^n \delta_{1i} \mathcal{E}_{1,i}^{(m)} \left\{ \sum_{l=1}^{J_{12,n}} c_{12,l} (W_i e^{-\beta_{12}^T X_i} - t_{12,l-1}) I(t_{12,l-1} \leq W_i e^{-\beta_{12}^T X_i} < t_{12,l}) \right. \\
& \quad \left. - \sum_{l=1}^{J_{12,n}} c_{12,l} (V_i e^{-\beta_{12}^T X_i} - t_{12,l-1}) I(t_{12,l-1} \leq V_i e^{-\beta_{12}^T X_i} < t_{12,l}) \right. \\
& \quad \left. + \frac{M_{12}}{J_{12,n}} \sum_{l=1}^{J_{12,n}} c_{12,l} I(W_i e^{-\beta_{12}^T X_i} \geq t_{12,l}) - \frac{M_{12}}{J_{12,n}} \sum_{l=1}^{J_{12,n}} c_{12,l} I(V_i e^{-\beta_{12}^T X_i} \geq t_{12,l}) \right\} \\
= & -\frac{1}{n} \sum_{i=1}^n \delta_{3i} \beta_{12}^T X_i + \frac{1}{n} \sum_{l=1}^{J_{12,n}} \log(c_{12,l}) \sum_{i=1}^n \delta_{3i} I(t_{12,l-1} \leq W_i e^{-\beta_{12}^T X_i} < t_{12,l}) \\
& -\frac{1}{n} \sum_{l=1}^{J_{12,n}} c_{12,l} \left[\sum_{i=1}^n \delta_{1i} \mathcal{E}_{1,i}^{(m)} (W_i e^{-\beta_{12}^T X_i} - t_{12,l-1}) I(t_{12,l-1} \leq W_i e^{-\beta_{12}^T X_i} < t_{12,l}) \right. \\
& \quad \left. - \sum_{i=1}^n \delta_{1i} \mathcal{E}_{1,i}^{(m)} (V_i e^{-\beta_{12}^T X_i} - t_{12,l-1}) I(t_{12,l-1} \leq V_i e^{-\beta_{12}^T X_i} < t_{12,l}) \right. \\
& \quad \left. + \frac{M_{12}}{J_{12,n}} \sum_{i=1}^n \delta_{1i} \mathcal{E}_{1,i}^{(m)} I(W_i e^{-\beta_{12}^T X_i} \geq t_{12,l}) - \frac{M_{12}}{J_{12,n}} \sum_{i=1}^n \delta_{1i} \mathcal{E}_{1,i}^{(m)} I(V_i e^{-\beta_{12}^T X_i} \geq t_{12,l}) \right]. \quad (19)
\end{aligned}$$

For a given value of β_{12} , maximizing Eq. (19) with respect to $c_{12,l}$, $l = 1, 2, \dots, J_n$, gives the following maximizers

$$\begin{aligned}
\hat{c}_{12,l}^{(m)} = & \left\{ \sum_{i=1}^n \delta_{3i} I(t_{12,l-1} \leq W_i e^{-\beta_{12}^T X_i} < t_{12,l}) \right\} \\
& \left[\sum_{i=1}^n \delta_{1i} \mathcal{E}_{1,i}^{(m)} (W_i e^{-\beta_{12}^T X_i} - t_{12,l-1}) I(t_{12,l-1} \leq W_i e^{-\beta_{12}^T X_i} < t_{12,l}) \right. \\
& \quad \left. - \sum_{i=1}^n \delta_{1i} \mathcal{E}_{1,i}^{(m)} (V_i e^{-\beta_{12}^T X_i} - t_{12,l-1}) I(t_{12,l-1} \leq V_i e^{-\beta_{12}^T X_i} < t_{12,l}) \right. \\
& \quad \left. + \frac{M_{12}}{J_{12,n}} \sum_{i=1}^n \delta_{1i} \mathcal{E}_{1,i}^{(m)} I(W_i e^{-\beta_{12}^T X_i} \geq t_{12,l}) - \frac{M_{12}}{J_{12,n}} \sum_{i=1}^n \delta_{1i} \mathcal{E}_{1,i}^{(m)} I(V_i e^{-\beta_{12}^T X_i} \geq t_{12,l}) \right]^{-1}. \quad (20)
\end{aligned}$$

The piecewise-based profile-likelihood function of β_{12} is obtained by plugging Eq. (20) into Eq. (19),

$$\begin{aligned}
l_{12}^{*p}(\beta_{12}) = & -\frac{1}{n} \sum_{i=1}^n \delta_{3i} \beta_{12}^T X_i - \frac{1}{n} \sum_{l=1}^{J_{12,n}} \sum_{i=1}^n \delta_{3i} I(t_{12,l-1} \leq W_i e^{-\beta_{12}^T X_i} < t_{12,l}) \\
& + \frac{1}{n} \sum_{l=1}^{J_{12,n}} \left(\left[\sum_{i=1}^n \delta_{3i} I(t_{12,l-1} \leq W_i e^{-\beta_{12}^T X_i} < t_{12,l}) \right] \log(\hat{c}_l^{12}) \right)
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{n} \sum_{i=1}^n \delta_{3i} \beta_{12}^T X_i - \frac{1}{n} \sum_{i=1}^n \delta_{3i} \\
&\quad + \frac{1}{n} \sum_{l=1}^{J_{12,n}} \left(\sum_{i=1}^n \delta_{3i} I(t_{12,l-1} \leq W_i e^{-\beta_{12}^T X_i} < t_{12,l}) \right. \\
&\quad \left[\log \left\{ \frac{J_{12,n}}{n M_{12}} \sum_{j=1}^n \delta_{3j} I(t_{12,l-1} \leq W_j e^{-\beta_{12}^T X_j} < t_{12,l}) \right\} \right. \\
&\quad \left. - \log \left\{ \frac{J_{12,n}}{n M_g} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \delta_{1j} (W_j e^{-\beta_{12}^T X_j} - t_{12,l-1}) I(t_{12,l-1} \leq W_j e^{-\beta_{12}^T X_j} < t_{12,l}) \right. \right. \\
&\quad \left. - \frac{J_{12,n}}{n M_{12}} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \delta_{1j} (V_j e^{-\beta_{12}^T X_j} - t_{12,l-1}) I(t_{12,l-1} \leq V_j e^{-\beta_{12}^T X_j} < t_{12,l}) \right. \\
&\quad \left. \left. + \frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \delta_{1j} I(W_j e^{-\beta_{12}^T X_j} \geq t_{12,l}) - \frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \delta_{1j} I(V_j e^{-\beta_{12}^T X_j} \geq t_{12,l}) \right\} \right] \right).
\end{aligned}$$

The term $I(W_j e^{-\beta_{12}^T X_j} \geq t_{12,l}) - I(V_j e^{-\beta_{12}^T X_j} \geq t_{12,l})$ consists of the following possible cases. If $V_j e^{-\beta_{12}^T X_j} < W_j e^{-\beta_{12}^T X_j} < t_{12,l}$, then $I(W_j e^{-\beta_{12}^T X_j} \geq t_{12,l}) - I(V_j e^{-\beta_{12}^T X_j} \geq t_{12,l}) = 0 - 0 = 0$. If $t_{12,l} \leq V_j e^{-\beta_{12}^T X_j} < W_j e^{-\beta_{12}^T X_j}$, then $I(W_j e^{-\beta_{12}^T X_j} \geq t_{12,l}) - I(V_j e^{-\beta_{12}^T X_j} \geq t_{12,l}) = 1 - 1 = 0$. If $V_j e^{-\beta_{12}^T X_j} < t_{12,l} \leq W_j e^{-\beta_{12}^T X_j}$, then $I(W_j e^{-\beta_{12}^T X_j} \geq t_{12,l}) - I(V_j e^{-\beta_{12}^T X_j} \geq t_{12,l}) = 1 - 0 = 1$. Thus, we can write

$$I(W_j e^{-\beta_{12}^T X_j} \geq t_{12,l}) - I(V_j e^{-\beta_{12}^T X_j} \geq t_{12,l}) = I(V_j e^{-\beta_{12}^T X_j} < t_{12,l} \leq W_j e^{-\beta_{12}^T X_j})$$

and as a consequence,

$$\begin{aligned}
&\frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \delta_{1j} I(W_j e^{-\beta_{12}^T X_j} \geq t_{12,l}) - \frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \delta_{1j} I(V_j e^{-\beta_{12}^T X_j} \geq t_{12,l}) \\
&= \frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \delta_{1j} I(V_j e^{-\beta_{12}^T X_j} < t_{12,l} \leq W_j e^{-\beta_{12}^T X_j}).
\end{aligned}$$

After plugging in this term into $l_{12}^p(\beta_{12})$, the profile log-likelihood becomes

$$\begin{aligned}
&l_{12}^{*p}(\beta_{12}) \\
&= -\frac{1}{n} \sum_{i=1}^n \delta_{3i} \beta_{12}^T X_i - \frac{1}{n} \sum_{i=1}^n \delta_{3i} \\
&\quad + \frac{1}{n} \sum_{l=1}^{J_{12,n}} \left(\sum_{i=1}^n \delta_{3i} I(t_{12,l-1} \leq W_i e^{-\beta_{12}^T X_i} < t_{12,l}) \right. \\
&\quad \left[\log \left\{ \frac{J_{12,n}}{n M_g} \sum_{j=1}^n \delta_{3j} I(t_{12,l-1} \leq W_j e^{-\beta_{12}^T X_j} < t_{12,l}) \right\} \right. \\
&\quad \left. - \log \left\{ \frac{J_{12,n}}{n M_{12}} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \delta_{1j} (W_j e^{-\beta_{12}^T X_j} - t_{12,l-1}) I(t_{12,l-1} \leq W_j e^{-\beta_{12}^T X_j} < t_{12,l}) \right. \right. \\
&\quad \left. - \frac{J_{12,n}}{n M_{12}} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \delta_{1j} (V_j e^{-\beta_{12}^T X_j} - t_{12,l-1}) I(t_{12,l-1} \leq V_j e^{-\beta_{12}^T X_j} < t_{12,l}) \right. \\
&\quad \left. \left. + \frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \delta_{1j} I(V_j e^{-\beta_{12}^T X_j} < t_{12,l} \leq W_j e^{-\beta_{12}^T X_j}) \right\} \right] \right).
\end{aligned}$$

By dropping the term $-\frac{1}{n} \sum_{i=1}^n \delta_{3i}$, and using similar arguments given earlier, as $n \rightarrow \infty$, $J_{12,n} \rightarrow \infty$ and $J_{12,n}/n \rightarrow 0$, $l_{12}^{*p}(\beta_{12})$ converges uniformly in β_{12} to a limiting function $l_{12}(\beta_{12})$,

$$l_{12}(\beta_{12}) = E \left[-\delta_3 \beta_{12}^T X + \delta_3 \log \left\{ \frac{\frac{\partial}{\partial t} P(\delta_3 = 1, We^{-\beta_{12}^T X} \leq t) \Big|_{t=we^{-\beta_{12}^T X}}}{E[\hat{\gamma} \delta_1 I(We^{-\beta_{12}^T X} < we^{-\beta_{12}^T X} \leq We^{-\beta_{12}^T X})]} \right\} \right].$$

The term $\frac{\partial}{\partial t} P(\delta_3 = 1, We^{-\beta_{12}^T X} \leq t)$ can be approximated by

$$\frac{1}{tna_{12}} \sum_{j=1}^n \delta_{3j} K \left(\frac{\log\{W_j\} - \beta_{12}^T X_j - \log t}{a_{12}} \right),$$

and the term $E \left[\hat{\gamma} \delta_1 I(We^{-\beta_{12}^T X} < t \leq We^{-\beta_{12}^T X}) \right]$ can be approximated by

$$\frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \delta_{1j} \int_{(\log\{V_j\} - \beta_{12}^T X_j - \log t)/a_{12}}^{(\log\{W_j\} - \beta_{12}^T X_j - \log t)/a_{12}} K(x) dx.$$

Clearly, this approximation takes into account the left truncation of the survival time by age at disease diagnosis. Thus, the term

$$E \left[\delta_3 \log \left\{ \frac{\frac{\partial}{\partial t} P(\delta_3 = 1, We^{-\beta_{12}^T X} \leq t) \Big|_{t=we^{-\beta_{12}^T X}}}{E[\hat{\gamma} \delta_1 I(We^{-\beta_{12}^T X} < we^{-\beta_{12}^T X} \leq We^{-\beta_{12}^T X})]} \right\} \right]$$

can be approximated by

$$\frac{1}{n} \sum_{i=1}^n \delta_{3i} \log \left\{ \frac{[W_i \exp\{-\beta_{12}^T X_i\} na_{12}]^{-1} \sum_{j=1}^n \delta_{3j} K \left(\frac{\log\{W_j\} - \beta_{12}^T X_j - \log\{W_i\} + \beta_{12}^T X_i}{a_{12}} \right)}{n^{-1} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \delta_{1j} \int_{(\log\{V_j\} - \beta_{12}^T X_j - \log\{W_i\} + \beta_{12}^T X_i)/a_{12}}^{(\log\{W_j\} - \beta_{12}^T X_j - \log\{W_i\} + \beta_{12}^T X_i)/a_{12}} K(x) dx} \right\}.$$

Combining the approximations above, $l_{12}^p(\beta_{12})$ is approximated by the following smooth function

$$\begin{aligned} l_{12}^s(\beta_{12}) &= -\frac{1}{n} \sum_{i=1}^n \delta_{3i} \beta_{12}^T X_i \\ &\quad + \frac{1}{n} \sum_{i=1}^n \delta_{3i} \log \left\{ \frac{[W_i \exp\{-\beta_{12}^T X_i\} na_{12}]^{-1} \sum_{j=1}^n \delta_{3j} K \left(\frac{\log\{W_j\} - \beta_{12}^T X_j - \log\{W_i\} + \beta_{12}^T X_i}{a_{12}} \right)}{n^{-1} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \delta_{1j} \int_{(\log\{V_j\} - \beta_{12}^T X_j - \log\{W_i\} + \beta_{12}^T X_i)/a_{12}}^{(\log\{W_j\} - \beta_{12}^T X_j - \log\{W_i\} + \beta_{12}^T X_i)/a_{12}} K(x) dx} \right\} \\ &= -\frac{1}{n} \sum_{i=1}^n \delta_{3i} \log\{W_i\} + \frac{1}{n} \sum_{i=1}^n \delta_{3i} \log \left\{ \frac{1}{na_{12}} \sum_{j=1}^n \delta_{3j} K \left(\frac{\log\{W_j\} - \beta_{12}^T X_j - \log\{W_i\} + \beta_{12}^T X_i}{a_{12}} \right) \right\} \\ &\quad - \frac{1}{n} \sum_{i=1}^n \delta_{3i} \log \left\{ \frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \delta_{1j} \int_{(\log\{V_j\} - \beta_{12}^T X_j - \log\{W_i\} + \beta_{12}^T X_i)/a_{12}}^{(\log\{W_j\} - \beta_{12}^T X_j - \log\{W_i\} + \beta_{12}^T X_i)/a_{12}} K(x) dx \right\}. \end{aligned}$$

Given the maximizer of $l_{12}^s(\beta_{12})$ at step m , denoted by $\hat{\beta}_{12}^{(m)}$, $h_{12}^0(t)$ is estimated by

$$\hat{h}_{12}^{o(m)}(t; \hat{\beta}_{12}^{(m)}) = \frac{\frac{1}{na_{12}t} \sum_{j=1}^n \delta_{3j} K \left(\frac{\log\{W_j\} - \hat{\beta}_{12}^{T(m)} X_j - \log(t)}{a_{12}} \right)}{\frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j}^{(m)} \delta_{1j} \int_{(\log\{V_j\} - \hat{\beta}_{12}^{T(m)} X_j - \log(t))/a_{12}}^{(\log\{W_j\} - \hat{\beta}_{12}^{T(m)} X_j - \log(t))/a_{12}} K(u) du}, \quad t > 0.$$

S5 Variance Estimation by weighted bootstrap

Let $G_i, i = 1, \dots, n$, be n independent copies of a positive random variable from a known distribution with mean and variance 1. Let

$$l_n^*(\Omega) = l_n^*(\sigma) + l_n^*(\beta_{01}, h_{01}^o) + l_n^*(\beta_{02}, h_{02}^o) + l_n^*(\beta_{12}, h_{12}^o)$$

be a weighted log-likelihood function of the complete data, where

$$\begin{aligned} l_n^*(\sigma) &= \frac{1}{n} \sum_{i=1}^n D_i \mathcal{E}_{2,i} G_i + \frac{1}{n} \sum_{i=1}^n \mathcal{E}_{3,i} G_i \\ &= \frac{1}{n} \sum_{i=1}^n \left(D_i + \frac{1}{\sigma} - 1 \right) \mathcal{E}_{2,i} G_i - \frac{1}{n\sigma} \sum_{i=1}^n \mathcal{E}_{1,i} G_i + \frac{1}{n} \left[\frac{1}{\sigma} \log \left\{ \frac{1}{\sigma} \right\} - \log \Gamma \left(\frac{1}{\sigma} \right) \right] \sum_{i=1}^n G_i, \\ l_n^*(\beta_{01}, h_{01}^o) &= \frac{1}{n} \sum_{i=1}^n \left\{ \delta_{1i} \left[\log h_{01}^o(V_i e^{-\beta_{01}^T X_i}) - \beta_{01}^T X_i \right] - \mathcal{E}_{1,i} H_{01}^o(V_i e^{-\beta_{01}^T X_i}) \right\} G_i, \\ l_n^*(\beta_{02}, h_{02}^o) &= \frac{1}{n} \sum_{i=1}^n \left\{ \delta_{2i} \left[\log h_{02}^o(V_i e^{-\beta_{02}^T X_i}) - \beta_{02}^T X_i \right] - \mathcal{E}_{1,i} H_{02}^o(V_i e^{-\beta_{02}^T X_i}) \right\} G_i, \\ l_n^*(\beta_{12}, h_{12}^o) &= \frac{1}{n} \sum_{i=1}^n \left\{ \delta_{3i} \left[\log h_{12}^o(W_i e^{-\beta_{12}^T X_i}) - \beta_{12}^T X_i \right] \right. \\ &\quad \left. - \delta_{1i} \mathcal{E}_{1,i} \left[H_{12}^o(W_i e^{-\beta_{12}^T X_i}) - H_{12}^o(V_i e^{-\beta_{12}^T X_i}) \right] \right\} G_i. \end{aligned}$$

Similarly, the weighted versions of $l_{jk}^s(\beta_{jk}), jk \in 01, 02, 12$, are given by

$$\begin{aligned} l_{0k}^{s*}(\beta_{0k}) &= -\frac{1}{n} \sum_{i=1}^n \delta_{ki} V_i G_i + \frac{1}{n} \sum_{i=1}^n \delta_{ki} G_i \log \left\{ \frac{1}{na_{0k}} \sum_{j=1}^n \delta_{kj} G_j K \left(\frac{\log\{V_j\} - \beta_{0k}^T X_j - \log\{V_i\} + \beta_{0k}^T X_i}{a_{0k}} \right) \right\} \\ &\quad - \frac{1}{n} \sum_{i=1}^n \delta_{ki} G_i \log \left\{ \frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j} G_i \int_{-\infty}^{[\log\{V_j\} - \beta_{0k}^T X_j - \log\{V_i\} + \beta_{0k}^T X_i]/a_{0k}} K(s) ds \right\}, \text{ for } k = 1, 2, \end{aligned} \quad (21)$$

and

$$\begin{aligned} l_{12}^{s*}(\beta_{12}) &= -\frac{1}{n} \sum_{i=1}^n \delta_{3i} w_i G_i + \frac{1}{n} \sum_{i=1}^n \delta_{3i} G_i \log \left\{ \frac{1}{na_{12}} \sum_{j=1}^n \delta_{3j} G_j K \left(\frac{\log\{W_j\} - \beta_{12}^T X_j - \log\{W_i\} + \beta_{12}^T X_i}{a_{12}} \right) \right\} \\ &\quad - \frac{1}{n} \sum_{i=1}^n \delta_{3i} G_i \log \left\{ \frac{1}{n} \sum_{j=1}^n \mathcal{E}_{1,j} \delta_{1j} G_i \int_{(\log\{V_j\} - \beta_{12}^T X_j - \log\{W_i\} + \beta_{12}^T X_i)/a_{12}}^{(\log\{W_j\} - \beta_{12}^T X_j - \log\{W_i\} + \beta_{12}^T X_i)/a_{12}} K(x) dx \right\}. \end{aligned} \quad (22)$$

The estimators of each bootstrap sample are derived by maximizing of Eq. (21) and (22).

Given the bootstrap estimates of the regression coefficients, the corresponding bootstrap estimators of the baseline hazard functions are given by

$$\widehat{h}_{0k}^{o*}(t; \widehat{\beta}_{0k}) = \frac{(na_0 t)^{-1} \sum_{j=1}^n \delta_{kj} G_j K \left(\frac{\log\{V_j\} - \widehat{\beta}_{0k}^{*T} X_j - \log(t)}{a_{0k}} \right)}{(n)^{-1} \sum_{j=1}^n \mathcal{E}_{1,j} G_i \int_{-\infty}^{[\log\{V_j\} - \widehat{\beta}_{0k}^{*T} X_j - \log(t)]/a_{0k}} K(u) du}, \quad t > 0, \quad k = 1, 2,$$

and

$$\widehat{h}_{12}^{o*}(t; \widehat{\beta}_{12}) = \frac{(na_{12} t)^{-1} \sum_{j=1}^n \delta_{3j} G_j K \left(\frac{\log\{W_j\} - \widehat{\beta}_{12}^{*T} X_j - \log(t)}{a_{12}} \right)}{(n)^{-1} \sum_{j=1}^n \mathcal{E}_{1,j} \delta_{1j} G_i \int_{(\log\{V_j\} - \widehat{\beta}_{12}^{*T} X_j - \log t)/a_{12}}^{(\log\{W_j\} - \widehat{\beta}_{12}^{*T} X_j - \log t)/a_{12}} K(u) du}, \quad t > 0.$$

S6 Estimation Procedure When Frailty is Not Required

Define $R_i^V(\beta) = \log V_i - \beta^T X_i$, $R_i^W(\beta) = \log W_j - \beta^T X_i$. Then, β_{0k} is estimated by maximization of $l_{0k}^s(\beta_{0k})$, $k = 1, 2$, where

$$\begin{aligned} l_{0k}^s(\beta_{0k}) &= -\frac{1}{n} \sum_{i=1}^n \delta_{ki} \log V_i + \frac{1}{n} \sum_{i=1}^n \delta_{ki} \log \left\{ \frac{1}{na_{0k,n}} \sum_{j=1}^n \delta_{kj} K \left(\frac{R_j^V(\beta_{0k}) - R_i^V(\beta_{0k})}{a_{0k,n}} \right) \right\} \\ &\quad - \frac{1}{n} \sum_{i=1}^n \delta_{ki} \log \left\{ \frac{1}{n} \sum_{j=1}^n \int_{-\infty}^{\{R_j^V(\beta_{0k}) - R_i^V(\beta_{0k})\}/a_{0k}} K(s) ds \right\}, \end{aligned}$$

β_{12} is estimated by maximization of

$$\begin{aligned} l_{12}^s(\beta_{12}) &= -\frac{1}{n} \sum_{i=1}^n \delta_{3i} \log W_i + \frac{1}{n} \sum_{i=1}^n \delta_{3i} \log \left\{ \frac{1}{na_{12,n}} \sum_{j=1}^n \delta_{3j} K \left(\frac{R_j^W(\beta_{12}) - R_i^W(\beta_{12})}{a_{12,n}} \right) \right\} \\ &\quad - \frac{1}{n} \sum_{i=1}^n \delta_{3i} \log \left\{ \frac{1}{n} \sum_{j=1}^n \delta_{1j} \int_{\{R_j^V(\beta_{12}) - R_i^V(\beta_{12})\}/a_{12}}^{\{R_j^W(\beta_{12}) - R_i^W(\beta_{12})\}/a_{12}} K(s) ds \right\}, \end{aligned}$$

and given $\hat{\beta}_{jk}^{(m)}$, $jk \in \{01, 02, 12\}$, the baseline hazard functions are estimated by

$$\begin{aligned} \hat{h}_{0k}^{o(m)}(t) &= \frac{(na_{0k,n}t)^{-1} \sum_{i=1}^n \delta_{ki} K \left(\{R_i^V(\hat{\beta}_{0k}^{(m)}) - \log t\}/a_{0k,n} \right)}{n^{-1} \sum_{i=1}^n \int_{-\infty}^{\{R_i^V(\hat{\beta}_{0k}^{(m)}) - \log t\}/a_{0k,n}} K(s) ds} \quad k = 1, 2, \\ \hat{h}_{12}^{o(m)}(t) &= \frac{(na_{12,n}t)^{-1} \sum_{i=1}^n \delta_{3i} K \left(\{R_i^W(\hat{\beta}_{12}^{(m)}) - \log t\}/a_{12,n} \right)}{n^{-1} \sum_{i=1}^n \delta_{1i} \int_{\{R_j^V(\hat{\beta}_{12}^{(m)}) - \log t\}/a_{12,n}}^{\{R_i^W(\hat{\beta}_{12}^{(m)}) - \log t\}/a_{12,n}} K(s) ds}, \end{aligned}$$

and $\hat{H}_{jk}^{o(m)}(t) = \int_0^t \hat{h}_{jk}^{o(m)}(s) ds$. No recursive algorithm is required.

S7 Simulation Results Without Frailty

To demonstrate the performance of the estimation method without frailty, data were generated similar to Section 4 but with $\mathcal{E}_{1i} \equiv 1$, $i = 1, \dots, 1000$. The regression coefficients and the hazard functions were estimated based on the procedure of Sections S5 above. In this simulation about 17% of the observations were censored at healthy state, and about 19% of those who diagnosed with the disease were censored before death. The results are summarized in Tables S1. As expected, the proposed estimators performs well in terms of bias and coverage rates.

S8 Additional Simulations Results

Table S2 presents simulation results for several baseline hazard functions with different values of ζ in the bandwidths used in the estimation of the regression coefficients.

S9 Marginal Survival Functions under the AFT Model

We begin with the probability $S_{0.}^M(t|X_i) = Pr(T_{1i} > t, T_{2i} > t|X_i)$.

$$S_{0.}^M(t|X_i) = Pr(T_{1i} > t, T_{2i} > t|X_i)$$

$$\begin{aligned}
&= \int_0^\infty Pr(T_{1i} > t, T_{2i} > t | X_i, \gamma_i) f(\gamma_i | X_i) d\gamma_i \\
&= \int_0^\infty Pr(T_{1i} > t, T_{2i} > t | X_i, \gamma_i) f(\gamma_i) d\gamma_i \\
&= \int_0^\infty Pr(T_{1i} > t | X_i, \gamma_i) Pr(T_{2i} > t | X_i, \gamma_i) f(\gamma_i) d\gamma_i \\
&= \int_0^\infty \exp\{-\gamma_i [H_{01}^o(t e^{-\beta_{01}^T X_i}) + H_{02}^o(t e^{-\beta_{02}^T X_i})]\} f(\gamma_i) d\gamma_i \\
&= \left\{1 + \sigma[H_{01}^o(t e^{-\beta_{01}^T X_i}) + H_{02}^o(t e^{-\beta_{02}^T X_i})]\right\}^{-1/\sigma}.
\end{aligned}$$

The marginal survival function $S_{12}^M(t | t_1, X_i) = Pr(T_{2i} > t | T_{1i} = t_1, T_{2i} > t_1, X_i)$ for $t > t_1 > 0$ equals

$$S_{12}^M(t | t_1, X_i) = \int_0^\infty Pr(T_{2i} > t | T_{1i} = t_1, T_{2i} > t_1, X_i, \gamma_i) f(\gamma_i | T_{1i} = t_1, T_{2i} > t_1, X_i) d\gamma_i.$$

For the conditional density of γ_i , we write

$$f(\gamma_i | T_{1i} = t_1, T_{2i} > t_1, X_i) = \frac{\lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(T_{1i} \in [t_1, t_1 + \Delta], T_{2i} > t_1 | X_i, \gamma_i) f(\gamma_i | X_i)}{\lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(T_{1i} \in [t_1, t_1 + \Delta], T_{2i} > t_1 | X_i)}.$$

where $f(\gamma_i | X_i) = f(\gamma_i)$ and under the gamma frailty model

$$\begin{aligned}
&\lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(T_{1i} \in [t_1, t_1 + \Delta], T_{2i} > t_1 | X_i, \gamma_i) = \\
&= \Pr(T_{1i} > t_1, T_{2i} > t_1 | X_i, \gamma_i) \lambda_{01}(t_1 | X_i, \gamma_i) \\
&= \exp\{-\gamma_i [H_{01}^o(t_1 e^{-\beta_{01}^T X_i}) + H_{02}^o(t_1 e^{-\beta_{02}^T X_i})]\} \gamma_i h_{01}^o(t_1 e^{-\beta_{01}^T X_i}) e^{-\beta_{01}^T X_i}.
\end{aligned}$$

Therefore,

$$\begin{aligned}
&\lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(T_{1i} \in [t_1, t_1 + \Delta], T_{2i} > t_1 | X_i) = \\
&= \int_0^\infty \lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(T_{1i} \in [t_1, t_1 + \Delta], T_{2i} > t_1 | X_i, \gamma_i) f(\gamma_i | X_i) d\gamma_i \\
&= \int_0^\infty \exp\{-\gamma_i [H_{01}^o(t_1 e^{-\beta_{01}^T X_i}) + H_{02}^o(t_1 e^{-\beta_{02}^T X_i})]\} \gamma_i h_{01}^o(t_1 e^{-\beta_{01}^T X_i}) e^{-\beta_{01}^T X_i} f(\gamma_i) d\gamma_i \\
&= h_{01}^o(t_1 e^{-\beta_{01}^T X_i}) e^{-\beta_{01}^T X_i} \int_0^\infty \exp\{-\gamma_i [H_{01}^o(t_1 e^{-\beta_{01}^T X_i}) + H_{02}^o(t_1 e^{-\beta_{02}^T X_i})]\} \gamma_i f(\gamma_i) d\gamma_i \\
&= h_{01}^o(t_1 e^{-\beta_{01}^T X_i}) e^{-\beta_{01}^T X_i} (1 + \sigma) \left\{1 + \sigma[H_{01}^o(t_1 e^{-\beta_{01}^T X_i}) + H_{02}^o(t_1 e^{-\beta_{02}^T X_i})]\right\}^{-1/\sigma+1}.
\end{aligned}$$

By combining the above expressions, we get

$$\begin{aligned}
S_{12}^M(t | t_1, X_i) &= Pr(T_{2i} > t | T_{1i} = t_1, T_{2i} > t_1, X_i) = \\
&= \int_0^\infty Pr(T_{2i} > t | T_{1i} = t_1, T_{2i} > t_1, X_i, \gamma_i) f(\gamma_i | T_{1i} = t_1, T_{2i} > t_1, X_i) d\gamma_i \\
&= \int_0^\infty \frac{\exp\left\{-\gamma_i H_{12}^o(t e^{-\beta_{12}^T X_i})\right\}}{\exp\left\{-\gamma_i H_{12}^o(t_1 e^{-\beta_{12}^T X_i})\right\}} \frac{\lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(T_{1i} \in [t_1, t_1 + \Delta], T_{2i} > t_1 | X_i, \gamma_i) f(\gamma_i | X_i)}{\lim_{\Delta \rightarrow 0} \frac{1}{\Delta} \Pr(T_{1i} \in [t_1, t_1 + \Delta], T_{2i} > t_1 | X_i)} d\gamma_i \\
&= \left\{ \frac{1 + \sigma[H_{01}^o(t_1 e^{-\beta_{01}^T X_i}) + H_{02}^o(t_1 e^{-\beta_{02}^T X_i})]}{1 + \sigma[H_{01}^o(t_1 e^{-\beta_{01}^T X_i}) + H_{02}^o(t_1 e^{-\beta_{02}^T X_i})] + H_{12}^o(t e^{-\beta_{12}^T X_i}) - H_{12}^o(t_1 e^{-\beta_{12}^T X_i})} \right\}^{1/\sigma+1}.
\end{aligned}$$

S10 Marginal Survival under the Conditional Cox Model

$$S_{0.}^{M,Cox}(t | X_i) = Pr(T_{1i} > t, T_{2i} > t | X_i)$$

Table S1: Simulation results without frailty

(a) The regression coefficients.

	$\hat{\beta}_{01,1}$	$\hat{\beta}_{01,2}$	$\hat{\beta}_{02,1}$	$\hat{\beta}_{02,2}$	$\hat{\beta}_{12,1}$	$\hat{\beta}_{12,2}$	$\hat{\beta}_{12,3}$
True values	1	0.5	1	1	0.5	0.5	1
mean	1.01	0.50	1.01	1.01	0.49	0.49	1.03
empirical SD	0.06	0.06	0.06	0.05	0.08	0.07	0.07
CR	0.95	0.97	0.93	0.92	0.93	0.96	0.90

(b) The cumulative baseline hazard functions.

t	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	1.00
$H_{01}^o(t)$	0.01	0.04	0.09	0.16	0.25	0.36	0.49	0.64	0.81	1.00
mean	0.01	0.04	0.09	0.16	0.25	0.36	0.49	0.64	0.81	1.00
empirical SD	0.00	0.01	0.01	0.02	0.02	0.03	0.04	0.06	0.08	0.10
CR	0.95	0.96	0.96	0.96	0.95	0.97	0.96	0.95	0.96	0.96
$H_{02}^o(t)$	0.02	0.06	0.14	0.24	0.38	0.54	0.74	0.96	1.22	1.50
mean	0.02	0.06	0.13	0.24	0.37	0.54	0.73	0.96	1.22	1.50
empirical SD	0.00	0.01	0.02	0.02	0.04	0.04	0.06	0.07	0.10	0.14
CR	0.94	0.97	0.95	0.95	0.97	0.94	0.95	0.97	0.96	0.96
$H_{12}^o(t)$	0.01	0.04	0.09	0.16	0.25	0.36	0.49	0.64	0.81	1.00
mean	0.01	0.04	0.09	0.16	0.26	0.37	0.50	0.65	0.83	1.01
empirical SD	0.03	0.04	0.05	0.06	0.06	0.07	0.08	0.09	0.10	0.13
CR	0.88	0.92	0.95	0.96	0.95	0.94	0.94	0.94	0.94	0.95

$$\begin{aligned}
 &= \int_0^\infty \Pr(T_{1i} > t, T_{2i} > t | X_i) f(\gamma_i | X_i, \gamma_i) d\gamma_i \\
 &= \int_0^\infty \exp \left\{ \gamma_i \left[H_{01}^o(t) e^{\beta_{01}^T X_i} + H_{02}^o(t) e^{\beta_{02}^T X_i} \right] \right\} f(\gamma_i) d\gamma_i \\
 &= \left\{ 1 + \sigma \left[H_{01}^o(t) e^{\beta_{01}^T X_i} + H_{02}^o(t) e^{\beta_{02}^T X_i} \right] \right\}^{-1/\sigma}.
 \end{aligned}$$

and

$$\begin{aligned}
 S_{12}^{M,Cox}(t|t_1, X_i) &= Pr(T_{2i} > t | T_{1i} = t_1, T_{2i} > t_1, X_i) \\
 &= \left\{ \frac{1 + \sigma [H_{01}^o(t_1) e^{\beta_{01}^T X_i} + H_{02}^o(t_1) e^{\beta_{02}^T X_i}]}{1 + \sigma [H_{01}^o(t_1) e^{\beta_{01}^T X_i} + H_{02}^o(t_1) e^{\beta_{02}^T X_i} + H_{12}^o(t) e^{\beta_{12}^T X_i} - H_{12}^o(t_1) e^{\beta_{12}^T X_i}]} \right\}^{1/\sigma+1}.
 \end{aligned}$$

S11 Analysis of the Rotterdam Breast Cancer Data Without Frailty

The analysis of the Rotterdam breast cancer data without frailty is presented in Table S3. Additionally, the RSPs of the proposed AFT model and the RSPs of the Cox model are presented in Figures S1-S2.

Table S2: Simulation results with frailty, various bandwidths ζ and various baseline hazard functions.

$h_{01}^o(t) = 2, h_{02}^o(t) = 3, h_{12}^o(t) = 2$										
σ	ζ		$\hat{\sigma}$	$\hat{\beta}_{01,1}$	$\hat{\beta}_{01,2}$	$\hat{\beta}_{02,1}$	$\hat{\beta}_{02,2}$	$\hat{\beta}_{12,1}$	$\hat{\beta}_{12,2}$	$\hat{\beta}_{12,3}$
True values				1.0	0.5	1.0	1.0	0.5	0.5	1.0
2	35	mean	1.98	0.98	0.45	0.97	1.00	0.46	0.41	0.92
		empirical SD	0.34	0.18	0.27	0.28	0.18	0.25	0.30	0.21
		CR	0.96	0.95	0.95	0.96	0.97	0.94	0.90	0.94
50	50	mean	1.91	1.04	0.51	1.03	1.03	0.53	0.51	1.04
		empirical SD	0.33	0.17	0.27	0.26	0.17	0.21	0.25	0.22
		CR	0.94	0.94	0.93	0.96	0.96	0.97	0.92	0.96
75	75	mean	1.88	1.06	0.51	1.05	1.05	0.54	0.54	1.08
		empirical SD	0.30	0.15	0.24	0.23	0.15	0.20	0.22	0.17
		CR	0.93	0.91	0.91	0.95	0.95	0.94	0.90	0.93
100	100	mean	1.81	1.10	0.52	1.08	1.08	0.56	0.54	1.14
		empirical SD	0.30	0.15	0.23	0.23	0.14	0.20	0.23	0.16
		CR	0.91	0.89	0.93	0.95	0.94	0.92	0.93	0.87
$h_{jk}^o(t) = a_{jk}/(1+t), a_{01} = 2, a_{02} = 3, a_{12} = 2$										
2	35	mean	2.05	0.93	0.48	0.96	0.96	0.37	0.40	0.73
		empirical SD	0.40	0.24	0.33	0.29	0.20	0.33	0.42	0.35
		CR	0.98	0.92	0.95	0.95	0.96	0.93	0.94	0.86
50	50	mean	1.95	1.01	0.50	1.01	1.01	0.48	0.51	0.94
		empirical SD	0.40	0.20	0.29	0.27	0.19	0.34	0.40	0.36
		CR	0.96	0.93	0.94	0.96	0.95	0.93	0.95	0.96
75	75	mean	1.92	1.04	0.48	1.01	1.02	0.52	0.50	1.09
		empirical SD	0.37	0.18	0.26	0.24	0.17	0.32	0.38	0.31
		CR	0.95	0.93	0.95	0.97	0.93	0.92	0.93	0.94
100	100	mean	1.90	1.05	0.48	1.02	1.04	0.53	0.49	1.13
		empirical SD	0.36	0.16	0.25	0.24	0.16	0.29	0.36	0.28
		CR	0.96	0.94	0.94	0.97	0.94	0.93	0.93	0.92
$h_{jk}^o(t) = t^{-1}\phi\{\log(t)\}/[1 - \Phi\{\log(t)\}], jk \in \{01, 02, 12\}$										
True values				1.0	0.5	1.0	1.0	0.5	0.5	1.0
2	35	mean	2.02	0.98	0.47	0.95	0.96	0.44	0.38	0.82
		empirical SD	0.35	0.14	0.18	0.19	0.15	0.24	0.25	0.26
		CR	0.96	0.97	0.95	0.95	0.95	0.95	0.91	0.88
50	50	mean	1.90	1.02	0.50	1.00	1.01	0.46	0.46	0.98
		empirical SD	0.32	0.12	0.16	0.18	0.14	0.26	0.25	0.30
		CR	0.94	0.98	0.95	0.96	0.95	0.96	0.95	0.94
75	75	mean	1.86	1.03	0.49	1.01	1.02	0.50	0.52	1.13
		empirical SD	0.33	0.10	0.14	0.16	0.12	0.27	0.27	0.32
		CR	0.90	0.94	0.96	0.96	0.92	0.97	0.94	0.93
100	100	mean	1.84	1.04	0.49	1.01	1.03	0.50	0.50	1.18
		empirical SD	0.32	0.10	0.12	0.15	0.11	0.22	0.25	0.31
		CR	0.92	0.91	0.96	0.95	0.92	0.97	0.96	0.93

Table S3: Rotterdam Tumor Bank Data Analysis: AFT model with our proposed estimation procedure without frailty and Cox illness-death model without frailty. Adjusted p-values in bold are less than 0.05.

	Proposed Model ($\zeta = 65$)					Cox Model				
	Est	exp	SE	P-value	Holm	Est	exp	SE	P-value	Holm
Transition: surgery → relapse										
Age at surgery*	0.13	1.14	0.05	0.016	0.052	-0.16	0.85	0.04	0.000	0.002
log of lymph nodes	-0.41	0.67	0.06	0.000	0.000	0.43	1.54	0.04	0.000	0.000
log of estrogen+1	0.08	1.08	0.03	0.024	0.052	-0.04	0.96	0.02	0.027	0.082
log of progesterone+1	0.08	1.08	0.03	0.007	0.035	-0.02	0.98	0.02	0.206	0.260
Postmenopausal (vs. premenopausal)	-0.30	0.74	0.16	0.059	0.059	0.18	1.19	0.12	0.130	0.260
Tumor size (vs. < 20mm)										
20-50mm	-0.30	0.74	0.10	0.004	0.021	0.21	1.24	0.08	0.006	0.026
> 50mm	-0.48	0.62	0.12	0.000	0.000	0.43	1.54	0.10	0.000	0.000
Hormone therapy	0.56	1.76	0.13	0.000	0.000	-0.42	0.66	0.09	0.000	0.000
Chemotherapy	0.47	1.60	0.10	0.000	0.000	-0.47	0.63	0.09	0.000	0.000
Tumor grade 3 (vs. 2)	-0.25	0.78	0.10	0.013	0.052	0.24	1.27	0.08	0.003	0.016
Transition: surgery → death										
Age at surgery*	-0.98	0.37	0.13	0.000	0.000	1.35	3.88	0.14	0.000	0.000
log of lymph nodes	-0.06	0.94	0.11	0.587	1.000	0.14	1.14	0.12	0.267	1.000
log of estrogen+1	0.01	1.01	0.05	0.919	1.000	-0.05	0.95	0.06	0.436	1.000
log of progesterone+1	-0.02	0.98	0.05	0.731	1.000	0.11	1.11	0.06	0.076	0.685
Postmenopausal (vs. premenopausal)	0.10	1.10	0.46	0.838	1.000	-0.31	0.73	0.63	0.624	1.000
Tumor size (ref. < 20mm)										
20-50mm	0.18	1.20	0.17	0.305	1.000	-0.10	0.91	0.24	0.682	1.000
> 50mm	-0.18	0.83	0.21	0.381	1.000	0.17	1.19	0.30	0.557	1.000
Hormone therapy	0.28	1.32	0.21	0.179	1.000	-0.27	0.76	0.24	0.262	1.000
Chemotherapy	0.06	1.06	0.37	0.881	1.000	-0.20	0.82	0.55	0.714	1.000
Tumor grade 3 (vs. 2)	0.04	1.04	0.23	0.866	1.000	0.01	1.01	0.23	0.978	1.000
Transition: relapse → death										
Age at surgery*	-0.10	0.90	0.06	0.081	0.407	0.12	1.13	0.05	0.017	0.137
Years from surgery to relapse*	1.08	2.94	0.26	0.000	0.000	-1.15	0.32	0.27	0.000	0.000
log of lymph nodes	0.04	1.04	0.06	0.527	0.854	0.08	1.08	0.04	0.091	0.548
log of estrogen+1	0.14	1.16	0.04	0.000	0.002	-0.01	0.99	0.02	0.529	1.000
log of progesterone+1	0.10	1.11	0.04	0.020	0.140	-0.12	0.89	0.02	0.000	0.000
Postmenopausal (vs. premenopausal)	-0.13	0.88	0.17	0.427	0.854	-0.15	0.86	0.14	0.292	0.876
Tumor size (ref. < 20mm)										
20-50mm	-0.24	0.79	0.11	0.029	0.176	0.24	1.27	0.10	0.012	0.111
> 50mm	-0.22	0.80	0.13	0.099	0.407	0.26	1.29	0.12	0.026	0.185
Hormone therapy	0.34	1.40	0.11	0.002	0.016	0.04	1.04	0.10	0.672	1.000
Chemotherapy	-0.17	0.85	0.15	0.267	0.800	0.17	1.18	0.11	0.129	0.644
Tumor grade 3 (vs. 2)	-0.36	0.70	0.08	0.000	0.000	0.12	1.13	0.09	0.200	0.800

*devided by 10




Figure S1: Goodness of fit plots for the illness-death models. Histograms of $\widehat{S}_0^M(V_i|X_i)$ (left of (a), (c)) $\widehat{S}_0^M(V_i, \delta_{1i}, \delta_{2i}, U_{1i})$ (right of (a), (c)), $\widehat{S}_{12}^M(W_i|V_i, X_i)$ (left of (b), (d)) and $\widehat{S}_{12}^M(W_i, V_i, \delta_{3i}, U_{2i})$ (right of (b), (d)). The dashed lines are the expected values under the uniform distribution.




Figure S2: Cox-Snell residuals of each transition, Cox model